ELEMENTARY GROUPS IN $PSL(3, \mathbb{C})$

W. BARRERA, A. CANO, J.P. NAVARRETE, AND J. SEADE

ABSTRACT. In this paper, we give a classification of the subgroups of $PSL(3, \mathbb{C})$ that act on $\mathbb{P}^2_{\mathbb{C}}$ in such a way that their Kulkarni limit set has finitely many lines in general position lines. These are the elementary groups.

;

To Ravi Kulkarni, with great admiration

INTRODUCTION

Kleinian groups are discrete subgroups of $PSL(2, \mathbb{C})$ acting on $\mathbb{S}^2 \cong \mathbb{P}^1_{\mathbb{C}}$ in such a way that their limit set is not all of \mathbb{S}^2 . These are classified into elementary and non-elementary groups. The elementary groups are those whose limit set consists of 0, 1 or 2 points, and they are classified, see for instance (see [20]).

In this work we look at complex Kleinian groups, that is, discrete subgroups of $PSL(3, \mathbb{C})$ acting properly and discontinuously on some open subset of $\mathbb{P}^2_{\mathbb{C}}$. In [11] it is proved that the Kulkarni limit set Λ_{Kul} (see Definition 1.1) of every infinite complex Kleinian group contains at least one complex projective line. In fact we known from [1] that under some generic hypothesis on the group, the Kulkarni limit set is a union of complex projective lines in general position, *i.e.* no three of them intersect. By definition, the exception are the elementary groups. These have finitely many lines in general position in their limit set, and there are two types efficients such groups of the second kind have a finite number of lines in their limit set, but only finitely many in general position.

As an example of elementary groups of the first kind, take the cyclic group generated by a 3×3 diagonal matrix with non-zero eigenvalues of different norms; this has two projective lines as limit set in $\mathbb{P}^2_{\mathbb{C}}$. On the other hand (see [9]), take a \mathbb{C} -Fuchsian group in PSL(3, \mathbb{C}). Then its limit set in $\mathbb{P}^2_{\mathbb{C}}$ consists of a cone of projective lines with base a circle (see [9]; it has infinitely many lines, but all of them pass through the vertex of the cone, so there are only two in general position. We remark that if we take an \mathbb{R} -Fuchsian group in PSL(3, \mathbb{C}), then it is non-elementary and its limit set actually has infinitely many lines in general position, see [9]

In this work we study and describe all the elementary groups in $PSL(3, \mathbb{C})$.

An interesting class of elementary groups are the discrete purely parabolic groups. One finds that the limit set of such a group consists of either one line, or a cone of lines over the circle, or the whole of $\mathbb{P}^2_{\mathbb{C}}$. We describe these in Section 2.

We know (see Theorem 1.5 below) that if $\Gamma \subset PSL(3, \mathbb{C})$ is an infinite discrete subgroup, then the number of complex projective lines in its limit set $\Lambda_{Kul}(\Gamma)$ is

²⁰¹⁰ Mathematics Subject Classification. Primary: 37F30, 32Q45; Secondary 37F45, 22E40. Key words and phrases. Kleinian groups, projective complex plane, discrete groups, limit set.

equal to 1,2,3 or ∞ , and the number of lines $\Lambda_{\text{Kul}}(\Gamma)$ in general position can be 1,2,3,4 or ∞ . Also, if the number of lines in $\Lambda_{\text{Kul}}(\Gamma)$ is exactly 3, then these lines are in general position.

It follows from Theorem 1.5 and the results in this note, that if $\Lambda_{\text{Kul}}(\Gamma)$ has more than one line, then it is a union of lines. The only case where $\Lambda_{\text{Kul}}(\Gamma)$ can have isolated points is when this set actually consists of one line and one point.

Section 1 contains the basic definitions and preliminary results that we use in the sequel. Sections 4 and 7 are expository and we cite in each the appropriate bibliography. The results in these sections are taken from [3] and [2] and we include them here for completeness. In [3] the authors discuss the groups with limit set a line, and in [2] those with four lines in general position.

Section 5 focuses on the case of two lines. These can be of the first kind, with exactly 2 lines in Λ_{Kul} , or of the second kind, with limit set a cone of lines over a circle, but only 2 in general position.

In section 6 we describe the groups with exactly three lines in their limit set, and section 8 discusses the only case where the limit set is not a union of lines. Then Λ_{Kul} consists of a line and a point.

1. Preliminaries

1.1. **Pseudo-projective transformations.** We let $\mathbb{P}^2_{\mathbb{C}} := (\mathbb{C}^3 \setminus \{0\})/\mathbb{C}^*$, be the complex projective plane. This is a compact connected complex 2-dimensional manifold. Let $[]: \mathbb{C}^3 \setminus \{0\} \to \mathbb{P}^2_{\mathbb{C}}$ be the quotient map. If $\beta = \{e_1, e_2, e_3\}$ is the standard basis of \mathbb{C}^3 , we write $[e_j] = e_j$ and if $w = (w_1, w_2, w_3) \in \mathbb{C}^3 \setminus \{0\}$ then we write $[w] = [w_1 : w_2 : w_3]$. Also, $\ell \subset \mathbb{P}^2_{\mathbb{C}}$ is said to be a complex line if $[\ell]^{-1} \cup \{0\}$ is a complex linear subspace of dimension 2. Given $p, q \in \mathbb{P}^2_{\mathbb{C}}$ distinct points, there is a unique complex projective line passing through p and q, such complex projective line is called a *line*, for short, and it is denoted by p, q.

Consider the action of \mathbb{Z}_3 (viewed as the cubic roots of the unity) on $SL(3, \mathbb{C})$ given by the usual scalar multiplication, then the Lie group

$$\mathrm{PSL}(3,\mathbb{C}) = \mathrm{SL}(3,\mathbb{C})/\mathbb{Z}_3$$

is the group of automorphisms of $\mathbb{P}^2_{\mathbb{C}}$. Let $[[\]] : \mathrm{SL}(3,\mathbb{C}) \to \mathrm{PSL}(3,\mathbb{C})$ be the quotient map, $\gamma \in \mathrm{PSL}(3,\mathbb{C})$ and $\tilde{\gamma} \in \mathrm{GL}(3,\mathbb{C})$, we say that $\tilde{\gamma}$ is a lift of γ if there is a cubic root τ of $Det(\gamma)$ such that $[[\tau \tilde{\gamma}]] = \gamma$. Also, we use the notation (γ_{ij}) to denote elements in $\mathrm{SL}(3,\mathbb{C})$.

We denote by $M_{3\times 3}(\mathbb{C})$ the space of all 3×3 matrices with entries in \mathbb{C} equipped with the standard topology. The quotient space

$$\operatorname{SP}(3,\mathbb{C}) := (M_{3\times 3}(\mathbb{C}) \setminus \{\mathbf{0}\})/\mathbb{C}^*$$

is called the space of *pseudo-projective maps of* $\mathbb{P}^2_{\mathbb{C}}$ and it is naturally identified with the projective space $\mathbb{P}^8_{\mathbb{C}}$. Since $\operatorname{GL}(3, \mathbb{C})$ is an open, dense, \mathbb{C}^* -invariant set of $M_{3\times 3}(\mathbb{C}) \setminus \{\mathbf{0}\}$, we obtain that the space of pseudo-projective maps of $\mathbb{P}^2_{\mathbb{C}}$ is a compactification of $\operatorname{PSL}(3, \mathbb{C})$. As in the case of projective maps, if $\mathbf{s} \in M_{3\times 3}(\mathbb{C}) \setminus \{\mathbf{0}\}$, then $[\mathbf{s}]$ denotes the equivalence class of the matrix \mathbf{s} in the space of pseudoprojective maps of $\mathbb{P}^2_{\mathbb{C}}$. Also, we say that $\mathbf{s} \in M_{3\times 3}(\mathbb{C}) \setminus \{\mathbf{0}\}$ is a lift of the pseudoprojective map S, whenever $[\mathbf{s}] = S$.

Let S be an element in $(M_{3\times 3}(\mathbb{C}) \setminus \{\mathbf{0}\})/\mathbb{C}^*$ and **s** a lift to $M_{3\times 3}(\mathbb{C}) \setminus \{\mathbf{0}\}$ of S. The matrix **s** induces a non-zero linear transformation $s : \mathbb{C}^3 \to \mathbb{C}^3$, which is not necessarily invertible. Let $Ker(s) \subsetneq \mathbb{C}^3$ be its kernel and let Ker(S) denote its projectivization to $\mathbb{P}^2_{\mathbb{C}}$, taking into account that $Ker(S) := \emptyset$ whenever $Ker(s) = \{(0,0,0)\}$. We refer to [10] for more details about this subject.

1.2. The limit set. There are two types of limit sets relevant for this work. These are the Kulkarni limit set Λ_{Kul} and the Myrberg limit set Λ_{Myr} . Let us define these. Let $\Gamma \subset \text{PSL}(n+1,\mathbb{C})$ be a discrete subgroup.

Definition 1.1. The *Kulkarni limit set* of Γ is:

$$\Lambda(\Gamma) = L_0(\Gamma) \cup L_1(\Gamma) \cup L_2(\Gamma),$$

where:

- $L_0(\Gamma)$ is the closure of the points in $\mathbb{P}^2_{\mathbb{C}}$ with infinite isotropy group.
- $L_1(\Gamma)$ is the closure of the set of accumulation points of the Γ -orbits of points in $\mathbb{P}^2_{\mathbb{C}} \setminus L_0(\Gamma)$.
- $L_2(\Gamma)$ is the closure of the union of accumulation points of $\{\gamma(K) : \gamma \in \Gamma\}$, where K runs over all the compact sets in $\mathbb{P}^2_{\mathbb{C}} \setminus (L_0(\Gamma) \cup L_1(\Gamma))$.

This is a closed Γ -invariant set. Its complement

$$\Omega_{\mathrm{Kul}}(\Gamma) = \mathbb{P}^n_{\mathbb{C}} \setminus \Lambda(\Gamma),$$

is the Kulkarni discontinuity region of Γ . We know from [18] that the Γ -action on $\Omega_{\text{Kul}}(\Gamma)$ is properly discontinuous, and we further know from [1] that $\Omega_{\text{Kul}}(\Gamma)$ contains the equicontinuity region Eq(Γ). However, as we will see later, $\Omega_{\text{Kul}}(\Gamma)$ generically is the largest open subset of $\mathbb{P}^2_{\mathbb{C}}$ where the group acts properly and discontinuously, but this is not always so.

Recall that $Eq(\Gamma)$ is the set of points in $\mathbb{P}^2_{\mathbb{C}}$ for which there is an open neighborhood where the family of transformations defined by Γ is a normal family.

Now, given $\Gamma \subset PSL(n + 1, \mathbb{C})$, a discrete subgroup, we let Γ' be the set of pseudo-projective maps of $\mathbb{P}^2_{\mathbb{C}}$ which are limits of sequences in Γ . That is:

 $\Gamma' := \{ S \in SP(3, \mathbb{C}) \mid S \text{ is an accumulation point of } \Gamma \}.$

The following notion is due to Myrberg [21] (cf. [1, Definition 3.3]):

Definition 1.2. The Myrberg limit set of Γ is:

$$\Lambda_{\mathrm{Mvr}}(\Gamma) := \bigcup_{S \in \Gamma'} Ker(S) \; ,$$

where Ker(S) is the kernel of the pseudoprojective transformation S defined above.

Notice that if a pseudoprojective map S is not in $PSL(3, \mathbb{C})$, then its kernel is either a line or a point. The following useful notion is introduced in [4]:

Definition 1.3. We say that a line $\ell \subset \mathbb{P}^2_{\mathbb{C}}$ is an effective line for Γ if there exists a pseudoprojective transformation $S \in \Gamma'$ with kernel ℓ .

The Myrberg limit set of Γ contains all the effective lines, and it is not hard to see that, by [22], it contains the Kulkarni limit set. It is immediate from [21] that Γ acts properly and discontinuously on the complement of $\Lambda_{Myr}(\Gamma)$. Furthermore, one has the following theorem.

Theorem 1.4. The Myrberg limit set is the complement of the equicontinuity region: $Eq(\Gamma) = \mathbb{P}^2_{\mathbb{C}} \setminus \Lambda_{Myr}(\Gamma)$. Also, the set $\Lambda_{Myr}(\Gamma)$ equals the union of all effective lines of Γ except when it is the disjoint union of one line and one point. Moreover, $\Lambda_{Kul}(\Gamma) \subset \Lambda_{Myr}(\Gamma)$. The first statement is a special case of [10, Lemma 4.1] (see also [4, Theorem 3.4]). The second statement is Corollary 4.5 in [4].

Throughout this work, when we say limit set we refer to the Kulkarni limit set, otherwise we will say it explicitly.

1.3. Classification of the elements in $PSL(3, \mathbb{C})$. Recall that the elements of $PSL(2, \mathbb{C})$ are classified as elliptic, parabolic or loxodromic: g is elliptic if regarded as a Möbius transformation in $S^2 \cong \mathbb{P}^1_{\mathbb{C}}$, up to conjugation it is a rotation; parabolic elements are translations up to conjugation and loxodromic elements are multiplication by a complex number of norm $\neq 1$. Equivalently, given $g \in PSL(2, \mathbb{C})$ and a lift \tilde{g} to $SL(2, \mathbb{C})$, g is elliptic if \tilde{g} is diagonalizable with unitary eigenvalues, it is parabolic if \tilde{g} is not diagonalizable and it has unitary eigenvalues, and g is loxodromic otherwise. The classification given in these terms extends to all dimensions (see [22, 8]).

The *elliptic* elements in $PSL(3, \mathbb{C})$ are those elements γ that have a lift to $SL(3, \mathbb{C})$ whose Jordan canonical form is

$$\left(\begin{array}{ccc} e^{i\theta_1} & 0 & 0 \\ 0 & e^{i\theta_2} & 0 \\ 0 & 0 & e^{i\theta_3} \end{array}\right).$$

The limit set for (the cyclic group generated by) γ elliptic is either empty or all of $\mathbb{P}^2_{\mathbb{C}}$, according to whether the order of γ is finite or infinite. The subgroups of $\mathrm{PSL}(3,\mathbb{C})$ containing an elliptic element of infinite order cannot be discrete.

The *parabolic* elements in $PSL(3, \mathbb{C})$ are the elements γ with limit set $\Lambda_{Kul}(\gamma)$ equal to one single complex line. If γ is parabolic then it has a lift to $SL(3, \mathbb{C})$ with Jordan canonical form one of the following matrices:

$$\left(\begin{array}{rrrr}1 & 1 & 0\\ 0 & 1 & 0\\ 0 & 0 & 1\end{array}\right), \left(\begin{array}{rrrr}1 & 1 & 0\\ 0 & 1 & 1\\ 0 & 0 & 1\end{array}\right), \left(\begin{array}{rrr}e^{2\pi i t} & 1 & 0\\ 0 & e^{2\pi i t} & 0\\ 0 & 0 & e^{-4\pi i t}\end{array}\right), e^{2\pi i t} \neq 1.$$

In the first case $\Lambda_{\text{Kul}}(\gamma)$ is the complex line consisting of all the fixed points of γ , in the second case $\Lambda_{\text{Kul}}(\gamma)$ is the unique γ -invariant complex line. In the last case $\Lambda_{\text{Kul}}(\gamma)$ is the complex line determined by the two fixed points of γ .

There are four kinds of *loxodromic* elements in $PSL(3, \mathbb{C})$:

• The *complex homotheties* are the elements that have a lift to SL(3, C) with Jordan canonical form:

$$\left(\begin{array}{ccc} \lambda & 0 & 0\\ 0 & \lambda & 0\\ 0 & 0 & \lambda^{-2} \end{array}\right), \quad |\lambda| \neq 1.$$

The limit set $\Lambda_{\text{Kul}}(\gamma)$ is the set of fixed points of γ , consisting of a complex line and a point.

• The screws are those elements $\gamma \in PSL(3, \mathbb{C})$ that have a lift to $SL(3, \mathbb{C})$ whose Jordan canonical form is

$$\left(\begin{array}{ccc}\lambda & 0 & 0\\ 0 & \mu & 0\\ 0 & 0 & (\lambda\mu)^{-1}\end{array}\right), \quad \lambda \neq \mu, \, |\lambda| = |\mu| \neq 1.$$

The limit set consists of a complex line l, on which γ acts as an elliptic transformation of $PSL(2, \mathbb{C})$, and the fixed point of γ not lying in l.

• The *loxoparabolic* elements $\gamma \in PSL(3, \mathbb{C})$ have a lift to $SL(3, \mathbb{C})$ whose Jordan canonical form is

$$\left(\begin{array}{ccc} \lambda & 1 & 0\\ 0 & \lambda & 0\\ 0 & 0 & \lambda^{-2} \end{array}\right), \quad |\lambda| \neq 1.$$

The limit set $\Lambda_{\text{Kul}}(\gamma)$ consists of two γ -invariant complex lines. The element γ acts on one of these complex lines as a parabolic element of $\text{PSL}(2,\mathbb{C})$ and on the other as a loxodromic element of $\text{PSL}(2,\mathbb{C})$.

• The strongly loxodromic elements $\gamma \in PSL(3, \mathbb{C})$ have a lift to $SL(3, \mathbb{C})$ whose Jordan canonical form is

$$\left(\begin{array}{ccc} \lambda_1 & 0 & 0\\ 0 & \lambda_2 & 0\\ 0 & 0 & \lambda_3 \end{array}\right), \quad |\lambda_1| < |\lambda_2| < |\lambda_3|.$$

This kind of transformation has three fixed points, one of them is attracting, another is repelling and the other point is a saddle. The limit set is the union of the complex line determined by the attracting and saddle points and the complex line determined by the saddle and repelling points.

1.4. Elementary groups in $PSL(3, \mathbb{C})$. In the previous section we have examples of elements in $PSL(3, \mathbb{C})$ where the limit set consists of one line, or one line and one point, or two lines. We have the following theorem from [4].

Theorem 1.5. Let $\Gamma \subset PSL(3, \mathbb{C})$ be an infinite discrete subgroup. Then:

- (i) The number of complex projective lines in $\Lambda_{\text{Kul}}(\Gamma)$ is equal to 1,2,3 or ∞ .
- (ii) The number of lines $\Lambda_{\text{Kul}}(\Gamma)$ in general position can be 1, 2, 3, 4 or ∞ .
- (iii) If the number of lines in Λ_{Kul}(Γ) is exactly 3, then these lines are in general position.
- (iv) If there are infinitely many lines in $\Lambda_{\text{Kul}}(\Gamma)$, then the effective lines form a perfect set in $\Lambda_{\text{Kul}}(\Gamma)$.
- (v) There can be at most one isolated point in $\Lambda_{\text{Kul}}(\Gamma)$, and in that case this limit set is the disjoint union of that point and one line.

Statements (i), (ii) and (iv) are theorems 1.1 and 1.2 in [4]. Statement (iii) is a corollary of [4, Proposition 5.4]. Statement (v) is not in the literature so we now give a short proof of it.

Recall that a discrete subgroup of $PSL(2, \mathbb{C})$ is elementary if its limit set has finite cardinality. We remark that in the case of $PSL(3, \mathbb{C})$ there are groups with infinitely many lines in their limit set Λ_{Kul} , but only two of them in general position. In view of the previous theorem, this leads to the following definition:

Definition 1.6. A discrete subgroup of $PSL(3, \mathbb{C})$ is elementary of the first kind if its limit set Λ_{Kul} has finitely many lines. The group is elementary of the second kind if Λ_{Kul} has finitely many lines in general position.

So every elementary group of the first kind also is of the second kind, but not conversely. In the following sections we describe the classification of elementary groups. 1.5. The control group. We refer to [8] for a discussion about this section. Consider $\Gamma \subset PSL(3, \mathbb{C})$ a (discrete or not) subgroup which acts on $\mathbb{P}^2_{\mathbb{C}}$ with a point p which is fixed by all of Γ . Choose an arbitrary line ℓ in $\mathbb{P}^2_{\mathbb{C}} \setminus \{p\}$, and notice we have a canonical projection:

$$\pi = \pi_{p,\ell} : \mathbb{P}^2_{\mathbb{C}} \setminus \{p\} \longrightarrow \ell \,,$$

given by $\pi(x) = \overleftarrow{x, p} \cap \ell$. It is clear that this map is holomorphic and it allows us to define a group homomorphism:

$$\Pi = \Pi_{p,\ell} : \Gamma \longrightarrow Bihol(\ell) \cong PSL(2,\mathbb{C}),$$

by $\Pi(g)(x) = \pi(g(x))$. If we choose another line, say ℓ' , one gets similarly a projection $\pi' = \pi_{p,\ell'} : \mathbb{P}^2_{\mathbb{C}} \setminus \{p\} \to \ell'$, and a group homomorphism $\Pi' = \Pi_{p,\ell'} : \Gamma \to \mathrm{PSL}(2,\mathbb{C})$. It is an exercise to see that Π and Π' are equivalent in the sense that there is a biholomorphism $h : \ell \to \ell'$ inducing an automorphism H of $\mathrm{PSL}(2,\mathbb{C})$ such that $H \circ \Pi = \Pi'$. As before, the line ℓ is called *the horizon*.

This leads to the following definition:

Definition 1.7. Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group as above. We call $\Pi = \Pi_{p,\ell}$ the control morphism (or map) and its image $\Pi(\Gamma) \subset PSL(2, \mathbb{C})$, is the *control group*. These are well-defined and independent of ℓ up to an automorphism of $PSL(2, \mathbb{C})$.

The control map and the control group allow us to get information about the dynamics of Γ by looking at a subgroup of $PSL(2, \mathbb{C})$, which is far easier to handle. The prize we pay is that the control group in $PSL(2, \mathbb{C})$ may not be discrete.

2. Purely parabolic groups

We now follow [5] and look at the discrete subgroups in $PSL(3, \mathbb{C})$ that, besides the identity, have only parabolic elements. These are called purely parabolic and there are five families of such groups; three of them split into various subfamilies according to their limit set (and their control group, see [8]). All of these are elementary.

The simplest purely parabolic groups are cyclic, generated by a parabolic element. As described above, there are three types of such elements in $PSL(3, \mathbb{C})$, described by the Jordan normal form of their lifts to $SL(3, \mathbb{C})$. Each of these belongs to a different type of the families we describe below. The first type generates torus groups (see definitions below), the second generates Abelian Kodaira groups and the ellipto-parabolic elements generate elliptic groups.

(i) Elliptic groups. These are the only purely parabolic groups that are not conjugate to subgroups of the Heisenberg group Heis(3, ℂ) and they are subgroups of fundamental groups of elliptic surfaces. These have limit set a single line. Up to conjugation these groups are of the form:

Ell(W,
$$\mu$$
) = $\left\{ \begin{bmatrix} \mu(w) & \mu(w)w & 0\\ 0 & \mu(w) & 0\\ 0 & 0 & \mu(w)^{-2} \end{bmatrix} : w \in W \right\},$

where $W \subset \mathbb{C}$ is an additive discrete subgroup and $\mu: W \to \mathbb{S}^1$ is a group morphism.

(ii) Torus groups. These are subgroups of fundamental groups of complex tori. These are of the form:

$$\mathcal{T}(\mathfrak{L}) = \left\{ \begin{bmatrix} 1 & 0 & a \\ 0 & 1 & b \\ 0 & 0 & 1 \end{bmatrix} : (a,b) \in \mathfrak{L} \right\} ,$$

where \mathfrak{L} is an additive discrete subgroup of \mathbb{C}^2 . These groups also have a single line as limit set, so they are elementary of the first kind.

(iii) Dual torus groups,

$$\mathcal{T}^*(\mathfrak{L}) = \left\{ \left[\begin{array}{rrr} 1 & a & b \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right] : (a,b) \in \mathfrak{L} \right\} \ .$$

These split into three types: the first have Kulkarni limit set a complex projective line, so these are elementary of the first kind. The second type have limit set a cone of projective lines over a circle, so these are of the second kind. The third type have all $\mathbb{P}^2_{\mathbb{C}}$ as limit set and they are non-elementary.

(iv) Inoue groups and their extensions. Inoue groups are proper subgroups of fundamental groups of Inoue surfaces. To define these, let $\mathfrak{L} \subset \mathbb{C}^2$ be an additive discrete subgroup and consider a dual torus group.

$$\mathcal{I} = \mathcal{I}(u, v) := \left\langle \begin{bmatrix} 1 & u & v \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} , (u, v) \in \mathfrak{L} \right\rangle .$$

In oue groups are obtained by taking a generator ${\cal I}$ as above and adding to it a generator of the form

$$\gamma_1 = \gamma_1(x, y, z) := \begin{bmatrix} 1 & x + z & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{bmatrix} \ x, y, z \in \mathbb{C} \ .$$

The limit set is a cone of lines over a circle, so these are elementary of the second kind.

Then one has the extended Inoue group, which are purely parabolic as well, with limit set all of $\mathbb{P}^2_{\mathbb{C}}$, so they are non-elementary and do not fall within the scope of this article. We refer to [5] for details.

(v) Finally one has the Kodaira groups Kod_0 , which are Abelian, and their extensions. A Kodaira group is a discrete group in $\text{PSL}(3, \mathbb{C})$ such that each element h in the group can be written as:

$$\left[\begin{array}{rrrr} 1 & a & b \\ 0 & 1 & a \\ 0 & 0 & 1 \end{array}\right] \,.$$

One can show that these are extensions of dual torus groups. Their limit set is a line, so they are elementary of the first kind.

There are five types of extensions $\widetilde{\mathcal{K}_i}$, $i = 1, \dots, 5$, which are purely parabolic and discrete. These are all non-Abelian and they split into five types according to their limit set and the control group. The first type $\widetilde{\mathcal{K}_1}$ have limit set a projective line, so they are elementary of the first kind; the second type $\widetilde{\mathcal{K}_2}$ have limit set a cone of projective lines over the circle,

so they are of the second kind, while the remaining three types have limit set all of $\mathbb{P}^2_{\mathbb{C}}$ and they are non-elementary. We refer to [5] for details.

3. Solvable groups

An essential step towards understanding elementary groups in $PSL(3, \mathbb{C})$ is studying the dynamics of solvable groups in $PSL(3,\mathbb{C})$. Unlike the classical case of $PSL(2, \mathbb{C})$, now there is great richness.

Definition 3.1. A group H is called virtually solvable if it contains a finite index subgroups G and subgroups G_i , i = 0, ..., k, such that:

- $e = G_0 \subset G_1 \subset \ldots \subset G_k = G$, and G_{j-1} is normal in G_j , and G_j/G_{j-1} is an abelian group, for $j = 1, \ldots, k$.

For instance, for a subgroup of $PSL(2, \mathbb{C})$, virtually solvable is equivalent to saying that the group is elementary, *i.e.* its limits set has finite cardinality, see[11] and [20], or in an equivalent way (via Tit's dichotomy) a Möbius subgroup is nonvirtually solvable if it contains either a non-Ciclyc Schottky group or a purely elliptic free group whose rank is at least two, see [7, 26] for a general discussion or [16, 20] for standard arguments in the one-dimensional case.

One has the following theorem from [27]:

Theorem 3.2. For subgroups of $PSL(3, \mathbb{C})$, solvable is equivalent to having a finite index subgroup which is conjugate to group of upper triangulable matrices.

So the following are examples of solvable groups:

- (i) All purely parabolic groups.
- (ii) Hyperbolic Toral groups (we discuss these later, in section 7).
- (iii) Fundamental groups of Inoue surfaces, see [2, 5, 11].

So, after our previous section 2, to have the complete picture of solvable groups we must look at solvable groups with loxodromic elements. This is done in [27,28]. In those articles the author determined the corresponding limit set and the representations of solvable groups in $PSL(3, \mathbb{C})$ containing loxodromic element. The main results can be summarized as follows. These summarize Theorems 1.1 to 1.4 in [28] and theorems 1 and 2 in [27]:

Theorem 3.3. The solvable groups in $PSL(3, \mathbb{C})$ can be of three main types:

- (i) Commutative.
- (ii) Cyclic extensions of torus groups by a strongly loxodromic element.
- (iii) Dual torus groups extensions.

Theorem 3.4. If a solvable group Γ is commutative, then its limit set consists of either two lines, three lines in general position, or it is a line and a point, and Γ can be either triangular or a fake Hopf group:

• Diagonal groups:

$$\{diag(\alpha^n, \alpha^m, 1) : n, m \in \mathbb{Z}\}\$$

here $\alpha, \beta \in \mathbb{C}^*$, $|\alpha| \neq 1$ or $|\beta| \neq 1$.

• Fake Hopf:

$$\left\{ \begin{bmatrix} \mu(w) & w\mu(w) & 0 \\ 0 & \mu(w) & 0 \\ 0 & 0 & \mu(w)^{-2} \end{bmatrix} \right\}$$

where $W \subset \mathbb{C}$ is a discrete group and $\mu : W \to \mathbb{C}^*$ is a group morphism, satisfying some technical conditions, see [28].

Theorem 3.5. If the group is an extension of either a torus group or a dual torus group, then its limit set can be a single projective line, or two lines, or else it has infinitely many lines but either it is a cone of lines over a circle or it has four lines in general position. And one has:

- (i) If the group Γ is a cyclic extension of a torus group, then it is a semidirect product of a torus group and the cyclic group generated by a strongly loxodromic element.
- (ii) If the group is an extension of a dual torus groups, then it can be:
 - An extension by a loxodromic element (of any type); or
 - An extension by two loxo-parabolic elements, and in this case the group is a semi-direct product of the dual torus group and the group generated by the two loxodromic elements.

In the sequel we describe all groups with limit set as stated in this theorem.

4. GROUPS WITH LIMIT SET EXACTLY ONE LINE

In the previous section we described several purely parabolic groups with limit set a single line. Here we follow [3] and describe all the discrete groups in $PSL(3, \mathbb{C})$ with limit set a line. In particular, every such group is virtually nilpotent.

Let Γ be complex Kleinian with limit set a line ℓ . Then Γ acts properly and discontinuously on the complement of ℓ . From the above described classification of the elements in PSL(3, \mathbb{C}) we known that the elements in Γ are all either elliptic, parabolic or loxoparabolic. If the group contains a loxoparabolic element, then one finds that Γ is conjugate to a group such that every element can be represented by a matrix of the form:

$$\left(\begin{array}{rrr} a & 0 & v \\ 0 & d & 0 \\ 0 & 0 & 1 \end{array}\right) \,,$$

where ℓ becomes the line $\overleftarrow{e_1, e_2}$. Moreover, Γ acts as an Euclidean group on the line $\overleftarrow{e_1, e_3}$. By well-known facts on Euclidean groups (see [20]), *a* is a root of unity of order 1, 2, 3, 4 or 6, and the rank of Γ considered as acting on $\overleftarrow{e_1, e_3}$ is equal to 1 or 2. One can show that *a* cannot be a root of unity of order 2, 3, 4 nor 6, otherwise Γ would contain a complex homothety. Finally, the rank of Γ considered as acting on $\overleftarrow{e_1, e_3}$ is not equal to 1, otherwise, the Kulkarni limit set of Γ would be equal to two lines.

On the other hand, if the group does not contain any loxoparabolic elements, then there are two cases. If it acts on ℓ without parabolic elements then Γ can be considered as a discrete group of Euclidean isometries of \mathbb{R}^4 . If some element acts on ℓ as parabolic element then the group can be identified with a group of triangular matrices. Then one can rule out the existence of irrational ellipto-parabolic elements and show that there exists a unipotent subgroup of finite index.

In this way we arrive to the following [3, Theorem 1.1]:

Theorem 4.1. Let Γ be a subgroup of $PSL(3, \mathbb{C})$ such that its Kulkarni limit consists of precisely one complex projective line ℓ . Then:

- (i) If Γ does not contain loxoparabolic elements nor an element which acts as a parabolic element on the line l, then Γ is a group of isometries of C² and it contains a free abelian normal subgroup of finite index and rank ≤ 4.
- (ii) If Γ does not contain loxoparabolic elements but it does contain an element which acts as a parabolic element on l, then Γ does not contain any irrational ellipto-parabolic elements and it is a finite extension of a unipotent subgroup that consists of unipotent parabolic maps. Hence it is a finite extension of a group of the form Z, Z², Z³, Z⁴, Δ_k or G_k, where

$$\Delta_k = \langle A, B, C, D : C, D \text{ are central and } [A, B] = C^k \rangle, k \in \mathbb{N},$$

 $G_k = \langle A, B, C : C \text{ is central and } [A, B] = C^k \rangle, k \in \mathbb{N},$

(iii) If Γ does contain a loxoparabolic element, then Γ is isomorphic to the group $\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}_{n_0}$, where $n_0 \in \mathbb{N}$ is arbitrary. The \mathbb{Z}_{n_0} summand is a group of complex reflections, while $\mathbb{Z} \oplus \mathbb{Z}$ is generated by a loxoparabolic element and another element which can be loxoparabolic or parabolic.

5. Groups with limit set two lines

We must distinguish two cases:

a) Groups that have exactly two lines in their limit set. These are elementary of the first kind; and

b) Groups with more than two lines in the limit set, but only two in general position. In this case, we know from Theorem 1.5 that the total number of lines actually is infinite. These are elementary of the second kind.

For instance, cyclic groups generated by a strongly loxodromic element have exactly two lines in their limit set. On the other hand, for instance, the dual torus groups of the second type, described before, have limit set a cone of projective lines over a circle. In this case, there are infinitely many lines in the limit set, but only two in general position.

Let us describe these elementary groups. As in [4], for every open Γ -invariant set U we denote by $\lambda(U)$ the maximum number of complex projective lines contained in $\mathbb{P}^2_{\mathbb{C}} \setminus U$, and by $\mu(U)$ the maximum number of such lines in general position. If U is the Kulkarni discontinuity region $\Omega_{\text{Kul}}(\Gamma)$, then its complement $\mathbb{P}^2_{\mathbb{C}} \setminus U$ is the limit set $\Lambda_{\text{Kul}} = \Lambda_{\text{Kul}}(\Gamma)$.

Theorem 5.1. Let $\Gamma \subset PSL(3, \mathbb{C})$ be a group satisfying $\mu(\Omega_{Kul}(\Gamma)) = 2$, then either the limit set is the union of two lines or it is a pencil of lines over a nowhere dense perfect set. Moreover, one of the following facts occurs:

- (i) if $\lambda(\Omega_{\text{Kul}}(\Gamma)) = 2$, then Γ is virtually solvable and contains a loxodromic element, see [27, 28] for a precise description.
- (ii) if $\lambda(\Omega_{\text{Kul}}(\Gamma)) = \infty$, then Γ either is virtually purely parabolic or contains a loxodromic element, and:
 - (a) If the group is virtually purely parabolic then it contains a finite index subgroup conjugate to one of the following groups: a dual torus group of type II, a Kleinian Inoue group, or an extended Kodaira group K₂.
 - (b) If the group contains a loxodromic element, then either Γ is virtually conjugate to a solvable group (described in [28]), or it is a weakly controllable group whose control group is non-elementary.

10

Proof. Since $\mu(\Omega_{\text{Kul}}(\Gamma)) = 2$, we have that Γ has a global fixed point p, the point where those two lines intersect, and $\lambda(\Omega_{\text{Kul}}(\Gamma))$ is either 2 or ∞ by Theorem 1.1 [4]. In fact, will see in the next section that if $\lambda(\Omega_{\text{Kul}}(\Gamma)) = 3$, then also $\mu(\Omega_{\text{Kul}}(\Gamma)) = 3$. And we know from Theorem 1.5 that if $\lambda(\Omega_{\text{Kul}}(\Gamma)) > 3$, then this number actually is ∞ .

Also, if $\lambda(\Omega_{\text{Kul}}(\Gamma)) = 2$, then, we claim, that Γ is virtually solvable. To see this notice that the group is weakly semi-controllable. Choose a line ℓ as the "horizon". Then Λ_{Kul} meets ℓ in two points which are invariant under the control group $\Pi(\Gamma)$. Hence $\Pi(\Gamma)$ is virtually diagonalizable and therefore Γ itself is virtually diagonalizable. Hence it is virtually solvable. Then, by Theorem 0.1 in [5], we deduce that Γ contains a loxodromic element. Then, by [27, 28], its limit set consists of exactly two lines.

In case $\lambda(\Omega_{\text{Kul}}(\Gamma)) = \infty$, we have that Γ is either virtually purely parabolic or it contains a loxodromic element. In the first case by part (b) item (2) of Theorem 0.1 in [5] we have that Γ contains a finite index subgroup conjugate to one of the following groups: a dual torus group of type II, a Kleinian Inoue group, or an extended Kodaira group \mathcal{K}_2 , thus, in this case, the limit set is a cone of lines over a Euclidean circle, see section 2 of [5]. Now the proof splits into the following cases:

Case 1. The group Γ is virtually solvable. In this case by Theorems 1.1, 1.2, 1.3, 1.4 in [28], the limit set of Γ is either two lines or a cone of lines over an Euclidean circle.

Case 2. The group Γ is non-virtually solvable. In this case it is clear that the control group $\Pi_{p,\ell}\Gamma$ is non-elementary, here ℓ is any line not containing p, and

$$\Lambda_{\mathrm{Kul}}(\Gamma) = \bigcup_{z \in \Lambda(\Pi(\Gamma))} \overleftarrow{p, z}.$$

That $\Pi_{p,\ell}\Gamma$ is non-elementary follows from the fact that, essentially by Borel's fixed point theory, a subgroup of $PSL(2, \mathbb{C})$ is elementary if and only if it is virtually solvable (see [11]).

6. Groups with limit set three lines

We now describe the subgroups of $PSL(3, \mathbb{C})$ with three lines in their limit set. As in the previous section, we denote by $\lambda = \lambda(\Omega_{Kul}(\Gamma))$ the maximum number of complex projective lines contained in Λ_{Kul} , and by $\mu = \mu(\Omega_{Kul}(\Gamma))$ the maximum number of such lines in general position. Then there are groups with $\mu = \lambda = 3$, and there are groups with $\lambda = \infty$ but $\mu = 3$. In the case $\lambda = \infty$ the groups in question are "suspensions" of non-elementary Kleinian groups in PSL(2, \mathbb{C}), see Chapter 5 in [8]:

Theorem 6.1. Let $\Gamma \subset PSL(3, \mathbb{C})$ be a group for which $\mu = 3$. Then λ can be either 3 or ∞ and:

 (i) If λ = 3, then these lines are in general position and Γ contains a finite index subgroup conjugate to:

$$G_0 = \{ diag[\alpha^n, \beta^m, 1] : n, m \in \mathbb{Z} \} ,$$

where $\alpha, \beta \in \mathbb{C}^*$ are non-unitary complex numbers.

 (ii) If λ = ∞, then the limit set is a cone of lines plus a non-concurrent line and Γ contains a finite index subgroup conjugate to:

$$H_{\Sigma,\rho,\alpha} = \left\{ \begin{bmatrix} \alpha^n \rho([g]) & 0\\ 0 & g \end{bmatrix} : n \in \mathbb{Z}, g \in \mathrm{SL}(2,\mathbb{C}), \ [g] \in \Sigma \right\}$$

where $\alpha \in \mathbb{C}^*$ is a non-unitary complex number, Σ is a non-elementary discrete group in $\text{PSL}(2,\mathbb{C})$ with non-empty discontinuity region, and ρ is a function $\Sigma \to \mathbb{C}^*$. In this case

$$\Lambda_{\mathrm{Kul}}(H_{\Sigma,\rho,\alpha}) = \overleftarrow{[e_2], [e_3]} \cup \bigcup_{p \in \Lambda(\Sigma)} \overleftarrow{[e_1], p}$$

Proof. We know from statement (iii) in Theorem 1.5 that if $\lambda = 3$ then $\mu = 3$. That is, if the group has exactly three lines in its limit set, then these lines are in general position.

Now we assume $\mu = 3$ and we choose three distinct lines in general position contained in $\Lambda_{\text{Kul}}(\Gamma)$, say ℓ_1, ℓ_2, ℓ_3 . We define

$$P = \{ p_{ij} = \ell_i \cap \ell_j : 1 \le i < j \le 3 \},\$$

then any line contained in $\Lambda_{\text{Kul}}(\Gamma)$ must intersect P.

We will say $p \in P$ is a vertex whenever there are infinite lines in $\Lambda_{\text{Kul}}(\Gamma)$ passing through it. If the set of vertices is empty, then by Proposition 5.6 in [4] the set P is a Γ -invariant set. Then $H = Isot(P, \Gamma)$ is a finite index subgroup of Γ . Moreover, $\Lambda_{\text{Kul}}(H) = \Lambda_{\text{Kul}}(\Gamma)$ and H is conjugate to a group G_0 where every element has a diagonal lift. So by Theorem 1.1 in [28] we conclude $G_0 = \{diag[\alpha^n, \beta^m, 1] : n, m \in \mathbb{Z}\}$ for some $\alpha, \beta \in \mathbb{C}^*$ non-unitary complex numbers.

If the set of vertices is non-empty then by the proof of Proposition 6.5 in [4] we must have that this set is Γ -invariant. Observe that in case the number of vertices is at least two, we must have four lines in general position (compare with Proposition 6.5 in [4]) which is not possible, therefore we have exactly one vertex, say v, which is Γ -invariant. Let $\ell \in \{\ell_1, \ell_2, \ell_3\}$ be a line not containing v. Since $\mu(\Omega_{\text{Kul}}(\Gamma)) = 3$, we conclude that ℓ is Γ -invariant. After conjugating with a projective transformation, if necessary, let us assume that $v = [e_1]$ and $\ell = [e_2], [e_3]$, as in Section 1.5 take $\pi = \pi_{v,\ell}$ and $\Pi = \Pi_{v,\ell}$, now we claim:

Claim 1. If $\Pi(\Gamma)$ is non-virtually solvable, then $\Pi(\Gamma)$ is discrete. On the contrary let us assume that $\Pi(\Gamma)$ is non-discrete, thus by Theorem 1 and Proposition 12 in [16] the principal component of $\overline{\Pi(\Gamma)}$ is PSL(2, \mathbb{C}) or conjugate to SO(3) or PSL(2, \mathbb{R}). In the first two cases, the orbit of any line passing through v is dense in ℓ , and this is not possible. In the latter case, let $(g_n) \subset \Pi(G)$ be a sequence such that $\lim_{n\to\infty} g_n = Id$ and $h_n = [g_1, g_n] \neq Id$ for $n \geq 2$. If $G_m \in \Gamma$ satisfies $\Pi(G_m) = g_m$, then a straightforward computation shows $\lim_{n\to\infty} [G_1, G_n] = Id$, which is a contradiction.

Claim 2. The group Γ cannot be virtually solvable. On the contrary, let us assume that Γ is solvable, then by Theorem 0.1 in [5] the group Γ contains a loxodromic element. By Theorems 1.1, 1.2, 1.3 and 1.4 in [28], there is $W \subset \mathbb{C}$ a discrete additive subgroup with $rank(W) \leq 2$ and $\alpha, \beta \in \mathbb{C}^*$ such that $\alpha \neq 1$ and $\alpha\beta^2 \notin \mathbb{R}$ and Γ is conjugate to:

12

$$\left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & w \\ 0 & 0 & 1 \end{bmatrix} : w \in W \right\} \rtimes \left\{ \begin{bmatrix} \alpha^{2n} \beta^n & 0 & 0 \\ 0 & \alpha^n \beta^{2n} & 0 \\ 0 & 0 & 1 \end{bmatrix} : n \in \mathbb{Z} \right\}.$$

Now a simple computation shows \boldsymbol{W} is non-discrete, which is a contradiction.

By the previous claims, we conclude that $\Pi(\Gamma)$ is a discrete non-elementary with a non-empty ordinary region. Now the rest of the proof is trivial.

7. GROUPS WITH LIMIT SET FOUR LINES IN GENERAL POSITION LINES

In this section we provide an algebraic characterization of the subgroups of $PSL(3, \mathbb{C})$ with exactly four lines lines in general position in their limit set Λ_{Kul} . Of course that by Theorem 1.5 this implies that Λ_{Kul} actually has infinitely many lines, but only four in general position. The basic reference is [2].

We recall first that an element $A \in SL(2, \mathbb{Z})$ is a hyperbolic toral automorphism if none of its eigenvalues has norm 1 (see [17]).

Definition 7.1. A subgroup Γ of $PSL(3, \mathbb{C})$ is a *hyperbolic toral group* if it is conjugate to a group of the form:

$$\Gamma_A = \left\{ \left(\begin{array}{cc} A^k & b \\ 0 & 1 \end{array} \right) \ \left| \ b = \left(\begin{array}{c} b_1 \\ b_2 \end{array} \right) \right., \ b_1, b_2, k \in \mathbb{Z} \right\} \right\}$$

where $A \in SL(2, \mathbb{Z})$ is a hyperbolic toral automorphism.

One has:

Theorem 7.2. Let $\Gamma \subset PSL(3, \mathbb{C})$ be a discrete group. The maximum number of complex lines in general position contained in its Kulkarni limit set is equal to four if, and only if, Γ contains a hyperbolic toral group whose index is at most 8. [In this case the limit set of the subgroup coincides with that of Γ , and the discontinuity region has four components each one diffeomorphic to $\mathbb{H} \times \mathbb{H}$, where \mathbb{H} denotes the half open plane in \mathbb{C} .]

To prove the theorem above, the first step is showing that a toral group Γ_A as above is the subgroup of $PSL(3, \mathbb{Z})$ generated by the unipotent parabolic elements P_1, P_2 and the strongly loxodromic element L given by:

$$P_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} , P_2 = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} , L = \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix} .$$

Therefore Γ_A is discrete. The dynamical properties of the hyperbolic toral group are used to show that $\Lambda_{\text{Kul}}(L)$ is contained in Kulkarni limit set of Γ_A . Also, a straightforward computation shows that $\Lambda_{\text{Kul}}(P_1)$ is contained in $\Lambda_{\text{Kul}}(\Gamma_A)$. Thus,

$$\Lambda_{\mathrm{Kul}}(\Gamma_A) = \overline{\bigcup_{\gamma \in \Gamma_A} \Lambda_{\mathrm{Kul}}(\gamma)},$$

because both sets in this equation contain three lines in general position (see Theorem 1.2 in [1]).

Now, it is possible to compute the sets $\Lambda_{\text{Kul}}(\gamma)$ for every $\gamma \in \Gamma_A$. One finds that $\Lambda_{\text{Kul}}(\Gamma_A)$ can be described as a union of two pencils of lines over two distinct circles, with vertices two distinct points, which share the line determined by the two vertices. Therefore, the maximum number of lines in general position contained in $\Lambda_{\text{Kul}}(\Gamma_A)$ is equal to four. Moreover, this description shows that the Kulkarni discontinuity region of Γ_A is a disjoint union of four copies of $\mathbb{H} \times \mathbb{H}$, where \mathbb{H} denotes the upper half plane in \mathbb{C} .

Conversely, if we assume that the maximum number of lines in general position contained in the Kulkarni limit set of a group Γ is equal to four, then one shows that there are two distinguished points (called vertices) contained in infinitely many lines in this limit set. The subgroup which stabilizes these two vertices is denoted by Γ_0 and it has index at most two in Γ . Furthermore, it can be proved that $\Omega_{\text{Kul}}(\Gamma_0)$ has four components, each one diffeomorphic to a copy of $\mathbb{H} \times \mathbb{H}$. Finally, the subgroup of Γ_0 stabilizing each of these components has finite index (at most four) in Γ_0 , and it can be shown that it is a hyperbolic toral group.

A description of the quotient spaces $\Omega_{\text{Kul}}(\Gamma_A)/\Gamma_A$ is given by the following theorem (see [6].)

Theorem 7.3. If Γ_A is a hyperbolic toral group then

- (i) The group Γ_A is isomorphic to a lattice of the group Sol (see Definition 7.4).
- (ii) The quotient space $\Omega_{Kul}(\Gamma_A)/\Gamma_A$ is a disjoint union of four copies of

$$(Sol/\Gamma_A) \times \mathbb{R}.$$

Definition 7.4. The group Sol is \mathbb{R}^3 equipped with the operation

$$\left(\left(\begin{array}{c} x_1 \\ y_1 \end{array} \right), t_1 \right) \left(\left(\begin{array}{c} x_2 \\ y_2 \end{array} \right), t_2 \right) = \left(\left(\begin{array}{c} x_1 + e^{t_1} x_2 \\ y_1 + e^{-t_1} y_2 \end{array} \right), t_1 + t_2 \right)$$

The proof of 7.3 is done by giving an explicit smooth foliation of $\mathbb{H} \times \mathbb{H}$ where the leaves are diffeomorphic to Sol, and by showing that $\mathbb{H} \times \mathbb{H}$ is diffeomorphic to Sol $\times \mathbb{R}$ by a Γ_A -equivariant diffeomorphism.

It can be shown that Γ_A is isomorphic to $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$, and this group is determined by the conjugacy class in $\operatorname{GL}(2,\mathbb{Z})$ of A (see [23] pages 96,97). It follows from Theorem 7.2 that there is a countable number of nonisomorphic Complex Kleinian groups such that the maximum number of lines in general position contained in its Kulkarni limit set is equal to four.

8. GROUPS WITH LIMIT SET HAVING AN ISOLATED POINT

Let us suppose now that $\Gamma \subset PSL(3, \mathbb{C})$ is a group with at least an isolated point in its limit set. Then the results explained in this article show that Γ must have only one line in its limit set, for otherwise Λ_{Kul} is a union of lines.

Let ℓ be the unique line in $\Lambda_{\text{Kul}}(\Gamma)$ and let $p \in \Lambda_{\text{Kul}}(\Gamma)$ be an isolated point. Then ℓ is an invariant set and we look at the action of Γ on ℓ to deduce from it consequences on the global dynamics of Γ .

We note that this is somehow similar to the controllable groups, where we have a global fix point and a line L that we call the horizon, which is not invariant but we get an action on it. Then we know a lot about the action of the group on $\mathbb{P}^2_{\mathbb{C}}$ by looking at the induced action on L. In the case we now envisage, we have the invariant line ℓ and the action on it says a lot about the dynamics of the group on $\mathbb{P}^2_{\mathbb{C}}$. There are two possibilities as the action of Γ on ℓ can be either:

- a. Virtually Solvable.
- b. Non- Virtually solvable.

The idea of the proof consists in showing first that the non-virtually solvable case cannot happen, so the action must be virtually solvable. This uses the Tits Alternative.

In the solvable case, the dynamics implies that on the invariant line ℓ we must have either one or two fixed points. Then one shows that in the first case, when there is only one fixed point, one actually has an invariant flag and up to conjugation the elements are all of the form

$$egin{bmatrix} lpha & 0 & 0 \ 0 & eta & \delta \ 0 & 0 & \gamma \end{bmatrix}$$
 .

In the second case we find that, up to conjugation, all the elements are diagonal, and the group consists of lodrodromic elements that are screws (see Section 1.3), so the limit set of each element consists of a point and a line. Then we show that the limit set of the whole group is a point and a line.

The above conisderations lead to the following:

Theorem 8.1. Let $\Gamma \subset PSL(3, \mathbb{C})$ be a group with at least an isolated point in its limit set. Then the limit set consists of exactly one line and one point, and Γ is conjugate to either:

(i)

$$G_0 = \left\{ \operatorname{diag}\left[\alpha^n, e^{2\pi i\theta}, e^{-2\pi i\theta}\right] : n, m \in \mathbb{Z} \right\}$$

where $\theta \in \mathbb{R}$ and $\alpha \in \mathbb{C}^*$ is non-unitary complex number.

(ii) or

$$H_0 = \left\{ \begin{bmatrix} \mu(w)^{-2} & 0 & 0\\ 0 & \mu(w) & w\mu(w)\\ 0 & 0 & \mu(w) \end{bmatrix} : w \in W \right\}$$

where $W \subset \mathbb{C}$ is a discrete additive group and $\mu : (\mathbb{C}, +) \to (\mathbb{C}^*, \cdot)$ is a group morphism satisfying that for every sequence of distinct elements $(w_n) \subset W$, the limit $\lim_{n\to\infty} \mu(w_n)$ is either 0 or ∞ .

Proof. By Theorem 1.2 in [2] we have $\mu(\Omega_{\text{Kul}}(\Gamma)) \leq 2$, and from Theorem 5.1 we conclude $\mu(\Omega_{\text{Kul}}(\Gamma)) = 1$. On the other hand, by Theorem 0.1 in [5], the group Γ contains a Loxodromic element. Let ℓ be the unique line in $\Lambda_{\text{Kul}}(\Gamma)$, $p \in \Lambda_{\text{Kul}}(\Gamma)$ be an isolated point and $\gamma \in \Gamma$ a loxodromic element. We must consider the following cases:

Case 1. The action of Γ restricted to ℓ is given by a virtually solvable group. In this case the group Γ is virtually solvable and the theorem follows from Claim 2 and Theorems 1,1, 1.2, 1.3 and 1.4 in [28], see also section 1 above.

Case 2. The action of Γ restricted to ℓ is given by a non- virtually solvable group. Under this assumption we claim:

Claim. There are $g_1, g_2 \in \Gamma$ such that the group generated by $\{\gamma_1|_{\ell}, \gamma_2|_{\ell}\}$ is a rank two free group which is either purely loxodromic or purely elliptic. If the action of Γ on ℓ is given by a discrete group H, then it is well know that H contains a rank two Shottky group, see Chapter X in [20], these will do the job. If the action on ℓ is given by a non discrete group G, thus by Theorem 1 and Proposition 12 in [16] the principal component of \overline{G} is PSL(2, \mathbb{C}), or conjugate to SO(3) or PSL(2, \mathbb{R}). In any such case we deduce that G contains a non-virtually solvable finitely generated subgroup, as a consequence of Tit's alternative we deduce that G contains a rank two free subgroup which is either purely loxodromic or purely elliptic. Which proves the claim.

Now let g_1, g_2 be the elements given by the previous claim. After conjugating with a projective transformation we can assume that $\ell = [e_2], [e_3]$ so $h = [g_1, g_2]$ is a loxodromic element and

$$h = \begin{bmatrix} 1 & 0 & 0 \\ c & d & e \\ f & g & h \end{bmatrix}$$

where dh - eg = 1 and $(d + h)^2 \in \mathbb{C} - \{4\}$. Thus h is either an elliptic element of infinite order, which cannot happen by the discreteness of Γ , or a strongly loxodromic element whose attracting and repelling points are contained in ℓ . We conclude that the attracting and repelling lines of h are distinct from ℓ . On the other hand, from the λ -Lemma (see Lemma 7.3.6 in [8]) we know that the attracting and repelling line of h are both contained in $\Lambda_{\text{Kul}}(\Gamma)$, which is a contradiction. \Box

References

- W. Barrera, A. Cano, J. P. Navarrete, *The Limit Set of Discrete Subgroups of* PSL(3, C), Mathematical Proceedings of the Cambridge Philosophical Society, Volume 150, Issue 1, January 2011, pp. 129 - 146.
- W. Barrera, A. Cano, J. P. Navarrete, Subgroups of PSL(3, C) with four lines in general position in its limit set, Conformal geometry and dynamics 15 (2011), 160-176.
- W. Barrera, A. Cano, J. P. Navarrete, One line complex Kleinian groups, Pacific J. Mathematics 272 (2014), 275-303.
- W. Barrera, A. Cano, J. P. Navarrete, On the number of lines in the limit set for discrete subgroups of PSL(3, C), Pacific J. Mathematics 281 (2016), 17-49.
- W. Barrera, A. Cano, J. P. Navarrete, J. Seade, Discrete parabolic groups in PSL(3, C), Linear Algebra and its Applications, Vol. 653, 15 November 2022, Pages 430-500.
- W. Barrera, R. García, J. P. Navarrete, Three-dimensional Sol Manifolds and Complex Kleinian Groups, Pacific J. Mathematics 294 (2018), 1-18.
- E. Breuillard, T. Gelander, A topological Tits alternative, Annals of Mathematics, 166 (2007), 427–474.
- A. Cano, J. P. Navarrete, J. Seade, *Complex Kleinian Groups*, Progress in Mathematics, 303, Birkäuser 2013.
- A. Cano, J. Parker, J. Seade, Action of ℝ-Fuchsian groups on CP², Asian journal of mathematics. 20 (2016), 449-474.
- A. Cano, J. Seade, On the Equicontinuity region of Discrete Subgroups of PU(1, n), J. Geom. Anal. (2010) 20, pp 291-305.
- A. Cano, J. Seade, On Discrete groups of Automorphisms of P²_C, Geom. Dedicata 168, 9–60 (2014).
- 12. H. Cartan, S. Eilenberg, Homological Algebra, Princeton University Press, 1999.
- 13. C. Series, A Crash Course on Kleinian Groups, Course notes for the Summer School and Conference on Geometry and Topology of 3-manifolds (6 -24 June 2005), ICTP.
- L. J. Corwin, F. P. Greenleaf, Representations of Nilpotent Lie groups and their Applications, Part I: Basic Theory and Examples, Cambridge University Press, 1989.
- J. P. Fillmore, J. Scheuneman, Fundamental groups of compact complete locally affine complex surfaces, Pacific Journal of Mathematics, Vol. 44, No. 2, 1973, pp. 487-496.
- 16. L. Greenberg, Discrete subgroups of the Lorentz group, Math. Scand. 10, 85–107 (1962)
- 17. A. Katok and B. Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems*, Encyclopedia Math. Appl., Vol. 54, Cambridge University Press, Cambridge (1995).
- R. S. Kulkarni, Groups with Domains of Discontinuity, Math. Ann. No 237, pp. 253-272 (1978).
- A. Marden, Outer Circles (An introduction to hyperbolic 3-manifolds), Cambridge University Press, 2007.

- B. Maskit, *Kleinian Groups*, Grundlehren der Mathematischen Wissenschaften 287, Springer, Berlin, 1988.
- P. J. Myrberg, Untersuchungen über die automorphen Funktionen beliebig vieler Variablen, Acta Math. 46: 3-4 (1925), 215-336.
- J.P. Navarrete, The Trace Function and Complex Kleinian Groups, International Journal of Mathematics. Volume 19, No. 7, August 2008.
- P. de la Harpe, *Topics in Geometric Group Theory*, Chicago Lectures in Mathematics, ISBN 0-226-31721-8.
- 24. J. Ratcliffe, Foundations of Hyperbolic Manifolds, Graduate Texts in Mathematics. 149.
- 25. T. Suwa, Compact Quotient Spaces of C², J. Differential Geometry, 10, 1975, pp. 239-252.
- 26. J. Tits, Free subgroups in linear groups, Journal of Algebra, 20 (2), 1972, 250-270.
- M. Toledo-Acosta, The Dynamics of Solvable Subgroups of PSL(3, C), Bull Braz Math Soc, New Series 53, 127–171 (2022).
- M. Toledo-Acosta, Representations of Solvable Subgroups of PSL(3, C), preprintarXiv:1912.09828, 2021.

FACULTAD DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA DE YUCATÁN, ANILLO PERIFÉRICO NORTE TABLAJE CAT 13615, MÉRIDA, YUCATÁN, MÉXICO. *Email address:* bvargas@uady.mx

INSTITUTO DE MATEMÁTICAS UNAM, AV. UNIVERSIDAD S/N, COL. LOMAS DE CHAMILPA, C.P. 62210, CUERNAVACA, MORELOS, MÉXICO.

 $Email \ address: \verb"angelcano@im.unam.mx"$

FACULTAD DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA DE YUCATÁN, ANILLO PERIFÉRICO NORTE TABLAJE CAT 13615, MÉRIDA, YUCATÁN, MÉXICO. Email address: jp.navarrete@uady.mx

Instituto de Matemáticas UNAM , Av. Universidad S/N, Col. Lomas de Chamilpa,

C.P. 62210, CUERNAVACA, MORELOS, MÉXICO. Email address: jseade@im.unam.mx