

SPORADIC SIMPLE GROUPS AS FLAG-TRANSITIVE AUTOMORPHISM GROUPS OF SYMMETRIC DESIGNS

SEYED HASSAN ALAVI AND ASHRAF DANESHKHAH

ABSTRACT. In this article, we study symmetric designs admitting flag-transitive, point-imprimitive almost simple automorphism groups with socle sporadic simple groups. As a corollary, we present a classification of symmetric designs admitting flag-transitive automorphism group whose socle is a sporadic simple group, and in conclusion, there are exactly seven such designs, one of which admits a point-imprimitive automorphism group and the remaining are point-primitive.

1. INTRODUCTION

A *symmetric* (v, k, λ) design is an incidence structure $\mathcal{D} = (\mathcal{P}, \mathcal{B})$ consisting of a set \mathcal{P} of v points and a set \mathcal{B} of v blocks such that every point is incident with exactly k blocks, and each pair of blocks is incident with exactly λ points. If $2 < k < v - 1$, then \mathcal{D} is called a *nontrivial* symmetric design. The complement of \mathcal{D} is a symmetric design whose points are the points of \mathcal{D} and blocks the complements of the blocks of \mathcal{D} . A *flag* of \mathcal{D} is an incident pair (α, B) , where α and B are a point and a block of \mathcal{D} , respectively. An *automorphism* of a symmetric design \mathcal{D} is a permutation of the points permuting the blocks and preserving the incidence relation. The automorphism group G of \mathcal{D} is called *flag-transitive* if it is transitive on the set of flags of \mathcal{D} , and G is called *anti-flag-transitive* if it is flag-transitive on the complement of \mathcal{D} . If G preserves a nontrivial G -invariant partitions on the point set \mathcal{P} , then G is said to be *point-imprimitive*, otherwise, G is called point-primitive. We use the standard notation for the groups in [4, 5, 10].

The sporadic simple groups as one of the classes of finite simple groups have always been thought as an important tool to study the symmetries of geometries. For example, Witt designs as the Steiner systems are better understood by their automorphism groups which are the sporadic simple Mathieu groups discovered 70 years earlier, see [4, Chapter IV]. This paper is devoted to studying the sporadic almost simple groups G as flag-transitive automorphism groups of symmetric designs. This problem have been studied for the point-primitive automorphism groups [16]. Here, we are interested in the case where G is point-imprimitive.

Theorem 1.1. *Let \mathcal{D} be a nontrivial symmetric design admitting a flag-transitive automorphism group G with socle a sporadic simple group. Then G is point-imprimitive if and only if \mathcal{D} has parameter set $(144, 66, 30)$, $G = M_{12}$, and the point-stabiliser and the block-stabiliser are isomorphic to $\text{PSL}_2(11)$, and both are the intersection of two non-conjugate maximal subgroups of G isomorphic to M_{11} .*

In [15], Praeger and Zhou studied point-imprimitive symmetric (v, k, λ) designs, and they improved the upper bound of k (and so v) in terms of λ given by O'Reilly-Regueiro [14], and a list of possible parameters for $\lambda \leq 10$ is given some of which is still open. This motivates Law, Praeger and Reichard [11] to classify flag-transitive symmetric $(96, 20, 4)$ designs, and as a result of their work, all symmetric designs with $\lambda \leq 4$ admitting a flag-transitive, point-imprimitive subgroup of automorphisms are known. In 2022, Mandić and Šubašić [12] have treated all other open possibilities reported in [15] excluding two sets of parameters. Recently, Montinaro [13] made a contribution to the study of point-imprimitive symmetric designs when $k > \lambda(\lambda - 3)/2$ with size of the intersection of block and imprimitivity class at least 3. In our study of point-imprimitive symmetric designs with sporadic almost simple automorphism groups, we observe that the candidate parameter sets fall into type “a” or “b”, the first two parts in [15, Theorem 1.1]. The outline and the method of proving Theorem 1.1 is described in Section 4.

Date: July 12, 2023.

2020 *Mathematics Subject Classification.* 05B05, 05B25, 20B25, 20D08.

Key words and phrases. Symmetric design, flag-transitive, sporadic simple groups.

Corresponding author: S.H. Alavi.

By Theorem 1.1 and the main result of [16], we are now able to present a classification result for symmetric designs admitting flag-transitive sporadic almost simple automorphism groups. The detailed information about these designs are given in Section 2.

Corollary 1.2. *Let \mathcal{D} be a nontrivial symmetric design admitting a flag-transitive automorphism group with socle a sporadic simple group. Then $v \in \{144, 176, 14080\}$ and (\mathcal{D}, G) is (up to isomorphism) as one of the rows in Table 1.*

TABLE 1. Sporadic simple groups and symmetric 2-designs.

Line	Design	v	k	λ	G	G_α	G_B	Comments	References
1	\mathcal{D}_1	144	66	30	M_{12}	$\text{PSL}_2(11)$	$\text{PSL}_2(11)$	point-imprimitive	Theorem 1.1
2	\mathcal{D}_2	144	66	30	$M_{12}, M_{12}:2$	$\text{PSL}_2(11)$	$\text{PSL}_2(11)$	point-primitive	[16]
3	\mathcal{D}_3	144	66	30	$M_{12}:2$	$\text{PSL}_2(11):2$	$\text{PSL}_2(11):2$	point-primitive	[16]
4	\mathcal{D}_4	176	126	90	M_{22}	A_7	A_7	point-primitive	[6, 16]
5	\mathcal{D}_5	176	50	14	HS	$\text{PSU}_3(5):2$	$\text{PSU}_3(5):2$	point-primitive	[9, 16]
6	\mathcal{D}_6	176	126	90	HS	$\text{PSU}_3(5):2$	$\text{PSU}_3(5):2$	point-primitive	[6, 16]
7	\mathcal{D}_7	14080	12636	11340	Fi_{22}	$\text{P}\Omega_7(3)$	$\text{P}\Omega_7(3)$	point-primitive	[6, 16]

Note: G_α is the point-stabiliser of α and G_B is the block-stabiliser of B .

2. EXAMPLES AND CONSTRUCTIONS

Here, we give the examples of symmetric (v, k, λ) designs admitting flag-transitive almost simple automorphism groups with socle sporadic simple groups. The point-primitive designs (Lines 2-7 of Table 1) are obtained in [16], see also [6, 8]. The point-imprimitive design (Line 1 of Table 1) is obtained in Theorem 1.1 and is constructed in Section 5. We use GAP [7] for the computations and in particular we use the software package “**design**” in GAP for design constructions. In what follows, “ $\text{nr}(G)$ ” is the position of the primitive group G in the list of the library of the primitive permutation groups in GAP. Moreover, the subgroups H and K are the point-stabiliser of a point and the block-stabiliser of a block of \mathcal{D} , respectively.

Line 1. The design \mathcal{D}_1 in this line is point-imprimitive with M_{12} as its automorphism group. Here, we follow the proof of Theorem 1.1, and present an explicit construction of this design. Let G , H and K be the groups generated by the permutations on 144 points listed in Table 2. Then G is the permutation group on 144 points which is isomorphic to M_{12} , and H and K are both subgroups of G of index 144. The subgroups H and K are conjugate in G which are isomorphic to $\text{PSL}_2(11)$. These subgroups are not maximal, and they are in fact contained in maximal subgroups isomorphic to M_{11} . The group G is point-imprimitive with point-stabiliser H fixing 1. The subgroup K has nontrivial orbits of length 11, 11, 55 and 66. The K -orbit B given in (5.1) has length 66 and it is a base block for the unique design \mathcal{D}_1 . As it is shown in the proof of Theorem 1.1, the only possibility for a subgroup to be a block-stabiliser of index 12 is when it is a subgroup of one of two non-conjugate maximal subgroups (isomorphic to) M_{11} . For any pairs of such subgroups with orbit length 66, the associated symmetric designs are isomorphic. The full automorphism group of this design is $M_{12}:2$ which is point-primitive, block-primitive and flag-transitive but not anti-flag-transitive. The same is true for $G = M_{12}$ and the group G has rank 5 on its action on 144 points. We note that the permutation representation of G on 144 points that we have given here is obtained by the coset action of G on the set of right cosets of the intersection of two non-conjugate maximal subgroups isomorphic to M_{11} . Indeed, we start with the permutation representation of M_{12} on 12 points, and once we obtain the subgroups H and K as explained in the proof of Theorem 1.1, we consider the right coset action of M_{12} on 144 right cosets of H . We then find the presentations of G , H and K as in Table 2 and construct the design as described.

Line 2. This design is one of the two point-primitive symmetric designs on 144 points. The full-automorphism group of this design is $M_{12}:2$. The groups M_{12} and $M_{12}:2$ are flag-transitive and point-primitive. Let $G = M_{12}$. The point-stabiliser is the maximal subgroup isomorphic to $\text{PSL}_2(11)$. This symmetric design was first constructed by Wirth [17]. This design is denoted by

TABLE 2. The generators of the group G and its subgroups H and K .

Group	Generators
G	$\alpha_1 = (2,11)(3,6)(4,12)(9,10)(13,109)(14,119)(15,120)(16,114)(17,116)(18,112)(19,115)(20,113)(21,117)(22,118)$ $(23,110)(24,111)(25,30)(26,29)(28,31)(33,34)(37,95)(38,91)(39,93)(40,90)(41,96)(42,86)(43,85)(44,89)(45,92)$ $(46,94)(47,87)(48,88)(49,54)(50,59)(55,60)(56,58)(61,129)(62,128)(63,130)(64,124)(65,131)(66,132)(67,125)$ $(68,127)(69,122)(70,126)(71,123)(72,121)(73,78)(74,83)(79,84)(80,82)(97,137)(98,144)(99,135)(100,133)$ $(101,139)(102,136)(103,140)(104,134)(105,142)(106,138)(107,141)(108,143)$ $\alpha_2 = (1,97,132)(2,107,128)(3,102,126)(4,108,123)(5,101,121)(6,99,129)(7,103,125)(8,104,127)(9,106,122)$ $(10,105,130)(11,98,124)(12,100,131)(13,78,36)(14,74,27)(15,83,26)(16,82,29)(17,80,28)(18,77,30)(19,79,32)$ $(20,84,31)(21,75,33)(22,81,34)(23,76,35)(24,73,25)(37,96,137)(38,92,141)(39,90,136)(40,87,143)(41,85,139)$ $(42,93,135)(43,89,140)(44,91,134)(45,86,138)(46,94,142)(47,88,144)(48,95,133)(49,61,111)(50,71,120)$ $(51,68,116)(52,62,119)(53,67,115)(54,66,109)(55,72,113)(56,70,114)(57,69,118)(58,63,117)(59,64,110)$ $(60,65,112)$
H	$\beta_1 = (2,11)(3,6)(4,12)(9,10)(13,109)(14,119)(15,120)(16,114)(17,116)(18,112)(19,115)(20,113)(21,117)(22,118)$ $(23,110)(24,111)(25,30)(26,29)(28,31)(33,34)(37,95)(38,91)(39,93)(40,90)(41,96)(42,86)(43,85)(44,89)(45,92)$ $(46,94)(47,87)(48,88)(49,54)(50,59)(55,60)(56,58)(61,129)(62,128)(63,130)(64,124)(65,131)(66,132)(67,125)$ $(68,127)(69,122)(70,126)(71,123)(72,121)(73,78)(74,83)(79,84)(80,82)(97,137)(98,144)(99,135)(100,133)$ $(101,139)(102,136)(103,140)(104,134)(105,142)(106,138)(107,141)(108,143)$ $\beta_2 = (3,6)(4,7)(5,11)(8,10)(13,128)(14,130)(15,122)(16,132)(17,121)(18,12)(19,124)(20,126)(21,127)(22,131)(23,125)$ $(24,129)(26,29)(27,30)(31,36)(32,33)(37,115)(38,109)(39,116)(40,114)(41,117)(42,111)(43,119)(44,113)(45,120)$ $(46,118)(47,112)(48,110)(49,51)(50,53)(52,60)(56,58)(61,86)(62,91)(63,85)(64,95)(65,94)(66,90)(67,88)(68,96)$ $(69,92)(70,89)(71,87)(72,93)(73,135)(74,139)(75,144)(76,143)(77,137)(78,134)(79,142)(80,141)(81,140)(82,136)$ $(83,138)(84,133)(99,104)(100,105)(101,106)(102,107)$ $\beta_3 = (2,9)(3,5)(4,11)(10,12)(14,16)(15,17)(19,20)(21,23)(25,69)(26,71)(27,64)(28,68)(29,63)(30,65)(31,67)(32,72)$ $(33,70)(34,61)(35,62)(36,66)(37,54)(38,57)(39,51)(40,50)(41,53)(42,55)(43,59)(44,52)(45,60)(46,49)(47,58)$ $(48,56)(73,113)(74,119)(75,111)(76,118)(77,112)(78,115)(79,116)(80,110)(81,109)(82,117)(83,114)(84,120)$ $(85,90)(86,92)(87,88)(94,95)(97,139)(98,134)(99,142)(100,135)(101,136)(102,137)(103,140)(104,143)(105,133)$ $(106,138)(107,141)(108,144)(122,131)(123,130)(125,127)(126,129)$
K	$\gamma_1 = (1,106)(2,97)(3,108)(4,102)(5,98)(6,104)(7,107)(8,103)(9,99)(10,101)(11,105)(12,100)(13,64)(14,71)(15,61)$ $(16,70)(17,65)(18,67)(19,68)(20,72)(21,63)(22,66)(23,69)(24,62)(25,120)(26,110)(27,109)(28,112)(29,114)$ $(30,117)(31,113)(32,115)(33,116)(34,119)(35,111)(36,118)(37,48)(38,41)(39,40)(45,47)(49,58)(51,59)(53,57)$ $(55,56)(73,121)(74,130)(75,128)(76,132)(77,127)(78,125)(79,122)(80,131)(81,124)(82,123)(83,129)(84,126)$ $(85,93)(86,88)(87,94)(91,96)(133,137)(135,139)(136,142)(138,140)$ $\gamma_2 = (1,136)(2,134)(3,135)(4,143)(5,142)(6,133)(7,137)(8,139)(9,138)(10,144)(11,141)(12,140)(13,20)(14,22)(15,17)$ $(21,24)(25,111)(26,116)(27,115)(28,109)(29,120)(30,117)(31,113)(32,112)(33,118)(34,119)(35,114)(36,110)$ $(37,75)(38,82)(39,77)(40,81)(41,74)(42,83)(43,78)(44,73)(45,76)(46,80)(47,79)(48,84)(49,66)(50,68)(51,72)$ $(52,67)(53,61)(54,69)(55,62)(56,63)(57,65)(58,71)(59,64)(60,70)(85,89)(86,96)(87,88)(91,94)(98,106)(99,100)$ $(103,108)(104,107)(122,132)(123,130)(124,127)(126,128)$ $\gamma_3 = (1,11)(3,4)(6,10)(9,12)(13,110)(14,109)(15,120)(16,112)(17,113)(18,117)(19,116)(20,115)(21,114)(22,118)$ $(23,119)(24,111)(25,40)(26,42)(27,41)(28,37)(29,46)(30,47)(31,39)(32,48)(33,38)(34,43)(35,45)(36,44)(49,128)$ $(50,121)(51,130)(52,132)(53,124)(54,125)(55,127)(56,131)(57,122)(58,129)(59,126)(60,123)(61,65)(62,67)$ $(68,71)(70,72)(73,103)(74,108)(75,106)(76,107)(77,99)(78,97)(79,101)(80,100)(81,104)(82,98)(83,105)(84,102)$ $(85,91)(86,94)(87,90)(93,95)(133,144)(136,139)(137,140)(142,143)$

\mathcal{D}_1 in [16]. Here $\text{nr}(G) = 3$, that is to say, $G = M_{12}$ is the 3rd group in the list of primitive permutation groups in GAP. The base block is

$$B_2 = \{6, 7, 9, 10, 13, 17, 18, 19, 20, 21, 32, 33, 34, 36, 37, 38, 46, 49, 51, 52, 53, 54, 58, 60, 61, 62, 63, \\ 64, 65, 66, 67, 68, 69, 71, 72, 74, 78, 79, 81, 85, 87, 89, 90, 92, 94, 95, 96, 98, 102, 105, 106, 109, \\ 110, 114, 118, 121, 122, 126, 127, 128, 129, 132, 134, 137, 141, 142\}.$$

The point-stabiliser H and the block-stabiliser K of \mathcal{D} are maximal subgroups of G , both isomorphic to $\text{PSL}_2(11)$ but not conjugate. The full automorphism group of \mathcal{D}_2 is $M_{12} : 2$ which is point-primitive, block-primitive and flag-transitive, and both G and $\text{Aut}(\mathcal{D}_2)$ are not anti-flag-transitive.

Line 3. The design in this line is the second symmetric design with parameter set $(144, 66, 30)$. Its automorphism group $M_{12}:2$ is flag-transitive and point-primitive with point-stabiliser (isomorphic to) $\text{PSL}_2(11):2$. This design is denoted by \mathcal{D}_2 in [16], and it is constructed by the 5th group $G = M_{12} : 2$ in the list of primitive permutation groups in GAP and the base block is

$$B_3 = \{3, 5, 6, 7, 8, 10, 11, 14, 16, 20, 21, 23, 25, 26, 27, 28, 29, 33, 39, 40, 44, 45, 47, 49, 51, 53, 54, \\ 56, 57, 59, 60, 61, 62, 64, 65, 71, 72, 73, 75, 76, 79, 82, 84, 85, 90, 95, 96, 97, 103, 104, 107, 109, \\ 111, 112, 113, 115, 116, 119, 126, 132, 134, 137, 138, 142, 143, 144\}.$$

The group G is flag-transitive, point-primitive and block-primitive but not anti-flag-transitive. The point-stabiliser and the block-stabiliser of \mathcal{D}_3 are isomorphic to $\text{PSL}_2(11) : 2$. We observe that if we take the socle of G and construct the design with the base block B_3 , then we obtain a design which is isomorphic to \mathcal{D}_1 .

Line 4. The design in this line arose from studying rank 3 automorphism groups of symmetric designs [6]. Here $G = \text{M}_{22}$ and $\text{nr}(G) = 3$, and if we take the base block B_4 below, then we obtain \mathcal{D}_4 as in Line 4 of Table 1. This design is denoted by \mathcal{D}_3 in [16].

$$B_4 = \{1, 3, 4, 5, 6, 7, 10, 12, 13, 14, 16, 17, 19, 23, 24, 25, 27, 28, 29, 31, 32, 33, 34, 35, 37, 38, 39, \\ 40, 41, 42, 43, 44, 45, 46, 47, 48, 50, 53, 54, 56, 60, 62, 63, 64, 65, 66, 68, 69, 70, 73, 74, 75, \\ 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 88, 89, 90, 91, 92, 94, 95, 96, 97, 99, 100, 101, 102, \\ 103, 105, 106, 107, 108, 109, 112, 114, 117, 119, 120, 123, 124, 125, 126, 128, 129, 132, \\ 134, 138, 139, 140, 141, 142, 143, 144, 147, 148, 149, 151, 152, 153, 154, 155, 156, 159, 160, \\ 161, 162, 163, 164, 165, 167, 168, 171, 172, 173, 174, 175, 176\}.$$

The full automorphism group of \mathcal{D}_4 is HS which is point-primitive, block-primitive, flag-transitive and anti-flag-transitive with both point-stabiliser and block-stabiliser isomorphic to $\text{PSU}_3(5):2$ but non-conjugate. The group $G = \text{M}_{22}$ is point-primitive, block-primitive, flag-transitive but not anti-flag-transitive, and both point-stabiliser and block-stabiliser are isomorphic to A_7 .

Line 5. The design in this line is the unique 2-transitive symmetric $(176, 50, 14)$ design [9]. This design is denoted by \mathcal{D}_4 in [16]. The point-stabiliser (block-stabiliser) of this design is isomorphic to $\text{PSU}_3(5):2$. The group $G = \text{HS}$ is flag-transitive, point-primitive, block-primitive and anti-flag-transitive, and the base block is

$$B_5 = \{1, 7, 10, 11, 15, 17, 25, 26, 30, 33, 36, 40, 41, 43, 51, 59, 64, 66, 69, 85, 90, 94, 97, \\ 99, 106, 109, 113, 114, 115, 119, 122, 132, 136, 142, 143, 148, 150, 152, 153, 154, 155, \\ 160, 162, 164, 166, 167, 168, 169, 171, 176\}. \quad (2.1)$$

Line 6. The design in this line is rank 3 symmetric $(176, 126, 14)$ design [6] with automorphism group HS. The point-stabiliser (block-stabiliser) is $\text{PSU}_3(11)$. The design \mathcal{D}_6 is the complement of the design in Line 5, and hence the base block of \mathcal{D}_6 is $\mathcal{P} \setminus B_5$, where $\mathcal{P} := \{1, \dots, 176\}$ and B_5 is as in 2.1. This design is denoted by \mathcal{D}_5 in [16].

Line 7. This design also arose from studying rank 3 automorphism groups of symmetric designs [6], and it has parameter set $(14080, 12636, 11340)$ admitting Fi_{22} as flag-transitive and point-primitive automorphism group with the point-stabiliser and block-stabiliser isomorphic to $\text{P}\Omega_7(3)$ (non-conjugate). This design is denoted by \mathcal{D}_6 in [16].

3. PRELIMINARY

In this section, we state some useful facts in both design theory and group theory, and then we establish our method in Section 4 in order to analyse symmetric designs admitting sporadic simple groups as their flag-transitive automorphism groups.

Lemma 3.1. [1, Lemma 2.1] *Let \mathcal{D} be a symmetric (v, k, λ) design, and let G be a flag-transitive automorphism group of \mathcal{D} . If α is a point of \mathcal{D} and $H = G_\alpha$, then*

- (a) $k(k-1) = \lambda(v-1)$;
- (b) $k \mid |H|$ and $\lambda v < k^2$;
- (c) $k \mid \lambda \gcd(v-1, |H|)$;
- (d) $|G| < |H|^3$;
- (e) $k \mid \lambda e$, for all nontrivial subdegrees e of G .

Lemma 3.2 below is an elementary result that is useful in computational matter for finding potential parameter sets. An algorithm based on this lemma is given in [2, Remark 3.8, Algorithm 1]. We remark here that in [2, Algorithm 1], we need to substitute the parameter “ d ” with parameter “ t ” as in this paper “ d ” is the number of the imprimitivity classes.

Lemma 3.2. [2, Lemma 3.7] *Let \mathcal{D} be a symmetric (v, k, λ) design \mathcal{D} , and let $mk = \lambda t$, where t is a divisor of $v-1$, for some positive integer m . Then*

- (a) $m \mid (k-1)$, and so $\gcd(m, k) = 1$;

- (b) $\lambda = \lambda_1 \lambda_2$, where $\lambda_1 = \gcd(\lambda, k-1)$ and $\lambda_2 = \gcd(\lambda, k)$;
(c) If $k_1 := (k-1)/\lambda_1$ and $k_2 := k/\lambda_2$, then $v-1 = k_1 k_2$, where k_2 divides t . Moreover, λ_1 divides m , $\lambda_1 < k_2$ and $\gcd(\lambda_1, k_2) = 1$.

Lemma 3.3. [15, Theorem 1.1] *Let $D = (\mathcal{P}, \mathcal{B})$ be a nontrivial symmetric (v, k, λ) design admitting a flag-transitive and point-imprimitive subgroup of automorphisms G that leaves invariant a nontrivial partition \mathcal{C} of \mathcal{P} with d classes of size c . Then there is a constant ℓ such that, for each $B \in \mathcal{B}$ and $\Delta \in \mathcal{C}$, the size $|B \cap \Delta|$ is either 0 or ℓ , and one of the following holds:*

- (a) $k \leq \lambda(\lambda-3)/2$;
(b) $(v, k, \lambda) = (\lambda^2(\lambda+2), \lambda(\lambda+1), \lambda)$ with $(c, d, \ell) = (\lambda^2, \lambda+2, \lambda)$ or $(\lambda+2, \lambda^2, 2)$;
(c) $(v, k, \lambda, c, d, \ell) = (\frac{(\lambda+2)(\lambda^2-2\lambda+2)}{4}, \frac{\lambda^2}{2}, \lambda, \frac{\lambda+2}{2}, \frac{\lambda^2-2\lambda+2}{2}, 2)$, and either $\lambda \equiv 0 \pmod{4}$, or $\lambda = 2u^2$, where u is odd, $u \geq 3$, and $2(u^2-1)$ is a square;
(d) $(v, k, \lambda, c, d, \ell) = (\frac{(\lambda+6)(\lambda^2+4\lambda-1)}{4}, \frac{\lambda(\lambda+5)}{2}, \lambda, \lambda+6, \frac{\lambda^2+4\lambda-1}{4}, 3)$, where $\lambda \equiv 1$ or $3 \pmod{6}$.

With the same notation as in Lemma 3.3, we have the following equalities:

$$v = cd, \tag{3.1}$$

$$k = \ell s, \tag{3.2}$$

$$\lambda(c-1) = k(\ell-1), \tag{3.3}$$

where s is the number of classes of \mathcal{C} that intersect a block B nontrivially, and $\ell > 1$ and $s > 1$.

Lemma 3.4. *Let $D = (\mathcal{P}, \mathcal{B})$ be a nontrivial symmetric (v, k, λ) design admitting a flag-transitive and point-imprimitive automorphism group G , and let (α, B) be a flag of \mathcal{D} . Let also M and N be the maximal subgroups of G containing the point-stabiliser $H := G_\alpha$ and the block-stabiliser $K := G_B$, respectively. Then*

- (a) $v = |G : H| = |G : K|$;
(b) $|G : M| \neq v$, and both $|G : M|$ and $|G : N|$ divide v ;
(c) $k = |H : H_B|$ and the index of a maximal subgroup of H containing H_B divides k ;
(d) $k = |K : K_\alpha|$ and the index of a maximal subgroup of K containing K_α divides k .

Proof. Part (a) follows from the fact that G is point-transitive and block-transitive. The fact that G is point-imprimitive implies that $M \neq H$, or equivalently, $|G : M| \neq |G : H| = v$, and so part (b) holds. Since G is flag-transitive, H is transitive on the set of blocks containing α and K is transitive on B , and hence parts (c)-(d) follow. \square

4. METHOD

In this section, we describe our method to find candidate parameters which are listed in Tables 6-7 in Appendix. In what follows, we suppose that $\mathcal{D} = (\mathcal{P}, \mathcal{B})$ is a nontrivial symmetric (v, k, λ) design admitting a flag-transitive and point-imprimitive subgroup G of automorphisms of \mathcal{D} . Suppose also that G is an almost simple group whose socle is a sporadic simple group. Let (α, B) be a flag of \mathcal{D} , and let M and N be the maximal subgroups of G containing the point-stabiliser $H := G_\alpha$ and the block-stabiliser $K := G_B$, respectively. The possibilities for M and N can be read of from GAP [7] and Atlas [5]. We use the following approach to obtain the possible parameter sets.

- (1) By Lemma 3.4(a), we have that $|H| = |K|$, and since G is flag-transitive and M and N are respectively maximal subgroups containing H and K , it follows that both M and N satisfy the large property as in Lemma 3.1(d), that is to say, $|G| \leq |M|^3$ and $|G| \leq |N|^3$. This enables us to obtain the possibilities for both M and N by [3, Proposition 6.2]. Here, we also use the database stored in GAP [7].
- (2) For a fixed M obtained in (1), since $v = |G : H| = |G : M| \cdot |M : H|$, we conclude that $v = z \cdot |G : M|$, for some divisor z of $|M|$. Therefore, for each M , we can find possible parameters v .
- (3) For each M and each (associated) v obtained in (2), we know by Lemma 3.1(b) that k divides $|H|$, so does $|M|$. We now use Lemma 3.2. Let $t := \gcd(v-1, |M|)$. Then for each divisor k_2 of t , let $k_1 := (v-1)/k_2$. For $\lambda_1 < k_2$ with $\gcd(\lambda_1, k_2) = 1$, let $k := 1 + k_1 \lambda_1$ and $\lambda := k(k-1)/(v-1)$. We check if λ is an integer, k divides M and $\lambda v < k^2$. This produces a list of possible parameters (v, k, λ) for each M . Here we use modified version of [2, Algorithm 1] with changing $\lambda_1 \leq k_2/2$ to $\lambda_1 \leq k_2$ and substituting the restrictive condition $2k < v$ to $k < v-1$. This is for finding potential complement parameter sets.

TABLE 3. The number of parameter sets for some sporadic almost simple groups.

Group	# parameters	Group	# parameters	Group	# parameters	Group	# parameters	Group	# parameters
M ₁₁	3	J ₂	0	Suz:2	51	O'N:2	52	Fi' ₂₄ :2	298
M ₁₂	26	J ₂ :2	9	McL	2	Co ₁	0	HN	0
M ₁₂ :2	11	J ₃	0	McL:2	35	Co ₂	0	HN:2	59
M ₂₂	6	J ₃ :2	11	Ru	0	Co ₃	1	Th	0
M ₂₂ :2	19	J ₄	0	He	1	Fi ₂₂	12	B	0
M ₂₃	1	HS	29	He:2	34	Fi ₂₂ :2	57	M	0
M ₂₄	4	HS:2	45	Ly	0	Fi ₂₃	0		
J ₁	0	Suz	0	O'N	0	Fi' ₂₄	0		

- (4) Since $k := |K : K_\alpha|$ divides $|K| = |H| = |G|/v$, it follows that k divides $|N|$, where N is a maximal subgroup of G containing K . Note also that $|G : N|$ divides $v = |G : K|$. This leaves 22 groups with 766 tuples $(M, N, (v, k, \lambda))$ listed in Tables 6-7 for further consideration. For each group, the number of such candidate parameter sets is recorded in Table 3. As we are also interested in point-imprimitive action, for each parameter set (v, k, λ) , we can also find parameters (c, d, ℓ) using (3.1), (3.2) and (3.3). By Lemma 3.3, we also can associate one of the type “a”, “b”, “c” and “d” to the parameter set $(v, k, \lambda, c, d, \ell)$ according to the part of Lemma 3.3 that the parameter set $(v, k, \lambda, c, d, \ell)$ satisfies.
- (5) For each pairs (M, N) , we check if M (respectively, N) has a subgroup of index $i_H := v/|G : M|$ (respectively, $i_K := v/|G : N|$). This suggests the pairs (H, K) of point-block stabilisers. In the case where M is of large order, we look for a maximal subgroup of M whose index divides i_H . This gives the possibilities for oversubgroups of H , and then we continue this process to find possible candidates for H . We do the same when N is of large order.
- (6) For each subgroup H obtained in (5), we consider the right coset action of G on the set of the right cosets of H in G , and obtain the H -orbits and their lengths (subdegrees). Since G is flag-transitive, we check if k is a divisor of λe , for all nontrivial subdegrees e of G , see Lemma 3.1(e). This also eliminates some cases.
- (7) For the remaining pairs (H, K) and parameters (v, k, λ) that have not been eliminated in (6), we consider the right coset action of G on the set of the right cosets of H in G , and obtain the K -orbits and their lengths in this action. Since G is flag-transitive, the length of one of the K -orbits has to be k . If such an orbit B exists, then we check if B is a base block for a symmetric design using the software package “design” in GAP [7].

5. PROOF OF THE MAIN RESULTS

In this section, we prove Theorem 1.1 and Corollary 1.2. In order to prove Theorem 1.1, we follow the steps described in Section 4.

Proof of Theorem 1.1 Suppose that $\mathcal{D} = (\mathcal{P}, \mathcal{B})$ is a nontrivial symmetric (v, k, λ) design admitting a flag-transitive, point-imprimitive subgroup G of automorphisms of \mathcal{D} . Suppose also that G is an almost simple group whose socle is a sporadic simple group. Let (α, B) be a flag of \mathcal{D} , and let M and N be the maximal subgroups of G containing the point-stabiliser $H := G_\alpha$ and the block-stabiliser $K := G_B$, respectively. We apply the method described in Section 4, and as it is mentioned in step 4, there are 22 groups with total number 766 of parameters which possibly give rise to a 2-design. We divide these groups in 2 classes:

- (A) M₁₁, M₁₂, M₁₂:2, M₂₂, M₂₂:2, M₂₃, M₂₄, J₂:2, J₃:2, HS, HS:2, McL, McL:2, He, He:2, Co₃;
 (B) Suz : 2, O'N : 2, HN : 2, Fi₂₂, Fi₂₂ : 2, Fi'₂₄ : 2.

(A) For the groups in class A, once we follow the steps in Section 4, all possibilities but 6 cases can be ruled out in steps 5 or 6. We record all the possibilities for the groups in Class A in Table 6. In the last column of Table 6, we use the notation “nsg” or “nsd”, or we leave it blank. We have 6 possible parameters in the latter case which are also separately recorded in Table 4.

The notation “nsg” (no subgroup) in Table 6 indicates that the case can be ruled out in step 6, that is to say, the maximal subgroup M (or N) has no subgroup of index i_H (or i_K). For example, if $G = M_{12}:2$, then the point-stabiliser H is contained in $M = M_{12}$. Then M_{12} should have a subgroup of index $v/2$, where $v \in \{16, 22, 40, 66, 88, 96, 144, 160, 396\}$, which is impossible.

TABLE 4. The remaining possible parameters for the groups in class A.

G	M	N	$(\text{nr}(M), \text{nr}(N))$	(i_H, i_K)	v	k	λ	c	d	ℓ
M_{12}	M_{11}	M_{11}	(1, 1)	(12, 12)	144	66	30	12	12	6
M_{12}	M_{11}	M_{11}	(1, 2)	(12, 12)	144	66	30	12	12	6
M_{12}	M_{11}	$\text{PSL}_2(11)$	(1, 5)	(12, 1)	144	66	30	12	12	6
M_{12}	M_{11}	M_{11}	(2, 1)	(12, 12)	144	66	30	12	12	6
M_{12}	M_{11}	M_{11}	(2, 2)	(12, 12)	144	66	30	12	12	6
M_{12}	M_{11}	$\text{PSL}_2(11)$	(2, 5)	(12, 1)	144	66	30	12	12	6

$i_H := |M : H|$ and $i_K := |N : K|$.

If “nsd” (no subdegree) is recorded in the last column of Table 6, then it means that there exist some subgroup H and K with $|M : H| = i_H$ and $|N : K| = i_K$. By considering H as a subgroup of the permutation group G on $v = |G : H|$ points, we observe that there exist some nontrivial subdegrees e of G such that λe is not divisible by k , which violates Lemma 3.1(e). In this case, we also present the smallest such a subdegree in this column. For example, let $G = \text{HS}$ and $(v, k, \lambda) = (8800, 420, 20)$, and $M = \text{PSU}_3(5) : 2$ and $N = \text{A}_8 : 2$. Then M has a subgroup $H = \text{S}_7$ of index $|M : H| = 50$. Then the permutation group G on the set of right cosets of H with 8800 points has nontrivial subdegrees $e \in \{7, 42, 126, 210, 252, 630, 1260, 2520\}$, and it is easy to see that $k = 420$ does not divide $\lambda e = 20e$ for any such e , which is a contradiction.

This leaves the possibilities recorded in Table 4 in which $G = M_{12}$. Moreover, $(v, k, \lambda) = (144, 66, 30)$ for which the point-stabiliser H is a subgroup of the maximal subgroup M_1 or M_2 , where $M_i \cong M_{11}$, and the block-stabiliser K is a subgroup of the maximal subgroup $N_1 \cong M_{11}$, $N_2 \cong M_{11}$ or $N_3 \cong \text{PSL}_2(11)$. We note here that M_1 and M_2 (respectively, N_1 and N_2) are not conjugate in G . It is also easy to check that $H = M_1 \cap M_2$ and it is isomorphic to $\text{PSL}_2(11)$. If $K \leq N_3$, then since $|G : N_3| = 144 = |G : K|$, we conclude in this case that $K = N_3 \cong \text{PSL}_2(11)$ is maximal in G , and the K -orbits are of length 12 and 132, which is a contradiction as $k = 66$.

Suppose that $K \leq N_1$. Then $K = N_1 \cap N_2$ which is conjugate to $H \cong \text{PSL}_2(11)$. Thus $K \cong \text{PSL}_2(11)$, and the length of K -orbits in the right coset action of G on the set of right cosets of H in G are 11, 11, 55 and 66. The K -orbit B of length 66 gives rise to the symmetric $(144, 66, 30)$ design $\mathcal{D} = (\mathcal{P}, B^G)$, where $G = M_{12}$, $\mathcal{P} = \{1, \dots, 144\}$ and B is the base block

$$\begin{aligned}
B := \{ & 1, 3, 4, 5, 8, 11, 13, 14, 16, 19, 20, 22, 26, 27, 28, 32, 33, 36, 37, 38, 41, 42, 44, \\
& 48, 49, 50, 51, 58, 59, 60, 64, 66, 68, 70, 71, 72, 73, 74, 75, 82, 83, 84, 98, 102, \\
& 103, 105, 106, 108, 109, 110, 112, 115, 116, 118, 121, 123, 126, 128, 129, 130, \\
& 135, 136, 139, 141, 142, 143 \}.
\end{aligned} \tag{5.1}$$

We denote this design by \mathcal{D}_1 which is the design in Line 1 of Table 1. We note here that the full automorphism group of \mathcal{D}_1 is $M_{12} : 2$ which is point-primitive while M_{12} is point-imprimitive with imprimitive class representatives $\{1, 13, 35, 38, 57, 62, 81, 91, 103, 109, 128, 140\}$ and $\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$. This shows that $(c, d, \ell) = (12, 12, 6)$. The case where $K \leq N_2$ can be treated similarly and we find the same design. In fact, each pair of designs obtained in this way (the orbit of length 66 of the intersection of two non-conjugate maximal subgroups isomorphic to M_{11}) are isomorphic.

(B) For the groups in class B, the possible parameters are listed in Table 7. Here, we prove that none of these groups gives rise to a 2-design. Indeed, for each group, we show that one of steps 5-6 fails, but for the computation matters, we need to check these cases separately.

Let $G = \text{Fi}_{22}$. In this case, the maximal subgroup M whose index divides v is one of the two non-conjugate subgroups (isomorphic to) $\text{P}\Omega_7(3)$. Then H is a subgroup of M of index 14, 40 or 105. However, $\text{P}\Omega_7(3)$ has no subgroup of such an index as the smallest index of the maximal subgroups of $\text{P}\Omega_7(3)$ is 351, see [5, p. 109].

Let now $G = \text{Fi}_{22}:2$. Then we have 190 possible parameter sets (v, k, λ) and H is contained in one of the two non-conjugate maximal subgroups of G being isomorphic to Fi_{22} . Let M be one of these subgroups. Then $i_H = |M : H|$ is 8, 20, 32, 33, 44, 48, 56, 60, 72, 80, 88, 108, 128, 143, 144, 165, 168, 182, 189, 198, 224, 243, 270, 273, 280, 320, 336, 350, 448, 455, 540, 585, 648, 704, 728, 864, 1008, 1320, 2048, 2800, 2912, 4032, 4050, 4752, 6804, 7040, 7488, 10368, 11200, 18480, 25344, 52800, 81648, 98560, 281600 or 739200. By inspecting the indices of the maximal subgroups of $M \cong \text{Fi}_{22}$ dividing i_H , we observe that H is contained in one of two non-conjugate maximal

TABLE 5. The remaining possible parameters for the groups in class B.

G	M	The indices of maximal subgroups of M
Suz : 2	Suz	1782, 22880, 32760, 135135, 232960, 370656, 405405, 926640, 1216215, 2358720, 3203200, 10378368, 17297280, 39916800, 57480192, 177914880
O'N : 2	O'N	122760, 2624832, 2857239, 17778376, 30968784, 42858585, 58183776, 182863296
HN : 2	HN	1140000, 1539000, 16500000, 74064375, 108345600, 136515456, 164587500, 263340000, 264515625, 364041216, 1436400000, 2926000000, 4681600000
Fi'_{24} : 2	Fi'_{24}	306936, 4860485028, 14081405184, 50177360142, 125168046080, 258870277120, 2503413946215, 5686767482760, 7819305288795, 70094105804800, 91122337546240, 100087107696576, 155717756992512, 633363728392395, 8212275503308800, 57650174033227776, 2306006961329111040, 11859464372549713920, 103770313259809996800, 574727888823563059200, 3091639677809511628800

subgroups $P\Omega_7(3)$ of M of index 7 or 20, and so $P\Omega_7(3)$ has to have a subgroup of index 7 or 20, which is impossible as the smallest index of the maximal subgroups of $P\Omega_7(3)$ is 351 by [5, p. 109].

For the remaining groups, we make the same argument. The required information about this groups are given in Table 5. If G is one of these groups as in the first column of Table 5, then H is contained in the maximal subgroup M of G (indeed the socle of G) as in the second column of the same table, and so and the index of a maximal subgroup of M containing H (listed in the third column) has to divide i_H which is recorded in Table 7, which is impossible.

Proof of Corollary 1.2 If G is point-imprimitive, then by Theorem 1.1, we obtain the design in Line 1 of Table 1. If G is point-primitive, then by [16, Theorem 1], we obtain one of the designs in Lines 2-7 of Table 1.

ACKNOWLEDGEMENTS

The authors are grateful to Alice Devillers and Cheryl E. Praeger for supporting their visit to UWA (The University of Western Australia) during February–June 2023. They also thank Bu-Ali Sina University for the support during their sabbatical leave. The authors thank the reviewers for their comments and suggestions.

REFERENCES

- [1] S. H. Alavi, M. Bayat, A. Daneshkhah, Symmetric designs admitting flag-transitive and point-primitive automorphism groups associated to two dimensional projective special groups, *Designs, Codes and Cryptography* 79 (2) (2016) 337–351.
URL <http://dx.doi.org/10.1007/s10623-015-0055-9> 4
- [2] S. H. Alavi, M. Bayat, A. Daneshkhah, Finite exceptional groups of Lie type and symmetric designs, *Discrete Mathematics* 345 (8) (2022) 112894.
URL <https://www.sciencedirect.com/science/article/pii/S0012365X22001005> 4, 5
- [3] S. H. Alavi, T. C. Burness, Large subgroups of simple groups, *J. Algebra* 421 (2015) 187–233.
URL <http://dx.doi.org/10.1016/j.jalgebra.2014.08.026> 5
- [4] T. Beth, D. Jungnickel, H. Lenz, Design theory. Vol. I, vol. 69 of *Encyclopedia of Mathematics and its Applications*, 2nd ed., Cambridge University Press, Cambridge, 1999.
URL <https://doi.org/10.1017/CB09780511549533> 1
- [5] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, R. A. Wilson, Atlas of finite groups, Oxford University Press, Eynsham, 1985, maximal subgroups and ordinary characters for simple groups, With computational assistance from J. G. Thackray. 1, 5, 7, 8
- [6] U. Dempwolff, Primitive rank 3 groups on symmetric designs, *Designs, Codes and Cryptography* 22 (2) (2001) 191–207.
URL <https://doi.org/10.1023/A:1008373207617> 2, 4
- [7] The GAP Group, GAP – Groups, Algorithms, and Programming, Version 4.11.0 (2020).
URL <http://www.gap-system.org> 2, 5, 6, 10
- [8] W. M. Kantor, 2-transitive designs, in: *Combinatorics*, Springer Netherlands, 1975, pp. 365–418. 2
- [9] W. M. Kantor, Classification of 2-transitive symmetric designs, *Graphs Combin.* 1 (2) (1985) 165–166.
URL <http://dx.doi.org/10.1007/BF02582940> 2, 4
- [10] P. Kleidman, M. Liebeck, The subgroup structure of the finite classical groups, vol. 129 of *London Mathematical Society Lecture Note Series*, Cambridge University Press, Cambridge, 1990.
URL <http://dx.doi.org/10.1017/CB09780511629235> 1
- [11] M. Law, C. E. Praeger, S. Reichard, Flag-transitive symmetric 2-(96, 20, 4)-designs, *J. Combin. Theory Ser. A* 116 (5) (2009) 1009–1022.
URL <http://dx.doi.org/10.1016/j.jcta.2008.12.005> 1

- [12] J. Mandić, A. Šubašić, Flag-transitive and point-imprimitive symmetric designs with $\lambda \leq 10$, J. Combin. Theory Ser. A 189 (2022) Paper No. 105620.
URL <https://doi.org/10.1016/j.jcta.2022.105620> 1
- [13] A. Montinaro, Flag-transitive, point-imprimitive symmetric $2-(v, k, \lambda)$ designs with $k > \lambda(\lambda - 3)/2$, arXiv e-prints (2022) arXiv:2203.09261. 1
- [14] E. O'Reilly-Regueiro, On primitivity and reduction for flag-transitive symmetric designs, J. Combin. Theory Ser. A 109 (1) (2005) 135–148.
URL <http://dx.doi.org/10.1016/j.jcta.2004.08.002> 1
- [15] C. E. Praeger, S. Zhou, Imprimitive flag-transitive symmetric designs, J. Combin. Theory Ser. A 113 (7) (2006) 1381–1395.
URL <http://dx.doi.org/10.1016/j.jcta.2005.12.006> 1, 5
- [16] D. Tian, S. Zhou, Flag-transitive $2-(v, k, \lambda)$ symmetric designs with sporadic socle, Journal of Combinatorial Designs 23 (4) (2015) 140–150.
URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/jcd.21385> 1, 2, 3, 4, 8
- [17] W. Wirth, Konstruktion symmetrischer designs, Ph.D. thesis, University of Mainz (2000). 2

APPENDIX

In this section, we present two tables namely Tables 6 and 7 which include all the possible parameter sets and the required information about the groups and their subgroups which we obtained by applying our method described in Section 4. In these tables, the first column shows the almost simple group G with socle a sporadic simple group. Recall that H and K are our potential point-stabiliser and block-stabiliser, respectively. The subgroups M and N in the second and third columns are the maximal subgroups of G containing H and K , respectively. In the fourth column, the first coordinate $\text{nr}(M)$ indicates the number associated to the maximal subgroup M of G in the library of software package “AtlasRep” in GAP [7], and similarly, the second coordinate $\text{nr}(N)$ shows the number associated to the subgroup N . In column 5, $i_H = |M : H|$ and $i_K = |N : K|$. The candidates of parameters v , k and λ are recorded in columns 6-8. In column 9, we give the type of imprimitivity of group G which is one of the parts (a)-(d) in Lemma 3.3 that the parameters (v, k, λ) satisfy. In last column of Table 6, we have two notation “nsg” and “nsd”, or it is left blank. The notation “nsg” (no subgroup) indicates that the maximal subgroup M (or N) has no subgroup of index i_H (or i_K). The notation “nsd” (no subdegree) means that there exist some subgroup H , but H as permutation subgroup of G on v points does not satisfy Lemma 3.1(e), that is to say, there exist a non-trivial H -orbit of length e such that λe is not divisible by k , and in such a case, the smallest subdegree e is presented. If the last column of a row in Table 6 is left blank, then the case has been treated separately in Section 5.

TABLE 6. The possible parameters for groups in class A.

G	M	N	$(\text{nr}(M), \text{nr}(N))$	(i_H, i_K)	v	k	λ	Type	Comments
M_{11}	$A_6.2_3$	$A_5.2$	(1, 4)	(6, 1)	66	40	24	a	nsg
M_{11}	$A_6.2_3$	$A_6.2_3$	(1, 1)	(6, 6)	66	40	24	a	nsg
M_{11}	$A_6.2_3$	$A_6.2_3$	(1, 1)	(8, 8)	88	30	10	a	nsg
M_{12}	M_{11}	$\text{PSL}_2(11)$	(1, 5)	(12, 1)	144	66	30	a	
M_{12}	M_{11}	M_{11}	(1, 1)	(12, 12)	144	66	30	a	
M_{12}	M_{11}	M_{11}	(1, 2)	(12, 12)	144	66	30	a	
M_{12}	M_{11}	$\text{PSL}_2(11)$	(2, 5)	(12, 1)	144	66	30	a	
M_{12}	M_{11}	M_{11}	(2, 1)	(12, 12)	144	66	30	a	
M_{12}	M_{11}	M_{11}	(2, 2)	(12, 12)	144	66	30	a	
M_{12}	$A_6.2^2$	$2 \times S_5$	(3, 8)	(6, 1)	396	80	16	a	nsg
M_{12}	$A_6.2^2$	$A_6.2^2$	(3, 3)	(6, 6)	396	80	16	a	nsg
M_{12}	$A_6.2^2$	M_{11}	(3, 1)	(6, 33)	396	80	16	a	nsg
M_{12}	$A_6.2^2$	M_{11}	(3, 2)	(6, 33)	396	80	16	a	nsg
M_{12}	$A_6.2^2$	$A_6.2^2$	(3, 4)	(6, 6)	396	80	16	a	nsg
M_{12}	M_{11}	$2 \times S_5$	(1, 8)	(33, 1)	396	80	16	a	nsg
M_{12}	M_{11}	$A_6.2^2$	(1, 3)	(33, 6)	396	80	16	a	nsg
M_{12}	M_{11}	M_{11}	(1, 1)	(33, 33)	396	80	16	a	nsg
M_{12}	M_{11}	M_{11}	(1, 2)	(33, 33)	396	80	16	a	nsg
M_{12}	M_{11}	$A_6.2^2$	(1, 4)	(33, 6)	396	80	16	a	nsg
M_{12}	M_{11}	$2 \times S_5$	(2, 8)	(33, 1)	396	80	16	a	nsg
M_{12}	M_{11}	$A_6.2^2$	(2, 3)	(33, 6)	396	80	16	a	nsg
M_{12}	M_{11}	M_{11}	(2, 1)	(33, 33)	396	80	16	a	nsg
M_{12}	M_{11}	M_{11}	(2, 2)	(33, 33)	396	80	16	a	nsg
M_{12}	M_{11}	$A_6.2^2$	(2, 4)	(33, 6)	396	80	16	a	nsg
M_{12}	$A_6.2^2$	$2 \times S_5$	(4, 8)	(6, 1)	396	80	16	a	nsg
M_{12}	$A_6.2^2$	$A_6.2^2$	(4, 3)	(6, 6)	396	80	16	a	nsg
M_{12}	$A_6.2^2$	M_{11}	(4, 1)	(6, 33)	396	80	16	a	nsg
M_{12}	$A_6.2^2$	M_{11}	(4, 2)	(6, 33)	396	80	16	a	nsg
M_{12}	$A_6.2^2$	$A_6.2^2$	(4, 4)	(6, 6)	396	80	16	a	nsg
$M_{12}:2$	M_{12}	M_{12}	(1, 1)	(8, 8)	16	6	2	b	nsg
$M_{12}:2$	M_{12}	M_{12}	(1, 1)	(20, 20)	40	27	18	a	nsg
$M_{12}:2$	M_{12}	M_{12}	(1, 1)	(33, 33)	66	40	24	a	nsg
$M_{12}:2$	M_{12}	M_{12}	(1, 1)	(44, 44)	88	30	10	a	nsg
$M_{12}:2$	M_{12}	M_{12}	(1, 1)	(48, 48)	96	20	4	b	nsg
$M_{12}:2$	M_{12}	$\text{PSL}_2(11).2$	(1, 2)	(72, 1)	144	66	30	a	nsg
$M_{12}:2$	M_{12}	M_{12}	(1, 1)	(72, 72)	144	66	30	a	nsg
$M_{12}:2$	M_{12}	$\text{PSL}_2(11).2$	(1, 3)	(72, 1)	144	66	30	a	nsg
$M_{12}:2$	M_{12}	M_{12}	(1, 1)	(80, 80)	160	54	18	a	nsg
$M_{12}:2$	M_{12}	$(2^2xA_5) : 2$	(1, 4)	(198, 1)	396	80	16	a	nsg
$M_{12}:2$	M_{12}	M_{12}	(1, 1)	(198, 198)	396	80	16	a	nsg
M_{22}	$\text{PSL}_3(4)$	$\text{PSL}_3(4)$	(1, 1)	(3, 3)	66	40	24	a	nsg
M_{22}	$\text{PSL}_3(4)$	$\text{PSL}_3(4)$	(1, 1)	(4, 4)	88	30	10	a	nsg
M_{22}	$\text{PSL}_3(4)$	A_7	(1, 3)	(8, 1)	176	126	90	a	nsg

Continued

TABLE 6 - continued

G	M	N	$(nr(M), nr(N))$	(i_H, i_K)	v	k	λ	Type	Comments
M ₂₂	PSL ₃ (4)	A ₇	(1, 4)	(8, 1)	176	126	90	a	nsg
M ₂₂	PSL ₃ (4)	PSL ₃ (4)	(1, 1)	(8, 8)	176	126	90	a	nsg
M ₂₂	PSL ₃ (4)	PSL ₃ (4)	(1, 1)	(18, 18)	396	80	16	a	nsg
M _{22:2}	M ₂₂	M ₂₂	(1, 1)	(8, 8)	16	6	2	b	nsg
M _{22:2}	M ₂₂	M ₂₂	(1, 1)	(32, 32)	64	28	12	a	nsg
M _{22:2}	PSL ₃ (4).2 ₂	PSL ₃ (4).2 ₂	(2, 2)	(3, 3)	66	40	24	a	nsg
M _{22:2}	PSL ₃ (4).2 ₂	M ₂₂	(2, 1)	(3, 33)	66	40	24	a	nsg
M _{22:2}	M ₂₂	PSL ₃ (4).2 ₂	(1, 2)	(33, 3)	66	40	24	a	nsg
M _{22:2}	M ₂₂	M ₂₂	(1, 1)	(33, 33)	66	40	24	a	nsg
M _{22:2}	PSL ₃ (4).2 ₂	PSL ₃ (4).2 ₂	(2, 2)	(4, 4)	88	30	10	a	nsg
M _{22:2}	PSL ₃ (4).2 ₂	M ₂₂	(2, 1)	(4, 44)	88	30	10	a	nsg
M _{22:2}	M ₂₂	PSL ₃ (4).2 ₂	(1, 2)	(44, 4)	88	30	10	a	nsg
M _{22:2}	M ₂₂	M ₂₂	(1, 1)	(44, 44)	88	30	10	a	nsg
M _{22:2}	M ₂₂	M ₂₂	(1, 1)	(48, 48)	96	20	4	b	nsg
M _{22:2}	PSL ₃ (4).2 ₂	PSL ₃ (4).2 ₂	(2, 2)	(8, 8)	176	126	90	a	nsg
M _{22:2}	PSL ₃ (4).2 ₂	M ₂₂	(2, 1)	(8, 88)	176	126	90	a	nsg
M _{22:2}	M ₂₂	PSL ₃ (4).2 ₂	(1, 2)	(88, 8)	176	126	90	a	nsg
M _{22:2}	M ₂₂	M ₂₂	(1, 1)	(88, 88)	176	126	90	a	nsg
M _{22:2}	PSL ₃ (4).2 ₂	PSL ₃ (4).2 ₂	(2, 2)	(18, 18)	396	80	16	a	nsg
M _{22:2}	PSL ₃ (4).2 ₂	M ₂₂	(2, 1)	(18, 198)	396	80	16	a	nsg
M _{22:2}	M ₂₂	PSL ₃ (4).2 ₂	(1, 2)	(198, 18)	396	80	16	a	nsg
M _{22:2}	M ₂₂	M ₂₂	(1, 1)	(198, 198)	396	80	16	a	nsg
M ₂₃	M ₂₂	M ₂₂	(1, 1)	(12, 12)	276	176	112	a	nsg
M ₂₄	M ₂₃	M ₂₃	(1, 1)	(4, 4)	96	20	4	b	nsg
M ₂₄	M ₂₃	M ₂₃	(1, 1)	(6, 6)	144	66	30	a	nsg
M ₂₄	M ₂₃	M ₂₃	(1, 1)	(12, 12)	288	42	6	b	nsg
M ₂₄	M ₂₃	M ₂₃	(1, 1)	(24, 24)	576	276	132	a	nsg
J _{2:2}	J ₂	J ₂	(1, 1)	(8, 8)	16	6	2	b	nsg
J _{2:2}	J ₂	J ₂	(1, 1)	(20, 20)	40	27	18	a	nsg
J _{2:2}	J ₂	J ₂	(1, 1)	(32, 32)	64	28	12	a	nsg
J _{2:2}	J ₂	J ₂	(1, 1)	(48, 48)	96	20	4	b	nsg
J _{2:2}	J ₂	J ₂	(1, 1)	(56, 56)	112	75	50	a	nsg
J _{2:2}	J ₂	J ₂	(1, 1)	(60, 60)	120	35	10	a	nsg
J _{2:2}	J ₂	J ₂	(1, 1)	(80, 80)	160	54	18	a	nsg
J _{2:2}	J ₂	J ₂	(1, 1)	(144, 144)	288	42	6	b	nsg
J _{2:2}	J ₂	J ₂	(1, 1)	(224, 224)	448	150	50	a	nsg
J _{3:2}	J ₃	J ₃	(1, 1)	(8, 8)	16	6	2	b	nsg
J _{3:2}	J ₃	J ₃	(1, 1)	(20, 20)	40	27	18	a	nsg
J _{3:2}	J ₃	J ₃	(1, 1)	(38, 38)	76	51	34	a	nsg
J _{3:2}	J ₃	J ₃	(1, 1)	(48, 48)	96	20	4	b	nsg
J _{3:2}	J ₃	J ₃	(1, 1)	(80, 80)	160	54	18	a	nsg
J _{3:2}	J ₃	J ₃	(1, 1)	(128, 128)	256	171	114	a	nsg
J _{3:2}	J ₃	J ₃	(1, 1)	(152, 152)	304	102	34	a	nsg
J _{3:2}	J ₃	J ₃	(1, 1)	(170, 170)	340	114	38	a	nsg
J _{3:2}	J ₃	J ₃	(1, 1)	(323, 323)	646	216	72	a	nsg
J _{3:2}	J ₃	J ₃	(1, 1)	(1292, 1292)	2584	288	32	a	nsg
J _{3:2}	J ₃	J ₃	(1, 1)	(1539, 1539)	3078	544	96	a	nsg
HS	M ₂₂	A _{8.2}	(1, 5)	(11, 1)	1100	315	90	a	nsg
HS	M ₂₂	PSL ₃ (4).2 ₁	(1, 4)	(11, 1)	1100	315	90	a	nsg
HS	M ₂₂	M ₂₂	(1, 1)	(11, 11)	1100	315	90	a	nsg
HS	M ₂₂	M ₂₂	(1, 1)	(12, 12)	1200	110	10	b	nsg
HS	A _{8.2}	A _{8.2}	(5, 5)	(8, 8)	8800	420	20	b	nsd, 7
HS	A _{8.2}	PSU ₃ (5).2	(5, 3)	(8, 50)	8800	420	20	b	nsd, 7
HS	A _{8.2}	PSL ₃ (4).2 ₁	(5, 4)	(8, 8)	8800	420	20	b	nsd, 7
HS	A _{8.2}	M ₂₂	(5, 1)	(8, 88)	8800	420	20	b	nsd, 7
HS	A _{8.2}	PSU ₃ (5).2	(5, 2)	(8, 50)	8800	420	20	b	nsd, 7
HS	PSU ₃ (5).2	A _{8.2}	(3, 5)	(50, 8)	8800	420	20	b	nsd, 7
HS	PSU ₃ (5).2	PSU ₃ (5).2	(3, 3)	(50, 50)	8800	420	20	b	nsd, 7
HS	PSU ₃ (5).2	PSL ₃ (4).2 ₁	(3, 4)	(50, 8)	8800	420	20	b	nsd, 7
HS	PSU ₃ (5).2	M ₂₂	(3, 1)	(50, 88)	8800	420	20	b	nsd, 7
HS	PSU ₃ (5).2	PSU ₃ (5).2	(3, 2)	(50, 50)	8800	420	20	b	nsd, 7
HS	PSL ₃ (4).2 ₁	A _{8.2}	(4, 5)	(8, 8)	8800	420	20	b	nsg
HS	PSL ₃ (4).2 ₁	PSU ₃ (5).2	(4, 3)	(8, 50)	8800	420	20	b	nsg
HS	PSL ₃ (4).2 ₁	PSL ₃ (4).2 ₁	(4, 4)	(8, 8)	8800	420	20	b	nsg
HS	PSL ₃ (4).2 ₁	M ₂₂	(4, 1)	(8, 88)	8800	420	20	b	nsg
HS	PSL ₃ (4).2 ₁	PSU ₃ (5).2	(4, 2)	(8, 50)	8800	420	20	b	nsg
HS	M ₂₂	A _{8.2}	(1, 5)	(88, 8)	8800	420	20	b	nsg
HS	M ₂₂	PSU ₃ (5).2	(1, 3)	(88, 50)	8800	420	20	b	nsg
HS	M ₂₂	PSL ₃ (4).2 ₁	(1, 4)	(88, 8)	8800	420	20	b	nsg
HS	M ₂₂	M ₂₂	(1, 1)	(88, 88)	8800	420	20	b	nsg

Continued

TABLE 6 - continued

G	M	N	$(nr(M), nr(N))$	(i_H, i_K)	v	k	λ	Type	Comments
HS	M_{22}	$PSU_3(5).2$	(1, 2)	(88, 50)	8800	420	20	b	nsg
HS	$PSU_3(5).2$	$A_8.2$	(2, 5)	(50, 8)	8800	420	20	b	nsd, 7
HS	$PSU_3(5).2$	$PSU_3(5).2$	(2, 3)	(50, 50)	8800	420	20	b	nsd, 7
HS	$PSU_3(5).2$	$PSL_3(4).2_1$	(2, 4)	(50, 8)	8800	420	20	b	nsd, 7
HS	$PSU_3(5).2$	M_{22}	(2, 1)	(50, 88)	8800	420	20	b	nsd, 7
HS	$PSU_3(5).2$	$PSU_3(5).2$	(2, 2)	(50, 50)	8800	420	20	b	nsd, 7
HS:2	HS	HS	(1, 1)	(8, 8)	16	6	2	b	nsg
HS:2	HS	HS	(1, 1)	(32, 32)	64	28	12	a	nsg
HS:2	HS	HS	(1, 1)	(33, 33)	66	40	24	a	nsg
HS:2	HS	HS	(1, 1)	(44, 44)	88	30	10	a	nsg
HS:2	HS	HS	(1, 1)	(48, 48)	96	20	4	b	nsg
HS:2	HS	HS	(1, 1)	(50, 50)	100	45	20	a	nsg
HS:2	HS	$M_{22}:2$	(1, 2)	(50, 1)	100	45	20	a	nsg
HS:2	HS	HS	(1, 1)	(56, 56)	112	75	50	a	nsg
HS:2	HS	HS	(1, 1)	(60, 60)	120	35	10	a	nsg
HS:2	HS	HS	(1, 1)	(88, 88)	176	126	90	a	nsg
HS:2	HS	HS	(1, 1)	(125, 125)	250	84	28	a	nsg
HS:2	HS	HS	(1, 1)	(198, 198)	396	80	16	a	nsg
HS:2	HS	HS	(1, 1)	(224, 224)	448	150	50	a	nsg
HS:2	HS	HS	(1, 1)	(320, 320)	640	72	8	b	nsg
HS:2	HS	HS	(1, 1)	(336, 336)	672	550	450	a	nsg
HS:2	HS	HS	(1, 1)	(448, 448)	896	180	36	a	nsg
HS:2	HS	HS	(1, 1)	(550, 550)	1100	315	90	a	nsg
HS:2	HS	$PSL_3(4).2^2$	(1, 3)	(550, 1)	1100	315	90	a	nsg
HS:2	HS	$M_{22}:2$	(1, 2)	(550, 11)	1100	315	90	a	nsg
HS:2	HS	$S_8 \times 2$	(1, 4)	(550, 1)	1100	315	90	a	nsg
HS:2	$M_{22}:2$	HS	(2, 1)	(11, 550)	1100	315	90	a	nsg
HS:2	$M_{22}:2$	$PSL_3(4).2^2$	(2, 3)	(11, 1)	1100	315	90	a	nsg
HS:2	$M_{22}:2$	$M_{22}:2$	(2, 2)	(11, 11)	1100	315	90	a	nsg
HS:2	$M_{22}:2$	$S_8 \times 2$	(2, 4)	(11, 1)	1100	315	90	a	nsg
HS:2	HS	HS	(1, 1)	(600, 600)	1200	110	10	b	nsg
HS:2	HS	$M_{22}:2$	(1, 2)	(600, 12)	1200	110	10	b	nsg
HS:2	$M_{22}:2$	HS	(2, 1)	(12, 600)	1200	110	10	b	nsg
HS:2	$M_{22}:2$	$M_{22}:2$	(2, 2)	(12, 12)	1200	110	10	b	nsg
HS:2	HS	HS	(1, 1)	(4032, 4032)	8064	2200	600	a	nsg
HS:2	HS	HS	(1, 1)	(4400, 4400)	8800	420	20	b	nsg
HS:2	HS	$PSL_3(4).2^2$	(1, 3)	(4400, 8)	8800	420	20	b	nsg
HS:2	HS	$M_{22}:2$	(1, 2)	(4400, 88)	8800	420	20	b	nsg
HS:2	HS	$S_8 \times 2$	(1, 4)	(4400, 8)	8800	420	20	b	nsg
HS:2	$PSL_3(4).2^2$	HS	(3, 1)	(8, 4400)	8800	420	20	b	nsg
HS:2	$PSL_3(4).2^2$	$PSL_3(4).2^2$	(3, 3)	(8, 8)	8800	420	20	b	nsg
HS:2	$PSL_3(4).2^2$	$M_{22}:2$	(3, 2)	(8, 88)	8800	420	20	b	nsg
HS:2	$PSL_3(4).2^2$	$S_8 \times 2$	(3, 4)	(8, 8)	8800	420	20	b	nsg
HS:2	$M_{22}:2$	HS	(2, 1)	(88, 4400)	8800	420	20	b	nsg
HS:2	$M_{22}:2$	$PSL_3(4).2^2$	(2, 3)	(88, 8)	8800	420	20	b	nsg
HS:2	$M_{22}:2$	$M_{22}:2$	(2, 2)	(88, 88)	8800	420	20	b	nsg
HS:2	$M_{22}:2$	$S_8 \times 2$	(2, 4)	(88, 8)	8800	420	20	b	nsg
HS:2	$S_8 \times 2$	HS	(4, 1)	(8, 4400)	8800	420	20	b	nsd, 7
HS:2	$S_8 \times 2$	$PSL_3(4).2^2$	(4, 3)	(8, 8)	8800	420	20	b	nsd, 7
HS:2	$S_8 \times 2$	$M_{22}:2$	(4, 2)	(8, 88)	8800	420	20	b	nsd, 7
HS:2	$S_8 \times 2$	$S_8 \times 2$	(4, 4)	(8, 8)	8800	420	20	b	nsd, 7
McL	$PSU_4(3)$	$PSU_4(3)$	(1, 1)	(4, 4)	1100	315	90	a	nsg
McL	$PSU_4(3)$	$PSU_4(3)$	(1, 1)	(32, 32)	8800	420	20	b	nsg
McL:2	McL	McL	(1, 1)	(8, 8)	16	6	2	b	nsg
McL:2	McL	McL	(1, 1)	(20, 20)	40	27	18	a	nsg
McL:2	McL	McL	(1, 1)	(32, 32)	64	28	12	a	nsg
McL:2	McL	McL	(1, 1)	(33, 33)	66	40	24	a	nsg
McL:2	McL	McL	(1, 1)	(44, 44)	88	30	10	a	nsg
McL:2	McL	McL	(1, 1)	(48, 48)	96	20	4	b	nsg
McL:2	McL	McL	(1, 1)	(50, 50)	100	45	20	a	nsg
McL:2	McL	McL	(1, 1)	(56, 56)	112	75	50	a	nsg
McL:2	McL	McL	(1, 1)	(60, 60)	120	35	10	a	nsg
McL:2	McL	McL	(1, 1)	(72, 72)	144	66	30	a	nsg
McL:2	McL	McL	(1, 1)	(80, 80)	160	54	18	a	nsg
McL:2	McL	McL	(1, 1)	(88, 88)	176	126	90	a	nsg
McL:2	McL	McL	(1, 1)	(125, 125)	250	84	28	a	nsg
McL:2	McL	McL	(1, 1)	(144, 144)	288	42	6	b	nsg
McL:2	McL	McL	(1, 1)	(165, 165)	330	189	108	a	nsg
McL:2	McL	McL	(1, 1)	(168, 168)	336	135	54	a	nsg
McL:2	McL	McL	(1, 1)	(198, 198)	396	80	16	a	nsg

Continued

TABLE 6 - continued

G	M	N	$(nr(M), nr(N))$	(i_H, i_K)	v	k	λ	Type	Comments
McL:2	McL	McL	(1, 1)	(224, 224)	448	150	50	a	nsg
McL:2	McL	McL	(1, 1)	(270, 270)	540	99	18	a	nsg
McL:2	McL	McL	(1, 1)	(336, 336)	672	550	450	a	nsg
McL:2	McL	McL	(1, 1)	(550, 550)	1100	315	90	a	nsg
McL:2	McL	PSU ₄ (3).2 ₃	(1, 2)	(550, 4)	1100	315	90	a	nsg
McL:2	PSU ₄ (3).2 ₃	McL	(2, 1)	(4, 550)	1100	315	90	a	nsg
McL:2	PSU ₄ (3).2 ₃	PSU ₄ (3).2 ₃	(2, 2)	(4, 4)	1100	315	90	a	nsg
McL:2	McL	McL	(1, 1)	(600, 600)	1200	110	10	b	nsg
McL:2	McL	McL	(1, 1)	(648, 648)	1296	630	306	a	nsg
McL:2	McL	McL	(1, 1)	(864, 864)	1728	1100	700	a	nsg
McL:2	McL	McL	(1, 1)	(1320, 1320)	2640	378	54	a	nsg
McL:2	McL	McL	(1, 1)	(1458, 1458)	2916	1750	1050	a	nsg
McL:2	McL	McL	(1, 1)	(2800, 2800)	5600	3564	2268	a	nsg
McL:2	McL	McL	(1, 1)	(3168, 3168)	6336	5250	4350	a	nsg
McL:2	McL	McL	(1, 1)	(4400, 4400)	8800	420	20	b	nsg
McL:2	McL	PSU ₄ (3).2 ₃	(1, 2)	(4400, 32)	8800	420	20	b	nsg
McL:2	PSU ₄ (3).2 ₃	McL	(2, 1)	(32, 4400)	8800	420	20	b	nsg
McL:2	PSU ₄ (3).2 ₃	PSU ₄ (3).2 ₃	(2, 2)	(32, 32)	8800	420	20	b	nsg
He	2 ² .PSL ₃ (4).S ₃	2 ² .PSL ₃ (4).S ₃	(2, 2)	(2, 2)	16660	4320	1120	a	nsd, 105
He:2	He	He	(1, 1)	(8, 8)	16	6	2	b	nsg
He:2	He	He	(1, 1)	(20, 20)	40	27	18	a	nsg
He:2	He	He	(1, 1)	(32, 32)	64	28	12	a	nsg
He:2	He	He	(1, 1)	(48, 48)	96	20	4	b	nsg
He:2	He	He	(1, 1)	(56, 56)	112	75	50	a	nsg
He:2	He	He	(1, 1)	(60, 60)	120	35	10	a	nsg
He:2	He	He	(1, 1)	(80, 80)	160	54	18	a	nsg
He:2	He	He	(1, 1)	(128, 128)	256	120	56	a	nsg
He:2	He	He	(1, 1)	(140, 140)	280	63	14	a	nsg
He:2	He	He	(1, 1)	(144, 144)	288	42	6	b	nsg
He:2	He	He	(1, 1)	(153, 153)	306	245	196	a	nsg
He:2	He	He	(1, 1)	(200, 200)	400	343	294	a	nsg
He:2	He	He	(1, 1)	(224, 224)	448	150	50	a	nsg
He:2	He	He	(1, 1)	(256, 256)	512	147	42	a	nsg
He:2	He	He	(1, 1)	(315, 315)	630	408	264	a	nsg
He:2	He	He	(1, 1)	(320, 320)	640	72	8	b	nsg
He:2	He	He	(1, 1)	(392, 392)	784	378	182	a	nsg
He:2	He	He	(1, 1)	(408, 408)	816	490	294	a	nsg
He:2	He	He	(1, 1)	(448, 448)	896	180	36	a	nsg
He:2	He	He	(1, 1)	(480, 480)	960	686	490	a	nsg
He:2	He	He	(1, 1)	(680, 680)	1360	756	420	a	nsg
He:2	He	He	(1, 1)	(816, 816)	1632	700	300	a	nsg
He:2	He	He	(1, 1)	(900, 900)	1800	1029	588	a	nsg
He:2	He	He	(1, 1)	(1568, 1568)	3136	210	14	b	nsg
He:2	He	He	(1, 1)	(1600, 1600)	3200	1372	588	a	nsg
He:2	He	He	(1, 1)	(2448, 2448)	4896	980	196	a	nsg
He:2	He	He	(1, 1)	(2975, 2975)	5950	1323	294	a	nsg
He:2	He	He	(1, 1)	(6800, 6800)	13600	1512	168	a	nsg
He:2	He	He	(1, 1)	(7200, 7200)	14400	2058	294	a	nsg
He:2	2 ² .PSL ₃ (4).D ₁₂	2 ² .PSL ₃ (4).D ₁₂	(3, 3)	(2, 2)	16660	4320	1120	a	nsd, 105
He:2	2 ² .PSL ₃ (4).D ₁₂	He	(3, 1)	(2, 8330)	16660	4320	1120	a	nsd, 105
He:2	He	2 ² .PSL ₃ (4).D ₁₂	(1, 3)	(8330, 2)	16660	4320	1120	a	nsd, 105
He:2	He	He	(1, 1)	(8330, 8330)	16660	4320	1120	a	nsd, 105
He:2	He	He	(1, 1)	(29988, 29988)	59976	2400	96	a	nsg
Co3	McL:2	McL:2	(1, 1)	(11, 11)	3036	1215	486	a	nsg

TABLE 7. The possible parameters for the groups in class B.

G	M	N	$(nr(M), nr(N))$	(i_H, i_K)	v	k	λ	Type
Suz:2	Suz	Suz	(1, 1)	(8, 8)	16	6	2	b
Suz:2	Suz	Suz	(1, 1)	(20, 20)	40	27	18	a
Suz:2	Suz	Suz	(1, 1)	(32, 32)	64	28	12	a
Suz:2	Suz	Suz	(1, 1)	(33, 33)	66	40	24	a
Suz:2	Suz	Suz	(1, 1)	(44, 44)	88	30	10	a
Suz:2	Suz	Suz	(1, 1)	(48, 48)	96	20	4	b
Suz:2	Suz	Suz	(1, 1)	(56, 56)	112	75	50	a
Suz:2	Suz	Suz	(1, 1)	(60, 60)	120	35	10	a
Suz:2	Suz	Suz	(1, 1)	(72, 72)	144	66	30	a
Suz:2	Suz	Suz	(1, 1)	(80, 80)	160	54	18	a

Continued

TABLE 7 - continued

G	M	N	$(nr(M), nr(N))$	(i_H, i_K)	v	k	λ	Type
Suz:2	Suz	Suz	(1, 1)	(88, 88)	176	126	90	a
Suz:2	Suz	Suz	(1, 1)	(108, 108)	216	130	78	a
Suz:2	Suz	Suz	(1, 1)	(128, 128)	256	120	56	a
Suz:2	Suz	Suz	(1, 1)	(143, 143)	286	96	32	a
Suz:2	Suz	Suz	(1, 1)	(144, 144)	288	42	6	b
Suz:2	Suz	Suz	(1, 1)	(165, 165)	330	189	108	a
Suz:2	Suz	Suz	(1, 1)	(168, 168)	336	135	54	a
Suz:2	Suz	Suz	(1, 1)	(182, 182)	364	243	162	a
Suz:2	Suz	Suz	(1, 1)	(189, 189)	378	117	36	a
Suz:2	Suz	Suz	(1, 1)	(198, 198)	396	80	16	a
Suz:2	Suz	Suz	(1, 1)	(224, 224)	448	150	50	a
Suz:2	Suz	Suz	(1, 1)	(243, 243)	486	195	78	a
Suz:2	Suz	Suz	(1, 1)	(270, 270)	540	99	18	a
Suz:2	Suz	Suz	(1, 1)	(273, 273)	546	110	22	a
Suz:2	Suz	Suz	(1, 1)	(280, 280)	560	130	30	a
Suz:2	Suz	Suz	(1, 1)	(320, 320)	640	72	8	b
Suz:2	Suz	Suz	(1, 1)	(336, 336)	672	550	450	a
Suz:2	Suz	Suz	(1, 1)	(350, 350)	700	234	78	a
Suz:2	Suz	Suz	(1, 1)	(448, 448)	896	180	36	a
Suz:2	Suz	Suz	(1, 1)	(455, 455)	910	405	180	a
Suz:2	Suz	Suz	(1, 1)	(540, 540)	1080	416	160	a
Suz:2	Suz	Suz	(1, 1)	(585, 585)	1170	168	24	a
Suz:2	Suz	Suz	(1, 1)	(648, 648)	1296	260	52	a
Suz:2	Suz	Suz	(1, 1)	(648, 648)	1296	630	306	a
Suz:2	Suz	Suz	(1, 1)	(704, 704)	1408	336	80	a
Suz:2	Suz	Suz	(1, 1)	(728, 728)	1456	486	162	a
Suz:2	Suz	Suz	(1, 1)	(864, 864)	1728	1100	700	a
Suz:2	Suz	Suz	(1, 1)	(1008, 1008)	2016	156	12	b
Suz:2	Suz	Suz	(1, 1)	(1320, 1320)	2640	378	54	a
Suz:2	Suz	Suz	(1, 1)	(2800, 2800)	5600	3564	2268	a
Suz:2	Suz	Suz	(1, 1)	(2912, 2912)	5824	648	72	a
Suz:2	Suz	Suz	(1, 1)	(4032, 4032)	8064	2200	600	a
Suz:2	Suz	Suz	(1, 1)	(4050, 4050)	8100	624	48	a
Suz:2	Suz	Suz	(1, 1)	(4752, 4752)	9504	2925	900	a
Suz:2	Suz	Suz	(1, 1)	(6804, 6804)	13608	2475	450	a
Suz:2	Suz	Suz	(1, 1)	(7040, 7040)	14080	12636	11340	a
Suz:2	Suz	Suz	(1, 1)	(7488, 7488)	14976	600	24	b
Suz:2	Suz	Suz	(1, 1)	(10368, 10368)	20736	10296	5112	a
Suz:2	Suz	Suz	(1, 1)	(11200, 11200)	22400	18954	16038	a
Suz:2	Suz	Suz	(1, 1)	(25344, 25344)	50688	3900	300	a
Suz:2	Suz	Suz	(1, 1)	(98560, 98560)	197120	75816	29160	a
O'N:2	O'N	O'N	(1, 1)	(8, 8)	16	6	2	b
O'N:2	O'N	O'N	(1, 1)	(20, 20)	40	27	18	a
O'N:2	O'N	O'N	(1, 1)	(32, 32)	64	28	12	a
O'N:2	O'N	O'N	(1, 1)	(33, 33)	66	40	24	a
O'N:2	O'N	O'N	(1, 1)	(44, 44)	88	30	10	a
O'N:2	O'N	O'N	(1, 1)	(48, 48)	96	20	4	b
O'N:2	O'N	O'N	(1, 1)	(62, 62)	124	42	14	a
O'N:2	O'N	O'N	(1, 1)	(72, 72)	144	66	30	a
O'N:2	O'N	O'N	(1, 1)	(80, 80)	160	54	18	a
O'N:2	O'N	O'N	(1, 1)	(88, 88)	176	126	90	a
O'N:2	O'N	O'N	(1, 1)	(98, 98)	196	66	22	a
O'N:2	O'N	O'N	(1, 1)	(110, 110)	220	147	98	a
O'N:2	O'N	O'N	(1, 1)	(128, 128)	256	171	114	a
O'N:2	O'N	O'N	(1, 1)	(140, 140)	280	63	14	a
O'N:2	O'N	O'N	(1, 1)	(144, 144)	288	42	6	b
O'N:2	O'N	O'N	(1, 1)	(165, 165)	330	189	108	a
O'N:2	O'N	O'N	(1, 1)	(168, 168)	336	135	54	a
O'N:2	O'N	O'N	(1, 1)	(198, 198)	396	80	16	a
O'N:2	O'N	O'N	(1, 1)	(209, 209)	418	279	186	a
O'N:2	O'N	O'N	(1, 1)	(228, 228)	456	196	84	a
O'N:2	O'N	O'N	(1, 1)	(256, 256)	512	147	42	a
O'N:2	O'N	O'N	(1, 1)	(320, 320)	640	72	8	b
O'N:2	O'N	O'N	(1, 1)	(341, 341)	682	228	76	a
O'N:2	O'N	O'N	(1, 1)	(392, 392)	784	378	182	a
O'N:2	O'N	O'N	(1, 1)	(440, 440)	880	294	98	a
O'N:2	O'N	O'N	(1, 1)	(448, 448)	896	180	36	a
O'N:2	O'N	O'N	(1, 1)	(480, 480)	960	686	490	a
O'N:2	O'N	O'N	(1, 1)	(539, 539)	1078	360	120	a
O'N:2	O'N	O'N	(1, 1)	(620, 620)	1240	532	228	a

Continued

TABLE 7 - continued

G	M	N	$(nr(M), nr(N))$	(i_H, i_K)	v	k	λ	Type
O'N:2	O'N	O'N	(1, 1)	(627, 627)	1254	896	640	a
O'N:2	O'N	O'N	(1, 1)	(770, 770)	1540	1216	960	a
O'N:2	O'N	O'N	(1, 1)	(798, 798)	1596	320	64	a
O'N:2	O'N	O'N	(1, 1)	(836, 836)	1672	558	186	a
O'N:2	O'N	O'N	(1, 1)	(992, 992)	1984	1323	882	a
O'N:2	O'N	O'N	(1, 1)	(1008, 1008)	2016	651	210	a
O'N:2	O'N	O'N	(1, 1)	(1026, 1026)	2052	294	42	a
O'N:2	O'N	O'N	(1, 1)	(1178, 1178)	2356	315	42	a
O'N:2	O'N	O'N	(1, 1)	(1320, 1320)	2640	378	54	a
O'N:2	O'N	O'N	(1, 1)	(1568, 1568)	3136	210	14	b
O'N:2	O'N	O'N	(1, 1)	(1568, 1568)	3136	1540	756	a
O'N:2	O'N	O'N	(1, 1)	(2090, 2090)	4180	1792	768	a
O'N:2	O'N	O'N	(1, 1)	(2790, 2790)	5580	798	114	a
O'N:2	O'N	O'N	(1, 1)	(3840, 3840)	7680	4389	2508	a
O'N:2	O'N	O'N	(1, 1)	(3968, 3968)	7936	2646	882	a
O'N:2	O'N	O'N	(1, 1)	(4480, 4480)	8960	868	84	a
O'N:2	O'N	O'N	(1, 1)	(6479, 6479)	12958	4320	1440	a
O'N:2	O'N	O'N	(1, 1)	(8085, 8085)	16170	4256	1120	a
O'N:2	O'N	O'N	(1, 1)	(9405, 9405)	18810	2688	384	a
O'N:2	O'N	O'N	(1, 1)	(11904, 11904)	23808	1254	66	a
O'N:2	O'N	O'N	(1, 1)	(25916, 25916)	51832	5760	640	a
O'N:2	O'N	O'N	(1, 1)	(32585, 32585)	65170	50688	39424	a
O'N:2	O'N	O'N	(1, 1)	(42532, 42532)	85064	49248	28512	a
Fi ₂₂	PΩ ₇ (3)	PΩ ₇ (3)	(3, 3)	(14, 14)	197120	75816	29160	a
Fi ₂₂	PΩ ₇ (3)	PΩ ₇ (3)	(3, 2)	(14, 14)	197120	75816	29160	a
Fi ₂₂	PΩ ₇ (3)	PΩ ₇ (3)	(2, 3)	(14, 14)	197120	75816	29160	a
Fi ₂₂	PΩ ₇ (3)	PΩ ₇ (3)	(2, 2)	(14, 14)	197120	75816	29160	a
Fi ₂₂	PΩ ₇ (3)	PΩ ₇ (3)	(3, 3)	(40, 40)	563200	101088	18144	a
Fi ₂₂	PΩ ₇ (3)	PΩ ₇ (3)	(3, 2)	(40, 40)	563200	101088	18144	a
Fi ₂₂	PΩ ₇ (3)	PΩ ₇ (3)	(2, 3)	(40, 40)	563200	101088	18144	a
Fi ₂₂	PΩ ₇ (3)	PΩ ₇ (3)	(2, 2)	(40, 40)	563200	101088	18144	a
Fi ₂₂	PΩ ₇ (3)	PΩ ₇ (3)	(3, 3)	(105, 105)	1478400	113724	8748	a
Fi ₂₂	PΩ ₇ (3)	PΩ ₇ (3)	(3, 2)	(105, 105)	1478400	113724	8748	a
Fi ₂₂	PΩ ₇ (3)	PΩ ₇ (3)	(2, 3)	(105, 105)	1478400	113724	8748	a
Fi ₂₂	PΩ ₇ (3)	PΩ ₇ (3)	(2, 2)	(105, 105)	1478400	113724	8748	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(8, 8)	16	6	2	b
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(20, 20)	40	27	18	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(32, 32)	64	28	12	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(33, 33)	66	40	24	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(44, 44)	88	30	10	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(48, 48)	96	20	4	b
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(56, 56)	112	75	50	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(60, 60)	120	35	10	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(72, 72)	144	66	30	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(80, 80)	160	54	18	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(88, 88)	176	126	90	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(108, 108)	216	130	78	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(128, 128)	256	120	56	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(143, 143)	286	96	32	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(144, 144)	288	42	6	b
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(165, 165)	330	189	108	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(168, 168)	336	135	54	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(182, 182)	364	243	162	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(189, 189)	378	117	36	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(198, 198)	396	80	16	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(224, 224)	448	150	50	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(243, 243)	486	195	78	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(270, 270)	540	99	18	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(273, 273)	546	110	22	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(280, 280)	560	130	30	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(320, 320)	640	72	8	b
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(336, 336)	672	550	450	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(350, 350)	700	234	78	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(448, 448)	896	180	36	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(455, 455)	910	405	180	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(540, 540)	1080	416	160	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(585, 585)	1170	168	24	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(648, 648)	1296	260	52	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(648, 648)	1296	630	306	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(704, 704)	1408	336	80	a

Continued

TABLE 7 - continued

G	M	N	$(nr(M), nr(N))$	(i_H, i_K)	v	k	λ	Type
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(728, 728)	1456	486	162	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(864, 864)	1728	1100	700	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(1008, 1008)	2016	156	12	b
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(1320, 1320)	2640	378	54	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(2048, 2048)	4096	2016	992	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(2800, 2800)	5600	3564	2268	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(2912, 2912)	5824	648	72	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(4032, 4032)	8064	2200	600	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(4050, 4050)	8100	624	48	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(4752, 4752)	9504	2925	900	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(6804, 6804)	13608	2475	450	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(7040, 7040)	14080	12636	11340	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(7488, 7488)	14976	600	24	b
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(10368, 10368)	20736	10296	5112	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(11200, 11200)	22400	18954	16038	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(18480, 18480)	36960	28431	21870	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(25344, 25344)	50688	3900	300	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(52800, 52800)	105600	56862	30618	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(81648, 81648)	163296	2970	54	b
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(98560, 98560)	197120	75816	29160	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(281600, 281600)	563200	101088	18144	a
Fi ₂₂ :2	Fi ₂₂	Fi ₂₂	(1, 1)	(739200, 739200)	1478400	113724	8748	a
HN:2	HN	HN	(1, 1)	(8, 8)	16	6	2	b
HN:2	HN	HN	(1, 1)	(20, 20)	40	27	18	a
HN:2	HN	HN	(1, 1)	(32, 32)	64	28	12	a
HN:2	HN	HN	(1, 1)	(33, 33)	66	40	24	a
HN:2	HN	HN	(1, 1)	(44, 44)	88	30	10	a
HN:2	HN	HN	(1, 1)	(48, 48)	96	20	4	b
HN:2	HN	HN	(1, 1)	(50, 50)	100	45	20	a
HN:2	HN	HN	(1, 1)	(56, 56)	112	75	50	a
HN:2	HN	HN	(1, 1)	(60, 60)	120	35	10	a
HN:2	HN	HN	(1, 1)	(72, 72)	144	66	30	a
HN:2	HN	HN	(1, 1)	(80, 80)	160	54	18	a
HN:2	HN	HN	(1, 1)	(88, 88)	176	126	90	a
HN:2	HN	HN	(1, 1)	(125, 125)	250	84	28	a
HN:2	HN	HN	(1, 1)	(128, 128)	256	120	56	a
HN:2	HN	HN	(1, 1)	(128, 128)	256	171	114	a
HN:2	HN	HN	(1, 1)	(144, 144)	288	42	6	b
HN:2	HN	HN	(1, 1)	(165, 165)	330	189	108	a
HN:2	HN	HN	(1, 1)	(168, 168)	336	135	54	a
HN:2	HN	HN	(1, 1)	(198, 198)	396	80	16	a
HN:2	HN	HN	(1, 1)	(200, 200)	400	190	90	a
HN:2	HN	HN	(1, 1)	(224, 224)	448	150	50	a
HN:2	HN	HN	(1, 1)	(270, 270)	540	99	18	a
HN:2	HN	HN	(1, 1)	(320, 320)	640	72	8	b
HN:2	HN	HN	(1, 1)	(336, 336)	672	550	450	a
HN:2	HN	HN	(1, 1)	(448, 448)	896	180	36	a
HN:2	HN	HN	(1, 1)	(512, 512)	1024	342	114	a
HN:2	HN	HN	(1, 1)	(550, 550)	1100	315	90	a
HN:2	HN	HN	(1, 1)	(600, 600)	1200	110	10	b
HN:2	HN	HN	(1, 1)	(627, 627)	1254	896	640	a
HN:2	HN	HN	(1, 1)	(648, 648)	1296	630	306	a
HN:2	HN	HN	(1, 1)	(704, 704)	1408	336	80	a
HN:2	HN	HN	(1, 1)	(770, 770)	1540	1216	960	a
HN:2	HN	HN	(1, 1)	(798, 798)	1596	320	64	a
HN:2	HN	HN	(1, 1)	(864, 864)	1728	1100	700	a
HN:2	HN	HN	(1, 1)	(960, 960)	1920	304	48	a
HN:2	HN	HN	(1, 1)	(1260, 1260)	2520	1375	750	a
HN:2	HN	HN	(1, 1)	(1320, 1320)	2640	378	54	a
HN:2	HN	HN	(1, 1)	(1458, 1458)	2916	1750	1050	a
HN:2	HN	HN	(1, 1)	(2048, 2048)	4096	2926	2090	a
HN:2	HN	HN	(1, 1)	(2090, 2090)	4180	1792	768	a
HN:2	HN	HN	(1, 1)	(2100, 2100)	4200	2432	1408	a
HN:2	HN	HN	(1, 1)	(2375, 2375)	4750	1584	528	a
HN:2	HN	HN	(1, 1)	(2800, 2800)	5600	3564	2268	a
HN:2	HN	HN	(1, 1)	(3168, 3168)	6336	5250	4350	a
HN:2	HN	HN	(1, 1)	(3240, 3240)	6480	342	18	b
HN:2	HN	HN	(1, 1)	(3840, 3840)	7680	4389	2508	a
HN:2	HN	HN	(1, 1)	(4032, 4032)	8064	2200	600	a
HN:2	HN	HN	(1, 1)	(4400, 4400)	8800	420	20	b

Continued

TABLE 7 - continued

G	M	N	$(nr(M), nr(N))$	(i_H, i_K)	v	k	λ	Type
HN:2	HN	HN	(1, 1)	(4608, 4608)	9216	4560	2256	a
HN:2	HN	HN	(1, 1)	(4950, 4950)	9900	3648	1344	a
HN:2	HN	HN	(1, 1)	(5000, 5000)	10000	4950	2450	a
HN:2	HN	HN	(1, 1)	(6080, 6080)	12160	9072	6768	a
HN:2	HN	HN	(1, 1)	(9405, 9405)	18810	2688	384	a
HN:2	HN	HN	(1, 1)	(12800, 12800)	25600	7315	2090	a
HN:2	HN	HN	(1, 1)	(15120, 15120)	30240	2750	250	a
HN:2	HN	HN	(1, 1)	(15400, 15400)	30800	4864	768	a
HN:2	HN	HN	(1, 1)	(30720, 30720)	61440	8778	1254	a
HN:2	HN	HN	(1, 1)	(51975, 51975)	103950	5472	288	a
HN:2	HN	HN	(1, 1)	(80000, 80000)	160000	79800	39800	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(8, 8)	16	6	2	b
Fi24':2	Fi24'	Fi24'	(1, 1)	(20, 20)	40	27	18	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(32, 32)	64	28	12	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(33, 33)	66	40	24	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(44, 44)	88	30	10	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(48, 48)	96	20	4	b
Fi24':2	Fi24'	Fi24'	(1, 1)	(56, 56)	112	75	50	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(60, 60)	120	35	10	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(65, 65)	130	87	58	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(72, 72)	144	66	30	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(80, 80)	160	54	18	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(88, 88)	176	126	90	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(98, 98)	196	66	22	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(98, 98)	196	91	42	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(108, 108)	216	130	78	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(110, 110)	220	147	98	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(116, 116)	232	78	26	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(128, 128)	256	120	56	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(138, 138)	276	176	112	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(140, 140)	280	63	14	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(143, 143)	286	96	32	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(143, 143)	286	115	46	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(144, 144)	288	42	6	b
Fi24':2	Fi24'	Fi24'	(1, 1)	(153, 153)	306	245	196	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(161, 161)	322	108	36	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(162, 162)	324	153	72	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(165, 165)	330	189	108	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(168, 168)	336	135	54	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(182, 182)	364	243	162	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(189, 189)	378	117	36	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(198, 198)	396	80	16	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(200, 200)	400	343	294	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(204, 204)	408	297	216	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(224, 224)	448	150	50	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(243, 243)	486	195	78	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(253, 253)	506	405	324	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(256, 256)	512	147	42	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(260, 260)	520	174	58	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(270, 270)	540	99	18	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(273, 273)	546	110	22	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(275, 275)	550	306	170	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(280, 280)	560	130	30	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(288, 288)	576	276	132	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(312, 312)	624	357	204	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(315, 315)	630	408	264	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(320, 320)	640	72	8	b
Fi24':2	Fi24'	Fi24'	(1, 1)	(336, 336)	672	550	450	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(348, 348)	696	140	28	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(350, 350)	700	234	78	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(368, 368)	736	540	396	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(377, 377)	754	252	84	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(391, 391)	782	143	26	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(392, 392)	784	378	182	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(408, 408)	816	490	294	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(425, 425)	850	567	378	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(440, 440)	880	294	98	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(448, 448)	896	180	36	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(455, 455)	910	405	180	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(480, 480)	960	686	490	a

Continued

TABLE 7 - continued

G	M	N	$(nr(M), nr(N))$	(i_H, i_K)	v	k	λ	Type
Fi24':2	Fi24'	Fi24'	(1, 1)	(483, 483)	966	580	348	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(506, 506)	1012	675	450	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(522, 522)	1044	448	192	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(539, 539)	1078	360	120	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(540, 540)	1080	416	160	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(575, 575)	1150	384	128	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(585, 585)	1170	168	24	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(598, 598)	1196	240	48	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(644, 644)	1288	352	96	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(648, 648)	1296	260	52	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(648, 648)	1296	630	306	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(680, 680)	1360	756	420	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(704, 704)	1408	336	80	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(704, 704)	1408	805	460	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(725, 725)	1450	162	18	c
Fi24':2	Fi24'	Fi24'	(1, 1)	(728, 728)	1456	486	162	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(735, 735)	1470	1131	870	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(748, 748)	1496	300	60	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(782, 782)	1564	522	174	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(816, 816)	1632	700	300	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(840, 840)	1680	1242	918	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(864, 864)	1728	1100	700	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(891, 891)	1782	1508	1276	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(900, 900)	1800	1029	588	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(910, 910)	1820	1071	630	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(918, 918)	1836	735	294	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(928, 928)	1856	1485	1188	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(935, 935)	1870	624	208	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(952, 952)	1904	693	252	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1008, 1008)	2016	156	12	b
Fi24':2	Fi24'	Fi24'	(1, 1)	(1040, 1040)	2080	595	170	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1078, 1078)	2156	1725	1380	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1080, 1080)	2160	255	30	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1218, 1218)	2436	975	390	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1242, 1242)	2484	1911	1470	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1274, 1274)	2548	567	126	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1320, 1320)	2640	378	54	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1386, 1386)	2772	816	240	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1392, 1392)	2784	2300	1900	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1518, 1518)	3036	1215	486	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1568, 1568)	3136	210	14	b
Fi24':2	Fi24'	Fi24'	(1, 1)	(1568, 1568)	3136	286	26	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1568, 1568)	3136	1540	756	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1600, 1600)	3200	1372	588	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1638, 1638)	3276	525	84	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1700, 1700)	3400	1134	378	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1755, 1755)	3510	2784	2208	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1760, 1760)	3520	783	174	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1840, 1840)	3680	2548	1764	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1856, 1856)	3712	2475	1650	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1944, 1944)	3888	3381	2940	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2024, 2024)	4048	1350	450	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2048, 2048)	4096	2016	992	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2100, 2100)	4200	2262	1218	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2304, 2304)	4608	272	16	b
Fi24':2	Fi24'	Fi24'	(1, 1)	(2349, 2349)	4698	672	96	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2415, 2415)	4830	440	40	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2448, 2448)	4896	891	162	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2448, 2448)	4896	980	196	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2450, 2450)	4900	4048	3344	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2464, 2464)	4928	2275	1050	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2496, 2496)	4992	714	102	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2548, 2548)	5096	1020	204	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2639, 2639)	5278	3519	2346	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2646, 2646)	5292	1925	700	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2668, 2668)	5336	4851	4410	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2695, 2695)	5390	952	168	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2737, 2737)	5474	2106	810	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2772, 2772)	5544	483	42	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2800, 2800)	5600	3564	2268	a

Continued

TABLE 7 - continued

G	M	N	$(nr(M), nr(N))$	(i_H, i_K)	v	k	λ	Type
Fi24':2	Fi24'	Fi24'	(1, 1)	(2898, 2898)	5796	1160	232	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2912, 2912)	5824	648	72	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2958, 2958)	5916	3381	1932	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2975, 2975)	5950	1323	294	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(3036, 3036)	6072	468	36	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(3120, 3120)	6240	1836	540	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(3128, 3128)	6256	5005	4004	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(3213, 3213)	6426	1800	504	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(3248, 3248)	6496	1300	260	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(3248, 3248)	6496	3465	1848	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(3315, 3315)	6630	2842	1218	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(3400, 3400)	6800	2093	644	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(3542, 3542)	7084	1575	350	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(3672, 3672)	7344	1050	150	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(3770, 3770)	7540	1078	154	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(3920, 3920)	7840	3016	1160	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(4032, 4032)	8064	2200	600	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(4048, 4048)	8096	1620	324	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(4050, 4050)	8100	624	48	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(4508, 4508)	9016	8415	7854	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(4752, 4752)	9504	2925	900	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(4784, 4784)	9568	8505	7560	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(4862, 4862)	9724	9261	8820	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(5148, 5148)	10296	1421	196	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(5440, 5440)	10880	990	90	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(5568, 5568)	11136	4455	1782	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(5632, 5632)	11264	1610	230	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(5670, 5670)	11340	783	54	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(5684, 5684)	11368	2106	390	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(5831, 5831)	11662	3888	1296	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(5880, 5880)	11760	5346	2430	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(6318, 6318)	12636	6860	3724	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(6468, 6468)	12936	5175	2070	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(6624, 6624)	13248	5096	1960	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(6800, 6800)	13600	1512	168	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(6804, 6804)	13608	2475	450	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(6912, 6912)	13824	9016	5880	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(7040, 7040)	14080	12636	11340	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(7200, 7200)	14400	2058	294	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(7200, 7200)	14400	5083	1794	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(7424, 7424)	14848	4950	1650	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(7488, 7488)	14976	600	24	b
Fi24':2	Fi24'	Fi24'	(1, 1)	(7820, 7820)	15640	2808	504	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(8096, 8096)	16192	1800	200	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(8096, 8096)	16192	6426	2550	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(8262, 8262)	16524	11440	7920	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(8280, 8280)	16560	7424	3328	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(8330, 8330)	16660	4320	1120	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(8568, 8568)	17136	3726	810	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(8613, 8613)	17226	1326	102	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(9240, 9240)	18480	1088	64	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(9315, 9315)	18630	5733	1764	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(9860, 9860)	19720	5635	1610	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(10005, 10005)	20010	14553	10584	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(10368, 10368)	20736	10296	5112	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(10556, 10556)	21112	7038	2346	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(11200, 11200)	22400	18954	16038	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(11339, 11339)	22678	7560	2520	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(11760, 11760)	23520	812	28	b
Fi24':2	Fi24'	Fi24'	(1, 1)	(13122, 13122)	26244	13041	6480	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(13524, 13524)	27048	7956	2340	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(14144, 14144)	28288	24696	21560	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(14168, 14168)	28336	1890	126	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(14848, 14848)	29696	5940	1188	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(15708, 15708)	31416	18850	11310	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(15778, 15778)	31556	25245	20196	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(17248, 17248)	34496	6900	1380	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(17408, 17408)	34816	1056	32	b
Fi24':2	Fi24'	Fi24'	(1, 1)	(18480, 18480)	36960	28431	21870	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(18768, 18768)	37536	15015	6006	a

Continued

TABLE 7 - continued

G	M	N	$(nr(M), nr(N))$	(i_H, i_K)	v	k	λ	Type
Fi24':2	Fi24'	Fi24'	(1, 1)	(19136, 19136)	38272	25515	17010	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(19575, 19575)	39150	32032	26208	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(19712, 19712)	39424	14688	5472	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(20735, 20735)	41470	13824	4608	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(20800, 20800)	41600	2448	144	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(21879, 21879)	43758	35721	29160	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(22050, 22050)	44100	8019	1458	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(22100, 22100)	44200	31104	21888	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(22880, 22880)	45760	37044	29988	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(23324, 23324)	46648	5184	576	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(23664, 23664)	47328	6762	966	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(24990, 24990)	49980	4347	378	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(25344, 25344)	50688	3900	300	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(25578, 25578)	51156	29120	16576	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(29400, 29400)	58800	4524	348	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(29988, 29988)	59976	2400	96	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(31556, 31556)	63112	42075	28050	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(32016, 32016)	64032	29106	13230	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(32292, 32292)	64584	11136	1920	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(32640, 32640)	65280	18009	4968	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(32768, 32768)	65536	6426	630	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(33408, 33408)	66816	11040	1824	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(33534, 33534)	67068	7371	810	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(34496, 34496)	68992	44226	28350	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(35343, 35343)	70686	28275	11310	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(35360, 35360)	70720	36432	18768	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(38896, 38896)	77792	55566	39690	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(39100, 39100)	78200	49764	31668	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(41216, 41216)	82432	39690	19110	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(45356, 45356)	90712	10080	1120	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(46656, 46656)	93312	20286	4410	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(49680, 49680)	99360	7644	588	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(50048, 50048)	100096	20020	4004	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(50600, 50600)	101200	9639	918	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(51968, 51968)	103936	6930	462	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(52800, 52800)	105600	56862	30618	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(57600, 57600)	115200	2352	48	b
Fi24':2	Fi24'	Fi24'	(1, 1)	(59840, 59840)	119680	74088	45864	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(61600, 61600)	123200	72471	42630	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(62640, 62640)	125280	37422	11178	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(66528, 66528)	133056	4095	126	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(71442, 71442)	142884	71253	35532	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(72128, 72128)	144256	67320	31416	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(72930, 72930)	145860	83349	47628	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(76544, 76544)	153088	51030	17010	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(79968, 79968)	159936	126846	100602	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(81648, 81648)	163296	2970	54	b
Fi24':2	Fi24'	Fi24'	(1, 1)	(82110, 82110)	164220	74646	33930	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(82940, 82940)	165880	18432	2048	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(90160, 90160)	180320	10608	624	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(94248, 94248)	188496	37700	7540	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(94668, 94668)	189336	75735	30294	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(98560, 98560)	197120	75816	29160	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(105705, 105705)	211410	96096	43680	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(119340, 119340)	238680	107163	48114	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(120060, 120060)	240120	43659	7938	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(126224, 126224)	252448	84150	28050	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(133952, 133952)	267904	59535	13230	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(135168, 135168)	270336	4160	64	b
Fi24':2	Fi24'	Fi24'	(1, 1)	(138600, 138600)	277200	85293	26244	a
Fi24':2	Fi24'	$2x\text{Fi}23$	(1, 2)	(153468, 1)	306936	242880	192192	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(153468, 153468)	306936	242880	192192	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(156800, 156800)	313600	9504	288	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(156800, 156800)	313600	88452	24948	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(165888, 165888)	331776	165600	82656	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(188370, 188370)	376740	12992	448	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(188416, 188416)	376832	368550	360450	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(211200, 211200)	422400	124236	36540	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(212992, 212992)	425984	351900	290700	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(216384, 216384)	432768	253692	148716	a

Continued

TABLE 7 - continued

G	M	N	$(nr(M), nr(N))$	(i_H, i_K)	v	k	λ	Type
Fi24':2	Fi24'	Fi24'	(1, 1)	(223074, 223074)	446148	34320	2640	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(234900, 234900)	469800	128128	34944	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(252448, 252448)	504896	100980	20196	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(269280, 269280)	538560	142884	37908	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(281600, 281600)	563200	101088	18144	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(306176, 306176)	612352	68040	7560	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(311168, 311168)	622336	148176	35280	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(328440, 328440)	656880	317115	153090	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(348160, 348160)	696320	24012	828	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(358904, 358904)	717808	135240	25480	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(367488, 367488)	734976	29400	1176	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(381024, 381024)	762048	420992	232576	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(384192, 384192)	768384	49896	3240	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(401408, 401408)	802816	255255	81158	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(583440, 583440)	1166880	166698	23814	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(739200, 739200)	1478400	113724	8748	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(852992, 852992)	1705984	451008	119232	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(853760, 853760)	1707520	51744	1568	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1085238, 1085238)	2170476	929475	398034	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1330056, 1330056)	2660112	809600	246400	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1378080, 1378080)	2756160	500480	90880	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(1379840, 1379840)	2759680	117936	5040	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2125764, 2125764)	4251528	2825550	1877850	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(2908224, 2908224)	5816448	1614600	448200	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(3503360, 3503360)	7006720	563760	45360	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(4138344, 4138344)	8276688	570807	39366	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(4667520, 4667520)	9335040	190512	3888	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(8977878, 8977878)	17955756	1092960	66528	a
Fi24':2	Fi24'	Fi24'	(1, 1)	(11351340, 11351340)	22702680	3403701	510300	a

SEYED HASSAN ALAVI, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, BU-ALI SINA UNIVERSITY, HAMEDAN, IRAN.

Email address: alavi.s.hassan@basu.ac.ir and alavi.s.hassan@gmail.com

ASHRAF DANESHKHAH, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, BU-ALI SINA UNIVERSITY, HAMEDAN, IRAN.

Email address: adanesh@basu.ac.ir and daneshkhah.ashraf@gmail.com