A convex optimization approach to the Lyapunov exponents

Christoph Kawan*

Abstract

The aim of this paper is to shed more light on some recent ideas about Lyapunov exponents and clarify the formal structures behind these ideas. In particular, we show that the vector of (averaged) Lyapunov exponents of a smooth measure-preserving dynamical system can be regarded as the solution of a vector-valued optimization problem on a space of Riemannian metrics. This result was first formulated and proved by Jairo Bochi in the language of linear cocycles and their conjugacies. We go a step further and prove that the optimization problem is geodesically convex with respect to the L^2 -metric. Moreover, we derive some consequences of this fact.

Keywords— Lyapunov exponents; Multiplicative Ergodic Theorem; Convex optimization; Space of Riemannian metrics

1 Introduction

Originally introduced as stability indicators for ordinary differential equations, Lyapunov exponents have become a cornerstone of the theory of smooth dynamical systems. The main reason is that they are related in numerous ways to other important characteristics of dynamical systems such as several notions of entropy and dimensions of invariant sets as well as invariant measures. The study of their properties and methods for their computation is an ongoing endeavor. Already in dimension two, examples of dynamical systems are known whose Lyapunov exponents up to this date cannot be effectively estimated on large subsets of the state space. For instance, for the Chirikov standard map, an area-preserving diffeomorphism of the 2-torus, it is a long-standing open problem to determine whether the maximal Lyapunov exponent is positive on a set of positive Lebesgue measure, see e.g. [2].

Lyapunov's first method for proving the asymptotic stability of a system consists in showing that the maximal Lyapunov exponent is negative. His second method uses a non-negative real-valued function on the state space with values decreasing along trajectories. Once a candidate for such a *Lyapunov function* has been found, the decrease condition can be checked by a simple computation involving only the candidate function and the right-hand side of the equation which defines the system.

^{*}Email: christoph.kawan@gmx.de

One benefit of this method thus consists in the reduction of the proof of a long-term property to a short-term computation.

Now, asymptotic stability is a comparatively simple property. Other properties, studied mainly in chaos theory, such as various forms of hyperbolicity, can most likely not be checked by evaluating a single scalar-valued function along trajectories. Since these properties are by nature multi-dimensional as their definitions involve a distinction of different state space directions, a structure which is itself multi-dimensional can help to verify them via simple short-term computations. The main example of such a structure is a Riemannian metric. Usually, a Riemannian metric which allows to verify a long-term property by a short-term computation is called an *adapted metric*. For instance, a classical result states that any uniformly hyperbolic set of a discrete-time system admits an adapted metric (also called a Lyapunov metric) in which contraction on the stable bundle and expansion on the unstable bundle can be seen in one time step, see e.g. [18, Ch. 6]. An analogous result for the more general property of supporting a dominated splitting is proved in [17]. In the context of non-uniformly hyperbolic systems, measurable adapted metrics can be constructed, see [19, p. 668]. Finally, in the context of stability analysis, besides Lyapunov functions, also Riemannian metrics are used to prove contractivity properties (implying asymptotic stability), which are known as *contraction metrics*, see, e.g., the recent survey [14].

In [3, 4]. Bochi and Navas introduced a novel method to construct Riemannian metrics adapted to continuous linear cocycles over topological dynamical systems. In this general context, they are not Riemannian metrics in the actual sense (as there is no smooth manifold involved), but rather continuous mappings assigning a positivedefinite matrix to each point in the state space. They can then be interpreted as conjugacies between cocycles. In the special case of the derivative cocycle of a smooth map, they become Riemannian metrics in the actual sense, which allow to approximate the average Lyapunov vector (with respect to an invariant probability) by integration over the vector of (log-) singular values of the right-hand side Jacobian. In this paper, we provide a self-contained proof of Bochi's result in this context. An important interpretation of the result is that the averaged Lyapunov vector is the infimum of a vector-valued function \mathcal{F} defined on the space of all continuous Riemannian metrics. Here, the infimum needs to be understood with respect to the so-called majorization order. It is important to note that it is not only a weak, i.e. a Pareto infimum, but an infimum in all components. Hence, the averaged Lyapunov vector appears as the solution of a vector-valued optimization problem.

We further prove that the optimization problem posed for \mathcal{F} is convex in a proper sense. First, one can restrict \mathcal{F} to the space \mathcal{M} of smooth Riemannian metrics which is a Fréchet manifold modeled on the Fréchet space of smooth symmetric (0, 2)-tensor fields. The most-studied Riemannian metric on this manifold is the L^2 -metric, which is defined by integration over the fiberwise trace metric (the metric of the symmetric space of positive-definite matrices). The L^2 -metric on \mathcal{M} has been studied for a long time in different contexts; the main references include [12, 13, 16, 9, 10, 7]. Though \mathcal{M} is only a weak Riemannian manifold, the central objects studied in Riemannian geometry (including geodesics) can be shown to exist and explicit formulas for them are available. An important submanifold of \mathcal{M} is the space \mathcal{M}_{ω} of metrics inducing the same (arbitrary) volume form ω . It was proved in [13] that \mathcal{M}_{ω} is a symmetric space with the inherited L^2 -metric; in particular, it is geodesically complete. The function \mathcal{F} has the property that its infimum is preserved under restriction to \mathcal{M}_{ω} . Hence, the search for a (nearly) optimal metric can be restricted to \mathcal{M}_{ω} . We prove that the restriction of \mathcal{F} to \mathcal{M}_{ω} is geodesically convex with respect to the L^2 -geodesics on its domain and the majorization order on its co-domain. In fact, this proof was already given in [20], but without putting it in the proper context and relating it to the geometry of \mathcal{M} .

In summary, we prove that the averaged Lyapunov vector is the infimum of a geodesically convex vector-valued function. The associated optimization problem has some serious disadvantages, however. It is not guaranteed that a minimizer exists, the problem is infinite-dimensional, defined on a Riemannian manifold (rather than a vector space) and non-smooth. We elaborate on some of these aspects and make suggestions how to mitigate the corresponding problems.

Several other works have proposed convex optimization to compute Lyapunov exponents or associated quantities. In particular, we point the reader to the paper [27], where SOS programming is used to find tight bounds on the maximal Lyapunov exponent of a system given by an ordinary differential equation whose right-hand side is polynomial. The optimization there is not over Riemannian metrics, but over auxiliary real-valued functions. We also mention the papers [15, 20, 24], in which the groundwork has been laid for the approach pursued in this paper, and numerical algorithms were developed to use it for the computation of contraction metrics [15], the estimation of Lyapunov dimension [24] and the estimation of restoration entropy [20].

Organization of the paper: In Section 2, we provide the necessary background on Lyapunov exponents. Section 3 is devoted to the proof of Bochi's result on the approximation of the Lyapunov vector by singular values of the time-one Jacobian, see Theorem 2. In Section 4, we introduce the L^2 -metric on the space of Riemannian metrics and present some results on the structure of this space. The subsequent Section 5 contains the convexity proof of the function \mathcal{F} , Theorem 3. In Theorem 4, we also prove the global Lipschitz continuity of the components of \mathcal{F} . In Section 6, we provide a formula for the directional derivatives of the components of \mathcal{F} in an idealized case, where the existence of these derivatives is guaranteed, see Theorem 5. The final Section 7 provides a summary and an outlook.

2 Preliminaries

2.1 Notation

By $\log(\cdot)$, we denote the natural logarithm. If A is a square matrix, we write A^{\top} and $\operatorname{tr}(A)$ for the transpose and the trace of A, respectively. If $x \in \mathbb{R}^d$, then $||x||_p$ denotes its p-norm for $p \in \{1, 2, \infty\}$. The components of x will always be denoted by x_1, \ldots, x_d . By 1, we denote the vector in \mathbb{R}^d whose components are all equal to 1. We write $\operatorname{GL}(d,\mathbb{R})$ for the general linear group of \mathbb{R}^d . If $f: M \leftarrow i$ is a differentiable self-map of a smooth manifold M, we write fx and df_x for its value and its derivative at $x \in M$, respectively. By I, we denote the identity matrix of any dimension. By a metric g on a smooth manifold M, we always mean a Riemannian metric, i.e. the assignment of an inner product g_x to each tangent space T_xM , which is at least continuous. We use the notation \mathcal{M}^0 (\mathcal{M}) for the space of all continuous (smooth) Riemannian metrics on a given manifold M.

2.2 Setup

We consider a discrete dynamical system given by a C^1 -diffeomorphism $f: M \leftrightarrow$ on a compact orientable¹ smooth manifold M of finite dimension d. We further assume that f preserves a Borel probability measure μ , i.e. $\mu(f^{-1}(A)) = \mu(A)$ for all Borel sets $A \subseteq M$.

2.3 Background on Lyapunov exponents

Our objects of interest are the Lyapunov exponents associated with the data (M, f, μ) . The existence of these numbers is guaranteed by the famous *Multiplicative Ergodic Theorem (MET)*, also known as *Oseledets Theorem*. This theorem includes quite a number of statements and can be formulated on different levels of generality. Here, we formulate a version that is adapted to our setup and our needs. First, we define the *Lyapunov exponent* at a point $x \in M$ in direction $0 \neq v \in T_x M$ as the limit

$$\lambda(x,v) := \lim_{n \to \infty} \frac{1}{n} \log \| \mathrm{d} f_x^n v \|,$$

provided that this limit exists. Here, $\|\cdot\|$ denotes the norm (on each tangent space) induced by a given Riemannian metric. The value of $\lambda(x, v)$, however, is independent of the chosen metric as a consequence of compactness.

Theorem 1 (MET) In the given setup, for μ -almost every $x \in M$ there exist numbers

$$\lambda_1(x) > \lambda_2(x) > \dots > \lambda_{r(x)}(x)$$

with multiplicities $m_1(x), m_2(x), \ldots, m_{r(x)}(x) \in \mathbb{N}$, respectively, such that the following statements hold:

- (i) For every tangent vector $0 \neq v \in T_x M$, $\lambda(x, v) = \lambda_i(x)$ for some *i*.
- (ii) $\sum_{i} m_i(x) = d$.
- (*iii*) $\sum_{i} m_i(x)\lambda_i(x) = \lim_{n \to \infty} (1/n) \log |\det df_x^n|.$
- (iv) The functions $r(\cdot)$, $m_i(\cdot)$ and $\lambda_i(\cdot)$ are μ -measurable and f-invariant.
- (v) Let $\alpha_1(\mathrm{d} f_x^n) \ge \alpha_2(\mathrm{d} f_x^n) \ge \cdots \ge \alpha_d(\mathrm{d} f_x^n)$ denote the singular values of the linear operator $\mathrm{d} f_x^n : T_x M \to T_{f^n x} M$. Then the limits

$$\lim_{n \to \infty} \frac{1}{n} \log \alpha_i(\mathrm{d} f_x^n), \quad i = 1, 2, \dots, d,$$

exist and each of them equals one of the numbers $\lambda_1(x), \ldots, \lambda_{r(x)}(x)$.

If μ is ergodic, then the functions $r(\cdot)$, $m_i(\cdot)$ and $\lambda_i(\cdot)$ are constant almost everywhere.

We base the analysis in this paper on statement (v) above, i.e. on the relation between the Lyapunov exponents and the growth rates of singular values. In contrast to eigenvalues, singular values are geometric quantities; they depend on the inner products on domain and co-domain of the linear operator in question. In our setup, this means

 $^{^1\}mathrm{The}$ orientability of M is a technical assumption which guarantees that we can work with volume forms.

that they depend on the Riemannian metric imposed on M, while their exponential growth rates, the Lyapunov exponents, do not. Hence, one can ask whether there exists a metric adapted to the given system such that in this metric the Lyapunov exponents can be computed (at least approximately) as the expansions rates in the first iterate, i.e.²

 $\lambda_i(x) \approx \log \alpha_i(\mathrm{d}f_x).$

The existence of such metrics follows from a result proved in [3, Prop. 4.1] about linear cocycles over topological dynamical systems. In the language of linear cocycles, switching to another metric corresponds to switching to a conjugate cocycle. For the convenience of the reader, we provide a complete proof of the result within the setup introduced above.

3 Bochi's result

We first need to introduce some technical concepts and the associated notation. Given an invertible linear operator $L: V \to W$ between finite-dimensional inner product spaces of dimension d, we write

$$\alpha_1(L) \ge \alpha_2(L) \ge \dots \ge \alpha_d(L)$$

for the singular values of L. Geometrically, these are the lengths of the semi-axes of the ellipsoid $\mathcal{E} := \{Lv : \langle v, v \rangle_V = 1\}$ measured in the norm induced by the inner product on W. We then write

$$\vec{\sigma}(L) := (\log \alpha_1(L), \log \alpha_2(L), \dots, \log \alpha_d(L))$$

By our assumption that L is invertible, the numbers $\alpha_i(L)$ are strictly positive, and hence $\vec{\sigma}(L) \in \mathbb{R}^d$. We can be more precise. Because of the ordering of the singular values, $\vec{\sigma}$ takes values in the closed convex cone³

$$\mathfrak{a}^+ := \{\xi \in \mathbb{R}^d : \xi_1 \ge \xi_2 \ge \cdots \ge \xi_d\}$$

A partial order on \mathfrak{a}^+ , which is very useful for our purposes, is defined by

$$\xi \preceq \eta \quad :\Leftrightarrow \quad \left\{ \begin{array}{ll} \xi_1 + \dots + \xi_k \leq \eta_1 + \dots + \eta_k & \text{for } k = 1, \dots, d-1, \\ \xi_1 + \dots + \xi_k = \eta_1 + \dots + \eta_k & \text{for } k = d. \end{array} \right.$$

This order is called the *majorization order*. It can be shown that it is the partial order induced by the dual cone of \mathfrak{a}^+ . If we relax the equality in the case k = d to an inequality, we obtain the *weak majorization order* that we denote by \leq_w . We refer to the book [26] for more information about majorization.

In terms of the order \leq , we can formulate *Horn's inequality* (see [5, Prop. I.7.4.3]) equivalently as

$$\vec{\sigma}(L_1 L_2) \preceq \vec{\sigma}(L_1) + \vec{\sigma}(L_2) \tag{1}$$

for any linear operators L_1 and L_2 that can be composed in the suggested way.

²Here, we allow multiple repeated values in the enumeration of the Lyapunov exponents.

 $^{^3 \}rm The notation \, \mathfrak{a}^+$ is adopted from the theory of semisimple Lie groups, where it is used for the so-called positive Weyl chamber.

Now, consider again our dynamical system (M, f). Given any Riemannian metric g on M, the linear operator $df_x^n : T_x M \to T_{f^n x} M$ has unique singular values with respect to the inner products g_x and $g_{f^n x}$. To highlight the dependence of these singular values on the metric g, we write $\vec{\sigma}^g(df_x^n) = \vec{\sigma}(df_x^n)$. By the MET, for μ -almost every $x \in M$, the Lyapunov vector

$$\vec{\lambda}(x) := \lim_{n \to \infty} \frac{1}{n} \vec{\sigma}^g(\mathrm{d} f_x^n) \in \mathfrak{a}^+$$

exists as a limit and equals the corresponding vector of Lyapunov exponents:

$$\dot{\lambda}(x) = (\underbrace{\lambda_1(x), \dots, \lambda_1(x)}_{m_1(x)}, \underbrace{\lambda_2(x), \dots, \lambda_2(x)}_{m_2(x)}, \dots, \underbrace{\lambda_{r(x)}(x), \dots, \lambda_{r(x)}(x)}_{m_{r(x)}}).$$

Observe that the almost-everyhwere existence of $\vec{\lambda}(x)$ already follows from Kingman's subadditive ergodic theorem, the subadditivity being a consequence of inequality (1). We also introduce the average over the Lyapunov vector

$$\vec{\lambda}(f) := \int \vec{\lambda}(x) \,\mathrm{d}\mu(x).$$

Observe that in the ergodic case the integral equals the integrand almost everywhere. As the Lyapunov exponents depend measurably on x, and the compactness of M together with the continuity of df easily implies their boundedness, the integral exists.

We now prove our first result.

_

Proposition 1 For every $g \in \mathcal{M}^0$, the following inequality holds:

$$\vec{\lambda}(f) \preceq \int \vec{\sigma}^g(\mathrm{d}f_x) \,\mathrm{d}\mu(x).$$

Proof: The inequality to be proven is mainly a consequence of Fekete's subadditivity lemma. For simplicity in notation, we write

$$\lambda_1(x) \ge \lambda_2(x) \ge \dots \ge \lambda_d(x)$$

for the Lyapunov exponents of f at x, i.e. we allow repetitions in this enumeration. Then, we also write

$$\lambda_i(f) := \int \lambda_i(x) \,\mathrm{d}\mu(x), \quad i = 1, 2, \dots, d.$$

With this notation at hand, the inequality to be proven is equivalent to

$$\lambda_1(f) + \dots + \lambda_k(f) \le \int \sum_{i=1}^k \log \alpha_i^g(\mathrm{d}f_x) \,\mathrm{d}\mu(x)$$

for k = 1, 2, ..., d with equality if k = d. For a fixed k, consider the sequence

$$a_n := \int \sum_{i=1}^k \log \alpha_i^g(\mathrm{d} f_x^n) \,\mathrm{d} \mu(x), \quad n = 1, 2, \dots$$

The sequence is subadditive, which easily follows from Horn's inequality and the *f*-invariance of μ . Hence, Fekete's lemma guarantees that a_n/n converges to $\inf\{a_n/n : n \in \mathbb{N}\}$. This implies

$$a_{1} = \int \sum_{i=1}^{k} \log \alpha_{i}^{g}(\mathrm{d}f_{x}) \,\mathrm{d}\mu(x) \ge \lim_{n \to \infty} \frac{1}{n} \int \sum_{i=1}^{k} \log \alpha_{i}^{g}(\mathrm{d}f_{x}^{n}) \,\mathrm{d}\mu(x)$$
$$= \sum_{i=1}^{k} \int \lim_{n \to \infty} \frac{1}{n} \log \alpha_{i}^{g}(\mathrm{d}f_{x}^{n}) \,\mathrm{d}\mu(x) = \sum_{i=1}^{k} \int \lambda_{i}(x) \,\mathrm{d}\mu(x) = \sum_{i=1}^{k} \lambda_{i}(f).$$

Interchanging the order of limit and integral is possible by the dominated convergence theorem, using that

$$\min_{z \in M} \log \alpha_d^g(\mathrm{d}f_z) \le \frac{1}{n} \log \alpha_i^g(\mathrm{d}f_x^n) \le \max_{z \in M} \log \alpha_1^g(\mathrm{d}f_z) \tag{2}$$

for all i, n and x. It remains to show the equality in the case k = d:

$$\sum_{i=1}^{d} \lambda_i(f) = \lim_{n \to \infty} \frac{1}{n} \int \log |\det_g df_x^n| d\mu(x)$$
$$= \lim_{n \to \infty} \frac{1}{n} \int \log \prod_{i=0}^{n-1} |\det_g df_{f^i x}| d\mu(x)$$
$$= \lim_{n \to \infty} \frac{1}{n} \sum_{i=0}^{n-1} \int \log |\det_g df_x| d\mu(x)$$
$$= \int \log |\det_g df_x| d\mu(x) = \int \sum_{i=1}^{d} \log \alpha_i^g (df_x) d\mu(x).$$

The first of these equalities follows from statement (iii) of the MET, the second is the multiplicativity of the determinant together with the chain rule, the third uses the invariance of the measure μ , the fourth is obvious, and the last one uses that the absolute determinant equals the product of the singular values.

Our next goal is to prove a converse result showing that $\vec{\lambda}(f)$ can be approximated by vectors of the form $\int \vec{\sigma}^g(df_x) d\mu(x)$. This result involves the construction of Riemannian metrics used to achieve the desired approximation. For this construction, we use geometric features of the space S^+ of positive-definite matrices. A good introduction to positive-definite matrices and the geometry of S^+ is the book [1]. In the following, we present some basic facts.

The space of symmetric $(d \times d)$ -matrices is $S_d := \{s \in \mathbb{R}^{d \times d} : s = s^{\top}\}$. It is a real vector space of dimension $\frac{d}{2}(d+1)$ and contains the open cone of positive-definite matrices $S_d^+ := \{p \in S_d : p > 0\}$. As an open subset, S_d^+ is a trivial smooth manifold of the same dimension as S_d . Its tangent space at any point $p \in S_d^+$ can be identified canonically with S_d . A very popular Riemannian metric on S_d^+ , useful in many different contexts, is the *trace metric*, given by

$$\langle v, w \rangle_p := \operatorname{tr}(p^{-1}vp^{-1}w) \text{ for all } p \in \mathcal{S}_d^+, v, w \in T_p\mathcal{S}_d^+ = \mathcal{S}_d.$$

Equipped with this metric, S_d^+ becomes a complete Riemannian manifold. By [1, Form. (6.14)], an explicit expression for the induced distance function is

$$d(p,q) = \left(\sum_{i=1}^{d} \log^2 \rho_i(p^{-1}q)\right)^{\frac{1}{2}},\tag{3}$$

where $\rho_i(p^{-1}q)$ are the eigenvalues of $p^{-1}q$ (which is similar to $p^{-\frac{1}{2}}qp^{-\frac{1}{2}} \in \mathcal{S}_d^+$).

For each two points $p, q \in S_d^+$, there is a unique geodesic joining p and q, denoted by $p \#_t q$ and parameterized on [0, 1]. An explicit formula is

$$p \#_t q = p^{\frac{1}{2}} \left(p^{-\frac{1}{2}} q p^{-\frac{1}{2}} \right)^t p^{\frac{1}{2}}, \quad t \in [0, 1].$$

If we instead specify a geodesic in S_d^+ by its initial point p and its initial direction v, an explicit formula is

$$\gamma_{p,v}(t) = p^{\frac{1}{2}} \exp(tp^{-\frac{1}{2}}vp^{-\frac{1}{2}})p^{\frac{1}{2}}, \quad t \in \mathbb{R}.$$

The group $\operatorname{GL}(d,\mathbb{R})$ acts transitively on \mathcal{S}_d^+ by isometries via

$$g * p := gpg^{\top}, \quad g \in \operatorname{GL}(d, \mathbb{R}), \ p \in \mathcal{S}_d^+$$

Another isometry of S_d^+ is the matrix inversion. Using the fact that isometries map geodesics to geodesics, the following identities become obvious:

$$g * (p\#_t q) = (g * p)\#_t (g * q)$$
 and $(p\#_t q)^{-1} = p^{-1}\#_t q^{-1}$. (4)

Another useful concept for the analysis on \mathcal{S}_d^+ is a vectorial distance given by 4

$$\vec{d}(p,q) := 2\vec{\sigma}(p^{-\frac{1}{2}}q^{\frac{1}{2}}), \quad p,q \in \mathcal{S}_d^+.$$

The vectorial distance is closely related to the distance (3) induced by the trace metric. The next proposition summarizes some of its properties, from which we will only use (i) and (ii). See also [3, Sec. 4].

Proposition 2 The vectorial distance $\vec{d}(\cdot, \cdot) : S_d^+ \times S_d^+ \to \mathfrak{a}^+$ has the following properties:

- (i) $\vec{d}(I,p) = \vec{\sigma}(p)$ and $\|\vec{d}(p,q)\|_2 = d(p,q)$ for all $p,q \in S_d^+$.
- (ii) $\vec{d}(p_1, q_1) = \vec{d}(p_2, q_2)$ if and only if there exists $g \in GL(d, \mathbb{R})$ with $g * p_1 = p_2$ and $g * q_1 = q_2$.
- (iii) $\vec{d}(p,q) \preceq \vec{d}(p,r) + \vec{d}(r,q)$ for all $p,q,r \in \mathcal{S}_d^+$.
- (*iv*) $\vec{d}(q,p) = i(\vec{d}(p,q))$ with $i(\xi) := -(\xi_d, \xi_{d-1}, \dots, \xi_1)$ for all $p, q \in \mathcal{S}_d^+$.

Another important ingredient for the proof of the converse of Proposition 1 is the following concept. The *barycenter* of l matrices $p_1, \ldots, p_l \in S_d^+$ $(l \in \mathbb{N})$ is defined as

$$\operatorname{bar}(p_1,\ldots,p_l) := \underset{p \in \mathcal{S}_d^+}{\operatorname{arg\,min}} \sum_{i=1}^l d(p,p_i)^2.$$

It can be shown that the minimizer exists and is unique, so $bar(p_1, \ldots, p_l)$ is well-defined. The following properties are well-known, see e.g. [21]:

⁴Caution: There is an overload of notation here. We use the letter d for the dimension of M, the distance function on M and the vectorial distance on S_d^+ . It is hopefully always clear from the context what is meant.

- Symmetry: $\operatorname{bar}(p_1,\ldots,p_l) = \operatorname{bar}(p_{\sigma(1)},\ldots,p_{\sigma(l)})$ for any permutation σ .
- For any $g \in \mathrm{GL}(d, \mathbb{R})$,

$$g * \operatorname{bar}(p_1, \ldots, p_l) = \operatorname{bar}(g * p_1, \ldots, g * p_l).$$

• If $u = bar(p_1, \ldots, p_{l-1}, p_l)$ and $v = bar(p_1, \ldots, p_{l-1}, p'_l)$, then

$$\vec{d}(u,v) \leq \frac{1}{l} \vec{d}(p_l, p_l').$$
(5)

• The mapping $(p_1, \ldots, p_l) \mapsto bar(p_1, \ldots, p_l)$ is continuous.

We need to lift some of the constructions introduced above from \mathcal{S}_d^+ to the space \mathcal{M}^0 of continuous metrics on M. If $F: M \leftrightarrow$ is a C^1 -diffeomorphism and $g \in \mathcal{M}^0$, then F acts on g by pulling back:

$$(F * g)_x(v, w) := g_{Fx}(\mathrm{d}F_x v, \mathrm{d}F_x w) \quad \text{for all } v, w \in T_x M.$$

Since F is C^1 , we have $F * g \in \mathcal{M}^0$. We will see below that this can be regarded an an extension of the action of $\operatorname{GL}(d,\mathbb{R})$ on \mathcal{S}_d^+ .

Next, we extend the barycenter map to Riemannian metrics. Given l metrics g^1, \ldots, g^l in \mathcal{M}^0 , we can define a new metric, denoted by $\operatorname{bar}(g^1, \ldots, g^l)$, and defined pointwise via the barycenter of positive-definite matrices. This is done as follows. Let (ϕ, U) be any chart of M around $x \in M$, i.e. $U \subset M$ is an open neighborhood of x and $\phi: U \to \phi(U) \subset \mathbb{R}^d$ a diffeomorphism. Then we identify an element $p \in \mathcal{S}^+_d$ with an inner product on $T_x M$ via⁵

$$\beta_{\phi}(p)(v,w) := \langle p^{-1} \mathrm{d}\phi_x v, \mathrm{d}\phi_x w \rangle \quad \text{for all } v, w \in T_x M, \tag{6}$$

where $\langle \cdot, \cdot \rangle$ denotes the standard Euclidean inner product on \mathbb{R}^d . The map β_{ϕ} is obviously invertible. If (ϕ_1, U) and (ϕ_2, V) are two charts around x, then the two matrix representations of the inner product g_x are related by

$$d(\phi_2 \circ \phi_1^{-1})_{\phi_1(x)} * \beta_{\phi_1}^{-1}(g_x) = \beta_{\phi_2}^{-1}(g_x),$$
(7)

which is an easy consequence of the definition in (6). From this and Proposition 2(ii), it follows that the vectorial distance of two inner products g_x^1 and g_x^2 on $T_x M$ can be defined independently of the chosen chart:

$$ec{d}(g^1_x,g^2_x) := ec{d}(eta^{-1}_\phi(g^1_x),eta^{-1}_\phi(g^2_x)).$$

The barycenter of l metrics g^1, \ldots, g^l is now defined by

$$\operatorname{bar}(g^{1},\ldots,g^{l})_{x} := \beta_{\phi}(\operatorname{bar}(\beta_{\phi}^{-1}(g_{x}^{1}),\ldots,\beta_{\phi}^{-1}(g_{x}^{l}))),$$
(8)

where for each $x \in M$ we choose an appropriate chart (ϕ, U) around x. Using (7), it can easily be shown that this definition is independent of the chosen chart. From (8), it follows that $\operatorname{bar}(g^1, \ldots, g^l)$ is continuous in the domain of every chart, since all involved functions on the right-hand side are continuous. Hence, $\operatorname{bar}(g^1, \ldots, g^l) \in \mathcal{M}^0$.

The proof of the following lemma is trivial, since all statements follow from a direct application of the definitions. Hence, we leave its proof to the reader.

⁵This identification is unusual because of the involved inversion of p. However, for our purposes it is more convenient as it allows to obtain more direct relations between the introduced operations on \mathcal{S}_d^+ and corresponding operations on \mathcal{M}^0 .

Lemma 1 For all $g, g^i \in \mathcal{M}^0$ and C^1 -diffeomorphisms $F, F_i : M \leftrightarrow$, the following properties hold:

(*i*)
$$F * \operatorname{bar}(g^1, \dots, g^l) = \operatorname{bar}(F * g^1, \dots, F * g^l)$$

(*ii*) $F_2 * (F_1 * g) = (F_1 \circ F_2) * g.$

(iii) Let (ϕ, U) be a chart around x and (ψ, V) one around Fx. Then

$$\beta_{\phi}^{-1}((F*g)_x) = d(\psi \circ F \circ \phi^{-1})_{\phi(x)}^{-1} * \beta_{\psi}^{-1}(g_{Fx}).$$

In this sense, the pullback of a metric via a diffeomorphism is an extension of the action of $\operatorname{GL}(d,\mathbb{R})$ on \mathcal{S}_d^+ .

 $(iv) \ \vec{d}(g_{Fx}^1,g_{Fx}^2) = \vec{d}((F*g^1)_x,(F*g^2)_x) \ \text{for all} \ x \in M.$

Lemma 2 Let $F: M \leftrightarrow$ be a C^1 -diffeomorphism and $g \in \mathcal{M}^0$. Then for any $x \in M$ we have

$$\vec{\sigma}^g(\mathrm{d}F_x) = \vec{\sigma}(p_2^{-\frac{1}{2}}A_x p_1^{\frac{1}{2}}),$$

where $p_1 = \beta_{\phi}^{-1}(g_x)$, $p_2 = \beta_{\psi}^{-1}(g_{Fx})$ and $A_x = d(\psi \circ F \circ \phi^{-1})_{\phi(x)}$ for charts (ϕ, U) and (ψ, V) around x and Fx, respectively.

Proof: The singular values of $dF_x : T_x M \to T_{Fx} M$ with respect to the metric g are the square roots of the eigenvalues of the self-adjoint operator

$$(\mathrm{d}F_x)(\mathrm{d}F_x)^*: T_{Fx}M \to T_{Fx}M.$$

Using the charts (ϕ, U) and (ψ, V) , we have for any $v \in T_x M$, $w \in T_{Fx} M$ that

$$g_{Fx}(\mathrm{d}F_{x}v,w) = \langle \beta_{\psi}^{-1}(g_{Fx})^{-1}\mathrm{d}\psi_{Fx}\mathrm{d}F_{x}v,\mathrm{d}\psi_{Fx}w\rangle$$

= $\langle \mathrm{d}\psi_{Fx}\mathrm{d}F_{x}v, [\beta_{\psi}^{-1}(g_{Fx})^{-1}]^{\top}\mathrm{d}\psi_{Fx}w\rangle$
= $\langle \mathrm{d}\phi_{x}v, [\mathrm{d}\psi_{Fx}\mathrm{d}F_{x}(\mathrm{d}\phi_{x})^{-1}]^{\top}[\beta_{\psi}^{-1}(g_{Fx})^{-1}]^{\top}\mathrm{d}\psi_{Fx}w\rangle$
= $\langle \beta_{\phi}^{-1}(g_{x})^{-1}\mathrm{d}\phi_{x}v, \beta_{\phi}^{-1}(g_{x})[\mathrm{d}\psi_{Fx}\mathrm{d}F_{x}(\mathrm{d}\phi_{x})^{-1}]^{\top}\beta_{\psi}^{-1}(g_{Fx})^{-1}\mathrm{d}\psi_{Fx}w\rangle.$

In the last identity, we used, in particular, that the matrix $\beta_{\phi}^{-1}(g_x)$ is symmetric. On the other hand,

$$g_x(v, (\mathrm{d}F_x)^* w) = \langle \beta_{\phi}^{-1}(g_x)^{-1} \mathrm{d}\phi_x v, \mathrm{d}\phi_x (\mathrm{d}F_x)^* w \rangle.$$

When we compare the two results, we see that

$$(\mathrm{d}F_x)^* = \mathrm{d}\phi_x^{-1}\beta_{\phi}^{-1}(g_x)[\mathrm{d}\psi_{Fx}\mathrm{d}F_x(\mathrm{d}\phi_x)^{-1}]^{\top}\beta_{\psi}^{-1}(g_{Fx})^{-1}\mathrm{d}\psi_{Fx}.$$

In the middle of the right-hand side term, we find the transpose of the local representation of dF_x , that we denote by A_x . Then

$$dF_x (dF_x)^* = d\psi_{Fx}^{-1} A_x \beta_{\phi}^{-1} (g_x) A_x^{\top} \beta_{\psi}^{-1} (g_{Fx})^{-1} d\psi_{Fx}$$

It follows that the singular values of dF_x with respect to the metric g are the square roots of the eigenvalues of the matrix $C_x := A_x \beta_{\phi}^{-1}(g_x) A_x^{\top} \beta_{\psi}^{-1}(g_{Fx})^{-1}$ which is similar to

$$[p_2^{-\frac{1}{2}}A_x p_1^{\frac{1}{2}}][p_1^{\frac{1}{2}}A_x^{\top} p_2^{-\frac{1}{2}}] = B_x B_x^{\top}, \quad B_x := p_2^{-\frac{1}{2}}A_x p_1^{\frac{1}{2}}.$$

It follows that the singular values of dF_x in the metric g are the eigenvalues of $(B_x B_x^{\top})^{\frac{1}{2}}$, or equivalently, the ordinary singular values of B_x , which directly implies the statement of the lemma.

Now, we can finally prove our converse result.

Proposition 3 For any $\varepsilon > 0$, there exists a metric $g^{\varepsilon} \in \mathcal{M}^0$ such that

$$\int \vec{\sigma}^{g^{\varepsilon}}(\mathrm{d}f_x) \,\mathrm{d}\mu(x) \preceq_w \vec{\lambda}(f) + \varepsilon \mathbb{1}$$

Proof: Using a fixed reference metric $g^0 \in \mathcal{M}^0$, for each $N \in \mathbb{N}$ we define the new metric

$$g^{N} := bar(g^{0}, f * g^{0}, f^{2} * g^{0}, \dots, f^{N-1} * g^{0})$$

via the iterates of f (in particular, $g^1 = g^0$). Then, using Lemma 1(i) and (ii) together with the symmetry of the barycenter map, we find that

$$f^{-1} * g^{N} = \operatorname{bar}(f^{-1} * g^{0}, g^{0}, \dots, f^{N-2} * g^{0})$$

=
$$\operatorname{bar}(g^{0}, f * g^{0}, \dots, f^{N-2} * g^{0}, f^{-1} * g^{0})$$

We now compare the two metrics g^N and $f^{-1} * g^N$ at the point fx (for an arbitrarily chosen $x \in M$), using inequality (5):

$$\begin{split} \vec{d}(g_{fx}^{N}, (f^{-1} * g^{N})_{fx}) \\ &= \vec{d}(\operatorname{bar}(\beta_{\phi}^{-1}(g_{fx}^{0}), \beta_{\phi}^{-1}([f * g^{0}]_{fx}), \dots, \beta_{\phi}^{-1}([f^{N-1} * g^{0}]_{fx})), \\ &\quad \operatorname{bar}(\beta_{\phi}^{-1}(g_{fx}^{0}), \beta_{\phi}^{-1}([f * g^{0}]_{fx}), \dots, \beta_{\phi}^{-1}([f^{-1} * g^{0}]_{fx}))) \\ &\preceq \frac{1}{N} \vec{d}(\beta_{\phi}^{-1}([f^{N-1} * g^{0}]_{fx}), \beta_{\phi}^{-1}([f^{-1} * g^{0}]_{fx})) \\ &= \frac{1}{N} \vec{d}((f^{N-1} * g^{0})_{fx}, (f^{-1} * g^{0})_{fx}) = \frac{1}{N} \vec{d}(g_{fNx}^{0}, (f^{-N} * g^{0})_{fNx}) \end{split}$$

For the last identity, we used Lemma 1(iv). Choose charts (ϕ, U) and (ψ, V) around the points $f^N x$ and x, respectively. Let $p(f^N x)$ be the matrix representation of g^0 at $f^N x$ and p(x) the matrix representation of g^0 at x. Then, using Lemma 1(iii), we find that

$$\vec{d}(g^0_{f^Nx},(f^{-N}\ast g^0)_{f^Nx})=\vec{d}(p(f^Nx),\mathbf{d}(\psi\circ f^{-N}\circ\phi^{-1})^{-1}_{\phi(f^Nx)}\ast p(x)).$$

We observe that $d(\psi \circ f^{-N} \circ \phi^{-1})_{\phi(f^N x)}$ is the inverse of the local representation of df^N at x, for which we write A_x^N . Hence, using Proposition 2(ii), we obtain

$$\begin{split} \vec{d}(g_{f^Nx}^0, (f^{-N} * g^0)_{f^Nx}) &= \vec{d}(p(f^Nx), A_x^N * p(x)) \\ &= \vec{d}(I, [p(f^Nx)^{-\frac{1}{2}} A_x^N p(x)^{\frac{1}{2}}] [p(f^Nx)^{-\frac{1}{2}} A_x^N p(x)^{\frac{1}{2}}]^\top). \end{split}$$

Writing $B_x^N := p(f^N x)^{-\frac{1}{2}} A_x^N p(x)^{\frac{1}{2}}$, Proposition 2(i) and Lemma 2 yield

$$\begin{split} \vec{d}(g^0_{f^Nx}, (f^{-N} * g^0)_{f^Nx}) &= \vec{\sigma}(B^N_x (B^N_x)^\top) = 2\vec{\sigma}(B^N_x) \\ &= 2\vec{\sigma}(p(f^Nx)^{-\frac{1}{2}}A^N_x p(x)^{\frac{1}{2}}) = 2\vec{\sigma}^{g^0}(\mathrm{d}f^N_x). \end{split}$$

Using the same arguments for $\vec{d}(g_{fx}^N, (f^{-1} * g^N)_{fx})$, we have thus proven that

$$\vec{\sigma}^{g^N}(\mathrm{d}f_x) \preceq \frac{1}{N} \vec{\sigma}^{g^0}(\mathrm{d}f_x^N) \quad \text{for all } x \in M.$$
(9)

To prove the statement of the proposition, observe that

$$\vec{\lambda}(f) = \int \lim_{n \to \infty} \frac{1}{n} \vec{\sigma}^{g^0}(\mathrm{d}f_x^n) \,\mathrm{d}\mu(x) = \lim_{n \to \infty} \int \frac{1}{n} \vec{\sigma}^{g^0}(\mathrm{d}f_x^n) \,\mathrm{d}\mu(x),$$

where we use again the dominated convergence theorem. Hence, for the given $\varepsilon > 0$, we can find N such that

$$\int \vec{\sigma}^{g^N}(\mathrm{d}f_x) \,\mathrm{d}\mu(x) \stackrel{(9)}{\preceq} \int \frac{1}{N} \vec{\sigma}^{g^0}(\mathrm{d}f_x^N) \,\mathrm{d}\mu(x) \preceq_w \vec{\lambda}(f) + \varepsilon \mathbb{1},$$

which completes the proof.

Remark 1 A careful inspection of the proof shows that the assumption of invertibility of f is actually not needed. It suffices to assume that the derivative df_x is an invertible linear operator for each $x \in M$. This is because, even though we are pulling back the metric g^N by f^{-1} , we are only evaluating the resulting metric at fx, and it holds that

$$(f^{-1} * g)_{fx} = g_x(\mathrm{d}f_{fx}^{-1}v, \mathrm{d}f_{fx}^{-1}w) = g_x((\mathrm{d}f_x)^{-1}v, (\mathrm{d}f_x)^{-1}w)$$

Also in [3], only the invertibility of the values of the linear cocycle is required.

Proposition 1 and 3 together show that the Lyapunov vector $\vec{\lambda}(f)$ can be regarded as a strong infimum⁶ of the vector-valued function

$$\vec{\mathcal{S}}_{f,\mu}: g \mapsto \int \vec{\sigma}^g(\mathrm{d}f_x) \,\mathrm{d}\mu(x), \quad \vec{\mathcal{S}}_{f,\mu}: \mathcal{M}^0 \to \mathfrak{a}^+.$$

The infimum needs to be understood with respect to the order \leq that was defined on the convex cone \mathfrak{a}^+ . Of course, the space \mathcal{M}^0 is huge, so it would be convenient if we could reduce this minimization problem to a smaller subspace. A first observation is that the multiplication with a positive scalar function does not change the value of $\vec{\mathcal{S}}_{f,\mu}$:

$$\vec{\mathcal{S}}_{f,\mu}(\gamma \cdot g) = \vec{\mathcal{S}}_{f,\mu}(g) \text{ for all } \gamma \in C^0(M,\mathbb{R}_{>0}), \ g \in \mathcal{M}^0.$$

Indeed, Lemma 2 implies

$$\log \alpha_i^{\gamma \cdot g}(\mathrm{d}f_x) = \frac{1}{2} (\log \gamma(fx) - \log \gamma(x)) + \log \alpha_i^g(\mathrm{d}f_x),$$

and the integral over $\log \gamma(fx) - \log \gamma(x)$ vanishes because μ is *f*-invariant. It follows that we can restrict the search for an optimal metric to the space of metrics inducing the same (arbitrary) volume form on *M*. To see this, let a volume form ω_0 be fixed and pick an arbitrary $g \in \mathcal{M}^0$. The metric *g* induces a volume form $\omega_g = \sqrt{\det g} \, dx^1 \wedge \dots \wedge dx^d$. Then, there exists a positive function γ such that $\omega_g = \gamma \cdot \omega_0$, and the metric $\tilde{g} := \gamma^{-2/d} \cdot g$ induces the volume form $\gamma^{-1} \cdot \omega_g = \omega_0$. In particular, in dimension d = 1there is no need to optimize, because here any metric is just a positive scalar function.

Furthermore, we can restrict $\vec{S}_{f,\mu}$ to the space of smooth metrics on M, since every continuous metric can be approximated uniformly (by compactness of M) by smooth metrics and the singular values depend continuously on the metric in the uniform topology.

Recall that \mathcal{M} denotes the space of all smooth Riemannian metrics on \mathcal{M} . Given a smooth volume form ω , we let $\mathcal{M}_{\omega} \subset \mathcal{M}$ denote the subspace of metrics inducing ω . We then have the following theorem which will be the basis of our further investigations.

⁶We use the term strong infimum in contrast to a weak (= Pareto) infimum.

Theorem 2 Let $f: M \leftrightarrow$ be a C^1 -diffeomorphism on a compact Riemannian manifold, preserving the probability measure μ . Then, for every smooth volume form ω on M, the following statements hold:

(i) For every $g \in \mathcal{M}_{\omega}$, it holds that

$$\dot{\lambda}(f) \preceq \vec{\mathcal{S}}_{f,\mu}(g)$$

(ii) For every $\varepsilon > 0$, there exists $g^{\varepsilon} \in \mathcal{M}_{\omega}$ such that

$$\hat{\mathcal{S}}_{f,\mu}(g^{\varepsilon}) \preceq_w \hat{\lambda}(f) + \varepsilon \mathbb{1}$$

Remark 2 It is a simple exercise to show that the inequality

$$a \preceq_w b \preceq_w a + \varepsilon \mathbb{1} \tag{10}$$

for vectors $a, b \in \mathfrak{a}^+$ implies $||a - b||_{\infty} \leq \varepsilon$. Hence, Theorem 2 really implies that $\vec{\lambda}(f)$ can be approximated by vectors of the form $\vec{S}_{f,\mu}(g)$ with $g \in \mathcal{M}_{\omega}$. However, the sequence of metrics employed in the proof is not very helpful for concrete computations, because it involves the derivatives of the iterates of f just like the definition of the Lyapunov exponents.

4 The space of Riemannian metrics

To explore deeper properties of the optimization problem for $g \mapsto \tilde{\mathcal{S}}_{f,\mu}(g)$, we need to gain a better understanding of the spaces \mathcal{M} and \mathcal{M}_{ω} and their natural geometric structures. In this section, we gather some facts from the literature about these spaces, mainly taken from [12, 13, 9]. We also refer to [28] for a comprehensive general treatment of manifolds of smooth mappings.

A Riemannian metric g assigns to each $x \in M$ an inner product on the tangent space $T_x M$, i.e. a positive-definite symmetric (0, 2)-tensor $g_x : T_x M \times T_x M \to \mathbb{R}$. We write $T_2^0 M$ for the tensor bundle of (0, 2)-tensors on M and $S_2^0 M \subset T_2^0 M$ for the subbundle of symmetric tensors. This subbundle attaches to each $x \in M$ the vector space of all symmetric (0, 2)-tensors on $T_x M$. In this formal setup, a Riemannian metric is a special section of the bundle $S_2^0 M$, namely one that is positive-definite. We write $S_2^0 M$ for the space of smooth sections of $S_2^0 M$, i.e., of symmetric (0, 2)-tensor fields of class C^{∞} . Obviously, $S_2^0 M$ is an infinite-dimensional vector space over the reals with the pointwise addition and scalar multiplication. Indeed, $S_2^0 M$ is a Fréchet space with the C^{∞} -topology, of which \mathcal{M} is an open subset. The fact that \mathcal{M} is open in the Fréchet space at any point can be identified canonically with $S_0^2 M$.

We define an inner product on each tangent space of \mathcal{M} as

$$\langle h, k \rangle_g := \int \operatorname{tr}(g_x^{-1}h_x g_x^{-1}k_x) \,\mathrm{d}\omega_g(x) \quad \text{for all } h, k \in T_g \mathcal{M} = \mathcal{S}_0^2 M,$$

where ω_g is the volume form induced by g, and g_x , h_x , k_x have to be understood as local representations. With similar arguments as used in the previous section, one

shows that the trace is independent of the charts used to obtain these representations. In terms of the trace metric $\langle \cdot, \cdot \rangle_{g_x}$, we can also write

$$\langle h, k \rangle_g = \int \langle h_x, k_x \rangle_{g_x} \,\mathrm{d}\omega_g(x) \,\mathrm{d}\omega_g(x)$$

For obvious reasons, this Riemannian metric is called the L^2 -metric on \mathcal{M} . It is only a *weak* metric, meaning that the induced topology on the tangent space does not coincide with the inherited topology of the manifold. Nevertheless, as shown in [9], the geodesic distance induced by the L^2 -metric turns \mathcal{M} into a (non-complete) metric space.⁷ It can also be seen easily (using Lemma 1(iii)) that the L^2 -metric is invariant under the diffeomorphism group of M acting by pullback. An explicit expression for the distance function on \mathcal{M} , induced by the L^2 -metric, was given in [10] (see also [8, App. B] for a more elegant proof):

$$d_{L^2}(g^1, g^2) = \left(\int_M d_x (g_x^1, g_x^2)^2 \,\mathrm{d}\omega_{g^0}(x)\right)^{\frac{1}{2}},\tag{11}$$

where g^0 is an arbitrary reference metric with $\operatorname{vol}(M, g^0) = 1$, and $d_x(\cdot, \cdot)$ is the fiber metric on the space of inner products on $T_x M$, induced by the Riemannian metric $\langle a, b \rangle_p = \operatorname{tr}(p^{-1}ap^{-1}b)\sqrt{\operatorname{det}(g^0(x)^{-1}p)}$.

The space \mathcal{M} admits the global product structure $\mathcal{M} \cong \mathcal{V} \times \mathcal{M}_{\omega}$, where \mathcal{V} denotes the space of smooth volume forms on M. The space \mathcal{V} is a Fréchet manifold, in fact an open subset of $\Omega^d(M)$, the Fréchet space of order-d differential forms on M. Given $\omega \in \mathcal{V}$ and $\nu \in \Omega^d(M)$, there exists a unique smooth function, denoted by $\frac{\nu}{\omega}$, such that

$$\nu = \left(\frac{\nu}{\omega}\right)\omega$$

In case that ν is also a volume form, the function $\frac{\nu}{\omega}$ is strictly positive. The space \mathcal{M}_{ω} is a smooth submanifold of \mathcal{M} with tangent space

$$T_g \mathcal{M}_{\omega} = \{ h \in \mathcal{S}_2^0 M : \operatorname{tr}(g_x^{-1} h_x) = 0 \text{ for all } x \in M \}.$$

The following result is proven in [13, Prop. 1.13 and Prop. 2.2].⁸

Proposition 4 For each $\omega \in \mathcal{V}$, the submanifold \mathcal{M}_{ω} is geodesically complete and the geodesic starting at $g^0 \in \mathcal{M}_{\omega}$ with initial tangent h is given by

$$g^{t} = g^{0} \exp(t(g^{0})^{-1}h).$$
(12)

Remark 3 The formula (12) needs to be understood pointwise. That is,

$$g_x^t = g_x^0 \exp(t(g_x^0)^{-1}h_x) \text{ for all } x \in M,$$

where we can use local representations of the involved objects g^0 and h with respect to a fixed chart, and $\exp(\cdot)$ is the usual matrix exponential.

⁷As on a finite-dimensional manifold, the geodesic distance of two points g^1, g^2 is defined as the infimum over the lengths of all piecewise differentiable curves connecting g^1 and g^2 . On general weak Riemannian manifolds, this is only a pseudo-metric.

⁸Actually, the result says much more, namely that \mathcal{M}_{ω} is a symmetric space.

Remark 4 The statement that \mathcal{M}_{ω} is geodesically complete does not imply that \mathcal{M}_{ω} is complete as a metric space, when equipped with the geodesic distance. In finite dimensions, this implication holds by the Hopf-Rinow theorem, but in infinite dimensions there are counter-examples. To the best of my knowledge, it is not known if \mathcal{M}_{ω} is also complete as a metric space (but most probably it is not, because \mathcal{M} is not complete).

Remark 5 The submanifold \mathcal{M}_{ω} is not totally geodesic in \mathcal{M} . Indeed, the formula for geodesics in \mathcal{M} is different, and will not be used in this paper.

Corollary 1 For each pair of metrics $g^a, g^b \in \mathcal{M}_{\omega}$, there exists a unique geodesic on [0, 1], connecting g^a and g^b . This geodesic is given by

$$g_x^t := (g_x^a \#_t g_x^b) \quad for \ all \ x \in M, \ t \in [0, 1],$$
(13)

where the operation $\#_t$ carries over from S_d^+ to the space of symmetric positive-definite (0,2)-tensors on $T_x M$ via local representations.

Proof: The proof consists of two parts. First, we show that the curve given by (13) is a geodesic in \mathcal{M}_{ω} which connects g^a and g^b . Second, we show that each such geodesic is of the given form.

(1) The curve $t \mapsto g^t$ is well-defined as a curve in \mathcal{M} because of (7) and (4). Obviously, $g^0 = g^a$ and $g^1 = g^b$. To show that $t \mapsto g^t$ is a geodesic in \mathcal{M}_{ω} , we define (in terms of local representations)

$$h_x := (g_x^a)^{\frac{1}{2}} \log((g_x^a)^{-\frac{1}{2}} g_x^b (g_x^a)^{-\frac{1}{2}}) (g_x^a)^{\frac{1}{2}} \quad \text{for all } x \in M.$$

Here, $\log(\cdot)$ denotes the matrix logarithm, which is well-defined and unique as an operator on S_d^+ . It is clear that h_x is symmetric for each x and easy to show that h_x is well-defined, i.e. independent of the representation. Hence, h is a section of $S_2^0 M$. It is a smooth section, because it is constructed from smooth sections via smooth operations. Hence, $h \in S_2^0 M$. Finally, we have

$$\operatorname{tr}((g_x^1)^{-1}h_x) = \operatorname{tr}(\log((g_x^a)^{-\frac{1}{2}}g_x^b(g_x^a)^{-\frac{1}{2}})) = \log\det((g_x^a)^{-\frac{1}{2}}g_x^b(g_x^a)^{-\frac{1}{2}}))$$

= $\log[\det(g_x^a)^{-1}\det(g_x^b)] = \log(1) = 0,$

because the fact that g^a and g^b induce the same volume form implies $\det(g_x^a) = \det(g_x^b)$ for all $x \in M$. This shows that $h \in T_{g^a} \mathcal{M}_{\omega}$. To see that $t \mapsto g^t$ is a geodesic, it remains to show that $g^t = g^a \exp(t(g^a)^{-1}h)$ and invoke Proposition 4. This, however, is a simple computation, left to the reader.

(2) To show the uniqueness, let $t \mapsto \tilde{g}^t$ be any geodesic in \mathcal{M}_{ω} with $\tilde{g}^0 = g^a$ and $\tilde{g}^1 = g^b$. By Proposition 4, it must be of the form $\tilde{g}^t = g^a \exp(t(g^a)^{-1}h)$ for some $h \in T_{g^a}\mathcal{M}_{\omega}$, implying $g_x^a \exp((g_x^a)^{-1}h_x) = g_x^b$ for all $x \in M$. Thus,

$$h_x = (g_x^a) \log((g_x^a)^{-1} g_x^b) = (g_x^a) \log((g_x^a)^{-\frac{1}{2}} (g_x^a)^{-\frac{1}{2}} g_x^b (g_x^a)^{-\frac{1}{2}} (g_x^a)^{\frac{1}{2}})$$

= $(g_x^a)^{\frac{1}{2}} \log((g_x^a)^{-\frac{1}{2}} g_x^b (g_x^a)^{-\frac{1}{2}}) (g_x^a)^{\frac{1}{2}}.$

Hence, by the first part of the proof, it follows that $\tilde{g}^t \equiv g^t$.

5 Convexity

In this section, we prove that the function $\vec{\mathcal{S}}_{f,\mu} : \mathcal{M}_{\omega} \to \mathfrak{a}^+$ is convex with respect to the structures introduced on its domain and co-domain. First, we need to prove continuity, which is implied by the following lemma.

Lemma 3 The function $(g, x) \mapsto \vec{\sigma}^g(\mathrm{d}f_x)$ is continuous on $\mathcal{M} \times M$.

Proof: We denote the function under consideration by β . By Lemma 2, we can write β locally as

$$\beta(q,x) = \vec{\sigma}(\operatorname{ev}(q,fx)^{-\frac{1}{2}} \cdot \mathrm{d}f_x \cdot \operatorname{ev}(q,x)^{\frac{1}{2}}),$$

where $\operatorname{ev}(g, x) := g_x$ is the evaluation map. By [28, Rem. 2.1.2], the evaluation map is continuous, implying that $(g, x) \mapsto \operatorname{ev}(g, fx)^{-\frac{1}{2}} \cdot \mathrm{d}f_x \cdot \operatorname{ev}(g, x)^{\frac{1}{2}}$ is continuous. Since the singular values depend continuously on the matrix, then also β is continuous. \Box

The notion of convexity to be used for $\vec{S}_{f,\mu}$ is a natural combination of the two wellstudied notions of *geodesic convexity* (see, e.g., [29, 6]) and *cone-convexity* (see, e.g., [25]). Recall that a function $\varphi: M \to \mathbb{R}$, defined on a finite-dimensional Riemannian manifold (M, g), is called *geodesically convex* if the composition $\varphi \circ \gamma$ with an arbitrary geodesic γ in M is convex in the usual sense. This can be reformulated as

$$\varphi(\gamma(t)) \le (1-t)\varphi(\gamma(0)) + t\varphi(\gamma(1))$$
 for all $t \in [0,1]$

If φ instead takes values in a vector space equipped with a partial order \preceq induced by a convex cone, and the inequality

$$\varphi(\gamma(t)) \preceq (1-t)\varphi(\gamma(0)) + t\varphi(\gamma(1))$$
 for all $t \in [0,1]$

holds for any geodesic in M, we call φ geodesically cone-convex. It makes sense to extend this definition to functions on infinite-dimensional Riemannian manifolds, which are well-behaved in terms of their geodesics. Since this is certainly the case for the manifold \mathcal{M}_{ω} with the L^2 -metric, as Proposition 4 shows, we can ask whether $\vec{\mathcal{S}}_{f,\mu}$ is geodesically cone-convex. The following proposition shows that this is already the case for the integrand in the definition of $\vec{\mathcal{S}}_{f,\mu}$.

Proposition 5 For each $x \in M$, the function $g \mapsto \vec{\sigma}^g(df_x)$, $\mathcal{M}_{\omega} \to \mathfrak{a}^+$, is geodesically cone-convex.

Proof: Let $g : [0,1] \to \mathcal{M}_{\omega}, t \mapsto g^t$, be a geodesic. By Corollary 1, we have $g_x^t \equiv g_x^0 \#_t g_x^1$. We prove the mid-point convexity of $g \mapsto \vec{\sigma}^g(\mathrm{d}f_x)$, i.e.

$$\vec{\sigma}^{g^{\frac{1}{2}}}(\mathrm{d}f_x) \leq \frac{1}{2}\vec{\sigma}^{g^0}(\mathrm{d}f_x) + \frac{1}{2}\vec{\sigma}^{g^1}(\mathrm{d}f_x) \quad \text{for all } x \in M.$$
(14)

To prove this, we choose charts (ϕ, U) and (ψ, V) around x and fx, respectively. Let A_x be a local representation of df_x with respect to this pair of charts. Further, let p_0 and p_1 (q_0 and q_1) represent g_x^0 and g_x^1 (g_{fx}^0 and g_{fx}^1), respectively. According to Lemma 2, then (14) is equivalent to⁹

$$\vec{\sigma}([q_0\#_{\frac{1}{2}}q_1]^{\frac{1}{2}}A_x[p_0\#_{\frac{1}{2}}p_1]^{-\frac{1}{2}}) \leq \frac{1}{2}\vec{\sigma}(q_0^{\frac{1}{2}}A_xp_0^{-\frac{1}{2}}) + \frac{1}{2}\vec{\sigma}(q_1^{\frac{1}{2}}A_xp_1^{-\frac{1}{2}}).$$

⁹Note that we are not using anymore the inversion of p to define the local representation as in (6).

A detailed proof of this inequality can be found in [20, Lem. 3.2]. The continuity, proven in Lemma 3, together with the mid-point convexity implies convexity by a standard argument in convex analysis, see also [20, Rem. 3.3]. \Box

The main result of this section is the following theorem.

Theorem 3 For every $\omega \in \mathcal{V}$, the function $\vec{\mathcal{S}}_{f,\mu} : \mathcal{M}_{\omega} \to \mathfrak{a}^+$ is continuous and geodesically cone-convex.

Proof: The continuity of $\vec{S}_{f,\mu}$ is a simple consequence of the continuity of $(g, x) \mapsto \vec{\sigma}^g(\mathrm{d}f_x)$ together with the compactness of M. The cone-convexity directly follows from the cone-convexity of the integrand together with the fact that the order \preceq is induced by a closed cone.

Observe that the geodesic cone-convexity of $\vec{S}_{f,\mu}$ is equivalent to the geodesic convexity of the functions

$$g \mapsto s_{k,f,\mu}(g) := \int \log \omega_k^g(\mathrm{d}f_x) \,\mathrm{d}\mu(x), \quad k = 1, 2, \dots, d$$

where

$$\omega_k^g(\mathrm{d}f_x) := \prod_{i=1}^k \log \alpha_i^g(\mathrm{d}f_x).$$

Moreover, a minimizer of the vector-valued optimization problem is the same as a common minimizer of the scalar optimization problems

$$\min_{g \in \mathcal{M}_{\omega}} s_{k,f,\mu}(g), \quad k = 1, 2, \dots d.$$

In particular, we obtain the following corollary.

Corollary 2 Let f be a C^2 -diffeomorphism and assume that μ is an ergodic SRB measure. Let k denote the number of positive Lyapunov exponents of (M, f, μ) . Then, the measure-theoretic entropy $h_{\mu}(f)$ can be written as the solution of a geodesically convex optimization problem as follows:

$$h_{\mu}(f) = \inf_{g \in \mathcal{M}_{\omega}} s_{k,f,\mu}(g).$$

Proof: If μ is an ergodic measure, the measure-theoretic entropy of f with respect to μ is given by

$$h_{\mu}(f) = \sum_{i=1}^{d} \max\{0, \lambda_i(f)\} = \sum_{i=1}^{k} \lambda_i(f)$$

according to the entropy-characterization of SRB, developed in [22] by Ledrappier and Young. Together with Theorem 2, this immediately implies the statement of the corollary. $\hfill \Box$

Convex optimization problems are more well-behaved than general (nonlinear) optimization problems in many respects. In particular, every local optimum is also a global optimum, the solution set is convex, there exist efficient numerical algorithms to solve them, and in the case of differentiability there exist first-order optimality criteria. However, the convex optimization problem considered in this paper still has a number of disadvantages which makes it hard to solve:

- The existence of a minimizer is not guaranteed.
- It is vector-valued.
- It is defined on an infinite-dimensional space.
- It is defined on a Riemannian manifold (rather than a vector space).
- It cannot be expected to be differentiable, since singular value functions are not differentiable, in general.

Despite these drawbacks, the optimization problem also has some properties which could turn out to be advantageous. For instance, we can prove that the functions $s_{k,f,\mu}$ satisfy global Lipschitz estimates on \mathcal{M}_{ω} with respect to the L^2 -metric.

Theorem 4 For each $k \in \{1, 2, ..., d\}$, the function $s_{k,f,\mu} : \mathcal{M}_{\omega} \to \mathbb{R}$ satisfies the global Lipschitz-like estimate

$$|s_{k,f,\mu}(g^1) - s_{k,f,\mu}(g^2)| \le \sqrt{k} \left(\int d(g_x^1, g_x^2)^2 \,\mathrm{d}\mu(x) \right)^{\frac{1}{2}},\tag{15}$$

where $d(\cdot, \cdot)$ is the distance function on S_d^+ . If μ is a volume measure induced by some Riemannian metric $g^0 \in \mathcal{M}_{\omega}$, then

$$|s_{k,f,\mu}(g^1) - s_{k,f,\mu}(g^2)| \le \sqrt{k} \, d_{L^2}(g^1, g^2).$$
(16)

That is, $s_{k,f,\mu}$ satisfies a global Lipschitz estimate with respect to the L^2 -distance.

Proof: First, observe that by Jensen's inequality we have

$$(s_{k,f,\mu}(g^1) - s_{k,f,\mu}(g^2))^2 \le \int_M \log^2 \frac{\omega_k^{g^1}(\mathrm{d}f_x)}{\omega_k^{g^2}(\mathrm{d}f_x)} \,\mathrm{d}\mu(x).$$

We split M into the two subsets

$$M_1 := \left\{ x \in M : \omega_k^{g^1}(\mathrm{d}f_x) > \omega_k^{g^2}(\mathrm{d}f_x) \right\}, \quad M_2 := M \setminus M_1$$

Then we have the following identities and inequalities:

$$\begin{split} &\int_{M_1} \log^2 \frac{\omega_k^{g^1}(\mathrm{d}f_x)}{\omega_k^{g^2}(\mathrm{d}f_x)} \,\mathrm{d}\mu(x) \\ &= \int_{M_1} \log^2 \frac{\omega_k((g_{fx}^1)^{-\frac{1}{2}} \mathrm{d}f_x(g_x^1)^{\frac{1}{2}})}{\omega_k((g_{fx}^2)^{-\frac{1}{2}} \mathrm{d}f_x(g_x^2)^{\frac{1}{2}})} \,\mathrm{d}\mu(x) \\ &= \int \log^2 \frac{\omega_k((g_{fx}^1)^{-\frac{1}{2}} (g_{fx}^2)^{\frac{1}{2}} (g_{fx}^2)^{-\frac{1}{2}} \mathrm{d}f_x(g_x^2)^{\frac{1}{2}} (g_x^2)^{-\frac{1}{2}} (g_x^1)^{\frac{1}{2}})}{\omega_k((g_{fx}^2)^{-\frac{1}{2}} \mathrm{d}f_x(g_x^2)^{\frac{1}{2}})} \,\mathrm{d}\mu(x) \\ &\leq \int_{M_1} \left(\log \omega_k((g_{fx}^1)^{-\frac{1}{2}} (g_{fx}^2)^{\frac{1}{2}}) + \log \omega_k((g_x^2)^{-\frac{1}{2}} (g_x^1)^{\frac{1}{2}}) \right)^2 \,\mathrm{d}\mu(x) \\ &\leq 2 \int_{M_1} \left(\log^2 \omega_k((g_{fx}^1)^{-\frac{1}{2}} (g_{fx}^2)^{\frac{1}{2}}) + \log^2 \omega_k((g_x^2)^{-\frac{1}{2}} (g_x^1)^{\frac{1}{2}}) \right) \,\mathrm{d}\mu(x). \end{split}$$

In the first equality, we simply use the definition of $s_{k,f,\mu}$. The second equality identifies df_x , g_x^1 and g_x^2 with their corresponding local representations and uses Lemma 2.

The third equality is trival. The subsequent inequality is Horn's inequality together with the fact that $\log^2(\cdot)$ is increasing on $(1, \infty)$. The last inequality simply uses that $(x+y)^2 \leq 2x^2 + 2y^2$ for any $x, y \in \mathbb{R}$.

Now, observe that for any $a, b \in \mathcal{S}_d^+$ we have

$$\log^2 \omega_k (a^{-\frac{1}{2}} b^{\frac{1}{2}}) = \left(\sum_{i=1}^k \log \alpha_i (a^{-\frac{1}{2}} b^{\frac{1}{2}}) \right)^2$$

$$\leq k \sum_{i=1}^k \log^2 \alpha_i (a^{-\frac{1}{2}} b^{\frac{1}{2}}) \leq k \sum_{i=1}^d \log^2 \alpha_i (a^{-\frac{1}{2}} b^{\frac{1}{2}})$$

$$= k \|\vec{\sigma} (a^{-\frac{1}{2}} b^{\frac{1}{2}})\|_2^2 = \frac{k}{4} \|\vec{d} (a, b)\|_2^2 = \frac{k}{4} d(a, b)^2.$$

Altogether, we obtain the estimate

$$\int_{M_1} \log^2 \frac{\omega_k^{g^1}(\mathrm{d}f_x)}{\omega_k^{g^2}(\mathrm{d}f_x)} \,\mathrm{d}\mu(x) \le \frac{k}{2} \int_{M_1} \left(d(g_{fx}^1, g_{fx}^2)^2 + d(g_x^1, g_x^2)^2 \right) \,\mathrm{d}\mu(x).$$

In a similar fashion, we derive that

$$\int_{M_2} \log^2 \frac{\omega_k^{g^1}(\mathrm{d}f_x)}{\omega_k^{g^2}(\mathrm{d}f_x)} \,\mathrm{d}\mu(x) \le \frac{k}{2} \int_{M_2} \left(d(g_{fx}^2, g_{fx}^1)^2 + d(g_x^2, g_x^1)^2 \right) \,\mathrm{d}\mu(x)$$

Because of the symmetry of the distance and the *f*-invariance of μ , summing the two integrals gives

$$(s_{k,f,\mu}(g^1) - s_{k,f,\mu}(g^2))^2 \le k \int_M d(g_x^1, g_x^2)^2 \,\mathrm{d}\mu(x),$$

which is equivalent to the claimed inequality (15).

To prove (16), assume that μ is the volume measure induced by some $g^0 \in \mathcal{M}_{\omega}$. According to (15) and (11), it suffices to show that $d_x(g_x^1, g_x^2) = d(g_x^1, g_x^2)$ for all $g^1, g^2 \in \mathcal{M}_{\omega}$ and $x \in M$. By definition $d_x(\cdot, \cdot)$ is the distance function on $(S_2^0 M)_x$ induced by the Riemannian metric

$$\langle a, b \rangle_p = \operatorname{tr}(p^{-1}ap^{-1}b)\sqrt{\operatorname{det}((g_x^0)^{-1}p)},$$

while $d(\cdot, \cdot)$ is the distance function induced by

$$\langle a,b\rangle'_p = \operatorname{tr}(p^{-1}ap^{-1}b).$$

We identify the space of inner products on $T_x M$ with S_d^+ . The submanifold of S_d^+ consisting of all matrices with the same determinant c > 0 is totally geodesic. Hence, if $\gamma : [0,1] \to S_d^+$ is a geodesic connecting g_x^1 with g_x^2 , then $\det(\gamma(t))$ is the same for all $t \in [0,1]$ and equals $\det(g_x^0)$. It follows that $\det((g_x^0)^{-1}\gamma(t)) = 1$ for all t, showing that $\langle \cdot, \cdot \rangle' = \langle \cdot, \cdot \rangle$, and hence $d_x(\cdot, \cdot) = d(\cdot, \cdot)$.

Remark 6 The proven fact that $s_{k,f,\mu}$ is globally Lipschitz on \mathcal{M}_{ω} if μ is a volume measure implies that there exists a unique extension of $s_{k,f,\mu}$ to the metric completion of \mathcal{M}_{ω} which obeys the same Lipschitz estimate. The completion of \mathcal{M} was studied in [11] and, roughly speaking, characterized as the space of measurable metrics inducing a finite volume. In fact, it is isometric to the space of L^2 -mappings from the manifold \mathcal{M} to the completion of the space of positive-definite symmetric matrices, as proven in [7].

6 Towards a first-order optimality criterion

Since singular value functions do not depend smoothly on parameters, we cannot expect the function $\vec{S}_{f,\mu}$ (or $s_{k,f,\mu}$) to be differentiable. However, the geodesic convexity implies that unilateral directional derivatives exist (see [29, Ch. 3, Thm. 4.2]) and possibly we can compute some generalized derivative, e.g., a subdifferential. This would lead to a first-order criterion for optimality.

In this section, we only take the first step and compute directional derivatives of the functions $s_{k,f,\mu}$. Moreover, we restrict ourselves to the simplest situation in which all singular values have multiplicity one at each point of the manifold. Although this assumption is not very realistic, it will give us some idea of what we can expect a subgradient of $s_{k,f,\mu}$ to look like.

Theorem 5 Let $g \in \mathcal{M}_{\omega}$ be a metric satisfying $\alpha_k^g(\mathrm{d}f_x) > \alpha_{k+1}^g(\mathrm{d}f_x)$ for all $x \in M$ and some $k \in \{1, 2, \ldots, d-1\}$. Then, all directional derivatives of $s_{k,f,\mu}$ at g exist. For any $h \in T_g \mathcal{M}_{\omega}$, the directional derivative of $s_{k,f,\mu}$ at g in direction h is given by

$$\partial_h s_{k,f,\mu}(g) = \int \langle Q_{k,x}^{\mathbf{l}} - Q_{k,x}^{\mathbf{r}}, h_x \rangle_{g_x} \, \mathrm{d}\mu(x),$$

where $Q_{k,x}^{l}$ $(Q_{k,x}^{r})$ denotes the orthogonal projection in $(T_{x}M, g_{x})$ onto the subspace spanned by the first k left (right) singular vectors of $df_{f^{-1}x}$ (df_{x}) .

Proof: The proof is subdivided into six steps.

Step 1: Let us write $s_{k,x}(g) := \log \omega_k^g(\mathrm{d}f_x)$, which is the integrand in the definition of $s_{k,f,\mu}(g)$. We first try to differentiate $s_{k,x}$ in the hope that the order of differentiation and integration can be interchanged. To this end, we choose a smooth curve γ in \mathcal{M}_{ω} with $\gamma(0) = g$, $\dot{\gamma}(0) = h$ and decompose the function $s_{k,x} \circ \gamma$ as

$$(s_{k,x} \circ \gamma)(t) = \ell_k \circ \alpha \circ \zeta_x(t),$$

where

$$\begin{aligned} \zeta_x(t) &= \gamma(t)_{f_x}^{\frac{1}{2}} \mathrm{d}f_x \gamma(t)_x^{-\frac{1}{2}}, \quad \zeta_x : \mathbb{R} \to \mathrm{GL}(d, \mathbb{R}), \\ \alpha(g) &:= (\alpha_1(g), \dots, \alpha_d(g)), \quad \alpha : \mathrm{GL}(d, \mathbb{R}) \to \mathbb{R}^d, \\ \ell_k(\xi) &:= \sum_{i=1}^k \log \hat{\xi}_i, \quad \ell_k : \mathbb{R}^d \to [-\infty, \infty). \end{aligned}$$

Here, some explanation is necessary. In the definition of $\zeta_x(t)$, we treat $\gamma(t)_x$, $\gamma(t)_{fx}$ and df_x as matrices, where we think of their local representations with respect to appropriate charts (with Lemma 2 in mind). The function α simply maps a matrix $g \in \operatorname{GL}(d, \mathbb{R})$ to the vector of its ordinary singular values. Finally, for a vector $\xi =$ $(\xi_1, \ldots, \xi_d) \in \mathbb{R}^d$, we write $\hat{\xi}_1 \geq \cdots \geq \hat{\xi}_d$ for the absolute values of its components, ordered from the largest to the smallest, and then define $\ell_k(\xi)$ as above. Here, the value $-\infty$ is possible if at least one component of ξ is zero. Using Lemma 2, we then see that $s_{k,x}$ can be decomposed in the suggested way (keeping mind that we are not using anymore the inversion of p to define the local representation as in (6)).

Step 2: We compute the derivative of ζ_x at t = 0. By the product rule, we have

$$\dot{\zeta}_x(t) = \left[\frac{\partial}{\partial t}\Big|_{t=0}\gamma(t)_{fx}^{\frac{1}{2}}\right] \mathrm{d}f_x\gamma(0)_x^{-\frac{1}{2}} + \gamma(0)_{fx}^{\frac{1}{2}}\mathrm{d}f_x\left[\frac{\partial}{\partial t}\Big|_{t=0}\gamma(t)_x^{-\frac{1}{2}}\right].$$

With the notation $\mathbf{r}: p \mapsto p^{\frac{1}{2}}$ and $\mathbf{i}: p \mapsto p^{-1}$ (both defined on \mathcal{S}_d^+), we can write this

$$\dot{\zeta}_x(t) = \left[\mathrm{d}\mathbf{r}_{\gamma(0)_{fx}} \dot{\gamma}(0)_{fx} \right] \mathrm{d}f_x \gamma(0)_x^{-\frac{1}{2}} + \gamma(0)_{fx}^{\frac{1}{2}} \mathrm{d}f_x \left[\mathrm{d}(\mathbf{i} \circ \mathbf{r})_{\gamma(0)_x} \dot{\gamma}(0)_x \right].$$

Using that $d\mathbf{i}_p h = -p^{-1}hp^{-1}$ and $d\mathbf{r}_p h = L_p^{-1}(h)$, where L_p is the Lyapunov operator¹⁰

$$L_p(X) := p^{\frac{1}{2}}X + Xp^{\frac{1}{2}},$$

we can finally write the derivative of ζ_x as

$$\dot{\zeta}_x(t) = L_{g_{fx}}^{-1}(h_{fx}) \mathrm{d}f_x g_x^{-\frac{1}{2}} - g_{fx}^{\frac{1}{2}} \mathrm{d}f_x g_x^{-\frac{1}{2}} L_{g_x}^{-1}(h_x) g_x^{-\frac{1}{2}}.$$
(17)

Observe that $X := L_{g_x}^{-1}(h_x)$ is a symmetric matrix, since it solves $g_x^{\frac{1}{2}}X + Xg_x^{\frac{1}{2}} = h_x$ and by symmetry of p and h_x , also X^{\top} solves this equation. Hence, $X = L_{g_x}^{-1}(h_x) = X^{\top}$. Step 3: Now, we consider the composed function $\ell_k \circ \alpha$. According to [23, Prop. 6.2], $\ell_k \circ \alpha$ is differentiable at $X \in \mathbb{R}^{d \times d}$ if and only if ℓ_k is differentiable at $\alpha(X)$. The latter is the case if $\alpha_k(X) > \alpha_{k+1}(X)$, which holds in our case by assumption, since $\alpha_k(X) = \alpha_k(\zeta_x(0)) = \alpha_k^g(\mathrm{d}f_x)$. In this case,

$$\nabla(\ell_k \circ \alpha)(X) = U^{\top} \operatorname{Diag}\left(\frac{1}{\alpha_1(X)}, \dots, \frac{1}{\alpha_k(X)}, 0, \dots, 0\right) V,$$

where $X = U^{\top} \text{Diag}(\alpha(X))V$ is a singular value decomposition.

Step 4: To compute the derivative of $s_{k,x} \circ \gamma$, we need to compose the two derivatives $\nabla(\ell_k \circ \alpha)(\zeta_x(0))$ and $\dot{\zeta}_x(0)$ in the proper way. This is the Euclidean inner product on $\mathbb{R}^{d \times d}$, which is given by $\langle X, Y \rangle = \operatorname{tr}[X^\top Y]$ (also known as *Frobenius inner product*). For brevity, let us write $S_x := \nabla(\ell_k \circ \alpha)(\zeta_x(0))$ and $Z_x := L_{q_x}^{-1}(h_x)$. Then

$$\begin{aligned} \frac{\partial}{\partial t}\Big|_{t=0} (s_{k,x} \circ \gamma)(0) &= \operatorname{tr}[S_x^\top \dot{\zeta}_x(0)] \\ \stackrel{(17)}{=} \operatorname{tr}\left[S_x^\top \left(Z_{fx} \mathrm{d} f_x g_x^{-\frac{1}{2}} - g_{fx}^{\frac{1}{2}} \mathrm{d} f_x g_x^{-\frac{1}{2}} Z_x g_x^{-\frac{1}{2}}\right)\right] \\ &= \operatorname{tr}\left[S_x^\top Z_{fx} \mathrm{d} f_x g_x^{-\frac{1}{2}}\right] - \operatorname{tr}\left[S_x^\top g_{fx}^{\frac{1}{2}} \mathrm{d} f_x g_x^{-\frac{1}{2}} Z_x g_x^{-\frac{1}{2}}\right] \\ &= \operatorname{tr}\left[Z_{fx} \mathrm{d} f_x g_x^{-\frac{1}{2}} S_x^\top\right] - \operatorname{tr}\left[Z_x g_x^{-\frac{1}{2}} S_x^\top g_{fx}^{\frac{1}{2}} \mathrm{d} f_x g_x^{-\frac{1}{2}}\right] \\ &= \operatorname{tr}\left[Z_{fx} g_{fx}^{-\frac{1}{2}} \zeta_x(0) S_x^\top\right] - \operatorname{tr}\left[Z_x g_x^{-\frac{1}{2}} S_x^\top \zeta_x(0)\right]. \end{aligned}$$

From the definition of S_x , it follows that

$$\zeta_x(0)S_x^{\top} = U_{\zeta_x(0)}^{\top}(I_{k\times k} \oplus 0_{(d-k)\times (d-k)})U_{\zeta_x(0)},$$

where $\zeta_x(0) = U_{\zeta_x(0)}^{\top} \text{Diag}(\alpha(\zeta_x(0))) V_{\zeta_x(0)}$ is the chosen singular value decomposition. Analogously,

$$S_x^{\top}\zeta_x(0) = V_{\zeta_x(0)}^{\top}(I_{k\times k} \oplus 0_{(d-k)\times (d-k)})V_{\zeta_x(0)}$$

We can write these matrices in a more compact way as $\bar{U}_{fx}\bar{U}_{fx}^{\top}$ and $\bar{V}_x\bar{V}_x^{\top}$, where \bar{U}_{fx} is the $(d \times k)$ -matrix whose columns are the first k left singular vectors of $\zeta_x(0)$, and \bar{V}_x

¹⁰Since $p^{\frac{1}{2}}$ has only strictly positive eigenvalues, the Lyapunov operator is invertible.

is defined in the same way via the right singular vectors. Observe that $\bar{U}_{fx}\bar{U}_{fx}^{\top}$ is the orthogonal projection onto the subspace spanned by the first k left singular vectors of $\zeta_x(0)$ (analogously for $\bar{V}_x\bar{V}_x^{\top}$). We thus have

$$\frac{\partial}{\partial t}\Big|_{t=0}(s_{k,x}\circ\gamma)(0) = \operatorname{tr}\left[Z_{fx}g_{fx}^{-\frac{1}{2}}\bar{U}_{fx}\bar{U}_{fx}^{\top}\right] - \operatorname{tr}\left[Z_{x}g_{x}^{-\frac{1}{2}}\bar{V}_{x}\bar{V}_{x}^{\top}\right]$$

Now, recall that $g_x^{\frac{1}{2}}Z_x + Z_x g_x^{\frac{1}{2}} = h_x$, implying $Z_x g_x^{-\frac{1}{2}} + g_x^{-\frac{1}{2}}Z_x = g_x^{-\frac{1}{2}}h_x g_x^{-\frac{1}{2}}$. Observe that for arbitrary symmetric matrices a, p, x, y with px + xp = y, we have $\operatorname{tr}[apx] = \operatorname{tr}[(apx)^{\top}] = \operatorname{tr}[xpa] = \operatorname{tr}[axp]$, and hence $\operatorname{tr}[ay] = \operatorname{tr}[a(px)] + \operatorname{tr}[a(xp)] = 2\operatorname{tr}[apx]$. This yields

$$\frac{\partial}{\partial t}\Big|_{t=0}(s_{k,x}\circ\gamma)(0) = \frac{1}{2}\operatorname{tr}\left[g_{fx}^{-\frac{1}{2}}h_{fx}g_{fx}^{-\frac{1}{2}}\bar{U}_{fx}\bar{U}_{fx}^{\top}\right] - \frac{1}{2}\operatorname{tr}\left[g_{x}^{-\frac{1}{2}}h_{x}g_{x}^{-\frac{1}{2}}\bar{V}_{x}\bar{V}_{x}^{\top}\right] \\
= \frac{1}{2}\operatorname{tr}\left[g_{fx}^{\frac{1}{2}}g_{fx}^{-1}h_{fx}g_{fx}^{-1}g_{fx}^{\frac{1}{2}}\bar{U}_{fx}\bar{U}_{fx}^{\top}\right] - \frac{1}{2}\operatorname{tr}\left[g_{x}^{\frac{1}{2}}g_{x}^{-1}h_{x}g_{x}^{-1}g_{x}^{\frac{1}{2}}\bar{V}_{x}\bar{V}_{x}^{\top}\right] \\
= \frac{1}{2}\langle g_{fx}^{\frac{1}{2}}\bar{U}_{fx}\bar{U}_{fx}^{\top}g_{fx}^{\frac{1}{2}}, h_{fx}\rangle_{g_{fx}} - \frac{1}{2}\langle g_{x}^{\frac{1}{2}}\bar{V}_{x}\bar{V}_{x}^{\top}g_{x}^{\frac{1}{2}}, h_{x}\rangle_{g_{x}}.$$
(18)

Step 5: We have to understand how the matrices $\bar{U}_{fx}\bar{U}_{fx}^{\top}$ and $\bar{V}_x\bar{V}_x^{\top}$ change, when we change the charts. Recall that \bar{U}_{fx} is a matrix built from left singular vectors of $\zeta_x(0)$, i.e. the eigenvectors of $\zeta_x(0)\zeta_x(0)^{\top}$. One can easily show that with respect to another pair of charts, the corresponding symmetric matrix becomes

$$\tilde{\zeta}_x(0)\tilde{\zeta}_x(0)^{\top} = o\zeta_x(0)\zeta_x(0)^{\top}o^{\top}$$

with an orthogonal matrix o of the form

$$o = (b * g_{fx})^{-\frac{1}{2}} b g_{fx}^{\frac{1}{2}},$$

where b is the derivative of the coordinate change at fx. Then, in the new pair of charts, $\bar{U}_{fx}\bar{U}_{fx}^{\top}$ becomes $o\bar{U}_{fx}\bar{U}_{fx}^{\top}o^{\top}$. Hence,

$$(b*g_{fx})^{\frac{1}{2}}o\bar{U}_{fx}\bar{U}_{fx}^{\top}o^{\top}(b*g_{fx})^{\frac{1}{2}} = bg_{fx}^{\frac{1}{2}}\bar{U}_{fx}\bar{U}_{fx}^{\top}g_{fx}^{\frac{1}{2}}b^{\top} = b*g_{fx}^{\frac{1}{2}}\bar{U}_{fx}\bar{U}_{fx}^{\top}g_{fx}^{\frac{1}{2}}$$

This computation shows that $g_{fx}^{\frac{1}{2}} \overline{U}_{fx} \overline{U}_{fx}^{\top} g_{fx}^{\frac{1}{2}}$ is the local representation of a global section of $S_2^0 M$, i.e. a symmetric (not necessarily smooth) (0, 2)-tensor field. The same can be shown for $g_x^{\frac{1}{2}} \overline{V}_x \overline{V}_x^{\top} g_x^{\frac{1}{2}}$. What are these global sections? Recall from the proof of Lemma 2 that a local representation of $df_x^* df_x$ is given by $p_x^{-1} A_x^{\top} p_{fx} A_x$, where A_x is a local representation of df_x and p_x (p_{fx}) one of g_x (g_{fx}). The eigenvectors v_1, \ldots, v_d of $df_x^* df_x$ then have local representations $\tilde{v}_1, \ldots, \tilde{v}_d$, which are the eigenvectors of $p_x^{-1} A_x^{\top} p_{fx} A_x =: C_x$. The matrix $\zeta_x(0)^{\top} \zeta_x(0)$ is then given by $p_x^{\frac{1}{2}} C_x p_x^{-\frac{1}{2}}$, and hence its eigenvectors are $p_x^{\frac{1}{2}} \tilde{v}_1, \ldots, p_x^{\frac{1}{2}} \tilde{v}_d$ (which form an orthogonal basis with respect to the standard Euclidean inner product). If Q_x is the orthogonal projection onto the linear subspace of \mathbb{R}^d spanned by the first k eigenvectors of $\zeta_x(0)^{\top} \zeta_x(0)$, then for $i = 1, \ldots, k$ we have $(p_x^{\frac{1}{2}} Q_x p_x^{\frac{1}{2}}) \tilde{v}_i = p_x \tilde{v}_i$ and for $i = k + 1, \ldots, d$ we have $(p_x^{\frac{1}{2}} Q_x p_x^{\frac{1}{2}}) \tilde{v}_i = 0$. Hence,

$$\langle (p_x^{\frac{1}{2}}Q_x p_x^{\frac{1}{2}})\tilde{v}_i, \tilde{v}_j \rangle = \begin{cases} \langle (p_x^{\frac{1}{2}}\tilde{v}_i), (p_x^{\frac{1}{2}}\tilde{v}_j) \rangle = \delta_{ij} & \text{for } i = 1, \dots, k, \\ 0 & \text{otherwise.} \end{cases}$$

It follows that $g_{fx}^{\frac{1}{2}} \bar{U}_{fx} \bar{U}_{fx}^{\top} g_{fx}^{\frac{1}{2}}$ is the local representation of the orthogonal projection $Q_{k,fx}^{l}$ onto the subspace of $T_{fx}M$ which is spanned by the first k left singular vectors of df_x . Analogously, $g_x^{\frac{1}{2}} \bar{V}_x \bar{V}_x^{\top} g_x^{\frac{1}{2}}$ represents the orthogonal projection $Q_{k,x}^{r}$ onto the subspace of T_xM spanned by the first k right singular vectors of df_x . Here, of course, the orthogonality is defined in terms of the metric $g = \gamma(0)$.

Step 6: Using the invariance of the measure μ , we obtain from (18) via integration that

$$\int \frac{\partial}{\partial t}\Big|_{t=0} (s_{k,x} \circ \gamma)(0) \,\mathrm{d}\mu(x) = \frac{1}{2} \int \langle Q_{k,x}^{\mathrm{l}} - Q_{k,x}^{\mathrm{r}}, h_x \rangle_{g_x} \,\mathrm{d}\mu(x).$$

It remains to prove that integration and derivative on the left-hand side can be interchanged. To show this, first observe that our assumption about the spectral gap carries over from g to nearby metrics $\gamma(t)$, $|t| \leq \varepsilon$, by Lemma 3. Hence, the above formula holds for all $(t, x) \in [-\varepsilon, \varepsilon] \times M$. Thus, the derivative of $s_{k,x} \circ \gamma$ at any $t \in [-\varepsilon, \varepsilon]$ has the same form as above, but the orthogonal projections are then defined in terms of $\gamma(t)$ instead of $\gamma(0)$:

$$\frac{\partial}{\partial t}\Big|_{t=0}(s_{k,x}\circ\gamma)(t) = \frac{1}{2}\langle Q_{k,x}^{l}(t) - Q_{k,x}^{r}(t), \dot{\gamma}(t)_{x}\rangle_{\gamma(t)_{x}}.$$

Here, we can estimate

$$\begin{aligned} \left| \langle Q_{k,x}^{l}(t) - Q_{k,x}^{r}(t), \dot{\gamma}(t)_{x} \rangle_{\gamma(t)_{x}} \right| \\ & \leq \left(\|Q_{k,x}^{l}(t)\|_{\gamma(t)_{x}} + \|Q_{k,x}^{r}(t)\|_{\gamma(t)_{x}} \right) \cdot \|\dot{\gamma}(t)_{x}\|_{\gamma(t)_{x}} \end{aligned}$$

The term $\|\dot{\gamma}(t)_x\|_{\gamma(t)_x}$ is continuous in (t, x), and thus can be estimated by a constant C on a compact set of the form $[-\varepsilon, \varepsilon] \times M$. The orthogonal projections have norm 1, hence we can estimate the complete term by a constant. We conclude that the Leibniz rule can be applied to interchange the order of integration and differentiation. The proof is complete.

Remark 7 In the case when μ is a volume measure, coming from a metric which induces the volume form ω , we would expect (in the setup of the theorem) that $s_{k,f,\mu}$ is differentiable at g with gradient $x \mapsto Q_{k,x}^{1} - Q_{k,x}^{r}$ (which may be non-smooth and live in some metric space completion). Hence, a vanishing gradient would mean that $Q_{k,x}^{1} = Q_{k,x}^{r}$, which is some sort of alignment of derivatives. If this holds for all k, it can also be interpreted as a symmetry property, since the left and right singular vectors of a real square matrix are the same if and only if the matrix is symmetric. This is only an analogy, however, because here we are not dealing with a single matrix but the composition of different linear operators (except at the fixed points of f).

7 Summary and outlook

We have shown that the vector of averaged Lyapunov exponents of a smooth measurepreserving system on a compact manifold M is the infimum of a geodesically coneconvex function, defined on the space of smooth Riemannian metrics on M which preserve a given volume form. Several research directions are conceivable to make this a fruitful approach for a better theoretical understanding or numerical computations. These include:

- (1) Finding criteria for the existence of minimizers; if such minimizers can be shown to exist only in the completion of \mathcal{M}_{ω} , an important question concerns their regularity.
- (2) Computation of subgradients for the derivation of a first-order optimality criterion. Maybe in some cases it is possible, to find an optimal metric by solving the equation defined by the first-order criterion.
- (3) Finding ways to restrict the optimization problem to one or several finitedimensional convex problems. The approach for doing this that comes first to mind is probably restricting the function $\vec{S}_{f,\mu}$ to finite-dimensional totally geodesic submanifolds of \mathcal{M}_{ω} . However, it is very likely that there are none except for possibly low-dimensional ones. One way to find such submanifolds is to consider fixed point components of isometries. If we consider \mathcal{M} instead of \mathcal{M}_{ω} , we know that the diffeomorphism group of M acts on \mathcal{M} by isometries. A fixed point of the action of one diffeomorphism φ is a metric in which φ is an isometry. So, one possible construction of a totally geodesic submanifold is

$$S := \{ g \in \mathcal{M} : f \in \operatorname{Isom}(M, g) \text{ for all } f \in F \},\$$

where $F \subset \text{Diff}(M)$ is any subset. As an example, consider a manifold M which is a Lie group and let F be the group of left translations. Then S is the space of left-invariant metrics which is finite-dimensional and can be identified with S_d^+ . In general, to obtain a finite-dimensional S by the above construction, one would expect that the orbits Fx all have to be dense in M. But then, any metric in S is determined by its value at a single point. Another way to obtain a finite-dimensional problem is to start with low-dimensional totally geodesic submanifold, solve the problem there, then use the solution as the initial guess for another finite-dimensional problem (in some other submanifold) and continue like this in the hope that the process converges to a globally (nearly) optimal metric.

Acknowledgements

I express my sincere gratitude to Dr. Nicola Cavallucci for his continuing interest in my paper and helping me to gain a better understanding of the space \mathcal{M} and its geometry.

References

- [1] R. Bhatia. Positive Definite Matrices. Princeton University Press, 2009.
- [2] A. Blumenthal, J. Xue, L.-S. Young. Lyapunov exponents for random perturbations of some area-preserving maps including the standard map. Annals of Mathematics 185(1), 285–310, 2017.
- J. Bochi. Ergodic optimization of Birkhoff averages and Lyapunov exponents. Proceedings of the International Congress of Mathematicians (ICM 2018), 1825–1846, 2018.
- [4] J. Bochi, A. Navas. A geometric path from zero Lyapunov exponents to rotation cocycles. Ergodic Theory and Dynamical Systems, vol. 35, no. 2, 374–402, 2015.

- [5] V. A. Boichenko, G. A. Leonov, V. Reitmann. Dimension Theory for Ordinary Differential Equations. Teubner, Stuttgart, 2005.
- [6] N. Boumal. An Introduction to Optimization on Smooth Manifolds. Cambridge University Press, 2023.
- [7] N. Cavallucci. The L²-completion of the space of Riemannian metrics is CAT (0): A shorter proof. Proceedings of the American Mathematical Society 151(7), 3183–3187, 2023.
- [8] N. Cavallucci, Z. Su. The metric completion of the space of vector-valued oneforms. arXiv preprint arXiv:2302.06840, 2023.
- B. Clarke. The metric geometry of the manifold of Riemannian metrics over a closed manifold. Calculus of Variations and Partial Differential Equations 39, 533-545, 2010.
- [10] B. Clarke. Geodesics, distance, and the CAT (0) property for the manifold of Riemannian metrics. Mathematische Zeitschrift 273(1-2), 55-93, 2013.
- [11] B. Clarke. The completion of the manifold of Riemannian metrics. Journal of Differential Geometry 93(2), 203–268, 2013.
- [12] D. G. Ebin. The manifold of Riemannian metrics. In: Global Analysis, Berkeley, Calif., 1968. Proc. Sympos. Pure Math., vol. 15, 1970.
- [13] D. S. Freed, D. Groisser. The basic geometry of the manifold of Riemannian metrics and of its quotient by the diffeomorphism group. Michigan Mathematical Journal 36(3), 323–344, 1989.
- [14] P. Giesl, S. Hafstein, C. Kawan. Review on contraction analysis and computation of contraction metrics. Journal of Computational Dynamics 10(1), 1–47, 2023.
- [15] P. Giesl, S. Hafstein, M. Haraldsdottir, D. Thorsteinsson, C. Kawan. Subgradient algorithm for computing contraction metrics for equilibria. Journal of Computational Dynamics 10(2), 281–303, 2023.
- [16] O. Gil-Medrano, P. W. Michor. The Riemannian manifold of all Riemannian metrics. Q. J. Math., Oxf. II. Ser. 42, No. 166, 183–202, 1991.
- [17] N. Gourmelon. Adapted metrics for dominated splittings. Ergodic Theory and Dynamical Systems 27(6), 1839–1849, 2007.
- [18] B. Hasselblatt, A. Katok. *Principal Structures*. Handbook of dynamical systems. Vol. 1. Elsevier Science, 2002. 1–203.
- [19] A. Katok, B. Hasselblatt. Introduction to the Modern Theory of Dynamical Systems. Vol. 54. Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge. 1995.
- [20] C. Kawan, S. Hafstein, P. Giesl. A subgradient algorithm for data-rate optimization in the remote state estimation problem. SIAM Journal on Applied Dynamical Systems 20(4), 2141–2173, 2021.
- [21] J. Lawson, Y. Lim. Monotonic properties of the least squares mean. Mathematische Annalen 351, 267–279, 2011.
- [22] F. Ledrappier, L.-S. Young. The metric entropy of diffeomorphisms: Part I: Characterization of measures satisfying Pesin's entropy formula. Annals of Mathematics, 509–539, 1985.

- [23] A. S. Lewis, H. S. Sendov. Nonsmooth analysis of singular values. Part I: Theory. Set-Valued Analysis 13(3), 213–241, 2005.
- [24] M. Louzeiro, C. Kawan, S. Hafstein, P. Giesl, J. Yuan. A Projected Subgradient Method for the Computation of Adapted Metrics for Dynamical Systems. SIAM Journal on Applied Dynamical Systems 21(4), 2610–2641, 2022.
- [25] D. T. Luc. Theory of Vector Optimization. Springer, 1989.
- [26] A. W. Marshall, I. Olkin, B. C. Arnold. Inequalities: Theory of Majorization and its Applications. 2nd ed. Springer, New York, 2011.
- [27] H. Oeri, D. Goluskin. Convex computation of maximal Lyapunov exponents. arXiv preprint arXiv:2212.07565, 2022.
- [28] A. Schmeding. An Introduction to Infinite-dimensional Differential Geometry. Cambridge University Press, vol. 202, 2022.
- [29] C. Udriste. Convex Functions and Optimization Methods on Riemannian Manifolds. Springer Science & Business Media 297, 1994.