# arXiv:2307.06373v1 [cond-mat.dis-nn] 12 Jul 2023

# Noise-induced transport in the Aubry-André-Harper model

Devendra Singh Bhakuni,<sup>1</sup> Talía L. M. Lezama,<sup>2</sup> and Yevgeny Bar Lev<sup>1</sup>

<sup>1</sup>Department of Physics, Ben-Gurion University of the Negev, Beer-Sheva 84105, Israel

<sup>2</sup>Department of Physics, Yeshiva University, New York, New York 10016, USA

We study quantum transport in a quasiperiodic Aubry-André-Harper (AAH) model induced by the coupling of the system to a Markovian heat bath. We find that coupling the heat bath locally does not affect transport in the delocalized and critical phases, while it induces logarithmic transport in the localized phase. Increasing the number of coupled sites at the central region introduces a transient diffusive regime, which crosses over to logarithmic transport in the localized phase and in the delocalized regime to ballistic transport. On the other hand, when the heat bath is coupled to equally spaced sites of the system, we observe a crossover from ballistic and logarithmic transport to diffusion in the delocalized and localized regimes, respectively. We propose a classical master equation, which captures our numerical observations for both coupling configurations on a qualitative level and for some parameters, even on a quantitative level. Using the classical picture, we show that the crossover to diffusion occurs at a time that increases exponentially with the spacing between the coupled sites, and the resulting diffusion constant decreases exponentially with the spacing.

## I. INTRODUCTION

Non-equilibrium dynamics of quantum systems has been a topic of great interest in condensed matter and statistical physics, particularly concerning the study of quantum transport [1-8]. While transport in generic many-body systems is typically diffusive, diffusion of a quantum particle can be suppressed by a random disordered potential due to localization of the single-particle states. This phenomenon, dubbed Anderson localization [9], occurs for any non-zero random disorder in one and two dimensions. At higher dimensions, *all* states are localized only for sufficiently high disorder. For disorder values smaller than a critical value, only eigenstates below a specific energy, called the mobility edge, are localized [10–12].

Localization also occurs for systems with non-random, quasi-periodic potentials. Such potentials can exhibit localization-delocalization transitions and mobility edges, even in one dimension [13–20], and are experimentally feasible [21–24]. In the quasi-periodic Aubry-André-Harper (AAH) potential, for example, all states are localized above a critical potential strength and delocalized otherwise, whereas the transition point features fractal states [13, 14]. Transport is absent in the localized phase, ballistic in the delocalized phase, and anomalous at the critical point due to the fractal structure of the eigenstates [25–28].

In recent years a number of theoretical works suggested that single-particle localization with or without quenched disorder is stable to the addition of sufficiently weak inter-particle interactions, giving rise to novel dynamical phase transitions away from thermal equilibrium [3, 29–34]. This phenomenon, dubbed many-body localization (MBL), manifested in the absence of all transport for a sufficiently large disorder, though logarithmically slow transport in the localized phase was reported recently and is still under debate [35–37]. For weak dis-



Figure 1. Schematic of the coupling of the heat bath to the system. (a) central region of l sites is coupled (b) sites at  $\lambda$  distance apart are coupled.

order, there has been numerical evidence of anomalous diffusion that is attributed to the existence of rare insulating regions [4-7]. The anomalous transport was also observed in quasiperiodic systems even though such systems do not exhibit rare regions due to the deterministic nature of the potential [38-41].

The stability of the many-body localized phase was recently challenged in a number of analytical and numerical works [42–46], which argue that the MBL phase is unstable in the thermodynamic limit and at long times. One proposed mechanism for destabilizing a many-body localized phase in disordered systems is through avalanche propagation [47–53], where small thermalizing regions known as thermal inclusions act as a bath, which grow and take over the localized regions, leading to delocalization of the system. The impact of these rare regions, also called ergodic bubbles, have also been investigated in cold-atom experiments [21, 53]. Various phenomenological models of ergodic bubbles were proposed: such as random matrices [54] or coupling to a bath [48, 49, 55–60]. Coupling a Markovian bath to *all* lattice sites of an Anderson insulator is known to destroy localization and results in diffusive transport [61– 65], while for correlated noise, a transient sub-diffusive regime can be seen [66]. On the other hand, it was demonstrated that local coupling to a Markovian bath induces logarithmic transport in an Anderson insulator [58].

The stationary current in a system with a finite density of sites coupled to a heat bath, was studied inRefs. [56, 57]. These works established a transition from a superdiffusive to diffusive stationary current as a function of the density of the coupled sites. However, Refs. [56, 57], do not provide insight on the temporal dependence of spreading excitations. Moreover, there are known cases where the behavior of the stationary current is different from temporal spreading of density excitations [25, 26]. It is therefore an open question if the spreading of density excitations exhibit one or several regimes of transport, and how these regimes depends on the coupling to the heat bath.

In this work, we answer these questions by studying the spreading of density excitations at infinite temperature for different couplings of the system to a Markovian bath.

The paper is organized as follows. We describe the model Hamiltonian and the methods in Section II. In Section III we present our results for different couplings of the model to a Markovian heat bath. Finally, we discuss our findings in the Section IV.

### II. MODEL AND METHODS

We consider a system of N spinless fermions in a chain of length L, which is described by the Hamiltonian,

$$\hat{H} = -J \sum_{m=1}^{L-1} \left( \hat{a}_{m+1}^{\dagger} \hat{a}_m + \hat{a}_m^{\dagger} \hat{a}_{m+1} \right)$$
(1)  
+ $W \sum_{m=1}^{L} \cos\left(2\pi\beta m + \phi\right) \hat{n}_m,$ 

where  $\hat{n}_m = \hat{a}_m^{\dagger} \hat{a}_m$ , and  $\hat{a}_m, \hat{a}_m^{\dagger}$  are annihilation and creation operators of a fermion on site m, J is the hopping strength, W is strength of the potential and  $\beta = (\sqrt{5}-1)/2$  is the Golden mean. The phases are taken uniformly from  $\phi \in [-\pi, \pi]$ . In contrast to the Anderson model, where all the single-particle eigenstates are localized for any non-zero value of W, the AAH model exhibits a delocalization-localization transition. For W < 2J, all the single-particle eigenstates are localized. At the critical point, W = 2J, the states are multifractal and transport is diffusive if characterized by the

mean squared displacement [26], while a characterization based on the stationary current suggests a sub-diffusive transport [25, 26].

We couple the system to a Markovian heat bath which does not affect the number of fermions in the system. Specifically, we assume that the density matrix of the system evolves via the Lindblad master equation [67],

$$\frac{\partial \hat{\rho}(t)}{\partial t} = -i \left[ \hat{H}, \hat{\rho}(t) \right] + \sum_{i} \left( \hat{L}_{i} \hat{\rho}(t) \hat{L}_{i}^{\dagger} - \frac{1}{2} \left\{ \hat{L}_{i}^{\dagger} \hat{L}_{i}, \hat{\rho}(t) \right\} \right)$$
(2)

where  $\{.\}$  is the anti-commutator, the first term represents the unitary evolution and the second term gives the non-unitary dynamics. The operators  $\hat{L}_i$  are Lindblad jump operators, which we take to be

$$\hat{L}_i = \sqrt{\gamma_i} \hat{n}_i, \tag{3}$$

where  $\gamma_i$  represents the strength of the dissipation on site i. The dimensions of the density matrix are  $\mathcal{N} \times \mathcal{N}$ , where  $\mathcal{N}$  is the Hilbert space dimension. This unfavorable scaling with the system size makes the numerical solution of the Lindblad equation computationally expensive. A more efficient approach is to unravel the evolution into a *unitary* evolution of wavefunctions in the presence of white noise and then to average over the realizations of the noise to obtain the quantities of interest [68–71].

For unitary unraveling, the time evolution operator is given by,

$$\hat{U}(t+dt,t) = e^{-i\hat{H}dt - i\sqrt{\gamma \, dt} \sum_i \eta_i(t)\hat{n}_i},\tag{4}$$

where  $\eta_i(t)$  are independent Gaussian random variables with mean zero and unit variance. The density matrix can be obtained by averaging over the realizations of the noise,

$$\hat{\rho}(t+dt) = \overline{|\psi(t+dt)\rangle\langle\psi(t+dt)|},\tag{5}$$

where  $|\psi(t + dt)\rangle = \hat{U}(t + dt, t)|\psi(t)\rangle$  and the over-bar represents averaging over the noise realizations. In this work, we set the noise strength to  $\gamma = 1$  and the timestep to, dt = 0.1. We have corroborated that our results do not change if dt is further reduced. We average over 10 noise trajectories and over 10 phase realizations ( $\phi$  in Eq. (1), see Ref. [58] for further numerical details). We have found this averaging sufficient to reduce the statistical uncertainty of the data. To study the transport properties, we consider the following observables:

Particle transport at infinite temperature. It is easy to check that irrespective of  $\hat{H}$  the RHS of Eq. (2) vanishes for Hermitian Lindblad jump operators and  $\hat{\rho} \propto \mathbb{I}$ , such that any initial state will reach an infinite temperature state. Therefore, to avoid transient effects we directly characterize particle transport at infinite temperature. For this purpose we look at the two-point density-density correlation function,

$$C_{i}(t) = \operatorname{Tr}\left[\hat{\rho}_{\infty}\left(\hat{n}_{i}(t) - \frac{1}{2}\right)\left(\hat{n}_{L/2} - \frac{1}{2}\right)\right], \quad (6)$$

where  $\hat{\rho}_{\infty} = \mathbb{I}/\mathcal{N}$  is the infinite-temperature density matrix, and  $\hat{n}_{L/2}$  corresponds to an initial density excitation at the center of the lattice. For non-interacting systems, Eq. (6) can be written as

$$C_{i}\left(t\right) = \left|\operatorname{Tr}\left[\hat{\rho}_{\infty}\hat{a}_{i}^{\dagger}\left(t\right)\hat{a}_{L/2}\right]\right|^{2} = \frac{1}{4}\left|U_{i,L/2}^{s}\left(t,0\right)\right|^{2},$$

where  $U_{i,L/2}^{s}(t,0)$  is the single-particle propagator. The second equality follows, since for unitary unraveling  $\hat{a}_{i}^{\dagger}(t)$  can be written as:  $\hat{a}_{i}^{\dagger}(t) = \sum_{k} U_{ik}^{s}(t,0) \hat{a}_{k}^{\dagger}$  and  $\operatorname{Tr}\left[\hat{\rho}_{\infty}\hat{a}_{i}^{\dagger}\hat{a}_{j}\right] = \frac{1}{2}\delta_{ij}$ . Since  $U_{i,L/2}^{s}(t,0)$  has dimensions  $L \times L$ , this allows us to efficiently study the system for large L.

We characterize the nature of transport using the root mean squared displacement (RMSD) [4, 6, 72, 73],

$$\tilde{R}(t) = \sqrt{\sum_{i=1}^{L} \left(i - \frac{L}{2}\right)^2 C_i(t)}.$$
(7)

Typically, the RMSD grows as a power law in time,  $\tilde{R}(t) \sim t^{\alpha}$ , where the dynamical exponent  $\alpha = 1/2$  corresponds to diffusive transport and  $\alpha = 1$  to systems with ballistic transport. Regimes characterized by subdiffusive and super-diffusive transport involve exponents ranging between  $0 \leq \alpha < 0.5$  and  $0.5 < \alpha < 1$ , respectively. For localized systems  $\alpha = 0$ .

Dynamics of entanglement entropy. While the entanglement entropy is not a good measure of quantum information for mixed states [74], for the unitary unraveling, it is well defined for each of the trajectories. Therefore, in situations when one can physically justify the specific form of the unraveling [75], the entanglement averaged over the various trajectories is a sensible quantity [58, 76]. Another advantage of the unitary unraveling is that the wavefunction  $|\psi(t)\rangle$ , is Gaussian through the entire evolution, which allows an efficient computation of the entanglement entropy, using the relation,

$$S(t) = -\sum_{\alpha} \left[ c_{\alpha}(t) \ln c_{\alpha}(t) + (1 - c_{\alpha}(t)) \ln (1 - c_{\alpha}(t)) \right].$$
(8)

Here,  $c_{\alpha}(t)$  are eigenvalues of the correlation function  $\langle \psi(t) | \hat{a}_i^{\dagger} \hat{a}_j | \psi(t) \rangle$ , restricted to the subsystem of interest [77, 78]. For ballistic transport, the entanglement entropy grows linearly with time, while anomalous transport is characterized by a sub-linear growth [79–82].



Figure 2. Root mean-squared displacement  $\tilde{R}(t)$  (left panels) and entanglement entropy S(t) (right panels) as a function of time; for system size L = 1000. The top panels, which are plotted on a log-log scale, correspond to the delocalized phase and the critical point. The bottom panels, which are plotted on a semi-log scale, correspond to the localized phase. The black dashed and dotted lines provide a guide to ballistic and diffusive transport, respectively. The statistical error is of the order of the line width.

### III. RESULTS

In this Section, we discuss our results for local coupling the heat bath to one site of the system and to a finite fraction of system sites. In finite fraction coupling we either couple the heat bath to a central region of the chain or to equally separated lattice sites through the entire chain (Fig.1). All the results below are obtained for a chain length of L = 1000.

# A. Local coupling

In the left panels of Figure 2 we plot the dynamics of the root mean squared displacement for the delocalized, critical, and localized phases of the AAH model. In the absence of coupling to the heat bath, particle transport in these phases is ballistic, anomalous and absent, respectively [25, 26]. As can be seen from Fig. 2(a), local heat bath does not affect the nature of transport in the delocalized and critical regimes. On the other hand, it induces logarithmic transport in the localized phase, similar to the case of the disordered Anderson model [58]. Within the time range considered, there is no sign of a crossover to diffusion, as opposed to the case where the noise is coupled to *all* the sites [61, 62, 65, 66, 83].

In the right panels of Fig. 2 we present the growth of the entanglement entropy starting from a random product state. We use a random product state so that initially the entanglement entropy is zero, which provides us with a finite regime of entanglement entropy growth. In the delocalized regime, the growth of the entangle-



Figure 3. Root mean-squared displacement  $\tilde{R}(t)$ , as a function of time for L = 1000. Left panels correspond to coupling of the heat bath to a central region of length l. Right panels to coupling the heat bath to system sites which are  $\lambda$  distance apart. Top panel correspond to W = 1 and bottom panels to W = 4. Dashed and dotted lines are guides to the eye for diffusive and ballistic transport, respectively. More intense colors stand for larger  $\lambda$  or l.

ment entropy is unaffected by the presence of the local heat bath and is consistent with a power-law dependence on time, observed in closed systems [84]. In the localized case, the heat bath induces logarithmic growth of entanglement entropy which is reminiscent of the entanglement entropy growth in the MBL phase [85–88]. However, here this growth is accompanied with logarithmic particle transport, while the presence of logarithmic particle transport in the MBL phase is under debate [35–37].

# B. Coupling to a finite part of the chain

We have seen that local coupling to the heat bath is not affecting transport in the delocalized phases and is inducing logarithmic transport in the localized phase. In this section we study how transport is affected when the heat bath is coupled to a finite fraction of system sites. Moreover, we will show that the spatial configuration of the coupled sites is important. Specifically we consider two different configurations: coupling l < L sites at the central region of the chain, or coupling sites which are separated a distance  $\lambda$  apart, such that their density is  $1/\lambda$ .

The left panels of Fig. 3 show the dynamics of R(t)in the delocalized phase (top panel) and the localized phase (bottom panel) for different widths l of the coupled central region. In both delocalized and localized phases  $\tilde{R}(t)$  initially grows diffusively. After this initial diffusive growth it crosses over to ballistic transport (Fig. 3(a)) in the delocalized phase, or to logarithmic transport in the localized phase (Fig. 3(c)). In both cases the crossover time, which we will designate by  $t_l$ , grows with the width



Figure 4. Root mean-squared displacement  $\tilde{R}(t)$ , as a function of time; for a number of coupled central region widths l (top panel) or the distance between the coupled sites,  $\lambda$  (bottom panel). Solid lines correspond to the the numerical solution of the Lindblad equation (2), and dashed lines to the solution of the classical master equation (9), with time rescaled by a factor of 20 to have the best fit with the solid lines. The parameters used are L = 1000 and W = 4.

of the coupled region, l.

The right panels of Fig. 3 show  $\hat{R}(t)$  for coupling sites which are a distance  $\lambda$  apart from each other. In the delocalized and localized phases, we observe a crossover to diffusion. At the critical phase, transport remains practically unaffected by the heat bath (not shown). In the delocalized phase, the initial transport is ballistic (see Fig. 3(b)) and in the localized phase, the initial transport is logarithmic (see Fig. 3(d)). In all cases the crossover time, which we denote by  $t_d$  increases with  $\lambda$ . In the next section we introduce a classical master equation, which provides an explanation to the dependence of the crossover times on the parameters of the system.

# C. Classical picture

The dephasing mechanism of the heat bath, diminishes the importance of interference effects, and gives hope that classical treatment might be sufficient to understand the underlying phenomenology. We therefore follow the variable-range hopping approach of Mott [89], and assume that coupling the system to the heat bath induces transitions between localized single-particle eigenstates.



Figure 5. Same as Fig. 4 but with the time axis rescaled by the crossover times  $t_l$  or  $t_d$  (see main text).

The probability to find a particle in a single-particle state  $\alpha$ , which we denote by  $p_{\alpha}$ , evolves according to a classical master equation [83, 90, 91],

$$\partial_t p_\alpha = \sum_\beta \left( \Gamma_{\alpha\beta} p_\beta - \Gamma_{\beta\alpha} p_\alpha \right). \tag{9}$$

where the transition rates  $\Gamma_{\alpha\beta}$  between the eigenstates  $\alpha$  and  $\beta$  can be calculated from first-order perturbation theory (see Appendix A),

$$\Gamma_{\alpha\beta} = \Gamma_{\beta\alpha} = \gamma \sum_{k \in \text{coupled sites}} \left| \phi_{\beta}^{*}\left(k\right) \phi_{\alpha}\left(k\right) \right|^{2}, \quad (10)$$

where  $\gamma$  is the strength of the coupling,  $\phi_{\alpha}(i)$  are the single particles states in the position basis, and the sum k runs over the sites coupled to the heat bath. In the localized phase, by definition,  $\phi_{\alpha}(i)$  decays exponentially. We order the single-particle eigenfunctions such that  $\phi_{\alpha}(i)$  has it center of mass around a site  $\alpha$ , which allows us to approximate the transition rates as,

$$\Gamma_{\alpha\beta} = \gamma \sum_{k \in \text{coupled sites}} e^{-|\alpha-k|/\xi} e^{-|\beta-k|/\xi}, \qquad (11)$$



Figure 6. Extraction of the diffusion coefficient for various  $\lambda$  by fitting  $Dt^{1/2}$  to  $\tilde{R}(t)$  obtained from the solution of the Lindblad equation (dashed lines, top panel). The bottom panel shows the diffusion coefficient as a function of  $\lambda$  for W = 4 and L = 1000. The dashed blue line corresponds to the theoretical prediction, with no fitting parameters.

where  $\xi$  is the localization, which for the AAH model is  $\xi = 1/\ln \frac{W}{2}$  [84]. The coupled site is, therefore, initiating transitions, or scattering, between localized states within its neighborhood. Transitions to far-lying states are exponentially suppressed with the distance from the coupled site. For local coupling, this leads to logarithmic transport,  $\tilde{R}(t) \sim \xi \ln (Jt)$  [58].

When the heat bath is coupled to a region of final width l, the transitions rates  $\Gamma_{\alpha\beta}$  are approximately constant in this region, and therefore a particle initiated in the coupled region is expected to diffuse. It takes the particle time,  $t_l \sim l^2$ , to leave this region. After leaving the coupled region, transitions rates exponentially decay with the distance from the region, and transport in the system is expected to be equivalent to the situation of local coupling. When the coupled sites are at equal distances  $\lambda$  apart, on a time-scale of moving between two nearby coupled sites, transport is logarithmic, but at larger time-scales the expected motion is diffusive. The crossover time can be obtained from  $\lambda = \xi \ln (Jt_d)$ , yielding  $t_d \sim J^{-1} \exp(\lambda/\xi)$ , and the diffusion constant as  $D \sim \lambda^2/t_d = \lambda^2 \exp(-\lambda/\xi)$ .

In Fig. 4 we compare the numerical solution of the classical master equation (9) to the numerical solution of the Lindblad equation (2). Since the classical rates (11) are obtained phenomenologically, the overall prefactor of the rates cannot be determined from microscopic considerations. We determine it by rescaling the time axis of the classical master equation such that the correspondence with the solution of the Lindblad equation is optimal (2). Apart from this trivial rescaling of the units of time, there are no fitting parameters. Remarkably, the agreement between the classical master equation and the Lindblad equation goes beyond the qualitative level for the central coupled region (top panel). For equal spacing coupling, there is still qualitative agreement, but the quantitative agreement is reasonable only for small  $\lambda/\xi$ .

In Fig. 5 we test the predictions of the classical theory for the crossover times, by rescaling of the time axis by  $t_d$ or  $t_l$ , respectively. We see that such a rescaling correctly identifies the crossover time for both couplings to the heat bath when either l or  $\lambda$  are varied. The prediction for the diffusion constant is verified in Fig. 6. The agreement is not quantitative, however the exponential decrease of the diffusion constant with  $\lambda$  is nicely captured.

# IV. DISCUSSION

Using the unitary unraveling of the Lindblad master equation, we study the dynamical properties of the AAH model coupled to a dephasing heat bath. We consider local coupling and coupling to a finite part of the chain. This setup is partly motivated by dynamics in MBL systems in the presence of finite density of ergodic bubbles [7, 48, 49, 92], with the crucial difference that it is dissipative.

For local coupling of the heat bath in the delocalized and critical phases of the AAH model, we didn't observe any qualitative effects of the heat bath on the dynamics of the particle. On the other hand, in the localized phase, the root mean square displacement, entanglement entropy, and average energy (not shown) show asymptotic logarithmic growth, as it occurs for the one-dimensional Anderson insulator in the presence of a local noise [58]. Suppose the region of the coupling to the bath is of finite width. In that case, we find a regime of transient diffusion, which crosses over to ballistic transport in the delocalized phase and logarithmic transport in the localized phase. We have shown that this crossover time increases as the square of the width of the region, as typical for diffusion. When the heat bath is coupled to the system on equally spaced sites, initial transport in the system is similar to the transport with local coupling. Eventually, it crosses over to diffusion in all phases. Specifically, in the localized phase, the crossover time increases exponentially with the distance between the coupled sites.

We have shown that a classical master equation with

transition rates that exponentially decay from the location of the coupled sites captures all the observed phenomenology. Moreover, it provides accurate predictions of crossover times for all studied couplings of the heat bath. This time scale is set by the time it takes for a particle to traverse the distance  $\lambda$  between two nearby coupled sites. In the delocalized phase, this time is proportional to  $\lambda$  (if the transport is ballistic) or  $\lambda^2$  (if it is diffusive). On the other hand, it scales as  $t_d \propto \exp(\lambda/\xi)$  in the localized phase. Within the classical model, the motion of the particle between the coupled sites can be viewed as a random walk with a spatial step of  $\lambda$  and a mean-free time of  $t_d$ , which means that the motion is diffusive, with diffusion constant given by  $D \sim \lambda^2/t_d = \lambda^2 \exp(-\lambda/\xi)$ .

In this work, we have focused only on fixed  $\lambda$ , such that D is also fixed. If  $\lambda$  is allowed to vary randomly, such that its distribution is unbounded, then the average time to transition between coupled sites,  $t_d$ , can diverge. In this case, the average diffusion coefficient will vanish, and transport will be subdiffusive (see Refs. [56, 57] and Appendix B). Since sites coupled to a heat bath can be thought as perfect ergodic bubbles, we argue that our results put an upper bound on transport in MBL systems due to the presence of ergodic bubbles [7, 48, 49, 92].

Here, we focus on a quasi-periodic model without mobility edges. In the presence of mobility edges, at least in principle, the coupling to the heat bath can create transitions between the localized and delocalized states. Several interesting questions arise for these models: would the coupling eliminate the intermittent logarithmic transport regime? How will the dynamics depend on the initial conditions? We leave these questions to future studies.

This research was supported by a grant from the United States-Israel Binational Foundation (BSF, Grant No. 2019644), Jerusalem, Israel, and the United States National Science Foundation (NSF, Grant No. DMR-1936006), and by the Israel Science Foundation (grants No. 527/19 and 218/19). D.S.B acknowledges funding from the Kreitman fellowship.

- [2] Anatoli Polkovnikov, Krishnendu Sengupta, Alessandro Silva, and Mukund Vengalattore, "Colloquium : Nonequilibrium dynamics of closed interacting quantum systems," Rev. Mod. Phys. 83, 863–883 (2011).
- [3] Rahul Nandkishore and David A. Huse, "Many-body localization and thermalization in quantum statistical mechanics," Annual Review of Condensed Matter Physics 6, 15–38 (2015).
- [4] Yevgeny Bar Lev, Guy Cohen, and David R. Reichman, "Absence of Diffusion in an Interacting System of Spinless Fermions on a One-Dimensional Disordered Lattice," Phys. Rev. Lett. 114, 100601 (2015).

Marcos Rigol, Vanja Dunjko, and Maxim Olshanii, "Thermalization and its mechanism for generic isolated quantum systems," Nature 452, 854–858 (2008).

- [5] Kartiek Agarwal, Sarang Gopalakrishnan, Michael Knap, Markus Müller, and Eugene Demler, "Anomalous Diffusion and Griffiths Effects Near the Many-Body Localization Transition," Phys. Rev. Lett. **114**, 160401 (2015).
- [6] David J. Luitz and Yevgeny Bar Lev, "The ergodic side of the many-body localization transition," Annalen der Physik 529, 1600350 (2017).
- [7] Kartiek Agarwal, Ehud Altman, Eugene Demler, Sarang Gopalakrishnan, David A. Huse, and Michael Knap, "Rare-region effects and dynamics near the manybody localization transition," Annalen der Physik 529, 1600326 (2017).
- [8] Kartiek Agarwal, Sriram Ganeshan, and R. N. Bhatt, "Localization and transport in a strongly driven Anderson insulator," Phys. Rev. B 96, 014201 (2017).
- [9] P. W. Anderson, "Absence of Diffusion in Certain Random Lattices," Phys. Rev. 109, 1492–1505 (1958).
- [10] E. Abrahams, P. W. Anderson, D. C. Licciardello, and T. V. Ramakrishnan, "Scaling Theory of Localization: Absence of Quantum Diffusion in Two Dimensions," Phys. Rev. Lett. 42, 673–676 (1979).
- [11] Ferdinand Evers and Alexander D. Mirlin, "Anderson transitions," Rev. Mod. Phys. 80, 1355–1417 (2008).
- [12] B Kramer and A MacKinnon, "Localization: theory and experiment," Rep. Prog. Phys. 56, 1469–1564 (1993).
- [13] Serge Aubry and Gilles André, "Analyticity breaking and anderson localization in incommensurate lattices," Ann. Israel Phys. Soc 3, 18 (1980).
- [14] P G Harper, "Single band motion of conduction electrons in a uniform magnetic field," Proceedings of the Physical Society. Section A 68, 874–878 (1955).
- [15] J. Biddle, D. J. Priour, B. Wang, and S. Das Sarma, "Localization in one-dimensional lattices with non-nearestneighbor hopping: Generalized anderson and aubryandré models," Phys. Rev. B 83, 075105 (2011).
- [16] Sriram Ganeshan, J. H. Pixley, and S. Das Sarma, "Nearest Neighbor Tight Binding Models with an Exact Mobility Edge in One Dimension," Phys. Rev. Lett. 114, 146601 (2015).
- [17] Dong-Ling Deng, Sriram Ganeshan, Xiaopeng Li, Ranjan Modak, Subroto Mukerjee, and J. H. Pixley, "Manybody localization in incommensurate models with a mobility edge," Annalen der Physik **529**, 1600399 (2017).
- [18] Xiao Li, Xiaopeng Li, and S. Das Sarma, "Mobility edges in one-dimensional bichromatic incommensurate potentials," Phys. Rev. B 96, 085119 (2017).
- [19] Yucheng Wang, Xu Xia, Long Zhang, Hepeng Yao, Shu Chen, Jiangong You, Qi Zhou, and Xiong-Jun Liu, "Onedimensional quasiperiodic mosaic lattice with exact mobility edges," Phys. Rev. Lett. **125**, 196604 (2020).
- [20] Alexander Duthie, Sthitadhi Roy, and David E. Logan, "Self-consistent theory of mobility edges in quasiperiodic chains," Phys. Rev. B 103, L060201 (2021).
- [21] Henrik P. Lüschen, Pranjal Bordia, Sebastian Scherg, Fabien Alet, Ehud Altman, Ulrich Schneider, and Immanuel Bloch, "Observation of Slow Dynamics near the Many-Body Localization Transition in One-Dimensional Quasiperiodic Systems," Phys. Rev. Lett. **119**, 260401 (2017).
- [22] Henrik P. Lüschen, Pranjal Bordia, Sean S. Hodgman, Michael Schreiber, Saubhik Sarkar, Andrew J. Daley, Mark H. Fischer, Ehud Altman, Immanuel Bloch, and Ulrich Schneider, "Signatures of Many-Body Localization in a Controlled Open Quantum System," Phys. Rev. X

**7**, 011034 (2017).

- [23] Henrik P. Lüschen, Sebastian Scherg, Thomas Kohlert, Michael Schreiber, Pranjal Bordia, Xiao Li, S. Das Sarma, and Immanuel Bloch, "Single-Particle Mobility Edge in a One-Dimensional Quasiperiodic Optical Lattice," Phys. Rev. Lett. **120**, 160404 (2018).
- [24] Fangzhao Alex An, Karmela Padavić, Eric J Meier, Suraj Hegde, Sriram Ganeshan, JH Pixley, Smitha Vishveshwara, and Bryce Gadway, "Interactions and mobility edges: Observing the generalized aubry-andré model," Phys. Rev. Lett. **126**, 040603 (2021).
- [25] Vipin Kerala Varma, Clélia de Mulatier, and Marko Žnidarič, "Fractality in nonequilibrium steady states of quasiperiodic systems," Phys. Rev. E 96, 032130 (2017).
- [26] Archak Purkayastha, Sambuddha Sanyal, Abhishek Dhar, and Manas Kulkarni, "Anomalous transport in the Aubry-André-Harper model in isolated and open systems," Phys. Rev. B 97, 174206 (2018).
- [27] Gim Seng Ng and Tsampikos Kottos, "Wavepacket dynamics of the nonlinear Harper model," Phys. Rev. B 75, 205120 (2007).
- [28] R. Ketzmerick, K. Kruse, S. Kraut, and T. Geisel, "What Determines the Spreading of a Wave Packet?" Phys. Rev. Lett. **79**, 1959–1963 (1997).
- [29] D.M. Basko, I.L. Aleiner, and B.L. Altshuler, "Metal-insulator transition in a weakly interacting many-electron system with localized single-particle states," Annals of Physics **321**, 1126–1205 (2006).
- [30] I. V. Gornyi, A. D. Mirlin, and D. G. Polyakov, "Interacting electrons in disordered wires: Anderson localization and low-t transport," Phys. Rev. Lett. 95, 206603 (2005).
- [31] Shankar Iyer, Vadim Oganesyan, Gil Refael, and David A. Huse, "Many-body localization in a quasiperiodic system," Phys. Rev. B 87, 134202 (2013).
- [32] John Z. Imbrie, "On Many-Body Localization for Quantum Spin Chains," J Stat Phys 163, 998–1048 (2016).
- [33] Fabien Alet and Nicolas Laflorencie, "Many-body localization: An introduction and selected topics," Comptes Rendus Physique 19, 498–525 (2018).
- [34] Dmitry A. Abanin and Zlatko Papić, "Recent progress in many-body localization," Annalen der Physik 529, 1700169 (2017).
- [35] Maximilian Kiefer-Emmanouilidis, Razmik Unanyan, Michael Fleischhauer, and Jesko Sirker, "Evidence for unbounded growth of the number entropy in many-body localized phases," Phys. Rev. Lett. **124**, 243601 (2020).
- [36] David J. Luitz and Yevgeny Bar Lev, "Absence of slow particle transport in the many-body localized phase," Phys. Rev. B 102, 100202 (2020).
- [37] Maximilian Kiefer-Emmanouilidis, Razmik Unanyan, Michael Fleischhauer, and Jesko Sirker, "Particle fluctuations and the failure of simple effective models for manybody localized phases," SciPost Phys. 12, 034 (2022).
- [38] Yevgeny Bar Lev, Dante M. Kennes, Christian Klöckner, David R. Reichman, and Christoph Karrasch, "Transport in quasiperiodic interacting systems: From superdiffusion to subdiffusion," EPL **119**, 37003 (2017).
- [39] Shenglong Xu, Xiao Li, Yi-Ting Hsu, Brian Swingle, and S. Das Sarma, "Butterfly effect in interacting Aubry-Andre model: Thermalization, slow scrambling, and many-body localization," Phys. Rev. Research 1, 032039 (2019).
- [40] Yongchan Yoo, Junhyun Lee, and Brian Swingle,

"Nonequilibrium steady state phases of the interacting Aubry-André-Harper model," Phys. Rev. B **102**, 195142 (2020).

- [41] Marko Žnidarič, "Comment on "Nonequilibrium steady state phases of the interacting Aubry-André-Harper model"," Phys. Rev. B 103, 237101 (2021).
- [42] Jan Šuntajs, Janez Bonča, Tomaž Prosen, and Lev Vidmar, "Quantum chaos challenges many-body localization," Phys. Rev. E 102, 062144 (2020).
- [43] Maximilian Kiefer-Emmanouilidis, Razmik Unanyan, Michael Fleischhauer, and Jesko Sirker, "Slow delocalization of particles in many-body localized phases," Phys. Rev. B 103, 024203 (2021).
- [44] Dries Sels and Anatoli Polkovnikov, "Dynamical obstruction to localization in a disordered spin chain," Phys. Rev. E 104, 054105 (2021).
- [45] Piotr Sierant and Jakub Zakrzewski, "Challenges to observation of many-body localization," Phys. Rev. B 105, 224203 (2022).
- [46] Dries Sels, "Bath-induced delocalization in interacting disordered spin chains," Phys. Rev. B 106, L020202 (2022).
- [47] Wojciech De Roeck, Francois Huveneers, Markus Müller, and Mauro Schiulaz, "Absence of many-body mobility edges," Phys. Rev. B 93, 014203 (2016).
- [48] Wojciech De Roeck and François Huveneers, "Stability and instability towards delocalization in many-body localization systems," Phys. Rev. B 95, 155129 (2017).
- [49] David J Luitz, François Huveneers, and Wojciech De Roeck, "How a small quantum bath can thermalize long localized chains," Phys. Rev. Lett. **119**, 150602 (2017).
- [50] Thimothée Thiery, François Huveneers, Markus Müller, and Wojciech De Roeck, "Many-body delocalization as a quantum avalanche," Phys. Rev. Lett. **121**, 140601 (2018).
- [51] P. J. D. Crowley and A. Chandran, "Avalanche induced coexisting localized and thermal regions in disordered chains," Phys. Rev. Res. 2, 033262 (2020).
- [52] Alan Morningstar, Luis Colmenarez, Vedika Khemani, David J. Luitz, and David A. Huse, "Avalanches and many-body resonances in many-body localized systems," Phys. Rev. B 105, 174205 (2022).
- [53] Julian Léonard, Sooshin Kim, Matthew Rispoli, Alexander Lukin, Robert Schittko, Joyce Kwan, Eugene Demler, Dries Sels, and Markus Greiner, "Probing the onset of quantum avalanches in a many-body localized system," Nature Physics 19, 481–485 (2023).
- [54] M. Schiró and M. Tarzia, "Toy model for anomalous transport and griffiths effects near the many-body localization transition," Phys. Rev. B 101, 014203 (2020).
- [55] Rahul Nandkishore, Sarang Gopalakrishnan, and David A. Huse, "Spectral features of a many-bodylocalized system weakly coupled to a bath," Phys. Rev. B 90, 064203 (2014).
- [56] Scott R. Taylor and Antonello Scardicchio, "Subdiffusion in a one-dimensional Anderson insulator with random dephasing: Finite-size scaling, Griffiths effects, and possible implications for many-body localization," Phys. Rev. B 103, 184202 (2021).
- [57] Xhek Turkeshi, Damien Barbier, Leticia Cugliandolo, Marco Schiro, and Marco Tarzia, "Destruction of localization by thermal inclusions: Anomalous transport and Griffiths effects in the Anderson and André-Aubry-

Harper models," SciPost Phys. 12, 189 (2022).

- [58] Talía L. M. Lezama and Yevgeny Bar Lev, "Logarithmic, noise-induced dynamics in the Anderson insulator," SciPost Phys. 12, 174 (2022).
- [59] Pietro Brighi, Alexios A. Michailidis, Dmitry A. Abanin, and Maksym Serbyn, "Propagation of many-body localization in an anderson insulator," Phys. Rev. B 105, L220203 (2022).
- [60] Bozhen Zhou, Xueliang Wang, and Shu Chen, "Exponential size scaling of the liouvillian gap in boundarydissipated systems with anderson localization," Phys. Rev. B 106, 064203 (2022).
- [61] Marko Žnidarič, "Dephasing-induced diffusive transport in the anisotropic heisenberg model," New Journal of Physics 12, 043001 (2010).
- [62] Marko Žnidarič and Martin Horvat, "Transport in a disordered tight-binding chain with dephasing," The European Physical Journal B 86, 67 (2013).
- [63] Jeremy M Moix, Michael Khasin, and Jianshu Cao, "Coherent quantum transport in disordered systems: I. the influence of dephasing on the transport properties and absorption spectra on one-dimensional systems," New Journal of Physics 15, 085010 (2013).
- [64] Mariya V Medvedyeva, Tomaž Prosen, and Marko Žnidarič, "Influence of dephasing on many-body localization," Phys. Rev. B 93, 094205 (2016).
- [65] Marko Žnidarič, Juan Jose Mendoza-Arenas, Stephen R. Clark, and John Goold, "Dephasing enhanced spin transport in the ergodic phase of a many-body localizable system," Annalen der Physik 529, 1600298 (2017).
- [66] Sarang Gopalakrishnan, K. Ranjibul Islam, and Michael Knap, "Noise-Induced Subdiffusion in Strongly Localized Quantum Systems," Phys. Rev. Lett. **119**, 046601 (2017).
- [67] G. Lindblad, "On the generators of quantum dynamical semigroups," Communications in Mathematical Physics 48, 119 – 130 (1976).
- [68] Todd A. Brun, "Continuous measurements, quantum trajectories, and decoherent histories," Phys. Rev. A 61, 042107 (2000).
- [69] Howard M Wiseman and Lajos Diósi, "Complete parameterization, and invariance, of diffusive quantum trajectories for markovian open systems," Chemical Physics 268, 91–104 (2001).
- [70] Crispin Gardiner and Peter Zoller, Quantum noise: a handbook of Markovian and non-Markovian quantum stochastic methods with applications to quantum optics (Springer Science & Business Media, 2004).
- [71] A. Chenu, M. Beau, J. Cao, and A. del Campo, "Quantum Simulation of Generic Many-Body Open System Dynamics Using Classical Noise," Phys. Rev. Lett. 118, 140403 (2017).
- [72] R. Steinigeweg, H. Wichterich, and J. Gemmer, "Density dynamics from current auto-correlations at finite timeand length-scales," Europhysics Letters 88, 10004 (2009).
- [73] R. Steinigeweg, F. Jin, D. Schmidtke, H. De Raedt, K. Michielsen, and J. Gemmer, "Real-time broadening of nonequilibrium density profiles and the role of the specific initial-state realization," Phys. Rev. B 95, 035155 (2017).
- [74] Jens Eisert and MB Plenio, "Introduction to the basics of entanglement theory in continuous-variable systems," Int. J. Quantum Inf. 01, 479–506 (2003).
- [75] For example, a system in a time-dependent noisy potential.

- [76] Xiangyu Cao, Antoine Tilloy, and Andrea De Luca, "Entanglement in a fermion chain under continuous monitoring," SciPost Phys. 7, 24 (2019).
- [77] Ingo Peschel, "Calculation of reduced density matrices from correlation functions," J. Phys. A: Math. Gen. 36, L205–L208 (2003).
- [78] Ingo Peschel and Viktor Eisler, "Reduced density matrices and entanglement entropy in free lattice models," J. Phys. A: Math. Theor. 42, 504003 (2009).
- [79] Pasquale Calabrese and John Cardy, "Evolution of entanglement entropy in one-dimensional systems," Journal of Statistical Mechanics: Theory and Experiment 2005, P04010 (2005).
- [80] Hyungwon Kim and David A. Huse, "Ballistic spreading of entanglement in a diffusive nonintegrable system," Phys. Rev. Lett. 111, 127205 (2013).
- [81] David J. Luitz, Nicolas Laflorencie, and Fabien Alet, "Extended slow dynamical regime close to the manybody localization transition," Phys. Rev. B 93, 060201 (2016).
- [82] Talía L. M. Lezama and David J. Luitz, "Power-law entanglement growth from typical product states," Phys. Rev. Res. 1, 033067 (2019).
- [83] Ariel Amir, Yoav Lahini, and Hagai B. Perets, "Classical diffusion of a quantum particle in a noisy environment," Phys. Rev. E 79, 050105 (2009).
- [84] Gergő Roósz, Uma Divakaran, Heiko Rieger, and Ferenc Iglói, "Nonequilibrium quantum relaxation across a localization-delocalization transition," Phys. Rev. B 90, 184202 (2014).
- [85] Marko Žnidarič, Tomaž Prosen, and Peter Prelovšek, "Many-body localization in the heisenberg xxz magnet in a random field," Phys. Rev. B 77, 064426 (2008).
- [86] Jens H. Bardarson, Frank Pollmann, and Joel E. Moore, "Unbounded growth of entanglement in models of manybody localization," Phys. Rev. Lett. 109, 017202 (2012).
- [87] Arun Nanduri, Hyungwon Kim, and David A. Huse, "Entanglement spreading in a many-body localized system," Phys. Rev. B 90, 064201 (2014).
- [88] Marko Žnidarič, "Entanglement in a dephasing model and many-body localization," Phys. Rev. B 97, 214202 (2018).
- [89] N. F. Mott, "Conduction in non-crystalline materials," Philos. Mag. 19, 835–852 (1969).
- [90] Ariel Amir, Yuval Oreg, and Yoseph Imry, "Localization, Anomalous Diffusion, and Slow Relaxations: A Random Distance Matrix Approach," Phys. Rev. Lett. 105, 070601 (2010).
- [91] Mark H Fischer, Mykola Maksymenko, and Ehud Altman, "Dynamics of a many-body-localized system coupled to a bath," Phys. Rev. Lett. 116, 160401 (2016).
- [92] Vedika Khemani, S. P. Lim, D. N. Sheng, and David A. Huse, "Critical properties of the many-body localization transition," Phys. Rev. X 7, 021013 (2017).

# Appendix A: Transition rates for white noise

In this section, we calculate the transition rates using the first order perturbation theory. Withing the unitary unraveling of the Lindblad equation the dynamics of the system is described by a the time-dependent Hamiltonian,

$$\hat{H}(t) = \hat{H}_0 + \hat{V}(t) \,,$$

where,  $\hat{V}(t) = \sqrt{\gamma} \sum_{i} \eta_i(t) |i\rangle \langle i|$  with  $\eta_i(t)$  being a Gaussian random variable with zero mean and unit variance. The noise is characterized by the correlation function,  $\langle \eta_i(t) \eta_j(t') \rangle = \delta_{ij} \delta(t-t')$ . The time evolution of the state  $|\psi(t)\rangle$  in the eigenbasis  $|\alpha\rangle$  of  $\hat{H}_0$  can be written as,

$$\partial_t c_\alpha = -i \sum_{\gamma} \left\langle \alpha \left| \hat{V}(t) \right| \gamma \right\rangle c_\gamma(t) \, e^{-i(\epsilon_\gamma - \epsilon_\alpha)t},\tag{A1}$$

where  $\epsilon_{\alpha}$  is the eigenvalue of  $\hat{H}_0$  which corresponds to  $|\alpha\rangle$ . In the integral form, this corresponds to,

$$c_{\alpha}(t) = c_{\alpha}(0) - i \int_{0}^{t} \sum_{\gamma} dt' \left\langle \alpha \left| \hat{V}(t) \right| \gamma \right\rangle c_{\gamma}(t) e^{-i(\epsilon_{\gamma} - \epsilon_{\alpha})t'}.$$
(A2)

Back substitution of the LHS into the RHS and ignoring terms of order,  $O(V^2)$  gives,

$$c_{\alpha}(t) = c_{\alpha}(0) - i \int_{0}^{t} \sum_{\gamma} dt' \left\langle \alpha \left| \hat{V}(t) \right| \gamma \right\rangle c_{\gamma}(0) e^{-i(\epsilon_{\gamma} - \epsilon_{\alpha})t'}.$$
(A3)

We assume that the system is initialized in state  $\beta$ , such that  $c_{\gamma}(0) = \delta_{\gamma\beta}$ , which yields,

$$c_{\alpha}(t) = \delta_{\alpha\beta} - i \int_{0}^{t} dt' \left\langle \alpha \left| \hat{V}(t) \right| \beta \right\rangle e^{-i(\epsilon_{\beta} - \epsilon_{\alpha})t'}.$$
(A4)

The transition probability rate between state  $|\alpha\rangle$  and  $|\beta\rangle$  is given by

$$\Gamma_{\alpha\beta} = \left| c_{\alpha}\left(t\right) \right|^{2}/t, = \frac{1}{t} \int_{0}^{t} dt' \int_{0}^{t} dt'' \left\langle \alpha \left| \hat{V}\left(t'\right) \right| \beta \right\rangle \left\langle \beta \left| \hat{V}\left(t''\right) \right| \alpha \right\rangle e^{-i(\epsilon_{\beta} - \epsilon_{\alpha})\left(t' - t''\right)}.$$
(A5)

Now since averaging over the noise gives,

$$\overline{\left\langle \alpha \left| \hat{V}\left(t'\right) \right| \beta \right\rangle \left\langle \beta \left| \hat{V}\left(t''\right) \right| \alpha \right\rangle} = \gamma \sum_{i,j} \overline{\eta_i\left(t'\right) \eta_j\left(t''\right)} \left\langle \alpha |i\rangle \left\langle i|\beta \right\rangle \left\langle \alpha |j\rangle \left\langle j|\beta \right\rangle = \gamma \sum_i \left| \left\langle \alpha |i\rangle \right|^2 \left| \left\langle \beta |i\rangle \right|^2 \delta\left(t' - t''\right), \quad (A6)$$

we obtain

$$\begin{split} \Gamma_{\alpha\beta} &= \frac{\gamma}{t} \sum_{i} |\langle \alpha | i \rangle|^{2} |\langle \beta | i \rangle|^{2} \int_{0}^{t} dt' \int_{0}^{t} dt'' \delta\left(t' - t''\right) e^{-i(\epsilon_{\beta} - \epsilon_{\alpha})\left(t' - t''\right)} \\ &= \gamma \sum_{i} |\langle \alpha | i \rangle|^{2} |\langle \beta | i \rangle|^{2} \,, \end{split}$$

where  $\langle \beta | i \rangle = \phi_{\beta}^{*}(i)$  and  $\langle k | \alpha \rangle = \phi_{\alpha}(k)$ .

# Appendix B: Coupling at random sites

In this section, we consider the case where a given site is coupled with probability p, which means that the distance between two coupled sites is distributed according to the Poisson distribution. In Fig. 7 we shows the dynamics of  $\tilde{R}(t)$  calculated using the classical master equation for W = 4.0 and and number of p. We average the data over 50 realizations of the couplings to the heat bath. As opposed to the case where the coupled sites are placed at an equal distance from each other leading to eventual diffusion, here we find sub-diffusive transport with a dynamical exponent which depends on p. Our results are consistent with Refs. [56, 57] where the stationary state current is calculated in a similar system.



Figure 7. Dynamics of  $\tilde{R}(t)$  for for W = 4.0, and the case where each site of the system can be coupled with probability p.