

Parameter estimation for contact tracing in graph-based models

Augustine Okolie^{a,*}, Johannes Müller^{a,b}, Mirjam Kretzschmar^c

^aCenter for Mathematical Sciences, Technische Universität München, 85748 Garching, Germany

^bInstitute for Computational Biology, Helmholtz Center Munich, 85764 Neuherberg, Germany

^cUniversity Medical Center Utrecht, Utrecht University, 3584CX Utrecht, The Netherlands

Abstract

We adopt a maximum-likelihood framework to estimate parameters of a stochastic susceptible-infected-recovered (SIR) model with contact tracing on a rooted random tree. Given the number of detectees per index case, our estimator allows to determine the degree distribution of the random tree as well as the tracing probability. Since we do not discover all infectees via contact tracing, this estimation is non-trivial. To keep things simple and stable, we develop an approximation suited for realistic situations (contact tracing probability small, or the probability for the detection of index cases small). In this approximation, the only epidemiological parameter entering the estimator is R_0 .

The estimator is tested in a simulation study and is furthermore applied to covid-19 contact tracing data from India. The simulation study underlines the efficiency of the method. For the empirical covid-19 data, we compare different degree distributions and perform a sensitivity analysis. We find that particularly a power-law and a negative binomial degree distribution fit the data well and that the tracing probability is rather large. The sensitivity analysis shows no strong dependency of the estimates on the reproduction number. Finally, we discuss the relevance of our findings.

Keywords: Stochastic SIR model on graph, contact tracing, epidemiology, branching process, parameter inference

2010 MSC: 92D30, 61D30

1. Introduction

Infectious disease models have been instrumental in the study of many infectious diseases. Usually, these models are dependent on several biological parameters which can be epidemiological such as transmission, recovery, etc, or intervention parameters such as contact tracing, screening, vaccination, and others. However, most of these parameters are not or only partially known and may cause predictions from these models to lack robustness [1] if not chosen appropriately. Missing data poses a major quantification challenge in epidemiology due to unobserved or partially observed events [2]. This makes parameter estimation essential in modelling disease spread. Often, the likelihood of parameters is maximised following the model predictions on sets of parameter values. In order to achieve parameter estimation, the model system property must be identifiable, i.e. estimating its parameters uniquely from the given data [3, 4, 5]. Several estimation methods, e.g statistically based techniques such as Approximate Bayesian computation (ABC) [6], Markov chain Monte Carlo (MCMC) integration [7], optimal control theory approach [8], classical least-squares method [9], and others (also see the review article [10]) have been instrumental in estimating parameters and making inferences in epidemic models. With respect to parameter estimation, contact tracing is particularly challenging, as we somehow need to estimate the fraction of contacts we miss to identify: We need to estimate something that is *per definitionem* unobserved.

*Corresponding author.

Email address: augustine.okolie@tum.de (A.Okolie).

Several estimation techniques have been proposed for estimating important intervention parameters in modelling the recent covid-19 pandemic. For instance, Manou-Abi et al. [11] obtained a best-fit model by proposing statistical methods for the underlying serial interval probability distribution for the covid-19 virus in Mayotte from March 2020 to January 2022. Their method was then used to estimate time-varying reproduction numbers and transmission rates observed from the collected data.

Only a few attempts have been made to identify parameters specific to contact tracing: Müller and Hösel [12] proposed a branching process approach for contact tracing in randomly mixing populations to estimate tracing probability from contact history at the onset of an epidemic, based on the theory introduced in Müller et al. [13]. The derived estimator was then applied to data from contact tracing for tuberculosis and chlamydia. Blum and Tran [6] use a Bayesian framework to estimate parameters for rates of contact tracing and detection by random screening, and the method of Dyson et al. [14] is based on fitting a Yaws and Trachoma contact tracing survey data to a stochastic household model. Tanaka et al. [15] took up the branching process approach to estimate the percentage of undiagnosed persons in the covid-19 pandemic with recursive full tracing.

In this paper, we propose methods for estimating parameters in graph-based models [16]. A stochastic SIR model on a tree-shaped contact graph is modelled such that the underlying contact structure is given by a fixed or random graph. Due to the nature of the problem, we adapt the branching process theory results for contact tracing on random trees [16] to formulate a likelihood estimator for estimating the tracing probability and expected number of contacts. We first performed a simulation study with a Poisson degree distribution to check the performance of the maximum likelihood estimator. Thereafter, we applied the model to contact tracing data collected during the covid-19 pandemic in Karnataka, India. Overall, we show that our estimator based on the branching process theory for contact tracing is well suited for estimating tracing probabilities and degree distribution of the underlying contact structure in tree-based models.

The remaining part of the paper is structured as follows: Section 2 outlines the tree model and model assumptions. Section 3 presents the distribution of ages since infection, while Section 4 discusses the distribution of detected cases from one index case. We set up a likelihood estimator for estimating the tracing probability and underlying contact structure using these results and simulated data in Section 5 followed by a sensitivity analysis in Section 6. Last, we discuss our findings in Section 7.

2. Model assumptions

For the convenience of the reader, we will first sketch the motivation and idea of the branching theory process for contact tracing on rooted random trees (tree model) in Okolie and Müller [16] for our estimation analysis. A contact network in most applications represents individuals as nodes and interaction links via edges. Interaction links are channels where individuals can have direct or indirect contact, e.g. family, school, work, etc. These contact networks are applicable and useful in analysing contact tracing because they hold information about individuals and their neighbours [17, 18]. However, applying contact networks to infectious disease dynamics is not straightforward as it requires a detailed understanding of the underlying network structure, e.g. the degree distribution and correlation, clustering coefficients, and properties defined by the network topology.

Once we have a defined contact network with pre-defined nodes and interacting links, we have a contact graph. The basic idea is to describe an epidemic by constructing a simple contact graph that is a rooted tree where only the root node is infected at the onset of the epidemic. The choice of this tree contact graph is for mathematical convenience as trees are not appropriate to describe more complex interactions for natural contact graphs. However, from a microscopic level, we can

gain a better understanding of the overall mechanism and functioning of larger and more complex graphs, as many graph models as the configuration model resemble locally a tree [19]. Then we assign independently on each edge connecting one infected and one uninfected node a probability of transmitting the disease. If we focus on edges that transmit the disease, then we have only the infection graph which is a subgraph of the contact graph. Contact tracing is also analysed on this infection graph such that upon recovery of an index case, direct neighbours of this index case are also removed with some tracing probability. From the number of detected cases by an index case via contact tracing, it is possible to estimate the degree distribution of the underlying contact network and also the tracing probability.

On the rooted random tree (see Fig. 1), the infection starts from the root node R and spreads downwards through the directed edges. Individual C which is a direct contact of the root node is infected and spreads the disease to the focal individual A . Furthermore, A also spreads the infection to B and D . Individuals B and D are the “downstreams” (infectees) of A while C is an “upstream” (infector) of A . We define K a random variable denoting the number of downstream edges of an individual with expectation $\mathbb{E}[K]$, where we assume that the downstream degree of each node is an i.i.d. realisation of K . We note that the root node R is special as it has no infector, or equivalently, it has no upstream edge. It follows that K coincides with the degree only of the root, while for all other nodes have a degree of $K + 1$ (infectees plus infector). At the onset of the epidemic, only the root is infected while other individuals are susceptible. We consider a susceptible-infected-recovered (SIR) model such that a recovered individual remains immune and does not get re-infected. Contact on one edge (between a susceptible and infected) will lead to infection. On a given edge, contacts happen at exponentially distributed waiting times at rate β . An infected individual recovers either unobserved at rate α , or observed and diagnosed at rate σ . Diagnosed individuals are immediately isolated or treated and classified as recovered. With probability $p_{obs} = \sigma / (\alpha + \sigma)$, an infected individual eventually is observed.

An observed/diagnosed infected individual not only becomes isolated but also is an index case that triggers contact tracing. That is, every adjacent edge has an independent probability p to be traced and consequently isolated if infected. In accordance with the data analysis we aim at, we focus here on one-step tracing, that is, traced individuals do not trigger further tracing events. We do, however, take into account forward and backward tracing as described in Okolie and Müller [16]. We do note this fact, as quite often, theoretical work solely focuses on forward tracing.

All in all, an infected individual can lose his or her infectivity in three possible ways; an unobserved recovery α , observed recovery σ , and a successful tracing event. It turns out that the central ingredient for the analysis is the probability for an infected individual to still be infectious at age a . Please note that “age” in the present paper always refers to the age of (or time since) infection, and never to chronological age. We define

$$\kappa(a) = \mathbb{P}(\text{a randomly chosen infected node of generation is infectious at age of infection } a), \quad (1)$$

which satisfies the following differential equation:

$$\frac{d}{da} \kappa(a) = -\kappa(a) (\alpha + \sigma + \text{tracing}(a)),$$

where $\kappa(0) = 1$. Without contact tracing, $\hat{\kappa}(a) := e^{-(\alpha+\sigma)a}$. With contact tracing, this probability $\kappa(a)$ is decreased and thus

$$\kappa(a) = \hat{\kappa}(a)[1 - p \times \text{tracing in the interval } [0, a]]. \quad (2)$$

In [16], expressions for $\kappa(a)$ are derived. As we do not use these results in the current paper, we only indicate the overall structure and refer the interested reader to that paper for the details.




Figure 1: The figure presents a schematic representation of the infection graph, illustrating the dynamics and interconnections in the process of disease transmission, and both forward and backward tracing. The root node is individual R . Focal individual A is infected by individual C and in turn, infects individuals B and D . In a forward tracing scenario starting from individual A , individuals B and D can be traced. In contrast, in a backward tracing scenario from A , individual C can be traced. The tracing probability is denoted in green.

3. Distribution of ages since infection

In order to work out the distribution of the number of detectees per index case, the age since infection of the index case at its diagnosis is required. Thereto, we consider the case without contact tracing, $p = 0$ (such that index cases are diagnosed but do not trigger contact tracing). This assumption simplifies the arguments and yields an approximation for the age distribution in case of $p > 0$, which is still appropriate if $p \ll 1$ or if $p_{obs} \ll 1$. It turns out, that the resulting approximation is sufficient for practical purposes.

Since the recovery rate α and the screening rate σ are constant, we have a Markovian model, and the age distribution of index cases coincides with the age distribution in the population.

Let $i(t, a)$ denote the age since infection-structured population size of infected individuals. As derived in Okolie and Müller [16], the age-structured model reads

$$(\partial_t + \partial_a)i(t, a) = -(\alpha + \sigma)i(t, a) \quad (3)$$

$$i(t, 0) = \int_0^\infty \theta(a) i(t, a) da. \quad (4)$$

where

$$\theta(a) = \mathbb{E}[K] \beta e^{-\beta a}$$

is the age-dependent rate at which an infected average individual produces (downstream) infecteds. At this point, it is crucial that the contact graph is a tree, and the downstream degree distribution of each node/individual is an i.i.d. realization of the degree distribution K . If we count the number of nodes with a certain distance to the root, this number of nodes is exponentially increasing (for $\mathbb{E}[K] > 1$), unless the tree is finite in a given realization. We can exclude the

case of finite trees since these realizations imply that we have a minor outbreak, and we are not interested in these minor outbreaks. As usual, the age-structured model will tend to an exponential growing solution with a stable age structure,

$$i(t, a) = I_0 e^{\lambda t} i(a)$$

with $i(a) = e^{-(\lambda+\alpha+\sigma)a}$ the probability to be infectious at age a . The exponent λ is the unique real root of

$$1 = \int_0^\infty \theta(a) e^{-(\lambda+\alpha+\sigma)a} da = \mathbb{E}[K] \beta \int_0^\infty e^{-(\lambda+\alpha+\sigma+\beta)a} da \quad \Rightarrow \quad \lambda = \beta(\mathbb{E}[K] - 1) - \alpha - \sigma.$$

The asymptotic age distribution of index cases (which are detected at rate σ) tends to

$$\varphi(a) = \lim_{t \rightarrow \infty} \frac{\sigma i(t, a)}{\int_0^\infty \sigma i(t, b) db} = \beta (\mathbb{E}[K] - 1) e^{-\beta(\mathbb{E}[K]-1)a}. \quad (5)$$

As a side remark, we also obtain the reproduction number from these considerations by

$$R_0 = \int_0^\infty \theta(a) e^{-(\alpha+\sigma)a} da = \frac{\mathbb{E}[K] \beta}{\alpha + \sigma + \beta}.$$

As shown in Fig. 2, the agreement of the age distribution with simulated data is still excellent, though we have in the simulation $p = 0.6$ and $p_{obs} = 1/2$. Furthermore, we have a higher density of lower age groups in the population. For any randomly chosen individual given by age since infection, it is not surprising to have a younger dominating age class. Due to the exponentially fast-growing population, this asymptotic age distribution is expected.




Figure 2: The figure displays the theoretical age-of-distribution $\varphi(a)$, represented by the solid line, in comparison with the distribution obtained from simulations, represented by bars. Parameters: $\beta = 1.5$, $\alpha = 0.5$, $\sigma = 0.5$, $p = 0.6$, and $\mathbb{E}[K] = 4$.

4. Distribution of detected cases

In this section, we derive the distribution of the number of detected cases per index case. That is, the fraction of contacts of an index case who is detected via a tracing event triggered by the index case. We start with forward tracing. We then

combine this result with backward tracing to yield full tracing.

Note that a central ingredient is the age distribution derived in the last section. We did not include contact tracing there. That is, all results in the present section are only a valid approximation if contact tracing does not crucially affect this age distribution. This is the case if either p or p_{obs} is small. All results are only valid under this assumption. However, the simulation study discussed below shows that this assumption is not too restrictive for practical purposes.

Proposition 4.1. *Let $\hat{p}(a)$ be the probability that an infected downstream node is successfully traced given that the focal individual becomes an index case at age since infection a .*

$$\hat{p}(a) = p \frac{\beta}{\alpha + \sigma - \beta} (e^{-\beta a} - e^{-(\alpha + \sigma)a}) \quad (6)$$

Proof. Note that an individual is only able to become an index case at the transition from I to R, that is, our focal individual is infectious in $[0, a)$. We consider one downstream individual. Let $s_1(c)$ represent the probability for this downstream individual to still be susceptible at age $c \in [0, a]$, $s_2(c)$ the probability to be infected, and $s_3(c)$ the probability for the downstream node to be removed. We have the following ODEs

$$\begin{aligned} \dot{s}_1 &= -\beta s_1 & s_1(0) &= 1 \\ \dot{s}_2 &= \beta s_1 - (\alpha + \sigma) s_2 & s_2(0) &= 0 \\ \dot{s}_3 &= (\alpha + \sigma) s_2 & s_3(0) &= 0. \end{aligned}$$




Figure 3: The figure depicts the transition states for a single edge in the infection process. These states represent the probabilities $s_1(c)$, $s_2(c)$, and $s_3(c)$ of an individual downstream from the index case to remain susceptible, infected, and be removed at age $c \in [0, a]$.

The probability for the downstream node to be infectious at the time the infector has age since infection a given by $s_2(a)$,

$$s_2 = \frac{\beta}{\alpha + \sigma - \beta} (e^{-\beta a} - e^{-(\alpha + \sigma)a}),$$

and $\hat{p}(a) = p s_2(a)$ establishes the result. \square

With this proposition and the age distribution $\varphi(a)$, we are able to find the distribution of the number of detected downstream individuals. For simplicity, we first consider a fixed degree distribution $K = k$ for some $k \in \mathbb{N}$, and then address the case of a random tree, where K is a random variable.

4.1. Fixed degree

In the present section, assume that the downstream degree of a node in the tree always is a deterministic number $K = k \in \mathbb{N}$. Particularly, $\mathbb{E}[K] = K$.

Proposition 4.2. *Let T be the random variable for the total number of successfully traced individuals by one index case and forward tracing only. The asymptotic probability distribution of T under forward tracing reads*

$$P(T = i) = \int_0^\infty \binom{k}{i} \hat{p}(a)^i (1 - \hat{p}(a))^{k-i} \beta(k-1) e^{-\beta(k-1)a} da. \quad (7)$$

Proof. As we assume that a tracing event acts independently on different edges, the random variable T , conditioned on the age of the index case at diagnosis a , follows a Binomial distribution with parameters k and an age-dependent tracing probability on one edge $\hat{p}(a)$, $T \sim \text{Binom}(k, \hat{p}(a))$. Thus, the probability of i downstream detectees given age a and k total downstream nodes is given as

$$P(T = i | a) = \binom{k}{i} \hat{p}(a)^i (1 - \hat{p}(a))^{k-i}. \quad (8)$$

Last we remove the condition a by integrating over all possible age of index cases $\varphi(a)$ (eqn. 5),

$$P(T = i) = \int_0^\infty P(T = i | a) \varphi(a) da = \int_0^\infty \binom{k}{i} \hat{p}(a)^i (1 - \hat{p}(a))^{k-i} \beta(k-1) e^{-\beta(k-1)a} da \quad (9)$$

□

Now we turn to full tracing. Thereto, we introduce the random characteristic I_a , which assumes the value 1 if the upstream individual of the index case (its infector) still is infected when the index case is identified (where the index case has age since infection a), and 0 else. Note that I_a is a Bernoulli random variable with

$$P(I_a = 1) = e^{-(\alpha+\sigma)a} + \mathcal{O}(p).$$

As before, in what follows we use the approximation

$$P(I_a = 1) = e^{-(\alpha+\sigma)a}$$

and drop the $\mathcal{O}(p)$ correction terms.

Proposition 4.3. *Let T_{tot} be the random variable for the total number of successfully traced individuals by one index case, under full tracing (forward and backward tracing). With $\varphi(a)$ and I_a as introduced above, the probability distribution of T_{tot} reads*

$$P(T_{tot} = i) = \int_0^\infty \left[pP(I_a = 1)P(T = i-1 | a) + (1 - pP(I_a = 1))P(T = i | a) \right] \varphi(a) da. \quad (10)$$

Proof. If the infector already is recovered ($I_a = 0$), then (conditioning in the age/time of infection of the index case a)

$$P(T_{tot} = i | a, I_a = 0) = P(T = i | a).$$

If the infector is still infectious, also the infector might be traced, such that one of the k detectees might be the upstream

individual (probability p), or not (probability $1 - p$),

$$P(T_{tot} = i|a, I_a = 1) = pP(T = i - 1|a) + (1 - p)P(T = i|a).$$

Taking these two cases together, we have

$$\begin{aligned} P(T_{tot} = i|a) &= P(I_a = 0)P(T = i|a) + P(I_a = 1) \left(pP(T = i - 1|a) + (1 - p)P(T = i|a) \right) \\ &= pP(I_a = 1)P(T = i - 1|a) + (1 - pP(I_a = 1))P(T = i|a). \end{aligned}$$

Integrating by $\varphi(a) da$ removed the condition on a and yields the result. □

4.2. Random degree

So far, the model is formulated for a fixed degree. In most applications, we do not always know individual contacts k due to randomness in contact structure. We now assume an arbitrary degree distribution such that the distribution of the contacts of a random node is defined by some probability distribution $P(K = k)$. The model for fixed case in eqn. 8 is adapted, we only have to take the expectation by summing over all possible numbers of contacts k multiplied by the corresponding probabilities. Thus,

$$P(T_{tot} = i) = \sum_{k=i}^{\infty} \int_0^{\infty} \left[pP(I_a = 1)P(T = i - 1|a, K = k) + (1 - pP(I_a = 1))P(T = i|a, K = k) \right] \varphi(a) da P(K = k). \quad (11)$$

As illustrated in Fig. 4, we have the distribution of the number of detected secondary cases via contact tracing. For the parameter chosen in our study (see Fig. 5), we find a good agreement between our theory results and simulation. We again emphasize that the age structure entering our estimation only is an approximation, as contact tracing is neglected. nevertheless, the results are more than acceptable, even for $p = 0.6$ and $p_{obs} = 0.5$.




Figure 4: The figure showcases theoretical predictions against simulated results. The left panel represents forward tracing for fixed degrees, while the right panel depicts full tracing for a Poisson degree distribution. In both panels, bullet points denote theoretical probabilities $P(T = i)$ or $P(T_{tot} = i)$, whereas the lines represent results from 100000 simulations. Additional parameters used during the simulation: $\beta = 1.5$, $\alpha = 0.5$, $\sigma = 0.5$, $p = 0.6$, and $\mathbb{E}[K] = 4$.

We note that our estimator is independent of time t . The only ‘‘time’’ that appears is the age since infection a . The probability $P(T_{tot} = i)$ consists of integrals as $\int_0^\infty g(a) \varphi(a) da$. Here, we are allowed to choose the time unit, resp. to define $a = \zeta b$ for $\zeta > 0$,

$$\int_0^\infty g(a) \varphi(a) da = \int_0^\infty g(b\zeta) \zeta \varphi(b\zeta) db.$$

If we choose one time unit to be $1/(\alpha + \sigma)$, which is, $\zeta = 1/(\alpha + \sigma)$, then in $P(T_{tot} = i)$ the epidemiological parameters β , α and σ always can be replaced by an expression of R_0 and $E[K]$. That is, the epidemiological parameters enter the estimator solely via R_0 . We only check that fact for one of the terms, as the argument is similar for the other terms.

$$\begin{aligned} & \int_0^\infty P(I_a = 1) P(T = i | a, K = k) \varphi(a) da \\ &= \binom{k}{i} \int_0^\infty e^{-(\alpha+\sigma)a} \hat{p}(a)^i (1 - \hat{p}(a))^{k-i} \beta (\mathbb{E}[K] - 1) e^{-\beta(\mathbb{E}[K]-1)a} da \\ &= \binom{k}{i} \int_0^\infty e^{-b} \hat{p}(b/(\alpha + \sigma))^i (1 - \hat{p}(a/(\alpha + \sigma)))^{k-i} \frac{\beta (\mathbb{E}[K] - 1)}{\alpha + \sigma} e^{-\frac{\beta(\mathbb{E}[K]-1)}{\alpha+\sigma} b} db. \end{aligned}$$

With $R_0 = \frac{\beta \mathbb{E}[K]}{\alpha + \sigma + \beta}$ we have (note that always $\mathbb{E}[K] > R_0$, as we have – in average – only $\mathbb{E}[K]$ downstream individuals who can get infected)

$$\frac{\beta}{\alpha + \sigma} = \frac{R_0}{\mathbb{E}[K] - R_0}$$

and hence

$$\frac{\beta (\mathbb{E}[K] - 1)}{\alpha + \sigma} = \frac{(\mathbb{E}[K] - 1) R_0}{\mathbb{E}[K] - R_0}, \quad \hat{p}(b/(\alpha + \sigma)) = \frac{\frac{\beta}{\alpha + \sigma}}{\frac{\beta}{\alpha + \sigma} - 1} \left(e^{-b} - e^{-\frac{\beta}{\alpha + \sigma} b} \right) = \frac{R_0}{2R_0 - \mathbb{E}[K]} \left(e^{-b} - e^{-\frac{R_0}{\mathbb{E}[K] - R_0} b} \right).$$

That is, all expressions only depend on R_0 , K , and p .

Corollary 4.4. $P(T_{tot} = i)$ and $P(T = i)$ only depend on the epidemiological parameters via R_0 and depends furthermore on the degree distribution given by K and on the tracing probability p .

We can use the formulas from above, where we pragmatically set $\alpha + \sigma$ to 1, and – given R_0 and $\mathbb{E}[K]$ – define $\beta = R_0/(\mathbb{E}[K] - R_0)$.

5. Estimation by maximum likelihood method

In this section, we will set up the likelihood estimator for our model. We assume that we have n observations of index cases, and where $i_\ell \in \mathbb{N}_0, \ell = 1, \dots, n$ denote the total number of detectees per index case (one step tracing only).

5.1. Likelihood estimator

We are able to set an estimator via $P(T_{tot} = i) = P(T_{tot} = i | \vec{\mu})$ for these data points where $\vec{\mu}$ are the parameters of the model we wish to estimate (tracing probability and parameters of the random variable K , e.g. $\mu = (p, \mathbb{E}[K])$ in case of a Poisson distribution for K). The likelihood for the data reads

$$\mathcal{L}(\vec{\mu} | i_\ell, \ell = 1, \dots, n) = \prod_{\ell=1}^n \int_0^\infty \left[p P(I_a = 1) P(T = i_\ell - 1 | a) + (1 - p P(I_a = 1)) P(T = i_\ell | a) \right] \varphi(a) da$$

and the log-likelihood is given by

$$\mathcal{L}\mathcal{L}(\vec{\mu} | i_\ell, \ell = 1, \dots, n) = \sum_{\ell=1}^n \ln \left(\int_0^\infty \left[pP(I_a = 1)P(T = i_\ell - 1 | a) + (1 - p)P(I_a = 1)P(T = i_\ell | a) \right] \varphi(a) da \right). \quad (12)$$

5.2. Simulated data

An agent-based stochastic model is used to simulate the data. To maximise the likelihood in eqn. 12, we plug into the likelihood function all independent observed data points and determine the arg max. As shown in Fig. 5, in our estimation we find back the true values we used in the simulation, namely the tracing probability $p = 0.6$ and the expected number of edges $\mathbb{E}[K] = 4$. The blue circle region contains a global maximum for the true parameter value of the estimated Poisson degree distribution. Other parameters β, α, σ are known and fixed. For the simulated data, we find a satisfying result based on our theory assumption.




Figure 5: Parameters estimated from the simulation data derived from 100000 iterations. The tracing probability, denoted as p , was determined to be 0.6 and the expected number of edges, represented by $\mathbb{E}[K]$, was 4. The blue region represents the 95% confidence region for the estimated parameters. Additional parameters used during the simulation: $\beta = 1.5, \alpha = 0.5, \sigma = 0.5$.

5.3. Covid-19 data

In the previous section, we used the maximum likelihood estimator with contact tracing on simulated data to estimate the tracing probability and expected number of edges. In the simulated dataset we analysed, we have information about the total number of contacts of index cases and also the number of infected contacts identified via contact tracing. In this section, we would like to see how this methods work with empirical data. We wish to estimate the tracing probability p and expected number of edges $\mathbb{E}[K]$ from a data set collected during the Covid-19 pandemic. We obtained a published data set on contact tracing conducted in 2020 from a remarkable extensive and nice study in Karnataka, India where 956 confirmed index cases were reported between the 9th of March 2020 and the 20th of May, 2020 [20]. A comprehensive description of the dataset including the data source, data handling and ethics approval can be found in Gupta et al. [20]. A summary of the number of detectees per index cases from the reported dataset is shown in Table 1.

We only look at one-step tracing at the moment because only detected secondary cases of primary (index) cases are accounted for in the likelihood estimator. Generally, in most epidemic modelling studies, secondary cases are defined as

close/direct contacts (e.g. household, family, etc) of index cases [21]. In the dataset, direct and indirect detected contacts of index cases were reported. For our study, we would focus on only the direct contacts, thus we are able to analyse this scenario with the estimator for only one-step forward tracing. For the reported reproduction number, we chose $R_0 = 3$ in accordance with Gupta et al. [20]. An extensive meta-analysis of covid-19 data from China encompassing 29 studies revealed an approximate R_0 value of 3.32 (95% CI: 2.81-3.82) [22]. In order to investigate the influence of R_0 on our estimates, we additionally carried out a sensitivity analysis.

Number of detectees	0	1	2	3	4	5	6	7	8	10	11	12	13	15	16	19	22	28	29
Frequency	766	87	34	19	16	12	3	4	3	2	1	1	1	1	1	2	1	1	1

Table 1: Total number and frequencies of the total number of detected cases.

We use standard random graph models (see Table 2) to get inspiration on which degree distribution might be appropriate for describing the data [19].

We performed (a) a maximum likelihood estimation and (b) for model comparison, we used the AIC and a chi-square (goodness-of-fit) test. The summary of the point estimates is shown in Table 3.

Optimization: We inspected the gradient of the result and the eigenvalues of the Hessian to ensure that we have (at least) an approximate local maximum. The estimator converged satisfyingly for all models except for the Poissonian degree model: In that case, the parameter of the distribution (the expectation) always increased. Seemingly, the optimum is either very large or even infinite.

Confidence intervals: The approximate 95% confidence intervals are based on the quadratic approximation of the log-likelihood at its maximum, respectively the approximation of the Fisher information matrix by the the inverse of the negative Hessian. We determined the confidence intervals for geometric, scale-free, and negative binomial distributions, as the other models turn out to be inappropriate for the data. In the case of the negative binomial distribution, however, the log-likelihood attains its maximum close to the theoretical lower boundary of $\mathbb{E}[K]$ which is R_0 . As this is numerically a delicate situation, we approximated the confidence interval for $\mathbb{E}[K]$ in that case by the maximum value of $\mathbb{E}[K]$ such that the log-likelihood, given the shape parameter, is larger than its maximum minus 2 (also see the blue curve in Fig. 6).

Chi square (goodness-of-fit) test: We bin the index cases with 5-7 detectees, and all index cases with more than 7 detectees to ensure that at least 10 observations are in one class.

We find that random mixing graphs and the Erdős-Renyi graph induce degree distributions with a rather lightweight tail. Therefore, these distributions do not fit the data appropriately. In the Poisson distribution, which is the degree distribution of an Erdős-Renyi graph, the optimization routine even does not obtain a local maximum: It seems as in this case, the optimum is only assumed for an expectation $\mathbb{E}[K]$ that is unreasonably large or even tends to infinity.

¹See Appendix A and Appendix B for further detail on full graph/random mixing and the optimization process respectively.

Table 2: Examples of standard random graph models.

Network Model	Degree Distribution for Large Population Size N
Full Graph/Random Mixing ¹	$K = N - 1 \rightarrow \infty, \beta \rightarrow 0, R_0$ constant
Erdős-Rényi	$K \sim$ Poisson
Configuration Model	Choice: $K \sim$ Geometric
Scale-free Network	Power-Law, $P(K = k) = ck^{-\gamma}, \gamma > 1$
Standard Degree Distribution	Negative Binomial

Distribution	p (95% CI)	$\mathbb{E}[K]$ (95% CI)	add. information	AIC	p-val (χ^2)
random mixing	0.98	-		2443	$< 10^{-20}$
Poisson	0.98	52	(not converged)	2464	$< 10^{-20}$
Geometric	0.87 (0.76, 0.97)	16.6 (11.3, 22.0)		1858	$< 10^{-20}$
Power-law	0.74 (0.61, 0.88)	11.4 (8.3, 14.5)	$\gamma = 1.48$	1687	0.003
negative Binomial	0.72 (0.59, 0.86)	4.5 (3, 9.5)	$r = 0.16$ (0.12, 0.20)	1675	0.14

Table 3: Parameter estimates and model comparison for five probability distributions fitted to a dataset, including the probability (p), the expected value ($\mathbb{E}[K]$), additional information (if available), Akaike Information Criterion (AIC), and p-value from a chi-squared test. Note that the estimator in the case of the Poissonian distribution did not converge, and we simply fixed a large expected value. The interval for r in the negative Binomial distribution is 95% CI.

The geometric distribution has a tail that is heavy enough to at least allow for a reasonable fit, but AIC as well as the chi-square (goodness of fit) test indicate that this model is rejected. The power-law (scale-free graph) is the first model that is at least weakly in line with the data: The tail of a power-law distribution may become heavy, and in this, there is a possibility to handle superspreading events appropriately. The AIC is worse but not too far from the winning model, and the p -value for the chi-square (goodness-of-fit) test is at least only in the range of 10^{-3} , and not less than 10^{-20} , as in the previous models.

The best model clearly is the negative binomial distribution, which is known to be an appropriate model for the number of contacts relevant for the transmission of respiratory infections [23]. The expected number of infectious contacts is small enough to be in the range of R_0 , while – as expected – the over-dispersion is distinct. This model has the best AIC among all models, and also the goodness-of-fit test does not reject this model.

It is interesting that the point estimate for p decreases if we choose models that have more mass in the tail. The reason is that the probability for k detectees scales with p^k , such that k small(er) needs to be balanced with more probability mass in the tail. A similar reason leads to smaller values for $\mathbb{E}[K]$ if the model distribution has more probability mass in the tail. However, this point estimate is rather similar for the power-law and the negative binomial and also does not heavily depend on the choice of R_0 (see “sensitivity analysis”). In that, the range of p -estimate seems to be trustworthy. Moreover, the information in the data is sufficiently strong to point to a specific degree distribution (negative binomial), which was not clear from the beginning. As the data are rather simple, the information content could also have been too little to allow for distinct conclusions. That the negative binomial distribution, which is well known to be appropriate in this situation, is selected, is another sign that the estimates are trustworthy.

We first draw a contour plot (fig. 6) indicating the point estimates (p and $\mathbb{E}[K]$) and also draw the cumulative empirical distribution vs. the cumulative theoretical distribution (fig. 7), that is, on the x -axis, we plot the number of infectees, and on the y -axis the percentage of index cases which is the number of infected contact persons or less.

Also, these graphics clearly indicates that the power-law and the negative binomial distribution yield the best fit, where the negative binomial distribution is superior to the power-law.

6. Sensitivity analysis

We carried out a sensitivity analysis on how the estimated parameters depend on R_0 . The sensitivity analysis revealed that the estimated parameters in the studied models are not highly sensitive to the choice of R_0 , at least for the power-law and negative distribution (the only degree distributions which meet the data satisfyingly). This observation is particularly noteworthy, as it underscores the robustness of these models in providing reliable estimates of epidemiological parameters, even when the initial assumptions about R_0 may vary. This characteristic is crucial in the context of real-world epidemiological studies, where the precise value of R_0 is often uncertain due to factors such as heterogeneous populations, changing contact patterns, and varying degrees of intervention measures.




Figure 6: A contour plot showing the point estimates p and $E(K)$ of the likelihood estimator in different network models. Distributions from left to right: Geometric, scale-free, and negative binomial. Parameters; $R_0 = 3$, for the negative binomial, $r = 0.17$.




Figure 7: Cumulative empirical distribution (blue bullets) compared with the cumulative theoretical distribution (black circles). The top row, from left to right, represents distributions for a homogeneous graph, a Poisson graph, and a geometric distribution, respectively. The bottom row, from left to right, represents distributions for a power-law distribution and a negative binomial distribution, respectively. Our proposed theory is indicated by the black circles.




Figure 8: Sensitivity analysis illustrating changes in the point estimates (p and $E(K)$), and log-likelihood for different choices of R_0 . Left: Power law (with the log-likelihood as inlay), middle and right: negative binomial distribution.

7. Discussion

In this paper, we present a graph-based method for estimating parameters in infectious disease models, offering valuable insights into the efficiency of contact tracing programs and some information about local contact structures, and their implications for the spread of infectious diseases. By comparing various degree distributions and assessing their suitability for modelling disease spread, our analysis contributes to the ongoing development of improved parameter estimation techniques in graph-based models. Our findings complement and expand upon the work of Müller and Hösel [12] who estimated tracing probability in homogeneous populations by a maximum likelihood estimator and applied it to tuberculosis and chlamydia contact tracing data. In contrast to focusing on homogeneous random mixing populations, our work explores contact graphs in form of trees, which enables us to capture the unique branching structure of infection transmission chains. This approach provides a more realistic representation of contact patterns at a microscopic level, allowing for a better understanding of the dynamics of infectious diseases and the effectiveness of contact tracing strategies.

Our comprehensive analysis of the COVID-19 contact tracing data from Karnataka, India, reveals that both scale-free network models and negative binomial distribution models offer a good fit for the data. The negative binomial distribution emerges as the most fitting model for the data, aligning with previous epidemiological research that has identified this distribution as a suitable assumption for the number of contacts relevant for the transmission of respiratory infections [23]. Furthermore, the observed over-dispersion in the number of secondary cases caused by individual index cases is accurately captured by the negative binomial distribution. This distribution is suitable for data where the variance exceeds the mean, reflective of scenarios where a small proportion of index cases are responsible for a disproportionate number of secondary infections. These findings resonate with the work of Gupta et al. [20] who had previously reported a clear over-dispersion in the data. Specifically, Gupta et al. [20] found that amongst 956 confirmed traced cases, just 8.7% of index cases, who had 14.4% of contacts, were responsible for 80% of all secondary cases. The power-law distribution also offers a reasonable fit, highlighting the potential relevance of scale-free networks in modelling infectious disease dynamics. In line with the principles of scale-free networks, our model highlights the role of a relatively small number of "super-spreader" individuals, who have a significantly larger number of contacts and thus a higher likelihood of transmitting the infection to a larger pool of people. This also validates the findings of Gupta et al. [20] who suggested that super-spreaders may have played a more dominant role in the covid-19 transmission in Karnataka, India.

Both the scale-free and negative binomial models allow for a thick or heavy tail, which is created by super-spreader events and it is known in the case of airborne infection such as covid-19 that super-spreader events and the over-dispersion of secondary cases have a significant impact on the effectiveness of contact tracing and surveillance schemes [24, 20]. Our findings also confirm that the effectiveness of tracing schemes is influenced by the degree of super-spreading events.

The estimated tracing probabilities from our chosen models reveal that as the intensity or frequency of super-spreading events increases, we observe a corresponding impact on the performance of tracing schemes. Our modelling study and findings highlight the importance of selecting appropriate models for estimating tracing probabilities and local contact structures in real-world scenarios. These estimates demonstrate the effectiveness of our graph-based method in capturing key epidemic parameters within heterogeneous and age-structured contact networks. The estimated degree ranges for the negative binomial and power-law distributions fall within plausible ranges found in the literature, supporting the validity of our approach in comparison to other studies that have examined contact tracing data [23, 25, 26]. While our model assumes fixed contact and recovery rates, it is worth noting that the high heterogeneity in individual-level contact networks could lead to underestimation or overestimation of these parameters [27, 28].

Furthermore, sensitivity analysis on the estimated parameters w.r.t. R_0 for the power-law and negative binomial distribution shows limited sensitivity to the choice of R_0 . This provides valuable insights into estimating key epidemiological and intervention parameters with greater confidence, enabling more effective public health strategies and interventions. All in all, our research contributes to the ongoing development of improved parameter estimation techniques in graph-based models for infectious disease dynamics. By utilizing a graph-based approach and building upon the methods of previous studies, we have demonstrated the value of incorporating contact graph structures, such as trees, for a more accurate representation of contact patterns and infectious disease dynamics. The results of our analysis highlight the need to consider heterogeneity in individual-level contact networks when designing and evaluating contact tracing strategies.

Future research in this area could explore the incorporation of additional data sources and model refinements. A key area of refinement could be incorporating temporal dynamics or considering other types of contact graphs that better represent real-world contact patterns. The use of tree-like graphs in the current study, while mathematically convenient, is a simplification of reality. Contact patterns, particularly within clusters such as households or other social groups, often exhibit significant clustering and interconnectedness that a tree structure may fail to accurately capture. This limitation of tree-based representations could lead to discrepancies in modelling infectious disease dynamics, and therefore, warrants attention. Integrating well-suited graph structures with modelling techniques such as agent-based models or compartmental models, could provide a more comprehensive understanding of infectious disease dynamics and inform the design of more effective public health interventions.

Acknowledgements

Portions of the work presented in this paper have been previously published in a PhD thesis by Okolie [29], which was submitted to the Technical University of Munich. The thesis is accessible online at <https://mediatum.ub.tum.de/1661774>. This current paper extends and improves the findings discussed in the thesis, particularly those related to the likelihood estimator models and their applications to simulated data and empirical COVID-19 contact tracing data, which were presented in chapters 3 – 5 of the thesis. The research was supported by a grant from the German Academic Exchange Service DAAD (AO), and by the Deutsche Forschungsgemeinschaft (DFG) through TUM International Graduate School of Science and Engineering (IGSSE), GSC 81, within the project GENOMIE_QADOP (JM). M.E.K. acknowledges support from Horizon 2020 research and innovation funding programme (Grant no. 101003480 (CORESMA)).

Declaration of Competing Interest

None

Supplementary material

Additional supplementary material, which includes the implementation of the algorithms and codes, is available.

References

- [1] Muhammad Altaf Khan et al. Parameter estimation and fractional derivatives of dengue transmission model. *AIMS Mathematics*, 5(3):2758–2779, 2020.
- [2] Roderick J Little, Ralph D’Agostino, Michael L Cohen, Kay Dickersin, Scott S Emerson, John T Farrar, Constantine Frangakis, Joseph W Hogan, Geert Molenberghs, Susan A Murphy, et al. The prevention and treatment of missing data in clinical trials. *New England Journal of Medicine*, 367(14):1355–1360, 2012.
- [3] Begoña Cantó, Carmen Coll, and Elena Sánchez. Structural identifiability of a model of dialysis. *Mathematical and computer modelling*, 50(5-6):733–737, 2009.
- [4] Begoña Cantó, Carmen Coll, and Elena Sánchez. Identifiability for a class of discretized linear partial differential algebraic equations. *Mathematical Problems in Engineering*, 2011, 2011.
- [5] Gheorghe Craciun and Casian Pantea. Identifiability of chemical reaction networks. *Journal of Mathematical Chemistry*, 44(1):244–259, 2008.
- [6] Michael GB Blum and Viet Chi Tran. Hiv with contact tracing: a case study in approximate bayesian computation. *Biostatistics*, 11(4):644–660, 2010.
- [7] PD O’Neill, GO Roberts, and Bradford Univ.(United Kingdom). Dept. of Mathematics;. *Bayesian inference for partially observed stochastic epidemics*. University of Bradford. School of Mathematical Sciences, 1997.
- [8] Thomas Götz, Nicole Altmeier, Wolfgang Bock, Robert Rockenfeller, Karunia Putra Wijaya, et al. Modeling dengue data from semarang, indonesia. *Ecological complexity*, 30:57–62, 2017.
- [9] FB Augusto and MA Khan. Optimal control strategies for dengue transmission in pakistan. *Mathematical biosciences*, 305:102–121, 2018.
- [10] N. Stollenwerk, M. Aguiar, S. Ballesteros, J. Boto, B. Kooi, and L. Mateus. Dynamic noise, chaos and parameter estimation in population biology. *Interface Focus*, 2(2):156–169, feb 2012. doi: 10.1098/rsfs.2011.0103.
- [11] Solym M Manou-Abi, Yousri Slaoui, and Julien Balicchi. Estimation of some epidemiological parameters with the covid-19 data of mayotte. *Frontiers in Applied Mathematics and Statistics* (), page 67, 2022.
- [12] Johannes Müller and Volker Hösel. Estimating the tracing probability from contact history at the onset of an epidemic. *Mathematical Population Studies*, 14(4):211–236, 2007.
- [13] Johannes Müller, Mirjam Kretzschmar, and Klaus Dietz. Contact tracing in stochastic and deterministic epidemic models. *Mathematical biosciences*, 164(1):39–64, 2000.
- [14] Louise Dyson, Michael Marks, Oliver M Crook, Oliver Sokana, Anthony W Solomon, Alex Bishop, David CW Mabey, and T Déirdre Hollingsworth. Targeted treatment of yaws with household contact tracing: how much do we miss? *American journal of epidemiology*, 187(4):837–844, 2018.
- [15] Takuma Tanaka, Takayuki Yamaguchi, and Yohei Sakamoto. Estimation of the percentages of undiagnosed patients of the novel coronavirus (SARS-CoV-2) infection in hokkaido, japan by using birth-death process with recursive full tracing. *PLOS ONE*, 15:e0241170, 2020. doi: 10.1371/journal.pone.0241170.
- [16] Augustine Okolie and Johannes Müller. Exact and approximate formulas for contact tracing on random trees. *Mathematical biosciences*, 321:108320, 2020.

- [17] Darren M Green and Istvan Z Kiss. Large-scale properties of clustered networks: Implications for disease dynamics. *Journal of biological dynamics*, 4(5):431–445, 2010.
- [18] Matthew J Keeling. The effects of local spatial structure on epidemiological invasions. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 266(1421):859–867, 1999.
- [19] István Z Kiss, Joel C Miller, Péter L Simon, et al. Mathematics of epidemics on networks. *Cham: Springer*, 598:31, 2017.
- [20] Mohak Gupta, Giridara G Parameswaran, Manraj S Sra, Rishika Mohanta, Devarsh Patel, Amulya Gupta, Bhavik Bansal, Vardhmaan Jain, Archisman Mazumder, Mehak Arora, et al. Contact tracing of covid-19 in karnataka, india: Superspreading and determinants of infectiousness and symptomatic infection. *Plos one*, 17(7):e0270789, 2022.
- [21] Beth P Bell, Marcia Goldoft, Patricia M Griffin, Margaret A Davis, Diane C Gordon, Phillip I Tarr, Charles A Bartleson, Jay H Lewis, Timothy J Barrett, Joy G Wells, et al. A multistate outbreak of escherichia coli o157:h7—associated bloody diarrhea and hemolytic uremic syndrome from hamburgers: the washington experience. *Jama*, 272(17):1349–1353, 1994.
- [22] Isabella Locatelli, Bastien Trächsel, and Valentin Rousson. Estimating the basic reproduction number for covid-19 in western europe. *Plos one*, 16(3):e0248731, 2021.
- [23] Joël Mossong, Niel Hens, Mark Jit, Philippe Beutels, Kari Auranen, Rafael Mikolajczyk, Marco Massari, Stefania Salmaso, Gianpaolo Scalia Tomba, Jacco Wallinga, et al. Social contacts and mixing patterns relevant to the spread of infectious diseases. *PLoS medicine*, 5(3):e74, 2008.
- [24] James O Lloyd-Smith, Sebastian J Schreiber, P Ekkehard Kopp, and Wayne M Getz. Superspreading and the effect of individual variation on disease emergence. *Nature*, 438(7066):355–359, 2005.
- [25] Loes Soetens, Don Klinkenberg, Corien Swaan, Susan Hahné, and Jacco Wallinga. Real-time estimation of epidemiologic parameters from contact tracing data during an emerging infectious disease outbreak. *Epidemiology*, 29(2):230–236, 2018.
- [26] Martyn Fyles, Elizabeth Fearon, Christopher Overton, University of Manchester COVID-19 Modelling Group, Tom Wingfield, Graham F Medley, Ian Hall, Lorenzo Pellis, and Thomas House. Using a household-structured branching process to analyse contact tracing in the sars-cov-2 pandemic. *Philosophical Transactions of the Royal Society B*, 376(1829):20200267, 2021.
- [27] Luca Ferretti, Chris Wymant, Michelle Kendall, Lele Zhao, Anel Nurtay, Lucie Abeler-Dörner, Michael Parker, David Bonsall, and Christophe Fraser. Quantifying sars-cov-2 transmission suggests epidemic control with digital contact tracing. *Science*, 368(6491):eabb6936, 2020.
- [28] Joel Hellewell, Sam Abbott, Amy Gimma, Nikos I Bosse, Christopher I Jarvis, Timothy W Russell, James D Munday, Adam J Kucharski, W John Edmunds, Fiona Sun, et al. Feasibility of controlling covid-19 outbreaks by isolation of cases and contacts. *The Lancet Global Health*, 8(4):e488–e496, 2020.
- [29] Augustine Okebunor Okolie. *Contact Tracing on Stochastic Graphs*. PhD thesis, Universitätsbibliothek der TU München, 2022.

Appendix A. Full graph/random mixing

In the case of a full graph, we start with a fixed degree $K = N - 1$, where N is the population size, and take the limit $N \rightarrow \infty$ and $\beta \rightarrow 0$ such that $R_0 = \frac{(N-1)\beta}{\mu+\sigma}$ is constant. Then,

$$(N-1)\hat{p}(a) = R_0 \frac{\alpha + \sigma}{\alpha + \sigma - \beta} \left(e^{-\beta a} - e^{-(\alpha+\sigma)a} \right) \rightarrow R_0 (1 - e^{-(\alpha+\sigma)a}).$$

Therefore the Binomial distribution approximates a Poissonian distribution, and eqn 7 becomes in the limit $N \rightarrow \infty$

$$P(T = i) = \int_0^\infty \text{dpois}(i, R_0 (1 - e^{-(\alpha+\sigma)a})) R_0 (\alpha + \sigma) e^{-R_0 (\alpha+\sigma)a} da$$

where $\text{dpois}(i, \mu) = \mu^i e^{-\mu} / i!$ is the probability function for the Poisson distribution.

Appendix B. Optimization Process

Given a function $f(x)$, the aim of optimization is to find an x that either maximizes or minimizes $f(x)$. This process involves the computation of gradients and the Hessian matrix.

The gradient of a function is a vector that points in the direction of the greatest increase of that function. It is calculated as the vector of the first derivatives of the function with respect to each variable. The gradient of a function $f(x)$, where $x = [x_1, x_2, \dots, x_n]$ is:

$$\nabla f(x) = \left[\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right] \quad (\text{B.1})$$

To ensure that the solution found is a local maximum and not a local minimum or a saddle point, the Hessian matrix is used. The Hessian matrix is the square matrix of second-order partial derivatives of the function. Each element in the Hessian matrix is the second derivative of the function with respect to different variables. The Hessian matrix for the function $f(x)$ is:

$$H(f(x)) = \begin{bmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \frac{\partial^2 f}{\partial x_2^2} & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \frac{\partial^2 f}{\partial x_n \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_n^2} \end{bmatrix} \quad (\text{B.2})$$

If the Hessian is positive definite (all eigenvalues are positive) at a point, then the function attains a local minimum at that point. If the Hessian is negative definite (all eigenvalues are negative), then the function attains a local maximum.

In the context of maximizing a likelihood function, we often convert the problem into a minimization problem by taking the negative of the likelihood function. This is due to the fact that many optimization algorithms are developed for minimization problems. The negative log-likelihood function becomes:

$$-\mathcal{L} \mathcal{L}(\vec{\mu} \mid i_\ell, \ell = 1, \dots, n) \quad (\text{B.3})$$

The goal now is to minimize this negative log-likelihood function, and the optimization problem becomes:

$$\vec{\mu}^* = \underset{\vec{\mu}}{\operatorname{argmin}} \{-\mathcal{L} \mathcal{L}(\vec{\mu} \mid i_\ell, \ell = 1, \dots, n)\} \quad (\text{B.4})$$

The same principles of gradients and Hessians apply to this minimization problem. The gradient of the negative log-likelihood function should point in the direction of greatest decrease of the function. The Hessian, on the other hand, should be negative definite at the point of minimum.

In the case of the log-likelihood function, the optimization problem can be solved using iterative methods such as Newton's method or Quasi-Newton methods, which make use of both the gradient and the Hessian of the function to find the minimum.

Appendix C. A Note on the Simulation Algorithm

The algorithm, implemented in Python and structured into four class objects, simulates the spread of an infectious disease with contact tracing.

The *Parameter* class stores all parameters of the simulation model, including the infection rate, recovery rate, tracing probability, and whether the degree is fixed or random, among other parameters. It also controls whether the tracing is forward, backward, or both.

The *Event* class stores all events. These include the event type (infection on an edge or recovery), event times, the individuals involved, and whether the event was deactivated (for instance, when an individual is traced during a recovery event). Event times are determined by an exponentially distributed waiting clock that triggers the events.

The *Population* class represents the entire population. It contains a list of individuals and manages the spread of infection and recovery events among them. The Population class also includes methods for adding individuals and events, updating individuals' statuses, and overseeing the infection and contact tracing processes.

The *Individual* class defines each individual or agent in the population. An individual has a unique ID, a list of downstream nodes, the upstream node, and disease states (S-I-R), among other individual attributes.

In the simulation, an infection spreads starting from the root node only (initially infected at time $t=0$) through the network. Upon recovery, an individual may trigger a contact tracing event, and infected nodes are removed with a probability of p . If forward tracing is enabled, the infected downstream nodes are identified and move to recovery. If backward tracing is enabled, the infector recovers. This data, representing the number of detectees per index case, is then collected and used for analysis.