# A test for counting sequences of integer-valued autoregressive models

Yuichi Goto\*

Faculty of Mathematics, Kyushu University, e-mail: yuichi.goto@math.kyushu-u.ac.jp and

#### Kou Fujimori\*

Department of Applied Economics, Faculty of Economics and Law, Shinshu University, e-mail: kfujimori@shinshu-u.ac.jp

Abstract: The integer autoregressive (INAR) model is one of the most commonly used models in nonnegative integer-valued time series analysis and is a counterpart to the traditional autoregressive model for continuousvalued time series. To guarantee the integer-valued nature, the binomial thinning operator or more generally the generalized Steutel and van Harn operator is used to define the INAR model. However, the distributions of the counting sequences used in the operators have been determined by the preference of analyst without statistical verification so far. In this paper, we propose a test based on the mean and variance relationships for distributions of counting sequences and a disturbance process to check if the operator is reasonable. We show that our proposed test has asymptotically correct size and is consistent. Numerical simulation is carried out to evaluate the finite sample performance of our test. As a real data application, we apply our test to the monthly number of anorexia cases in animals submitted to animal health laboratories in New Zealand and we conclude that binomial thinning operator is not appropriate.

**MSC2020 subject classifications:** Primary 60G10, 62M10; secondary 62F12.

Keywords and phrases: INAR model, non-negative integer-valued time series, thinning operator.

# Contents

1	Introduction
2	Preliminary
3	Main results
	3.1 Asymptotic properties of the proposed test
	3.2 Example of the estimators
4	Numerical study
5	Empirical application 12
А	General model and estimation
В	Proof of Theorem A.1
	*This research was supported by JSPS Grant-in-Aid for Early-Career Scientists

<sup>\*</sup>This research was supported by JSPS Grant-in-Aid for Early-Career Scientists JP23K16851 (Y.G.) and JP21K13271 (K.F.).

2

Acknowledgements																22
References																22

## 1. Introduction

Non-negative integer-valued time series are ubiquitous, and the analysis of this type of data has received much attention over the past two decades. Applications of this research include the analysis of the number of infected people (Schmidt and Pereira, 2011, Xu et al., 2023), crimes (Chen and Lee, 2017, Böhning and van der Heijden, 2019), and so on. A review of recent developments in this area can be found in Davis et al. (2021).

Integer autoregressive (INAR) models and integer-valued generalized autoregressive conditional heteroskedastic (INGARCH) models are popular models for non-negative integer-valued time series. INGARCH was introduced by Ferland, Latour and Oraichi (2006) and the conditional expectation of the process has autoregressive structures. On the other hand, INAR model was proposed by McKenzie (1985) and Al-Osh and Alzaid (1987), and is a natural extension of the classical autoregressive model for continuous-valued time series to count time series. To ensure that the process takes non-negative integer values, the binomial thinning operator is often used in the model. The binomial thinning operator is defined as the sum of independent Bernoulli random variables called counting sequences. Latour (1997) extended the binomial thinning operator to the generalized Steutel and van Harn operator in multivariate settings, which allows the use of non-binomial counting sequences in the operator. Zhu and Joe (2003) suggested an extended thinning operator by using the distribution including binomial distribution as a special case. Weiß (2008) defined the new operator based on the convolution of the Bernoulli and the geometric distributions. Ristić, Bakouch and Nastić (2009) and Yang et al. (2019) studied the operator based on geometric and generalized Poisson distributions, respectively. Aly and Bouzar (2019) advocated the thinning operator based on a distribution with a linear fractional probability generating function.

Many operators have been proposed so far. However, operators have been completely determined by the analyst without statistical guarantees. In this paper, we propose a test for distributions of counting sequences and a distribution of the disturbance process. To construct the test, we make use of the important feature of the non-negative integer-valued distributions that the variance of the distribution takes the form of the function of its mean. Note that Goto and Fujimori (2023) considered a test for conditional variances based on this feature. Their setting is essentially for INGARCH models and does not include INAR models since nuisance parameters in the conditional variance do not be allowed. To encompass INAR models, in Appendix we consider the integervalued time series { $Z_t$ } satisfying

$$\mathbf{E}\left(Z_t \mid \mathcal{F}_{t-1}\right) = m_t(\boldsymbol{\mu}_0) \text{ and } \operatorname{Var}\left(Z_t \mid \mathcal{F}_{t-1}\right) = v_t(\boldsymbol{\theta}_0), \tag{1.1}$$

where  $\mathcal{F}_{t-1}$  is the  $\sigma$ -field generated by  $\{Z_s, s \leq t-1\}$ , for a known function mon  $[0, \infty)^{\infty} \times \mathbb{R}^d$  to  $(\delta, +\infty)$  for some  $\delta > 0$ ,

$$m_t(\boldsymbol{\mu}_0) := m(Z_{t-1}, Z_{t-2}, \ldots; \boldsymbol{\mu}_0),$$

for a known function v on  $[0,\infty)^{\infty} \times \mathbb{R}^{d'}$  to  $(\delta, +\infty)$ ,

$$v_t(\boldsymbol{\theta}_0) := v(Z_{t-1}, Z_{t-2}, \ldots; \boldsymbol{\theta}_0),$$

and  $(\boldsymbol{\mu}_0^{\top}, \boldsymbol{\theta}_0^{\top})^{\top} \in \mathbb{R}^{d+d'}$  are unknown parameters, and establish the strong consistency and the asymptotic normality of the M-estimator of  $\boldsymbol{\mu}_0$  and  $\boldsymbol{\theta}_0$ . Since INAR models defined later in (2.1) have the conditional expectation and conditional variance of  $\{Z_t\}$  given  $\mathcal{F}_{t-1}$  of the form

$$m_t(\mu_0) = \sum_{i=1}^p \mu_i Z_{t-i} + \mu_\epsilon \text{ and } v_t(\theta_0) = \sum_{i=1}^p \sigma_i^2 Z_{t-i} + \sigma_\epsilon^2,$$
 (1.2)

where  $\boldsymbol{\mu}_0 := (\mu_1, \ldots, \mu_p, \mu_{\epsilon})^{\top}$  and  $\boldsymbol{\theta}_0 := (\sigma_1^2, \ldots, \sigma_p^2, \sigma_{\epsilon}^2)^{\top}$ , the above setting includes INAR models. In this article, we consider the conditional least squares estimator with a simple form of the asymptotic variance as a special case of the M-estimator. Based on the asymptotic normality of estimators for  $\boldsymbol{\mu}_0$  and  $\boldsymbol{\theta}_0$ , we show that our proposed test has asymptotically size  $\phi$  and is consistent.

As a related work, several goodness of fit tests for integer-valued time series have been developed. For instance, tests for the intensity function of IN-GARCH models were considered by Neumann (2011), Fokianos and Neumann (2013), and Leucht and Neumann (2013). For the distributions of INGARCH and INAR, tests relied on probability generating functions were investigated by Meintanis and Karlis (2014), Hudecová, Hušková and Meintanis (2015), Schweer (2016) and Hudecová, Hušková and Meintanis (2021). Tests based on the Pearson statistic and the Stein-Chen identity were studied by Weiß (2018)) and Aleksandrov, Weiß and Jentsch (2022), respectively. A test for overdispersion and zero inflation was proposed by Weiß, Homburg and Puig (2019). However, tests related to operators for INAR models have not been considered so far.

The remainder of this paper is organized as follows. In Section 2, we define generalized INAR models and formulate the test for distributions of counting sequences and a distribution of the disturbance process. A test statistic is also defined. In Section 3.1, we show the asymptotic null distribution of the proposed test statistic and the consistency of our test. In Section 3.2, we show the strong consistency and asymptotic normality of the conditional least squares estimators. In Section 4, we illustrate the finite sample performance of our test. In Section 5, we apply our test to the monthly number of skin lesions and anorexia cases in animals submitted to animal health laboratories in New Zealand. In Appendix, we derive sufficient conditions for the strong consistency and asymptotic normality of the M-estimator.

#### 2. Preliminary

In this section, we define a generalized integer autoregressive process, a testing problem, and a test statistic. Let  $\{Z_t\}$  follow a generalized integer autoregressive process of order p (GINAR(p)) defined as

$$Z_t := \sum_{i=1}^p \mu_i \circ Z_{t-i} + \epsilon_t, \qquad (2.1)$$

where  $\circ$  is so called the generalized Steutel and van Harn operator (see Latour (1997)) given by  $\mu_i \circ Z_{t-i} := \sum_{j=1}^{Z_{t-i}} W_{i,t,j}$ ,  $\{W_{i,t,j}\}$  is a counting sequence, which is an i.i.d. (with respect to t and j) non-negative integer-valued random variable with mean  $\mu_i$  and variance  $\sigma_i^2$ ,  $\{W_{i,t,j}\}$  is independent of  $\{W_{i',t',j'}\}$  for  $i \neq i'$ , and  $\{\epsilon_t\}$  is an i.i.d. non-negative integer-valued random variable with mean  $\mu_{\epsilon}$  and variance  $\sigma_{\epsilon}^2$ , which is independent of  $\{W_{i,t,j}\}$ .

To ensure the strictly stationarity of  $\{Z_t\}$ , we suppose that  $1 - \mu_1 z - \mu_2 z^2 - \cdots - \mu_p z^p \neq 0$  for all  $z \in \mathbb{C}$  such that  $|z| \leq 1$ . It is also known that  $\{Z_t\}$  is ergodic. See Latour (1997).

In this paper, we are interested in the following hypothesis testing problem:

$$\tilde{H}_0: (W_{1,t,j}, \dots, W_{p,t,j}, \epsilon_t)$$
 follows a target joint distribution  
v.s.  $\tilde{K}_0: \tilde{H}_0$  does not hold. (2.2)

Before constructing the test statistic, we present some examples of a counting sequence  $\{W_{i,t,j}\}$ .

**Example 2.1.** The most popular choice of a counting sequence  $\{W_{i,t,j}\}$  might be the Bernoulli random variable. In this case, the generalized Steutel and van Harn operator reduces to the binomial thinning.

The second example of  $\{W_{i,t,j}\}$  is the extended thinning proposed by Zhu and Joe (2003), which is defined as the generalized Steutel and van Harn operator with  $\{W_{i,t,j}\}$  following a distribution with probability generating function

$$\phi_{\rm ZJ}(k) := \frac{(1 - \mu_{\rm ZJ}) + (\mu_{\rm ZJ} - \gamma_{\rm ZJ})k}{1 - \mu_{\rm ZJ}\gamma_{\rm ZJ} - (1 - \mu_{\rm ZJ})\gamma_{\rm ZJ}k} \quad \text{for } \mu_{\rm ZJ} \in [0, 1] \text{ and } \gamma_{\rm ZJ} \in [0, 1].$$

See also Remark A.9 of Weiß (2008). The distribution has mean  $\mu_{\rm ZJ}$  and variance  $\mu_{\rm ZJ}(1-\mu_{\rm ZJ})(1+\gamma_{\rm ZJ})/(1-\gamma_{\rm ZJ})$ . According to Zhu and Joe (2003, Remark in p.239 and 240), the random variable can be generated by  $B_{\rm ZJ}G_{\rm ZJ}$ , where  $B_{\rm ZJ}$  and  $G_{\rm ZJ}$  follow the Bernoulli distribution with the success probability  $(1-\gamma_{\rm ZJ})\mu_{\rm ZJ}/(1-\gamma_{\rm ZJ}\mu_{\rm ZJ})$  and the shifted geometric distribution with a parameter  $1-(1-\mu_{\rm ZJ})\gamma_{\rm ZJ}/[\{(1-\mu_{\rm ZJ})\gamma_{\rm ZJ}\}+(1-\gamma_{\rm ZJ})]$ , respectively, and  $B_{\rm ZJ}$  is independent of  $G_{\rm ZJ}$ .

The third example of  $\{W_{i,t,j}\}$  is the  $\text{BerG}(\pi_{\text{BW}}, \xi_{\text{BW}})$  distribution, where  $\pi_{\text{BW}}$  and  $\xi_{\text{BW}}$  are parameters satisfying  $0 < \pi_{\text{BW}} < 1$  and  $\xi_{\text{BW}} > 0$ , investigated by Bourguignon and Weiß (2017). The probability generating function, the mean, and the variance of the distribution are given by  $\phi_{\text{BW}}(s) := \{1 - 1\}$ 

 $\pi_{\rm BW}(1-s)\}/\{1+\xi_{\rm BW}(1-s)\}, \mu_{\rm BW} := \pi_{\rm BW} + \xi_{\rm BW}, \text{ and } \mu_{\rm BW}(1-\mu_{\rm BW}+2\xi_{\rm BW}),$ respectively. The random variable  $B_{\rm BW}+G_{\rm BW}$ , where  $B_{\rm BW}$  and  $G_{\rm BW}$  follow the Bernoulli distribution with the success probability  $\pi_{\rm BW}$  and the geometric distribution with a parameter  $1/(1+\xi_{\rm BW})$ , respectively, follows the BerG( $\pi_{\rm BW}, \xi_{\rm BW}$ ) distribution. Aly and Bouzar (2019) considered a different parametrization from Bourguignon and Weiß (2017) and elucidated that the family of distributions encompasses at least nine distributions as a special case including the second example above.

As mentioned in the introduction, the variances of many non-negative distributions can be written in term of their mean of the distributions. For example, the mean and variance of the binomial distribution with the number of trials n = 1 and the success probability  $p \in (0, 1)$  take the form of  $\mu_{\text{bin}} := p$ ,  $\kappa_{\text{bin}}(\mu_{\text{bin}}) := \mu_{\text{bin}}(1 - \mu_{\text{bin}})$ , respectively. For the Poisson distribution with parameter  $\lambda$ , the mean and variance are given by  $\mu_{\text{Pois}} := \lambda$  and  $\kappa_{\text{Pois}}(\mu_{\text{Pois}}) := \mu_{\text{Pois}}$ . Similarly, for the negative binomial distribution with success probability p and given parameter r, the mean and variance can be expressed as  $\mu_{\text{NB}} := r(1 - p)/p$  and  $\kappa_{NB}(\mu_{\text{NB}}) := (\mu_{\text{NB}} + r)\mu_{\text{NB}}/r$ .

Based on this fact, we rephrase the hypothesis defined in (2.2) as

$$H_0: \text{ the variance of } (W_{1,t,j}, \dots, W_{p,t,j}, \epsilon_t) \text{ takes the form of } \boldsymbol{\kappa} (\boldsymbol{\mu}_0)$$
  
v.s.  $K_0: H_0 \text{ does not hold},$  (2.3)

where  $\boldsymbol{\kappa}(\boldsymbol{\mu}_0) := (\kappa_1(\mu_1), \ldots, \kappa_p(\mu_p), \kappa_{\epsilon}(\mu_{\epsilon}))^{\top}$  and  $\kappa_1, \ldots, \kappa_p$ , and  $\kappa_{\epsilon}$  are the functions of means corresponding to variances of the target marginal distributions for  $W_{1,t,j}, \ldots, W_{p,t,j}$ , and  $\epsilon_t$ , respectively. The hypothesis  $H_0$  is a necessary condition for  $\tilde{H}_0$ .

**Remark 2.1.** In this paper, we specifically focused on the property of nonnegative discrete random variables, where the variance is a function of its mean, and reframed the problem from the distributions of counting sequences to their second-order moments, leveraging the characteristic of non-negative discrete random variables. Importantly, our proposed test does not have any power for counting processes that deviate from the target distribution but share the same first and second-order moments with the target distribution. To our best knowledge, non-negative discrete-valued random variables with separately parametrized mean and variance have not been introduced. Such variables would be highly desirable, though.

To define a test statistic, we consider estimators  $\hat{\boldsymbol{\mu}}_n$  and  $\hat{\boldsymbol{\theta}}_n$  of  $\boldsymbol{\mu}_0$  and  $\boldsymbol{\theta}_0$  endowed with the strong consistency and the asymptotic normality with asymptotic variance  $\boldsymbol{V} := (\boldsymbol{v}_{ij})_{i,j=1,2}$ , that is,

$$\begin{pmatrix} \hat{\boldsymbol{\mu}}_n \\ \hat{\boldsymbol{\theta}}_n \end{pmatrix} \to \begin{pmatrix} \boldsymbol{\mu}_0 \\ \boldsymbol{\theta}_0 \end{pmatrix} \quad \text{almost surely as } n \to \infty \tag{2.4}$$

and

$$\sqrt{n} \begin{pmatrix} \hat{\boldsymbol{\mu}}_n - \boldsymbol{\mu}_0 \\ \hat{\boldsymbol{\theta}}_n - \boldsymbol{\theta}_0 \end{pmatrix} \Rightarrow N(\mathbf{0}, \boldsymbol{V}) \quad \text{as } n \to \infty.$$
(2.5)

The concrete examples of the estimators including the conditional least squares estimators and the M-estimators will be presented in Section 3.2. Motivated by the mean and variance relationships, we consider the asymptotic distribution of  $\kappa(\hat{\mu}_n) - \hat{\theta}_n$ . By Theorem 3.1 and Proposition 6.4.3 of Brockwell and Davis (2009), we know that  $\kappa(\hat{\mu}_n) - \hat{\theta}_n$  converges in distribution to the (p + 1)dimensional centered normal distribution with variance

$$W := K v_{11} K - K v_{12} - v_{21} K + v_{22}, \qquad (2.6)$$

where  $\mathbf{K}$  is a (p+1)-diagonal matrix whose *i*-th diagonal entry is  $(\partial/\partial \mu_i)\kappa_i(\mu_i)$ for  $i \in \{1, \ldots, p\}$  and (p+1)-th diagonal entry is  $(\partial/\partial \mu_{\epsilon})\kappa_{\epsilon}(\mu_{\epsilon})$ , provided that  $\boldsymbol{\kappa}(\cdot)$  is continuously differentiable in a neighborhood of  $\boldsymbol{\mu}_0$ . The quantities  $\boldsymbol{\kappa}(\boldsymbol{\mu}_0)$  and  $\boldsymbol{\theta}_0$  correspond to the variance of  $(W_{1,t,j},\ldots, W_{p,t,j}, \epsilon_t)$  for the null hypothesis we wish to test and for the underlying distributions, respectively.

Thus, under the assumption that V and K are non-singular matrices we propose the following test statistic

$$T_{n} := n \left( \boldsymbol{\kappa} \left( \hat{\boldsymbol{\mu}}_{n} \right) - \hat{\boldsymbol{\theta}}_{n} \right)^{\top} \widehat{\boldsymbol{W}}^{-1} \left( \boldsymbol{\kappa} \left( \hat{\boldsymbol{\mu}}_{n} \right) - \hat{\boldsymbol{\theta}}_{n} \right),$$

where  $\widehat{W}^{-1}$  is the inverse matrix of  $\widehat{W}$ ,  $\widehat{W}$  is the consistent estimator of W. In Section 3.1, we derive asymptotic properties of the test based on  $T_n$ .

#### 3. Main results

#### 3.1. Asymptotic properties of the proposed test

In this subsection, we derive the asymptotic distribution of  $T_n$  and the consistency of the test based on  $T_n$ .

Based on the above discussion, the asymptotic null distribution of the test statistic is given in the following theorem.

**Theorem 3.1.** Suppose that estimators  $\hat{\boldsymbol{\mu}}_n$  and  $\hat{\boldsymbol{\theta}}_n$  of  $\boldsymbol{\mu}_0$  and  $\boldsymbol{\theta}_0$  satisfy the strong consistency and the asymptotic normality with asymptotic variance  $\boldsymbol{V} := (\boldsymbol{v}_{ij})_{i,j=1,2}, \boldsymbol{\kappa}(\cdot)$  is continuously differentiable in a neighborhood of  $\boldsymbol{\mu}_0$ , matrices  $\boldsymbol{V}$  and  $\boldsymbol{K}$  are non-singular, and  $\widehat{\boldsymbol{W}}$  is the consistent estimator of  $\boldsymbol{W}$ . Then, under the null hypothesis  $H_0$ ,  $T_n$  converges in distribution to the chi-square distribution with p+1 degrees of freedom as  $n \to \infty$ .

*Proof.* The discussion in Section 2 and the assumption of this theorem yield the conclusion.  $\Box$ 

**Remark 3.1.** Examples of the estimators  $\hat{\boldsymbol{\mu}}_n$  and  $\hat{\boldsymbol{\theta}}_n$  of  $\boldsymbol{\mu}_0$  and  $\boldsymbol{\theta}_0$  satisfying the strong consistency and the asymptotic normality are given in Section 3.2. For the least squared estimator stated in Section 3.2, the moment conditions  $EW_{i,t,j}^8 < \infty$  and  $E\epsilon_t^8 < \infty$  suffice to construct the consistent estimator for  $\boldsymbol{W}$  (see also Remark 3.2).

Therefore, the test which rejects the null hypothesis  $H_0$  whenever  $T_n \geq \chi^2_{p+1}[1-\phi]$  has the asymptotic size  $\phi$ , where  $\phi$  is a significance level and  $\chi^2_{p+1}$  denotes the upper  $\phi$ -percentile of the chi-square distribution with p+1 degrees of freedom.

The following theorem shows the consistency of the test.

**Theorem 3.2.** Suppose that estimators  $\hat{\mu}_n$  and  $\hat{\theta}_n$  of  $\mu_0$  and  $\theta_0$  satisfy the strong consistency and the asymptotic normality with asymptotic variance  $\mathbf{V} := (\mathbf{v}_{ij})_{i,j=1,2}, \kappa(\cdot)$  is continuously differentiable in a neighborhood of  $\mu_0$ , matrices  $\mathbf{V}$  and  $\mathbf{K}$  are non-singular, and  $\widehat{\mathbf{W}}$  is the consistent estimator of  $\mathbf{W}$ . Then, the test based on  $T_n$  is consistent, that is, (under the alternative hypothesis  $K_0$ ,) the power of the test tends to one as  $n \to \infty$ .

*Proof.* Denote  $\kappa_0$  and  $\kappa_1$  as the vector-valued functions of means corresponding to variances of the target marginal distributions for  $W_{1,t,j}, \ldots, W_{p,t,j}$ , and  $\epsilon_t$  under the null hypothesis and under the alternative, respectively. The consistency of  $\widehat{W}$  and  $\widehat{\mu}_n$  and  $\widehat{\theta}_n$  gives that, under the alternative hypothesis  $K_0$ ,

$$P\left(\frac{T_n}{n} \ge \frac{\chi_{p+1}^2[1-\phi]}{n}\right)$$
  
=P $\left(\left(\kappa_0\left(\mu_0\right) - \kappa_1\left(\mu_0\right) + \kappa_1\left(\mu_0\right) - \theta_0\right)^\top W^{-1}$   
 $\times \left(\kappa_0\left(\mu_0\right) - \kappa_1\left(\mu_0\right) + \kappa_1\left(\mu_0\right) - \theta_0\right) \ge \frac{\chi_{p+1}^2[1-\phi]}{n}\right)$   
=P $\left(\left(\kappa_0\left(\mu_0\right) - \kappa_1\left(\mu_0\right)\right)^\top W^{-1}\left(\kappa_0\left(\mu_0\right) - \kappa_1\left(\mu_0\right)\right) \ge 0\right) + o_p(1).$ 

Here we used the relationship  $\kappa_1(\mu_0) = \theta_0$ . The definition of the nonnegativedefinite matrix yields

$$P\left(\left(\boldsymbol{\kappa}_{0}\left(\boldsymbol{\mu}_{0}\right)-\boldsymbol{\kappa}_{1}\left(\boldsymbol{\mu}_{0}\right)\right)^{\top}\boldsymbol{W}^{-1}\left(\boldsymbol{\kappa}_{0}\left(\boldsymbol{\mu}_{0}\right)-\boldsymbol{\kappa}_{1}\left(\boldsymbol{\mu}_{0}\right)\right)\geq0\right)=1.$$

#### 3.2. Example of the estimators

In this subsection, we derive the example of the estimators  $\hat{\mu}_n$  and  $\hat{\theta}_n$  of  $\mu_0$  and  $\theta_0$  endowed with the strong consistency and the asymptotic normality.

The conditional least squares estimators for  $\mu$  and  $\theta$  are defined as

$$\hat{\boldsymbol{\mu}}_n := \operatorname*{arg\,min}_{\boldsymbol{\mu} \in \boldsymbol{H}} \frac{1}{n} \sum_{t=1+p}^n (Z_t - \boldsymbol{\mu}^\top \boldsymbol{Y}_{t-1})^2$$

imsart-generic ver. 2020/08/06 file: Goto\_Fujimori\_arXiv.tex date: February 5, 2024

and

$$\hat{\boldsymbol{\theta}}_n := \operatorname*{arg\,min}_{\boldsymbol{\theta}\in\boldsymbol{\Theta}} \frac{1}{n} \sum_{t=1+p}^n \left( (Z_t - \boldsymbol{\mu}^\top \boldsymbol{Y}_{t-1})^2 - \boldsymbol{\theta}^\top \boldsymbol{Y}_{t-1} \right)^2,$$

where  $\mathbf{Y}_{t-1} := (Y_{t-1}, \ldots, Y_{t-p}, 1)^{\top}$ ,  $\mathbf{H}$  is a compact subspace of  $\boldsymbol{\mu} \in \mathbb{R}^{p+1}$  such that  $1 - \mu_1 z - \mu_2 z^2 - \cdots - \mu_p z^p \neq 0$  for all  $z \in \mathbb{C}$  such that  $|z| \leq 1$  and  $\mu_i > 0$  for any  $i \in \{1, \ldots, p\}$  and  $\boldsymbol{\Theta}$  is a compact subspace of  $\boldsymbol{\theta} \in \mathbb{R}^{p+1}$  such that  $\sigma_i^2 > 0$  for any  $i \in \{1, \ldots, p\}$  (see Klimko and Nelson, 1978). The least squared estimator can be written explicitly as

$$\hat{\boldsymbol{\mu}}_n := \left(\frac{1}{n} \sum_{t=1+p}^n \boldsymbol{Y}_{t-1} \boldsymbol{Y}_{t-1}^\top\right)^{-1} \frac{1}{n} \sum_{t=1+p}^n Z_t \boldsymbol{Y}_{t-1}$$

and

$$\hat{\theta}_n := \left(\frac{1}{n} \sum_{t=1+p}^n Y_{t-1} Y_{t-1}^\top\right)^{-1} \frac{1}{n} \sum_{t=1+p}^n \left(Z_t - \hat{\mu}_n^\top Y_{t-1}\right)^2 Y_{t-1}.$$

The strong consistency and asymptotic normality of  $\hat{\mu}_n$  and  $\hat{\theta}_n$  are shown in the following theorem.

**Lemma 3.1.** Suppose that  $EW_{i,t,j}^8 < \infty$  and  $E\epsilon_t^8 < \infty$ . Then,  $\hat{\mu}_n$  and  $\hat{\theta}_n$  tend almost surely to  $\mu_0$  and  $\theta_0$  as  $n \to \infty$ , respectively. Moreover, it holds that

$$\sqrt{n} \begin{pmatrix} \hat{\boldsymbol{\mu}}_n - \boldsymbol{\mu}_0 \\ \hat{\boldsymbol{\theta}}_n - \boldsymbol{\theta}_0 \end{pmatrix} \Rightarrow N \left( \boldsymbol{0}, \boldsymbol{J^{-1}} \boldsymbol{I} \boldsymbol{J^{-1}} \right) \quad as \ n \to \infty,$$

where  $\mathcal{O}_{p+1}$  is a  $(p+1) \times (p+1)$  zero matrix,

$$oldsymbol{J} := egin{pmatrix} oldsymbol{J}_m & oldsymbol{\mathcal{O}}_{p+1} \ oldsymbol{\mathcal{O}}_{p+1} & oldsymbol{J}_v \end{pmatrix}, \ and \ oldsymbol{I} := egin{pmatrix} oldsymbol{I}_m & oldsymbol{I}_m v \ oldsymbol{I}_m v & oldsymbol{I}_v \end{pmatrix}$$

with  $\boldsymbol{J}_m := \mathrm{E}\left(\boldsymbol{Y}_{t-1}\boldsymbol{Y}_{t-1}^{\top}\right), \, \boldsymbol{J}_v := \mathrm{E}\left(\boldsymbol{Y}_{t-1}\boldsymbol{Y}_{t-1}^{\top}\right), \, \boldsymbol{I}_m := \mathrm{E}\left(\left(\boldsymbol{\theta}_0^{\top}\boldsymbol{Y}_{t-1}\right)\boldsymbol{Y}_{t-1}\boldsymbol{Y}_{t-1}^{\top}\right), \\ \boldsymbol{I}_{mv} := \mathrm{E}\left(\left(\boldsymbol{Z}_t - \boldsymbol{\mu}_0^{\top}\boldsymbol{Y}_{t-1}\right)^3\boldsymbol{Y}_{t-1}\boldsymbol{Y}_{t-1}^{\top}\right), \, and \\ \boldsymbol{I}_v := \mathrm{E}\left(\left(\boldsymbol{Z}_t - \boldsymbol{\mu}_0^{\top}\boldsymbol{Y}_{t-1}\right)^4 - \left(\boldsymbol{\theta}_0^{\top}\boldsymbol{Y}_{t-1}\right)^2\right)\boldsymbol{Y}_{t-1}\boldsymbol{Y}_{t-1}^{\top}\right).$ 

*Proof.* It is easy to check the conditions (B1)-(B12) and (C1)-(C13) in Appendix. Thus, Theorem A.1 in Appendix yields the conclusion.

**Remark 3.2.** The conditions  $EW_{i,t,j}^8 < \infty$  and  $E\epsilon_t^8 < \infty$  ensure that  $EZ_t^8 < \infty$  (See Theorem 2.2 of Drost, Van den Akker and Werker 2008). The condition  $EZ_t^8 < \infty$  is used when we show  $I_v$  is finite. To see this, it suffices to show that, for any  $s, k \in \{1, \ldots, p\}$ ,

$$\operatorname{E}\left(\left(\left(Z_{t}-\boldsymbol{\mu}_{0}^{\top}\boldsymbol{Y}_{t-1}\right)^{4}-\left(\boldsymbol{\theta}_{0}^{\top}\boldsymbol{Y}_{t-1}\right)^{2}\right)Z_{t-s}Z_{t-k}\right)<\infty$$

since the proofs of

$$\mathbb{E}\left(\left((Z_t - \boldsymbol{\mu}_0^{\top} \boldsymbol{Y}_{t-1})^4 - \left(\boldsymbol{\theta}_0^{\top} \boldsymbol{Y}_{t-1}\right)^2\right) Z_{t-s}\right) < \infty$$
$$\mathbb{E}\left(\left((Z_t - \boldsymbol{\mu}_0^{\top} \boldsymbol{Y}_{t-1})^4 - \left(\boldsymbol{\theta}_0^{\top} \boldsymbol{Y}_{t-1}\right)^2\right)\right) < \infty$$

and

$$\mathbb{E}\left(\left(\left(Z_t - \boldsymbol{\mu}_0^\top \boldsymbol{Y}_{t-1}\right)^4 - \left(\boldsymbol{\theta}_0^\top \boldsymbol{Y}_{t-1}\right)^2\right)\right) < \infty$$

are similar. By the Cauchy-Schwarz inequality and the definition of the convex function, we obtain

$$\left\{ E\left(\left((Z_{t}-\boldsymbol{\mu}_{0}^{\top}\boldsymbol{Y}_{t-1})^{4}-\left(\boldsymbol{\theta}_{0}^{\top}\boldsymbol{Y}_{t-1}\right)^{2}\right)Z_{t-s}Z_{t-k}\right)\right\}^{2} \\ \leq E\left((Z_{t}-\boldsymbol{\mu}_{0}^{\top}\boldsymbol{Y}_{t-1})^{4}-\left(\boldsymbol{\theta}_{0}^{\top}\boldsymbol{Y}_{t-1}\right)^{2}\right)^{2} E\left(Z_{t-s}^{2}Z_{t-k}^{2}\right) \\ \leq 2E\left((Z_{t}-\boldsymbol{\mu}_{0}^{\top}\boldsymbol{Y}_{t-1})^{8}+\left(\boldsymbol{\theta}_{0}^{\top}\boldsymbol{Y}_{t-1}\right)^{4}\right)\left\{E\left(Z_{t-s}^{4}\right)\right\}^{1/2}\left\{E\left(Z_{t-k}^{4}\right)\right\}^{1/2} \\ \leq E\left(2^{8}Z_{t}^{8}+2^{8}(\boldsymbol{\mu}_{0}^{\top}\boldsymbol{Y}_{t-1})^{8}+2\left(\boldsymbol{\theta}_{0}^{\top}\boldsymbol{Y}_{t-1}\right)^{4}\right)\left\{E\left(Z_{t-s}^{4}\right)\right\}^{1/2}\left\{E\left(Z_{t-k}^{4}\right)\right\}^{1/2},$$

which is finite since the repeated applications of the definition of the convex function yields that  $E\left((\boldsymbol{\mu}_{0}^{\top}\boldsymbol{Y}_{t-1})^{8}\right)$  is finite. The estimator of  $\boldsymbol{J}^{-1}\boldsymbol{I}\boldsymbol{J}^{-1}$  can be constructed by replacing  $\boldsymbol{\mu}_{0}$  and  $\boldsymbol{\theta}_{0}$  with

 $\hat{\mu}_n$  and  $\hat{\theta}_n$ , respectively, using empirical sums instead of expectations. Under the condition  $EZ_t^8 < \infty$ , the consistency of the estimator of  $J^{-1}IJ^{-1}$  can be shown by the uniform law of large numbers (see, e.g., Nishiyama 2022, Theorem 7.3.1 (ii)).

Remark 3.3. In Appendix, we derive the asymptotic normality of the Mestimator for the general model defined in (1.1). Since  $m_t(\boldsymbol{\mu})$  and  $v_t(\boldsymbol{\mu},\boldsymbol{\theta})$  include unobservable variables  $Z_0, Z_{-1}, \ldots$ , define computable quantities

$$\tilde{m}_t(\boldsymbol{\mu}) := m(Z_{t-1}, Z_{t-2}, \dots, Z_1, \boldsymbol{x}_0; \boldsymbol{\mu})$$

and

$$\tilde{v}_t(\boldsymbol{\theta}) := v(Z_{t-1}, Z_{t-2}, \dots, Z_1, \boldsymbol{x}_0; \boldsymbol{\theta})$$

for some constant  $x_0$ . The M-estimator is defined as

$$\check{\boldsymbol{\mu}}_n := \arg\max_{\boldsymbol{\mu}\in\boldsymbol{H}} \frac{1}{n} \sum_{t=1}^n \ell_m(Z_t, \tilde{m}_t(\boldsymbol{\mu}))$$

and

$$\check{\boldsymbol{\theta}}_n := \operatorname*{arg\,max}_{\boldsymbol{\theta}\in\boldsymbol{\Theta}} \frac{1}{n} \sum_{t=1}^n \ell_v(Z_t, \tilde{m}_t(\check{\boldsymbol{\mu}}_n), \tilde{v}_t(\boldsymbol{\theta})),$$

where  $\ell_m$  and  $\ell_v$  are some estimating functions.

The asymptotic variance of the M-estimator is given by  $\tilde{V} := (\tilde{v}_{ij})_{i,j=1,2}$ , where

$$ilde{oldsymbol{v}}_{11} := ilde{oldsymbol{J}}_m^{-1} ilde{oldsymbol{J}}_m^{-1}, \quad ilde{oldsymbol{v}}_{12} := ilde{oldsymbol{J}}_m^{-1} ( ilde{oldsymbol{I}}_m v - ilde{oldsymbol{I}}_m^{-1} ilde{oldsymbol{J}}_v^{-1}, \quad ilde{oldsymbol{v}}_{21} := ilde{oldsymbol{v}}_{12}^{-1},$$

$$ilde{v}_{22} := ilde{J}_v^{-1} ( ilde{I}_v + ilde{J}_{vm} ilde{J}_m^{-1} ilde{I}_m ilde{J}_m^{-1} ilde{J}_{vm}^ op - ilde{I}_{mv}^ op ilde{J}_m^{-1} ilde{J}_{vm}^ op - ilde{J}_{vm} ilde{J}_m^{-1} ilde{I}_{mv}) ilde{J}_v^{-1}$$

with

$$\begin{split} \tilde{\boldsymbol{I}}_{m} &:= \mathbf{E} \left( \frac{\partial}{\partial \boldsymbol{\mu}} \ell_{m}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\mu}_{0})) \frac{\partial}{\partial \boldsymbol{\mu}^{\top}} \ell_{m}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\mu}_{0})) \right), \\ \tilde{\boldsymbol{J}}_{m} &:= \mathbf{E} \left( \frac{\partial^{2}}{\partial \boldsymbol{\mu} \partial \boldsymbol{\mu}^{\top}} \ell_{m}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\mu}_{0})) \right), \\ \tilde{\boldsymbol{I}}_{v} &:= \mathbf{E} \left( \frac{\partial}{\partial \boldsymbol{\theta}} \ell_{v}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\mu}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})) \frac{\partial}{\partial \boldsymbol{\theta}^{\top}} \ell_{v}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\mu}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})) \right), \\ \tilde{\boldsymbol{J}}_{v} &:= \mathbf{E} \left( \frac{\partial^{2}}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^{\top}} \ell_{v}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\mu}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})) \right), \\ \tilde{\boldsymbol{I}}_{mv} &:= \mathbf{E} \left( \frac{\partial}{\partial \boldsymbol{\mu}} \ell_{m}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\mu}_{0})) \frac{\partial}{\partial \boldsymbol{\theta}^{\top}} \ell_{v}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\mu}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})) \right), \\ \text{and } \tilde{\boldsymbol{J}}_{vm} &:= \mathbf{E} \left( \frac{\partial^{2}}{\partial \boldsymbol{\theta} \partial \boldsymbol{\mu}^{\top}} \ell_{v}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\mu}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})) \right). \end{split}$$

The asymptotic variance of  $\check{\boldsymbol{\theta}}_n$  is complicated due to the fact that  $\ell_v$  includes  $\tilde{m}_t(\check{\boldsymbol{\mu}}_n)$ . In the proof, it can be shown that

$$\begin{pmatrix} \sqrt{n}(\check{\boldsymbol{\mu}}_n - \boldsymbol{\mu}_0) \\ \sqrt{n}(\check{\boldsymbol{\theta}}_n - \boldsymbol{\theta}_0) \end{pmatrix} = \begin{pmatrix} \tilde{\boldsymbol{J}}_m^{-1} & \mathcal{O}_{p+1} \\ -\tilde{\boldsymbol{J}}_v^{-1} \tilde{\boldsymbol{J}}_v^{-n} \tilde{\boldsymbol{J}}_m^{-1} & -\tilde{\boldsymbol{J}}_v^{-1} \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{n}} \sum_{t=1}^n \frac{\partial}{\partial \boldsymbol{\mu}} \ell_m(Z_t, m_t(\boldsymbol{\mu}_0)) \\ \frac{1}{\sqrt{n}} \sum_{t=1}^n \frac{\partial}{\partial \boldsymbol{\theta}} \ell_v(Z_t, m_t(\boldsymbol{\mu}_0), v_t(\boldsymbol{\theta}_0)) \end{pmatrix} + o_p(1),$$

where  $\mathcal{O}_{p+1}$  is a  $(p+1) \times (p+1)$  zero matrix, and the martingale central limit theorem yields the asymptotic normality.

On the other hand, the asymptotic variance of the least squares estimators is considerably simple due to the fact that

$$\mathbb{E}\left(\frac{\partial^2}{\partial\boldsymbol{\theta}\partial\boldsymbol{\eta}^{\top}}((Z_t - m_t(\boldsymbol{\mu}_0))^2 - v_t(\boldsymbol{\theta}_0))^2\right) = \boldsymbol{\mathcal{O}}_{p+1},$$

which corresponds to  $\tilde{J}_{vm} := \mathcal{O}_{p+1}$ .

One may consider that the alternative definition of the least squares estimator as

$$\breve{\boldsymbol{\theta}}_n := \operatorname*{arg\,min}_{\boldsymbol{\theta}\in\boldsymbol{\Theta}} \frac{1}{n} \sum_{t=1}^n (Z_t^2 - m_t(\hat{\boldsymbol{\mu}}_n)^2 - v_t(\boldsymbol{\theta}))^2.$$

However in this case, we have

$$\mathbb{E}\left(\frac{\partial^2}{\partial\boldsymbol{\theta}\partial\boldsymbol{\eta}^{\top}}(Z_t^2 - m_t(\boldsymbol{\mu}_0)^2 - v_t(\boldsymbol{\theta}_0))^2\right)$$
  
=  $\mathbb{E}\left(4m_t(\boldsymbol{\mu}_0)\frac{\partial}{\partial\boldsymbol{\theta}}v_t(\boldsymbol{\theta}_0)\frac{\partial}{\partial\boldsymbol{\mu}^{\top}}m_t(\boldsymbol{\mu}_0))\right),$ 

which is not equal to a zero matrix. Therefore, our choise for  $\ell_v$  is essential to obtain a simple form for the asymptotic variance of the estimator.

**Remark 3.4.** It is worth to highlight that we established the asymptotic normality of the estimator for the variance of counting sequences. Therefore, it is reasonable to expect that the eighth order moments of the process to estimate the asymptotic variance of the variance for counting sequences. In that case one still think that the assumption of eighth order moment is strong, we suggest to apply the weighted least square estimator (e.g. Section 3.5 of Aknouche and Francq (2021) and Remark below Theorem 2 of Kachour and Truquet 2011), which has  $\sqrt{n}$ -convergence rate, to relax the moment assumption.

On the other hand, the technique of Carlstein (1986) or Peligrad and Shao (1994) might be applied to relax the moment assumption. Nonetheless, a fundamental trade-off exists in convergence rate. Specifically, their estimator converges at some rate slower than  $\sqrt{n}$ .

**Remark 3.5.** Our approach can be applied to the integer-valued bilinear process defined as

$$Z_t := \sum_{i=1}^p \alpha_i \circ Z_{t-i} + \sum_{j=1}^q \beta_j \circ \epsilon_{t-i} + \sum_{k=1}^m \sum_{\ell=1}^n \gamma_{k,\ell} \circ (X_{t-k}\epsilon_{t-\ell}) + \epsilon_t,$$

the integer-valued ARCH process (see Doukhan, Latour and Oraichi 2006, Drost, Van den Akker and Werker 2008, Latour and Truquet 2009, Doukhan, Fokianos and Li 2012), the signed INAR model (see Kachour and Truquet 2011, Li, Chen and Zhu 2023, and references therein).

#### 4. Numerical study

In this section, we investigate the finite sample performance of our test. We consider INAR(1) model whose counting sequence and disturbance process follow  $BerG(\pi_{BW}, \xi_{BW})$  and Poisson distributions with parameter  $\lambda = 1$ , respectively. See Example 2.1.

We set the null hypothesis as the counting sequence following the Bernoulli distribution, which corresponds to the case  $\xi_{\rm BW} = 0$ , and the disturbance process following the Poisson distribution. For this null hypothesis, the functions  $\kappa_1(\mu_1)$  and  $\kappa_{\epsilon}(\mu_{\epsilon})$  are given by  $\kappa_1(\mu_1) := \mu_1(1 - \mu_1)$  and  $\kappa_{\epsilon}(\mu_{\epsilon}) := \mu_{\epsilon}$ , respectively, where  $\mu_1 := \pi_{\rm BW}$  and  $\lambda := \mu_{\epsilon}$ . Then, the matrix  $\boldsymbol{K}$  in  $\boldsymbol{W}$  is given by  $\boldsymbol{K} := \begin{pmatrix} 1 - 2\mu_1 & 0 \\ 0 & 1 \end{pmatrix}$ .

Let the significance level of the test be  $\phi := 0.05$ , the number of iterations R := 1000, the time series length  $n \in \{500, 1000, 2000\}$ , burn-in period 1000. We use the parameters  $(\pi_{BW}, \xi_{BW})$  on the sets  $\{0.2, 0.3, \ldots, 0.8\} \times \{0\}$  for the null hypothesis and  $\{0.2, 0.3, \ldots, 0.6\} \times \{0.05, 0.1, \ldots, 0.3\}$  for the alternative hypothesis. Note that the parameters  $\pi_{BW}$  and  $\xi_{BW}$  need to satisfy the condition  $\pi_{BW} + \xi_{BW} < 1$  to ensure stationarity of the process. The least squares estimators are used to estimate unknown parameters.

$\pi_{\rm BW}$	n									
	500	1000	2000							
0.2	0.066	0.070	0.059							
0.3	0.070	0.062	0.050							
0.4	0.093	0.063	0.057							
0.5	0.087	0.065	0.056							
0.6	0.086	0.054	0.059							
0.7	0.076	0.070	0.070							
0.8	0.090	0.075	0.075							

TABLE 1 The empirical size at the nominal size 0.05

TABLE 2												
The en	npirical pou	ver at a	the nominal	power	0.05							

$\pi_{\rm BW}$	n	$\xi_{ m BW}$											
		0.05	0.1	0.15	0.2	0.25	0.3						
	500	0.061	0.084	0.180	0.389	0.643	0.903						
0.2	1000	0.049	0.128	0.411	0.758	0.968	0.999						
	2000	0.088	0.327	0.775	0.978	1.000	1.000						
	500	0.047	0.097	0.302	0.603	0.865	0.983						
0.3	1000	0.073	0.215	0.640	0.933	0.998	1.000						
	2000	0.111	0.510	0.948	1.000	1.000	1.000						
	500	0.064	0.165	0.486	0.806	0.971	1.000						
0.4	1000	0.078	0.401	0.840	0.994	1.000	1.000						
	2000	0.178	0.748	0.990	1.000	1.000	1.000						
	500	0.072	0.303	0.694	0.949	0.999	0.993						
0.5	1000	0.129	0.598	0.970	1.000	1.000	0.998						
	2000	0.289	0.912	1.000	1.000	1.000	1.000						
	500	0.086	0.418	0.870	0.981	0.974	0.951						
0.6	1000	0.190	0.785	0.993	0.999	0.983	0.957						
	2000	0.445	0.987	1.000	1.000	0.998	0.973						

We generate INAR(1) process with length n and apply our test. Then, we iterate 1000 times to calculate empirical size or power. Table 1 shows the empirical size of the test. We observe that our test has reasonable size overall although our test has slightly over-rejection for the large values of  $\pi_{BW}$ . This corresponds to the fact that when  $\pi_{BW}$  takes large values,  $\pi_{BW} + \xi_{BW}$  is close to the boundary of the stationary region of INAR(1).

Table 2 presents the empirical power of the test. As n or  $\xi_{BW}$  gets larger, as the power of the test increases. As expected, it is more difficult to reject the null hypothesis when  $\xi_{BW}$  is small, which is the case that the null and alternative hypotheses are close.

#### 5. Empirical application

In this section, we apply our method to the monthly number of skin lesions and anorexia cases in animals submitted to animal health laboratories in New Zealand (84 observations were collected from January 2003 to December 2009). These data can be found in Tables 3 and 4 of Mohammadpour, Bakouch and Shirozhan (2018). Jazi, Jones and Lai (2012) and Mohammadpour, Bakouch and Shirozhan

(2018) fitted these datasets to binomial thinning base INAR(1) model with zero inflated Poisson innovations and innovations induced by the Poisson–Lindley distribution, respectively. Our objective of this section is to confirm if the binomial thinning operator is appropriate for the datasets.

The estimated values of the least squares estimators  $\hat{\boldsymbol{\mu}}_n = (\hat{\mu}_1, \hat{\mu}_{\epsilon})$  and  $\hat{\boldsymbol{\theta}}_n = (\hat{\sigma}_1^2, \hat{\sigma}_{\epsilon}^2)$  are (0.325, 0.964) and (0.841, 1.97) for skin-lesions and (0.680, 0.263) and (2.40, 0.102) for anorexia, respectively. The estimation shows that the counting series and disturbance process have overdispersion property for skin-lesions and overdispersion and underdispersion properties for anorexia, respectively.

We consider the null hypothesis that the datasets follow INAR(1) model with the binomial thinning operator. Our test only for the thinning operator (not for both thinning operator and innovation) gives p-values 0.182 for skinlesions and 0.00277 for anorexia. We conclude there is no evidence to reject the null hypothesis for skin-lesions but that binomial thinning operator is not appropriate for anorexia. Therefore, we suggest that the counting series should be chosen carefully, rather than using the binomial thinning operator that has been used as customary.

#### Appendix A: General model and estimation

In this section, we introduce the general model including the INAR(p) model and show the strong consistency and the asymptotic normality of the M-estimator.

Let  $\{Z_t\}$  be integer-valued time series satisfying

$$\mathbb{E}(Z_t \mid \mathcal{F}_{t-1}) = m_t(\boldsymbol{\eta}_0) \text{ and } \operatorname{Var}(Z_t \mid \mathcal{F}_{t-1}) = v_t(\boldsymbol{\theta}_0),$$

where  $\mathcal{F}_{t-1}$  is the  $\sigma$ -field generated by  $\{Z_s, s \leq t-1\}$ , for a known function mon  $[0, \infty)^{\infty} \times \mathbb{R}^d$  to  $(\delta, +\infty)$  for some  $\delta > 0$ ,

$$m_t(\boldsymbol{\eta}_0) := m(Z_{t-1}, Z_{t-2}, \ldots; \boldsymbol{\eta}_0),$$

for a known function v on  $[0,\infty)^{\infty} \times \mathbb{R}^{d'}$  to  $(\delta, +\infty)$ ,

$$v_t(\boldsymbol{\theta}_0) := v(Z_{t-1}, Z_{t-2}, \ldots; \boldsymbol{\theta}_0),$$

and  $(\boldsymbol{\eta}_0^{\top}, \boldsymbol{\theta}_0^{\top})^{\top} \in \mathbb{R}^{d+d'}$  is unknown parameter.

Since  $m_t(\boldsymbol{\eta})$  and  $v_t(\boldsymbol{\theta})$  include unobservable variables  $Z_0, Z_{-1}, \ldots$ , define

$$\tilde{m}_t(\boldsymbol{\eta}) := m(Z_{t-1}, Z_{t-2}, \dots, Z_1, \boldsymbol{x}_0; \boldsymbol{\eta})$$

and

$$\tilde{v}_t(\boldsymbol{\theta}) := v(Z_{t-1}, Z_{t-2}, \dots, Z_1, \boldsymbol{x}_0; \boldsymbol{\theta})$$

for some constant  $x_0$ . We assume the following condition throughout this Appendix.

Assumption A.1. (A)  $\{Z_t\}$  is strictly stationary and ergodic.

The M-estimator of  $\eta_0$  is defined as

$$\hat{\boldsymbol{\eta}}_n := rg \max_{\boldsymbol{\eta} \in \boldsymbol{H}} \tilde{L}_{n,m}(\boldsymbol{\eta}),$$

where  $\tilde{L}_{n,m}(\eta) := \frac{1}{n} \sum_{t=1}^{n} \ell_m(Z_t, \tilde{m}_t(\eta))$  and  $\ell_m$  is a measurable function. We assume the next condition to prove the strong consistency and the asymptotic equation  $\tilde{L}_{n,m}(\eta) = 0$ .

We assume the next condition to prove the strong consistency and the asymptotic normality of  $\hat{\eta}_n$ .

Assumption A.2. (B1) It holds that

$$\sup_{\boldsymbol{\eta}\in\boldsymbol{H}} |\ell_m(Z_t,\tilde{m}_t(\boldsymbol{\eta})) - \ell_m(Z_t,m_t(\boldsymbol{\eta}))| \to 0 \quad \text{a.s. as } t \to \infty.$$

(B2) The function m is almost surely continuous with respect to  $\eta$  and  $\ell_m(\cdot, \cdot)$  is almost surely continuous with respect to its second component.

(B3) The expectation of  $\ell_m(Z_t, m_t(\eta_0))$  is finite.

(B4) The expectation of  $\ell_m(Z_t, m_t(\eta_0))$  has a unique maximum at  $\eta_0$ .

(B5) The parameter space H is compact.

(B6) The function m is twice continuously differentiable with respect to  $\eta$  and  $\ell_m(\cdot, \cdot)$  is twice continuously differentiable with respect to its second component.

(B7) It holds that

$$\sup_{\boldsymbol{\eta}\in\boldsymbol{H}} \left\| \frac{\partial}{\partial\boldsymbol{\eta}} \ell_m(Z_t, \tilde{m}_t(\boldsymbol{\eta})) - \frac{\partial}{\partial\boldsymbol{\eta}} \ell_m(Z_t, m_t(\boldsymbol{\eta})) \right\| = O(t^{-\frac{1}{2}-\delta}) \quad \text{a.s. as } t \to \infty.$$

(B8) The true parameter  $\eta_0$  belongs to the interior of H.

(B9) There exists a neighborhood  $B(\eta_0)$  of  $\eta_0$  such that

$$\mathbb{E}\left(\sup_{\boldsymbol{\eta}\in B(\boldsymbol{\eta}_0)}\left\|\frac{\partial^2}{\partial\boldsymbol{\eta}\partial\boldsymbol{\eta}^{\top}}\ell_m(Z_t,m_t(\boldsymbol{\eta}))\right\|\right)<\infty.$$

(B10) It holds that  $\mathbb{E}\left(\ell'_m(Z_t, m_t(\eta_0)) \mid \mathcal{F}_{t-1}\right) = 0$  a.s., where  $\ell'_m(\cdot, \cdot)$  is the first derivative of  $\ell_m(\cdot, \cdot)$  with respect to its second component.

(B11) The following holds true:  $E(\ell''_m(Z_t, m_t(\boldsymbol{\eta}_0)) | \mathcal{F}_{t-1})$  is almost surely negative, where  $\ell''_m(\cdot, \cdot)$  is the second derivative of  $\ell_m(\cdot, \cdot)$  with respect to its second component, and if  $\boldsymbol{s}^{\top} \frac{\partial}{\partial \boldsymbol{\eta}} m_t(\boldsymbol{\eta}_0) = 0$ , then  $\boldsymbol{s} = \boldsymbol{0}$ .

(B12) It holds that

$$\mathbb{E} \left\| \frac{\partial}{\partial \boldsymbol{\eta}} \ell_m(Z_t, m_t(\boldsymbol{\eta}_0)) \right\|^{2+\delta} < \infty.$$

The following lemma is due to Theorems 1 and 2 of Goto and Fujimori (2023).

**Lemma A.1.** Under Assumptions (A) and (B1)–(B5),  $\hat{\eta}_n$  tends almost surely to  $\eta_0$  as  $n \to \infty$ . Moreover, under Assumptions (A) and (B1)–(B12),

$$\sqrt{n}(\hat{\boldsymbol{\eta}}_n - \boldsymbol{\eta}_0) \Rightarrow N(0, {\boldsymbol{J}_m}^{-1} {\boldsymbol{I}_m} {\boldsymbol{J}_m}^{-1}) \quad as \ n \to \infty,$$

where

$$\begin{aligned} \boldsymbol{J}_{m} &:= \mathbb{E}\left(\frac{\partial^{2}}{\partial\boldsymbol{\eta}\partial\boldsymbol{\eta}^{\top}}\ell_{m}(\boldsymbol{Z}_{t},\boldsymbol{m}_{t}(\boldsymbol{\eta}_{0}))\right) \\ &= \mathbb{E}\left(\ell_{m}^{\prime\prime}(\boldsymbol{Z}_{t},\boldsymbol{m}_{t}(\boldsymbol{\eta}_{0}))\frac{\partial}{\partial\boldsymbol{\eta}}\boldsymbol{m}_{t}(\boldsymbol{\eta}_{0})\frac{\partial}{\partial\boldsymbol{\eta}^{\top}}\boldsymbol{m}_{t}(\boldsymbol{\eta}_{0})\right) \\ and \ \boldsymbol{I}_{m} &:= \mathbb{E}\left(\frac{\partial}{\partial\boldsymbol{\eta}}\ell_{m}(\boldsymbol{Z}_{t},\boldsymbol{m}_{t}(\boldsymbol{\eta}_{0}))\frac{\partial}{\partial\boldsymbol{\eta}^{\top}}\ell_{m}(\boldsymbol{Z}_{t},\boldsymbol{m}_{t}(\boldsymbol{\eta}_{0}))\right) \\ &= \mathbb{E}\left(\left(\ell_{m}^{\prime}(\boldsymbol{Z}_{t},\boldsymbol{m}_{t}(\boldsymbol{\eta}_{0}))\right)^{2}\frac{\partial}{\partial\boldsymbol{\eta}}\boldsymbol{m}_{t}(\boldsymbol{\eta}_{0})\frac{\partial}{\partial\boldsymbol{\eta}^{\top}}\boldsymbol{m}_{t}(\boldsymbol{\eta}_{0})\right) \end{aligned}$$

Next, we consider the estimation of  $\theta_0$ . The M-estimator of  $\theta_0$  is defined by

$$\hat{\boldsymbol{\theta}}_n := \underset{\boldsymbol{\theta}\in\boldsymbol{\Theta}}{\arg\max} \tilde{L}_{n,v}(\hat{\boldsymbol{\eta}}_n, \boldsymbol{\theta}),$$

where  $\tilde{L}_{n,v}(\hat{\eta}_n, \theta) := \frac{1}{n} \sum_{t=1}^n \ell_v(Z_t, \tilde{m}_t(\hat{\eta}_n), \tilde{v}_t(\theta))$  and  $\ell_v$  is a measurable function.

We suppose the following conditions.

Assumption A.3. (C1) It holds that

$$\sup_{(\boldsymbol{\eta},\boldsymbol{\theta})\in\boldsymbol{H}\times\boldsymbol{\Theta}} |\ell_v(Z_t,\tilde{m}_t(\boldsymbol{\eta}),\tilde{v}_t(\boldsymbol{\theta})) - \ell_v(Z_t,m_t(\boldsymbol{\eta}),v_t(\boldsymbol{\theta}))| \to 0 \quad \text{a.s. as } t \to \infty.$$

(C2) The function  $\ell_v(\cdot, \cdot, \cdot)$  is differentiable with respect to its second and third components and, for some neighborhood  $B(\eta_0)$  of  $\eta_0$ ,

$$\mathbb{E}\left(\sup_{\boldsymbol{\eta}\in B(\boldsymbol{\eta}_0)}\sup_{\boldsymbol{\theta}\in\Theta}\left\|\frac{\partial}{\partial\boldsymbol{\eta}}\ell_v(Z_t,m_t(\boldsymbol{\eta}),v_t(\boldsymbol{\theta}))\right\|\right)<\infty.$$

(C3) The function v is almost surely continuous with respect to  $\theta$  and  $\ell_v(\cdot, \cdot, \cdot)$  is almost surely continuous with respect to its third component.

(C4) The expectation of  $\ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}_0))$  is finite.

(C5) The expectation of  $\ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}))$  has a unique maximum at  $\boldsymbol{\theta}_0$ .

(C6) The parameter space  $\Theta$  is compact.

(C7) The function v is twice continuously differentiable with respect to  $\theta$ , and

$$\frac{\partial^2}{\partial z \partial y} \ell_v(x, y, z)$$
 and  $\frac{\partial^2}{\partial z \partial z} \ell_v(x, y, z)$ 

exist and are continuous with respect to their second and third components.

(C8) It holds that

$$\sup_{\substack{(\boldsymbol{\eta},\boldsymbol{\theta})\in\boldsymbol{H}\times\boldsymbol{\Theta}}} \left\| \frac{\partial}{\partial\boldsymbol{\theta}} \ell_v(Z_t, \tilde{m}_t(\boldsymbol{\eta}), \tilde{v}_t(\boldsymbol{\theta})) - \frac{\partial}{\partial\boldsymbol{\theta}} \ell_v(Z_t, m_t(\boldsymbol{\eta}), v_t(\boldsymbol{\theta})) \right\|$$
$$= O(t^{-\frac{1}{2}-\delta}) \quad \text{a.s. as } t \to \infty.$$

(C9) The true parameter  $\theta_0$  belongs to the interior of  $\Theta$ .

(C10) There exists neighborhoods  $B_1(\eta_0)$  of  $\eta_0$  and  $B_2(\theta_0)$  of  $\theta_0$  such that

$$\mathbb{E}\left(\sup_{(\boldsymbol{\eta},\boldsymbol{\theta})\in B_1(\boldsymbol{\eta}_0)\times B_2(\boldsymbol{\theta}_0)}\left\|\frac{\partial^2}{\partial\boldsymbol{\theta}\partial\boldsymbol{\eta}^{\top}}\ell_v(Z_t,m_t(\boldsymbol{\eta}),v_t(\boldsymbol{\theta}))\right\|\right) < \infty$$

and

$$\mathbb{E}\left(\sup_{(\boldsymbol{\eta},\boldsymbol{\theta})\in B_1(\boldsymbol{\eta}_0)\times B_2(\boldsymbol{\theta}_0)}\left\|\frac{\partial^2}{\partial\boldsymbol{\theta}\partial\boldsymbol{\theta}^{\top}}\ell_v(Z_t,m_t(\boldsymbol{\eta}),v_t(\boldsymbol{\theta}))\right\|\right)<\infty.$$

(C11) It holds that  $E(\ell'_v(Z_t, m_t(\eta_0), v_t(\theta_0)) | \mathcal{F}_{t-1}) = 0$  a.s., where  $\ell'_v(\cdot, \cdot, \cdot)$  is the first derivative of  $\ell_v(\cdot, \cdot, \cdot)$  with respect to its third component.

(C12) The following holds true:  $\mathbb{E}\left(\ell_v''(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}_0)) \mid \mathcal{F}_{t-1}\right)$  is almost surely negative, where  $\ell_v''(\cdot, \cdot, \cdot)$  is the second derivative of  $\ell_v(\cdot, \cdot, \cdot)$  with respect to its third component, and if  $\boldsymbol{s}^\top \frac{\partial}{\partial \boldsymbol{\theta}} v_t(\boldsymbol{\theta}_0) = 0$ , then  $\boldsymbol{s} = \mathbf{0}$ .

(C13) It holds that

$$\mathbb{E} \left\| \frac{\partial}{\partial \boldsymbol{\theta}} \ell_{v}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0}), v_{t}(\boldsymbol{\theta}_{0})) \right\|^{2+\delta} < \infty$$

The following theorem shows that the strong consistency of  $\hat{\theta}_n$  and asymptotic joint normality of  $\hat{\eta}_n$  and  $\hat{\theta}_n$ .

**Theorem A.1.** Under (A), (B1)–(B12), and (C1)–(C6),  $\hat{\theta}_n$  tends almost surely to  $\theta_0$  as  $n \to \infty$ . Moreover, under (A), (B1)–(B12), and (C1)–(C14), it holds that

$$\sqrt{n} \begin{pmatrix} \hat{\boldsymbol{\eta}}_n - \boldsymbol{\eta}_0 \\ \hat{\boldsymbol{\theta}}_n - \boldsymbol{\theta}_0 \end{pmatrix} \Rightarrow N(\mathbf{0}, \mathbf{V}) \quad as \ n \to \infty,$$

17

where  $V := (v_{ij})_{i,j=1,2}$  and

$$\begin{split} \boldsymbol{v}_{11} &:= \boldsymbol{J}_m^{-1} \boldsymbol{I}_m \boldsymbol{J}_m^{-1}, \\ \boldsymbol{v}_{12} &:= -\boldsymbol{J}_m^{-1} \boldsymbol{I}_m \boldsymbol{J}_m^{-1} \boldsymbol{J}_{vm}^{\top} \boldsymbol{J}_v^{-1} + \boldsymbol{J}_m^{-1} \boldsymbol{I}_{mv} \boldsymbol{J}_v^{-1} \\ &= \boldsymbol{J}_m^{-1} (\boldsymbol{I}_{mv} - \boldsymbol{I}_m \boldsymbol{J}_m^{-1} \boldsymbol{J}_{vm}^{\top}) \boldsymbol{J}_v^{-1}, \\ \boldsymbol{v}_{21} &:= -\boldsymbol{J}_v^{-1} \boldsymbol{J}_{vm} \boldsymbol{J}_m^{-1} \boldsymbol{I}_m \boldsymbol{J}_m^{-1} + \boldsymbol{J}_v^{-1} \boldsymbol{I}_{mv}^{\top} \boldsymbol{J}_m^{-1} \\ &= \boldsymbol{v}_{12}^{\top}, \\ \boldsymbol{v}_{22} &:= \boldsymbol{J}_v^{-1} \boldsymbol{J}_{vm} \boldsymbol{J}_m^{-1} \boldsymbol{I}_m \boldsymbol{J}_m^{-1} \boldsymbol{J}_{vm}^{\top} \boldsymbol{J}_v^{-1} - \boldsymbol{J}_v^{-1} \boldsymbol{I}_{mv}^{\top} \boldsymbol{J}_m^{-1} \boldsymbol{J}_v^{\top} \boldsymbol{J}_v^{-1} \\ &\quad - \boldsymbol{J}_v^{-1} \boldsymbol{J}_{vm} \boldsymbol{J}_m^{-1} \boldsymbol{I}_m \boldsymbol{J}_v^{-1} + \boldsymbol{J}_v^{-1} \boldsymbol{I}_v \boldsymbol{J}_v^{-1} \\ &= \boldsymbol{J}_v^{-1} (\boldsymbol{I}_v + \boldsymbol{J}_{vm} \boldsymbol{J}_m^{-1} \boldsymbol{I}_m \boldsymbol{J}_w^{-1} \boldsymbol{J}_v^{\top} - \boldsymbol{I}_m^{\top} \boldsymbol{J}_m^{-1} \boldsymbol{J}_v^{\top} \boldsymbol{J}_w^{-1} \boldsymbol{J}_v^{-1} \\ \end{split}$$

with

$$\begin{split} \boldsymbol{J}_{vm} &:= \mathbb{E}\left(\frac{\partial^2}{\partial \boldsymbol{\theta} \partial \boldsymbol{\eta}^{\top}} \ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}_0))\right), \\ \boldsymbol{J}_v &:= \mathbb{E}\left(\frac{\partial^2}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^{\top}} \ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}_0))\right), \\ \boldsymbol{I}_{mv} &:= \mathbb{E}\left(\frac{\partial}{\partial \boldsymbol{\eta}} \ell_m(Z_t, m_t(\boldsymbol{\eta}_0)) \frac{\partial}{\partial \boldsymbol{\theta}^{\top}} \ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}_0))\right), \\ and \ \boldsymbol{I}_v &:= \mathbb{E}\left(\frac{\partial}{\partial \boldsymbol{\theta}} \ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}_0)) \frac{\partial}{\partial \boldsymbol{\theta}^{\top}} \ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}_0))\right). \end{split}$$

# Appendix B: Proof of Theorem A.1

In this section, we prove Theorem A.1. We follow the proof of Ahmad and Francq (2016, Theorems 2.1 and 2.2) and Francq and Zakoian (2010, Theorem 7.1, p.159).

First, we prove the strong consistency of  $\hat{\theta}_n$ . Assumption (C1) gives that the effect of initial value is asymptotically negligible, that is,

$$\begin{split} \sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}} \left| \tilde{L}_{n,v}(\boldsymbol{\hat{\eta}}_{n},\boldsymbol{\theta}) - L_{n,v}(\boldsymbol{\hat{\eta}}_{n},\boldsymbol{\theta}) \right| \\ \leq & \frac{1}{n} \sum_{t=1}^{n} \sup_{(\boldsymbol{\eta},\boldsymbol{\theta})\in\boldsymbol{H}\times\boldsymbol{\Theta}} \left| \ell_{v}(Z_{t},\tilde{m}_{t}(\boldsymbol{\eta}),\tilde{v}_{t}(\boldsymbol{\theta})) - \ell_{v}(Z_{t},m_{t}(\boldsymbol{\eta}),v_{t}(\boldsymbol{\theta})) \right| \to 0 \quad \text{a.s.} \end{split}$$

as  $n \to \infty$ . Assumptions (B8) and (C2) and the strong consistency of  $\eta_0$  yield that

$$\sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}} |L_{n,v}(\hat{\boldsymbol{\eta}}_{n},\boldsymbol{\theta}) - L_{n,v}(\boldsymbol{\eta}_{0},\boldsymbol{\theta})|$$
  
$$\leq \frac{1}{n} \sum_{t=1}^{n} \sup_{\boldsymbol{\theta}\in\boldsymbol{\Theta}} |\ell_{v}(Z_{t}, m_{t}(\hat{\boldsymbol{\eta}}_{n}), v_{t}(\hat{\boldsymbol{\eta}}_{n},\boldsymbol{\theta})) - \ell_{v}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0}), v_{t}(\boldsymbol{\theta}))|$$

$$\leq \|\hat{\boldsymbol{\eta}}_n - \boldsymbol{\eta}_0\|_{\ell_1} \frac{1}{n} \sum_{t=1}^n \sup_{\boldsymbol{\theta} \in \boldsymbol{\Theta}} \left\| \frac{\partial}{\partial \boldsymbol{\eta}} \ell_v(Z_t, m_t(\hat{\boldsymbol{\eta}}_n^*), v_t(\boldsymbol{\theta})) \right\|_{\ell_1} \to 0 \quad \text{a.s. as } n \to \infty,$$

where  $\boldsymbol{\eta}_0 \leq \hat{\boldsymbol{\eta}}_n^* \leq \hat{\boldsymbol{\eta}}_n$ . Now, we have, for any  $\boldsymbol{\theta}_1$ ,

$$\sup_{\boldsymbol{\theta}\in B(\boldsymbol{\theta}_{1};r)} L_{n,v}(\hat{\boldsymbol{\eta}}_{n},\boldsymbol{\theta})$$

$$= \sup_{\boldsymbol{\theta}\in B(\boldsymbol{\theta}_{1};r)} \tilde{L}_{n,v}(\hat{\boldsymbol{\eta}}_{n},\boldsymbol{\theta}) - \sup_{\boldsymbol{\theta}\in B(\boldsymbol{\theta}_{1};r)} L_{n,v}(\hat{\boldsymbol{\eta}}_{n},\boldsymbol{\theta}) + \sup_{\boldsymbol{\theta}\in B(\boldsymbol{\theta}_{1};r)} L_{n,v}(\hat{\boldsymbol{\eta}}_{n},\boldsymbol{\theta})$$

$$- \sup_{\boldsymbol{\theta}\in B(\boldsymbol{\theta}_{1};r)} L_{n,v}(\boldsymbol{\eta}_{0},\boldsymbol{\theta}) + \sup_{\boldsymbol{\theta}\in B(\boldsymbol{\theta}_{1};r)} L_{n,v}(\boldsymbol{\eta}_{0},\boldsymbol{\theta}),$$

where  $B(\theta_1; r) := \{\theta : \|\theta - \theta_1\|_{\ell_1} < 1/r\}$ , and so by the above argument and the ergodic theorem for non-integrable processes (see Francq and Zakoian, 2010, p.181, Problem 7.3), we observe

$$\limsup_{n\to\infty}\sup_{\boldsymbol{\theta}\in B(\boldsymbol{\theta}_1;r)}\tilde{L}_{n,v}(\hat{\boldsymbol{\eta}}_n,\boldsymbol{\theta})\leq \mathbb{E}\left(\sup_{\boldsymbol{\theta}\in B(\boldsymbol{\theta}_1;r)}\ell_v(Z_t,m_t(\boldsymbol{\eta}_0),v_t(\boldsymbol{\theta}))\right).$$

Here we used the inequality

$$\sup_{\boldsymbol{\theta}\in B(\boldsymbol{\theta}_1;r)} L_{n,v}(\boldsymbol{\eta}_0,\boldsymbol{\theta}) \leq \frac{1}{n} \sum_{t=1}^n \sup_{\boldsymbol{\theta}\in B(\boldsymbol{\theta}_1;r)} \ell_v(Z_t,m_t(\boldsymbol{\eta}_0),v_t(\boldsymbol{\theta})).$$

The Beppo-Levi theorem and Assumption (C3) yield that

$$\operatorname{E}\left(\sup_{\boldsymbol{\theta}\in B(\boldsymbol{\theta}_{1};r)}\ell_{v}(Z_{t},m_{t}(\boldsymbol{\eta}_{0}),v_{t}(\boldsymbol{\theta}))\right)\to\operatorname{E}\left(\ell_{v}(Z_{t},m_{t}(\boldsymbol{\eta}_{0}),v_{t}(\boldsymbol{\theta}_{1}))\right)\quad\text{as }r\to\infty.$$

Therefore, for any  $\theta_1 \neq \theta_0$ , there exists a neighborhood  $B(\theta_1)$  satisfying

$$\begin{split} \limsup_{n \to \infty} \sup_{\boldsymbol{\theta} \in B(\boldsymbol{\theta}_1)} \tilde{L}_{n,v}(\boldsymbol{\hat{\eta}}_n, \boldsymbol{\theta}) &\leq \operatorname{E}\left(\ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\eta}_0, \boldsymbol{\theta}_1))\right) \quad a.s. \\ &< \operatorname{E}\left(\ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}_0))\right) \quad a.s. \\ &= \lim_{n \to \infty} \tilde{L}_{n,v}(\boldsymbol{\hat{\eta}}_n, \boldsymbol{\theta}_0) \quad a.s. \end{split}$$

Here we used Assumption (C5). By Assumption (C6), there exists, for any covering set  $\{B(\theta_i); i \in \{0\} \cup \mathbb{N}\}$  of  $\Theta$  such that  $\theta_i \in \Theta \setminus B(\theta_0)$  for all  $i \in \mathbb{N}$  and

$$\limsup_{n\to\infty}\sup_{\boldsymbol{\theta}\in B(\boldsymbol{\theta}_1)}\tilde{L}_{n,v}(\hat{\boldsymbol{\eta}}_n,\boldsymbol{\theta})<\lim_{n\to\infty}\tilde{L}_{n,v}(\hat{\boldsymbol{\eta}}_n,\boldsymbol{\theta}_0),$$

the finite covering set  $\{B'(\boldsymbol{\theta}_i); i = 0, 1, \dots, s\}$  such that  $\Theta := \bigcup_{i=0}^s B'(\boldsymbol{\theta}_i)$  and  $B'(\boldsymbol{\theta}_0) = B(\boldsymbol{\theta}_0)$ . Then, we observe that, for sufficiently large n,

$$\sup_{\boldsymbol{\theta}\in\Theta}\tilde{L}_{n,v}(\hat{\boldsymbol{\eta}}_n,\boldsymbol{\theta}) = \max_{i=0,1,\dots,s} \sup_{\boldsymbol{\theta}\in B'(\boldsymbol{\theta}_i)}\tilde{L}_{n,v}(\hat{\boldsymbol{\eta}}_n,\boldsymbol{\theta})$$

Y.Goto and K.Fujimori/A test for counting sequences of INAR model

$$= \sup_{\boldsymbol{\theta} \in B'(\boldsymbol{\theta}_0)} \tilde{L}_{n,v}(\boldsymbol{\hat{\eta}}_n, \boldsymbol{\theta}) \quad \text{a.s.}$$

Since  $B(\theta_0)$  can be chosen an arbitrary small set, we obtain the strong consistency of  $\hat{\theta}_n$ .

Next, we show the asymptotic normality of  $(\hat{\boldsymbol{\eta}}_n^{\top}, \hat{\boldsymbol{\theta}}_n^{\top})^{\top}$ . From Assumptions (B6), (B8), (C7), (C8), and (C9), we have

$$\begin{aligned} \mathbf{0} &= \frac{1}{\sqrt{n}} \sum_{t=1}^{n} \frac{\partial}{\partial \theta} \ell_{v}(Z_{t}, \tilde{m}_{t}(\hat{\boldsymbol{\eta}}_{n}), \tilde{v}_{t}(\hat{\boldsymbol{\theta}}_{n})) \\ &= \frac{1}{\sqrt{n}} \sum_{t=1}^{n} \frac{\partial}{\partial \theta} \ell_{v}(Z_{t}, m_{t}(\hat{\boldsymbol{\eta}}_{n}), v_{t}(\hat{\boldsymbol{\theta}}_{n})) + o_{p}(1) \\ &= \frac{1}{\sqrt{n}} \sum_{t=1}^{n} \frac{\partial}{\partial \theta} \ell_{v}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0}), v_{t}(\boldsymbol{\theta}_{0})) \\ &+ \frac{1}{n} \sum_{t=1}^{n} \frac{\partial^{2}}{\partial \theta \partial \boldsymbol{\eta}^{\top}} \ell_{v}(Z_{t}, m_{t}(\hat{\boldsymbol{\eta}}_{n}^{**}), v_{t}(\hat{\boldsymbol{\theta}}_{n}^{**})) \sqrt{n}(\hat{\boldsymbol{\eta}}_{n} - \boldsymbol{\eta}_{0}) \\ &+ \frac{1}{n} \sum_{t=1}^{n} \frac{\partial^{2}}{\partial \theta \partial \theta^{\top}} \ell_{v}(Z_{t}, m_{t}(\hat{\boldsymbol{\eta}}_{n}^{**}), v_{t}(\hat{\boldsymbol{\theta}}_{n}^{**})) \sqrt{n}(\hat{\boldsymbol{\theta}}_{n} - \boldsymbol{\theta}_{0}) + o_{p}(1) \\ &= \frac{1}{\sqrt{n}} \sum_{t=1}^{n} \frac{\partial}{\partial \theta} \ell_{v}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0}), v_{t}(\boldsymbol{\theta}_{0})) + \boldsymbol{J}_{vm}^{*} \sqrt{n}(\hat{\boldsymbol{\eta}}_{n} - \boldsymbol{\eta}_{0}) \\ &+ \boldsymbol{J}_{v}^{*} \sqrt{n}(\hat{\boldsymbol{\theta}}_{n} - \boldsymbol{\theta}_{0}) + o_{p}(1), \end{aligned}$$
(B.2)

where  $\boldsymbol{\eta}_0 \stackrel{<}{_{>}} \hat{\boldsymbol{\eta}}_n^{**} \stackrel{\leq}{_{>}} \hat{\boldsymbol{\eta}}_n, \, \boldsymbol{\theta}_0 \stackrel{\leq}{_{>}} \hat{\boldsymbol{\eta}}_n^{**} \stackrel{\leq}{_{>}} \hat{\boldsymbol{\theta}}_n,$ 

$$\boldsymbol{J}_{vm}^* := \frac{1}{n} \sum_{t=1}^n \frac{\partial^2}{\partial \boldsymbol{\theta} \partial \boldsymbol{\eta}^\top} \ell_v(\boldsymbol{Z}_t, \boldsymbol{m}_t(\hat{\boldsymbol{\eta}}_n^{**}), \boldsymbol{v}_t(\hat{\boldsymbol{\theta}}_n^{**})),$$

and

$$\boldsymbol{J}_v^* := \frac{1}{n} \sum_{t=1}^n \frac{\partial^2}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^\top} \ell_v(\boldsymbol{Z}_t, \boldsymbol{m}_t(\boldsymbol{\hat{\eta}}_n^{**}), \boldsymbol{v}_t(\boldsymbol{\hat{\theta}}_n^{**})).$$

The assumption (C10) gives that, for some r such that  $B(\eta_0; r) \times B(\theta_0; r) \subset B_1(\eta_0) \times B_2(\theta_0)$  and for large n such that  $\hat{\eta}_n^{**} \times \hat{\theta}_n^{**} \in B(\eta_0; r) \times B(\theta_0; r)$ ,

$$\begin{aligned} & \left| \frac{1}{n} \sum_{t=1}^{n} \frac{\partial^{2}}{\partial \theta_{ij} \partial \eta_{ij}^{\top}} \ell_{v}(Z_{t}, m_{t}(\hat{\boldsymbol{\eta}}_{n}^{**}), v_{t}(\hat{\boldsymbol{\theta}}_{n}^{**})) - \mathrm{E} \frac{\partial^{2}}{\partial \theta_{ij} \partial \eta_{ij}^{\top}} \ell_{v}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0}), v_{t}(\boldsymbol{\theta}_{0})) \right| \\ \leq & \frac{1}{n} \sum_{t=1}^{n} \sup_{(\boldsymbol{\eta}, \boldsymbol{\theta}) \in B(\boldsymbol{\eta}_{0}; r) \times B(\boldsymbol{\theta}_{0}; r)} \\ & \left| \frac{\partial^{2}}{\partial \theta_{ij} \partial \eta_{ij}^{\top}} \ell_{v}(Z_{t}, m_{t}(\boldsymbol{\eta}), v_{t}(\boldsymbol{\theta})) - \mathrm{E} \frac{\partial^{2}}{\partial \theta_{ij} \partial \eta_{ij}^{\top}} \ell_{v}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0}), v_{t}(\boldsymbol{\theta}_{0})) \right| \end{aligned}$$

imsart-generic ver. 2020/08/06 file: Goto\_Fujimori\_arXiv.tex date: February 5, 2024

$$\rightarrow \mathbf{E} \sup_{(\boldsymbol{\eta}, \boldsymbol{\theta}) \in B(\boldsymbol{\eta}_0; r) \times B(\boldsymbol{\theta}_0; r)} \\ \left| \frac{\partial^2}{\partial \theta_{ij} \partial \eta_{ij}^{\top}} \ell_v(Z_t, m_t(\boldsymbol{\eta}), v_t(\boldsymbol{\theta})) - \mathbf{E} \frac{\partial^2}{\partial \theta_{ij} \partial \eta_{ij}^{\top}} \ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}_0)) \right|$$

a.s. as  $n \to \infty$ . By applying the Beppo-Levi theorem, we have

$$\begin{split} & \underset{(\boldsymbol{\eta},\boldsymbol{\theta})\in B(\boldsymbol{\eta}_{0};r)\times B(\boldsymbol{\theta}_{0};r)}{\sup} \\ & \left| \frac{\partial^{2}}{\partial\theta_{ij}\partial\eta_{ij}^{\top}} \ell_{v}(Z_{t},m_{t}(\boldsymbol{\eta}),v_{t}(\boldsymbol{\theta})) - \mathrm{E}\frac{\partial^{2}}{\partial\theta_{ij}\partial\eta_{ij}^{\top}} \ell_{v}(Z_{t},m_{t}(\boldsymbol{\eta}_{0}),v_{t}(\boldsymbol{\theta}_{0})) \right| \to 0 \end{split}$$

a.s. as  $r \to \infty$ , and thus,  $J_{vm}^*$  converges almost surely to  $J_{vm}$  as  $n \to \infty$ , where

$$\boldsymbol{J}_{vm} := \mathrm{E}\left(\frac{\partial^2}{\partial \boldsymbol{\theta} \partial \boldsymbol{\eta}^{\top}} \ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}_0))\right).$$

Similarly, we obtain  $J_v^*$  converges almost surely to  $J_v$  as  $n \to \infty$ , where

$$\boldsymbol{J}_{v} := \mathrm{E} \frac{\partial^{2}}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^{\top}} \ell_{v}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0}), v_{t}(\boldsymbol{\theta}_{0})).$$

The non-singularity of  $J_v$  is ensured by Assumptions (C12). Actually, we observe

$$\begin{aligned} \boldsymbol{J}_{v} &:= \mathbb{E}\left(\frac{\partial^{2}}{\partial\boldsymbol{\theta}\partial\boldsymbol{\theta}^{\top}}\ell_{v}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\eta}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0}))\right) \\ &= \mathbb{E}\left[\frac{\partial}{\partial\boldsymbol{\theta}^{\top}}\left\{\left(\frac{\partial}{\partial\boldsymbol{\theta}}\boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})\right)\ell_{v}'(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\eta}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0}))\right\}\right] \\ &= \mathbb{E}\left\{\left(\frac{\partial}{\partial\boldsymbol{\theta}}\boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})\right)\left(\frac{\partial}{\partial\boldsymbol{\theta}^{\top}}\boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})\right)\ell_{v}''(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\eta}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0}))\right. \\ &\left.+\left(\frac{\partial^{2}}{\partial\boldsymbol{\theta}\partial\boldsymbol{\theta}^{\top}}\boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})\right)\ell_{v}'(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\eta}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0}))\right\} \\ &= \mathbb{E}\left\{\left(\frac{\partial}{\partial\boldsymbol{\theta}}\boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})\right)\left(\frac{\partial}{\partial\boldsymbol{\theta}^{\top}}\boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})\right)\ell_{v}''(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\eta}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0}))\right\}. \end{aligned}$$

For any  $\boldsymbol{s} \in \mathbb{R}^{d'}$ , it holds

$$\boldsymbol{s}^{\top} \boldsymbol{J}_{\boldsymbol{v}} \boldsymbol{s} = \mathbb{E} \left\{ \left| \boldsymbol{s}^{\top} \frac{\partial}{\partial \boldsymbol{\theta}} \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0}) \right|^{2} \mathbb{E} \left( \ell_{\boldsymbol{v}}^{\prime\prime}(\boldsymbol{Z}_{t}, m_{t}(\boldsymbol{\eta}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})) \mid \mathcal{F}_{t-1} \right) \right\} \geq 0,$$

which shows that  $J_v$  is a non-negative definite matrix. Suppose  $\mathbf{s}^\top J_v \mathbf{s} = 0$ . Then, we have  $\mathbf{s}^\top \frac{\partial}{\partial \theta} v_t(\theta_0) = 0$ , which by Assumption (C11) and (C12), gives  $\mathbf{s} = \mathbf{0}$ . This implies that  $J_v$  is a positive definite matrix. Similarly,  $J_m$  is a positive definite matrix by Assumption (B11). From (B.1) and (B.2), it follows that

$$\begin{pmatrix} \sqrt{n}(\hat{\boldsymbol{\eta}}_n - \boldsymbol{\eta}_0) \\ \sqrt{n}(\hat{\boldsymbol{\theta}}_n - \boldsymbol{\theta}_0) \end{pmatrix}$$

imsart-generic ver. 2020/08/06 file: Goto\_Fujimori\_arXiv.tex date: February 5, 2024

Y.Goto and K.Fujimori/A test for counting sequences of INAR model

$$= \begin{pmatrix} \sqrt{n}(\hat{\eta}_n - \eta_0) \\ -J_v^{*-1}\frac{1}{\sqrt{n}}\sum_{t=1}^n \frac{\partial}{\partial\theta}\ell_v(Z_t, m_t(\eta_0), v_t(\theta_0)) - J_v^{*-1}J_{vm}^*\sqrt{n}(\hat{\eta}_n - \eta_0) \end{pmatrix} \\ + o_p(1) \\ = \begin{pmatrix} \mathcal{I}_p & \mathcal{O}_p \\ -J_v^{*-1}J_{vm}^* & -J_v^{*-1} \end{pmatrix} \begin{pmatrix} \sqrt{n}(\hat{\eta}_n - \eta_0) \\ \frac{1}{\sqrt{n}}\sum_{t=1}^n \frac{\partial}{\partial\theta}\ell_v(Z_t, m_t(\eta_0), v_t(\theta_0)) \end{pmatrix} + o_p(1) \\ = \begin{pmatrix} \mathcal{I}_p & \mathcal{O}_p \\ -J_v^{-1}J_{vm} & -J_v^{-1} \end{pmatrix} \begin{pmatrix} -J_m^{-1}\frac{1}{\sqrt{n}}\sum_{t=1}^n \frac{\partial}{\partial\theta}\ell_w(Z_t, m_t(\eta_0)) \\ \frac{1}{\sqrt{n}}\sum_{t=1}^n \frac{\partial}{\partial\theta}\ell_v(Z_t, m_t(\eta_0), v_t(\theta_0)) \end{pmatrix} + o_p(1), \end{cases}$$

where  $\mathcal{I}_p$  is an identity matrix of size p and  $\mathcal{O}_p$  is a p-by-p zero matrix. The proof will be complete by

$$\begin{pmatrix} \mathcal{I}_p & \mathcal{O}_p \\ -\mathbf{J}_v^{-1}\mathbf{J}_{vm} & -\mathbf{J}_v^{-1} \end{pmatrix} \begin{pmatrix} \mathbf{J}_m^{-1}\mathbf{I}_m\mathbf{J}_m^{-1} & -\mathbf{J}_m^{-1}\mathbf{I}_{mv} \\ -\mathbf{I}_{mv}^{\top}\mathbf{J}_m^{-1} & \mathbf{I}_v \end{pmatrix} \begin{pmatrix} \mathcal{I}_p & \mathcal{O}_p \\ -\mathbf{J}_v^{-1}\mathbf{J}_{vm} & -\mathbf{J}_v^{-1} \end{pmatrix}^{\top} = \mathbf{V},$$

provided we can show that

$$\begin{pmatrix} \boldsymbol{J}_{m}^{-1} \frac{1}{\sqrt{n}} \sum_{t=1}^{n} \frac{\partial}{\partial \boldsymbol{\eta}} \ell_{m}(\boldsymbol{Z}_{t}, m_{t}(\boldsymbol{\eta}_{0})) \\ \frac{1}{\sqrt{n}} \sum_{t=1}^{n} \frac{\partial}{\partial \boldsymbol{\theta}} \ell_{v}(\boldsymbol{Z}_{t}, m_{t}(\boldsymbol{\eta}_{0}), v_{t}(\boldsymbol{\theta}_{0})) \end{pmatrix}$$
(B.3)

$$\Rightarrow N\left(\mathbf{0}_{d+d'}, \begin{pmatrix} \boldsymbol{J}_m^{-1}\boldsymbol{I}_m\boldsymbol{J}_m^{-1} & -\boldsymbol{J}_m^{-1}\boldsymbol{I}_{mv} \\ -\boldsymbol{I}_{mv}^{\top}\boldsymbol{J}_m^{-1} & \boldsymbol{I}_v \end{pmatrix}\right) \tag{B.4}$$

as  $n \to \infty$ , where

$$\boldsymbol{I}_{mv} := \mathbf{E} \left( \frac{\partial}{\partial \boldsymbol{\eta}} \ell_m(Z_t, m_t(\boldsymbol{\eta}_0)) \frac{\partial}{\partial \boldsymbol{\theta}^{\top}} \ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}_0)) \right)$$
  
and  $\boldsymbol{I}_v := \mathbf{E} \left( \frac{\partial}{\partial \boldsymbol{\theta}} \ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}_0)) \frac{\partial}{\partial \boldsymbol{\theta}^{\top}} \ell_v(Z_t, m_t(\boldsymbol{\eta}_0), v_t(\boldsymbol{\theta}_0)) \right).$ 

From Assumptions (B12) and (C11), we know that

$$\begin{pmatrix} \boldsymbol{J}_m^{-1} \frac{1}{\sqrt{n}} \sum_{t=1}^n \frac{\partial}{\partial \boldsymbol{\eta}} \ell_m(\boldsymbol{Z}_t, \boldsymbol{m}_t(\boldsymbol{\eta}_0)) \\ \frac{1}{\sqrt{n}} \sum_{t=1}^n \frac{\partial}{\partial \boldsymbol{\theta}} \ell_v(\boldsymbol{Z}_t, \boldsymbol{m}_t(\boldsymbol{\eta}_0), \boldsymbol{v}_t(\boldsymbol{\theta}_0)) \end{pmatrix}$$

is a martingale difference sequence. Assumptions (B12) and (C13) ensures the Lindeberg condition: for any constant  $(\boldsymbol{c}_1^{\top}, \boldsymbol{c}_2^{\top})^{\top} \in \mathbb{R}^{d+d'}$ ,

$$\frac{1}{\sqrt{n}} \sum_{t=1}^{n} \mathbb{E} \left[ \left( c_{1}^{\top} \boldsymbol{J}_{m}^{-1} \sum_{t=1}^{n} \frac{\partial}{\partial \boldsymbol{\eta}} \ell_{m}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\eta}_{0})) + c_{2}^{\top} \sum_{t=1}^{n} \frac{\partial}{\partial \boldsymbol{\theta}} \ell_{v}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\eta}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})) \right)^{2} \times \mathbb{I} \left\{ \frac{1}{\sqrt{n}} \middle| c_{1}^{\top} \boldsymbol{J}_{m}^{-1} \sum_{t=1}^{n} \frac{\partial}{\partial \boldsymbol{\eta}} \ell_{m}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\eta}_{0})) \right\}$$

imsart-generic ver. 2020/08/06 file: Goto\_Fujimori\_arXiv.tex date: February 5, 2024

Y.Goto and K.Fujimori/A test for counting sequences of INAR model

$$+ c_{2}^{\top} \sum_{t=1}^{n} \frac{\partial}{\partial \theta} \ell_{v}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0}), v_{t}(\boldsymbol{\theta}_{0})) \bigg| > \epsilon \bigg\} \bigg]$$

$$= \mathbb{E} \bigg[ \bigg( c_{1}^{\top} \boldsymbol{J}_{m}^{-1} \sum_{t=1}^{n} \frac{\partial}{\partial \eta} \ell_{m}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0})) + c_{2}^{\top} \sum_{t=1}^{n} \frac{\partial}{\partial \theta} \ell_{v}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0}), v_{t}(\boldsymbol{\theta}_{0})) \bigg)^{2}$$

$$\times \mathbb{I} \bigg\{ \frac{1}{(\epsilon \sqrt{n})^{\delta}} \bigg| c_{1}^{\top} \boldsymbol{J}_{m}^{-1} \sum_{t=1}^{n} \frac{\partial}{\partial \eta} \ell_{m}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0}))$$

$$+ c_{2}^{\top} \sum_{t=1}^{n} \frac{\partial}{\partial \theta} \ell_{v}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0}), v_{t}(\boldsymbol{\theta}_{0})) \bigg|^{\delta} > 1 \bigg\} \bigg],$$

$$\frac{1}{(\epsilon \sqrt{n})^{\delta}} \mathbb{E} \bigg| c_{1}^{\top} \boldsymbol{J}_{m}^{-1} \sum_{t=1}^{n} \frac{\partial}{\partial \eta} \ell_{m}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0})) + c_{2}^{\top} \sum_{t=1}^{n} \frac{\partial}{\partial \theta} \ell_{v}(Z_{t}, m_{t}(\boldsymbol{\eta}_{0}), v_{t}(\boldsymbol{\theta}_{0})) \bigg|^{2+\delta},$$

which tends to zero as  $n \to \infty$ .

The ergodic theorem yields that

$$\frac{1}{\sqrt{n}} \sum_{t=1}^{n} \mathbf{E} \left\{ \left( c_{1}^{\top} \boldsymbol{J}_{m}^{-1} \sum_{t=1}^{n} \frac{\partial}{\partial \boldsymbol{\eta}} \ell_{m}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\eta}_{0})) + c_{2}^{\top} \sum_{t=1}^{n} \frac{\partial}{\partial \boldsymbol{\theta}} \ell_{v}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\eta}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})) \right)^{2} \mid \mathcal{F}_{t-1} \right\}$$
$$\rightarrow \mathbf{E} \left( c_{1}^{\top} \boldsymbol{J}_{m}^{-1} \sum_{t=1}^{n} \frac{\partial}{\partial \boldsymbol{\eta}} \ell_{m}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\eta}_{0})) + c_{2}^{\top} \sum_{t=1}^{n} \frac{\partial}{\partial \boldsymbol{\theta}} \ell_{v}(\boldsymbol{Z}_{t}, \boldsymbol{m}_{t}(\boldsymbol{\eta}_{0}), \boldsymbol{v}_{t}(\boldsymbol{\theta}_{0})) \right)^{2} \right\}$$

as  $n \to \infty$ . Therefore, the Cramer–Wold device and the martingale central limit theorem yield (B.3) and (B.4).

#### Acknowledgements

Authors would like to express sincere thanks to anonymous reviewers, which greatly improved the earlier version of this paper.

# References

- AHMAD, A. and FRANCQ, C. (2016). Poisson QMLE of count time series models. J. Time Series Anal. 37 291–314.
- AKNOUCHE, A. and FRANCQ, C. (2021). Two-stage weighted least squares estimator of the conditional mean of observation-driven time series models. *J. Econometrics* **237** 105174.
- AL-OSH, M. A. and ALZAID, A. A. (1987). First-order integer-valued autoregressive (INAR (1)) process. J. Time Ser. Anal. 8 261–275.

imsart-generic ver. 2020/08/06 file: Goto\_Fujimori\_arXiv.tex date: February 5, 2024

- ALEKSANDROV, B., WEISS, C. H. and JENTSCH, C. (2022). Goodness-of-fit tests for Poisson count time series based on the Stein-Chen identity. *Stat. Neerl.* **76** 35–64.
- ALY, E.-E. A. and BOUZAR, N. (2019). Expectation thinning operators based on linear fractional probability generating functions. J. Indian Soc. Probab. Stat. 20 89–107.
- BÖHNING, D. and VAN DER HEIJDEN, P. G. M. (2019). The identity of the zero-truncated, one-inflated likelihood and the zero-one-truncated likelihood for general count densities with an application to drink-driving in Britain. Ann. Appl. Stat. 13 1198 – 1211.
- BOURGUIGNON, M. and WEISS, C. H. (2017). An INAR (1) process for modeling count time series with equidispersion, underdispersion and overdispersion. *Test* 26 847–868.
- BROCKWELL, P. J. and DAVIS, R. A. (2009). *Time series: theory and methods*. Springer science & business media.
- CARLSTEIN, E. (1986). The use of subseries values for estimating the variance of a general statistic from a stationary sequence. Ann. Statist. 1171–1179.
- CHEN, C. W. and LEE, S. (2017). Bayesian causality test for integer-valued time series models with applications to climate and crime data. J. R. Stat. Soc. Ser. C 66 797–814.
- DAVIS, R. A., FOKIANOS, K., HOLAN, S. H., JOE, H., LIVSEY, J., LUND, R., PIPIRAS, V. and RAVISHANKER, N. (2021). Count time series: A methodological review. J. Amer. Statist. Assoc. 116 1533–1547.
- DOUKHAN, P., FOKIANOS, K. and LI, X. (2012). On weak dependence conditions: The case of discrete valued processes. *Stat. Probab. Lett.* 82 1941–1948.
- DOUKHAN, P., LATOUR, A. and ORAICHI, D. (2006). A simple integer-valued bilinear time series model. *Adv. Appl. Probab.* **38** 559–578.
- DROST, F. C., VAN DEN AKKER, R. and WERKER, B. J. (2008). Note on integer-valued bilinear time series models. *Stat. Probab. Lett.* **78** 992–996.
- FERLAND, R., LATOUR, A. and ORAICHI, D. (2006). Integer-valued GARCH process. J. Time Ser. Anal. 27 923–942.
- FOKIANOS, K. and NEUMANN, M. H. (2013). A goodness-of-fit test for Poisson count processes. *Electron. J. Statist.* 7 793–819.
- FRANCQ, C. and ZAKOIAN, J. M. (2010). GARCH models: structure, statistical inference and financial applications. John Wiley & Sons.
- GOTO, Y. and FUJIMORI, K. (2023). Test for conditional variance of integervalued time series. *Stat. Sin.* **32**.
- HUDECOVÁ, Ś., HUŠKOVÁ, M. and MEINTANIS, S. G. (2015). Tests for time series of counts based on the probability-generating function. *Statistics* **49** 316–337.
- HUDECOVÁ, Ś., HUŠKOVÁ, M. and MEINTANIS, S. G. (2021). Goodness-of-fit tests for bivariate time series of counts. *Econometrics* **9** 10.
- JAZI, M. A., JONES, G. and LAI, C.-D. (2012). First-order integer valued AR processes with zero inflated Poisson innovations. J. Time Ser. Anal. 33 954–963.
- KACHOUR, M. and TRUQUET, L. (2011). A p-Order signed integer-valued au-

toregressive (SINAR (p)) model. J. Time Ser. Anal. 32 223–236.

- KLIMKO, L. A. and NELSON, P. I. (1978). On conditional least squares estimation for stochastic processes. Ann. Statist. 629–642.
- LATOUR, A. (1997). The multivariate GINAR (p) process. *Adv. Appl. Probab.* **29** 228–248.
- LATOUR, A. and TRUQUET, L. (2009). An integer-valued bilinear type model.
- LEUCHT, A. and NEUMANN, M. H. (2013). Degenerate U-and V-statistics under ergodicity: asymptotics, bootstrap and applications in statistics. Ann. Inst. Statist. Math. 65 349–386.
- LI, Q., CHEN, H. and ZHU, F. (2023). Z-valued time series: Models, properties and comparison. J. Statist. Plann. Inference 106099.
- MCKENZIE, E. (1985). Some simple models for discrete variate time series. J. Am. Water Resour. Assoc. 21 645–650.
- MEINTANIS, S. G. and KARLIS, D. (2014). Validation tests for the innovation distribution in INAR time series models. *Comput. Statist.* **29** 1221–1241.
- MOHAMMADPOUR, M., BAKOUCH, H. S. and SHIROZHAN, M. (2018). Poisson– Lindley INAR(1) model with applications. *Braz. J. Probab. Stat.* **32** 262 – 280.
- NEUMANN, M. H. (2011). Absolute regularity and ergodicity of poisson count processes. *Bernoulli* **17** 1268–1284.
- NISHIYAMA, Y. (2022). Martingale methods in statistics. Monographs on Statistics and Applied Probability **170**. CRC Press, Boca Raton, FL.
- PELIGRAD, M. and SHAO, Q.-M. (1994). Self-normalized central limit theorem for sums of weakly dependent random variables. J. Theor. Probab. 7 309–338.
- RISTIĆ, M. M., BAKOUCH, H. S. and NASTIĆ, A. S. (2009). A new geometric first-order integer-valued autoregressive (NGINAR (1)) process. J. Statist. Plann. Inference 139 2218–2226.
- SCHMIDT, A. M. and PEREIRA, J. B. M. (2011). Modelling time series of counts in epidemiology. Int. Stat. Rev. 79 48–69.
- SCHWEER, S. (2016). A Goodness-of-Fit Test for Integer-Valued Autoregressive Processes. J. Time Ser. Anal. 37 77–98.
- WEISS, C. H. (2008). Thinning operations for modeling time series of counts—a survey. Adv. Stat. Anal. 92 319–341.
- WEISS, C. H. (2018). Goodness-of-fit testing of a count time series' marginal distribution. *Metrika* 81 619–651.
- WEISS, C. H., HOMBURG, A. and PUIG, P. (2019). Testing for zero inflation and overdispersion in INAR (1) models. *Stat. Pap.* **60** 823–848.
- XU, X., . YIJIONG, Z., Y., C., LIU, Y., GOTO, Y. and TANIGUCHI, M. (2023). Long-memory log-linear zero-inflated generalized Poisson autoregression for Covid-19 pandemic modeling.
- YANG, K., KANG, Y., WANG, D., LI, H. and DIAO, Y. (2019). Modeling overdispersed or underdispersed count data with generalized Poisson integervalued autoregressive processes. *Metrika* 82 863–889.
- ZHU, R. and JOE, H. (2003). A new type of discrete self-decomposability and its application to continuous-time Markov processes for modeling count data time series. *Stoch. Models* **19** 235–254.