

# $A_\infty$ -algebra Structure in Cohomology and its Applications

Tornike Kadeishvili

A. Razmadze Mathematical Institute of Tbilisi State University

LECTURE NOTES

Algebra, Topology and Analysis:  $C^*$  and  $A_\infty$ -Algebras

Summer School/Conference

Ivane Javakhishvili Tbilisi State University Press, 2023

The main method of algebraic topology is to assign to a topological space a certain algebraic object (model) and to study this relatively simple algebraic object instead of complex geometric one.

Examples of such models are chain and cochain complexes, homology and homotopy groups, cohomology algebras, etc.

The main problem here is to find models that classify spaces up to some equivalence relation, such as homeomorphism, homotopy equivalence, rational homotopy equivalence, etc.

Usually such models are not *complete*: the equivalence of models does not guarantee the equivalence of spaces. They can just distinguish spaces.

The models which carry richer algebraic structure contain more information about the space. For example the model "cohomology algebra" allows to distinguish spaces which cannot be distinguished by the model "cohomology groups".

Here we are going to present one more additional algebraic structure on cohomology, which was constructed in [16], [17], namely, we show that on cohomology  $H^*(X, R)$  there exists Stasheff's  $A_\infty$ -algebra structure (minimality

theorem). This structure consists of a collection of operations

$$\{m_i : H^*(X, R) \otimes \dots (i \text{ times}) \dots \otimes H^*(X, R) \rightarrow H^*(X, R), i = 1, 2, 3, \dots\}.$$

In fact this structure extends the usual structure of the cohomology algebra: the first operation  $m_1 : H^*(X, R) \rightarrow H^*(X, R)$ , the differential, is trivial (minimality) and  $m_2 : H^*(X, R) \otimes H^*(X, R) \rightarrow H^*(X, R)$  coincides with the cohomology multiplication.

Stasheff's  $A_\infty$  algebra is a sort of Strong Homotopy Associative Algebra, the operation  $m_3$  is a homotopy which measures the nonassociativity of the product  $m_2$ . So the existence of an  $A_\infty$  algebra structure on a *strictly associative* cohomology algebra  $H^*(X, R)$  looks a bit strange, but although the product on  $H^*(X, R)$  is associative, there appears a structure of a (generally nondegenerate) *minimal*  $A_\infty$ -algebra, which can be considered as an  $A_\infty$  *deformation* of the classical cohomology  $(H^*(X, R), \mu^*)$ , [26].

The cohomology algebra equipped with this additional structure

$$(H^*(X, R), \{m_i : H^*(X, R)^{\otimes i} \rightarrow H^*(X, R), i = 2, 3, \dots\}, m_2 = \mu^*),$$

which we call cohomology  $A_\infty$ -algebra, carries more information about the space than the cohomology algebra. For example just the cohomology algebra  $H^*(X, R)$  does not determine the cohomology of the loop space  $H^*(\Omega X, R)$ , but the cohomology  $A_\infty$ -algebra  $(H^*(X, R), \{m_i\})$  does. Dually, the Pontrjagin ring  $H_*(G)$  does not determine the homology  $H_*(B_G)$  of the classifying space, but the homology  $A_\infty$ -algebra  $(H_*(G), \{m_i\})$  does.

These  $A_\infty$ -algebras have several applications in the cohomology theory of fibre bundles too, see [17].

But this invariant also is not complete. One cannot expect the existence of a more or less simple complete algebraic invariant in general, but for the rational homotopy category there are various complete homotopy invariants (algebraic models):

- (i) The model of Quillen  $L_X$ , which is a differential graded Lie algebra;
- (ii) The minimal model of Sullivan  $M_X$ , which is a commutative graded differential algebra;
- (iii) The filtered model of Halperin and Stasheff  $\Lambda X$ , which is a filtered commutative graded differential algebra.

The rational cohomology algebra  $H^*(X, Q)$  is not a complete invariant even for rational spaces: two spaces might have isomorphic cohomology algebras, but different rational homotopy types. Bellow we also present the main

result of [21]: There is the notion of  $C_\infty$ -algebra which is the commutative version of Stasheff's notion of  $A_\infty$ -algebra, and in [21] we have shown that in the rational case on cohomology  $H^*(X, Q)$  arises a structure of  $C_\infty$ -algebra  $(H^*(X, Q), \{m_i\})$ . The main application of this structure is the following: it completely determines the rational homotopy type, that is, 1-connected spaces  $X$  and  $X'$  have the same rational homotopy type if and only if their cohomology  $C_\infty$ -algebras  $(H^*(X, Q), \{m_i\})$  and  $(H^*(X', Q), \{m'_i\})$  are isomorphic.

We present also several applications of this complete rational homotopy invariant to some problems of rational homotopy theory.

The  $C_\infty$ -algebra structure in cohomology and the applications of this structure in rational homotopy theory were already presented in the hardly available small book [23] (see also the preprint [22]).

Applications of cohomology  $C_\infty$ -algebra in rational homotopy theory are inspired by the existence of Sullivan's commutative cochains  $A(X)$  in this case. The cohomology  $C_\infty$ -algebra  $(H^*(X, Q), \{m_i\})$  carries the same amount of information as  $A(X)$  does. Actually these two objects are equivalent in the category of  $C_\infty$ -algebras.

We want to remark that for simplicity in these lectures signs are ignored. Of course they can be reconstructed using the Koszul sign rule.

The organization is as follows.

In Section 1 the notions of chain and cochain complexes are presented. In Section 2 the differential algebras and coalgebras are defined. In Section 3 the bar and cobar constructions are introduced. Twisting cochains and Berikashvili's functor  $D$  are presented in Section 4. In Section 5 the Stasheff's  $A_\infty$ -algebras are discussed. Hochschild cochains which are used for description of  $A_\infty$ -algebras are presented in Section 6. Section 7 is dedicated to our central topic, the Minimality Theorem. In the next Section 8 its applications are given. And the last Section 9 is dedicated to applications of the cohomology  $C_\infty$  algebra in rational homotopy theory.

## 1 Differential Graded Modules

### 1.1 Chain and Cochain Complexes

#### 1.1.1 Graded Modules

We work over a commutative associative ring with unit  $R$ .

A *graded module* is a collection of  $R$ -modules

$$M_* = \{ \dots, M_{-1}, M_0, M_1, \dots, M_n, M_{n+1}, \dots \} .$$

A *morphism of graded modules*  $M_* \rightarrow M'_*$  is a collection of homomorphisms  $\{f_i : M_i \rightarrow M'_i, i \in Z\}$ .

Sometimes we use the following notion: a *morphism of graded modules of degree  $n$*  is a collection of homomorphisms  $\{f_i : M_i \rightarrow M'_{i+n}, i \in Z\}$ . So a morphism of graded modules has the degree 0.

### 1.1.2 Chain Complexes

**Definition 1** . A *differential graded (dg) module (or a chain complex)* is a sequence of  $R$  modules and homomorphisms

$$\dots \xleftarrow{d_{-1}} C_{-1} \xleftarrow{d_0} C_0 \xleftarrow{d_1} C_1 \xleftarrow{d_2} \dots \xleftarrow{d_{n-1}} C_{n-1} \xleftarrow{d_n} C_n \xleftarrow{d_{n+1}} C_{n+1} \xleftarrow{d_{n+2}} \dots$$

such that  $d_i d_{i+1} = 0$ .

Elements of  $C_n$  are called  *$n$ -dimensional chains*; the homomorphisms  $d_i$  are called *boundary operators*, or *differentials*; elements of  $Z_n = \text{Ker } d_n \subset C_n$  are called  *$n$ -dimensional cycles* and elements of  $B_n = \text{Im } d_{n+1} \subset C_n$  are called  *$n$ -dimensional boundaries*.

It follows from the condition  $d_i d_{i+1} = 0$  that  $B_n \subset Z_n$ .

**Definition 2** . The  $n$ -th homology module  $H_n(C_*)$  of a dg module  $(C_*, d_*)$  is defined as the quotient  $Z_n/B_n$ .

A sequence  $C_{n-1} \xleftarrow{d_n} C_n \xleftarrow{d_{n+1}} C_{n+1}$  is *exact*, that is,  $B_n = Z_n$ , iff  $H_n(C_*) = 0$ . Thus homology measures the deviation from exactness.

### 1.1.3 Cochain Complexes

The notion of *cochain complex* differs from the notion of chain complex by the direction of the differential

$$\dots \xrightarrow{d^{-1}} C^{-1} \xrightarrow{d^0} C^0 \xrightarrow{d^1} C^1 \xrightarrow{d^2} \dots \xrightarrow{d^{n-1}} C^{n-1} \xrightarrow{d^n} C^n \xrightarrow{d^{n+1}} C^{n+1} \xrightarrow{d^{n+2}} \dots .$$

Corresponding terms here are cochains, cocycles  $Z^n = \text{Ker } d_{n+1} \subset C^n$ , coboundaries  $B^n = \text{Im } d^n \subset C^n$ , cohomology  $H^n(C^*) = Z^n/B^n$ .

Changing indices  $C^n = C_{-n}$ ,  $d^n = d_{-n}$  we convert a chain complex  $(C_*, d_*)$  to a cochain complex  $(C^*, d^*)$ .

### 1.1.4 Dual Cochain Complex

For a chain complex  $(C_*, d_*)$  and an  $R$ -module  $A$  the *dual cochain complex*  $C^* = (\text{Hom}(C_*, A), \delta^*)$  is defined as

$$C^n = \text{Hom}(C_n, A), \quad \delta^*(\phi) = \phi d.$$

### 1.1.5 Chain maps

**Definition 3** . A chain map of chain complexes  $f : (C_*, d_*) \rightarrow (C'_*, d'_*)$  is defined as a sequence of homomorphisms  $\{f_i : C_i \rightarrow C'_i\}$  such that  $d'_n f_n = f_{n-1} d_n$ .

This condition means the commutativity of the diagram

$$\begin{array}{cccccccc} \dots & \xleftarrow{d_{n-1}} & C_{n-1} & \xleftarrow{d_n} & C_n & \xleftarrow{d_{n+1}} & C_{n+1} & \xleftarrow{d_{n+2}} & \dots \\ & & \downarrow f_{n-1} & & \downarrow f_n & & \downarrow f_{n+1} & & \\ \dots & \xleftarrow{d'_{n-1}} & C'_{n-1} & \xleftarrow{d'_n} & C'_n & \xleftarrow{d'_{n+1}} & C'_{n+1} & \xleftarrow{d'_{n+2}} & \dots \end{array}$$

**Proposition 1** . The composition of chain maps is a chain map.

Chain complexes and chain maps form a category, which we denote by  $DGMod$ .

**Proposition 2** . If  $\{f_i\} : (C_*, d_*) \rightarrow (C'_*, d'_*)$  is a chain map, then  $f_n$  sends cycles to cycles and boundaries to boundaries, i.e.,  $f_n(Z_n) \subset Z'_n$  and  $f_n(B_n) \subset B'_n$ .

**Proposition 3** . A chain map  $\{f_i\} : (C_*, d_*) \rightarrow (C'_*, d'_*)$  induces the well defined homomorphism of homology groups

$$f_n^* : H_n(C_*) \rightarrow H_n(C'_*).$$

Homology is a functor from the category of  $dg$  modules to the category of graded modules

$$H : DGMod \rightarrow GMod.$$

### 1.1.6 Hom Complex

For two chain complexes  $C, C'$  define the chain complex  $(\text{Hom}(C, C'), D)$  as

$$\text{Hom}(C, C')_n = \text{Hom}^n(C, C')$$

where  $\text{Hom}^m(C, C') = \{\phi : C_* \rightarrow C'_{*+m}\}$  is the module of homomorphisms of degree  $m$ , and the differential  $D : \text{Hom}^n(C, C') \rightarrow \text{Hom}^{n-1}(C, C')$  is given by  $D(\phi) = d'\phi + (-1)^{\text{deg}} \phi d$ .

### 1.1.7 Tensor Product

For two chain complexes  $A$  and  $B$  the tensor product  $A \otimes B$  is defined as the following chain complex:

$$(A \otimes B)_n = \sum_{p+q=n} A_p \otimes B_q,$$

with differential  $d_{\otimes} : (A \otimes B)_n \rightarrow (A \otimes B)_{n-1}$  given by

$$d_{\otimes}(a_p \otimes b_q) = d_p(a_p) \otimes b_q + (-1)^p a_p \otimes d'_q(b_q).$$

If  $f : A \rightarrow A'$  and  $g : B \rightarrow B'$  are chain maps then there is a chain map

$$f \otimes g : A \otimes B \rightarrow A' \otimes B'$$

defined as  $(f \otimes g)(a \otimes b) = f(a) \otimes g(b)$ .

### 1.1.8 Chain Homotopy

**Definition 4** . Two chain maps  $\{f_i\}, \{g_i\} : (C_*, d_*) \rightarrow (C'_*, d'_*)$  are called chain homotopic, if there exists a sequence of homomorphisms  $D_n : C_n \rightarrow C'_{n+1}$ ,

$$\begin{array}{ccccccccccc} \dots & \xleftarrow{d_{n-1}} & C_{n-1} & & \xleftarrow{d_n} & C_n & & \xleftarrow{d_{n+1}} & C_{n+1} & & \xleftarrow{d_{n+2}} & \dots \\ & & f_{n-1} \downarrow \downarrow g_{n-1} & \searrow & D_{n-1} & f_n \downarrow \downarrow g_n & \searrow & D_n & f_{n+1} \downarrow \downarrow g_{n+1} & & & \\ \dots & \xleftarrow{d'_{n-1}} & C'_{n-1} & & \xleftarrow{d'_n} & C'_n & & \xleftarrow{d'_{n+1}} & C'_{n+1} & & \xleftarrow{d'_{n+2}} & \dots \end{array}$$

such that  $f_n - g_n = d'_{n+1}D_n + D_{n-1}d_n$ . In this case we write  $f \sim_D g$ .

**Proposition 4** . *Chain homotopy is an equivalence relation:*

- (a)  $f \sim_0 f$ ;
- (b)  $f \sim_D g \implies g \sim_{-D} f$ ;
- (c)  $f \sim_D g, g \sim_{D'} h \implies f \sim_{D+D'} h$ .

**Proposition 5** . *Chain homotopy is compatible with compositions:*

- (a)  $f \sim_D g \implies hf \sim_{hD} hg$ ;
- (b)  $f \sim_D g \implies fk \sim_{Dk} gk$ .

Thus there is a category  $hoDGM od$  whose objects are chain complexes and morphisms are chain homotopy classes

$$Hom_{hoDGM od}(C, C') = [C, C'] = Hom_{DGM od}(C, C') / \sim .$$

**Proposition 6** . *If two chain maps  $\{f_i\}, \{g_i\} : (C_*, d_*) \rightarrow (C'_*, d'_*)$  are chain homotopic, then the induced homomorphisms of homology groups coincide:  $f_n^* = g_n^* : H_n(C_*) \rightarrow H_n(C'_*)$ .*

*Thus we have the commutative diagram of functors*

$$\begin{array}{ccc} DGM od & \longrightarrow & hoDGM od \\ H \searrow & & \swarrow H \\ & GMod. & \end{array}$$

### 1.1.9 Chain Equivalence

Chain complexes  $C$  and  $C'$  are called *chain equivalent*  $C \sim C'$ , if there exist chain maps

$$f : C \xleftarrow{\sim} C' : g$$

such that  $gf \sim id_C, fg \sim id_{C'}$ . This means that  $C$  and  $C'$  are isomorphic in  $hoDGM od$ .

A chain complex  $C$  is called *contractible* if  $C \sim 0$ , equivalently if  $id_C \sim 0 : C \rightarrow C$ .

**Proposition 7** . *Each contractible  $C$  is acyclic, i.e.,  $H_i(C) = 0$  for all  $i$ .*

**Proposition 8** . *If all  $C_i$  are free and  $C$  is acyclic, then  $C$  is contractible.*

## 1.2 Algebraic and Topological Examples

### 1.2.1 Algebraic Example

Let  $(A, \mu : A \otimes A \rightarrow A)$  be an associative algebra, then

$$C(A) = (A \xleftarrow{\mu} A \otimes A \xleftarrow{\mu \otimes id - id \otimes \mu} A \otimes A \otimes A \xleftarrow{\mu \otimes id \otimes id - id \otimes \mu \otimes id + id \otimes id \otimes \mu} \dots)$$

is a chain complex: the associativity condition guarantees that  $dd = 0$ .

If  $A$  has a unit  $e \in A$  then this complex is contractible, that is  $id : C(A) \rightarrow C(A)$  and  $0 : C(A) \rightarrow C(A)$  are homotopic: the suitable chain homotopy is given by  $D(a_1 \otimes \dots \otimes a_n) = (e \otimes a_1 \otimes \dots \otimes a_n)$ . This immediately implies that  $C(A)$  is acyclic, that is  $H_i(C(A)) = 0$  for all  $i > 0$ .

This example is a particular case of more general chain complex called the bar construction, see later.

### 1.2.2 Simplicial Complexes

Simplicial complex is a formal construction, which models topological spaces.

**Definition 5** . A simplicial complex is a set  $V$  with a given family of finite subsets, called simplices, so that the following conditions are satisfied:

- (1) all points (called vertices) of  $V$  are simplices;
- (2) any nonempty subset of a simplex is a simplex.

A simplex consisting of  $(n + 1)$  points is called  $n$ -dimensional simplex. The 0-dimensional simplices, i.e., the points of  $V$  are called vertices.

**Definition 6** . A simplicial map of simplicial complexes  $V \rightarrow V'$  is a map of vertices  $f : V \rightarrow V'$  such that the image of any simplex of  $V$  is a simplex in  $V'$ .

**Proposition 9** . The composition of simplicial maps is a simplicial map.

Simplicial complexes and simplicial maps form a category which we denote as  $SC$ .

**Proposition 10** . To any simplicial set  $V$  corresponds a topological space  $|V|$  (called its realization) and to any simplicial map  $f : V \rightarrow V'$  corresponds a continuous map of realizations  $|f| : |V| \rightarrow |V'|$ .

So the realization is a functor from the category of simplicial complexes to the category of topological spaces

$$|-| : SC \rightarrow Top.$$

### 1.2.3 Homology Modules of a Simplicial Complex

In this section we consider *ordered* simplicial complexes: we assume that the set of vertices  $V$  is ordered by a certain order.

We assign to such an ordered simplicial complex the following chain complex  $(C_*(V), d_*)$ : Let  $C_n(V)$  be the free  $R$ -module, generated by all ordered  $n$ -simplices  $\sigma_n = (v_{k_0}, v_{k_1}, \dots, v_{k_n})$ , where  $v_{k_0} < v_{k_1} < \dots < v_{k_n}$ ; the differential  $d_n : C_n(V) \rightarrow C_{n-1}(V)$  on a generator  $\sigma_n = (v_{k_0}, v_{k_1}, \dots, v_{k_n}) \in C_n(V)$  is given by

$$d_n(v_{k_0}, v_{k_1}, \dots, v_{k_n}) = \sum_{i=0}^n (-1)^i (v_{k_0}, \dots, \widehat{v_{k_i}}, \dots, v_{k_n}),$$

where  $(v_{k_0}, \dots, \widehat{v_{k_i}}, \dots, v_{k_n})$  is the  $(n-1)$ -simplex obtained by omitting  $v_{k_i}$ , and is extended on the whole  $C_n(V)$  linearly.

**Proposition 11** . *The composition  $d_{n-1}d_n$  is zero, thus  $(C_*(V), d_*)$  is a chain complex.*

**Definition 7** . *The  $n$ -th homology group  $H_n(V)$  of an ordered simplicial set  $V$  is defined as the  $n$ -th homology group  $H_n(C_*(V))$ .*

### 1.2.4 Cohomology Modules of a Simplicial Complex

Let  $A$  be an  $R$ -module. The cochain complex of  $V$  with coefficients in  $A$  is defined as the dual to the chain complex  $C_*(V)$ :  $C^*(V, A) = \text{Hom}(C_*(V), A)$ . The  $n$ -th cohomology module of  $V$  with coefficients in  $A$  is just the  $n$ -th homology of this cochain complex.

Below we show that the cohomology  $H^*(V, A)$  is more interesting than the homology  $H_*(V)$ , since cohomology possesses richer algebraic structure: it is a ring.

## 2 Differential Graded Algebras and Coalgebras

### 2.1 Differential Graded Algebras

#### 2.1.1 Graded algebras

A *graded algebra* is a graded module

$$A_* = \{ \dots, A_{-1}, A_0, A_1, \dots, A_n, A_{n+1}, \dots \}$$

equipped with an associative multiplication  $\mu : A_* \otimes A_* \rightarrow A_*$  of degree 0, i.e.,  $\mu(\mu \otimes id) = \mu(id \otimes \mu)$  and  $\mu : A_p \otimes A_q \rightarrow A_{p+q}$ . We denote  $a \cdot b = \mu(a \otimes b)$ .

For a graded algebra  $A_*$  the component  $A_0$  is an associative algebra.

A *morphism of graded algebras*  $f : (A, \mu) \rightarrow (A', \mu')$  is a morphism of graded modules  $\{f_k : A_k \rightarrow A'_k\}$  which is multiplicative, that is,  $f\mu = \mu'(f \otimes f)$ , i.e.,  $f(a \cdot b) = f(a) \cdot f(b)$ .

Let  $f, g : (A, \mu) \rightarrow (A', \mu')$  be two morphisms of graded algebras. An  $(f, g)$ -*derivation* of degree  $k$  is defined as a morphism of graded modules of degree  $k$   $D : A_* \rightarrow A'_{*+k}$ , i.e., a collection of homomorphisms  $\{D_i : A_i \rightarrow A'_{i+k}, i \in \mathbb{Z}\}$  which satisfies the condition

$$D(a \cdot b) = D(a) \cdot g(b) + (-1)^{k \cdot |a|} f(a) \cdot D(b).$$

An essential particular case of this notion is a  $k$ -*derivation*  $D : (A, \mu) \rightarrow (A, \mu)$  which is an  $(id, id)$   $k$ -*derivation*, i.e., it satisfies the condition  $D(a \cdot b) = D(a) \cdot b + a \cdot D(b)$ .

It is easy to see that if  $D, D' : A_* \rightarrow A'_{*+k}$  are two  $(f, g)$  derivation, then their sum  $D + D'$  is also an  $(f, g)$  derivation.

Moreover, for graded algebra morphisms  $h : B \rightarrow A$ ,  $l : A' \rightarrow C$  and an  $(f, g)$  derivations  $D : A \rightarrow A'$  the composition  $Dh$  is an  $(fh, gh)$  derivation and  $lD$  is an  $(lf, lg)$ -derivation.

#### 2.1.2 Differential graded algebras

**Definition 8** . A *differential graded algebra* (*dga in short*)  $(A, d, \mu)$  is a *dg module*  $(A, d)$  equipped additionally with a *multiplication*

$$\mu : A \otimes A \rightarrow A$$

so that  $(A, \mu)$  is a graded algebra, and the multiplication  $\mu$  is a chain map, that is, the differential  $d$  and  $\mu$  are connected by the condition

$$d(a \cdot b) = da \cdot b + (-1)^{|a|} a \cdot db.$$

This condition means simultaneously that  $\mu$  is a chain map, and that  $d$  is an  $(id, id)$ -derivation of degree  $-1$ .

A morphism of dga-s  $f : (A, d, \mu) \rightarrow (A, d, \mu)$  is defined as a multiplicative chain map:

$$df = fd, \quad f(a \cdot b) = f(a) \cdot f(b).$$

We denote the obtained category as  $DGAlg$ .

For a dg algebra  $(A, d, \mu)$  its homology  $H_*(A)$  is a graded algebra with the following multiplication:

$$H_*(A) \otimes H_*(A) \xrightarrow{\phi} H_*(A \otimes A) \xrightarrow{H_*(\mu)} H_*(A),$$

where  $\phi : H_*(A) \otimes H_*(A) \rightarrow H_*(A \otimes A)$  is the standard map

$$\phi(h_1 \otimes h_2) = cl(z_{h_1} \otimes z_{h_2}).$$

In other words the multiplication on  $H(A)$  is defined as follows: For  $h_1, h_2 \in H(A)$  the product  $h_1 \cdot h_2$  is the homology class of the cycle  $z_{h_1} \cdot z_{h_2}$ .

Furthermore, a dga map induces a multiplicative map of homology graded algebras.

Thus homology is a functor from the category of dg algebras to the category of graded algebras.

### 2.1.3 Derivation homotopy

Two dg algebra maps  $f, g : A \rightarrow A'$  are called homotopic if there exists a chain homotopy  $D : A \rightarrow A'$ ,  $f - g = dD + Dd$  which, in addition is a  $(f, g)$ -derivation, that is

$$D(a \cdot b) = D(a) \cdot g(b) + (-1)^{|a|} f(a) \cdot D(b).$$

Note that generally this is not an equivalence relation.

## 2.2 Differential Graded Colgebras

### 2.2.1 Graded coalgebras

A graded coalgebra  $(C, \Delta)$  is a graded module

$$C = \{ \dots, C_{-1}, C_0, C_1, \dots, C_n, C_{n+1}, \dots \}$$

equipped with a comultiplication

$$\Delta : C \otimes C \rightarrow C$$

which is coassociative, that is  $(\Delta \otimes id)\Delta = (id \otimes \Delta)\Delta$ , i.e., the following diagram commutes:

$$\begin{array}{ccccc} & C & \xrightarrow{\Delta} & C \otimes C & \\ \Delta & \downarrow & & \downarrow & \Delta \otimes id \\ & C \otimes C & \xrightarrow{id \otimes \Delta} & C \otimes C \otimes C & . \end{array}$$

A morphism of graded coalgebras  $f : (C, \Delta) \rightarrow (C', \Delta')$  is a morphism of graded modules  $\{f_k : C_k \rightarrow C'_k\}$  which is comultiplicative, that is  $\Delta' f = (f \otimes f)\Delta$ , i.e., the following diagram commutes:

$$\begin{array}{ccccc} & C & \xrightarrow{f} & C' & \\ \Delta & \downarrow & & \downarrow & \Delta' \\ & C \otimes C & \xrightarrow{f \otimes f} & C' \otimes C' & . \end{array}$$

If  $f, g : C \rightarrow C'$  are two morphisms of graded coalgebras, then an  $(f, g)$ -coderivation of degree  $k$  is defined as a collection of homomorphisms  $\{D_i : C_i \rightarrow C_{i+k}\}$  which satisfies the condition  $\Delta' D = (f \otimes D + D \otimes g)\Delta$ , i.e., the following diagram commutes:

$$\begin{array}{ccccc} & C & \xrightarrow{D} & C' & \\ \Delta & \downarrow & & \downarrow & \Delta' \\ & C \otimes C & \xrightarrow{f \otimes D + D \otimes g} & C' \otimes C' & . \end{array}$$

An essential particular case of this notion is that of a  $k$ -coderivation  $D : (C, \Delta) \rightarrow (C, \Delta)$ , which is an  $(id, id)$ - $k$ -coderivation, i.e., it satisfies the condition  $\Delta D = (id \otimes D + D \otimes id)\Delta$ .

It is easy to see that if  $D, D' : C_* \rightarrow C'_{*+k}$  are two  $(f, g)$ -coderivations, then their sum  $D + D'$  is also an  $(f, g)$ -coderivation.

Moreover, for graded coalgebra morphisms  $h : A \rightarrow C$ ,  $l : C' \rightarrow B$  and an  $(f, g)$ -coderivation  $D : C \rightarrow C'$  the composition  $Dh$  is an  $(fh, gh)$ -coderivation and  $lD$  is an  $(lf, lg)$  coderivation.

### 2.2.2 Differential Graded Coalgebras

**Definition 9** . A differential graded coalgebra (dgc in short)  $(C, d, \Delta)$  is a dg module  $(C, d)$  equipped additionally with a comultiplication  $\Delta : C \rightarrow C \otimes C$  so that  $(C, \Delta)$  is a graded coalgebra and the comultiplication  $\Delta$  and the differential  $d$  are related by the condition

$$\Delta d = (d \otimes id + id \otimes d)\Delta.$$

This condition means simultaneously that  $\Delta$  is a chain map, and that  $d$  is a  $(id, id)$ -coderivation of degree  $-1$ .

A morphism of dgc-s  $f : (C, d, \Delta) \rightarrow (C', d', \Delta')$  is defined as a morphism of graded coalgebras which is a chain map.

We denote the obtained category as  $DGC\text{alg}$ .

Generally for a dg coalgebra  $(C, d, \mu)$  its homology  $H_*(C)$  is not a graded coalgebra:

$$H_*(C) \otimes H_*(C) \xrightarrow{\phi} H_*(C \otimes C) \xleftarrow{H(\Delta)} H_*(C),$$

the map  $\phi : H_*(C) \otimes H_*(C) \rightarrow H_*(C \otimes C)$  has the wrong direction, but if all  $H_i(C)$  are free then  $\phi$  is invertible and  $H_*(C)$  is a graded coalgebra.

### 2.2.3 Coderivation homotopy

In the category of dg coalgebras there is the following notion of homotopy: two dg coalgebra maps  $f, g : (C, d_C, \Delta_C) \rightarrow (C', d_{C'}, \Delta_{C'})$  are homotopic, if there exists  $D : C \rightarrow C'$  of degree  $+1$  such that  $d_{C'}D + Dd_C = f - g$ , i.e., the chain maps  $f$  and  $g$  are chain homotopic, and additionally the homotopy  $D$  is a  $f - g$ -coderivation, that is  $\Delta_{C'}D = (f \otimes D + D \otimes g)\Delta_C$ .

### 2.2.4 Dual cochain algebra

Let  $(C_*, d_C, \Delta_C)$  be a dg coalgebra and let  $(A, \mu_A : A \otimes A \rightarrow A)$  be a (nondifferential and nongraded) algebra. Then the dual cochain complex  $(C^* = Hom(C_*, A), \delta^*)$ ,  $\delta^*(\phi) = \phi d_C$ , becomes a dg algebra (of cochain type, i.e., the degree of the differential is  $+1$ ) with the multiplication (cup product)

$$\mu^* : Hom(C_*, A) \otimes Hom(C_*, A) \rightarrow Hom(C_*, A)$$

given by

$$\phi \smile \psi = \mu^*(\phi \otimes \psi) = \mu_A(\phi \otimes \psi)\Delta_C.$$

The obtained object  $(C^* = \text{Hom}(C_*, A), \delta^*(\phi) = \phi d_C, \smile)$  is called the cochain dg algebra of the chain dg coalgebra  $(C_*, d_C, \Delta_C)$  with coefficients in  $A$ .

## 2.3 Applications in Topology

### 2.3.1 Alexander-Whitney diagonal

Let  $V$  be an ordered simplicial complex (1.2.3) and let  $(C_*(V), d_*)$  be its chain complex. There exists a comultiplication

$$\Delta : C_*(V) \rightarrow C_*(V) \otimes C_*(V),$$

the so called Alexander-Whitney diagonal, which turns  $(C_*(V), d_*, \Delta)$  into a dg coalgebra. This diagonal is defined by

$$\Delta(v_{i_0}, \dots, v_{i_n}) = \sum_{k=0}^n (v_{i_0}, \dots, v_{i_k}) \otimes (v_{i_k}, \dots, v_{i_n}).$$

### 2.3.2 Cohomology algebra

Let again  $(A, \mu_A : A \otimes A \rightarrow A)$  be a (nondifferential and nongraded) associative algebra. The Alexander-Whitney diagonal of  $C_*(V)$  induces on the dual cochain complex  $C^*(V, A) = \text{Hom}(C_*(V), A), \delta^*$  the cup product

$$\smile : C^*(V) \otimes C^*(V) \rightarrow C^*(V)$$

which for  $\phi \in C^p(V, A), \psi \in C^q(V, A)$  looks as

$$\phi \smile \psi(v_{i_0}, \dots, v_{i_{p+q}}) = \phi(v_{i_0}, \dots, v_{i_p}) \cdot \psi(v_{i_p}, \dots, v_{i_{p+q}}).$$

This turns  $(C^*(V, A), \delta^*, \smile)$  into a dg algebra.

This structure induces on the cohomology  $H^*(V, A)$  a structure of graded algebra.

The cohomology algebra  $H^*(V, A)$  is a more powerful invariant than the cohomology groups: the two spaces  $X = S^1$  times  $S^1$  and  $Y = S^1 \vee S^1 \vee S^2$  have the same cohomology groups

$$H^0 = R, \quad H^1 = R \cdot a \oplus R \cdot b, \quad H^2 = R \cdot c,$$

with generators  $a, b$  in dimension 1 and  $c$  in dimension 2, but they have different cohomology algebras, namely  $a \cdot b = 0$  in  $H^*(Y)$  and  $a \cdot b = c$  in  $H^*(X)$ .

### 3 Bar and Cobar Functors

Here we describe two classical adjoint functors

$$B : DGA\!l\!g \overleftarrow{\hspace{1cm}} DGCoalg : \Omega,$$

the bar functor  $B : DGA\!l\!g \rightarrow DGCoalg$  from the category of dg algebras to the category of dg coalgebras, and the cobar functor  $\Omega : DGCoalg \rightarrow DGA\!l\!g$  in the opposite direction.

We start with the definitions of free (in the category of graded algebras) and cofree (in the category of graded coalgebras) objects.

#### 3.1 Tensor Algebra and Tensor Coalgebra

##### 3.1.1 Tensor Algebra

Let  $V = \{V_i\}$  be a graded  $R$ -module. The tensor algebra generated by  $V$  is defined as

$$T(V) = R \oplus V \oplus (V \otimes V) \oplus (V \otimes V \otimes V) \oplus \dots = \sum_{i=0}^{\infty} V^{\otimes i}$$

with grading  $\dim(a_1 \otimes \dots \otimes a_m) = \dim a_1 + \dots + \dim a_m$ , and with *multiplication*  $\mu : T(V) \otimes T(V) \rightarrow T(V)$  given by

$$\mu((a_1 \otimes \dots \otimes a_m) \otimes (a_{m+1} \otimes \dots \otimes a_{m+n})) = a_1 \otimes \dots \otimes a_{m+n}.$$

The unit element for this multiplication is  $1 \in R = V^{\otimes 0}$ .

By  $i_k$  we denote the obvious inclusion  $i_k : V^{\otimes k} \rightarrow T(V)$ .

##### 3.1.2 Universal Property of $T(V)$

The tensor algebra  $T(V)$  is the free object in the category of graded algebras: for an arbitrary graded algebra  $(A, \mu_A)$  and a map of graded modules  $\alpha : V \rightarrow A$  there exists a unique morphism of graded algebras  $f_\alpha : T(V) \rightarrow A$  such that  $f_\alpha(v) = \alpha(v)$  (i.e.,  $f_\alpha i_1 = \alpha$ ).

This morphism  $f_\alpha$  (which is called *multiplicative extension* of  $\alpha$ ) is defined as  $f_\alpha(a_1 \otimes \dots \otimes a_m) = \alpha(a_1) \cdot \dots \cdot \alpha(a_m)$ . Or, equivalently  $f_\alpha$  is described by:

$$f_\alpha i_k = \sum_k \mu_A^k(\alpha \otimes \dots \otimes \alpha),$$

where  $\mu_A^k : A^{\otimes k} \rightarrow A$  is the  $k$ -fold iteration of the multiplication  $\mu_A : A \otimes A \rightarrow A$ , namely,  $\mu_A^1 = id$ ,  $\mu_A^2 = \mu_A$ ,  $\mu_A^k = \mu_A(\mu_A^{k-1} \otimes id)$ .

So, to summarize, we have the following universal property

$$\begin{array}{ccc} V & \xrightarrow{i_1} & T(V) \\ \alpha & \searrow & \downarrow f_\alpha = \sum_k \mu_A^k(\alpha \otimes \dots \otimes \alpha) \\ & & A. \end{array}$$

### 3.1.3 Universal Property for Derivations

The tensor algebra has an analogous universal property also for derivations: for a graded algebra  $(A, \mu)$ , two homomorphisms  $\alpha, \alpha' : V \rightarrow A$  of degree 0 and a homomorphism  $\beta : V \rightarrow A$  of degree  $k$  there exist: morphisms of graded algebras  $f_\alpha, f_{\alpha'} : T(V) \rightarrow A$  and a unique  $(f_\alpha, f_{\alpha'})$  derivation of degree  $k$

$$D_\beta : T(V) \rightarrow A$$

such that  $f_\alpha(v) = \alpha(v)$ ,  $f_{\alpha'}(v) = \alpha'(v)$  and  $D(v) = \beta(v)$ , i.e., the following diagram commutes.

$$\begin{array}{ccc} V & \xrightarrow{i_1} & T(V) \\ \beta & \searrow & \downarrow D \\ & & A. \end{array}$$

The derivation  $D$  is defined as

$$D(a_1 \otimes \dots \otimes a_n) = \sum_{k=1}^n \mu^n(\alpha(a_1) \otimes \dots \otimes \alpha(a_{k-1}) \otimes \beta(a_k) \otimes \alpha'(a_{k+1}) \otimes \dots \otimes \alpha'(a_n)).$$

Or, equivalently  $D \cdot i_n = \sum_k \mu^n(\alpha^{\otimes(k-1)} \otimes \beta \otimes \alpha'^{\otimes(n-k)})i_n$ .

### 3.1.4 Tensor Coalgebra

Here is the dualization of the previous notion.

Let  $V = \{V_i\}$  be a graded  $R$ -module. The *tensor coalgebra* cogenerated by  $V$  is defined (again) as

$$T^c(V) = R \oplus V \oplus V \otimes V \oplus V \otimes V \otimes V \oplus \dots = \sum_{i=0}^{\infty} V^{\otimes i}$$

with the same grading  $\dim(a_1 \otimes \dots \otimes a_m) = \dim a_1 + \dots + \dim a_m$ , but now with *comultiplication*  $\Delta : T^c(V) \rightarrow T^c(V) \otimes T^c(V)$  given by

$$\Delta(a_1 \otimes \dots \otimes a_n) = \sum_{i=0}^n (a_1 \otimes \dots \otimes a_i) \otimes (a_{i+1} \otimes \dots \otimes a_n),$$

here  $( ) = 1 \in R = V^{\otimes 0}$ .

By  $p_k$  we denote the obvious projection  $p_k : T^c(V) \rightarrow V^{\otimes k}$ .

### 3.1.5 Universal Property of $T^c(V)$

In order to formulate the about universal property in this case we have to introduce some dimensional restrictions in this case.

Let  $V = \{\dots, 0, 0, V_1, V_2, \dots\}$  be a *connected* graded module, that is,  $V_i = 0$  for  $i \leq 0$ .

The tensor coalgebra of such  $V$  is the cofree object in the category of *connected* graded coalgebras: for a map of graded modules  $\alpha : C \rightarrow V$  there exists a unique morphism of graded coalgebras  $f_\alpha : C \rightarrow T^c(V)$  such that  $p_1 f_\alpha = \alpha$ , i.e., the following diagram commutes

$$\begin{array}{ccc} V & \xleftarrow{p_1} & T^c(V) \\ \alpha \searrow & & \uparrow f_\alpha \\ & & C. \end{array}$$

The coalgebra map  $f_\alpha$  (which is called *comultiplicative coextension* of  $\alpha$ ) is defined as  $f_\alpha = \sum_k (\alpha \otimes \dots \otimes \alpha) \Delta^k$ , where  $\Delta^k : C \rightarrow C^{\otimes k}$  is the  $k$ -th iteration of the comultiplication  $\Delta : C \rightarrow C \otimes C$ , i.e.,  $\Delta^1 = id$ ,  $\Delta^2 = \Delta$ ,  $\Delta^k = (\Delta^{k-1} \otimes id) \Delta$ .

### 3.1.6 Universal Property for Coderivations

The tensor coalgebra has a similar universal property also for coderivations.

Namely, for a graded coalgebra  $(C, \Delta)$ , two homomorphisms of degree 0  $\alpha, \alpha' : C \rightarrow V$  and a homomorphism of degree  $k$   $\beta : C \rightarrow V$  there exist: morphisms of graded coalgebras  $f_\alpha, f_{\alpha'} : C \rightarrow T(V)$  and a unique  $(f_\alpha, f_{\alpha'})$ -coderivation of degree  $k$   $\partial_\beta : C \rightarrow T^c(V)$  such that  $p_1 \partial_\beta = \beta$ , i.e., commutes

the following diagram:

$$\begin{array}{ccc} V & \xleftarrow{p_1} & T^c(V) \\ \beta & \nwarrow & \uparrow \partial_\beta \\ & & C. \end{array}$$

The coderivation  $\partial_\beta$  is defined as

$$\partial_\beta = \sum_{n=0}^{\infty} \sum_{k=1}^n (\alpha^{\otimes(k-1)} \otimes \beta \otimes \alpha'^{(n-k)} \Delta^k).$$

### 3.2 Shuffle Comultiplication and Shuffle Comultiplication — Bialgebra Structures on $T(V)$ and $T'(V)$

In fact,  $T(V)$  and  $T^c(V)$  coincide as graded modules, but the multiplication of  $T(V)$  and the comultiplication of  $T^c(V)$  are not compatible with each other, so they do not define a graded bialgebra structure on the graded module  $T(V) = T^c(V)$ .

Nevertheless there exists the *shuffle* comultiplication

$$\nabla_{sh} : T(V) \rightarrow T(V) \otimes T(V)$$

introduced by Eilenberg and MacLane [?], which turns  $(T(V), \Delta_{sh}, \mu)$  into a graded bialgebra.

And dually, there exists the *shuffle* multiplication  $\mu_{sh} : T^c(V) \otimes T^c(V) \rightarrow T^c(V)$  which turns  $(T^c(V), \Delta, \mu_{sh})$  into a graded bialgebra.

#### 3.2.1 Shuffle Comultiplication – Bialgebra Structure on Tensor Algebra

There exists on this free graded algebra  $(T(V), \mu)$  a comultiplication  $\nabla_{sh} : T(V) \rightarrow T(V) \otimes T(V)$  which is a morphism of graded algebras, and, consequently turns  $(T(V), \mu, \nabla_{sh})$  into a graded bialgebra.

Namely, the *shuffle comultiplication*  $\nabla_{sh} : T(V) \rightarrow T(V) \otimes T(V)$  is a graded coalgebra map induced by the universal property of  $T(V)$  (3.1.2) by  $\alpha : T(V) \rightarrow T(V) \otimes T(V)$  given by  $\alpha(v) = v \otimes 1 + 1 \otimes v$ ,  $\alpha(1) = 1 \otimes 1$ .

This comultiplication is associative and in fact is given by

$$\begin{aligned} \nabla(v_1 \otimes \dots \otimes v_n) &= 1 \otimes (v_1 \otimes \dots \otimes v_n) + (v_1 \otimes \dots \otimes v_n) \otimes 1 + \\ &\quad \sum_p \sum_{\sigma \in sh(p, n-p)} (v_{\sigma(1)} \otimes \dots \otimes v_{\sigma(p)}) \otimes (v_{\sigma(p+1)} \otimes \dots \otimes v_{\sigma(n)}) \end{aligned}$$

where  $sh(p, n - p)$  consists of all  $(p, n - p)$ -shuffles, that is all permutations of  $1, \dots, n$  such that  $\sigma(1) < \dots < \sigma(p)$  and  $\sigma(p + 1) < \dots < \sigma(n)$ .

### 3.2.2 Shuffle Multiplication – Bialgebra Structure on Tensor Coalgebra

The *shuffle* multiplication  $\mu_{sh} : T^c(V) \otimes T^c(V) \rightarrow T^c(V)$ , introduced by Eilenberg and MacLane [7], turns  $(T^c(V), \Delta, \mu_{sh})$  into a graded bialgebra.

This multiplication is defined as the graded coalgebra map induced by the universal property of  $T^c(V)$  (3.1.5) by  $\alpha : T^c(V) \otimes T^c(V) \rightarrow V$  given by  $\alpha(v \otimes 1) = \alpha(1 \otimes v) = v$  and  $\alpha = 0$  otherwise. This multiplication is associative and in fact is given by

$$\mu_{sh}((v_1 \otimes \dots \otimes v_n) \otimes (v_{n+1} \otimes \dots \otimes v_{n+m})) = \sum \pm v_{\sigma(1)} \otimes \dots \otimes v_{\sigma(n+m)}, \quad (1)$$

where summation is taken over all  $(m, n)$ -shuffles, that is, over all permutations of the set  $(1, 2, \dots, n + m)$  which satisfy the condition:  $i < j$  if  $1 \leq \sigma(i) < \sigma(j) \leq n$  or  $n + 1 \leq \sigma(i) < \sigma(j) \leq n + m$ .

Particularly,

$$\begin{aligned} \mu_{sh}((v_1) \otimes (v_2)) &= v_1 \otimes v_2 \pm v_2 \otimes v_1, \\ \mu_{sh}((v_1) \otimes (v_2 \otimes v_3)) &= v_1 \otimes v_2 \otimes v_3 \pm v_2 \otimes v_1 \otimes v_3 \pm v_2 \otimes v_3 \otimes v_1, \\ &\dots \end{aligned}$$

## 3.3 Bar and Cobar Constructions

### 3.3.1 Cobar Construction

Let  $(C, d, \Delta)$  be a dg coalgebra with  $C_i = 0$ ,  $i \leq 1$  and let  $s^{-1}C$  be the desuspension of  $C$ , that is,  $(s^{-1}C)_k = C_{k+1}$ .

The cobar construction  $\Omega C$  is defined as the tensor algebra  $T(s^{-1}C)$ . We use the following notation for elements of this tensor coalgebra:

$$s^{-1}a_1 \otimes \dots \otimes s^{-1}a_n = [a_1, \dots, a_n].$$

So the dimension of  $[a_1, \dots, a_n]$  is  $\sum_i \dim a_i - n$ .

The differential  $d_\Omega : \Omega C \rightarrow \Omega C$  is defined as

$$\begin{aligned} d_\Omega[a_1, \dots, a_n] &= \\ \sum_i \pm [a_1, \dots, a_{i-1}, da_i, a_{i+1}, \dots, a_n] &+ \sum_i \pm [a_1, \dots, a_{i-1}, \Delta(a_i), a_{i+1}, \dots, a_n]. \end{aligned}$$

In fact  $d_\Omega = \partial_\beta$  where  $\partial_\beta$  is the derivation defined by the above universal property (3.1.3) for

$$\beta[a] = da + \Delta a.$$

Besides, the properties of  $d$  and  $\Delta$  from the definition of a dg coalgebra (2.2.2) guarantee that the restriction  $d_\Omega d_\Omega | V = d_\Omega d_\Omega i_1$  is 0 and this, by the universal property (3.1.3), implies  $d_\Omega d_\Omega = 0$ . Thus  $\Omega C \in DGAlg$ .

### 3.3.2 Bar Construction

Let  $(A, d, \mu = \cdot)$  be a dg algebra with  $A_i = 0$  for  $i \leq 1$  and let  $sA$  be the suspension of  $A$ , that is  $(sA)_k = A_{k-1}$ .

As a graded coalgebra the bar construction  $BA$  is defined as the tensor coalgebra:  $T^c(sA)$ . We use the following notation for elements of this tensor coalgebra

$$sa_1 \otimes \dots \otimes sa_n = [a_1, \dots, a_n].$$

So the dimension of  $[a_1, \dots, a_n]$  is  $\sum_i \dim a_i + n$ .

The differential  $d_B : BA \rightarrow BA$  is defined as

$$d_B[a_1, \dots, a_n] = \sum_i \pm [a_1, \dots, a_{i-1}, da_i, a_{i+1}, \dots, a_n] + \sum_i \pm [a_1, \dots, a_i \cdot a_{i+1}, \dots, a_n].$$

In fact,  $d_B = D_\beta$  where  $D_\beta$  is the coderivation defined by the above universal property (3.1.6) for

$$\beta[a_1, \dots, a_n] = \begin{cases} [da_1] & \text{for } n = 1; \\ [a_1 \cdot a_2] & \text{for } n = 2 \\ 0 & \text{for } n > 2. \end{cases}$$

Besides, the properties of  $d$  and  $\mu$  from the definition of a dg algebra (2.1.2) guarantee that the projection  $p_1 d_B d_B$  is 0 and this, by the universal property (3.1.6), implies  $d_B d_B = 0$ . Thus  $BA \in DGCoalg$ .

### 3.3.3 Bar Construction of a Commutative dg Algebra

Assume now that  $(A, d, \mu = \cdot)$  is a commutative dg algebra. How the commutativity reflects on the bar construction  $BA$ ?

By definition, the differential  $d_B : BA \rightarrow BA$  is a coderivation with respect to the standard graded coalgebra structure of the tensor coalgebra  $BA = T^c(sA)$ . As it was mentioned above in (3.2.2),  $BA$  carries also the

shuffle product  $\mu_{sh} : T^c(sV) \otimes T^c(sA) \rightarrow T^c(sA)$  which turns it into a graded bialgebra.

**Proposition 12** . *If a dg algebra  $(A, d, \mu = \cdot)$  is commutative, the differential  $d_\beta : BA \rightarrow BA$  is not only a coderivation with respect to multiplication  $\Delta$  but also a derivation with respect to the shuffle product  $\mu_{sh}$ . So in this case the bar construction  $(BA, \Delta, \mu_{sh})$  is a dg bialgebra.*

**Proof.** The map  $\Phi : BA \otimes BA \rightarrow BA$  defined as

$$\Phi = d_\beta \mu_{sh} - \mu_{sh}(d_\beta \otimes id + id \otimes d_\beta)$$

is a coderivation, see the arguments in (2.2.1). Thus, according to universal the property (3.1.6) of  $T^c(s^{-1}A)$ , the map  $\Phi$  is trivial if and only if  $p_1 \Phi = 0$ . This is equivalent to the commutativity of  $A$ .

### 3.3.4 Adjunction

Let  $(C, d, \Delta)$  be a dg coalgebra and  $(A, d, \mu)$  a dg algebra. A twisting cochain [5] is a homomorphism  $\tau : C \rightarrow A$  of degree +1 satisfying Browns' condition

$$d\tau + \tau d = \tau \smile \tau, \tag{2}$$

where  $\tau \smile \tau' = \mu_A(\tau \otimes \tau')\Delta$ . We denote by  $T(C, A)$  the set of all twisting cochains  $\tau : C \rightarrow A$ .

There are universal twisting cochains  $C \rightarrow \Omega C$  and  $BA \rightarrow A$ , namely, the obvious inclusion and projection, respectively. Here are essential consequences of the condition (3):

- (i) The multiplicative extension  $f_\tau : \Omega C \rightarrow A$  is a map of dg algebras, so there is a bijection  $T(C, A) \leftrightarrow Hom_{DG-Alg}(\Omega C, A)$ ;
- (ii) The comultiplicative coextension  $f_\tau : C \rightarrow BA$  is a map of dg coalgebras, so there is a bijection  $T(C, A) \leftrightarrow Hom_{DG-Coalg}(C, BA)$ .

Thus we have two bijections

$$Hom_{DG-Alg}(\Omega C, A) \leftrightarrow T(C, A) \leftrightarrow Hom_{DG-Coalg}(C, BA).$$

Besides, there are two weak equivalences (homology isomorphisms)

$$\alpha_A : \Omega B(A) \rightarrow A, \quad \beta_C : C \rightarrow B\Omega(C).$$

## 4 Twisting Cochains and Functor $D$

### 4.1 Brown's twisting cochains

#### 4.1.1 Definition of twisting cochain

Let  $(K, d_K : K_* \rightarrow K_{*-1}, \nabla_K : K \rightarrow K \otimes K)$  be a dg coalgebra and let  $(A, d_A : A_* \rightarrow A_{*-1}, \mu_A : A \otimes A \rightarrow A)$  be a dg algebra. Then  $C^*(K, A) = \text{Hom}(K, A)$  with differential  $\delta\alpha = \alpha d_K + d_A \alpha$  and multiplication  $\alpha \smile \beta = \mu_A(\alpha \otimes \beta) \nabla_K$  is a dg algebra.

**Definition 10** . *A Brown twisting cochain is a homomorphism*

$$\phi : K_* \rightarrow A_{*-1},$$

*i.e.,  $\text{deg}\phi = -1$ , satisfying*

$$d\phi + \phi d = \phi \smile \phi. \quad (3)$$

Brown's condition sometimes is called also *Maurer-Cartan equation*. It can be rewritten as  $d_A \phi + \phi d_K = \mu_A(\phi \otimes \phi) \nabla_K$ .

The set of all twisting cochains  $\phi : K \rightarrow A$  we denote as  $\text{Tw}(K, A)$ .

#### 4.1.2 Twisted tensor product

Let  $(M, d_M, \nu : A \otimes M \rightarrow M)$  be a dg  $A$ -module. Then any twisting cochain  $\phi : K \rightarrow A$  determines a homomorphism  $d_\phi : K \otimes M \rightarrow K \otimes M$  by

$$d_\phi(k \otimes m) = d_K k \otimes m + k \otimes d_M m + (k \otimes m) \cap \phi$$

where  $(k \otimes m) \cap \phi = (id_K \otimes \nu)(id_K \otimes \phi \otimes id_M)(\nabla_K \otimes id_M)(k \otimes m)$ . Brown's condition  $d\phi = \phi \smile \phi$  implies that  $d_\phi d_\phi = 0$ .

**Definition 11** . *The twisted tensor product  $(K \otimes_\phi M, d_\phi)$  is defined as the chain complex  $(K \otimes M, d_\phi)$ .*

#### 4.1.3 Application: model of fibration

Let  $(E, p, B, F, G)$  be a fibre bundle with base  $B$ , fibre  $F$ , structure group  $G$ . So  $K = C_*(B)$  is a dg coalgebra,  $A = C_*(G)$  is a dg algebra,  $M = C_*(F)$

is a dg module over  $A$ . Then, by Brown's theorem there exists a twisting cochain  $\phi : K = C_*(B) \rightarrow A = C_*(G)$  such that the twisted tensor product

$$(K \otimes_\phi M, d_\phi) = (C_*(B) \otimes_{\phi: C_*(B) \rightarrow C_{*-1}(G)} C_*(F), d_\phi)$$

gives the homology of fibre space  $H_*(E)$ . It is clear that  $\phi$  determines all differentials of the Serre spectral sequence also.

Brown's twisting cochain  $\phi$  is not determined uniquely, so it would be useful to have a possibility to choose one convenient for computations.

#### 4.1.4 Twisting cochains and the Bar and cobar constructions

By the universal property of the tensor coalgebra from (B1.3) any homomorphism  $\phi : K \rightarrow A$  of degree  $-1$  induces a graded coalgebra map  $f_\phi : K \rightarrow T^c(sA)$  by

$$f_\phi = \sum_i (\phi \otimes \dots \otimes \phi) \nabla_K^i.$$

If, in addition,  $\phi : K \rightarrow A$  is a twisting cochain, that is, it satisfies Brown's condition  $\delta\phi = \phi \smile \phi$ , then  $f_\phi : K \rightarrow T^c(sA)$  is a chain map, i.e., is a map of dg coalgebras: since the tensor coalgebra is cofree one has the equality  $d_B f_\phi = f_\phi d_K$  if and only if equal the projections  $pd_B f_\phi$  and  $= pf_\phi d_K$  are equal, and this is exactly Brown's condition  $\delta\phi = \phi \smile \phi$ .

Conversely, any dg coalgebra map  $f : K \rightarrow T^c(sA)$  is  $f_\phi$  for  $\phi = pf : K \rightarrow A$ . In fact we have a bijection  $Mor_{DGC_{\text{coal}}}(K, T^c(sA)) \leftrightarrow Tw(K, A)$ .

Dually, by the universal property of the tensor algebra (B1.2), any homomorphism  $\phi : K \rightarrow A$  of degree  $-1$  induces a graded algebra map  $g_\phi : T(s^{-1}K) \rightarrow A$  by

$$g_\phi = \sum_k \mu^k(\alpha \otimes \dots \otimes \alpha).$$

If, in addition,  $\phi : K \rightarrow A$  is a twisting cochain, that is, it satisfies Brown's condition  $\delta\phi = \phi \smile \phi$ , then  $g_\phi : T(s^{-1}K) \rightarrow A$  is a chain map, i.e., it is a map of dg algebras: since of freeness of tensor algebra one has the equality  $d_A g_\phi = g_\phi d_\Omega$  if and only if equal the restrictions  $d_A g_\phi i = g_\phi d_\Omega i$  and this is exactly the Brown's condition  $\delta\phi = \phi \smile \phi$ .

Conversely, any dg algebra map  $g : T(s^{-1}K) \rightarrow A$  is  $g_\phi$  for  $\phi = gi : K \rightarrow A$ . In fact we have a bijection  $Mor_{DGA_{\text{alg}}}(T(s^{-1}K), A) \leftrightarrow Tw(K, A)$

So we have bijections

$$\begin{array}{ccc} & Tw(K, A) & \\ \approx \swarrow & & \searrow \approx \\ Mor_{DGA}(\Omega K, A) & & Mor_{DGC}(\Omega K, A) \end{array}$$

#### 4.1.5 (Co)universal twisting cochains

The standard inclusion  $i : K \rightarrow \Omega K$  satisfies Brown's condition. So it is a (so called universal) twisting cochain. Thus it induces two chain maps

$$f_i : \Omega K \rightarrow \Omega K \quad \text{and} \quad g_i : K \rightarrow B\Omega K.$$

The first one is the identity map and the second one a quasi isomorphism (homology isomorphism).

Dually, the standard projection  $p : BA \rightarrow A$  satisfies the Brown's condition, so it is a (so called couniversal) twisting cochain. Thus it induces two chain maps

$$f_p : \Omega BA \rightarrow A \quad \text{and} \quad g_p : BA \rightarrow BA.$$

The first one is a quasi isomorphism (homology isomorphism) and the second one is the identity map.

## 4.2 Berikashvili's Functor $D$

### 4.2.1 Equivalence of twisting cochains

Two twisting cochains  $\phi, \psi : K \rightarrow A$  are equivalent (Berikashvili [3]) if there exists  $c : K \rightarrow A$ ,  $\deg c = 0$ ,  $c = \sum_i c_i$ ,  $c_i : C_i \rightarrow A_i$  with  $c_0 = c|_{C_0} = 0$ , such that

$$\psi = \phi + \delta c + \psi \smile c + c \smile \phi, \tag{4}$$

notation  $\phi \sim_c \psi$ .

This is an equivalence relation:

$$\phi \sim_{c=0} \phi; \quad \phi \sim_c \phi', \phi' \sim_{c'} \phi'' \Rightarrow \phi \sim_{c+c'+c' \smile c} \phi''$$

and  $\phi \sim_c \phi' \Rightarrow \phi' \sim_{c'} \phi$  where  $c'$  can be solved from  $c + c' + c' \smile c = 0$  inductively.

This notion of equivalence allows to *perturb* twisting cochains. Let

$$\phi = \phi_2 + \phi_3 + \dots + \phi_n + \dots : K \rightarrow A, \quad \phi_n : K_n \rightarrow A_{n-1}$$

be a twisting cochain, and let's take an arbitrary cochain  $c = c_n : K_n \rightarrow A_n$ . Then there exists a twisting cochain  $F_{c_n}\phi = \psi : K \rightarrow A$  such that  $\phi \sim_{c_n} \psi$ . Actually the components of the perturbed twisting cochain

$$F_{c_n}\phi = \psi = \psi_2 + \psi_3 + \dots + \psi_n + \dots$$

can be solved from (4) inductively, and the solution particularly gives that the perturbation  $F_{c_n}\phi$  does not change the first components, i.e.  $\psi_i = \phi_i$  for  $i < n$  and  $\psi_n = \phi_n + d_A c_n$ .

The main benefit of this notion is that the equivalent twisting cochains define isomorphic twisted tensor products:

**Theorem 1** . *If  $\phi \sim_c \psi$  then*

$$(K \otimes_\phi M, d_\phi) \xrightarrow{F_c} (K \otimes_\psi M, d_\psi)$$

*given by  $F_c(k \otimes m) = (k \otimes m) + (k \otimes m) \cap c$  is an isomorphism of dg modules.*

The inverse  $(K \otimes_\psi M, d_\psi) \xrightarrow{F_{c'}} (K \otimes_\phi M, d_\phi)$  is defined by  $c' : K \rightarrow A$  which, as above, can be solved inductively from  $c + c' + c' \smile c = 0$ .

Let

$$Tw(K, A) = \{\phi : K \rightarrow A, \delta\phi = \phi \circ \phi\}.$$

be the set of all twisting cochains.

**Definition 12** . *Berikashvili's functor  $D(K, A)$  is defined as the factorset  $D(K, A) = \frac{Tw(K, A)}{\sim}$ .*

Berikashvili's relation of equivalence of twisting cochains allows to perturb a given twisting cochain so as to get the simplest one to simplify calculations.

#### 4.2.2 Equivalence of twisting cochains and homotopy of induced maps

How does the equivalence of twisting cochains affect induced maps  $K \rightarrow BA$  and  $\Omega K \rightarrow A$ ?

**Theorem 2** . *If  $\phi \sim_c \psi$  then  $f_\phi$  and  $f_\psi$  are homotopic as dg coalgebra maps: the coderivation chain homotopy  $D(c) : K \rightarrow BA$  is given by*

$$D(c) = \sum_{i,j} (\psi \otimes \dots (j \text{ times}) \dots \otimes \psi \otimes c \otimes \phi \otimes \dots \otimes \phi) \nabla_K^i. \quad (5)$$

**Proof.** The homotopy  $D(c)$  satisfies  $f_\phi - f_\psi = d_B D(c) + D(c)d_K$  because the projection of this condition on  $A$  gives  $pf_\phi - pf_\psi = pd_B D(c) + pD(c)d_K$  and this is exactly the condition (4). Besides,  $D(c)$  is a  $f_\phi - f_\psi$ -coderivation, that is,

$$\nabla_B D(c) = (f_\psi \otimes D(c) + D(c) \otimes f_\phi) \nabla_K.$$

This also follows from the universal property of the tensor coalgebra and the extension rule (14).

The converse is also true: if  $f_\phi$  and  $f_\psi$  are homotopic by a coderivation homotopy  $D : K \rightarrow BA$  then  $\phi \sim_c \psi$  by  $c = p_1 D$ .

The dual statement is also true:  $g_\phi, g_\psi : \omega K \rightarrow A$  are homotopic in the category of dg algebras if and only if  $\phi \sim \psi$ .

So we have bijections

$$[\Omega K, A]_{DGA} \underset{\approx}{\swarrow} \begin{matrix} D(K, A) \\ \searrow \approx \end{matrix} [(K, BA)]_{DGC}.$$

this means that  $B$  and  $\Omega$  are adjoint functors.

### 4.2.3 Lifting of Twisting Cochains

Any dg algebra map  $f : A \rightarrow A'$  induces the map  $Tw(K, A) \rightarrow Tw(K, A')$ : if  $\phi : K \rightarrow M$  is a twisting cochain so is the composition  $f\phi : K \rightarrow A \rightarrow A'$ . Moreover, if  $\phi \sim_c \phi'$  then  $f\phi \sim_{fc} f\phi'$ . Thus we have a map  $D(f) : D(K, A) \rightarrow D(K, A')$ .

**Theorem 3** . (Berikashvili [3]). *Let  $(K, d_K, \nabla_K)$  be a dg colagebra with free  $K_i$  and let  $(A, d_A, \mu_A)$  be a connected dg algebra. If  $f : A \rightarrow A'$  is a weak equivalence of connected dg algebras (i.e., a homology isomorphism), then*

$$D(f) : D(K, A) \rightarrow D(K, A')$$

*is a bijection.*

Particularly this theorem means that  $[K, BA] \rightarrow [K, BA']$  is a bijection.

Below we'll need the surjectivity part of this theorem whose proof we sketch here.

**Theorem 4 .** Let  $(K, d_K, \nabla_K)$  be a dg colagebra with free  $K_i$ s and let  $f : (A, d_A, \mu_A) \rightarrow (A', d_{A'}, \mu_{A'})$  be a weak equivalence of connected dg algebras. Then for an arbitrary twisting cochain

$$\phi = \phi_2 + \phi_3 + \dots + \phi_n + \dots : K \rightarrow A'$$

there exists a twisting cochain

$$\psi = \psi_2 + \psi_3 + \dots + \psi_n + \dots : K \rightarrow A$$

such that  $\phi \sim f\psi$ .

**Proof.** Start with a twisting cochain  $\phi = \phi_2 + \phi_3 + \dots + \phi_n + \dots : K \rightarrow A'$ . Brown's defining condition (3) gives  $d_A\phi_2 = 0$ , and since  $f : A \rightarrow A'$  is a homology isomorphism there exist  $\psi_2 : K_2 \rightarrow A_1$  and  $c'_2 : K_2 \rightarrow A'_2$  such that  $d_A\psi_2 = 0$  and  $f\psi_2 = \phi_2 + d_{A'}c'_2$  (we assume all  $K_n$  are free modules). Perturbing  $\phi$  by this  $c'_2$  we obtain a new twisting cochain  $F_{c'_2}\phi$  for which  $(F_{c'_2}\phi)_2 = \phi_2 + d_{A'}c'_2 = f\psi_2$ . So we can assume that  $\phi_2 = f\psi_2$ .

Assume now that we already have  $\psi_2, \psi_3, \dots, \psi_{n-1}$  which satisfy (3) in appropriate dimensions and  $f\psi_k = \phi_k, k = 2, 3, \dots, n-1$ . We need the next component  $\psi_n : K_n \rightarrow A_{n-1}$  such that

$$d_A\psi_n = \psi_{n-1}d_K + \sum_{i=2}^{n-2} \psi_i \smile \psi_{n-i}$$

and  $c'_n : K_n \rightarrow A'_n$  such that  $f\psi_n = \phi_n + d_{A'}c'_n$ . Then perturbing  $\phi$  by  $c'_n$  we obtain new  $\phi_n$  for which  $f\psi_n = \phi_n$ , and this will complete the proof.

Let us write

$$\begin{aligned} U_n &= \psi_{n-1}d_K + \sum_{i=2}^{n-2} \psi_i \smile \psi_{n-i} : K_n \rightarrow A_{n-2}, \\ U'_n &= \phi_{n-1}d_K + \sum_{i=2}^{n-2} \phi_i \smile \phi_{n-i} : K_n \rightarrow A'_{n-2}. \end{aligned}$$

So we have  $U'_n = d_{A'}\phi_n$  and we want  $\psi_n : K_n \rightarrow A_{n-1}, c'_n : K_n \rightarrow A'_n$  such that  $U_n = d_A\psi_n$  and  $f\psi_n = \phi_n + d_{A'}c'_n$ .

First, it is not hard to check that  $d_A U_n = 0$ , that is,  $U_n$  maps  $K_n$  to cycles  $Z(A_{n-2}) \subset A_{n-2}$ .

Then

$$\begin{aligned} fU_n &= f(\psi_{n-1}d_K + \sum_{i=2}^{n-2} \psi_i \smile \psi_{n-i}) = f\psi_{n-1}d_K + \sum_{i=2}^{n-2} f\psi_i \smile f\psi_{n-i} \\ &= \phi_{n-1}d_K + \sum_{i=2}^{n-2} \phi_i \smile \phi_{n-i} = U'_n = d_{A'}\phi_n, \end{aligned}$$

thus,  $fU_n$  maps  $K_n$  to boundaries  $B(A_{n-2}) \subset A_{n-2}$ . Since  $f : A \rightarrow A'$  is a homology isomorphism, there is  $\overline{\psi}_n : K_n \rightarrow A_{n-1}$  such that  $d_A \overline{\psi}_n = U_n$ .

Now we must take care of the condition  $f\psi_n = \phi_n$ . It is clear that  $d_A f\overline{\psi}_n = d_A \phi_n$ . Let us denote by  $z'_n = f\overline{\psi}_n - \phi_n$ ; this is the homomorphism which maps  $K_n$  to cycles  $Z(A'_{n-1})$ . Again, since  $f : A \rightarrow A'$  is a homology isomorphism there exist  $z_n : K_n \rightarrow Z(A_{n-1})$  and  $c'_n : K_n \rightarrow A'_n$  such that  $fz_n = z'_n - d_{A'} c'_n$ . Let us define  $\psi_n = \overline{\psi}_n - z_n$ . Then  $d_A \psi_n = d_A \overline{\psi}_n$  and

$$f\psi_n = f\overline{\psi}_n - fz_n = (\phi_n + z'_n) - (z'_n - d_{A'} c'_n) = \phi_n + d_{A'} c'_n.$$

Perturbing  $\phi$  by this  $c'_n$  we obtain  $F_{c'_n} \phi$  with  $(F_{c'_n} \phi)_n = \phi_n + d_{A'} c'_n = f\psi_n$ .

## 5 Stasheff's $A_\infty$ -algebras

### 5.1 Category of $A_\infty$ -algebras

The notion of  $A_\infty$ -algebra was introduced by J. Stasheff [32]. This notion generalizes the notion of differential graded algebra and in fact it is so called *strong homotopy associative algebra* where the strict associativity is replaced with *associativity up to higher coherent homotopies*.

#### 5.1.1 Notion of $A_\infty$ -algebra

**Definition 13** . An  $A_\infty$ -algebra is a graded module  $M = \{M^k\}_{k \in \mathbb{Z}}$  equipped with a sequence of operations

$$\{m_i : M \otimes \dots (i \text{ times}) \dots \otimes M \rightarrow M, i = 1, 2, 3, \dots\}$$

satisfying the conditions  $m_i((\otimes^i M)^q) \subset M^{q-i+2}$ , that is  $\deg m_i = 2 - i$ , and

$$\sum_{k=0}^{i-1} \sum_{j=1}^{i-k} \pm m_{i-j+1}(a_1 \otimes \dots \otimes a_k \otimes m_j(a_{k+1} \otimes \dots \otimes a_{k+j}) \otimes \dots \otimes a_i) = 0. \quad (6)$$

For  $i = 1$  this condition reads

$$m_1 m_1 = 0.$$

For  $i = 2$  this condition reads

$$m_1 m_2(a_1 \otimes a_2) \pm m_2(m_1(a_1) \otimes a_2) \pm m_2(a_1 \otimes m_1(a_2)) = 0.$$

For  $i = 3$  this condition reads

$$m_1 m_3(a_1 \otimes a_2 \otimes a_3) \pm m_3(m_1(a_1) \otimes a_2 \otimes a_3) \pm m_3(a_1 \otimes m_1(a_2) \otimes a_3) \pm m_3(a_1 \otimes a_2 \otimes m_1(a_3)) \pm m_2(m_2(a_1 \otimes a_2) \otimes a_3) \pm m_2(a_1 \otimes m_2(a_2 \otimes a_3)) = 0.$$

These three conditions mean that for an  $A_\infty$ -algebra  $(M, \{m_i\})$  the first two operations form a *nonassociative* dga  $(M, m_1, m_2)$  with differential  $m_1$  and multiplication  $m_2$  which is associative just up to homotopy and the suitable homotopy is the operation  $m_3$ .

*Spacial case:* An  $A_\infty$ -algebra  $(M, \{m_1, m_2, m_3 = 0, m_4 = 0, \dots\})$  is a strictly associative dg algebra.

### 5.1.2 Morphism of $A_\infty$ -algebras

This is the notion from ([17]).

**Definition 14** . *A morphism of  $A_\infty$ -algebras*

$$\{f_i\} : (M, \{m_i\}) \rightarrow (M', \{m'_i\})$$

is a sequence  $\{f_i : \otimes^i M \rightarrow M', i = 1, 2, \dots, \deg f_1 = 1 - i\}$  such that

$$\begin{aligned} & \sum_{k=0}^{i-1} \sum_{j=1}^{i-k} \pm f_{i-j+1}(a_1 \otimes \dots \otimes a_k \otimes m_j(a_{k+1} \otimes \dots \otimes a_{k+j}) \otimes \dots \otimes a_i) = \\ & \sum_{t=1}^i \sum_{k_1+\dots+k_t=i} \pm m'_t(f_{k_1}(a_1 \otimes \dots \otimes a_{k_1}) \otimes \dots \otimes f_{k_t}(a_{i-k_t+1} \otimes \dots \otimes a_i)). \end{aligned} \quad (7)$$

In particular for  $n = 1$  this condition gives  $f_1 m_1(a) = m'_1 f_1(a)$ , i.e.,  $f_1 : (M, m_1) \rightarrow (M', m'_1)$  is a chain map; for  $n = 2$  it gives

$$f_1 m_2(a_1 \otimes a_2) + m'_2(f_1(a_1) \otimes f_1(a_2)) = m'_1 f_2(a_1 \otimes a_2) + f_2(m_1 a_1 \otimes a_2) + f_2(a_1 \otimes m_1 a_2),$$

thus  $f_1 : (M, m_1, m_2) \rightarrow (M', m'_1, m'_2)$  is multiplicative just up to the chain homotopy  $f_2$ .

The composition of  $A_\infty$  morphisms

$$\{h_i\} : (M, \{m_i\}) \xrightarrow{\{f_i\}} (M', \{m'_i\}) \xrightarrow{\{g_i\}} (M'', \{m''_i\})$$

is defined as

$$\begin{aligned} h_n(a_1 \otimes \dots \otimes a_n) &= \sum_{t=1}^n \sum_{k_1+\dots+k_t=n} \\ g_n(f_{k_1}(a_1 \otimes \dots \otimes a_{k_1}) \otimes \dots \otimes f_{k_t}(a_{n-k_t+1} \otimes \dots \otimes a_n)). \end{aligned} \quad (8)$$

The bar construction argument (see (5.1.3) below) allows to show that so defined composition satisfies the condition (7).

*Special case:* A morphism of  $A_\infty$ -algebras  $\{f_1, f_2 = 0, f_3 = 0, \dots\} : (M, \{m_1, m_2, m_3 = 0, m_4 = 0, \dots\}) \rightarrow (M, \{m_1, m_2, m_3 = 0, m_4 = 0, \dots\})$  is ordinary map of  $DG$ -algebras. In fact, the category of dg algebras is a subcategory of the category of  $A_\infty$ -algebras.

### 5.1.3 Bar construction of an $A_\infty$ -algebra

Let  $(M, \{m_i\})$  be an  $A_\infty$ -algebra. The structure maps  $m_i$  define the map  $\beta : T^c(s^{-1}M) \rightarrow s^{-1}M$  by  $\beta[a_1, \dots, a_n] = [s^{-1}m_n(a_1 \otimes \dots \otimes a_n)]$ . Extending this  $\beta$  as a coderivation (see the universal property (3.1.6)) we obtain  $d_\beta : T^c(s^{-1}M) \rightarrow T^c(s^{-1}M)$  which in fact looks as

$$d_\beta[a_1, \dots, a_n] = \sum_k \pm [a_1, \dots, a_k, m_j(a_{k+1} \otimes \dots \otimes a_{k+j}), a_{k+j+1}, \dots, a_n].$$

The defining condition (28) of an  $A_\infty$ -algebra guarantees that  $d_\beta d_\beta = 0$ : the composition of coderivations  $d_\beta d_\beta : T^c(s^{-1}M) \rightarrow T^c(s^{-1}M)$  is a coderivation and the defining condition (28) is nothing else than that  $p_1 d_\beta d_\beta = 0$  and this is equivalent to  $d_\beta d_\beta = 0$  since the tensor coalgebra is cofree.

The obtained dg coalgebra  $(T^c(s^{-1}M), d_\beta, \Delta)$  is called *bar construction* of the  $A_\infty$ -algebra  $(M, \{m_i\})$  and is denoted by  $B(M, \{m_i\})$ .

For an  $A_\infty$ -algebra of type  $(M, \{m_1, m_2, 0, 0, \dots\})$ , i.e., for a strictly associative dg algebra, this bar construction coincides with the ordinary bar construction of this dg algebra.

### 5.1.4 Bar interpretation of a morphism of $A_\infty$ -algebras

A morphism of  $A_\infty$ -algebras  $\{f_i\} : (M, \{m_i\}) \rightarrow (M', \{m'_i\})$  defines a dg coalgebra map of bar constructions

$$F = B(\{f_i\}) : B(M, \{m_i\}) \rightarrow B(M', \{m'_i\})$$

as follows: The collection  $\{f_i\}$  defines the map  $\alpha : T^c(s^{-1}M) \rightarrow s^{-1}M$  by  $\alpha[a_1, \dots, a_n] = [s^{-1}f_n(a_1 \otimes \dots \otimes a_n)]$ . Extending this  $\alpha$  as a coalgebra map (3.1.2)) we obtain  $F : T^c(s^{-1}M) \rightarrow T^c(s^{-1}M)$  which in fact looks like

$$F[a_1, \dots, a_n] = \sum \pm [f_{k_1}(a_1 \otimes \dots \otimes a_{k_1}), \dots, f_{k_t}(a_{n-k_t+1} \otimes \dots \otimes a_n)].$$

The defining condition (7) of an  $A_\infty$ -morphism is nothing else than  $p_1 d_{\beta'} F = p_1 F d_\beta$ , and this is equivalent to  $d_{\beta'} F = F d_\beta$ , so  $F$  is a chain map.

Now we are able to show that the composition of  $A_\infty$ -morphisms is well defined: to the composition of morphisms (8) corresponds the composition of dg coalgebra maps

$$B((M, \{m_i\})) \xrightarrow{B(\{f_i\})} B((M', \{m'_i\})) \xrightarrow{B(\{g_i\})} B((M'', \{m''_i\}))$$

which is a dg coalgebra map. Thus for the projection  $p_1 B(\{g_i\}) B(\{f_i\})$ , i.e., for the collection  $\{h_i\}$ , the condition (7) is satisfied.

### 5.1.5 Homotopy in the category of $A_\infty$ -algebras

Two morphisms of  $A_\infty$ -algebras

$$\{f_i\}, \{g_i\} : (M, \{m_i\}) \rightarrow (M', \{m'_i\})$$

we call *homotopic* if there exists a collection of homomorphisms

$$\{h_i : (\otimes^i M) \rightarrow M', \quad i = 1, 2, \dots, \quad \deg h_i = -i\},$$

which satisfy the following condition

$$\begin{aligned} & f_n(a_1 \otimes \dots \otimes a_n) - g_n(a_1 \otimes \dots \otimes a_n) \\ &= \sum_{i+j=n+1} \sum_{k=0}^{n-j} h_i(a_1 \otimes \dots \otimes a_k \otimes m_j(a_{k+1} \otimes \dots \otimes a_{k+j}) \otimes \dots \otimes a_n) \\ &+ \sum_{k_1+\dots+k_t=n} m'_t(f_{k_1}(a_1 \otimes \dots \otimes a_{k_1}) \otimes \dots \otimes f_{k_{i-1}}(a_{k_1+\dots+k_{i-2}+1} \otimes \dots \otimes a_{k_1+\dots+k_{i-1}}) \\ &\quad \otimes h_{k_i}(a_{k_1+\dots+k_{i-1}+1} \otimes \dots \otimes a_{k_1+\dots+k_i}) \otimes g_{k_{i+1}}(a_{k_1+\dots+k_i+1} \otimes \dots \otimes a_{k_1+\dots+k_{i+1}}) \\ &\quad \otimes g_{k_t}(a_{k_1+\dots+k_{t-1}+1} \otimes \dots \otimes a_{k_1+\dots+k_t}). \end{aligned} \tag{9}$$

In particular for  $n = 1$  this condition means

$$f_1(a) - g_1(a) = m'_1 h_1(a) + h_1(m_1 a),$$

that is, the chain maps  $f_1, g_1 : (M, m_1) \rightarrow (M', m'_1)$  are chain homotopic.

For  $n = 2$  this condition means

$$\begin{aligned} & f_2(a_1 \otimes a_2) - g_2(a_1 \otimes a_2) = m'_1 h_2(a_1 \otimes a_2) + m'_2 (f_1(a_1) \otimes h_1(a_2)) \\ &+ m'_2 (h_1(a_1) \otimes g_1(a_2)) + h_1(m_2(a_1 \otimes a_2)) + h_2(m_1 a_1 \otimes a_2) + h_2(a_1 \otimes m_1 a_2). \end{aligned} \tag{10}$$

### 5.1.6 Bar interpretation of homotopy

The collections  $\{f_i\}, \{g_i\}, \{h_i\}$  define a homomorphism

$$D : B(M, \{m_i\}) \rightarrow B(M', \{m'_i\})$$

by

$$\begin{aligned} & D(a_1 \otimes \dots \otimes a_n) \\ = & \sum_{k_1 + \dots + k_t = n} f_{k_1}(a_1 \otimes \dots \otimes a_{k_1}) \otimes \dots \otimes f_{k_{i-1}}(a_{k_1 + \dots + k_{i-2} + 1} \otimes \dots \otimes a_{k_1 + \dots + k_{i-1}}) \\ & \otimes h_{k_i}(a_{k_1 + \dots + k_{i-1} + 1} \otimes \dots \otimes a_{k_1 + \dots + k_i}) \otimes g_{k_{i+1}}(a_{k_1 + \dots + k_i + 1} \otimes \dots \otimes a_{k_1 + \dots + k_{i+1}}) \\ & \otimes g_{k_t}(a_{k_1 + \dots + k_{t-1} + 1} \otimes \dots \otimes a_{k_1 + \dots + k_t}), \end{aligned}$$

which is a  $(B(\{f_i\}), B(\{g_i\}))$ -coderivation.

Besides, the condition (9) means nothing else than

$$p_1(B(\{f_i\}) - B(\{g_i\})) = p_1(d_{m'}D + Dd_m),$$

and this, again since the tensor coalgebra is cofree, gives

$$B(\{f_i\}) - B(\{g_i\}) = d_{m'}D + Dd_m,$$

that is, the dg coalgebra maps  $B(\{f_i\})$  and  $B(\{g_i\})$  are homotopic in the category of dg coalgebras.

### 5.1.7 Category *DASH*

The category *DASH* (Differential Algebras and Strongly Homotopy Multiplicative Maps) was first considered in Halperin and Stasheff's article [13]. The objects are dg algebras, and a morphism  $\{f_i\} : (A, d, \mu) \rightarrow (A', d', \mu')$  is defined as a collection of homomorphisms

$$\{f_i : \otimes^i A \rightarrow A; , i = 1, 2, \dots, \deg f_i = 1 - i\}$$

which satisfies the following conditions

$$\begin{aligned} & \sum_i f_n(a_1 \otimes \dots \otimes a_{i-1} \otimes da_i \otimes a_{i+1} \otimes \dots \otimes a_n) \\ + & \sum_i f_{n-1}(a_1 \otimes \dots \otimes a_{i-1} \otimes a_i \cdot a_{i+1} \otimes a_{i+2} \otimes \dots \otimes a_n) \\ = & \sum_i f_i(a_1 \otimes \dots \otimes a_i) \cdot f_{n-i}(a_{i+1} \otimes \dots \otimes a_n). \end{aligned} \quad (11)$$

In particular, for  $n = 1$  this condition gives  $f_1 d(a) = df_1(a)$ , i.e.,  $f_1 : (A, d) \rightarrow (A', d')$  is a chain map; for  $n = 2$  it gives

$$\begin{aligned} & f_1(a_1 \cdot a_2) - f_1(a_1) \cdot f_1(a_2) \\ = & d' f_2(a_1 \otimes a_2) + f_2(da_1 \otimes a_2) + f_2(a_1 \otimes da_2), \end{aligned}$$

thus  $f_1 : (A, d, \mu) \rightarrow (A', d', \mu')$  is multiplicative up to the homotopy  $f_2$ .

The existence of higher components  $\{f_i, i = 1, 2, 3, 4, \dots\}$  is the reason why such a morphism is called a *Strongly Homotopy Multiplicative Map* and the category is called DASH: *Differential Algebras and Strongly Homotopy multiplicative maps*.

In fact this is a *full subcategory* of the category of  $A_\infty$ -algebras whose objects are ordinary dg-algebras  $(A, d, \mu) = (A, \{m_1 = d, m_2 = \mu, m_3 = 0, m_4 = 0, \dots\})$ : the defining condition of an  $A_\infty$ -morphism (7) here looks like (11).

Thus we have hierarchy of categories

$$DGA_{lg} \subset DASH \subset A_\infty,$$

all three with notions of homotopies, and we have the commutative diagram of functors

$$\begin{array}{ccccc} DGA_{lg} & \subset & DASH & \subset & A_\infty \\ B \searrow & & B \downarrow & & \swarrow B \\ & & DGCoalg. & & \end{array}$$

### 5.1.8 Minimality

Let  $\{f_i\} : (M, \{m_i\}) \rightarrow (M', \{m'_i\})$  be a morphism of  $A_\infty$ -algebras. It follows from (7) that the first component  $f_1 : (M, m_1) \rightarrow (M', m'_1)$  is a chain map.

Let us define a weak equivalence of  $A_\infty$ -algebras as a morphism  $\{f_i\}$  for which  $B(\{f_i\})$  is a weak equivalence (homology isomorphism) of dg coalgebras. The standard spectral sequence argument allows to prove the following

**Proposition 13** . *A morphism of  $A_\infty$ -algebras is a weak equivalence if and only if its first component  $f_1 : (M, m_1) \rightarrow (M', m'_1)$  is a weak equivalence of chain complexes.*

Furthermore, it is easy to see that if  $\{f_i\} : (M, \{m_i\}) \rightarrow (M', \{m'_i\})$  is an isomorphism of  $A_\infty$  algebras then its first component  $f_1 : (M, m_1) \rightarrow (M', m'_1)$  is an isomorphism of chain complexes. Conversely, if  $f_1$  is an isomorphism, then  $\{f_i\}$  is an isomorphism of  $A_\infty$  algebras: The components of the opposite morphism  $\{g_i\} : (M', \{m'_i\}) \rightarrow (M, \{m_i\})$  can be solved inductively from the equation  $\{g_i\}\{f_i\} = \{id_M, 0, 0, \dots\}$ . Thus we have the

**Proposition 14** . *A morphism of  $A_\infty$ -algebras is an isomorphism if and only if its first component  $f_1 : (M, m_1) \rightarrow (M', m'_1)$  is an isomorphism of dg modules.*

**Definition 15** . An  $A_\infty$ -algebra  $(M, \{m_i\})$  we call *minimal* if  $m_1 = 0$ .

In this case  $(M, m_2)$  is a *strictly* associative graded algebra.

The above propositions easily imply:

**Proposition 15** . Each weak equivalence of minimal  $A_\infty$ -algebras is an isomorphism.

**Proof.** Suppose  $f = \{f_i\} : (M, \{m_i\}) \rightarrow (M', \{m'_i\})$  is a weak equivalence of  $A_\infty$ -algebras. Then by (13) the chain map  $f_1 : (M, m_1 = 0) \rightarrow (M', m'_1 = 0)$  induces an isomorphism of homology modules  $f_1^* : H(M, m_1 = 0) \rightarrow H(M', m'_1 = 0)$ , but  $H(M, m_1 = 0) = M$  and  $H(M', m_1 = 0) = M'$ . So  $f_1$  is an isomorphism and by of (14)  $f$  is an isomorphism too.

This fact motivates the word *minimal* in this notion since Sullivan's minimal model has a similar property.

## 5.2 $A_\infty$ Twisting Cochains

Here we are going to generalize the above material about twisting cochains and the functor  $D$  from the case of dg algebras to the case of  $A_\infty$  algebras, see [19], see also [30].

Start with a dg coalgebra  $(C, d_C, \nabla : C \rightarrow C \otimes C)$  and an  $A_\infty$ -algebra  $(M, \{m_i\})$ . We'll recall the following multicooperations

$$\begin{aligned} \nabla^i : C \rightarrow C \otimes \dots (i \text{ times}) \dots \otimes C \rightarrow C, \quad i = 1, 2, \dots, \\ \nabla^1 = id, \quad \nabla^2 = \nabla, \quad \nabla^i = (\nabla^{i-1} \otimes id)\nabla. \end{aligned}$$

### 5.2.1 The notion of $A_\infty$ -twisting cochain

An  $A_\infty$ -twisting cochain [19], [20] we have defined as a homomorphism  $\phi : C \rightarrow M$  of degree  $-1$  (that is  $\phi : C^{*} \rightarrow M^{*-1}$ ) satisfying the condition

$$\phi d_C = \sum_{i=1}^{\infty} m_i(\phi \otimes \dots \otimes \phi)\nabla^i. \quad (12)$$

Let  $Tw(C, M)$  be the set of all such twisting cochains.

Note that if for  $(M, \{m_i\})$  one has  $m_{>2} = 0$ , i.e., it is an ordinary dg algebra  $(M, \{m_1, m_2, 0, 0, \dots\})$  then this notion coincides with Brown' classical notion.

### 5.2.2 Bar interpretation

Any  $A_\infty$  twisting cochain  $\phi : C \rightarrow M$  induces a dg-coalgebra map  $B(\phi) : C \rightarrow BM$  by

$$B(\phi) = \sum_i (\phi \otimes \dots \otimes \phi) \nabla^i.$$

So defined  $B(\phi)$  automatically is a coalgebra map and the condition (12) guarantees that it is a chain map.

Conversely, any dg coalgebra map  $f : C \rightarrow BM$  is  $B(\phi)$  for  $\phi = p \circ f : C \rightarrow BM \rightarrow M$ . In fact we have a bijection

$$Mor_{DGC\text{alg}}(C, BM) \leftrightarrow Tw(C, M).$$

### 5.2.3 Equivalence of $A_\infty$ -twisting cochains.

Two  $A_\infty$ -twisting cochains  $\phi, \psi : C \rightarrow M$  are called equivalent [19] if there exists  $c : C \rightarrow M$ ,  $deg\ c = 0$ , such that

$$\phi - \psi = cd_C + \sum_{i=1}^{\infty} \sum_{k=1}^{i-1} m_i(\phi \otimes \dots (k\ \text{times}) \dots \phi \otimes c \otimes \psi \otimes \dots \psi) \nabla^i, \quad (13)$$

notation  $\phi \sim_c \psi$ .

Again, if for  $(M, \{m_i\})$  one has  $m_{>2} = 0$ , i.e., it is an ordinary dg algebra  $(M, \{m_1, m_2, 0, 0, \dots\})$  then this notion coincides with Berikashvili's described above equivalence.

### 5.2.4 Bar interpretation

If  $\phi \sim_c \psi$  then  $B(\phi)$  and  $B(\psi)$  are homotopic in the category of dg coalgebras: a chain homotopy  $D(c) : C \rightarrow BA$  is given by

$$D(c) = \sum_{i=1}^{\infty} \sum_{k=1}^{i-1} (\phi \otimes \dots (k\ \text{times}) \dots \phi \otimes c \otimes \psi \otimes \dots \psi) \nabla^i. \quad (14)$$

So defined  $D$  automatically is a  $B(\phi), B(\psi)$ -coderivation and the condition (13) guarantees that it realizes chain homotopy, that is,  $B(\phi) - B(\psi) = d_B D(c) + D(c) d_C$ .

### 5.2.5 The Functor $\tilde{D}$ .

The functor  $\tilde{D}(C, M)$  ([19]) is defined as the factorset

$$\tilde{D}(C, A) = \frac{Tw(C, M)}{\sim}.$$

Any  $A_\infty$ -algebra map  $f = \{f_i\} : M \rightarrow M'$  induces the map  $Tw(C, M) \rightarrow Tw(C, M')$ : if  $\phi : C \rightarrow M$  is an  $A_\infty$ -twisting cochain then so is

$$f\phi = p'B(f)B(\phi) = \sum_i f_i(\phi \otimes \dots \otimes \phi)\nabla^i : C \rightarrow B(M) \rightarrow B(M') \rightarrow M'. \quad (15)$$

Furthermore, if  $\phi \sim_c \psi$  then  $f\phi \sim_C f\psi$  with

$$C = \sum_{i=1}^{\infty} \sum_{k=1}^{i-1} (\phi \otimes \dots (k \text{ times}) \dots \phi \otimes c \otimes \psi \otimes \dots \psi)\nabla^i.$$

Thus we have a map  $\tilde{D}(f) : \tilde{D}(C, M) \rightarrow \tilde{D}(C, M')$ .

Dually, any dg coalgebra map  $g : C' \rightarrow C$  induces a map  $Tw(C, M) \rightarrow Tw(C', M)$ : if  $\phi : C \rightarrow M$  is an  $A_\infty$ -twisting cochain so is the composition  $\phi g : C' \rightarrow C \rightarrow M$ . Moreover, if  $\phi \sim_c \psi$  then  $\phi g \sim_{cg} \psi g$ . Thus we have a map  $D(g) : D(C, M) \rightarrow D(C', M)$ .

### 5.2.6 Bar interpretation

Assigning to an  $A_\infty$ -twisting cochain  $\phi : C \rightarrow M$  the dg coalgebra map  $B(\phi) : C \rightarrow BM$  and having in mind that  $\phi \sim_c \psi$  implies  $B(f\phi) \sim_{D(c)} B(\psi)$  we obtain the

**Theorem 5** . *There is a bijection  $\tilde{D}(C, M) \leftrightarrow [C, BM]_{DGC\text{alg}}$ .*

Taking  $C = B(M')$ , the bar construction of an  $A_\infty$ -algebra  $(M', \{m'_i\})$  and having in mind that  $[B(M'), B(M)]_{DGC\text{alg}}$  is bijective to  $[M', M]_{A_\infty}$ , the set of homotopy classes in the category of  $A_\infty$ -algebras, we obtain the

**Proposition 16** . *There is a bijection  $D(B(M'), M) \leftrightarrow [M', M]_{A_\infty}$ .*

### 5.2.7 Bijections.

The following property of the functor  $\tilde{D}$  makes it useful in the homotopy classification of maps.

**Theorem 6** . (See [19]) (a) If  $\{f_i\} : M \rightarrow M'$  is a weak equivalence of  $A_\infty$ -algebras, then

$$D(\{f_i\}) : D(C, M) \rightarrow D(C, M')$$

is a bijection.

(b) If  $g : C \rightarrow C'$  is a weak equivalence of connected dg coalgebras (i.e., a homology isomorphism), then

$$D(g) : D(C', M') \rightarrow D(C, M')$$

is a bijection.

Combining we obtain

**Proposition 17** . A weak equivalence of  $A_\infty$ -algebras  $\{f_i\} : M \rightarrow M'$  and a weak equivalence of dg coalgebras  $g : C \rightarrow C'$  induce bijections

$$D(C, M) \xrightarrow{D(\{f_i\})} D(C, M') \xleftarrow{D(g)} D(C', M').$$

The following theorem is an analogue of Berikashvilis's Theorem 3 for  $A_\infty$ -algebras and was proved in [19].

**Theorem 7** . Let  $(K, d_K, \nabla_K)$  be a dg coalgebra with free  $K_i$  and let  $(M, \{m_i\})$  be a connected dg algebra. If  $f = \{f_i\} : (M, \{m_i\}) \rightarrow (M', \{m'_i\})$  is a weak equivalence of  $A_\infty$ -algebras then

$$D_\infty(f) : D_\infty(K, M) \rightarrow D_\infty(K, M')$$

is a bijection.

In fact this theorem means that  $[K, BM] \rightarrow [K, BM']$  is a bijection. The theorem consists of two parts:

**Surjectivity.** Any  $A_\infty$ -twisting cochain  $\phi : K \rightarrow M'$  can be lifted to an  $A_\infty$ -twisting cochain  $\psi : K \rightarrow M$  so that  $\phi \sim f\psi$ .

**Injectivity.** If  $\psi \sim \psi' \in T_\infty(K, M)$  then  $f\psi \sim f\psi' \in T_\infty(K, M')$ .

### 5.2.8 Lifting of $A_\infty$ -Twisting Cochains

Below we'll use the surjectivity part of this theorem whose proof we sketch here.

**Theorem 8 .** *Let  $(K, d_K, \nabla_K)$  be a dg coalgebra with free  $K_i$  and let  $f : (M, \{m_i\}) \rightarrow (M', \{m'_i\})$  be a weak equivalence of connected  $A_\infty$ -algebras (that is,  $M_0 = M'_0 = 0$ ). Then for an arbitrary  $A_\infty$ -twisting cochain*

$$\phi = \phi_2 + \phi_3 + \dots + \phi_n + \dots : K \rightarrow M'$$

there exists an  $A_\infty$ -twisting cochain

$$\psi = \psi_2 + \psi_3 + \dots + \psi_n + \dots : K \rightarrow M$$

such that  $\phi \sim f\psi$ .

**Proof.** Start with an  $A_\infty$ -twisting cochain

$$\phi = \phi_2 + \phi_3 + \dots + \phi_n + \dots : K \rightarrow M'.$$

The condition (12) gives  $m_1\phi_2 = 0$ , and since  $f : (M, m_1) \rightarrow (M', m'_1)$  is homology isomorphism then there exist  $\psi_2 : K_2 \rightarrow M_1$  and  $c'_2 : K_2 \rightarrow M'_2$  such that  $m_1\psi_2 = 0$  and  $f\psi_2 = \phi_2 + m'_1c'_2$ . Perturbing  $\phi$  by this  $c'_2$  we obtain  $F_{c'_2}\phi$  for which  $(F_{c'_2}\phi)_2 = \phi_2 + m'_1c'_2 = f\psi_2$ .

Assume now that we already have  $\psi_2, \psi_3, \dots, \psi_{n-1}$  which satisfy (12) and (15) in appropriate dimensions, that is

$$m_1\psi_k = \psi_{k-1}d_K + \sum_{i=2}^{\text{int}(k/2)} \sum_{k_1+\dots+k_i=k} m_i(\psi_{k_1} \otimes \dots \otimes \psi_{k_i})\nabla_K^k, \quad k = 2, 3, \dots, n-1 \quad (16)$$

and

$$\phi_k = \sum_{i=1}^{\text{int}(k/2)} \sum_{k_1+\dots+k_i=k} f_i(\psi_{k_1} \otimes \dots \otimes \psi_{k_i})\nabla_K^i, \quad k = 1, 2, \dots, n-1. \quad (17)$$

We need the next component  $\psi_n : K_n \rightarrow A_{n-1}$  satisfying the condition (16) for  $k = n$

$$m_1\psi_n = \psi_{n-1}d_K + \sum_{i=2}^{\text{int}(n/2)} \sum_{k_1+\dots+k_i=n} m_i(\psi_{k_1} \otimes \dots \otimes \psi_{k_i})\nabla_K^k, \quad (18)$$

and  $c'_n : K_n \rightarrow A'_n$  such that

$$m_1 c'_n + \phi_n = \sum_{i=1}^{\text{int}(n/2)} \sum_{k_1+\dots+k_i=n} f_i(\psi_{k_1} \otimes \dots \otimes \psi_{k_i}) \nabla_K^i. \quad (19)$$

Then perturbing  $\phi$  by  $c'_n$  we obtain new  $\phi_n$  for which the condition (17) will be satisfied for  $k = n$ , and this will complete the proof.

Let us put

$$\begin{aligned} U_n &= \psi_{n-1} d_K + \sum_{i=2}^{\text{int}(n/2)} \sum_{k_1+\dots+k_i=n} m_i(\psi_{k_1} \otimes \dots \otimes \psi_{k_i}) \nabla_K^k, \\ U'_n &= \phi_{n-1} d_K + \sum_{i=2}^{\text{int}(n/2)} \sum_{k_1+\dots+k_i=n} m_i(\phi_{k_1} \otimes \dots \otimes \phi_{k_i}) \nabla_K^k, \\ V_n &= \sum_{i=2}^{\text{int}(n/2)} \sum_{k_1+\dots+k_i=n} f_i(\psi_{k_1} \otimes \dots \otimes \psi_{k_i}) \nabla_K^i. \end{aligned}$$

Then the needed conditions (18) and (19) become

$$U_n = m_1 \psi_n, \quad m'_1 c'_n + \phi_n = f_1 \psi_n + V_n.$$

First, it is possible to check that  $m_1 U_n = 0$  and  $f_1 U_n = U'_n + m'_1 V_n$ . Having in mind that  $U'_n = m'_1 \phi_n$  the last condition means  $f_1 U_n = m'_1(\phi_n + V_n)$ . So  $U_n$  is a map to  $m_1$ -cycles

$$U_n : K_n \rightarrow Z(M_{n-2}) \subset M_{n-2}$$

and  $f_1 U_n$  maps  $K_n$  to boundaries

$$f_1 U_n : K_n \rightarrow B(M'_{n-2}) \subset M_{n-2}.$$

Then since  $f_1 : (M, m_1) \rightarrow (M', m'_1)$  is a homology isomorphism, there exist  $\overline{\psi}_n : K_n \rightarrow M_{n-1}$  such that  $m_1 \overline{\psi}_n = U_n$ . So our  $\overline{\psi}_n$  satisfies (18). Now let us perturb this  $\overline{\psi}$  in order to catch the condition (19) too.

Using the above equality  $f_1 U_n = m'_1(\phi_n + V_n)$ , we have

$$m'_1 f_1 \overline{\psi}_n = f_1 m_1 \overline{\psi}_n = f_1 U_n = m'_1(\phi_n + V_n)$$

i.e.,  $m'_1(f_1 \overline{\psi}_n - (\phi_n + V_n)) = 0$ . Thus

$$z'_n = (f_1 \overline{\psi}_n - (\phi_n + V_n)) : K_n \rightarrow Z(M'_{n-1}),$$

and again, since  $f_1 : (M, m_1) \rightarrow (M', m'_1)$  is a homology isomorphism there exist  $z_n : K_n \rightarrow Z(M_{n-1})$  and  $c'_n : K_n \rightarrow M'_n$  such that  $z'_n = f_1 z_n - m'_1 c'_n$ . Let's define  $\psi_n = \overline{\psi}_n - z_n$ . Then

$$f_1 \psi_n = f_1 \overline{\psi}_n - f_1 z_n = (z'_n + \phi_n + V_n) - (z'_n - m'_1 c'_n) = \phi_n + V_n.$$

### 5.3 $C_\infty$ -algebras

This is the commutative version of the notion of  $A_\infty$ -algebra. For an  $A_\infty$ -algebra  $(M, \{m_i\})$  it is clear what it means for the operation  $m_2 : M \otimes M \rightarrow M$ , but what about the commutativity of the higher operations  $m_i : M \otimes \dots \otimes M \rightarrow M$ ,  $i \geq 3$ ? We are going to describe this now.

#### 5.3.1 Shuffle product

It was mentioned in (3.2) that the tensor algebra  $(T(V), \mu)$  and the tensor coalgebra  $(T^c(V), \Delta)$  coincide as graded modules, but the multiplication  $\mu$  of  $T(V)$  and the comultiplication  $\Delta$  of  $T^c(V)$  are not compatible with each other, so they do not define a graded bialgebra structure on  $T(V) = T^c(V)$ .

But, as it was indicated in (3.2) there exists the *shuffle* multiplication  $\mu_{sh} : T^c(V) \otimes T^c(V) \rightarrow T^c(V)$  introduced by Eilenberg and MacLane which turns  $(T^c(V), \Delta, \mu_{sh})$  into a graded bialgebra. This multiplication is defined as a graded coalgebra map induced by the universal property of  $T^c(V)$  by  $\alpha : T^c(V) \otimes T^c(V) \rightarrow V$  given by  $\alpha(v \otimes 1) = \alpha(1 \otimes v) = v$  and  $\alpha = 0$  otherwise. This multiplication is associative and in fact is given by

$$\mu_{sh}([a_1, \dots, a_m] \otimes [a_{i+1}, \dots, a_n]) = \sum \pm [a_{\sigma(1)}, \dots, a_{\sigma(n)}],$$

where the summation is taken over all  $(m, n)$ -shuffles. That is, over all permutations of the set  $(1, 2, \dots, n + m)$  which satisfy the condition:  $i < j$  if  $1 \leq \sigma(i) < \sigma(j) \leq n$  or  $n + 1 \leq \sigma(i) < \sigma(j) \leq n + m$ . In particular

$$[a] *_{sh} [b] = [a, b] \pm [b, a],$$

$$[a] *_{sh} [b, c] = [a, b, c] \pm [b, a, c] \pm [b, c, a].$$

#### 5.3.2 The notion of $C_\infty$ -algebra

Now we can define the notion of  $C_\infty$ -algebra, which is a commutative version of Stasheff's notion of  $A_\infty$ -algebra.

**Definition 16** . ([30], [21],[28], [11]) *A  $C_\infty$ -algebra is an  $A_\infty$ -algebra  $(M, \{m_i\})$  which additionally satisfies the following condition: each operation  $m_i$  vanishes on shuffles, that is, for  $a_1, \dots, a_i \in M$  and  $k = 1, 2, \dots, i - 1$*

$$m_i(\mu_{sh}((a_1 \otimes \dots \otimes a_k) \otimes (a_{k+1} \otimes \dots \otimes a_i))) = 0. \quad (20)$$

In particular, this gives

$$\begin{aligned} m_2(a \otimes b) + m_2(b \otimes a) &= 0, \\ m_3(a \otimes b \otimes c) + m_3(a \otimes c \otimes b) + m_3(c \otimes a \otimes b) &= 0, \\ \dots \end{aligned}$$

i.e., the maps  $f_i$  vanish on shuffles.

**Definition 17** . *A morphism of  $C_\infty$ -algebras is defined as a morphism of  $A_\infty$ -algebras  $\{f_i\} : (M, \{m_i\}) \rightarrow (M', \{m'_i\})$  whose components  $f_i$  vanish on shuffles, that is*

$$f_i((\mu_{sh}(a_1 \otimes \dots \otimes a_k) \otimes (a_{k+1} \otimes \dots \otimes a_i))) = 0. \quad (21)$$

The composition is defined as in the  $A_\infty$ -case and the bar construction argument (see (5.3.3) below) allows to show that the composition is a  $C_\infty$ -morphism.

In particular, this gives

$$\begin{aligned} f_2(a \otimes b) + m_2(b \otimes a) &= 0, \\ f_3(a \otimes b \otimes c) + m_3(a \otimes c \otimes b) + m_3(c \otimes a \otimes b) &= 0, \\ \dots \end{aligned}$$

i.e., the maps  $f_i$  vanish on shuffles.

In particular, for the operation  $m_2$  we have  $m_2(a \otimes b \pm b \otimes a) = 0$ , so a  $C_\infty$ -algebra of type  $(M, \{m_1, m_2, 0, 0, \dots\})$  is a commutative dg algebra (cdga) with the differential  $m_1$  and strictly associative and commutative multiplication  $m_2$ . Thus the category of commutative dg algebras is a subcategory of the category of  $C_\infty$ -algebras.

### 5.3.3 Bar interpretation of a $C_\infty$ -algebra

The notion of  $C_\infty$ -algebra is motivated by the following observation. It is well known that if a dg algebra  $(A, d, \mu)$  is graded commutative then the differential of the bar construction  $BA$  is not only a coderivation but also a derivation with respect to the shuffle product, so the bar construction  $(BA, d_\beta, \Delta, \mu_{sh})$  of a cdga is a dg bialgebra, see (3.3.3).

By definition the bar construction of an  $A_\infty$ -algebra  $(M, \{m_i\})$  is a dg coalgebra  $\tilde{B}(M) = (T^c(s^{-1}M), d_\beta, \Delta)$ .

But if  $(M, \{m_i\})$  is a  $C_\infty$ -algebra, then  $\tilde{B}(M)$  becomes a dg bialgebra:

**Proposition 18** . For an  $A_\infty$ -algebra  $(M, \{m_i\})$  the differential of the bar construction  $d_\beta$  is a derivation with respect to the shuffle product if and only if each operation  $m_i$  vanishes on shuffles, that is  $(M, \{m_i\})$  is a  $C_\infty$ -algebra.

**Proof.** The map  $\Phi : T^c(s^{-1}M) \otimes T^c(s^{-1}M) \rightarrow T^c(s^{-1}M)$  defined by  $\Phi = d_\beta \mu_{sh} - \mu_{sh}(d_\beta \otimes id + id \otimes d_\beta)$  is a coderivation, see the arguments in (2.2.1). Thus, according to the universal property of the tensor coalgebra (3.1.6) the map  $\Phi$  is trivial if and only if  $p_1 \Phi = 0$ , and the last condition means exactly (20).

### 5.3.4 Bar interpretation of a morphism of $C_\infty$ -algebras

A morphism of  $C_\infty$  algebras has an analogous interpretation.

**Proposition 19** . Let  $\{f_i\} : (M, \{m_i\}) \rightarrow (M', \{m'_i\})$  be an  $A_\infty$ -algebra morphism of  $C_\infty$ -algebras. Then the induced map of bar constructions  $\tilde{B}\{f_i\}$  is a map of dg bialgebras if and only if each  $f_i$  vanishes on shuffles, that is  $\{f_i\}$  is a morphism of  $C_\infty$ -algebras.

**Proof.** The map  $\Psi = \tilde{B}\{f_i\} \mu_{sh} - \mu_{sh}(\tilde{B}\{f_i\} \otimes \tilde{B}\{f_i\})$  is a coderivation, see the arguments in (2.2.1). Thus, according to the universal property of the tensor coalgebra (3.1.6) the map  $\Psi$  is trivial if and only if  $p_1 \Psi = 0$ , and the last condition means exactly (21).

Thus the bar functor maps the subcategory of  $C_\infty$ -algebras to the category of dg bialgebras.

### 5.3.5 Adjunctions

The bar and cobar functors

$$B : DGAlg \rightarrow DGCoalg, \quad \Omega : DGCoalg \rightarrow DGAlg$$

are adjoint and there exist standard weak equivalences  $\Omega B(A) \rightarrow A$ ,  $C \rightarrow B\Omega C$ . So  $\Omega B(A) \rightarrow A$  is a *free resolution* of a dga  $A$ .

If  $A$  is commutative, the cobar-bar resolution is out of the category:  $\Omega B(A)$  is not commutative.

In this case instead the cobar-bar functors we must use the adjoint functors  $\Gamma, \mathcal{A}$ , see [33], which we describe now.

For a commutative dg algebra the bar construction is a dg bialgebra, so the restriction of the bar construction is the functor  $B : CDGAlg \rightarrow$

*DGBialg*. Furthermore, the functor of indecomposables  $Q : DGBialg \rightarrow DGLieCoalg$  maps the category of dg bialgebras to the category of dg Lie coalgebras. Let  $\Gamma$  be the composition

$$\Gamma : CDGAlg \xrightarrow{B} DGBialg \xrightarrow{Q} DGLieCoalg.$$

There is the adjoint of  $\Gamma$   $\mathcal{A} : DGLieCoalg \rightarrow CDGAlg$ , which is dual to the Chevalley-Eilenberg functor. There is the standard weak equivalence  $\mathcal{A}\Gamma A \rightarrow A$ . This is the commutative analogue of the standard weak equivalence  $\Omega B(A) \rightarrow A$  from (3.3.4).

## 6 Minimal $A_\infty$ and $C_\infty$ -Algebras and Hochschild and Harrison Cohomology

Here we present the connection of the notion of minimal  $A_\infty$  (resp.  $C_\infty$ ) algebra with the Hochschild (resp. Harrison) cochain complexes, studied in [21], see also [26].

### 6.1 Hochschild Cohomology

Below we describe the classical Hochschild cochain complex [15] and present also some additional structures on it which will be essential to describe minimal  $A_\infty$  algebra structures.

#### 6.1.1 Hochschild Cochain Complex

Let  $A$  be an algebra and let  $M$  be a bimodule over  $A$ . The Hochschild cochain complex is defined as

$$C^*(A, M) = \sum C^n(A, M), \quad C^n(A, M) = Hom(\otimes^n A, M),$$

the coboundary operator  $\delta : C^n(A, M) \rightarrow C^{n+1}(A, M)$  is given by

$$\begin{aligned} \delta f(a_1 \otimes \dots \otimes a_{n+1}) &= a_1 \cdot f(a_2 \otimes \dots \otimes a_{n+1}) \\ &+ \sum_k \pm f(a_1 \otimes \dots \otimes a_k \cdot a_{k+1} \otimes \dots \otimes a_{n+1}) \pm f(a_1 \otimes \dots \otimes a_n) \cdot a_{n+1}. \end{aligned}$$

The Hochschild cohomology of  $A$  with coefficients in  $M$  is defined as the homology of this cochain complex and is denoted by  $Hoch^*(A, M)$ .

If  $A$  and  $M$  are graded then each  $C^n(A, M)$  is graded too:  $C^n(A, M) = \sum_k C^{n,k}(A, M)$  where  $C^{n,k}(A, M) = \text{Hom}^k(\otimes^n A, M)$ ; here  $\text{Hom}^k$  means homomorphisms of degree  $k$ . So the Hochschild cochain complex is bigraded in this case.

It is clear that  $\delta$  maps  $C^{n,k}(A, M)$  to  $C^{n+1,k}(A, M)$ , so  $(C^{*,k}(A, M), \delta)$  is a subcomplex in  $(C^*(A, M), \delta)$ . Thus Hochschild cohomology in this case is bigraded:  $Hoch^n(A, M) = \sum_k Hoch^{n,k}(A, M)$ , where  $Hoch^{n,k}(A, M)$  is the  $n$ -th homology module of  $(C^{*,k}(A, M), \delta)$ .

For our needs instead of  $M$  we take the algebra  $A$  itself. The complex  $C^{*,*}(A, A)$  is a bigraded differential algebra with respect to the following cup product: for  $f \in C^{m,k}(A, A)$  and  $g \in C^{n,t}(A, A)$  the product  $f \smile g \in C^{m+n,k+t}(A, A)$  is defined by

$$f \smile g(a_1 \otimes \dots \otimes a_{m+n}) = f(a_1 \otimes \dots \otimes a_m) \cdot g(a_{m+1} \otimes \dots \otimes a_{m+n}),$$

and the Hochschild differential is a derivative:

$$\delta(f \smile g) = \delta f \smile g \pm f \smile \delta g. \quad (22)$$

### 6.1.2 Gerstenhabers circle product

Besides this product  $C^{*,*}(A, A)$  there exist much richer and important algebraic operations, finally forming a hGa (homotopy Gerstenhaber algebra) structure. First of all, there is Gerstenhaber's so called circle product [8]  $f \circ g$ , sometimes called Gerstenhabers brace  $f\{g\}$ , but let us denote it as  $f \smile_1 g$  since it has properties very similar to Steenrod's  $\smile_1$  product in the cochain complex of a topological space. Here is the definition of this Gerstenhabers product: for  $f \in C^{m,s}(A, A)$ ,  $g \in C^{n,t}(A, A)$  their  $\smile_1$  product is defined as

$$f \smile_1 g(a_1 \otimes \dots \otimes a_{m+n-1}) = \sum_{k=0}^{m-1} \pm f(a_1 \otimes \dots \otimes a_k \otimes g(a_{k+1} \otimes \dots \otimes a_{k+n}) \otimes \dots \otimes a_{m+n-1}) \in C^{m+n-1,s+t}(A, A). \quad (23)$$

In [21], see also [26], it is shown that Gerstenhabers circle = brace =  $\smile_1$  product satisfies Steenrod's condition

$$\delta(f \smile_1 g) = \delta f \smile_1 g \pm f \smile_1 \delta g \pm f \smile g \pm g \smile f. \quad (24)$$

What is more amazing,  $\smile_1$  also has the following property:

$$(f \smile g) \smile_1 h = f \smile (g \smile_1 h) \pm (f \smile_1 h) \smile g, \quad (25)$$

which means that  $\dots \smile_1 h$  is a derivation. This is an analogue of Hirsch's formula in the cochain complex of a topological space.

Besides, although the  $\smile_1$  is not associative, it is possible to show that it satisfies the so called pre-Jacobi identity

$$f \smile_1 (g \smile_1 h) - (f \smile_1 g) \smile_1 h = f \smile_1 (h \smile_1 g) - (f \smile_1 h) \smile_1 g$$

which guarantees that the commutator

$$[f, g] = f \smile_1 g - g \smile_1 f$$

satisfies the Jacobi identity. Besides, the condition (24) implies that  $[\smile_1, \smile_1] : C^{m,s}(A, A) \otimes C^{n,t}(A, A) \rightarrow C^{m+n-1, s+t}(A, A)$  is a chain map, and this implies that  $(C^{*,*}(A, A), \delta, [\smile_1, \smile_1])$  is a dg Lie algebra.

By the way, the Hochschild differential  $\delta$  can be expressed as

$$\delta f = f \smile_1 \mu + \mu \smile_1 f = [f, \mu]$$

where  $\mu : A \otimes A \rightarrow A$  is the product operation of  $A$ .

### 6.1.3 Higher operations in the Hochschild complex

In order to perturb  $\smile_1$ -twisting cochains (i.e., minimal  $A_\infty$  algebra structures, as we'll see below) we need a kind of analog of the Berikishvili equivalence of Brown's twisting cochains, but now for this nonassociative  $\smile_1$  case. Unfortunately for this only Gerstenhaber's  $\circ = \smile_1$  is not enough, so in [21] we have introduced higher order multioperations, the  $\smile_1$  products  $f \smile_1 (g_1, \dots, g_i)$  of one Hochschild cochain  $f$  and a collection of Hochschild cochains  $g_1, \dots, g_i$ . Now these operations are called Getzler-Kadeishvili brace operations and are denoted as  $f\{g_1, \dots, g_i\}$ . Here is the definition:

$$\begin{aligned} f \smile_1 (g_1, \dots, g_i)(a_1 \otimes \dots \otimes a_{n+n_1+\dots+n_i-i}) = \\ \sum_{k_1, \dots, k_i} f(a_1 \otimes \dots \otimes a_{k_1} \otimes g_1(a_{k_1+1} \otimes \dots \otimes a_{k_1+n_1}) \otimes \\ \dots \otimes a_{k_2} \otimes g_2(a_{k_2+1} \otimes \dots \otimes a_{k_2+n_2}) \otimes a_{k_2+n_2+1} \otimes \dots \\ \otimes a_{k_i} \otimes g_i(a_{k_i+1} \otimes \dots \otimes a_{k_i+n_i}) \otimes \dots \otimes a_{n+n_1+\dots+n_i-i}). \end{aligned} \quad (26)$$

These higher braces  $f\{g_1, \dots, g_i\}$  form on the Hochschild cochain complex  $C^*(A, A)$  a so called homotopy Gerstenhaber algebra (hGa) structure.

### 6.1.4 Hochschild cochain complex as a Homotopy Gerstenhaber Algebra

Let  $H$  be a graded algebra. Consider the Hochschild cochain complex of this graded algebra with coefficients in itself,  $C^{*,*}(H, H)$  which is bigraded in this case:

$$C^{n,m}(H, H) = \text{Hom}^m(H^{\otimes n}, H),$$

where  $\text{Hom}^m$  means homomorphisms of degree  $m$ .

This bigraded complex carries a structure of homotopy Gerstenhaber algebra, see [21], [11], [10], [26]  $C^{*,*}(H, H), \delta, \smile, \{E_{1,k}, k = 1, 2, \dots\}$ , which consists of the following structure maps:

(i) The Hochschild differential  $\delta : C^{n-1,m}(H, H) \rightarrow C^{n,m}(H, H)$  given by

$$\begin{aligned} \delta f(a_1 \otimes \dots \otimes a_n) = & a_1 \cdot f(a_2 \otimes \dots \otimes a_n) \\ & + \sum_k \pm f(a_1 \otimes \dots \otimes a_{k-1} \otimes a_k \cdot a_{k+1} \otimes \dots \otimes a_n) \\ & \pm f(a_1 \otimes \dots \otimes a_{n-1}) \cdot a_n; \end{aligned}$$

(ii) The  $\smile$  product defined by

$$f \smile g(a_1 \otimes \dots \otimes a_{n+m}) = f(a_1 \otimes \dots \otimes a_n) \cdot g(a_{n+1} \otimes \dots \otimes a_{n+m}).$$

(iii) The *brace* operations  $f\{g_1, \dots, g_i\}$  from [21], which we write here as

$$f\{g_1, \dots, g_i\} = E_{1,i}(f; g_1, \dots, g_i),$$

$$E_{1,i} : C^{n,m} \otimes C^{m_1,m_1} \otimes \dots \otimes C^{m_i,m_i} \rightarrow C^{m+\sum n_t-i, m+\sum m_t},$$

given by

$$\begin{aligned} & E_{1,i}(f; g_1, \dots, g_i)(a_1 \otimes \dots \otimes a_{n+n_1+\dots+n_i-i}) \\ & = \sum_{k_1, \dots, k_i} \pm f(a_1 \otimes \dots \otimes a_{k_1} \otimes g_1(a_{k_1+1} \otimes \dots \otimes a_{k_1+n_1}) \otimes \\ & \quad \dots \otimes a_{k_2} \otimes g_2(a_{k_2+1} \otimes \dots \otimes a_{k_2+n_2}) \otimes a_{k_2+n_2+1} \otimes \dots \\ & \quad \otimes a_{k_i} \otimes g_i(a_{k_i+1} \otimes \dots \otimes a_{k_i+n_i}) \otimes \dots \otimes a_{n+n_1+\dots+n_i-i}). \end{aligned} \tag{27}$$

The first brace operation  $E_{1,1}$  has the properties of Steenrod's  $\smile_1$  product, so we use the notation  $E_{1,1}(f, g) = f \smile_1 g$ . In fact this is Gerstenhaber's  $f \circ g$  product [8], [9].

The name *Homotopy G-algebra* is motivated by the fact that this structure induces on homology  $H(A)$  the structure of Gerstenhaber algebra (G-algebra).

The sequence  $\{E_{1,k}\}$  defines a twisting cochain

$$E : BC^{*,*}(H, H) \otimes BC^{*,*}(H, H) \rightarrow C^{*,*}(H, H),$$

and, consequently, defines a strictly associative product on the bar construction  $BC^{*,*}(H, H)$

$$\mu_E : BC^{*,*}(H, H) \otimes BC^{*,*}(H, H) \rightarrow BC^{*,*}(H, H)$$

which turns it into a DG-bialgebra.

## 6.2 Description of minimal $A_\infty$ -algebra structure as a twisting cochain in Hochschild complex

Here we present the connection of the notion of minimal  $A_\infty$ -algebra with Hochschild cochain complexes, studied in [21], see also [26].

### 6.2.1 Minimal $A_\infty$ structure as a Hochschild twisting cochain

Now let

$$(H, \{m_1 = 0, m_2, m_3, \dots, m_n, \dots\})$$

be a minimal  $A_\infty$ -algebra. So  $(H, m_2)$  is an associative graded algebra with multiplication  $a \cdot b = m_2(a \otimes b)$  and we can consider the Hochschild cochain complex  $C^{*,*}(H, H)$ .

Each operation  $m_i$  can be considered as a Hochschild cochain  $m_i \in C^{i, 2-i}(H, H)$ . Let  $m = m_3 + m_4 + \dots \in C^{*, 2-*}(H, H)$ . Stasheff's defining condition of  $A_\infty$ -algebra (13)

$$\sum_{k=0}^{n-1} \sum_{j=1}^{n-k} m_{n-j+1}(a_1 \otimes \dots \otimes a_k \otimes m_j(a_{k+1} \otimes \dots \otimes a_{k+j}) \otimes \dots \otimes a_n) = 0. \quad (28)$$

in our case looks like this:

$$\begin{aligned} & m_2(a_1 \otimes m_{n-1}(a_2 \otimes \dots \otimes a_n)) + m_2(m_{n-1}(a_1 \otimes \dots \otimes a_{n-1}) \otimes a_n) \\ & + \sum_k m_{n-1}(a_1 \otimes \dots \otimes a_k \otimes m_2(a_{k+1} \otimes a_{k+2}) \otimes a_{k+3} \otimes \dots \otimes a_n) \\ & = \sum_{k=0}^{n-1} \sum_{j=3}^{n-k} m_{n-j+1}(a_1 \otimes \dots \otimes a_k \otimes m_j(a_{k+1} \otimes \dots \otimes a_{k+j}) \otimes \dots \otimes a_n). \end{aligned}$$

This can be rewritten as

$$\delta m_n = \sum_{j=3, \dots, n-2} m_{n-j+1} \smile_1 m_j.$$

And this means exactly that  $\delta m = m \smile_1 m$  for  $m = m_3 + m_4 + \dots$ ,  $m_i \in C^{i,2-i}(H, H)$ .

So a minimal  $A_\infty$ -algebra structure on  $H$  is, in fact, a *twisting cochain*

$$m = m_3 + m_4 + \dots \in C^{*,2-*}(H, H)$$

in the Hochschild complex with respect to the (nonassociative)  $\smile_1$  product.

### 6.2.2 Perturbations of minimal $A_\infty$ algebras

Now let  $(H, \{m_i\})$  be a minimal  $A_\infty$ -algebra, so  $(H, m_2)$  is an associative graded algebra with multiplication  $a \cdot b = m_2(a \otimes b)$ .

As it was mentioned in section 6.2.1, each operation  $m_i$  can be considered as a Hochschild cochain  $m_i \in C^{i,2-i}(H, H)$ . Let  $m = m_3 + m_4 + \dots \in C^{*,2-*}(H, H)$ . The defining condition of an  $A_\infty$ -algebra (13) means exactly  $\delta m = m \smile_1 m$ . So a minimal  $A_\infty$ -algebra structure on  $H$  is, in fact, a *twisting cochain* in the Hochschild complex with respect to the  $\smile_1$  product.

There is the notion of equivalence of such twisting cochains:  $m \sim m'$  if there exists  $p = p^{2,-1} + p^{3,-2} + \dots + p^{i,1-i} + \dots$ ,  $p^{i,1-i} \in C^{i,1-i}(H, H)$  such that

$$\begin{aligned} m - m' &= \delta p + p \smile p + p \smile_1 m + \\ m' \smile_1 p &+ E_{1,2}(m'; p, p) + E_{1,3}(m'; p, p, p) + \dots \end{aligned} \quad (29)$$

**Proposition 20** . *Twisting cochains  $m, m' \in C^{*,2-*}(H, H)$  are equivalent if and only if  $(H, \{m_i\})$  and  $(H', \{m'_i\})$  are isomorphic  $A_\infty$ -algebras.*

**Proof.** Indeed,

$$\{p_i\} : (H, \{m_i\}) \rightarrow (H, \{m'_i\})$$

with  $p_1 = id$ ,  $p_i = p^{i,1-i}$  is the needed isomorphism: the condition (29) coincides with the defining condition (7) of a morphism of  $A_\infty$ -algebras and the Proposition 14 implies that this morphism is an isomorphism.

This gives the possibility of perturbing twisting cochains without changing their equivalence class:

**Proposition 21** . *Let  $m$  be a twisting cochain (i.e., a minimal  $A_\infty$ -algebra structure on  $H$ ) and let  $p \in C^{n,1-n}(H, H)$  be an arbitrary cochain. Then there exists a twisting cochain  $\bar{m}$ , equivalent to  $m$ , such that  $m_i = \bar{m}_i$  for  $i \leq n$  and  $\bar{m}_{n+1} = m_{n+1} + \delta p$ .*

**Proof.** The twisting cochain  $\bar{m}$  can be solved inductively from the equation (29).

**Theorem 9 .** *Suppose for a graded algebra  $H$  that its Hochschild cohomology groups are trivial,  $Hoch^{n,2-n}(H, H) = 0$  for  $n \geq 3$ . Then each  $m \sim 0$ , that is each minimal  $A_\infty$ -algebra structure on  $H$  is degenerate, i.e.,  $(H, \{m_i\})$  is isomorphic to a (strictly associative)  $A_\infty$  algebra  $(H, \{m_1 = 0, m_2, m_3 = 0, \dots\})$ .*

**Proof.** From the equality  $\delta m = m \smile_1 m$  in dimension 4 we obtain  $\delta m_3 = 0$ , that is,  $m_3$  is a cocycle. Since  $Hoch^{3,-1}(H, H) = 0$  there exists  $p^{2,-1}$  such that  $m_3 = \delta p^{2,-1}$ . Perturbing our twisting cochain  $m$  by  $p^{2,-1}$  we obtain a new twisting cochain  $\bar{m} = \bar{m}_3 + \bar{m}_4 + \dots$  equivalent to  $m$  and with  $\bar{m}_3 = 0$ . Now the component  $\bar{m}_4$  becomes a cocycle, which can be killed using  $Hoch^{4,-2}(H, H) = 0$ , etc.

### 6.2.3 Minimal $C_\infty$ -algebra structure and Harrison cohomology

Suppose now  $(H, \mu)$  is a commutative graded algebra. The Harrison cochain complex  $\bar{C}^*(H, H)$  is defined as a subcomplex of the Hochschild complex consisting of cochains which disappear on shuffles. If  $(H, \{m_i\})$  is a  $C_\infty$ -algebra then the twisting element  $m = m_3 + m_4 + \dots$  belongs to Harrison subcomplex  $\bar{C}^*(H, H) \subset C^*(H, H)$  and we have the

**Theorem 10 .** *Suppose for a graded commutative algebra  $H$  Harrison cohomology  $Harr^{n,2-n}(H, H) = 0$  for  $n \geq 3$ . Then each  $m \sim 0$ , that is each minimal  $C_\infty$ -algebra structure on  $H$  is degenerate.*

## 7 Minimality theorem, $A_\infty$ -algebra structure in homology

### 7.1 $A_\infty$ -algebra structure in homology

Let  $(A, d, \mu)$  be a dg algebra and  $(H(A), \mu^*)$  be its homology algebra.

Although the dg algebras  $(A, d, \mu)$  and  $(H(A), d = 0, \mu^*)$  have same homology algebras, the homology algebra  $H(A)$  carries less information than the dg algebra  $A$ .

Generally, there does not exist a map of dg algebras  $H(A) \rightarrow A$  which induces an isomorphism of homology algebras. Of course, stepping from  $A$  to the smaller object  $H(A)$  one loses part of the information. To compensate this loss, it is natural to enrich the algebraic structure on the smaller object  $H(A)$ . The classical examples of such enrichments are Steenrod squares, Massey products, ... .

### 7.1.1 Minimality Theorem

Below we present one sort of such additional algebraic structure, namely  $A_\infty$ -algebra structure on the cohomology algebra, the so called minimality theorem.

It was mentioned above (5.1.1) Stasheff's  $A_\infty$ -algebras are sort of Strong Homotopy Associative Algebras, the operation  $m_3$  is a homotopy which measures the nonassociativity of the product  $m_2$ . So its existence on homology the  $H(A)$ , which is strictly associative looks a bit strange, but although the product on  $H(A)$  is associative, there appears a structure of a (generally nondegenerate) *minimal*  $A_\infty$ -algebra, which can be considered as an  $A_\infty$  *deformation* of  $(H(A), \mu^*)$ , [26] . Namely, in [16], [17] the following *minimality theorem* was proved:

**Theorem 11** . *Suppose that for a dg algebra  $(A, d, \mu)$  all homology modules  $H_i(A)$  are free.*

(a) *Then there exist a structure of minimal  $A_\infty$ -algebra on  $H(A)$*

$$(H(A), \{m_1 = 0, m_2 = \mu^*, m_3, \dots, m_i\})$$

*and a weak equivalence of  $A_\infty$ -algebras*

$$\{f_i\} : (H(A), \{m_i\}) \rightarrow (A, \{d, \mu, 0, 0, \dots\})$$

*such that  $m_1 = 0, m_2 = \mu^*, f_1^* = id_{H(A)}$ .*

(b) *Furthermore, for a dga map  $g : A \rightarrow A'$  there exists a morphism of  $A_\infty$ -algebras  $\{g_i\} : (H(A), \{m_i\}) \rightarrow (H(A'), \{m'_i\})$  with  $g_1 = g^*$  and such that the diagram*

$$\begin{array}{ccc} (H(A), \{m_i\}) & \xrightarrow{\{g_i\}} & (H(A'), \{m'_i\}) \\ \{f_i\} \downarrow & & \downarrow \{f'_i\} \\ A & \xrightarrow{g} & A' \end{array}$$

commutes up to homotopy in the category of  $A_\infty$  algebras.

(c) Such a structure is unique up to isomorphism in the category of  $A_\infty$ -algebras: if  $(H(A), \{m_i\})$  and  $(H(A), \{m'_i\})$  are two minimal  $A_\infty$ -algebra structures on  $H(A)$  which satisfy the conditions from (a) then there exists an isomorphism of  $A_\infty$ -algebras  $\{g_i\} : (H(A), \{m_i\}) \rightarrow (H(A), \{m'_i\})$  with  $g_1 = id$ .

**Proof of (a).** We are going to construct the components  $f_i$ ,  $m_i$  inductively satisfying the defining condition of an  $A_\infty$ -morphism

$$\begin{aligned} df_n(a_1 \otimes \dots \otimes a_n) &= f_1 m_n(a_1 \otimes \dots \otimes a_n) \\ &+ \sum_{j=2}^{n-1} \sum_{k=0}^{n-j} f_{n-j+1}(a_1 \otimes \dots \otimes a_k \otimes m_j(a_{k+1} \otimes \dots \otimes a_{k+j}) \otimes \dots \otimes a_n) \\ &+ \sum_{k=1}^{n-1} f_k(a_1 \otimes \dots \otimes a_k) \cdot f_{n-k}(a_{k+1} \otimes \dots \otimes a_n). \end{aligned} \tag{30}$$

Let us start with a cycle-choosing homomorphism  $f_1 : H(A) \rightarrow A$ , that is  $f_1(a) \in a \in H(A)$  which can be constructed using the freeness of the modules  $H_i(A)$ . This map is not multiplicative but  $f_1(a_1 \cdot a_2) - f_1(a_1) \cdot f_1(a_2) \sim 0 \in A$ . So there exists  $f_2 : H(A) \otimes H(A) \rightarrow A$  s.t.  $f_1(a_1 \cdot a_2) - f_1(a_1) \cdot f_1(a_2) = df_2(a_1 \otimes a_2)$ , and this, assuming  $m_2(a_1 \otimes a_2) = a_1 \cdot a_2$ , is exactly the condition (30) for  $n = 2$ .

Let us denote

$$\begin{aligned} U_3(a_1 \otimes a_2 \otimes a_3) &= \\ f_1(a_1) \cdot f_2(a_2 \otimes a_3) &+ f_2(a_1 \cdot a_2 \otimes a_3) + f_2(a_1 \otimes a_2 \cdot a_3) + f_2(a_1 \otimes a_2) \cdot f_1(a_3). \end{aligned}$$

Note that the main condition (30) for  $n = 3$  looks as follows

$$df_3(a_1 \otimes a_2 \otimes a_3) = f_1 m_3(a_1 \otimes a_2 \otimes a_3) - U_3(a_1 \otimes a_2 \otimes a_3).$$

Direct calculation shows that  $dU_3(a_1 \otimes a_2 \otimes a_3) = 0$ , so  $U_3(a_1 \otimes a_2 \otimes a_3)$  is a cycle. Let us define  $m_3(a \otimes b \otimes c) \in H(A)$  as the homology class of this cycle

$$\begin{aligned} m_3(a \otimes b \otimes c) &= \{U_3(a \otimes b \otimes c)\} = \\ f_1(a_1) \cdot f_2(a_2 \otimes a_3) &+ f_2(a_1 \cdot a_2 \otimes a_3) + f_2(a_1 \otimes a_2 \cdot a_3) + f_2(a_1 \otimes a_2) \cdot f_1(a_3). \end{aligned} \tag{31}$$

Then, since  $f_1$  is a *cycle choosing* homomorphism,

$$f_1 m_3(a_1 \otimes a_2 \otimes a_3) - U_3(a_1 \otimes a_2 \otimes a_3)$$

is homological to zero. Thus, again using the freeness of  $H_i(A)$ , it is possible to construct a homomorphism  $f_3 : H(A) \otimes H(A) \otimes H(A) \rightarrow A$  such that

$$df_3(a_1 \otimes a_2 \otimes a_3) = f_1 m_3(a_1 \otimes a_2 \otimes a_3) - U_3(a_1 \otimes a_2 \otimes a_3),$$

and this is exactly the condition (30) for  $n = 3$ .

Assume now that  $f_i, m_i, i \leq n - 1$  are already constructed and they satisfy (30).

Let us denote

$$\begin{aligned} & U_n(a_1 \otimes \dots \otimes a_n) \\ = & \sum_{j=2}^{n-1} \sum_{k=0}^{n-j} f_{n-j+1}(a_1 \otimes \dots \otimes a_k \otimes m_j(a_{k+1} \otimes \dots \otimes a_{k+j})) \otimes \dots \otimes a_n \\ & + \sum_{k=1}^{n-1} f_k(a_1 \otimes \dots \otimes a_k) \cdot f_{n-k}(a_{k+1} \otimes \dots \otimes a_n). \end{aligned}$$

Then the defining condition (30) can be rewritten as

$$df_n(a_1 \otimes \dots \otimes a_n) = f_1 m_n(a_1 \otimes \dots \otimes a_n) + U_n(a_1 \otimes \dots \otimes a_n). \quad (32)$$

Direct calculation shows that  $dU_n = 0$ , so we define  $m_n(a_1 \otimes \dots \otimes a_n)$  as the homology class of the cycle  $U_n(a_1 \otimes \dots \otimes a_n)$ . Then using the freeness of modules the  $H_i(A)$  one can construct a homomorphism  $f_n : H_*(A) \otimes \dots \otimes H(A) \rightarrow A$  satisfying (32), and this completes the proof of (a).

**Proof of (b).** The morphism of  $A_\infty$ -algebras  $(H(A), \{m_i\}) \xrightarrow{\{f_i\}} A \xrightarrow{g} A'$  induces a twisting cochain  $\phi : B(H(A), \{m_i\}) \rightarrow A$ .

Furthermore, since

$$(H(A'), \{m'_i\}) \xrightarrow{\{f'_i\}} A'$$

is a weak equivalence of  $A_\infty$ -algebras, by the lifting theorem (8)  $\phi$  can be lifted to the  $A_\infty$ -twisting cochain  $\psi : B(H(A), \{m_i\}) \rightarrow (H(A'), \{m'_i\})$  which in fact represents an  $A_\infty$ -morphism  $\{g_i\} : (H(A), \{m_i\}) \rightarrow (H(A'), \{m'_i\})$ , such that  $\{f'_i\} \circ \psi \sim \phi$ . This equivalence means that  $A_\infty$ -morphisms  $\{f'_i\} \circ \{g_i\}$  and  $g \circ \{g_i\}$  are homotopic. This completes the proof.

**Proof of (c).** Suppose for a dg algebra  $(A, d, \mu)$  we have two homology  $A_\infty$ -algebras  $(H(A), \{m_i\})$  and  $(H(A), \{m'_i\})$ , i.e., there exist  $A_\infty$  weak equivalences  $\{f_i\} : (H(A), \{m_i\}) \rightarrow A, \{f'_i\} : (H(A), \{m_i\}) \rightarrow A$  with  $(f_1)^* = (f'_1)^* = id_{H(A)}$ . Then, using the part (b) for  $d = id_A$  we obtain  $\{g_i\} : (H(A), \{m_i\}) \rightarrow (H(A), \{m'_i\})$  with  $g_1 = g^* = id_{H(A)}$ . So,  $\{g_i\}$  is an isomorphism of  $A_\infty$  algebras by (13).

**Corollary 1 .** *The mapping of differential coalgebras*

$$B(\{f_i\}) : \tilde{B}(H(A), \{m_i\}) \rightarrow B(A)$$

*induces an isomorphism in homology.*

### 7.1.2 Connection with Massey Products.

The first new operation  $m_3(a_1 \otimes a_2 \otimes a_3) \in H(A)$  was defined as the homology class of the cycle

$$U_3(a_1 \otimes a_2 \otimes a_3) = f_1(a_1) \cdot f_2(a_2 \otimes a_3) + f_2(a_1 \cdot a_2 \otimes a_3) + f_2(a_1 \otimes a_2 \cdot a_3) + f_2(a_1 \otimes a_2) \cdot f_1(a_3).$$

This description immediately induces the connection of  $m_3$  with the Massey product: If  $a, b, c \in H(A)$  is a Massey triple, i.e., if  $a_1 \cdot a_2 = a_2 \cdot a_3 = 0$ , then

$$U_3(a_1 \otimes a_2 \otimes a_3) = f_1(a_1) \cdot f_2(a_2 \otimes a_3) + f_2(a_1 \otimes b_2) \cdot f_1(a_3)$$

and this is exactly the combination which defines the Massey product  $\langle a_1, a_2, a_3 \rangle$ , thus  $m_3(a_1 \otimes a_2 \otimes a_3) \in \langle a_1, a_2, a_3 \rangle$ . This gives examples of dg algebras with essentially nontrivial homology  $A_\infty$ -algebras.

### 7.1.3 Special cases

Taking  $(A, d, \mu) = C^*(X)$ , the cochain complex of a topological space  $X$ , the theorem defines on the cohomology algebra  $H^*(X)$  the structure of a minimal  $A_\infty$ -algebra  $(H^*(X), \{m_i\})$ , which carries more information about  $X$  than just the graded algebra structure. In particular, we shall show later that this cohomology  $A_\infty$ -algebra  $(H^*(X), \{m_i\})$  determines the cohomology modules of the loop space  $\Omega X$  whereas the bare the cohomology algebra  $(H^*(X), m_2)$  does not.

Furthermore, taking  $(A, d, \mu) = C_*(G)$ , the chain complex of a topological group, the theorem defines on the Pontriagin ring  $H_*(G)$  the structure of a minimal  $A_\infty$ -algebra  $(H_*(G), \{m_i\})$  which will carry more information about  $G$  than just the ring structure. In particular, we shall show later that this homology  $A_\infty$ -algebra  $(H_*(G), \{m_i\})$  determines homology modules of classifying space  $B_G$  whereas the bare Pontriagin ring  $(H_*(G), m_2)$  does not.

### 7.1.4 Minimal $A_\infty$ -algebra structure on the homology of an $A_\infty$ -algebra

The Minimality Theorem is true also when, instead of a dg algebra  $(A, d, \mu)$  we take an arbitrary  $A(\infty)$ -algebra  $(M, \{m_i\})$ .

In this case, minimal  $A_\infty$ -algebra structure appears on the homology  $H(M)$  of the dg module  $(M, m_1 : M \rightarrow M)$ , see [18].

This structure had applications in string theory, see for example [1].

## 8 Applications of the Minimality Theorem

### 8.1 Application: Cohomology $A_\infty$ algebra of a space and Cohomology modules of Loop Space.

Taking  $A = C^*(X)$ , the cochain dg algebra of a 1-connected space  $X$ , we obtain an  $A_\infty$ -algebra structure  $(H^*(X), \{m_i\})$  on the cohomology algebra  $H^*(X)$ .

The cohomology algebra equipped with this additional structure carries more information than just the cohomology algebra. Some applications of this structure are given in [17] , [23]. For example the cohomology  $A_\infty$ -algebra  $(H^*(X), \{m_i\})$  determines the cohomology of the loop space  $H^*(\Omega X)$  whereas the bare algebra  $(H^*(X), m_2)$  does not:

**Theorem 12** .  $H(B(H^*(X), \{m_i\})) = H^*(\Omega X)$ .

Firstly, according to the minimality theorem the homology of the dg algebra  $(C^*(X), d, \smile)$  carries a structure of minimal  $A_\infty$ -algebra  $(H^*(X), \{m_i\})$ , such that there is and a weak equivalence of  $A_\infty$ -algebras

$$\{f_i\} : (H^*(X), \{m_i\}) \rightarrow (C^*(X), \{m_1 = d, m_2 = \smile, m_3 = 0, \dots\}),$$

which induces weak equivalences of their bar constructions and an isomorphism of graded modules

$$H(B(H^*(X), \{m_i\})) \approx H(BC^*(X)) \approx H^*(\Omega X).$$

Thus the object  $(H^*(X), \{m_i\})$ , which is called the cohomology  $A_\infty$ -algebra of  $X$ , determines the cohomology *modules* of the loop space  $H^*(\Omega X)$ . But not the cohomology *algebra*  $H^*(\Omega X)$ .

### 8.2 Application: Homology modules of the Classifying Space of a Topological Group.

Taking  $A = C_*(G)$ , the chain dg algebra of a topological group  $G$ , we obtain an  $A_\infty$ -algebra structure  $(H_*(G), \{m_i\})$  on the Pontryagin algebra  $H_*(G)$ . The homology  $A_\infty$ -algebra  $(H_*(G), \{m_i\})$  determines the homology of the classifying space  $H_*(B_G)$  when just the Pontryagin algebra  $(H_*(G), m_2)$  does not:

**Theorem 13** .  $H(B(H_*(G), \{m_i\})) = H_*(B_G)$ .

### 8.3 Application: $A_\infty$ -model of a Fibre Bundle

The minimality theorem (11) and the theorem (8) about lifting of twisting cochains allow to construct effective models of fibre bundles. Actually this model and higher operations  $\{m_i\}$  and  $\{p_i\}$  were constructed in [16]. Later we have recognized that they form Stasheff's  $A_\infty$ -structures and the model in these terms was presented in [17]. A similar model was also presented in [30].

**Topological level.** Let  $\xi = (X, p, B, G)$  be a principal  $G$ -fibration. Let  $F$  be a  $G$ -space. Then the action  $G \text{ times } F \rightarrow F$  determines the *associated fibre bundle*  $\xi(F) = (E, p, B, F, G)$  with fiber  $F$ . Thus  $\xi$  and the action  $G \text{ times } F \rightarrow F$  on the topological level determine  $E$ .

**Chain level.** Let  $K = C_*(B)$ ,  $A = C_*(G)$ ,  $P = C_*(F)$ . The classical result of E. Brown [5] states that the principal fibration  $\xi$  determines a twisting cochain  $\phi : K = C_*(B) \rightarrow A = C_*(G)$ , and the action on the chain level  $C_*(G) \otimes C_*(F) \rightarrow C_*(F)$  defines the twisted tensor product  $K \otimes_\phi P = C_*(B) \otimes_\phi C_*(F)$  which gives the homology modules of the total space  $H_*(E)$ . Thus  $\xi$  and the action on the chain level  $C_*(G) \otimes C_*(F) \rightarrow C_*(F)$  determine  $H_*(E)$ .

The twisting cochain  $\phi$  is not uniquely determined and it can be perturbed by the above equivalence relations for computational reasons.

**Homology level.** Nodar Berikashvili stated the problem to lift the previous "chain level" model of associated fibration to "homology level", i.e., to construct a "twisted differential" on  $C_*(B) \otimes H_*(F)$ . Investigation had shown that the principal fibration  $\xi$  and the action of of Pontryagin ring  $H_*(G)$  on  $H_*(F)$ , that is the pairing  $H_*(G) \otimes H_*(F) \rightarrow H_*(F)$  *do not determine*  $H_*(E)$ . But by the minimality theorem it appeared that  $H_*(G)$  carries not only the Pontryagin product  $H_*(G) \otimes H_*(G) \rightarrow H_*(G)$  but also richer algebraic structure, namely the structure of a minimal  $A_\infty$ -algebra  $(H_*(G), \{m_i\})$ .

Furthermore, the action  $G \text{ times } F \rightarrow F$  induces not only the pairing  $H_*(G) \otimes H_*(F) \rightarrow H_*(F)$  but, by the modular analog of the Minimality Theorem [17] also the structure of minimal  $A_\infty$ -module  $(H_*(F), \{p_i\})$ ,

$$p_i : H_*(G) \otimes \dots((i-1) \text{ times})\dots \otimes H_*(G) \otimes H_*(F) \rightarrow H_*(F),$$

and all these operations allow to define the correct differential on  $C_*(B) \otimes H_*(F)$ : there is a weak equivalence, homology isomorphism

$$C_*(B) \otimes_\psi H_*(F) = K \otimes_\psi H(P) \rightarrow K \otimes_\phi P = C_*(B) \otimes_\phi C_*(F) \sim C_*(E).$$

## 9 $C_\infty$ -algebra structure in the homology of a commutative dg algebra, applications in Rational Homotopy Theory

There is a commutative version of the above Minimality Theorem, see [22], [23], [28]:

**Theorem 14** . *Suppose that for a commutative dg algebra  $A$  all homology  $R$ -modules  $H^i(A)$  are free.*

*Then there exist: a structure of minimal  $C_\infty$ -algebra  $(H(A), \{m_i\})$  on  $H(A)$  and a weak equivalence of  $C_\infty$ -algebras*

$$\{f_i\} : (H(A), \{m_i\}) \rightarrow (A, \{d, \mu, 0, 0, \dots\})$$

*such that  $m_1 = 0$ ,  $m_2 = \mu^*$ ,  $f_1^* = id_{H(A)}$ .*

*Furthermore, for a cdga map  $f : A \rightarrow A'$  there exists a morphism of  $C_\infty$ -algebras  $\{f_i\} : (H(A), \{m_i\}) \rightarrow (H(A'), \{m'_i\})$  with  $f_1 = f^*$ .*

Such a structure is unique up to isomorphism in the category of  $C_\infty$ -algebras.

Below we present some applications of this  $C_\infty$ -algebra structure in rational homotopy theory.

### 9.1 Classification of rational homotopy types

Let  $X$  be a 1-connected space. In the case of rational coefficients there exist Sullivan's *commutative* cochain complex  $A(X)$  of  $X$ . It is well known that the weak equivalence type of a cdg algebra  $A(X)$  determines the rational homotopy type of  $X$ : 1-connected  $X$  and  $Y$  are *rationally homotopy equivalent* if and only if  $A(X)$  and  $A(Y)$  are weakly equivalent cdg algebras. Indeed, in this case  $A(X)$  and  $A(Y)$  have *isomorphic* minimal models  $M_X \approx M_Y$ , and this implies that  $X$  and  $Y$  are rationally homotopy equivalent. This is the key geometrical result of Sullivan which we are going to exploit below.

Now we take  $A = A(X)$  and apply Theorem 14. Then we obtain on  $H(A) = H^*(X, Q)$  a structure of minimal  $C_\infty$  algebra  $(H^*(X, Q), \{m_i\})$  which we call the *rational cohomology  $C_\infty$ -algebra of  $X$* .

Generally, an isomorphism of rational cohomology algebras  $H^*(X, Q)$  and  $H^*(Y, Q)$  does not imply a homotopy equivalence  $X \sim Y$ , not even rationally. We claim that  $(H^*(X, Q), \{m_i\})$  is a *complete* rational homotopy invariant:

**Theorem 15** . *1-connected  $X$  and  $X'$  are rationally homotopy equivalent if and only if  $(H^*(X, Q), \{m_i\})$  and  $(H^*(X', Q), \{m'_i\})$  are isomorphic as  $C_\infty$ -algebras.*

**Proof.** Suppose  $X \sim X'$ . Then  $A(X)$  and  $A(X')$  are weakly equivalent, that is there exists a cgda  $A$  and weak equivalences  $A(X) \leftarrow A \rightarrow A(X')$ . This implies weak equivalences of the corresponding homology  $C_\infty$ -algebras

$$(H^*(X, Q), \{m_i\}) \leftarrow (H^*(A), \{m_i\}) \rightarrow (H^*(X', Q), \{m'_i\}),$$

which by of minimality are both isomorphisms.

Conversely, suppose  $(H^*(X, Q), \{m_i\}) \approx (H^*(X', Q), \{m'_i\})$ . Then

$$\mathcal{AQB}(H^*(X, Q), \{m_i\}) \approx \mathcal{AQB}(H^*(X', Q), \{m'_i\}).$$

Denote this cdga as  $A$ . Then we have weak equivalences of CGD algebras (see (5.3.5))

$$A(X) \leftarrow \mathcal{A}\Gamma A(X) \leftarrow A \rightarrow \mathcal{A}\Gamma A(X') \rightarrow A(X').$$

This theorem in fact classifies rational homotopy types with given cohomology algebra  $H$  as all possible minimal  $C_\infty$ -algebra structures on  $H$  modulo  $C_\infty$  isomorphisms.

**Example.** Here we describe an example which we will use to illustrate the results of this and forthcoming sections.

We consider the following commutative graded algebra. Its underlying graded  $Q$ -vector space has the following generators: a generator  $e$  of dimension 0, generators  $x, y$  of dimension 2, and a generator  $z$  of dimension 5, so

$$H^* = \{H^0 = Q_e, 0, H^2 = Q_x \oplus Q_y, 0, 0, H^5 = Q_z, 0, 0, \dots\}, \quad (33)$$

and the multiplication is trivial by dimensional reasons, with unit  $e$ . In fact

$$H^* = H^*(S^2 \vee S^2 \vee S^5, Q).$$

This example was considered in [13]. It was shown there that there are just two rational homotopy types with such cohomology algebra.

The same result can be obtained from our classification.

What minimal  $C_\infty$ -algebra structures are possible on  $H^*$ ?

For dimensional reasons only one nontrivial operation  $m_3 : H^2 \otimes H^2 \otimes H^2 \rightarrow H^5$  is possible.

The specific condition of a  $C_\infty$ -algebra, namely the disappearance on shuffles implies that

$$m_3(x, x, x) = 0, \quad m_3(y, y, y) = 0, \quad m_3(x, y, x) = 0, \quad m_3(y, x, y) = 0$$

and

$$m_3(x, x, y) = m_3(y, x, x), \quad m_3(x, y, y) = m_3(y, y, x).$$

Thus each  $C_\infty$ -algebra structure on  $H^*$  is characterized by a pair of rational numbers  $p, q$  and

$$m_3(x, x, y) = pz, \quad m_3(x, y, y) = qz.$$

So let us write an arbitrary minimal  $C_\infty$ -algebra structure on  $H^*$  as a column vector  $\begin{pmatrix} p \\ q \end{pmatrix}$ .

Now let us look at the structure of an isomorphism of  $C_\infty$ -algebras

$$\{f_i\} : (H^*, m_3) \rightarrow (H^*, m'_3).$$

Again for dimensional reasons just one component  $f_1 : H^* \rightarrow H^*$  is possible, which in its turn consists of two isomorphisms

$$f_1^2 : H^2 = Q_x \oplus Q_y \rightarrow H^2 = Q_x \oplus Q_y, \quad f_1^5 : H^5 = Q_z \rightarrow H^5 = Q_z.$$

The first one is represented by a nondegenerate matrix  $A = \begin{pmatrix} a & c \\ b & d \end{pmatrix}$ ,

$$f_1^2(x) = ax \oplus by, \quad f_1^2(y) = cx \oplus dy,$$

and the second one by a nonzero rational number  $r, f_1^5(z) = rz$ .

A calculation shows that the condition  $f_1^5 m_3 = m'_3 (f_1^2 \otimes f_1^2 \otimes f_1^2)$ , to which the defining condition of an  $A_\infty$ -algebra morphism (7) degenerates looks as follows

$$r \begin{pmatrix} p \\ q \end{pmatrix} = \det A \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} p' \\ q' \end{pmatrix}.$$

This condition shows that two minimal  $C_\infty$ -algebra structures  $m_3 = \begin{pmatrix} p \\ q \end{pmatrix}$  and  $m'_3 = \begin{pmatrix} p' \\ q' \end{pmatrix}$  are isomorphic if and only if they are related by a nondegenerate linear transformation.

Thus there exist just two isomorphism classes of minimal  $C_\infty$ -algebras on  $H^*$ : the trivial one  $(H^*, m_3 = 0)$  and the nontrivial one  $(H^*, m_3 \neq 0)$ . So we have just two rational homotopy types whose rational cohomology is  $H^*$ . We denote them by  $X$  and  $Y$ , respectively and analyze them in the next sections.

Below we give some applications of the cohomology  $C_\infty$ -algebra in various problems of rational homotopy theory.

## 9.2 Formality

Among rational homotopy types with given cohomology algebra, there is one called *formal* which is a "formal consequence of its cohomology algebra" (Sullivan). Explicitly, this is the type whose minimal model  $M_X$  is isomorphic to the minimal model of the cohomology  $H^*(X, Q)$ .

Our  $C_\infty$  model implies the following criterion of formality:

**Theorem 16** .  *$X$  is formal if and only if its cohomology  $C_\infty$ -algebra is degenerate, i.e., it is  $C_\infty$  isomorphic to one with  $m_{\geq 3} = 0$ .*

Below we deduce some known results about formality using this criterion.

1. A commutative graded 1-connected algebra  $H$  is called *intrinsically formal* if there is only one homotopy type with this cohomology algebra  $H$ , of course the formal one.

The above Theorem 10 immediately implies the following sufficient condition for formality due to Tanre [34]:

**Theorem 17** . *If for a 1-connected graded  $Q$ -algebra  $H$  one has*

$$Harr^{k,k-2}(H, H) = 0, \quad k = 3, 4, \dots$$

*then  $H$  is intrinsically formal, that is there exists only one rational homotopy type with  $H^*(X, Q) \approx H$ .*

2. The following theorem of Halperin and Stasheff from [13] is an immediate result of our criterion:

**Theorem 18** . *A commutative graded  $Q$ -algebra of type*

$$H = \{H^0 = Q, 0, 0, \dots, 0, H^n, H^{n+1}, \dots, H^{3n-2}, 0, 0, \dots\}$$

*is intrinsically formal*

**Proof.** Since  $\deg m_i = 2 - i$  there is no room for operations  $m_{i>2}$ , indeed the shortest range is  $m_3 : H^n \otimes H^n \otimes H^n \rightarrow H^{3n-1} = 0$ .

3. Theorem 16 easily implies the

**Theorem 19 .** *Any 1-connected commutative graded algebra  $H$  with  $H^{2k} = 0$  for all  $k$  is intrinsically formal.*

**Proof.** Any  $A_\infty$ -operation  $m_i$  has degree  $2 - i$ , thus

$$m_i : H^{2k_1+1} \otimes \dots \otimes H^{2k_i+1} \rightarrow H^{2(k_1+\dots+k_i+1)} = 0.$$

Thus any  $C_\infty$ -operation is trivial too.

This implies a result of Baues: any space whose even-dimensional cohomologies are trivial has the rational homotopy type of a wedge of spheres. Indeed, such an algebra is realized by wedge of spheres and because of intrinsic formality this is the only homotopy type.

**Example.** The algebra  $H^*$  from the example of the previous section is not intrinsically formal since there are two homotopy types,  $X$  and  $Y$ , with  $H^*(X, Q) = H^* = H^*(Y)$ . The space  $X$  is formal (and actually  $X = S^2 \vee S^2 \vee S^5$ ), since its cohomology  $C_\infty$ -algebra  $(H^*, m_3 = 0)$  is trivial. But the space  $Y$  is not: its cohomology  $C_\infty$ -algebra  $(H^*, m_3 \neq 0)$  is not degenerate.

We remark here that the formal type is represented by  $X = S^2 \vee S^2 \vee S^5$  and it is possible to show that the nonformal one is represented by  $Y = S^2 \vee S^2 \cup_{f: S^4 \rightarrow S^2 \vee S^2} e^5$ , where the attaching map  $f$  is a nontrivial element from  $\pi_4(S^2 \vee S^2) \otimes Q$ .

### 9.3 Rational homotopy groups

Since the cohomology  $C_\infty$ -algebra  $(H^*(X, Q), \{m_i\})$  determines the rational homotopy type it must determine the rational homotopy groups  $\pi_i(X) \otimes Q$  too. We present a chain complex whose homology is  $\pi_i(X) \otimes Q$ . Moreover, the Lie algebra structure is determined as well.

For a cohomology  $C_\infty$ -algebra  $(H^*(X, Q), \{m_i\})$  the bar construction  $B(H^*(X, Q), \{m_i\})$  is a dg bialgebra, see (5.3.3). Acting on this bialgebra by the functor  $Q$  of indecomposables we obtain a dg Lie coalgebra.

On the other hand the rational homotopy groups  $\pi_*(\Omega X) \otimes Q$  form a graded Lie algebra with respect to Whitehead product. Thus its dual cohomotopy groups  $\pi^*(\Omega X, Q) = (\pi_*(\Omega X) \otimes Q)^*$  form a graded Lie coalgebra.

**Theorem 20** . Homology of a dg Lie coalgebra  $QB(H^*(X, Q), \{m_i\})$  is isomorphic to the cohomotopy Lie coalgebra  $\pi^*(\Omega X, Q)$ .

**Proof.** The theorem follows from the sequence of graded Lie coalgebra isomorphisms:

$$\begin{aligned} \pi^*(\Omega X, Q) &\approx (\pi_*(\Omega X, Q))^* \approx (PH_*(\Omega x, Q))^* \approx QH^*(\Omega X, Q) \\ &\approx QH(B(A(X))) \approx QH(\tilde{B}(H^*(X, Q), \{m_i\})) \\ &\approx H(Q\tilde{B}(H^*(X, Q), \{m_i\})). \end{aligned}$$

**Example.** For the algebra  $H^*$  from the previous examples the complex  $QB(H^*)$  in low dimensions looks as

$$0 \rightarrow Q_x \oplus Q_y \xrightarrow{0} Q_{x \otimes x} \oplus Q_{x \otimes y} \oplus Q_{y \otimes y} \xrightarrow{0} Q_{x \otimes x \otimes y} \oplus Q_{x \otimes y \otimes y} \xrightarrow{d=m_3} Q_z \oplus \dots$$

The differential  $d = m_3$  is trivial for the formal space  $X$  and is nontrivial for  $Y$ . Thus for both rational homotopy types we have

$$\pi^2 = H^1(QB(H^*)) = 2Q, \quad \pi^3 = H^2(QB(H^*)) = 3Q,$$

and

$$\begin{aligned} \pi^4(X) &= H^3(QB(H^*), d = 0) = 2Q, \\ \pi^4(Y) &= H^3(QB(H^*), d \neq 0) = \text{Ker } d = Q. \end{aligned}$$

## 9.4 Realization of homomorphisms

Let  $G : H^*(X, Q) \rightarrow H^*(Y, Q)$  be a homomorphism of cohomology algebras. When this homomorphism is realizable as a map of rationalizations  $g : Y_Q \rightarrow X_Q$ ,  $g^* = G$ ? In the case when  $G$  is an isomorphism this question was considered in [13]. It was considered also in [35]. The following theorem gives the complete answer:

**Theorem 21** . A homomorphism  $G$  is realizable if and only if it is extendable to a  $C_\infty$ -map

$$\{g_1 = G, g_2, g_3, \dots\} : (H^*(X, Q), \{m_i\}) \rightarrow (H^*(Y, Q), \{m'_i\}).$$

**Proof.** One direction of this is consequence of the last part of Theorem 14.

To show the other direction we use Sullivan's minimal models  $M_X$  and  $M_Y$  of  $A(X)$  and  $A(Y)$ . It is enough to show that the existence of  $\{g_i\}$  implies the existence of cdg algebra map  $g : M_Y \rightarrow M_X$ .

So we have  $C_\infty$ -algebra maps

$$M_X \xleftarrow{\{f_i\}} (H^*(X, Q), \{m_i\}) \xrightarrow{\{g_i\}} (H^*(y, Q), \{m'_i\}) \xrightarrow{\{f'_i\}} M_Y.$$

Recall the following property of a minimal cdg algebra  $M$ : for a weak equivalence of cdg algebras  $\phi : A \rightarrow B$  and a cdg algebra map  $f : M \rightarrow B$  there exists a cdg algebra map  $F : M \rightarrow A$  such that  $\phi F$  is homotopic to  $f$ . Using this property it is easy to show the existence of a cdga map  $\beta : M_X \rightarrow \mathcal{AQB}(M_X)$ , the right inverse of the standard map  $\alpha : \mathcal{AQB}(M_X) \rightarrow M$ . Composing this map with  $\mathcal{AQB}(\{f'_i\})\mathcal{AQB}(\{g_i\})$  we obtain a cdga map

$$\mathcal{AQB}(\{f'_i\})\mathcal{AQB}(\{g_i\})\beta : M_X \rightarrow M_Y.$$

This theorem immediately implies the

**Corollary 2** *For formal  $X$  and  $Y$  each  $G : H^*(X, Q) \rightarrow H^*(Y, Q)$  is realizable.*

**Proof.** In this case  $\{G, 0, 0, \dots\}$  is a  $C_\infty$ -extension of  $G$ .

**Example.** Consider the homomorphism

$$G : H^*(X) = H^*(Y) \rightarrow H^*(S^5)$$

induced by the standard imbedding  $g : S^5 \rightarrow X = S^2 \vee S^2 \vee S^5$ . Of course  $G$  is realizable as  $g : S^5 \rightarrow X$  but not as  $S^5 \rightarrow Y$ . Indeed, for such realizability, according to Theorem 21, we need a  $C_\infty$ -algebra morphism

$$\{g_i\} : (H^*, \{0, 0, m_3, 0, \dots\}) \rightarrow (H^5(S^5, Q), \{0, 0, 0, \dots\})$$

with  $g_1 = G$ . For dimensional reasons all the components  $g_2, g_3, \dots$  all are trivial, so this morphism has the form  $\{G, 0, 0, \dots\}$ . But this collection is not a morphism of  $C_\infty$ -algebras since the condition  $Gm_3 = 0$ , to which the defining condition (7) of an  $A_\infty$ -algebra morphism degenerates, is not satisfied.

## References

- [1] Paul S. Aspinwall and Sheldon Katz. Computation of superpotentials for D-branes. *Comm. Math. Phys.*, 264(1) (2006), 227–253.
- [2] C. Berger, B. Fresse, Combinatorial operad action on cochains, *Math. Proc. Cambridge Philos. Soc.* 137 (2004), 135–174.

- [3] N. Berikashvili, On differentials of spectral sequences, *Bull. Georg. Acad. Sci.*, 51, 1 (1968), 9–14.
- [4] N. Berikashvili, On the homology theory of spaces, *Bull. Georg. Acad. Sci.*, 86, 3 (1977), 529–532.
- [5] E. H. Brown, Twisted tensor products, I, *Ann. Of Math.* (2) 69 (1959), 223–246.
- [6] P. Deligne, P. Griffiths, J. Morgan, D. Sullivan, Real homotopy theory of Kähler manifolds, *Inv. math.*, 29 (1975), 245–275.
- [7] S. Eilenberg, S. MacLane, On the groups  $H(\pi, n)$ , *Ann. of Math.* 58 (1953), 55–106.
- [8] M. Gerstenhaber, The cohomology structure of an associative ring, *Ann. of Math.*, 78 (1963), 267–288.
- [9] M. Gerstenhaber, On the deformations of rings and algebras, *Ann. of Math.*, 79, 1 (1964), 59–103.
- [10] M. Gerstenhaber and A. Voronov, Higher operations on Hochschild complex, *Functional Anal. Appl.* 29 (1995), 1–6.
- [11] E. Getzler and J.D. Jones, Operads, homotopy algebra, and iterated integrals for double loop spaces, preprint (1994), hep-th/9403055.
- [12] V. Gugenheim and J. Stasheff, On perturbations and  $A(\infty)$ -structures, *Bull. Soc. Math. Belgique*, 38 (1986), 237–246.
- [13] S. Halperin, J. Stasheff, Obstructions to homotopy equivalences, *Adv. in Math.*, 23 (1979), 233–279.
- [14] V. Hinich, V. Schechtman, On homotopy limit of homotopy algebras, in “K-theory, arithmetic and geometry (Moscow, 1984–1986)”, *Lecture Notes in Mathematics* 1289, Springer-Verlag (1987), 240–264.
- [15] G. Hochschild, On the cohomology groups of an associative algebra *The Annals of Mathematics*, 2nd ser., 46, No. 1 (Jan., 1945)

- [16] T. Kadeishvili, On the Differentials of Spectral Sequence of Fiber Bundle, *Bull. of Georg. Acad. Sci.*, 82 1976, 285–288.
- [17] T. Kadeishvili, On the Homology Theory of Fibrations, *Russian Math. Surveys*, 35, 3 (1980), 231–238.
- [18] T. Kadeishvili, Algebraic Structure in Homology of an  $A(\infty)$ -algebra, *Bull. of Georg. Acad. Sci.*, 108, 2 1982, 309–312.
- [19] T. Kadeishvili, Predifferential of a Fiber Bundle, *Russian Math. Surveys*, 41, 6 1986, 135–147.
- [20] T. Kadeishvili, Functor D for the Category of  $A(\infty)$ -algebras, *Bull. of Georg. Acad. Sci.*, 125, 2 1987, 273–276.
- [21] T. Kadeishvili, The  $A_\infty$ -algebra Structure and Cohomology of Hochschild and Harrison, *Proc. of Tbil. Math. Inst.*, 91, (1988), 19–27.
- [22] T. Kadeishvili,  $A(\infty)$ -algebra Structure in Cohomology and Rational Homotopy Type, *Forschungsswerpunkt Geometrie, Universitaet Heidelrg, Math. Inst.*, Heft 37, Oktober 1988, 1–64.
- [23] T. Kadeishvili,  $A_\infty$  -algebra Structure in Cohomology and Rational Homotopy Type, (in Russian), *Proc. of Tbil. Mat. Inst.*, 107, (1993), 1–94.
- [24] T. Kadeishvili, Cochain operations defining Steenrod  $\smile_i$ -products in the bar construction, *Georgian Math. Journal*, v.10, n. 1 2003, 115–125, preprint at math.AT/0207010.
- [25] T. Kadeishvili, On the cobar construction of a bialgebra, *Homology, Homotopy and Appl.*, v. 7(2) 2005, 109–122.
- [26] T. Kadeishvili, Twisting Elements in Homotopy G-algebra, arXiv:0709.3130.
- [27] J. McLure, J. Smith, Multivariable cochain operations and little n-cubes, preprint (2001), math.QA/0106024.
- [28] M. Markl, A cohomology theory for  $A(m)$ -algebras and applications. *Journ. Pure Appl. Algebra*, 83 (1992), 141—175.

- [29] D. Quillen, Rational homotopy theory, *Ann. of Math.*, 90(1969), 205–295.
- [30] V. Smirnov, Homology of fiber spaces, *Russian Math. Surveys*, 35, 3 (1980), 183–188.
- [31] V. Smirnov, Homotopy theory of coalgebras, *Izv. Akad. Nauk SSSR Ser. Mat.* 49 (1985), 1302-1321. English translation in *Math. USSR-Izv.* 27 (1986), 575–592.
- [32] J.D. Stasheff, Homotopy associativity of H-spaces. I, II, *Trans. Amer. Math. Soc.*, 108 (1963), 275–312.
- [33] M. Shlessinger and J. Stasheff, The Lie algebra structure on tangent cohomology and deformation theory, *J. Pure Appl. Algebra* 89 (1993), 231–235.
- [34] D. Tanre, Cohomologie de Harrison et type d’homotopie rationnelle, *J. of Pure and appl. Alg.*, 38(1985), 353–366.
- [35] M. Vigue-Poirrier, Realisation de morphismes donnees en cohomologie et suite spectral d’Eilenberg-Moore, *TAMS*, 265, 2 (1981), 447–483.

A. Razmadze Mathematical Institute,  
 1, M. Alexidze Str., Tbilisi, 0193, Georgia  
 tornike.kadeishvili@tsu.ge