THE GROWTH OF THE GREEN FUNCTION FOR RANDOM WALKS AND POINCARE SERIES ´

MATTHIEU DUSSAULE, LONGMIN WANG, AND WENYUAN YANG

ABSTRACT. Given a probability measure μ on a finitely generated group Γ , the Green function $G(x, y|r)$ encodes many properties of the random walk associated with μ . Finding asymptotics of $G(x, y|r)$ as y goes to infinity is a common thread in probability theory and is usually referred as renewal theory in literature. Endowing Γ with a word distance, we denote by $H_r(n)$ the sum of the Green function $G(e, x|r)$ along the sphere of radius n. This quantity appears naturally when studying asymptotic properties of branching random walks driven by μ on Γ and the behavior of $H_r(n)$ as n goes to infinity is intimately related to renewal theory. Our motivation in this paper is to construct various examples of particular behaviors for $H_r(n)$. First, our main result exhibits a class of relatively hyperbolic groups with convergent Poincaré series generated by $H_r(n)$, which answers some questions raised in [\[DWY22\]](#page-50-0). Along the way, we investigate the behavior of $H_r(n)$ for several classes of finitely generated groups, including abelian groups, certain nilpotent groups, lamplighter groups, and Cartesian products of free groups.

1. INTRODUCTION

Let Γ be a finitely generated group endowed with a finite generating set. Denote by | \cdot | the induced word distance and by $S_n = \{x \in \Gamma : |x| = n\}$ the sphere of radius n centered at the neutral element e of Γ. We set

$$
v = \limsup_{n \to \infty} \frac{1}{n} \log |S_n|.
$$

The number v is called the volume growth of Γ with respect to the chosen generating set. Note that v is also the growth rate of balls, i.e. $v = \limsup_{n \to \infty} \frac{1}{n} \log \sharp B_n$, where B_n is the ball of radius n centered at e, see [\[Man12,](#page-50-1) Section 2.2] for more details. The group Γ has exponential growth (respectively sub-exponential growth) if $v > 0$ (respectively $v = 0$) and this does not depend on the choice of the generating set, although the exact value of v does, when non-zero.

Consider an admissible probability measure μ on Γ. Let $p_n(x, y)$ be the *n*-step transition probability for the random walk (X_k) with step distribution μ . Define for $0 < r \leq R$ the Green function

$$
G(x,y|r) = \sum_{n=0}^{\infty} r^n p_n(x,y),
$$

where R is the radius of convergence of the series, which is also the inverse of the spectral radius of the random walk. When $r = 1$ we write $G(x, y) = G(x, y|1)$ for simplicity. Note that we have $G(x, y|r) = G(e, e|r)F(x, y|r)$, where $F(x, y|r)$ is

L.W. is supported by National Natural Science Foundation of China (12171252). W.Y. is supported by National Key R & D Program of China (SQ2020YFA070059).

the first return Green function [\[Woe00,](#page-51-0) Lemma 1.13]. Define the growth function associated with the Green function $H_r(n)$ as

$$
H_r(n) = \sum_{x \in S_n} G(e, x|r)
$$

and the growth rate of the Green function $\omega_{\Gamma}(r)$ as

$$
\omega_{\Gamma}(r) = \limsup_{n \to \infty} \frac{1}{n} \log H_r(n).
$$

We will always consider random walks that are transient at the spectral radius, i.e. $G(x, y|r)$ is finite for every $x, y \in \Gamma$ and for every $r \leq R$. By Varapoulos Theorem [\[Woe00,](#page-51-0) Theorem 7.8], only groups with at most quadratic growth can carry a random walk which is not transient at the spectral radius. In other words, we will always assume that Γ is not virtually \mathbb{Z}^d , $d \leq 2$. In particular, this ensures that $H_r(n)$ and $\omega_{\Gamma}(r)$ are well defined for every $r \leq R$.

Notations. In all the paper, given two functions f and g, we write $f \simeq g$ if the difference between f and g is bounded from above and below, that is $|f - g| \leq C$ for some constant C. We write $f \simeq_L g$ if the constant C depends on a parameter L. Assuming further that f and g are positive, we write $f \approx g$ if the ratio of f and g is bounded from above and below, that is $\frac{1}{C} f \leq g \leq C f$ for some positive constant C. Similarly, if C depends on L, we write $f \succeq_L g$. Also, if $f \leq Cg$, we write $f \prec g$ and $f \prec_L g$ if C depends on L. If the dependency is not clear from the context, we will avoid these notations.

This paper mostly deals with the asymptotics of $H_r(n)$ as n goes to infinity. A fruitful line of research in the theory of random walks is to compute asymptotics in space of the Green function, that is asymptotics of $G(e, x|r)$ as x goes to infinity. This is referred as renewal theory and goes back to Blackwell's renewal theorem for drifted random walks on R, see [\[Bla53\]](#page-49-0) and earlier references therein. The terminology renewal comes from an interpretation of the Green function $G(e, x)$ as the probability that a renewal event takes place at x for a suited process, see [\[Spi76,](#page-51-1) Chapter II.9].

In fact, the terminology renewal theory is used in a much broader setting and we refer to [\[Fel66,](#page-50-2) Chapter XI] for a more complete exposition within the scope of probability theory. Let us also mention that Lalley [\[Lal89\]](#page-50-3) generalized classical renewal theory, with a new approach to deal with asymptotics of certain counting functions arising in geometric group theory. This led to significant research in dynamical systems, where renewal theory is now a common thread.

Finally, note that Ledrappier interpreted the computation of the limit of $H_1(n)$ on the free group as a renewal theorem for the distance [\[Led01\]](#page-50-4), see also [\[Led93\]](#page-50-5) for related results concerning the Brownian motion on the universal cover of compact negatively curved manifolds. However, besides such specific examples, the behaviors of $H_r(n)$ and $\omega_{\Gamma}(r)$ have not been investigated much in literature, so let us first explain in which context these quantities occur.

Consider a probability measure ν on $\mathbb{Z}_{\geq 0} = \{0, 1, 2, ...\}$. The branching random walk driven by ν and μ on Γ , denoted by BRW(Γ , ν , μ) is described as follows. One starts with a single particle at e . For every n, every alive particle at time n dies after giving birth to an independent number of children, according to the distribution of ν, each of which independently moves on Γ according to the distribution of µ. The measure ν is called the offspring distribution.

In recent years, there has been a large body of work dedicated to understanding the asymptotic behavior of BRW(Γ, ν , μ) in terms of geometric features of Γ. For such study, one usually condition on non-extinction of the system, which boils down to considering an offspring measure distributed on $\mathbb{N} = \{1, 2, ...\}$, see [\[AN04,](#page-49-1) Chapter 1. It is thus natural to assume that $\mathbf{E}[\nu] > 1$, otherwise, conditioned on non-extinction, ν is the Dirac measure at 1 and the branching random walk is nothing but the usual random walk whose step distribution is given by μ . One of the cornerstone result is that letting $\mathbf{E}[\nu] = r$, then $1 < r \leq R$ if and only if the branching random walk is transient, i.e. almost surely it does not visit every vertex infinitely many times, see [\[BP94\]](#page-49-2) and [\[GM06\]](#page-50-6). Furthermore, if $1 < r \leq R$, then the branching random walk has exponential volume growth by [\[BM12\]](#page-49-3). Precisely, letting M_n be the cardinality of the number of elements of S_n that are ever visited by the branching random walk, we have that almost surely,

$$
\limsup_{n \to \infty} \frac{1}{n} \log M_n > 0.
$$

In some cases, such as hyperbolic groups [\[SWX22,](#page-51-2) Theorem 1.1] and relatively hyperbolic groups [\[DWY22,](#page-50-0) Theorem 1.1], it was shown that this growth rate coincides almost surely with the growth rate of the Green function at $1 < r \leq R$. That is,

$$
\limsup_{n \to \infty} \frac{1}{n} \log M_n = \omega_{\Gamma}(r).
$$

Moreover, for hyperbolic groups, we have $H_r(n) \approx e^{n\omega_{\Gamma}(r)}$, i.e. the Green function has purely exponential growth, see [\[SWX22,](#page-51-2) Theorem 3.1].

In fact, our original motivation for this paper was to answer some questions raised in [\[DWY22\]](#page-50-0) about branching random walks on relatively hyperbolic groups. We introduce the following Poincaré series associated with μ on Γ. For $r \leq R$ and $s \in \mathbb{R}$, we set

(1.1)
$$
\Theta_{\Gamma}(r,s) = \sum_{x \in \Gamma} G(e,x|r) e^{-sd(e,x)} = \sum_{n \ge 0} H_r(n) e^{-sn}.
$$

The growth rate $\omega_{\Gamma}(r)$ is thus the critical exponent of this Poincaré series, i.e. for $s < \omega_{\Gamma}(r)$, $\Theta_{\Gamma}(r,s)$ diverges and for $s > \omega_{\Gamma}(r)$, $\Theta_{\Gamma}(r,s)$ converges.

Several criteria were found in [\[DWY22\]](#page-50-0) to ensure that this Poincaré series diverges at $s = \omega_{\Gamma}(r)$. One of the missing pieces was whether there exists an example with convergent Poincaré series. Our main result exhibits such an example by actually constructing a relatively hyperbolic group for which divergence of the Poincaré series depends on r. Precisely, we prove the following.

Theorem 1.1 (Theorem [7.9\)](#page-37-0). There exists a relatively hyperbolic group Γ , endowed with a finitely supported symmetric admissible probability measure μ , and there exist $1 < r_* < r_{\text{t}} < R$ such that

- (1) for any $r \leq r_*, \Theta_{\Gamma}(r, \omega_{\Gamma}(r))$ diverges,
- (2) at $r = r_{\sharp}, \Theta_{\Gamma}(r, \omega_{\Gamma}(r))$ converges.

Remark 1.2. We are unable to determine whether the second assertion is true for any $r > r_*$. If this were true, we would obtain a phase transition for the divergence of the Poincare series and also for all the assertions in Corollary [1.3](#page-3-0) below. However, up to changing the measure μ , we have a weak form of a phase transition at the level of the parabolic subgroups, see Remark [7.11](#page-39-0) for more details.

We are now moving on explaining several applications of Theorem [1.1.](#page-2-0) Before that, let us put into context the series $\Theta_{\Gamma}(r,s)$ under consideration. The question whether $\Theta_{\Gamma}(r,s)$ converges or diverges at $s = \omega_{\Gamma}(r)$ is of particular importance for many properties such as the construction of boundary measures and growth problems. It is also related to the parabolic gap property coined in [\[DWY22\]](#page-50-0) that we discuss next.

Consider a relatively hyperbolic group Γ and a maximal parabolic subgroup P . The *growth rate of the Green function* induced on P is defined as

$$
\omega_P(r) = \limsup_{n \to \infty} \frac{1}{n} \log \sum_{\substack{x \in P \\ |x| = n}} G(e, x|r).
$$

Following [\[DWY22\]](#page-50-0), we say that Γ has a parabolic gap along P for the Green function at $r \in [1, R]$ if $\omega_P(r) < \omega_{\Gamma}(r)$. If this holds along every parabolic subgroups, we say Γ has a parabolic gap for the Green function at r and if this in turn holds for every $r \in [1, R]$ we say that Γ has a parabolic gap for the Green function.

This notion is analogous to the parabolic gap condition introduced in [\[DOP00\]](#page-49-4), where first examples of convergent (standard) Poincaré series were constructed without this parabolic gap condition. As an interesting consequence, the authors of [\[DOP00\]](#page-49-4) produced Patterson-Sullivan measures having atoms at parabolic points in the visual boundary of Hadamard manifolds.

In $[DWY22]$, we proved that if there exists a relatively hyperbolic group with convergent Poincaré series, then the parabolic gap condition fails. On the other hand, having a parabolic gap has various applications related to asymptotic properties of branching random walks, see in particular [\[DWY22,](#page-50-0) Theorem 1.8, Remark 5.4].

Motivated by this discussion and by the work of [\[DOP00\]](#page-49-4), we make use of the above theorem to further develop various properties with a similar behavior. We introduce a family of Patterson-Sullivan type measures $\nu_e(r)$ associated with the Poincaré series defined in [\(1.1\)](#page-2-1) and a family of proper distances \mathfrak{d}_r which are quasiisometric to the word distance. We summarize here the different results we obtain, see Theorem [7.9,](#page-37-0) Corollary [7.13](#page-40-0) and Theorem [7.19.](#page-43-0)

Corollary 1.3. The pair (Γ, μ) in Theorem [1.1](#page-2-0) has the following properties:

- (1) the parabolic gap for the Green functions holds for $r \leq r_*$ but fails at $r = r_{\sharp}$,
- (2) the Patterson-Sullivan measure $\nu_e(r)$ is supported on conical limit points for $r \leq r_*$ and is purely atomic and supported on parabolic limit points at $r = r_{\sharp}$,
- (3) the growth tightness for the proper distance \mathfrak{d}_r holds for $r \leq r_*$ but fails at $r = r_{\text{H}}$.

Remark 1.4. In [\[Pei11\]](#page-50-7), Peigné constructed a divergent Schottky group without the parabolic gap condition. In view of our examples, it is relevant to ask here whether there exist examples of divergent Poincaré series (1.1) without a parabolic gap for the Green function.

Remark 1.5. The proper metric in (3) is defined by

$$
\mathfrak{d}_r(x,y) := \omega_{\Gamma}(r)|x^{-1}y| + |x^{-1}y|_r
$$

so it is a linear combination of the word metric $|x^{-1}y|$ and the r-Green metric $|x^{-1}y|_r := -\log \frac{G(x,y|r)}{G(e,e|r)}$. Cashen-Tao [\[CT16\]](#page-49-5) have shown examples of product groups with growth tightness for one generating set but not for another generating set. The above examples within the class of relatively hyperbolic groups are new.

These theorems are the conclusion of several results that we prove along the paper. If we restrict our attention to branching random walks, the study of $H_r(n)$ and $\omega_{\Gamma}(r)$ seems to be relevant only for $r > 1$. However, as we now observe, even for $r \leq 1$, both these quantities appear naturally and are worth being studied.

For any subset A of Γ, the growth rate $\omega_A(r)$ of the Green function restricted to A at r can be defined similarly to $\omega_{\Gamma}(r)$. Namely, we set

$$
H_{A,r}(n) = \sum_{x \in S_n \cap A} G(e, x | r)
$$

and

$$
\omega_A(r) = \limsup_{n \to \infty} \frac{1}{n} \log H_{A,r}(n).
$$

We can interpret $\omega_A(r)$ via the first return kernel to A associated with $r\mu$, which is defined by

$$
p_{r,A}(x,y) = \sum_{n \ge 1} \sum_{z_1, \dots, z_{n-1} \notin A} r^n p(x, z_1) p(z_1, z_2) \dots p(z_{n-1}, y), \ x, y \in A.
$$

Letting $G_{r,A}$ denote the corresponding Green function, it holds that for every x, y in A,

$$
G_{r,A}(x,y|1) = G(x,y|r)
$$

by [\[DG21,](#page-49-6) Lemma 4.4]. Consequently, the growth rate $\omega_A(r)$ coincides with the growth rate at 1 of this new Green function $G_{r,A}$. In general, there is no reason for $p_{r,A}$ to be a Markov transition kernel, that is, $\sum_{y \in A} p_{r,A}(x, y)$ might not be a constant equal to 1. When A is a subgroup of Γ , then $p_{r,A}$ is A-invariant, so $\sum_{y\in A} p_{r,A}(x,y)$ is independent of x, but there is still no reason for this transition kernel to be Markov. As a matter of fact, in various interesting cases, $p_{r,A}$ is a sub-Markov transition kernel. Letting t be its total mass and setting $\tilde{p}_{r,A} = t^{-1}p_{r,A}$ and $G_{r,A}$ for the corresponding Green function, we see that $G_{r,A}(\cdot,\cdot|1) = G_{r,A}(\cdot,\cdot|t)$. Therefore, the growth rate $\omega_A(r)$ restricted to A at r coincides in this case with the growth rate at some $t < 1$ of a Markov transition kernel on A. As explained above, in the context of relatively hyperbolic groups, letting A be a maximal parabolic subgroup, the relation between $\omega_A(r)$ and $\omega_{\Gamma}(r)$ is of particular importance in understanding the asymptotic behavior of $H_r(n)$. In particular, this discussion motivates the study of $H_r(n)$ and $\omega_{\Gamma}(r)$ also for $r \leq 1$.

In fact, a large part of our work is devoted to understanding the behavior of $H_1(n)$ for symmetric admissible and finitely supported random walks on amenable groups and our study goes beyond applications to relatively hyperbolic groups. Here is a summary of the different results we obtain.

Theorem 1.6 (Theorem [3.1,](#page-9-0) Theorem [4.1\)](#page-15-0). The function $H_1(n)$ is asymptotically linear in n for the following classes of groups:

(1) if Γ is a free abelian group of finite rank endowed with a finitely supported symmetric admissible probability measure, then $H_1(n) \sim Cn$ for any finite generating set,

- (2) if Γ is a finitely generated nilpotent group of nilpotency class $N_{\Gamma} \leq 2$ endowed with a finitely supported symmetric admissible probability measure, then $H_1(n) \simeq n$ for any finite generating set,
- (3) if Γ is a lamplighter group endowed with the set of generators considered in [\[BW05\]](#page-49-7), then for the simple random walk, $H_1(n) \sim Cn$.

Finally, elaborating on the work of Picardello and Woess [\[PW94\]](#page-50-8), we prove the following phase transition result for random walks on bi-trees. Consider the Cartesian product of two regular trees T_1 and T_2 of respective degree l_1 and l_2 . Let μ_i be finitely supported admissible symmetric probability measures on T_i and set $\mu = \alpha \mu_1 + (1 - \alpha) \mu_2$ for $0 < \alpha < 1$.

Theorem 1.7 (Theorem [5.1\)](#page-18-0). The probability measure μ on $T_1 \times T_2$ satisfies the following. If $l_1 = l_2$, then for every fixed $r < R$, $H_r(n) \approx e^{n\omega_{\Gamma}(r)}$ for all $n \geq 1$. If $l_1 > l_2$, then there exists a phase transition at some $r_0 \in (1, R)$ such that the following holds.

- (1) For every $r < r_0$, we have $H_r(n) \ge e^{n\omega_{\Gamma}(r)}$ for all $n \ge 1$.
- (2) At $r = r_0$, we have $H_r(n) \approx n^{-1} e^{n\omega_{\Gamma}(r)}$ for all $n \geq 1$.
- (3) For every $r_0 < r < R$, we have $H_r(n) \approx n^{-3/2} e^{n\omega_{\Gamma}(r)}$ for all $n \geq 1$.

This theorem is a pivotal result in our paper. It is the conclusion of our general study of $H_r(n)$ on finitely generated group and is also one of the main pieces in proving Theorem [1.1.](#page-2-0)

Let us now give more details on how the paper is organized. It has seven sections including the introduction. In the first half which consists of Sections [2,](#page-6-0) [3,](#page-9-1) [4](#page-14-0) and [5,](#page-18-1) we give general statements on $H_r(n)$ and $\omega_{\Gamma}(r)$, as well as various examples of different situations. The second half focuses on applications to relatively hyperbolic groups and is divided into Sections [6](#page-24-0) and [7.](#page-33-0) It is based both on the first half and on previous results of [\[DWY22\]](#page-50-0).

More specifically, in Section [2,](#page-6-0) we consider the growth rate $\omega_{\Gamma}(r)$ at the special value $r = 1$. We prove that $\omega_{\Gamma}(1) = 0$ and study further continuity of the function $r \mapsto \omega_{\Gamma}(r)$ at 1.

We then study the asymptotic behavior of $H_1(n)$ on amenable groups Γ and we prove Theorem [1.6](#page-4-0) in Sections [3](#page-9-1) and [4.](#page-14-0) We show in particular that $H_1(n)$ is asymptotically linear in n for abelian groups and provide partial results for nilpotent groups, see Theorem [3.1](#page-9-0) and Conjecture [3.5.](#page-13-0) This raises the question whether $H_1(n)$ always behaves linearly in amenable groups. We prove that it is the case for the simple random walk on the lamplighter group in Theorem [4.1.](#page-15-0) The main strategy in these two sections is basically to combine previously known renewal results, i.e. about the asymptotics in space of the Green function and known results on the large scale properties of the spheres S_n .

Then, in Section [5,](#page-18-1) we prove Theorem [1.7.](#page-5-0) The behavior of $H_r(n)$ that we exhibit is of an again different form than the one described above, for (relatively) hyperbolic groups and for amenable groups respectively. As a particular consequence, we see that the quantity $H_r(n)$ may fail to be sub-multiplicative. This disproves an argument due to Candellero, Gilch and Müller [\[CGM12\]](#page-49-8) as well as its consequences, see Remark [7.14](#page-40-1) for more details.

In Section [6,](#page-24-0) we focus on relatively hyperbolic groups. We introduce a proper distance \mathfrak{d}_r on Γ , $1 \leq r \leq R$, and its associated Poincaré series $\mathcal{P}(s)$. For $r < R$, \mathfrak{d}_r is quasi-isometric to the word distance. The critical exponent of $\mathcal{P}(s)$ is 1 and we prove that divergence, respectively convergence of $\mathcal{P}(s)$ at 1 is equivalent to divergence, respectively convergence of the Poincaré series $\Theta_{\Gamma}(r, s)$ at $s = \omega_{\Gamma}(r)$. Then, we use $P(s)$ to construct Patterson-Sullivan type measures ν_x on the Bowditch boundary of the group. This allows us to find a characterization of divergence of $\Theta_{\Gamma}(r,s)$ at the growth rate $\omega_{\Gamma}(r)$ in terms of the support of the measures ν_x , see in particular Theorem [6.17.](#page-32-0)

We then prove Theorem [1.1](#page-2-0) in Section [7](#page-33-0) and answer some questions raised in [\[DWY22\]](#page-50-0). In particular, we show that whenever maximal parabolic subgroups are amenable, Γ necessarily has a parabolic gap for the Green function, see Corol-lary [7.4.](#page-34-0) Along the way, we study further the distance \mathfrak{d}_r and prove that the parabolic gap for the Green function is equivalent to growth tightness for the distance \mathfrak{d}_r . In particular, we find that for suited r, Γ is not growth tight for \mathfrak{d}_r , which allows us to partially answer a question raised in [\[ACT15\]](#page-49-9), see Corollary [7.22.](#page-44-0)

2. THE GROWTH RATE OF THE GREEN FUNCTION AT $r = 1$

We study $\omega_{\Gamma}(1)$ in this section. We first show that $\omega_{\Gamma}(r)$ always vanishes at 1 and then study continuity at this special value.

2.1. Nullity of the growth rate at $r = 1$. In this section, we do not need to assume that the random walk is symmetric, nor that it is finitely supported. We start with the following lower bound.

Lemma 2.1. Let Γ be a finitely generated group and μ an admissible probability measure on Γ . Then, $\omega_{\Gamma}(1) > 0$.

Proof. Since the series $\sum_{n=0}^{\infty} H_1(n) = \sum_{k=0}^{\infty} \sum_{x \in \Gamma} p_k(e, x) = \sum_{k=0}^{\infty} 1 = \infty$, we have that

$$
\omega_{\Gamma}(1) = \limsup_{n \to \infty} \frac{1}{n} \log H_1(n) \ge 0.
$$

Remark 2.2. If μ is finitely supported, then there is a constant $C > 0$ such that $H_n(1) \geq C$ for all $n \geq 1$. In fact, since the random walk driven by μ is transient and has bounded jumps at most $C_0 > 0$, it will eventually visit the annulus

$$
A(n, n + C_0) = \{x, n \le |x| \le n + C_0\}.
$$

Note that

$$
\sum_{x \in A(n,n+C_0)} G(e,x) = \sum_{x \in A(n,n+C_0)} \sum_{k \ge 0} p_k(e,x)
$$

=
$$
\sum_{x \in A(n,n+C_0)} \sum_{k \ge 0} \mathbf{P}(X_k = x).
$$

Thus,

$$
\sum_{x \in A(n, n+C_0)} G(e, x) \ge \mathbf{P}(\text{the random walk ever visits } A(n, n+C_0))
$$

\n
$$
\ge 1.
$$

Now by [\[DWY22,](#page-50-0) Lemma 3.1 (1)], $H_1(n) \approx H_1(n+1)$, hence

$$
\sum_{x \in A(n,n+C_0)} G(e,x) \asymp H_1(n),
$$

which concludes the proof that $H_n(1) \geq C$.

Proposition 2.3. Let Γ be a finitely generated group and μ be an admissible probability measure on Γ. Then $\omega_{\Gamma}(1) \leq 0$. Moreover, if Γ has exponential growth, there exists $C > 0$ such that $H_1(n) \leq Cn^3$.

Proof. Note that

$$
H_1(n) \le G(e, e) \sharp S_n,
$$

hence, if the volume growth rate

$$
v = \limsup_{n \to \infty} \frac{1}{n} \log |S_n| = 0,
$$

then $\omega_{\Gamma}(1) \leq 0$.

We assume that $v > 0$ in the remainder of the proof. Then there exist $c_1 > 1$ and $v_1, v_2 > 0$ such that

(2.1)
$$
c_1^{-1}e^{v_1 n} \leq \sharp B_n \leq c_1 e^{v_2 n},
$$

where $\sharp B_n$ is the cardinality of the ball of radius n. By [\[Var91,](#page-51-3) Theorem 1],

$$
p_n(x, y) \le c_2 e^{-c_3 n^{1/3}}
$$

for some $c_2, c_3 > 0$. Choose c_4 so that $c_4^{1/3}c_3 > v_2$. Then

$$
H_1(n) \leq \sum_{k=0}^{c_4 n^3} \sum_{x \in S_n} p_k(e, x) + c_2 \sum_{x \in S_n} \sum_{k=c_4 n^3 + 1}^{\infty} c_3 e^{-c_3 k^{1/3}}
$$

$$
\leq c_4 n^3 + 1 + c_5 \sum_{k=c_4 n^3 + 1}^{\infty} e^{v_2 n} e^{-c_3 k^{1/3}}.
$$

Note that

$$
\sum_{k=c_4n^3+1}^{\infty} e^{-c_3 k^{1/3}} \le \int_{c_4n^3}^{\infty} e^{-c_3 t^{1/3}} dt \le c_6 n^2 e^{-c_3 c_4^{1/3} n}.
$$

Thus we have

$$
H_1(n) \le c_4 n^3 + 1 + c_7 n^2 e^{\left(v_2 - c_3 c_4^{1/3}\right)n} \le c_8 n^3
$$

and hence $\omega_{\Gamma}(1) \leq 0$.

Lemma [2.1](#page-6-1) and Proposition [2.3](#page-7-0) together yield the following corollary.

Corollary 2.4. Let Γ be a finitely generated group and let μ be an admissible probability measure on Γ . Then, $\omega_{\Gamma}(1) = 0$.

We also have the following result.

Proposition 2.5. Let Γ be a finitely generated group and let μ be an admissible probability measure on Γ . Then, $r \mapsto \omega_{\Gamma}(r)$ is monotonically non-decreasing.

Proof. Since the Green function itself is non-decreasing in r, $H_n(s) \leq H_n(r)$, hence $\omega_{\Gamma}(s) \leq \omega_{\Gamma}(r)$ if $s \leq r$. $\omega_{\Gamma}(s) \leq \omega_{\Gamma}(r)$ if $s \leq r$.

2.2. Continuity of the growth rate. We start with the following result which holds for every finitely generated group, without assuming that the random walk is finitely supported. We assume however that it is symmetric.

Proposition 2.6. Let Γ be a finitely generated group and let μ be a symmetric admissible probability measure on Γ. Then, the function $\omega_{\Gamma}(r)$ is continuous for $0 < r < R$.

Proof. We modify the arguments in [\[DWY22,](#page-50-0) Lemma 3.1], where μ was assumed to be finitely supported and only the case $1 < r < R$ was treated. There are constants $c_1 > 0$ and $v_2 \ge 0$ such that $\sharp S_n \le c_1 e^{v_2 n}$. Fix $\delta > 0$. We choose c_2 so that for every $\delta \leq r \leq R - \delta$,

$$
v_2 - c_2 \left(\log R - \log(R - r)\right) < \omega_\Gamma(r).
$$

Note that since the underlying random walk is symmetric, for every x and every k , we have $p_k(e, x)p_k(e, x) \leq p_{2k}(e, e)$ and by [\[Woe00,](#page-51-0) Lemma 1.9], $p_{2k}(e, e) \leq R^{-2k}$. Thus,

$$
(2.2) \t\t\t\t\t\t p_k(e,x) \le R^{-k}
$$

for every $x \in \Gamma$ and $k \geq 0$. Consequently, we have for $\delta \leq r \leq R - \delta$,

$$
\sum_{x \in S_n} \sum_{k > c_2 n} r^k p_k(e, x) \le c_1 e^{v_2 n} \sum_{k > c_2 n} \left(\frac{r}{R}\right)^k \le c_1 \delta^{-1} R e^{v_2 n - c_2 (\log R - \log r) n}.
$$

By the choice of c_2 we have that

$$
\omega_{\Gamma}(r) = \limsup_{n \to \infty} \frac{1}{n} \log \sum_{x \in S_n} \sum_{k=0}^{c_2 n} r^k p_k(e, x).
$$

Now for $0 < s < r \leq R - \delta$,

$$
H_n(s) \ge \sum_{x \in S_n} \sum_{k=0}^{c_2 n} s^k p_k(e, x) \ge \left(\frac{s}{r}\right)^{c_2 n} \sum_{x \in S_n} \sum_{k=0}^{c_2 n} r^k p_k(e, x),
$$

and hence $0 \leq \omega_{\Gamma}(r) - \omega_{\Gamma}(s) \leq c_2 (\log r - \log s)$, since ω_{Γ} is non-decreasing by Proposition [2.5.](#page-7-1) Now $\delta > 0$ is arbitrary, so we prove that $\omega_{\Gamma}(r)$ is continuous in $0 < r < R$.

The continuity at the inverse of the spectral radius seems to be a difficult problem in general. It is already known that ω_{Γ} is continuous at R in hyperbolic groups [\[SWX22,](#page-51-2) Theorem 1.1] and in relatively hyperbolic groups [\[DWY22,](#page-50-0) Theorem 1.1]. However, we do not know much beyond these classes of groups.

We now investigate further continuity at $r = 1$. By [\[Woe00,](#page-51-0) Theorem 12.5], $R = 1$ can only happen if Γ is amenable, hence the following discussion only applies to amenable groups.

Assume that the random walk satisfies the following Gaussian lower bound. There exists a sub-exponential function f, i.e. $\frac{1}{n} \log f(n) \to 0$, $n \to \infty$, such that for every k, for every $x \in \Gamma$,

$$
(2.3) \t\t\t p_k(e,x) \gtrsim f(k)e^{-c\frac{|x|^2}{k}}.
$$

Then,

$$
H_r(n) = \sum_{x \in S_n} \sum_{k=0}^{\infty} r^k p_k(e, x) \gtrsim \sharp S_n \sum_{k=0} f(k) e^{-k \log r^{-1} - c \frac{n^2}{k}}
$$

$$
\gtrsim f\left(\left\lfloor \frac{n}{\sqrt{\log r^{-1}}}\right\rfloor\right) \sharp S_n e^{-n(1+c)\sqrt{\log r^{-1}}}.
$$

It follows that

$$
\omega_{\Gamma}(r) \ge -(1+c)\sqrt{\log r^{-1}} + v.
$$

Letting $r \uparrow 1$, we see that $0 \geq \limsup_{r \uparrow 1} \omega_{\Gamma}(r) \geq v$. Therefore $v = 0$ and

$$
\lim_{r \uparrow 1} \omega_{\Gamma}(r) = 0.
$$

Remark 2.7. We thus recover the fact that a Gaussian lower bound like [\(2.3\)](#page-8-0) cannot hold for groups of exponential volume growth, which was already noticed in literature, see for instance the comments after [\[Ale92,](#page-49-10) (0.3)].

Proposition 2.8. Let Γ be a finitely generated virtually nilpotent group endowed with a finite generating set and let μ be a finitely supported symmetric admissible probability measure on Γ. Then, the function ω_{Γ} is left-continuous at 1.

Proof. The fact that a Gaussiam lower bound [\(2.3\)](#page-8-0) with $f(k) = k^{-d/2}$ holds for virtually nilpotent groups is well known, see for instance [\[Ale02,](#page-49-11) Corollary 1.9]. \Box

3. Abelian and nilpotent groups

Our goal in this section is to prove the following result.

Theorem 3.1. Let Γ be a finitely generated nilpotent group of nilpotency class N_{Γ} at most 2. Consider a finitely generating set S and a finitely supported admissible symmetric probability measure μ on Γ . Then,

$$
\sum_{x \in S_n} G(e, x) \asymp n.
$$

In general, there exists $C \geq 1$ and $0 < \beta \leq 1$ that only depends on N_{Γ} such that

$$
\frac{1}{C}n \le \sum_{x \in S_n} G(e, x) \le Cn^{2-\beta}.
$$

Moreover, if $\Gamma = \mathbb{Z}^d$, $d \geq 3$ then there exists $C > 0$ such that

$$
\sum_{x \in S_n} G(e, x) \sim Cn.
$$

3.1. Random walks in \mathbb{Z}^d . We start with the following result.

Proposition 3.2. Let μ be a finitely supported symmetric admissible probability measure on \mathbb{Z}^d , $d \geq 3$. Endow \mathbb{Z}^d with a finite generating set S. Then, there exists $C > 0$ such that

$$
\sum_{x \in S_n} G(e, x) \sim Cn.
$$

Proof. Classical estimates show that $G(e, x) \sim C_0 ||x||^{-d+2}$, see [\[Woe00,](#page-51-0) Theorem 25.11. Here, $||x||$ is the norm of x given by the inverse of the covariance matrix, which is symmetric and defines a positive definite quadratic form Q associated with the random walk. These asymptotics actually hold in a more general setting, namely it is only required that μ has suited finite polynomial moments related to the dimension d, as proved by Uchiyama, see in particular [\[Uch98,](#page-51-4) Theorem 2]. The function $x \mapsto ||x||^{-d+2}$ is $(-d+2)$ -homogeneous in the sense that for $a > 0$, $\|ax\|^{-d+2} = a^{-d+2} \|x\|^{-d+2}$. Let us fix a generating set S for \mathbb{Z}^d . Then, letting S_n be the sphere of radius n centered at e with respect to S, we have by [\[DLM12,](#page-49-12) Theorem 1.1]

$$
\frac{1}{\sharp S_n} \sum_{x \in S_n} ||x||^{-d+2} \sim C_1 n^{-d+2}.
$$

Moreover, by [\[DLM12,](#page-49-12) Theorem 1.4],

$$
\sharp S_n \sim C_2 n^{d-1},
$$

hence

$$
\sum_{x \in S_n} ||x||^{-d+2} \sim C_3 n.
$$

Consequently,

$$
\sum_{x \in S_n} G(e, x) \sim C_4 n.
$$

For sake of completeness, let us consider the case of a (possibly lazy) simple random walk on \mathbb{Z}^d , endowed with the standard set of generators. The quadratic form Q associated with the covariance matrix introduced above is the Hessian at $u = 0$ of the function

$$
\Phi(u) = \sum_{x \in \mathbb{Z}^d} \mu(x) e^{u \cdot x},
$$

see [\[Woe00,](#page-51-0) Section 8 B, Section 13] for more details. Letting $(e_1, ..., e_d)$ be standard generators defined by $e_i = (0, ..., 0, 1, 0..., 0)$, where 1 is at the *i*th position and assuming that μ is of the form

$$
\mu = \alpha \delta_e + \frac{1 - \alpha}{2d} \sum_{i=1}^d (\delta_{e_i} + \delta_{-e_i}),
$$

we have

$$
\Phi(u) = \alpha + \frac{1-\alpha}{d} \sum_{i=1}^{d} \cosh(u_i),
$$

where u_i are the coordinates of u. In particular,

$$
Q(x) = \frac{1 - \alpha}{d} \langle x, x \rangle
$$

and so the explicit computations of [\[Woe00,](#page-51-0) Theorem 25.11] yield

$$
G(e,x) \sim \frac{\Gamma(\frac{d-2}{2})\pi^{-\frac{d}{2}}(1-\alpha)^{-d+3/2}}{2d^{-d+3/2}} \|x\|_2^{-d+2},
$$

where $\|\cdot\|_2$ is the Euclidean norm.

Furthermore, the constant in [\[DLM12,](#page-49-12) Theorem 1.1] is given by

$$
\int_L \|x\|^{-d+2} d\mu_L,
$$

where L is the boundary of the convex hull C of the generating set S and $d\mu$ _L is the cone volume on L . Applying this to our context where S is the standard generating set and $\|\cdot\|$ is the Euclidean norm $\|\cdot\|_2$, we get

$$
\frac{1}{\sharp S_n}\sum_{x\in S_n}\|x\|^{-d+2}\sim \int_{\partial B_1}\|x\|_2^{-d+2}d\mu_1 n^{-d+2},
$$

where B_1 is the unit ball for the $\|\cdot\|_1$ -norm $\{x,\sum |x_i|\leq 1\}$, ∂B_1 is the unit sphere $\{x,\sum |x_i| = 1\}$ and μ_1 is the uniform measure on ∂B_1 . Also, the constant in [\[DLM12,](#page-49-12) Theorem 1.4] is given by $d \cdot \text{vol}(\mathbf{C})$ and the volume of the ball B_1 is given by $\frac{2^d}{d!}$ $\frac{2^{\alpha}}{d!}$, see for instance [\[Bal97,](#page-49-13) Lecture 1]. We finally get

$$
\sum_{x \in S_n} G(e, x) \sim n \cdot \frac{2^d}{d!} \int_{\partial B_1} ||x||_2^{-d+2} d\mu_1 \frac{\Gamma(\frac{d-2}{2})\pi^{-\frac{d}{2}} (1-\alpha)^{-d+3/2}}{2d^{-d+1/2}}.
$$

3.2. Non-abelian nilpotent groups. In general, for nilpotent groups, we cannot prove such precise asymptotics. Let Γ be a finitely generated nilpotent group. Set $\Gamma^1 = \Gamma, \Gamma^2 = [\Gamma, \Gamma], ..., \Gamma^n = [\Gamma^{n-1}, \Gamma].$ Let N_{Γ} be the nilpotency class of Γ , that is N_{Γ} is the smallest n such that Γ^{n+1} is trivial. Note that the groups Γ^{n}/Γ^{n+1} are all finitely generated abelian groups and so have a well defined rank. The homogeneous dimension of Γ is defined as

$$
D = \sum_{n=1}^{N_{\Gamma}} n \operatorname{rank} \left(\Gamma^n / \Gamma^{n+1} \right).
$$

By $[Ale02, Theorem 1.8, Corollary 1.9], there exists c such that$

(3.1)
$$
\frac{1}{c} n^{-D/2} e^{-c\frac{|x|^2}{2n}} \leq p_n(e, x) \leq c n^{-D/2} e^{-\frac{1}{c}\frac{|x|^2}{2n}},
$$

where D is the homogeneous dimension of Γ. This is enough to deduce rough asymptotics for the Green function as we now show.

Lemma 3.3. For any positive a, b, c, there exists $K > 0$ such that as u tends to infinity,

$$
\sum_{n\geq 1} c n^{-a} e^{-bu/n} \sim K u^{-(a-1)}.
$$

Proof. The proof is adapted from [\[Woe00,](#page-51-0) Theorem 25.11]. We set $t_n = \frac{n}{u}$, so that

$$
\Delta_n = t_n - t_{n-1} = \frac{1}{u} \underset{u \to \infty}{\longrightarrow} 0.
$$

Then,

$$
\frac{1}{c}u^{a-1}\sum_{n\geq 1}cn^{-a}e^{-bu/n} = \sum_{n\geq 1}t_n^{-a}e^{-\frac{b}{t_n}}\Delta_n.
$$

The right hand-side is a Riemannian sum of

$$
\int_0^\infty t^{-a} \mathrm{e}^{-b/t} dt
$$

so it converges to a constant that only depends on a and b . \Box

Applying this lemma to $a = D/2$ and $u = |x|^2$, we deduce from [\(3.1\)](#page-11-0) that for every symmetric finitely supported admissible probability measure on a finitely generated nilpotent group,

(3.2)
$$
G(e, x) \asymp |x|^{-D+2}
$$
.

Guivarc'h [\[Gui73\]](#page-50-9) and Bass [\[Bas72\]](#page-49-14) independently proved that the homogeneous dimension D is the degree of the polynomial growth of Γ, that is, letting B_n be the ball of radius n ,

$$
\sharp B_n\asymp n^D.
$$

Conversely, by the landmark paper of Gromov [\[Gro81\]](#page-50-10), a finitely generated group of polynomial growth is virtually nilpotent. Extending the work of Gromov, Pansu [\[Pan83\]](#page-50-11) proved deep results about the asymptotic behavior of the word distance in terms of the geometry of the universal cover of Γ . As a particular case of his results, we have

$$
\sharp B_n \sim cn^D,
$$

see [\[Pan83,](#page-50-11) Proposition (5), Section (51)]. Although one might expect from these asymptotics that $\sharp S_n \sim c'n^{D-1}$, it is in fact much more difficult to obtain precise asymptotics for $\sharp S_n$. However, Breuillard and Le Donne proved the following.

Proposition 3.4. [\[BD13,](#page-49-15) Corollary 9, Corollary 11] Let Γ be a finitely generated nilpotent group endowed with a finite generating set. Let N_{Γ} be its nilpotency class.

(1) If $N_{\Gamma} \leq 2$, then

$$
\sharp B_n = Cn^D + O(n^{D-1})
$$

and

$$
\sharp S_n \asymp n^{D-1}.
$$

(2) In general, there exists $0 < \beta \leq 1$ that only depends on N_{Γ} such that

$$
\sharp B_n = Cn^D + O(n^{D-\beta})
$$

and there exists $C > 0$ such that

$$
\frac{1}{C}n^{D-1} \le \sharp S_n \le Cn^{D-\beta}.
$$

This proposition allows us to conclude the proof of Theorem [3.1.](#page-9-0)

Proof of Theorem [3.1.](#page-9-0) Combining [\(3.2\)](#page-12-0) and Proposition [3.4,](#page-12-1) we have for $N_{\Gamma} \leq 2$

$$
\sum_{x \in S_n} G(e, x) \asymp n
$$

and in general we have

$$
\frac{1}{C}n \le \sum_{x \in S_n} G(e, x) \le Cn^{2-\beta}.
$$

Together with Proposition [3.2,](#page-9-2) this concludes the proof of the theorem. \Box

It is also conjectured that one can always take $\beta = 1$ in the above result of Breuillard and Le Donne, see [\[BD13,](#page-49-15) Conjecture 10]. In such case, we would always have $H_1(n) \simeq n$. Conversely, note that since we already know that Gaussian estimates [\(3.1\)](#page-11-0) hold, the following conjecture is a reformulation of that of Breuillard and Le Donne.

Conjecture 3.5. Let Γ be a finitely generated nilpotent group and let μ be a symmetric admissible finitely supported probability measure and S be a finite generating set. Then,

$$
\sum_{x \in S_n} G(e, x) \asymp n.
$$

Finally, note that for the particular case of the Heisenberg group $\mathbb{H}_{3}(\mathbb{Z})$ whose homogeneous dimension is 4, the fact that $\sharp S_n \sim Cn^3$ for the standard set of generators is well known and was first proved by Shapiro, see [\[Sha89,](#page-51-5) Theorem, p.607]. See also [\[DM14,](#page-49-16) Theorem 25] which is analogous to [\[DLM12,](#page-49-12) Theorem 1.1] for abelian groups, which in turn is a key result that we used within the proof of Proposition [3.2.](#page-9-2) However, even for the Heisenberg group, it is unknown to our knowledge if one can replace $\frac{\alpha}{\alpha}$ with \sim in the asymptotics of the Green function [\(3.2\)](#page-12-0).

3.3. Polycyclic groups and groups of intermediate growth. Let us conclude this section with a small discussion on possible ways to generalize our results for nilpotent groups.

By a seminal result of Mal'cev [\[Mal51\]](#page-50-12), finitely generated torsion free nilpotent groups are exactly the lattices in simply connected nilpotent Lie groups, see also [\[Rag72,](#page-50-13) Theorem 2.18]. A further topic of interest would be the growth rate of the Green function for lattices in simply connected solvable Lie groups. By [\[Rag72,](#page-50-13) Theorem 4.28], these are exactly torsion free polycyclic groups. Moreover, every finitely generated polycyclic group has a subgroup of finite index which is torsion free by [\[Rag72,](#page-50-13) Lemma 4.6].

This gives motivation for studying $H_1(n)$ for polycyclic groups. The exponent 1/3 in Varapoulos' on-diagonal upper bound $p_n(e, e) \prec e^{-c_1 n^{1/3}}$ that we used within the proof of Proposition [2.3](#page-7-0) is known to be optimal for polycyclic groups of expo-nential growth. Indeed, by [\[Ale92,](#page-49-10) Theorem 1], we also have $p_n(e,e) \gtrsim e^{-c_2 n^{1/3}}$ for such groups. In order to get asymptotics of $H_1(n)$, optimal off-diagonal upper and lower bounds would be needed. However, as already mentioned, Gaussian-type estimates like [\(2.3\)](#page-8-0) cannot hold for groups of exponential volume growth and finding optimal upper and lower bounds would have to involve somehow the growth rate v. This in turn would require new material.

In another direction, it is not hard to see that the rough asymptotics of the Green function in Theorem [3.1](#page-9-0) hold for any virtually nilpotent groups. Indeed, let Γ be a virtually nilpotent and N be a finite index nilpotent subgroup of Γ . Consider a word distance $|\cdot|_{\Gamma}$ on Γ and a word distance $|\cdot|_{N}$ on N. Then, the restriction of $|\cdot|_{\Gamma}$ to N is bi-Lipschitz to $|\cdot|_N$. Moreover, if $|x|_{\Gamma}=n$, then x is within a uniform bounded distance of a point $y \in N$. In particular, $G(e, x) \approx G(e, y)$ and $|y|_N \approx n$. We find that

$$
\sum_{x \in \Gamma, |x|_{\Gamma} = n} G(e, x) \asymp \sum_{x \in N, |x|_{N} \asymp n} G(e, x).
$$

Moreover, $|\cdot|_{\Gamma}^{-D+2} \times |\cdot|_{N}^{-D+2}$ and given the asymptotics of Proposition [3.4,](#page-12-1) the sphere $S_n(\Gamma)$ for $|\cdot|_{\Gamma}$ and $S_n(N)$ for $|\cdot|_N$ satisfy that $\sharp S_n(\Gamma) \asymp \sharp S_n(N)$. Thus we have

$$
\sum_{x \in \Gamma, |x|_{\Gamma} = n} G(e, x) \asymp n^{-D+2} \sharp S_n(\Gamma) \asymp n^{-D+2} \sharp S_n(N).
$$

As explained above, virtually nilpotent groups are exactly the groups of polynomial growth, by a celebrated result of Gromov. Answering Conjecture [3.5](#page-13-0) would thus settle the case of polynomial growth. It would be interesting to see if the linear asymptotics of $H_1(n)$ also hold for groups of intermediate growth.

4. The lamplighter group and DL graphs

Beyond nilpotent groups, studying the growth rate and the asymptotics of $H_1(n)$ for general finitely generated solvable groups seems to be a difficult task. Our next goal is to compute precise such asymptotics for the lamplighter group, which is the first step into this direction.

The lampligther group L_q is the restricted wreath product $\mathbb{Z}_q \wr \mathbb{Z}$. The study of this group is a classical topic in geometric group theory and we refer to [\[CM17,](#page-49-17) Lecture 15 for an overview of its properties. The usual description of L_q is as follows. Consider the line $\mathbb Z$ and suppose that there is a lamp at each site of $\mathbb Z$. Every lamp can be either off or lit with $q-1$ different colors, so that it has q different possible states. At the beginning, every lamp is switched off. The lampligther is at the origin of Z and can either change the state of the lamp where they stand or take one step to the left or to the right in Z. Those possible actions are considered as the generators of the group. Elements of L_q are then described by the position of the lamplighter on $\mathbb Z$ and the states of the lamps, which can be seen as a function $\mathbb{Z} \to \mathbb{Z}_q$ with finite support.

There is a one-to-one correspondence between the lamplighter group $\mathbb{Z}_q \wr \mathbb{Z}$ and the vertices of the Diestel-Leader graph $DL(q, q)$, which is defined as the following horocycle product. Consider the $(q+1)$ -regular tree \mathbb{T}_q and let h be a horofunction based at some fixed chosen point at infinity (see [\[BW05,](#page-49-7) Section 2] for a precise definition and also [\[CM17,](#page-49-17) Section 15.4], where h is called a height function). Then,

$$
DL(q, q) = \{(x_1, x_2), x_i \in \mathbb{T}_q, h(x_1) = -h(x_2)\}.
$$

Thus, any element x of $\mathbb{Z}_q \wr \mathbb{Z}$ can be written as $x = (x_1, x_2), x_i \in \mathbb{T}_q$.

Following [\[BW05\]](#page-49-7), what we call here the simple random walk on L_q is the simple random walk on $DL(q, q)$ endowed with the product graph distance. In other words, it is the simple random walk on the graph whose vertices are elements of $DL(q, q)$ and two vertices (x_1, x_2) and (y_1, y_2) are connected by an edge if x_i and y_i are connected with an edge in \mathbb{T}_q for $i = 1, 2$. This is exactly the simple random walk on L_q endowed with an appropriate set of generators described in [\[BW05,](#page-49-7) Section 2], see in particular [\[BW05,](#page-49-7) (2.3)] for a formula for the probability measure μ (with $\alpha = \frac{1}{2}$ since we do not consider biased random walks here).

In terms of lamplighting, we need to change a bit the above description to understand this random walk. Think of the lamps not placed at each vertex of $\mathbb Z$ but at the middle of each edge. Suppose the current position of the lamplighter is $k \in \mathbb{Z}$. They first toss a coin. If "head" comes up, they move to $k+1$ and switch the lamp on the transverse edge to a state chosen at random in \mathbb{Z}_q . Otherwise, they move to $k-1$ and also switch the lamp on the transverse edge to a random state.

Lamplighter groups provide a great source of examples of particular asymptotic behaviors of random walks, see [\[KV83\]](#page-50-14), [\[Ers01\]](#page-50-15), [\[Ers03\]](#page-50-16), [\[Rev03a\]](#page-50-17), [\[Rev03b\]](#page-50-18), [\[BNW06\]](#page-49-18) just to name a few. We prove here that $H_1(n)$ is still asymptotically linear in n for the simple random walk. Our study is based on the renewal theorems of [\[BW05\]](#page-49-7), see also references therein for other significant results related to random walks on these groups.

Theorem 4.1. Consider the simple random walk on L_q . There exists C such that

$$
\sum_{x \in S_n} G(e, x) \sim Cn.
$$

We identify $L_q = \mathbb{Z}_q \wr \mathbb{Z}$ with $DL(q, q)$ as above, with h a fixed horofunction. By [\[BW05,](#page-49-7) (4.1)], the word distance in L_q is given by

$$
d(x, y) = d(x_1, y_1) + d(x_2, y_2) - |h(y_1) - h(x_1)|.
$$

Note that $h(y_2) - h(x_2) = -h(y_1) + h(x_1)$, so

$$
d(x, y) = d(x_1, y_1) + d(x_2, y_2) - |h(y_2) - h(x_2)|.
$$

The identity element e is identified with (o_1, o_2) in $DL(q, q)$, where o_i is the root of \mathbb{T}_q . In particular, we have $h(o_1) = h(o_2) = 0$. For $(x_1, x_2) \in DL(q, q)$, we now set, following [\[BW05\]](#page-49-7),

$$
|x_i| = d(o_i, x_i) = \mathfrak{d}_i + \mathfrak{u}_i,
$$

$$
h(x_i) = \mathfrak{d}_i - \mathfrak{u}_i
$$

and

$$
\mathfrak{s} := \mathfrak{u}_1 + \mathfrak{u}_2 = \mathfrak{d}_1 + \mathfrak{d}_2.
$$

The quantities $\mathfrak d$ and $\mathfrak u$ are defined in terms of geometric features of $\mathbb T_q$, see [\[BW05,](#page-49-7) Section 2, Figure 1 for more details. The elements x_1 and x_2 are completely determined by $|x_1|, |x_2|$ and $h(x_1) = -h(x_2)$. Thus, any element (x_1, x_2) is completely determined by u_1 , \mathfrak{d}_2 and \mathfrak{s} .

Let us now describe the sphere of radius n in L_q . By what precedes,

$$
d(e,x) = |x_1| + |x_2| - |h(x_1)| = \mathfrak{u}_1 + \mathfrak{d}_1 + \mathfrak{u}_2 + \mathfrak{d}_2 - |\mathfrak{d}_1 - \mathfrak{u}_1|.
$$

We first find the $x \in S_n$ with $h(x_1) > 0$. For such x,

$$
d(e,x)=2\mathfrak{u}_1+\mathfrak{u}_2+\mathfrak{d}_2=\mathfrak{u}_1+\mathfrak{d}_1+2\mathfrak{d}_2=\mathfrak{u}_1+\mathfrak{d}_2+\mathfrak{s}.
$$

We write $\tilde{S}(n,m)$ the set of $z \in \mathbb{T}_q$ with $|z|=n$ and $h(z)=m, -n \leq m \leq n$, and so $|n-m|$ needs to be even. Let us compute the cardinality of $\tilde{S}(n, m)$ for small values of n, m, when $q = 3$, i.e. \mathbb{T}_q is the 4-regular tree, which is the Cayley graph of F_2 . We write $F_2 = \langle a, b \rangle$ and we fix the point at infinity defining the horofunction h to be a^{∞} . We have $\tilde{S}(1,1) = \{a^{-1}, b, b^{-1}\}$ and $\tilde{S}(1,-1) = \{a\}$, so

$$
\sharp \tilde{S}(1,1) = 3, \; \sharp \tilde{S}(1,-1) = 1.
$$

For $n = 2$, we have $\tilde{S}(2, 2) = \{a^{-2}, a^{-1}b, a^{-1}b^{-1}, ba, ba^{-1}, b^2, b^{-1}a, b^{-1}a^{-1}, b^{-2}\},$ $\tilde{S}(2,0) = \{ab, ab^{-1}\}\$ and $\tilde{S}(2,-2) = \{a^2\}$. Therefore,

$$
\sharp \tilde{S}(2,2) = 3^2, \; \sharp \tilde{S}(2,0) = 2, \; \sharp \tilde{S}(2,-2) = 1.
$$

Similarly, we prove that

$$
\sharp \tilde{S}(3,3) = 3^3, \; \sharp \tilde{S}(3,1) = 2 \times 3, \; \sharp \tilde{S}(3,-1) = 2, \; \sharp \tilde{S}(3,-3) = 1,
$$

 $\sharp \tilde{S}(4,4) = 3^4$, $\sharp \tilde{S}(4,2) = 2 \times 3^2$, $\sharp \tilde{S}(4,0) = 2 \times 3$, $\sharp \tilde{S}(4,-2) = 2$, $\sharp \tilde{S}(4,-4) = 1$. In general, we have for $0 \leq k \leq n$

$$
\sharp \tilde{S}(n,n) = q^n, \; \sharp \tilde{S}(n,-n) = 1
$$

and

$$
\sharp \tilde{S}(n, n-2k) = (q-1) \times q^{n-1-k}, 1 \le k \le n-1.
$$

Note that if $z \in \tilde{S}(n, n-2k)$, then $\mathfrak{u} = k$ and $\mathfrak{d} = n-k$. In fact, \mathfrak{u} is the number of steps upward in direction to a^{∞} and $\mathfrak d$ is the number of steps downward.

We now consider the set $A(i, j, k) = \{(x_1, x_2), u_1 = i, \mathfrak{d}_2 = j, \mathfrak{s} = k\}$, with $i \leq k$, $j \leq k$ and $i < k - j$ (the last condition being equivalent to $h(x_1) > 0$). Then, by what precedes, for $0 < j < k$,

$$
\sharp A(0,j,k) = (q-1)q^{k-1}, \; \sharp A(k,j,k) = (q-1)q^{k-1}
$$

and for $0 < i < k$,

$$
\sharp A(i,j,k) = (q-1)^2 q^{k-2}.
$$

Now, for $j = 0$, we have

$$
\sharp A(0,0,k) = q^k, \; \sharp A(k,0,k) = 1
$$

and for $0 < i < k$,

$$
\sharp A(i,0,k) = (q-1)q^{k-1}.
$$

Finally, for $j = k$, we have

$$
\sharp A(0,k,k) = (q-1)q^{k-1}, \; \sharp A(k,k,k) = q^k,
$$

and for $0 < i < k$,

$$
\sharp A(i,k,k) = (q-1)q^{k-1}.
$$

We are now ready to prove Theorem [4.1.](#page-15-0)

Proof. The following asymptotics of the Green function are proven in [\[BW05,](#page-49-7) Theorem 4.2]. We have, as $|x|$ tends to infinity,

$$
(4.1) \ \ G(e,x)\sim \frac{C}{\mathfrak{s}^4 q^{\mathfrak{s}}}\bigg(\frac{q+1}{q-1}\big(\mathfrak{u}_1(\mathfrak{s}-\mathfrak{d}_2)+(\mathfrak{s}-\mathfrak{u}_1)\mathfrak{d}_2\big)+\mathfrak{s}\mathfrak{u}_1\mathfrak{d}_2+\mathfrak{s}(\mathfrak{s}-\mathfrak{u}_1)(\mathfrak{s}-\mathfrak{d}_2)\bigg).
$$

We fix $(x_1, x_2) \in S_{2n}$ and $h(x_1) > 0$, so $2n = \mathfrak{s} + \mathfrak{d}_2 + \mathfrak{u}_1$ and $\mathfrak{d}_2 + \mathfrak{u}_1 < \mathfrak{s}$. In particular, $s > n$. Also, since $\mathfrak{d}_2, \mathfrak{u}_1 \geq 0$, we have $s \leq 2n$. Conversely, choose any couple $(\mathfrak{d}_2, \mathfrak{s})$ satisfying $n + 1 \leq \mathfrak{s} \leq 2n$ and $0 \leq \mathfrak{d}_2 \leq 2n - \mathfrak{s}$. Set $\mathfrak{u}_1 = 2n - \mathfrak{s} - \mathfrak{d}_2$. Note that $\mathfrak{s} > n$ implies that $2n - \mathfrak{s} \leq \mathfrak{s} - 1$, so $0 \leq \mathfrak{u}_1 \leq \mathfrak{s} - 1 - \mathfrak{d}_2$ and $0 \leq \mathfrak{d}_2 < \mathfrak{s}$. We find that $u_1 < \mathfrak{s} - \mathfrak{d}_2$ and so any such triple $(u_1, \mathfrak{d}_2, \mathfrak{s})$ defines a point (x_1, x_2) with $h(x_1) > 0$. By [\(4.1\)](#page-16-0), setting $u_1 = i$, $v_2 = j$, $v_3 = k$ we get the following. The sum of the Green function along points x in the sphere S_{2n} satisfying $h(x_1) > 0$ can be written as

$$
\sum_{\substack{(x_1,x_2)\in S_{2n},\\h(x_1)>0}} G(e,x) \sim \sum_{k=n+1}^{2n} \sum_{j=0}^{2n-k} \frac{C}{k^4 q^k} \sharp A(i,j,k)
$$

$$
\left(\frac{q+1}{q-1} (i(k-j)+(k-i)j) + kij + k(k-i)(k-j)\right).
$$

Thus, replacing u_1 with $2n - s - \mathfrak{d}_2$, we find

$$
\sum_{\substack{(x_1,x_2)\in S_{2n},\\h(x_1)>0}} G(e,x) \sim \sum_{k=n+1}^{2n} \sum_{j=0}^{2n-k} \frac{C}{k^4 q^k} \sharp A(2n-k-j,j,k)
$$

$$
\left(\frac{q+1}{q-1}((2n-k-j)(k-j)+(2k+j-2n)j)+k(2n-k-j)(k-j)\right).
$$

Since the dominant term in $\sharp A(i,j,k)$ is q^k , we find

$$
\sum_{\substack{(x_1, x_2) \in S_{2n}, \ x_1 \ge 0}} G(e, x) \sim C_1 \sum_{k=n+1}^{2n} \sum_{j=0}^{2n-k} \frac{1}{k^4} \left((2n - k - j)(k - j) + (2k + j - 2n)j + k(2k + j - 2n)(k - j) \right)
$$

$$
\sim C_2 n.
$$

By symmetry, we have the same asymptotics for $h(x_1) < 0$. Now, if $h(x_1) = 0$, then $u_1 = \mathfrak{s} - \mathfrak{d}_2$. Combining this with $u_1 + \mathfrak{d}_2 + \mathfrak{s} = 2n$, we get $\mathfrak{s} = n$, so the sum of the Green function along S_{2n} , assuming further $h(x_1) = 0$ is asymptotic to a constant C₃. Thus, we find that $\sum_{x \in S_{2n}} G(e, x)$ is linear in n. The same proof with more delicate sums to handle, when considering the integer part of $n/2$, shows that the same is true for S_{2n+1} . This concludes the proof of Theorem [4.1.](#page-15-0)

Lamplighter groups are solvable and thus amenable. They are actually classical examples of non-polycyclic solvable groups, see for instance the comments at the end of [\[DK18,](#page-49-19) Chapter 13]. In light of the discussion in Section [3.3,](#page-13-1) this raises the following question.

Question 4.2. Is it true that

$$
\sum_{x \in S_n} G(e, x) \asymp n
$$

holds for every symmetric finitely supported admissible probability measure on a finitely generated amenable group ? If not, can a counter-example be found among finitely generated solvable groups ?

Also, let us conclude this section with further comments on horocycle products. We refer to [\[Woe13\]](#page-51-6) for a more complete exposition. Given two hyperbolic spaces X_1, X_2 , one can perform a construction similar to that of $DL(q, q)$ by choosing two Busemann functions h_1 , respectively h_2 , based at infinity on X_1 , respectively X_2 . The horocycle product of X_1 and X_2 is the set of pairs $(x_1, x_2) \in X_1 \times X_2$ with $h_1(x_1) + h_2(x_2) = 0$. Interesting examples of such spaces are the horocycle product $HT(p, q)$ of $\mathbb{H}(p)$ with a homogeneous tree of degree $q+1$ and the horocycle product $Sol(p, q)$ of $\mathbb{H}(p)$ and $\mathbb{H}(q)$. In both cases, $\mathbb{H}(p)$ is the standard Poincaré upper halfspace with suited rescaling of the hyperbolic metric. Amenable Baumslag-Solitar groups $BS(1, q)$ act properly discontinuously and co-compactly via isometries on $HT(q, q)$, see [\[FM98\]](#page-50-19). On the other hand, the spaces $Sol(p, q)$ are classical examples of solvable Lie groups and play an important role in Thurston's geometrization theorem. We refer to [\[BSW12\]](#page-49-20) and references therein for further details. A next

subject of interest would be the study of $H_1(n)$ on amenable Baumslag-Solitar groups and on lattices of $Sol(p, q)$. Note that the later are examples of polycyclic groups.

5. Cartesian products of trees

In this section, we study the behavior of $H_r(n)$ for random walks on $T \times T'$, where T, T' are regular trees. The asymptotics for the Green functions are given by the work of Picardello and Woess [\[PW94\]](#page-50-8), see also [\[Woe00,](#page-51-0) Section 28] and references therein for the particular case of $T \times \mathbb{Z}$.

Let T_1, T_2 be regular trees of degree $l_1, l_2 \geq 3$. We consider the lazy simple random walk μ_i on T_i whose transition kernel $p_i(x, y)$ is defined by

$$
p_i(x, y) = \begin{cases} \frac{1}{2l_i} \text{ if } x, y \text{ are connected with an edge in } T_i, \\ \frac{1}{2} \text{ if } x = y, \\ 0 \text{ otherwise.} \end{cases}
$$

In particular, μ_i is an admissible symmetric finitely supported probability measures on T_i . For every $\alpha_1, \alpha_2 \geq 0$, $\alpha_1 + \alpha_2 = 1$, we let μ be the probability measure on $T_1 \times T_2$ given by

$$
\mu = \alpha_1 \mu_1 + \alpha_2 \mu_2.
$$

In terms of Markov operators, this means that

$$
P_{\mu} = \alpha_1 P_{\mu_1} \otimes I + \alpha_2 I \otimes P_{\mu_2}.
$$

As noted in [\[PW94,](#page-50-8) Section 3], the lazy simple random walk on $T_1 \times T_2$ with $\mu(x,x) = 1/2$ is given by $\alpha_i = \frac{l_i}{l_1+l_2}$. Set $\rho_i = \frac{1}{2} + \frac{\sqrt{l_i-1}}{l_i}$ and $R = \frac{1}{\alpha_1 \rho_1 + \alpha_2 \rho_2}$. Then, ρ_i is the spectral radius of μ_i , see [\[PW94,](#page-50-8) (2.2)] and $\rho = R^{-1}$ is the spectral radius of μ , see [\[PW94,](#page-50-8) Section 3]. We prove here the following.

Theorem 5.1. If $l_1 = l_2$, then for every $r < R$, we have $H_r(n) \approx e^{n\omega_{\Gamma}(r)}$. If $l_1 > l_2$, then there exists a phase transition at some $r_0 \in (1, R)$ such that the following holds.

- For every $r < r_0$, we have $H_r(n) \approx e^{n\omega_{\Gamma}(r)}$.
- At $r = r_0$, we have $H_r(n) \asymp n^{-1} e^{n\omega_{\Gamma}(r)}$.
- For every $r_0 < r < R$, we have $H_r(n) \approx n^{-3/2} e^{n\omega_{\Gamma}(r)}$.

The remainder of the section is devoted to the proof of this theorem. By [\[PW94,](#page-50-8) Theorem 3.1], for every $r < R$, for every $\lambda_0 \geq 0$, there exist r_1, r_2 such that as $x = (x_1, x_2) \in T_1 \times T_2$ tends to infinity and $|x_2|/|x_1| = \lambda$ converges λ_0 , we have

$$
G(e, x|r) \sim G_1(e, x_1|r_1)G_2(e, x_2|r_2)\left(|x_1| + \frac{l_1}{l_1 - 2}\right)\left(|x_2| + \frac{l_2}{l_2 - 2}\right)\frac{1}{|x_1|^{5/2}}C(\lambda),
$$

where $C(\lambda)$ is a continuous positive function.

The numbers r_1 and r_2 are the unique solutions of the system

(5.2)
$$
\begin{cases} \alpha_1 r_1^{-1} + \alpha_2 r_2^{-1} = r^{-1} \\ \alpha_2 \sqrt{(r_2^{-1} - \frac{1}{2})^2 - \frac{l_2 - 1}{l_2^2}} = \lambda \alpha_1 \sqrt{(r_1^{-1} - \frac{1}{2})^2 - \frac{l_1 - 1}{l_1^2}} \end{cases}
$$

By symmetry, the same holds when $|x_1|/|x_2| = \lambda'$ converges to $\lambda'_0 \in [0, +\infty)$, switching the indices 1 and 2 and replacing the function C with a function C' which is also continuous and positive.

Set
$$
\beta_i = \frac{\sqrt{l_i - 1}}{l_i}
$$
 and
\n
$$
F_i(r) = \frac{l_i}{l_i - 1} \left(r^{-1} - \frac{1}{2} - \sqrt{\left(r^{-1} - \frac{1}{2} \right)^2 - \beta_i^2} \right),
$$
\n
$$
G_i(r) = \frac{r^{-1}}{r^{-1} - (1/2)(1 + F_i(r))}.
$$

By $[PW94, (2.4)]$, we have that

$$
G_i(e_i, x_i|r) = G_i(r)F_i(r)^{|x_i|},
$$

hence in particular $G_i(e_i, e_i | r) = G_i(r)$. Since $1 \le G_i(r) \le G_i(\rho_i^{-1})$, we have (5.3) $n^{i}(r) = \sum$ $x_i \in T_i : |x_i|=n$ $G_i(e_i, x_i|r) = l_i(l_i - 1)^{n-1} G_i(r) F_i(r)^n \approx e^{n\omega_{T_i}(r)},$

where

$$
\omega_{T_i}(r) = \log(l_i - 1) + \log F_i(r).
$$

We now fix $r < R$ and we consider $n \geq 0$. The sphere S_n in $T_1 \times T_2$ can be decomposed as

$$
S_n = \bigcup_{k=0}^n S_k^1 \times S_{n-k}^2.
$$

Then, for every $k \leq n$, there exists a unique couple $(r_1(\lambda), r_2(\lambda))$ satisfying [\(5.2\)](#page-18-2) with $\lambda = \frac{n-k}{k}$. Moreover,

$$
\frac{1}{|x_1|^{5/2}}C(\lambda) = \frac{1}{(|x_1| + |x_2|)^{5/2}}(1 + \lambda)^{5/2}C(\lambda)
$$

and for $k \geq n/2$, $\lambda \leq 1$, hence by the continuity of $C(\lambda)$,

$$
\frac{1}{|x_1|^{5/2}}C(\lambda) \asymp \frac{1}{(|x_1| + |x_2|)^{5/2}}.
$$

Similarly,

$$
\frac{1}{|x_2|^{5/2}}C'(\lambda') = \frac{1}{(|x_1|+|x_2|)^{5/2}}(1+\lambda')^{5/2}C'(\lambda')
$$

and so for $k \leq n/2$,

$$
\frac{1}{|x_2|^{5/2}}C'(\lambda') \asymp \frac{1}{(|x_1|+|x_2|)^{5/2}}.
$$

Setting $\kappa_i = \frac{l_i}{l_i - 2}$, we thus have

$$
\sum_{x \in S_n} G(e, x|r) \asymp \frac{1}{n^{5/2}} \sum_{k=0}^n \sum_{x_1 \in S_k^1} \sum_{x_2 \in S_{n-k}^2} G_1(e, x_1|r_1(\lambda))G_2(e, x_2|r_2(\lambda))
$$
\n
$$
\asymp \frac{1}{n^{5/2}} \sum_{k=0}^n (k + \kappa_1)((n - k) + \kappa_2)H_k^1(r_1(\lambda))H_{n-k}^2(r_2(\lambda)).
$$

Applying [\(5.3\)](#page-19-0), we see that

(5.4)
$$
H_n(r) \approx \frac{1}{n^{5/2}} \sum_{k=0}^n (k + \kappa_1)((n - k) + \kappa_2) \exp(n\Psi(\lambda)),
$$

with $\lambda = \frac{n-k}{k}$ and

$$
\Psi(\lambda)=\frac{1}{1+\lambda}\omega_{T_1}(r_1(\lambda))+\frac{\lambda}{1+\lambda}\omega_{T_2}(r_2(\lambda)).
$$

In order to find the asymptotics of $H_n(r)$, we thus need to find where the function $\Psi(\lambda)$ takes its maximum value.

Now let $t = r^{-1} - \frac{1}{2}$ and $t_i = t_i(\lambda) = r_i(\lambda)^{-1} - \frac{1}{2}$. Then (t_1, t_2) solves the system of equations

(5.5)
$$
\begin{cases} \alpha_1 t_1 + \alpha_2 t_2 = t, \\ \alpha_2 \sqrt{t_2^2 - \beta_2^2} = \lambda \alpha_1 \sqrt{t_1^2 - \beta_1^2}. \end{cases}
$$

Lemma 5.2. The functions $\lambda \mapsto t_1(\lambda)$ and $\lambda \mapsto t_2(\lambda)$ are continuously differentiable. Furthermore,

$$
t'_{1}(\lambda) = -\frac{\lambda \alpha_{1}(t_{1}^{2} - \beta_{1}^{2})}{\alpha_{2}t_{2} + \lambda^{2} \alpha_{1}t_{1}},
$$

$$
t'_{2}(\lambda) = -\frac{\alpha_{1}}{\alpha_{2}}t'_{1}(\lambda) = \frac{\lambda^{-1} \alpha_{2}(t_{2}^{2} - \beta_{2}^{2})}{\alpha_{2}t_{2} + \lambda^{2} \alpha_{1}t_{1}}.
$$

Proof. Let U be the open set $(\beta_1, +\infty) \times (\beta_2, +\infty) \times (\beta, +\infty) \times (0, +\infty)$ with $\beta = \rho - \frac{1}{2}$. We set

$$
\Upsilon : (t_1, t_2, t, \lambda) \in U \mapsto \left(\alpha_1 t_1 + \alpha_2 t_2 - 2t, \alpha_2 \sqrt{t_2^2 - \beta_2^2} - \lambda \alpha_1 \sqrt{t_1^2 - \beta_1^2}\right).
$$

Then

$$
\frac{\partial \Upsilon}{\partial t} = (-2,0)
$$

and

$$
\frac{\partial \Upsilon}{\partial \lambda} = \left(0, -\alpha_1 \sqrt{t_1^2 - \beta^2}\right).
$$

For $t_1 > \beta_1$, the matrix

$$
\begin{pmatrix}\n-2 & 0 \\
0 & -\alpha_1 \sqrt{t_1^2 - \beta^2}\n\end{pmatrix}
$$

is invertible. The implicit function theorem shows that the solution (t_1, t_2) of (5.5) is continuously differentiable in the variables (t, λ) . The formulas for $t'_1(\lambda)$ and $t'_{2}(\lambda)$ are then derived from [\(5.5\)](#page-20-0).

Define

$$
\varphi_i(t) = \log l_i + \log \left(t - \sqrt{t^2 - \beta_i^2} \right).
$$

Then

$$
\Psi(\lambda) = \frac{1}{1+\lambda}\varphi_1(t_1(\lambda)) + \frac{\lambda}{1+\lambda}\varphi_2(t_2(\lambda)).
$$

Since

(5.6)
$$
\varphi_i'(t) = -\frac{1}{\sqrt{t^2 - \beta_i^2}},
$$

we have that

$$
\lambda \varphi_2'(t_2(\lambda)) = \frac{\alpha_2}{\alpha_1} \varphi_1'(t_1(\lambda)),
$$

and hence

(5.7)
$$
\Psi'(\lambda) = \frac{\varphi'_1(t_1(\lambda))}{1+\lambda} t'_1(\lambda) + \frac{\lambda \varphi'_2(t_2(\lambda))}{1+\lambda} t'_2(\lambda) + \frac{\varphi_2(t_2(\lambda)) - \varphi_1(t_1(\lambda))}{(1+\lambda)^2} \n= \frac{\alpha_1 t'_1(\lambda) + \alpha_2 t'_2(\lambda)}{\alpha_1 (1+\lambda)} \varphi'_1(t_1(\lambda)) + \frac{\varphi_2(t_2(\lambda)) - \varphi_1(t_1(\lambda))}{(1+\lambda)^2} \n= \frac{\varphi_2(t_2(\lambda)) - \varphi_1(t_1(\lambda))}{(1+\lambda)^2}.
$$

Furthermore,

(5.8)
$$
\Psi''(\lambda) = -\frac{2}{(1+\lambda)^3} \left[\varphi_2(t_2(\lambda)) - \varphi_1(t_1(\lambda)) \right] - \frac{\alpha_1 \sqrt{t_1^2 - \beta_1^2}}{(1+\lambda) (\alpha_2 t_2 + \lambda^2 \alpha_1 t_1)}.
$$

By Lemma [5.2](#page-20-1) and [\(5.6\)](#page-20-2), we see that $\varphi_1(t_1(\lambda))$ (resp. $\varphi_2(t_2(\lambda))$) is strictly increasing (resp. decreasing) in λ . It follows that there is at most one $\lambda_0 \in [0, +\infty)$ such that $\Psi'(\lambda_0) = 0$. Note that $t_2(0) = \beta_2$, $t_1(0) = \alpha_1^{-1} (t - \alpha_2 \beta_2) > \beta_1$, and

$$
\varphi_2(t_2(0)) = \frac{1}{2} \log (l_2 - 1).
$$

On the other hand, $\lim_{\lambda \to +\infty} t_1(\lambda) = \beta_1$, $\lim_{\lambda \to +\infty} t_2(\lambda) = \alpha_2^{-1}(t - \alpha_1 \beta_1) > \beta_2$, and

$$
\lim_{\lambda \to +\infty} \varphi_1(t_1(\lambda)) = \frac{1}{2} \log (l_1 - 1).
$$

Assume $l_1 = l_2 = l$. Then for $\lambda_0 = \frac{\alpha_2}{\alpha_1}$ we have that $t_1(\lambda_0) = t_2(\lambda_0) = t$ and $\Psi'(\lambda_0)=0$. Note that

$$
\Psi''(\lambda_0) = -\frac{\alpha_1^2 \sqrt{t^2 - \beta^2}}{\alpha_2 t} < 0
$$

by Lemma [5.2](#page-20-1) and [\(5.6\)](#page-20-2). By Lemma [5.3](#page-22-0) (ii) below, we can deduce that

(5.9)
$$
H_n(r) \approx e^{n\Psi(\lambda_0)} = e^{n \log \left[l\left(r^{-1} - \frac{1}{2} - \sqrt{\left(r^{-1} - \frac{1}{2}\right)^2 - \beta^2}\right)\right]}.
$$

This concludes the proof of Theorem [5.1](#page-18-0) for the case $l_1 = l_2$.

Assume now that $l_1 > l_2$. Then, we have that $\beta_1 < \beta_2$ and $\varphi_1(\beta_1) > \varphi_2(\beta_2)$. Since $\lim_{s\to\infty}\varphi_1(s)=-\infty$, there exists $t_0>\alpha_1\beta_1+\alpha_2\beta_2>\beta_1$ such that

$$
\varphi_1\left(\alpha_1^{-1}\left(t_0-\alpha_2\beta_2\right)\right)=\varphi_2(\beta_2).
$$

If $t < t_0$, then

$$
\varphi_1(t_1(0)) = \varphi_1(\alpha_1^{-1} (t - \alpha_2 \beta_2)) > \varphi_1(\alpha_1^{-1} (t_0 - \alpha_2 \beta_2)) = \varphi_2(t_2(0))
$$

and hence $\Psi'(\lambda) < 0$ for all $\lambda \geq 0$. It follows that $\Psi(\lambda)$ takes its unique maximum at $\lambda = 0$. By Lemma [5.3](#page-22-0) (i) below,

(5.10)
$$
H_n(r) \approx n^{-3/2} e^{n \log \varphi_1(\alpha_1^{-1}(t-\alpha_2\beta_2))}.
$$

Similarly, if $t = t_0$, then 0 is also the unique maximum point of $\Psi(\lambda)$, and we have further that $\Psi'(0) = 0$,

$$
\Psi''(0) = -\frac{\sqrt{(t_0 - \alpha_2 \beta_2)^2 - \alpha_1 \beta_1^2}}{\alpha_2 \beta_2} < 0.
$$

Thus, Lemma [5.3](#page-22-0) (iii) below shows that

(5.11)
$$
H_r(n) \asymp n^{-1} e^{n\Psi(0)} = n^{-1} e^{n \log \varphi_1(\alpha_1^{-1}(t - \alpha_2 \beta_2))}.
$$

It remains to consider the case that $t > t_0$. Since $t_0 > \alpha_1 \beta_1 + \alpha_2 \beta_2$ we have

$$
\lim_{\lambda \to +\infty} \varphi_2(t_2(\lambda)) = \varphi_2\left(\alpha_2^{-1} \left(t - \alpha_1 \beta_1\right)\right) < \varphi_2(\beta_2) < \varphi_1(\beta_1) = \lim_{\lambda \to +\infty} \varphi_1(t_1(\lambda)),
$$

and

$$
\varphi_1(t_1(0)) = \varphi_1\left(\alpha_1^{-1}\left(t - \alpha_2\beta_2\right)\right) < \varphi_1\left(\alpha_1^{-1}\left(t_0 - \alpha_2\beta_2\right)\right) = \varphi_2(t_2(0)).
$$

Thus there exists $\lambda_0 > 0$ such that $\varphi_1(t_1(\lambda_0)) = \varphi_2(t_2(\lambda_0))$. By [\(5.8\)](#page-21-0), $\Psi''(\lambda_0) < 0$, and we see from Lemma [5.3](#page-22-0) (ii) below that

(5.12)
$$
H_r(n) \approx e^{n\varphi_1(t_1(\lambda_0))}.
$$

To conclude, what is left to do is proving that $r_0 > 1$, i.e. $t_0 < 1/2$. Lengthily computations would prove that

$$
\varphi_1\left(\alpha_1^{-1}\left(\frac{1}{2}-\alpha_2\beta_2\right)\right) < \varphi_2(\beta_2),
$$

hence we necessarily have $t_0 < 1/2$. However, we see that at t_0 , we have by (5.11) $H_r(n) \approx n^{-1} e^{n\Psi(0)} = n^{-1} e^{n \log \varphi_1(\alpha_1^{-1}(t-\alpha_2\beta_2))}$. In particular,

$$
\omega_{\Gamma}(r_0) = \log \varphi_1 \left(\alpha_1^{-1} \left(t_0 - \alpha_2 \beta_2 \right) \right) = \log \varphi_2(\beta_2) > 0.
$$

Since $\omega_{\Gamma}(1) = 0$ and ω_{Γ} is increasing, we see directly that $r_0 > 1$. This ends the proof of Theorem [5.1.](#page-18-0) \Box

Lemma 5.3. Assume that $\Phi \in C^2([0, +\infty))$ is eventually decreasing and has a unique maximum point at $0 \leq \lambda_0 < \infty$. Define

$$
f(n) = \sum_{k=0}^{n} k(n-k)e^{n\Phi\left(\frac{n-k}{k}\right)}.
$$

(i) If $\lambda_0 = 0$ and $\Phi'(0) < 0$, then

$$
f(n) \asymp n e^{n\Phi(0)}.
$$

(ii) If $\lambda_0 > 0$ and $\Phi''(\lambda_0) < 0$, then

$$
f(n) \asymp n^{5/2} e^{n\Phi(\lambda_0)}.
$$

(iii) If $\lambda_0 = 0$, $\Phi'(0) = 0$ and $\Phi''(0) < 0$, then $f(n) \approx n^{3/2} e^{n\Phi(0)}$.

Proof. (i) For any $0 < \varepsilon < -\Phi'(0)$, there exists $\delta > 0$ such that $|\Phi'(\lambda) - \Phi'(0)| <$ $ε$ for every $0 ≤ λ ≤ δ$. By the mean value theorem, for $λ ≤ δ$,

$$
(\Phi'(0)-\varepsilon)\lambda \leq \Phi(\lambda)-\Phi(0) \leq (\Phi'(0)+\varepsilon)\lambda.
$$

If $\frac{n-k}{k} \leq \delta$, then we have $k \geq (1+\delta)^{-1}n$. Thus

$$
f(n) \geq \sum_{k \geq (1+\delta)^{-1}n} k(n-k)e^{n\Phi(\frac{n-k}{k})}
$$

\n
$$
\succ n e^{n\Phi(0)} \sum_{k \geq (1+\delta)^{-1}n} (n-k)e^{n(\Phi'(0)-\varepsilon)\frac{n-k}{k}}
$$

\n
$$
\geq n e^{n\Phi(0)} \sum_{k \leq \frac{\delta}{1+\delta}n} k e^{(\Phi'(0)-\varepsilon)k}
$$

\n
$$
\succ n e^{n\Phi(0)}.
$$

Since 0 is the unique maximum point of $\Phi(\lambda)$, there exists $\eta > 0$ such that $\Phi(\lambda) \leq \Phi(0) - \eta$ for all $\lambda \geq \delta$. Thus

.

$$
\sum_{k < (1+\delta)^{-1}n} k(n-k) e^{n\Phi(\frac{n-k}{k})} \le n^3 e^{n(\Phi(0)-\eta)}
$$

Also,

$$
\sum_{k\geq (1+\delta)^{-1}n} k(n-k)e^{n\Phi\left(\frac{n-k}{k}\right)} \leq n e^{n\Phi(0)} \sum_{k\geq (1+\delta)^{-1}n} (n-k)e^{n\Phi'(0)+\varepsilon\left(\frac{n-k}{k}\right)^{-1}\frac{n-k}{k}} \\
\leq n e^{n\Phi(0)} \sum_{k\leq \frac{\delta}{1+\delta}n} k e^{\Phi'(0)+\varepsilon\left(\frac{n}{1+\delta}\right)^{-1}k} \\
\leq n e^{n\Phi(0)} \sum_{k=0}^{\infty} k e^{\Phi'(0)+\varepsilon\left(\frac{n-k}{1+\delta}\right)^{-1}k}
$$

Combining the last two displays yields the desired upper-bound. (ii) For $0 < c_1 < -\frac{\Phi''(\lambda_0)}{2} < c_2$, there exists $\delta > 0$ such that

(5.13)
$$
-c_2(\lambda - \lambda_0)^2 \leq \Phi(\lambda) - \Phi(\lambda_0) \leq -c_1(\lambda - \lambda_0)^2
$$

for $|\lambda - \lambda_0| \leq \delta$. Clearly, we have that

$$
\sum_{k \colon \left| \frac{n-k}{k} - \lambda_0 \right| > \delta} k(n-k) e^{n\Phi\left(\frac{n-k}{k}\right)} \prec n^3 e^{n(\Phi(\lambda_0) - \eta)}
$$

for some $\eta > 0$. Now,

$$
\sum_{k \colon \left| \frac{n-k}{k} - \lambda_0 \right| \le \delta} k(n-k) e^{n\Phi\left(\frac{n-k}{k}\right)} \prec n^3 e^{n\Phi(\lambda_0)} \sum_{k \colon \left| \frac{n-k}{k} - \lambda_0 \right| \le \delta} \frac{1}{n} e^{-c_1 n \left(\frac{n}{k} - 1 - \lambda_0\right)^2}
$$

$$
\asymp n^3 e^{n\Phi(\lambda_0)} \int_{(1+\lambda_0+\delta)^{-1}}^{(1+\lambda_0-\delta)^{-1}} e^{-c_1 n \left(x^{-1} - 1 - \lambda_0\right)^2} dx.
$$

By a change of variables, we get

$$
\sum_{k:\ |\frac{n-k}{k}-\lambda_0|\leq \delta} k(n-k)e^{n\Phi(\frac{n-k}{k})} = n^3 e^{n\Phi(\lambda_0)} \int_{-\delta}^{\delta} e^{-c_1 ny^2} \frac{dy}{(y+1+\lambda_0)^2}
$$

$$
\lesssim n^{5/2} e^{n\Phi(\lambda_0)} \int_{-\delta\sqrt{n}}^{\delta\sqrt{n}} e^{-c_1 z^2} dz
$$

$$
\lesssim n^{5/2} e^{n\Phi(\lambda_0)}.
$$

The lower bound can be proved by the same arguments, changing c_1 with c_2 . (iii) The proof of (iii) is similar to that of (ii), except that we need to replace [\(5.13\)](#page-23-0) with

$$
-c_2\lambda^2 \le \Phi(\lambda) - \Phi(0) \le -c_1\lambda^2
$$

and the order of magnitude is not $\frac{n-k}{k} \approx \lambda_0 \pm \delta$ anymore, but $k \geq (1+\delta)^{-1} \frac{n-k}{k}$ as in (i).

6. Twisted Patterson-Sullivan measures

In this and next sections, we will give applications of our results to relatively hyperbolic groups. We start here by investigating the properties of Patterson-Sullivan like measures on the Bowditch boundary of such groups, which we define by using suited Poincaré series.

6.1. Some background on relatively hyperbolic groups. Since their introduction by Gromov, relatively hyperbolic groups were studied by many authors through several equivalent definitions. We will mainly use the viewpoint of Bowditch [\[Bow12\]](#page-49-21) and Gerasimov-Potyagailo [\[GP13\]](#page-50-20), [\[GP15\]](#page-50-21), [\[GP16\]](#page-50-22) in the sequel. Consider a finitely generated group Γ acting properly via isometries on a proper geodesic Gromov hyperbolic space X. Define the *limit set* Λ_{Γ} as the closure of Γ in the Gromov boundary ∂X of X, that is, fixing a base point x_0 in X, Λ_{Γ} is the set of all possible limits of sequences $g_n \cdot x_0$ in ∂X , $g_n \in \Gamma$. The proper action of Γ on X by isometries extends to a convergence group action on Λ_{Γ} by homeomorphisms, which means that the induced action on the space of distinct triples is properly continuous (see [\[Bow99,](#page-49-22) Proposition 1.11] for example). If Λ_{Γ} contains at least three points, then Γ acts minimally on $Λ$ _Γ.

A loxodromic element $x \in \Gamma$ is an infinite order element with exactly two fixed points $x_-\neq x_+$ in Λ_{Γ} . Moreover, x acts via North-South dynamics on Λ_{Γ} in the sense that for any $\xi \neq x_{\pm}$, $x^{\mp n}\xi$ converges to x_{\mp} as n goes to $+\infty$. Then x_{+} is called the attracting fixed point of x and $x_$ is called its repelling fixed point. The fixed points of any two loxodromic elements are either the same or disjoint. So Γ contains infinitely many loxodromic elements with pairwise disjoint fixed points.

A point $\xi \in \Lambda_{\Gamma}$ is called *conical* if there is a sequence g_n of Γ and distinct points ξ_1, ξ_2 in Λ_{Γ} such that $g_n\xi$ converges to ξ_1 and $g_n\zeta$ converges to ξ_2 for all $\zeta \neq \xi$ in Λ_{Γ} . A point $\xi \in \Lambda_{\Gamma}$ is called *parabolic* if its stabilizer in Γ is infinite, fixes exactly ξ in $Λ_Γ$ and contains no loxodromic element. If, in addition, its stabilizer in Γ acts co-compactly on $\Lambda_{\Gamma} \setminus {\{\xi\}}$, then ξ is called *bounded parabolic*. Say that the action of Γ on X is *geometrically finite* if the induced convergence group action on the limit set Λ_{Γ} is geometrically finite: Λ_{Γ} only consists of conical limit points and bounded parabolic limit points. See [\[Bow99\]](#page-49-22) for more general facts on convergence groups.

A group Γ is called *relatively hyperbolic* with respect to a collection of subgroups $\mathbb P$ if it acts geometrically finitely on a proper geodesic hyperbolic space X such that the stabilizers of parabolic limit points are exactly the conjugates of the elements of $\mathbb P$. Elements of $\mathbb P$ are called maximal parabolic subgroups. We will write $\mathbb P_0$ for the choice of a set of representatives of conjugacy classes of elements of \mathbb{P} . By [\[Bow12,](#page-49-21) Proposition 6.15, such a set \mathbb{P}_0 is finite.

The limit set Λ_{Γ} in the Gromov boundary of X is called the *Bowditch boundary* of Γ. By [\[Bow12,](#page-49-21) Theorem 9.4], it is unique up to equivariant homeomorphism and in particular does not depend on the choice of a proper geodesic Gromov hyperbolic space X on which Γ acts geometrically finitely. We will write $\partial \Gamma$ for the Bowditch boundary of Γ in the sequel. A relatively hyperbolic group is called non-elementary if its Bowditch boundary is infinite; equivalently, $\sharp \partial \Gamma > 2$.

We now fix a finite set \mathbb{P}_0 of representatives of conjugacy classes of maximal parabolic subgroups. When $P \in \mathbb{P}_0$ and $q \in \Gamma$, we call qP a maximal parabolic coset.

Definition 6.1. Let gP be a maximal parabolic coset and $\eta, L > 0$ be fixed constants. A point p on a geodesic α is called (η, L) -deep in qP if

$$
B(p, 2L) \cap \alpha \subseteq N_{\eta}(gP).
$$

It is called an (η, L) -transition point if it is not (η, L) -deep in any maximal parabolic coset qP .

The following set of inequalities is called weak relative Ancona inequalities and will be helpful below. They extend similar inequalities proved for hyperbolic groups by Ancona [\[Anc88\]](#page-49-23) for $r = 1$, by Gouëzel-Lalley [\[GL13\]](#page-50-23) on co-compact Fuchsian groups for $r \leq R$ and by Gouezel [\[Gou14\]](#page-50-24) in full generality for $r \leq R$. The version for relatively hyperbolic groups that we use here were first proved for $r = 1$ by Gekhtman-Gerasimov-Potyagailo-Yang [\[GGPY21\]](#page-50-25) and then by Dussaule-Gekhtman [\[DG21\]](#page-49-6) for $r \leq R$.

Lemma 6.2. [\[GGPY21,](#page-50-25) Theorem 1.1][\[DG21,](#page-49-6) Theorem 1.6] Let Γ be a relatively hyperbolic group and let μ be a finitely supported admissible and symmetric probability measure on Γ. Then, for every $c, \eta, L \geq 0$ there exists $C > 0$ such that for every $r \leq R$, the following holds. For every $x, y, z \in \Gamma$, if y has a distance at most c to an (η, L) -transition point on a geodesic from x to z, then

$$
\frac{1}{C}G(x,y|r)G(y,z|r) \leq G(x,z|r) \leq CG(x,y|r)G(y,z|r).
$$

Note that the constant C is independent of $r \in [1, R]$. In other words, the Green function is roughly multiplicative along transition points on a geodesic. In both [\[GGPY21\]](#page-50-25) and [\[DG21\]](#page-49-6), these inequalities are formulated in terms of the Floyd distance, which is a suited rescaling of the word distance. However, the statement for transition points directly follows from [\[GP15,](#page-50-21) Corollary 5.10] which relates transition points with the Floyd distance. We also refer to [\[GGPY21,](#page-50-25) Section 9] for more details.

We will also use the following at some point.

Lemma 6.3. [\[Yan22,](#page-51-7) Lemma 2.14] There exist universal constants η , L with the following property. Let γ be a geodesic ray ending at a conical point $\xi \in \partial \Gamma$. Then γ contains a unbounded sequence of (η, L) -transition points x_n .

In the remainder of this section, we consider a finitely generated relatively hyperbolic group Γ. When speaking of a transition point, we mean an (η, L) -transition point with (η, L) satisfying Lemma [6.3.](#page-25-0) We also fix a finitely supported symmetric and admissible probability measure μ on Γ .

6.2. Busemann cocyles. Given $x, y, z \in \Gamma$, let $B_z(x, y) := d(x, z) - d(y, z)$ and $K_z(x,y|r) = \frac{G(x,z|r)}{G(y,z|r)}$. The function B_z is called the Busemann function associated with the distance d at z .

Following [\[BB07\]](#page-49-24), we define the r-Green distance by

$$
d_r(x,y) = -\log F_r(x,y) = -\log \frac{G(x,y|r)}{G(e,e|r)}.
$$

Then $K_z(x, y|r) = e^{-[d_r(x,z)-d_r(y,z)]}$ is the exponential of the Busemann function for the r-Green distance. We also write $|x^{-1}y|_r = d_r(x, y)$, and $|x^{-1}y| = d(x, y)$.

Consider the distance for $x, y \in \Gamma$:

$$
\mathfrak{d}_r(x,y) := \omega_{\Gamma}(r)|x^{-1}y| + |x^{-1}y|_r.
$$

Lemma 6.4. If $1 \leq r < R$, then the distance \mathfrak{d}_r is proper and quasi-isometric to the word distance.

Proof. The proof is standard, but we write a complete argument for sake of completeness. First we prove that for every $r < R$, there exist $C_1 > 0$ and $\alpha_1 > 0$ such that for every $x \in \Gamma$

(6.1)
$$
G(e, x|r) \leq C_1 e^{-\alpha|x|}.
$$

Since μ is finitely supported, there exists $c > 0$ such that

$$
G(e, x|r) = \sum_{n \ge c|x|} r^n p_n(e, x).
$$

Moreover, by (2.2) ,

$$
p_n(e,x) \le R^{-n}
$$

.

Therefore,

$$
G(e, x|r) \leq \sum_{n \geq c|x|} \left(\frac{r}{R}\right)^n \leq C_1 \left(\frac{r}{R}\right)^{c|x|}.
$$

This proves [\(6.1\)](#page-26-0).

Second, we prove that for every $r \geq 1$, there exists $C_2 > 0$ and $\alpha_2 > 0$ such that for every $x \in \Gamma$,

(6.2)
$$
G(e, x|r) \geq C_2 e^{-\alpha_2|x|}.
$$

Indeed, since the support of μ generates Γ, there exists a path $x_0 = e, x_1, ..., x_n = x$ such that $n \asymp |x|$ and $\mu(x_k^{-1}x_{k+1}) > 0$. In particular, we find that

 $G(e, x|r) \ge G(e, x|1) \ge \mu(x_0^{-1}x_1) \cdots \mu(x_{n-1}^{-1}x_n),$

which proves [\(6.2\)](#page-26-1). We conclude that for $1 \leq r < R$, the Green distance is quasiisometric to the word distance and that $G(e, x|r)$ vanishes at infinity. Consequently, the distance \mathfrak{d}_r also is quasi-isometric to the word distance and satisfies that as x goes to infinity, $\mathfrak{d}_r(e, x)$ tends to infinity. In particular, any ball for \mathfrak{d}_r is contained in a larger ball for the word distance and thus is finite, so \mathfrak{d}_r is proper.

Remark 6.5. According to [\[GL13,](#page-50-23) Lemma 2.1], for any non-amenable group Γ and any finitely supported symmetric admissible probability measure μ , $G(e, x|R)$ converges to 0 as |x| goes to infinity. As a consequence, the distance \mathfrak{d}_R is proper, although it might not be quasi-isometric to the word distance.

Define the corresponding Busemann cocycle

(6.3)
$$
\mathfrak{B}_{\xi}(x,y;r) = \omega_{\Gamma}(r)B_{\xi}(x,y) - \log K_{\xi}(x,y|r)
$$

Lemma 6.6. There exists a constant $C > 0$ with the following property. Let $\xi \in \partial \Gamma$ be a conical point, and $x, y \in \Gamma$. There exists a neighborhood $V = V(x, y)$ of ξ in $Γ ∪ ∂Γ$ such that for any $z ∈ V ∩ Γ$:

$$
|B_{\xi}(x, y) - B_{z}(x, y)| \le C,
$$

$$
|\log K_{\xi}(x, y|r) - \log K_{z}(x, y|r)| \le C.
$$

Proof. The statement for B_{ξ} is proved in [\[Yan22,](#page-51-7) Lemma 2.20]. Also, by [\[DG21,](#page-49-6) Proposition 4.1, the Martin kernel $K_z(\cdot, e|r) = G(\cdot, z|r)/G(e, z|r)$ extends continuously to $K_{\xi}(\cdot, e|r)$ as z converges to a conical limit point ξ . This follows from weak relative Ancona inequalities. In particular, $K_z(x, y|r)$ converges to $K_{\xi}(x, y|r)$ as z converges to ξ , so the statement for K_{ξ} also holds. converges to ξ , so the statement for K_{ξ} also holds.

As a consequence, the Busemann cocyle $\mathfrak{B}_{z}(x, y)$ extends to a coarse cocycle $\mathfrak{B}_{\xi}(x, y)$ at a conical point ξ. That is,

$$
\mathfrak{B}_\xi(x,y):=\limsup_{z\to\xi}\mathfrak{B}_\xi(x,y)
$$

does not depend on the choice of $z \to \xi$, up to a bounded additive error C independent of ξ .

6.3. Quasi-conformal densities. A Borel measure μ on a topological Hausdorff space T is regular if $\mu(A) = \inf \{ \mu(U) : A \subset U, U \text{ is open} \}$ for any Borel set A in T. It is called tight if $\mu(A) = \sup \{ \mu(K) : K \subset A, K \text{ is compact} \}$ for any Borel set A in T . A finite Borel measure is called $Radon$ if it is tight and regular. It is well-known that all finite Borel measures on compact metric spaces are Radon, see [\[Bil99,](#page-49-25) Theorem 1.1, Theorem 1.3].

Denote by $\mathcal{M}(\partial \Gamma)$ the set of finite positive Radon measures on $\partial \Gamma$. Then Γ acts on $\mathcal{M}(\partial \Gamma)$ via $g_* \nu(A) = \nu(g^{-1}A)$ for any Borel set A in $\partial \Gamma$.

Definition 6.7. We call a map $x \mapsto \nu_x$ equivariant if for every $x, g \in \Gamma$, we have

$$
\nu_{gx}(A) = \nu_x(g^{-1}A)
$$

for every Borel set $A \subset \partial \Gamma$.

Definition 6.8. Let $\omega \in [0, \infty]$. We call a Γ-equivariant map

$$
\nu : \Gamma \to \mathcal{M}(\partial \Gamma), \ x \mapsto \nu_x
$$

an ω -dimensional quasi-conformal density if for any $x, y \in \Gamma$ the following holds

(6.4)
$$
\frac{d\nu_x}{d\nu_y}(\xi) \approx e^{-\omega B_{\xi}(x,y)} K_{\xi}(x,y|r),
$$

for ν_y -a.e. points $\xi \in \partial \Gamma$, where the implicit constant does neither depend on x, y, nor on ξ .

By the equivariant property of μ , we see the following result.

Lemma 6.9. Let $\{\nu_x\}_{x \in \Gamma}$ be a σ-dimensional quasi-conformal density on $\partial \Gamma$. Then the support of any ν_x is $\partial \Gamma$.

Proof. By definition, the support $\text{supp}(\mu_x)$ is a maximal closed subset such that any point in $\partial\Gamma\setminus \text{supp}(\mu_x)$ has an open neighborhood which is ν_x -null. It is well-known that $\partial \Gamma$ is a minimal Γ-invariant closed set, see for instance [\[Bow99,](#page-49-22) Section 2]. Thus, it suffices to prove that the support of ν_x is G-invariant. This follows from equivariance and quasi-conformality, since $\nu_x(A) = \nu_{gx}(gA)$ for any Borel subset $A \subset \partial \Gamma$ and ν_x and ν_{gx} are absolutely continuous with respect to each other.

As explained in the introduction, we associate the following Poincaré series to μ and to the word distance, by setting

$$
\Theta_{\Gamma}(r,s) := \sum_{x \in \Gamma} G(e,x|r) e^{-s|x|} = \sum_{n \ge 0} H_r(n) e^{-sn} = G(e,e|r) \cdot \sum_{x \in \Gamma} e^{-s|x| - |x|_r}
$$

where we recall that $H_r(n) = \sum_{x \in S_n} G(e, x|r)$ and that the *critical exponent* $\omega_{\Gamma}(r)$ is defined by

$$
\omega_{\Gamma}(r) = \limsup_{n \to \infty} \frac{1}{n} \log H_r(n).
$$

The group Γ is of *divergent* (resp. *convergent*) type for the Green function if $\Theta_{\Gamma}(r, s)$ is divergent (resp. convergent) at $s = \omega_{\Gamma}(r)$.

Recall $\mathfrak{d}_r(x,y) := \omega_{\Gamma}(r)|x^{-1}y| + |x^{-1}y|_r.$

Lemma 6.10. The series defined as follows

$$
\forall s > 0, \ \mathcal{P}_{\Gamma}(s) := \sum_{x \in \Gamma} e^{-s \mathfrak{d}_r(x, y)}
$$

has critical exponent 1, and the divergence of $\mathcal{P}_{\Gamma}(s)$ at $s = 1$ is equivalent to that of $\Theta_{\Gamma}(r,s)$ at $s=\omega_{\Gamma}(r)$.

Proof. By [\[GL13,](#page-50-23) Lemma 2.1], $G(e, x|R)$ goes to 0 as |x| tends to infinity. Thus, $G(e, x|r) \leq G(e, x|R) \leq 1$ for large enough x. In particular, we see that for $s > 1$,

$$
\sum_{x \in \Gamma} G(e, x|r)^s e^{-s\omega_{\Gamma}(r)|x|} \le C \sum_{x \in \Gamma} G(e, x|r) e^{-s\omega_{\Gamma}(r)|x|}.
$$

The right-hand side converges, so the left-hand side also converges. Conversely, for $s < 1$,

$$
\sum_{x \in \Gamma} G(e, x|r) e^{-s\omega_{\Gamma}(r)|x|} \le C \sum_{x \in \Gamma} G(e, x|r)^s e^{-s\omega_{\Gamma}(r)|x|}.
$$

The left-hand side diverges, so the right-hand side also diverges. Thus, the critical exponent of $\mathcal{P}_{\Gamma}(s)$ is 1.

Also, note that $\mathcal{P}_{\Gamma}(1) = \Theta_{\Gamma}(r, \omega_{\Gamma}(r))$, so the second conclusion of the lemma follows. follows. \Box

Write $\mu(f) = \int f d\mu$ for a continuous function $f \in C(\partial \Gamma)$. We endow $\mathcal{M}(\partial \Gamma)$ with the weak topology. That is, a sequence $\mu_n \in \mathcal{M}(\partial \Gamma)$ converges to μ if $\mu_n(f)$ converges to $\mu(f)$ for any $f \in C(\partial \Gamma)$. Equivalently, by the Portmanteau Theorem [\[Bil99,](#page-49-25) Theorem 2.1], μ_n converges to μ if $\liminf_{n\to\infty} \mu_n(U) \geq \mu(U)$ for any open set U ⊂ ∂ Γ.

We start by constructing a family of measures $\{\nu_x^s\}_{x \in \Gamma}$ supported on Γ for any s > 1. First, assume that $\mathcal{P}_{\Gamma}(s)$ is divergent at $s = 1$. Set

$$
\nu_x^s = \frac{1}{\mathcal{P}_\Gamma(s)} \sum_{z \in \Gamma} e^{-s \mathfrak{d}_r(x,z)} \cdot \text{Dirac}(z),
$$

where $s > 1$ and $x \in \Gamma$. Note that ν_x^s is a probability measure.

On the contrary, assume that $\mathcal{P}_{\Gamma}(s)$ is convergent at $s = 1$, Patterson introduced in [\[Pat76,](#page-50-26) Lemma 3.1] a monotonically increasing function $H : \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ with the following property:

(6.5)
$$
\forall \epsilon > 0, \ \exists t_{\epsilon}, \ \forall t > t_{\epsilon}, \ \forall a > 0: \quad H(a+t) \leq \exp(a\epsilon)H(t).
$$

and such that the following modified series

$$
\mathcal{P}'_{\Gamma}(s) := \sum_{x \in \Gamma} H(\mathfrak{d}_r(x, z)) \cdot e^{-s \mathfrak{d}_r(x, z)}
$$

is divergent for $s \leq 1$ and convergent for $s > 1$. Then define measures as follows:

$$
\nu_x^s = \frac{1}{\mathcal{P}'_{\Gamma}(s)} \sum_{z \in \Gamma} e^{-s \mathfrak{d}_r(x,z)} \cdot H(\mathfrak{d}_r(x,z)) \cdot \text{Dirac}(z),
$$

where $s > 1$ and $x \in \Gamma$.

Choose $s_i \searrow 1$ such that $\nu_x^{s_i}$ are convergent in $\mathcal{M}(\partial \Gamma \cup \Gamma)$ for all $x \in \Gamma$. Let $\nu_x = \lim_i \nu_x^{s_i}$ be the limit measures, which are called *Patterson-Sullivan measures* associated with the Poincaré series \mathcal{P}_{Γ} . Note that forcing the Poincaré series to be divergent at 1, we have $\nu_x(\partial \Gamma) = 1$. In the sequel, we write PS-measures as shorthand for Patterson-Sullivan measures. We also write $\partial \Gamma^{\text{con}}$ for the set of conical limit points in the Bowditch boundary.

Proposition 6.11. The PS-measures $\{\nu_x\}_{x \in \Gamma}$ on the Bowditch boundary are absolutely continuous with respect to each other and satisfy

(6.6)
$$
\forall \xi \in \partial \Gamma: \quad \frac{d\nu_x}{d\nu_y}(\xi) \geq e^{-\mathfrak{d}_r(x,y)},
$$

(6.7)
$$
\forall \nu_y \ a.e. \ \xi \in \partial \Gamma^{\text{con}}: \quad \frac{d\nu_x}{d\nu_y}(\xi) \asymp e^{-\omega_{\Gamma}(r)B_{\xi}(x,y)}K_{\xi}(x,y|r),
$$

where the implicit constant is independent of x, y and ξ .

Remark 6.12. If Γ is of divergent type for Green function, then Theorem [6.17](#page-32-0) below says that PS-measures have no atoms on Bowditch boundary and give full measure to conical limit points, so [\(6.7\)](#page-29-0) holds for ν_y -a.e. $\xi \in \partial \Gamma$. In this case, ν is an $\omega_{\Gamma}(r)$ -dimensional quasi-conformal density.

Proof. Since \mathfrak{d}_r satisfies the triangle inequality and $\lim_{t\to\infty} \frac{H(a+t)}{H(t)} = 1$, we see that $\{\nu_x : x \in \Gamma\}$ are absolutely continuous with respect to each other,

$$
e^{-\mathfrak{d}_r(x,y)} \leq \frac{d\nu_x}{d\nu_y}(\xi) \leq e^{\mathfrak{d}_r(x,y)}
$$
.

We now verify the quasi-conformality at conical limit points. We only consider the case where Γ is of convergent type, the divergent type being simpler. Let $\epsilon > 0$ and t_{ϵ} the number be given by [\(6.5\)](#page-28-0) for the function H. Let $\phi = \frac{dv_x}{dv_y}$ be the Radon-Nikodym derivative, uniquely defined up to a ν_y -null set. Let $\xi \in \partial \Gamma$ be a conical limit point and consider the open neighborhood V of ξ and the uniform constant C given by Lemma [6.6.](#page-26-2)

Let f be a continuous function supported in V . One can choose V also such that $\mathfrak{d}_r(y, z) > t_\epsilon$ for all $z \in V$. If $z \in V$ satisfies $\mathfrak{d}_r(x, z) > \mathfrak{d}_r(y, z)$, then we have

$$
H(\mathfrak{d}_r(x, z)) = H(\mathfrak{d}_r(x, z) - \mathfrak{d}_r(y, z) + \mathfrak{d}_r(y, z))
$$

\n
$$
\leq e^{\epsilon[\mathfrak{d}_r(x, z) - \mathfrak{d}_r(y, z)]} \cdot H(\mathfrak{d}_r(y, z))
$$

\n
$$
\leq e^{\epsilon(C + \mathfrak{B}_{\xi}(x, y))} \cdot H(\mathfrak{d}_r(y, z)).
$$

Since H is increasing, we have

(6.8)
$$
C_{\epsilon}^{-1}H(\mathfrak{d}_r(y,z)) \leq H(\mathfrak{d}_r(x,z)) \leq C_{\epsilon}H(\mathfrak{d}_r(y,z)),
$$

where $C_{\epsilon} = e^{\epsilon(C + \mathfrak{B}_{\xi}(x,y))} > 1$ depends on ϵ, C and (x, y) , but not on $z \in V$. Note that $C_{\epsilon} \to 1$ as $\epsilon \to 0$. By symmetry, the conclusion [\(6.8\)](#page-29-1) also holds if $\mathfrak{d}_r(x, z) < \mathfrak{d}_r(y, z)$ for $z \in V$.

Using [\(6.8\)](#page-29-1), we get the following estimates. First,

$$
\nu_x^s(f) = \frac{1}{\mathcal{P}_{\Gamma}(s)} \sum_{z \in V} e^{-s \mathfrak{d}_r(x,z)} H(\mathfrak{d}_r(x,z)) f(z)
$$

$$
\leq C_{\epsilon} e^{-s \mathfrak{B}_z(x,y))} \cdot \frac{1}{\mathcal{P}_{\Gamma}(s)} \sum_{z \in V} e^{-s \mathfrak{d}_r(y,z)} H(\mathfrak{d}_r(y,z)) f(z)
$$

$$
\asymp C_{\epsilon} e^{-s \mathfrak{B}_{\xi}(x,y)} \nu_y^s(f)
$$

and second

$$
\nu_x^s(f) = \frac{1}{\mathcal{P}_\Gamma(s)} \sum_{z \in V} e^{-s \mathfrak{d}_r(x,z)} H(\mathfrak{d}_r(x,z)) f(z)
$$

\n
$$
\geq C_{\epsilon}^{-1} e^{-s \mathfrak{B}_z(x,y))} \cdot \frac{1}{\mathcal{P}_\Gamma(s)} \sum_{z \in V} e^{-s \mathfrak{d}_r(y,z)} H(\mathfrak{d}_r(y,z)) f(z)
$$

\n
$$
\asymp C_{\epsilon} e^{-s \mathfrak{B}_{\xi}(x,y)} \nu_y^s(f)
$$

where the implicit constants depend only on C but not on ϵ . Letting $s \to 1$, we get

$$
C_{\epsilon}^{-1} e^{-\mathfrak{B}_{\xi}(x,y)} \nu_{y}(f) \prec \nu_{x}(f) \prec C_{\epsilon} e^{-\mathfrak{B}_{\xi}(x,y)} \nu_{y}(f)
$$

for any continuous function f supported in V. As $\epsilon \to 0$, $C_{\epsilon} \to 1$, hence it follows that

$$
\phi(\xi) \asymp e^{-\mathfrak{B}_{\xi}(x,y)}
$$

for ν_v -a.e. conical limit point $\xi \in \partial \Gamma$.

6.4. Shadow Lemma. The key observation in the theory of PS-measures is the Sullivan Shadow Lemma shedding some light on the relation between the PSmeasure and the geometric properties of the boundary.

Recall that $\partial \Gamma$ is a visual boundary for the Cayley graph Cay(G, S): any two distinct points $x, y \in \text{Cay}(G, S) \cup \partial \Gamma$ can be connected by a geodesic. See [\[GP13,](#page-50-20) Proposition 2.4].

Definition 6.13. Let $C > 0$ and $x \in \Gamma$. The shadow $\Pi_C(x)$ at x is the set of points $\xi \in \partial \Gamma$ such that there exists some geodesic $\gamma = [e, \xi]$ intersecting $B(x, C)$.

The partial shadow $\Psi_C(x)$ at x is the set of points $\xi \in \partial \Gamma$ such that some geodesic $[e, \xi]$ contains a transition point C-close to x.

We now state the Shadow Lemma in our context, whose proof follows closely the proofs of [\[Yan22,](#page-51-7) Lemmas 4.1 $\&$ 4.2] with Lemma [6.6](#page-26-2) replacing Lemma 2.19 there. Let us denote by $\Psi_C^{\text{con}}(g)$ the set of all conical limit points in $\Psi_C(g)$.

Lemma 6.14 (Shadow Lemma). Let $\{\nu_x\}_{x \in \Gamma}$ be an $\omega_{\Gamma}(r)$ -dimensional PS measures on the Bowditch boundary $\partial \Gamma$. Then there exists $C_0 > 0$ such that for any $C \geq C_0$ and $x \in \Gamma$ the following two inequalities hold

(6.9)
$$
e^{-\omega_{\Gamma}(r)|x|}G(e,x|r)\prec_C\nu_e(\Psi_C(x))\leq\nu_e(\Pi_C(x)),
$$

(6.10)
$$
\nu_e(\Psi_C^{\text{con}}(x)) \prec_C e^{-\omega_{\Gamma}(r)|x|} G(e,x|r).
$$

Remark 6.15. If ν_e has no atoms at parabolic points which form a countable subset of the Bowditch boundary, then we obtain the full strength of the partial shadow lemma without having to restrict our attention to conical points. The upper bound [\(6.10\)](#page-30-0) for the partial shadow uses the relative Ancona inequalities

(Lemma [6.2\)](#page-25-1), while it is unknown whether the upper bound holds for the usual shadow.

Proof. Let F be a set of three loxodromic elements with pairwise disjoint fixed points. For each $f \in F$, let $\alpha := \bigcup_{i \in \mathbb{Z}} f^i[e, f]$ be an $\langle f \rangle$ -invariant quasi-geodesic between two fixed points $f_-, f_+ \in \partial \Gamma$. Let $U_f \subset \partial \Gamma$ be an open neighborhood of f_+ so that for any $\eta \in U_f$, the projection of η to the axis α has a distance to e at least C. By [\[DWY22,](#page-50-0) Lemma 2.4], for any $x \in \Gamma$, there exists $f \in F$ so that $[x^{-1}, \eta]$ contains a transition point *C*-close to *e*. Thus, $U_f \subset x^{-1} \Psi_C(x)$.

As Γ acts minimally with a dense orbit in $\partial \Gamma$, the Γ-orbit of any open set $U \subset \partial \Gamma$ covers $\partial \Gamma$, so U have positive ν_e -measure. Hence, setting

$$
D = \min\{\nu_e(U_f) : f \in F\} > 0
$$

which is independent of x , we have

$$
\nu_e(x^{-1}\Psi_C(x)) \ge D.
$$

Since ν_x is equivariant, the lower bound in [\(6.6\)](#page-29-2) implies

$$
\nu_e(\Psi_C(x)) = \nu_{x^{-1}}(x^{-1}\Psi_C(x)) \ge e^{-\omega_\Gamma(r)|x|} \frac{G(e, x|r)}{G(e, e|r)} \cdot \nu_e(x^{-1}\Psi_C(x))
$$

$$
\ge De^{-\omega_\Gamma(r)|x|} \frac{G(e, x|r)}{G(e, e|r)}.
$$

Since $G(e, e|r)$ is bounded by $G(e, e|R)$, this concludes the proof of the lower bound.

For any $\xi \in x^{-1} \Psi_C^{\text{con}}(x)$, there is a geodesic γ from x^{-1} to ξ which intersects $B(e, C)$ and contains a transition point. Thus, $|B_{\xi}(x^{-1}, e) - d(x^{-1}, e)| \leq 2C$. By the relative Ancona inequalities (Lemma [6.2\)](#page-25-1), there is a constant C_1 independent of r such that

$$
K_{\xi}(x^{-1}, e|r) = \lim_{z \to \xi} \frac{G(x^{-1}, z|r)}{G(e, z|r)} \le C_1 G(e, x|r).
$$

Also, by [\(6.7\)](#page-29-0) there is a constant $C_2 > 0$ such that for ν_e -a.e. conical limit point $\xi \in \partial \Gamma^{\rm con},$

$$
\frac{d\nu_{x^{-1}}}{d\nu_e}(\xi) \le C_2 e^{-\omega_{\Gamma}(r)B_{\xi}(x^{-1},e)} K_{\xi}(x^{-1},e|r).
$$

Combining together the above estimates, we have

$$
\nu_e(\Psi_C^{\text{con}}(x)) = \nu_{x^{-1}}(x^{-1}\Psi_C^{\text{con}}(x))
$$

\n
$$
\leq C_2 \int_{x^{-1}\Psi_C^{\text{con}}(x)} e^{-\omega_{\Gamma}(r)B_{\xi}(x^{-1}, e)} K_{\xi}(x^{-1}, e|r) d\nu_e(\xi)
$$

\n
$$
\leq C_1 C_2 e^{2C\omega_{\Gamma}(r)} \cdot e^{-\omega_{\Gamma}(r)|x|} G(e, x|r),
$$

which finishes the proof of the upper bound. \square

Proposition 6.16. Suppose that ν_e gives positive measure to the set of conical limit points. Then Γ is of divergent type for the Green function.

Proof. List $\Gamma = \{x_1, \ldots, x_i, \ldots\}$ such that for all $i, |x_i| \leq |x_{i+1}|$. Let C_0 be given by Lemma [6.14.](#page-30-1) For any $C > C_0$, set

(6.11)
$$
A_C := \bigcap_{n=1}^{\infty} \bigcup_{i=n}^{\infty} \Psi_C^{\text{con}}(x_i).
$$

$$
\mathbb{L}^2
$$

By Lemma [6.3,](#page-25-0) we have $\partial^{\text{con}} \Gamma = A_C$. In other words, any conical limit point is shadowed infinitely many times by elements of Γ.

We claim that $\Theta_{\Gamma}(r, \omega_{\Gamma}(r))$ is divergent. Recall that

$$
\Theta_{\Gamma}(r,\omega_{\Gamma}(r)) = \sum_{x \in \Gamma} e^{-\omega_{\Gamma}(r)|x|} G(e,x|r).
$$

By Lemma [6.14,](#page-30-1) we see that

$$
\sum_{|x| \ge n} e^{-\omega_{\Gamma}(r)|x|} G(e, x|r) \succ \sum_{|x| \ge n} \nu_e(\Psi_C^{\text{con}}(x))
$$

$$
\succ \nu_e\left(\bigcup_{|x| \ge n} \Psi_C^{\text{con}}(x)\right) \succ \nu_e(A_C) > 0.
$$

This is true for all $n > 0$ and $\nu_e(A_C)$ is independent of n. Thus, $\Theta_{\Gamma}(r,s)$ is indeed divergent at $s = \omega_{\Gamma}(r)$.

Theorem 6.17. If Γ is of divergent type for Green function, then ν_e has no atoms. Otherwise, ν_e is purely atomic and supported on the set of parabolic points.

Proof. First of all, the otherwise statement follows from Proposition [6.16.](#page-31-0) Indeed, if Γ is of convergent type, then ν_e cannot give positive measure to conical limit points, hence it is supported on the set of parabolic limit points. Since this set is countable, ν_e is necessarily purely atomic.

Assume now that Γ is of divergent type. Let $q \in \partial \Gamma$ be a bounded parabolic point so that the stabilizer $P \in \mathbb{P}$ acts co-compactly on $\partial \Gamma \cup \Gamma \setminus \{q\}$. Let $K \subset \Gamma \cup \partial \Gamma \setminus \{q\}$ be a compact fundamental domain. For a point $y \in \Gamma$, we let $\pi_P(y)$ be the set of nearest point-projections

$$
\pi_P(y) = \{ p \in P : d(y, p) = d(y, P) \}.
$$

Define $\pi_P(A) := \bigcup_{a \in A} \pi_P(a)$. As ∂K is disjoint with $\partial P = \{q\}$, the shortest projection $Z := \pi_P (K \cap \Gamma)$ has bounded diameter by [\[GP16,](#page-50-22) Proposition 3.3]. By enlarging Z, assume without loss of generality that $1 \in Z$.

Note that a maximal parabolic P has the contracting property by [\[GP16,](#page-50-22) Proposition 8.5]: any geodesic [x, y] with large projection to P is uniformly close to $\pi_P(x)$ and $\pi_P(y)$. Thus, for any $y \in \Gamma \cap K$ and $p \in P$, any geodesic $[e, py]$ passes within uniformly bounded distance of $p \in pZ$. Consequently, $|py| \simeq |p| + |y|$. Moreover, $[e, py]$ exits P with bounded distance to pZ , so p is within a bounded distance of a transition point on $[e, py]$. Now, by the relative Ancona inequalities in Lemma [6.2,](#page-25-1) $G(e, py|r) \simeq G(e, p|r)G(e, y|r)$. Here the implicit constants in \simeq and \simeq are independent of y and p .

We can estimate the $\nu_e^s\text{-measure}$ of an open neighborhood

$$
U_n = \{q\} \cup \{pK : p \in P; |p| \ge n\}
$$

as follows

$$
\nu_e^s(U_n) \le \sum_{|p|>n} \nu_e^s(pK) \le \frac{1}{\mathcal{P}_\Gamma(s)} \sum_{|p|>n} \sum_{y \in K} e^{-s\omega_\Gamma(r)|p y|} [G(e, py|r)]^s,
$$

hence

$$
\nu_e^s(U_n) \prec \frac{1}{\mathcal{P}_\Gamma(s)} \sum_{|p|>n} \sum_{y \in K} e^{-s\omega_\Gamma(r)|p|} e^{-s\omega_\Gamma(r)|y|} [G(e, y|r)]^s [G(e, p|r)]^s
$$

$$
\prec \nu_e^s(K) \sum_{|p|>n} e^{-s\omega_\Gamma(r)|p|} [G(e, p|r)]^s.
$$

By [\[DWY22,](#page-50-0) Corollary 3.9], $\Theta_P(r, \omega_{\Gamma}(r))$ is convergent. Also, by the Portman-teau Theorem [\[Bil99,](#page-49-25) Theorem 2.1], $\limsup_{s\to 1} \nu_e^s(K) \leq \nu_e(K)$. Letting $s \to 1$ and then $n \to \infty$, we see that $\nu_e(U_n) \to 0$. Thus, ν_e has no atoms at parabolic limit points. By Lemma [6.3,](#page-25-0) a conical limit point ξ is contained in infinitely many partial shadows $\Psi_C(x_n)$. By Lemma [6.14,](#page-30-1) as $x_n \to \xi$, the ν_e -measure of $\Psi_C(x_n)$ tends to 0. so conical limit points are not atoms as well. 0, so conical limit points are not atoms as well.

7. CONVERGENT POINCARÉ SERIES AND APPLICATIONS

This final section is devoted to answering some questions raised in [\[DWY22\]](#page-50-0) where we initiated the study of branching random walks on relatively hyperbolic groups. In particular, we end here the proof of our main result, Theorem [1.1.](#page-2-0)

One important notion that was coined in [\[DWY22\]](#page-50-0) is the parabolic gap for the Green function whose definition we now recall. Let μ be a probability measure on Γ, let \mathbb{P}_0 be a finite set of representatives of conjugacy classes of maximal parabolic subgroups and let $P \in \mathbb{P}_0$. We set

$$
H_{P,r}(n) = \sum_{x \in S_n \cap P} G(e, x|r)
$$

and

$$
\omega_P(r) = \limsup_{n \to \infty} \frac{1}{n} \log H_{P,r}(n).
$$

Definition 7.1. If $\omega_P(r) < \omega_\Gamma(r)$, we say that Γ has a *parabolic gap* along P for the Green function at r. If for every P, for every $r \in (1, R]$, $\omega_P(r) < \omega_\Gamma(r)$, then we say that Γ has a *parabolic gap* for the Green function.

One of the consequences of having a parabolic gap for the Green function is that $H_r(n)$ is roughly multiplicative and the Green function has purely exponential growth. Namely, by [\[DWY22,](#page-50-0) Theorem 1.8], if Γ is a non-elementary relatively hyperbolic group and if μ is a finitely supported admissible and symmetric probability measure on Γ , then for every $1 < r \leq R$, there exist C and C' such that for all n,

(7.1)
$$
\frac{1}{C}H_r(n+m) \leq H_r(n)H_r(m) \leq CH_r(n+m)
$$

and

(7.2)
$$
\frac{1}{C'}e^{n\omega_{\Gamma}(r)} \leq H_r(n) \leq C'e^{n\omega_{\Gamma}(r)}.
$$

As proved in [\[SWX22\]](#page-51-2), these two properties [\(7.1\)](#page-33-1) and [\(7.2\)](#page-33-2) hold for all hyperbolic groups. It was proved in [\[DWY22\]](#page-50-0) that if maximal parabolic subgroups are amenable and if $r < R$, then the parabolic gap condition holds, hence so do the properties [\(7.1\)](#page-33-1) and [\(7.2\)](#page-33-2). Under additional assumptions on the random walk, this was also proved at R. Among the unanswered problems in $DWY22$ are the following questions. Does the parabolic gap condition holds at R as soon as maximal

parabolic subgroups are amenable ? Does there exist examples of relatively hyperbolic groups endowed with a finitely supported admissible symmetric probability measures such that the properties [\(7.1\)](#page-33-1) and [\(7.2\)](#page-33-2) fail ? As particular cases of our work, we answer these two questions here.

7.1. First return to maximal parabolic subgroups. We gather the results of Section [2](#page-6-0) to prove that whenever maximal parabolic subgroups are amenable, the parabolic gap for the Green functions holds.

Consider the first return kernel $p_{r,P}$ to P for $r\mu$ defined by

$$
p_{r,P}(x,y) = \sum_{n \geq 1} \sum_{z_1, ..., z_{n-1} \notin P} r^n p(x, z_1) p(z_1, z_2) \cdots p(z_{n-1}, y).
$$

Denote by $p_{r,P}^{(n)}$ the *n*th convolution power of this transition kernel and by $G_{r,P}$ the associated Green function. As explained in the introduction, for every $x, y \in P$, $G_{r,P}(x,y|1) = G(x,y|r)$ by [\[DG21,](#page-49-6) Lemma 4.4]. Set $t_{r,P} = \sum_{x \in P} p_{r,P}(e,x)$. Then $t_{r,P}^{-1} p_{r,P}$ is a symmetric admissible and P-invariant transition kernel, thus defines a random walk on P.

Proposition 7.2. Let Γ be a relatively hyperbolic group and let $P \in \mathbb{P}_0$. Consider an admissible and symmetric probability measure μ on Γ . If $t_{r,P} \leq 1$, then we have $\omega_P(r) \leq 0$; in particular, $\omega_P(r) < \omega_\Gamma(r)$.

Proof. Denote $t = t_{r,P}$ for simplicity and let G_t be the Green function associated with $t^{-1}p_{r,P}$. By Proposition [2.3,](#page-7-0) $\sum_{x \in P, |x|=n} G_t(e,x|1)$ has growth rate at most 0. Since $t \leq 1$ we have that

$$
G(e, x|r) = G_{r,P}(e, x|1) = G_t(e, x|t) \le G_t(e, x|1).
$$

Therefore $\omega_P(r) \le 0$.

Proposition 7.3. Let Γ be a relatively hyperbolic group and let P be a maximal parabolic subgroup. Consider an admissible and symmetric probability measure μ on Γ. If P is amenable, then for every $r \leq R$, $\omega_P(r) \leq 0$.

Proof. Since $G(e, x|r) < \infty$, we deduce that the spectral radius of $p_{r,P}$ is at most 1. By [\[Woe00,](#page-51-0) Corollary 12.5] and the fact that P is amenable, the spectral radius of $t_{r,P}^{-1} p_{r,P}$ is 1 and hence $t_{r,P} \leq 1$. The result follows from Proposition [7.2.](#page-34-1) □

Note that we do not need to assume that μ is finitely supported in this proposition, although we need this assumption in the following corollary, which also relies on [\[DWY22,](#page-50-0) Theorem 1.8] mentionned above, where the assumption is crucially used.

Corollary 7.4. Let Γ be a relatively hyperbolic group endowed with a finitely supported admissible and symmetric probability measure μ . Assume that maximal parabolic subgroups of Γ are amenable. Then Γ has a parabolic gap for the Green function and so (7.1) and (7.2) hold.

Note that $t_{1,P}$ is the probability that the random walk associated to μ eventually returns to P. We see that $t_{1,P} < 1$ for all $P \in \mathbb{P}_0$. Otherwise, the random walk would visit P infinitely many times with positive probability, which in turn would imply that it accumulates at the parabolic limit points fixed by P . This would contradict the fact that the random walk almost surely converges to a conical limit point [\[GGPY21,](#page-50-25) Theorem 9.14].

More generally, consider a branching random walk on Γ whose step distribution is given by μ and with mean offspring r. Following [\[CGM12\]](#page-49-8), by collecting all particles returning to P, one gets a Galton-Watson process with mean offspring $t_{r,P}$, see precisely the proof of [\[CGM12,](#page-49-8) Proposition 4.3]. Consequently, the branching random walk returns to P infinitely many times if and only if $t_{r,P} > 1$. In such case, the branching random walk must accumulate in P . Moreover, the same arguments as in [\[CGM12,](#page-49-8) Proposition 4.4] show that there are almost surely infinitely many cosets gP , such that the branching random walk accumulates in gP . On the contrary, if $t_{r,P} \leq 1$, then for all coset gP, the branching random walk eventually leaves gP . This follows from the fact that g is visited finitely many time almost surely and that starting a branching random walk at g , it comes back to gP only finitely many times, see also [\[CGM12,](#page-49-8) Proposition 4.5] for a more detailed proof.

Define now

$$
r_{\mathcal{P}} = \sup\{r > 1 \colon t_{r,P} \le 1 \text{ for all } P \in \mathbb{P}_0\}.
$$

Then for $1 < r \leq r_{\mathcal{P}}$, we have $\omega_P(r) \leq 0$ for all $P \in \mathbb{P}_0$, hence Γ has a parabolic gap for the Green function and $H_r(n)$ has purely exponential growth. Moreover, r_p is the transition for the branching random walk spending infinite times in maximal parabolic subgroups : if $r \leq r_{\mathcal{P}}$, then the branching random walk eventually leaves every coset $gP, P \in \mathbb{P}_0$, while if $r > r_{\mathcal{P}}$, it accumulates in infinitely many cosets gP for at least one of the $P \in \mathbb{P}_0$.

Recall that the limit set Λ of a branching random walk inside the Bowditch boundary $\partial \Gamma$ is the set of accumulation points in $\partial \Gamma$ of the trace of BRW(Γ, ν, μ), which is the set of elements of Γ that are ever visited by the branching random walk. We take the occasion to derive the following consequence which sheds some light on the geometry of the limit set.

Proposition 7.5. Let Γ be a relatively hyperbolic group endowed with a finitely supported admissible and symmetric probability measure μ . Consider a probability measure ν on N with mean $r \leq R$. Let Λ be the limit set inside the Bowditch boundary of Γ of the branching random walk associated with μ and ν .

- (1) If $r < r_{\mathcal{P}}$, then almost surely, Λ does not contain any parabolic limit point.
- (2) If $r > r_{\mathcal{P}}$, then almost surely, Λ contains an infinite number of parabolic limit points.

Note that Λ cannot contain all parabolic limit points in case (2). Otherwise, since parabolic limit points are dense, Λ would coincide with the whole Bowditch boundary, but this is impossible since its Hausdorff dimension with respect to the shortcut distance is at most half the Hausdorff dimension of the whole boundary by [\[DWY22,](#page-50-0) Theorem 1.1, Theorem 1.2].

This result is a consequence of the following one. For $x \in \Gamma$ and $C \geq 0$, we denote by $\Omega(x, C)$ the partial cone at x of width C, which is the set of points $y \in \Gamma$ such that x is within C of a transition point on a geodesic from e to y.

Proposition 7.6. Let Γ be a relatively hyperbolic group endowed with a finitely supported admissible and symmetric probability measure μ . For every $r \in [1, R]$, there exist $\beta > 0$ with the following property. Consider a probability measure ν on N with mean $r \leq R$. For any $x \in \Gamma$, the probability that the branching random walk visits $\Omega(x, C)$ is at most $C_1(1+|x|^{\beta})G(e, x|r)$, where C_1 is a constant.

The proof of this proposition relies mostly on material from [\[DWY22\]](#page-50-0) and its proof is postponed to the Appendix.

Proof of Proposition [7.5.](#page-35-0) As we saw above, if $r > r_p$, then the branching random walk accumulates in infinitely many cosets gP for at least one $P \in \mathbb{P}_0$. In particular, the limit set Λ contains all parabolic limit points fixed by gPg^{-1} , for every such coset gP. Thus, we only need to prove (1) to conclude and we assume that $r < r_P$.

Recall that a sequence x_n in Γ converges to a parabolic limit point ξ fixed by gPg^{-1} , $P \in \mathbb{P}_0$, if and only if the sequence of projections of x_n on gP tends to infinity. Thus, $\xi \in \Lambda$ if and only if the branching random walk visits infinitely many $\Omega(x, C)$, with $x \in gP$. Fix $g \in \Gamma$ and $P \in \mathbb{P}_0$. Denote by A_n the event

$$
A_n = \{ \text{BRW}(\Gamma, \nu, \mu) \text{ visits } \Omega(x, C) \text{ for some } x \in gP, \text{ with } |g^{-1}x| = n \}.
$$

Then, by Proposition [7.6,](#page-35-1)

$$
\mathbf{P}(A_n) \le C_1 (1+n)^{\beta} \sum_{x \in P, |x|=n} G(e, gx|r).
$$

Since g is a transition point on a relative geodesic from e to gx , by relative Ancona inequalities we get

$$
\mathbf{P}(A_n) \le C_2 (1+n)^{\beta} H_{P,r}(n).
$$

Since $t_{r,P} < 1$, $\omega_P(r) < 0$ by Proposition [7.2](#page-34-1) and so $H_{P,r}(n)$ decays exponentially fast as n goes to infinity. This proves that

$$
\sum_n \mathbf{P}(A_n) < \infty.
$$

By the Borel-Cantelli lemma, we deduce that almost surely, the branching random walk only visits finitely many $\Omega(x, C)$, $x \in gP$, hence $\xi \notin \Lambda$. Since parabolic limit points are countable, this settles the proof. points are countable, this settles the proof.

The critical case $r = r_{\mathcal{P}}$ remains open. In the context of free products, the branching random walk needs to visit qP infinitely many times in order to accumulate at ξ . As a consequence, the authors of [\[CGM12\]](#page-49-8) prove in Propositions 4.3, 4.4 and 4.5 that for $r = r_{\mathcal{P}}$, almost surely, Λ does not contain any parabolic limit point. However, for arbitrary relatively hyperbolic groups, the branching random walk can visit infinitely many partial cones $\Omega(x, C)$, $x \in gP$, without entering gP at all. Thus, we might need new material to figure this critical case.

7.2. Convergent Poincaré series. Let Γ be a relatively hyperbolic group and let μ be a finitely supported symmetric admissible probability measure on Γ. We consider the Poincaré series $\Theta_{\Gamma}(r, s)$ defined above and for P a maximal parabolic subgroup, the Poincaré series $\Theta_P (r, s)$ defined by

$$
\Theta_P(r,s) = \sum_{x \in P} G(e,x|r) e^{-sd(e,x)}.
$$

In [\[DWY22,](#page-50-0) Example C], we proved that if a finitely generated group Γ_0 can be endowed with a symmetric finitely supported admissible probability measure μ_0 such that $\Theta_{\Gamma_0}(r_0, \omega_{\Gamma_0}(r_0))$ is convergent for some $r_0 < R_0$, then the free product $\Gamma = \Gamma_0 * \mathbb{Z}^d$ can also be endowed with a symmetric finitely supported admissible probability measure μ such that $\Theta_{\Gamma}(r, \omega_{\Gamma}(r))$ is convergent for some $r \leq R$ depending on r_0 . Here, R_0 denotes the inverse of the spectral radius of μ_0 and R the inverse of the spectral radius of μ .

Note that in this situation, Γ is relatively hyperbolic with respect to the conjugates of Γ_0 and \mathbb{Z}^d . When considered as a maximal parabolic subgroup of Γ , we will write P for Γ_0 in the sequel, for sake of consistency with the previous sections.

The question of whether such a couple (Γ_0, μ_0) exists was left unanswered, but we announced that it was possible to construct one. We provide the details of this construction here and prove a more precise result.

Let \mathbb{F}_n be the free group with n generators and let $\Gamma_0 = \mathbb{F}_n \times \mathbb{F}_m$, $n \neq m$. The Cayley graph of Γ_0 is a Cartesian product of two regular trees T_{l_1}, T_{l_2} with respective degrees $l_1 = 2n$ and $l_2 = 2m$. We consider the measure μ_0 on Γ_0 defined by [\(5.1\)](#page-18-3). We deduce the following from Theorem [5.1.](#page-18-0)

Proposition 7.7. There exists $1 < r_0 < R_0$ such that for $r \leq r_0$, the Poincaré series $\Theta_{\Gamma_0}(r, \omega_{\Gamma_0}(r))$ is divergent and for $r_0 < r < R_0$, it is convergent.

We now recall how the measure μ is constructed on the free product $\Gamma = \Gamma_0 * \Gamma_1$ where $\Gamma_1 = \mathbb{Z}^d$. Let μ_1 be any finitely supported symmetric admissible probability measure on \mathbb{Z}^d . Following [\[DWY22\]](#page-50-0), we assume that $d \geq 3$ for convenience, so that the random walk associated with μ_1 is transient at the spectral radius, i.e. the Green function $G_{\mu_1}(e, e|R_1)$ is finite, where R_1 is the inverse of the spectral radius of μ_1 .

For $0 \leq \alpha \leq 1$, we set

$$
\mu_{\alpha} = \alpha \mu_1 + (1 - \alpha) \mu_0.
$$

For simplicity, we write $\mu = \mu_{\alpha}$ below. We write G for the Green function associated with μ and G_i for the Green function associated with μ_i . By [\[DWY22,](#page-50-0) (10),(11)], there exist two numbers $w_{0,\alpha,r}$ and $w_{1,\alpha,r}$ and continuous non-decreasing functions $\zeta_{i,\alpha}$ of $r \leq R$ such that

(7.3)
$$
G(e, x|r) = \frac{1}{1 - w_{0,\alpha,r}} G_0(e, x | \zeta_{0,\alpha}(r)), \ x \in \Gamma_0
$$

and

(7.4)
$$
G(e, x|r) = \frac{1}{1 - w_{1,\alpha,r}} G_1(e, x | \zeta_{1,\alpha}(r)), \ x \in \Gamma_1 = \mathbb{Z}^d.
$$

Furthermore, we have $\zeta_{\alpha,0}(r) = \frac{(1-\alpha)r}{1-w_{0,\alpha,r}}$, and

(7.5)
$$
w_{0,\alpha,r} = \sum_{n\geq 1} \mathbf{P}(X_n = e, X_k \neq e, 1 \leq k < n, \text{ first step chosen using } \alpha \mu_1) r^n.
$$

Similar expressions hold for $w_{1,\alpha,r}$ and $\zeta_{\alpha,1}(r)$.

Lemma 7.8. For fixed α , the functions $w_{i,\alpha,r}$ and $\zeta_{\alpha,i}(r)$ are (strictly) increasing in r.

Proof. The functions $w_{i,\alpha,r}$ are power series in r with positive coefficients, so they are increasing. It follows from the above expression of $\zeta_{\alpha,i}$ that these functions are increasing too.

We prove here the following.

Theorem 7.9. If α is small enough, then there exist $r_*(\alpha) < r_{\sharp}(\alpha) < R$ so that the following holds. For $r \leq r_*(\alpha)$, the Poincaré series $\Theta_{\Gamma}(r, \omega_{\Gamma}(r))$ is divergent and Γ has a parabolic gap for the Green function. On the other hand, at $r = r_{\rm \#}(\alpha)$, it is convergent and $\omega_P(r) = \omega_\Gamma(r)$, where $P = \Gamma_0$.

In the proof of Theorem [7.9,](#page-37-0) we shall also use the following result which is an enhanced version of [\[DWY22,](#page-50-0) (14)].

Lemma 7.10. For every $\epsilon > 0$, there exists α_0 such that for $\alpha \leq \alpha_0$, the following holds. For every $r \in [0, R]$ and for every $x \in \Gamma_1 \setminus \{e\}$, we have

$$
G(e, x|r) \le \epsilon.
$$

Proof. By [\[DWY22,](#page-50-0) Lemma 3.15], if $\alpha \leq \alpha_0$, then $w_{1,\alpha,r}$ stays bounded away from 1 and $\zeta_{1,\alpha}(r)$ converges to 0 as α tends to 0 uniformly over $r \leq R$. We conclude from [\(7.4\)](#page-37-1) that for small enough α , independently of r, for every $x \neq e \in \Gamma_1$ we have

$$
G(e, x|r) \le \frac{G_1(e, x|\zeta_{1,\alpha}(r))}{1 - w_{1,\alpha,r}} \le \epsilon.
$$

We are ready to complete the proof of Theorem [7.9.](#page-37-0) Let us first explain briefly how the quantities α , $r_*(\alpha)$ and $r_\sharp(\alpha)$ are chosen.

By [\[DWY22,](#page-50-0) Lemma 3.14], as α converges to 0, $\zeta_{0,\alpha}(R)$ converges to R_0 . Let us now fix any $r_1 > r_0$, where r_0 is given in Proposition [7.7.](#page-37-2) Thus, there exists α_1 so that $\zeta_{\alpha,0}(R) > r_1$ holds for any $\alpha \leq \alpha_1$. Since $\zeta_{0,\alpha}(r)$ is increasing in r, there exist $r_* < r_{\sharp} < R$ depending on α such that $\zeta_{\alpha,0}(r_*) = r_0$ and $\zeta_{0,\alpha}(r_{\sharp}) = r_1$.

We will also need to choose some $\epsilon > 0$ only depending on μ_0 , μ_1 and r_1 such that Equation [\(7.10\)](#page-39-1) holds below. Then, we choose α small enough such that the conclusions of Lemma [7.10](#page-38-0) holds for such ϵ and such that there exist $r_* < r_{\sharp} < R$ with $\zeta_{\alpha,0}(r_*)=r_0$ and $\zeta_{0,\alpha}(r_{\sharp})=r_1$.

Proof of Theorem [7.9.](#page-37-0) The proof follows the lines of [\[DWY22,](#page-50-0) Example C]. We will write as announced above P for Γ_0 when it is considered as a maximal parabolic subgroup of Γ. In particular, we write

$$
\omega_{\Gamma_0}(s) = \limsup_{n \to \infty} \frac{1}{n} \log \sum_{x \in \Gamma_0, |x| = n} G_0(e, x | s)
$$

for the growth rate of the Green function G_0 associated with μ_0 and

$$
\omega_P(r) = \limsup_{n \to \infty} \frac{1}{n} \log \sum_{x \in P, |x| = n} G(e, x|r)
$$

for the growth rate of the Green function G associated with μ , induced on P. By [\(7.3\)](#page-37-3), we see that

(7.6)
$$
\omega_P(r) = \omega_{\Gamma_0}(\zeta_{0,\alpha}(r)).
$$

We will also write $\Theta_{\Gamma_0}(r,s)$ for the Poincaré series associated with μ_0 on Γ_0 and $\Theta_P(r, s)$ for the Poincaré series associated with μ induced on P, defined as above by

$$
\Theta_P(r,s) = \sum_{x \in P} G(e,x|r) e^{-s|x|}.
$$

If α is small enough, there exist $r_* < r_{\sharp} < R$ with $\zeta_{\alpha,0}(r_*) = r_0$ and $\zeta_{0,\alpha}(r_{\sharp}) = r_1$. First, if $r \leq r_*$, then $\zeta_{0,\alpha}(r) \leq r_0$. By Proposition [7.7,](#page-37-2) $\Theta_{\Gamma_0}(\zeta_{0,\alpha}(r), \omega_{\Gamma_0}(\zeta_{0,\alpha}(r)))$ diverges, hence $\Theta_P(r, \omega_P(r))$ also diverges. Consequently, according to [\[DWY22,](#page-50-0) Corollary 3.9, $\omega_P(r) < \omega_\Gamma(r)$. Moreover, since Γ_1 is amenable, we also deduce from Proposition [7.3](#page-34-2) that the induced growth rate on Γ_1 considered as a maximal parabolic subgroup is smaller than $\omega_{\Gamma}(r)$. In other words, Γ has a parabolic gap for the Green function on $(0, r_*]$. Therefore, [\[DWY22,](#page-50-0) Lemma 3.7] yields that the Poincaré series $\Theta_{\Gamma}(r, \omega_{\Gamma}(r))$ is divergent for $r \leq r_*$.

Next, fix $r = r_{\sharp}$ so that $\zeta_{0,\alpha}(r) = r_1 > r_0$. According to [\[DWY22,](#page-50-0) (15)],

$$
\Theta_{\Gamma}(r,s) \leq G(e,e|r) \sum_{k\geq 0} \left(\sum_{x \in \Gamma_0 \setminus \{e\}} \frac{G(e,x|r)}{G(e,e|r)} e^{-s|x|} \right)^k \left(\sum_{y \in \Gamma_1 \setminus \{e\}} \frac{G(e,y|r)}{G(e,e|r)} e^{-s|y|} \right)^k
$$

 $\text{By (7.6)}, \ \omega_P(r) = \omega_{\Gamma_0}(\zeta_{0,\alpha}(r)) = \omega_{\Gamma_0}(r_1) \text{ and so } \omega_{\Gamma}(r) \geq \omega_{\Gamma_0}(\zeta_{0,\alpha}(r)).$ $\text{By (7.6)}, \ \omega_P(r) = \omega_{\Gamma_0}(\zeta_{0,\alpha}(r)) = \omega_{\Gamma_0}(r_1) \text{ and so } \omega_{\Gamma}(r) \geq \omega_{\Gamma_0}(\zeta_{0,\alpha}(r)).$ $\text{By (7.6)}, \ \omega_P(r) = \omega_{\Gamma_0}(\zeta_{0,\alpha}(r)) = \omega_{\Gamma_0}(r_1) \text{ and so } \omega_{\Gamma}(r) \geq \omega_{\Gamma_0}(\zeta_{0,\alpha}(r)).$ Moreover, by Theorem [5.1,](#page-18-0) $r_0 > 1$, so that $r_1 > 1$ and so we deduce from Lemma [2.1](#page-6-1) and Proposition [2.6](#page-8-2) that $\omega_{\Gamma_0}(r_1) \geq c > 0$. Therefore,

$$
\omega_{\Gamma}(r) \geq c > 0.
$$

Using [\(7.3\)](#page-37-3) and Proposition [7.7,](#page-37-2) we see that

$$
(7.8)\qquad \sum_{x\in\Gamma_0\setminus\{e\}}\frac{G(e,x|r)}{G(e,e|r)}e^{-\omega_{\Gamma}(r)|x|} \le \frac{1}{G_0(e,e|r_1)}\Theta_{\Gamma_0}(r_1,\omega_{\Gamma_0}(r_1)) < \infty.
$$

Also, \mathbb{Z}^d has polynomial growth and so by [\(7.7\)](#page-39-2)

$$
(7.9) \qquad \sum_{y \in \Gamma_1 \setminus \{e\}} \frac{1 - w_{1,\alpha,r}}{G(e,e|r)} e^{-\omega_{\Gamma}(r)|y|} \le \frac{1}{G_1(e,e|\zeta_{1,\alpha}(r_{\sharp}))} \sum_{y \in \Gamma_1 \setminus \{e\}} e^{-c|y|} \le C_1.
$$

We then choose $\epsilon > 0$ such that

(7.10)
$$
C_2 := \epsilon \frac{1}{G_0(e, e|r_1)} \Theta_{\Gamma_0}(r_1, \omega_{\Gamma_0}(r_1)) C_1 < 1.
$$

Note that ϵ does not depend on μ but only on C_1 , μ_0 and r_1 . In particular, it does not depend on α and so by Lemma [7.10,](#page-38-0) we can choose α small enough so that for every $y \in \Gamma_1 \setminus \{e\},\$

$$
G(e, y|r) \le \epsilon.
$$

Combining [\(7.8\)](#page-39-3), [\(7.9\)](#page-39-4), [\(7.10\)](#page-39-1) and [\(7.2\)](#page-38-0), we get

$$
\Theta_{\Gamma}(r, \omega_{\Gamma}(r)) \le G(e, e|r) \sum_{k \ge 0} \left(\epsilon \frac{1}{G_0(e, e|\zeta_{0,\alpha}(r))} \Theta_{\Gamma_0}(\zeta_{0,\alpha}, \omega_{\Gamma_0}(\zeta_{0,\alpha}(r))) C_1 \right)^k
$$

$$
\le G(e, e|r) \sum_{k \ge 0} C_2^k,
$$

so that $\Theta_{\Gamma}(r, \omega_{\Gamma}(r))$ is finite.

Finally, we deduce from [\[DWY22,](#page-50-0) Lemma 3.7] that at $r = r_{\sharp}$, Γ does not have a parabolic gap for the Green function. Since Γ_1 is amenable, we necessarily have $\omega_{\Gamma}(r) = \omega_{P}(r)$ by Proposition [7.3.](#page-34-2) This concludes the proof.

Remark 7.11. In the proof of Theorem [7.9,](#page-37-0) note that we can choose r_1 arbitrarily close to r_0 . Unfortunately, as r_1 goes to r_0 , $\Theta_{\Gamma_0}(r_1,\omega_{\Gamma_0}(r_1))$ tends to infinity, so ϵ satisfying [\(7.10\)](#page-39-1) converges to 0. Consequently, the parameter α also tends to 0. In other words, as r_1 tends to r_0 , we need to choose a measure μ that tends to the measure μ_0 distributed on Γ_0 . Now, since the functions $\zeta_{\alpha,i}$ of r depend on α , we cannot guarantee that $r_{\text{#}}$ tends to r_* . In particular, we cannot prove that there is a true phase transition for the convergence of the Poincaré series $\Theta_{\Gamma}(r, \omega_{\Gamma}(r))$ at r_* . **Corollary 7.12.** If α is small enough, then the following holds. For $r = r_{\sharp}(\alpha)$, Γ does not have a parabolic gap for the Green function and [\(7.1\)](#page-33-1) and [\(7.2\)](#page-33-2) do not hold.

Proof. By Theorem [7.9,](#page-37-0) if $r = r_{\text{t}}(\alpha)$, then the Poincaré series $\Theta_{\Gamma}(r, \omega_{\Gamma}(r))$ is convergent and Γ does not have a parabolic gap. Assume by contradiction that there exists C such that

$$
H_r(m+n) \leq CH_r(n)H_r(m).
$$

Then, the quantity $CH_r(n)$ is sub-multiplicative, hence by Fekete's lemma,

$$
\omega_{\Gamma}(r) = \lim_{n \to \infty} \frac{1}{n} \log CH_r(n) = \inf_{n \ge 1} \frac{1}{n} \log CH_r(n).
$$

Thus, for every n , we have

$$
CH_r(n) \ge e^{n\omega_{\Gamma}(r)}.
$$

This implies that $\Theta_{\Gamma}(r, \omega_{\Gamma}(r))$ diverges, which is a contradiction. In particular, we see that [\(7.1\)](#page-33-1) fails. Now, (7.1) is a direct consequence of [\(7.2\)](#page-33-2), with $C = (C')^3$, hence (7.2) also fails.

We also deduce the following from Theorem [6.17](#page-32-0) and Theorem [7.9.](#page-37-0)

Corollary 7.13. If α is small enough, then the following holds. For $r \leq r_*(\alpha)$, the measure ν_e on the Bowditch boundary has no atom and is supported on the set of conical limit points. For $r = r_{\sharp}(\alpha)$, it is purely atomic and is supported on the set of parabolic limit points.

Remark 7.14. In [\[CGM12\]](#page-49-8), the authors prove that in the context of free products, we always have $\omega_P(r) < \omega_\Gamma(r)$ for every maximal parabolic subgroup P, i.e. Γ has a parabolic gap for the Green function. However, their proof relies on an unproved statement, namely that the quantity $H_r(n)$ is sub-multiplicative for every finitely generated group and then apply this property to the maximal parabolic subgroup P, see precisely the proof of [\[CGM12,](#page-49-8) Lemma 4.7] and also [\[DWY22,](#page-50-0) Remark 3.17]. However, by Theorem [5.1,](#page-18-0) we see that sub-multiplicativity fails for the Cartesian product of two regular trees if $r \ge r_0$. Moreover, by Corollary [7.12,](#page-40-2) in the above example, if $r = r_{\text{t}}(\alpha)$, then Γ does not have a parabolic gap for the Green function.

7.3. The growth tightness property. Let d be a proper left invariant distance on Γ. The *growth rate* of Γ for d is defined as follows:

$$
\delta(\Gamma, d) := \limsup_{n \to \infty} \frac{\log \sharp \{ x \in \Gamma : d(e, x) \le n \}}{n}
$$

A nontrivial quotient $\bar{\Gamma}$ of Γ means that the kernel of the canonical projection $\Gamma \to \bar{\Gamma}$ is an infinite normal subgroup of Γ. We say that Γ is *growth tight* for the distance d if for every nontrivial quotient Γ of Γ , endowed with the quotient distance d from d, we have $\delta(\bar{\Gamma}, \bar{d}) < \delta(\Gamma, d)$.

Let us assume that Γ is a relatively hyperbolic group. Whenever a maximal parabolic group P has growth rate $\delta(P, d)$ strictly less than $\delta(\Gamma, d)$, we say that Γ has a parabolic gap along P fir the distance d. When Γ has parabolic gap along every maximal parabolic subgroup, we say that Γ has the parabolic gap property.

Recall that by Lemma [6.4,](#page-26-3) $\mathfrak{d}_r(x,y) = \omega_{\Gamma}(r)|x^{-1}y| + |x^{-1}y|_r$ is quasi-isometric to the word distance for $r < R$. By Lemma [6.10,](#page-28-1) we have $\delta(\Gamma, \mathfrak{d}_r) = 1$. The following result relates the gap property for Green functions to the gap property for the distance $\delta(\Gamma, \mathfrak{d}_r)$.

Proposition 7.15. Let Γ be a group endowed with a probability measure μ such that the μ -random walk is transient at the spectral radius, i.e. $G(e, e|R)$ is finite. Let $A \subset \Gamma$ be any subset. If $\omega_A(r) < \omega_{\Gamma}(r)$ for some $1 < r \leq R$, then $\delta(A, \mathfrak{d}_r) < 1$.

We need the following lemma, which generalizes [\[Tan17,](#page-51-8) Lemma 3.1] in hyperbolic groups with a similar proof.

Lemma 7.16. Under the assumption of Proposition [7.15.](#page-41-0) For any $\theta \in \mathbb{R}$, $r \leq R$ and $A \subset \Gamma$, define

$$
\omega_{A,r}(\theta) := \limsup_{n \to \infty} \frac{1}{n} \log \sum_{x \in A, |x| = n} \left[G(e, x|r) \right]^{\theta}.
$$

Then $\omega_{A,r}(\theta)$ is a convex function on R. If Γ is a relatively hyperbolic group, then $\omega_{\Gamma,r}(\theta)$ is a true limit.

Proof. Denote $H_r^{\theta}(n) := \sum_{x \in A, |x|=n} \left[G(e, x|r) \right]^{\theta}$. For $\theta_0, \theta_1 \in \mathbb{R}$ and $0 < t < 1$, by the Hölder inequality,

$$
H_r^{t\theta_0+(1-t)\theta_1}(n) \leq (H_r^{\theta_0}(n))^t (H_r^{\theta_1}(n))^{1-t}.
$$

Thus $\omega_{A,r}$ is convex:

$$
\omega_{A,r}(t\theta_0 + (1-t)\theta_1) \leq t\omega_{A,r}(\theta_0) + (1-t)\omega_{A,r}(\theta_1).
$$

If Γ is a relatively hyperbolic group, the same proof as in [\[DWY22,](#page-50-0) Lemma 3.2] shows (there with $\theta = 1$) that for $A = \Gamma$, the sequence $H_r^{\theta}(n)$ is sub-multiplicative, that is, $H_r^{\theta}(n+m) \leq CH_r^{\theta}(n)H_r^{\theta}(m)$ for some $C > 0$. Thus, the limit exists by Feketa's lemma.

As a convex function on \mathbb{R} , $\omega_{\Gamma,r}(\theta)$ is a continuous function of $\theta \in \mathbb{R}$, and is differentiable, except maybe at countably infinitely many points.

Proof of Proposition [7.15.](#page-41-0) To show $\delta(A, \mathfrak{d}_r) < 1$, it suffices to find some $\epsilon > 0$ such that

$$
\sum_{x \in A} e^{-(1-\epsilon)\mathfrak{d}_r(e,x)} = \sum_{x \in A} e^{-(1-\epsilon)\omega_{\Gamma}(r)|x|} \left[G(e,x|r) \right]^{1-\epsilon} < \infty.
$$

By Lemma [7.16,](#page-41-1) the function $\omega_{A,r}(\theta)$ is continuous in $\theta \in \mathbb{R}$. If $\omega_A(r) < \omega_{\Gamma}(r)$, we can choose $\epsilon, \eta > 0$ small enough so that

$$
\omega_A(r) + \eta < (1 - \epsilon)\omega_\Gamma(r),
$$

and at the same time, by continuity of $\omega_{A,r}(\theta)$ at $\theta = 1$, the following holds : for large enough n ,

$$
\sum_{x \in A, |x|=n} \left[G(e, x|r) \right]^{1-\varepsilon} \le e^{n(\omega_{A,r}(1)+\eta)}.
$$

By definition, $\omega_{A,r}(1) = \omega_A(r)$, so the two inequalities above yield

$$
\sum_{x \in A} e^{-(1-\epsilon)\omega_{\Gamma}(r)|x|} G(e, x|r)^{1-\epsilon} < \infty,
$$

which is the desired inequality. \Box

Given $f \in \Gamma$ and $\epsilon > 0$, let $\mathcal{V}_{\epsilon,f}$ be the set of *barrier-free* elements $x \in \Gamma$, that is, elements for which the ϵ -neighborhood of some geodesic $[e, x]$ contains a geodesic segment representing f. The following is analogous to [\[Yan19,](#page-51-9) Theorem C].

Lemma 7.17. Let Γ be a relatively hyperbolic group with parabolic gap property for Green function. Then there exists some $\epsilon > 0$ such that the set $V := V_{\epsilon,f}$ has growth rate strictly less than 1 for any element $f \in \Gamma: \delta(\mathcal{V}, \mathfrak{d}_r) < 1$.

Proof. By Proposition [7.15,](#page-41-0) it suffices to prove $\omega_{\mathcal{V}}(r) < \omega_{\Gamma}(r)$. Set

$$
a^{\omega}(n) = e^{-\omega n} \sum_{x \in \mathcal{V}, |x|=n} G(e, x; r).
$$

Assume that $\omega_{\mathcal{V}}(r) > \omega_{\mathcal{P}}(r)$ for every maximal parabolic subgroup P; otherwise the parabolic gap concludes the proof. If $\omega_{\mathcal{V}}(r) > \omega > \omega_{P}(r)$, one obtains

$$
a^{\omega}(n+m) \le c_0 \sum_{1 \le i \le n} a^{\omega}(i) \sum_{1 \le j \le n} a^{\omega}(j)
$$

by the same argument of [\[DWY22,](#page-50-0) Lemma 3.7] where $a^{\omega}(n)$ is summed up over Γ instead of $\mathcal V$. This implies via a variant of Feketa's lemma in [\[DPPS11,](#page-50-27) Lemma 4.3] that the series $\sum_{x \in \mathcal{V}} e^{-s|x|} G(e, x; r)$ diverges at $s = \omega_{\mathcal{V}}(r)$.

Fix any $L > 0$. We choose an *L-separated net* $A \subset V$ in word distance: if for any $x, y \in A$ we have $|x^{-1}y| > L$ and for any $y \in V$, there exists $x \in A$ such that $|x^{-1}y| \leq L$. Note that if $|x^{-1}y| \leq L$, then $G(e, x; r) \approx_L G(e, y; r)$. Since any ball of radius L in word distance contains a fixed number of elements, we deduce that $\Theta_A(r,s) \asymp_L \Theta_A(r,s)$ whenever they are finite. Thus, $\omega_A(r) = \omega_{\mathcal{V}}(r)$.

Following [\[Yan19,](#page-51-9) Section 4.2], we use a ping-pong argument to construct a free product of sets inside Γ : if L is large enough, there exist a finite set of elements $B \subset \Gamma$ such that the set $\mathcal{W}(A, B)$ of alternating words over A and B embeds into Γ as a free semi-group under the evaluation map. This construction uses only the word distance. Now, by [\[DWY22,](#page-50-0) Lemma 3.8], we have $\omega_A(r) < \omega_{\Gamma}(r)$ and then $\delta(\mathcal{V}, \mathfrak{d}_r) < 1$ by Proposition [7.15.](#page-41-0)

Proposition 7.18. If a relatively hyperbolic group Γ has parabolic gap for the Green function, then it is growth tight for the distance \mathfrak{d}_r . Otherwise, there exists a nontrivial quotient Γ such that $\delta(\overline{\Gamma}, \overline{\mathfrak{d}}_{\mathfrak{r}}) = \delta(\Gamma, \mathfrak{d}_r)$.

Proof. (1). We follow the proof of [\[Yan19,](#page-51-9) Corollary 4.6] in our setup. Let N be the infinite kernel of $\Gamma \to \Gamma$. We form a set A by choosing a shortest representative $h \in hN$ for each $hN \in G$ so that $\mathfrak{d}_r(e, h) = \mathfrak{d}_r(e, hN)$. By definition of the quotient distance, the growth rate of the set A for \mathfrak{d}_r is exactly the growth rate of Γ for $\bar{\mathfrak{d}}_r$.

We now choose a sufficiently long loxodromic element $f \in N$, which exists since N is infinite. If $|f|/\epsilon$ is large enough, we see that any geodesic $[e, h]$ cannot contain f in its ϵ -neighborhood. Indeed, if not, the loxodromic element f produces two transition points on some $[e, h]$ with a distance comparable with $|f|$. Now, we use the following fact given by Lemma [6.2:](#page-25-1) if u, v are two transition points in this order on a word geodesic $[x, y]$, then

(7.11)
$$
\mathfrak{d}_r(x,y) \simeq \mathfrak{d}_r(x,u) + \mathfrak{d}_r(u,v) + \mathfrak{d}_r(v,y)
$$

where \simeq denote the equality up to a uniform additive constant. We could then shorten $\mathfrak{d}_r(e, h)$ by an amount $\mathfrak{d}_r(e, f) = \omega_{\Gamma}(r) |f| + |f|_r$, giving a contradiction with the above choice of $h \in hN$ as the shortest one.

In other words, we proved that $A \subset \mathcal{V}_{\epsilon,f}$. Hence, $\delta(A, \mathfrak{d}_r) \leq \delta(\mathcal{V}_{\epsilon,f}, \mathfrak{d}_r) < 1$ by Lemma [7.17.](#page-42-0) The growth tightness follows.

(2). Assume that $\omega_P(r) = \omega_\Gamma(r)$ for a maximal parabolic subgroup P. Then,

$$
\sum_{x \in P} e^{-\mathfrak{d}_r(e,x)} = \sum_{x \in P} e^{-\omega_P(r)} G(e,x|r)
$$

hence, we see that the growth rate for \mathfrak{d}_r induced on P equals 1.

Fix a loxodromic element $f \in G$. For any large enough n, the quotient group $\overline{\Gamma}$ defined as $G/\langle \langle f^n \rangle \rangle$ is again a relatively hyperbolic group, and $P \cap \langle \langle f^n \rangle \rangle$ is trivial (see [\[Yan22,](#page-51-7) Lemma 8.9]). Thus, the set of elements in P embeds into Γ whose image we denote by \overline{P} , so $\delta(\overline{P}, \overline{\mathfrak{d}}_{\mathfrak{r}}) \geq \delta(P, \mathfrak{d}_r) = 1$. Therefore, $\delta(\overline{\Gamma}, \overline{\mathfrak{d}}_{\mathfrak{r}}) = 1$.

Relatively hyperbolic groups endowed with a word distance are always growth tight by [\[Yan14,](#page-51-10) [ACT15\]](#page-49-9). In fact, any co-compact action of a relatively hyperbolic group on a proper geodesic space contains a contracting element and thus is growth tight. Here, the existence of a contracting element in the co-compact action follows from the fact that in a relatively hyperbolic group, a loxodromic element is contracting with respect to all word quasi-geodesics: any c-quasi-geodesic outside the C-neighborhood of the axis has C-bounded projection for some $C = C(c)$. See [\[GP16,](#page-50-22) Proposition 8.5].

On the contrary, as a corollary of Theorem [7.9](#page-37-0) and Proposition [7.18,](#page-42-1) growth tightness for \mathfrak{d}_r may fail and depends on r.

Theorem 7.19. There exists a relatively hyperbolic group Γ endowed with a finitely supported symmetric and admissible probability measure μ such that the following holds. There exist $1 < r_* < r_{\sharp} < R$ such that Γ endowed with the distance \mathfrak{d}_r is growth tight for $r \leq r_*$, but is not for $r = r_{\sharp}$.

Note that the proper distance \mathfrak{d}_r is quasi-isometric to any word distance for $r < R$ by Lemma [6.4.](#page-26-3) We say that a metric space (X, d) is *D*-coarsely geodesic for some $D > 0$ if for any two points $x, y \in X$, there exists a $(1, D)$ -quasi-isometric embedding $\phi : [0, l] \to X$ for $l := d(x, y)$ so that $\phi(0) = x, \phi(l) = y$, and

$$
|d(\phi(m), \phi(n)) - |m - n|| \le D
$$

for any $0 \leq m \leq n \leq l$. It is an open question whether the Green distance is a geodesic distance on hyperbolic groups, see [\[BHM11,](#page-49-26) Section 1.7]. We shall however derive the following corollary from Theorem [7.19.](#page-43-0)

Corollary 7.20. For $r = r_{\sharp}$, (Γ, \mathfrak{d}_r) is not a coarsely geodesic metric space.

The proof requires the following observation of independent interest. Recall that an element of infinite order q in a finitely generated group Γ is called contracting for a distance d on Γ if any d-metric ball in Γ disjoint with the subgroup $\langle q \rangle$ has C-bounded projection to $\langle q \rangle$ for some universal constant $C > 0$. In a D-coarsely geodesic metric space, this is equivalent to the bounded image property: there exists $C = C(D) > 0$ such that any D-coarse geodesic outside the C-neighborhood of $\langle g \rangle$ has shortest projection of diameter at most C to it. For simplicity, we can take the same C for both statements.

Lemma 7.21. Any loxodromic element in a relatively hyperbolic group is contracting with respect to \mathfrak{d}_r where $1 \leq r \leq R$.

Proof. By [\(7.11\)](#page-42-2), the proper distance \mathfrak{d}_r is coarsely additive along the set of transition points on the word geodesic. That is, if z is a transition point on $[x, y]$ we have $\mathfrak{d}_r(x,y) \geq \mathfrak{d}_r(x,z)+\mathfrak{d}_r(z,y)-D$ for some universal $D > 0$. Let γ be a quasi-geodesic preserved by a loxodromic element g. Then γ is C_0 - contracting with respect to word distance for some $C_0 > 0$. We claim that the shortest projection z of any point x to γ for the distance \mathfrak{d}_r is D₀-close to the shortest projection w of x to γ for the word distance. Indeed, as w is uniformly close to a transition point on $[x, z]$, we see that $\mathfrak{d}_r(x, z) + D_0 \geq \mathfrak{d}_r(x, w) + \mathfrak{d}_r(w, z)$ for some $D_0 = D_0(C_0) > 0$. By the definition of \mathfrak{d}_r -shortest projection, we have $\mathfrak{d}_r(x, z) \leq \mathfrak{d}_r(x, w)$ and thus the claim follows.

We now prove that γ is contracting for \mathfrak{d}_r . Pick any \mathfrak{d}_r -distance ball B centered at x disjoint with γ . Let $y \in B$ so that the projections denoted by u, v respectively of x, y to γ realizes the \mathfrak{d}_r -diameter of that of B to γ . By Lemma [6.4,](#page-26-3) \mathfrak{d}_r is quasiisometric to word metric for $1 \leq r < R$. For $r = R$, by [\[GL13,](#page-50-23) Lemma 2.1], $f(n) := \max\{G(e, x|R) : x \in S_n\} \to 0$ as $n \to \infty$, so $f(n)$ is a proper function. We can thus choose $\mathfrak{d}_r(u, v)$ large enough so that $|u^{-1}y| \geq C_0$, hence the contracting property of γ in word distance implies that u, v are uniformly close to transition points on [x, y]. By the additive property of $\mathfrak{d}_r(x, y)$ along transition points, we obtain $\mathfrak{d}_r(x,y) \geq \mathfrak{d}_r(x,u) + \mathfrak{d}_r(u,v) + \mathfrak{d}_r(v,y) - D_1$ for some $D_1 > 0$. As B is a \mathfrak{d}_r -distance ball disjoint with γ , we have $\mathfrak{d}_r(x, y) \leq \mathfrak{d}_r(x, u)$. We then obtain a contradiction if $\mathfrak{d}_r(u, v) > D_1$. Thus, the contracting property for \mathfrak{d}_r follows. contradiction if $\mathfrak{d}_r(u, v) > D_1$. Thus, the contracting property for \mathfrak{d}_r follows.

Proof of Corollary [7.20.](#page-43-1) Assume on the contrary that \mathfrak{d}_r is a coarsely geodesic distance for $r > r₊$. By Lemma [7.21,](#page-43-2) the axis of any loxodromic element satisfies the bounded image property for coarse geodesics, so any loxodromic element is contracting in the sense of [\[Yan14\]](#page-51-10). As the action on Γ is co-compact, the same argument as in [\[Yan14\]](#page-51-10) holds verbatim by replacing word geodesics with coarse \mathfrak{d}_r geodesics and we can show that \mathfrak{d}_r on Γ is growth tight. However, this contradicts Theorem [7.19.](#page-43-0) Thus the corollary follows.

In [\[ACT15,](#page-49-9) Question 1], Arzhantseva-Cashen-Tao asked whether growth tightness is invariant among cocompact actions on geodesic metric spaces. Cashen-Tao [\[CT16\]](#page-49-5) showed the first examples of product groups with growth tightness for one generating set but not for another generating set. Examples of non-growth tight relatively hyperbolic groups with non-cocompact actions were already considered in [\[ACT15,](#page-49-9) Obs. 7.9]. As the action is not-cocompact, the induced pseudo-distance on the group pulled back from the action is not quasi-isometric to the word distance. It is natural to ask the following variant of Arzhantseva-Cashen-Tao's question for relatively hyperbolic groups about quasi-isometry invariance of growth tightness : if a proper (pseudo-)distance d on Γ is quasi-isometric to the word distance, does the growth tightness for d hold ? In our last corollary, we produce examples of non-geodesic distances on relatively hyperbolic groups Γ that answer negatively this question.

Corollary 7.22. There exists a relatively hyperbolic group with a proper left invariant distance quasi-isometric to the word distance which does not have the growth tightness property.

Appendix A. Probability that the branching random walk visits partial cones

We consider a relatively hyperbolic group Γ. Our goal is to prove Proposition [7.6.](#page-35-1) The proof basically consists of a reorganization of arguments of [\[DWY22\]](#page-50-0). We first recall some notations and definitions.

Given a subset A of Γ, we write $G(x, y; A|r)$ for the Green function restricted to paths staying in A, expect maybe the first and last point. That is,

$$
G(x, y; A|r) = \sum_{n \ge 0} \sum_{z_1, ..., z_{n-1} \in A} r^n \mu(x^{-1}z_1) \mu(z_1^{-1}z_2) ... \mu(z_{n-1}^{-1}y).
$$

Fix $C > 0$ and $x \in \Gamma$. The C-partial cone $\Omega(x, C)$ consists of points $z \in G$ such that x is within C of an (η, L) -transition point on the geodesic $[e, z]$. Let $C > 0$ be any sufficiently large constant given by [\[DWY22,](#page-50-0) Lemma 2.9] so that the relative thin triangle property holds for (η, L) -transition points : for every triple (x, y, z) of Γ, any (η, L) -tranition point on $[x, y]$ is within C of either an (η, L) -transition point on $[x, z]$ or an (η, L) -transition point on $[y, z]$.

Let $B([x, z])$ be the ball centered at the middle point of $[x, z]$ of radius $d(x, z)/2$. Define $U(x)$ to be the union of the balls $B([x, z])$ for all geodesics $[x, z]$ between x and $z \in \Omega(x, C)$. That is,

$$
U(x) := \bigcup \{ B([x,z]) : \forall [x,z], \forall z \in \Omega(x,C) \}.
$$

It is clear that $\Omega(x, C)$ is contained in $U(x)$.

Fix $\epsilon \in (0, 1/2)$. Let $U_{\epsilon}(x)$ be the set of points $z \in U(x)$ such that $[x, z]$ contains a transition point w being at distance at least $\epsilon d(x, z)$ to one of the endpoints:

$$
\max\{d(w, x), d(w, z)\} \ge \epsilon d(x, z).
$$

For any $m \geq 1$, let $U_{\epsilon}(x, m)$ be the set of elements $z \in U_{\epsilon}(x)$ such that $d(x, z) \geq m$.

We now consider a finitely supported symmetric and admissible probability measure μ on Γ and a probability measure ν on N. We denote by BRW(Γ , ν , μ) the branching random walk associated with ν and μ . In what follows, we shall often use the following estimates proved in [\(6.2\)](#page-26-1). There exists $\alpha > 0$ such that for any $x \in \Gamma$, for every $r \geq 1$,

$$
(A.1) \tG(e, x|r) \ge e^{-\alpha |x|}.
$$

Also, by Lemma [7.2](#page-33-2) there exists $c_1 > 0$ such that for any $n \geq 1$,

♯Sⁿ ≤ c1e vn (A.2)

Lemma A.1. For any $\epsilon \in (0, 1/2)$, there exists $\kappa_0 > 0$ such that for every $\kappa \geq \kappa_0$, the following holds. For all but finitely many $x \in \Gamma$: the following event

$$
E_1 := \{ BRW(\Gamma, \nu, \mu) \text{ first enters } U(x) \text{ at a point } z \in U_{\epsilon}(x, \kappa \log |x|) \}
$$

has probability at most $G(e, x|r)$.

Proof. Let us freeze all particles of BRW(Γ, ν , μ) when the event E_1 happens, and denote by $\mathcal Z$ the collection of frozen particles. Set $m := \kappa \log n$ for simplicity, where $n = |x|$. Then for $z \in \mathcal{Z}$ we have

- (1) $d(x, z) > m$,
- (2) max $\{d(y, x), d(y, z)\} > \epsilon d(x, z) 3C$ where y is an (η, L) -transition point on $[e, z]$ given by [\[DWY22,](#page-50-0) Lemma 6.6].

 $\overline{3}$ $\overline{3}$ $\overline{2}$ $\overline{3}$ $\overline{$

As the genealogy path from e to z does not intersect $B(y, \epsilon d(x, z) - 3C)$, the expected number of particles frozen at $z \in U_{\epsilon}(x)$ is upper bounded by

$$
G(e, z; [U_{\epsilon}(x)]^c | r) \le G(e, z; [B(y, \epsilon d(x, z) - 3C)]^c | r) \le e^{-e^{o[\epsilon d(x, z) - 3C]}}.
$$

where $\delta = \delta(\eta, L)$ be given by [\[DG21,](#page-49-6) Proposition 3.5]. As a consequence, there exist $\epsilon_1 = \epsilon_1(\epsilon, \delta, v)$ and $n_0 > 0$ such that for any $m > \kappa \log n_0$, we have by [\(A.2\)](#page-45-0) that

$$
\mathbf{E}[\sharp \mathcal{Z}] \leq \sum_{z \in U_{\epsilon}(x,m)} G\big(e, z; [B(y, \epsilon d(x, z) - 3C]^c \big| r\big) \leq \sum_{k=m}^{\infty} c_1 \cdot e^{kv - e^{\delta(\epsilon k - 3C)}} \leq e^{-e^{\epsilon_1 m}}.
$$

Choose κ so that $n^{\kappa \epsilon_1} > \alpha n$ holds for any $n > n_0$. Then,

$$
\mathbf{P}(E_1) \le \mathbf{E}(\sharp \mathcal{Z} \ge 1) \le e^{-n^{\epsilon_1 \kappa}} \le e^{-\alpha n} \le G(e, x|r)
$$

where the last inequality uses $(A.1)$.

Similarly, we prove the following.

Lemma A.2. For every $K \geq 1$ and $\hat{C} \geq 0$, there exists $\kappa_0 > 0$ such that for all $\kappa \geq \kappa_0$, the following holds. For any sufficiently large $n \geq 1$, the following event

 $E_2 := \{\text{BRW}(\Gamma, \nu, \mu) \text{ eventually visits a point } z \in \Omega(x, \hat{C}) \text{ with } d(e, z) \leq K d(e, x)$ but without entering $B(x, \kappa \log |x|)$ where $|x| \geq n$ }

has probability at most $G(e, x|r)$.

Proof. We freeze particles when the event E_2 happens and denote by $\mathcal Z$ the set of frozen particles. By [\[DWY22,](#page-50-0) Lemma 2.11], if $y \in [e, z]$ is a transition point C-close to x, the expected number of particles frozen at $z \in B(e, K|x|)$ is upper bounded by

$$
G(e, z; [B(y, \kappa \log |x| - \hat{C})]^c | r) \le e^{-n^{\delta \kappa}}
$$

where c_2 depends on \hat{C} , δ . Thus, we have

$$
\mathbf{E}[\sharp \mathcal{Z}] \leq \sum_{m \geq n} \sum_{z \in S_{Km}} G(e, z; [B(y, \kappa \log m - \hat{C})]^c | r) \leq \sum_{m \geq n} c_1 e^{vKm} e^{-c_2 m^{\delta \kappa}}.
$$

Choose κ , $n_0 > 0$ so that $c_2 m^{\delta \kappa} - vKm > \alpha m$ holds for any $m \ge n_0$. The conclusion follows again. follows again.

Lemma A.3. For any $K > 1$, there exist ϵ_0 , κ_0 such that for all $\epsilon \leq \epsilon_0$ and $\kappa \geq \kappa_0$, the following holds. There exists $c < 0$ such that for all but finitely many $x \in \Gamma$, the following event

$$
E_3 := \{ BRW(\Gamma, \nu, \mu) \text{ first enters } U(x) \text{ at a point}
$$

$$
z \in U(x) \setminus U_{\epsilon}(x, \kappa \log |x|) \text{ with } |z| \ge K|x| \}
$$

has probability at most $c \cdot G(e, x|r)$.

Proof. Let V be the set of $z \in U(x) \setminus U_{\epsilon}(x, \kappa \log |x|)$ satisfying $|z| \geq K|x|$. By definition,

- (1) either $d(z, x) \leq \kappa \log |x|$,
- (2) or the $[\epsilon, 1 \epsilon]$ -percentage of $[x, z]$ does not contain any (η, L) -transition point.

If K and κ are fixed, noticing that

$$
(K-1)|x| \le d(x, z) \le (1 + 1/K)|z|,
$$

the case (1) is impossible for sufficiently large $|x|$. Thus, it suffices to consider the case (2). Set $K_1 = \epsilon(1 + 1/K)$ and $K_2 = (1 - 2\epsilon)(1 - 1/K)$. By [\[DWY22,](#page-50-0)

Lemma 6.1], there exist a unique coset $P_z \in \mathbb{P}$ such that if y_1, y_2 are the entrance and exit points of $[x, z]$ in $N_n(P_z)$, then

(A.3)
$$
\max\{d(x, y_1), d(y_2, z)\} \le \epsilon d(x, z) \le \epsilon (1 + 1/K)|z| \le K_1|z|,
$$

and

(A.4)
$$
d(y_1, y_2) \ge (1 - 2\epsilon)d(x, z) \ge (1 - 2\epsilon)(1 - 1/K)|z| \ge K_2|z|.
$$

Before moving on, we need the following facts about y_1 . By relative thin triangle property, there exists a constant \hat{C}_1 depending only on C so that x is \hat{C} -close to a transition point on $[e, y_1]$. Moreover, there exists \hat{C}_2 depending on C, η so that the projection $\pi_{N_{\eta}(P_z)}(e)$ of e to $N_{\eta}(P_z)$ is within \hat{C}_2 of y_1 and y_1 is \hat{C}_2 -close to a transition point on $[e, w]$ for any $w \in N_n(P_z)$. For given $P \in \mathbb{P}$, let $P(y_1)$ denote the set of $w \in N_{\eta}(P)$ with $d(w, y_1) \leq d(e, y_1)$, where y_1 is \hat{C}_2 -close to $\pi_{N_{\eta}(P)}(e)$.

Consider first the sub-event E_{30} of E_3 , where BRW(Γ, ν , μ) enters $N_n(P_z)$ at a point $w \in P_z(y_1)$. Thus, $d(e, w) \leq d(w, y_1) + d(e, y_1) \leq 2|y_1|$. The same proof of [\[DWY22,](#page-50-0) Lemma 6.6] implies that $B(y_1, d(y_1, x) - 3C)$ is contained in $U(x)$. Thus, the particle does not visit $B(y_1, d(y_1, x) - 3C)$. Assume first that $|y_1| > 2|x|$. Note that there exists $n_0, \kappa_0 > 0$ so that for any $|x| \geq n_0$, we have

$$
\kappa_0 \log |y_1| \le |y_1| - |x| - 3C \le d(y_1, x) - 3C.
$$

In particular, the branching random walk does not visit $B(y_1, \kappa_0 \log |y_1|)$ before arriving at w. Now, if $|y_1| \leq 2|x|$, then $d(e, w) \leq 2|y_1| \leq 4|x|$. By definition of E_3 , the branching random walk does not visit $B(x, \kappa_0 \log |x|)$. In summary, this sub-event E_{30} is included into the event E_2 in Lemma [A.2](#page-46-0) with constants \hat{C} and $K = 4$, so there exists $\kappa_0 \geq \kappa > 0$ so that the probability of E_{30} is at most $G(e, x|r)$.

Now, it remains to consider the particles of BRW(Γ, ν , μ) in the event E_3 that do not enter $N_{\eta}(P_z)$ at some point $w \in P_z(y_1)$. Then, we have the following two sub-events denoted by E_{31} and E_{32} respectively: the particles either do not visit $N_{\eta}(P_z)$ at all or do visit $N_{\eta}(P_z)$ but at a first entrance point w not in $P_z(y_1)$. Let us denote by W the set of points $w \in N_n(P) \setminus P(y_1)$ for all $P \in \mathbb{P}$ where y_1 is \hat{C}_2 -close to $\pi_{N_{\eta}(P)}(e)$ and $|y_1|>|x|$. In the first case, we freeze particles when they first enter the set V . In the second case, we freeze particles when they first enter the set W. We denote by \mathcal{Z}_{31} and \mathcal{Z}_{32} respectively the sets of frozen particles. We have

(A.5)
$$
\mathbf{E}[\sharp \mathcal{Z}_{31}] \leq \sum_{z \in V} G(e, z; [N_{\eta}(P_z)]^c | r),
$$

(A.6)
$$
\mathbf{E}[\sharp \mathcal{Z}_{32}] \leq \sum_{w \in W} G(e, w; [N_{\eta}(P_z)]^c | r),
$$

We first bound $\mathbf{E}[\sharp \mathcal{Z}_{32}]$ in [\(A.6\)](#page-47-0). Recall that $\pi_{N_{\eta}(P_z)}(e)$ is \hat{C}_2 -close to y_1 . If BRW(Γ, ν , μ) enters $N_{\eta}(P_z) \setminus P_z(y_1)$ at a point w, by [\[DWY22,](#page-50-0) Lemma 2.12], for every $M \geq 0$, there exists η_0 such that for all $\eta \geq \eta_0$,

$$
G(e, w; [N_{\eta}(P_z)]^c | r) \le e^{-Md(\pi_{N_{\eta}(P_z)}(e), w)} \le c_0 e^{-Md(w, y_1)}
$$

for some $c_0 = c_0(\hat{C}_2) > 0$. Using $(A.1)$, we first sum up over y_1 with $|y_1| > |x|$ and then $w \in P_z(y_1)$ with $d(y_1, w) > d(e, y_1)$:

$$
\mathbf{E}[\sharp \mathcal{Z}_{32}] \leq \sum_{n \geq |x|} c_0 \cdot e^{vn} \sum_{m \geq n} e^{(v-M)m}.
$$

Choosing $M > 2v + \alpha$ that is is independent of η , we have that for $\eta \geq \eta_0$,

$$
\mathbf{E}[\sharp \mathcal{Z}_{32}] \le c e^{-\alpha|x|} \le cG(e, x|r)
$$

where c depends on c_0 , thus on η .

We are left to bound $\mathbf{E}[\sharp \mathcal{Z}_{31}]$ in [\(A.5\)](#page-47-1). As the support of μ is finite, we can replace each edge in the geodesic from z to $y_2 \in N_{\eta}(P_z)$ by a μ -trajectory with uniformly bounded length. Moving possibly the endpoint y_2 up to a bounded distance depending on supp (μ) , this produces a trajectory outside $N_{\eta}(P_z)$ for the μ -random walk from z to y_2 so that its length is bounded above by a linear function of $d(y_2, z)$. This implies the existence of a positive β independent on z, y_1, y_2 and η such that

$$
G(z, y_2; [N_{\eta}(P_z)]^c | r) \ge e^{-\beta d(y_2, z)} \ge e^{-\beta K_1 |z|}.
$$

Taking into account that

$$
G(e, z; [N_{\eta}(P_z)]^c|r) \cdot G(z, y_2; [N_{\eta}(P_z)]^c|r) \le G(e, e|r) \cdot G(e, y_2; [N_{\eta}(P_z)]^c|r)
$$

we obtain

(A.7)
$$
G(e, z; [N_{\eta}(P_z)]^c | r) \le G(e, y_2; [N_{\eta}(P_z)]^c | r) \cdot e^{\beta K_1 |z|},
$$

As above, the projection $\pi_{N_{\eta}(P_z)}(e)$ has a distance at most \hat{C}_2 depending on η to y_1 . By [\[DWY22,](#page-50-0) Lemma 2.12], for every $M \geq 0$, there exists η_0 such that for all $\eta \geq \eta_0$

$$
G(e, y_2; [N_{\eta}(P_z)]^c | r) \le e^{-Md(\pi_{N_{\eta}(P_z)}(e), y_2)} \le c_0 e^{-Md(y_2, y_1)} \le c_1 e^{-K_2 M |z|}
$$

for some $c_0 = c_0(\eta) > 0$. Summing over $z \in V$ with $|z| > K|x|$, choosing $M > 0$ so that $MK_2 > \beta K_1 + v + \alpha$, we have by [\(A.7\)](#page-48-0) and [\(A.1\)](#page-45-1) that for $\eta \ge \eta_0$,

$$
\mathbf{E}[\sharp \mathcal{Z}_{31}]\leq \sum_{n\geq K|x|}c_0c_1\cdot \mathrm{e}^{(v+\beta K_1-MK_2)n}\leq c\mathrm{e}^{-\alpha|x|}\leq cG(e,x|r).
$$

where c depends on c_0, c_1 . The lemma is proved.

$$
\Box
$$

We can now finish the proof of Proposition [7.6.](#page-35-1)

Proof of Proposition [7.6.](#page-35-1) Using the thin-triangle property, we see that if z is contained in $U(x) \setminus U_{\epsilon}(x, \kappa \log |x|)$, then $z \in \Omega(x, C)$ for some uniform \tilde{C} .

Fix any $K > 1$. Let ϵ be small enough and κ be large enough such that the conclusions of Lemmas [A.1,](#page-45-2) [A.2](#page-46-0) and [A.3](#page-46-1) hold. Using [\(A.1\)](#page-45-1), the probability that the branching random walk visits $B(x, \kappa \log |x|)$ is bounded by

$$
\sum_{z \in B(x,\kappa \log|x|)} G(e,z|r) \le c e^{v\kappa \log|x|} G(e,x|r) e^{\alpha \kappa \log|x|} \le |x|^\beta G_r(e,x)
$$

for some β depending on α , κ and v .

Now assume that the branching random walk visits $U(x)$ through paths outside $B(x, \kappa \log |x|)$. If the events E_1 and E_3 do not happen, then the event E_2 happens: BRW(Γ, ν, μ) first visits $U(x)$ at a point $z \in U(x) \setminus U_{\epsilon}(x, \kappa \log |x|)$ with $|z| < K |x|$, without entering $B(x, \kappa \log |x|)$. Therefore,

$$
\mathbf{P}(\text{BRW}(\Gamma, \nu, \mu) \text{ visits } \Omega(x, C)) \leq \mathbf{P}(\text{BRW}(\Gamma, \nu, \mu) \text{ visits } U(x))
$$

\n
$$
\leq \mathbf{P}(\text{BRW}(\Gamma, \nu, \mu) \text{ visits } B(x, \kappa \log|x|)) + \mathbf{P}(E_1 \cup E_2 \cup E_3)
$$

\n
$$
\leq C(1+|x|^{\beta})G(e, x|r) \qquad \Box
$$

REFERENCES

- [ACT15] Goulnara N. Arzhantseva, Christopher H. Cashen, and Jing Tao. Growth tight actions. Pacific J. Math., 278(1):1–49, 2015.
- [Ale92] Georgios Alexopoulos. A lower estimate for central probabilities on polycyclic groups. Canadian Journal of Mathematics, 44:897–910, 1992.
- [Ale02] Georgios Alexopoulos. Random walks on discrete groups of polynomial volume growth. Ann. Probab., 30:723–801, 2002.
- [AN04] Krishna Athreya and Peter Ney. Branching processes. Dover Publications, Inc., Mineola, NY, 2004. Reprint of the 1972 original [Springer, New York; MR0373040].
- [Anc88] Alano Ancona. Positive harmonic functions and hyperbolicity. In Potential theorysurveys and problems, pages 1–23. Lecture notes in mathematics, Springer, 1988.
- [Bal97] Keith Ball. An elementary introduction to modern convex geometry. In Flavors of Geometry, volume 31. MSRI publications, 1997.
- [Bas72] Hyman Bass. The degree of polynomial growth of finitely generated nilpotent groups. Proc. London Math. Soc., 25:603–614, 1972.
- [BB07] Sébastien Blachère and Sara Brofferio. Internal diffusion limited aggregation on discrete groups having exponential growth. Probab. Theory Relat. Fields, 137:323–343, 2007.
- [BD13] Emmanuel Breuillard and Enrico Le Donne. On the rate of convergence to the asymptotic cone for nilpotent groups and subFinsler geometry. PNAS, 110:19220–19226, 2013.
- [BHM11] Sébastien Blachère, Peter Haïssinsky, and Pierre Mathieu. Harmonic measures versus quasiconformal measures for hyperbolic groups. Ann. Sci. Ec. Norm. Supér., 44:683–721, 2011.
- [Bil99] Patrick Billingsley. Convergence of probability measure. John Wiley and Sons, second edition, 1999.
- [Bla53] David Blackwell. Extension of a renewal theorem. Pacific J. Math., 33:315–320, 1953.
- [BM12] Itai Benjamini and Sebastian Müller. On the trace of branching random walks. *Groups* Geom. Dyn., 6:231–247, 2012.
- [BNW06] Laurent Bartholdi, Markus Neuhauser, and Wolfgang Woess. Horocyclic products of trees. J. Eur. Math. Soc., 10:771–816, 2006.
- [Bow99] B. H. Bowditch. Convergence groups and configuration spaces. In Geometric group theory down under (Canberra, 1996), pages 23–54. de Gruyter, Berlin, 1999.
- [Bow12] Brian Bowditch. Relatively hyperbolic group. Int. J. Algebra Comput., 22:1–66, 2012. [BP94] Itai Benjamini and Yuval Peres. Markov chains indexed by trees. Ann. Probab., 22(1):219–243, 1994.
- [BSW12] Sara Brofferio, Maura Salvatori, and Wolfgang Woess. Brownian motion and harmonic functions on $Sol(p, q)$. Int. Math. Res., 2012:5182-5218, 2012.
- [BW05] Sara Brofferio and Wolfgang Woess. Green kernel estimates and the full Martin boundary for random walks on lamplighter groups and Diestel–Leader graphs. ANN I H POINCARE-PR, 41:1101–1123, 2005.
- [CGM12] Elisabetta Candellero, Lorenz Gilch, and Sebastian Müller. Branching random walks on free products of groups. Proc. London Math. Soc., 104:1085–1120, 2012.
- [CM17] Matt Clay and Dan Margalit. Office Hours with a Geometric Group Theorist. Princeton University Press, 2017.
- [CT16] Christopher H. Cashen and Jing Tao. Growth tight actions of product groups. Groups Geom. Dyn., 10(2):753–770, 2016.
- [DG21] Matthieu Dussaule and Ilya Gekhtman. Stability phenomena for Martin boundaries of relatively hyperbolic groups. Probab. Theory Relat. Fields, 179:201–259, 2021.
- [DK18] Cornelia Drutu and Michael Kapovich. Geometric Group Theory. American Mathematical Society, 2018.
- [DLM12] Moon Duchin, Samuel Lelièvre, and Christopher Mooney. The geometry of spheres in free abelian groups. Geom. Dedicata, 161:169–187, 2012.
- [DM14] Moon Duchin and Christopher Mooney. Fine asymptotic geometry in the Heisenberg group. Indiana Univ. Math. J., 63:885–916, 2014.
- [DOP00] Françoise Dal'bo, Jean-Pierre Otal, and Marc Peigné. Séries de Poincaré des groupes géométriquement finis. Isr. J. Math., 118:109-124, 2000.
- [DPPS11] Françoise Dal'Bo, Marc Peigné, Jean-Claude Picaud, and Andrea Sambusetti. On the growth of quotients of Kleinian groups. Ergodic Theory Dynam. Systems, 31(3):835– 851, 2011.
- [DWY22] Matthieu Dussaule, Longmin Wang, and Wenyuan Yang. Branching random walks on relatively hyperbolic groups. arXiv:2211.07213, 2022.
- [Ers01] Anna Erschler (Dyubina). On the asymptotics of the rate of departure to infinity. Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Steklov (POMI), 283:251–257, 2001.
- [Ers03] Anna Erschler. On drift and entropy growth for random walks on groups. Ann. Probab., 31:1193–1204, 2003.
- [Fel66] William Feller. An Introduction to Probability Theory and Its Applications, vol. 2. John Wiley and Sons, 1966.
- [FM98] Benson Farb and Lee Mosher. A rigidity theorem for the solvable Baumslag-Solitar groups. Invent. Math., 131:419–451, 1998.
- [GGPY21] Ilya Gekhtman, Victor Gerasimov, Leonid Potyagailo, and Wenyuan Yang. Martin boundary covers Floyd boundary. Invent. Math., 223:759–809, 2021.
- [GL13] Sébastien Gouëzel and Steven Lalley. Random walks on co-compact Fuchsian groups. Ann. Sci. Éc. Norm. Supér., 46:129–173, 2013.
- [GM06] Nina Gantert and Sebastian Müller. The critical branching Markov chain is transient. Markov Process. Related Fields, 12:805–814, 2006.
- [Gou14] Sébastien Gouëzel. Local limit theorem for symmetric random walks in Gromovhyperbolic groups. JAMS, 27:893–928, 2014.
- [GP13] Victor Gerasimov and Leonid Potyagailo. Quasi-isometric maps and Floyd boundaries of relatively hyperbolic groups. JEMS, 15:2115–2137, 2013.
- [GP15] Victor Gerasimov and Leonid Potyagailo. Non-finitely generated relatively hyperbolic groups and Floyd quasiconvexity. Groups, Geom. Dyn., 9:369–434, 2015.
- [GP16] Victor Gerasimov and Leonid Potyagailo. Quasiconvexity in relatively hyperbolic groups. J. Reine Angew. Math., 710:95–135, 2016.
- [Gro81] Mikhael Gromov. Groups of polynomial growth and expanding maps. Publ. Mathématiques, 53:53-73, 1981.
- [Gui73] Yves Guivarc'h. Croissance polynomiale des groupes de Lie et périodes des fonctions harmoniques. Bull. de la Soc. Math. de France, 101:333-379, 1973.
- [KV83] Vadim Kaimanovich and Anatoly Vershik. Random walks on discrete groups: boundary and entropy. Ann. Probab., 11:457–490, 1983.
- [Lal89] Steven Lalley. Renewal theorems in symbolic dynamics, with applications to geodesic flows, noneuclidean tessellations and their fractal limits. Acta Math., 163:1-55, 1989.
- [Led93] François Ledrappier. A renewal theorem for the distance in negative curvature. In Stochastic analysis, volume 57 of Proc. Sympos. Pure Math., pages 351–360. Amer. Math. Soc., 1993.
- [Led01] François Ledrappier. Some asymptotic properties of random walks in free groups. In Topics in Probability and Lie Groups: Boundary Theory, pages 117–152. CRM Proceedings and Lecture Notes, 2001.
- [Mal51] Anatoly Mal'cev. On a class of homogeneous spaces. Amer. Math. Soc. Translation, 39, 1951.
- [Man12] Avinoam Mann. How Groups Grow, volume 395 of London Mathematical Society Lecture Note Series. Cambridge University Press, 2012.
- [Pan83] Pierre Pansu. Croissance des boules et des géodésiques fermées dans les nivariétés. Ergod. Theory Dyn., 3:415–445, 1983.
- [Pat76] S. J. Patterson. The limit set of a Fuchsian group. Acta Math., 136(3-4):241–273, 1976.
- [Pei11] Marc Peign´e. On some exotic Schottky groups. Discrete Contin. Dyn. Syst., 31(2):559– 579, 2011.
- [PW94] Massimo Picardello and Wolfgang Woess. The full Martin boundary of the bi-tree. Ann. Probab., 22:2203–2222, 1994.
- [Rag72] Madabusi Santanam Raghunathan. Discrete subgroups of Lie groups. Springer, 1972. [Rev03a] David Revelle. Heat kernel asymptotics on the lamplighter group. Electron. Commun.
- Probab., 8:142–154, 2003. [Rev03b] David Revelle. Rate of escape of random walks on wreath products and related groups.
- Ann. Probab., 31:1917–1934, 2003.
- [Sha89] Michael Shapiro. A geometric approach to the almost convexity and growth of some nilpotent groups. Math. Ann., 285:601–624, 1989.
- [Spi76] Frank Spitzer. Principles of Random Walks. Springer, 1976.
- [SWX22] Vladas Sidoravicius, Longmin Wang, and Kainan Xiang. Limit set of branching random walks on hyperbolic groups. to appear in Comm. Pure Appl. Math, 2022. https://doi.org/10.1002/cpa.22088.
- [Tan17] Ryokichi Tanaka. Hausdorff spectrum of harmonic measure. Ergodic Theory Dynam. Systems, 37(1):277–307, 2017.
- [Uch98] Kôhei Uchiyama. Green's functions for random walks on \mathbb{Z}^n . Proc. London Math. Soc., 77:215–240, 1998.
- [Var91] Nicholas Theodoros Varapoulos. Groups of superpolynomial growth. In Satoru Igari, editor, ICM-90 Satellite Conference Proceedings, pages 194–200, Tokyo, 1991. Springer Japan.
- [Woe00] Wolfgang Woess. Random Walks on Infinite Graphs and Groups. Cambridge Press University, 2000.
- [Woe13] Wolfgang Woess. What is a horocyclic product, and how is it related to lamplighters? Internat. Math. Nachr., 224:1–27, 2013.
- [Yan14] Wenyuan Yang. Growth tightness for groups with contracting elements. Math. Proc. Cambridge Philos. Soc., 157(2):297–319, 2014.
- [Yan19] Wenyuan Yang. Statistically convex-cocompact actions of groups with contracting elements. Int. Math. Res., 23:7259–7323, 2019.
- [Yan22] Wenyuan Yang. Patterson-Sullivan measures and growth of relatively hyperbolic groups. Peking Math. J., 5:153–212, 2022.

Matthieu Dussaule : Cogitamus Laboratory, France Email address: matthieu.dussaule@hotmail.fr

Longmin Wang : School of Statistics and Data Science, LPMC, Nankai University, 94 Weijin Road, Nankai District, Tianjin, China

Email address: wanglm@nankai.edu.cn

Wenyuan Yang : Beijing International Center for Mathematical Research (BICMR), Beijing University, No. 5 Yiheyuan Road, Haidian District, Beijing, China Email address: yabziz@gmail.com