PySDTest: a Python/Stata Package for Stochastic Dominance Tests

Kyungho Lee^{*} Yoon-Jae Whang[†]

August 7, 2024

Abstract

We introduce PySDTest, a Python/Stata package for statistical tests of stochastic dominance. PySDTest implements various testing procedures such as Barrett and Donald (2003), Linton et al. (2005), Linton et al. (2010), and Donald and Hsu (2016), along with their extensions. Users can flexibly combine several resampling methods and test statistics, including the numerical delta method (Dümbgen, 1993; Hong and Li, 2018; Fang and Santos, 2019). The package allows for testing advanced hypotheses on stochastic dominance relations, such as stochastic maximality among multiple prospects. We first provide an overview of the concepts of stochastic dominance and testing methods. Then, we offer practical guidance for using the package and the Stata command **pysdtest**. We apply PySDTest to investigate the portfolio choice problem between the daily returns of Bitcoin and the S&P 500 index as an empirical illustration. Our findings indicate that the S&P 500 index returns second-order stochastically dominate the Bitcoin returns.

Keywords: pysdtest, stochastic dominance, statistical testing, resampling, bitcoin

^{*}Email: kyungho.lee@yale.edu Department of Economics, Yale University

[†]Email: whang@snu.ac.kr. Department of Economics, Seoul National University. This work was supported by the 2021 Development Fund of the Department of Economics at Seoul National University.

1 Introduction

Stochastic dominance (SD) is an ordering rule of distribution functions, originally suggested by Hadar and Russell (1969) and Hanoch and Levy (1969). It is based on the expected utility paradigm and gives a uniform ordering of prospects that does not depend on a specific preference structure (or utility function) of a decision maker. Because of its generality and nonparametric feature, academics and practitioners in various areas, such as economics, finance, insurance, medicine and statistics, have regarded SD ordering as a fundamental concept of decision-making. They have empirically examined SD relations in diverse areas. Examples include the portfolio choice problem (Post, 2003; Bali et al., 2009), income inequality and welfare analysis (Amin et al., 2003; Anderson, 2003), financial market efficiency (Cho et al., 2007), auction bids (De Silva et al., 2003), firm productivity and export decision (Delgado et al., 2002), obesity inequality (Pak et al., 2016), and agricultural productivity (Langyintuo et al., 2005) among many others. The number of applications has been growing along with recent advances in inference methods for testing SD hypotheses.

There exists a big literature for statistical tests of SD; see Whang (2019) for an extensive survey. McFadden (1989) pioneered statistical tests of SD, proposing a Kolmogorov-Smirnov type test for the first and second order SD. Other early works are Klecan et al. (1991); Kaur et al. (1994); Anderson (1996); Davidson and Duclos (2000). Barrett and Donald (2003, BD) introduce resampling procedures for consistent testing of arbitrary SD order between two distributions under i.i.d. and mutually independent sampling schemes. Linton, Maasoumi and Whang (2005, LMW) propose a consistent subsampling method which is valid under general sampling schemes, including serially dependent observations and mutually dependent prospects. The LMW test also allows for prospects that may depend on some unknown finitedimensional parameters, enabling comparison between residuals after controlling for observed confounders. Linton, Song and Whang (2010, LSW) introduce the *contact set approach* to improve the power performance of SD tests and show large sample uniform validity of the test. Also, they allow the prospects to depend on unknown *infinite* dimensional parameters. Donald and Hsu (2016, DH) provide an alternative method to enhance the performance by suggesting the *selective recentering approach*. However, to our knowledge, comprehensive statistical software for conducting SD tests is not available.

This paper introduces a Python package PySDTest and Stata command pysdtest for testing SD hypotheses, hopefully to help practitioners implement SD tests in a convenient and flexible way. The package implements various SD tests developed in the recent literature. Supported testing procedures include, but are not limited to, the least favorable case approach of BD, subsampling approach of LMW, contact set approach of LSW, and selective recentering method of DH. In addition, we include the numerical delta method (NDM) based on the results of Dümbgen (1993), Fang and Santos (2019), and Hong and Li (2018) in a more general context to compute critical values, as the NDM may have computational advantages in some contexts. This is a novel result since there has been no practical software tailored to SD testing. We provide guidance on how to use our package in Python and Stata environments with practical examples.

PySDTest also allows for flexibly testing a complex and advanced null hypothesis based on SD relations. For example, practitioners can compare more than 2 prospects simultaneously. To be specific, given $K(\geq 3)$ prospects, they can perform a joint test whether there exists

at least one prospect stochastically dominating some of the others, which is called *stochastic non-maximality* among multiple prospects (Klecan et al., 1991). If they reject the hypothesis, then they may confer that the set of prospects is *stochastically maximal*, in the sense that no prospect in the set is dominated by any of the others. This practice is possible because the package offers additional features such as diverse resampling methods and functional type of test statistics. We describe them in detail with an illustrative example.

As an empirical illustration, we apply PySDTest to the portfolio choice problem in the cryptocurrency and stock markets. We investigate the SD relation between the daily returns of Bitcoin price and S&P 500 index. Our testing result shows that the S&P 500 returns second order stochastically dominate the Bitcoin returns. This implies that a risk-averse economic agent with a monotone increasing utility function would prefer S&P 500 index to Bitcoin as a financial asset.

This paper is organized as follows. In Section 2, we introduce the concepts of SD and their testing methods. In Section 3, we provide a full description of the Python package PySDTest and Stata command pysdtest. In Section 4, we apply PySDTest to compare the daily return distributions of Bitcoin price and S&P 500 index. In Section 5, we conclude the paper.

2 Concepts and Tests of Stochastic Dominance

2.1 Concepts of SD

Let X_1 and X_2 be two (continuous) random variables with cumulative distribution functions (CDF) F_1 and F_2 , respectively. Let $Q_k = \inf\{x : F_k(x) \ge \tau\}$ denote the quantile function of X_k for k = 1, 2, respectively. In addition, let \mathcal{U}_1 be the class of all monotone increasing utility (or social welfare) functions. If the functions are differentiable, then \mathcal{U}_1 can be written as:

$$\mathcal{U}_1 = \{u(\cdot) : u' \ge 0\}$$

Definition 2.1.1 (FSD) The random variable X_1 is said to first-order stochastically dominates the random variable X_2 , denoted by $F_1 \succeq_1 F_2$ if any of the following equivalent conditions holds: (1) $F_1(x) \leq F_2(x)$ for all $x \in \mathbb{R}$; (2) $E[u(X_1)] \geq E[u(X_2)]$ for all $u \in \mathcal{U}_1$; and (3) $Q_1(\tau) \geq Q_2(\tau)$ for all $\tau \in [0, 1]$.

This definition can be understood by considering an investor's portfolio choice problem. Let the random variables be returns of some financial assets. The condition $F_1(x) \leq F_2(x)$ for all $x \in \mathbb{R}$ can be denoted as:

$$P(X_1 > x) \ge P(X_2 > x) \text{ for all } x \in \mathbb{R}$$
(1)

which means, for all $x \in \mathbb{R}$, the probability of attaining returns larger than x is larger under F_1 than F_2 . An investor with a monotone increasing utility function would therefore prefer F_1 to F_2 . If the two CDFs intersect, however, FSD does not hold. This means that there exist some u and v both in \mathcal{U}_1 such that $E[u(X_1)] > E[u(X_2)]$, but $E[v(X_1)] < E[v(X_2)]$.

The second-order stochastic dominance can be defined in a similar manner. Let \mathcal{U}_2 be the class of all monotone increasing and concave (utility or social welfare) functions. If functions are differentiable, then we can write the class \mathcal{U}_2 as:

$$\mathcal{U}_2 = \{ u(\cdot) : u' \ge 0, u'' \le 0 \}$$

Definition 2.1.2 (SSD) The random variable X_1 is said to second-order stochastically dominates the random variable X_2 , denoted by $F_1 \succeq_2 F_2$ if any of the following equivalent conditions holds: (1) $\int_{-\infty}^x F_1(t) dx \leq \int_{-\infty}^x F_2(t) dx$ for all $x \in \mathbb{R}$; (2) $E[u(X_1)] \geq E[u(X_2)]$ for all $u \in \mathcal{U}_2$; and (3) $\int_0^\tau Q_1(p) dp \geq \int_0^\tau Q_2(p) dp$ for all $\tau \in [0, 1]$.

Consider the portfolio choice problem of an investor again. The concavity of utility function reflects risk-aversion of the investor. If $X_1 \succeq_2 X_2$, which represents $E[u(X_1)] \ge E[u(X_2)]$ for all $u \in \mathcal{U}_2$, the risk-averse investor would prefer X_1 to X_2 .

Higher-order SD relations also can be defined. Define *integrated CDF* and *integrated quantile function* as (Davidson and Duclos, 2000):

$$F_k^{(s)}(x) = \begin{cases} F_k(x) & \text{for } s = 1\\ \int_{-\infty}^x F_k^{(s-1)}(t) dt & \text{for } s \ge 2 \end{cases}$$
(2)

$$Q_k^{(s)}(x) = \begin{cases} Q_k(x) & \text{for } s = 1\\ \int_0^x Q_k^{(s-1)}(t) dt & \text{for } s \ge 2 \end{cases}$$
(3)

The higher SD order corresponds to the smaller class of utility (or social welfare) functions. That is, for $s \ge 1$, the class of utility functions is defined as:

$$\mathcal{U}_s = \{ u(\cdot) : u' \ge 0, u'' \le 0, ..., (-1)^{(s+1)} u^{(s)} \ge 0 \}.$$

Definition 2.1.3 The random variable X_1 is said to s-order stochastically dominates the random variable X_2 , denoted by $F_1 \succeq_s F_2$ if any of the following equivalent conditions holds: (1) $F_1^{(s)}(x) \leq F_2^{(s)}(x)$ for all $x \in \mathbb{R}$ and $F_1^{(r)}(\infty) \leq F_2^{(r)}(\infty)$ for all r = 1, ..., s - 1; (2) $E[u(X_1)] \geq E[u(X_2)]$ for all $u \in \mathcal{U}_s$; and (3) $Q_1^{(s)}(\tau) \geq Q_2^{(s)}(\tau)$ for all $\tau \in [0, 1]$ and $Q_1^{(r)}(1) \geq Q_2^{(r)}(1)$ for all r = 1, ..., s - 1.

For example, the third-order stochastic dominance (s = 3) reflects preference toward positive skewness of a distribution (Whitmore (1970), Whitmore and Findlay (1978), and Levy (2016)). That is, investors would prefer positively skewed distributions to negatively skewed distributions.

2.2 Tests of SD

2.2.1 Hypothesis of interest

For a given stochastic order s, consider the following hypotheses of interest:

$$H_0^s: F_1^{(s)}(x) \le F_2^{(s)}(x)$$
 for all x v.s. $H_1^s: F_1^{(s)}(x) > F_2^{(s)}(x)$ for some x (4)

The null hypothesis means that random variable X_1 s-th order stochastically dominates X_2 and the alternative hypothesis is the negation of the null hypothesis.¹

2.2.2 BD Test: Least Favorable Case Approach

Let $\{X_{1,i} : i = 1, 2, ..., N_1\}$ and $\{X_{2,i} : i = 1, 2, ..., N_2\}$ be random samples from F_1 and F_2 , respectively. We assume $N_1 = N_2 \equiv N$ for simplification.² Let \mathcal{X} denote the common compact support of X_1 and X_2 .

We first introduce and define frequently used notions in the SD testing literature. The integrated CDF, $F_k^{(s)}(\cdot)$, can be written through integration by parts as (Davidson and Duclos, 2000):

$$F_k^{(s)}(x) \equiv \frac{1}{(s-1)!} \int_0^x (x-t)^{s-1} dF_k(t) = \frac{1}{(s-1)!} E\left[x - X_k\right]_+^{s-1},\tag{5}$$

where $[\cdot]_{+} = \max\{\cdot, 0\}$. The natural empirical counterpart, *empirical integrated CDF*, of (5) is given by

$$\bar{F}_{k}^{(s)}(x) = \frac{1}{N(s-1)!} \sum_{i=1}^{N} \left[x - X_{k,i} \right]_{+}^{s-1}.$$
(6)

Define the difference between integrated CDFs and empirical integrated CDFs as:

$$D_{k,l}^{(s)}(x) = F_k^{(s)}(x) - F_l^{(s)}(x)$$
(7)

$$\bar{D}_{k,l}^{(s)}(x) = \bar{F}_k^{(s)}(x) - \bar{F}_l^{(s)}(x).$$
(8)

BD propose the supremum (or Kolmogorov-Smirnov) type test statistic

$$BD_N = \sqrt{N} \sup_{x \in \mathcal{X}} \left(\bar{D}_{1,2}^{(s)}(x) \right)$$

to test the hypothesis (4). Note that we can write (5) as:

$$\mathcal{J}_s(x, F_k) := F_k^{(s)}(x) = \frac{1}{(s-1)!} \int_0^x (x-t)^{s-1} dF_k(t), \tag{9}$$

where $\mathcal{J}_s(x, F_k)$ is a linear and continuous functional with respect to F_k . Under i.i.d. random sampling and compact support \mathcal{X} , the following weak convergence result holds:

$$\sqrt{N}(\bar{F}_k^{(s)}(\cdot) - F_k^{(s)}(\cdot)) \Rightarrow \mathcal{J}_s(\cdot, \mathcal{B}_k \circ F_k) \text{ for } k = 1, 2$$
(10)

as $N \to \infty$, where \mathcal{B}_1 and \mathcal{B}_2 are independent standard Brownian bridges and \Rightarrow denotes the weak convergence. This result comes from applying Donsker theorem to the case s = 1

¹This type of hypotheses has mainly been considered for statistical inference of the SD relations in the literature. There are two other types of hypotheses that are also considered. The first one is the null hypothesis of non-dominance and the alternative hypothesis of dominance. The other type is the null hypothesis of equivalent distributions and the alternative hypothesis of (strict) stochastic dominance. In this paper, we do not cover these hypotheses. For further discussions, see Section 2.1 of Whang (2019).

²PySDTest allows for the case with different sample sizes $N_1 \neq N_2$.

and continuous mapping theorem for the case $s \ge 2$. The limit process, $\mathcal{J}_s(\cdot, \mathcal{B}_k \circ F_k)$, is a Gaussian process with the covariance function given by

$$C_s(x_1, x_2; F_k) = E\mathcal{J}_s(x_1, \mathcal{B}_k \circ F_k)\mathcal{J}_s(x_2, \mathcal{B}_k \circ F_k).$$

Define

$$\hat{T}_{s}(\cdot) := \sqrt{N} \left[(\bar{F}_{1}^{(s)}(\cdot) - F_{1}^{(s)}(\cdot)) - (\bar{F}_{2}^{(s)}(\cdot) - F_{2}^{(s)}(\cdot)) \right]$$
$$\nu_{s}(\cdot) := \mathcal{J}_{s}(\cdot, \mathcal{B}_{1} \circ F_{1}) - \mathcal{J}_{s}(\cdot, \mathcal{B}_{2} \circ F_{2}).$$

Equation (10) implies

$$\hat{T}_s(\cdot) \Rightarrow \nu_s(\cdot).$$
 (11)

Then, under the null hypothesis, we have the following result:

$$BD_N = \sup_{x \in \mathcal{X}} \left[\hat{T}_s(x) + D_{1,2}^{(s)}(x) \right]$$

$$\leq \sup_{x \in \mathcal{X}} \hat{T}_s(x) \text{ under } H_0$$

$$\Rightarrow \sup_{x \in \mathcal{X}} \nu_s(x) \text{ by (11) and the continuous mapping theorem.}$$

Let $c_{1-\alpha}^{BD}$ is the $(1-\alpha)$ quantile of $\sup_{x\in\mathcal{X}}\nu_s(x)$ for $\alpha\in[0,1]$. BD show the following results:

$$\lim_{N \to \infty} \sup P \left[BD_N \ge c_{1-\alpha}^{BD} \right] \le \alpha \text{ under } H_0^s$$

$$\lim_{N \to \infty} P \left[BD_N \ge c_{1-\alpha}^{BD} \right] = 1 \text{ under } H_1^s$$
(12)

The equality of (12) holds when the least favorable case (LFC), $F_1 = F_2$, holds. Note that the limit distribution $\sup_{x \in \mathcal{X}} \nu_s(x)$ depends on the true CDF, F_k , hence it cannot be tabulated.

BD propose three procedures to compute the critical values. All these methods impose the LFC restriction under the null hypothesis and try to mimic the distribution $\sup_{x \in \mathcal{X}} \nu_s(x)$. The first one is the multiplier method. It first generates two independent random samples $\{U_{k,i} : i = 1, 2, ..., N\}$ from N(0, 1) for k = 1, 2. Then, simulate the process that mimics the F_k -Brownian bridge $\mathcal{B}_k \circ F_k(x)$:

$$\mathcal{B}_{k}^{*} \circ \bar{F}_{k}(x) := \frac{1}{\sqrt{N}} \sum_{i=1}^{N} \left[1(X_{i,k} \le x) - \bar{F}_{k}(x) \right] U_{k,i} \text{ for } k = 1, 2.$$

The multiplier method generates the following simulated process:

$$\mathcal{J}_s(x, \mathcal{B}_k^* \circ \bar{F}_k) = \frac{1}{\sqrt{N}(s-1)!} \sum_{i=1}^N \left[(x - X_{k,i})^{s-1} \mathbb{1} \left[X_{k,i} \le x \right] - \frac{1}{N} \sum_{j=1}^N (x - X_{k,j})^{s-1} \mathbb{1} \left[X_{k,j} \le x \right] \right] U_{k,i}$$

for k = 1, 2. Then, the critical value for the significance level α is defined as the $(1 - \alpha)$ quantile of the following simulated distribution:

$$BD_{N,MP}^* := \sup_{x \in \mathcal{X}} \{ \mathcal{J}_s(x, \mathcal{B}_1^* \circ \bar{F}_1) - \mathcal{J}_s(x, \mathcal{B}_2^* \circ \bar{F}_2) \}.$$
(13)

Note that $BD_{N,MP}^*$ approximates the distribution of $\sup_{x \in \mathcal{X}} \nu_s(x)$ conditional on the original sample.

The second method is the pooled-sample bootstrap method. This method draws bootstrap samples $S_k^* = \{X_{k,i}^* : i = 1, 2, ..., N\}$ for k = 1, 2 from the pooled original sample $\{X_{k,i} : k = 1, 2 \text{ and } i = 1, 2, ..., N\}$ and calculates the empirical CDFs by using the bootstrap samples as:

$$\bar{F}_k^*(x) = \frac{1}{N} \sum_{i=1}^N \mathbb{1}(X_{k,i}^* \le x) \text{ for } k = 1, 2.$$
(14)

Then, we compute the following distribution of the bootstrap test statistic:

$$BD_{N,pool}^{*} := \sup_{x \in \mathcal{X}} \{ \mathcal{J}_{s}(x, \bar{F}_{1}^{*}) - \mathcal{J}_{s}(x, \bar{F}_{2}^{*}) \}.$$
(15)

The critical value for the significance level α is defined to be the $(1 - \alpha)$ quantile of the $BD^*_{N,pool}$ distribution. Note that the LFC restriction is imposed by pooling the original samples.

The third procedure is the recentered bootstrap method. This method draws bootstrap samples S_1^* and S_2^* from $\{X_{1,i} : i = 1, 2, ..., N\}$ and $\{X_{2,i} : i = 1, 2, ..., N\}$, respectively. Then, the empirical CDFs are calculated using each bootstrap sample as in (14). Define

$$BD_{N,recenter}^* := \sup_{x \in \mathcal{X}} \{ (\mathcal{J}_s(x, \bar{F}_1^*) - \mathcal{J}_s(x, \bar{F}_1)) - (\mathcal{J}_s(x, \bar{F}_2^*) - \mathcal{J}_s(x, \bar{F}_2)) \},$$
(16)

where the LFC restriction is imposed by recentering the bootstrap statistic $\mathcal{J}_s(x, \bar{F}_k^*)$ by subtracting its mean $\mathcal{J}_s(x, \bar{F}_k)$ for k = 1, 2. The critical values are defined to be the $(1 - \alpha)$ quantile of the bootstrap distribution $BD_{N,recenter}^*$.

2.2.3 LMW Test: Subsampling Apporach for General Sampling Scheme

For a given set of K prospects, LMW considered the following hypotheses:

$$H_0: \min_{k \neq l} \{D_{k,l}^{(s)}(x)\} \le 0 \text{ for all } x \in \mathcal{X}$$

$$\tag{17}$$

$$H_1: \min_{k \neq l} \{D_{k,l}^{(s)}(x)\} > 0 \text{ for some } x \in \mathcal{X},$$

$$(18)$$

where $D_{k,l}^{(s)}(x)$ is defined as in (7) and \mathcal{X} is the union of the supports of X_k for k = 1, 2, ..., K. The null hypothesis H_0 is the hypothesis of *stochastic non-maximality* (Klecan et al. (1991)). It says that there exists at least one prospect that *s*-th order stochastically dominates some of the others in the set of the *K*-prospects. The alternative hypothesis is that of *stochastic maximality*, that is there does not exist any prospect in the set which *s*-th order stochastic dominates any of the others.

LMW propose a supremum type test for testing (17) with the following features: (i) SD of *arbitrary* order, (ii) Subsampling critical values allowing for *general sampling scheme* such as serial dependence of observations and arbitrary dependence amongst prospects at a specific time, (iii) Prospects may depend on finite dimensional parameters, enabling *conditional ranking*. For instance, the test can be applied to residuals from regressions to control for systematic factors.

Let $X_{k,i}(\theta)$ be the prospect that may depend on a finite dimensional parameter θ for k = 1, 2..., K. If we consider the residual from the linear regression, we may take $X_{k,i}(\theta) =$ $Y_{k,i} - Z_{k,i}^{\dagger} \theta$. Define

$$F_k^{(1)}(x,\theta) := F_k(x,\theta) = P(X_{k,i}(\theta) \le x)$$

$$F_k^{(s)}(x,\theta) := \int_{-\infty}^x F_k^{(s-1)}(t,\theta) dt \text{ for } s \ge 2.$$

Likewise, we define $\bar{F}_k^{(s)}(x,\theta)$ as the sample analogue of $F_k^{(s)}(x,\theta)$ for $s \ge 1$. Let $F_k^{(s)}(x) := F_k^{(s)}(x,\theta_{k,0})$ and $\bar{F}_k^{(s)}(x) := \bar{F}_k^{(s)}(x,\theta_{k,0})$, where $\theta_{k,0}$ denotes the true value, and define

$$d_s := \min_{k \neq l} \sup_{x \in \mathcal{X}} \left[D_{k,l}^{(s)}(x) \right]$$

We can represent the hypotheses (17) and (18) as:

$$H_0: d_s \le 0$$
 v.s. $H_1: d_s > 0$.

LMW use the sample analogue of d_s as the basis of their test statistic:

$$LMW_N = \sqrt{N} \min_{k \neq l} \sup_{x \in \mathcal{X}} \left[\bar{F}_k^{(s)}(x, \hat{\theta}_k) - \bar{F}_l^{(s)}(x, \hat{\theta}_l) \right],$$

where $\hat{\theta}_k$ is a consistent estimator of $\theta_{k,0}$ for k = 1, 2, ..., K. The pointwise asymptotic mull distribution of LMW_N is a functional of a mean zero Gaussian process when $d_s = 0$ (the "boundary" case), and degenerates to $-\infty$ when $d_s < 0$ (the "interior" case) under some regularity conditions. For illustration, let K = 2 and there be no parameter to be estimated.³ Then, the test statistic LMW_N can be denoted as:

$$LMW_N = \sqrt{N} \min\{\sup_{x \in \mathcal{X}} \bar{D}_{1,2}^{(s)}(x), \sup_{x \in \mathcal{X}} \bar{D}_{2,1}^{(s)}(x)\}.$$

To describe the asymptotic null distribution, define the *contact set* to be:

$$\mathcal{C}_0 := \{ x \in \mathcal{X} : D_{1,2}^{(s)}(x) = 0 \}$$
(19)

Then, under the null hypothesis H_0 ,

$$LMW_N \Rightarrow \begin{cases} \min\{\sup_{x \in \mathcal{C}_0} \nu_{1,2}^{(s)}(x), \sup_{x \in \mathcal{C}_0} \nu_{2,1}^{(s)}(x)\} & \text{if } d_s = 0\\ -\infty & \text{if } d_s < 0, \end{cases}$$
(20)

where $\nu_{k,l}^{(s)}(\cdot)$ is a mean zero Gaussian process for k, l = 1, 2 and $k \neq l$.⁴ Under the alternative hypothesis H_1 ($d_s > 0$), $LMW_N \rightarrow_p \infty$ where \rightarrow_p denotes convergence in probability. Note that the asymptotic null distribution depends on the true distribution hence is not pivotal.

³The function pysdtest.test_sd() in the PySDTest package with the 'subsampling' argument implements the routine of the LMW test for a two-sample and no parameter case. However, users can implement the test with multiple prospects and parameters by using the additional features provided by PySDTest.

⁴This result is based the pointwise asymptotics and hence does not fully describe the behavior of the test statistic in finite samples. See Section 2.2.4 for a result based on the uniform asymptotics.

LMW suggest a subsampling procedure to compute the critical values.⁵ This method not only circumvents the non-pivotal issue but also yields consistent critical values under a general sampling scheme. Let b be the size of subsamples, satisfying the condition $b \to \infty$ and $b/N \to 0$ as $N \to \infty$. Let $\alpha \in [0, 1]$ be the nominal significance level. Then the subsampling procedure can be conducted as the following steps:

- 1. Compute LMW_N using the full original sample $\mathcal{S}_N = \{W_1, ..., W_N\}$ where $W_i = \{(Y_{k,i}, Z_{k,i}) : k = 1, 2, ..., K\}$.
- 2. Generate subsamples $S_{i,b} = \{W_i, ..., W_{i+b-1}\}$ for i = 1, 2, ..., N b + 1.
- 3. Compute $\{LMW_{i,b} : i = 1, 2, ..., N b + 1\}$ by using the subsamples.
- 4. Compute the critical value $g_{N,b}(1-\alpha)$ which is defined to be $(1-\alpha)$ quantiles of the distribution $\{LMW_{i,b} : i = 1, 2, ..., N b + 1\}$.
- 5. Reject H_0 if $LMW_N > g_{N,b}(1-\alpha)$.

Under Assumptions 1-4 of LMW, the subsampling test is asymptotically valid in the sense

$$\lim_{N \to \infty} P \left[LMW_N \ge g_{N,b} (1 - \alpha) \right] = \begin{cases} \alpha & \text{if } d_s = 0\\ 0 & \text{if } d_s < 0\\ 1 & \text{if } d_s > 0. \end{cases}$$
(21)

In practice, we need to pick the subsample size b which might affect the test result. Given that there is no compelling theory on the optimal choice of b in the literature, LMW propose three data dependent methods for choosing b: minimum volatility method (Politis et al., 1999), mean critical value and median critical value. Let B_N be a set of subsampling size candidates. Then, the minimum volatility method suggests practitioners select one in B_N that minimizes local variations of critical values. The mean and median critical value methods recommend to pick the mean and median of critical values, respectively. However, LMW also suggest computing and drawing a plot of p-values, which checks whether p-values are insensitive to the choice of b. This is because, if so, inferences would be robust regardless of the specific value of b.

2.2.4 Contact Set Approach

The classical approach imposing the LFC restriction to obtain critical values can be too conservative in practice. This is because it tries to estimate the upper bound of the asymptotic null distributions. To see this, one can establish that under the null hypothesis,

$$BD_N \Rightarrow \sup_{x \in \mathcal{C}_0} \nu_s(x),$$
 (22)

where \mathcal{C}_0 is the *contact set* defined in (19). Clearly, $\sup_{x \in \mathcal{C}_0} \nu_s(x)$ is stochastically dominated by $\sup_{x \in \mathcal{X}} \nu_s(x)$ because $\mathcal{C}_0 \subseteq \mathcal{X}$. Therefore, the asymptotic size of LFC-based tests achieves

 $^{{}^{5}}$ The subsampling method has been proposed by Politis and Romano (1994), see Politis et al. (1999) for a comprehensive review.

the nominal level only when prospects of comparison have equal distributions. This leaves the room for power enhancement.

LSW instead propose the *contact set approach* which estimates the contact set, C_0 , and directly mimics the asymptotic null distribution (22) for inference. This leads to enhanced power property, compared to the LFC-based procedure.

LSW consider an integral-type test for arbitrary SD orders with the hypotheses of interests given by (4). In addition to these features, LSW allow unbounded support and prospects depending on finite and infinite dimensional parameters and establish uniform size validity. Their tests can be applied to serially independent observations yet allow cross-sectional dependence between prospects.

Consider the one-sided Cramér-von Mises type functional:

$$d_s := \int_{\mathcal{X}} \max\{q(x)D_{1,2}^{(s)}(x), 0\}^2 dx,$$

where $q(\cdot)$ is a bounded weight function and $D_{1,2}^{(s)}(\cdot)$ is defined as in (7).⁶ The weight function here is to accommodate unbounded supports, which may entail the integrability issue of $F_k^{(s)}$ for $s \geq 2$. If \mathcal{X} is bounded or s = 1, q(x) = 1 is allowed. Otherwise, LSW suggest

$$q(x) = \begin{cases} 1 & \text{if } x \in [z_1, z_2] \\ a/(a + |x - z_2|^{(s-1)\vee(1+\delta)}) & \text{if } x > z_2 \\ a/(a + |x - z_1|^{(s-1)\vee(1+\delta)}) & \text{if } x < z_1 \end{cases}$$

for $z_1 < z_2$ and $a, \delta > 0$ where $a \lor b$ denotes max $\{a, b\}$. Then the hypotheses of interests (4) can be represented as

$$H_0: d_s = 0$$
 v.s. $H_1: d_s > 0$.

Non/semi-parametric models are allowed with the following specification of prospects:

$$X_k(\theta, \tau) = \varphi_k(W; \theta, \tau)$$
 for $k = 1, 2$

where $W = (Y, Z) \in \mathbb{R}^{d_W}$ is a random vector and $\varphi(\cdot; \theta, \tau)$ is a real-valued function with parameters $(\theta, \tau) \in \Theta \times \mathcal{T}$. Θ is a finite-dimensional Euclidean space and \mathcal{T} is an infinitedimensional space. For example, $X_k := X_k(\theta_0, \tau_0)$ may be specified as the residual from the partially linear regression, i.e. $X_k(\theta, \tau) = Y_k - Z_{k,1}^{\mathsf{T}}\theta - \tau(Z_{k,2})$. Also, we may consider the single-index model such as $X_k = Y_k - \tau(Z_{k,1}^{\mathsf{T}}\theta)$.

Let $X_{k,i}(\theta,\tau) = \varphi_k(W_i;\theta,\tau)$ and $\{W_i: i = 1, 2, ..., N\}$ be a random sample. Define the integrated empirical CDF as

$$\bar{F}_k^{(s)}(x,\theta,\tau) := \frac{1}{N(s-1)!} \sum_{i=1}^N \left[x - X_{k,i}(\theta,\tau) \right]_+^{s-1} \text{ for } k = 1,2$$

$$d_s := \int_{\mathcal{X}} \max\{q(x)D_{1,2}^{(s)}(x), 0\}^2 w(x) dx$$

where w(x) is a nonnegative integrable weight function.

 $^{^{6}}$ LSW also allow practitioners to use a weight function for empirical analysis focusing on a specific part of distributions. For example, one may be interested in comparing income levels below the median of distributions. In this case, the functional would be defined to be

and their differences as

$$\bar{D}_{1,2}^{(s)}(x,\theta,\tau) := \bar{F}_1^{(s)}(x,\theta,\tau) - \bar{F}_2^{(s)}(x,\theta,\tau).$$

LSW propose the integral type test statistic

$$LSW_N = N \int_{\mathcal{X}} \max\left\{q(x)\bar{D}_{1,2}^{(s)}(x,\hat{\theta},\hat{\tau}),0\right\}^2 dx,\tag{23}$$

where $(\hat{\theta}, \hat{\tau})$ is a consistent estimator of (θ, τ) .

For brevity, we describe the testing procedure under the case when there are no estimated parameters, q(x) = 1, and \mathcal{X} is bounded. Thus, we consider the following simplified test statistic

$$LSW_N = N \int_{\mathcal{X}} \max\left\{\bar{D}_{1,2}^{(s)}(x), 0\right\}^2 dx$$

where $\bar{D}_{1,2}^{(s)}(\cdot)$ and $\bar{F}_k^{(s)}(\cdot)$ are defined to be the same as in (8) and (6), respectively. Denote the set of probabilities that satisfy the null hypothesis as \mathcal{P}_0 . Also, let $\mathcal{P}_{00} := \{P \in \mathcal{P}_0 :$ $\lambda(C_0) > 0\}$ where λ is the Lebesgue measure. Put it differently, \mathcal{P}_{00} is the subset of \mathcal{P}_0 with positive Lebesgue measure on the contact set.

We shall describe the asymptotic behavior of LSW_N for a given $P \in \mathcal{P}_0$. With some regularity conditions (Assumptions 1-3 of LSW),

$$\begin{split} LSW_N &= \int_{\mathcal{X}} \max\left\{\sqrt{N} \left[\bar{D}_{1,2}^{(s)}(x) - D_{1,2}^{(s)}(x)\right] + \sqrt{N} D_{1,2}^{(s)}(x), 0\right\}^2 dx \\ &= \int_{\mathcal{C}_0} \max\left\{\sqrt{N} \left[\bar{D}_{1,2}^{(s)}(x) - D_{1,2}^{(s)}(x)\right], 0\right\}^2 dx \\ &+ \int_{\mathcal{X}\setminus\mathcal{C}_0} \max\left\{\sqrt{N} \left[\bar{D}_{1,2}^{(s)}(x) - D_{1,2}^{(s)}(x)\right] + \sqrt{N} D_{1,2}^{(s)}(x), 0\right\}^2 dx \\ &= \int_{\mathcal{C}_0} \max\left\{\sqrt{N} \left[\bar{D}_{1,2}^{(s)}(x) - D_{1,2}^{(s)}(x)\right], 0\right\}^2 dx + o_p(1) \\ &\Rightarrow \int_{\mathcal{C}_0} \max\{\nu_s(x), 0\}^2 dx, \end{split}$$

where $\nu_s(\cdot)$ is a mean zero Gaussian process. The second equality comes from the definition of the contact set. The third equality holds because $\sqrt{N}D_{1,2}^{(s)}(x) \to -\infty$ on $\mathcal{X} \setminus \mathcal{C}_0$ under the null hypothesis. Lastly, the weak convergence follows from $\sqrt{N}\left[\bar{D}_{1,2}^{(s)}(\cdot) - D_{1,2}^{(s)}(\cdot)\right] \Rightarrow \nu_s(\cdot)$ and the continuous mapping theorem. On the other hand, for $P \in \mathcal{P}_0 \setminus \mathcal{P}_{00}, \lambda(\mathcal{C}_0) = 0$ gives us $LSW_N \Rightarrow 0$. Therefore, we can summarize the asymptotic results under the null hypothesis as

$$LSW_N \Rightarrow \begin{cases} \int_{\mathcal{C}_0} \max\{\nu_s(x), 0\}^2 dx & \text{if } P \in \mathcal{P}_{00} \\ 0 & \text{if } P \in \mathcal{P}_0 \setminus \mathcal{P}_{00} \end{cases}$$

LSW call \mathcal{P}_{00} as the set of *boundary* points and $\mathcal{P}_0 \setminus \mathcal{P}_{00}$ as the set of *interior* points. The notable thing is that there is a discontinuity in the asymptotic null distributions, depending on the true probability. Since there is no discontinuity in finite samples, the above discontinuity indicates that the *uniform* asymptotics, rather than *pointwise* asymptotics, would provide better approximation and complete asymptotic behaviors.

Let α be the nominal significance level and E_P be the expectation under the probability P. Then, the definitions are

Definition 2.2.1 (a) A test ρ_{α} has an asymptotically (uniformly) valid size if

$$\limsup_{N \to \infty} \sup_{P \in \mathcal{P}_0} E_P \rho_\alpha \le \alpha \tag{24}$$

(b) A test ρ_{α} has an asymptotically exact size if it satisfies (24) and there exists a non-empty set $\mathcal{P}'_0 \subset \mathcal{P}_0$ such that

$$\limsup_{N \to \infty} \sup_{P \in \mathcal{P}'_0} |E_P \rho_\alpha - \alpha| = 0 \tag{25}$$

(c) A test ρ_{α} is called asymptotically similar on \mathcal{P}'_0 if it satisfies (25).

The notable thing is that LFC-based testing procedures might be asymptotically nonsimilar hence biased in large samples against a large set of alternatives. The bootstrap procedure proposed by LSW is asymptotically similar on a larger subset of \mathcal{P}_0 in which the LFC is a special case. Therefore, the procedure is asymptotically biased against a smaller class of alternatives than LFC-based tests and hence is more powerful.

Let B denote the number of bootstrap repetitions and α be the nominal significance level. The procedure can be described as follows:

- 1. Compute LSW_N using the original sample $\{W_1, ..., W_N\}$, where $W_i = \{(Y_{k,i}, Z_{k,i}) : k = 1, 2\}$.
- 2. Draw $\{W_{i,b}^*: i = 1, ..., N\}$ with replacement from $\{W_i: i = 1, ..., N\}$.
- 3. Compute estimates θ_b^* and τ_b^* using $\{W_{i,b}^* : i = 1, ..., N\}$.
- 4. Compute $X_{k,i,b}^* = \varphi_k(W_{i,b}^*; \theta_b^*, \tau_b^*)$ for k = 1, 2.
- 5. Compute

$$\bar{D}_{1,2,b}^{(s),*}(x) = \frac{1}{N} \sum_{i=1}^{N} \left[h_x(X_{1,i,b}^*) - h_x(X_{2,i,b}^*) - \frac{1}{N} \sum_{i=1}^{N} \{ h_x(\hat{X}_{1,i}) - h_x(\hat{X}_{2,i}) \} \right],$$

where $\hat{X}_{k,i} = \varphi_k(W_i; \hat{\theta}, \hat{\tau})$ for $k = 1, 2$ and $h_x(z) := \frac{(x-z)^{(s-1)} 1\{z \le x\}q(x)}{(s-1)!}$.

6. Take a sequence c_N such that $c_N \to 0$ and $c_N \sqrt{N} \to \infty$ and estimate the contact set as

$$\hat{\mathcal{C}} = \{ x \in \mathcal{X} : q(x) | \bar{D}_{1,2}(x) | < c_N \}.$$

7. Compute

$$LSW_{N,b}^{*} = \begin{cases} \int_{\hat{\mathcal{C}}} \max\{q(x)\sqrt{N}\bar{D}_{1,2,b}^{(s),*}(x), 0\}^{2}dx, & \text{if } \lambda(\hat{\mathcal{C}}) > 0\\ \int_{\mathcal{X}} \max\{q(x)\sqrt{N}\bar{D}_{1,2,b}^{(s),*}(x), 0\}^{2}dx, & \text{if } \lambda(\hat{\mathcal{C}}) = 0 \end{cases}$$

8. Repeat step 2-7 for b = 1, 2, ..., B and compute the critical value

$$c_{\alpha,N,B}^* := \inf\{B^{-1}\sum_{b=1}^B \mathbb{1}\{LSW_{N,b}^* \le t\} \ge 1 - \alpha\}.$$

Note that $W_i = (X_{1,i}, X_{2,i})^{\intercal}$ and step 3-4 should be omitted if there are no estimated parameters.

We shall introduce some asymptotic properties of the contact set approach. Suppose that Assumptions 1-4 of LSW hold. Under H_0 ,

$$\limsup_{N \to \infty} \sup_{P \in \tilde{\mathcal{P}}_0} P(LSW_N > c^*_{\alpha, N, \infty}) \le \alpha$$
(26)

for some $\tilde{\mathcal{P}}_0 \subset \mathcal{P}_0$ where $c^*_{\alpha,N,\infty}$ is the critical value from the ideal bootstrap $(B = \infty)$. In addition, for some $\tilde{\mathcal{P}}_{00,N} \subset \tilde{\mathcal{P}}_0$,

$$\limsup_{N \to \infty} \sup_{P \in \tilde{\mathcal{P}}_{00,N}} |P(LSW_N > c^*_{\alpha,N,\infty}) - \alpha| = 0.$$
(27)

Equation (26) shows that the test has asymptotically valid size uniformly over a class of "regular" probabilities $\tilde{\mathcal{P}}_0 \subset \mathcal{P}_0$. In addition, equation (27) indicates that test has asymptotically exact size on $\tilde{\mathcal{P}}_{00,N} \subset \tilde{\mathcal{P}}_0$. On the other hand, the test is consistent against a fixed alternative. That is, for fixed alternative $P \notin \mathcal{P}_0$ such that $\int_{\mathcal{X}} \max\{q(x)D_{1,2}(x), 0\}^2 dx > 0$,

$$\lim_{N \to \infty} P(LSW_N > c^*_{\alpha, N, \infty}) = 1$$

In addition, Theorem 4 of LSW shows that the contact set approach is locally more powerful than the LFC-based tests.

2.2.5 Selective Recentering Approach

DH propose the selective recentering approach to enhance the power performance of BD tests. The main idea is similar to the contact set approach: DH try to directly approximate the limit distribution of BD_N , $\sup_{x \in C_0} \nu_s(x)$ defined in (22), to attain the critical values. For this end, DH extend the recentering method introduced by Hansen (2005). It is also closely related to the generalized moment selection approach of Andrews and Shi (2013) in the moment inequality literature.

DH also consider the hypotheses of interest given by (4) and they consider the Barrett-Donald test statistic BD_N . They consider a sampling scheme similar to that of BD, but further discuss extension of their method to deal with weakly dependent data and mutual dependence between X_1 and X_2 .

Let us briefly describe the selective recentering approach. Let

$$\mu(x) := \min\{D_{1,2}^{(s)}(x), 0\}.$$
(28)

Then, we have that the asymptotic null distribution of

$$BD_N^0 = \sup_{x \in \mathcal{X}} \left[\sqrt{N} \left(\bar{D}_{1,2}^{(s)}(x) - D_{1,2}^{(s)}(x) \right) + \sqrt{N} \mu(x) \right]$$
(29)

is the same as that of BD_N because $\mu(x) = D_{1,2}^{(s)}(x)$ under the null. DH propose a resampling procedure that tries to mimic (29) using the sample analogue of (28). That is, first take a

sequence a_N of negative numbers such that $a_N \to -\infty$ and $a_N/\sqrt{N} \to 0$ and approximate (28) via the recentering function:

$$\hat{\mu}_N(x) := \bar{D}_{1,2}^{(s)} \cdot 1\left\{\bar{D}_{1,2}^{(s)} < a_N/\sqrt{N}\right\}$$

Under regularity conditions, it is a uniformly consistent estimator in the sense that

$$\sup_{x \in \mathcal{X}} |\hat{\mu}_N(x) - \mu(x)| \to_p 0$$

Write the simulated processes defined in (13), (15), and (16) as

$$\nu_{s,MP}^{*} = \mathcal{J}_{s}(x, \mathcal{B}_{1}^{*} \circ \bar{F}_{1}) - \mathcal{J}_{s}(x, \mathcal{B}_{2}^{*} \circ \bar{F}_{2}),$$

$$\nu_{s,pool}^{*} = \mathcal{J}_{s}(x, \bar{F}_{1}^{*}) - \mathcal{J}_{s}(x, \bar{F}_{2}^{*}),$$

$$\nu_{s,recenter}^{*} = (\mathcal{J}_{s}(x, \bar{F}_{1}^{*}) - \mathcal{J}_{s}(x, \bar{F}_{1})) - (\mathcal{J}_{s}(x, \bar{F}_{2}^{*}) - \mathcal{J}_{s}(x, \bar{F}_{2})).$$

To compute the critical values, DH suggest using the bootstrap distributions

$$DH_{N,k}^* = \sup_{x \in \mathcal{X}} \{\nu_{s,k}^*(x) + \sqrt{N}\hat{\mu}_N(x)\} \text{ for } k = \{MP, pool, recenter\}.$$
(30)

Letting $\hat{c}_{\alpha,k}$ be the $(1-\alpha)$ quantile of $DH^*_{N,k}$, DH suggest the following critical value:

$$\hat{c}_{\alpha,\eta,k} := \max\{\hat{c}_{\alpha,k},\eta\},\$$

where η is a small positive number such as 10^{-6} . η is introduced here to accommodate the situation where both the original test statistic and the bootstrap test statistic degenerate to zero, such as the interior case $D_{1,2}^{(2)}(x) < 0$ for all $x \in \mathcal{X}$.

In practice, two tuning parameters, a_N and η , need to be selected. Based on the simulation experiments conducted by DH, the specific choice of η is not very important as long as η is a small positive number. However, a_N should be chosen carefully because the choice of a_N may affect the finite sample performance of the test. DH suggest using $a_N = -0.1\sqrt{\log \log(N)}$ which performed well in their simulations and empirical studies.

As regard to the asymptotic properties, DH show that under the null hypothesis,

$$\lim_{N \to \infty} P(BD_N > \hat{c}_{\alpha,\eta,k}) = 0 \text{ if } D_{1,2}^{(s)}(x) < 0 \text{ for all } x \in \mathcal{X}$$
$$\lim_{N \to \infty} P(BD_N > \hat{c}_{\alpha,\eta,k}) = \alpha \text{ if } D_{1,2}^{(s)}(x) = 0 \text{ for all } x \in \mathcal{X}$$

For a fixed alternative, the test is shown to be consistent:

$$\lim_{N \to \infty} P(BD_N > \hat{c}_{\alpha,\eta,k}) = 1.$$

2.2.6 Numerical Delta Method

As an alternative method for attaining critical values, we can apply an inference method applicable to directionally differentiable functions. Fang and Santos (2019) show the asymptotic validity of an inference method that employs a consistent estimate of the first order

directional derivative when the target parameters are Hadamard directionally differentiable. Complementing their idea, Hong and Li (2018) propose the method using numerical derivatives for approximating the limit distributions which eases the burden of analytical calculation.⁷ Even though Fang and Santos (2019) discuss the testing for SD as an application of their methods, they do not provide a formal result let alone the software for the SD tests.

We briefly discuss the NDM, which also has enhanced power properties compared to the classical LFC-based tests. The power enhancement is possible because the NDM also directly approximates the asymptotic null distribution, as in the contact set approach and selective recentering approach. Theoretical validity of applying the NDM to SD tests is discussed by Lee et al. (2023). To be specific, Lee et al. (2023) introduce the test for time stochastic dominance (time SD) with application of the NDM and the static SD concept is regarded as a special case of the time SD concept with a single time period.

To describe testing for SD by the NDM, we first introduce the definition of *Hadamard* directionally differentiability of a map between Banach spaces (Shapiro, 1990).

Definition 2.2.2 Let \mathbb{D} and \mathbb{E} be Banach spaces and $\phi : \mathbb{D}_{\phi} \subset \mathbb{D} \to \mathbb{E}$. The map ϕ is said to be Hadamard directionally differentiable at $\theta \in \mathbb{D}_{\phi}$ tangentially to a set $\mathbb{D}_0 \subset \mathbb{D}$ if there exists a continuous map $\phi'_{\theta} : \mathbb{D}_0 \to \mathbb{E}$ such that

$$\lim_{N \to \infty} \left\| \frac{\phi(\theta + t_N h_N) - \phi(\theta)}{t_N} - \phi'_{\theta}(h) \right\|_{\mathbb{E}} = 0$$

for all sequences $\{h_N\} \subset \mathbb{D}$ and $\{t_N\} \subset \mathbb{R}_+$ such that $t_N \downarrow 0$, $h_N \to h \in \mathbb{D}_0$ as $N \to \infty$ and $\theta + t_N h_N \in \mathbb{D}_{\phi}$ for all N.

We say $\phi'_{\theta}(h)$ is (first-order) Hadamard directional derivative at θ in direction h. In addition, if ϕ'_{θ} is linear, then ϕ is said to be Hadamard differentiable at $\theta \in \mathbb{D}_{\phi}$ tangentially to a set $\mathbb{D}_0 \subset \mathbb{D}$.

Theorem 2.1 of Fang and Santos (2019) shows that if $r_N(\hat{\theta}_N - \theta_0) \Rightarrow \mathbb{G}_0$ in \mathbb{D} for some $r_N \to \infty$ where \mathbb{G}_0 is tight, then $r_N\left[\phi(\hat{\theta}_N) - \phi(\theta_0)\right] \Rightarrow \phi'_{\theta_0}(\mathbb{G}_0)$. On top of this result, Theorem 3.2 of Fang and Santos (2019) shows that $\phi'_{\theta_0}(\mathbb{G}_0)$ can be approximated by $\phi'_N(\mathbb{Z}_N^*)$ where \mathbb{Z}_N^* and ϕ'_N are consistent estimators of \mathbb{G}_0 and ϕ'_{θ_0} , respectively. Hong and Li (2018) suggest the NDM which numerically approximates $\phi'_{\theta_0}(\mathbb{G}_0)$ hence does not need analytical calculation of ϕ'_{θ_0} . Hong and Li (2018) suggest the estimation of ϕ'_{θ_0} by using the following numerical derivatives

$$\tilde{\phi}_N'(\mathbb{Z}_N^*) := \frac{\phi(\hat{\theta}_N + \epsilon_N \mathbb{Z}_N^*) - \phi(\hat{\theta}_N)}{\epsilon_N}$$
(31)

where ϵ_N is a step size satisfying $\epsilon_N \to 0$ and $r_N \epsilon_N \to \infty$ as $N \to \infty$.

The NDM can be applied to the SD testing problem. Consider the testing with the L_1 -type test statistic and our hypothesis of interest is the same as (4). First, denote

$$l^{\infty}(\mathcal{X}) := \{ f : \mathcal{X} \to \mathbb{R} \text{ such that } \|f\|_{\infty} < \infty \} \text{ and } \|f\|_{\infty} = \sup_{x \in \mathcal{X}} f(x).$$

⁷Both Fang and Santos (2019) and Hong and Li (2018) are extensions of the pioneering work of Dümbgen (1993) that studies a validity of the bootstrap under a lack of differentiability.

For any $\theta \in l^{\infty}(\mathcal{X}) \times l^{\infty}(\mathcal{X}), \phi : l^{\infty}(\mathcal{X}) \times l^{\infty}(\mathcal{X}) \to \mathbb{R}$ is defined as

$$\phi(\theta) := \int_{\mathcal{X}} \max\{\theta_1(x) - \theta_2(x), 0\} dx.$$

We then may write

$$\phi(\theta_0) = \int_{\mathcal{X}} \max\{F_1^{(s)}(x) - F_2^{(s)}(x), 0\} dx,$$

where $\theta_0 = (F_1^{(s)}(x), F_2^{(s)}(x)) \in l^{\infty}(\mathcal{X}) \times l^{\infty}(\mathcal{X})$. It can be seen that $\phi(\cdot)$ is Hadamard directionally differentiable with its derivative given as

$$\phi_{\theta}'(h) = \int_{\mathcal{C}_{+}(\theta)} (h_{1}(x) - h_{2}(x)) dx + \int_{\mathcal{C}_{0}(\theta)} \max\{h_{1}(x) - h_{2}(x), 0\} dx,$$

where

$$\mathcal{C}_{+}(\theta) := \{ x \in \mathcal{X} : \theta_{1}(x) > \theta_{2}(x) \}$$
$$\mathcal{C}_{0}(\theta) := \{ x \in \mathcal{X} : \theta_{1}(x) = \theta_{2}(x) \}.$$

Note that $C_0(\theta_0)$ is equivalent to the contact set C_0 in (19). In addition, we have $C_+(\theta_0) = \emptyset$ under the null hypothesis.

Letting $\hat{\theta}_N = (\bar{F}_1^{(s)}(\cdot), \bar{F}_2^{(s)}(\cdot))$, we may write the L_1 -type test statistic as $r_N \phi(\hat{\theta}_N)$, where $r_N = \sqrt{N}$. In addition, we know that $r_N(\bar{F}_k^{(s)}(\cdot) - F_k^{(s)}(\cdot))$ weakly converges to a tight Gaussian process. If we let $\mathbb{Z}_N^* = r_N(\bar{F}_k^{(s),*}(\cdot) - \bar{F}_k^{(s)}(\cdot))$, where $\bar{F}_k^{(s),*}$ is the bootstrap sample analogue of $\bar{F}_k^{(s)}$, then we can approximate the limit distribution of $r_N \phi(\hat{\theta}_N)$ by employing (31).

There are some cases in which the first-order derivatives may degenerate, such as the the L_2 -type test statistic. In this case, we can employ the second-order NDM (Hong and Li, 2018). $\phi''_{\theta_0}(h)$ is said to be the second order Hadamard directional derivative at $\theta_0 \in \mathbb{D}_{\phi}$ in the direction h if

$$\lim_{N \to \infty} \left\| \frac{\phi(\theta_0 + t_N h_N) - \phi(\theta_0) - t_N \phi'_{\theta_0}(h_N)}{\frac{1}{2} t_N^2} - \phi''_{\theta_0}(h) \right\|_{\mathbb{E}} = 0$$

for all sequences $\{h_N\} \subset \mathbb{D}$ and $\{t_N\} \subset \mathbb{R}_+$ such that $t_N \downarrow 0$, $h_N \to h \in \mathbb{D}_0$ as $N \to \infty$ and $\theta + t_N h_N \in \mathbb{D}_{\phi}$ for all N. Theorem 4.1 of Hong and Li (2018) shows that if $r_N(\hat{\theta}_n - \theta_0) \Rightarrow \mathbb{G}_0$ for some tight process \mathbb{G}_0 , then

$$r_N^2 \left[\phi(\hat{\theta}_N) - \phi(\theta_0) - \phi'_{\theta_0}(\hat{\theta}_N - \theta_0) \right] \Rightarrow \frac{1}{2} \phi''_{\theta_0}(\mathbb{G}_0)$$

When the first-order derivatives degenerate, $\phi'_{\theta_0}(h) = 0$, it is possible to approximate $\frac{1}{2}\phi''_{\theta_0}(\mathbb{G}_0)$ by

$$\frac{1}{2}\tilde{\phi}_{\theta_0}''(\mathbb{Z}_N^*) := \begin{cases} \frac{\phi(\hat{\theta}_N + \epsilon_N \mathbb{Z}_N^*) - \phi(\hat{\theta}_N)}{\epsilon_N^2} \text{ or} \\ \frac{\phi(\hat{\theta}_N + 2\epsilon_N \mathbb{Z}_N^*) - 2\phi(\hat{\theta}_N - \epsilon_N \mathbb{Z}_N^*) + \phi(\hat{\theta}_N)}{2\epsilon_N^2} \end{cases}$$
(32)

where ϵ_N satisfies $\epsilon_N \to 0$ and $r_N \epsilon_N \to \infty$ as $N \to \infty$. In the case of the L₂-type test statistic, which is the simplified version of LSW_N in (23), we may take

$$\phi(\theta) := \int_{\mathcal{X}} \max\{\theta_1(x) - \theta_2(x), 0\}^2 dx$$

$$\phi(\theta_0) = \int_{\mathcal{X}} \max\{F_1^{(s)}(x) - F_2^{(s)}(x), 0\}^2 dx$$

Then we have $\phi'_{\theta_0}(h) = 0$ under the null hypothesis but

$$\phi_{\theta_0}''(h) = \int_{\mathcal{C}_0(\theta_0)} \max\{h_1(x) - h_2(x), 0\}^2 dx$$

holds. Thus, similarly to the first-order NDM case, we can approximate the limit distribution of $r_N^2 \phi(\hat{\theta}_N)$ using (32).

3 The Package: PySDTest

3.1 Installation

It is required to install the package, PySDTest, for both Python and Stata environments. The package is listed on the Python Package Index (PyPI) which is a repository of softwares for the Python programming language. If Python is installed on your computer, a user can install PySDTest by entering:

pip install PySDTest

in Windows cmd or Mac (or Linux) terminal.⁸ For details about installing a Python package, see Python Package User Guide in PyPI.

The package is written based on Python numpy module with version $\geq 1.26.4$. Therfore, it is also required to install numpy for employing the package.

To install the Stata module, pysdtest, users can use the following command in Stata:

net install pysdtest, from("https://raw.githubusercontent.com/lee-kyungho/ pysdtest/main/Stata") replace

3.2 Implementation in Python

This section describes how to use the package PySDTest in the Python environment. Notably, Stata with version ≥ 16.0 supports running the package within Stata. For more details on using Python in Stata, please refer to the Python integration feature in Stata.

First, users need to import the package through the following code in the Python environment in order to use the package:

>>> import pysdtest

 $^{^{8} \}rm Alternatively,$ users can access source codes of the package through https://github.com/lee-kyungho/pysdtest

PySDTest implements comprehensive SD tests developed in the literature. In addition, users can freely choose procedures suitable to their objects of study. In this section, we describe the details of how to implement each testing procedure.

We first generate simulated data as:

```
>>> import numpy as np
>>> mu1 , mu2 , sigma1 , sigma2 = 0, 0 , 1, 1
>>> n = 500
>>> np.random.seed(0)
>>> sample1 = mu1 + sigma1 * np.random.randn(n)
>>> sample2 = mu2 + sigma2 * np.random.randn(n)
```

3.2.1 BD and LMW tests by pysdtest.test_sd()

The function pysdtest.test_sd() enables the user to implement the tests proposed by BD and LMW. The structure of the function pysdtest.test_sd is given as:

pysdtest.test_sd(sample1, sample2, ngrid, s, resampling, b1 = None, b2 = None, nboot = 200, quiet = False)

- sample1 and sample2 are 1-dimensional NumPy arrays.
- ngrid determines the number of grid points for computing the value on the support.⁹
- **s** argument specifies the stochastic order of the hypothesis.
- resampling argument defines the resampling method:
 - bootstrap: Implements the recentered bootstrap method as described in BD.
 - subsampling: Follows the method suggested by LMW, requiring subsampling sizes b_1 and b_2 for each sample.
 - paired_bootstrap: Uses a paired recentered bootstrap (X_i, Y_i) to consider dependencies.
- nboot: Sets the number of bootstrapping iterations. The default value is 200 for both bootstrap and paired_bootstrap
- quiet: Specifies whether to print output results or not.

The following lines of codes implement the testing procedure by BD:

```
>>> testing_normal = pysdtest.test_sd(sample1, sample2, ngrid = 100,
s = 1, resampling = 'bootstrap')
>>> testing_normal.testing()
```

⁹Increasing ngrid enhances precision but raises computational costs.

#--- Testing for Stochastic Dominance ----# * H0 : sample1 first order SD sample2 #-----# *** Test Setting *** * Resampling method = bootstrap * SD order = 1 * # of (sample1) = 500 * # of (sample2) = 500 * # of bootstrapping = 200 * # of grid points = 100 #-----# *** Test Result *** * Test statistic
* Significance level
-= 0.2214 = 0.05 * Critical-value = 1.2017 * P-value = 0.8600 * Time elapsed = 0.54 Sec

Users can run the subsampling procedure by giving subsampling as an input for the resampling argument:

```
>>> testing_normal_sub = pysdtest.test_sd(sample1, sample2, ngrid = 100,
s = 1, resampling = 'subsampling', b1=40, b2=40)
>>> testing_normal_sub.testing()
#--- Testing for Stochastic Dominance ----#
* H0 : sample1 first order SD sample2
#-----#
*** Test Setting ***
* Resampling method
                     = subsampling
* SD order
                      =
                             1
* # of (sample1)
                      =
                           500
* # of (sample2)
                      = 500
* # of (subsample1)
                           40
                      =
* # of (subsample2)
                    =
                           40
```

#-----#

**	** Test Result ***		
*	Test statistic	=	0.2214
*	Significance level	=	0.20
*	Critical-value	=	0.8944
*	P-value	=	0.8482
*	Time elapsed	=	0.11 Sec

3.2.2 Contact Set Approach by pysdtest.test_sd_contact()

The function pysdtest.test_sd_contact() in PySDTest implements the contact set approach proposed by LSW.¹⁰ The structure of the function is written as

```
pysdtest.test_sd_contact(sample1, sample2, ngrid, s, resampling,
nboot=200, c=0.75, quiet = False)
```

Arguments sample1, sample2, ngrid, s, resampling, b1, b2, nboot, and quiet are the same as pysdtest.test_sd().

We need to take a sequence c_N for the contact set estimation in the step 6 of LSW. Following the recommendation by LSW, we set

$$c_N = \frac{c \log(\log(N))}{\sqrt{N}}$$

where c is some positive constant. If two sample sizes are different, $N = (N_1 + N_2)/2$ holds. The argument c designates the constant c. We set the default value of c to be 0.75.

Users can implement the LSW test by:

```
>>> testing_normal_contact = pysdtest.test_sd_contact(sample1, sample2,
ngrid = 100, s = 1, resampling = 'bootstrap')
>>> testing normal contact.testing()
#--- Testing for Stochastic Dominance ----#
* H0 : sample1 first order SD sample2
* Contact Set Approach
#-----#
*** Test Setting ***
* Resampling method
                       = bootstrap
* SD order
                        =
                              1
* # of (sample1)
                        =
                            500
```

¹⁰To be specific, the case when there is no estimated parameters and q(x) = 1 for all x is implemented by pysdtest.test_sd_contact().

```
=
* # of (sample2)
                        500
* # of bootstrapping
                   =
                        200
* # of grid points
                 =
                        100
# Tuning parameter ------
* с
                   = 0.7500
#-----#
*** Test Result ***
* Test statistic
                  = 0.1495
* Significance level
                  = 0.05
* Critical-value = 18.9272
* P-value
                  = 0.8850
* Time elapsed
                  = 0.52 Sec
```

3.2.3 Selective Recentering Approach by pysdtest.test_sd_SR()

The selective recentering approach of DH with k = recenter (Equation (30)) can be implemented by pysdtest.test_sd_SR(). The structure of the function is

```
pysdtest.test_sd_SR(sample1, sample2, ngrid, s, resampling, nboot=200,
a=0.1, eta=1e-06, quiet = False)
```

Arguments sample1, sample2, ngrid, s, resampling, b1, b2, nboot, and quiet are the same as pysdtest.test_sd().

The selective recentering approach additionally requires two tuning parameters: a_N and η . We follow the recommendation by DH and set a_N to be

$$a_N = -a\sqrt{\log(\log(N))},$$

where a is some positive constant and $N = N_1 + N_2$. Arguments **a** and **eta** set the values of a and η with the default given by a = 0.1 and $\eta = 10^{-6}$.

```
>>> testing_normal_SR = pysdtest.test_sd_SR(sample1, sample2,
ngrid = 100, s = 1, resampling = 'bootstrap')
>>> testing_normal_SR.testing()
#--- Testing for Stochastic Dominance -----#
* H0 : sample1 first order SD sample2
* Selective Recentering Approach
#------#
*** Test Setting ***
```

```
* Resampling method
                          bootstrap
                    =
* SD order
                              1
                        =
* # of (sample1)
                             500
                        =
* # of (sample2)
                             500
                        =
* # of bootstrapping
                            200
                        =
* # of grid points
                        =
                             100
# Tuning paremeters ------
* a
                        = 0.1000
                        = 0.00001
* eta
#-----
                 -----#
*** Test Result ***
* Test statistic
                        = 0.2214
* Significance level
                      = 0.05
* Critical-value
                       = 1.0768
* P-value
                        = 0.8200
* Time elapsed
                        = 0.52 Sec
```

3.2.4 Numerical Delta Method by pysdtest.test_sd_NDM()

Users can approximate the limit distribution based on the NDM approach using the function pysdtest.test_sd_NDM(). The function has the following structure:

pysdtest.test_sd_NDM(sample1, sample2, ngrid, s, resampling, nboot = 200, epsilon = None, form = "L1", quiet = False)

Three different types of functional are available for users: Kolmogorov-Smirnov (supremum), L_1 , and L_2 type functionals. This can be set by giving input 'KS', 'L1', and 'L2' for the 'form' argument, respectively. We apply the second-order NDM for L_2 type testing and the limit distribution is approximated by $\frac{\phi(\hat{\theta}_N + \epsilon_N \mathbb{Z}_N^*) - \phi(\hat{\theta}_N)}{\epsilon_N^2}$.

In addition, the NDM requires users to specify the step size (ϵ_N) as a tuning parameter. The user can set the step size by the argument 'epsilon'. We set $\epsilon = r_N^{-1/16}$ as a default value when nothing is specified by the user. The user can implement the NDM for SD tests as

```
>>> testing_normal_NDM_KS = pysdtest.test_sd_NDM(sample1, sample2, ngrid = 100,
s = 1, resampling = 'bootstrap', form = 'KS')
>>> testing_normal_NDM_KS.testing()
#--- Testing for Stochastic Dominance -----#
* H0 : sample1 first order SD sample2
* Numerical Delta Method
* KS Type Test Statistic
```

#-----#

```
*** Test Setting ***
* Resampling method
                 = bootstrap
* SD order
                           1
                     =
* # of (sample1)
                          500
                     =
* # of (sample2)
                         500
                     =
* # of bootstrapping
                         200
                     =
* # of grid points
                         100
                     =
# Tuning paremeter ------
* epsilon
                     = 0.8415
#-----#
*** Test Result ***
* Test statistic
                    = 3.1939
* Significance level = 0.05
* Critical-value
                    = 0.9805
* P-value
                     = 0.0000
* Time elapsed
                     = 0.51 Sec
```

Table 1 contains common arguments for all testing procedures: BD, LMW, LSW, DH, and NDM. In addition, we summarize additional arguments for each estimation in Table 2.

Argument	Description	Data type
sample1	X_1	(1-dim) numpy array
sample2	X_2	(1-dim) numpy array
S	Stochastic dominance order	int
ngrid	Number of grid points	int
resampling	Resampling methods (bootstrap, subsampling, paired_bootstrap)	str
nboot	Number of bootstrapping	int
b1	Size of subsampling for sample1	int
b2	Size of subsampling for sample2	int
quiet	whether to print output results	boolean

Ta	ble	1:	Common	Input	Arguments	for	Testing
----	-----	----	--------	-------	-----------	-----	---------

Tal	ble 2	: Ad	lditional	Arguments	for	each	Procee	lure
-----	-------	------	-----------	-----------	-----	------	--------	------

Argument	Description	Method	Data type
с	Tuning parameter for a sequence c_N	Contact set approach (LSW)	float
a	Tuning parameter for a sequence a_N	Selective recentering (DH)	float
eta	Tuning parameter η for attaining a critical value	Selective recentering (DH)	float
epsilon	Step size ϵ_N for numerical approximation	Numerical delta method (NDM)	float
form	Type of functional (KS, $L1$, and $L2$)	Numerical delta method (NDM)	str

3.2.5 Additional Features

Every implementation of testing through PySDTest saves results as an object hence users can save the test results. For example, if the user runs the following code

>>> results = testing_normal.result

then the 'results' in the above code is the variable in which the Python dictionary of the test result is allocated. The dictionary contains calculated test statistics, resampled test statistics, critical-value and p-value.

In addition, every function for testing provides a feature for plotting integrated empirical CDFs. This functionality may help users to visually compare testing objects. The plotting is based on the Python Matplotlib package. The method plot_CDF() has the following structure

```
plot_CDF(save=False, title=None, label1='sample1', label2='sample2', xlabel='x')
```

In table 3, arguments for plot_CDF() are listed.

Argument	Description	Data type
save	Whether to save figure (True or False)	boolean
title	Title of saving image file	str
label1	Name of sample1 for legend	str
label2	Name of sample2 for legend	str
xlabel	Name of x-axis	str

Table 3: Input Arguments for plot_CDF()

For instance, plotting can be conducted by

```
>>> testing_normal.plot_CDF(save=True, title="Normal distributions.png")
```

and we show the corresponding result in Figure 1. Plotting a higher-order CDF can be conducted by

```
>>> testing_normal_2 = pysdtest.test_sd(sample1, sample2,
ngrid = 100, s = 2, resampling = 'bootstrap')
>>> testing_normal_2.plot_CDF(save=True, title="Normal distributions2.png")
```

Figure 2 shows the corresponding result.

PySDTest provides three resampling procedures: bootstrap, paired bootstrap, and subsampling. The subsampling() function generates and returns subsamples $S_{i,b}$. The argument b means the length of block, and nsub reflects the number of resampled test statistics. In testing codes, we set nsub as N - b + 1.

pytestsd.subsampling(sample, b, nsub)

The **bootstrap()** function generates and returns bootstrap samples. The argument b means the number of observations for one boostrapping, which is used to calculate test statistics. In testing codes, we set b = N. The nboot means the number of bootstrapping.





Figure 2: (Second order) Plotting Result of $plot_D()$



pytestsd.bootstrap(sample, b, nboot)

The paired_bootstrap() function conducts bootstrapping in a pair $(X_{1,i}, X_{2,i})$, and the description of arguments is the same as bootstrap().

```
pytestsd.paired_bootstrap(sample1, sample2, b, nboot)
```

PySDTest provides two functions that help users calculate the test statistics by themselves. The set_grid() function returns the 'ngrid' number of grid points which are equally divided from the minimum to the maximum value of samples. For instance,

```
>>> # Concatenate samples
>>> samples = [sample1, sample2]
>>> # Set grid
>>> pytestsd.set_grid(samples, ngrid)
```

The CDF() calculates the (integrated) empirical CDF of the sample. For example,

```
>>> pytestsd.CDF(sample1, grid, s = 1)
```

3.2.6 Advanced Hypothesis Testing

Through PySDTest, the user can perform advanced hypothesis regarding SD. For example, consider the case when there are K distributions. Then, the hypothesis of our interest can be $H_0^{(s)}$: There exists at least one distribution that s-th order stochastically dominates some of the other distributions. $H_1^{(s)}$ is the negation of $H_0^{(s)}$. This can be expressed as

$$H_0^{(s)} : \min_{k \neq l} \left\{ D_{k,l}^{(s)}(x) \right\} \le 0 \text{ for all } x \in \mathcal{X}$$

$$(33)$$

$$H_1^{(s)}: \min_{k \neq l} \left\{ D_{k,l}^{(s)}(x) \right\} > 0 \text{ for some } x \in \mathcal{X}$$

$$(34)$$

We describe testing procedures step-by-step and conduct testing via PySDTest. Before testing, we generate random samples.

```
>>> # Parameters
>>> mu1, mu2, mu3 = 0, 0.5, 1
>>> sigma1, sigma2, sigma3 = 1, 1.5, 2
>>> # Sample size
>>> N = 1000
>>>
>>> # set the seed
>>> np.random.seed(0)
>>>
>>> # Random numbers from a normal distribution
>>> sample1 = mu1 + sigma1 * np.random.randn(N)
>>> sample2 = mu2 + sigma2 * np.random.randn(N)
>>> sample3 = mu3 + sigma3 * np.random.randn(N)
```

First, we need to set options for the testing; the user should specify the SD order, the number of grid points, and grid. In this example, we test first order SD and set the number of grid points as 100.

```
>>> " Step 0: Setting Options "
>>> # SD order
>>> s = 1
>>> # Setting grid
>>> samples = [sample1, sample2, sample3]
>>> ngrid = 100
>>> grid = pysdtest.set_grid(samples, ngrid)
>>> # Setting the Number of Bootstrapping
>>> nsamp = 200
```

Then, the user has to compute test statistics. This can be done by using the CDF() function in PySDTest. We consider the supremum-type test statistics, T_N , which may be written as

$$T_N = \sqrt{N} \min_{k \neq l} \left\{ \sup_{x \in \mathcal{X}} D_{k,l}^{(s)}(x) \right\} \text{ for all } k, l \in \{1, 2, ..., K\}$$

For simplicity, we let K = 3. Then, the sample analogue of the test statistics can be computed as:

Then we conduct resampling. In here, we generate bootstrap samples as

```
>>> " Step 2: Generate Bootstrap Sample (or Subsample) "
>>> # Resampling (bootstrap)
>>> btspsample1 = pysdtest.bootstrap(sample1, b = N, nboot = nsamp)
>>> btspsample2 = pysdtest.bootstrap(sample2, b = N, nboot = nsamp)
>>> btspsample3 = pysdtest.bootstrap(sample3, b = N, nboot = nsamp)
```

The next step is to calculate bootstrap test statistics and attain bootstrap distribution. We conduct recentered bootstrap as

```
>>> " Step 3: Compute Test Statistics by Bootstrap Sample (or subsample) "
>>> # Calculating D by bootstrap samples
>>> D_b_12 = CDF(btspsample1, grid, s) - CDF(btspsample2, grid, s)
>>> D_b_21 = CDF(btspsample2, grid, s) - CDF(btspsample1, grid, s)
>>> D_b_23 = CDF(btspsample2, grid, s) - CDF(btspsample3, grid, s)
>>> D_b_32 = CDF(btspsample3, grid, s) - CDF(btspsample2, grid, s)
>>> D_b_13 = CDF(btspsample1, grid, s) - CDF(btspsample3, grid, s)
>>> D_b_31 = CDF(btspsample3, grid, s) - CDF(btspsample1, grid, s)
>>> b_b_31 = CDF(btspsample3, grid, s) - CDF(btspsample1, grid, s)
>>> # Calculating Test Stat by bootstrap samples
>>> # Calculating Test Stat by bootstrap samples
>>> D_b_recentered = np.array(D_b_collection) - np.array(D_collection)
>>> resampled_stat = np.sqrt(N) * np.min(np.max(D_b_recentered,1),0)
```

Then we compute the critical value under the significance level α . Let $\alpha = 0.05$, and

```
>>> " Step 4: Compute the Critical Value of the Bootstrap Distribution "
>>> # Critical value and P-value (alpha = 0.05)
>>> alpha = 0.05
>>> critical_value = np.quantile(resampled_stat, 1 - alpha)
>>> pval = (resampled_stat >= test_stat).mean(0)
```

Lastly, we decide whether to reject the null hypothesis or not. If the computed test statistics is larger than the critical value, reject $H_0^{(s)}$.

```
>>> " Step 5: Reject H0 if Test stat > Critical Value "
>>> print("Test Statistic: %6.3f" % test_stat)
>>> print("Critical value: %6.3f" % critical_value)
>>> print("P-value: %6.3f" % pval)
Test Statistic: 0.791
Critical value: 0.506
P-value: 0.000
```

3.3 Stata Command: pysdtest

In this section, we introduce the Stata command pysdtest. The command is based on the python package PySDTest. Therefore, Stata with version ≥ 16.0 , Python 3 software and installation of the Python package are necessary for using the command. Supported testing procedures include, but are not limited to, tests through the least favorable case approach of BD, subsampling approach of LMW, contact set approach of LSW and selective recentering method of DH. In addition, the NDM can be used to compute the critical value. pysdtest

also supports the combination of various resampling methods, test statistics and procedures for approximating the limiting distribution.

Our hypothesis of interest is (4); for given stochastic order s, the null hypothesis H_0^s is that X_1 (Sample 1) s-th order stochastic dominates X_2 (Sample 2) and the alternative hypothesis H_1^s is the negation of H_0^s . pysdtest allows arbitrary s-th SD order for testing.

3.3.1 Syntax and Options

pysdtest varlist(max = 2) [if] [, options]

varlist specifies up to two variables for comparison. If only one variable is specified, a by() option must be used to divide the sample. The command supports several options to customize the testing procedure:

options	Description
by(varname)	Specify a binary variable for dividing the sample. Without this, two variables are required for testing.
<u>sw</u> itch	Whether to switch the order of the sample when using by().
resampling(<i>string</i>)	Resampling method (<i>bootstrap</i> , <i>subsampling</i> , or <i>paired_bootstrap</i>). Default is <i>bootstrap</i> . For <i>subsampling</i> , specify b1() and b2().
approach(<i>string</i>) Define a testing method. Default is the Kolmogorov-Smirnov type statist BD and LMW. Alternatives include <i>contact</i> , <i>SR</i> (selective recentering), and (numerical delta method).	
<u>functional(string)</u>	For NDM , specify a functional type (KS, L1, L2); default is L1.
ngrid(#)	Number of grid points for test statistics; default is 100.
$\mathbf{s}(integer)$	Stochastic order of the null hypothesis; default is 1.
b1(integer)	Number of subsamples for the first variable.
b2(integer)	Number of subsamples for the second variable.
nboot(integer)	Number of bootstrapping iterations.
c(#)	Tuning parameter for the contact set approach; default is 0.75.
a(#)	Tuning parameter for the selective recentering approach; default is 0.1.
eta(#)	Tuning parameter for the selective recentering approach; default is 10^{-6} .
epsilon(#)	Tuning parameter for the numerical delta method; default is 0.75.
alpha(#)	Significance level for the statistical test; default is 0.05.

3.3.2 Stored Results

The following scalars, macros and matrices are stored as a result of running pysdtest: Scalars

	r(N1)	# of observations of the first variable	r(N2)	# of observations of the second variable
	r(b1)	# of subsamples of the first variable	r(b2)	# of subsamples of the second variable
	r(s)	stochastic order	r(alpha)	significance level
	r(ngrid)	# of grid points	r(test_stat)	value of the test statistic
	r(p_val)	p-value	r(critic_val)	critical value
Mac	ros			
	r(resampling)	resampling method	r(approach)	testing approach
	r(form)	type of functional for the NDM		
Mat	rices			
	r(grid)	matrix of grid values	r(limit_dist)	matrix of the limit distribution of test statistics approximated by resampling

3.3.3 Examples

In this section, we describe example usages of **pysdtest** in Stata. We first generate the simulated data from the standard normal distribution.

```
* Simulate the data
* Clear the existing dataset
. clear
.
* Set the number of observations
. set obs 100
Number of observations (_N) was 0, now 100.
.
* Set the seed
. set seed 2024
.
* Generate a normally distributed variable
. gen X1 = rnormal(0, 1)
. gen X2 = rnormal(0, 1)
```

If there is no option given, **pysdtest** implements Kolmogorov-Smirnov type test statistics as in BD and LMW. The default resampling method is bootstrap and recentering as in BD.

```
. * Run pysdtest
. pysdtest X1 X2
Running PySDTest
Sample1: X1
Sample2: X2
#--- Testing for Stochastic Dominance -----#
* HO : sample1 first order SD sample2
#-----#
*** Test Setting ***
* Resampling method
                    = bootstrap
                   = 1
* SD order
* # of (sample1)
* # of (sample2)
                    =
                         100
                         100
* # of bootstrapping = 200
* # of grid points = 100
#-----#
*** Test Result ***
* Test statistic
                   = 0.2121
```

* Significance level	= 0.05
* Critical-value	= 1.0607
* P-value	= 0.8600
* Time elapsed	= 0.13 Sec

The subsampling method as in LMW can be used by giving "subsampling" as an argument to the option resampling(). In this case, the user needs to set the number of subsamples by b1() and b2() options.

```
. * Run pysdtest with subsampling
. pysdtest X1 X2, resampling("subsampling") b1(40) b2(40)
Running PySDTest
Sample1: X1
Sample2: X2
#--- Testing for Stochastic Dominance -----#
* H0 : sample1 first order SD sample2
#-----#
*** Test Setting ***
*** lest setting ***

* Resampling method = subsampling

* SD order = 1
* SD order = 1
* # of (sample1) = 100
* # of (sample2) = 100
* # of (subsample1) = 40
* # of (subsample2) = 40
#-----#
*** Test Result ***

    Test statistic

                          = 0.2121
* Test stat1st1c - 0.2221
* Significance level = 0.05
* Critical-value = 1.0062
* P-value = 0.9180
* Time elapsed = 0.02 Sec
```

In addition, users can employ the paired bootstrap method by giving "*paired_bootstrap*" as an argument to the option **resampling()**. The number of observations for both sample1 and sample2 must be the same.

Also, the contact set approach can be utilized by giving "contact" to approach() option. The additional tuning parameter can be chosen by c() option. For the contact set approach, the integral type test statistic is employed as in (23).

```
. * Run pysdtest with paired bootstrapping and contact set approach
. pysdtest X1 X2, resampling("paired_bootstrap") approach("contact") c(0.5)
Running PySDTest
Sample1: X1
Sample2: X2
#--- Testing for Stochastic Dominance -----#
* HO : sample1 first order SD sample2
* Contact Set Approach
#_____#
*** Test Setting ***
* Resampling method = paired_bootstrap
* SD order = 1
* # of (sample1) = 100
* # of (sample2) = 100
* # of bootstrapping = 200
* # of grid points = 100
# Tuning parameter -----
* c
                     = 0.5000
#-----#
```

```
*** Test Result ***
* Test statistic = 0.3800
* Significance level = 0.05
* Critical-value = 22.5270
* P-value = 0.8400
* Time elapsed = 0.13 Sec
```

pysdtest provides **by()** option to compare distributions divided by a binary variable. As an illustration, we use automobile data stored in Stata and compare price distributions by the foreign dummy.

```
sysuse auto, clear
(1978 automobile data)
       foreign_str = "Domestic" if foreign == 0
. gen
(22 missing values generated)
. replace foreign_str = "Foreign" if foreign == 1
(22 real changes made)
. * Run pysdtest with by( ) option
. pysdtest price, by(foreign_str)
Running PySDTest
Groups:
"Domestic" `"Foreign"
Sample1: price of Foreign
Sample2: price of Domestic
#--- Testing for Stochastic Dominance -----#
* HO : sample1 first order SD sample2
#-----#
*** Test Setting ***
*** lest Setting ***
* Resampling method = bootstrap
* SD order = 1
* # of (sample1) = 22
* # of (sample2) = 52
* # of prid points = 100
#-----#
*** Test Result ***
                      = 0.3025
* Test statistic
* Significance level = 0.05
* Critical-value = 1.1215
* P-value = 0.7650
* Time elapsed = 0.05 Sec
```

If users want to switch the order of samples, they can use the option switch to swap the order of samples.

```
. * Run pysdtest with by( ) option and switch
. pysdtest price, by(foreign_str) switch
Running PySDTest
Groups:
    "Domestic"´ "Foreign"´
Sample1: price of Domestic
Sample2: price of Foreign
#---- Testing for Stochastic Dominance -----#
* H0 : sample1 first order SD sample2
#-----#
*** Test Setting ***
* Resampling method = bootstrap
* SD order = 1
```

<pre>* # of (sample1)</pre>	= 52					
* # of (sample2)	= 22					
<pre>* # of bootstrapping</pre>	= 200					
<pre>* # of grid points</pre>	= 100					
#	#					
*** Test Result ***						
* Test statistic = 0.8867						
* Significance level = 0.05						
* Critical-value = 1.0184						
* P-value	= 0.1150					
* Time elapsed	= 0.05 Sec					

NDM can be used by giving "NDM" to approach() option. We employ Kolmogorov-Smirnov type test statistic in this example. We can also set the number of grid points by ngrid() option.

```
. * Run pysdtest with by( ) option, NDM
. pysdtest price, by(foreign_str) switch ///
> approach("NDM") functional("KS") ngrid(200)
Running PySDTest
Groups:
"Domestic" `"Foreign"
Sample1: price of Domestic
Sample2: price of Foreign
#--- Testing for Stochastic Dominance -----#
* HO : sample1 first order SD sample2
* Numerical Delta Method
* KS Type Test Statistic
#-----#
*** Test Setting ***
* Resampling method
                      = bootstrap
* SD order
                      =
                            1
* # of (sample1)
                      =
                            52
* # of (sample2)
* # of (sampiez,
* # of bootstrapping =
_______
                           22
                           200
                           200
# Tuning paremeter -----
          = 0.9180
* epsilon
  -----#
*** Test Result ***
* Test statistic
                      = 0.9623
* Significance level = 0.05
                      = 1.0114
* Critical-value
* P-value
                      = 0.0600
                      = 0.06 Sec
* Time elapsed
```

4 Empirical Illustration

In this section, we compare historical daily returns of Bitcoin price and S&P 500 index and investigate SD relations between them. This paper applies PySDTest to illustrate how portfolio comparison based on SD criteria can be done. SD concept is useful for portfolio comparison because SD provides a general rule for the portfolio selection under certain expected utility paradigm. We can interpret testing results in terms of preference to prospects such as monotonicity and risk aversion of an economic agent. The replication code is shown in Appendix.

In this paper, daily returns, R_t^k , are defined as:

$$R_t^k = \log(\frac{P_t^k}{P_{t-1}^k}) \text{ for } t = 1, ..., N \text{ and } k \in \{\text{Bitcoin}, \text{ S\&P 500}\}$$
(35)

where P_t^k is price of Bitcoin or S&P 500 index at day t.¹¹ Table 4 summarizes the time length, number of observations and descriptive statistics of daily return data. Because the cryptocurrency market operates in weekends and holidays, the number of observations of Bitcoin price is more than S&P 500 index.

Name	Time	Observations	Mean	Std	Min	Max
Bitcoin S&P 500	Apr.29.2013 - Feb.27.2021 Apr.29.2013 - Feb.27.2021	$2861 \\ 1995$	$0.0020 \\ 0.0004$	$0.0426 \\ 0.0109$	-0.4647 -0.1277	$0.3575 \\ 0.0897$

Table 4: Descriptive Statistics of Bitcoin price and S&P 500

Standard mean-variance analysis cannot determine superiority of daily returns between Bitcoin and S&P 500 index. Although the mean return of Bitcoin price is larger than that of S&P 500 index, standard deviation of Bitcoin is also larger. In addition, the distribution of Bitcoin price seems to be more dispersed than that of S&P 500 index. Figure 3 shows cumulative distribution functions of daily return of Bitcoin and S&P 500 index.¹²

We conduct testing for the following null hypotheses to investigate SD relations:

$$H_{0,1}^s: R^{\text{Bitcoin}} \succeq_s R^{\text{S\&P 500}}$$
(36)

$$H_{0,2}^s: R^{\text{S\&P 500}} \succeq_s R^{\text{Bitcoin}}$$
(37)

for s = 1, 2 The alternative hypotheses $H_{1,i}^s$ are the negation of $H_{0,i}^s$ for i, s = 1, 2, respectively. We use the subsampling method by LMW in order to account for time-series dependence. We set 100 grid points. 1,000 and 900 are set as a size of subsamples for daily returns of Bitcoin and S&P 500 index, respectively. Table 5 summarizes the testing results.

H_0	$R^{\text{Bitcoin}} \geq$	$\leq_s R^{\text{S&P 500}}$	$R^{S\&P 500}$	$\succeq_s R^{\text{Bitcoin}}$
s	1	2	1	2
p-value	0.0000	0.0000	0.0000	0.3349

Table 5:	Testing	Result	s
----------	---------	--------	---

Notes. We use the subsampling method and set 100 grid points. 1,000 and 900 subsamples are set for daily return of Bitcoin and S&P 500 index, respectively.

In Table 5, all hypotheses of the first SD order (s = 1) are rejected. This means that an economic agent with monotone increasing utility function strictly prefer neither Bitcoin nor

¹¹We use closing price of Bitcoin and S&P 500. Because the cryptocurrency market does not actually 'close', we alternatively use price at 11:59 p.m. provided by the CoinMarketCap

¹²This is plotted by using the function $plot_CDF()$ in PySDTest





Notes This figure shows cumulative distribution functions of daily returns of Bitcoin price and S&P 500 index. The red dash line indicates the distribution function of daily return of Bitcoin. The blue solid line indicates that of S&P 500 index.

S&P 500 to the other. In contrast, it is shown that daily returns of S&P 500 second order (s = 2) stochastically dominate those of Bitcoin. This implies that an agent with risk-averse utility function prefers investing in S&P 500 index to Bitcoin.

5 Conclusion

In this paper, we have introduced a Python package, PySDTest, and a Stata command, **pysdtest**. The package implements various Stochastic Dominance (SD) tests, including those proposed by BD, LMW, LSW, and DH. PySDTest includes testing using the NDM, which applies modern developments in statistical inference methods to directionally differentiable functions. We provide guidance for using PySDTest and **pysdtest**, including testing for extended notions of the SD hypothesis. We applied PySDTest to compare the daily return distributions between Bitcoin prices and the S&P 500 index. The results show that the S&P 500 second-order stochastically dominates Bitcoin returns.

Empirical implementation of the SD concept has not been as extensive as it could be despite the fundamental role of the SD concept in diverse research areas and the recent advances in inference procedures. This may be partly due to the lack of publicly available software. In this paper, we provide a comprehensive Python package and Stata command for SD tests to assist empirical practitioners and academics. We look forward to seeing many applications of our software that help practitioners as intended.

References

- Amin, Sajeda, Ashok S Rai, and Giorgio Topa, "Does microcredit reach the poor and vulnerable? Evidence from northern Bangladesh," *Journal of development Economics*, 2003, 70 (1), 59–82.
- Anderson, Gordon, "Nonparametric tests of stochastic dominance in income distributions," *Econometrica: Journal of the Econometric Society*, 1996, pp. 1183–1193.
- _ , "Poverty in America 1970–1990: Who did gain ground? An application of stochastic dominance criteria employing simultaneous inequality tests in a partial panel," *Journal of Applied Econometrics*, 2003, 18 (6), 621–640.
- Andrews, Donald WK and Xiaoxia Shi, "Inference based on conditional moment inequalities," *Econometrica*, 2013, 81 (2), 609–666.
- Bali, Turan G, K Ozgur Demirtas, Haim Levy, and Avner Wolf, "Bonds versus stocks: Investors' age and risk taking," *Journal of Monetary Economics*, 2009, 56 (6), 817–830.
- Barrett, Garry F and Stephen G Donald, "Consistent tests for stochastic dominance," *Econometrica*, 2003, 71 (1), 71–104.
- Cho, Young-Hyun, Oliver Linton, and Yoon-Jae Whang, "Are there Monday effects in stock returns: A stochastic dominance approach," *Journal of Empirical Finance*, 2007, 14 (5), 736–755.
- **Davidson, Russell and Jean-Yves Duclos**, "Statistical inference for stochastic dominance and for the measurement of poverty and inequality," *Econometrica*, 2000, 68 (6), 1435–1464.
- **Delgado, Miguel A, Jose C Farinas, and Sonia Ruano**, "Firm productivity and export markets: a non-parametric approach," *Journal of international Economics*, 2002, 57 (2), 397–422.
- **Donald, Stephen G and Yu-Chin Hsu**, "Improving the power of tests of stochastic dominance," *Econometric Reviews*, 2016, *35* (4), 553–585.
- Dümbgen, Lutz, "On nondifferentiable functions and the bootstrap," *Probability Theory* and Related Fields, 1993, 95 (1), 125–140.
- Fang, Zheng and Andres Santos, "Inference on directionally differentiable functions," The Review of Economic Studies, 2019, 86 (1), 377–412.
- Hadar, Josef and William R Russell, "Rules for ordering uncertain prospects," The American economic review, 1969, 59 (1), 25–34.
- Hanoch, G. and H. Levy, "The Efficiency Analysis of Choices Involving Risk," *The Review of Econonic Studies*, 1969, *36* (3), 335–346.

- Hansen, Peter Reinhard, "A test for superior predictive ability," Journal of Business & Economic Statistics, 2005, 23 (4), 365–380.
- Hong, Han and Jessie Li, "The numerical delta method," *Journal of Econometrics*, 2018, 206 (2), 379–394.
- Kaur, Amarjot, BLS Prakasa Rao, and Harshinder Singh, "Testing for second-order stochastic dominance of two distributions," *Econometric theory*, 1994, 10 (5), 849–866.
- Klecan, Lindsey, Raymond McFadden, and Daniel McFadden, "A robust test for stochastic dominance," Technical Report, working paper, Department of Economics, MIT 1991.
- Langyintuo, Augustine S, Emmanuel K Yiridoe, Wilson Dogbe, and James Lowenberg-Deboer, "Yield and income risk-efficiency analysis of alternative systems for rice production in the Guinea Savannah of Northern Ghana," Agricultural economics, 2005, 32 (2), 141–150.
- Lee, Kyungho, Oliver Linton, and Yoon-Jae Whang, "Testing for time stochastic dominance," Journal of Econometrics, 2023, 235 (2), 352–371.
- Levy, Haim, Stochastic dominance: Investment decision making under uncertainty, Springer, 2016.
- Linton, Oliver, Esfandiar Maasoumi, and Yoon-Jae Whang, "Consistent testing for stochastic dominance under general sampling schemes," *The Review of Economic Studies*, 2005, 72 (3), 735–765.
- _ , Kyungchul Song, and Yoon-Jae Whang, "An improved bootstrap test of stochastic dominance," Journal of Econometrics, 2010, 154 (2), 186–202.
- McFadden, Daniel, "Testing for stochastic dominance," in "Studies in the Economics of Uncertainty," Springer, 1989, pp. 113–134.
- Pak, Tae-Young, Susana Ferreira, and Gregory Colson, "Measuring and tracking obesity inequality in the United States: evidence from NHANES, 1971-2014," *Population Health Metrics*, 2016, 14 (1), 1–13.
- Politis, Dimitris N and Joseph P Romano, "The stationary bootstrap," Journal of the American Statistical association, 1994, 89 (428), 1303–1313.
- _ , _ , and Michael Wolf, Subsampling, Springer Science & Business Media, 1999.
- Post, Thierry, "Empirical tests for stochastic dominance efficiency," The Journal of Finance, 2003, 58 (5), 1905–1931.
- Shapiro, Alexander, "On concepts of directional differentiability," Journal of optimization theory and applications, 1990, 66 (3), 477–487.

- Silva, Dakshina G De, Timothy Dunne, and Georgia Kosmopoulou, "An empirical analysis of entrant and incumbent bidding in road construction auctions," *The Journal of Industrial Economics*, 2003, 51 (3), 295–316.
- Whang, Yoon-Jae, Econometric analysis of stochastic dominance: Concepts, methods, tools, and applications, Cambridge University Press, 2019.
- Whitmore, George A, "Third-degree stochastic dominance," The American Economic Review, 1970, 60 (3), 457–459.
- **and M Chapman Findlay**, Stochastic dominance: an approach to decision-making under risk, Lexington Books, 1978.

A. Appendix

5.1 Replication Code for Section 4.

5.1.1 Replication by Python codes

```
1 """ import modules """
2
3 import pandas as pd
4 import numpy as np
5 import pysdtest
6 import matplotlib.pyplot as plt
8 """ Descriptive Analysis """
9
10 BTC_daily_rr = pd.read_excel("BTC_daily_rr.xlsx")
11 sp_daily_rr = pd.read_excel("sp_daily_rr.xlsx")
12
13 BTC_daily_rr.describe()
14 sp_daily_rr.describe()
15
16 # Datatype for tests should 1-dim numpy array
17 BTC_daily_rr = np.array(BTC_daily_rr['d_ln_Close'])
18 sp_daily_rr = np.array(sp_daily_rr['d_ln_Close'])
19
  """ Plotting """
20
21
22 # Plotting D
23 crypto_testing_ct_1 = pysdtest.test_sd(BTC_daily_rr, sp_daily_rr, ngrid =
     100, s= 1, b1 = 1000, b2 = 900, resampling = 'subsampling')
24 crypto_testing_ct_1.plot_CDF(save=True, title="CDFs of Bitcoin and S&P
     Index.png",label1="Bitcoin",label2="S&P 500", xlabel = "Daily Return")
25
26 """ Testing HO: BTC >= S&P500 """
27
28 # First order
29
30 crypto_testing_1 = pysdtest.test_sd(BTC_daily_rr, sp_daily_rr, ngrid =
     100, s= 1, b1 = 1000, b2 = 900, resampling = 'subsampling')
31 crypto_testing_1.testing()
32
33 # Second order
34
35 crypto_testing_2 = pysdtest.test_sd(BTC_daily_rr, sp_daily_rr, ngrid =
     100, s= 2, b1 = 1000, b2 = 900, resampling = 'subsampling')
36 crypto_testing_2.testing()
37
  """ Switching samples
38
39
40 Testing HO : S&P500 >= BTC
41
42 пп
43 # First order
```

5.1.2 Replication by the Stata Command

```
. clear
. use bitcoin_sp500_daily_rr, replace
.
. * Specify the python environment
. set python_exec /opt/anaconda3/bin/python
. python query // Check the python version
```

```
Python Settings
     set python_exec
                       /opt/anaconda3/bin/python
     set python_userpath
   Python system information
     initialized no
     version
                       3.11.8
     architecture
                      64-bit
                       /opt/anaconda3/lib/libpython3.11.dylib
     library path
. * Run
. pysdtest BTC_daily_rr SP500_daily_rr, ///
> resampling("subsampling") s(1) b1(1000) b2(900)
Running PySDTest
Sample1: BTC_daily_rr
Sample2: SP500_daily_rr
#--- Testing for Stochastic Dominance -----#
* HO : sample1 first order SD sample2
#-----#
*** Test Setting ***
* Resampling method
                    = subsampling
* SD order
                     = 1
* # of (sample1)
                     = 2861
                    = 1995
= 1000
* # of (sample2)
* # of (subsample1)
* # of (subsample2)
                    =
                        900
#-----#
*** Test Result ***
                    = 6.1859
* Test statistic
* Significance level
                   = 0.05
* Critical-value
                     = 4.0054
                     = 0.0000
* P-value
* Time elapsed
                     = 6.83 Sec
. pysdtest BTC_daily_rr SP500_daily_rr, ///
> resampling("subsampling") s(2) b1(1000) b2(900)
Running PySDTest
Sample1: BTC_daily_rr
Sample2: SP500_daily_rr
#--- Testing for Stochastic Dominance -----#
* HO : sample1 second order SD sample2
#-----#
```

```
*** Test Setting ***
                              = subsampling
* Resampling method
* SD order
                                  = 2

      * # of (sample1)
      =
      2861

      * # of (sample2)
      =
      1995

      * # of (subsample1)
      =
      1000

      * # of (subsample2)
      =
      900

#-----#
*** Test Result ***
* Test statistic
                                = 0.3177
* Significance level = 0.05
* Critical-value = 0.2147
* P-value
                                  = 0.0000
* Time elapsed
                           = 6.88 Sec
. pysdtest SP500_daily_rr BTC_daily_rr, ///
> resampling("subsampling") s(1) b1(900) b2(1000)
Running PySDTest
Sample1: SP500_daily_rr
Sample2: BTC_daily_rr
#--- Testing for Stochastic Dominance -----#
* HO : sample1 first order SD sample2
#-----#
*** Test Setting ***
                         od = subsampling
= 1
= 1995
* Resampling method
* SD order
* # of (sample1)
* # of (sample2)

      * # of (sample1)
      -
      1995

      * # of (sample2)
      =
      2861

      * # of (subsample1)
      =
      900

      * # of (subsample2)
      =
      1000

#-----#
*** Test Result ***
                                = 8.2384

    * Test statistic

* Significance level = 0.05

      * Critical-value
      = 6.7111

      * P-value
      = 0.0000

      * Time elapsed
      = 6.80 Sec

. pysdtest SP500_daily_rr BTC_daily_rr, ///
 > resampling("subsampling") s(2) b1(900) b2(1000)
Running PySDTest
Sample1: SP500_daily_rr
Sample2: BTC_daily_rr
#--- Testing for Stochastic Dominance ----#
* HO : sample1 second order SD sample2
#-----#
*** Test Setting ***
* Resampling method = subsampling
* Kesampring method - Subsamp

* SD order = 2

* # of (sample1) = 1995

* # of (sample2) = 2861

* # of (subsample1) = 900

* # of (subsample2) = 1000
#-----#
*** Test Result ***

      * Test statistic
      = 0.0541

      * Significance level
      = 0.05

      * Critical-value
      = 0.0783

      * P-value
      = 0.3349

      * Time elapsed
      = 6.88 Sec
```