

A NEW METRIC ON THE CONTACTOMORPHISM GROUP OF ORDERABLE CONTACT MANIFOLDS

LUKAS NAKAMURA

Abstract

We introduce a pseudo-metric on the contactomorphism group of any contact manifold (M, ξ) with a cooriented contact structure ξ . It is the contact analogue of a corresponding semi-norm in Hofer's geometry, and on certain classes of contact manifolds, its lift to the universal cover can be viewed as a continuous version of the integer valued bi-invariant metric introduced by Fraser, Polterovich, and Rosen. We show that it is non-degenerate if and only if (M, ξ) is strongly orderable and that its metric topology agrees with the interval topology introduced by Chernov and Nemirovski. In particular, the interval topology is Hausdorff whenever it is non-trivial, which answers a question of Chernov and Nemirovski. We discuss analogous results for isotopy classes of Legendrians and universal covers.

1. INTRODUCTION

Let (M, ξ) be a connected, cooriented contact manifold of dimension $2n+1$, and let $\text{Cont}_0(M, \xi)$ denote identity component of the space of compactly supported contactomorphisms on M . Let α be a contact form for ξ so that the induced coorientation agrees with the given one. The contact form α induces a one-to-one correspondence between compactly supported contact vector fields on (M, ξ) and functions $H : M \rightarrow \mathbb{R}$ by associating to a contact vector field X the function $H = \alpha(X)$, called the *Hamiltonian*. Conversely, to a (time-dependent) Hamiltonian H_t we can associate a (time-dependent) contact vector field X_t^H , and by integrating X_t^H we obtain a contact isotopy ϕ_t^H starting at the identity.

With the goal of studying the geometry of $\text{Cont}_0(M, \xi)$, Eliashberg and Polterovich [EP00] introduced the binary relation \leq on the universal cover $\widetilde{\text{Cont}}_0(M, \xi)$ of the contactomorphism group $\text{Cont}_0(M, \xi)$, where $f \leq g$ if and only if there exists a *non-negative* contact isotopy from f to g in the sense that the associated Hamiltonian is non-negative. It is clear that \leq is transitive and reflexive, and that it is antisymmetric if and only if there does not exist any contractible non-trivial non-negative loop of contactomorphisms on (M, ξ) . In the case that any contractible non-negative loop of contactomorphisms is the constant loop, (M, ξ) is called *orderable*. If M is compact, one similarly defines the quasi-order¹ \ll on $\widetilde{\text{Cont}}_0(M, \xi)$ via *positive* contact isotopies generated by a *positive* Hamiltonian. Eliashberg and Polterovich showed that (M, ξ) is orderable if and only if there does not exist a contractible positive loop of contactomorphisms in which case also \ll defines a partial order on $\widetilde{\text{Cont}}_0(M, \xi)$.

Using this partial order, they defined *the relative growth* of two elements $f, g \in \widetilde{\text{Cont}}_0(M, \xi)$ with $f \gg Id_M$ as

$$(1) \quad \gamma(f, g) := \lim_{k \rightarrow \infty} \frac{\inf\{p \in \mathbb{Z} \mid f^p \geq g^k\}}{k}$$

and showed that the limit exists in \mathbb{R} whenever (M, ξ) is orderable.

These constructions extend naturally to $\text{Cont}_0(M, \xi)$ to give quasi-orders \leq and \ll and the relative growth $\gamma(\phi, \psi)$ of two contactomorphisms $\phi, \psi \in \text{Cont}_0(M, \xi)$ with $\phi \gg Id_M$. On $\text{Cont}_0(M, \xi)$, both \leq and \ll are anti-symmetric if and only if there does not exist *any* positive loop of contactomorphisms on (M, ξ) in which case (M, ξ) is called *strongly orderable*, and in

¹Note that \ll is not reflexive unless there exists a positive loop of contactomorphisms. We will still refer to \ll as a quasi-order.

this case $\gamma(\phi, \psi) \in \mathbb{R}$ always exists. Following [EP00], (strong) orderability or non-orderability was shown for many contact manifolds (see for example [AF12], [AFM15], [AM18], [Bhu01], [BZ15], [CFP16], [CN10b], [CN16], [EKP06], [Wei13]).

Now fix a closed Legendrian $L \subseteq M$, and recall that any isotopy of L through Legendrians is induced by an ambient contact isotopy. As was already noted by Eliashberg and Polterovich, this implies that \leq also induces a quasi-order on the Legendrian isotopy class $\mathcal{L}(L)$ of L and its universal cover $\widetilde{\mathcal{L}}(L)$, and if L is closed, also \ll induces a quasi-order. We will denote the induced quasi-orders by the same symbols \leq and \ll . L is called (strongly) orderable if \leq is a partial order on $\widetilde{\mathcal{L}}(L)$ (resp. $\mathcal{L}(L)$). As before, in the compact case L is (strongly) orderable if and only if there does not exist a positive loop of Legendrians in $\widetilde{\mathcal{L}}(L)$ (resp. $\mathcal{L}(L)$). These quasi-orders were first studied in detail in [CFP16] where among other results it was shown that the fibre of the unit cotangent bundle ST^*N of any manifold N that is smoothly covered by \mathbb{R}^n is strongly orderable. In [CN10b] this result was extended to the case that the universal cover of N is an open manifold, and in [CN16] it was shown for arbitrary N that the fibre in ST^*N is orderable.

In analogy to the Hofer metric on the group of Hamiltonian diffeomorphisms of a symplectic manifold, one can associate to a contact isotopy an energy which is given by the $L^{1,\infty}$ -norm of the corresponding Hamiltonian:

$$(2) \quad \|\phi_t^H\|_\alpha := \|H_t\| := \int_0^1 \max_M |H_t| dt.$$

It follows from the behaviour of the Hamiltonians under left-/right-composition and inversion of contact isotopies that the induced *Shelukhin-Hofer metric*

$$(3) \quad d_{SH}^\alpha(\phi_0, \phi_1) := \inf\{\|H_t\| \mid H_t : M \rightarrow \mathbb{R}, \phi_1^H \phi_0 = \phi_1\}$$

defines a right-invariant pseudo-metric on $\text{Cont}_0(M, \xi)$ which behaves naturally under left-composition:

$$(4) \quad d_{SH}^\alpha(\psi\phi_0, \psi\phi_1) = d_{SH}^{\psi^*\alpha}(\phi_0, \phi_1).$$

Furthermore, it follows easily that d_{SH}^α and $d_{SH}^{\alpha'}$ are equivalent for any two choices of contact forms α and α' for ξ which agree outside of a compact subset.

In [She17], Shelukhin showed that d_{SH}^α is, in fact, non-degenerate on any contact manifold. Consequently, it defines a genuine metric on $\text{Cont}_0(M, \xi)$.

The energy of a contact isotopy also naturally induces a right-invariant and left-natural pseudo-metric $\widetilde{d}_{SH}^\alpha$ on the universal cover $\widetilde{\text{Cont}}_0(M, \xi)$ of $\text{Cont}_0(M, \xi)$, but as was pointed out by Shelukhin [She17], it is only known that $\widetilde{d}_{SH}^\alpha(f, g) > 0$ for $f, g \in \widetilde{\text{Cont}}_0(M, \xi)$ if $\pi(f) \neq \pi(g)$ where $\pi : \widetilde{\text{Cont}}_0(M, \xi) \rightarrow \text{Cont}_0(M, \xi)$ denotes the universal covering map.

d_{SH}^α (and $\widetilde{d}_{SH}^\alpha$) induce pseudo-metrics d_{SCH}^α (and $\widetilde{d}_{SCH}^\alpha$) on orbits of parametrized or unparametrized subsets A of M (and their universal cover) under the action of $\text{Cont}_0(M, \xi)$ (and $\widetilde{\text{Cont}}_0(M, \xi)$), which we will refer to as the *Shelukhin-Chekanov-Hofer metric*. d_{SCH}^α was studied for closed unparametrized submanifolds L by Rosen and Zhang in [RZ20] where they observed the following. The pseudo-metric d_{SCH}^α can only be non-degenerate if L is coisotropic, and if $\dim(L) < n$, where $\dim(M) = 2n + 1$, then d_{SCH}^α vanishes identically. In the case that $\dim(L) = n$, d_{SCH}^α vanishes identically unless L has a Legendrian component, and if L is a connected Legendrian, then d_{SCH}^α either vanishes identically or is non-degenerate. For disjoint unions of connected, parametrized non-Legendrian submanifolds of dimension n , Dimitroglou Rizell and Sullivan [DRS22] proved that d_{SCH}^α vanishes identically as well.

Motivated by the Alexandrov topology in Lorentzian geometry, Chernov and Nemirovski [CN20] used the quasi-order \ll to define the *interval topologies* $\widetilde{\mathcal{O}}_{\ll}$ and \mathcal{O}_{\ll} on $\widetilde{\text{Cont}}_0(M, \xi)$ and $\text{Cont}_0(M, \xi)$ respectively, which are generated by intervals

$$(5) \quad I_{f_0, f_1}^{\ll} = \{f \in \widetilde{\text{Cont}}_0(M, \xi) \mid f_0 \ll f \ll f_1\}$$

in $\widetilde{\text{Cont}}_0(M, \xi)$ with respect to \ll , and similarly for \mathcal{O}_{\ll} . In fact, it is straightforward to see that the intervals $I_{f_t^\alpha f, f_t^\alpha f}$ and $I_{\phi_{-\varepsilon}^\alpha \phi, \phi_\varepsilon^\alpha \phi}$, $\varepsilon > 0$, $f \in \widetilde{\text{Cont}}_0(M, \xi)$, $\phi \in \text{Cont}_0(M, \xi)$, form a basis of $\widetilde{\mathcal{O}}_{\ll}$ and \mathcal{O}_{\ll} , respectively, where ϕ_t^α denotes the Reeb flow and $f_t^\alpha := [s \mapsto \phi_{st}^\alpha]$ denotes its lift to $\widetilde{\text{Cont}}_0(M, \xi)$.

It follows easily that the interval topologies are trivial if there exists a (contractible) positive loop of contactomorphisms. In the absence of positive loops, Chernov and Nemirovski asked whether the interval topologies are necessarily Hausdorff.

Question 1.1. [CN20] *If (M, ξ) is (strongly) orderable, is $\widetilde{\mathcal{O}}_{\ll}$ (\mathcal{O}_{\ll}) Hausdorff?*

They proved that this is indeed the case on spherical cotangent bundles of closed manifolds which can be smoothly covered by an open subset of \mathbb{R}^n .

Hedicke [Hed21] showed that the interval topology is coarser than the topology $\mathcal{O}_{d_{SH}^\alpha}$ induced by d_{SH}^α , i.e. $\mathcal{O}_{\ll} \subseteq \mathcal{O}_{d_{SH}^\alpha}$, which leads to the following natural question.

Question 1.2. (Question 2 in [Hed21]) *Are \mathcal{O}_{\ll} and $\mathcal{O}_{d_{SH}^\alpha}$ equal?*

Using the quasi-order \ll on $\mathcal{L}(L)$ and $\widetilde{\mathcal{L}}(L)$, Chernov and Nemirovski [CN20] again defined intervals I_{L_0, L_1}^{\ll} and interval topologies \mathcal{O}_{\ll}^L and $\widetilde{\mathcal{O}}_{\ll}^L$ for a closed Legendrian $L \subseteq M$. As above, these can only be non-trivial if L is (strongly) orderable, and intervals whose endpoints are related via the Reeb flow form a basis. Hedicke [Hed21] proved also in this case that $\mathcal{O}_{\ll}^L \subseteq \mathcal{O}_{d_{CSH}^\alpha}^L$.

Further inspired by Lorentzian geometry, Hedicke [Hed21] recently introduced the Lorentzian distance function

$$(6) \quad \begin{aligned} \tau_\alpha : \text{Cont}_0(M, \xi) \times \text{Cont}_0(M, \xi) &\rightarrow [0, \infty] \\ (\phi_0, \phi_1) &\mapsto \begin{cases} \sup \left\{ \int_0^1 \min_M H_t dt \mid \phi_1^H \phi_0 = \phi_1, H \geq 0 \right\} & \text{if } \phi_1 \geq \phi_0, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

and showed that if (M, ξ) is strongly orderable, then $\tau_\alpha < \infty$, and that $\tau_\alpha(\phi_0, \phi_1) > 0$ if and only if $\phi_0 \ll \phi_1$. Furthermore, he proved that τ_α is continuous with respect to \mathcal{O}_{\ll} and thus also with respect to d_{SH}^α .

As explained in [Hed21], τ_α can also be defined on $\mathcal{L}(L)$ for any closed Legendrian L , and it satisfies similar properties as on $\text{Cont}_0(M, \xi)$.

The definition of τ_α extends naturally also to $\widetilde{\text{Cont}}_0(M, \xi)$ and $\widetilde{\mathcal{L}}(L)$, and it is easy to check Hedicke's results mentioned above still apply with the same proofs.

The goal of this article is to further analyse the above questions. Similarly to the semi-norms in symplectic geometry which measure only the positive or negative part of a normalized Hamiltonian, we introduce below ‘‘distances’’ δ_α^\pm on $\text{Cont}_0(M, \xi)$ which are bounded by the Shelukhin-Hofer metric and can be used to express the Lorentzian distance function. It turns out that their maximum $d_\alpha := \max\{\delta_\alpha^\pm, 0\}$ defines a pseudo-metric on $\text{Cont}_0(M, \xi)$. In fact, we prove the following, which answers Question 1.1 above (see Section 2.2 below).

Theorem 1.3. d_α either vanishes identically or is non-degenerate, and it satisfies $d_\alpha \leq d_{SH}^\alpha$. If M is compact, then the topology on $\text{Cont}_0(M, \xi)$ induced by d_α agrees with the interval topology \mathcal{O}_{\ll} , and d_α vanishes identically if and only if there exists a positive loop of contactomorphisms on M . \square

We also introduce versions of δ_α^\pm and d_α on $\widetilde{\text{Cont}}_0(M, \xi)$ and spaces of parametrized or unparametrized subspaces and their universal covers and discuss similar results for these spaces, but similarly to the Shelukhin-Hofer metric, our methods do not suffice to prove genuine non-degeneracy on the universal covers.

Remark 1.4. Allais and Arlove [AA23] independently also studied the pseudo-metrics considered here and obtained proofs of our main results. After prepublication of the first version of this article, it came to our attention that the definition of the pseudo-norms associated to $\widetilde{\delta}_\alpha^\pm$ and \widetilde{d}_α on $\widetilde{\text{Cont}}_0(M, \xi)$ together with Example 2.9 below were already introduced in Section 5 of Arlove's PhD thesis [Arl21].

Acknowledgements: I am grateful to Georgios Dimitroglou Rizell for introducing me to Question 1.1 which was the starting point for this project, and to Stefan Nemirovski for explaining to me part of his work on orderability. I would also like to thank my advisor Tobias Ekholm for valuable discussions and for careful reading of a preliminary version of this article. I want to thank Simon Allais and Pierre-Alexandre Arlove for directing my attention to Arlove's PhD thesis.

This work is partially supported by the Knut and Alice Wallenberg Foundation, grant KAW2020.0307.

2. MAIN RESULTS

2.1. Distances on the universal cover $\widetilde{\text{Cont}}_0(M, \xi)$ of $\text{Cont}_0(M, \xi)$.

For any two $f_0, f_1 \in \widetilde{\text{Cont}}_0(M, \xi)$, we define

$$(7) \quad \begin{aligned} \widetilde{\delta}_\alpha^-(f_0, f_1) &:= \inf\{\varepsilon \in \mathbb{R} \mid \exists H_t : M \rightarrow [-\varepsilon, \infty) : [\phi_t^{H_t}] \cdot f_0 = f_1\}, \\ \widetilde{\delta}_\alpha^+(f_0, f_1) &:= \inf\{\varepsilon \in \mathbb{R} \mid \exists H_t : M \rightarrow (\infty, \varepsilon] : [\phi_t^{H_t}] \cdot f_0 = f_1\}. \end{aligned}$$

Recall that we write $f \ll g$ for $f, g \in \widetilde{\text{Cont}}_0(M, \xi)$ if there exists a positive contact isotopy from f to g . Using the class $f_t^\alpha = [s \mapsto \phi_{st}^\alpha, s \in [0, 1]] \in \widetilde{\text{Cont}}_0(M, \xi)$ of the time- t Reeb flow ϕ_t^α and the cones

$$(8) \quad I_f^\pm := \left\{ g \in \widetilde{\text{Cont}}_0(M, \xi) \mid f^{\pm 1} \ll g^{\pm 1} \right\}$$

in the case that M is compact, $\widetilde{\delta}_\alpha^\pm$ can also be expressed as

$$(9) \quad \widetilde{\delta}_\alpha^\pm(f_0, f_1) = \inf \left\{ \varepsilon \in \mathbb{R} \mid f_1 \in I_{f_{\pm\varepsilon}^\mp f_0}^\mp \right\},$$

since any contact isotopy ϕ_t can be written as $\phi_{-\varepsilon t}^\alpha \circ (\phi_{\varepsilon t}^\alpha \circ \phi_t) = \phi_{\varepsilon t}^\alpha \circ (\phi_{-\varepsilon t}^\alpha \circ \phi_t)$ and the Reeb flow is a strict contact isotopy. Then the Hamiltonian associated to ϕ_t is $\geq -\varepsilon$ if and only if the Hamiltonian associated to $\phi_{\varepsilon t}^\alpha \circ \phi_t$ is ≥ 0 , and it is $\leq \varepsilon$ if and only if the Hamiltonian associated to $\phi_{-\varepsilon t}^\alpha \circ \phi_t$ is ≤ 0 .

This together with the observation that for any $\eta > 0$ and $f_1 \in I_{f_{\pm\varepsilon}^\mp f_0}^\mp$, we have $I_{f_{-\eta}^\alpha f_1, f_\eta^\alpha f_1}^{\ll} = I_{f_\eta^\alpha f_1}^- \cap I_{f_{-\eta}^\alpha f_1}^+ \subseteq I_{f_{\pm(\varepsilon+\eta)}^\mp f_0}^\mp$ (see (5) above for the definition of I^{\ll}) shows that $\widetilde{\delta}_\alpha^\pm$ is continuous with respect to the interval topology. Note also that $\widetilde{\delta}_\alpha^-(f_0, f_1) < 0$ if and only if $f_1 \gg f_0$.

Let $\pi : \widetilde{\text{Cont}}_0(M, \xi) \rightarrow \text{Cont}_0(M, \xi)$ denote the natural projection. $\widetilde{\delta}_\alpha^+$ and $\widetilde{\delta}_\alpha^-$ satisfy the following properties.

Proposition 2.1. *Let $f_0, f_1, f_2 \in \widetilde{\text{Cont}}_0(M, \xi)$. Then*

(i) (triangle inequality) $\widetilde{\delta}_\alpha^\pm(f_0, f_2) \leq \widetilde{\delta}_\alpha^\pm(f_0, f_1) + \widetilde{\delta}_\alpha^\pm(f_1, f_2)$.

(ii) (symmetry) $\widetilde{\delta}_\alpha^+(f_0, f_1) = \widetilde{\delta}_\alpha^-(f_1, f_0)$.

(iii) (left naturality) $\widetilde{\delta}_\alpha^\pm(f_2 f_0, f_2 f_1) = \widetilde{\delta}_{(\pi(f_2))^* \alpha}^\pm(f_0, f_1)$.

(iv) (inverse) $\widetilde{\delta}_\alpha^+(id_M, f_0^{-1}) = \widetilde{\delta}_\alpha^-(id_M, f_0)$.

(v) (right invariance) $\widetilde{\delta}_\alpha^\pm(f_0 \circ f_2, f_1 \circ f_2) = \widetilde{\delta}_\alpha^\pm(f_0, f_1)$.

(vi) (orderability) *Assume that M is compact. If (M, ξ) is not orderable, then $\widetilde{\delta}_\alpha^\pm \equiv -\infty$. If (M, ξ) is orderable, then $\widetilde{\delta}_\alpha^\pm > -\infty$, and at least one of $\widetilde{\delta}_\alpha^+(f_0, f_1)$ and $\widetilde{\delta}_\alpha^-(f_0, f_1)$ is non-negative, and if one is negative, then the other one is positive.*

Proof. The properties (ii)–(v) follow similarly to the corresponding properties of the Shelukhin-Hofer metric d_{SH}^α from the behavior of the Hamiltonians (see the proof of Theorem A in [She17]). (i) is a consequence of Proposition 2.5 below. To see (vi), first note that if there exists a contractible positive loop of contactomorphisms, then for any $f \in \widetilde{\text{Cont}}_0(M, \xi)$, $I_f^\pm = \widetilde{\text{Cont}}_0(M, \xi)$, so $\widetilde{\delta}_\alpha^\pm = -\infty$. Conversely, assume that $\widetilde{\delta}_\alpha^\pm(f_0, f_1) = -\infty$. Then by the triangle inequality, also $\widetilde{\delta}_\alpha^\pm(f_1, f_1) = -\infty$. If $\widetilde{\delta}_\alpha^+(f_0, f_1) < 0$ and $\widetilde{\delta}_\alpha^-(f_0, f_1) = \widetilde{\delta}_\alpha^+(f_1, f_0) \leq 0$ or $\widetilde{\delta}_\alpha^+(f_0, f_1) \leq 0$ and $\widetilde{\delta}_\alpha^-(f_0, f_1) = \widetilde{\delta}_\alpha^+(f_1, f_0) < 0$, then again by the triangle inequality, $\widetilde{\delta}_\alpha^\pm(f_1, f_1) < 0$. In either case $f_1 \gg f_1$, so (M, ξ) is not orderable. \square

Now we define

$$(10) \quad \widetilde{d}_\alpha(f_0, f_1) := \max\{\widetilde{\delta}_\alpha^-(f_0, f_1), \widetilde{\delta}_\alpha^+(f_0, f_1), 0\}.$$

Remark 2.2. In the case that the Reeb flow of (M, α) is 1-periodic, this definition was previously considered in the form of the metric balls of \widetilde{d}_α in [FPR18] where it was shown that the boundary of the balls

$$(11) \quad \mathcal{B}(k) := \{f \in \widetilde{\text{Cont}}_0(M, \xi) \mid f_{-k}^\alpha \leq f \leq f_k^\alpha\}, \quad k \in \mathbb{N},$$

with respect to the interval topology $\widetilde{\mathcal{O}}_\ll$ only contains lifts of contactomorphisms with k - or $-k$ -translated points.

In the case that M is compact, it follows from Equation 9 that $\widetilde{d}_\alpha(f_0, f_1)$ is also equal to

$$(12) \quad \widetilde{d}_\alpha(f_0, f_1) = \inf\{\varepsilon > 0 \mid f_1 \in I_{f_{-\varepsilon}^\alpha f_0, f_\varepsilon^\alpha f_0}^\ll\},$$

or equivalently,

$$(13) \quad \{f_1 \in \widetilde{\text{Cont}}_0(M, \xi) \mid \widetilde{d}_\alpha(f, f_1) < \varepsilon\} = I_{f_{-\varepsilon}^\ll f, f_\varepsilon^\alpha f}^\ll.$$

With Proposition 2.1, it is immediate that \widetilde{d}_α is a pseudo-metric, and by (13), its metric topology agrees with the interval topology $\widetilde{\mathcal{O}}_\ll$ since the intervals $I_{f_{-\varepsilon}^\ll f, f_\varepsilon^\alpha f}^\ll$ form a basis.

In fact, the following holds.

Proposition 2.3. *Let $f_0, f_1, f_2 \in \widetilde{\text{Cont}}_0(M, \xi)$. Then*

(i) (triangle inequality) $\widetilde{d}_\alpha(f_0, f_2) \leq \widetilde{d}_\alpha(f_0, f_1) + \widetilde{d}_\alpha(f_1, f_2)$.

(ii) (symmetry) $\widetilde{d}_\alpha(f_0, f_1) = \widetilde{d}_\alpha(f_1, f_0)$.

(iii) (left naturality) $\widetilde{d}_\alpha(f_2 f_0, f_2 f_1) = d_{(\pi(f_2))^* \alpha}(f_0, f_1)$.

(iv) (inverse) $\widetilde{d}_\alpha(f_0^{-1}, id_M) = \widetilde{d}_\alpha(f_0, id_M)$.

(v) (right invariance) $\widetilde{d}_\alpha(f_0 \circ f_2, f_1 \circ f_2) = \widetilde{d}_\alpha(f_0, f_1)$.

(vi) (topology) *If M is compact, then the metric topology of \widetilde{d}_α agrees with the interval topology $\widetilde{\mathcal{O}}_\ll$.*

Proof. Immediate from Proposition 2.1 and the above discussion. \square

The properties in Proposition 2.3 together with the following straightforward lemma imply that $(\widetilde{\text{Cont}}_0(M, \xi), \widetilde{d}_\alpha)$ is a topological group.

Lemma 2.4. *Let $\alpha' = \lambda\alpha$ be another contact form for ξ where $\lambda : M \rightarrow \mathbb{R}_{>0}$ is a strictly positive function that is equal to the constant function 1 outside of a compact set. Then*

$$(14) \quad \min_M(\lambda) \widetilde{d}_\alpha \leq \widetilde{d}_{\alpha'} \leq \max_M(\lambda) \widetilde{d}_\alpha.$$

In particular, \widetilde{d}_α and $\widetilde{d}_{\alpha'}$ induce the same topology on $\widetilde{\text{Cont}}_0(M)$.

Proof. Clear. \square

It is also easy to see that \widetilde{d}_α is always bounded by the Shelukhin-Hofer metric $\widetilde{d}_{SH}^\alpha$. In fact, if we denote by $\widetilde{\delta}_{SH}^+(f_0, f_1)$, resp. $\widetilde{\delta}_{SH}^-(f_0, f_1)$, the infimum of $\|H_t\|^+$, resp. $\|H_t\|^-$, over all Hamiltonians H_t whose associated contact isotopy ϕ_t^H satisfies $[\phi_t^H] \cdot f_0 = f_1$, where $\|H_t\|^\pm := \int_0^1 \max_M \{\pm H_t\} dt$, we have the following.

Proposition 2.5. *For any $f_0, f_1 \in \widetilde{\text{Cont}}_0(M, \alpha)$,*

$$(15) \quad \widetilde{\delta}_{SH}^\pm = \widetilde{\delta}_\alpha^\pm.$$

In particular,

$$(16) \quad \widetilde{d}_\alpha(f_0, f_1) = \max \left\{ \widetilde{\delta}_{SH}^-(f_0, f_1), \widetilde{\delta}_{SH}^+(f_0, f_1), 0 \right\} \leq d_{SH}(f_0, f_1).$$

Proof. Note that if M is non-compact, $\max_M \{\pm H_t\} \geq 0$ for any compactly supported Hamiltonian H_t , in which case Proposition 2.5 follows directly from Lemma 3.1 below applied to $\max_M \{\pm H_t\}$. In the compact case, the formula in Proposition 2.5 is just (40) in Section 2.4 below. \square

An obvious question to ask is when \widetilde{d}_α is non-degenerate. Of course, part (vi) of Proposition 2.1 shows that \widetilde{d}_α vanishes identically if M is compact and (M, ξ) is not orderable. It turns out that the converse is true as well, and that \widetilde{d}_α is always non-degenerate in a weaker sense unless it vanishes identically. We say that a pseudo-metric d on $\widetilde{\text{Cont}}_0(M, \xi)$ is π -non-degenerate if for any $f_0, f_1 \in \widetilde{\text{Cont}}_0(M, \xi)$ with $\pi(f_0) \neq \pi(f_1)$, we have that $d(f_0, f_1) > 0$. Similarly, we say that a topology \mathcal{O} on $\widetilde{\text{Cont}}_0(M, \xi)$ is π -Hausdorff if any $f_0, f_1 \in \widetilde{\text{Cont}}_0(M, \xi)$ with $\pi(f_0) \neq \pi(f_1)$ have disjoint open neighbourhoods, where $\pi : \widetilde{\text{Cont}}_0(M, \xi) \rightarrow \text{Cont}_0(M, \xi)$ is the covering map. Clearly, these notions are weaker than Hausdorffness and, respectively, non-degeneracy in general.

Theorem 2.6. *\widetilde{d}_α either vanishes identically or is π -non-degenerate. If M is compact, then \widetilde{d}_α vanishes identically if and only if there exists a contractible positive loop of contactomorphisms on M .*

The proof of Theorem 2.6 is given in Section 4.

Since $\widetilde{\mathcal{O}}_\ll$ is the metric topology of \widetilde{d}_α , it is clear that $\widetilde{\mathcal{O}}_\ll$ is (π) -Hausdorff if \widetilde{d}_α is (π) -non-degenerate. An immediate consequence is the following.

Corollary 2.7. *Let (M, ξ) be a closed contact manifold. Then the interval topology $\widetilde{\mathcal{O}}_\ll$ on $\widetilde{\text{Cont}}_0(M, \xi)$ is π -Hausdorff if and only if there does not exist a contractible positive loop of contactomorphisms on (M, ξ) .*

In the case that \tilde{d}_α is π -non-degenerate, we now have the two natural metrics $\tilde{d}_\alpha \leq \tilde{d}_{SH}^\alpha$ on $\widetilde{\text{Cont}}_0(M, \xi)$.

Question 2.8. *Is \tilde{d}_α equivalent to \tilde{d}_{SH}^α if it is π -non-degenerate?*

We expect that $\tilde{d}_\alpha \neq \tilde{d}_{SH}^\alpha$ in general (see Remark 2.14 below).

Example 2.9. (strict, autonomous contact flows in ST^*T^n) (see also [Arl21, Theorem 5.8]) Consider the unit cotangent bundle ST^*T^n of the n -torus T^n . We use coordinates $(q, p) \in ST^*T^n \subseteq T^*T^n \approx T^n \times \mathbb{R}^n$ where $q \in T^n = ([0, 1]/(0 = 1))^n$ and $p \in ST_q^*T^n \approx S^{n-1} \subseteq \mathbb{R}^n$. The contact form is the standard contact form given by $\alpha = pdq$. Let $H = H(p)$ and $G = G(p)$ be autonomous Hamiltonians depending only the p -coordinate. We claim that

$$(17) \quad \tilde{\delta}_\alpha^+(f_t^H, f_t^G) = t \max_M(G - H), \quad \tilde{\delta}_\alpha^-(f_t^H, f_t^G) = -t \min_M(G - H) \quad \text{for } t \geq 0.$$

In particular, it follows that

$$(18) \quad \tilde{d}_{SH}^\alpha(Id, f_t^H) = \tilde{d}_\alpha(Id, f_t^H) = t \max_M |H|,$$

which means that f_t^H is length minimizing for \tilde{d}_{SH} and \tilde{d}_α for all times.

To see this, we need to understand the relation \ll for contact isotopies of the above form. Let $H(p)$ and $G(p)$ be Hamiltonians as above. Then $f_t^H \ll f_t^G$ if and only if $H(p) < G(p)$ for all $p \in S^{n-1}$. Indeed, first note that the contactomorphism ϕ_t^H is strict since H is preserved by the Reeb flow $\phi_t^\alpha(q, p) = (q + tp, p)$. Thus, the contact isotopy $(\phi_t^H)^{-1} \circ \phi_t^G$ is generated by the Hamiltonian

$$(19) \quad F_t = -H \circ \phi_t^H + G \circ \phi_t^H,$$

which is positive if $H < G$. Conversely, if $f_t^H \ll f_t^G$, then in the notation of Section 3.2 in [EP00], we have that

$$(20) \quad H(p) = r_\pm(p, f_t^H) \leq r_\pm(p, f_t^G) = G(p)$$

for any $p \in S^{n-1}$ (viewed now also as a non-zero cohomology class of T^n). In fact, if $f_t^H \ll f_t^G$ for some fixed t , then also $f_t^H \ll f_t^{G-\varepsilon}$ for all sufficiently small $\varepsilon > 0$, so we see that $H(p) \leq G(p) - \varepsilon < G(p)$.

Since $s \mapsto \phi_{st}^H, s \in [0, 1]$, and $s \mapsto \phi_{s\varepsilon}^\alpha \phi_{st}^G, s \in [0, 1]$, are generated by the Hamiltonians tH and $tG + \varepsilon$, resp., it follows that $f_t^H \ll f_\varepsilon^\alpha f_t^G$ if and only if $tH < tG + \varepsilon$, and $f_t^H \gg f_{-\varepsilon}^\alpha f_t^G$ if and only if $tH > tG - \varepsilon$. In view of (9), this is equivalent to (17) if $t \geq 0$.

Remark 2.10. One may ask about the continuity properties of the relation \leq with respect to the interval topology $\widetilde{\mathcal{O}}_\ll$, e.g. are the intervals

$$(21) \quad I_{f_0, f_1}^\leq := \{f \in \widetilde{\text{Cont}}_0(M, \xi) \mid f_0 \leq f \leq f_1\}$$

or more generally the cones

$$(22) \quad I_f^{\pm, \leq} := \{g \in \widetilde{\text{Cont}}_0(M, \xi) \mid f^{\pm 1} \leq g^{\pm 1}\}$$

closed with respect to $\widetilde{\mathcal{O}}_\ll$? It is easy to see that $I_f^{\pm, \leq} \subseteq \overline{I_f^\pm}$ is always true, but it is unclear when equality holds.

Example 2.9 shows that for an autonomous Hamiltonian H on ST^*T^n that depends only on the p -coordinate, $f^H \in \overline{I_{id}^\pm}$ if and only if $f^H \in I_{id}^{\pm, \leq}$: If, say, $f^H \not\leq id$, then for some $p \in S^{n-1}$, $H(p) < 0$. It follows that for any $\varepsilon > 0$ with $H(p) + \varepsilon < 0$, $f_\varepsilon^\alpha f \not\leq id$. In particular, $I_{f_\varepsilon^\alpha f, f_\varepsilon^\alpha f}^\leq \cap I_{id}^+ = \emptyset$. Since the intervals $I_{f_\varepsilon^\alpha f, f_\varepsilon^\alpha f}^\leq$ form a neighbourhood basis of f with respect to $\widetilde{\mathcal{O}}_\ll$, this is equivalent to $f \notin \overline{I_{id}^+}$. The case $f^H \not\leq id$ works similarly.

The same question can be asked about the relation \leq and the interval topology on the other spaces considered below. In the case of the path-connected component of the zero section in

the space of Legendrians in the 1-jet bundle J^1L of a closed manifold L , it follows in a similar way from Example 2.24 below that for a function $f : L \rightarrow \mathbb{R}$, $j^1f \in \overline{I_{j^1_0}^\pm}$ if and only if $f \in I_{j^1_0}^{\pm, \leq}$ if and only if $\pm f \geq 0$.

2.2. Distances on $\text{Cont}_0(M, \xi)$.

On $\text{Cont}_0(M, \xi)$, one similarly defines δ_α^\pm as before, except that we now take the infimum over *all* contact isotopies with the given endpoints and not just those in a fixed homotopy class. The distances δ_α^\pm then satisfy the analogue of Proposition 2.1 by the same proof, where in (vi) we replace orderability by strong orderability. Thus,

$$(23) \quad d_\alpha := \max\{\delta_\alpha^-, \delta_\alpha^+, 0\}$$

defines a pseudo-metric on $\text{Cont}_0(M, \xi)$ satisfying the analogues of (i) – (v) in Proposition 2.3. By essentially the same proof as before, \mathcal{O}_\ll agrees with the metric topology of d_α , and by the simplicity of $\text{Cont}_0(M, \xi)$ (see [Ryb10]) we have the following version of Theorem 2.6 (see Section 4 for the proof).

Theorem 2.11. *On $\text{Cont}_0(M, \xi)$, d_α either vanishes identically or is non-degenerate. If M is compact, then d_α vanishes identically if and only if there exists a positive loop of contactomorphisms on M . \square*

Corollary 2.12. *Let (M, ξ) be a closed contact manifold. Then the interval topology \mathcal{O}_\ll on $\text{Cont}_0(M, \xi)$ is Hausdorff if and only if there does not exist a positive loop of contactomorphisms on (M, ξ) . \square*

Clearly, d_α is bounded by the Shelukhin-Hofer metric. Again, it is unclear whether d_α is equivalent to the Shelukhin-Hofer metric on strongly orderable contact manifolds.

Remark 2.13. \tilde{d}_α also induces the pseudo-metric

$$(24) \quad \widehat{d}_\alpha(\phi_0, \phi_1) := \inf\{\tilde{d}_\alpha(f_0, f_1) \mid \pi(f_0) = \phi_0, \pi(f_1) = \phi_1\}$$

on $\text{Cont}_0(M, \xi)$ where $\pi : \widetilde{\text{Cont}}_0(M, \xi) \rightarrow \text{Cont}_0(M, \xi)$ denotes the covering map. It again satisfies the analogues of properties (i) – (v) in Proposition 2.3, and it is either non-degenerate or vanishes identically. But in contrast to d_α , the proof of Theorem 2.6 actually shows that \tilde{d}_α vanishes identically if \widehat{d}_α is degenerate (see Remark 4.1). It follows that \widehat{d}_α is non-degenerate if and only if (M, ξ) is orderable. In particular, \widehat{d}_α is different from d_α in general. For example, ST^*S^2 with its standard contact structure has periodic Reeb flow and is orderable by [CN16, Theorem 1.1].

Clearly, we always have

$$(25) \quad d_\alpha \leq \widehat{d}_\alpha \leq d_{SH}^\alpha.$$

There is also a version of the interval topology which agrees with the metric topology of \widehat{d}_α . Namely, it is generated by sets of the form

$$(26) \quad I_{\phi_t}^\ll := \bigcap_f \pi(I_{f, [\phi_t]f}^\ll),$$

where $\phi_t \in \text{Cont}_0(M, \xi)$, $t \in [0, 1]$, is a contact isotopy (not necessarily starting at the identity), and the intersection is taken over all $f \in \widetilde{\text{Cont}}_0(M, \xi)$ with $\pi(f) = \phi_0$.

Remark 2.14. The definitions of δ_α^\pm , d_α , and \widehat{d}_α are reminiscent of the semi-norms ρ^\pm , $\rho^+ + \rho^-$, and ρ_f in Hofer's geometry (see [EP93], [McD02]). It was shown by McDuff [McD02] that ρ_f and $\rho^+ + \rho^-$ may differ, so we expect that $\widehat{d}_\alpha \neq d_\alpha$ is possible even if M is strongly orderable.

Remark 2.15. Assume now that (M, ξ) is compact and orderable, but not strongly orderable, i.e. \widetilde{d}_α and \widehat{d}_α define (π) -non-degenerate pseudo-metrics on $\widetilde{\text{Cont}}_0(M, \xi)$ and $\text{Cont}_0(M, \xi)$, respectively, and there exists a positive loop $\phi_t \in \text{Cont}_0(M, \xi)$, $t \in [0, 1]$. By Lemma 3.1.A in [EP00] we may assume that the generating Hamiltonian F_t of ϕ_t is 1-periodic. Denote by f the element of $\widetilde{\text{Cont}}_0(M, \xi)$ represented by the loop ϕ_t . Following the constructions in [FPR18], we can define

$$(27) \quad \begin{aligned} \widetilde{\nu}_{FPR}^{f,+}(g) &:= \min\{k \in \mathbb{Z} \mid f^k \geq g\}, \\ \widetilde{\nu}_{FPR}^{f,-}(g) &:= \max\{k \in \mathbb{Z} \mid f^k \leq g\}, \\ \widetilde{\nu}_{FPR}^f(g) &:= \max\{|\widetilde{\nu}_{FPR}^{f,+}(g)|, |\widetilde{\nu}_{FPR}^{f,-}(g)|\} \\ &= \min\{k \in \mathbb{Z}_{\geq 0} \mid f^{-k} \leq g \leq f^k\}, \end{aligned}$$

for any $g \in \widetilde{\text{Cont}}_0(M, \xi)$. If the Reeb flow of the contact form α is 1-periodic and $f = f_1^\alpha$, this definition reduces to the norms ν_\pm and ν considered in [FPR18]. By the same arguments that were used in [FPR18] it is straightforward to check that $\widetilde{\nu}_{FPR}^f$ defines a pseudo-norm on $\widetilde{\text{Cont}}_0(M, \xi)$ that is non-degenerate if and only if (M, ξ) is orderable. Furthermore, $\widetilde{\nu}_{FPR}^{f,\pm}$ and $\widetilde{\nu}_{FPR}^f$ are conjugation invariant since f is in the center of $\widetilde{\text{Cont}}_0(M, \xi)$ and \geq is bi-invariant.

We claim that

$$(28) \quad (\widetilde{\nu}_{FPR}^f(g) - 1)\widetilde{\tau}_\alpha([id_M], f) \leq \widetilde{d}_\alpha([id_M], g) \leq \widetilde{\nu}_{FPR}^f(g)\widetilde{d}_\alpha([id_M], f),$$

where $\widetilde{\tau}_\alpha$ denotes the natural extension of τ_α to the universal cover $\widetilde{\text{Cont}}_0(M, \xi)$ (see Section 2.4).

Indeed, let G_t be a Hamiltonian that generates g . If $\int_0^1 \max G_t dt < k \int_0^1 \min F_t dt$ for $k \geq 0$, then $f^k \gg g$, and if $\int_0^1 \min G_t dt > k \int_0^1 \min F_t dt$ for $k \leq 0$, then $f^k \ll g$. This implies the first inequality. On the other hand, if $f^k \geq g$ for $k \geq 0$, then it follows from Lemma 2.5 in [FPR18] that g can be generated by a Hamiltonian G_t that satisfies $\int_0^1 \max G_t dt \leq k \int_0^1 \max F_t dt$, and if $f^k \leq g$ for $k \leq 0$, then g can be generated by a Hamiltonian G_t that satisfies $\int_0^1 \min G_t dt \geq k \int_0^1 \max F_t dt$. This implies the second inequality.

In the special case that the Reeb flow of the contact form α is 1-periodic and $f = f_1^\alpha$, (28) simply reads

$$(29) \quad \widetilde{\nu}_{FPR}^{f_1^\alpha}(g) - 1 \leq \widetilde{d}_\alpha([id_M], g) \leq \widetilde{\nu}_{FPR}^{f_1^\alpha}(g)$$

since $\widetilde{\tau}_\alpha([id_M], f_1^\alpha) = \widetilde{d}_\alpha([id_M], f_1^\alpha) = 1$ (see Section 2.5).

Note that the left-most and the right-most terms in (28) are invariant under conjugating g , whereas the middle term may change. In particular, these inequalities give bounds on $\widetilde{d}_\alpha([id_M], g)$ for all g in a fixed conjugacy class.

Following [FPR18], we define

$$(30) \quad \nu_{FPR}^f(\psi) := \min\{\widetilde{\nu}_{FPR}^f(g) \mid \pi(g) = \psi\}$$

for $\psi \in \text{Cont}_0(M, \xi)$, where the minimum is achieved since $\widetilde{\nu}_{FPR}^f(g) \geq 0$ is integer valued.

By taking the infimum over all g with $\pi(g) = \psi$ in (28), we obtain

$$(31) \quad (\nu_{FPR}^f(\psi) - 1)\tilde{\tau}_\alpha([id_M], f) \leq \widehat{d}_\alpha(id_M, \psi) \leq \nu_{FPR}^f(\psi)\tilde{d}_\alpha([id_M], f).$$

As before the right-most and the left-most terms are invariant under conjugating ψ , so that we get bounds on $\widehat{d}_\alpha(id_M, \psi)$ for all ψ in a fixed conjugacy class.

2.3. Distances on orbit spaces of subsets.

Let $A \subseteq M$ be a closed subset, and let

$$(32) \quad \mathcal{L}(A) := \{\phi(A) \subseteq M \mid \phi \in \text{Cont}_0(M, \xi)\}$$

be the orbit of A under $\text{Cont}_0(M)$. Denote the inclusion of $A \hookrightarrow M$ by i . An isotopy $i_t, t \in [0, 1]$, starting at i is a *positive* (resp. *non-negative*) isotopy if it is induced by an ambient contact isotopy ϕ_t with Hamiltonian H_t that satisfies $H_t|_{\phi_t(A)} > 0$ (resp. $H_t|_{\phi_t(A)} \geq 0$). Note that the condition on H_t does not depend on the choice of ϕ_t since $H_t \circ i_t = \alpha(\frac{di_t}{dt})$, and that the existence of a positive or non-negative isotopy does not depend on the parametrization i of A . As before, non-negative and positive isotopies induce quasi-orders \leq and (if A is compact) \ll on $\mathcal{L}(A)$. A is called *strongly orderable* if \leq defines a partial order. For compact $A_0, A_1 \in \mathcal{L}(A)$, we define the interval $I_{A_0, A_1}^{\ll} := \{B \in \mathcal{L}(A) \mid A_0 \ll B \ll A_1\}$, and the collection of such intervals generates a topology \mathcal{O}_{\ll}^A on $\mathcal{L}(A)$.

The distances $\delta_\alpha^{\pm, A}$ also induce distances on $\mathcal{L}(A)$ via

$$(33) \quad \begin{aligned} \delta_\alpha^{\pm, A}(A_0, A_1) &:= \inf\{\delta_\alpha^\pm(id_M, \phi) \mid \phi \in \text{Cont}_0(M), \phi(A_0) = A_1\}, \quad A_0, A_1 \in \mathcal{L}(A), \\ d_\alpha^A(A_0, A_1) &:= \max\{\delta_\alpha^{+, A}(A_0, A_1), \delta_\alpha^{-, A}(A_0, A_1), 0\}. \end{aligned}$$

Also, $\delta_\alpha^{\pm, A}$ and d_α^A on $\mathcal{L}(A)$ satisfy the analogues of the properties in Propositions 2.1 and 2.3.

Remark 2.16. One can also define the pseudo-metric

$$(34) \quad \bar{d}_\alpha^A(A_0, A_1) := \inf\{d_\alpha(id_M, \phi) \mid \phi(A_0) = A_1\}$$

induced by d_α on $\mathcal{L}(A)$. It is easy to check that it satisfies the properties in Proposition 2.3 where we use the interval topology $\bar{\mathcal{O}}_{\ll}$ generated by sets of the form

$$(35) \quad I_{\phi_0, \phi_1}^{\ll}(A) := \{\phi(A) \mid \phi_0 \ll \phi \ll \phi_1\}, \quad \phi_0, \phi_1 \in \text{Cont}_0(M, \xi).$$

An analogous remark also applies to the spaces $\mathcal{L}(i)$, $\widetilde{\mathcal{L}}(A)$ and $\widetilde{\mathcal{L}}(i)$ considered below.

Similar constructions also work for *parametrized* subsets. Let

$$(36) \quad \mathcal{L}(i) := \{\phi \circ i : A \hookrightarrow M \mid \phi \in \text{Cont}_0(M, \xi)\}$$

denote the orbit of i under the action of $\text{Cont}_0(M, \xi)$. \leq and (if A is compact) \ll on $\mathcal{L}(i)$ are defined as before. If A is compact, we again define the intervals $I_{i_0, i_1}^{\ll} := \{j \in \mathcal{L}(i) \mid i_0 \ll j \ll i_1\}$ for $i_0, i_1 \in \mathcal{L}(i)$, and the collection of such intervals generates a topology \mathcal{O}_{\ll}^i on $\mathcal{L}(i)$.

Then $\delta_\alpha^{\pm, i}$ and d_α^i on $\mathcal{L}(i)$ induced by δ_α^\pm satisfy the analogues of the properties in Proposition 2.3 as well.

In particular, this means that if A is compact, the metric topologies of d_α^A and d_α^i agree with the interval topologies \mathcal{O}_{\ll}^A and \mathcal{O}_{\ll}^i . The proof for d_α^i if A is non-empty and the proof for d_α^A if A is non-empty and not equal to M differs only notationally from the proof for $\text{Cont}_0(M, \xi)$. If A is empty then $\mathcal{L}(i)$ and $\mathcal{L}(A)$ both only consist of a single point, and if $A = M$, $\mathcal{L}(A)$ only consists of a single point, so the assertion is satisfied trivially.

Following [RZ20] (see also [Ush15]), we show in Section 5 that d_α^A and d_α^i satisfy a version of Chekanov's dichotomy.

Theorem 2.17. *Let $i : L \hookrightarrow M^{2n+1}$ be a proper embedding of a submanifold $L \subseteq M$. If d_α^L is non-degenerate or if for all $i_0, i_1 \in \mathcal{L}(i)$ with $i_0(L) \neq i_1(L)$ we have that $d_\alpha^i(i_0, i_1) > 0$, then L is coisotropic. If $\dim L \leq n$, then d_α^L and d_α^i vanish identically unless L has a Legendrian component.*

If L is a connected Legendrian submanifold, then either d_α^L vanishes identically or d_α^L is non-degenerate, and either d_α^i vanishes identically or for any $i_0, i_1 \in \mathcal{L}(i)$ with $i_0(L) \neq i_1(L)$, we have that $d_\alpha^i(i_0, i_1) > 0$. If in addition L is compact, then d_α^L vanishes identically if and only if L admits a positive loop, and d_α^i vanishes identically if and only if i admits a positive loop.

As an immediate consequence, we get

Corollary 2.18. *Let $L \subseteq (M, \xi)$ be a closed, connected Legendrian. Then the interval topology \mathcal{O}_\ll^L on $\mathcal{L}(L)$ is Hausdorff if and only if L does not admit a positive loop. \square*

Remark 2.19. Note that if $i : L \hookrightarrow M$ is a proper Legendrian embedding, then d_α^i is never non-degenerate since isotopies of $i(L)$ are induced by ambient contact isotopies whose Hamiltonian vanishes along the image of $i(L)$. In this case, we can define an equivalence relation on $\mathcal{L}(i)$ where i_0 and i_1 are equivalent if and only if they are related by an isotopy of L which starts at the identity. Denote the set of equivalence classes by $\widehat{\mathcal{L}}(i)$. Then it is easy to see that d_α^i descends to a pseudo-metric \widehat{d}_α^i on $\widehat{\mathcal{L}}(i)$ satisfying properties (i) – (v) in Proposition 2.3. Similarly, the quasi-orders \leq and (if L is closed) \ll descend to quasi-orders on $\widehat{\mathcal{L}}(i)$, so we can again define an interval topology $\widehat{\mathcal{O}}_\ll^i$. As before, the metric topology induced by \widehat{d}_α^i agrees with $\widehat{\mathcal{O}}_\ll^i$ if L is closed.

Question 2.20. *Is \widehat{d}_α^i non-degenerate if it does not vanish identically?*

Question 2.20 seems to be hard to answer and its nature appears to be very different from the results discussed above. For example, assume that the answer is affirmative for the isotopy class of the zero-section in the 1-jet bundles J^1L of a manifold L . Then this would imply that there does not exist a compactly supported Legendrian isotopy $i_t : L \rightarrow J^1L$ starting at the embedding of the zero-section so that $i_1(L) = i_0(L)$, when i_1 and i_0 represent different elements of the mapping class group of L , since the energy of such an isotopy can be made arbitrarily small by conjugation with the contactomorphism $(q, p, z) \mapsto (q, \lambda p, \lambda z)$ for small $\lambda > 0$. Conversely, if the answer is no, then there has to exist such an isotopy i_t .

It is clear that if d_α^i vanishes identically then also d_α^L vanishes identically as $d_\alpha^i(i_0, i_1) \geq d_\alpha^L(i_0(L), i_1(L))$.

Question 2.21. *If d_α^L vanishes identically, then does d_α^i vanish identically as well?*

Remark 2.22. Note that for the parametrized and unparametrized Shelukhin-Chekanov-Hofer metrics the answer to the corresponding question is yes. Indeed, assume that for two fixed Legendrians $L_0 \neq L_1$ and any $\varepsilon > 0$, there exists $\phi_t \in \text{Cont}_0(M, \xi)$, $t \in [0, 1]$, with $\phi_0 = id_M$, $\phi_1(L_0) = L_1$, and $\|\phi_t\|_\alpha < \varepsilon$. Let $\psi_t \in \text{Cont}_0(M, \xi)$, $t \in [0, 1]$, $\psi_0 = id_M$, be so that $\psi_1(L_0) \neq L_0$ and $\text{supp } \psi_t \cap L_1 = \emptyset$. Then the concatenation $(\phi_t * (\psi_1 \circ \phi_{1-t}))|_{L_0}$ is well-defined and its energy is bounded by $(1 + \max \chi)\varepsilon$, where χ is defined by $\psi_1^* \alpha = \chi \alpha$. Taking $\varepsilon \rightarrow 0$ while fixing ψ_t , we obtain contact isotopies θ_t , $t \in [0, 1]$, starting at the identity which satisfy $\theta_1|_{L_0} = \psi_1|_{L_0}$ and have arbitrarily small energy. By the analogue of Theorem 2.17 for the Shelukhin-Chekanov-Hofer metric (which in the unparametrized case is proven in [RZ20] and for parametrized Legendrians follows by the same proof that we use in Section 5 for d_α^i), it follows that the parametrized metric vanishes identically as well.

This proof does however not work for d_α^L and d_α^i as we can only assume one-sided bounds on the generating Hamiltonian of ϕ_t , so the Hamiltonian of $\psi_1 \circ \phi_{1-t}$ may not be small.

Assume that $i_L \hookrightarrow M$ is a proper Legendrian embedding of a connected manifold L . Like d_α , d_α^L and d_α^i are bounded from above by the unparametrized or parametrized Shelukhin-Chekanov-Hofer metrics on $\mathcal{L}(L)$ and $\mathcal{L}(i)$, respectively.

Question 2.23. *If d_α^L (resp. d_α^i) does not vanish identically, then is it equivalent to the unparametrized (resp. parametrized) Shelukhin-Chekanov-Hofer metric?*

Example 2.24. (1-jets in J^1L) Consider the 1-jet bundle J^1L of a closed manifold L with the canonical contact form $\lambda = dz - pdq$. We identify L with the zero section in J^1L . Following Viterbo [Vit92], one defines a function $c_- : \mathcal{L}(L) \rightarrow \mathbb{R}$, which is non-decreasing along non-negative isotopies by Lemma 5.2 in [CN10a] and on 1-jets of functions $f : L \rightarrow \mathbb{R}$ it is equal to $c_-(j^1f) = \min f$ (see Example 3.1 in [CN20]). Using this, we can compute d_α^L on 1-jets. First note that if $i : L \rightarrow J^1L$ has a generating function S , then $\phi_t^\alpha \circ i$ has the generating function² $S + t$ where ϕ_t^α denotes the time- t Reeb flow on J^1L . From this it follows immediately that $c_-(\phi_t^\alpha(i(L))) = c_-(i(L)) + t$. Let $H_t : J^1L \rightarrow (-\varepsilon, \infty)$ be a Hamiltonian, and denote its associated contact isotopy by ϕ_t^H . Then $\phi_{\varepsilon t}^\alpha \circ \phi_t^H$ is positive. Thus, $c_-(\phi_t^H(L')) = c_-(\phi_{\varepsilon t}^\alpha(\phi_t^H(L'))) - \varepsilon \geq c_-(L') - \varepsilon$ for any $L' \in \mathcal{L}(L)$. This together with $c_-(j^1f) = \min f$ implies that $\delta_\alpha^{-,L}(L, j^1f) \geq -\min f$. Since the Hamiltonian $H(q, p, z) = f(q)$ generates the contact isotopy $(q, p, z) \mapsto (q, p + tdf, z + tf)$, we actually have equality

$$(37) \quad \delta_\alpha^{-,L}(L, j^1f) = -\min f.$$

Using right invariance, we obtain

$$(38) \quad \delta_\alpha^{+,L}(L, j^1f) = \delta_\alpha^{-,L}(j^1f, L) = \delta_\alpha^{-,L}(L, j^1(-f)) = -\min(-f) = \max f.$$

It follows that $d_\alpha^L(L, j^1f) = \max |f| = d_{SCH}(L, j^1f)$. Again using right invariance this implies that $d_\alpha^L(j^1f, j^1g) = \max |f - g| = d_{SCH}(j^1f, j^1g)$.

Remark 2.25. We can also consider the interval topologies $\tilde{\mathcal{O}}_\ll^i$ and $\tilde{\mathcal{O}}_\ll^A$ and the pseudo-metrics \tilde{d}_α^i and \tilde{d}_α^A on the universal covers $\tilde{\mathcal{L}}(i)$ and $\tilde{\mathcal{L}}(A)$ of $\mathcal{L}(i)$ and $\mathcal{L}(A)$, respectively. By the same proofs as before (see Remark 5.15) we see that the following holds.

Theorem 2.26. *Let $i : L \hookrightarrow M^{2n+1}$ be the embedding of a properly embedded submanifold $L \subseteq M$. If \tilde{d}_α^L on $\tilde{\mathcal{L}}(L)$ is π -non-degenerate or if for all $\iota_0, \iota_1 \in \tilde{\mathcal{L}}(i)$ with $\pi(\iota_0)(L) \neq \pi(\iota_1)(L)$ we have that $\tilde{d}_\alpha^L(\iota_0, \iota_1) > 0$, then L is coisotropic. If $\dim L \leq n$, then \tilde{d}_α^L and \tilde{d}_α^i vanish identically unless L has a Legendrian component.*

If L is a connected Legendrian submanifold, then either \tilde{d}_α^L vanishes identically or \tilde{d}_α^L is π -non-degenerate, and either \tilde{d}_α^i vanishes identically or for any $\iota_0, \iota_1 \in \tilde{\mathcal{L}}(i)$ with $\pi(\iota_0)(L) \neq \pi(\iota_1)(L)$, $\tilde{d}_\alpha^i(\iota_0, \iota_1) > 0$. If in addition L is compact, then \tilde{d}_α^L vanishes identically if and only if L admits a contractible positive loop, and \tilde{d}_α^i vanishes identically if and only if i admits a contractible positive loop. \square

Corollary 2.27. *Let $L \subseteq (M, \xi)$ be a closed, connected Legendrian. Then the interval topology \mathcal{O}_\ll^L on $\tilde{\mathcal{L}}(L)$ is π -Hausdorff if and only if L does not admit a contractible positive loop.* \square

2.4. Hedicke's Lorentzian distance function.

The definitions in [Hed21] of the Lorentzian distance functions τ_α and τ_α^A on $\text{Cont}_0(M, \xi)$ and $\mathcal{L}(A)$ for a closed Legendrian A extend naturally also to $\mathcal{L}(A)$ and $\mathcal{L}(i)$ for inclusions $i : A \rightarrow M$ of arbitrary closed subsets and to the universal covers $\widetilde{\text{Cont}}_0(M, \xi)$, $\tilde{\mathcal{L}}(A)$, and $\tilde{\mathcal{L}}(i)$ of these spaces. These will be denoted by τ_α^A , τ_α^i , $\tilde{\tau}_\alpha$, $\tilde{\tau}_\alpha^A$, and $\tilde{\tau}_\alpha^i$.

²Technically, $S + t$ is not quadratic at infinity, but this causes no issue (cf. the notion of equivalence of generating functions in [Vit92]). We could also cut-off t outside of a sufficiently large compact subset of the fibres of $L \times \mathbb{R}^N \rightarrow L$ where $S : L \times \mathbb{R}^N \rightarrow \mathbb{R}$, to obtain a generating function that is quadratic at infinity.

These different versions of the Lorentzian distance functions can be expressed in terms of the different versions of δ_α^+ . For this, recall the following lemma from [Hed21]³.

Lemma 2.28. *Let $H_t : M \rightarrow \mathbb{R}, t \in [0, 1]$, be a Hamiltonian on a closed contact manifold $(M, \ker \alpha)$. Then for any $\varepsilon > 0$ there exists another Hamiltonian G_t such that $[\phi_t^G] = [\phi_t^H] \in \widetilde{\text{Cont}}_0(M, \xi)$ and*

$$(39) \quad \min_M G_t \in \left(\int_0^1 \min_M H_s ds - \varepsilon, \int_0^1 \min_M H_s ds + \varepsilon \right).$$

With this lemma, we can rewrite the definitions of $\tilde{\tau}_\alpha$ and $\tilde{\delta}_\alpha^\pm$ if M is compact:

$$(40) \quad \begin{aligned} \tilde{\delta}_\alpha^-(f_0, f_1) &= \inf \left\{ - \int_0^1 \min_M H_t dt \mid f^H f_0 = f_1 \right\} \\ &= - \sup \left\{ \int_0^1 \min_M H_t dt \mid f^H f_0 = f_1 \right\} \\ \tilde{\delta}_\alpha^+(f_0, f_1) &= \inf \left\{ \int_0^1 \max_M H_t dt \mid f^H f_0 = f_1 \right\}, \\ \tilde{\tau}_\alpha(f_0, f_1) &= \max \left\{ \sup \left\{ \int_0^1 \min_M H_t dt \mid f^H f_0 = f_1 \right\}, 0 \right\}. \end{aligned}$$

In particular, we see that

$$(41) \quad \tilde{\tau}_\alpha = \max \{ -\tilde{\delta}_\alpha^-, 0 \},$$

using that $\tilde{\delta}_\alpha^-(f_0, f_1) < 0$ if and only if $f_0 \ll f_1$. The same formulas hold also for τ_α and δ_α^\pm . For $\tau_\alpha^A, \tau_\alpha^i, \delta_\alpha^{\pm, A}, \delta_\alpha^{\pm, i}, \tilde{\tau}_\alpha^A, \tilde{\tau}_\alpha^i, \tilde{\delta}_\alpha^{\pm, A},$ and $\tilde{\delta}_\alpha^{\pm, i}$, the corresponding formulas where the min/max is taken over A instead hold under the weaker assumption that the Reeb flow in (M, α) is complete, as is clear from the proof of Lemma 3.1 in [Hed21].

Using this observation, it follows that Theorem 2.7 and 5.8 in [Hed21] hold for the constant $C_\alpha = 1$ whenever (M, ξ) is closed and orderable, resp. when the Reeb flow is complete and the Legendrian is orderable.

Proposition 2.29. *If \tilde{d}_α is non-degenerate, then $\tilde{\delta}_\alpha^\pm$ is 1-Lipschitz with respect to \tilde{d}_α in both arguments. If in addition M is closed⁴, then also $\tilde{\tau}_\alpha$ is 1-Lipschitz with respect to \tilde{d}_α in both arguments. In particular, $\tilde{\tau}_\alpha(f_0, f_1) \leq \tilde{d}_\alpha(f_0, f_1) \leq \tilde{d}_{SH}(f_0, f_1)$ for any $f_0, f_1 \in \widetilde{\text{Cont}}_0(M, \xi)$.*

Proof. By (41) and property (ii) in Proposition 2.1, it is enough to prove the claim for $\tilde{\delta}_\alpha^-$. We only show that $\tilde{\delta}_\alpha^-$ is 1-Lipschitz in the second argument, the proof for the first argument is analogous.

For this, let $f, g, \tilde{g} \in \widetilde{\text{Cont}}_0(M, \xi)$ be so that $\tilde{d}_\alpha(g, \tilde{g}) = \varepsilon > 0$. Denote $a := \tilde{\delta}_\alpha^-(f, \tilde{g})$ and $b := \tilde{\delta}_\alpha^-(f, g)$. We will show that $|a - b| \leq \varepsilon$. Without loss of generality, we may assume that $a \leq b$. Then we see that

$$(42) \quad b = \tilde{\delta}_\alpha^-(f, g) \leq \tilde{\delta}_\alpha^-(f, \tilde{g}) + \tilde{\delta}_\alpha^-(\tilde{g}, g) \leq \tilde{\delta}_\alpha^-(f, \tilde{g}) + \tilde{d}_\alpha(\tilde{g}, g) = a + \varepsilon,$$

or equivalently, $|b - a| = b - a \leq \varepsilon$.

³The lemma in [Hed21] was stated in a weaker form not fixing the homotopy class, but it is clear from its proof that one may also assume that ϕ_t^G and $\phi_t^H, t \in [0, 1]$, are homotopic relative to the endpoints.

⁴If we apply the definition of $\tilde{\tau}_\alpha$ to open manifolds, then it necessarily vanishes identically. In particular, it is 1-Lipschitz.

The second part of the proposition is an immediate consequence of the first part since $\tilde{\tau}(f_0, f_0) = 0$ by the orderability of (M, ξ) . \square

The corresponding result applies with the same proof to the other variants of τ_α and δ_α^\pm .

Remark 2.30. The above results imply that for a closed, strongly orderable Legendrian L , $(\mathcal{L}(L), d_\alpha^L, \ll, \leq, \tau_\alpha^L)$ is a strongly causal Lorentzian pre-length space in the sense of [KS18]. However, it is in general not a Lorentzian length space. Indeed, any Lorentzian length space is by definition localizable (see [KS18] for the definitions). In particular, if $(\mathcal{L}(L), d_\alpha^L, \ll, \leq, \tau_\alpha^L)$ were a Lorentzian length space, every $L \in \mathcal{L}(L)$ would have an open neighbourhood U (with respect to d_α^L) together with a number $C > 0$ so that for any non-negative isotopy $L_t \subseteq U$, we have that $L^{d_\alpha^L}(L_t) < C$, where $L^{d_\alpha^L}(L_t)$ denotes the length of L_t viewed as a rectifiable curve in the metric space $(\mathcal{L}(L), d_\alpha^L)$. This is not satisfied already for the zero section in the 1-jet space of any closed manifold L . In fact, even any C^k -small neighbourhood of the zero section contains arbitrarily long non-negative isotopies. To see this, choose an open cover $\{U_i\}_{i=1}^N$, $N \in \mathbb{N}$, of L so that for all $j \in \{1, \dots, N\}$, $U_j \setminus \bigcup_{i \neq j} U_i$ is non-empty. Pick a partition of unity $\{\lambda_i\}_{i=1}^N$ subordinate to this cover. Note that $\lambda_i^{-1}(\{1\})$ is non-empty. Then for any $\varepsilon > 0$ and $k \in \mathbb{N}_0$, the non-negative Legendrian isotopies given by the 1-jets

$$(43) \quad j^1 \left(\varepsilon \left(t\lambda_j + \sum_{i=1}^{j-1} \lambda_i + k \right) \right), \quad j \in \{1, \dots, N\}, t \in [0, 1],$$

have length ε by Example 2.24 above. Fix some $\delta > 0$. Let $\varepsilon > 0$ be so small that for any $t \in [0, 1]$, $j^1(t\varepsilon)$ is $\frac{\delta}{2}$ -close to the zero section in the C^k -norm. Then for any sufficiently large number $K \in \mathbb{N}$, the Legendrian isotopies

$$(44) \quad j^1 \left(\frac{\varepsilon}{K} \left(t\lambda_j + \sum_{i=1}^{j-1} \lambda_i + k \right) \right), \quad j \in \{1, \dots, N\}, k \in \{0, \dots, K-1\}, t \in [0, 1],$$

are δ -close to the zero section in the C^k -norm. Then the Legendrian isotopy $L_t, t \in [0, NK]$, from $j^1 0$ to $j^1 \varepsilon$ which is equal to

$$(45) \quad j^1 \left(\frac{\varepsilon}{K} \left(t\lambda_j + \sum_{i=1}^{j-1} \lambda_i + k \right) \right)$$

for $t \in [j-1+kN, j+kN]$, $j \in \{1, \dots, N\}$, $k \in \{0, \dots, K-1\}$, is δ -close to the zero section in the C^k -norm and has length $\frac{\varepsilon}{K}NK = \varepsilon N$. Taking the limit $N \rightarrow \infty$ (while potentially increasing K as well), we find arbitrarily large non-negative isotopies in any C^k -neighbourhood of the zero section as claimed.

As any Legendrian admits a Weinstein neighbourhood, we expect that $(\mathcal{L}(L), d_\alpha^L, \ll, \leq, \tau_\alpha^L)$ is not localizable for any closed, strongly orderable Legendrian L in any contact manifold.

Similarly, for a closed, strongly orderable contact manifold (M, ξ) with contact form α , $(\text{Cont}_0(M, \xi), d_\alpha, \ll, \leq, \tau_\alpha)$ is a strongly causal Lorentzian pre-length space. Again, we expect it to be non-localizable. One reason is that the graph of a contactomorphism in $M \times M \times \mathbb{R}$ is Legendrian, and any Legendrian C^1 -close to a graph is again graphical, so the above construction performed in a Weinstein neighbourhood of the graph gives a contact isotopy of arbitrarily large energy C^k -close to the identity.

2.5. Reeb flow.

Generalizing Corollaries 2.4 and 2.5 in [Hed21], we observe that the Reeb flow is always a “length-minimizing geodesic” for $\tilde{\delta}_\alpha^\pm$ if (M, ξ) is orderable. In fact, as an immediate consequence of (9), the corresponding formula on $\text{Cont}_0(M, \xi)$, and the left-invariance of \ll , we have the following result.

Proposition 2.31. *Let $f \in \widetilde{\text{Cont}}_0(M, \xi)$, $\phi \in \text{Cont}_0(M, \xi)$, and assume that M is closed. If (M, ξ) is orderable, then*

$$(46) \quad \widetilde{\delta}_\alpha^\pm(\text{Id}_M, f_t^\alpha f) = \widetilde{\delta}_\alpha^\pm(\text{Id}_M, f) \pm t.$$

If (M, ξ) is strongly orderable,

$$(47) \quad \delta_\alpha^\pm(\text{Id}_M, \phi_t^\alpha \phi) = \delta_\alpha^\pm(\text{Id}_M, \phi) \pm t.$$

□

Similarly, the following holds.

Proposition 2.32. *Assume that the Reeb flow on (M, α) is complete. Let $i : A \rightarrow M$ be the embedding of a closed subset, and let $j \in \mathcal{L}(i)$, $\widetilde{j} \in \widetilde{\mathcal{L}}(i)$, $B \in \mathcal{L}(A)$, and $\widetilde{B} \in \widetilde{\mathcal{L}}(A)$. If i is orderable, then*

$$(48) \quad \widetilde{\delta}^{\pm, i}([i], f_t^\alpha \widetilde{j}) = \widetilde{\delta}^{\pm, i}([i], \widetilde{j}) \pm t.$$

If i is strongly orderable, then

$$(49) \quad \delta^{\pm, i}(i, \phi_t^\alpha \circ j) = \delta^{\pm, i}(i, j) \pm t.$$

If A is orderable, then

$$(50) \quad \widetilde{\delta}^{\pm, A}([A], f_t^\alpha \widetilde{B}) = \widetilde{\delta}^{\pm, A}([A], \widetilde{B}) \pm t.$$

If A is strongly orderable, then

$$(51) \quad \delta^{\pm, A}(A, \phi_t^\alpha(B)) = \delta^{\pm, A}(A, B) \pm t.$$

□

2.6. The relative growth of contactomorphisms.

The relative growth of two contact isotopies can be estimated in terms of generating Hamiltonians of the two isotopies.

Proposition 2.33. *Assume that M is compact and orderable, and let $f, g \in \widetilde{\text{Cont}}_0(M, \xi)$ with $f \gg \text{Id}_M$. If $\widetilde{\delta}_\alpha^+(\text{Id}_M, g) \geq 0$, then*

$$(52) \quad \gamma(f, g) \leq \frac{\widetilde{\delta}_\alpha^+(\text{Id}_M, g)}{\widetilde{\tau}_\alpha(\text{Id}_M, f)},$$

if $\widetilde{\delta}_\alpha^+(\text{Id}_M, g) < 0$, then

$$(53) \quad \gamma(f, g) \leq \frac{\widetilde{\delta}_\alpha^+(\text{Id}_M, g)}{\widetilde{\delta}_\alpha^+(\text{Id}_M, f)},$$

if $\widetilde{\delta}_\alpha^-(\text{Id}_M, g) < 0$, then

$$(54) \quad \gamma(f, g) \geq \frac{\widetilde{\tau}_\alpha(\text{Id}_M, g)}{\widetilde{\delta}_\alpha^+(\text{Id}_M, f)},$$

and if $\widetilde{\delta}_\alpha^-(\text{Id}_M, g) \geq 0$, then

$$(55) \quad \gamma(f, g) \geq \frac{-\widetilde{\delta}_\alpha^-(\text{Id}_M, g)}{\widetilde{\tau}_\alpha(\text{Id}_M, f)}.$$

Proof. Let F_t and G_t denote generating Hamiltonians of f and g so that $\int_0^1 \min_M F_t dt > 0$. Assume that for $k \in \mathbb{N}$ and $p \in \mathbb{Z}_{\geq 0}$, we have that

$$(56) \quad \frac{p}{k} > \frac{\int_0^1 \max_M G_t dt}{\int_0^1 \min_M F_t dt}.$$

Then $f^{-p}g^k$ can be generated by a Hamiltonian H_t which satisfies

$$(57) \quad \int_0^1 \max_M H_t dt = k \int_0^1 \max_M G_t dt - p \int_0^1 \min_M F_t dt < 0,$$

i.e. $Id_M \gg f^{-p}g^k$, or $f^p \gg g^k$. This shows that

$$(58) \quad \gamma(f, g) \leq \max \left\{ 0, \inf_{F, G} \frac{\int_0^1 \max_M G_t dt}{\int_0^1 \min_M F_t dt} \right\},$$

which in the case that $\tilde{\delta}_\alpha^+(Id_M, g) \geq 0$, means that

$$(59) \quad \gamma(f, g) \leq \frac{\tilde{\delta}_\alpha^+(Id_M, g)}{\tilde{\tau}_\alpha(Id_M, f)}.$$

Now assume that $\tilde{\delta}_\alpha^+(Id_M, g) < 0$, or equivalently $g \ll Id_M$, and that for $k \in \mathbb{N}$ and $p \in \mathbb{Z}_{<0}$,

$$(60) \quad \frac{p}{k} > \frac{\int_0^1 \max_M G_t dt}{\int_0^1 \max_M F_t dt}.$$

Then $f^{-p}g^k$ can be generated by a Hamiltonian H_t which satisfies

$$(61) \quad \int_0^1 \max_M H_t dt = k \int_0^1 \max_M G_t dt - p \int_0^1 \max_M F_t dt < 0,$$

which implies that $f^p \gg g^k$. It follows that

$$(62) \quad \gamma(f, g) \leq \inf_{F, G} \frac{\int_0^1 \max_M G_t dt}{\int_0^1 \max_M F_t dt},$$

which implies that

$$(63) \quad \gamma(f, g) \leq \frac{\tilde{\delta}_\alpha^+(Id_M, g)}{\tilde{\delta}_\alpha^+(Id_M, f)}.$$

On the other hand assume that for some $p \in \mathbb{Z}_{\geq 0}$ and $k \in \mathbb{N}$, we have that

$$(64) \quad \frac{p}{k} < \frac{\int_0^1 \min_M G_t dt}{\int_0^1 \max_M F_t dt}.$$

Then $g^{-k}f^p$ can be generated by a Hamiltonian H_t that satisfies

$$(65) \quad \int_0^1 \max_M H_t dt = p \int_0^1 \max_M F_t dt - k \int_0^1 \min_M G_t dt < 0,$$

which implies that $f^p \ll g^k$. This shows that

$$(66) \quad \gamma(f, g) \geq \sup_{F, G} \frac{\int_0^1 \min_M G_t dt}{\int_0^1 \max_M F_t dt}$$

whenever $\int_0^1 \min_M G_t dt > 0$, which means that

$$(67) \quad \gamma(f, g) \geq \frac{\tilde{\tau}_\alpha(Id_M, g)}{\tilde{\delta}_\alpha^+(Id_M, f)}.$$

If

$$(68) \quad \frac{p}{k} < \frac{\int_0^1 \min_M G_t dt}{\int_0^1 \min_M F_t dt}.$$

for some $p \in \mathbb{Z}_{<0}$ and $k \in \mathbb{N}$, then $g^{-k} f^p$ can be generated by a Hamiltonian H_t that satisfies

$$(69) \quad \int_0^1 \max_M H_t dt = p \int_0^1 \min_M F_t dt - k \int_0^1 \min_M G_t dt < 0,$$

which again implies that $f^p \ll g^k$. It follows that

$$(70) \quad \gamma(f, g) \geq \min \left\{ 0, \sup_{F, G} \frac{\int_0^1 \min_M G_t dt}{\int_0^1 \min_M F_t dt} \right\},$$

where the supremum is taken over all generating Hamiltonians so that $\int_0^1 \min_M F_t dt > 0$. In the case that $\tilde{\delta}_\alpha^-(Id_M, g) \geq 0$ this implies that

$$(71) \quad \gamma(f, g) \geq \frac{-\tilde{\delta}_\alpha^-(Id_M, g)}{\tilde{\tau}_\alpha(Id_M, f)}.$$

□

For autonomous Hamiltonians F, G on ST^*T^n which depend only on the coordinate on the fibre (see Example 2.9 above) Eliashberg and Polterovich [EP00] showed that

$$(72) \quad \gamma(f_t^F, f_t^G) = \max_{p \in S^{n-1}} \frac{G(p)}{F(p)},$$

whenever $F(p) > 0$ for all p and $G(p) > 0$ for some p . In fact, this remains true even if $G(p) \leq 0$ for all p . Indeed, $(f_t^G)^k$ is generated by kG and $(f_t^F)^q$ is generated by qF , so by the characterization of \ll given in Example 2.9, $(f_t^F)^q \gg (f_t^G)^k$ if and only if $qF(p) > kG(p)$ for all $p \in S^{n-1}$.

In this case the above inequalities, say in the case that $G > 0$ (i.e. $\tilde{\delta}^+(Id_M, g) > 0$ and $\tilde{\delta}^-(Id_M, g) < 0$), read

$$(73) \quad \frac{\min G}{\max F} \leq \max \frac{G}{F} \leq \frac{\max G}{\min F},$$

so we see that the inequalities cannot be replaced by equalities in general.

If on the other hand $f = f_t^\alpha, t > 0$, and $g = f_s^\alpha, s \in \mathbb{R}$, then in view of Proposition 2.31 the inequalities become

$$(74) \quad \frac{s}{t} \leq \gamma(f_t^\alpha, f_s^\alpha) \leq \frac{s}{t},$$

which means that they are, in fact, equalities.

3. REPARAMETRIZATIONS OF HAMILTONIANS

In this section we show the following lemma, which allows us to use reparametrizations of contact isotopies to get bounds on the maximum or minimum of the generating Hamiltonian.

Lemma 3.1. *Let $\lambda : [0, 1] \rightarrow [0, \infty)$ be a continuous function, and let $\varepsilon > 0$. Then there exists a diffeomorphism $\sigma : [0, 1] \rightarrow [0, 1]$ with $\sigma(0) = 0$ so that*

$$(75) \quad \int_0^1 \lambda(t) dt + \varepsilon > \max_t (\lambda(\sigma(t))\sigma'(t)).$$

Proof. Denote $l := \int_0^1 \lambda(t) dt$. For $\delta > 0$, let $U := \{t \in [0, 1] \mid \frac{\lambda(t)}{l} < \delta\}$. Let $\phi : [0, \infty) \rightarrow [0, \infty)$ be a smooth function so that $\phi(x) = x$ for $x \leq \frac{1}{\delta}$ and $\phi(x) \in [\frac{1}{\delta}, \frac{1}{\delta} + 1]$ for $x > \frac{1}{\delta}$.

There exists a number $\tau > 0$ and a diffeomorphism $\tilde{\sigma} : [0, \tau] \rightarrow [0, 1]$, $\tau > 0$, which are uniquely determined by the differential equation $\tilde{\sigma}'(t) = \phi(\frac{l}{\lambda(\tilde{\sigma}(t))}) \geq \min\{\frac{1}{\delta}, \frac{l}{\max \lambda}\} > 0$ with the initial condition $\tilde{\sigma}(0) = 0$.

Note that

$$(76) \quad \int_0^1 \frac{1}{\phi(\frac{l}{\lambda(t)})} dt = \int_0^\tau \frac{1}{\phi(\frac{l}{\lambda(\tilde{\sigma}(t))})} \tilde{\sigma}'(t) dt = \tau.$$

Note further that

$$(77) \quad \begin{aligned} \tau &= \int_0^1 \frac{1}{\phi(\frac{l}{\lambda(t)})} dt = \int_{[0,1] \setminus U} \frac{\lambda(t)}{l} dt + \int_U \frac{1}{\phi(\frac{l}{\lambda(t)})} dt \\ &= 1 + \int_U \frac{1}{\phi(\frac{l}{\lambda(t)})} - \frac{\lambda(t)}{l} dt \in \left[1 - \delta \int_U 1 dt, 1 + \delta \int_U 1 dt \right] \end{aligned}$$

by definition of U and ϕ . In particular, $\tau \rightarrow 1$ as $\delta \rightarrow 0$. Now we define the diffeomorphism $\sigma : [0, 1] \rightarrow [0, 1]$, $\sigma(t) = \tilde{\sigma}(\frac{t}{\tau})$. Then it follows that $\sigma'(t) = \frac{1}{\tau} \phi(\frac{l}{\lambda(\sigma(t))})$, which implies that $\lambda(\sigma(t))\sigma'(t) = \lambda(\sigma(t))\frac{1}{\tau} \phi(\frac{l}{\lambda(\sigma(t))})$. If $\sigma(t) \in [0, 1] \setminus U$, this is equal to $\frac{l}{\tau}$, and if $\sigma(t) \in U$, $\lambda(\sigma(t)) \in [0, \delta l)$ and $\phi(\frac{l}{\lambda(\sigma(t))}) \in [\frac{1}{\delta}, \frac{1}{\delta} + 1]$ which implies that $\lambda(\sigma(t))\sigma'(t) \in [0, (1 + \delta)\frac{l}{\tau})$. In particular, for any $\varepsilon > 0$, $\lambda(\sigma(t))\sigma'(t) < l + \varepsilon$ for all $t \in [0, 1]$ if δ is sufficiently small. \square

4. PROOFS OF THEOREMS 2.6 AND 2.11

4.1. Proof of the first part of Theorem 2.6.

To prove the first part of Theorem 2.6, assume that \tilde{d}_α is not π -non-degenerate, i.e. by right-invariance of \tilde{d}_α , there exists $h \in \widetilde{\text{Cont}}_0(M, \xi)$, $\pi(h) \neq id_M$, with $\tilde{d}_\alpha(h, id_M) = 0$. Let $U \subseteq M$ be a open subset so that $\pi(h)(U) \cap U = \emptyset$. Using an argument by Eliashberg and Polterovich [EP93] (see also [She17], Remark 7), it is straightforward to show that for any $f \in \widetilde{\text{Cont}}_0(M, \xi)$ with support in U , $\tilde{d}_\alpha(f, id_M) = 0$. By the naturality of \tilde{d}_α under conjugation, there exists for any point $x \in M$ a neighbourhood U_x and $h_x \in \widetilde{\text{Cont}}_0(M, \xi)$ so that $\tilde{d}_\alpha(h_x, id_M) = 0$ and $\pi(h_x)(U_x) \cap U_x = \emptyset$. By the fragmentation lemma for contact isotopies (see [Ban97]) and right-invariance, it then follows that \tilde{d}_α vanishes identically.

To prove the claim, let h and U be as above. We first show that \tilde{d}_α vanishes on commutators of elements in $\widetilde{\text{Cont}}_0(U, \xi)$. Let $f_1, f_2 \in \widetilde{\text{Cont}}_0(U, \xi)$. We define $g := [h, f_1^{-1}] := f_1 h^{-1} f_1^{-1} h$ as the commutator of h and f_1^{-1} . Then an easy argument shows that $[g, f_2] = [f_1, f_2]$. Furthermore,

$$(78) \quad \begin{aligned} \tilde{d}_\alpha([g, f_2], id_M) &= \tilde{d}_\alpha(f_2^{-1} g^{-1} f_2 g, id_M) \leq \tilde{d}_\alpha(f_2^{-1} g^{-1} f_2, id_M) + \tilde{d}_\alpha(g, id_M) \\ &\leq (1 + C(\pi(f_2^{-1}))) \tilde{d}_\alpha(g, id_M) = (1 + C(\pi(f_2^{-1}))) \tilde{d}_\alpha(f_1 h^{-1} f_1^{-1} h, id_M) \\ &\leq (1 + C(\pi(f_2^{-1}))) (1 + C(\pi(f_1))) \tilde{d}_\alpha(h, id_M) = 0, \end{aligned}$$

where for any contactomorphism $\psi \in \text{Cont}_0(M, \alpha)$, $C(\psi) := \max \chi$, where $\chi : M \rightarrow (0, \infty)$ denotes the conformal factor of ψ defined via $\psi^* \alpha = \chi \alpha$. In particular, $\tilde{d}_\alpha([f_1, f_2], id_M) = 0$. Since $\widetilde{\text{Cont}}_0(U, \xi)$ is perfect (see [Ryb10, Appendix]), it follows that $\tilde{d}_\alpha(-, id_M)$ vanishes on the image of $\widetilde{\text{Cont}}_0(U, \xi)$ in $\widetilde{\text{Cont}}_0(M, \xi)$, as claimed. \square

Remark 4.1. Above we proved that for any $f_1, f_2, h \in \widetilde{\text{Cont}}_0(M, \xi)$ so that

$$(79) \quad (\text{supp}(f_1) \cup \text{supp}(f_2)) \cap \pi(h)(\text{supp}(f_1) \cup \text{supp}(f_2)) = \emptyset,$$

we have that

$$(80) \quad \widetilde{d}_\alpha([f_1, f_2], id_M) \leq (1 + C(\pi(f_2^{-1}))) (1 + C(\pi(f_1))) \widetilde{d}_\alpha(h, id_M).$$

Let $U \subseteq M$ be so that $\pi(h)(U) \cap U = \emptyset$. Taking the infimum over all h with fixed $\phi := \pi(h)$, we obtain that

$$(81) \quad \widetilde{d}_\alpha([f_1, f_2], id_M) \leq (1 + C(\pi(f_2^{-1}))) (1 + C(\pi(f_1))) \widehat{d}_\alpha(\phi, id_M).$$

If $\widehat{d}_\alpha(\phi, id_M) = 0$, then it follows as above that \widetilde{d}_α vanishes identically.

4.2. Proof of the first part of Theorem 2.11.

The first part of Theorem 2.11 could be proven in the same way since also $\text{Cont}_0(M, \xi)$ is perfect, but $\widetilde{\text{Cont}}_0(M, \xi)$ is, in fact, simple (see [Ryb10], Corollary 1.2), so the claim also follows immediately from the observation that $\{\phi \in \text{Cont}_0(M, \xi) \mid d_\alpha(Id_M, \phi) = 0\}$ is a normal subgroup of $\text{Cont}_0(M, \xi)$. \square

4.3. Proof of the second part of Theorems 2.6 and 2.11.

To prove the last part of Theorem 2.6, assume that M is compact.

Denote the Reeb flow on (M, α) by ϕ_t^α and its lift to $\widetilde{\text{Cont}}_0(M, \xi)$ by f_t^α . If \widetilde{d}_α vanishes identically on $\widetilde{\text{Cont}}_0(M, \xi)$, then there exists a Hamiltonian $H_t : M \rightarrow (-1, \infty)$ so that $[\phi_t^H] = f_{-1}^\alpha$, where ϕ_t^H denotes the contact isotopy associated to H . Then the contractible loop of contactomorphisms $\phi_t^\alpha \circ \phi_t^H$ is generated by the Hamiltonian $H_t + 1$ which is positive.

Conversely, let $\phi_t : M \rightarrow M, t \in [0, 1]$, be a contractible positive loop of contactomorphisms. Without loss of generality, we may assume that $\phi_0 \neq \phi_{\frac{1}{2}}$. Then $\widetilde{d}_\alpha([\phi_0], [\phi_{\frac{1}{2}}]) = 0$ since there exist both positive and negative contact isotopies from $[\phi_0]$ to $[\phi_{\frac{1}{2}}]$. By what we already proved above, this implies that \widetilde{d}_α vanishes identically. \square

The proof of the second part of Theorem 2.11 works in the same way, except that we do not keep track of the homotopy classes of contact isotopies and positive loops. \square

5. PROOF OF THEOREM 2.17

In this section, we prove Theorem 2.17. For the most part, the proof follows the arguments in [RZ20] and is an adaption of Appendix D of the author's master's thesis [Nak19] to the present setting. In fact, the degeneracy of the unparametrized metric d_α^A for non-coisotropic submanifolds and the vanishing of the parametrized and unparametrized metrics follow immediately from the corresponding result for the Shelukhin-Chekanov-Hofer metric that were proven in [RZ20] and [DRS22], and the dichotomy for unparametrized Legendrians is a special case of Lemma 7 in [Can23], but since the literature does not cover all of the different cases we consider here, we include a proof for completeness.

Let $i : A \hookrightarrow M$ be the inclusion of a closed subset. We write

$$(82) \quad \Sigma_A := \{\phi \in \text{Cont}_0(M, \xi) \mid \phi(A) = A\}$$

and

$$(83) \quad \Sigma_i := \{\phi \in \text{Cont}_0(M, \xi) \mid \phi \circ i = i\}$$

for the stabilizer subgroups of A and i .

Proposition 5.1. *The closure of Σ_A with respect to the topology on $\text{Cont}_0(M, \xi)$ induced by d_α is given by*

$$(84) \quad \overline{\Sigma}_A = \{\phi \in \text{Cont}_0(M, \xi) \mid d_\alpha^A(A, \phi(A)) = 0\}.$$

The closure of Σ_i with respect to the topology on $\text{Cont}_0(M, \xi)$ induced by d_α is given by

$$(85) \quad \overline{\Sigma}_i = \{\phi \in \text{Cont}_0(M, \xi) \mid d_\alpha^i(i, \phi \circ i) = 0\}.$$

Proof. Let $\phi \in \overline{\Sigma}_A$ and let $\varepsilon > 0$ be a number. Then there exists $\psi \in \Sigma_A$ with $d_\alpha(\phi\psi^{-1}, id_M) = d_\alpha(\phi, \psi) < \varepsilon$. Since $\psi(A) = A$ and thus also $\psi^{-1}(A) = A$, we see that

$$(86) \quad d_\alpha^A(A, \phi(A)) = d_\alpha^A(A, \phi(\psi^{-1}(A))) < \varepsilon.$$

Because $\varepsilon > 0$ was arbitrary, it follows that $d_\alpha^A(A, \phi(A)) = 0$.

Conversely, let $\phi \in \text{Cont}_0(M, \xi)$ be a contactomorphism with $d_\alpha^A(A, \phi(A)) = 0$, i.e. given any number $\varepsilon > 0$ there exists a contactomorphism $\psi \in \text{Cont}_0(M, \xi)$ with $\psi(A) = \phi(A)$ and $d_\alpha(id_M, \psi) < \varepsilon$. Thus, $\psi^{-1}\phi \in \Sigma_A$ and

$$(87) \quad d_\alpha(\psi^{-1}\phi, \phi) = d_\alpha(\psi^{-1}, id_M) = d_\alpha(\psi, id_M) < \varepsilon.$$

Since $\varepsilon > 0$ was arbitrary, it follows that $\phi \in \overline{\Sigma}_A$.

The proof of the second part of the proposition is only notationally different. \square

In particular, this implies that $\overline{\Sigma}_A$ and $\overline{\Sigma}_i$ are groups as the closure of any subgroup of a topological group is again a subgroup.

We define the rigid loci of A and i by

$$(88) \quad R_A := \bigcap_{\phi \in \overline{\Sigma}_A} \phi^{-1}(A)$$

and

$$(89) \quad R_i := \bigcap_{\phi \in \overline{\Sigma}_i} \phi^{-1}(A),$$

which both are closed subsets of A .

For any open subset $U \subseteq M$, $\text{Cont}_U(M, \xi)$ denotes the subgroup of all contactomorphisms $\phi \in \text{Cont}_0(M, \xi)$ that are generated by a Hamiltonian having support in U .

Proposition 5.2. *We have*

$$(90) \quad \text{Cont}_{M \setminus R_A}(M, \xi) \subseteq \overline{\Sigma}_A$$

and

$$(91) \quad \text{Cont}_{M \setminus R_i}(M, \xi) \subseteq \overline{\Sigma}_i$$

Proof. Let $x \in M \setminus R_A$ be a point. By definition of R_A there exists a contactomorphism $\phi \in \overline{\Sigma}_A$ with $\phi(x) \notin A$. Since A is closed, there exists a neighbourhood U_x of x with $\phi(U_x) \cap A = \emptyset$. Then it follows that

$$(92) \quad \phi \text{Cont}_{U_x}(M, \xi) \phi^{-1} = \text{Cont}_{\phi(U_x)}(M, \xi) \subseteq \overline{\Sigma}_A.$$

Because $\overline{\Sigma}_A$ is a subgroup of $\text{Cont}_0(M, \xi)$ and $\phi \in \overline{\Sigma}_A$, we see that $\text{Cont}_{U_x}(M, \xi) \subseteq \overline{\Sigma}_A$. Note that the sets U_x , $x \in M \setminus R_A$, cover $M \setminus R_A$. By the fragmentation lemma (see [Ban97]), $\text{Cont}_{M \setminus R_A}(M, \xi)$ is generated by $\bigcup_{x \in M \setminus R_A} \text{Cont}_{U_x}(M, \xi)$ which proves that $\text{Cont}_{M \setminus R_A}(M, \xi) \subseteq \overline{\Sigma}_A$.

The proof of the second inclusion is only notationally different. \square

Remark 5.3. Proposition 5.2 implies that if $R_A = \emptyset$ (resp. $R_i = \emptyset$), then $\overline{\Sigma}_A = \text{Cont}_0(M, \xi)$ (resp. $\overline{\Sigma}_i = \text{Cont}_0(M, \xi)$) which means that d_α^A (resp. d_α^i) vanishes identically by Proposition 5.1.

On the other hand, it is easy to check that $R_A = A$ if and only if d_α is non-degenerate on $\mathcal{L}(A)$. Indeed, we see that

$$(93) \quad \begin{aligned} R_A = A &\Leftrightarrow \forall \phi \in \overline{\Sigma}_A : A \subseteq \phi^{-1}(A) \quad (\Leftrightarrow \phi(A) \subseteq A) \\ &\Leftrightarrow \forall \phi \in \overline{\Sigma}_A : A = \phi(A) \\ &\Leftrightarrow d_\alpha^A \text{ is non-degenerate on } \mathcal{L}(A), \end{aligned}$$

where we used that for any $\phi \in \overline{\Sigma}_A$ also $\phi^{-1} \in \overline{\Sigma}_A$ in order to get the equivalence between the first and the second line. The equivalence between the second and the third line follows from Proposition 5.1.

Similarly,

$$(94) \quad \begin{aligned} R_i = A &\Leftrightarrow \phi \in \overline{\Sigma}_i : A \subseteq \phi^{-1}(A) \\ &\Leftrightarrow \forall \phi \in \overline{\Sigma}_i : A = \phi(A) \\ &\Leftrightarrow \forall j \in \mathcal{L}(i), j(A) \neq A : d_\alpha^i(i, j) > 0 \\ &\Leftrightarrow \forall i_0, i_1 \in \mathcal{L}(i), i_0(A) \neq i_1(A) : d_\alpha^i(i_0, i_1) > 0. \end{aligned}$$

Proposition 5.2 shows that a contactomorphism in $\text{Cont}_0(M, \xi)$ that is equal to the identity in a neighbourhood of R_A (resp. R_i) lies in $\overline{\Sigma}_A$ (resp. $\overline{\Sigma}_i$). But for this to be true we actually only have to demand that the contactomorphism is generated by a contact Hamiltonian that vanishes along R_A (resp. R_i), as the following proposition shows:

Proposition 5.4. *Let $H \in C_c^\infty(M \times [0, 1])$ be a compactly supported Hamiltonian. If $H|_{R_A \times [0, 1]} \equiv 0$, then its contact Hamiltonian flow ϕ_t^H satisfies $\phi_s^H \in \overline{\Sigma}_A$ for all $s \in [0, 1]$.*

If $H|_{R_i \times [0, 1]} \equiv 0$, then $\phi_s^H \in \overline{\Sigma}_i$ for all $s \in [0, 1]$.

Proof. If we define $H^s(t, x) := sH(st, x)$, then $\phi_1^{H^s} = \phi_s^H$. Therefore, it will be enough to prove the proposition in the case that $s = 1$.

Let $H_t : M \rightarrow \mathbb{R}$ be a function with $H_t|_{R_A} \equiv 0$. For any number $n \in \mathbb{N}$, let $f_n : \mathbb{R} \rightarrow \mathbb{R}$ be a non-decreasing function such that $f_n(x) = x$ if $|x| \geq \frac{2}{n}$ and $f_n(x) = 0$ if $|x| \leq \frac{1}{n}$. Then $\phi_1^{f_n \circ H}$ converges to ϕ_1^H in the topology defined by d_α since $\phi_{-t}^H \phi_t^{f_n \circ H}$ is generated by the Hamiltonian $\chi_t(f_n \circ H_t - H_t) \circ \phi_t^H$, where χ_t is the positive function defined by $(\phi_t^H)^* \alpha = \chi_t \alpha$. Because H_t vanishes on R_A for all $t \in [0, 1]$, $f_n \circ H_t$ vanishes in a neighbourhood of R_A and hence,

$$(95) \quad \phi_1^{f_n \circ H} \in \text{Cont}_{M \setminus R_A}(M, \xi) \subseteq \overline{\Sigma}_A$$

by Proposition 5.2. As $\overline{\Sigma}_A$ is closed, it follows that also $\phi_1^H \in \overline{\Sigma}_A$.

The proof of the second part is the same. □

In the following, we will only consider autonomous contact Hamiltonians $H \in C_c^\infty(M)$. For any closed subset $B \subseteq M$ we define

$$(96) \quad I_B := \{H \in C_c^\infty(M) \mid H|_B = 0\}.$$

Let $\{\cdot, \cdot\}$ denote the Lie bracket defined by

$$(97) \quad \begin{aligned} \{F, G\} &:= \alpha([X_F, X_G]) \\ &= (dG)(X_F) - (dF)(X_G) - d\alpha(X_F, X_G) \\ &= (dG)(X_F) - (dF)(R_\alpha)G. \end{aligned}$$

for $F, G \in C_c^\infty(M)$, where X_F and X_G denote the contact vector fields associated to F and G , respectively.

Corollary 5.5. $I_{R_A}, I_{R_i} \subseteq C_c^\infty(M)$ are closed under $\{\cdot, \cdot\}$.

Proof. Let $F, G \in I_{R_A}$. It follows immediately from the definition of R_A that R_A is preserved by all elements of $\overline{\Sigma}_A$. By Proposition 5.4 we know that $\phi_t^F \in \overline{\Sigma}_A$ and thus $\phi_t^F(R_A) = R_A$ for all $t \in [0, 1]$. Because $G \in I_{R_A}$, this implies that $G \circ \phi_t^F|_{R_A} \equiv 0$ for all $t \in [0, 1]$. Hence, for any $x \in R_A$ we have

$$(98) \quad \{F, G\}(x) = (dG)(X_F)(x) = \left. \frac{d}{dt} \right|_{t=0} (G \circ \phi_t^F)(x) = 0.$$

The proof for I_{R_i} is the same. \square

As the following lemma shows, Corollary 5.5 restricts the possible structures of the rigid locus.

Lemma 5.6. [RZ20] *If $N \subseteq M$ is a submanifold and $x \in N$ is a point, then there exist functions $F, G \in I_N$ with $\{F, G\}(x) \neq 0$ if and only if*

$$(99) \quad (T_x N \cap \xi_x)^{\perp_{d\alpha}} \not\subseteq T_x N \cap \xi_x$$

is satisfied. \square

Remark 5.7. If $\dim(M) = 2n + 1$ and A is a submanifold of M of dimension $\dim(A) = n$, then the condition in the lemma is equivalent to A being non-Legendrian at x .

As an immediate consequence of Corollary 5.5 and Lemma 5.6 we obtain the following.

Corollary 5.8. *If $i : A \hookrightarrow M$ is the inclusion of a properly embedded submanifold and R_A (resp. R_i) is a union of components of A , then*

$$(100) \quad (T_x A \cap \xi_x)^{\perp_{d\alpha}} \subseteq T_x A \cap \xi_x$$

for all $x \in R_A$ (resp. for all $x \in R_i$). \square

Combined with Remark 5.3, this implies the first assertion of Theorem 2.17:

Corollary 5.9. *If $i : A \hookrightarrow M$ is the inclusion of a properly embedded submanifold and d_α^A is non-degenerate or for any $i_0, i_1 \in \mathcal{L}(i)$ with $i_0(A) \neq i_1(A)$ we have that $d_\alpha^i(i_0, i_1) > 0$, then*

$$(101) \quad (T_x A \cap \xi_x)^{\perp_{d\alpha}} \subseteq T_x A \cap \xi_x$$

for all $x \in A$. \square

Remark 5.10. Submanifolds satisfying the condition in the corollary are also called *coisotropic* (see [Hua15]).

Corollary 5.11. *Let $x \in R_A$ be a point and let $F_1, \dots, F_k \in I_{R_A}$ (resp. $F_1, \dots, F_k \in I_{R_i}$) be functions such that $dF_1|_{\xi_x}, \dots, dF_k|_{\xi_x} \in \xi_x^*$ are linearly independent. Then the map*

$$(102) \quad \begin{aligned} \psi : \mathbb{R}^k &\rightarrow M \\ (a_1, \dots, a_k) &\mapsto \phi_1^{\sum_i a_i F_i}(x) \end{aligned}$$

has image contained in R_A (resp. R_i) and its restriction to a sufficiently small ball around the origin is an embedding.

Proof. Because $\sum_i a_i F_i \in I_{R_A}$, we see that ψ has image contained in R_A (cf. the proof of Corollary 5.5).

The differential of ψ at $0 \in \mathbb{R}^k$ maps the standard basis vectors e_1, \dots, e_k to the vectors $X_{F_1}(x), \dots, X_{F_k}(x)$. The assumption on the $dF_i|_{\xi_x}$ implies that these vectors are linearly independent. Thus, the restriction of ψ to a sufficiently small neighbourhood of the origin is an embedding.

The proof for R_i is the same. \square

Corollary 5.12. *Let $i : A \hookrightarrow M^{2n+1}$ be the inclusion of a closed subset of M with $R_A \neq \emptyset$ (resp. $R_i \neq \emptyset$) and let $L \subseteq M$ be a properly embedded submanifold. If $\dim(L) < n$, then R_A (resp. R_i) is not contained in L and if $\dim(L) = n$ and R_A is contained in L , then R_A (resp. R_i) is a non-empty union of connected components of L .*

Proof. Suppose that $R_A \subseteq L$ and denote $k := \dim(M) - \dim(L)$. Let $x \in R_A$ be a point. Using a coordinate chart around x in which L looks like $\{0\} \times \mathbb{R}^{\dim(M)-k}$ we can find functions F_1, \dots, F_k that vanish on L such that $dF_1(x), \dots, dF_k(x) \in T_x^*M$ are linearly independent. This means that the linear map

$$(103) \quad (dF_1(x), \dots, dF_k(x)) : T_x M \rightarrow \mathbb{R}^k$$

has rank k which implies that the linear map

$$(104) \quad (dF_1, \dots, dF_k)|_{\xi_x} : \xi_x \rightarrow \mathbb{R}^k$$

has at least rank $k - 1$. After reordering the F_i we may assume that the linear map

$$(105) \quad (dF_1, \dots, dF_{k-1})|_{\xi_x} : \xi_x \rightarrow \mathbb{R}^k$$

has rank $k - 1$. Corollary 5.11 now implies that we can embed a $(k - 1)$ -dimensional ball into $R_A \subseteq L$.

In the case that $\dim(L) < n$ this immediately gives a contradiction since

$$(106) \quad k - 1 = 2n + 1 - \dim(L) - 1 > n > \dim(L).$$

If $\dim(L) = n$, then $k - 1 = \dim(L)$ and we have shown that a ball-shaped neighbourhood of $x \in L$ is contained in R_A . This means that $R_A \subseteq L$ is open and since it is also closed and non-empty, the corollary follows.

The proof for R_i is the same. \square

Corollary 5.9 and Remark 5.7 already show that d_α^A is degenerate and that d_α^i is “degenerate on embeddings with different images” if $A \subseteq M^{2n+1}$ is a non-Legendrian submanifold of dimension n . But an even stronger statement holds, which gives the second and third assertion of Theorem 2.17:

Corollary 5.13. *Let $i : A \subseteq M^{2n+1}$ be the inclusion of a properly embedded submanifold of dimension $\leq n$. If A has no Legendrian component, then d_α^A and d_α^i vanish identically. If A is a connected Legendrian submanifold, then d_α^A either vanishes identically or is non-degenerate, and d_α^i either vanishes identically or for any $i_0, i_1 \in \mathcal{L}(i)$ with $i_0(A) \neq i_1(A)$ we have that $d_\alpha^i(i_0, i_1) > 0$.*

Proof. Corollary 5.12 implies that R_A is empty if $\dim(A) < n$ and that R_A is a union of components of A if $\dim(A) = n$. In the later case we know by Corollary 5.8 (and Remark 5.7) that every component of R_A is Legendrian. Then Remark 5.3 finishes the proof.

The proof for d_α^i is analogous. \square

The next proposition finishes the proof of Theorem 2.17.

Proposition 5.14. *Let $i : L \hookrightarrow M$ be the inclusion of a compact and connected Legendrian submanifold. Then d_α^L vanishes identically if and only if there exists a positive loop of Legendrians starting at L . d_α^i vanishes identically if and only if there exists a positive loop of parametrized Legendrians starting at i .*

The proof differs only notationally from the proof of the second part of Theorem 2.6.

Remark 5.15. On $\widetilde{L}(A)$ and $\widetilde{L}(i)$ one can similarly define

$$(107) \quad \begin{aligned} \widetilde{\Sigma}_A &:= \{f \in \widetilde{\text{Cont}}_0(M, \xi) \mid f([A]) = [A]\}, \\ \widetilde{\Sigma}_i &:= \{f \in \widetilde{\text{Cont}}_0(M, \xi) \mid f([i]) = [i]\} \end{aligned}$$

and

$$(108) \quad \begin{aligned} \widetilde{R}_A &:= \bigcap_{f \in \widetilde{\Sigma}_A} (\pi(f))^{-1}(A), \\ \widetilde{R}_i &:= \bigcap_{f \in \widetilde{\Sigma}_i} (\pi(f))^{-1}(A), \end{aligned}$$

where $[A]$ and $[i]$ denote the elements of $\widetilde{L}(A)$ and $\widetilde{L}(i)$ represented by the constant paths at $[A]$ and $[i]$, and $\pi : \widetilde{\text{Cont}}_0(M, \xi) \rightarrow \text{Cont}_0(M, \xi)$ is the projection. The proof of Proposition 5.1 still applies to show that

$$(109) \quad \widetilde{\Sigma}_A = \{f \in \widetilde{\text{Cont}}_0(M, \xi) \mid \widetilde{d}_\alpha^A([A], f([A])) = 0\}$$

and

$$(110) \quad \widetilde{\Sigma}_i = \{f \in \widetilde{\text{Cont}}_0(M, \xi) \mid \widetilde{d}_\alpha^i([i], f([i])) = 0\}.$$

Assume that $i : A \rightarrow M$ is the proper embedding of a submanifold. Then the above arguments apply to show that \widetilde{R}_A and \widetilde{R}_i are empty if $\dim(A) < n$, that $\widetilde{R}_A = A$ or $\widetilde{R}_i = A$ only if A is coisotropic, and that \widetilde{R}_A and \widetilde{R}_i are unions of Legendrian components of A if $\dim(A) = n$.

As before, we have that \widetilde{d}_α^A vanishes identically if $\widetilde{R}_A = \emptyset$ and \widetilde{d}_α^i vanishes identically if $\widetilde{R}_i = \emptyset$.

On the other hand, observe that

$$(111) \quad \begin{aligned} \widetilde{R}_A = A &\Leftrightarrow \forall f \in \widetilde{\Sigma}_A : A \subseteq (\pi(f))^{-1}(A) \\ &\Leftrightarrow \forall f \in \widetilde{\Sigma}_A : A = (\pi(f))(A) \\ &\Leftrightarrow \widetilde{d}_\alpha^A \text{ is } \pi\text{-non-degenerate on } \widetilde{\mathcal{L}}(A), \end{aligned}$$

where the equivalence between the second and the third line follows from (109).

Similarly,

$$(112) \quad \begin{aligned} \widetilde{R}_i = A &\Leftrightarrow \forall f \in \widetilde{\Sigma}_i : A \subseteq (\pi(f))^{-1}(A) \\ &\Leftrightarrow \forall f \in \widetilde{\Sigma}_i : A = (\pi(f))(A) \\ &\Leftrightarrow \forall \iota \in \widetilde{\mathcal{L}}(i), (\pi(\iota))(A) \neq A : \widetilde{d}_\alpha^i([i], \iota) > 0 \\ &\Leftrightarrow \forall \iota_0, \iota_1 \in \widetilde{\mathcal{L}}(i), (\pi(\iota_0))(A) \neq (\pi(\iota_1))(A) : \widetilde{d}_\alpha^i(\iota_0, \iota_1) > 0, \end{aligned}$$

where here, π denotes the projection $\pi : \widetilde{\mathcal{L}}(i) \rightarrow \mathcal{L}(i)$, and the equivalence between the second and the third line follows from (110).

TODO update reference cant

REFERENCES

- [AA23] Simon Allais and Pierre-Alexandre Arlove, *Spectral selectors and contact orderability*, preprint, arXiv:2309.10578, 9 2023.
- [AF12] Peter Albers and Urs Frauenfelder, *A Variational Approach to Givental's nonlinear Maslov Index*, Geometric and Functional Analysis **22** (2012), 1033–1050.
- [AFM15] Peter Albers, Urs Fuchs, and Will J. Merry, *Orderability and the Weinstein conjecture*, Compositio Mathematica **151** (2015), no. 12, 2251–2272.

- [AM18] Peter Albers and Will Merry, *Orderability, contact non-squeezing, and Rabinowitz Floer homology*, Journal of Symplectic Geometry **16** (2018).
- [Arl21] Pierre-Alexandre Arlove, *Normes sur le groupe des contactomorphismes et contactisation de domaines étoilés, leurs géodésiques et leurs caractéristiques translitées.*, PhD thesis, Université de Strasbourg, 2021.
- [Ban97] Augustin Banyaga, *The structure of classical diffeomorphism groups*, first ed., Mathematics and Its Applications, Springer, Boston, MA, 1997.
- [Bhu01] Mohan Bhupal, *A partial order on the group of contactomorphisms of \mathbb{R}^{2n+1} via generating functions*, Turkish Journal of Mathematics **25** (2001).
- [BZ15] Matthew Borman and Frol Zapolsky, *Quasi-morphisms on contactomorphism groups and contact rigidity*, Geometry & Topology **19** (2015).
- [Can23] Dylan Cant, *Remarks on the oscillation energy of Legendrian isotopies*, Geometriae Dedicata **217** (2023).
- [CFP16] Vincent Colin, Emmanuel Ferrand, and Petya Pushkar, *Positive Isotopies of Legendrian Submanifolds and Applications*, International Mathematics Research Notices **2017** (2016), no. 20, 6231–6254.
- [CN10a] Vladimir Chernov and Stefan Nemirovski, *Legendrian Links, Causality, and the Low Conjecture*, Geometric and Functional Analysis **19** (2010).
- [CN10b] ———, *Nonnegative Legendrian isotopy in ST^*M* , Geometry & Topology - GEOM TOPOL **14** (2010), 611–626.
- [CN16] ———, *Universal orderability of legendrian isotopy classes*, Journal of Symplectic Geometry **14** (2016), 149–170.
- [CN20] ———, *Interval topology in contact geometry*, Communications in Contemporary Mathematics **22** (2020), no. 5.
- [DRS22] Georgios Dimitroglou Rizell and Michael Sullivan, *C^0 -limits of Legendrians and positive loops*, preprint, arXiv:2212.09190v3, 2022.
- [EKP06] Yakov Eliashberg, Sang Seon Kim, and Leonid Polterovich, *Geometry of contact transformations and domains: orderability versus squeezing*, Geom. Topol. **10** (2006), 1635–1747.
- [EP93] Ya. M. Eliashberg and L. Polterovich, *Biinvariant metrics on the group of hamiltonian diffeomorphisms*, International Journal of Mathematics **04** (1993).
- [EP00] ———, *Partially ordered groups and geometry of contact transformations*, Geom. funct. anal. **10** (2000), 1448–1476.
- [FPR18] Maia Fraser, Leonid Polterovich, and Daniel Rosen, *On Sandon-type metrics for contactomorphism groups*, Annales mathématiques du Québec **42** (2018).
- [Hed21] Jakob Hedicke, *Lorentzian distance functions in contact geometry*, Journal of Topology and Analysis (2021).
- [Hua15] Yang Huang, *On Legendrian foliations in contact manifolds I: Singularities and neighborhood theorems*, Math. Res. Lett. **22** (2015), no. 5, 1373–1400. MR 3488380
- [KS18] Michael Kunzinger and Clemens Sämann, *Lorentzian length spaces*, Annals of Global Analysis and Geometry **54** (2018).
- [McD02] Dusa McDuff, *Geometric Invariants of the Hofer norm*, Journal of Symplectic Geometry **1** (2002).
- [Nak19] Lukas Nakamura, *Non-rigidity Results for non-Legendrian and C^0 -limits of Legendrian Submanifolds*, Master’s thesis, Ludwig-Maximilians-Universität München, 2019.
- [Ryb10] Tomasz Rybicki, *Commutators of contactomorphisms*, Adv. Math. **225** (2010), no. 6, 3291–3326. MR 2729009
- [RZ20] Daniel Rosen and Jun Zhang, *Chekanov’s dichotomy in contact topology*, Mathematical Research Letters (2020).
- [She17] Egor Shelukhin, *The Hofer norm of a contactomorphism*, J. Symplectic Geom. **15** (2017), no. 4, 1173–1208. MR 3734612
- [Ush15] Michael Usher, *Observations on the Hofer distance between closed subsets*, Math. Res. Lett. **22** (2015), no. 6, 1805–1820. MR 3507263
- [Vit92] Claude Viterbo, *Symplectic topology as the geometry of generating functions*, Mathematische Annalen **292** (1992), 685–710.
- [Wei13] Peter Weigel, *Orderable contact structures on liouville-fillable contact manifolds*, Journal of Symplectic Geometry **13** (2013).