

# ON FINITE FIELD ANALOGUES OF DETERMINANTS INVOLVING THE BETA FUNCTION

HAI-LIANG WU, LI-YUAN WANG, HAO PAN\*

ABSTRACT. Motivated by the works of L. Carlitz, R. Chapman and Z.-W. Sun on cyclotomic matrices, in this paper, we investigate certain cyclotomic matrices concerning the Jacobi sums over finite fields, which can be viewed as finite field analogues of certain matrices involving the Beta function. For example, let  $q > 1$  be a prime power and let  $\chi$  be a generator of the group of all multiplicative characters of  $\mathbb{F}_q$ . Then we prove that

$$\det [J_q(\chi^i, \chi^j)]_{1 \leq i, j \leq q-2} = (q-1)^{q-3},$$

where  $J_q(\chi^i, \chi^j)$  is the Jacobi sum over  $\mathbb{F}_q$ . This is a finite analogue of

$$\det [B(i, j)]_{1 \leq i, j \leq n} = (-1)^{\frac{n(n-1)}{2}} \prod_{r=0}^{n-1} \frac{(r!)^3}{(n+r)!},$$

where  $B$  is the Beta function. Also, if  $q = p \geq 5$  is an odd prime, then we show that

$$\det [J_p(\chi^{2i}, \chi^{2j})]_{1 \leq i, j \leq (p-3)/2} = \frac{1 + (-1)^{\frac{p+1}{2}} p}{4} \left( \frac{p-1}{2} \right)^{\frac{p-5}{2}}.$$

## 1. INTRODUCTION

**1.1. Notations.** Let  $q = p^n$  be a prime power with  $p$  prime and  $n \in \mathbb{Z}^+$ . Let  $\mathbb{F}_q$  be the finite field of  $q$  elements, and let  $\mathbb{F}_q^\times$  be the group of all nonzero elements of  $\mathbb{F}_q$ . Let  $\widehat{\mathbb{F}_q^\times}$  denote the group of all multiplicative characters of  $\mathbb{F}_q$ . In addition, for any  $\psi \in \widehat{\mathbb{F}_q^\times}$  we define  $\psi(0) = 0$ . Throughout out this paper, we let  $\chi$  be a generator of  $\widehat{\mathbb{F}_q^\times}$ . Also, For any  $\chi^i, \chi^j \in \widehat{\mathbb{F}_q^\times}$ , the Jacobi sum of  $\chi^i, \chi^j$  is defined by

$$J_q(\chi^i, \chi^j) := \sum_{x \in \mathbb{F}_q} \chi^i(x) \chi^j(1-x).$$

---

2020 *Mathematics Subject Classification.* Primary 11L05, 11C20; Secondary 15B33, 11R18.

*Keywords.* finite fields, determinants, Jacobi sums.

This research was supported by the National Natural Science Foundation of China (Grant Nos. 12101321, 12201291 and 12071208).

\*Corresponding author.

**1.2. Background and Motivations.** We first introduce some related works involving cyclotomic matrices. The earliest research on cyclotomic matrices came from Lehmer [8] and Carlitz [1]. For example, given a nontrivial character  $\psi \in \widehat{\mathbb{F}_p^\times}$ , Carlitz studied the characteristic polynomial of the cyclotomic matrix

$$C_p(\psi) := [\psi(j-i)]_{1 \leq i, j \leq p-1}.$$

In particular, when  $\psi$  is a quadratic character, Carlitz showed that the characteristic polynomial of  $C_p(\psi)$  is

$$\left(t^2 - (-1)^{\frac{p-1}{2}} p\right)^{\frac{p-3}{2}} \left(t^2 - (-1)^{\frac{p-1}{2}}\right).$$

Along this line, Chapman [2] initiated the study of several variants of Carlitz's results. Surprisingly, these variants have close connections with the real quadratic field  $\mathbb{Q}(\sqrt{p})$ . For example, let  $\left(\frac{\cdot}{p}\right)$  be the Legendre symbol and let

$$\varepsilon_p^{(2 - (\frac{2}{p}))h_p} = a_p + b_p\sqrt{p} \quad (a_p, b_p \in \mathbb{Q}),$$

where  $\varepsilon_p > 1$  and  $h_p$  are the fundamental unit and class number of  $\mathbb{Q}(\sqrt{p})$  respectively. Chapman conjectured that

$$\det \left[ \left( \frac{j-i}{p} \right) \right]_{1 \leq i, j \leq \frac{p+1}{2}} = \begin{cases} -a_p & \text{if } p \equiv 1 \pmod{4}, \\ 1 & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

This challenging conjecture was later known as Chapman's "evil determinant" conjecture, and was confirmed by Vsemirnov [12, 13].

In recent years, Sun [11] and Krachun and his collaborators [6] studied some variants of the above determinant. For example, Sun proved that

$$-\det \left[ \left( \frac{i^2 + j^2}{p} \right) \right]_{1 \leq i, j \leq (p-1)/2}$$

is a quadratic residue modulo  $p$ . Later the first author [14] proved that for any  $c \in \mathbb{F}_p$  with  $c \neq \pm 2$ ,

$$\det \left[ \left( \frac{i^2 + cij + j^2}{p} \right) \right]_{0 \leq i, j \leq p-1} = 0$$

whenever the curve defined by the equation  $y^2 = x(x^2 + cx + 1)$  is a supersingular elliptic curve over  $\mathbb{F}_p$ . Readers may refer to [6, 11, 14, 15] for recent progress on this topic.

After introducing the above relevant research results, we now describe our research motivations.

For any complex numbers  $x, y$  with  $\operatorname{Re}(x) > 0$  and  $\operatorname{Re}(y) > 0$ , the Beta function is defined by

$$B(x, y) = \int_0^1 t^{x-1}(1-t)^{y-1} dt.$$

If we define a function

$$J_q : \widehat{\mathbb{F}}_q^\times \times \widehat{\mathbb{F}}_q^\times \rightarrow \mathbb{Q}(e^{2\pi i/(q-1)})$$

by sending  $(\chi^i, \chi^j)$  to the Jacobi sum  $J_q(\chi^i, \chi^j)$ , then it is known that  $J_q$  is indeed a finite analogue of the Beta function. Readers may refer to [4, 5] for the detailed introduction on this topic.

Let  $n$  be a positive integer. We first consider the following matrix

$$B_n := [B(i, j)]_{1 \leq i, j \leq n}.$$

It is known that for any positive integers  $i, j$  we have

$$B(i, j) = \frac{\Gamma(i)\Gamma(j)}{\Gamma(i+j)},$$

where  $\Gamma(\cdot)$  is the Gamma function. By this and [10, (4.8)] we obtain

$$\det B_n = (-1)^{\frac{n(n-1)}{2}} \prod_{r=0}^{n-1} \frac{(r!)^3}{(n+r)!}. \quad (1.1)$$

Recall that  $\chi$  is a generator of  $\widehat{\mathbb{F}}_q^\times$ . Since  $J_q$  is a finite field analogue of the Beta function and  $\mathbb{F}_q^\times \cong \widehat{\mathbb{F}}_q^\times$ , the determinant

$$\det [J_q(\chi^i, \chi^j)]_{1 \leq i, j \leq q-2} \quad (1.2)$$

can be naturally viewed as a finite field analogue of (1.1).

On the other hand, we turn to the following matrix involving the binomial coefficients:

$$C_n := \left[ \binom{i+j}{i} \right]_{1 \leq i, j \leq n} = \left[ \frac{\Gamma(i+j+1)}{\Gamma(i+1)\Gamma(j+1)} \right]_{1 \leq i, j \leq n}.$$

One can easily verify that (we will give a detailed proof in the appendix)

$$\det C_n = n+1. \quad (1.3)$$

Let  $\zeta_{q-1} = e^{2\pi i/(q-1)}$  and let  $\zeta_p = e^{2\pi i/p}$ . Consider the cyclotomic field  $L = \mathbb{Q}(\zeta_{q-1}, \zeta_p)$  and let  $\mathcal{O}_L$  be the ring of all algebraic integers over  $L$ . Let  $\mathfrak{p}$  be a prime ideal of  $\mathcal{O}_L$  with  $p \in \mathfrak{p}$ . Then it is easy to verify that

$$\mathcal{O}_L/\mathfrak{p} \cong \mathbb{F}_q.$$

Let  $\chi_{\mathfrak{p}}$  be the Teichmüller character of  $\mathfrak{p}$ , i.e.,

$$\chi_{\mathfrak{p}}(x \bmod \mathfrak{p}) \equiv x \pmod{\mathfrak{p}}$$

for any  $x \in \mathcal{O}_L$ . Then for any integer  $1 \leq i, j \leq q-2$ , the following congruence holds (see [3, Proposition 3.6.4]).

$$J_q(\chi_{\mathfrak{p}}^{-i}, \chi_{\mathfrak{p}}^{-j}) \equiv -\binom{i+j}{i} \equiv -\frac{\Gamma(i+j+1)}{\Gamma(i+1)\Gamma(j+1)} \pmod{\mathfrak{p}}. \quad (1.4)$$

Now congruence (1.4) establishes a connection between binomial coefficients and Jacobi sums, and hence (1.2) can be also viewed as a finite field analogue of (1.3).

Motivated by the above results, it is natural to consider the matrices with Jacobi sums as their entries. In general, for any positive integer  $k \mid q - 1$ , we define the cyclotomic matrix

$$J_q(k) := [J_q(\chi^{ki}, \chi^{kj})]_{1 \leq i, j \leq (q-1-k)/k} \quad (1.5)$$

which concerns the Jacobi sums of nontrivial  $k$ -th multiplicative characters.

**1.3. Main Results.** Now we state our main results of this paper.

**Theorem 1.1.** *Let  $q = p^n$  be a prime power with  $p$  prime and  $n \in \mathbb{Z}^+$ . Let  $k \mid q - 1$  be a positive integer and let  $m = (q - 1)/k$ . Then the following results hold.*

- (i)  $\det J_q(k) \in \mathbb{Z}$  and is independent of the choice of the generator  $\chi$ .
- (ii) We have the congruence

$$\det J_q(k) \equiv (-1)^{\frac{(k+1)(m^2-m)}{2}} \pmod{p}. \quad (1.6)$$

*Remark 1.1.* For an arbitrary positive divisor  $k$  of  $q - 1$ , determining the explicit value of  $\det J_q(k)$  seems very difficult. However, for  $k = 1, 2$  we can obtain some satisfactory results.

The following result follows Theorem 1.1 and (1.4) directly.

**Corollary 1.1.** *Let  $q = p^n$  be a prime power with  $p$  prime and  $n \in \mathbb{Z}^+$ . Let  $k \mid q - 1$  be a positive integer and let  $m = (q - 1)/k$ . Then*

$$\det \left[ \begin{pmatrix} ki + kj \\ ki \end{pmatrix} \right]_{1 \leq i, j \leq m-1} \equiv (-1)^{\frac{(k+1)m^2 - (k-1)m-2}{2}} \pmod{p}.$$

Our next result concerns the explicit value of  $\det J_q(1)$ .

**Theorem 1.2.** *Let  $q = p^n \geq 3$  with  $p$  prime and  $n \in \mathbb{Z}^+$ . Then*

$$\det J_q(1) = (q - 1)^{q-3}.$$

The next theorem determines the explicit value of  $J_p(2)$ .

**Theorem 1.3.** *Let  $p \geq 5$  be an odd prime. Then*

$$\det J_p(2) = \frac{1 + (-1)^{\frac{p+1}{2}} p}{4} \left( \frac{p-1}{2} \right)^{\frac{p-5}{2}}.$$

*Remark 1.2.* Unlike Theorem 1.2, it is difficult to generalize this result to the finite field  $\mathbb{F}_{p^n}$  with  $n \geq 2$ . In Remark 4.3, we will provide a detailed explanation of the computational obstacles.

**1.4. Outline of This Paper.** We will prove our main results in Sections 2–4 respectively. In Section 5, we will pose some open problems for further research. Finally, the proof of (1.3) will be given in the appendix.

## 2. PROOF OF THEOREM 1.1

Throughout this section, we let  $m = (q - 1)/k$ . For an integer  $a$  with  $\gcd(a, m) = 1$ , the map  $x \bmod m \mapsto ax \bmod m$  induces a permutation  $\tau_m(a)$  of  $\mathbb{Z}/m\mathbb{Z}$ . We begin with the following result due to Lerch [7].

**Lemma 2.1.** *Let  $\text{sign}(\tau_m(a))$  denote the sign of  $\tau_m(a)$ . Then*

$$\text{sign}(\tau_m(a)) = \begin{cases} \left(\frac{a}{m}\right) & \text{if } m \equiv 1 \pmod{2}, \\ 1 & \text{if } m \equiv 2 \pmod{4}, \\ (-1)^{(a-1)/2} & \text{if } m \equiv 0 \pmod{4}, \end{cases}$$

where  $\left(\frac{\cdot}{m}\right)$  is the Jacobi symbol if  $m$  is odd. In particular,

$$\text{sign}(\tau_m(-1)) = (-1)^{\frac{(m-1)(m-2)}{2}}.$$

We also need the well-known Lucas congruence.

**Lemma 2.2.** *Let  $a, b, c, d \in \mathbb{N} = \{0, 1, 2, \dots\}$  with  $0 \leq b, d \leq p - 1$ . Then*

$$\binom{ap + b}{cp + d} \equiv \binom{a}{c} \binom{b}{d} \pmod{p}.$$

Now we are in a position to prove our first theorem.

**Proof of Theorem 1.1.** (i) It is known that

$$\text{Gal}(\mathbb{Q}(\zeta_{q-1})/\mathbb{Q}) \cong (\mathbb{Z}/(q-1)\mathbb{Z})^\times.$$

Hence

$$\text{Gal}(\mathbb{Q}(\zeta_{q-1})/\mathbb{Q}) = \{\sigma_r : r \in \mathbb{Z} \text{ and } \gcd(r, q-1) = 1\},$$

where  $\sigma_r(\zeta_{q-1}) = \zeta_{q-1}^r$ . Now for any  $\sigma_r \in \text{Gal}(\mathbb{Q}(\zeta_{q-1})/\mathbb{Q})$ , by Lemma 2.1 we have

$$\begin{aligned} \sigma_r(\det J_q(k)) &= \det [J_q(\chi^{rki}, \chi^{rkj})]_{1 \leq i, j \leq m-1} \\ &= \text{sign}(\tau_m(r))^2 \cdot \det J_q(k) \\ &= \det J_q(k). \end{aligned}$$

By the Galois theory and noting that  $\det J_q(k)$  is an algebraic integer, we have  $\det J_q(k) \in \mathbb{Z}$ . Also, for any integer  $r$ , the character  $\chi^r$  is a generator of  $\widehat{\mathbb{F}_q^\times}$  if and only if  $\gcd(r, q-1) = 1$ . Thus, for any generator  $\chi'$ , there exists an  $r$  with  $\gcd(r, q-1) = 1$  such that

$$\det [J_q(\chi'^{ki}, \chi'^{kj})]_{1 \leq i, j \leq m-1} = \sigma_r(\det J_q(k)) = \det J_q(k).$$

Hence  $\det J_q(k)$  is independent of the choice of the generator  $\chi$ .

(ii) Now we adopt the notations of (1.4). By (i) we see that  $\det J_q(k)$  is independent of the choice of the generator  $\chi$ . Hence we let  $\chi = \chi_{\mathfrak{p}}$  be the Teichmüller character of  $\mathfrak{p}$  (note that  $\chi_{\mathfrak{p}}$  is a generator of  $\widehat{\mathbb{F}_q^\times}$ ).

By Lemma 2.1 we first have

$$\det J_q(k) = (-1)^{\frac{(m-1)(m-2)}{2}} \det [J_q(\chi_{\mathfrak{p}}^{ki}, \chi_{\mathfrak{p}}^{-kj})]_{1 \leq i, j \leq m-1}. \quad (2.7)$$

For any  $1 \leq i < j \leq m-1$ , by (1.4) and Lemma 2.2 we have

$$\begin{aligned} J_q(\chi_{\mathfrak{p}}^{ki}, \chi_{\mathfrak{p}}^{-kj}) &= J_q(\chi_{\mathfrak{p}}^{-(q-1-ki)}, \chi_{\mathfrak{p}}^{-kj}) \\ &\equiv -\binom{q-1+kj-ki}{kj} \\ &\equiv -\binom{kj-ki-1}{kj} \\ &\equiv 0 \pmod{\mathfrak{p}}. \end{aligned}$$

Combining this with (2.7) and noting that  $J_q(\chi_{\mathfrak{p}}^{ki}, \chi_{\mathfrak{p}}^{-ki}) = (-1)^{ki+1}$  for  $1 \leq i \leq q-2$ , we obtain

$$\begin{aligned} \det J_q(k) &\equiv (-1)^{\frac{(m-1)(m-2)}{2}} \det [J_q(\chi_{\mathfrak{p}}^{ki}, \chi_{\mathfrak{p}}^{-kj})]_{1 \leq i, j \leq m-1} \\ &\equiv (-1)^{\frac{(m-1)(m-2)}{2}} \prod_{1 \leq i \leq m-1} J_q(\chi_{\mathfrak{p}}^{ki}, \chi_{\mathfrak{p}}^{-ki}) \\ &\equiv (-1)^{\frac{(m-1)(m-2)}{2}} \prod_{1 \leq i \leq m-1} (-1)^{ki+1} \\ &\equiv (-1)^{\frac{(k+1)m^2 - (k+1)m}{2}} \pmod{\mathfrak{p}}. \end{aligned}$$

Since  $\det J_q(k) \in \mathbb{Z}$ , we finally obtain

$$\det J_q(k) \equiv (-1)^{\frac{(k+1)m^2 - (k+1)m}{2}} \pmod{p}.$$

In view of the above, we have completed the proof of Theorem 1.1.  $\square$

### 3. PROOF OF THEOREM 1.2

In this section, we set  $\mathbb{F}_q = \{a_0, a_1, a_2, \dots, a_{q-1}\}$  with  $a_0 = 0$  and  $a_1 = 1$ . Also, we define

$$\Delta_q = \prod_{1 \leq i < j \leq q-1} (\chi(a_j) - \chi(a_i)). \quad (3.8)$$

We begin with the following result.

**Lemma 3.3.** *Let notations be as above. Then*

$$\Delta_q^2 = (-1)^{\frac{(q-1)(q-2)+2q}{2}} (q-1)^{q-1}.$$

*Proof.* As  $\chi$  is a generator of  $\widehat{\mathbb{F}_q^\times}$ , we have

$$G(t) := t^{q-1} - 1 = \prod_{1 \leq j \leq q-1} (t - \chi(a_j)). \quad (3.9)$$

By (3.9) we have

$$\prod_{1 \leq j \leq q-1} \chi(a_j) = (-1)^q. \quad (3.10)$$

Let  $G'(t) = (q-1)t^{q-2}$  be the derivative of  $G(t)$ . Then by (3.9) and (3.10)

$$\begin{aligned}
 \Delta_q^2 &= (-1)^{\frac{(q-1)(q-2)}{2}} \prod_{1 \leq i \neq j \leq q-1} (\chi(a_j) - \chi(a_i)) \\
 &= (-1)^{\frac{(q-1)(q-2)}{2}} \prod_{1 \leq j \leq q-1} \prod_{i \neq j} (\chi(a_j) - \chi(a_i)) \\
 &= (-1)^{\frac{(q-1)(q-2)}{2}} \prod_{1 \leq j \leq q-1} G'(\chi(a_j)) \\
 &= (-1)^{\frac{(q-1)(q-2)}{2}} (q-1)^{q-1} \prod_{1 \leq j \leq q-1} \chi(a_j)^{-1} \\
 &= (-1)^{\frac{(q-1)(q-2)+2q}{2}} (q-1)^{q-1}.
 \end{aligned}$$

This completes the proof.  $\square$

We also need the following result.

**Lemma 3.4.** *Let notations be as above. Then for any  $y \in \mathbb{F}_q^\times$  we have*

$$D_y := \prod_{x \in \mathbb{F}_q^\times \setminus \{y\}} (\chi(y) - \chi(x)) = (q-1)\chi(y)^{-1}.$$

*Proof.* By (3.9) we have

$$D_y = G'(\chi(y)) = (q-1)\chi(y)^{-1}.$$

This completes the proof.  $\square$

The next lemma concerns the sign of a permutation of  $a_2, a_3, \dots, a_{q-1}$ .

**Lemma 3.5.** *Let  $\pi_q$  be the permutation of the sequence  $a_2, a_3, \dots, a_{q-1}$  induced by the map  $x \mapsto -x + 1$ . Then the sign of  $\pi_q$  is equal to*

$$\text{sign}(\pi_q) = (-1)^{\frac{(q-2)(q-3)}{2}}.$$

*Proof.* We first consider the case  $2 \mid q$ . In this case, clearly  $\pi_q$  can be written as a product of  $(q-2)/2$  transpositions. Hence we have

$$\text{sign}(\pi_q) = (-1)^{\frac{q-2}{2}} = (-1)^{\frac{(q-2)(q-3)}{2}}.$$

Suppose now  $2 \nmid q$ . Then  $\{1, -1\}$  can be viewed as a subset of  $\mathbb{F}_q$ . Hence

$$\text{sign}(\pi_q) = \prod_{2 \leq i < j \leq q-1} \frac{(1-a_j) - (1-a_i)}{a_j - a_i} = (-1)^{\frac{(q-2)(q-3)}{2}}.$$

This completes the proof.  $\square$

Now we are in a position to prove our second theorem.

**Proof of Theorem 1.2.** For any  $1 \leq i, j \leq q-2$ , we clearly have

$$J_q(\chi^i, \chi^j) = \sum_{2 \leq k \leq q-1} \chi^i(a_k) \chi^j(1-a_k).$$

By this we obtain the matrix decomposition

$$J_q(1) = M_q N_q, \quad (3.11)$$

where  $M_q$  is a  $(q-2) \times (q-2)$  matrix defined by

$$M_q := [\chi^i(a_k)]_{1 \leq i \leq q-2, 2 \leq k \leq q-1},$$

and  $N_q$  is also a  $(q-2) \times (q-2)$  matrix defined by

$$N_q := [\chi^j(1-a_k)]_{2 \leq k \leq q-1, 1 \leq j \leq q-2}.$$

We first consider  $\det M_q$ . Recall that  $\Delta_q$  and  $D_y$  are defined by (3.8) and Lemma 3.4 respectively. By definition and (3.10) we have

$$\begin{aligned} \det M_q &= \begin{vmatrix} \chi^1(a_2) & \chi^1(a_3) & \cdots & \chi^1(a_{q-1}) \\ \chi^2(a_2) & \chi^2(a_3) & \cdots & \chi^2(a_{q-1}) \\ \vdots & \vdots & \ddots & \vdots \\ \chi^{q-2}(a_2) & \chi^{q-2}(a_3) & \cdots & \chi^{q-2}(a_{q-1}) \end{vmatrix} \\ &= \prod_{2 \leq j \leq q-1} \chi(a_j) \prod_{2 \leq i < j \leq q-1} (\chi(a_j) - \chi(a_i)) \\ &= \frac{(-1)^q \Delta_q}{D_1} \\ &= \frac{(-1)^q \Delta_q}{q-1}. \end{aligned} \quad (3.12)$$

We now turn to  $\det N_q$ . By Lemma 3.5 we have

$$\begin{aligned} \det N_q &= \begin{vmatrix} \chi^1(1-a_2) & \chi^2(1-a_2) & \cdots & \chi^{q-2}(1-a_2) \\ \chi^1(1-a_3) & \chi^2(1-a_3) & \cdots & \chi^{q-2}(1-a_3) \\ \vdots & \vdots & \ddots & \vdots \\ \chi^1(1-a_{q-1}) & \chi^2(1-a_{q-1}) & \cdots & \chi^{q-2}(1-a_{q-1}) \end{vmatrix} \\ &= \text{sign}(\pi_q) \cdot \begin{vmatrix} \chi^1(a_2) & \chi^2(a_2) & \cdots & \chi^{q-2}(a_2) \\ \chi^1(a_3) & \chi^2(a_3) & \cdots & \chi^{q-2}(a_3) \\ \vdots & \vdots & \ddots & \vdots \\ \chi^1(a_{q-1}) & \chi^2(a_{q-1}) & \cdots & \chi^{q-2}(a_{q-1}) \end{vmatrix}. \end{aligned}$$

Now similar to the calculation of  $\det M_q$ , one can verify that

$$\det N_q = (-1)^{\frac{(q-2)(q-3)+2q}{2}} \Delta_q / (q-1). \quad (3.13)$$

Combining (3.12) with (3.13) and by Lemma 3.3, we obtain

$$\det J_q(1) = \det M_q \cdot \det N_q = (q-1)^{q-3}.$$

In view of the above, we have completed the proof of Theorem 1.2.  $\square$

## 4. PROOF OF THEOREM 1.3

Let  $M$  be an  $r \times n$  complex matrix with  $r \leq n$  and let  $N$  be an  $n \times r$  complex matrix. Set

$$\mathcal{S}_r = \{\mathbf{s} = (j_1, j_2, \dots, j_r) : 1 \leq j_1 < j_2 < \dots < j_r \leq n\}.$$

For any  $\mathbf{s} \in \mathcal{S}_r$ , we define  $M_{\mathbf{s}}$  (respectively  $N^{\mathbf{s}}$ ) to be the  $r \times r$  submatrix of  $M$  (respectively submatrix of  $N$ ) obtained by deleting all columns (respectively all rows) except those with indices in  $\mathbf{s}$ . We begin with the well-known Cauchy-Binet formula.

**Lemma 4.6.** *Let  $M, N$  be two complex matrices of sizes  $r \times n$  and  $n \times r$  respectively with  $r \leq n$ . Then*

$$\det(MN) = \sum_{\mathbf{s} \in \mathcal{S}_r} \det(M_{\mathbf{s}}) \det(N^{\mathbf{s}}).$$

In the remaining part of this section, we let  $n = (p-1)/2$ . Then it is clear that  $1^2, 2^2, \dots, n^2$  are exactly all nonzero squares of  $\mathbb{F}_p$ . Also, we define

$$S_p := \prod_{1 \leq i < j \leq n} (\chi(j^2) - \chi(i^2)). \quad (4.14)$$

The next result determines the explicit value of  $S_p^2$ .

**Lemma 4.7.** *Let notations be as above. Then*

$$S_p^2 = (-1)^{\frac{n^2+n+2}{2}} n^n.$$

*Proof.* Since  $\chi$  is a generator of  $\widehat{\mathbb{F}_p^\times}$ , we have

$$H(t) := t^n - 1 = \prod_{1 \leq j \leq n} (t - \chi(j^2)). \quad (4.15)$$

By (4.15) we obtain

$$\prod_{1 \leq j \leq n} \chi(j^2) = (-1)^{n+1}. \quad (4.16)$$

Now by (4.15) and (4.16) one can verify that

$$\begin{aligned} S_p^2 &= (-1)^{\frac{n(n-1)}{2}} \prod_{1 \leq i \neq j \leq n} (\chi(j^2) - \chi(i^2)) \\ &= (-1)^{\frac{n(n-1)}{2}} \prod_{1 \leq j \leq n} \prod_{i \neq j} H'(\chi(j^2)) \\ &= (-1)^{\frac{n(n-1)}{2}} n^n \prod_{1 \leq j \leq n} \chi(j^2)^{-1} \\ &= (-1)^{\frac{n^2+n+2}{2}} n^n \end{aligned}$$

This completes the proof. □

Let  $t_1, t_2, \dots, t_n$  be variables. For any positive integer  $k$ , we let

$$P_k(t_1, t_2, \dots, t_n) = t_1^k + t_2^k + \dots + t_n^k.$$

Also, the  $k$ -th elementary symmetric polynomial of  $t_1, \dots, t_n$  is denoted by  $e_k(t_1, \dots, t_n)$ , i.e.,

$$e_k(t_1, \dots, t_n) = \sum_{1 \leq j_1 < \dots < j_k \leq n} \prod_{r=1}^k t_{j_r}.$$

Also, we let  $e_0(t_1, \dots, t_n) = 1$ .

We need the following result.

**Lemma 4.8.** *Let  $\chi(i^2) = x_i$  for  $1 \leq i \leq n$ . Then for any  $1 \leq k \leq n-1$ ,*

$$P_k(x_1, x_2, \dots, x_n) = x_1^k + x_2^k + \dots + x_n^k = 0$$

*Proof.* By (4.15) it is easy to see that

$$e_k := e_k(x_1, x_2, \dots, x_n) = 0$$

for any  $1 \leq k \leq n-1$ . By the Newton identities,  $P_k(x_1, \dots, x_n)$  can be written as a polynomial of  $e_1, e_2, \dots, e_k$ . This implies that

$$P_k(x_1, x_2, \dots, x_n) = x_1^k + x_2^k + \dots + x_n^k = 0$$

for any  $1 \leq k \leq n-1$ . This completes the proof.  $\square$

Now we are in a position to prove our last theorem.

**Proof of Theorem 1.3.** Note first that for any  $1 \leq i, j \leq n-1$  we have

$$\begin{aligned} J_q(\chi^{2i}, \chi^{2j}) &= \sum_{k=1}^{p-1} \chi^i(k^2) \chi^j((1-k)^2) \\ &= \sum_{k=1}^n \chi^i(k^2) (\chi^j((1-k)^2) + \chi^j((1+k)^2)). \end{aligned}$$

By this one can verify the following matrix decomposition

$$J_p(2) = AB, \tag{4.17}$$

where  $A$  is an  $(n-1) \times n$  matrix defined by

$$A := [\chi^i(k^2)]_{1 \leq i \leq n-1, 1 \leq k \leq n},$$

and  $B$  is an  $n \times (n-1)$  matrix defined by

$$B := [\chi^j((1-k)^2) + \chi^j((1+k)^2)]_{1 \leq k \leq n, 1 \leq j \leq n-1}.$$

Also, for any  $1 \leq k \leq n$ , we let  $A_{(k)}$  be the submatrix of  $A$  obtained by deleting the  $k$ -th column of  $A$  and let  $B^{(k)}$  be the submatrix of  $B$  obtained by deleting the  $k$ -th row of  $B$ . By Lemma 4.6 we have

$$\det J_p(2) = \sum_{k=1}^n \det A_{(k)} \det B^{(k)}. \tag{4.18}$$

We first consider  $\det A_{(k)}$ . It is clear that

$$\det A_{(k)} = \begin{vmatrix} \chi^1(1^2) & \cdots & \chi^1((k-1)^2) & \chi^1((k+1)^2) & \cdots & \chi^1(n^2) \\ \chi^2(1^2) & \cdots & \chi^2((k-1)^2) & \chi^2((k+1)^2) & \cdots & \chi^2(n^2) \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \chi^{n-1}(1^2) & \cdots & \chi^{n-1}((k-1)^2) & \chi^{n-1}((k+1)^2) & \cdots & \chi^{n-1}(n^2) \end{vmatrix}.$$

Combining this with (4.15) and (4.16), one can verify that

$$\begin{aligned} \det A_{(k)} &= \prod_{i \neq k} \chi(i^2) \cdot \frac{\prod_{1 \leq i < j \leq n} (\chi(j^2) - \chi(i^2))}{\prod_{1 \leq i \leq k-1} (\chi(k^2) - \chi(i^2)) \prod_{k+1 \leq i \leq n} (\chi(i^2) - \chi(k^2))} \\ &= \frac{S_p \cdot \prod_{1 \leq i \leq n} \chi(i^2)}{\chi(k^2)(-1)^{n-k} \prod_{i \neq k} (\chi(k^2) - \chi(i^2))} \\ &= \frac{(-1)^{k+1}}{n} S_p. \end{aligned} \quad (4.19)$$

The last equality follows from

$$\prod_{i \neq k} (\chi(k^2) - \chi(i^2)) = H'(\chi(k^2)) = n\chi(k^2)^{-1}.$$

Applying (4.19) to (4.18) and by the determinant expansion formula, we obtain

$$\det J_p(2) = \frac{S_p}{n} \sum_{k=1}^n (-1)^{k+1} \det B^{(k)} = \frac{S_p}{n} \det \tilde{B}, \quad (4.20)$$

where  $\tilde{B}$  is an  $n \times n$  matrix obtained by adding an  $n \times 1$  column vector with all elements 1 before the first column of  $B$ .

We now turn to  $\det \tilde{B}$ . For simplicity, we let  $\chi(i^2) = x_i$  for  $0 \leq i \leq p-1$ . Note that  $x_0 = 0, x_1 = 1$  and  $x_i = x_{p-i}$  for any  $1 \leq i \leq p-1$ . Using these notations we have

$$\det \tilde{B} = \begin{vmatrix} 1 & x_0 + x_2 & x_0^2 + x_2^2 & \cdots & x_0^{n-1} + x_2^{n-1} \\ 1 & x_1 + x_3 & x_1^2 + x_3^2 & \cdots & x_1^{n-1} + x_3^{n-1} \\ 1 & x_2 + x_4 & x_2^2 + x_4^2 & \cdots & x_2^{n-1} + x_4^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_{n-1} + x_{n+1} & x_{n-1}^2 + x_{n+1}^2 & \cdots & x_{n-1}^{n-1} + x_{n+1}^{n-1} \end{vmatrix}. \quad (4.21)$$

By Lemma 4.8 and noting that  $x_n = x_{n+1}$ , we have

$$\sum_{r=0}^{n-1} (x_r^k + x_{r+2}^k) = -x_1^k + 2 \sum_{r=1}^n x_r^k = -x_1^k \quad (4.22)$$

for any  $1 \leq k \leq n-1$ .

In (4.21), adding the 2nd row, 3rd row,  $\dots$ ,  $n$ -th row to the 1st row and by (4.22), we obtain

$$\det \tilde{B} = \begin{vmatrix} n & -x_1 & -x_1^2 & \cdots & -x_1^{n-1} \\ 1 & x_1 + x_3 & x_1^2 + x_3^2 & \cdots & x_1^{n-1} + x_3^{n-1} \\ 1 & x_2 + x_4 & x_2^2 + x_4^2 & \cdots & x_2^{n-1} + x_4^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_{n-1} + x_{n+1} & x_{n-1}^2 + x_{n+1}^2 & \cdots & x_{n-1}^{n-1} + x_{n+1}^{n-1} \end{vmatrix}. \quad (4.23)$$

In (4.23), by adding the 1st row to the 2nd row we have

$$\det \tilde{B} = -\frac{1}{2} \begin{vmatrix} -2n & 2x_1 & 2x_1^2 & \cdots & 2x_1^{n-1} \\ n+1 & x_3 & x_3^2 & \cdots & x_3^{n-1} \\ 1 & x_2 + x_4 & x_2^2 + x_4^2 & \cdots & x_2^{n-1} + x_4^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_{n-1} + x_{n+1} & x_{n-1}^2 + x_{n+1}^2 & \cdots & x_{n-1}^{n-1} + x_{n+1}^{n-1} \end{vmatrix}. \quad (4.24)$$

By Lemma 4.8 again, for  $1 \leq k \leq n-1$  we have

$$2x_1^k + x_3^k + \sum_{r=2}^{n-1} (x_r^k + x_{r+2}^k) = -x_2^k + 2 \sum_{r=1}^n x_r^k = -x_2^k. \quad (4.25)$$

In (4.24), adding the 2nd row, 3rd row,  $\dots$ ,  $n$ th row to the 1st row and by (4.25), we have

$$\det \tilde{B} = \frac{1}{2} \begin{vmatrix} 1 & x_2 & x_2^2 & \cdots & x_2^{n-1} \\ n+1 & x_3 & x_3^2 & \cdots & x_3^{n-1} \\ 1 & x_2 + x_4 & x_2^2 + x_4^2 & \cdots & x_2^{n-1} + x_4^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_{n-1} + x_{n+1} & x_{n-1}^2 + x_{n+1}^2 & \cdots & x_{n-1}^{n-1} + x_{n+1}^{n-1} \end{vmatrix}. \quad (4.26)$$

In (4.26), by using the row operations recursively, one can verify that

$$\det \tilde{B} = \frac{1}{2} \begin{vmatrix} 1 & x_2 & x_2^2 & \cdots & x_2^{n-1} \\ n+1 & x_3 & x_3^2 & \cdots & x_3^{n-1} \\ \alpha_4 & x_4 & x_4^2 & \cdots & x_4^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \alpha_n & x_n & x_n^2 & \cdots & x_n^{n-1} \\ \alpha_{n+1} & x_{n+1} & x_{n+1}^2 & \cdots & x_{n+1}^{n-1} \end{vmatrix}, \quad (4.27)$$

where

$$\alpha_k = \begin{cases} \frac{1-(-1)^{k/2}}{2} & \text{if } k \equiv 0 \pmod{2}, \\ (-1)^{(k+1)/2} \cdot (n+1) + \frac{1+(-1)^{(k-1)/2}}{2} & \text{otherwise.} \end{cases} \quad (4.28)$$

In (4.27), by subtracting the  $(n-1)$ -th row from the  $n$ -th row and noting that  $x_n = x_{n+1}$ , we obtain

$$\begin{aligned}
 \det \tilde{B} &= \frac{1}{2} \begin{vmatrix} 1 & x_2 & x_2^2 & \cdots & x_2^{n-1} \\ n+1 & x_3 & x_3^2 & \cdots & x_3^{n-1} \\ \alpha_4 & x_4 & x_4^2 & \cdots & x_4^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \alpha_n & x_n & x_n^2 & \cdots & x_n^{n-1} \\ \alpha_{n+1} - \alpha_n & 0 & 0 & \cdots & 0 \end{vmatrix} \\
 &= \frac{1}{2} \cdot (-1)^{n+1} \cdot \beta_p \cdot \begin{vmatrix} x_2 & x_2^2 & \cdots & x_2^{n-1} \\ x_3 & x_3^2 & \cdots & x_3^{n-1} \\ x_4 & x_4^2 & \cdots & x_4^{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ x_n & x_n^2 & \cdots & x_n^{n-1} \end{vmatrix}, \tag{4.29}
 \end{aligned}$$

where  $\beta_p = \alpha_{n+1} - \alpha_n$  and by (4.28) one can verify that

$$\beta_p = (-1)^{\frac{n(n-1)}{2} + n + 1} \cdot \frac{p + (-1)^{n+1}}{2}. \tag{4.30}$$

Now by (4.29) and (4.16) we finally obtain

$$\begin{aligned}
 \det \tilde{B} &= \frac{1}{2} \cdot (-1)^{n+1} \cdot \beta_p \left( \prod_{2 \leq i \leq n} x_i \right) \prod_{2 \leq i < j \leq n} (x_j - x_i) \\
 &= \frac{1}{2} \cdot \beta_p \cdot \frac{S_p}{\prod_{2 \leq i \leq n} (x_i - x_1)} \\
 &= \frac{(-1)^{n-1}}{2n} \beta_p S_p. \tag{4.31}
 \end{aligned}$$

The last equality follows from

$$\prod_{2 \leq i \leq n} (x_i - x_1) = (-1)^{n-1} H'(x_1) = (-1)^{n-1} n.$$

By (4.20), (4.30), (4.31) and Lemma 4.7, we finally obtain

$$\det J_p(2) = \frac{S_p}{n} \det \tilde{B} = \frac{1 + (-1)^{\frac{p+1}{2}} p}{4} \left( \frac{p-1}{2} \right)^{\frac{p-5}{2}}.$$

In view of above, we have completed the proof of Theorem 1.3. □

*Remark 4.3.* Now we explain why our method used in the above proof cannot be applied to  $\mathbb{F}_{p^l}$  with  $l \geq 2$ . In fact, in the prime field  $\mathbb{F}_p$ , we can naturally list all nonzero squares as  $1^2, 2^2, \dots, n^2$ . Let  $x_i = \chi(i^2)$  for  $0 \leq i \leq q-1$ . Then it is easy to verify that the  $(i, j)$ -entry of the matrix  $B$  defined by (4.17) can be simplified as  $x_{i-1}^j + x_{i+1}^j$ .

However, for  $\mathbb{F}_{p^l}$  with  $l \geq 2$ , if we choose elements  $a_1, \dots, a_{(p^l-1)/2}$  such that

$$a_1^2, \dots, a_{(p^l-1)/2}^2$$

are exactly all the nonzero squares of  $\mathbb{F}_{p^l}$ . Then we can easily get the matrix decomposition

$$J_q(2) = A'B',$$

where

$$A' = [\chi^i(a_k^2)]_{1 \leq i \leq \frac{p^l-3}{2}, 1 \leq k \leq \frac{p^l-1}{2}},$$

and

$$B' = [\chi^j((1-a_k)^2) + \chi^j((1+a_k)^2)]_{1 \leq k \leq \frac{p^l-1}{2}, 1 \leq j \leq \frac{p^l-3}{2}}.$$

Similar to (4.18), we also have

$$\det J_q(2) = \sum_{k=1}^{\frac{p^l-1}{2}} \det A'_{(k)} \det B'^{(k)},$$

and it is easy to verify that

$$\det A'_{(k)} = (-1)^{k+1} \frac{2}{p^l-1} S_q,$$

where

$$S_q = \prod_{1 \leq i < j \leq (p^l-1)/2} (\chi(a_j^2) - \chi(a_i^2)).$$

Using this we obtain

$$\det J_q(2) = \frac{2}{p^l-1} S_q \det \widetilde{B}',$$

where  $\widetilde{B}'$  is an  $(p^l-1)/2 \times (p^l-1)/2$  matrix obtained by adding an  $(p^l-1)/2 \times 1$  column vector with all elements 1 before the first column of  $B'$ .

Now our main obstacle is the calculation of  $\det \widetilde{B}'$ . This is because we cannot represent the  $(i, j)$ -entry of  $B'$  in a simple form like that of  $B$ .

## 5. CONCLUDING REMARKS

Greene [5, Definition 2.4] posed an analogue of binomial coefficients. For any  $A, B \in \widehat{\mathbb{F}_q^\times}$ , Greene defined

$$\binom{A}{B} := \frac{B(-1)}{q} J(A, \overline{B}),$$

where  $\overline{B} \in \widehat{\mathbb{F}_q^\times}$  such that  $\overline{B}(x) = B(x)^{-1}$  for any  $x \in \mathbb{F}_q^\times$ . Motivated by Greene's analogue and our theorems, investigating

$$\det \left[ \begin{pmatrix} \chi^{i+j} \\ \chi^i \end{pmatrix} \right]_{1 \leq i, j \leq q-2} \quad \text{and} \quad \det \left[ \begin{pmatrix} \chi^{2i+2j} \\ \chi^{2i} \end{pmatrix} \right]_{1 \leq i, j \leq (q-3)/2}$$

might be also meaningful. However, we cannot solve this problem currently.

## 6. APPENDIX

In this appendix we prove (1.3). It is known that for any positive integers  $r, l$ , we have

$$\binom{r}{l} = \binom{r-1}{l-1} + \binom{r-1}{l}. \quad (6.32)$$

Now we calculate  $\det C_n = \det \left[ \binom{i+j}{i} \right]_{1 \leq i, j \leq n}$ . The case  $n = 1$  is trivial. Assume now  $n \geq 2$ . Subtracting the  $(n-1-i)$ -th row from the  $(n-i)$ -th row for  $i = 0, 1, \dots, n-2$  sequentially and by (6.32), one can verify that

$$\det C_n = \det \left[ \binom{i+j}{i} \right]_{1 \leq i, j \leq n} = \begin{vmatrix} 2 & 3 & \cdots & n+1 \\ \binom{2}{2} & \binom{3}{2} & \cdots & \binom{n+1}{2} \\ \vdots & \vdots & \ddots & \vdots \\ \binom{n}{n} & \binom{n+1}{n} & \cdots & \binom{2n-1}{n} \end{vmatrix}. \quad (6.33)$$

We next consider the right hand side of (6.33). Subtracting the  $(n-1-i)$ -th column from the  $(n-i)$ -th column for  $i = 0, 1, \dots, n-2$  sequentially and by (6.32) again, we obtain

$$\det C_n = \begin{vmatrix} 1+1 & 1+0 & \cdots & 1+0 & 1+0 \\ 1 & \binom{2}{1} & \cdots & \binom{n-1}{1} & \binom{n}{1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & \binom{n-1}{n-2} & \cdots & \binom{2n-4}{n-2} & \binom{2n-3}{n-2} \\ 1 & \binom{n}{n-1} & \cdots & \binom{2n-3}{n-1} & \binom{2n-2}{n-1} \end{vmatrix} = \det D_n + \det C_{n-1}, \quad (6.34)$$

where

$$\det D_n = \begin{vmatrix} 1 & 1 & \cdots & 1 & 1 \\ 1 & \binom{2}{1} & \cdots & \binom{n-1}{1} & \binom{n}{1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & \binom{n-1}{n-2} & \cdots & \binom{2n-4}{n-2} & \binom{2n-3}{n-2} \\ 1 & \binom{n}{n-1} & \cdots & \binom{2n-3}{n-1} & \binom{2n-2}{n-1} \end{vmatrix}.$$

For  $\det D_n$ , subtracting the  $(n-1-i)$ -th row from the  $(n-i)$ -th row for  $i = 0, 1, \dots, n-2$  sequentially and using (6.32), we see that

$$\det D_n = \begin{vmatrix} 1 & 1 & \cdots & 1 & 1 \\ 0 & \binom{1}{1} & \cdots & \binom{n-2}{1} & \binom{n-1}{1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & \binom{n-2}{n-2} & \cdots & \binom{2n-5}{n-2} & \binom{2n-4}{n-2} \\ 0 & \binom{n-1}{n-1} & \cdots & \binom{2n-4}{n-1} & \binom{2n-3}{n-1} \end{vmatrix} = \begin{vmatrix} 1 & \cdots & \binom{n-2}{1} & \binom{n-1}{1} \\ \vdots & \ddots & \vdots & \vdots \\ 1 & \cdots & \binom{2n-5}{n-2} & \binom{2n-4}{n-2} \\ 1 & \cdots & \binom{2n-4}{n-1} & \binom{2n-3}{n-1} \end{vmatrix}.$$

Repeating the above procedure, one can easily verify that

$$\det D_n = 1. \quad (6.35)$$

Combining (6.35) with (6.34) and noting that  $\det C_1 = 2$ , we finally obtain  $\det C_n = n + 1$ . This completes the proof of (1.3).

## REFERENCES

- [1] L. Carlitz, Some cyclotomic matrices, *Acta Arith.* 5 (1959), 293–308.
- [2] R. Chapman, Determinants of Legendre symbol matrices, *Acta Arith.* 115 (2004), 231–244.
- [3] H. Cohen, *Number Theory, Vol. I: Tools and Diophantine Equations*, Springer, New York, 2007.
- [4] J. Fuselier, L. Long, R. Ramakrishna, H. Swisher and F.-T. Tu, Hypergeometric functions over finite fields. *Mem. Amer. Math. Soc.* 280 (2022), no. 1382, 124 pp.
- [5] J. Greene, Hypergeometric functions over finite fields, *Trans. Am. Math. Soc.* 301 (1987), 77–101.
- [6] D. Krachun, F. Petrov, Z.-W. Sun, M. Vsemirnov, On some determinants involving Jacobi symbols, *Finite Fields Appl.* 64 (2020) 101672.
- [7] M. Lerch, Sur un théorème de Zolotarev, *Bull. Intern. Acad. François Joseph* 3 (1896), 34–37.
- [8] D. H. Lehmer, On certain character matrices, *Pacific J. Math.* 6 (1956), 491–499.
- [9] R. Lidl and H. Niederreiter, *Finite Fields*, 2nd ed., Cambridge Univ. Press, Cambridge, 1997.
- [10] J.-M. Normand, Calculation of some determinants using the  $s$ -shifted factorial, *J. Phys. A: Math. Gen.* 37 (2004) 5737–5762.
- [11] Z.-W. Sun, On some determinants with Legendre symbols entries, *Finite Fields Appl.* 56 (2019), 285–307.
- [12] M. Vsemirnov, On the evaluation of R. Chapman’s “evil determinant”, *Linear Algebra Appl.* 436 (2012), 4101–4106.
- [13] M. Vsemirnov, On R. Chapman’s “evil determinant”: case  $p \equiv 1 \pmod{4}$ , *Acta Arith.* 159 (2013), 331–344.
- [14] H.-L. Wu, Elliptic curves over  $\mathbb{F}_p$  and determinants of Legendre matrices, *Finite Fields Appl.* 76 (2021), 101929.
- [15] H.-L. Wu, Y.-F. She and L.-Y. Wang, Cyclotomic matrices and hypergeometric functions over finite fields, *Finite Fields Appl.* 82 (2022), 102054.

(HAI-LIANG WU) SCHOOL OF SCIENCE, NANJING UNIVERSITY OF POSTS AND TELECOMMUNICATIONS, NANJING 210023, PEOPLE’S REPUBLIC OF CHINA  
*Email address:* whl.math@smail.nju.edu.cn

(HAO PAN) SCHOOL OF APPLIED MATHEMATICS, NANJING UNIVERSITY OF FINANCE AND ECONOMICS, NANJING 210046, PEOPLE’S REPUBLIC OF CHINA  
*Email address:* haopan79@zoho.com

(LI-YUAN WANG) SCHOOL OF PHYSICAL AND MATHEMATICAL SCIENCES, NANJING TECH UNIVERSITY, NANJING 211816, PEOPLE’S REPUBLIC OF CHINA  
*Email address:* wly@smail.nju.edu.cn