

Operator Norm Bounds on the Correlation Matrix of the SK Model at High Temperature

Christian Brennecke* Changji Xu† Horng-Tzer Yau‡

July 25, 2023

Abstract

We prove that the two point correlation matrix $\mathbf{M} = (\langle \sigma_i; \sigma_j \rangle)_{1 \leq i, j \leq N} \in \mathbb{R}^{N \times N}$ of the Sherrington-Kirkpatrick model has the property that for every $\epsilon > 0$ there exists $K_\epsilon > 0$, that is independent of N , such that

$$\mathbb{P}(\|\mathbf{M}\|_{\text{op}} \leq K_\epsilon) \geq 1 - \epsilon$$

for N large enough, for suitable interaction and external field parameters (β, h) in the replica symmetric region. In other words, the operator norm of \mathbf{M} is of order one with high probability. Our results are in particular valid for all $(\beta, h) \in (0, 1) \times (0, \infty)$ and thus complement recently obtained results in [16, 7] that imply the operator norm boundedness of \mathbf{M} for all $\beta < 1$ in the special case of vanishing external field.

1 Setup and Main Results

We consider systems of N interacting spins $\sigma_i \in \{-1, 1\}, i \in [N] = \{1, \dots, N\}$, described by the Sherrington-Kirkpatrick [25] Hamiltonian $H_N : \{-1, 1\}^N \rightarrow \mathbb{R}$ which is defined by

$$H_N(\sigma) = \beta \sum_{1 \leq i < j \leq N} g_{ij} \sigma_i \sigma_j + h \sum_{i=1}^N \sigma_i = \frac{\beta}{2} (\sigma, \mathbf{G} \sigma) + h (\mathbf{1}, \sigma). \quad (1.1)$$

The symmetric matrix $\mathbf{G} = (g_{ij})_{1 \leq i, j \leq N}$ is a GOE matrix, that is, up to the symmetry constraint its entries are i.i.d. centered Gaussian random variables of variance N^{-1} for $i \neq j$ and we set $g_{ii} = 0$ for simplicity. The standard Euclidean inner product in \mathbb{R}^N and its induced norm are denoted by (\cdot, \cdot) and $\|\cdot\|$, respectively. We assume $\beta > 0, h > 0$ and

*Institute for Applied Mathematics, University of Bonn, Endenicher Allee 60, 53115 Bonn, Germany

†Center of Mathematical Sciences and Applications, Harvard University, Cambridge MA 02138, USA

‡Department of Mathematics, Harvard University, One Oxford Street, Cambridge MA 02138, USA

we assume the $\{g_{ij}\}$ to be realized in some probability space $(\Omega, \mathcal{F}, \mathbb{P})$. The expectation with regards to \mathbb{P} is denoted by $\mathbb{E}(\cdot)$ and the $L^p(\Omega, \mathcal{F}, \mathbb{P})$ norms are $\|\cdot\|_p = (\mathbb{E}|\cdot|^p)^{1/p}$.

Based on a novel representation of the entries of the two point correlation matrix

$$\mathbf{M} = \mathbf{M}_{\beta,h} = (m_{ij})_{1 \leq i,j \leq N} = (\langle \sigma_i; \sigma_j \rangle)_{1 \leq i,j \leq N} \in \mathbb{R}^{N \times N}$$

as sums over weights of self-avoiding paths, which is motivated by the results of [1], we have recently proved in [7] for the special case $h = 0$ that at high temperature, \mathbf{M} is asymptotically close to a resolvent of \mathbf{G} , in the sense that

$$\lim_{N \rightarrow \infty} \|\mathbf{M}_{\beta,h=0} - (1 + \beta^2 - \beta \mathbf{G})^{-1}\|_{\text{op}} = 0 \quad (1.2)$$

in probability. Here, $\|\mathbf{A}\|_{\text{op}} = \sup_{\mathbf{u} \in \mathbb{R}^N: \|\mathbf{u}\|=1} \|\mathbf{A}\mathbf{u}\|$ denotes the standard operator norm, for $\mathbf{A} \in \mathbb{R}^{N \times N}$, and $\langle \cdot \rangle = \langle \cdot \rangle_{\beta,h}$ denotes the Gibbs measure induced by H_N , i.e.

$$\langle f \rangle = \frac{1}{Z_N} \sum_{\sigma \in \{-1,1\}^N} f(\sigma) e^{H_N(\sigma)} \quad \text{with} \quad Z_N = \sum_{\sigma \in \{-1,1\}^N} e^{H_N(\sigma)}$$

for $f : \{-1,1\}^N \rightarrow \mathbb{R}$ and $\langle f; g \rangle = \langle fg \rangle - \langle f \rangle \langle g \rangle$. By standard properties of GOE matrices, the validity of (1.2) naturally suggests the (well-known) existence of a phase transition at $\beta = 1$ and [7] verifies the validity of (1.2) indeed in the full replica symmetric region at $h = 0$, that is, for all $\beta < 1$. It implies in particular that

$$\lim_{N \rightarrow \infty} \|\mathbf{M}\|_{\text{op}} = (1 - \beta)^{-2},$$

so that the operator norm of \mathbf{M} is typically of order one as long as $\beta < 1$. The boundedness of $\|\mathbf{M}\|_{\text{op}}$ has also been proved independently in [16], in the sense that $\mathbb{E} \|\mathbf{M}\|_{\text{op}} = O(1)$.

A natural question is whether \mathbf{M} has bounded norm in the general case $h \neq 0$ as well and whether this is connected to the replica symmetric phase, like in the case $h = 0$. Although a complete proof is to date still lacking, it is generally expected that the SK model is replica symmetric for all (β, h) satisfying the AT [15] condition¹

$$\beta^2 E \operatorname{sech}^4(h + \beta \sqrt{q} Z) < 1, \quad (1.3)$$

where here and in the following $q = q_{\beta,h}$ denotes the unique solution to

$$q = E \tanh^2(h + \beta \sqrt{q} Z). \quad (1.4)$$

Moreover, $Z \sim \mathcal{N}(0, 1)$ denotes a standard Gaussian independent of the disorder $\{g_{ij}\}$.

¹Expectations over effective random variables, like for instance Z in Eq. (1.3) and Eq. (1.4), that are independent of the disorder $\{g_{ij}\}$ in (1.1), are denoted by $E(\cdot)$, which is to be distinguished from expectations $\mathbb{E}(\cdot)$ over the disorder in (1.1).

Proving the norm boundedness of \mathbf{M} (or an analogue of (1.2)) for $h \neq 0$ under (1.3) is a challenging problem. In this work instead, we impose further assumptions on (β, h) which ensure exponential concentration of the overlap, based on results obtained by Talagrand in [28, 29]. More precisely, we assume in the sequel the AT condition (1.3) together with

$$(\partial_m \Phi)(m, q')|_{m=1} < 0 \text{ for all } q < q' \leq 1, \quad (1.5)$$

where for $0 \leq m \leq 1$ and $q \leq q' \leq 1$ we set

$$\Phi(m, q') = \log 2 + \frac{\beta^2}{4}(1-q')^2 - \frac{\beta^2}{4}m(q'^2 - q^2) + \frac{1}{m}E \log E_{Z'} \cosh^m(h + \beta\sqrt{q}Z + \beta\sqrt{q' - q}Z').$$

Here, $(Z, Z') \sim \mathcal{N}(0, \text{id}_{\mathbb{R}^2})$ and $E_{Z'}(\cdot)$ denotes the expectation over Z' . Conditions (1.3) & (1.5) ensure that we consider parameters (β, h) inside a suitable interior region of the replica symmetric phase of the model; see [29, Chapter 13] for more details on this. They imply in particular locally uniform in β exponential concentration of the overlap, which in turn enables an efficient computation of many observables.

While we recall the results from [28, 29] that are based on (1.3) & (1.5) and that are used in the sequel in Section 2, let us already point out that (1.3) & (1.5) are satisfied for every external field strength $h > 0$ whenever $\beta < 1$: indeed following [29] one has

$$\begin{aligned} (\partial_m \Phi)(m, q')|_{m=1}(q) &= 0, \quad \partial_{q'} \left((\partial_m \Phi)(m, q')|_{m=1} \right)(q) = 0, \\ \partial_{q'}^2 \left((\partial_m \Phi)(m, q')|_{m=1} \right) &= \frac{\beta^2}{2} \left(\beta^2 E \frac{E_{Z'} \text{sech}^3(h + \beta\sqrt{q}Z + \beta\sqrt{q' - q}Z')}{E_{Z'} \cosh(h + \beta\sqrt{q}Z + \beta\sqrt{q' - q}Z')} - 1 \right) \end{aligned}$$

and simple monotonicity arguments as in [13, Section 3] imply

$$\beta^2 \frac{E_{Z'} \text{sech}^3(h + \beta\sqrt{q}Z + \beta\sqrt{q' - q}Z')}{E_{Z'} \cosh(h + \beta\sqrt{q}Z + \beta\sqrt{q' - q}Z')} \leq \beta^2 E_{Z'} \text{sech}^4(h + \beta\sqrt{q}Z + \beta\sqrt{q' - q}Z').$$

Thus, $(\partial_m \Phi)(m, q')|_{m=1} < 0$ for all $q < q' \leq 1$ whenever $\beta < 1$.

Assuming from now on (1.3) & (1.5), our strategy to prove the norm boundedness of \mathbf{M} combines recent ideas from [5, 6, 1, 14, 9, 7] and provides several results of independent interest. In the first step of our analysis, we compute \mathbf{M} up to an error which has vanishing Frobenius norm in the limit $N \rightarrow \infty$. To state this more precisely, we set from now on

$$\mathbf{m} = (m_1, \dots, m_N) = (\langle \sigma_1 \rangle, \dots, \langle \sigma_N \rangle) \in \mathbb{R}^N$$

and we denote by $\langle \cdot \rangle^{(i)}$ the Gibbs measure obtained from $\langle \cdot \rangle$ by removing the spin σ_i .

Proposition 1.1. *Assume that (β, h) satisfy (1.3) & (1.5). Then, there exists a constant $C = C_{\beta, h} > 0$ such that for all $1 \leq i \neq j \leq N$, we have that*

$$\mathbb{E} \left(m_{ij} - \beta(1 - m_i^2) \sum_{k=1}^N g_{ik} m_{kj}^{(i)} \right)^2 \leq CN^{-5/2}. \quad (1.6)$$

Remark 1.1. Eq. (1.6) shows that $m_{ij} \approx \beta(1 - m_i^2) \sum_{k=1}^N g_{ik} m_{kj}^{(i)}$. Iterating the r.h.s., one obtains a representation of m_{ij} as a sum over weights $w(\gamma)$ of self-avoiding paths γ from i to j : each edge $e = \{i_1, i_2\} \in \gamma$ contributes a factor $\beta g_{i_1 i_2}$ to $w(\gamma)$ and each vertex a factor $(1 - (m_i^{(S)})^2)$, for suitable $S \subset [N]$. Since we do not use this representation to control the operator norm $\|\mathbf{M}\|_{\text{op}}$, we omit further details on its derivation, but only mention that it generalizes the path representation of $\mathbf{M}_{\beta, h=0}$ obtained in [7] in the range $\beta < 1$.

As discussed in detail in [1] (for β sufficiently small), after expressing the $m_{kj}^{(i)}$ through the original two point functions m_{kj} , (1.6) suggests the validity of the resolvent equation

$$\mathbf{M} \approx \frac{1}{\mathbf{D} + \beta^2(1 - q) - 2\beta^2 N^{-1} \mathbf{m} \otimes \mathbf{m} - \beta \mathbf{G}}, \quad (1.7)$$

where $\mathbf{D} \in \mathbb{R}^{N \times N}$ is the diagonal matrix with entries

$$(\mathbf{D})_{ij} = m_{ii}^{-1} \delta_{ij} = \frac{1}{1 - m_i^2} \delta_{ij}. \quad (1.8)$$

Motivated by this approximation, the second step of our analysis proves the following.

Proposition 1.2. Assume that (β, h) satisfy (1.3) & (1.5), let q denote the solution of (1.4) and let $\mathbf{D} \in \mathbb{R}^{N \times N}$ be defined as in (1.8). Then we have that

$$\|(\mathbf{D} + \beta^2(1 - q) - \beta \mathbf{G})\mathbf{M} - \text{id}_{\mathbb{R}^N}\|_{\text{op}} \leq o(1)\|\mathbf{M}\|_{\text{op}} + O(1)$$

for an error $o(1) \rightarrow 0$ as $N \rightarrow \infty$ in probability and where the error $O(1) \geq 0$ is of order one with high probability: there exists a constant $C = C_{\beta, h} > 0$ such that

$$\mathbb{P}(O(1) \leq K) \geq 1 - CK^{-2}$$

for every $K > 0$.

Remark 1.2. In view of the resolvent heuristics (1.7), it seems natural to expect that

$$\lim_{N \rightarrow \infty} \|(\mathbf{D} + \beta^2(1 - q) - 2\beta^2 N^{-1} \mathbf{m} \otimes \mathbf{m} - \beta \mathbf{G})\mathbf{M} - \text{id}_{\mathbb{R}^N}\|_{\text{op}} = 0,$$

with high probability. While the methods presented in this paper do not seem to allow a simple proof of this norm convergence, we plan to address this point in a follow-up work.

Finally, the norm boundedness of \mathbf{M} follows by deriving a uniform lower bound on the matrix $\mathbf{D} + \beta^2(1 - q) - \beta \mathbf{G}$ appearing in Prop. 1.2. This follows from the next result which is a direct consequence of combining recent results and ideas from [5, 6, 14, 9]. In its statement, $(\mathbf{m}^{(k)})_{k \geq 1}$ denotes Bolthausen's iterative TAP solution [5, 6], whose construction is recalled in detail in Section 4 below.

Proposition 1.3. *Assume that (β, h) satisfy the AT condition (1.3) and let q denote the solution of (1.4). Then, there exists $c = c_{\beta, h} > 0$, which is independent of $N \in \mathbb{N}$, so that*

$$\mathbf{D}_{\mathbf{m}^{(k)}} + \beta^2(1 - q) - \frac{2\beta^2}{N} \mathbf{m}^{(k)} \otimes \mathbf{m}^{(k)} - \beta \mathbf{G} \geq c \quad (1.9)$$

for k large enough, with probability tending to one as $N \rightarrow \infty$. Here, $\mathbf{m}^{(k)}$ denotes Bolthausen's iterative TAP solution at step k and $\mathbf{D}_{\mathbf{m}^{(k)}} \in \mathbb{R}^{N \times N}$ has entries

$$(\mathbf{D}_{\mathbf{m}^{(k)}})_{ij} = \frac{1}{1 - (m_i^{(k)})^2} \delta_{ij}.$$

Moreover, assuming (β, h) to satisfy both (1.3) & (1.5), it follows that

$$\mathbf{D} + \beta^2(1 - q) - \beta \mathbf{G} \geq \mathbf{D} + \beta^2(1 - q) - \frac{2\beta^2}{N} \mathbf{m} \otimes \mathbf{m} - \beta \mathbf{G} \geq c$$

with probability tending to one as $N \rightarrow \infty$, where $\mathbf{D} \in \mathbb{R}^{N \times N}$ is defined as in (1.8). In particular, $\mathbf{D} + \beta^2(1 - q) - \beta \mathbf{G}$ is invertible with high probability and satisfies

$$\|(\mathbf{D} + \beta^2(1 - q) - \beta \mathbf{G})^{-1}\|_{\text{op}} \leq c^{-1} < \infty.$$

Remark 1.3. *We point out that the proof of (1.9) only requires (β, h) to satisfy (1.3). The additional assumption (1.5) is used to show that \mathbf{m} is close to $\mathbf{m}^{(k)}$, applying the main result of [14] (see Prop. 4.5 for the details). The proof of (1.9) follows, up to a few modifications, from translating the arguments in [9] to the present setting. Related to this point, notice that the matrix on the l.h.s. in (1.9) is, up to a negligible error², equal to the negative of the Hessian of the TAP free energy functional [2] at $\mathbf{m}^{(k)}$. In particular, the lower bound in (1.9) resolves a recent open question from [21], which studies the limiting spectral distribution of the Hessian at $\mathbf{m}^{(k)}$ and which provides an interesting characterization of Plefka's conditions [24]. For further recent work related to the TAP approach, see also [4, 12, 11, 10, 8, 17, 3, 23, 19] and the references therein.*

Collecting Propositions 1.1, 1.2 and 1.3, we arrive at the following main result.

Theorem 1.4. *Assume that (β, h) satisfy (1.3) & (1.5). Then, for every $\epsilon > 0$ there exists a constant $K_\epsilon > 0$, that is independent of N , such that*

$$\mathbb{P}(\|\mathbf{M}\|_{\text{op}} \leq K_\epsilon) \geq 1 - \epsilon$$

for all N large enough.

²Under (1.3), it holds true that $q = N^{-1} \sum_{i=1}^N (m_i^{(k)})^2 + o(1)$ for an error $o(1)$ which is such that $\lim_{k \rightarrow \infty} \limsup_{N \rightarrow \infty} o(1) \rightarrow 0$, in probability.

Proof. On a set of probability tending to one as $N \rightarrow \infty$, Prop. 1.3 shows that

$$\|(\mathbf{D} + \beta^2(1 - q)) - \beta \mathbf{G}\|_{\text{op}}^{-1} \leq c_{\beta, h}^{-1}$$

for some constant $c_{\beta, h} > 0$, that is independent of N . Using Prop. 1.2 and the norm bound $\|\mathbf{A}\mathbf{B}\|_{\text{op}} \leq \|\mathbf{A}\|_{\text{op}}\|\mathbf{B}\|_{\text{op}}$ for all $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{N \times N}$, this implies

$$\begin{aligned} \|\mathbf{M}\|_{\text{op}} &\leq c_{\beta, h}^{-1} (1 + \|(\mathbf{D} + \beta^2(1 - q)) - \beta \mathbf{G}\|_{\text{op}}) \|\mathbf{M}\|_{\text{op}} \\ &\leq c_{\beta, h}^{-1} (1 + O(1)) + c_{\beta, h}^{-1} o(1) \|\mathbf{M}\|_{\text{op}} \end{aligned}$$

so that

$$\|\mathbf{M}\|_{\text{op}} \leq \frac{c_{\beta, h}^{-1} + O(1)}{1 - c_{\beta, h}^{-1} o(1)}.$$

The claim now follows from the fact that $\lim_{N \rightarrow \infty} o(1) = 0$ in the sense of probability and the bound $\mathbb{P}(O(1) > K) \leq CK^{-2}$, for some $C > 0$ and every $K > 0$. \square

The rest of this paper is devoted to the proofs of Prop. 1.1, Prop. 1.2 and Prop. 1.3. Section 2 recalls several useful results from [28, 29] and uses them to derive decay estimates on suitable correlation functions. Applying tools from stochastic calculus as in [1], this implies Prop. 1.1 and Prop. 1.2, which is explained in Section 3. Finally, in Section 4 we use the results of [5, 6, 14, 9] to deduce Prop. 1.3.

2 Bounds on (k, p) -Point Correlation Functions

In this section, we derive suitable decay bounds on a certain class of correlation functions that occur naturally in the proofs of Propositions 1.1 and 1.2. Our bounds combine ideas from [1] and results by Talagrand [28, 29] that follow from the assumptions (1.3) & (1.5).

To efficiently employ the results from [28, 29] that are recalled below, it is convenient to work in a slightly more general setting compared to the previous Section 1. To be more precise, we consider in this section spin systems with Hamiltonian of the form

$$H_N(\sigma) = \beta \sum_{1 \leq i < j \leq N} g_{ij} \sigma_i \sigma_j + \sum_{i=1}^N h_i \sigma_i = \frac{\beta}{2} (\sigma, \mathbf{G} \sigma) + (\mathbf{h}, \sigma), \quad (2.1)$$

where $\mathbf{h} \in \mathbb{R}^N$ is assumed to be a Gaussian random vector whose components are i.i.d. copies of a Gaussian random variable h with mean and variance denoted by

$$h \sim \mathcal{N}(\mu_h, \sigma_h^2).$$

We allow for the possibility that $\sigma_h^2 = 0$ (so that all of the following results apply in particular to systems with deterministic field as in (1.1)) and we assume that $E h^2 > 0$ (in

particular, the Gaussian field \mathbf{h} need not be centered). Under the latter assumption, recall from [20] and [28, Chapter 1] that there is a unique solution

$$q = q_{\beta, \mu_h, \sigma_h^2} \in (0, 1)$$

to (1.4). Let us point out that in this slightly more general setting, the expectation $E(\cdot)$ in (1.3), (1.4) and (1.5) is taken over all of the independent random variables Z, Z' and h .

In the following, we use dynamical methods to control the correlation functions of SK models with Hamiltonians as in (2.1), but with possibly modified interaction coupling β' and Gaussian field \mathbf{h}' (whose entries are i.i.d. copies of a modified Gaussian random variable $h' \sim \mathcal{N}(\mu_{h'}, \sigma_{h'}^2)$). All considered parameters (β', h') satisfy, however, conditions (1.3) & (1.5) and this motivates us to define the parameter set

$$\mathcal{A}_{RS^-} = \{(\beta, \mu_h, \sigma_h^2) \in (0, \infty) \times \mathbb{R} \times [0, \infty) : (\beta, h) \text{ satisfy (1.3) \& (1.5), for } h \sim \mathcal{N}(\mu_h, \sigma_h^2)\}.$$

Using the continuous dependence $(\beta, \mu_h, \sigma_h^2) \mapsto q_{\beta, h} = q_{\beta, \mu_h, \sigma_h^2}$, it is straightforward to check that the set $\mathcal{A}_{RS^-} \subset (0, \infty) \times \mathbb{R} \times [0, \infty)$ is open, which is frequently used below.

We denote by $(\sigma^l)_{l \in \mathbb{Z}}$ a sequence of i.i.d. samples from $\langle \cdot \rangle$, called replicas. Following standard conventions, the l -th replica σ^l corresponds to the l -th coordinate in $\prod_{j \in \mathbb{Z}} \{-1, 1\}^N$ with product Gibbs measure $\bigotimes_{j \in \mathbb{Z}} \langle \cdot \rangle$. By slight abuse of notation, we write $\langle \cdot \rangle$ also for expectations over functions of several replicas. Consider e.g. the overlap $R : \{-1, 1\}^N \times \{-1, 1\}^N \rightarrow \mathbb{R}$ which equals the normalized inner product of two replicas:

$$R(\sigma^l, \sigma^{l'}) = N^{-1} \sum_{i=1}^N \sigma_i^l \sigma_i^{l'} = N^{-1}(\sigma^l, \sigma^{l'}).$$

Below, we abbreviate for simplicity $R_{l, l'} = R(\sigma^l, \sigma^{l'})$ when computing Gibbs expectations of functions of the overlap of multiple replicas. Following the previous remark, we thus write for instance $\langle f(R_{1,2}, R_{3,4}) \rangle = \langle f(N^{-1}(\sigma^1, \sigma^2), N^{-1}(\sigma^3, \sigma^4)) \rangle^{\otimes 4}$ for every $f : \mathbb{R} \rightarrow \mathbb{R}$.

Theorem 2.1 ([29, Theorem 13.7.1.]). *Assume $(\beta, \mu_h, \sigma_h^2) \in \mathcal{A}_{RS^-}$ and $h \sim \mathcal{N}(\mu_h, \sigma_h^2)$. Let $q_{\beta, h}$ denote the unique solution of (1.4). Then, there is a constant $K_{\beta, h} > 0$ such that*

$$\mathbb{E} \langle \exp(N(R_{1,2} - q_{\beta, h})^2 / K_{\beta, h}) \rangle_{\beta, h} \leq 2.$$

Moreover, $K_{\beta, h} = K_{\beta, \mu_h, \sigma_h^2}$ is locally bounded in $(\beta, \mu_h, \sigma_h^2)$: for $\delta = \delta_{\beta, h} > 0$ small enough s.t. $(\beta - \delta, \beta + \delta) \times (\mu_h - \delta, \mu_h + \delta) \times [(\sigma_h^2 - \delta)_+, \sigma_h^2 + \delta) \subset \mathcal{A}_{RS^-}$, we find $K > 0$ such that

$$\sup_{\substack{\beta' \in (\beta - \delta, \beta + \delta), \\ \mu_{h'} \in (\mu_h - \delta, \mu_h + \delta), \\ \sigma_{h'}^2 \in [(\sigma_h^2 - \delta)_+, \sigma_h^2 + \delta)}} \mathbb{E} \langle \exp(N(R_{1,2} - q_{\beta', h'})^2 / K) \rangle_{\beta', h'} \leq 2, \quad (2.2)$$

where $(\sigma_h^2 - \delta)_+ = \max(0, \sigma_h^2 - \delta)$ and where $h' \sim \mathcal{N}(\mu_{h'}, \sigma_{h'}^2)$.

Remark 2.1. That $K_{\beta,h} > 0$ is locally bounded in $(\beta, \mu_h, \sigma_h^2)$ is not explicitly stated in [29, Theorem 13.7.1.], but it readily follows from the arguments in [29, Sections 13.4-13.7].

Theorem 2.1 implies useful decay bounds on a large class of correlation functions. Here, we rely on a direct consequence of (2.2) explained in detail in [28, Sections 1.8 & 1.10]. For the precise statement, we recall from [28] for $l, l' \in \mathbb{Z}$ the notation

$$T_{l,l'} = N^{-1}(\sigma^l - \mathbf{m}, \sigma^{l'} - \mathbf{m}), \quad T_l = N^{-1}(\sigma^l - \mathbf{m}, \mathbf{m}), \quad T = N^{-1}(\mathbf{m}, \mathbf{m}) \quad (2.3)$$

for replicas $(\sigma^l)_{l \in \mathbb{Z}}$ as well as

$$\begin{aligned} \nu_1^2 &= \frac{1 - 2q + q_4}{1 - \beta^2(1 - 2q + q_4)}, & \nu_2^2 &= \frac{q - q_4}{(1 - \beta^2(1 - 2q + q_4))(1 - \beta^2(1 - 4q + 3q_4))}, \\ \nu_3^2 &= \frac{q_4 - q^2}{1 - \beta^2(1 - 4q + 3q_4)} + \frac{\beta^2(q_4 - q^2)A^2}{1 - \beta^2(1 - 4q + 3q_4)} + \frac{2\beta^2(2q + q^2 - q_4)B^2}{1 - \beta^2(1 - 4q + 3q_4)}, \end{aligned} \quad (2.4)$$

where q is as in (1.4) and where $q_4 = E \tanh^4(h + \beta\sqrt{q}Z)$.

Theorem 2.2 ([28, Theorem 1.10.1]). Assume $(\beta, \mu_h, \sigma_h^2) \in \mathcal{A}_{RS^-}$ and $h \sim \mathcal{N}(\mu_h, \sigma_h^2)$. Denote by $U_{l,l'}, U_l$ and U independent centered Gaussian random variables with variance ν_1^2, ν_2^2 and, respectively, ν_3^2 , as in (2.4). Let $T_{l,l'}, T_l$ and T be defined as in (2.3). Then, for fixed $n \in \mathbb{N}_0$, $k(l, l') \in \mathbb{N}_0$ for $1 \leq l < l' \leq n$, $k(l) \in \mathbb{N}_0$ for $1 \leq l \leq n$, $k \in \mathbb{N}_0$ as well as

$$m = \sum_{1 \leq l < l' \leq n} k(l, l') + \sum_{1 \leq l \leq n} k(l) + k,$$

we have that

$$\left| \mathbb{E} \left\langle \prod_{1 \leq l < l' \leq n} T_{l,l'}^{k(l,l')} \prod_{1 \leq l \leq n} T_l^{k(l)} T^k \right\rangle - N^{-\frac{m}{2}} E \prod_{1 \leq l < l' \leq n} U_{l,l'}^{k(l,l')} \prod_{1 \leq l \leq n} U_l^{k(l)} U^k \right| \leq CN^{-\frac{m+1}{2}}.$$

Moreover, the constant $C = C_{\beta,h}$ is locally bounded in $(\beta, \mu_h, \sigma_h^2)$.

Remark 2.2. We remark that [28, Theorem 1.10.1] assumes $\beta < 1/2$ in order to apply [28, Eq. (1.88)] to obtain the error bounds in [28, Eq. (1.296)]. The latter is the key result that implies [28, Lemma 1.10.2 & Corollary 1.10.3], which in turn implies [28, Theorem 1.10.1]. Instead of assuming $\beta < 1/2$, we can equally assume $(\beta, \mu_h, \sigma_h^2) \in \mathcal{A}_{RS^-}$ and $h \sim \mathcal{N}(\mu_h, \sigma_h^2)$ to get [28, Eq. (1.296)], by Theorem 2.1. Given [28, Eq. (1.296)], Theorem 2.2 can then be proved exactly as in [28, Section 1.10]. Moreover, the fact that the constant $C_{\beta,h}$ is locally bounded in $(\beta, \mu_h, \sigma_h^2)$ is not explicitly explained in [28, Section 1.10], but it readily follows as a consequence of Theorem 2.1 and the relevant tools from [28, Sections 1.6, 1.8 & 1.10], in particular [28, Eq. (1.151), (1.154), (1.215) & (1.216)].

We are now ready to apply the preceding results to derive suitable decay bounds on correlation functions. For $k \in \mathbb{N}$ and $i_1, \dots, i_k \in [N]$, we define the k -point functions by

$$m_{i_1 i_2 \dots i_k} = \partial_{h_{i_1}} \partial_{h_{i_2}} \dots \partial_{h_{i_k}} \log Z_N(\mathbf{h}). \quad (2.5)$$

To derive our main results, it is useful to consider a certain class of (k, p) -point correlation functions, which are defined as follows: fixing $k, p \in \mathbb{N}_0$ and $j_1, \dots, j_p \in [N]$, we define

$$m_{j_1 \dots j_p; k} = N^{-k} \sum_{i_1, \dots, i_k=1}^N m_{i_1 i_2 i_2 \dots i_k i_k j_1 j_2 \dots j_p}. \quad (2.6)$$

Notice that each of the indices i_1, \dots, i_k over which we average occurs exactly twice.

Lemma 2.3. *Assume $(\beta, \mu_h, \sigma_h^2) \in \mathcal{A}_{RS^-}$ and $h \sim \mathcal{N}(\mu_h, \sigma_h^2)$. Set $A_m = m/2$ if $m \leq 2$ and $A_m = m/2 + 1/2$ if $m \geq 3$. Then, we have for every $k \geq 0, p \geq 1$ with $2k + p \geq 2$ that*

$$N^{-p} \sum_{j_1, \dots, j_p=1}^N \mathbb{E} m_{j_1 \dots j_p; k}^2 \leq C_{\beta, h} N^{-A_{2k+p}}$$

for some $C_{\beta, h} > 0$ that is locally bounded in $(\beta, \mu_h, \sigma_h^2)$.

Proof. For simplicity, we give a detailed proof for the cases $k = 0$ and $k = 1$; the remaining cases $k \geq 2$ are proved with analogous arguments. We use the representation

$$m_{j_1 \dots j_p} = \left\langle \sigma_{j_1}^1 \prod_{u=1}^{p-1} \left(\sum_{v=1}^u (\sigma_{j_{u+1}}^v - \sigma_{j_{u+1}}^{u+1}) \right) \right\rangle, \quad (2.7)$$

where we recall that $(\sigma^l)_{l \in \mathbb{N}}$ denote i.i.d. replicas sampled from $\langle \cdot \rangle$. Eq. (2.7) follows from

$$\begin{aligned} \partial_{h_i} \langle f(\sigma^1, \dots, \sigma^u) \rangle &= \sum_{\sigma^1, \dots, \sigma^u \in \{-1, 1\}^N} f(\sigma^1, \dots, \sigma^u) \partial_{h_i} \frac{e^{H_N(\sigma^1) + \dots + H_N(\sigma^u)}}{Z_N^u} \\ &= \left\langle f(\sigma^1, \dots, \sigma^u) \left(\sum_{v=1}^u \sigma_i^v - u \sigma_i^{u+1} \right) \right\rangle \\ &= \left\langle f(\sigma^1, \dots, \sigma^u) \sum_{v=1}^u (\sigma_i^v - \sigma_i^{u+1}) \right\rangle \end{aligned}$$

and induction. Observe that the previous identity implies in particular that

$$\left\langle \prod_{u=1}^{p-1} \left(\sum_{v=1}^u (\sigma_{j_{u+1}}^v - \sigma_{j_{u+1}}^{u+1}) \right) \right\rangle = \partial_{h_{j_2}} \dots \partial_{h_{j_p}} \langle 1 \rangle = 0. \quad (2.8)$$

Now, consider first the case $k = 0$. Recalling that $(\sigma^l)_{l \in \mathbb{Z}}$ denotes a sequence of i.i.d. replicas sampled from $\langle \cdot \rangle$, Eq. (2.7) and (2.8) imply that we can express $m_{j_1 \dots j_p}^2$ as

$$m_{j_1 \dots j_p}^2 = \left\langle (\sigma_{j_1}^1 \sigma_{j_1}^{-1} - N^{-1} \|\mathbf{m}\|^2) \prod_{u=1}^{p-1} \left(\sum_{v=1}^u (\sigma_{j_{u+1}}^v - \sigma_{j_{u+1}}^{u+1}) \right) \left(\sum_{v'=-1}^{-u} (\sigma_{j_{u+1}}^{v'} - \sigma_{j_{u+1}}^{-(u+1)}) \right) \right\rangle$$

so that

$$\begin{aligned} & N^{-p} \sum_{j_1, \dots, j_p=1}^N m_{j_1 \dots j_p}^2 \\ &= \left\langle (R_{1,-1} - N^{-1} \|\mathbf{m}\|^2) \prod_{u=1}^{p-1} \sum_{v=1}^u \sum_{v'=-1}^{-u} (R_{vv'} - R_{v,-(u+1)} - R_{u+1,v'} + R_{u+1,-(u+1)}) \right\rangle \\ &= \left\langle (T_{1,-1} + T_1 + T_{-1}) \prod_{u=1}^{p-1} \sum_{v=1}^u \sum_{v'=-1}^{-u} (T_{vv'} - T_{v',-(u+1)} - T_{u+1,v'} + T_{u+1,-(u+1)}) \right\rangle. \end{aligned}$$

Here, we used in the second step that $R_{l,l'} = T_{l,l'} + T_l + T_{l'} + N^{-1} \|\mathbf{m}\|^2$ and that

$$\begin{aligned} & R_{vv'} - R_{v,-(u+1)} - R_{u+1,v'} + R_{u+1,-(u+1)} \\ &= T_{vv'} + T_v + T_{v'} + N^{-1} \|\mathbf{m}\|^2 - T_{v,-(u+1)} - T_v - T_{-(u+1)} - N^{-1} \|\mathbf{m}\|^2 \\ &\quad - T_{u+1,v'} - T_{u+1} - T_{v'} - N^{-1} \|\mathbf{m}\|^2 + T_{u+1,-(u+1)} + T_{u+1} + T_{-(u+1)} + N^{-1} \|\mathbf{m}\|^2 \\ &= T_{vv'} - T_{v',-(u+1)} - T_{u+1,v'} + T_{u+1,-(u+1)}. \end{aligned}$$

Now, applying and using the same notation as in Theorem 2.2, we get

$$\left| N^{-p} \sum_{j_1, \dots, j_p=1}^N \mathbb{E} m_{j_1 \dots j_p}^2 - N^{-\frac{p}{2}} E \prod_{u=0}^{p-1} Z_u \right| \leq C N^{-\frac{p+1}{2}}, \quad (2.9)$$

where $(Z_u)_{0 \leq u \leq p-1}$ is a Gaussian random vector whose entries are defined by

$$Z_u = \begin{cases} U_{1,-1} + U_1 + U_{-1} & : u = 0, \\ \sum_{v=1}^u \sum_{v'=-1}^{-u} (U_{vv'} - U_{v,-(u+1)} - U_{u+1,v'} + U_{u+1,-(u+1)}) & : u \geq 1. \end{cases}$$

This proves the lemma for the case $k = 0$ and $p = 2$. To obtain an additional decay factor $N^{-1/2}$ for the cases $p \geq 3$, notice that the entries of $(Z_u)_{1 \leq u \leq p-1}$ are independent:

$$\begin{aligned} EZ_u Z_{u'} &= E \sum_{v=1}^u \sum_{v'=-1}^{-u} \sum_{w=1}^{u'} \sum_{w'=-1}^{-u'} (U_{vv'} - U_{v,-(u+1)} - U_{u+1,v'} + U_{u+1,-(u+1)}) U_{ww'} \\ &= \sum_{v=1}^u \sum_{v'=-1}^{-u} (E U_{vv'}^2 - E U_{v,-(u+1)}^2 - E U_{u+1,v'}^2 + E U_{u+1,-(u+1)}^2) = 0 \end{aligned}$$

for all $1 \leq u < u' \leq p-1$. Since $(Z_u)_{0 \leq u \leq p-1}$ is a Gaussian vector, we can then write

$$Z_0 = \sum_{u=1}^{p-1} x_u Z_u + Z',$$

where $\mathbf{x} = (x_1, \dots, x_{p-1}) \in \mathbb{R}^{p-1}$ and where Z' is a Gaussian random variable independent of the remaining entries Z_1, \dots, Z_{p-1} . This implies for $p \geq 3$ that

$$E \prod_{u=0}^{p-1} Z_u = \sum_{u=1}^{p-1} x_u (E Z_u^2) E \prod_{\substack{u'=1, \\ u' \neq u}}^{p-1} Z_{u'} = 0$$

and combining this with (2.9), we conclude that

$$\left| N^{-p} \sum_{j_1, \dots, j_p=1}^N \mathbb{E} m_{j_1 \dots j_p}^2 \right| \leq C N^{-\frac{p+1}{2}}.$$

Next, consider the case $k = 1$. By (2.7), we have that

$$\begin{aligned} m_{j_1 \dots j_p; 1} &= \frac{1}{N} \sum_{i_1=1}^N m_{i_1 i_1 j_1 \dots j_p} = \frac{1}{N} \sum_{i_1=1}^N \left\langle \sigma_{i_1}^1 (\sigma_{i_1}^1 - \sigma_{i_1}^2) \prod_{u=2}^{p+1} \left(\sum_{v=1}^u (\sigma_{j_{u-1}}^v - \sigma_{j_{u-1}}^{u+1}) \right) \right\rangle \\ &= \left\langle (1 - R_{1,2}) \prod_{u=2}^{p+1} \left(\sum_{v=1}^u (\sigma_{j_{u-1}}^v - \sigma_{j_{u-1}}^{u+1}) \right) \right\rangle \\ &= \left\langle (N^{-1} \|\mathbf{m}\|^2 - R_{1,2}) \prod_{u=2}^{p+1} \left(\sum_{v=1}^u (\sigma_{j_{u-1}}^v - \sigma_{j_{u-1}}^{u+1}) \right) \right\rangle, \end{aligned}$$

where we used (2.8) in the last step. Hence

$$\begin{aligned} m_{j_1 \dots j_p; 1}^2 &= \left\langle (N^{-1} \|\mathbf{m}\|^2 - R_{12}) (N^{-1} \|\mathbf{m}\|^2 - R_{-1, -2}) \right. \\ &\quad \left. \times \prod_{u=2}^{p+1} \left(\sum_{v=1}^u (\sigma_{j_{u-1}}^v - \sigma_{j_{u-1}}^{u+1}) \right) \left(\sum_{v'=-1}^{-u} (\sigma_{j_{u-1}}^{v'} - \sigma_{j_{u-1}}^{-(u+1)}) \right) \right\rangle \end{aligned}$$

and thus, arguing as in the first step,

$$\begin{aligned} N^{-p} \sum_{j_1, \dots, j_p=1}^N \mathbb{E} m_{j_1 \dots j_p; 1}^2 &= \mathbb{E} \left\langle (T_1 + T_2 + T_{12}) (T_{-1} + T_{-2} + T_{-1, -2}) \right. \\ &\quad \left. \times \prod_{u=2}^{p+1} \left(\sum_{v=1}^u \sum_{v'=-1}^{-u} (T_{vv'} - T_{v, -(u+1)} - T_{u+1, v'} + T_{u+1, -(u+1)}) \right) \right\rangle. \end{aligned}$$

Applying and using the same notation as in Theorem 2.2, we obtain that

$$\left| N^{-p} \sum_{j_1, \dots, j_p=1}^N \mathbb{E} m_{j_1 \dots j_p; 1}^2 - E(U_1 + U_2 + U_{12})(U_{-1} + U_{-2} + U_{-1, -2}) Y_{1,p} \right| \leq CN^{-\frac{p+3}{2}},$$

where we set

$$Y_{1,p} = \prod_{u=2}^{p+1} \left(\sum_{v=1}^u \sum_{v'=-1}^{-u} (U_{vv'} - U_{v, -(u+1)} - U_{u+1, v'} + U_{u+1, -(u+1)}) \right).$$

Notice in particular that $Y_{1,p}$ is independent of the random variables $(U_{l,l'})_{l,l' \in \mathbb{Z}: ll' > 0}$ and $(U_l)_{l \in \mathbb{Z}}$ and that $EY_{1,p} = 0$, arguing as in the previous step. We therefore conclude that

$$\left| N^{-p} \sum_{j_1, \dots, j_p=1}^N \mathbb{E} m_{j_1 \dots j_p; 1}^2 \right| \leq CN^{-\frac{p+3}{2}} = CN^{-A_{2+p}}.$$

For the remaining cases $k \geq 2$, using the same arguments as above, we find that

$$\left| N^{-p} \sum_{j_1, \dots, j_p=1}^N \mathbb{E} m_{j_1 \dots j_p; k}^2 - E X_{k,p} Y_{k,p} \right| \leq CN^{-\frac{2k+p+1}{2}} = N^{-A_{2k+p}},$$

where

$$Y_{k,p} = \prod_{u=2k}^{p+2k-1} \left(\sum_{v=1}^u \sum_{v'=-1}^{-u} (U_{vv'} - U_{v, -(u+1)} - U_{u+1, v'} + U_{u+1, -(u+1)}) \right)$$

so that by independence $EY_{k,p} = 0$, and where $X_{k,p}$ is a finite polynomial of the random variables $(U_{l,l'})_{l,l' \in \mathbb{Z}: ll' > 0}$ and $(U_l)_{l \in \mathbb{Z}}$ so that $E[X_{k,p} Y_{k,p}] = 0$. \square

We conclude this section with a few corollaries of Lemma 2.3 that provide useful bounds on the correlation functions, conditionally on a fixed spin σ_i . Recall that $\mathbf{m} \in \mathbb{R}^N$ denotes the magnetization vector (i.e. $m_i = \langle \sigma_i \rangle$) and that $\mathbf{m}^{(i)}$ denotes the magnetization of the system after removing σ_i . We write $\langle \cdot \rangle^{(i)}$ for the corresponding Gibbs measure so that

$$\mathbf{m}^{(i)} = (m_1^{(i)}, \dots, m_{i-1}^{(i)}, m_{i+1}^{(i)}, \dots, m_N^{(i)}) = (\langle \sigma_1 \rangle^{(i)}, \dots, \langle \sigma_{i-1} \rangle^{(i)}, \langle \sigma_{i+1} \rangle^{(i)}, \dots, \langle \sigma_N \rangle^{(i)}).$$

Note that $\langle \cdot \rangle^{(i)}$ is a Gibbs measure induced by the SK Hamiltonian as in (1.1), but with $N - 1$ particles and with coupling $\beta' = (1 - 1/N)^{1/2} \beta$ (which is close to β for $N \gg 1$).

More generally, for fixed $i \in [N]$, let us identify $(g_{ij})_{j:j \neq i}$ with $(g_{ij}(1))_{j:j \neq i}$, where the components of $((g_{ij}(t))_{j:j \neq i})_{t \in [0,1]}$ denote $N - 1$ independent Brownian motions of speed

$1/N$, i.e. $\mathbb{E}(g_{ij}(t))^2 = t/N$. Then, for $\sigma_i \in \{-1, 1\}$ and $t \in [0, 1]$, we denote by $\langle \cdot \rangle^{[i]}(\sigma_i, t)$ the Gibbs measure induced by the SK Hamiltonian

$$\begin{aligned} H_{N-1}^{[i]}(\sigma) &= \sum_{\substack{1 \leq u < v \leq N, \\ u, v \neq i}} \beta' g'_{uv} \sigma_u \sigma_v + \sum_{\substack{1 \leq u \leq N: \\ u \neq i}}^N (h_u + \beta g_{iu}(t) \sigma_i) \sigma_u \\ &= \sum_{\substack{1 \leq u < v \leq N, \\ u, v \neq i}} \beta' g'_{uv} \sigma_u \sigma_v + \sum_{\substack{1 \leq u \leq N: \\ u \neq i}}^N h'_u(\sigma_i, t) \sigma_u \end{aligned}$$

for $\sigma = (\sigma_j)_{j \in [N], j \neq i} \in \{-1, 1\}^{N-1}$. Here, the interactions $g'_{uv} = (1 - 1/N)^{-1/2} g_{uv}$ define a GOE matrix $\mathbf{G}' \in \mathbb{R}^{N-1 \times N-1}$ and the random field $(h'_j(\sigma_i, t))_{j \in [N], j \neq i}$ consists of independent Gaussian copies of $h' \sim \mathcal{N}(\mu_{h'}, \sigma_{h'}^2) = \mathcal{N}(\mu_h, \sigma_h^2 + \beta^2 t/N)$. Observe that

$$|\mu_h - \mu_{h'}| = 0, \quad |\beta - \beta'| \rightarrow 0, \quad |\sigma_h^2 - \sigma_{h'}^2| \rightarrow 0 \quad (2.10)$$

as $N \rightarrow \infty$ and that $\langle \cdot \rangle^{[i]}(\sigma_i, 0) = \langle \cdot \rangle^{(i)}$. Moreover, observe that $\langle \cdot \rangle^{[i]}(\cdot, 1)$ corresponds to the Gibbs expectation $(\langle \cdot \rangle | \sigma_i)$ conditionally on σ_i .

In the sequel, we denote by $m_{j_1 \dots j_p}^{[i]} = m_{j_1 \dots j_p}^{[i]}(\sigma_i, t)$ the k -point and by $m_{j_1 \dots j_p; k}^{[i]} = m_{j_1 \dots j_p; k}^{[i]}(\sigma_i, t)$ the (k, p) -point correlation functions with regards to $\langle \cdot \rangle^{[i]}$, defined as in (2.5) and (2.6). Following [1], we finally define for observables $f : \{-1, 1\}^{N-1} \rightarrow \mathbb{R}$

$$\begin{aligned} \delta_i \langle f \rangle^{[i]} &= (\delta_i \langle f \rangle^{[i]})(t) = \frac{1}{2} \langle f \rangle^{[i]}(1, t) - \frac{1}{2} \langle f \rangle^{[i]}(-1, t), \\ \epsilon_i \langle f \rangle^{[i]} &= (\epsilon_i \langle f \rangle^{[i]})(t) = \frac{1}{2} \langle f \rangle^{[i]}(1, t) + \frac{1}{2} \langle f \rangle^{[i]}(-1, t), \\ \Delta_i \langle f \rangle^{[i]} &= (\Delta_i \langle f \rangle^{[i]})(t) = (\epsilon_i \langle f \rangle^{[i]})(t) - \langle f \rangle^{(i)}. \end{aligned}$$

The next lemmas estimate certain correlation functions, conditionally on σ_i . To ease the notation, we write \sum and understand implicitly that all variables are averaged over $[N] \setminus \{i\}$.

Lemma 2.4. *Assume $(\beta, \mu_h, \sigma_h^2) \in \mathcal{A}_{RS}$ and $h \sim \mathcal{N}(\mu_h, \sigma_h^2)$. Then, for every $k \geq 0, p \geq 1$ with $2k + p \geq 1$, there exists a constant $C > 0$ such that*

$$\begin{aligned} \sup_{t \in [0, 1]} N^{-p} \sum_{j_1, \dots, j_p} \left\| (\Delta_i m_{j_1 \dots j_p; k}^{[i]})(t) \right\|_2^2 &\leq C N^{-A_{2k+p+2}}, \\ \sup_{t \in [0, 1]} N^{-p} \sum_{j_1, \dots, j_p} \left\| (\delta_i m_{j_1 \dots j_p; k}^{[i]})(t) \right\|_2^2 &\leq C N^{-A_{2k+p+1}}. \end{aligned}$$

Proof. Applying Itô's lemma w.r.t. $((g_{ij}(t))_{j \in [N]: j \neq i})_{t \in [0,1]}$, we find by definition (2.6) that

$$\begin{aligned} d(\Delta_i m_{j_1 \dots j_p; k}^{[i]}) &= \beta \sum_l (\delta_i m_{l j_1 \dots j_p; k}^{[i]}) dg_{il} + \frac{\beta^2}{2} (1 - N^{-1}) (\epsilon_i m_{j_1 \dots j_p; k+1}^{[i]}) dt, \\ d(\delta_i m_{j_1 \dots j_p; k}^{[i]}) &= \beta \sum_l (\epsilon_i m_{l j_1 \dots j_p; k}^{[i]}) dg_{il} + \frac{\beta^2}{2} (1 - N^{-1}) (\delta_i m_{j_1 \dots j_p; k+1}^{[i]}) dt. \end{aligned}$$

Notice that the integrands on the r.h.s. in the previous equations are linear combinations of suitable (k, p) -point functions with interaction coupling and external field parameters all satisfying (1.3) & (1.5) for N large (by the remarks around (2.10)) s.t. by Lemma 2.3

$$\begin{aligned} \sup_{t \in [0,1]} N^{-p-1} \sum_{l, j_1, \dots, j_p} \|(\epsilon_i m_{l j_1 \dots j_p; k}^{[i]})(t)\|_2^2 &\leq CN^{-A_{2k+p+1}}, \\ \sup_{t \in [0,1]} N^{-p} \sum_{j_1, \dots, j_p} \|(\delta_i m_{j_1 \dots j_p; k+1}^{[i]})(t)\|_2^2 &\leq CN^{-A(2k+2+p)}. \end{aligned}$$

Hence, employing the Itô isometry yields

$$\sup_{t \in [0,1]} N^{-p} \sum_{j_1, \dots, j_p} \|(\delta_i m_{j_1 \dots j_p; k}^{[i]})(t)\|_2^2 \leq CN^{-A_{2k+p+1}}$$

and plugging this into the dynamical equation for $\Delta_i m_{j_1 \dots j_p; k}^{[i]}$, we get

$$\sup_{t \in [0,1]} N^{-p} \sum_{j_1, \dots, j_p} \|(\Delta_i m_{j_1 \dots j_p; k}^{[i]})(t)\|_2^2 \leq CN^{-A_{2k+p+2}}.$$

□

Lemma 2.5. *Assume $(\beta, \mu_h, \sigma_h^2) \in \mathcal{A}_{RS-}$ and $h \sim \mathcal{N}(\mu_h, \sigma_h^2)$. Then, for every $k \geq 0, p \geq 1$ with $2k + p \geq 1$, there exists a constant $C > 0$ such that*

$$\begin{aligned} \sup_{t \in [0,1]} N^{-p} \sum_{j_1, \dots, j_p} \left\| \sum_j g_{ij}(t) (\Delta_i m_{j_1 \dots j_p; k}^{[i]})(t) \right\|_2^2 &\leq CN^{-A_{2k+p+3}}, \\ \sup_{t \in [0,1]} N^{-p} \sum_{j_1, \dots, j_p} \left\| \sum_j g_{ij}(t) (\delta_i m_{j_1 \dots j_p; k}^{[i]})(t) \right\|_2^2 &\leq CN^{-A_{2k+p+2}}. \end{aligned} \tag{2.11}$$

Proof. The bounds follow from integration by parts applied to $g_{ij}(t)$ and $g_{ij'}(t)$ in

$$N^{-p} \sum_{j_1, \dots, j_p} \sum_{j, j'} \mathbb{E} g_{ij} g_{ij'} (\Delta_i m_{j_1 \dots j_p; k}^{[i]})(\Delta_i m_{j_1 \dots j_p; k}^{[i]})$$

and applying Lemma 2.4. It will be clear that the bounds are uniform in $t \in [0, 1]$ so for simplicity, let us ignore the t -dependence in the notation. Applying Gaussian integration by parts first to the $g_{ij'}$ implies

$$\begin{aligned}
& N^{-p} \sum_{j_1, \dots, j_p} \sum_{j, j'} \mathbb{E} g_{ij} g_{ij'} (\Delta_i m_{jj_1 \dots j_p; k}^{[i]}) (\Delta_i m_{j'j_1 \dots j_p; k}^{[i]}) \\
&= N^{-p} \sum_{j_1, \dots, j_p=1}^N \left(\frac{t}{N} \sum_{j=1}^N \mathbb{E} (\Delta_i m_{jj_1 \dots j_p; k}^{[i]})^2 \right. \\
&\quad + \beta^2 t (1 - N^{-1}) \sum_{j=1}^N \mathbb{E} g_{ij} (\Delta_i m_{jj_1 \dots j_p; k}^{[i]}) (\Delta_i m_{j_1 \dots j_p; k+1}^{[i]}) \\
&\quad \left. + \beta^2 \frac{t}{N} \sum_{j, j'=1}^N \mathbb{E} g_{ij} (\Delta_i m_{jj'j_1 \dots j_p; k}^{[i]}) (\Delta_i m_{j'j_1 \dots j_p; k}^{[i]}) \right)
\end{aligned}$$

and then, applying Gaussian integration by parts to the g_{ij} , we arrive at

$$\begin{aligned}
& N^{-p} \sum_{j_1, \dots, j_p} \sum_{j, j'} \mathbb{E} g_{ij} g_{ij'} (\Delta_i m_{jj_1 \dots j_p; k}^{[i]}) (\Delta_i m_{j'j_1 \dots j_p; k}^{[i]}) \\
&= N^{-p} \sum_{j_1, \dots, j_p} \left(\frac{t}{N} \sum_j \mathbb{E} (\Delta_i m_{jj_1 \dots j_p; k}^{[i]})^2 + \beta^4 t^2 (1 - N^{-1}) \mathbb{E} (\Delta_i m_{j_1 \dots j_p; k+1}^{[i]})^2 \right. \\
&\quad + \frac{\beta^4 t^2}{N} (1 - N^{-1}) \sum_j \mathbb{E} (\Delta_i m_{jj_1 \dots j_p; k}^{[i]}) (\Delta_i m_{jj_1 \dots j_p; k+1}^{[i]}) \\
&\quad + \frac{\beta^4 t^2}{N} (1 - N^{-1}) \sum_{j'} \mathbb{E} (\Delta_i m_{j'j_1 \dots j_p; k+1}^{[i]}) (\Delta_i m_{j'j_1 \dots j_p; k}^{[i]}) \\
&\quad \left. + \frac{\beta^4 t^2}{N^2} \sum_{j, j'} \mathbb{E} (\Delta_i m_{jj'j_1 \dots j_p; k}^{[i]})^2 \right).
\end{aligned}$$

Applying Lemma 2.4 and Cauchy-Schwarz, we thus obtain

$$\sup_{t \in [0, 1]} N^{-p} \sum_{j_1, \dots, j_p} \left\| \sum_j g_{ij}(t) (\Delta_i m_{jj_1 \dots j_p; k}^{[i]})(t) \right\|_2^2 \leq C N^{-A_{2k+p+3}}.$$

Repeating the computation while replacing Δ_i by δ_i in all integrands, we obtain (2.11). \square

3 Proof of Propositions 1.1 and 1.2

We start with the proof of Prop. 1.1 which is a simple consequence of Lemma 2.4.

Proof of Prop. 1.1. A direct computation shows that

$$m_{ij} = (1 - m_i^2) \delta_i m_j^{[i]}.$$

Now, applying Itô's lemma as in the previous section on the i -th row of \mathbf{G} , we find

$$\begin{aligned} \frac{m_{ij}}{1 - m_i^2} - \beta \sum_l g_{il} m_{lj}^{(i)} &= \beta \sum_k \int_0^1 (\Delta_i m_{jk}^{[i]})(t) dg_{ik}(t) + \frac{\beta^2}{2N} \sum_k \int_0^1 (\delta_i m_{kkj}^{[i]})(t) dt \\ &= \beta \sum_k \int_0^1 (\Delta_i m_{jk}^{[i]})(t) dg_{ik}(t) + \frac{\beta^2}{2} (1 - N^{-1}) \int_0^1 (\delta_i m_{j;1}^{[i]})(t) dt. \end{aligned}$$

Using that $(1 - m_i^2) \leq 1$ and applying Lemma 2.4, we conclude that

$$\begin{aligned} &\left\| \frac{m_{ij}}{1 - m_i^2} - \beta \sum_{l \neq i} g_{il} m_{lj}^{(i)} \right\|_2^2 \\ &\leq \frac{C}{N} \sum_k \sup_{t \in [0,1]} \|(\Delta_i m_{jk}^{[i]})(t)\|_2^2 + C \sup_{t \in [0,1]} \|(\delta_i m_{j;1}^{[i]})(s)\|_2^2 \\ &= C \sup_{t \in [0,1]} \left(\frac{1}{(N-1)^2} \sum_{j_1, j_2} \|(\Delta_i m_{j_1 j_2}^{[i]})(t)\|_2^2 + \frac{1}{N-1} \sum_j \|(\delta_i m_{j;1}^{[i]})(s)\|_2^2 \right) \leq CN^{-5/2}. \end{aligned} \tag{3.1}$$

Notice that in the second step of the last bound we used the symmetry among the sites $j \in [N] \setminus \{i\}$, in order to average over both indices of the two point functions. \square

We conclude this section with the proof of Prop. 1.2. Our goal is to show that

$$\mathbf{Y} = (\mathbf{D} + \beta^2(1 - q) - \beta \mathbf{G}) \mathbf{M} - \text{id}_{\mathbb{R}^N} \in \mathbb{R}^{N \times N}$$

with entries

$$y_{ik} = \begin{cases} (1 - m_i^2)(\beta^2(1 - q) - \beta \sum_j g_{ij} \delta_i m_j^{[i]}) & : i = k, \\ (1 - m_i^2)^{-1} m_{ik} + \beta^2(1 - q) m_{ik} - \beta \sum_j g_{ij} m_{jk} & : i \neq k, \end{cases} \tag{3.2}$$

has operator norm $\|\mathbf{Y}\|_{\text{op}}$ bounded by $o(1)\|\mathbf{M}\|_{\text{op}}$, up to a quantity $O(1)$ which is of order one with high probability. Our proof relies on Prop. 1.1. In the following, the standard Frobenius (or Hilbert-Schmidt) norm is denoted by $\|\cdot\|_{\text{F}}$; recall that $\|\cdot\|_{\text{op}} \leq \|\cdot\|_{\text{F}}$.

Proof of Prop. 1.2. A direct computation shows that

$$m_{jk} = \langle m_j^{[i]}; m_k^{[i]} \rangle + \langle m_{jk}^{[i]} \rangle = (\delta_i m_j^{[i]}) m_{ik} + \epsilon_i m_{jk}^{[i]} + m_i (\delta_i m_{jk}^{[i]})$$

so that, motivated by Prop. 1.1, we can decompose y_{ik} for $i \neq k$ into

$$\begin{aligned} y_{ik} &= \frac{m_{ik}}{1 - m_i^2} - \beta \sum_j g_{ij} m_{jk}^{(i)} - \beta \sum_j g_{ij} (m_{jk} - m_{jk}^{(i)}) + \beta^2 (1 - q) m_{ik} \\ &= \left(\beta^2 (1 - q) - \beta \sum_j g_{ij} \delta_i m_j^{[i]} \right) m_{ik} + \left(\frac{m_{ik}}{1 - m_i^2} - \beta \sum_j g_{ij} m_{jk}^{(i)} \right) \\ &\quad - \beta \sum_j g_{ij} (\Delta_i m_{jk}^{[i]}) - \beta m_i \sum_j g_{ij} (\delta_i m_{jk}^{[i]}). \end{aligned}$$

Comparing this with the diagonal entries of \mathbf{Y} in Eq. (3.2), we can split

$$\mathbf{Y} = \mathbf{Y}_1 \mathbf{M} + \mathbf{Y}_2 - \mathbf{Y}_3 - \mathbf{Y}_4,$$

where the $\mathbf{Y}_j \in \mathbb{R}^{N \times N}$, for $j \in \{1, 2, 3, 4\}$, are defined by

$$\begin{aligned} (\mathbf{Y}_1)_{ik} &= \left(\beta^2 (1 - q) - \beta \sum_j g_{ij} \delta_i m_j^{[i]} \right) \delta_{ik}, & (\mathbf{Y}_2)_{ik} &= \begin{cases} 0 & : i = k, \\ \frac{m_{ik}}{1 - m_i^2} - \beta \sum_j g_{ij} m_{jk}^{(i)} & : i \neq k, \end{cases} \\ (\mathbf{Y}_3)_{ik} &= \begin{cases} 0 & : i = k, \\ \beta \sum_j g_{ij} (\Delta_i m_{jk}^{[i]}) & : i \neq k, \end{cases} & (\mathbf{Y}_4)_{ik} &= \begin{cases} 0 & : i = k, \\ \beta m_i \sum_j g_{ij} (\delta_i m_{jk}^{[i]}) & : i \neq k. \end{cases} \end{aligned}$$

The matrix \mathbf{Y}_2 vanishes in norm when $N \rightarrow \infty$, in the sense of probability, by Eq. (3.1) and Markov's inequality: for every $\delta > 0$, we find that

$$\mathbb{P}(\|\mathbf{Y}_2\|_{\text{op}} > \delta) \leq \delta^{-2} \mathbb{E} \|\mathbf{Y}_2\|_{\text{op}}^2 \leq \delta^{-2} \mathbb{E} \|\mathbf{Y}_2\|_{\text{F}}^2 = N^2 \max_{i,j \in [N]: i \neq j} \mathbb{E} (\mathbf{Y}_2)_{ij}^2 \leq C \delta^{-2} N^{-1/2}.$$

By Lemma 2.5, the same argument implies $\lim_{N \rightarrow \infty} \|\mathbf{Y}_3\|_{\text{op}} = 0$, and we have that

$$\begin{aligned} \mathbb{P}(\|\mathbf{Y}_4\|_{\text{op}} > K) &\leq K^{-2} \beta^2 N^2 \max_{i,j \in [N]: i \neq j} \mathbb{E} (\mathbf{Y}_4)_{ij}^2 \\ &\leq C K^{-2} N^2 N^{-1} \sum_{j_1} \left\| \sum_j g_{1j} (\delta_1 m_{jj_1}^{[1]}) \right\|_2^2 \leq C K^{-2}, \end{aligned}$$

for some $C > 0$, independent of N , where we used symmetry among the sites to reduce the computation to the first row of \mathbf{G} . Combining these remarks, Prop. 1.2 follows if

$$\lim_{N \rightarrow \infty} \|\mathbf{Y}_1\|_{\text{op}} = 0 \tag{3.3}$$

in probability.

To prove (3.3), we argue first as in (3.1) to obtain that

$$\mathbb{E} \left| \sum_j g_{ij} \left(\delta_i m_j^{[i]} - \beta \sum_k g_{ik} m_{kj}^{(i)} \right) \right|^2 \leq C N^{-5/2}.$$

Indeed, this bound is a direct consequence of Itô's lemma, which shows that

$$\begin{aligned} d \sum_j g_{ij} \left(\delta_i m_j^{[i]} - \beta \sum_k g_{ik} m_{kj}^{(i)} \right) &= \beta \sum_j \left(\delta_i m_j^{[i]} - \beta \sum_k g_{ik} m_{kj}^{(i)} \right) dg_{ij} + \beta \sum_{j,k} g_{ij} (\Delta_i m_{jk}^{[i]}) dg_{ik} \\ &\quad + \frac{\beta^2}{2} (1 - N^{-1}) \sum_j g_{ij} (\delta_i m_{j;1}^{[i]}) dt + \frac{\beta}{N} \sum_j (\Delta_i m_{jk}^{[i]}) dt, \end{aligned}$$

and estimating the different contributions on the r.h.s. of the previous identity using Lemmas 2.4 and 2.5. Combining the previous bound with Markov thus implies

$$\lim_{N \rightarrow \infty} \sup_{i \in [N]} \left| (\mathbf{Y}_1)_{ii} - \left(\beta^2 (1 - q) - \beta^2 \sum_j g_{ij} \sum_k g_{ik} m_{kj}^{(i)} \right) \right| = 0$$

in probability.

We now split

$$\begin{aligned} &\beta^2 (1 - q) - \beta^2 \sum_j g_{ij} \sum_k g_{ik} m_{kj}^{(i)} \\ &= \beta^2 \langle R_{1,2} - q \rangle^{(i)} - \beta^2 \sum_j (g_{ij}^2 - (N - 1)^{-1}) (1 - (m_j^{(i)})^2) - \beta^2 \sum_{j,k:j \neq k} g_{ij} g_{ik} m_{kj}^{(i)}, \end{aligned}$$

and it is straightforward to deduce from Theorem 2.1 that

$$\lim_{N \rightarrow \infty} \sup_{i \in [N]} \left| \langle R_{1,2} - q \rangle^{(i)} \right| = 0$$

in probability. Indeed, this follows from Markov's inequality combined with Theorem 2.1, the continuity of $(\beta, \mu_h, \sigma_h^2) \rightarrow q_{\beta,h} = q_{\beta, \mu_h, \sigma_h^2}$ and the fact that $\langle \cdot \rangle^{(i)}$ is a SK measure with interaction coupling $\beta' = \beta'_N$ and Gaussian external field $h' = h'_N$ such that $|\beta - \beta'| \rightarrow 0$, $|\sigma_h^2 - \sigma_{h'}^2| \rightarrow 0$ as $N \rightarrow \infty$ and $|\mu_h - \mu_{h'}| = 0$, as remarked already at (2.10).

On the other hand, a standard exponential concentration bound yields

$$\lim_{N \rightarrow \infty} \sup_{i \in [N]} \left| \sum_j (g_{ij}^2 - (N - 1)^{-1}) (1 - (m_j^{(i)})^2) \right| = \lim_{N \rightarrow \infty} \sup_{i \in [N]} \left| \sum_j (g_{ij}^2 - N^{-1}) \right| = 0$$

and, finally, applying once more Theorem 2.1, a straightforward computation shows that

$$\mathbb{E} \left(\sum_{j,k:j \neq k} g_{ij} g_{ik} m_{kj}^{(i)} \right)^4 \leq CN^{-4} \sum_{j_1, j_2, j_3, j_4} \mathbb{E} (m_{j_1 j_2}^{(i)})^2 (m_{j_3 j_4}^{(i)})^2.$$

Then, using the identity

$$(m_{j_1 j_2}^{(i)})^2 = \langle (\sigma_{j_1}^1 - \sigma_{j_1}^2) (\sigma_{j_2}^1 - \sigma_{j_2}^3) (\sigma_{j_1}^{-1} - \sigma_{j_1}^{-2}) (\sigma_{j_2}^{-1} - \sigma_{j_2}^{-3}) \rangle^{(i)}$$

together with Theorem 2.1 and Cauchy-Schwarz, we arrive at

$$\mathbb{E} \left(\sum_{j,k:j \neq k} g_{ij} g_{ik} m_{kj}^{(i)} \right)^4 \leq C \mathbb{E} \left(\langle (R_{1,-1} + R_{2,-2} - R_{1,-2} - R_{2,-1})^2 \rangle^{(i)} \right)^2 \leq CN^{-2}.$$

Hence, by Markov's inequality, we obtain that

$$\lim_{N \rightarrow \infty} \sup_{i \in [N]} \left| \sum_{j,k:j \neq k} g_{ij} g_{ik} m_{kj}^{(i)} \right| = 0,$$

in probability. This proves $\lim_{N \rightarrow \infty} \|\mathbf{Y}_1\|_{\text{op}} = 0$ and concludes Prop. 1.2. \square

4 Proof of Proposition 1.3

In this section we conclude Theorem 1.4 by proving Prop. 1.3. Our arguments rely on the main results of [5, 6, 14] and translate, up to a few modifications, the main ideas from [9] to the present context. For the rest of this paper, $h > 0$ denotes a deterministic field strength as in the definition of H_N in Eq. (1.1) in Section 1.

We first recall the main result of [14], which states that, in a suitable subregion of the replica symmetric phase, the magnetization vector \mathbf{m} is well approximated by an iterative solution to the TAP [2] equations introduced by Bolthausen in [5, 6]. To state this precisely, we need to recall Bolthausen's construction of the TAP solution and collect some of its properties. Here, we find it convenient to follow the conventions and notation used in [6, 8]. We start with the sequences $(\alpha_k)_{k \in \mathbb{N}}$, $(\gamma_k)_{k \in \mathbb{N}}$ and $(\Gamma_k)_{k \in \mathbb{N}}$ which have initializations

$$\alpha_1 = \sqrt{q}\gamma_1, \quad \gamma_1 = E \tanh(h + \beta\sqrt{q}Z), \quad \Gamma_1^2 = \gamma_1^2$$

and we define $\psi : [0, q] \rightarrow [0, q]$ by

$$\psi(t) = E \tanh(h + \beta\sqrt{t}Z + \beta\sqrt{q-t}Z') \tanh(h + \beta\sqrt{t}Z + \beta\sqrt{q-t}Z'')$$

Then, we set recursively

$$\alpha_k = \psi(\alpha_{k-1}), \quad \gamma_k = \frac{\alpha_k - \Gamma_{k-1}^2}{\sqrt{q - \Gamma_{k-1}^2}}, \quad \Gamma_k^2 = \sum_{j=1}^k \gamma_j^2.$$

Lemma 4.1. (*[5, Lemma 2.2, Corollary 2.3, Lemma 2.4], [6, Lemma 2]*)

- 1) ψ is strictly increasing and convex in $[0, q]$ with $0 < \psi(0) < \psi(q) = q$. If (1.3) is satisfied, then q is the unique fixed point of ψ in $[0, q]$.
- 2) $(\alpha_k)_{k \in \mathbb{N}}$ is increasing and $\alpha_k > 0$ for all $k \in \mathbb{N}$. If (1.3) is satisfied, then $\lim_{k \rightarrow \infty} \alpha_k = q$.

3) For $k \geq 2$, we have that $0 < \Gamma_{k-1}^2 < \alpha_k < q$ and that $0 < \gamma_k < \sqrt{q - \Gamma_{k-1}^2}$. If (1.3) is satisfied, then $\lim_{k \rightarrow \infty} \Gamma_k^2 = q$ and, consequently, $\lim_{k \rightarrow \infty} \gamma_k = 0$.

Next, we recall Bolthausen's decomposition of the interaction matrix \mathbf{G} , which yields a convenient representation of the iterative TAP solution. Without loss of generality, we may assume that the interaction matrix $\mathbf{G} \in \mathbb{R}^{N \times N}$ in (1.1) is equal to

$$\mathbf{G} = \frac{\mathbf{W} + \mathbf{W}^T}{\sqrt{2}} = \bar{\mathbf{W}}$$

for a random matrix $\mathbf{W} = (w_{ij})_{1 \leq i, j \leq N} \in \mathbb{R}^{N \times N}$ with zero diagonal and i.i.d. Gaussian entries $w_{ij} \sim \mathcal{N}(0, N^{-1})$ (without symmetry constraint). Here and in the following, we abbreviate $\bar{\mathbf{X}} = (\mathbf{X} + \mathbf{X}^T)/\sqrt{2}$ for $\mathbf{X} \in \mathbb{R}^{N \times N}$. Then, we set

$$\mathbf{W}^{(1)} = \mathbf{W}, \quad \mathbf{G}^{(1)} = \bar{\mathbf{W}}^{(1)}, \quad \phi^{(1)} = N^{-1/2} \mathbf{1} \in \mathbb{R}^N, \quad \mathbf{m}^{(1)} = \sqrt{q} \mathbf{1} \in \mathbb{R}^N$$

and, assuming $\mathbf{W}^{(s)}, \mathbf{G}^{(s)} = \bar{\mathbf{W}}^{(s)}, \phi^{(s)}, \mathbf{m}^{(s)}$ are defined for $1 \leq s \leq k$, we set³

$$\zeta^{(s)} = \mathbf{G}^{(s)} N^{1/2} \phi^{(s)}.$$

Moreover, \mathcal{G}_k denotes the σ -algebra

$$\mathcal{G}_k = \sigma(\mathbf{W}^{(s)} N^{1/2} \phi^{(s)}, (\mathbf{W}^{(s)})^T N^{1/2} \phi^{(s)} : 1 \leq s \leq k).$$

Conditional expectations and respectively probabilities with respect to \mathcal{G}_k are denoted by \mathbb{E}_k and \mathbb{P}_k . We then define the iterative cavity field $\mathbf{z}^{(k+1)} \in \mathbb{R}^N$ and the iterative TAP solution $\mathbf{m}^{(k+1)} \in \mathbb{R}^N$ at step $k+1$ by

$$\mathbf{z}^{(k+1)} = \sum_{s=1}^{k-1} \gamma_s \zeta^{(s)} + \sqrt{q - \Gamma_{k-1}^2} \zeta^{(k)}, \quad \mathbf{m}^{(k+1)} = \tanh(h \mathbf{1} + \beta \mathbf{z}^{(k+1)}) \quad (4.1)$$

and we also set

$$\phi^{(k+1)} = \frac{\mathbf{m}^{(k+1)} - \sum_{s=1}^k (\mathbf{m}^{(k+1)}, \phi^{(s)}) \phi^{(s)}}{\|\mathbf{m}^{(k+1)} - \sum_{s=1}^k (\mathbf{m}^{(k+1)}, \phi^{(s)}) \phi^{(s)}\|},$$

recalling that $\phi^{(k+1)}$ is well-defined for all $k < N$ [6, Lemma 5]. Finally, we define

$$\begin{aligned} \mathbf{W}^{(k+1)} &= \mathbf{W}^{(k)} - \rho^{(k)}, \quad \text{for} \\ \rho^{(k)} &= \mathbf{W}^{(k)} \phi^{(k)} \otimes \phi^{(k)} + \phi^{(k)} \otimes (\mathbf{W}^{(k)})^T \phi^{(k)} - (\mathbf{W}^{(k)} \phi^{(s)}, \phi^{(k)}) \phi^{(k)} \otimes \phi^{(k)}, \end{aligned}$$

³Notice that here we use the convention that $(\phi^{(s)})_{s=1}^k$ forms an orthonormal sequence in $(\mathbb{R}^N, \langle \cdot, \cdot \rangle)$. In contrast to that, in [6, 8] the inner product $\langle \cdot, \cdot \rangle$ and the tensor product \otimes are rescaled by a factor N^{-1} .

where $(\mathbf{x} \otimes \mathbf{y})(\mathbf{z}) = (\mathbf{y}, \mathbf{z}) \mathbf{x}$, and by symmetrization

$$\begin{aligned} \mathbf{G}^{(k+1)} &= \bar{\mathbf{W}}^{(k+1)} = \mathbf{G}^{(k)} - \bar{\rho}^{(k)}, \quad \text{for} \\ \bar{\rho}^{(k)} &= N^{-1/2} \zeta^{(k)} \otimes \phi^{(k)} + N^{-1/2} \phi^{(k)} \otimes \zeta^{(k)} - N^{-1/2} (\zeta^{(k)}, \phi^{(k)}) \phi^{(k)} \otimes \phi^{(k)}. \end{aligned}$$

$(\phi^{(s)})_{s=1}^k$ is orthonormal in $(\mathbb{R}^N, (\cdot, \cdot))$ and $\mathbf{P}^{(k)}, \mathbf{Q}^{(k)}$ denote the orthogonal projections

$$\mathbf{P}^{(k)} = \sum_{s=1}^k \phi^{(s)} \otimes \phi^{(s)} = (P_{ij}^{(k)})_{1 \leq i, j \leq N}, \quad \mathbf{Q}^{(k)} = \mathbf{1}_{\mathbb{R}^N} - \mathbf{P}^{(k)} = (Q_{ij}^{(k)})_{1 \leq i, j \leq N}.$$

Notice that $\mathbf{P}^{(k)}$ equals the orthogonal projection onto

$$\text{span}(\mathbf{m}^{(s)} : 1 \leq s \leq k) = \text{span}(\phi^{(s)} : 1 \leq s \leq k).$$

In the next result we collect several useful facts about $(\mathbf{z}^{(s)})_{s=1}^k, (\mathbf{m}^{(s)})_{s=1}^k$ and $(\mathbf{G}^{(s)})_{s=1}^k$. We say for $(X_N)_{N \geq 1}, (Y_N)_{N \geq 1}$ that may depend on parameters like β, h , etc. that

$$X_N \simeq Y_N$$

if and only if there exist positive constants $c, C > 0$, which may depend on the parameters, but which are independent of N , such that for every $t > 0$ we have

$$\mathbb{P}(|X_N - Y_N| > t) \leq C e^{-cNt^2}.$$

Proposition 4.2. ([5, Prop. 2.5], [6, Prop. 4, Prop. 6, Lemmas 3, 11, 14 & 16])

1) $\mathbf{m}^{(k)}$ and $\phi^{(k)}$ are \mathcal{G}_{k-1} -measurable for every $k \geq 1$ and

$$\mathbf{W}^{(k)} \phi^{(s)} = (\mathbf{W}^{(k)})^T \phi^{(s)} = \mathbf{G}^{(k)} \phi^{(s)} = 0, \quad \forall s < k.$$

2) Conditionally on \mathcal{G}_{k-2} , $\mathbf{W}^{(k)}$ and $\mathbf{W}^{(k-1)}$ are Gaussian, and we have that

$$\mathbb{E}_{k-2} w_{ij}^{(k)} w_{st}^{(k)} = \frac{1}{N} Q_{is}^{(k-1)} Q_{jt}^{(k-1)}$$

and, consequently, that

$$\mathbb{E}_{k-2} g_{ij}^{(k)} g_{st}^{(k)} = \frac{1}{N} Q_{is}^{(k-1)} Q_{jt}^{(k-1)} + \frac{1}{N} Q_{it}^{(k-1)} Q_{js}^{(k-1)}.$$

3) Conditionally on \mathcal{G}_{k-2} , $\mathbf{W}^{(k)}$ is independent of \mathcal{G}_{k-1} . In particular, conditionally on \mathcal{G}_{k-1} , $\mathbf{W}^{(k)}$ and $\mathbf{G}^{(k)}$ are Gaussian with the same covariance as in 2).

4) For every $k \geq 1$, $N^{-1/2}(\zeta^{(k)}, \phi^{(k)})$ is unconditionally Gaussian with variance $2/N$.

5) Conditionally on \mathcal{G}_{k-1} , the random variables $\zeta^{(k)}$ are Gaussian with

$$\mathbb{E}_{k-1} \zeta_i^{(k)} \zeta_j^{(k)} = Q_{ij}^{(k-1)} + \phi_i^{(k)} \phi_j^{(k)}.$$

6) For every $k \geq 1$ and $s < k$, one has

$$\begin{aligned} N^{-1/2}(\mathbf{m}^{(k)}, \phi^{(s)}) &\simeq \gamma_s, & N^{-1/2}(\mathbf{m}^{(k)}, \phi^{(k)}) &\simeq \sqrt{q - \Gamma_{k-1}^2}, \\ N^{-1}(\mathbf{m}^{(k)}, \mathbf{m}^{(s)}) &\simeq \alpha_s, & N^{-1}(\mathbf{m}^{(k)}, \mathbf{m}^{(k)}) &\simeq q. \end{aligned}$$

In particular, $\lim_{N \rightarrow \infty} N^{-1} \mathbb{E} \|\mathbf{m}^{(k)}\|^2 = q$ and, assuming (1.3), we have that

$$\limsup_{j, k \rightarrow \infty} \lim_{N \rightarrow \infty} N^{-1} \mathbb{E} \|\mathbf{m}^{(j)} - \mathbf{m}^{(k)}\|^2 = 0.$$

7) For every $k \geq 1$, we have that

$$\|N^{-1/2}(\zeta^{(k)}, \mathbf{m}^{(k+1)}) - \beta(1-q)\sqrt{q - \Gamma_{k-1}^2}\|_2 = 0,$$

and, for $1 \leq s \leq k-1$, we have that

$$\|N^{-1/2}(\zeta^{(s)}, \mathbf{m}^{(k+1)}) - \beta(1-q)\gamma_s\|_2 = 0.$$

In addition to the properties listed in Prop. 4.2 it is well-known that the sequences $(\mathbf{m}^{(s)})_{s=1}^k$ and $(\mathbf{z}^{(s)})_{s=2}^{k+1}$ have an explicit joint limiting law, which we recall in the next proposition. In its statement, convergence of a sequence $(\mu_n)_{n \in \mathbb{N}}$ of probability measures on \mathbb{R}^d , $\mu_n \in \mathcal{P}(\mathbb{R}^d)$ for each $n \in \mathbb{N}$, to a limiting measure $\mu \in \mathcal{P}(\mathbb{R}^d)$ in $\mathcal{W}_2(\mathbb{R}^d)$ means that

$$\lim_{n \rightarrow \infty} \mathcal{W}_2(\mu_n, \mu) = 0, \quad \text{where} \quad \mathcal{W}_2(\mu, \nu) = \inf_{\Pi} \sqrt{E\|\mathbf{X} - \mathbf{Y}\|^2}$$

denotes the usual Wasserstein 2-distance between two probability measures $\mu, \nu \in \mathcal{P}(\mathbb{R}^d)$ (the infimum is taken over all couplings $\Pi \in \mathcal{P}(\mathbb{R}^{2d})$ of μ and ν , and (\mathbf{X}, \mathbf{Y}) has joint distribution $(\mathbf{X}, \mathbf{Y}) \sim \Pi$). The next theorem follows from combining [22, 5, 6].

Theorem 4.3. *In the sense of probability (w.r.t. the disorder \mathbf{G}), we have that*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \delta_{m_i^{(1)}, \dots, m_i^{(k)}, z_i^{(2)}, \dots, z_i^{(k+1)}} = \mathcal{L}_{(M_1, \dots, M_k, Z_2, \dots, Z_{k+1})}$$

in $\mathcal{W}_2(\mathbb{R}^{2k})$, where $M_1 = \sqrt{q}$, $M_s \stackrel{s \geq 1}{=} \tanh(h + \beta Z_s)$, $(Z_2, \dots, Z_{k+1}) \sim \mathcal{N}(\mathbf{0}, \mathbf{K}_{\leq k})$ is a Gaussian vector in \mathbb{R}^k and $\mathcal{L}_{(M_1, \dots, M_k, Z_2, \dots, Z_{k+1})}$ is the law of $(M_1, \dots, M_k, Z_2, \dots, Z_{k+1})$. The covariance $\mathbf{K}_{\leq k} \in \mathbb{R}^{k \times k}$ equals $K_{st} = \text{Cov}(Z_{s+1}, Z_{t+1}) = EM_s M_t$; in particular

$$K_{st} \stackrel{s, t \geq 1}{=} E \tanh(h + \beta Z_s) \tanh(h + \beta Z_t) = EM_s M_t = \begin{cases} q & : s = t, \\ \alpha_{s \wedge t} & : s \neq t. \end{cases}$$

Remark 4.3. In [22, 5], the iterative TAP sequence is defined differently compared to (4.1). Namely, in the language of [22, 5] one sets $\tilde{\mathbf{m}}^{(0)} = \mathbf{0}$, $\tilde{\mathbf{m}}^{(1)} = \sqrt{q}\mathbf{1}$ and for $k \geq 1$

$$\tilde{\mathbf{z}}^{(k+1)} = \mathbf{G}\tilde{\mathbf{m}}^{(k)} - \beta(1-q)\tilde{\mathbf{m}}^{(k-1)}, \quad \tilde{\mathbf{m}}^{(k+1)} = \tanh(h\mathbf{1} + \beta\tilde{\mathbf{z}}^{(k+1)}).$$

The validity of Theorem 4.3 then follows from

$$\lim_{N \rightarrow \infty} N^{-1} \mathbb{E} \|\tilde{\mathbf{z}}^{(k)} - \mathbf{z}^{(k)}\|^2 = 0, \quad \lim_{N \rightarrow \infty} N^{-1} \mathbb{E} \|\tilde{\mathbf{m}}^{(k)} - \mathbf{m}^{(k)}\|^2 = 0$$

for every $k \geq 2$, which can be proved inductively (note that $\tilde{\mathbf{m}}^{(1)} = \mathbf{m}^{(1)}$, $\tilde{\mathbf{m}}^{(2)} = \mathbf{m}^{(2)}$), based on Prop. 4.2 and the decomposition $\mathbf{G} = \mathbf{G}^{(k)} + \sum_{s=1}^{k-1} \tilde{\rho}^{(s)}$, and which implies that

$$\lim_{N \rightarrow \infty} \mathcal{W}_2 \left(\frac{1}{N} \sum_{i=1}^N \delta_{m_i^{(1)}, \dots, m_i^{(k)}, z_i^{(2)}, \dots, z_i^{(k+1)}}, \frac{1}{N} \sum_{i=1}^N \delta_{\tilde{m}_i^{(1)}, \dots, \tilde{m}_i^{(k)}, \tilde{z}_i^{(2)}, \dots, \tilde{z}_i^{(k+1)}} \right) = 0.$$

in probability.

The properties of Bolthausen's iterative solution $(\mathbf{m}^{(k)})_{k \geq 1}$ suggest that it is close to the magnetization vector \mathbf{m} of the SK model, at least in a suitable subregion of the RS phase. This is proved in [14], based on a locally uniform overlap concentration assumption.

Theorem 4.4 ([14]). Assume that $\beta, h > 0$ are such that for some $\delta > 0$ we have

$$\lim_{N \rightarrow \infty} \sup_{\beta - \delta \leq \beta' \leq \beta} \mathbb{E} \langle |R_{12} - q_{\beta', h}|^2 \rangle_{\beta', h} = 0. \quad (4.2)$$

Then we have that

$$\lim_{k \rightarrow \infty} \lim_{N \rightarrow \infty} N^{-1} \mathbb{E} \|\mathbf{m} - \mathbf{m}^{(k)}\|^2 = 0. \quad (4.3)$$

Remark 4.4. Observe that (4.2) is satisfied under (1.3) & (1.5), by Eq. (2.2) in Theorem 2.1. Notice furthermore that [14] defines the TAP iteration differently compared to (4.1):

$$(\mathbf{z}')^{(k+1)} = \mathbf{G}(\mathbf{m}')^{(k)} - \beta(1 - N^{-1} \|(\mathbf{m}')^{(k)}\|^2) (\mathbf{m}')^{(k-1)}, \quad (\mathbf{m}')^{(k+1)} = \tanh(h + \beta(\mathbf{z}')^{(k+1)}).$$

Similar remarks as for Theorem 4.3 apply: based on Prop. 4.2 and induction, one obtains

$$\lim_{N \rightarrow \infty} N^{-1} \mathbb{E} \|(\mathbf{z}')^{(k)} - \mathbf{z}^{(k)}\|^2 = 0, \quad \lim_{N \rightarrow \infty} N^{-1} \mathbb{E} \|(\mathbf{m}')^{(k)} - \mathbf{m}^{(k)}\|^2 = 0,$$

which implies (4.3) with $\mathbf{m}^{(k)}$ as defined in (4.1).

Equipped with the above preparations, we now turn to the proof of Prop. 1.3, which is a direct consequence of the next proposition. For a sequence $(X_N)_{N \in \mathbb{N}}$, we set

$$\begin{aligned} \text{p-lim inf } X_N &= \sup \left\{ t \in \mathbb{R} : \lim_{N \rightarrow \infty} \mathbb{P}(X_N \leq t) = 0 \right\}, \\ \text{p-lim sup } X_N &= \inf \left\{ t \in \mathbb{R} : \lim_{N \rightarrow \infty} \mathbb{P}(X_N \geq t) = 0 \right\}. \end{aligned}$$

The next result follows by translating the main ideas [9, Section 4] to the present context. For completeness, we carry out the key steps, with a few modifications, in detail below.

Proposition 4.5. Assume that (β, h) satisfy (1.3), and define for $\mathbf{v} \in (-1, 1)^N$ the diagonal matrix $\mathbf{D}_v \in \mathbb{R}^{N \times N}$ and the set $S_{N, \varepsilon, k} \subset \mathbb{R}^N \times (-1, 1)^N$ by

$$(\mathbf{D}_v)_{ij} = \frac{\delta_{ij}}{1 - v_i^2}, \quad S_{N, \varepsilon, k} = \left\{ (\mathbf{u}, \mathbf{v}) \in \mathbb{R}^N \times (-1, 1)^N : \|\mathbf{u}\|^2 = 1, \|\mathbf{v} - \mathbf{m}^{[k]}\|^2 \leq N\varepsilon \right\}. \quad (4.4)$$

Then, there exists a constant $c = c_{\beta, h} > 0$, that is independent of N , such that

$$\liminf_{\varepsilon \rightarrow 0} \liminf_{k \rightarrow \infty} \text{p-lim inf}_{N \rightarrow \infty} \left[\inf_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left(\mathbf{u}, (\mathbf{D}_v + \beta^2(1 - q) - \frac{2\beta^2}{N} \mathbf{v} \otimes \mathbf{v} - \beta \mathbf{G}) \mathbf{u} \right) \right] \geq c.$$

Remark 4.5. As already remarked in Section 1, note that Prop. 4.5 only requires (β, h) to satisfy the AT condition (1.3). The additional assumption (1.5) in Prop. 1.3 is used to approximate the magnetization \mathbf{m} of the SK model by the iterative TAP solution $\mathbf{m}^{(k)}$.

Before giving the proof of Prop. 4.5, let us first record that it implies Prop. 1.3.

Proof of Prop. 1.3, given Prop. 4.5. Observe, first of all, that $\mathbf{D} = \mathbf{D}_m$, by (1.8) & (4.4). Now, choosing $\varepsilon > 0$ sufficiently small and $k \in \mathbb{N}$ sufficiently large, Prop. 4.5 implies that

$$\inf_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left(\mathbf{u}, (\mathbf{D}_v + \beta^2(1 - q) - \frac{2\beta^2}{N} \mathbf{v} \otimes \mathbf{v} - \beta \mathbf{G}) \mathbf{u} \right) \geq \tilde{c}/2$$

for some constant $\tilde{c} > 0$, with probability tending to one as $N \rightarrow \infty$. By the assumptions (1.3) & (1.5), we can apply Theorem 4.4 which combined with Markov implies

$$\|\mathbf{m} - \mathbf{m}^{(k)}\|^2 \leq N\varepsilon$$

with probability tending to one as $N \rightarrow \infty$. Thus

$$\begin{aligned} & \inf_{\|\mathbf{u}\|^2=1} \left(\mathbf{u}, (\mathbf{D} + \beta^2(1 - q) - \beta \mathbf{G}) \mathbf{u} \right) \\ & \geq \inf_{\|\mathbf{u}\|^2=1} \left(\mathbf{u}, (\mathbf{D}_m + \beta^2(1 - q) - \frac{2\beta^2}{N} \mathbf{m} \otimes \mathbf{m} - \beta \mathbf{G}) \mathbf{u} \right) \\ & \geq \inf_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left(\mathbf{u}, (\mathbf{D}_v + \beta^2(1 - q) - \frac{2\beta^2}{N} \mathbf{v} \otimes \mathbf{v} - \beta \mathbf{G}) \mathbf{u} \right) \geq \tilde{c}/2, \end{aligned}$$

i.e. $\mathbf{D} + \beta^2(1 - q) - \beta \mathbf{G} \geq c$ for $c = \tilde{c}/2$, with probability tending to one as $N \rightarrow \infty$. \square

The rest of this section is devoted to the proof of Prop. 4.5.

Proof of Prop. 4.5. Step 1: We adapt the main idea of [9] and apply the Sudakov-Fernique inequality [27, 18, 26], conditionally on \mathcal{G}_k , to reduce the problem to a solvable variational problem. To this end, we first write

$$\beta(\mathbf{u}, \mathbf{G}\mathbf{u}) = \beta(\mathbf{u}, \mathbf{G}^{(k+1)}\mathbf{u}) + \frac{2\beta}{\sqrt{N}} \sum_{s=1}^k (\mathbf{u}, \zeta^{(s)} \otimes \phi^{(s)}, \mathbf{u}) + o_{\mathbf{u}}(1)$$

for an error $o_{\mathbf{u}}(1) = -N^{-1/2} \sum_{s=1}^k (\zeta^{(s)}, \phi^{(s)}) |(\phi^{(s)}, \mathbf{u})|^2$ which satisfies $\sup_{\|\mathbf{u}\|=1} o_{\mathbf{u}}(1) \rightarrow 0$ as $N \rightarrow \infty$ almost surely, by Prop. 4.2 5). This means that

$$\begin{aligned} & \text{p-} \limsup_{N \rightarrow \infty} \left[\inf_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left(\mathbf{u}, \left(\beta \mathbf{G} + \frac{2\beta^2}{N} \mathbf{v} \otimes \mathbf{v} - \mathbf{D}_{\mathbf{v}} - \beta^2(1-q) \right) \mathbf{u} \right) \right] \\ &= \text{p-} \limsup_{N \rightarrow \infty} \left[\inf_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left(\mathbf{u}, \left(\beta \mathbf{G}^{(k+1)} + \frac{2\beta}{\sqrt{N}} \sum_{s=1}^k \zeta^{(s)} \otimes \phi^{(s)} + \frac{2\beta^2}{N} \mathbf{v} \otimes \mathbf{v} - \mathbf{D}_{\mathbf{v}} - \beta^2(1-q) \right) \mathbf{u} \right) \right]. \end{aligned}$$

Now, comparing the (conditionally on \mathcal{G}_k) Gaussian processes

$$\begin{aligned} (\mathbf{X}_{\mathbf{u}, \mathbf{v}})_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} &= \left(\beta(\mathbf{u}, \mathbf{G}^{(k+1)} \mathbf{u}) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right)_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}}, \\ (\mathbf{Y}_{\mathbf{u}, \mathbf{v}})_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} &= \left(\frac{2\beta}{\sqrt{N}} \|\mathbf{Q}^{(k)} \mathbf{u}\| (\mathbf{Q}^{(k)} \mathbf{u}, \xi) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right)_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}}, \end{aligned}$$

where $\xi \sim \mathcal{N}(0, \text{id}_{\mathbb{R}^N})$ denotes a Gaussian vector independent of the remaining disorder and where we abbreviate

$$\mathbf{f}(\mathbf{u}, \mathbf{v}) = \frac{2\beta}{\sqrt{N}} \sum_{s=1}^k (\mathbf{u}, \zeta^{(s)}) (\phi^{(s)}, \mathbf{u}) + \frac{2\beta^2}{N} (\mathbf{u}, \mathbf{v})^2 - (\mathbf{u}, \mathbf{D}_{\mathbf{v}} \mathbf{u}) - \beta^2(1-q),$$

we have $\mathbb{E}_k \mathbf{X}_{\mathbf{u}, \mathbf{v}} = \mathbb{E}_k \mathbf{Y}_{\mathbf{u}, \mathbf{v}} = \mathbf{f}(\mathbf{u}, \mathbf{v})$ and an application of Prop. 4.2, part 4), shows that

$$\mathbb{E}_k (\mathbf{X}_{\mathbf{u}, \mathbf{v}} - \mathbf{X}_{\mathbf{u}', \mathbf{v}'})^2 \leq \mathbb{E}_k (\mathbf{Y}_{\mathbf{u}, \mathbf{v}} - \mathbf{Y}_{\mathbf{u}', \mathbf{v}'})^2$$

for every $(\mathbf{u}, \mathbf{v}), (\mathbf{u}', \mathbf{v}') \in S_{N, \varepsilon, k}$. Thus, by Vitale's extension [30] of the Sudakov-Fernique inequality [27, 18, 26] we obtain that a.s.

$$\mathbb{E}_k \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\beta(\mathbf{u}, \mathbf{G}^{(k+1)} \mathbf{u}) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] \leq \mathbb{E}_k \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\frac{2\beta}{\sqrt{N}} \|\mathbf{Q}^{(k)} \mathbf{u}\| (\mathbf{Q}^{(k)} \mathbf{u}, \xi) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right]. \quad (4.5)$$

Next, observing that conditionally on \mathcal{G}_k , we have the distributional equality

$$\mathbf{G}^{(k+1)} \stackrel{d}{=} \mathbf{Q}^{(k)} \frac{1}{\sqrt{2N}} (\mathbf{U} + \mathbf{U}^T) \mathbf{Q}^{(k)}$$

for some random matrix $\mathbf{U} = (u_{ij})_{1 \leq i, j \leq N} \in \mathbb{R}^{N \times N}$ with i.i.d. entries $u_{ij} \sim \mathcal{N}(0, 1)$ (without symmetry constraint) for $i \neq j$ and $u_{ii} = 0$, a standard application of Gaussian concentration (see, for instance, [28, Theorem 1.3.4]) implies that

$$\begin{aligned} & \mathbb{P}_k \left(\left| \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\beta(\mathbf{u}, \mathbf{G}^{(k+1)} \mathbf{u}) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] \right. \right. \\ & \quad \left. \left. - \mathbb{E}_k \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\beta(\mathbf{u}, \mathbf{G}^{(k+1)} \mathbf{u}) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] \right| > t \right) \leq 2e^{-C N t^2}. \end{aligned}$$

Indeed, for $\mathbf{X}, \mathbf{Y} \in \mathbb{R}^{N \times N}$, notice that

$$\begin{aligned}
& \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\frac{\beta}{\sqrt{N}} (\mathbf{Q}^{(k)} \mathbf{u}, \bar{\mathbf{X}} \mathbf{Q}^{(k)} \mathbf{u}) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] - \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\frac{\beta}{\sqrt{N}} (\mathbf{Q}^{(k)} \mathbf{u}, \bar{\mathbf{Y}} \mathbf{Q}^{(k)} \mathbf{u}) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] \\
&= \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\frac{\sqrt{2}\beta}{\sqrt{N}} (\mathbf{Q}^{(k)} \mathbf{u}, (\mathbf{X} - \mathbf{Y}) \mathbf{Q}^{(k)} \mathbf{u}) + \frac{\sqrt{2}\beta}{\sqrt{N}} (\mathbf{Q}^{(k)} \mathbf{u}, \mathbf{Y} \mathbf{Q}^{(k)} \mathbf{u}) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] \\
&\quad - \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\frac{2\beta}{\sqrt{N}} (\mathbf{Q}^{(k)} \mathbf{u}, \mathbf{Y} \mathbf{Q}^{(k)} \mathbf{u}) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] \\
&\leq \sup_{\mathbf{u}' \in \mathbb{R}^N: \|\mathbf{u}'\|^2 \leq 1} \frac{\sqrt{2}\beta}{\sqrt{N}} (\mathbf{u}', (\mathbf{X} - \mathbf{Y}) \mathbf{u}') \leq \frac{C}{\sqrt{N}} \left(\sum_{i,j=1}^N (\mathbf{X} - \mathbf{Y})_{ij}^2 \right)^{1/2} = \frac{C}{\sqrt{N}} \|\mathbf{X} - \mathbf{Y}\|
\end{aligned}$$

so that, upon switching the roles of $\mathbf{X} \in \mathbb{R}^{N \times N}$ and $\mathbf{Y} \in \mathbb{R}^{N \times N}$, we get

$$\begin{aligned}
& \left| \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\frac{\beta}{\sqrt{N}} (\mathbf{Q}^{(k)} \mathbf{u}, \bar{\mathbf{X}} \mathbf{Q}^{(k)} \mathbf{u}) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] - \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\frac{\beta}{\sqrt{N}} (\mathbf{Q}^{(k)} \mathbf{u}, \bar{\mathbf{Y}} \mathbf{Q}^{(k)} \mathbf{u}) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] \right| \\
&\leq \frac{C}{\sqrt{N}} \|\mathbf{X} - \mathbf{Y}\|
\end{aligned}$$

for the deterministic constant $C = \sqrt{2}\beta > 0$. Arguing along the same lines, we also find

$$\begin{aligned}
& \mathbb{P}_k \left(\left| \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\frac{2\beta}{\sqrt{N}} \|\mathbf{Q}^{(k)} \mathbf{u}\| (\mathbf{Q}^{(k)} \mathbf{u}, \xi) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] \right. \right. \\
&\quad \left. \left. - \mathbb{E}_k \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\frac{2\beta}{\sqrt{N}} \|\mathbf{Q}^{(k)} \mathbf{u}\| (\mathbf{u}, \xi) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] \right| > t \right) \leq 2e^{-CNt^2},
\end{aligned}$$

and since $C > 0$ is independent of \mathcal{G}_k , we can take the expectation $\mathbb{E}(\cdot)$ over the previous two tail bounds to see that they hold true unconditionally. Thus, we conclude

$$\lim_{N \rightarrow \infty} \left| \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\beta(\mathbf{u}, \mathbf{G}^{(k+1)} \mathbf{u}) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] - \mathbb{E}_k \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\beta(\mathbf{u}, \mathbf{G}^{(k+1)} \mathbf{u}) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] \right| = 0$$

and

$$\begin{aligned}
& \lim_{N \rightarrow \infty} \left| \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\frac{2\beta}{\sqrt{N}} \|\mathbf{Q}^{(k)} \mathbf{u}\| (\mathbf{Q}^{(k)} \mathbf{u}, \xi) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] \right. \\
&\quad \left. - \mathbb{E}_k \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left[\frac{2\beta}{\sqrt{N}} \|\mathbf{Q}^{(k)} \mathbf{u}\| (\mathbf{Q}^{(k)} \mathbf{u}, \xi) + \mathbf{f}(\mathbf{u}, \mathbf{v}) \right] \right| = 0
\end{aligned}$$

in the sense of probability. Combining these observations with (4.5), we finally arrive at

$$\begin{aligned}
& \limsup_{\varepsilon \rightarrow 0} \limsup_{k \rightarrow \infty} \text{p-} \limsup_{N \rightarrow \infty} \left[\sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \left(\mathbf{u}, \left(\beta \mathbf{G} + \frac{2\beta^2}{N} \mathbf{v} \otimes \mathbf{v} - \mathbf{D}_{\mathbf{v}} - \beta^2(1 - q) \right) \mathbf{u} \right) \right] \\
&\leq \limsup_{\varepsilon \rightarrow 0} \limsup_{k \rightarrow \infty} \text{p-} \limsup_{N \rightarrow \infty} \left[\sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} \mathbf{F}(\mathbf{u}, \mathbf{v}, \xi, \mathbf{M}^{(k)}, \mathbf{Z}^{(k)}) \right],
\end{aligned} \tag{4.6}$$

where $\mathbf{M}^{(k)}$ and $\mathbf{Z}^{(k)}$ denote the matrices

$$\mathbf{M}^{(k)} = (\mathbf{m}^{(1)} \mathbf{m}^{(2)} \dots \mathbf{m}^{(k)}) \in \mathbb{R}^{N \times k}, \quad \mathbf{Z}^{(k)} = (\mathbf{z}^{(2)}, \dots, \mathbf{z}^{(k+1)}) \in \mathbb{R}^{N \times k}$$

and where

$$\begin{aligned} F(\mathbf{u}, \mathbf{v}, \xi, \mathbf{M}^{(k)}, \mathbf{Z}^{(k)}) &= \frac{2\beta}{\sqrt{N}} \|\mathbf{Q}^{(k)} \mathbf{u}\|(\mathbf{u}, \xi) + 2\beta(\mathbf{u}, \mathbf{Z}^{(k)} (\mathbf{M}^{(k)T} \mathbf{M}^{(k)})^{-1} \mathbf{M}^{(k)T} \mathbf{u}) \\ &\quad + \frac{2\beta^2}{N} (\mathbf{u}, \mathbf{m}^{(k)})^2 - (\mathbf{u}, \mathbf{D}_v \mathbf{u}) - \beta^2(1 - q). \end{aligned}$$

Here, we used additionally that

$$\sup_{\|\mathbf{v} - \mathbf{m}^{(k)}\|^2 \leq N\varepsilon} N^{-1} \|\mathbf{v} \otimes \mathbf{v} - \mathbf{m}^{(k)} \otimes \mathbf{m}^{(k)}\|_{\text{op}} \leq 2\sqrt{\varepsilon}$$

and that, in the sense of probability, we have

$$\begin{aligned} \lim_{N \rightarrow \infty} \left| \sup_{\|\mathbf{u}\|^2=1} N^{-1/2} (\mathbf{P}^{(k)} \mathbf{u}, \xi) \right| &\leq \lim_{N \rightarrow \infty} N^{-1/2} \|\mathbf{P}^{(k)} \xi\| = 0, \\ \lim_{N \rightarrow \infty} \left\| \frac{1}{\sqrt{N}} \sum_{s=1}^k \zeta^{(s)} \otimes \phi^{(s)} - \mathbf{Z}^{(k)} (\mathbf{M}^{(k)T} \mathbf{M}^{(k)})^{-1} \mathbf{M}^{(k)T} \right\|_{\text{op}} &= 0. \end{aligned}$$

The last two identities readily follow e.g. from a simple second moment bound and, respectively, from Prop. 4.2. Notice that both matrices $N^{-1/2} \sum_{s=1}^k \zeta^{(s)} \otimes \phi^{(s)}$ and $\mathbf{Z}^{(k)} (\mathbf{M}^{(k)T} \mathbf{M}^{(k)})^{-1} \mathbf{M}^{(k)T}$ are at most of rank k ($\mathbf{Q}^{(k)} \mathbb{R}^N$ contained in their kernels) so that the norm convergence follows from pointwise convergence on the vectors $(\mathbf{m}^{(s)})_{s=1}^k$.

Step 2: The remainder of the proof is based on analyzing the optimization problem on the r.h.s. of Eq. (4.6). This can be done with the same arguments as in [9, Sections 4.1 to 4.4]. For completeness, we translate the arguments from [9] to the present setting. In particular, we point out at which steps the AT condition (1.3) enters the analysis.

First, we control the r.h.s. of Eq. (4.6) through an optimization problem that only involves the limiting distributions of $\mathbf{M}^{(k)}$ and $\mathbf{Z}^{(k)}$, based on Theorem 4.3. Setting

$$\mu_N^{(k)} = \frac{1}{N} \sum_{i=1}^N \delta_{\sqrt{N} u_i, v_i, \xi_i, \mathbf{M}_i^{(k)}, \mathbf{z}_i^{(k)}} \in \mathcal{P}(\mathbb{R}^{2k+3}),$$

where by $\mathbf{X}_i \in \mathbb{R}^n$ we denote the i -th row of $\mathbf{X} \in \mathbb{R}^{m \times n}$, and setting

$$\mathbb{E}_\mu(f) = \int \mu(dx) f(x), \quad \mathbb{P}_\mu(S) = \int \mu(dx) \mathbf{1}_S(x) = \mu(S), \quad \langle f, g \rangle_{L^2(d\mu)} = \int \mu(dx) f(x)g(x)$$

for $\mu \in \mathcal{P}(\mathbb{R}^{2k+3})$, we have that

$$F(\mathbf{u}, \mathbf{v}, \xi, \mathbf{M}^{(k)}, \mathbf{Z}^{(k)}) = \Phi(\mu_N^{(k)}),$$

where for every $\mu \in \mathcal{P}(\mathbb{R}^{2k+3})$ with finite second moment, we define

$$\begin{aligned}\Phi(\mu) &= 2\beta \mathbb{E}_\mu(\mathbf{U}\mathbf{Z}^T) [\mathbb{E}_\mu(\mathbf{M}\mathbf{M}^T)]^{-1} \mathbb{E}_\mu(\mathbf{M}\mathbf{U}) + 2\beta \|\mathbf{Q}_{\mathbf{M}}\mathbf{U}\|_{L^2(d\mu)} \langle \mathbf{U}, \Xi \rangle_{L^2(d\mu)} \\ &\quad + 2\beta^2 \langle M_k, \mathbf{U} \rangle_{L^2(d\mu)}^2 - \mathbb{E}_\mu((1 - \mathbf{V}^2)^{-1} \mathbf{U}^2) - \beta^2(1 - q).\end{aligned}$$

Here, the coordinates in the integrals over \mathbb{R}^{2k+3} w.r.t. to μ are denoted by $(\mathbf{U}, \mathbf{V}, \Xi, \mathbf{M}, \mathbf{Z})$ (or, in other words, $\mathcal{L}(\mathbf{U}, \mathbf{V}, \Xi, \mathbf{M}, \mathbf{Z}) = \mu$) and $\mathbf{Q}_{\mathbf{M}}$ denotes the projection onto the orthogonal complement of the coordinates of $\mathbf{M} = (M_1, \dots, M_k)$, i.e.

$$\mathbf{Q}_{\mathbf{M}}\mathbf{U} = \mathbf{U} - \mathbf{M}^T (\mathbb{E}_\mu \mathbf{M} \mathbf{M}^T)^{-1} \mathbb{E}_\mu \mathbf{M} \mathbf{U}.$$

To stay consistent with the construction in (4.1) and with the previous notation, we write in the following $\mathbf{Z} = (Z_2, \dots, Z_{k+1})$. Now, applying [9, Lemma 1] (whose proof in [9, Appendix B.2] carries over directly to the present context upon replacing the functional $F_{x,k}$ from [9] by Φ defined above, based on Prop. 4.2), we obtain the upper bound

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{k \rightarrow \infty} \text{p-} \limsup_{N \rightarrow \infty} \sup_{(\mathbf{u}, \mathbf{v}) \in S_{N, \varepsilon, k}} F(\mathbf{u}, \mathbf{v}, \xi, \mathbf{M}^{(k)}, \mathbf{Z}^{(k)}) \leq \limsup_{\varepsilon \rightarrow 0} \limsup_{k \rightarrow \infty} \sup_{\mu \in \mathcal{S}_{\varepsilon, k}} \Phi(\mu), \quad (4.7)$$

where

$$\begin{aligned}\mathcal{S}_{\varepsilon, k} &= \{ \mu = \mathcal{L}(\mathbf{U}, \mathbf{V}, \Xi, \mathbf{M}, \mathbf{Z}) : \|\mathbf{U}\|_{L^2(d\mu)}^2 = 1, \|\mathbf{V} - M_k\|_{L^2(d\mu)}^2 \leq \varepsilon, \mathbb{P}_\mu(|\mathbf{V}| < 1) = 1, \\ &\quad \mathbf{Z} = (Z_2, \dots, Z_{k+1}) \sim \mathcal{N}(\mathbf{0}, \mathbf{K}_{\leq k}), \mathbf{M} = (M_1, \dots, M_k) \text{ with } M_1 = \sqrt{q}, \\ &\quad M_s \stackrel{s \geq 1}{\cong} \tanh(h + \beta Z_{s-1}), \Xi \sim \mathcal{N}(0, 1) \text{ independent of } \mathbf{Z} \}.\end{aligned}$$

Next, using that $\mathbb{E}_\mu(\mathbf{Z}\mathbf{Z}^T) = \mathbb{E}_\mu(\mathbf{M}\mathbf{M}^T) = \mathbf{K}_{\leq k}$, we observe that

$$\begin{aligned}&\|\mathbf{Z}^T [\mathbb{E}_\mu(\mathbf{M}\mathbf{M}^T)]^{-1} \mathbb{E}_\mu(\mathbf{M}\mathbf{U})\|_{L^2(d\mu)}^2 + \|\mathbf{Q}_{\mathbf{M}}\mathbf{U}\|_{L^2(d\mu)}^2 \|\Xi\|_{L^2(d\mu)}^2 \\ &= \mathbb{E}_\mu(\mathbf{M}\mathbf{U})^T [\mathbb{E}_\mu(\mathbf{M}\mathbf{M}^T)]^{-1} \mathbb{E}_\mu(\mathbf{Z}\mathbf{Z}^T) [\mathbb{E}_\mu(\mathbf{M}\mathbf{M}^T)]^{-1} \mathbb{E}_\mu(\mathbf{M}\mathbf{U}) + \|\mathbf{Q}_{\mathbf{M}}\mathbf{U}\|_{L^2(d\mu)}^2 = 1\end{aligned}$$

so that, by the independence of \mathbf{Z} and Ξ , we have that under μ

$$\mathbf{Z}^T [\mathbb{E}_\mu(\mathbf{M}\mathbf{M}^T)]^{-1} \mathbb{E}_\mu(\mathbf{M}\mathbf{U}) + \|\mathbf{Q}_{\mathbf{M}}\mathbf{U}\|_{L^2(d\mu)} \Xi \sim \mathcal{N}(0, 1).$$

Arguing similarly that

$$\langle Z_{k+1}, \mathbf{Z}^T [\mathbb{E}_\mu(\mathbf{M}\mathbf{M}^T)]^{-1} \mathbb{E}_\mu(\mathbf{M}\mathbf{U}) \rangle_{L^2(d\mu)} = \langle M_k, \mathbf{U} \rangle_{L^2(d\mu)}$$

and using that $\|Z_{k+1}\|_{L^2(d\mu)} = \|M_k\|_{L^2(d\mu)} = \sqrt{q}$, we can thus write

$$\mathbf{Z}^T [\mathbb{E}_\mu(\mathbf{M}\mathbf{M}^T)]^{-1} \mathbb{E}_\mu(\mathbf{M}\mathbf{U}) + \|\mathbf{Q}_{\mathbf{M}}\mathbf{U}\|_{L^2(d\mu)} \Xi = q^{-1} \langle M_k, \mathbf{U} \rangle_{L^2(d\mu)} Z_{k+1} + \|\mathbf{Q}_{M_k}\mathbf{U}\|_{L^2(d\mu)} \Xi'$$

for some Gaussian $\Xi' \sim \mathcal{N}(0, 1)$, which is independent of Z_{k+1} , and where Q_{M_k} denotes the projection onto the orthogonal complement of M_k in $L^2(d\mu)$. In particular, we have

$$\begin{aligned}\Phi(\mu) &= \frac{2\beta}{q} \langle \mathbf{U}, M_k \rangle_{L^2(d\mu)} \langle Z_{k+1}, \mathbf{U} \rangle_{L^2(d\mu)} + 2\beta \|\mathbf{Q}_{M_k} \mathbf{U}\|_{L^2(d\mu)} \langle \mathbf{U}, \Xi' \rangle_{L^2(d\mu)} \\ &\quad + 2\beta^2 \langle M_k, \mathbf{U} \rangle_{L^2(d\mu)}^2 - \mathbb{E}_\mu((1 - V^2)^{-1} \mathbf{U}^2) - \beta^2(1 - q),\end{aligned}$$

and setting $M_{k+1} = \tanh(h + \beta Z_{k+1})$, such that under the AT condition (1.3) we have $\|M_{k+1} - M_k\|_{L^2(d\mu)}^2 = 2q - 2\alpha_k \rightarrow 0$ as $k \rightarrow \infty$ by Lemma 4.1 and Prop. 4.2 4), we find

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{k \rightarrow \infty} \sup_{\mu \in \mathcal{S}_{\varepsilon, k}} \Phi(\mu) \leq \limsup_{\varepsilon \rightarrow 0} \sup_{\mu \in \mathcal{S}_\varepsilon} \Psi(\mu), \quad (4.8)$$

where Ψ is defined by

$$\begin{aligned}\Psi(\mu) &= 2\beta q^{-1/2} \langle \mathbf{U}, \mathbf{M} \rangle_{L^2(d\mu)} \langle \mathbf{Z}, \mathbf{U} \rangle_{L^2(d\mu)} + 2\beta \sqrt{1 - q^{-1} \langle \mathbf{M}, \mathbf{U} \rangle_{L^2(d\mu)}^2} \langle \mathbf{W}, \mathbf{U} \rangle_{L^2(d\mu)} \\ &\quad + 2\beta^2 \langle \mathbf{M}, \mathbf{U} \rangle_{L^2(d\mu)}^2 - \mathbb{E}_\mu((1 - V^2)^{-1} \mathbf{U}^2) - \beta^2(1 - q),\end{aligned}$$

for every $\mu = \mathcal{L}(\mathbf{M}, \mathbf{U}, \mathbf{V}, \mathbf{W}, \mathbf{Z}) \in \mathcal{S}_\varepsilon$, with \mathcal{S}_ε defined by

$$\begin{aligned}\mathcal{S}_\varepsilon &= \{ \mathcal{L}(\mathbf{M}, \mathbf{U}, \mathbf{V}, \mathbf{W}, \mathbf{Z}) : \|\mathbf{U}\|_{L^2(d\mu)}^2 = 1, \|\mathbf{V} - \mathbf{M}\|_{L^2(d\mu)}^2 \leq \varepsilon, \\ &\quad \mathbb{P}_\mu(|V| < 1) = 1, (\mathbf{W}, \mathbf{Z}) \sim \mathcal{N}(0, \text{id}_{\mathbb{R}^2}), \mathbf{M} = \tanh(h + \beta \sqrt{q} \mathbf{Z}) \}.\end{aligned}$$

Step 3: Finally, we need to analyze the optimization problem on the r.h.s. in Eq. (4.8). Here, we introduce Lagrange multipliers and simply upper bound the r.h.s. by

$$\begin{aligned}\Psi(\mu) &= 2\beta q^{-1/2} \langle \mathbf{U}, \mathbf{M} \rangle_{L^2(d\mu)} \langle \mathbf{Z}, \mathbf{U} \rangle_{L^2(d\mu)} + 2\beta \sqrt{1 - q^{-1} \langle \mathbf{M}, \mathbf{U} \rangle_{L^2(d\mu)}^2} \langle \mathbf{W}, \mathbf{U} \rangle_{L^2(d\mu)} \\ &\quad + 2\beta^2 \langle \mathbf{M}, \mathbf{U} \rangle_{L^2(d\mu)}^2 - \mathbb{E}_\mu((1 - V^2)^{-1} \mathbf{U}^2) - \beta^2(1 - q) \\ &= 2\beta q^{-1/2} x \langle \mathbf{Z}, \mathbf{U} \rangle_{L^2(d\mu)} + 2\beta \sqrt{1 - q^{-1} x^2} \langle \mathbf{W}, \mathbf{U} \rangle_{L^2(d\mu)} + 2\beta^2 x^2 - \beta^2(1 - q) \\ &\quad + \lambda_u (\|\mathbf{U}\|_{L^2(d\mu)}^2 - 1) + \lambda_x (\langle \mathbf{U}, \mathbf{M} \rangle_{L^2(d\mu)} - x) - \mathbb{E}_\mu((1 - V^2)^{-1} \mathbf{U}^2) \\ &\leq 2\beta^2 x^2 - \beta^2(1 - q) - \lambda_u - \lambda_x x + \mathbb{E}_\mu \Theta(\mathbf{M}, \mathbf{V}, \mathbf{W}, \mathbf{Z}),\end{aligned}$$

where we set $x = x(\mathbf{U}, \mathbf{M}) = \langle \mathbf{U}, \mathbf{M} \rangle_{L^2(d\mu)}$ as well as

$$\Theta(m, v, w, z) = \sup_{u \in \mathbb{R}} \left[(2\beta q^{-1/2} x z + 2\beta \sqrt{1 - q^{-1} x^2} w + \lambda_x m) u + (\lambda_u - (1 - v^2)^{-1}) u^2 \right].$$

Fixing from now on $\lambda_u < 1$ and assuming w.l.o.g. $|v| < 1$ (in accordance to the fact that $\mathbb{P}_\mu(|V| < 1) = 1$), strict concavity implies that

$$\Theta(m, v, w, z) = \frac{1}{4} \frac{(2\beta q^{-1/2} x z + 2\beta \sqrt{1 - q^{-1} x^2} w + \lambda_x m)^2}{((1 - v^2)^{-1} - \lambda_u)}.$$

Together with the global Lipschitz continuity of $[-1, 1] \ni v \mapsto ((1-v^2)^{-1} - \lambda_u)^{-1} \in [0, \infty)$, we thus obtain for every fixed $\lambda_u < 1$ and $K > 0$ the simple upper bound

$$\begin{aligned}
& \limsup_{\varepsilon \rightarrow 0} \sup_{\mu \in \mathcal{S}_\varepsilon} \Psi(\mu) \\
& \leq \max_{|x| \leq 1} \min_{|\lambda_x| \leq K} \left[2\beta^2 x^2 - \beta^2(1-q) - \lambda_u - \lambda_x x \right. \\
& \quad \left. + \mathbb{E}_\mu \frac{(2\beta q^{-1/2} x Z + 2\beta \sqrt{1-q^{-1}x^2} W + \lambda_x M)^2}{4(\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u)} \right] \\
& = \max_{|x| \leq 1} \min_{|\lambda_x| \leq K} \left[2\beta^2 x^2 - \beta^2(1-q) - \lambda_u + \mathbb{E}_\mu \frac{\beta^2(1-q^{-1}x^2)}{(\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u)} \right. \\
& \quad \left. - \lambda_x x + \mathbb{E}_\mu \frac{\beta^2 q^{-1} x^2 Z^2 + \beta q^{-1/2} x \lambda_x M Z + \lambda_x^2 M^2 / 4}{(\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u)} \right].
\end{aligned} \tag{4.9}$$

Notice that, in the last step, we used the independence of W of Z and, consequently, of $M = \tanh(h + \beta\sqrt{q}Z)$. Applying Gaussian integration by parts w.r.t. Z and rescaling x and λ_x by a factor $1/\sqrt{q}$ and respectively \sqrt{q} , we find that

$$\begin{aligned}
& \max_{|x| \leq 1} \min_{|\lambda_x| \leq K} \left[2\beta^2 x^2 - \beta^2(1-q) - \lambda_u + \mathbb{E}_\mu \frac{\beta^2(1-q^{-1}x^2)}{(\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u)} \right. \\
& \quad \left. - \lambda_x x + \mathbb{E}_\mu \frac{\beta^2 q^{-1} x^2 Z^2 + \beta q^{-1/2} x \lambda_x M Z + \lambda_x^2 M^2 / 4}{(\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u)} \right] \\
& = \max_{|x| \leq q^{-1/2}} \left\{ 2\beta^2 q x^2 - \beta^2(1-q) - \lambda_u + \mathbb{E}_\mu \frac{\beta^2}{(\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u)} \right. \\
& \quad - \mathbb{E}_\mu \frac{2\beta^4 q (\cosh^2(h + \beta\sqrt{q}Z) + \sinh^2(h + \beta\sqrt{q}Z))}{(\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u)^2} x^2 \\
& \quad + \mathbb{E}_\mu \frac{8\beta^4 q \cosh^2(h + \beta\sqrt{q}Z) \sinh^2(h + \beta\sqrt{q}Z)}{(\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u)^3} x^2 \\
& \quad + \min_{|\lambda_x| \leq \sqrt{q}K} \mathbb{E}_\mu \left[\frac{\beta^2}{\cosh^2(h + \beta\sqrt{q}Z) (\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u)} x \lambda_x \right. \\
& \quad \quad \left. - \frac{2\beta^2 \cosh^2(h + \beta\sqrt{q}Z) \tanh^2(h + \beta\sqrt{q}Z)}{(\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u)^2} x \lambda_x \right. \\
& \quad \quad \left. - x \lambda_x + \frac{1}{4} \frac{M^2}{(\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u)} \lambda_x^2 \right] \Big\} \\
& = \max_{|x| \leq q^{-1/2}} \Sigma_{\lambda_u}(x).
\end{aligned} \tag{4.10}$$

Now, evaluating the Hessian of Σ_{λ_u} for $\lambda_u = 0$, one obtains with $q_4 = \mathbb{E}_\mu M^4$ that

$$\begin{aligned} \frac{1}{2} \frac{d^2 \Sigma_0}{dx^2} &= 2\beta^2 q - 2\beta^4 q(1 - 4q + 3q_4) - (q - q_4)^{-1} (1 - \beta^2(1 - 4q + 3q_4))^2 \\ &= \frac{(1 - \beta^2(1 - 4q + 3q_4))}{(q - q_4)} (2\beta^2 q(q - q_4) - (1 - \beta^2(1 - 2q + q_4)) - 2\beta^2(q - q_4)) \\ &= \frac{(1 - \beta^2(1 - 4q + 3q_4))}{(q - q_4)} (- (1 - \beta^2(1 - 2q + q_4)) - 2\beta^2(1 - q)(q - q_4)). \end{aligned}$$

Finally, by the AT condition (1.3) and $q - q_4 = \mathbb{E}_\mu \left(\frac{\tanh^2}{\cosh^2} \right) (h + \beta\sqrt{q}Z) \geq 0$, we get

$$\begin{aligned} 1 - \beta^2(1 - 2q + q_4) &= 1 - \beta^2 \mathbb{E}_\mu \operatorname{sech}^4(h + \beta\sqrt{q}Z) > 0, \\ 1 - \beta^2(1 - 4q + 3q_4) &= 1 - \beta^2(1 - 2q + q_4) + 2\beta^2(q - q_4) > 0, \end{aligned}$$

and together with $1 - q = \mathbb{E}_\mu \operatorname{sech}^2(h + \beta\sqrt{q}Z) \geq 0$ and smoothness in λ_u , this implies

$$\frac{d^2 \Sigma_{\lambda_u}}{dx^2} < 0$$

for $\lambda_u \in (0, 1)$ sufficiently small. By concavity, we obtain for $\lambda_u > 0$ small enough that

$$\begin{aligned} \max_{|x| \leq 1} \min_{|\lambda_x| \leq K} &\left[2\beta^2 x^2 - \beta^2(1 - q) - \lambda_u + \mathbb{E}_\mu \frac{\beta^2(1 - q^{-1}x^2)}{(\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u)} \right. \\ &\quad \left. - \lambda_x x + \mathbb{E}_\mu \frac{\beta^2 q^{-1} x^2 Z^2 + \beta q^{-1/2} x \lambda_x MZ + \lambda_x^2 M^2/4}{(\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u)} \right] \\ &= -\beta^2(1 - q) - \lambda_u + \beta^2 \mathbb{E}_\mu \frac{1}{\cosh^2(h + \beta\sqrt{q}Z) - \lambda_u} \\ &= - \int_0^{\lambda_u} dt \left[1 - \beta^2 \mathbb{E}_\mu \frac{1}{(\cosh^2(h + \beta\sqrt{q}Z) - t)^2} \right] \leq -c < 0 \end{aligned}$$

for some positive constant $c = c_{\beta, h} > 0$. Here, the last inequality follows from the assumption that $\lambda_u > 0$ is sufficiently small and from the AT condition (1.3), noting that

$$- \left[1 - \beta^2 \mathbb{E}_\mu \frac{1}{(\cosh^2(h + \beta\sqrt{q}Z) - t)^2} \right]_{t=0} = -(1 - \beta^2 \mathbb{E}_\mu \operatorname{sech}^4(h + \beta\sqrt{q}Z)) < 0.$$

Combining this with (4.6), (4.7), (4.8) and (4.10), this proves Prop. 4.5. \square

Acknowledgements. C. B. acknowledges support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – GZ 2047/1, Projekt-ID 390685813. The work of C. X. is partially funded by a Simons Investigator award. The work of H.-T. Y. is partially supported by the NSF grant DMS-1855509 and DMS-2153335 and a Simons Investigator award.

References

- [1] A. Adhikari, C. Brennecke, P. von Soosten, and H.-T. Yau. Dynamical Approach to the TAP Equations for the Sherrington-Kirkpatrick Model. *J. Stat. Phys.*, 183, 2021.
- [2] P. W. Anderson, R. G. Palmer, and D. J. Thouless. Solution of 'solvable model in spin glasses'. *Philos. Magazin*, 35:593–601, 1977.
- [3] D. Belius. High temperature TAP upper bound for the free energy of mean field spin glasses. *Preprint: arXiv:2204.00681*.
- [4] D. Belius and N. Kistler. The TAP-Plefka variational principle for the spherical SK model. *Comm. Math. Physics*, 367, 2019.
- [5] E. Bolthausen. An Iterative Construction of Solutions of the TAP Equations for the Sherrington-Kirkpatrick Model. *Comm. Math. Phys.*, 325(1):333–366, 2014.
- [6] E. Bolthausen. A Morita type proof of the replica-symmetric formula for SK. *Statistical Mechanics of Classical and Disordered Systems, Springer Proceedings in Mathematics & Statistics*, pages 63–93, 2018.
- [7] C. Brennecke, A. Schertzer, C. Xu, and H.-T. Yau. The Two Point Function of the SK Model without External Field at High Temperature. *Preprint: arXiv:2212.14476*.
- [8] C. Brennecke and H.-T. Yau. The Replica Symmetric Formula for the SK Model Revisited. *J. Math. Phys.*, 63, 2022.
- [9] M. Celentano. Sudakov-ferniqne post-AMP, and a new proof of the local convexity of the TAP free energy. *Preprint: arXiv:2208.09550*.
- [10] M. Celentano, Z. Fan, and S. Mei. Local convexity of the TAP free energy and AMP convergence for Z2-synchronization. *Preprint: arXiv:2106.11428*.
- [11] W. Chen, D. Panchenko, and E. Subag. The generalized TAP free energy II. *Comm. Math. Physics*, 381, 2021.
- [12] W. Chen, D. Panchenko, and E. Subag. The generalized TAP free energy. *Comm. Pure Appl. Math.*, 76, 2023.
- [13] W.-K. Chen. On the Almeida-Thouless transition line in the Sherrington-Kirkpatrick model with centered Gaussian external field. *Electron. Commun. Probab.*, 26:1–9, 2021.
- [14] W.-K. Chen and S. Tang. On convergence of the cavity and Bolthausen's TAP iterations to the local magnetization. *Comm. Math. Phys.*, 386(2):1209–1242, 2021.
- [15] J. R. L. de Almeida and D. Thouless. Stability of the Sherrington-Kirkpatrick solution of a spin glass model. *J. Phys. A: Math. Gen.*, 11:983–990, 1978.
- [16] A. El Alaoui and J. Gaitonde. Bounds on the covariance matrix of the Sherrington-Kirkpatrick model. *Preprint arXiv:2212.02445*.
- [17] A. El Alaoui, A. Montanari, and M. Sellke. Sampling from the Sherrington-Kirkpatrick Gibbs measure via algorithmic stochastic localization. *63th Annual Conference on Foundations of Computer Science (FOCS)*, 2022.

- [18] X. Fernique. Régularité des trajectoires des fonctions aléatoires gaussiennes. *Ecole d'Eté de Probabilités de Saint-Flour IV*, pages 1–96, 1975.
- [19] V. Gayraud. Emergence of near-TAP free energy functional in the SK model at high temperature. *Preprint: arXiv:2306.02402*.
- [20] F. Guerra. Sum rules for the free energy in the mean field spin glass model. *Fields Institute Communications*, 30, 2001.
- [21] S. Gufler, A. Schertzer, and M. A. Schmidt. On concavity of TAP free energy in the SK model. *Preprint: arXiv:2209.08985*.
- [22] A. Javanmard and A. Montanari. State evolution for general approximate message passing algorithms, with applications to spatial coupling. *Information and Inference: A Journal of the IMA*, 2(2):115–144, 2013.
- [23] N. Kistler, M. A. Schmidt, and G. Bastiani. On the REM approximation of TAP free energies. *Preprint: arXiv:2212.11718*.
- [24] T. Plefka. Convergence condition of the tap equation for the infinite- ranged ising spin glass model. *J. Phys. A: Math. Gen.*, 15:1971–1978, 1982.
- [25] D. Sherrington and S. Kirkpatrick. Solvable model of a spin glass. *Phys. Rev. Lett.*, 35:1792–1796, 1975.
- [26] V. N. Sudakov. Gaussian random processes and measures of solid angles in hilbert space. *Doklady Akademii Nauk*, 197:43–45, 1971.
- [27] V. N. Sudakov. *Geometric Problems in the Theory of Infinite-Dimensional Probability Distributions*, volume 141. American Mathematical Society, 1979.
- [28] M. Talagrand. *Mean Field Models for Spin Glasses: Volume I: Basic Examples*. Springer Science & Business Media, 2010.
- [29] M. Talagrand. *Mean Field Models for Spin Glasses: Volume II: Advanced Replica-Symmetry and Low Temperature*. Springer Science & Business Media, 2011.
- [30] R. A. Vitale. Some comparisons for gaussian processes. *Proc. Amer. Math. Soc.*, 128, 2000.