

---

---

# A general Temperature-Guided Language model to engineer enhanced Stability and Activity in Proteins

Pan Tan<sup>1,2†</sup>, Mingchen Li<sup>2,6†</sup>, Yuanxi Yu<sup>2,3†</sup>, Fan Jiang<sup>2,3†</sup>, Lirong Zheng<sup>2†</sup>, Banghao Wu<sup>2,4†</sup>, Xinyu Sun<sup>5†</sup>, Yufeng Pei<sup>5</sup>, Jie Song<sup>5</sup>, Liqi Kang<sup>2,3</sup>, Liang Zhang<sup>3</sup>, Yi Xiong<sup>4</sup>, Wanli Ouyang<sup>1</sup>, Zhiqiang Hu<sup>7</sup>, Guisheng Fan<sup>6</sup>, Yan Feng<sup>4</sup>, Guangyu Yang<sup>4</sup>, Qian Liu<sup>4</sup>, Liang Hong<sup>1,2,3,8\*</sup>.

1. Shanghai Artificial Intelligence Laboratory, Shanghai 200240, China
2. Shanghai National Center for Applied Mathematics (SJTU Center), & Institute of Natural Sciences, Shanghai Jiao Tong University
3. School of Physics and Astronomy & School of Pharmacy, Shanghai Jiao Tong University, Shanghai 200240, China
4. School of Life Sciences and Biotechnology, Shanghai Jiao Tong University, Shanghai 200240, China
5. Department of Instrument science and engineering, Shanghai Jiao Tong University, Shanghai 200240, China
6. School of Information Science and Engineering, East China University of Science and Technology, Shanghai 200240, China
7. SenseTime Research, Shanghai, 201103, China
8. Zhang jiang Institute for Advanced Study, Shanghai Jiao Tong University, Shanghai 200240, China

## Abstract

Designing protein mutants of both high stability and activity is a critical yet challenging task in protein engineering. Here, we introduce PRIME, a deep learning zero-shot model, which can suggest protein mutants of improved stability and activity without any prior experimental mutagenesis data. By leveraging temperature-guided language modelling, PRIME demonstrated superior predictive power compared to current state-of-the-art models on the public mutagenesis dataset over 33 proteins. Furthermore, we carried out wet experiments to test PRIME on five distinct proteins to engineer certain physicochemical properties, including thermal stability, rates of RNA polymerization and DNA cleavage, hydrolase activity, antigen-antibody binding affinity, or even the nonnatural properties, e.g., the ability to polymerize non-natural nucleic acid or resilience to extreme alkaline conditions. Surprisingly, about 40% AI-designed mutants show better performance than the one before mutation for all five proteins studied and for all properties targeted for engineering. Hence, PRIME demonstrates the general applicability in protein engineering.

## Introduction

Proteins are the fundamental constituents of living systems, playing a crucial role in a vast

---

---

array of biological processes, spanning from enzyme catalysis and cellular metabolism to immune responses, signal transduction, and transport, among others. Beyond their biological significance, proteins are critical to numerous industries. In biomedicine, they serve as therapeutic agents and targets; in the food industry, they are involved in food processing and preservation; in brewing, they are essential to the production process; and in chemical engineering, they act as key catalysts for various reactions. Additionally, proteins are the cornerstone of in vitro diagnostic (IVD) tests, being instrumental in the detection and monitoring of numerous diseases. However, proteins extracted from biological organisms, namely “wild type”, often require modifications to make them suitable for industrial applications. This is primarily because the physicochemical environments (for example, temperature) in which these proteins need to function in industrial settings are often drastically different from their native biological contexts<sup>1,2</sup>. Therefore, to meet the demands of these diverse application scenarios, the proteins need to be engineered through mutations to improve their physicochemical properties. These modifications could aim to enhance stability under extreme temperature or pH conditions, or to increase enzymatic activity and specificity. The process of optimizing proteins for such industrial applications typically involves iterative cycles of mutation, screening, and selection - a labor-intensive and time-consuming endeavor.

Indeed, a primary focus in protein engineering lies in enhancing the robustness of proteins to function effectively under extreme conditions. The initial target often is the improvement of thermostability, allowing proteins to retain their structure and function at high temperatures. Additionally, improving resistance to extreme pH environments (acidic or alkaline) and harsh solvents is equally crucial. Such enhancements enable proteins to be utilized in a broader range of industrial processes, many of which involve conditions that can denature or deactivate proteins. In addition to stability, augmenting protein activity is another major goal in protein engineering. This could involve optimizing the efficiency of gene editing enzymes, increasing the rate of biocatalytic reactions, enhancing the binding affinity of antibodies, and so on. All these modifications aim at boosting the performance of proteins in their respective applications. Hence, the core mission of protein engineering involves designing protein mutants that not only exhibit heightened activity but also display increased stability under a range of conditions.

A sophisticated interplay exists between protein stability and activity, both of which are fundamental to their overall performance. Protein stability is crucial for maintaining the proteins' native structure and function<sup>3</sup>. Adverse conditions can provoke alterations in protein folding states, leading to the loss of native conformation and function. Notably, proteins are prone to denaturation under extreme circumstances, including elevated temperatures or exposure to potent amino acids and alkalis<sup>4,5</sup>. Improving protein stability can increase evolvability by allowing a protein to tolerate a broader range of beneficial mutations while maintaining its native structure<sup>6</sup>. However, overemphasis on stability at the expense of protein flexibility may inhibit enzymatic activity<sup>7</sup>. Therefore, achieving an optimal balance between stability and activity is vital for optimizing protein efficacy across diverse contexts.

As computational simulation and related technologies continue to advance, various software tools have emerged to enhance protein thermostability, including Rosetta<sup>8</sup>, ABACUS<sup>9</sup>, and FoldX<sup>10</sup>, which employ physical or statistical potential functions. While these computational methods often provide relatively accurate stability predictions, their capacity to predict protein biological activity

---

---

is limited. Typically, modifying the biological activity of proteins requires long-term (~years) meticulous experimental research into their working mechanisms, which is the primary method of rational protein design. However, mechanistic research is time-consuming and labor-intensive, and it increasingly fails to meet the modification needs of many important industrial enzymes commonly used in everyday applications. In recent years, deep learning has been extensively applied in protein engineering. Large-scale protein language models<sup>11-15</sup>, such as those utilizing self-supervised learning of protein sequence to understand protein sequence semantics and grammar, have demonstrated high predictive performance for protein fitness<sup>16</sup>, even in zero-shot settings<sup>15,17,18</sup>. Zero-shot setting here means the model can predict the mutation sites of a protein to improve its properties without relying on any prior experimental mutagenesis data. However, most of these models, pre-trained on extensive protein sequence databases, lack interpretation for distinct protein properties, such as thermostability and enzymatic activity. These specific properties indeed form the real goals of protein engineering. Other supervised deep learning methods often exhibit high accuracy in predicting protein function but rely on high-throughput experiments to generate hundreds or even thousands of data points<sup>19,20</sup>. This approach may not be practical for many proteins due to resource limitations. In this study, we amassed a comprehensive dataset comprising 96 million sequence-host bacterial strain optimal growth temperatures (OGT)<sup>21</sup>. Host bacterial strain optimal growth temperature has been shown to strongly correlate with information such as protein optimal enzymatic activity temperature and melting temperature<sup>22</sup>. Leveraging this dataset, we developed an interpretable deep learning-based methodology, termed PRIME, Protein language model for Intelligent Masked pretraining and Environment (temperature) prediction. In its training process, PRIME utilizes a masked language modeling (MLM) task, a methodology inspired by the transformer-based language models<sup>23</sup>. This task involves artificially modifying protein sequences based on the natural probability distribution of amino acids, followed by attempting to restore the sequences to their original state. Such a procedure allows PRIME to learn and comprehend the semantic and grammatical features inherent in protein sequences. Alongside this, PRIME capitalizes on a multi-task learning paradigm to capture the temperature traits associated with these sequences. This approach fosters an inherent predisposition in PRIME to assign superior scores to protein sequences exhibiting enhanced temperature tolerance and exhibiting conformity to natural biological principles. PRIME is trained with the objective of predicting optimal growth temperatures (OGTs) across a wide range of bacterial strains. As a result, PRIME naturally correlates higher scores with sequences that are more likely to contribute to robustness and survivability in varied environmental conditions, including extreme temperature scenarios. Therefore, PRIME proves particularly proficient in the design and optimization of industrial enzymes and proteins that often demand high-temperature tolerance and resilience for practical applications. Our model has demonstrated exceptional predictive performance relative to other state-of-the-art (SOTA) models, especially in forecasting the thermostability (change of  $T_m$ ) and enzymatic activity of protein mutated sequences.

To further evaluate the efficacy of our model, we applied it to five distinct proteins and subjected the results to wet experimental validation. The proteins under study included T7 RNA polymerase, creatinase, KmAgo, the variable domain of the heavy chain of a nano-antibody against growth hormone (VHH) and DNA polymerase. Each protein was engineered with a distinct objective for the real practical purpose: the T7 RNA polymerase and creatinase were modified to enhance thermostability, the prokaryotic Argonaute protein was modified to augment its activity at

---

---


ambient temperatures, the VHH was engineered to increase stability under extreme alkaline pH conditions, while the DNA polymerase was aimed to enhance the polymerizing rate of non-natural nucleic acids. Without any prior experimental mutagenesis data, we use the PRIME model to pick up a few tens (~30) of best-ranking single-site mutants for experimental testing. Our results revealed that approximately 40% of these mutants displayed significant improvement of the physicochemical property targeted for engineering. We note that certain improvement occurs on the protein properties, which are rarely seen in nature, such as the tolerance to extreme basic conditions (pH > 13) and the ability to polymerize non-natural nucleic acids. This indeed underlines the superior predictive performance and general applicability of PRIME, facilitating the design of mutation sequences with enhanced stability and activity.

## Results

### PRIME Architecture

PRIME is a pre-trained model based on the Transformer architecture<sup>24</sup>, as illustrated in the Figure 1.A. PRIME consists of three main components. The first is the encoder module for sequence feature extraction, which is a Transformer encoder model to extract the latent representation of the sequence. The second component is the masked language modeling (MLM) module, which is designed to learn the contextual representation of amino acids. The third component is the Optimal Growth Temperature (OGT) prediction module, which can predict the OGT of the organism in which the protein is located, based on the latent representation. The model and training details of PRIME are described in the supplementary information.

---



**Figure 1. Overview of the PRIME architecture and its applications.** **A:** The architecture design of PRIME. PRIME incorporates a BERT-oriented transformer encoder, augmented by two domain-specific modules, one for Masked Language Modeling (MLM) and another for Optimal Growth Temperature (OGT) prediction tasks. The learning objectives comprise three distinct loss functions: MLM loss, quantified via cross-entropy; OGT loss, assessed through the mean squared error criterion; and Correlation loss, evaluated by the inverse Pearson correlation coefficient. **B:** Elucidation of the Correlation Loss. The stochastically generated single-point mutant sequences are scored by the MLM module while their corresponding OGTs are predicted by the OGT module. A subsequent Pearson correlation between the predicted OGT values and their corresponding MLM scores is computed and optimized, thereby prompting an alignment between mutant scores and thermal profiles. The objective of this loss function is the maximization of the Pearson correlation coefficient. **C:** The use of PRIME for single-point mutation scoring. The wild-type sequence is reconstructed via the MLM module, generating a probabilistic distribution for amino acid identity at the mutation locus. The mutational impact is subsequently quantified by the log-odds ratio between the mutated and wild-type amino acids. **D:** The strategy for generating multi-point mutants involves several steps. First, PRIME is used to assess the impact of single-point mutations, from which the top-K mutants are selected for

---

experimental fitness evaluation in the laboratory. Subsequently, this experimental data serves as the training set to fine-tune PRIME. The fine-tuned model is then employed to predict the fitness of multi-point mutants. Based on these predictions, the top-K multi-point mutants are selected for further experimentation.

### 1) The pre-training objectives of PRIME

The learning of PRIME is structured into three alternative objectives: the Masked Language Modeling (MLM) objective, the Optimal Growth Temperature (OGT) prediction objective, and the correlation objective. The details of these objectives are as follows:

#### 1.1) Masked language modeling

Masked Language Modeling (MLM) is often used as a pre-training method for sequential data representation. In this objective, noised protein sequences serve as the input, wherein parts of tokens are masked as '[MASK]' or substituted with alternative tokens. The training objective is to reconstruct these noised tokens. This approach facilitates the model's ability to capture dependencies among amino acids as well as contextual information along the sequence. The details can be found in the supplementary information. Moreover, we can use this reconstruction process to score mutations (Figure 1C).

#### 1.2) Optimal growth temperature (OGT) prediction

The second training objective is optimized under supervised conditions. We utilize a dataset containing 96 million protein sequences annotated with OGT to train the PRIME model. The input of this objective is protein sequence, and the OGT module generates a temperature value ranging between 0 and 100°C. Notably, the OGT and MLM modules operate with a shared encoder. This architecture enables the model to simultaneously capture amino acid contextual information and temperature-related sequence characteristics.

#### 1.3) Correlation objective

To facilitate feedback from the predicted OGTs to the MLM scores, we introduce a learning objective aimed at aligning these two metrics. Specifically, for a protein sequence, random single-point mutants are generated. We utilized wild-type sequences as input to the Encoder and MLM (Masked Language Model) module of PRIME. Subsequently, the MLM module outputs a probabilistic distribution matrix for the reconstruction matrix  $M$ , denoted by dimensions  $[L, K]$ , where  $L$  denotes the length of the sequence and  $K$  represents the count of possible amino acids, specifically 20. The element in the  $k_{th}$  row and  $a_{th}$  column of this matrix,  $M_k^a$ , represents the probability, as inferred by PRIME, of the appearance of the amino acid  $a$  at the  $k_{th}$  position. Given any sequence  $S = (s_1, s_2, \dots, s_L)$  of the same length, its sequence log-likelihood score is computed by:  $\sum_{i=1}^L \log p(M_i^{s_i})/L$ . This value is referred to as the "sequence score" for that particular sequence. For mutants generated, their sequence scores are equivalent to these computed log-likelihoods, or called "MLM score". And their corresponding OGTs are assessed by the OGT module. The Pearson correlation between these mutant sequence scores and predicted OGT values, is subsequently optimized serving to align the mutant thermal profiles with their corresponding MLM scores (Figure 1B.). The goal of this objective is the maximization of the Pearson correlation coefficient. We use Pearson correlation as our learning objective due to its differentiable properties (for backpropagation), in contrast to the non-differentiable nature of Spearman correlation.

### 2) Zero-Shot Single-Point Mutation Scoring.

Models trained with the Masked Language Modeling objective can output the likelihood of amino

---

---

acids appearing at a specific position based on surrounding context. We use this to score single-point mutations. Given a mutation, we treat the amino acid in the wild-type protein as a reference and compare its likelihood to that of the mutated amino acid. The mutations are then scored using the log-odds ratio at the mutated position. (See in Figure 1C, details are in supplementary information).

3) Augmentation of single-point mutation prediction performance in PRIME through fine-tuning on homologous sequences via the MLM learning objective.

While PRIME exhibits commendable performance in zero-shot mutant effect prediction, we observed that additional unsupervised fine-tuning of the language modeling module on homologous protein sequences of target proteins yields improved results, without adding supervision from experimental data. Explicitly, for the fine-tuning process, we deploy homologous sequences of the proteins of interest as an unsupervised dataset, optimizing both the encoder and masked language modeling (MLM) modules of PRIME. Evaluation's result substantiate that this method significantly amplifies PRIME's predictive accuracy for single-point mutations.


### **PRIME outperforms state-of-the-art methods in predicting thermostability of mutated protein sequence**

We conducted a comparison of the zero-shot prediction capacity on thermostability between our model, PRIME, and several current state-of-the-art models, including deep learning models ESM-1v<sup>17</sup>, MSA-transformer<sup>13</sup>, Tranception<sup>25</sup> CARP<sup>26</sup> and MIF-ST<sup>27</sup>, as well as the traditional computational method, Rosetta<sup>28</sup>. Notably, among these methods, MIF-ST and Rosetta incorporates protein structure information, whereas the others rely solely on protein sequence. Our analysis utilized a dataset derived from MPTherm<sup>29</sup>, FireProtDB<sup>30</sup> and ProThermDB<sup>31</sup>, featuring single-point mutations in proteins with  $\Delta T_m$ , i.e., changing of melting temperature as compared to the wild type, collected under the same experimental pH and ensuring a minimum of 10 data points per protein, which amassing a total of 67 assays. Concurrently, the analysis also incorporated assays from Deep Mutational Scanning (DMS), specifically those housed within ProteinGym<sup>25</sup>. ProteinGym presents a meticulously constructed substitution benchmark, characterized by the experimental delineation and assessment of approximately 1.5 million missense variants. These variants are dispersed across 87 distinct DMS assays and encompass a range of protein properties including, but not limited to, enzymatic catalysis, binding affinity, stability and fluorescence intensity. Such a comprehensive assembly of missense variants within the substitution benchmark of ProteinGym provides a robust and expansive repository, thereby facilitating the nuanced evaluative study of the myriad documented missense variants. This repository thus serves as a valuable asset for the systematic examination and interpretation of the diverse and intricate landscape of protein mutations and their associated properties.

These comprehensive datasets enabled a systematic investigation of the impact of specific mutations on protein fitness and thermostability, supporting the development and validation of advanced predictive models such as PRIME. The comparison provides valuable insights into the relative performance of different modeling approaches and highlights the potential of PRIME for

---

predicting protein mutations in a zero-shot setting. The results are illustrated in Figure 2A (Detailed data for every assay can be found in the separated Excel file in the Supplementary Information). As can be seen, PRIME demonstrates superior performance over all the other methods in predicting protein fitness and thermostability. In the ProteinGym benchmark, PRIME exhibits superior performance over its competitor, ESM-1v, registering a score of 0.451 against 0.411 ( $p=0.047$ , Wilcoxon). This lead is further accentuated when PRIME is finetuned on ProteinGym's homologous sequences using only the MLM objective, scoring 0.523 versus ESM-1v's 0.411 ( $p=0.0001$ , Wilcoxon). In the  $\Delta T_m$  dataset, PRIME's performance significantly surpasses its closest competitor, MIF-ST, with scores of 0.472 and 0.377 respectively ( $p=0.044$ , Wilcoxon). It's crucial to note that the OGT employed by PRIME isn't a direct representation of protein  $T_m$ . Instead, a correlation exists between them<sup>22</sup>. There are some enzymes from thermophiles turn out to be not very thermostable<sup>32</sup>. However, even when leveraging the slightly imprecise OGT as a stand-in for protein sequences'  $T_m$  attribute, PRIME markedly outshines models that don't incorporate OGT. For instance, the similar-architecture counterpart, ESM-1v, achieves only 0.278 in the  $\Delta T_m$  dataset. We posit that PRIME's performance would witness a significant boost with access to a vast dataset of accurate  $T_m$  values for natural proteins. These findings underscore PRIME's potential in protein engineering endeavors, particularly in crafting protein sequences with enhanced thermostability and other fitness attributes. Across the board, PRIME outclasses both traditional computational strategies and other deep learning models, underscoring its unparalleled effectiveness.



**Figure 2.** Comparison of performance between PRIME and other methods. A: Unsupervised Model Benchmarking on  $\Delta T_m$  and ProteinGym Datasets. When we refer to "PRIME (homologous sequences)," it denotes the fine-tuning of the PRIME model using the MLM loss on homologous sequences of the target proteins present in either ProteinGym or  $\Delta T_m$  datasets. B: Supervised Prediction of  $T_m$  (Melting Temperature). C: Supervised Prediction of  $T_{opt}$  (Optimal enzymatic activity Temperature). For supervised benchmarks, three metrics are employed to gauge the models' accuracy and predictive power: Spearman correlation, Pearson correlation, and  $R^2$  (coefficient of



---

determination). Where the datasets and data splits for  $T_m$  (~39,000 protein sequences) and  $T_{opt}$  (~1900 sequences) are referenced from Ref<sup>33</sup>. We obtained the wild-type protein structure from the Protein Data Bank and employed AlphaFold2<sup>34</sup> to construct structures absent in PDB for the input to Rosetta MIF-ST. In Figures B and C, the error bars represent the standard deviation, averaged over three random seeds used for finetuning both ESM2 and PRIME, keeping the pre-trained model parameters fixed. The data of DeepET was from the reference<sup>33</sup>. We were unable to obtain the finetuning code for DeepET, which is why there are no error bars associated with it. Notably, DeepET failed to generate scores for 27 assays within the ProteinGym dataset. Due to this limitation, we excluded DeepET from the benchmarking process on the ProteinGym dataset. The data points and the p-value test associated with Figure 2A was shown in Table S1.

In addition to the zero-shot assignment, we also tested the representational capacity and transferability of PRIME. Specifically, we conduct supervised fine-tuning on two temperatures related downstream tasks, keeping the pre-trained model parameters fixed. (Details can be found in SI). As the pretraining of PRIME incorporates the optimum growth temperature of the bacterial where the protein lives in, it is anticipated that PRIME can also perform better in predicting other properties of proteins associated with temperature. As exhibited in Figure 2B and 2C, PRIME also outperforms other supervised methods in the task of predicting melting temperature ( $T_m$ ) of a native protein and its optimal enzymatic activity temperature ( $T_{opt}$ ). We delved deeper into understanding the individual contributions of the three core modules within PRIME: the OGT prediction module, the mask language modeling module (MLM), and the correlation term. Our findings, detailed Table S2, highlight that relying solely on either the OGT prediction or the MLM module leads to a dip in PRIME's performance. Among these, the MLM module stands out as having the most pronounced effect across all zero-shot benchmarks. The OGT module plays a pivotal role in  $\Delta T_m$  prediction, with the standard PRIME achieving a score of 0.472, in contrast to PRIME/-OGT which scores 0.323 ( $p=9e-10$ , Wilcoxon). Similarly, the correlation term significantly influences  $\Delta T_m$  prediction, with PRIME/-correlation registering a score of 0.434 ( $p=8e-5$ , Wilcoxon). In the context of the ProteinGym benchmark, both the OGT and correlation terms continue to exert a significant influence ( $p<0.05$ ). This is likely attributed to the presence of tasks in ProteinGym that are associated with protein stability. This finding highlights the significance of combining both the OGT prediction, MLM, and correlation modules in the PRIME model to achieve optimal performance. The synergistic effect of these three modules allow the model to better understand the complex relationships between protein sequences and their thermostability properties, ultimately resulting in improved predictive capabilities. The integration of all these modules in the PRIME model ensures a more comprehensive understanding of the protein sequence information, which in turn contributes to its superior performance compared to other state-of-the-art models. Further, we assessed PRIME's performance in other supervised protein engineering tasks. Specifically, in the FLIP benchmark<sup>35</sup>, PRIME outperformed ESM-2 and CARP in 11 out of the 12 tasks, as shown in Table S3. Notably, in the Meltome<sup>36</sup> dataset task of FLIP, which involves predicting the  $T_m$  of human-derived proteins, PRIME, integrated with OGT information, consistently surpassed models with similar architectures like ESM2. This indicates that even though PRIME's pre-training process only learned the OGT information of bacterial-derived protein sequences, it still excels in predicting the  $T_m$  temperature attributes of proteins from other species. This demonstrates PRIME's remarkable generalization capabilities.

---

---

---

## Wet experimental testing of AI designed mutants of various proteins for different engineering purpose

In practical applications of protein engineering, the prevailing approach involves the identification of the good positive single-site mutations with which the protein will become better (more active or more stable), and then combines them to form multisite mutants of desired property probably through a greedy search method<sup>20</sup>. Thus, the successful identification of these greatly performing single-site mutations form the cornerstone of successful protein engineering. To further substantiate the effectiveness and generosity of our methodology, we tested the PRIME model on designing single-site mutant for five distinct proteins, namely T7-RNA polymerase, creatinase, prokaryotic Argonaute protein, the variable domain of the heavy chain of a nano-antibody against growth hormone (VHH) and a DNA polymerase. The specific engineering goals for these proteins varied: enhancing thermostability for T7-RNA polymerase and creatinase; augmenting activity at ambient temperatures for prokaryotic Argonaute protein; increasing stability under extreme alkaline pH conditions for VHH; and increasing the polymerizing rates of non-natural nucleic acid for DNA polymerase (2'-fluoroarabino nucleic acid). Briefly, we deployed PRIME to rank all single-site mutants within the landscape, based on the likelihood of the mutated sequences compared to their wildtype counterparts (see Mutated protein sequence scoring strategy). Subsequently, we selected a few tens of top-ranked mutant for wet-lab experimental verification. Detailed results are elucidated below.

### T7 RNA polymerase

T7 RNA polymerase is a monomeric enzyme derived from T7 bacteriophage, comprising a total of 883 amino acids. Since its initial utilization in RNA synthesis in the early 1980s, T7 RNA polymerase has become a crucial tool in the fields of molecular biology and genetic engineering<sup>37</sup>. It is now commonly employed in various applications such as in vitro transcription experiments, mRNA vaccine production<sup>38</sup>, cell-free protein expression<sup>39</sup> and isothermal amplification detection techniques<sup>40,41</sup>, etc. However, T7 RNA polymerase also has some application drawbacks, for example, it produces by-product double-stranded RNA (dsRNA) during the transcription process<sup>42</sup>, which poses a great cost for mRNA vaccine production in terms of purifying<sup>43</sup>. Recent studies have indicated that increasing the reaction temperature to above 48°C effectively reduces the by-products<sup>44</sup>. Nevertheless, the wild-type T7 RNA polymerase unfolds at temperatures around 45°C, resulting in decreased enzymatic activity and an inability to transcribe the desired target products at higher temperatures. Therefore, there is a critical need to enhance the thermal stability of T7 RNA polymerase.

In this study, we utilized PRIME to forecast mutation sites in T7 RNA polymerase and chose the top 28 single-site mutants for subsequent experimental verification. As illustrated in Figure 3B, our experimental findings reveal that 46% of mutants exhibit a  $T_m$  value exceeding that of the wildtype. Intriguingly, 50% of thermally positive mutants simultaneously demonstrate enhancements in both thermostability and activity. Notably, the highest-performing mutant showcased a 3.5°C increase in the melting temperature and presented a 40-fold augmentation in enzymatic activity after a 5-minute heat treatment at 48°C, compared to the wildtype (Detailed data can be found in Table S3).

---

---

---

## Creatinase

Creatinase, a dimeric proteinase, is widely employed in enzymatic assays for measuring creatinine levels<sup>45,46</sup>. It is primarily derived from microorganisms such as *Pseudomonas.*, *Bacillus.*, and *Alcaligenes*<sup>47-49</sup>. Creatinase is crucial in medical diagnostics and plays a role in quantifying creatinine in serum and urine<sup>50</sup>. Elevated creatinine levels indicate impaired kidney or muscle function. Nevertheless, the optimal catalytic temperature for creatinase typically falls within the range of 30-40°C, which constraints both the industrial and clinical diagnostic applications. Enhancing the thermal stability of creatinase not only improves the efficiency of clinical creatinine detection but also facilitates enzyme production, storage, and transportation.

Here, we employed the unsupervised model PRIME to predict single-point mutations in creatinase obtained from *Alcaligenes faecalis*<sup>51</sup>. At the end 28 single-site mutants were selected for experimental validation. As depicted in Figure 3D, among the 28 mutants, 9 exhibited improved thermal stability. The positive rate is about 32%. Remarkably, about 44% of these positive mutants also demonstrated an enhancement in activity. (Detailed data can be found in Table S4). It demonstrates that our model can provide mutants of good enzymatic rate with improved thermostability.

## Argonaute protein

Prokaryotic Argonaute proteins (pAgos) have emerged as a compelling research focus within the field of molecular biology, presenting promising alternatives or complementary approaches to established biotechnologies, such as the CRISPR/Cas system<sup>52-55</sup>. Propositions have been advanced concerning the role of pAgos in the creation of accurate and efficient nucleic acid diagnostic tools, exploiting their unique DNA-guided DNA/RNA targeting mechanisms.<sup>56</sup> At present, the low enzymatic activity of pAgos at ambient temperatures presents a substantial impediment to their utilization in nucleic acid detection. Consequently, the augmentation of protein activity is of pressing scientific interest. In this study, we selected *Kurthia massiliensis* Ago (*KmAgo*) - an Argonaute protein which retains functionality at ambient temperatures as a prospective candidate for room-temperature nucleic acid detection applications<sup>57</sup>. Through employing PRIME prediction model, we chose 28 Single-site mutants, among which ~39% exhibit enhanced cleavage activity (~39% positive rate, see Figure. 3F, and comprehensive data are presented in Table S5). And the best mutant's activity is enhanced by 4.7-fold.

## VHH

VHH antibody is the antigen binding fragment of heavy chain only antibodies<sup>58</sup>. Due to the advantages of small size, monomer state, robust structure and easy tailoring, VHH has been employed as an important tool in medical research and clinical antibody drug development<sup>59</sup>, which have been developed as an affinity ligand to selectively purify biopharmaceutical, for example prothrombin, tetrabromobisphenol A, intercellular adhesion molecule 1 and so on<sup>60-62</sup>. In the practical production of biological products, the most widely used method of clean-in-place (CIP) is 0.5 M NaOH cleaning for 24 hours. Hence, VHH antibodies used for biopharmaceutical purification need mutational engineering to tolerate the harsh alkaline condition, which is rarely seen in nature<sup>63,64</sup>.

In this study, we employed our unsupervised model PRIME to predict mutation sites for an

---

---


VHH antibody against growth hormone which we select from an immunized camelid. The top 29 mutants were chosen for further testing, 11 of 29 mutants enhanced stability after incubation at 0.3 M NaOH for 24 h, as shown in Figure 3H and Table S6. Among these, the M1(A57D) mutation displayed a remarkable twelve-fold enhancement in alkali tolerance. Besides, ~73% of positive mutations with better alkali tolerance than the wildtype also shows increased affinity for antigen without the alkaline treatment. Again, this demonstrates that our model can suggest good mutation sites of both high activity, i.e., the binding affinity without alkaline treatment, and stability, i.e., tolerance to the extreme alkaline condition.

### DNA polymerase

Tgo is a DNA polymerase that has been identified in the thermophilic bacterium *Thermococcus gorgonarius*, which was isolated from a geothermal vent in New Zealand<sup>65</sup>. Tgo has been found to accurately replicate 2'-fluoroarabino nucleic acid (FANA), a genetic polymer with 2'-fluoroarabino residues in deoxyribonucleotides<sup>66,67</sup>. However, Tgo DNA polymerase can only catalyze the synthesis of FANA on the DNA template at a rate of ~15 nt/min<sup>67</sup>, which is much lower than that of Tgo for DNA synthesis (~400 nt/min)<sup>68</sup>, limiting the application of FANA as a substitute for DNA in information storage<sup>69</sup>, disease treatment<sup>70,71</sup> and other fields<sup>72,73</sup>. Vitor et al. employed the compartmentalized self-tagging (CST) strategy to screen the engineered FANA polymerase Tgo-D4K (TgoT: L403P, P657T, E658Q, K659H, Y663H, E664K, D669A, K671N, T676I) from the TgoT (V93Q, D141A, E143A, A485L) library<sup>74</sup>. The polymerase was able to extend FANA on the DNA template at a rate of ~80±27 nt/min, while the rate of DNA extension on the DNA template was reduced to 16±3 nt/min<sup>68</sup>. Furthermore, the synthesis rate of Tgo-D4K for FANA is still lower than that of Tgo for DNA synthesis. Therefore, new methods are required to modify existing polymerases to screen for polymerases with higher FANA synthesis rates.

In this study, we commenced our investigation with Tgo-D4K as the starting point. Utilizing PRIME, we systematically screened potential mutation sites across various domains of Tgo-D4K. Ultimately, we selected 27 promising mutations for subsequent experimental validation. The polymerase kinetic profiling (PKPro) strategy was used to detect the FANA synthesis rate of the mutants as previously described<sup>68</sup>. The experimental results (Figure 3J and Table S7) showed that more than 40% of the polymerases had a higher FANA synthesis rate than Tgo-D4K, and the single-point M27 mutation was identified, which can significantly increase the extension rate to ~3.2-fold that of the Tgo-D4K enzyme.

---



**Figure 3.** The structure and experimental results of single-site mutants predicted by PRIME for T7 RNA polymerase (A,B), creatinase (C,D), KmAgo (E,F), VHH (G,H) and DNA polymerase (I,J) are depicted. The data points representing the mutations were arranged systematically in ascending order, with the corresponding value for the wild-type protein delineated by a black bar for comparative purposes. Mutants that exhibited superior performance compared to their wild-type counterparts in terms of targeted attributes are emphasized in yellow. While the negative mutants are shown in blue. The engineering goals varied between proteins for the real practical purpose: for T7 RNA polymerase and creatinase, the objective was enhanced thermostability ( $T_m$ ); for KmAgo, it was to increase the nucleic acid cleavage activity; for VHH, the goal was to improve the tolerance ability under extreme alkaline pH conditions (EC50 of VHH binding to the antigen); and for DNA polymerase, the aim was to accelerate the synthesis rate of 2'-fluoroarabino nucleic acid.

---

---

**Some positive mutants predicted by PRIME fall outside the knowledge space of conventional rational design by human experience.**

In rational protein engineering, chemically plausible mutations are typically deemed beneficial<sup>2,75</sup>. Conventional wisdom posits that mutations should not disrupt hydrogen-bonding or salt-bridge interactions, especially for designing thermal-stable proteins<sup>76</sup>. Moreover, earlier experience all tells that better mutations should be evolutionarily conserved<sup>77</sup>. Yet, the positive mutation sites predicted by PRIME challenged this conventional wisdom. We conducted structural and evolutionary conservation analyses on all positive mutations for T7 RNA polymerase, creatinase, and KmAgo (Detailed data were shown in Table S8, S9, S10). We discovered that most of these positive mutation sites, which render enhanced thermostability or enzymatic activity to the proteins, did not impact the number of hydrogen bonds or salt bridges. Surprisingly, our study uncovered that some mutations, such as S430P in T7 RNA polymerase, which reduced one hydrogen bond (between S430 and F432) post-mutation (Figure S1A), resulted in an enhancement of  $T_m$  by 2.7°C and a 29-fold increase in activity. Similarly, mutation D17V in creatinase, leading to a decrease in a salt bridge (between D17 and R91, Figure S1B), contributed to a  $T_m$  increase by 2.4°C. These intriguing observations underscore the complex interplay between molecular modifications and functional enhancements in proteins, an understanding that defies conventional wisdom in rational design based on human experience. The exact mechanisms elucidating why these mutations enhance functionality and how our model identifies them remain elusive and beyond the scope of this present investigation. This remarkable contradiction between PRIME predictions and traditional protein engineering understanding certainly merits deeper exploration.

**Conclusion and discussion**

In conclusion, we unveil PRIME, a pioneering deep learning methodology, which deftly exploits an expansive dataset, comprising sequence-host bacterial strain optimal growth temperatures. Utilizing an adapted masked language model (MLM) for Optimal Growth Temperature (OGT) prediction, PRIME capably assimilates semantic, grammatical, and temperature-related characteristics of protein sequences. Our systematic *in silico* experimental validations unequivocally establish PRIME's superior performance over other leading models, such as ESM-1v, ESM2, MSA-transformer, Tranception, and Rosetta, in predicting thermostability and activity of mutation sequences in proteins. Facilitated by PRIME, we engineered five single-site mutated proteins, achieving better performance on targeted physicochemical properties, with an approximate ~40% success rate among ~30 total mutants designed by AI. This underscores PRIME's potential in the realm of protein engineering.

Traditional protein engineering strategies typically rely on high-throughput experimental screening to identify beneficial mutations. This approach involves generating a large library of protein variants, each with different mutations, and then screening these variants for desired traits. These traits can include enhanced stability under certain conditions (such as high temperature or extreme pH) or increased enzymatic activity<sup>78</sup>. These techniques, however, are labor and time-intensive and costly. For many important proteins, designing a high-throughput experimental protocol is challenging, rendering low-throughput experimental testing a more common and practical approach. If we can identify a sufficient number of superior single-site mutations via low-throughput experimentation alone, we can then build on these single-site mutations to generate cumulative multi-site mutations. In such cases, a more targeted approach based on computational

---

---

prediction can be valuable. Tools like PRIME can predict the impact of specific mutations on protein stability and activity, enabling a more focused and efficient approach to engineering improved protein variants. By reducing the reliance on extensive experimental screening, such computational tools can make the protein engineering process more efficient and accessible, potentially broadening the range of proteins that can be effectively engineered.

Furthermore, the adaptable nature of PRIME's general language model representation offers scope for its deployment in other prediction tasks such as determining the melting temperature ( $T_m$ ) or optimal enzymatic activity temperature ( $T_{opt}$ ) of native proteins. By lowering the thresholds for protein modification, PRIME allows for improvements in the stability and biological activity of proteins, bypassing the need for extensive mechanistic investigations. Additionally, PRIME's multi-task learning paradigm, which incorporates domain knowledge into the language model, could catalyze the development of more specialized AI methodologies across diverse fields of specialty.

To our surprise, the PRIME model can predict mutation sites to improve the properties of a protein, which are rarely seen in nature. For example, improve the stability of the antibody in extreme alkaline condition and the polymerization rate of a DNA polymerase on nonnative nucleic acids. This highlights the general capability of the model on protein engineering.

### **Acknowledgements**

This work was supported by the grants from the National Science Foundation of China (grant number 12104295). The engineering of VHH was supported by GeneScience Pharmaceuticals. We acknowledge Shanghai Artificial Intelligence Laboratory for computing resources.

### **Author contributions**

P.T. and L.H. designed and supervised this project. P.T. and M.L. designed the model architecture of PRIME and designed all the mutations. Y.Y. and F.J. conducted the experiment of T7-RNA polymerase and Creatinase. L.Z. conducted the experiment of KmAgo. B.W. performed the experiment of VHH. X.S. Y.P. and J.S. conducted the experiment of DNA polymerase. L.K., L.Z., Y.X., W.O., Z.H., G.F., collected the datasets and performed the benchmark jobs. Y.F., G.Y. and Q.L. supervised all the wet experiments. All authors reviewed and accepted the manuscript. P.T, M.L, Y.Y., F.J., L.Z., B.W. and X.S. contributed equally to this work.

### **Competing interests**

Patents have been filed in 2022 for all these mutations in the present work. Among which, GeneScience Pharmaceuticals, a company that discovered the native sequence of VHH, own the patent of VHH.

### **Reference**

- 1 Xia, Y. *et al.* Development of thermostable sucrose phosphorylase by semi-rational design for efficient biosynthesis of alpha-D-glucosylglycerol. *Applied Microbiology and Biotechnology* **105**, 7309-7319 (2021).
  - 2 Reetz, M. T., Sun, Z. & Qu, G. *Enzyme Engineering: Selective Catalysts for Applications in Biotechnology, Organic Chemistry, and Life Science.* (John Wiley & Sons, 2023).
  - 3 Pinney, M. M. *et al.* Parallel molecular mechanisms for enzyme temperature adaptation. *Science* **371**, eaay2784 (2021).
-

- 
- 4 Kauzmann, W. in *Advances in Protein Chemistry* Vol. 14 (eds C. B. Anfinsen, M. L. Anson, Kenneth Bailey, & John T. Edsall) 1-63 (Academic Press, 1959).
- 5 Jaenicke, R. Protein stability and molecular adaptation to extreme conditons. *European Journal of Biochemistry* **202**, 715-728 (1991). [https://doi.org:https://doi.org/10.1111/j.1432-1033.1991.tb16426.x](https://doi.org/https://doi.org/10.1111/j.1432-1033.1991.tb16426.x)
- 6 Bloom, J. D., Labthavikul, S. T., Otey, C. R. & Arnold, F. H. Protein stability promotes evolvability. *Proceedings of the National Academy of Sciences* **103**, 5869-5874 (2006). <https://doi.org:10.1073/pnas.0510098103>
- 7 Liu, Z. *et al.* Entropic contribution to enhanced thermal stability in the thermostable P450 CYP119. *Proceedings of the National Academy of Sciences* **115**, E10049-E10058 (2018). <https://doi.org:10.1073/pnas.1807473115>
- 8 Das, R. & Baker, D. Macromolecular modeling with rosetta. *Annu. Rev. Biochem.* **77**, 363-382 (2008).
- 9 Xiong, P., Chen, Q. & Liu, H. Computational protein design under a given backbone structure with the ABACUS statistical energy function. *Computational Protein Design*, 217-226 (2017).
- 10 Schymkowitz, J. *et al.* The FoldX web server: an online force field. *Nucleic Acids Research* **33**, W382-W388 (2005). <https://doi.org:10.1093/nar/gki387>
- 11 Madani, A. *et al.* Large language models generate functional protein sequences across diverse families. *Nature Biotechnology* (2023). <https://doi.org:10.1038/s41587-022-01618-2>
- 12 Hie, B., Zhong Ellen, D., Berger, B. & Bryson, B. Learning the language of viral evolution and escape. *Science* **371**, 284-288 (2021). <https://doi.org:10.1126/science.abd7331>
- 13 Rao, R. M. *et al.* MSA Transformer. *Proceedings of the 38th International Conference on Machine Learning* **139**, 8844--8856 (2021).
- 14 Rives, A. *et al.* Biological structure and function emerge from scaling unsupervised learning to 250 million protein sequences. *Proceedings of the National Academy of Sciences* **118**, e2016239118 (2021). <https://doi.org:10.1073/pnas.2016239118>
- 15 Lin, Z. *et al.* Evolutionary-scale prediction of atomic-level protein structure with a language model. *Science* **379**, 1123-1130 (2023). <https://doi.org:10.1126/science.ade2574>
- 16 Li, M. *et al.* SESNet: sequence-structure feature-integrated deep learning method for data-efficient protein engineering. *Journal of Cheminformatics* **15**, 12 (2023). <https://doi.org:10.1186/s13321-023-00688-x>
- 17 Meier, J. *et al.* Language models enable zero-shot prediction of the effects of mutations on protein function. *bioRxiv*, 2021.2007.2009.450648 (2021). <https://doi.org:10.1101/2021.07.09.450648>
- 18 Hsu, C., Nisonoff, H., Fannjiang, C. & Listgarten, J. Learning protein fitness models from evolutionary and assay-labeled data. *Nature Biotechnology* **40**, 1114-1122 (2022). <https://doi.org:10.1038/s41587-021-01146-5>
- 19 Luo, Y. *et al.* ECNet is an evolutionary context-integrated deep learning framework for protein engineering. *Nature Communications* **12**, 5743 (2021). <https://doi.org:10.1038/s41467-021-25976-8>
- 20 Wu, Z., Kan, S. B. J., Lewis Russell, D., Wittmann Bruce, J. & Arnold Frances, H. Machine learning-assisted directed protein evolution with combinatorial libraries. *Proceedings of the National Academy of Sciences* **116**, 8852-8858 (2019). <https://doi.org:10.1073/pnas.1901979116>
-



- 
- 21 Engqvist, M. K. M. Correlating enzyme annotations with a large set of microbial growth temperatures reveals metabolic adaptations to growth at diverse temperatures. *BMC Microbiology* **18**, 177 (2018). <https://doi.org/10.1186/s12866-018-1320-7>
- 22 Li, G., Rabe, K. S., Nielsen, J. & Engqvist, M. K. M. Machine Learning Applied to Predicting Microorganism Growth Temperatures and Enzyme Catalytic Optima. *ACS Synthetic Biology* **8**, 1411-1420 (2019). <https://doi.org/10.1021/acssynbio.9b00099>
- 23 Devlin, J., Chang, M.-W., Lee, K. & Toutanova, K. Bert: Pre-training of deep bidirectional transformers for language understanding. *arXiv preprint arXiv:1810.04805* (2018).
- 24 Vaswani, A. *et al.* Attention is all you need. *Advances in neural information processing systems* **30** (2017).
- 25 Notin, P. *et al.* Tranception: Protein Fitness Prediction with Autoregressive Transformers and Inference-time Retrieval. *Proceedings of the 39th International Conference on Machine Learning* **162**, 16990--17017 (2022).
- 26 Yang, K. K., Fusi, N. & Lu, A. X. Convolutions are competitive with transformers for protein sequence pretraining. *bioRxiv*, 2022-2005 (2022).
- 27 Yang, K. K., Zanichelli, N. & Yeh, H. Masked inverse folding with sequence transfer for protein representation learning. *bioRxiv*, 2022.2005.2025.493516 (2022). <https://doi.org/10.1101/2022.05.25.493516>
- 28 Das, R. & Baker, D. Macromolecular modeling with rosetta. *Annual Review of Biochemistry* **77**, 363-382 (2008).
- 29 Kulandaisamy, A., Sakthivel, R. & Gromiha, M. M. MPTherm: database for membrane protein thermodynamics for understanding folding and stability. *Briefings in Bioinformatics* **22**, 2119-2125 (2021).
- 30 Stourac, J. *et al.* FireProtDB: database of manually curated protein stability data. *Nucleic Acids Research* **49**, D319-D324 (2021). <https://doi.org/10.1093/nar/gkaa981>
- 31 Nikam, R., Kulandaisamy, A., Harini, K., Sharma, D. & Gromiha, M. M. ProThermDB: thermodynamic database for proteins and mutants revisited after 15 years. *Nucleic Acids Research* **49**, D420-D424 (2021). <https://doi.org/10.1093/nar/gkaa1035>
- 32 Turner, P., Mamo, G. & Karlsson, E. N. Potential and utilization of thermophiles and thermostable enzymes in biorefining. *Microbial cell factories* **6**, 1-23 (2007).
- 33 Levin, I., Liu, M., Voigt, C. A. & Coley, C. W. Merging enzymatic and synthetic chemistry with computational synthesis planning. *Nature Communications* **13**, 7747 (2022). <https://doi.org/10.1038/s41467-022-35422-y>
- 34 Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589 (2021). <https://doi.org/10.1038/s41586-021-03819-2>
- 35 Dallago, C. *et al.* FLIP: Benchmark tasks in fitness landscape inference for proteins. *NeurIPS Datasets and Benchmarks* **11** (2021).
- 36 Jarzab, A. *et al.* Meltome atlas—thermal proteome stability across the tree of life. *Nature Methods* **17**, 495-503 (2020). <https://doi.org/10.1038/s41592-020-0801-4>
- 37 Borkotoky, S. & Murali, A. The highly efficient T7 RNA polymerase: a wonder macromolecule in biological realm. *Int J Biol Macromol* **118**, 49-56 (2018).
- 38 Dousis, A., Ravichandran, K., Hobert, E. M., Moore, M. J. & Rabideau, A. E. An engineered T7 RNA polymerase that produces mRNA free of immunostimulatory byproducts. *Nature Biotechnology* **41**, 560-568 (2023).
-

- 
- 39 An, W. & Chin, J. W. Synthesis of orthogonal transcription-translation networks. *Proceedings of the National Academy of Sciences* **106**, 8477-8482 (2009).
- 40 Leone, G., van Gemen, B., Schoen, C. D., van Schijndel, H. & Kramer, F. R. Molecular beacon probes combined with amplification by NASBA enable homogeneous, real-time detection of RNA. *Nucleic Acids Research* **26**, 2150-2155 (1998).
- 41 Cui, Z. *et al.* Novel real-time simultaneous amplification and testing method to accurately and rapidly detect Mycobacterium tuberculosis complex. *Journal of Clinical Microbiology* **50**, 646-650 (2012).
- 42 Nelson, J. *et al.* Impact of mRNA chemistry and manufacturing process on innate immune activation. *Science advances* **6**, eaaz6893 (2020).
- 43 Karikó, K., Muramatsu, H., Ludwig, J. & Weissman, D. Generating the optimal mRNA for therapy: HPLC purification eliminates immune activation and improves translation of nucleoside-modified, protein-encoding mRNA. *Nucleic acids research* **39**, e142-e142 (2011).
- 44 Wu, M. Z., Asahara, H., Tzertzinis, G. & Roy, B. Synthesis of low immunogenicity RNA with high-temperature in vitro transcription. *Rna* **26**, 345-360 (2020).
- 45 Killard, A. J. & Smyth, M. R. Creatinine biosensors: principles and designs. *Trends in biotechnology* **18**, 433-437 (2000).
- 46 Zhi, Q. *et al.* Biochemical and molecular characterization of a novel high activity creatine amidinohydrolase from *Arthrobacter nicotianae* strain 02181. *Process Biochemistry* **44**, 460-465 (2009).
- 47 Bai, X. *et al.* Improved thermostability of creatinase from *Alcaligenes Faecalis* through non-biased phylogenetic consensus-guided mutagenesis. *Microb. Cell Fact.* **19**, 1-13 (2020).
- 48 Afshari, E., Amini-Bayat, Z., Hosseinkhani, S. & Bakhtiari, N. Cloning, expression and purification of *Pseudomonas putida* ATCC12633 Creatinase. *Avicenna journal of medical biotechnology* **9**, 169 (2017).
- 49 Suzuki, K., Sagai, H., Sugiyama, M. & Imamura, S. Molecular cloning and high expression of the *Bacillus creatinase* gene in *Escherichia coli*. *Journal of fermentation and bioengineering* **76**, 77-81 (1993).
- 50 Berberich, J. A., Yang, L. W., Bahar, I. & Russell, A. J. A stable three enzyme creatinine biosensor. 2. Analysis of the impact of silver ions on creatine amidinohydrolase. *Acta Biomaterialia* **1**, 183-191 (2005).
- 51 Jiang, F. *et al.* Creatinase: Using Increased Entropy to Improve the Activity and Thermostability. *The Journal of Physical Chemistry B* **127**, 2671-2682 (2023).
- 52 Hegge, J. W., Swarts, D. C. & van der Oost, J. Prokaryotic Argonaute proteins: novel genome-editing tools? *Nature Reviews Microbiology* **16**, 5-11 (2018).
- 53 Song, J. *et al.* Highly specific enrichment of rare nucleic acid fractions using *Thermus thermophilus* argonaute with applications in cancer diagnostics. *Nucleic Acids Research* **48**, e19-e19 (2020). <https://doi.org:10.1093/nar/gkz1165>
- 54 He, R. *et al.* *Pyrococcus furiosus* Argonaute-mediated nucleic acid detection. *Chemical communications* **55**, 13219-13222 (2019).
- 55 Liu, Q. *et al.* Argonaute integrated single-tube PCR system enables supersensitive detection of rare mutations. *Nucleic acids research* **49**, e75-e75 (2021).
- 56 van Beljouw, S. P. B., Sanders, J., Rodríguez-Molina, A. & Brouns, S. J. J. RNA-targeting CRISPR–Cas systems. *Nature Reviews Microbiology* **21**, 21-34 (2023).
-

- 
- <https://doi.org/10.1038/s41579-022-00793-y>
- 57 Kropocheva, E., Kuzmenko, A., Aravin, A. A., Esyunina, D. & Kulbachinskiy, A. A programmable pAgo nuclease with universal guide and target specificity from the mesophilic bacterium *Kurthia massiliensis*. *Nucleic acids research* **49**, 4054-4065 (2021).
- 58 Tang, H., Gao, Y. & Han, J. Application Progress of the Single Domain Antibody in Medicine. *International Journal of Molecular Sciences* **24**, 4176 (2023).
- 59 Muyldermans, S. Applications of Nanobodies. *Annual Review of Animal Biosciences* **9**, 401-421 (2021). <https://doi.org/10.1146/annurev-animal-021419-083831>
- 60 Wang, J. *et al.* Heterologous antigen selection of camelid heavy chain single domain antibodies against tetrabromobisphenol A. *Anal Chem* **86**, 8296-8302 (2014). <https://doi.org/10.1021/ac5017437>
- 61 Zettl, I. *et al.* Generation of high affinity ICAM-1-specific nanobodies and evaluation of their suitability for allergy treatment. *Front Immunol* **13**, 1022418 (2022). <https://doi.org/10.3389/fimmu.2022.1022418>
- 62 Pabst, T. M. *et al.* Camelid V(H) H affinity ligands enable separation of closely related biopharmaceuticals. *Biotechnol J* **12** (2017). <https://doi.org/10.1002/biot.201600357>
- 63 Palmer, B., Angus, K., Taylor, L., Warwicker, J. & Derrick, J. P. Design of stability at extreme alkaline pH in streptococcal protein G. *J Biotechnol* **134**, 222-230 (2008). <https://doi.org/10.1016/j.jbiotec.2007.12.009>
- 64 Laughlin, T. M. & Horn, J. R. Engineering pH-Sensitive Single-Domain Antibodies. *Methods Mol Biol* **2446**, 269-298 (2022). [https://doi.org/10.1007/978-1-0716-2075-5\\_13](https://doi.org/10.1007/978-1-0716-2075-5_13)
- 65 Bult, C. J. *et al.* Complete Genome Sequence of the Methanogenic Archaeon. *Science* **273**, 1058-1073 (1996). <https://doi.org/doi:10.1126/science.273.5278.1058>
- 66 Kois, P. *et al.* Synthesis and Some Properties of Modified Oligonucleotides. II. Oligonucleotides Containing 2'-Deoxy-2'-fluoro-β-D-arabinofuranosyl Pyrimidine Nucleosides. *Nucleosides and Nucleotides* **12**, 1093-1109 (1993). <https://doi.org/10.1080/07328319308016207>
- 67 Wang, Y., Ngor, A. K., Nikoomezar, A. & Chaput, J. C. Evolution of a General RNA-Cleaving FANA Enzyme. *Nature Communications* **9** (2018). <https://doi.org/10.1038/s41467-018-07611-1>
- 68 Nikoomezar, A., Dunn, M. R. & Chaput, J. C. Evaluating the Rate and Substrate Specificity of Laboratory Evolved XNA Polymerases. *Analytical Chemistry* **89**, 12622-12625 (2017). <https://doi.org/10.1021/acs.analchem.7b03807>
- 69 Yan, S. *et al.* Direct sequencing of 2'-deoxy-2'-fluoroarabinonucleic acid (FANA) using nanopore-induced phase-shift sequencing (NIPSS). *Chemical Science* **10**, 3110-3117 (2019). <https://doi.org/10.1039/C8SC05228J>
- 70 Harshe, R. P. *et al.* Endogenous antisense RNA curbs CD39 expression in Crohn's disease. *Nat. Commun.* **11**, 5894 (2020).
- 71 Pelisch, N., Rosas Almanza, J., Stehlik, K. E., Aperi, B. V. & Kroner, A. Use of a Self-Delivering Anti-CCL3 FANA Oligonucleotide as an Innovative Approach to Target Inflammation after Spinal Cord Injury. *eNeuro* **8**, ENEURO.0338-0320.2021 (2021).
- 72 El-Khoury, R. & Damha, M. J. 2'-Fluoro-arabinonucleic Acid (FANA): A Versatile Tool for Probing Biomolecular Interactions. *Accounts of Chemical Research* **54**, 2287-2297 (2021). <https://doi.org/10.1021/acs.accounts.1c00125>
- 73 Wang, Q. *et al.* 2'-Fluoroarabinonucleic Acid Nanostructures as Stable Carriers for Cellular
-

- 
- Delivery in the Strongly Acidic Environment. *ACS Appl. Mater. Interfaces* **12**, 53592 (2020).
- 74 Pinheiro, V. B. *et al.* Synthetic Genetic Polymers Capable of Heredity and Evolution. *Science* **336**, 341-344 (2012). <https://doi.org:10.1126/science.1217622>
- 75 Lu, H. *et al.* Machine learning-aided engineering of hydrolases for PET depolymerization. *Nature* **604**, 662-667 (2022).
- 76 Cui, Y. *et al.* Computational Redesign of a PETase for Plastic Biodegradation under Ambient Condition by the GRAPE Strategy. *ACS Catalysis* **11**, 1340-1350 (2021). <https://doi.org:10.1021/acscatal.0c05126>
- 77 Sternke, M., Tripp, K. W. & Barrick, D. Consensus sequence design as a general strategy to create hyperstable, biologically active proteins. *Proceedings of the National Academy of Sciences* **116**, 11275-11284 (2019).
- 78 Arnold, F. H. Design by directed evolution. *Accounts of chemical research* **31**, 125-131 (1998).
-