Recursions and ODEs for correlations in integrable systems and random matrices.

B. Eynard^{1,2} D. Mitsios¹ S. Oukassi¹

¹ Institut de Physique Théorique de Saclay,
 F-91191 Gif-sur-Yvette Cedex, France.
 ² CRM, Centre de recherches mathématiques de Montréal,
 Université de Montréal, QC, Canada.

Abstract

An integrable system is often formulated as a flat connection, satisfying a Lax equation. It is given in terms of compatible systems having a common solution called the "wave function" Ψ living in a Lie group G, which satisfies some differential equations with rational coefficients. From this wave function, it is usual to define a sequence of "correlators" W_n , that play an important role in many applications in mathematical physics. Here, we show how to systematically obtain ordinary differential equations (ODE) with polynomial coefficients for the correlators.

An application is random matrix theory, where the wave functions are the expectation value of the characteristic polynomial, they form a family of orthogonal polynomials, and are known to satisfy an integrable system. The correlators are then the correlation functions of resolvents or of eigenvalue densities. We give the ODE and recursion on the matrix size that they satisfy.

1 Introduction

In short, we consider an invertible matrix $\Psi(x)$ that satisfies a linear differential equation with rational coefficients

$$\frac{d}{dx}\Psi(x) = \mathcal{D}(x) \ \Psi(x) \qquad \mathcal{D}(x) \in M_r(\mathbb{C}(x)).$$

where $M_r(\mathbb{C}(x))$ denotes the set of $r \times r$ matrices with entries rational functions with complex coefficients. Invertibility of $\Psi(x)$ suggests that it belongs to a representation of the Lie group $GL(r,\mathbb{C})$ for each $x \in \mathbb{C}$. Therefore, it is locally a holomorphic section of a bundle over $\mathbb{C}P^1 = \mathbb{C} \cup \{\infty\}$, whose fiber is the Lie group $G = GL(r,\mathbb{C})$, $\nabla = d - \mathcal{D}(x)dx$ is a rational connexion on this bundle, and $\Psi(x)$ is a flat section.

1.1 Differential systems and flat sections of Lie groups

More generally, we consider $\Psi(x)$ a flat section of a reductive Lie group G trivial holomorphic bundle over $\mathbb{C}P^1$, with a meromorphic connection:

$$\nabla \Psi = 0$$
 , $\nabla = d - \mathcal{D}(x)dx$ (1-1)

where $\mathcal{D}(x)$ is valued in the adjoint bundle, which implies that for each x, $\mathcal{D}(x)$ is in the Lie algebra \mathfrak{g} of the group G, and its entries are meromorphic functions of x (i.e. rational) with possibly some poles, i.e.

$$\mathcal{D}(x) \in \mathfrak{g} \otimes \mathbb{C}(x). \tag{1-2}$$

 $\Psi(x)$ being a flat section can be written in some faithful matrix representation of G, as:

$$\frac{d}{dx}\Psi(x) = \mathcal{D}(x)\Psi(x) \qquad \Psi(x) \in G. \tag{1-3}$$

Example 1.1 ($\Psi \in GL(r,\mathbb{C})$) Consider the case $G = GL(r,\mathbb{C})$, then $\Psi(x)$ is an invertible $r \times r$ matrix, and $\mathfrak{g} = \mathfrak{gl}(r,\mathbb{C}) = M_r(\mathbb{C})$, and $\mathcal{D}(x)$ is an $r \times r$ matrix, whose elements are rational functions of x (polynomial if the only pole is at ∞).

Example 1.2 (Airy differential system) Choose $G = SL(2, \mathbb{C})$ (2 × 2 matrices of determinant 1), with $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{C})$ (2 × 2 matrices of trace 0), and

$$\mathcal{D}(x) = \begin{pmatrix} 0 & 1 \\ x & 0 \end{pmatrix} \tag{1-4}$$

¹In this article we restrict ourselves to trivial G-bundles over $\mathbb{C}P^1$. All the formalism can be extended to more general cases but requires more background, and we leave it for future work. The case studied here has already a large number of practical applications.

A flat section $\Psi(x)$ is solution of

$$\nabla \Psi = 0 \quad \Leftrightarrow \quad \frac{d}{dx} \Psi(x) = \mathcal{D}(x) \Psi(x)$$
 (1-5)

it is expressed in terms of two linearly independent solutions of the Airy differential equation $(\frac{d^2}{dx^2} - x)f(x) = 0$ called the Airy and Bairy functions, denoted as Ai(x) and Bi(x) respectively.

$$\Psi(x) = \begin{pmatrix} Ai(x) & Bi(x) \\ Ai'(x) & Bi'(x) \end{pmatrix} \in G = SL(2, \mathbb{C}).$$
 (1-6)

additionally, the Wronskian of Ai(x) and Bi(x) is 1

$$\det \Psi(x) = Ai(x)Bi'(x) - Ai'(x)Bi(x) = 1.$$
 (1-7)

Example 1.3 (Hermite differential system, also called GUE) Let N > 0 be an integer. In the case where $G = GL(2, \mathbb{C})$, with $\mathfrak{g} = \mathfrak{gl}(2, \mathbb{C})$, and

$$\mathcal{D}(x) = \begin{pmatrix} x & -1\\ N & 0 \end{pmatrix} \tag{1-8}$$

The flat section $\Psi(x)$ is a solution of

$$\nabla \Psi = 0 \quad \Leftrightarrow \quad \frac{d}{dx} \Psi(x) = \mathcal{D}(x) \Psi(x) \tag{1-9}$$

given by

$$\Psi(x) = \begin{pmatrix} H_{N-1}(x) & \tilde{H}_{N-1}(x) \\ H_{N}(x) & \tilde{H}_{N}(x) \end{pmatrix}$$

$$\tag{1-10}$$

where $H_N(x)$ is the Nth Hermite polynomial, and \tilde{H}_N is the Nth Hermite function (which is not a polynomial of x). Both H_N and \tilde{H}_N are solutions of the equation:

$$f''(x) - xf'(x) + Nf(x) = 0. (1-11)$$

1.2 Adjoint Flat sections

The connection ∇ acts in the adjoint bundle by the commutator, we define $\nabla_{\text{adj}} = d - [\mathcal{D}dx, .]$, and we shall from now on denote the adjoint connection $\nabla_{\text{adj}} = \nabla$ when there is no risk of confusion. A flat section M(x) in the adjoint bundle $(M(x) \in \mathfrak{g})$ for each x, must satisfy:

$$\nabla_{\text{adj}} M = 0 \tag{1-12}$$

in other words

$$\frac{d}{dx}M(x) = [\mathcal{D}(x), M(x)]. \tag{1-13}$$

Lemma 1.1 Let $\Psi(x)$ a flat section of the group bundle. Every adjoint flat section M(x) must be of the form:

$$M(x) = \Psi(x)E\Psi(x)^{-1} = Ad_{\Psi(x)}E$$
 (1-14)

where $E \in \mathfrak{g}$ is constant dE/dx = 0.

proof: This is a classical result. Let us redo it here, in a matrix representation:

$$\frac{d}{dx}M(x) = \frac{d}{dx}\left(\Psi(x)E\Psi(x)^{-1}\right)
= \frac{d}{dx}\Psi(x) \quad E\Psi(x)^{-1} + \Psi(x) \quad E\frac{d}{dx}\Psi(x)^{-1} + \Psi(x)\frac{dE}{dx}\Psi(x)^{-1}
= \frac{d}{dx}\Psi(x) \quad E\Psi(x)^{-1} - \Psi(x) \quad E\Psi(x)^{-1}\frac{d}{dx}\Psi(x) \quad \Psi(x)^{-1} + \Psi(x)\frac{dE}{dx}\Psi(x)^{-1}
= \left[\frac{d}{dx}\Psi(x)\Psi(x)^{-1}, \Psi(x) \quad E\Psi(x)^{-1}\right] + \Psi(x)\frac{dE}{dx}\Psi(x)^{-1}
= \left[\mathcal{D}(x), \quad M(x)\right] + \Psi(x)\frac{dE}{dx}\Psi(x)^{-1}$$
(1-15)

It satisfies $dM/dx = [\mathcal{D}, M]$ if and only if dE/dx = 0. \square

Definition 1.1 (M(x.E) Adjoint flat sections) For each $E \in \mathfrak{g}$, we shall denote this section as M(x.E):

$$M(x.E) = \operatorname{Ad}_{\Psi(x)} E. \tag{1-16}$$

We can also write it:

$$M(X) \tag{1-17}$$

where $X = x.E \in \mathbb{C}P^1 \times \mathfrak{g}$, is a pair of a point and a Lie algebra element, i.e. a Lie algebra weighted point.

The intuitive way of thinking of it, is that E is a vector in the Lie algebra \mathfrak{g} , more precisely the \mathfrak{g} -fiber of the adjoint bundle at the "base point" used to define the universal cover of $\mathbb{C}P^1 \setminus \text{poles}$, and then M(x.E) is the parallel transport of E from the base point to the fiber over the point x.

1.3 Definition of Correlators

We define the following functions, as they are very useful in many applications of these frameworks, for example in random matrix theory or many integrable systems. They were first introduced in [BE09] and subsequent works, and popularized by [BDY16b; BDY16a].

For this, we choose a faithful matricial representation of the group G into $GL(r,\mathbb{C})$ (invertible $r \times r$ matrices), and \mathfrak{g} represented as $r \times r$ matrices in $\mathfrak{g}l(r,\mathbb{C})$. We consider the Trace Tr in this representation.

Definition 1.2 (Correlators) Let $\Psi(x)$ a once for all chosen flat section of the group bundle. Let $M(x.E) = \Psi(x)E\Psi(x)^{-1}$ a flat section of the adjoint bundle, parametrized by $E \in \mathfrak{g}$. We define:

$$W_1(x.E) = \operatorname{Tr} \mathcal{D}(x) M(x.E) \tag{1-18}$$

$$W_2(x_1.E_1, x_2.E_2) = \frac{1}{(x_1 - x_2)^2} \operatorname{Tr} M(x_1.E_1) M(x_2.E_2)$$
 (1-19)

and for $n \geq 3$

$$W_n(x_1.E_1, \dots, x_n.E_n) = \sum_{\sigma \in \mathfrak{S}_n^{1-cycle}} (-1)^{\sigma} \frac{\text{Tr} \prod_{i=1}^n M(x_{\sigma^i(1)}.E_{\sigma^i(1)})}{\prod_{i=1}^n (x_i - x_{\sigma(i)})}$$
(1-20)

where $\mathfrak{S}_n^{1-cycle}$ is the subset of \mathfrak{S}_n of permutations having just 1-cycle.

Example 1.4 (Correlator W_3) Recalling that $X_i = x_i.E_i$ we have

$$W_{3}(X_{1}, X_{2}, X_{3}) = \frac{\operatorname{Tr} M(X_{1})M(X_{2})M(X_{3})}{(x_{1} - x_{2})(x_{2} - x_{3})(x_{3} - x_{1})} + \frac{\operatorname{Tr} M(X_{1})M(X_{3})M(X_{2})}{(x_{1} - x_{3})(x_{3} - x_{2})(x_{2} - x_{1})}$$

$$= \frac{\operatorname{Tr} M(X_{1})[M(X_{2}), M(X_{3})]}{(x_{1} - x_{2})(x_{2} - x_{3})(x_{3} - x_{1})}$$
(1-21)

and so on for higher n.

Our goal now is to exhibit the differential equations satisfied by these W_n s.

2 ODEs for Correlators

2.1 ODE for W_1

Let us warm up with $W_1(X)$ with X = x.E. Let ' denote d/dx. We have

$$W'_{1}(X) = \operatorname{Tr} \mathcal{D}'(x)M(X) + \operatorname{Tr} \mathcal{D}(x)M'(X)$$

$$= \operatorname{Tr} \mathcal{D}'(x)M(X) + \operatorname{Tr} \mathcal{D}(x)[\mathcal{D}(x), M(X)]$$

$$= \operatorname{Tr} \mathcal{D}'(x)M(X) + \operatorname{Tr} [\mathcal{D}(x), \mathcal{D}(x)]M(X)$$

$$= \operatorname{Tr} \mathcal{D}'(x)M(X).$$
(2-1)

where the previous to last equality follows due to cyclicity of the trace.

Then

$$W_1''(X) = \operatorname{Tr} \mathcal{D}''(x)M(X) + \operatorname{Tr} \mathcal{D}'(x)M'(X)$$

$$= \operatorname{Tr} \mathcal{D}''(x)M(X) + \operatorname{Tr} \mathcal{D}'(x)[\mathcal{D}(x), M(X)]$$

$$= \operatorname{Tr} \mathcal{D}''(x)M(X) + \operatorname{Tr} [\mathcal{D}'(x), \mathcal{D}(x)]M(X)$$

$$= \operatorname{Tr} (\mathcal{D}''(x) + [\mathcal{D}'(x), \mathcal{D}(x)])M(X).$$
(2-2)

We can repeat this and get

$$\frac{d^k}{dx^k}W_1(X) = \operatorname{Tr} \mathcal{D}_k(x)M(X)$$
(2-3)

with the matrix $\mathcal{D}_k(x)$ computed as follows

$$\mathcal{D}_0 = \mathcal{D}$$
 , $\mathcal{D}_1 = \mathcal{D}'$, $\mathcal{D}_2 = \mathcal{D}'' + [\mathcal{D}', \mathcal{D}]$, $\mathcal{D}_3 = \mathcal{D}''' + 2[\mathcal{D}'', \mathcal{D}] + [[\mathcal{D}', \mathcal{D}], \mathcal{D}]$ (2-4)

and recursively

$$\mathcal{D}_{k+1} = \mathcal{D}_k' + [\mathcal{D}_k, \mathcal{D}]. \tag{2-5}$$

Now notice that each $\mathcal{D}_k(x)$ is a rational function of x valued in the Lie algebra \mathfrak{g} , that is a finite dimensional vector space of dimension dim \mathfrak{g} . Therefore, at most dim \mathfrak{g} of them can be linearly independent over the field $\mathbb{C}(x)$ which implies that there must exist some rational coefficients $\alpha_k(x)$, such that

$$\sum_{k=0}^{\dim \mathfrak{g}} \alpha_k(x) \mathcal{D}_k(x) = 0. \tag{2-6}$$

And upon multiplying by the common denominator, we can assume that $\alpha_k(x) \in \mathbb{C}[x]$ are polynomial. Therefore,

Theorem 2.1 $W_1(x)$ satisfies a linear differential equation of order dim \mathfrak{g} , with polynomial coefficients $\alpha_k(x)$:

$$\sum_{k=0}^{\dim \mathfrak{g}} \alpha_k(x) \frac{d^k}{dx^k} W_1(x.E) = 0.$$
 (2-7)

The coefficients $\alpha_k(x)$ are determined by

$$\sum_{k=0}^{\dim \mathfrak{g}} \alpha_k(x) \mathcal{D}_k(x) = 0 \quad \text{where} \quad \mathcal{D}_0 = \mathcal{D} \quad , \quad \mathcal{D}_{k+1} = \mathcal{D}'_k + [\mathcal{D}_k, \mathcal{D}]. \tag{2-8}$$

Notice that the matrices $\mathcal{D}_k(x)$ are independent of E, and so are the $\alpha_k(x)$. This implies that for any choice of $E \in \mathfrak{g}$ we have a solution of this differential equation. Therefore,

Theorem 2.2 The space of solutions of

$$\left(\sum_{k=0}^{\dim \mathfrak{g}} \alpha_k(x) \frac{d^k}{dx^k}\right) w(x) = 0 \tag{2-9}$$

is the vector space of dimension $\dim \mathfrak{g}$ generated by

$$W_1(x.E) = \operatorname{Tr} \mathcal{D}(x)M(x.E) \quad \text{for } E \in \mathfrak{g}.$$
 (2-10)

Example 2.1 (Airy) We have $\mathfrak{g} = \mathfrak{sl}(2)$ which has dimension $\dim \mathfrak{sl}(2) = 3$:

$$\mathcal{D}_0(x) = \begin{pmatrix} 0 & 1 \\ x & 0 \end{pmatrix} , \quad \mathcal{D}_1(x) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} , \quad \mathcal{D}_2(x) = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} , \quad \mathcal{D}_3(x) = \begin{pmatrix} 0 & -2 \\ 2x & 0 \end{pmatrix}$$
(2-11)

They satisfy the linear relation

$$\frac{1}{2}\mathcal{D}_3 = 2x\mathcal{D}_1 - \mathcal{D}_0,\tag{2-12}$$

by the use of Theorem 2.1

$$\frac{1}{2}W_1''' = 2xW_1' - W_1. (2-13)$$

The 3 linearly independent solutions to this equation are:

$$W_1(x.E) \tag{2-14}$$

with E any linear combination of the 3 Chevaley-basis² vectors:

$$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \quad , \quad \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \quad , \quad \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{pmatrix}. \tag{2-15}$$

Example 2.2 (Hermite/GUE) We have $\mathfrak{g} = \mathfrak{gl}(2)$ with dimension $\dim \mathfrak{gl}(2) = 4$:

$$\mathcal{D}_0(x) = \begin{pmatrix} x & -1 \\ N & 0 \end{pmatrix} , \quad \mathcal{D}_1(x) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} , \quad \mathcal{D}_2(x) = \begin{pmatrix} 0 & -1 \\ -N & 0 \end{pmatrix}$$
 (2-16)

$$\mathcal{D}_3(x) = \begin{pmatrix} -2N & x \\ -Nx & 2N \end{pmatrix} , \quad \mathcal{D}_4(x) = \begin{pmatrix} 0 & 4N - x^2 + 1 \\ 4N^2 - Nx^2 - N & 0 \end{pmatrix}. \tag{2-17}$$

The linear relation is

$$\mathcal{D}_4 = (x^2 - 4N)\mathcal{D}_2 + x\mathcal{D}_1 - \mathcal{D} \tag{2-18}$$

using Theorem 2.1

$$W_1'''' = (x^2 - 4N)W_1'' + xW_1' - W_1. (2-19)$$

We shall come back to this equation and its consequences in Example 5.7.

 $^{^2{\}rm In}$ fact any basis of ${\mathfrak g}$ would be good. The Chevaley basis is a cannonical basis defined for any semi-simple Lie algebra.

2.2 ODE for W_2

Let $X_1 = x_1.E_1$ and $X_2 = x_2.E_2$. We have defined

$$W_2(X_1, X_2) = \operatorname{Tr} \frac{M(X_2)}{(x_1 - x_2)^2} M(X_1).$$
 (2-20)

Taking the derivative with respect to x_1 we get

$$\frac{d}{dx_1}W_2(X_1, X_2) = \text{Tr}\left(\frac{[M(X_2), \mathcal{D}(x_1)]}{(x_1 - x_2)^2} - \frac{2M(X_2)}{(x_1 - x_2)^3}\right) M(X_1). \tag{2-21}$$

and in general

$$\frac{d^k}{dx_1^k} W_2(X_1, X_2) = \operatorname{Tr} \mathcal{D}_k(x_1; X_2) \ M(X_1)$$
 (2-22)

where

$$\mathcal{D}_0(x_1; X_2) = \frac{M(X_2)}{(x_1 - x_2)^2} \quad , \quad \mathcal{D}_{k+1}(x_1; X_2) = \frac{d}{dx_1} \mathcal{D}_k(x_1; X_2) + [\mathcal{D}_k(x_1; X_2), \mathcal{D}(x_1)].$$
(2-23)

Remark that for each k, $\mathcal{D}_k(x_1; X_2)$ is a rational function of x_1 , but it may be a transcendantal function of x_2 (in the Airy example it involves $Ai(x_2)$). Again, no more than dim \mathfrak{g} of them can be linearly independent, and therefore there exist some coefficients $\alpha_k(x_1; X_2)$ polynomial in x_1 , such that

$$\sum_{k=0}^{\dim \mathfrak{g}} \alpha_k(x_1; X_2) \mathcal{D}_k(x_1; X_2) = 0.$$
 (2-24)

This implies,

Theorem 2.3 $W_2(X_1, X_2)$ satisfies an x_1 -differential equation of order dim \mathfrak{g} , with coefficients $\alpha_k(x_1; X_2)$ polynomials of x_1 :

$$\sum_{k=0}^{\dim \mathfrak{g}} \alpha_k(x_1; X_2) \frac{d^k}{dx_1^k} W_2(X_1, X_2) = 0.$$
 (2-25)

Example 2.3 (Airy) We find the following expression for $\alpha_k s$ in the case of Airy:

$$\alpha_{0}(x_{1}, x_{2}) = -6x_{1}^{4} \text{Ai}^{4}(x_{2}) + 4x_{1}^{3}x_{2} \text{Ai}^{4}(x_{2}) + 16x_{1}^{3} \text{Ai}^{2}(x_{2}) \text{Ai}^{'2}(x_{2})$$

$$+ 2x_{1}^{2}x_{2}^{2} \text{Ai}^{4}(x_{2}) - 16x_{1}^{2}x_{2} \text{Ai}^{2}(x_{2}) \text{Ai}^{'2}(x_{2}) - 2x_{1}^{2} \text{Ai}^{3}(x_{2}) \text{Ai}^{'}(x_{2})$$

$$- 10x_{1}^{2} \text{Ai}^{'4}(x_{2}) - 4x_{1}x_{2} \text{Ai}^{3}(x_{2}) \text{Ai}^{'}(x_{2}) + 12x_{1}x_{2} \text{Ai}^{'4}(x_{2})$$

$$+ 8x_{1} \text{Ai}(x_{2}) \text{Ai}^{'3}(x_{2}) + 6x_{2}^{2} \text{Ai}^{3}(x_{2}) \text{Ai}^{'}(x_{2}) - 2x_{2}^{2} \text{Ai}^{'4}(x_{2})$$

$$- 6x_{2} \text{Ai}^{4}(x_{2}) - 8x_{2} \text{Ai}(x_{2}) \text{Ai}^{'3}(x_{2}) + 6 \text{Ai}^{2}(x_{2}) \text{Ai}^{'2}(x_{2})$$

$$- 6x_{2} \text{Ai}^{4}(x_{2}) + 8x_{1}^{4}x_{2} \text{Ai}^{4}(x_{2}) + 8x_{1}^{4} \text{Ai}^{2}(x_{2}) \text{Ai}^{'2}(x_{2})$$

$$- 4x_{1}^{5} \text{Ai}^{4}(x_{2}) + 8x_{1}^{4}x_{2} \text{Ai}^{4}(x_{2}) + 8x_{1}^{4} \text{Ai}^{2}(x_{2}) \text{Ai}^{'2}(x_{2})$$

$$- 4x_{1}^{3}x_{2}^{2} \text{Ai}^{4}(x_{2}) + 16x_{1}^{3}x_{2} \text{Ai}^{2}(x_{2}) \text{Ai}^{'2}(x_{2}) + 4x_{1}^{3} \text{Ai}^{3}(x_{2}) \text{Ai}^{'}(x_{2})$$

$$- 4x_{1}^{3}x_{2}^{2} \text{Ai}^{4}(x_{2}) + 8x_{1}^{2}x_{2}^{2} \text{Ai}^{2}(x_{2}) \text{Ai}^{'2}(x_{2}) + 8x_{1}^{2}x_{2} \text{Ai}^{3}(x_{2}) \text{Ai}^{'}(x_{2})$$

$$+ 8x_{1}^{2}x_{2} \text{Ai}^{4}(x_{2}) + 8x_{1}^{2}x_{2}^{2} \text{Ai}^{2}(x_{2}) \text{Ai}^{'2}(x_{2}) + 4x_{1}^{2}x_{2} \text{Ai}^{3}(x_{2}) \text{Ai}^{'}(x_{2})$$

$$- 4x_{1}x_{2}^{2} \text{Ai}^{3}(x_{2}) \text{Ai}^{'}(x_{2}) - 3x_{1}^{2} \text{Ai}^{4}(x_{2}) + 4x_{1}^{2}x_{2} \text{Ai}^{2}(x_{2}) \text{Ai}^{'3}(x_{2})$$

$$- 8x_{1}x_{2} \text{Ai}^{2}(x_{2}) \text{Ai}^{'3}(x_{2}) + 3x_{2}^{2} \text{Ai}^{4}(x_{2}) + 4x_{2}^{2} \text{Ai}^{2}(x_{2}) \text{Ai}^{'3}(x_{2})$$

$$- 12x_{2} \text{Ai}^{2}(x_{2}) \text{Ai}^{'2}(x_{2}) + 6x_{1} \text{Ai}^{'4}(x_{2})$$

$$+ 6x_{1}x_{2} \text{Ai}^{2}(x_{2}) \text{Ai}^{'2}(x_{2}) + 6x_{1} \text{Ai}^{'4}(x_{2})$$

$$+ 6x_{1}x_{2} \text{Ai}^{2}(x_{2}) \text{Ai}^{'2}(x_{2}) + 6x_{1} \text{Ai}^{'4}(x_{2}) + 3x_{2}^{2} \text{Ai}^{2}(x_{2}) \text{Ai}^{'2}(x_{2})$$

$$- 6x_{2} \text{Ai}^{'4}(x_{2})$$

$$- 6x_{2} \text{Ai}^{'4}(x_{2})$$

$$- 2x_{1}^{4} \text{Ai}^{4}(x_{2}) - 2x_{1}^{3}x_{2} \text{Ai}^{4}(x_{2}) - 2x_{1}^{3} \text{Ai}^{2}(x_{2}) \text{Ai}^{'2}(x_{2})$$

$$- 2x_{1}^{4} \text{Ai}^{2}(x_{2}) + 4x_{1}^{2}$$

2.3 ODE for W_2 with polynomial coefficients

In fact, we can also find an x_1 -ODE for $W_2(X_1, X_2)$, whose coefficients are polynomial functions of both x_1 and x_2 .

We begin by writing $\operatorname{Tr} M(X_1)M(X_2)$ as a quadratic form of the coefficients of $M(X_1)$ and $M(X_2)$. Let us warm up with the example $\mathfrak{g} = \mathfrak{g}l(r,\mathbb{C})$, where $M(X_1)$ and $M(X_2)$ are $r \times r$ matrices:

$$W_2(X_1, X_2) = \sum_{i,j,i',j'=1}^r M_{i',j'}(X_2) Q_{0;i',j',i,j}(x_1, x_2) M_{i,j}(X_1)$$
 (2-27)

with $Q_0(x_1, x_2)$ the $r^2 \times r^2$ matrix

$$Q_{0;i',j',i,j}(x_1,x_2) = \frac{\delta_{i,j'}\delta_{i',j}}{(x_1 - x_2)^2}.$$
 (2-28)

Hence, we get,

$$\frac{d^k}{dx_1^k} W_2(x_1, x_2) = \sum_{i,j,i',j'=1}^r M_{i',j'}(X_2) Q_{k;i',j',i,j}(x_1, x_2) M_{i,j}(X_1)$$
 (2-29)

with the recursion relation

$$Q_{k+1;i',j',i,j}(x_1, x_2) = \frac{d}{dx_1} Q_{k;i',j',i,j}(x_1, x_2) + \sum_{l} (Q_{k;i',j',l,j}(x_1, x_2)D(x_1)_{l,i} - Q_{k;i',j',i,l}(x_1, x_2)D(x_1)_{j,l})$$
(2-30)

For each k, Q_k is an $r^2 \times r^2$ matrix, and there are at most r^4 such matrices that can be linearly independent. Therefore, there exist some polynomial coefficients $\tilde{\alpha}_k(x_1, x_2)$, such that

$$\sum_{k=0}^{r^4} \tilde{\alpha}_k(x_1, x_2) Q_k(x_1, x_2) = 0.$$
 (2-31)

This implies

$$\sum_{k=0}^{r^4} \tilde{\alpha}_k(x_1, x_2) \frac{d^k}{dx_1^k} W_2(X_1, X_2) = 0.$$
 (2-32)

General case: Let $Q_k \in \mathfrak{g}^* \otimes \mathfrak{g}^*$ the rational quadratic form such that

$$\frac{d^k}{dx_1^k} W_2(X_1, X_2) = Q_k(M(X_1), M(X_2))$$
(2-33)

We have the recursion

$$Q_{k+1} = \frac{d}{dx_1} Q_k + [Q_k, \mathcal{D}(x_1)]_1$$
 (2-34)

where the commutator acts only on the 1st factor of $\mathfrak{g}^* \otimes \mathfrak{g}^*$ and in terms of components is given in eq. 2-30. The space $\mathfrak{g}^* \otimes \mathfrak{g}^*$ has dimension $(\dim \mathfrak{g})^2$ and thus there must exist some polynomial coefficients $\tilde{\alpha}_k(x_1, x_2) \in \mathbb{C}[x_1, x_2]$, such that

$$\sum_{k=0}^{(\dim \mathfrak{g})^2} \tilde{\alpha}_k(x_1, x_2) Q_k(x_1, x_2) = 0.$$
 (2-35)

This implies,

Theorem 2.4 W_2 satisfies a polynomial ODE of order $(\dim \mathfrak{g})^2$. There exist some coefficients $\tilde{\alpha}_k(x_1, x_2) \in \mathbb{C}[x_1, x_2]$, such that

$$\sum_{k=0}^{(\dim \mathfrak{g})^2} \tilde{\alpha}_k(x_1, x_2) \frac{d^k}{dx_1^k} W_2(X_1, X_2) = 0.$$
 (2-36)

2.4 ODEs for higher correlators

Let $X_i = x_i \cdot E_i$ n points in $\mathbb{C}P^1 \times \mathfrak{g}$.

We write,

$$W_n(X_1, \dots, X_n) = \text{Tr } D_{0,n}(x_1; X_2, \dots, X_n) M(X_1)$$
 (2-37)

where

$$D_{0,n}(x_1; X_2, \dots, X_n) = \sum_{\sigma \in \mathfrak{S}_{-\text{cycle}}^{1-\text{cycle}}} (-1)^{\sigma} \frac{\text{Tr} \prod_{i=1}^{n-1} M(X_{\sigma^i(1)})}{\prod_{i=1}^n (x_i - x_{\sigma(i)})}$$
(2-38)

We then have

$$\frac{d^k}{dx_1^k} W_n(X_1, \dots, X_n) = \text{Tr } D_{k,n}(x_1; X_2, \dots, X_n) M(X_1)$$
(2-39)

where $D_{k,n}(x_1; X_2, \dots, X_n)$ is obtained by the recursion

$$D_{k+1,n}(x_1; X_2, \dots, X_n) = \frac{d}{dx_1} D_{k,n}(x_1; X_2, \dots, X_n) + [D_{k,n}(x_1; X_2, \dots, X_n), \mathcal{D}(x_1)].$$
(2-40)

Each $D_{k,n}(x_1; X_2, ..., X_n)$ is rational in x_1 , but transcendental in $x_2, ..., x_n$. At most dim \mathfrak{g} of them can be linearly independent, hence there exist some coefficients $\alpha_{k,n}(x_1; X_2, ..., X_n)$ polynomial in x_1 , such that

$$\sum_{k=0}^{\dim \mathfrak{g}} \alpha_{k,n}(x_1; X_2, \dots, X_n) D_{k,n}(x_1; X_2, \dots, X_n) = 0.$$
 (2-41)

This implies

Theorem 2.5 There exist some coefficients $\alpha_{k,n}(x_1; X_2, \ldots, X_n)$, polynomial in x_1 , such that

$$\sum_{k=0}^{\dim \mathfrak{g}} \alpha_{k,n}(x_1; X_2, \dots, X_n) \frac{d^k}{dx_1^k} W_n(X_1, \dots, X_n) = 0.$$
 (2-42)

The coefficients $\alpha_{k,n}$ are found by eq. 2-40 and eq. 2-41.

2.5 ODE for higher correlators with polynomial coefficients

Moreover, there is a tensor $Q_{0,n} \in (\mathfrak{g}^*)^{\otimes n}$, such that

$$W_n(X_1, \dots, X_n) = Q_{0,n}(M(X_1), M(X_2), \dots, M(X_n))$$
(2-43)

whose coefficients are rational functions of x_1, \ldots, x_n . For $\mathfrak{gl}(r, \mathbb{C})$, the expression of $Q_{0,n}$ can be written as:

$$Q_{0,n} = \sum_{\sigma \in \mathfrak{S}_n^{1-cycle}} (-1)^{\sigma} \sum_{l_1,\dots,l_n} \frac{e_{l_1 l_{\sigma(1)}}^1 \otimes \dots \otimes e_{l_n l_{\sigma(n)}}^n}{\prod_i (x_i - x_{\sigma(i)})}$$
(2-44)

with $e_{i,j} \in \mathfrak{gl}(r,\mathbb{C})^*$ and satisfy $e_{i,j}(M) = M_{i,j}$ for any $M \in \mathfrak{gl}(r,\mathbb{C})$. Moreover, we get,

$$\frac{d^k}{dx_1^k} W_n(X_1, \dots, X_n) = Q_{k,n}(M(X_1), M(X_2), \dots, M(X_n))$$
(2-45)

where

$$Q_{k+1,n} = \frac{d}{dx_1} Q_{k,n} + [Q_{k,n}, \mathcal{D}(x_1)]_1.$$
 (2-46)

The commutator³

acts only on the 1st factor of $(\mathfrak{g}^*)^{\otimes n}$. The space $(\mathfrak{g}^*)^{\otimes n}$ has dimension $(\dim \mathfrak{g})^n$, and so at most $(\dim \mathfrak{g})^n$ such tensors can be linearly independent. Consequently, there exist some polynomial coefficients $\tilde{\alpha}_{k,n}(x_1,\ldots,x_n) \in \mathbb{C}[x_1,\ldots,x_n]$, such that

$$\sum_{k=0}^{\dim \mathfrak{g}^n} \tilde{\alpha}_{k,n}(x_1, \dots, x_n) \ Q_{k,n}(x_1; x_2, \dots, x_n) = 0.$$
 (2-47)

This implies

Theorem 2.6 There exist some coefficients $\tilde{\alpha}_{k,n}(x_1; x_2, \dots, x_n) \in \mathbb{C}[x_1, \dots, x_n]$, such that

$$\sum_{k=0}^{\dim \mathfrak{g}^n} \tilde{\alpha}_{k,n}(x_1,\dots,x_n) \frac{d^k}{dx_1^k} W_n(X_1,X_2,\dots,X_n) = 0.$$
 (2-48)

The coefficients $\tilde{\alpha}_{k,n}$ are found by eq. 2-46 and eq. 2-47. The $(\dim \mathfrak{g})^n$ linearly independent solutions are obtained by choosing $(E_1, E_2, \ldots, E_n) \in \mathfrak{g}^{\otimes n}$.

3 Differential equations with respect to other parameters

Consider a compatible **family** of such systems, parametrized by times $\mathbf{t} = (t_1, t_2, \dots)$, i.e. we have a time-dependent family of sections $\Psi(\mathbf{t}; x)$ of a Lie group G principal bundle over $\mathbb{C}P^1$ with rational connections:

$$\frac{d}{dx}\Psi(\mathbf{t};x) = \mathcal{D}(\mathbf{t};x)\Psi(\mathbf{t};x), \tag{3-1}$$

$$\frac{d}{dt_k}\Psi(\mathbf{t};x) = R_k(\mathbf{t};x)\Psi(\mathbf{t};x), \tag{3-2}$$

where each $R_k(\mathbf{t};x)$ is rational in x (but can be a transcendantal function of \mathbf{t}).

$$[Q_{k,n}, \mathcal{D}(x_1)]_{1;i',j',i,j} = \sum_{l} (Q_{k;i',j',l,j}(x_1, x_2)D(x_1)_{l,i} - Q_{k;i',j',i,l}(x_1, x_2)D(x_1)_{j,l})$$

³in terms of components,

Compatibility of these equations, is equivalent to saying that there exists (locally) a common solution which, itself, is equivalent to the Lax equations:

$$\frac{d}{dt_k}\mathcal{D}(\mathbf{t};x) = [R_k(\mathbf{t};x), \mathcal{D}(\mathbf{t};x)] + \frac{d}{dx}R_k(\mathbf{t};x)$$
(3-3)

$$\frac{d}{dt_k}R_l(\mathbf{t};x) = [R_k(\mathbf{t};x), R_l(\mathbf{t};x)] + \frac{d}{dt_l}R_k(\mathbf{t};x). \tag{3-4}$$

The adjoint flat sections are defined as in Definition 1.1 by replacing $\Psi(x) \to \Psi(\mathbf{t}; x)$

$$M(\mathbf{t}; x.E) = \Psi(\mathbf{t}; x)E\Psi(\mathbf{t}; x)^{-1}$$
(3-5)

with $E \in \mathfrak{g}$, independent of x and t.

They satisfy the adjoint ODE

$$\frac{d}{dt_j}M(\mathbf{t};x.E) = [R_j(\mathbf{t};x), M(\mathbf{t};x.E)]. \tag{3-6}$$

3.1 Time Differential equation for W_1

Let us choose a time t_j . For each $k \geq 0$ we can write

$$\frac{d^k}{dt_i^k} W_1(x.E) = \operatorname{Tr} R_{j,k}(\mathbf{t}; x) \ M(\mathbf{t}; x.E)$$
(3-7)

where the matrix $R_{j,k}(\mathbf{t};x)$ is defined by the following recursion:

initial term:
$$R_{j,0}(\mathbf{t};x) = \mathcal{D}(\mathbf{t};x)$$
recursion:
$$R_{j,k+1}(\mathbf{t};x) = \frac{d}{dt_j} R_{j,k}(\mathbf{t};x) + [R_{j,k}(\mathbf{t};x), R_j(\mathbf{t};x)]$$
(3-8)

Again at most dim $\mathfrak g$ of them are independent, there must exist some coefficients polynomial in x such that

$$\sum_{k=0}^{\dim \mathfrak{g}} \alpha_{j,k,1}(\mathbf{t};x) R_{j,k}(\mathbf{t};x) = 0.$$
(3-9)

This implies

Theorem 3.1 (Time differential equation for W_1) there exist coefficients $\alpha_{j,k,1}(\mathbf{t};x)$ polynomial in x, such that

$$\sum_{k=0}^{\dim \mathfrak{g}} \alpha_{j,k,1}(\mathbf{t};x) \frac{d^k}{dt_j^k} W_1(\mathbf{t};x.E) = 0.$$
 (3-10)

The coefficients $\alpha_{j,k,1}$ are determined by eq. 3-8 and eq. 3-9.

3.2 Time Differential equation for W_n , $n \ge 2$

Results of subsection 3.1 directly generalise to the case of n-point correlators for $n \geq 2$

Theorem 3.2 there exist coefficients $\alpha_{j,k,n}(\mathbf{t}; x_1; X_2, \dots, X_n)$ polynomial in x_1 , such that

$$\sum_{k=0}^{\dim \mathfrak{g}} \alpha_{j,k,n}(\mathbf{t}; x_1; X_2, \dots, X_n) \frac{d^k}{dt_j^k} W_n(\mathbf{t}; X_1, X_2, \dots, X_n) = 0.$$
 (3-11)

where $\alpha_{j,k,n}$ are determined by

Initial term:
$$R_{j,0,n}(\mathbf{t}; x_1, X_2, \dots, X_n) = D_{0,n}(\mathbf{t}; x_1, X_2, \dots, X_n).$$

Recursion: $R_{j,k+1,n}(\mathbf{t}; x_1, X_2, \dots, X_n) = \frac{d}{dt_j} R_{j,k,n}(\mathbf{t}; x_1, X_2, \dots, X_n) + [R_{j,k,n}(\mathbf{t}; x_1, X_2, \dots, X_n), R_j(\mathbf{t}; x_1)].$

Linear relation: $\sum_{k=0}^{\dim \mathfrak{g}} \alpha_{j,k,n}(\mathbf{t}; x_1, X_2, \dots, X_n) R_{j,k,n}(\mathbf{t}; x_1, X_2, \dots, X_n) = 0.$
(3-12)

Theorem 3.3 there exist coefficients $\tilde{\alpha}_{j,k,n}(\mathbf{t};x_1;x_2,\ldots,x_n) \in \mathbb{C}[x_1,\ldots,x_n]$, such that

$$\sum_{k=0}^{\dim \mathfrak{g}^n} \tilde{\alpha}_{j,k,n}(\mathbf{t}; x_1; x_2, \dots, x_n) \frac{d^k}{dt_j^k} W_n(\mathbf{t}; x_1.E_1, x_2.E_2, \dots, x_n.E_n) = 0.$$
 (3-13)

where $\tilde{\alpha}_{i,k,n}$ are determined by

Initial term:
$$\tilde{Q}_{j,0,n}(\mathbf{t}; x_1, \dots, x_n) = Q_{0,n}(\mathbf{t}; x_1, \dots, x_n).$$

$$Recursion: \qquad \tilde{Q}_{j,k+1,n}(\mathbf{t}; x_1, \dots, x_n) = \frac{d}{dt_j} \tilde{Q}_{j,k,n}(\mathbf{t}; x_1, \dots, x_n)$$

$$+ \sum_{i=1}^n [\tilde{Q}_{j,k,n}(\mathbf{t}; x_1, \dots, x_n), R_j(\mathbf{t}; x_i)]_i. \qquad (3-14)$$

$$Linear relation: \qquad \sum_{k=0}^{\dim \mathfrak{g}^n} \tilde{\alpha}_{j,k,n}(\mathbf{t}; x_1, x_2, \dots, x_n) \tilde{Q}_{j,k,n}(\mathbf{t}; x_1, \dots, x_n) = 0.$$

4 Recursion relations

Now consider a compatible **sequence** of such systems indexed by an integer N, i.e. we have a section $\Psi^{(N)}(x)$ of a Lie group G principal bundle over $\mathbb{C}P^1$ with rational connections:

$$\frac{d}{dx}\Psi^{(N)}(x) = \mathcal{D}^{(N)}(x)\Psi^{(N)}(x), \tag{4-1}$$

that satisfies a recursion

$$\Psi^{(N+1)}(x) = \mathcal{R}^{(N)}(x)\Psi^{(N)}(x), \tag{4-2}$$

where each $R^{(N)}(x)$ is rational in x (but can be any function of N).

In analogy to section 3, compatibility of all these equations, is equivalent to saying that there exists (locally) a common solution, and this is equivalent to the discrete Lax equations:

Proposition 4.1 (Compatibility) The ODE and recursion are compatible:

$$\mathcal{D}^{(N+1)}(x)\mathcal{R}^{(N)}(x) - \mathcal{R}^{(N)}(x)\mathcal{D}^{(N)}(x) = \frac{d}{dx}\mathcal{R}^{(N)}(x)$$
(4-3)

We write it as

$$\mathcal{D}^{(N+1)}(x) = \mathcal{R}^{(N)}(x)\mathcal{D}^{(N)}(x)\mathcal{R}^{(N)}(x)^{-1} + \frac{d}{dx}\mathcal{R}^{(N)}(x)\mathcal{R}^{(N)}(x)^{-1}$$

$$= \mathcal{R}^{(N)}(x)\left(\mathcal{D}^{(N)}(x) + \mathcal{R}^{(N)}(x)^{-1}\frac{d}{dx}\mathcal{R}^{(N)}(x)\right)\mathcal{R}^{(N)}(x)^{-1}$$

$$= \mathcal{R}^{(N)}(x)\mathcal{D}^{(N,1)}(x)\mathcal{R}^{(N)}(x)^{-1}.$$
(4-4)

where we defined

$$\mathcal{D}^{(N,1)}(x) = \mathcal{D}^{(N)}(x) + \mathcal{R}^{(N)}(x)^{-1} \frac{d}{dx} \mathcal{R}^{(N)}(x). \tag{4-5}$$

We also define adjoint flat sections

$$M^{(N)}(x.E) = \Psi^{(N)}(x)E\Psi^{(N)}(x)^{-1} \qquad , \qquad E = \begin{pmatrix} 1 & 0\\ 0 & 0 \end{pmatrix}$$
 (4-6)

Notice that

Tr
$$M^{(N)}(x) = 1$$
 , det $M^{(N)}(x) = 0$, $(M^{(N)}(x))^2 = M^{(N)}(x)$. (4-7)

Proposition 4.2 (Recursion and ODE for the adjoint section)

$$M^{(N+1)}(x) = \mathcal{R}^{(N)}(x)M^{(N)}(x)\mathcal{R}^{(N)}(x)^{-1}.$$
(4-8)

$$\frac{d}{dx}M^{(N)}(x) = [\mathcal{D}^{(N)}(x), M^{(N)}(x)]. \tag{4-9}$$

4.1 Recursion for $W_1^{(N)}$

We have,

$$W_1^{(N)}(x.E) = \text{Tr}\,\mathcal{D}^{(N)}(x)M^{(N)}(x.E) \tag{4-10}$$

and thus

$$W_{1}^{(N+1)}(x.E) = \operatorname{Tr} \mathcal{D}^{(N+1)}(x) M^{(N+1)}(x.E)$$

$$= \operatorname{Tr} \mathcal{D}^{(N+1)}(x) \mathcal{R}^{(N)}(x) M^{(N)}(x.E) R^{(N)}(x)^{-1}$$

$$= \operatorname{Tr} \mathcal{R}^{(N)}(x)^{-1} \mathcal{D}^{(N+1)}(x) \mathcal{R}^{(N)}(x) M^{(N)}(x.E)$$

$$= \operatorname{Tr} \left(\mathcal{D}^{(N)}(x) + \mathcal{R}^{(N)}(x)^{-1} \mathcal{R}'^{(N)}(x) \right) M^{(N)}(x.E)$$

$$(4-11)$$

Notice that, $W_1^{(N+1)}(x.E) - W_1^{(N)}(x.E) = \operatorname{Tr} \mathcal{R}^{(N)}(x)^{-1} \mathcal{R}'^{(N)}(x) M^{(N)}(x.E)$.

Then, in general we shall have

Lemma 4.1 $(W_1^{(N+k)}(x.E))$

$$W_1^{(N+k)}(x.E) = \text{Tr } D_k^{(N)}(x) \ M^{(N)}(x.E)$$
(4-12)

with $D_0^{(N)}(x) = \mathcal{D}^{(N)}(x)$ and by recursion

$$D_k^{(N)}(x) = \mathcal{R}^{(N)}(x)^{-1}\mathcal{R}^{(N+1)}(x)^{-1}\dots\mathcal{R}^{(N+k-1)}(x)^{-1} \mathcal{D}^{(N+k)}(x) \mathcal{R}^{(N+k-1)}(x)\dots\mathcal{R}^{(N)}(x)$$
(4-13)

is rational in x.

At most dim \mathfrak{g} of these matrices can be independent, and therefore, there exist some coefficients $\alpha_k^{(N)}(x) \in \mathbb{C}[x]$, such that:

$$\sum_{k=0}^{\dim \mathfrak{g}} \alpha_k^{(N)}(x) D_k^{(N)}(x) = 0. \tag{4-14}$$

This implies

Theorem 4.1 (Recursion equation for W_1) $W_1(x.E)$ satisfies a linear recursion relation of order dim \mathfrak{g} , with polynomial coefficients $\alpha_k^{(N)}(x)$, independent of E:

$$\sum_{k=0}^{\dim \mathfrak{g}} \alpha_k^{(N)}(x) W_1^{(N+k)}(x) = 0. \tag{4-15}$$

4.2 Recursion for $W_n^{(N)}$, $n \ge 2$

The rest proceeds in the same way as for differential equations:

Theorem 4.2 (Recursion equation for $W_n^{(N)}$) $W_n^{(N)}(x_1.E_1,...,x_n.E_n)$ satisfies a linear recursion relation of order $(\dim \mathfrak{g})^n$, with coefficients $\tilde{\alpha}_k^{(N)}(x_1,...,x_n) \in \mathbb{C}[x_1,...,x_n]$, independent of $E_1,...,E_n$:

$$\sum_{k=0}^{(\dim \mathfrak{g})^n} \tilde{\alpha}_{k,n}^{(N)}(x_1,\dots,x_n) W_n^{(N+k)}(x_1.E_1,\dots,x_n.E_n) = 0.$$
 (4-16)

where $\tilde{\alpha}_{k,n}^{(N)}$ are determined by:

Initial term:
$$\tilde{Q}_{0,n}^{(N)}(x_1,\ldots,x_n) = Q_{0,n}(x_1,\ldots,x_n).$$

$$\tilde{Q}_{k+1,n}^{(N)}(x_1,\ldots,x_n) = \left(\left(\operatorname{Ad}_{R^{(N+k)}(x_1)}\otimes\cdots\otimes\operatorname{Ad}_{R^{(N+k)}(x_n)}\right)\tilde{Q}_{k,n}^{(N)}\right)(x_1,\ldots,x_n).$$

$$\operatorname{Cdim}\mathfrak{g}^{(N)}(x_1,\ldots,x_n)\tilde{Q}_{k,n}^{(N)}(x_1,\ldots,x_n) = 0.$$

$$\operatorname{Linear relation}: \sum_{k=0}^{\operatorname{ddim}\mathfrak{g}^{(N)}}\tilde{\alpha}_{k,n}^{(N)}(x_1,\ldots,x_n)\tilde{Q}_{k,n}^{(N)}(x_1,\ldots,x_n) = 0.$$

Notice that the initial term $\tilde{Q}_{0,n}^{(N)}(x_1,\ldots,x_n)=Q_{0,n}(x_1,\ldots,x_n)$ is actually independent of N.

proof: By definition $W_n^{(N+k)}(x_1,\ldots,x_n) = \tilde{Q}_{k,n}^{(N)}(M(x_1.E_1),\ldots,M(x_n.E_n))$. Therefore, proving the recursion and linear relation will conclude the proof.

The tensor $Q_{0,n}$ is given by the application:

$$Q_{0,n} \colon \mathfrak{g}^{\otimes n} \to \mathbb{C}$$

$$(M_1, ..., M_n) \mapsto \sum_{\sigma \in \mathfrak{S}_n^{1-\text{cycle}}} (-1)^{\sigma} \frac{\text{Tr} \prod_{i=1}^n M_{\sigma^i(1)}}{\prod_{i=1}^n (x_i - x_{\sigma(i)})}$$

Due to Definition 1.2 and Proposition 4.2 we get,

$$\begin{split} W_n^{(N+1)} &= Q_{0,n}(M^{(N+1)}(X_1), \cdots, M^{(N+1)}(X_n)) \\ &= Q_{0,n}(\operatorname{Ad}_{R^{(N)}(x_1)} M^{(N)}(X_1), \cdots, \operatorname{Ad}_{R^{(N)}(x_n)} M^{(N)}(X_n)) \\ &=: \left(\left(\operatorname{Ad}_{R^{(N)}(x_1)} \otimes \cdots \otimes \operatorname{Ad}_{R^{(N)}(x_n)} \right) Q_{0,n} \right) (M^{(N)}(X_1), \cdots, M^{(N)}(X_n)) \end{split}$$

By induction, we conclude that,

$$\tilde{Q}_{k+1,n}^{(N)}(x_1,\dots,x_n) = \left(\left(\operatorname{Ad}_{R^{(N+k)}(x_1)} \otimes \dots \otimes \operatorname{Ad}_{R^{(N+k)}(x_n)} \right) \tilde{Q}_{k,n}^{(N)} \right) (x_1,\dots,x_n). \quad (4-18)$$

Finally, since $Q_{k,n}^{(N)} \in (\mathfrak{g}^*)^{\otimes n}$ at most $(\dim \mathfrak{g})^n$ of them can be linearly independent. Therefore, the exist coefficients $\tilde{\alpha}_{k,n}^{(N)} \in \mathbb{C}[x_1,\ldots,x_n]$ such that

$$\sum_{k=0}^{(\dim \mathfrak{g})^n} \tilde{\alpha}_{k,n}^{(N)}(x_1,\dots,x_n)\tilde{Q}_{k,n}^{(N)}(x_1,\dots,x_n) = 0.$$
 (4-19)

Remark, that in the case of $\mathfrak{g} = \mathfrak{g}l(r,\mathbb{C})$ the presented proof for the recursion could be formulated using index notation, eq. 2-44 reads

$$Q_{0,n} = \sum_{\sigma \in \mathfrak{S}_n^{1-cycle}} (-1)^{\sigma} \sum_{l_1,\dots,l_n} \frac{e_{l_1 l_{\sigma(1)}}^1 \otimes \dots \otimes e_{l_n l_{\sigma(n)}}^n}{\prod_i (x_i - x_{\sigma(i)})}$$

Moreover, with the use of Definition 1.2 and Proposition 4.2

$$W_n^{(N+1)} = \sum_{\sigma \in \mathfrak{S}_n^{1-\text{cycle}}} (-1)^{\sigma} \frac{\text{Tr} \prod_{i=1}^n M(x_{\sigma^i(1)}.E_{\sigma^i(1)})}{\prod_{i=1}^n (x_i - x_{\sigma(i)})}$$

$$= \sum_{\sigma \in \mathfrak{S}_n^{1-\text{cycle}}} (-1)^{\sigma} \frac{\text{Tr} \left(\prod_i R^{(N)}(x_{\sigma^i(1)}) M^{(N)}(x_{\sigma^i(1)}) R^{(N)}(x_{\sigma^i(1)})^{-1}\right)}{\prod_i (x_i - x_{\sigma(i)})}$$

$$= \left(Q_{0,n}|_{e_{l_i,l_{\sigma(i)}}^i \to \text{Ad}_{R^{(N)}(x_i)}} e_{l_i,l_{\sigma(i)}}^i\right) (M^{(N)}(x_1), \dots, M^{(N)}(x_N))$$

which leads to

$$\tilde{Q}_{1,n}^{(N)} = \left(\left(\operatorname{Ad}_{R^{(N)}(x_1)} \otimes \cdots \otimes \operatorname{Ad}_{R^{(N)}(x_n)} \right) Q_{0,n} \right) (x_1, \dots, x_n).$$

Finally, $\tilde{Q}_{k,n}^{(N)}$ is obtained by induction.

5 Application: 1-Matrix model

Expectation values of spectral densities of random matrices, typically fit in the framwork developed here. So let us see how it applies to random matrices to get differential equations and recursions.

5.1 Introduction

Usually, random matrix theory is defined with a probability law on a set of Hermitian (resp. Normal) matrices with a measure continuous with respect to the Lebesgue measure.

However, since most of the time the random variables of interest, concern the eigenvalues, we may also directly define random matrix theory as its marginal law of eigenvalues.

The eigenvalues induced measure (possibly complex measure) takes the following form for the diagonal matrix Λ of eigenvalues:

$$\Lambda = \operatorname{diag}(\lambda_1, \lambda_2, \dots, \lambda_N) \in \gamma^N$$
(5-1)

$$d\mu(\Lambda) = \frac{1}{N!Z} \Delta(\Lambda)^2 e^{-\text{Tr} V(\Lambda)} d\Lambda$$
 (5-2)

where the function V is called the potential, $d\Lambda$ is the product measure $d\Lambda = d\lambda_1 \dots d\lambda_N$ on γ^N , and the Vandermonde $\Delta(\Lambda)$ is defined as

$$\Delta(\Lambda) = \prod_{i < j} (\lambda_i - \lambda_j), \tag{5-3}$$

and the partition function Z is

$$Z = \frac{1}{N!} \int_{\gamma^N} \Delta(\Lambda)^2 e^{-\text{Tr} V(\Lambda)} d\lambda_1 \dots d\lambda_N.$$
 (5-4)

The integration domain γ is the space to which the λ_i s belong. It needs not be the real axis, it may be any space on which this measure can be defined, it could be for example a Polish subspace of \mathbb{R} , but it can also be a Jordan arc in the complex plane, as long as Z is an absolutely convergent integral.

Example 5.1 (Gaussian Unitary Ensemble (GUE)) The Gaussian Unitary Ensemble is the case $\gamma = \mathbb{R}$ and $V(x) = \frac{1}{2}x^2$. It is, thus, a probability law on \mathbb{R}^N :

$$d\mu(\Lambda) = \frac{1}{Z^{(N)}} \Delta(\Lambda)^2 e^{-\frac{1}{2} \text{Tr } \Lambda^2} d\Lambda$$
 (5-5)

 $Z^{(N)}$ is known to be proportional to the Barnes function

$$Z^{(N)} = (2\pi)^{\frac{N}{2}} \prod_{k=0}^{N-1} k!.$$
 (5-6)

5.2 Potentials and Jordan arcs

It is customary to consider potentials to be polynomials or rational, and to choose γ a Jordan arc such that the measure vanishes at the boundaries of γ (the boundaries may be at ∞).

5.2.1 Polynomial case

Let $V(x) \in \mathbb{C}[x]$ a polynomial with deg $V \geq 1$ called the potential. Notice that $e^{-V(x)}$ can vanish only at ∞ .

An integral

$$\int_{\gamma} f(x)e^{-V(x)}dx\tag{5-7}$$

where $f(x) \in \mathbb{C}[x]$, is invariant under homotopic deformations of the smooth Jordan arc γ , and it can be absolutely convergent only if γ goes to ∞ in directions in which $\Re V(x) \to +\infty$.

There are exactly deg V angular sectors around ∞ in which $\Re V(x) \to +\infty$, and thus there are deg $V-1=\deg V'$ independent possible Jordan arcs (up to addition by concatenation), that go from an allowed sector to another one. We shall say that the homology space of "admissible" paths, is of dimension

$$\dim(\text{Homology space of admissible paths}) = \deg V'.$$
 (5-8)

If V(x) is an even degree real polynomial whose leading coefficient is > 0, then in that case, $\gamma = \mathbb{R}$ is an admissible path.

For example if $V(x) = \frac{1}{2}x^2$, then $\deg V' = 1$ and thus $\gamma = \mathbb{R}$ is the only (up to homotopy and multiplication by a non-zero scalar) admissible path.

Otherwise admissible paths are often complex paths.

5.2.2 Rational case

We repeat the same considerations if $V'(x) \in \mathbb{C}(x)$ is a rational function. This means that V(x) may have logarithmic singularities. e^{-V} may vanish only at the poles of V'.

An integral

$$\int_{\gamma} f(x)e^{-V(x)}dx\tag{5-9}$$

where $f(x) \in \mathbb{C}[x]$, is invariant under homotopic deformations of the smooth Jordan arc γ , and it can be convergent only if γ approaches singularities of V in directions in which $\Re V(x) \to +\infty$.

It is explained in [BE06b] how to construct the homology space of all admissible paths, and its dimension is found to be

$$\dim = \deg V' = \text{sum of degrees of all poles, including the pole at } \infty.$$
 (5-10)

5.3 Definitions

5.3.1 Wave functions and orthogonal polynomials

Definition 5.1 (Wave functions) Let

$$p^{(N)}(x) = \int_{\gamma^N} d\mu(\Lambda) \, \det(x - \Lambda) \tag{5-11}$$

It is a polynomial of x of degree N (it is the expectation value of the characteristic polynomial).

Proposition 5.1 (Orthogonal polynomials) The polynomials $p^{(N)}$ form a family of orthogonal polynomials for the measure $e^{-V(x)}dx$ on γ :

$$\int_{\gamma} dx \ p^{(N)}(x) \ p^{(N')}(x)e^{-V(x)} = h^{(N)} \ \delta_{N,N'}. \tag{5-12}$$

proof: It is a well known result [Hei78; Sze39; Meh04] □

Definition 5.2 (Normalized wave functions) We thus define

$$h^{(N)} = \int_{\gamma} dx \ p^{(N)}(x)p^{(N)}(x)e^{-V(x)} \qquad , \quad \gamma_N = \sqrt{\frac{h^{(N)}}{h^{(N-1)}}}$$
 (5-13)

$$\psi^{(N)}(x) = \frac{e^{-\frac{1}{2}V(x)}}{\sqrt{h^{(N)}}} p^{(N)}(x)$$
 (5-14)

They form an orthonormal $L_2(\gamma)$ family:

$$\int_{\gamma} dx \ \psi^{(N)}(x)\psi^{(N')}(x) = \delta_{N,N'}. \tag{5-15}$$

Definition 5.3 (Dual wave functions) We also define

$$\phi^{(N)}(x) = \frac{e^{\frac{1}{2}V(x)}}{\sqrt{h^{(N)}}} \int_{\gamma} dx' e^{-V(x')} \frac{p^{(N)}(x')}{x - x'}$$
(5-16)

Definition 5.4 (Matrix of wave functions)

$$\Psi^{(N)}(x) = \begin{pmatrix} \psi^{(N-1)}(x) & \phi^{(N-1)}(x) \\ \psi^{(N)}(x) & \phi^{(N)}(x) \end{pmatrix}$$
 (5-17)

Example 5.2 (Gaussian (GUE))

$$p^{(N)}(x) = H_N(x) = Hermite\ polynomial$$
 (5-18)

$$\int_{\mathbb{R}} H_N(x) H_M(x) e^{-\frac{1}{2}x^2} dx = \sqrt{2\pi} N! \ \delta_{N,M}$$
 (5-19)

$$h^{(N)} = \sqrt{2\pi}N!$$
 , $\gamma^{(N)} = \sqrt{N}$. (5-20)

5.3.2 Resolvents

Definition 5.5 (Resolvents) Let

$$W_1^{(N)}(x) = \int_{\gamma^N} d\mu(\Lambda) \operatorname{Tr} \frac{1}{x - \Lambda}$$
 (5-21)

and in general

$$\hat{W}_n^{(N)}(x_1, \dots, x_n) = \frac{\delta_{n,2}}{(x_1 - x_2)^2} + \int_{\gamma^N} d\mu(\Lambda) \prod_{i=1}^n \text{Tr} \frac{1}{x_i - \Lambda}$$
 (5-22)

Let W_n their cumulants, defined by inverting the formula:

$$\hat{W}_{n}^{(N)}(x_{1},\ldots,x_{n}) = \sum_{\mu \vdash \{x_{1},\ldots,x_{n}\}} \prod_{i=1}^{\ell(\mu)} W_{|\mu_{i}|}^{(N)}(\mu_{i})$$
 (5-23)

i.e. the $\hat{W}_n^{(N)}$ are the sums over all partitions of the n variables, of products of cumulants of each part.

The system of equations (5-23) that define the cumulants is a triangular system easy to invert, it gives for example $W_1^{(N)} = \hat{W}_1^{(N)}$, and the n = 2 cumulant is the covariance of resolvents:

$$W_2^{(N)}(x_1, x_2) = \hat{W}_2^{(N)}(x_1, x_2) - W_1^{(N)}(x_1)W_1^{(N)}(x_2).$$
 (5-24)

Similarly

$$W_{3}^{(N)}(x_{1}, x_{2}, x_{3}) = \hat{W}_{3}^{(N)}(x_{1}, x_{2}, x_{3}) - W_{1}^{(N)}(x_{1})\hat{W}_{2}^{(N)}(x_{2}, x_{3}) - W_{1}^{(N)}(x_{2})\hat{W}_{2}^{(N)}(x_{1}, x_{3}) - W_{1}^{(N)}(x_{3})\hat{W}_{2}^{(N)}(x_{1}, x_{2}) + 2W_{1}^{(N)}(x_{1})W_{1}^{(N)}(x_{2})W_{1}^{(N)}(x_{3}).$$
 (5-25)

5.3.3 Spectral densities and correlations

Definition 5.6 (Spectral density and correlations) Let $x \in \gamma$, define the "expected spectral density"

$$\rho_1^{(N)}(x) = \int_{\gamma^N} d\mu(\Lambda) \sum_{i=1}^N \delta(x - \lambda_i)$$
 (5-26)

In other words:

 $\frac{1}{N}\rho_1^{(N)}(x)$ is the probability that x is an eigenvalue of Λ .

and in general for $x_1, \ldots, x_n \in \gamma^n$, define

$$\hat{\rho}_n^{(N)}(x_1,\dots,x_n) = \int_{\gamma^N} d\mu(\Lambda) \prod_{i=1}^n \left(\sum_{j=1}^N \delta(x_i - \lambda_j) \right)$$
 (5-27)

Let ρ_n their cumulants, defined by the formula:

$$\hat{\rho}_n^{(N)}(x_1, \dots, x_n) = \sum_{\mu \vdash \{x_1, \dots, x_n\}} \prod_{i=1}^{\ell(\mu)} \rho_{|\mu_i|}^{(N)}(\mu_i)$$
 (5-28)

i.e. the $\hat{\rho}_n^{(N)}$ are the sums over all partitions of the n variables of products of cumulants of each part.

In other words: $\frac{1}{N^n}\hat{\rho}_n^{(N)}(x_1,\ldots,x_n)$ is the expectation that x_1,\ldots,x_n are simultaneously eigenvalues, it is interpretted as the correlation function of n eigenvalues.

The cumulants $\frac{1}{N^n}\rho_n^{(N)}(x_1,\ldots,x_n)$ are called the connected correlation function of n eigenvalues, they generalize the covariance for $n \geq 3$.

Resolvents and spectral correlations are closely related.

• Resolvents are "Stieltjes transforms" of spectral correlations:

$$W_1^{(N)}(x) = \int_{x' \in \gamma} \frac{1}{x - x'} \,\rho_1^{(N)}(x') dx' \tag{5-29}$$

$$W_n^{(N)}(x_1, \dots, x_n) = \int_{\gamma^n} \prod_{i=1}^n \frac{1}{x_i - x_i'} \, \rho_n^{(N)}(x_1', \dots, x_n') dx_1' \dots dx_n'.$$
 (5-30)

• Spectral correlations are discontinuities of resolvents. Notice that the resolvents are ill-defined whenever some x_i s belong to the Jordan arc γ . In fact they have a discontinuity across γ , and this discontinuity is proportional to the spectral density:

$$\rho_1^{(N)}(x) = \frac{1}{2\pi i} \left(W_1^{(N)}(x_{\text{left}}) - W_1^{(N)}(x_{\text{right}}) \right) = \frac{1}{2\pi i} \operatorname{Disc} W_1^{(N)}(x).$$
 (5-31)

$$\rho_n^{(N)}(x_1, \dots, x_n) = \frac{1}{(2\pi i)^n} \operatorname{Disc}_{x_1} \dots \operatorname{Disc}_{x_n} W_n^{(N)}(x_1, \dots, x_n).$$
 (5-32)

Corollary 5.1 (Same linear ODE and recursions) As a consequence of eq. 5-32, if W_n satisfies a linear relation (ODE, PDE or recursion) with rational coefficients (in the variables x, x_1, \ldots, x_n), then the spectral correlation ρ_n satisfy the same linear relation.

5.3.4 Determinantal formulae

The following relation is well known [BE09; EKR18]

Proposition 5.2 (Determinantal formulae) Let

$$E = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} , \quad M^{(N)}(x) = \Psi^{(N)}(x)E\Psi^{(N)}(x)^{-1}.$$
 (5-33)

We have

$$W_1^{(N)}(x) = \text{Tr}\,\Psi^{(N)'}(x)E\Psi^{(N)}(x)^{-1} \tag{5-34}$$

$$W_2^{(N)}(x_1, x_2) = \frac{1}{(x_1 - x_2)^2} \operatorname{Tr} M^{(N)}(x_1) M^{(N)}(x_2)$$
 (5-35)

and in general we have

$$W_n^{(N)}(x_1, \dots, x_n) = W_n(x_1.E, \dots, x_n.E)$$
(5-36)

where the right hand side is eq. 1-20 of Definition 1.2.

Remark 5.1 In fact, the initial motivation of [BE09] for introducing all this formalism, was that it applied to random matrices. So this proposition was actually not a consequence, but the origin.

5.4 Recursion and ODE for the wave functions

5.4.1 3-terms recursion relations

The following is well known [Sze39; Meh04]

Proposition 5.3 (Jacobi 3-term recursion relation) The orthogonal polynomials satisfy a 3 terms recursion relation

$$xp^{(N)}(x) = p^{(N+1)}(x) + S_N p^{(N)}(x) + \frac{h^{(N)}}{h^{(N-1)}} p^{(N-1)}(x)$$
 (5-37)

which gives

$$x\psi^{(N)}(x) = \gamma^{(N+1)}\psi^{(N+1)}(x) + S_N\psi^{(N)}(x) + \gamma^{(N)}\psi^{(N-1)}(x).$$
 (5-38)

We write it:

$$x\psi^{(N)}(x) = \sum_{m \in \mathbb{N}} Q_{N,m}\psi^{(m)}(x)$$
 (5-39)

where Q is an infinite matrix, with only 3 bands:

$$Q_{n,m} = S_n \delta_{n,m} + \gamma^{(n)} \delta_{n,m+1} + \gamma^{(m)} \delta_{n+1,m}.$$
 (5-40)

It is symmetric, $Q_{n,m} = Q_{m,n}$ i.e.

$$Q = Q^T. (5-41)$$

It implies a recursion for the matrix Ψ :

$$\Psi^{(N+1)}(x) = \mathcal{R}^{(N)}(x)\Psi^{(N)}(x) \tag{5-42}$$

where

$$\mathcal{R}^{(N)}(x) = \begin{pmatrix} 0 & 1\\ \frac{-\gamma^{(N)}}{\gamma^{(N+1)}} & \frac{x-S_N}{\gamma^{(N+1)}} \end{pmatrix}$$
 (5-43)

is called the shift operator, or sometimes the "ladder" operator.

 $\mathcal{R}^{(N)}(x)$ is a matrix polynomial of x of degree 1. Its inverse is also polynomial of degree 1:

$$\mathcal{R}^{(N)}(x)^{-1} = \begin{pmatrix} \frac{x - S_N}{\gamma^{(N)}} & -\frac{\gamma^{(N+1)}}{\gamma^{(N)}} \\ 1 & 0 \end{pmatrix}$$
 (5-44)

Example 5.3 (GUE) Hermite polynomials satisfy

$$p^{(N+1)}(x) = xp^{(N)}(x) - Np^{(N-1)}(x)$$
(5-45)

$$\implies \gamma^{(N+1)}\psi^{(N+1)}(x) = x\psi^{(N)}(x) - \gamma^{(N)}\psi^{(N-1)}(x) \tag{5-46}$$

Definition 5.7 (Iterated shift operator) Let

$$\mathcal{R}^{(N,k)}(x) := \mathcal{R}^{(N+k-1)}(x)\mathcal{R}^{(N+k-2)}(x)\cdots\mathcal{R}^{(N)}(x)$$
 (5-47)

which is a matrix, polynomial of x of degree at most k.

It is such that

$$\Psi^{(N+k)}(x) = \mathcal{R}^{(N,k)}(x)\Psi^{(N)}(x). \tag{5-48}$$

5.4.2 ODE for the wave function

Proposition 5.4 (ODE for Ψ) If $V'(x) \in \mathbb{C}(x)$ is a rational function, then the orthogonal polynomials satisfy an ODE with rational coefficients, which in matrix form can be written

$$\frac{d}{dx}\Psi^{(N)}(x) = \mathcal{D}^{(N)}(x)\Psi^{(N)}(x)$$
 (5-49)

where

$$\mathcal{D}^{(N)}(x) = \begin{pmatrix} \frac{V'(x)}{2} & 0\\ 0 & -\frac{V'(x)}{2} \end{pmatrix} + \begin{pmatrix} w_{N-1,N-1}(x) & w_{N-1,N}(x)\\ w_{N,N-1}(x) & w_{N,N}(x) \end{pmatrix} \begin{pmatrix} 0 & -\gamma^{(N)}\\ \gamma^{(N)} & 0 \end{pmatrix}$$
(5-50)

where

$$w_{N,N'}(x) = \frac{1}{\sqrt{h^{(N)}h^{(N')}}} \int_{\gamma} e^{-V(x')} dx' \ p^{(N)}(x') \frac{V'(x) - V'(x')}{x - x'} p^{(N')}(x')$$

$$= \left(\frac{V'(x) - V'(Q)}{x - Q}\right)_{N,N'}$$
(5-51)

with the matrix Q defined in eq. 5-40.

We have

$$\operatorname{Tr} \mathcal{D}^{(N)}(x) = 0. \tag{5-52}$$

proof: See [BE06b; EKR18]. \square

Example 5.4 (GUE) Hermite polynomials satisfy

$$\frac{d}{dx}p^{(N)}(x) = Np^{(N-1)}(x) = xp^{(N)}(x) - p^{(N+1)}(x)$$
(5-53)

$$\frac{d}{dx}p^{(N+1)}(x) = (N+1)p^{(N)}(x) \tag{5-54}$$

therefore

$$\frac{d}{dx} \begin{pmatrix} p^{(N-1)}(x) \\ p^{(N)}(x) \end{pmatrix} = \begin{pmatrix} x & -1 \\ N & 0 \end{pmatrix} \begin{pmatrix} p^{(N-1)}(x) \\ p^{(N)}(x) \end{pmatrix}$$
 (5-55)

which implies

$$\frac{d}{dx}\Psi^{(N)}(x) = \mathcal{D}^{(N)}(x)\Psi^{(N)}(x)$$
 (5-56)

$$\mathcal{D}^{(N)}(x) = \begin{pmatrix} \frac{1}{2}x & -\gamma^{(N)} \\ \gamma^{(N)} & -\frac{1}{2}x \end{pmatrix}$$
 (5-57)

Notice that

$$\operatorname{Tr} \mathcal{D}^{(N)}(x) = 0 \tag{5-58}$$

$$\det\left(y - \mathcal{D}^{(N)}(x)\right) = y^2 - \frac{1}{4}x^2 + (\gamma^{(N)})^2 = y^2 - \frac{1}{4}x^2 + N. \tag{5-59}$$

The eigenvalues of $\mathcal{D}^{(N)}(x)$ are $y = \pm \frac{1}{2}\sqrt{x^2 - 4N}$.

Proposition 5.5 (Some properties of the shift operator)

$$\mathcal{R}^{(N)}(x) = \mathcal{R}^{(N)}(Id + x\mathcal{R}^{(N)'}) \tag{5-60}$$

$$\mathcal{R}^{(N)}(x)^{-1} = (Id - x\mathcal{R}^{(N)'})(\mathcal{R}^{(N)})^{-1}$$
(5-61)

with $\mathcal{R}^{(N)} = \mathcal{R}^{(N)}(0)$, and $(\mathcal{R}^{(N)'})^2 = 0$ nilpotent.

$$\mathcal{R}^{(N)'} = \begin{pmatrix} 0 & \frac{-1}{\gamma_N} \\ 0 & 0 \end{pmatrix} \tag{5-62}$$

proof: Both properties follow from direct computation. \Box This implies

$$\mathcal{R}^{(N)}(x_2)\mathcal{R}^{(N)}(x_1)^{-1} = \mathcal{R}^{(N)}(\mathrm{Id} + (x_2 - x_1)\mathcal{R}^{(N)'})(\mathcal{R}^{(N)})^{-1}$$
(5-63)

Proposition 5.6 (Recursion and ODE for the adjoint section)

$$M^{(N+1)}(x) = \mathcal{R}^{(N)}(x)M^{(N)}(x)\mathcal{R}^{(N)}(x)^{-1}.$$
 (5-64)

$$\frac{d}{dx}M^{(N)}(x) = [\mathcal{D}^{(N)}(x), M^{(N)}(x)]. \tag{5-65}$$

5.5 Recursion and ODE for $W_1^{(N)}(x)$

We will now merely apply the formalism of section 4 and section 2.

Notice that we have

$$W_1^{(N+k)}(x) = \text{Tr}\,\mathcal{D}_k^{(N)}(x)M^{(N)}(x)$$
(5-66)

where

$$\mathcal{D}_k^{(N)}(x) := \mathcal{R}^{(N,k)}(x)^{-1} \mathcal{D}^{(N+k)}(x) \mathcal{R}^{(N,k)}(x)$$
(5-67)

Theorem 5.1 (Recursion) At most 3 of the matrices $\mathcal{D}_0^{(N)}(x), \ldots, \mathcal{D}_3^{(N)}(x)$ can be linearly independent (use eq. 5-52), therefore there exist some coefficients $C^{(N,k)}(x)$, polynomials of x, such that

$$\sum_{k=0}^{3} C^{(N,k)}(x) \mathcal{D}_k^{(N)}(x) = 0$$
 (5-68)

This implies

$$\sum_{k=0}^{3} C^{(N,k)}(x)W_1^{(N+k)}(x) = 0 (5-69)$$

i.e. $W_1^{(N)}(x)$ satisfies a linear recursion relation of order 3, with coefficients polynomials of x.

Example 5.5 (Recursion of $W_1^{(N)}$ in GUE) In the case of GUE $W_1^{(N)}$ satisfies eq. 5-69 with

$$C^{(N,0)} = -n - 1$$
, $C^{(N,1)} = x^2 - n - 3$, $C^{(N,2)} = -x^2 + n$, $C^{(N,3)} = n + 2$. (5-70)

This gives

$$(N+2)W_1^{(N+3)}(x)-(x^2-N)W_1^{(N+2)}(x)+(x^2-N-3)W_1^{(N+1)}(x)-(N+1)W_1^{(N)}(x)=0. \eqno(5-71)$$

Theorem 5.2 (ODE) Let

$$\hat{\mathcal{D}}^{(N,0)}(x) = \mathcal{D}^{(N)}(x), \qquad \hat{\mathcal{D}}^{(N,k+1)}(x) = \frac{d}{dx}\hat{\mathcal{D}}^{(N,k)}(x) + [\hat{\mathcal{D}}^{(N,k)}(x), \mathcal{D}^{(N)}(x)]$$
(5-72)

At most 3 of the matrices $\hat{\mathcal{D}}^{(N,0)}(x), \ldots, \hat{\mathcal{D}}^{(N,3)}(x)$ can be linearly independent, therefore there exist some coefficients $\hat{C}^{(N,k)}(x)$, polynomials of x, such that

$$\sum_{k=0}^{3} \hat{C}^{(N,k)}(x)\hat{\mathcal{D}}^{(N,k)}(x) = 0$$
 (5-73)

This implies

$$\sum_{k=0}^{3} \hat{C}^{(N,k)}(x) \frac{d^k}{dx^k} W_1^{(N)}(x) = 0$$
 (5-74)

 $W_{\rm l}^{(N)}$ satisfies an ODE of order 3 with polynomial coefficients.

Example 5.6 (GUE) $W_1^{(N)}(x)$ satisfies an ODE of order 3:

$$\left(x + (4N - x^2)\frac{d}{dx} + \frac{d^3}{dx^3}\right)W_1^{(N)}(x) = 0$$
(5-75)

Example 5.7 (Consequence for moments of GUE) Let the moments of traces of powers:

$$t_k^{(N)} = \int_{\gamma^N} d\mu(\Lambda) \operatorname{Tr} \Lambda^k = \int_{\gamma} x^k \rho_1^{(N)}(x) dx.$$
 (5-76)

We have $t_0^{(N)} = N$, and it is clear that for the Gaussian measure, which is symmetric under $\Lambda \to -\Lambda$, all the odd moments vanish $t_{2k+1}^{(N)} = 0$. The resolvent can be written

$$W_1^{(N)}(x) = -\frac{x}{2} + \sum_{k>0} t_k^{(N)} x^{-k-1}$$
 (5-77)

The ODE (5-75) implies a recursion relation for the even moments t_{2k} .

$$t_{2k}^{(N)} = 2N \frac{2k-1}{k+1} t_{2k-2}^{(N)} + \frac{1}{2} \frac{(2k-3)(2k-2)(2k-1)}{k+1} t_{2k-4}^{(N)}, \quad k \ge 2$$
 (5-78)

with initial terms $t_0^{(N)} = N, t_2^{(N)} = N^2$.

The recursion (5-71) implies:

$$t_{k+2}^{(N+2)} - t_{k+2}^{(N+1)} = (N+2)t_k^{(N+3)} + Nt_k^{(N+2)} - (N+3)t_k^{(N+1)} - (N+1)t_k^{(N)}.$$
 (5-79)

5.6 Recursion and ODE for $W_2^{\left(N ight)}$

We have

$$W_2^{(N+k)}(x_1, x_2) = \frac{1}{(x_1 - x_2)^2} \text{Tr} \quad M^{(N)}(x_1) \mathcal{R}^{(N,k)}(x_1)^{-1} \mathcal{R}^{(N,k)}(x_2)$$

$$M^{(N)}(x_2) \mathcal{R}^{(N,k)}(x_2)^{-1} \mathcal{R}^{(N,k)}(x_1)$$
(5-80)

At most 4 of the 2×2 matrices $\mathcal{R}^{(N,k)}(x_1)^{-1}\mathcal{R}^{(N,k)}(x_2)M^{(N)}(x_2)\mathcal{R}^{(N,k)}(x_2)^{-1}\mathcal{R}^{(N,k)}(x_1)$ for k=0,1,2,3,4 can be linearly independent. By Definition 5.7 $\mathcal{R}^{(N,k)}(x)$ and its inverse are polynomial in x. Therefore,

Proposition 5.7 There exist some coefficients $C_2^{(N,k)}(x_1;x_2)$, polynomials of x_1 , such that

$$\sum_{k=0}^{4} \alpha_2^{(N,k)}(x_1, x_2) \mathcal{R}^{(N,k)}(x_1)^{-1} \mathcal{R}^{(N,k)}(x_2) M^{(N)}(x_2) \mathcal{R}^{(N,k)}(x_2)^{-1} \mathcal{R}^{(N,k)}(x_1) = 0 \quad (5-81)$$

i.e. $W_2^{(N)}(x_1, x_2)$ satisfies a linear recursion relation of order 3, with coefficients polynomials of x_1 (possibly transcendental of x_2).

Moreover, we get

Proposition 5.8 There exist some coefficients $\alpha_2^{(N,k)}(x_1;x_2) \in \mathbb{C}[x_1,x_2]$, such that

$$\sum_{k=0}^{16} \alpha_2^{(N,k)}(x_1, x_2) W_2^{(N+k)}(x_1, x_2) = 0.$$
 (5-82)

proof: We apply Theorem 4.2 for $\mathfrak{g} = \mathfrak{gl}(2,\mathbb{C})$

5.7 Recursion and ODE for $W_n^{(N)}, n \geq 3$

We have

$$W_n^{(N+k)}(x_1, \dots, x_n) = \sum_{\sigma \in \mathfrak{S}_n^{1-cycle}} (-1)^{\sigma} \frac{\operatorname{Tr} \prod_{i=1}^n \mathcal{R}^{(N,k)}(x_{\sigma^i(1)}) M^{(N)}(x_{\sigma^i(1)}) \mathcal{R}^{(N,k)}(x_{\sigma^i(1)})^{-1}}{\prod_{i=1}^n (x_i - x_{\sigma(i)})}$$
(5-83)

We can write it:

$$W_n^{(N)}(x_1, \dots, x_n) = \text{Tr } M^{(N)}(x_1) H_n^{(N)}(x_1; x_2, \dots, x_n)$$
 (5-84)

where $H_n^{(N)}(x_1; x_2, \dots, x_n)$ is a 2×2 matrix, rational function of x_1 .

Theorem 5.3 At most 4 of the 2 × 2 matrices $\mathcal{R}^{(N,k)}(x_1)^{-1}H_n^{(N+k)}(x_1;x_2,\ldots,x_n)\mathcal{R}^{(N,k)}(x_1)$ for $k=0,1,2,\ldots,4$ can be linearly independent, therefore there exist some coefficients $C_n^{(N,k)}(x_1;x_2,\ldots,x_n)$, polynomials of x_1 , such that

$$\sum_{k=0}^{4} C_n^{(N,k)}(x_1; x_2, \dots, x_n) W_n^{(N+k)}(x_1, x_2, \dots, x_n) = 0$$
 (5-85)

Following Theorem 4.2, we can also write:

$$W_n^{(N+k)}(x_1, \dots, x_n) = \mathcal{H}_n^{(N+k)}(M^{(N)}(x_1), M^{(N)}(x_2), \dots, M^{(N)}(x_n))$$
 (5-86)

where $\mathcal{H}_n^{(N)}(x_1; x_2, \ldots, x_n)$ is a linear form on $\mathfrak{gl}(2, \mathbb{C})^{\otimes n}$, it belongs to the dual $\mathfrak{gl}(2, \mathbb{C})^{\otimes n^*}$, which has dimension 4^n , therefore at most 4^n of $\mathcal{H}_n^{(N+k)}(x_1; x_2, \ldots, x_n)$ for $k = 0, 1, \ldots, 4^n$ can be independent.

This implies

Theorem 5.4 There exist some coefficients $\tilde{C}_n^{(N,k)}(x_1, x_2, \dots, x_n) \in \mathbb{C}[x_1, x_2, \dots, x_n]$, such that

$$\sum_{k=0}^{4^n} \tilde{C}_n^{(N,k)}(x_1, x_2, \dots, x_n) W_n^{(N+k)}(x_1, x_2, \dots, x_n) = 0$$
 (5-87)

Similarly, we get an ODE for $W_n^{(N)}(x_1,\ldots,x_n)$ as shown in subsection 2.5. We begin with,

$$\frac{d^k}{dx_1^k} W_n^{(N)}(x_1, \dots, x_n) = \operatorname{Tr} M^{(N)}(x_1) \ H_{k,n}^{(N)}(x_1; x_2, \dots, x_n)$$
 (5-88)

where $H_{k,n}^{(N)}(x_1; x_2, \ldots, x_n)$ is computed recursively

$$H_{0,n}^{(N)}(x_1; x_2, \dots, x_n) = H_n^{(N)}(x_1; x_2, \dots, x_n)$$

$$H_{k+1,n}^{(N)}(x_1; x_2, \dots, x_n) = [H_{k,n}^{(N)}(x_1; x_2, \dots, x_n), \mathcal{D}^{(N)}(x_1)]$$

$$+ \frac{d}{dx_1} H_{k,n}^{(N)}(x_1; x_2, \dots, x_n)$$
(5-89)

Hence,

Theorem 5.5 There exist some coefficients $\hat{C}_n^{(N,k)}(x_1; x_2, \ldots, x_n)$, polynomials of x_1 , such that

$$\sum_{k=0}^{4} \hat{C}_n^{(N,k)}(x_1; x_2, \dots, x_n) \frac{d^k}{dx_1^k} W_n(x_1, \dots, x_n) = 0.$$
 (5-90)

Theorem 5.6 There exist some coefficients $\tilde{C}_n^{(N,k)}(x_1,\ldots,x_n) \in \mathbb{C}[x_1,\ldots,x_n]$, such that

$$\sum_{k=0}^{4^n} \tilde{C}_n^{(N,k)}(x_1,\dots,x_n) \frac{d^k}{dx_1^k} W_n(x_1,\dots,x_n) = 0.$$
 (5-91)

5.8 Spectral density and correlations

As we mentionned in Corollary 5.1, the densities ρ_n obey the same linear relations (ODE and recursion) as resolvents W_n .

6 2-matrix model

All the steps mentioned so far can be applied to the 2-matrix model. The only difference with the 1-matrix model is that the Lie algebra is not $\mathfrak{sl}(2)$ but $\mathfrak{sl}(\tilde{d})$ for some $\tilde{d} \geq 2$, i.e. matrices can have higher dimensions than 2.

The link between the 2-matrix model and the formalism needed here, i.e. the wave functions $\Psi(x)$, the connection $\nabla = d - \mathcal{D}(x)dx$, the adjoint sections M(x.E) and the determinantal correlators $W_n(x_1.E_1,\ldots,x_n.E_n)$ can all be found in the literature [BE06b; BE06a; EKR18; Meh04], let us recall them in short.

The following defines what is called the "2-matrix model" measure, actually its marginal probability law of eigenvalues.

Definition 6.1 (Matrix-eigenvalues measure) Let two potentials V and \tilde{V} , such that $V' \in \mathbb{C}(x)$, and $\tilde{V}' \in \mathbb{C}(x)$. Let $\Lambda = \operatorname{diag}(\Lambda_1, \ldots, \Lambda_N)$ and $\tilde{\Lambda} = \operatorname{diag}(\tilde{\Lambda}_1, \ldots, \tilde{\Lambda}_N)$, two $N \times N$ diagonal matrices. We define

$$Z^{(N)} = \int_{\gamma^N \times \tilde{\gamma}^N} d\Lambda d\tilde{\Lambda} \ \Delta(\Lambda) \Delta(\tilde{\Lambda}) \ \det\left(e^{\Lambda_i \tilde{\Lambda}_j}\right) e^{-\text{Tr}\left(V(\Lambda) + \tilde{V}(\tilde{\Lambda})\right)}$$
(6-1)

and we define the marginal measure for $\Lambda \in \gamma^N$:

$$d\mu^{(N)}\Lambda = \frac{1}{Z^{(N)}}d\Lambda \int_{\tilde{\gamma}^N} d\tilde{\Lambda} \ \Delta(\Lambda)\Delta(\tilde{\Lambda}) \ \det\left(e^{\Lambda_i\tilde{\Lambda}_j}\right)e^{-\operatorname{Tr}\left(V(\Lambda)+\tilde{V}(\tilde{\Lambda})\right)}$$
(6-2)

and the marginal measure for $\tilde{\Lambda} \in \tilde{\gamma}^N$:

$$d\tilde{\mu}^{(N)}\tilde{\Lambda} = \frac{1}{Z^{(N)}}d\tilde{\Lambda} \int_{\gamma^N} d\Lambda \ \Delta(\Lambda)\Delta(\tilde{\Lambda}) \ \det\left(e^{\Lambda_i\tilde{\Lambda}_j}\right)e^{-\operatorname{Tr}\left(V(\Lambda)+\tilde{V}(\tilde{\Lambda})\right)}. \tag{6-3}$$

We denote

$$d = 1 + \deg V' \quad , \quad \tilde{d} = 1 + \deg \tilde{V}' \tag{6-4}$$

where $\deg V'$ is the sum of degrees of all poles, including the pole at ∞ . (if V, resp. \tilde{V} is a polynomial, then $d = \deg V$, resp. $\tilde{d} = \deg \tilde{V}$).

Definition 6.2 (Wave functions) Let

$$p^{(N)}(x) := \int_{\gamma^N} d\mu^{(N)} \Lambda \quad \det(x - \Lambda) \quad , \quad q^{(N)}(y) := \int_{\tilde{\gamma}^N} d\tilde{\mu}^{(N)} \tilde{\Lambda} \quad \det(y - \tilde{\Lambda}), \quad (6-5)$$

Let

$$h^{(N)} := \int_{\gamma \times \tilde{\gamma}} dx dy \, p^{(N)}(x) q^{(N)}(y) e^{-(V(x) + \tilde{V}(y) - xy)} \quad , \quad \gamma^{(N)} = \sqrt{h^{(N)}/h^{(N-1)}}. \quad (6-6)$$

It is well known [Meh04; BE06a] that $p^{(N)}$, $q^{(N)}$ form a biorthogonal family:

$$\int_{\gamma \times \tilde{\gamma}} dx dy \, p^{(N)}(x) q^{(\tilde{N})}(y) \, e^{-(V(x) + \tilde{V}(y) - xy)} = h^{(N)} \delta_{N,\tilde{N}}. \tag{6-7}$$

We define:

$$\psi^{(N)}(x) := \frac{e^{-V(x)}}{\sqrt{h^{(N)}}} p^{(N)}(x) \quad , \quad \tilde{\psi}^{(N)}(y) := \frac{e^{-\tilde{V}(y)}}{\sqrt{h^{(N)}}} q^{(N)}(y). \tag{6-8}$$

and we define the vector of dimension \tilde{d} :

$$\vec{\psi}^{(N)}(x) = \begin{pmatrix} \psi_j^{(N-\tilde{d}+1)}(x) \\ \vdots \\ \psi_j^{(N)}(x) \end{pmatrix}$$

$$(6-9)$$

Definition 6.3 (Dual wave functions) We define the Hilbert transform

$$\phi^{(N)}(x) := \frac{e^{V(x)}}{\sqrt{h^{(N)}}} \int_{\gamma \times \tilde{\gamma}} dx' dy' \frac{1}{x - x'} q^{(N)}(y') \ e^{-(V(x') + \tilde{V}(y') - x'y')}$$
(6-10)

and the vector

$$\vec{\phi}^{(N)}(x) = \begin{pmatrix} \phi^{(N-1)}(x) \\ \phi^{(N)}(x) \\ \vdots \\ \phi^{(N+\tilde{d}-2)}(x) \end{pmatrix}$$
(6-11)

We also define the Fourier transforms,

$$\phi^{(N,i)}(x) := \int_{\tilde{\gamma}_i} dy' \phi^{(N)}(y') \ e^{x'y'} \tag{6-12}$$

where $\tilde{\gamma}_i$, $i = 1, ..., \tilde{d} - 1$ form a basis of homology paths on which the integral can be defined.

We define the $\tilde{d} \times \tilde{d}$ matrix by its matrix elements $i, j = 1, \dots, \tilde{d}$:

$$(\Phi^{(N)}(x))_{j,i} = \phi^{(N-\tilde{d}+i,j-1)}(x)$$
(6-13)

where we defined $\phi^{(N,0)}(x) = \phi^{(N)}(x)$.

6.1 Recursion and ODE for wave functions

Proposition 6.1 (Recursion relation) The wave function satisfies a recursion relation of order \tilde{d} , written:

$$x\psi^{(N)}(x) = \gamma^{(N+1)}\psi^{(N+1)}(x) + \sum_{k=0}^{\tilde{d}-1} Q_{N,N-k}\psi^{(N-k)}(x).$$
 (6-14)

or written in matrix form

$$\vec{\psi}^{(N+1)}(x) = \mathcal{R}^{(N)}(x)\vec{\psi}^{(N)}(x) \tag{6-15}$$

$$\mathcal{R}^{(N)}(x) = \begin{pmatrix} 0 & 1 & & & \\ 0 & & \ddots & & \\ 0 & & & 1 \\ \frac{-Q_{N,N-\tilde{d}+1}}{\gamma^{(N+1)}} & \dots & \frac{-Q_{N,N-1}}{\gamma^{(N+1)}} & \frac{x-Q_{N,N}}{\gamma^{(N+1)}} \end{pmatrix}$$
(6-16)

 $\mathcal{R}^{(N)}(x)$ is a matrix of size $\tilde{d} \times \tilde{d}$, polynomial of x of degree 1. Its inverse is also polynomial of degree 1.

Similarly for the dual wave functions we have, for $N \geq 1$

$$x\phi^{(N)}(x) = \gamma^{(N)}\phi^{(N-1)}(x) + \sum_{k=0}^{\tilde{d}-1} Q_{N+k,N}\phi^{(N+k)}(x).$$
 (6-17)

which can be written

$$\vec{\phi}^{(N+1)}(x) = \tilde{\mathcal{R}}^{(N)}(x)\vec{\phi}^{(N)}(x) \tag{6-18}$$

$$\tilde{\mathcal{R}}^{(N)}(x) = \begin{pmatrix} \frac{x - Q_{N,N}}{\gamma^{(N)}} & \dots & \frac{-Q_{N+\tilde{d}-2,N}}{\gamma^{(N)}} & \frac{-Q_{N+\tilde{d}-1,N}}{\gamma^{(N)}} \\ 1 & & 0 \\ & \ddots & & \vdots \\ & & 1 & 0 \end{pmatrix}^{-1}$$
(6-19)

Definition 6.4 (Iterated shift operator) Let

$$\mathcal{R}^{(N,k)}(x) := \mathcal{R}^{(N+k-1)}(x)\mathcal{R}^{(N+k-2)}(x)\dots\mathcal{R}^{(N)}(x)$$
(6-20)

which is a matrix, polynomial of x of degree at most k. It is such that

$$\vec{\psi}^{(N+k)}(x) = \mathcal{R}^{(N,k)}(x)\vec{\psi}^{(N)}(x).$$
 (6-21)

similarly

$$\tilde{\mathcal{R}}^{(N,k)}(x) := \tilde{\mathcal{R}}^{(N+k-1)}(x)\tilde{\mathcal{R}}^{(N+k-2)}(x)\cdots\tilde{\mathcal{R}}^{(N)}(x)$$
(6-22)

which is a matrix, polynomial of x of degree at most k. It is such that

$$\vec{\phi}^{(N+k)}(x) = \tilde{\mathcal{R}}^{(N,k)}(x)\vec{\phi}^{(N)}(x). \tag{6-23}$$

The wave functions satisfy a differential system of order \tilde{d} (see [BE06b]):

Proposition 6.2 (ODE)

$$\frac{d}{dx}\vec{\psi}^{(N)}(x) = \mathcal{D}^{(N)}(x)\vec{\psi}^{(N)}(x) \tag{6-24}$$

where $\mathcal{D}^{(N)}(x)$ is a matrix of size $\tilde{d} \times \tilde{d}$, polynomial of x of degree less or equal to degree of V'. The matrix $\mathcal{D}^{(N)}(x)$ is given explicitly in [BE06b; BE06a].

Similarly, there is a $\tilde{d} \times \tilde{d}$, matrix $\tilde{\mathcal{D}}^{(N)}(x)$ polynomial of x of degree less or equal to degree of V' such that

$$\frac{d}{dx}\vec{\phi}^{(N)}(x) = \tilde{\mathcal{D}}^{(N)}(x)\vec{\phi}^{(N)}(x). \tag{6-25}$$

6.2 Matrix Wave function

In this short presentation, we bypass the definition of the full $\tilde{d} \times \tilde{d}$ matrix $\Psi^{(N)}(x)$, we shall admit the following propositions, and refer the reader to [BE06a; EKR18] in case of interest.

Proposition 6.3 (Matrix wave function) there exists a square invertible matrix of size $\tilde{d} \times \tilde{d}$

$$\Psi^{(N)}(x) \in GL(\tilde{d}, \mathbb{C}) \tag{6-26}$$

whose first column is $(\psi^{(N-\tilde{d}+1)}(x), \dots, \psi^{(N)}(x))$, and that satisfies for N large enough:

$$\frac{d}{dx}\Psi^{(N)}(x) = \mathcal{D}^{(N)}(x)\Psi^{(N)}(x). \tag{6-27}$$

and

$$\Psi^{(N+1)}(x) = \mathcal{R}^{(N)}(x)\Psi^{(N)}(x). \tag{6-28}$$

And there exists a square invertible matrix of size $\tilde{d} \times \tilde{d}$ (given in eq. 6-13)

$$\Phi^{(N)}(x) \in GL(\tilde{d}, \mathbb{C}) \tag{6-29}$$

whose first column is $(\phi^{(N-1)}(x), \dots, \phi^{(N+\tilde{d}-2)}(x))$, and that satisfies for N large enough:

$$\frac{d}{dx}\Phi^{(N)}(x) = \tilde{\mathcal{D}}^{(N)}(x)\Phi^{(N)}(x). \tag{6-30}$$

and

$$\Phi^{(N+1)}(x) = \tilde{\mathcal{R}}^{(N)}(x)\Phi^{(N)}(x). \tag{6-31}$$

proof: Proved in [BE06b; BE06a]. \square

Theorem 6.1 (Duality)

$$\Phi^{(N)^t}(x) \mathbb{A}^{(N)} \Psi^{(N)}(x) = Id \tag{6-32}$$

where $\mathbb{A}^{(N)}$ is the $\tilde{d} \times \tilde{d}$ matrix

$$\mathbb{A}_{i,j}^{(N)} = A_{N-2+i,N-2+j} \tag{6-33}$$

where

$$A_{N-1+i,N-j} = \begin{cases} Q_{N-1+i,N-j} & \text{if } 0 < i+j < \tilde{d} \\ -\gamma^{(N-1)} & \text{if } i = j = 0 \\ 0 & \text{otherwise} \end{cases}$$
 (6-34)

This implies

$$\Psi^{(N)^{-1}}(x) = \Phi^{(N)^t}(x)A^{(N)}.$$
(6-35)

$$\mathbb{A}^{(N)}\mathcal{D}^{(N)}(x) = -\tilde{\mathcal{D}}^{(N)}(x)\mathbb{A}^{(N)}.$$
(6-36)

proof: Proved in [BE06b; BE06a].

6.3 Adjoint section

Definition 6.5

$$M^{(N)}(x) = \Psi^{(N)}(x)E\Psi^{(N)}(x)^{-1}$$
 , $E = \text{diag}(1, 0, \dots, 0)$ (6-37)

or equivalently

$$M^{(N)}(x) = \Psi^{(N)}(x)E\Phi^{(N)}(x)^{t}\mathbb{A}^{(N)}.$$
(6-38)

Proposition 6.4 (proved in [BEH02])

$$(M^{(N)}(x))_{i,j} = \psi^{(N-\tilde{d}+i)}(x) \sum_{k=0}^{\tilde{d}-1} \phi^{(N-2+k)}(x) A_{N-2+k,N-\tilde{d}+j}$$
 (6-39)

where

$$A_{N-1+i,N-j} = \begin{cases} Q_{N-1+i,N-j} & \text{if } 0 < i+j < \tilde{d} \\ -\gamma^{(N-1)} & \text{if } i = j = 0 \\ 0 & \text{otherwise} \end{cases}$$
 (6-40)

proof: proved in [BEH02]. \square

Theorem 6.2 (adjoint equations)

$$\frac{d}{dx}M^{(N)}(x) = [\mathcal{D}^{(N)}(x), M^{(N)}(x)] \tag{6-41}$$

$$M^{(N+1)}(x) = \mathcal{R}^{(N)}(x)M^{(N)}(x)\mathcal{R}^{(N)}(x)^{-1}$$
(6-42)

proof: Obvious. \square

6.4 Correlations

Definition 6.6 Let the expectation of products of resolvents

$$\hat{W}_{n}^{(N)}(x_{1},...,x_{n}) = \int_{\gamma^{N}} d\mu^{(N)} \Lambda \quad \prod_{i=1}^{n} \operatorname{Tr} \frac{1}{x_{i} - \Lambda}$$
 (6-43)

and $W_n^{(N)}$ their cumulants.

Let also the densities

$$\hat{\rho}_n^{(N)}(x_1,\dots,x_n) = \int_{\gamma^N} d\mu^{(N)} \Lambda \quad \prod_{i=1}^n \operatorname{Tr} \delta(x_i - \Lambda)$$
 (6-44)

and $\rho^{(N)}$ their cumulants.

Proposition 6.5 The correlations of resolvents are the same as given by Definition 1.2:

$$W_1^{(N)}(x.E) = \text{Tr } \mathcal{D}^{(N)}(x)M^{(N)}(x.E)$$
(6-45)

$$W_2^{(N)}(x_1.E_1, x_2.E_2) = \frac{1}{(x_1 - x_2)^2} \operatorname{Tr} M^{(N)}(x_1.E_1) M^{(N)}(x_2.E_2)$$
 (6-46)

and for $n \geq 3$

$$W_n^{(N)}(x_1, \dots, x_n) = \sum_{\sigma \in \mathfrak{S}_n^{1-cycle}} (-1)^{\sigma} \frac{\text{Tr} \prod_{i=1}^n M^{(N)}(x_{\sigma^i(1)})}{\prod_{i=1}^n (x_i - x_{\sigma(i)})}$$
(6-47)

proof: See [BE06a]. \square

6.5 Densities

The same formulas as in subsubsection 5.3.3 apply also in the 2-matrix model.

Notice from eq. 6-12 that $\phi^{(N)}(x)$ is discontinuous across γ :

Disc
$$\phi^{(N)}(x) = \frac{1}{\sqrt{h^{(N)}}} \int_{\tilde{\gamma}} e^{-\tilde{V}(y') + xy'} q^{(N)}(y') dy'.$$
 (6-48)

This implies that resolvents W_n are discontinuous, and due to the general Stieltjes theory, the density of eigenvalues and correlations, are the discontinuities of resolvents. This gives the same formulas as in subsubsection 5.3.3.

6.6 Linear ODE, Recursions

As a consequence:

Proposition 6.6 $W_1^{(N)}$ satisfy an ODE and a linear recursion of order r with polynomial coefficients, where $r = \tilde{d}^2 = \dim \mathfrak{gl}(\tilde{d})$.

proof: The ODE follows from Theorem 2.1 where we set $\mathfrak{g} = \mathfrak{g}l(\tilde{d})$ and $\mathcal{D}(x) \to \mathcal{D}^{(N)}(x)$. The linear recursion is given by Theorem 4.1 with $\mathfrak{g} = \mathfrak{g}l(\tilde{d})$.

Proposition 6.7 We have the following properties:

- 1. $W_n^{(N)}(x_1,...,x_n)$ satisfies an ODE of order r with respect to the variable x_1 with coefficients polynomials of x_1 .
- 2. $W_n^{(N)}(x_1,...,x_n)$ satisfy an ODE with respect to the variable x_1 and a linear recursion of order r^n with coefficients polynomials of $(x_1,...,x_n)$.

proof: The first property follows from Theorem 2.5. For the second statement, the ODE follows from Theorem 2.6 and the linear recursion follows from Theorem 4.2. \Box

7 Minimal models

Our formalism is also well suited for minimal models.

Minimal models arise as limits of matrix models. They can also be defined independently of matrix models, from integrable hierarchies, KdV, KP, Toda, see [BBT03]. They are classified by two integers (p,q), in short, the (p,q) minimal model can be formulated in terms of a differential system of order q, with polynomial coefficients whose degree is bounded by p.

7.1 Minimal models q = 2

This is the case of order 2 ODE, i.e. $\mathfrak{s}l(2,\mathbb{C})$.

Let U(t) a function of time t that we shall precise below, but let us already mention that it will be solution to some non-linear ODE having the Painlevé property. In all what follows, there will be functions of x and of time t. We shall denote f' = df/dx and $\dot{f} = df/dt$.

Definition 7.1 (Gelfand-Dikii polynomials) We define by recursion the following differential polynomials $R_n(U) \in \mathbb{Q}[U, \dot{U}, \ddot{U}, \ddot{U}, \dots]$

$$R_0(U) = 2 \tag{7-1}$$

$$\dot{R}_{n+1} = -2U\dot{R}_n - \dot{U}R_n + \frac{1}{4}\ddot{R}_n \tag{7-2}$$

and we choose the primitive that is homogeneous (in U and ").

The first few are

$$R_0 = 2 \quad , \quad R_1 = -2U \quad , \quad R_2 = 3U^2 - \frac{1}{2}\ddot{U} \quad , \quad R_3 = -5U^3 + \frac{5}{2}U\ddot{U} + \frac{5}{4}\dot{U}^2 - \frac{1}{8}\dddot{U}$$
(7-3)

Definition 7.2 (Lax pair) Let

$$\mathcal{R}(x,t) := \begin{pmatrix} 0 & 1\\ x + 2U(t) & 0 \end{pmatrix} \tag{7-4}$$

Define,

$$B_n(x,U) := \frac{1}{2} \sum_{j=0}^n x^{n-j} R_j(U)$$
 (7-5)

Define

$$\mathcal{D}_n(x,t) := \begin{pmatrix} -\frac{1}{2}\dot{B}_n & B_n \\ (x+2U)B_n - \frac{1}{2}\ddot{B}_n & \frac{1}{2}\dot{B}_n \end{pmatrix}$$
 (7-6)

Notice that $\mathcal{R}(x,t)$ and $\mathcal{D}_n(x,t)$ are traceless, so that they belong to the Lie algebra $\mathfrak{sl}(2,\mathbb{C})$.

Proposition 7.1 (Gelfand Dikii) We have

$$\dot{\mathcal{D}}_n(x,t) + \left[\mathcal{D}_n(x,t), \mathcal{R}(x,t)\right] = -\dot{R}_{n+1} \begin{pmatrix} 0 & 0\\ 1 & 0 \end{pmatrix}$$
 (7-7)

proof: It is the classical Gelfand-Dikii theorem [GD75; BBT03]. □

Proposition 7.2 (Compatibility and wave function) Let $m \geq 0$ an integer. Let $\vec{t} = (t_0, t_1, t_2, \dots, t_m)$ a set of parameters (called higher times). Let

$$\mathcal{D}(x,t,\vec{t}) = \sum_{k=0}^{m} t_k \mathcal{D}_k(x,t). \tag{7-8}$$

 $\mathcal{D}(x,t;\vec{t})$ satisfies the Lax equation

$$\frac{d}{dt}\mathcal{D}(x,t;\vec{t}) + [\mathcal{D}(x,t;\vec{t}),\mathcal{R}(x,t)] = \frac{d}{dx}\mathcal{R}(x,t)$$
 (7-9)

if and only if U is solution to the equation

$$\sum_{k=0}^{m} t_k R_{k+1}(U) = -t. (7-10)$$

In this case, there exists a matrix $\Psi(x,t;\vec{t})$ such that

$$\frac{d}{dx}\Psi = \mathcal{D}\Psi \quad , \quad \frac{d}{dt}\Psi = \mathcal{R}\Psi \tag{7-11}$$

proof: This is an immediate corollary of Proposition 7.1. \square

Example 7.1 (m = 0 : Airy) take $\vec{t} = (1)$, eq. 7-10 reduces to:

$$-2U = -t \tag{7-12}$$

We have

$$\mathcal{R}(x,t) = \begin{pmatrix} 0 & 1 \\ x+t & 0 \end{pmatrix} = \mathcal{D}(x,t) \tag{7-13}$$

The matrix $\Psi(x,t) = \Psi(x+t)$ satisfies the Airy equation, and is worth:

$$\Psi(x,t) = \Psi(x+t) = \begin{pmatrix} Ai(x+t) & Bi(x+t) \\ Ai'(x+t) & Bi'(x+t) \end{pmatrix}$$
(7-14)

Example 7.2 (m = 1 : Painlevé 1) [Pai00] take $\vec{t} = (0, 1)$, then eq. 7-10 is the Painlevé 1 equation:

$$3U^2 - \frac{1}{2}\ddot{U} = -t\tag{7-15}$$

and

$$B_1(x,t) = x - U(t), (7-16)$$

and

$$\mathcal{D}_1(x,t) = \begin{pmatrix} \frac{1}{2}\dot{U} & x - U\\ (x + 2U)(x - U) + \frac{1}{2}\ddot{U} & -\frac{1}{2}\dot{U} \end{pmatrix}$$
(7-17)

Its characteristic polynomial is

$$\det(y - \mathcal{D}_1(x,t)) = y^2 - \frac{1}{4}\dot{U}^2 - (x+2U)(x-U)^2 - \frac{1}{2}\ddot{U}(x-U)$$

$$= y^2 - x^3 - (3U^2 - \frac{1}{2}\ddot{U})x - 2U^3 - \frac{1}{4}\dot{U}^2 + \frac{1}{2}U\ddot{U}$$

$$= y^2 - x^3 + tx - 2U^3 - \frac{1}{4}\dot{U}^2 + \frac{1}{2}U\ddot{U}$$
(7-18)

Remark that

$$\frac{d}{dt} \left(-2U^3 - \frac{1}{4}\dot{U}^2 + \frac{1}{2}U\ddot{U} \right) = -6U^2\dot{U} - \frac{1}{2}\dot{U}\ddot{U} + \frac{1}{2}\dot{U}\ddot{U} + \frac{1}{2}U\ddot{U}
= U(-6U\dot{U} + \frac{1}{2}\ddot{U})
= U\frac{d}{dt}(-3U^2 + \frac{1}{2}\ddot{U})
= U$$
(7-19)

7.2 ODE with respect to time for W_1

Proposition 7.3 (ODE for W_1) W_1 satisfies the following ODE of order 3:

$$W_1 = \sum_{k=0}^{m} t_k \left((2(x+2U)B_k - \frac{1}{2}\ddot{B}_k)\dot{W}_1 + \frac{1}{2}\dot{B}_k\ddot{W}_1 - \frac{1}{2}B_k\ddot{W}_1 \right)$$
(7-20)

proof: Using the formula $W_1 = \text{Tr}(DM)$ we have:

$$\frac{d}{dt}W_1 = \operatorname{Tr}\left(\frac{d}{dt}DM + D\frac{d}{dt}M\right)
\dot{W}_1 = \operatorname{Tr}\left((\dot{D} + [D, R])M\right)
\dot{W}_1 = \operatorname{Tr}\left(R'M\right) \quad (By \text{ using the compatibility equation})$$
(7-21)

Therefore:

$$\frac{d^k}{dt^k}W_1 = \text{Tr}(D_k M) = \text{Tr}(\dot{D}_{k-1} + [D_{k-1}, R])M, \quad D_0 = D = \sum_{k=0}^m t_k D_k$$
 (7-22)

we have:

$$D_1 = R' = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$
 $D_2 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ $D_3 = \begin{pmatrix} 0 & -2 \\ 2(x+2U) & 0 \end{pmatrix}$

Also we remark that

$$(x+2U)D_1 - \frac{1}{2}D_3 = \begin{pmatrix} 0 & 1\\ 0 & 0 \end{pmatrix}$$

Therefore

$$D_0 = \sum_{k=0}^{m} t_k \left[B_k((x+2U)D_1 - \frac{1}{2}D_3) + \frac{1}{2}\dot{B}_k D_2 + ((x+2U)B_k - \frac{1}{2}\ddot{B}_k)D_1 \right]$$

Hence

$$W_1 = \sum_{k=0}^{m} t_k \left[(2(x+2U)B_k - \frac{1}{2}\ddot{B}_k)\dot{W}_1 + \frac{1}{2}\dot{B}_k\dot{W}_1 - \frac{1}{2}B_k\ddot{W}_1 \right]$$

Example 7.3 (Airy m = 0, ODE W_1)

$$W_1 - 2(x+t)\dot{W}_1 + \frac{1}{2}\ddot{W}_1 = 0 (7-23)$$

Example 7.4 (Painlevé 1, m = 1, ODE W_1)

$$W_1 - (2x^2 + 2Ux - U^2 + t)\dot{W}_1 + \frac{1}{2}\dot{U}\ddot{W}_1 + \frac{1}{2}(x - U)\ddot{W}_1 = 0$$
 (7-24)

7.3 Minimal models general case

Minimal models can also be defined for rank $q \ge 2$ systems. Let us give the example (4,3), also called "Ising model":

$$R(x,U) = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ x + \frac{3}{2}\hbar\dot{U} & 3U & 0 \end{pmatrix}$$
 (7-25)

$$\mathcal{D}(x,U) = \begin{pmatrix} 2U^2 - \frac{1}{6}\hbar^2\ddot{U} & x + \frac{1}{2}\hbar\dot{U} & -U \\ -Ux + \frac{5}{2}\hbar\dot{U}\dot{U} - \frac{1}{6}\hbar^3\ddot{U} & -U^2 + \frac{1}{3}\hbar^2\ddot{U} & x - \frac{1}{2}\hbar\dot{U} \\ x^2 + \hbar^2(\frac{7}{4}\dot{U}^2 + \frac{5}{2}U\ddot{U}) - \frac{1}{6}\hbar^4\dddot{U} & 2Ux - \hbar\dot{U}\dot{U} + \frac{1}{6}\hbar^3\ddot{U} & -U^2 - \frac{1}{6}\hbar^2\ddot{U} \end{pmatrix}$$
(7-26)

The Lax equation implies

$$-\frac{1}{6}\hbar^4 \ddot{U} + 3\hbar^2 U \ddot{U} + \frac{3}{2}\hbar^2 \dot{U}^2 - 4U^3 = t \tag{7-27}$$

The ODE, with respect to time, satisfied by W_1 is

$$\sum_{k=0}^{8} \alpha_k(x,t) \frac{d^k}{dt^k} W_1(x,t) = 0$$
 (7-28)

where

$$\begin{split} \alpha_0(x,t) &= 9\hbar^2 \dot{U} \\ \alpha_1(x,t) &= -\frac{3\hbar}{4} \left(4\hbar^4 U U^{(5)} - 6\hbar^4 \dot{U} \, \ddot{U} - 60\hbar^2 U^2 \ddot{U} \right. \\ &\qquad \qquad - 54\hbar^2 U \dot{U} \ddot{U} + 63\hbar^2 \dot{U}^3 + 36x^2 \dot{U} + 72U^3 \dot{U} \right) \\ \alpha_2(x,t) &= -\frac{39\hbar^4 U \, \dddot{U}}{2} + 15\hbar^4 \dot{U} \, \dddot{U} + \frac{369\hbar^2 U^2 \ddot{U}}{2} \\ &\qquad \qquad + \frac{189\hbar^2 U \dot{U}^2}{4} + 27Ux^2 - 108U^4 \\ \alpha_3(x,t) &= -\frac{105\hbar^3 U \, \dddot{U}}{2} + \frac{75\hbar^3 \dot{U} \, \ddot{U}}{2} + 189\hbar U^2 \dot{U} \\ \alpha_4(x,t) &= -\frac{147\hbar^2 U \, \ddot{U}}{2} + 36\hbar^2 \dot{U}^2 + 81U^3 \\ \alpha_5(x,t) &= -36\hbar U \, \dot{U} \\ \alpha_6(x,t) &= -18U^2 \\ \alpha_7(x,t) &= -\hbar \dot{U} \\ \alpha_8(x,t) &= U \end{split}$$

8 Schlesinger systems

8.1 Definition

The Schlesinger system [Sch12] with Lie algebra \mathfrak{g} and N points z_1, \ldots, z_N , is defined as follows:

$$\mathcal{D}(x) = \sum_{i=1}^{N} \frac{A_i}{x - z_i} \qquad A_i \in \mathfrak{g}, \ \sum_{i=1}^{N} A_i = 0.$$
 (8-1)

(remark: dropping the constraint $\sum_{i=1}^{N} A_i = 0$ is equivalent to choosing a pole $z_{N+1} = \infty$ and take $A_{\infty} = -\sum_{i=1}^{N} A_i$).

We choose fixed monodromies:

eigenvalues
$$(A_i) = diag(\theta_{i,1}, \dots, \theta_{i,r}) = \theta_i \in \mathfrak{h} = Cartan algebra.$$
 (8-2)

and we fix the eigenvectors of A_i by the Schlesinger equations:

$$\frac{dA_i}{dz_j} = [A_j, A_i]. (8-3)$$

8.2 Schlesinger $\mathfrak{sl}(2)$, N=3

Take the points $0, 1, \infty$ and

$$\mathcal{D}(x) = \frac{A_0}{x} + \frac{A_1}{x - 1} \tag{8-4}$$

with

$$A_0 = \begin{pmatrix} \theta_0 & 0\\ 0 & -\theta_0 \end{pmatrix} \tag{8-5}$$

$$A_{1} = \frac{1}{2\theta_{0}} \begin{pmatrix} \theta_{\infty}^{2} - \theta_{1}^{2} - \theta_{0}^{2} & 2\theta_{0}\theta_{1} + \theta_{\infty}^{2} - \theta_{1}^{2} - \theta_{0}^{2} \\ 2\theta_{0}\theta_{1} - (\theta_{\infty}^{2} - \theta_{1}^{2} - \theta_{0}^{2}) & -\theta_{\infty}^{2} + \theta_{1}^{2} + \theta_{0}^{2} \end{pmatrix}$$
(8-6)

$$A_{\infty} = -A_0 - A_1 = \frac{-1}{2\theta_0} \begin{pmatrix} \theta_{\infty}^2 - \theta_1^2 + \theta_0^2 & 2\theta_0\theta_1 + \theta_{\infty}^2 - \theta_1^2 - \theta_0^2 \\ 2\theta_0\theta_1 - (\theta_{\infty}^2 - \theta_1^2 - \theta_0^2) & -\theta_{\infty}^2 + \theta_1^2 - \theta_0^2 \end{pmatrix}$$
(8-7)

They are such that

eigenvalues
$$(A_i) = diag(\theta_i, -\theta_i).$$
 (8-8)

We find

$$\sum_{k=0}^{3} \alpha_k(x) \frac{d^k}{dx^k} W_1(x) = 0 \tag{8-9}$$

with coefficients:

$$\alpha_0(x,t) = 6\theta_0^2 x^2 - 10\theta_0^2 x + 4\theta_0^2 - 6\theta_1^2 x^2 + 2\theta_1^2 x - 4\theta_\infty^2 x^3 + 6\theta_\infty^2 x^2 - 2\theta_\infty^2 x + 4x^3 - 6x^2 + 2x$$

$$\alpha_1(x,t) = 2x(x-1) \left(2\theta_0^2 x - 2\theta_0^2 - 2\theta_1^2 x - 2\theta_\infty^2 x^2 + 2\theta_\infty^2 x + 7x^2 - 7x + 1 \right)$$

$$\alpha_2(x,t) = 4(2x-1)x^2(1-x)^2$$

$$\alpha_3(x,t) = x^3(1-x)^3.$$
(8-10)

9 Conclusion

We have shown how to exploit the ODE satisfied by the wave function (flat section of a Lie group bundle) or equivalently the adjoint ODE satisfied by M (flat section of the adjoint Lie algebra bundle) to derive, in a systematic way, linear ODE and recursions with polynomial coefficients for every correlator W_n .

Acknowledgments

This work is supported by the ERC synergy grant ERC-2018-SyG 810573, "Re-NewQuantum". This paper is a result of the ERC-SyG project, Recursive and Exact New Quantum Theory (ReNewQuantum) which received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under grant agreement No 810573. D. M. acknowledges support by Onassis Foundation under scholarship ID: F ZR 038/1-2021/2022

References

- [BBT03] O. Babelon, D. Bernard, and M. Talon. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2003. DOI: 10.1017/CB09780511535024.001.
- [BE06a] M. Bergère and B. Eynard. "Mixed correlation function and spectral curve for the 2-matrix model". *Journal of Physics A: Mathematical and General* 39.49 (Nov. 2006), p. 15091. DOI: 10.1088/0305-4470/39/49/004. arXiv: math-ph/0605010.
- [BE09] M. Bergère and B. Eynard. "Determinantal formulae and loop equations" (2009). arXiv: 0901.3273.
- [BEH02] M. Bertola, B. Eynard, and J. Harnad. "Duality, Biorthogonal Polynomials and Multi-Matrix Models". Communications in Mathematical Physics 229.1 (Aug. 2002), pp. 73–120. DOI: 10.1007/s002200200663. arXiv: nlin/0108049.

- [BDY16a] M. Bertola, B. Dubrovin, and D. Yang. "Correlation functions of the KdV hierarchy and applications to intersection numbers over $\overline{M}_{g,n}$ ". Physica D: Nonlinear Phenomena 327 (2016), pp. 30–57. ISSN: 0167-2789. DOI: https://doi.org/10.1016/j.physd.2016.04.008. arXiv: 1504.06452.
- [BDY16b] M. Bertola, B. Dubrovin, and D. Yang. "Simple Lie Algebras and Topological ODEs". International Mathematics Research Notices 2018.5 (Dec. 2016), pp. 1368–1410. ISSN: 1073-7928. DOI: 10.1093/imrn/rnw285. arXiv: 1508.03750.
- [BE06b] M. Bertola and B. Eynard. "The PDEs of Biorthogonal Polynomials Arising in the Two-Matrix Model". *Mathematical Physics, Analysis and Geometry* 9.1 (Feb. 2006), pp. 23–52. ISSN: 1572-9656. DOI: 10.1007/s11040-005-9000-x. arXiv: nlin/0311033.
- [EKR18] B. Eynard, T. Kimura, and S. Ribault. *Random matrices*. 2018. arXiv: 1510.04430 [math-ph].
- [GD75] I. M. Gel'fand and L. A. Dikii. "Asymptotic behaviour of the resolvent of Sturm-Liouville equations and the algebra of the Korteweg-De Vries equations". Russian Mathematical Surveys 30.5 (Oct. 1975), p. 77. DOI: 10.1070/RM1975v030n05ABEH001522.
- [Hei78] E. Heine. Handbuch der Kugelfunctionen, Theorie und Anwendungen. De Gruyter, 1878.
- [Meh04] M. L. Mehta, ed. *Random Matrices*. Vol. 142. Pure and Applied Mathematics. Elsevier, 2004. DOI: https://doi.org/10.1016/S0079-8169(04) 80122-3.
- [Pai00] P. Painlevé. "Mémoire sur les équations différentielles dont l'intégrale générale est uniforme". fr. Bulletin de la Société Mathématique de France 28 (1900), pp. 201–261. DOI: 10.24033/bsmf.633.
- [Sch12] L. Schlesinger. "Über eine Klasse von Differentialsystemen beliebiger Ordnung mit festen kritischen Punkten." 1912.141 (1912), pp. 96–145. DOI: doi:10.1515/crll.1912.141.96.
- [Sze39] G. Szegö. Orthogonal polynomials. Vol. 23. Third edition. American Mathematical Society Colloquium Publications, 1939. DOI: http://dx.doi.org/10.1090/coll/023.