CELL DECOMPOSITION AND DUAL BOUNDARY COMPLEXES OF CHARACTER VARIETIES

TAO SU

Dedicated to the memory of my high school math teacher Zhu, Yuwen (1975-2023)

ABSTRACT. The weak geometric P=W conjecture of L. Katzarkov, A. Noll, P. Pandit, and C. Simpson asserts that for any smooth Betti moduli space \mathcal{M}_B of complex dimension d over a punctured Riemann surface, the dual boundary complex $\mathbb{D}\partial \mathcal{M}_B$ is homotopy equivalent to a (d - 1)-dimensional sphere. Here, we consider \mathcal{M}_B as a generic $GL_n(\mathbb{C})$ -character variety defined on a Riemann surface of genus g, with local monodromies specified by generic semisimple conjugacy classes at k punctures.

In this article, we establish the weak geometric P=W conjecture for all *very generic* \mathcal{M}_B in the sense that at least one conjugacy class is regular semisimple. A crucial step is to establish a stronger form of A. Mellit's cell decomposition theorem, i.e. we decompose \mathcal{M}_B (without passing to a vector bundle) into locally closed subvarieties of the form $(\mathbb{C}^{\times})^{d-2b} \times \mathcal{A}$, where \mathcal{A} is stably isomorphic to \mathbb{C}^b . A second ingredient involves a motivic characterization of the integral cohomology of dual boundary complexes developed in a subsequent article [59]. Following C. Simpson's strategy, the proof is now an inductive computation of the dual boundary complexes from such a cell decomposition.

Contents

Introduction		1
1. Setup		4
2. Cell decomposition of character varieties		12
3. Dual boundary complexes of character varieties		26
Appendix A.	Cell decomposition of braid varieties	31
Appendix B.	Quotients of varieties	34
References		38

INTRODUCTION

Let Σ be a genus g closed Riemann surface with k punctures $\sigma = \{q_1, \dots, q_k\}, k \ge 1, 2g + k \ge 3$, and $G = GL_n(\mathbb{C})$. Modulo extra input, the tame nonabelian Hodge correspondence over noncompact curves [51, 34] induces a diffeomorphism

NAH : $\mathcal{M}_{\text{Dol}} \simeq \mathcal{M}_B$

between two moduli spaces: the Dolbeault moduli space \mathcal{M}_{Dol} of stable filtered regular (parabolic) *G*-Higgs bundles on (Σ, σ) with parabolic degree 0; and the Betti moduli space \mathcal{M}_B of stable filtered *G*-local systems on $\Sigma \setminus \sigma$ with parabolic degree 0. For more on nonabelian Hodge theory, see [9, 52, 53, 54, 2, 42, 43, 26, 31].

The geometric P=W conjecture of L. Katzarkov, A. Noll, P. Pandit and C. Simpson [32, 55] predicts that, under NAH, the "Hitchin fibration at infinity" of \mathcal{M}_{Dol} matches, up to homotopy, with a "fibration at infinity" intrinsic to the algebraic variety \mathcal{M}_B . More concretely, on the Dolbeault side, the Hitchin fibration $h : \mathcal{M}_{Dol} \to \mathbb{A}$ induces:

$$\overline{h}: N_{\text{Dol}}^* = \mathcal{M}_{\text{Dol}} \setminus h^{-1}(B_R(0)) \xrightarrow{n} \mathbb{A} \setminus B_R(0) \to (\mathbb{A} \setminus B_R(0))/\text{scaling} = S^{d-1}, R \gg 0, d = \dim \mathcal{M}_{\text{Dol}};$$

On the Betti side, there is a fibration

$$\alpha: N_B^* \to \mathbb{D}\partial \mathcal{M}_B,$$

well-defined up to homotopy. Here, we fix any log compactification $\overline{\mathcal{M}}_B$ of \mathcal{M}_B with simple normal crossing boundary divisor $\partial \mathcal{M}_B$. Then, N_B^* is a punctured tubular neighborhood of $\partial \mathcal{M}_B$ in $\overline{\mathcal{M}}_B$. Moreover, $\mathbb{D}\partial \mathcal{M}_B$

is the dual boundary complex of \mathcal{M}_B , i.e. the dual complex of the irreducible components of $\partial \mathcal{M}_B$. Notice that the dual boundary complex is defined for any smooth quasi-projective variety, see Definition 3.2. For the moment, we skip the definition of α . For the details, see Remark 3.9. Then

Conjecture 0.1 ([32, 55], Geometric P=W). There is a homotopy commutative square

$$\begin{array}{ccc} N_{\text{Dol}}^{*} & \xrightarrow{\simeq} & N_{B}^{*} \\ & & & \downarrow^{\alpha} \\ S^{d-1} & \xrightarrow{\simeq} & \mathbb{D}\partial\mathcal{M}_{B} \end{array}$$

As a weak form of the geometric P=W conjecture, we in particular have

Conjecture 0.2 ([32, 55], weak geometric P=W conjecture). *The dual boundary complex* $\mathbb{D}\partial \mathcal{M}_B$ *is homotopy equivalent to the sphere* S^{d-1} *, where* $d = \dim_{\mathbb{C}} \mathcal{M}_B$.

By a **folklore conjecture**, all (smooth) Betti moduli spaces (i.e. character varieties) \mathcal{M}_B are expected to be log Calabi-Yau (**CY**). This has been verified for the case $G = SL_2(\mathbb{C})$ [62, 17]. Then, the weak geometric P=W conjecture is closely related to the following conjecture:

Conjecture 0.3 (Kontsevich(-Kollár-Xu)). *The dual boundary complex of a log CY variety is (a finite quotient of) a sphere.*

So far, the only known general result is due to J. Kollár and C. Xu [37]: If X is log Calabi-Yau of dimension ≤ 5 , then $\mathbb{D}\partial X$ is a finite quotient of a sphere.

The geometric P=W conjecture was originally inspired by and aimed at a geometric interpretation of the (cohomological) P=W conjecture of M. de Cataldo, T. Hausel and L. Migliorini [12]. The latter states that, NAH exchanges the weight filtration (algebraic geometry) on $H^*(\mathcal{M}_B, \mathbb{Q})$ with the Perverse-Leray filtration (topology) on $H^*(\mathcal{M}_{Dol}, \mathbb{Q})$:

$$\mathrm{NAH}^*(W_{2k}H^*(\mathcal{M}_B,\mathbb{Q})=W_{2k+1}H^*(\mathcal{M}_B,\mathbb{Q}))=P_kH^*(\mathcal{M}_{\mathrm{Dol}},\mathbb{Q}).$$

After the results for rank 2 [12] and respectively for genus 2 [8], the cohomological P=W conjecture has now been resolved independently by three groups [45, 29, 46] for the major case of twisted $GL_n(\mathbb{C})$ -character varieties (in particular, k = 1). See also [13, Question 4.1.7], [11, Conj.B], [39, 19], [41, Conj.4.2.7], for the extensions of the P=W conjectures to the singular or stacky character varieties.

The geometric P=W conjecture is known to recover the cohomoloical P=W conjecture for the weight in top degree [41, Thm.6.2.6]. In general, it's not sufficient to imply the latter [41, Rmk.6.2.11]. Nevertheless, the geometric P=W conjecture does contain some information beyond the cohomological P=W conjecture. For example, the latter concerns only the cohomology with rational coefficients, hence captures only $H^*(\mathbb{D}\partial \mathcal{M}_B, \mathbb{Q})$ via the identification (see e.g. [48])

$$\widetilde{H}_{i-1}(\mathbb{D}\partial\mathcal{M}_B,\mathbb{Q})\cong\operatorname{Gr}^W_{2d}H^{2d-i}(\mathcal{M}_B,\mathbb{Q}),\quad d=\dim_{\mathbb{C}}\mathcal{M}_B$$

On the other hand, the former knows $\widetilde{H}^*(\mathbb{D}\partial \mathcal{M}_B,\mathbb{Z})$.

As explained above, to interpret the cohomogical P=W conjecture in all weights, a refinement of the geometric P=W conjecture is required. Let's make a complementary remark. Assuming the folklore conjecture, we may consider only (dlt) log **CY** compactifications. Then we obtain a *refined dual boundary complex* DMR($\overline{\mathcal{M}}_B, \partial \mathcal{M}_B$), well-defined up to PL-homeomorphism [14, Prop.11]. By Conjecture 0.2, DMR($\overline{\mathcal{M}}_B, \partial \mathcal{M}_B$) is a PL-sphere of dimension d - 1. It's expected that [55, §1.2] this PL-sphere is closed related to the Kontsevich-Soibelman picture [36]: the Kontsevich-Soibelman chambers in the Hitchin base \mathbb{A} of \mathcal{M}_{Dol} should correspond to the cells in DMR($\overline{\mathcal{M}}_B, \partial \mathcal{M}_B$).

We haven't said anything about the current state of the geometric P=W conjecture. Here we are. The full geometric P=W conjecture is known for: the Painlevé cases [47, 60, 61]; the case (g, k) = (1, 0) or (k, n) = (0, 1) [41, Thm.B]. Our major interest in this paper is its weak form, i.e. the weak geometric P=W conjecture 0.2. Previously, this is only known in a few cases: $G = SL_2(\mathbb{C})$ [33, 55, 17]; the Painlevé cases as above; singular character variety of any rank with g = 1 and k = 0 [41]; smooth wild character variety of any rank with g = 0 and k = 1 [58].

Results. As a complement to the discussion above, our main result is the following:

Theorem 0.4 (Theorem 3.10). Let $(\Sigma, \sigma = \{q_1, \dots, q_k\})$ be a k-punctured genus g Riemann surface, and $\mathcal{M}_B = \mathcal{M}_\mu$ be its G-character variety of type $\mu \in \mathcal{P}_n^k$. If \mathcal{M}_μ is very generic, then the weak geometric P=W conjecture 0.2 holds for \mathcal{M}_μ .

Here, \mathcal{M}_{μ} is a $GL_n(\mathbb{C})$ -character variety on $\Sigma \setminus \sigma$ with fixed semisimple conjugacy class C_i around q_i ; $\mu = (\mu^1, \dots, \mu^k)$, where $\mu^i = (\mu_1^i \ge \mu_2^i \ge \dots) \in \mathcal{P}_n$ encodes the multiplicities of the eigenvalues of C_i . *'generic'* means: (C_1, \dots, C_k) is *generic* in the sense of [27, Def.2.1.1] (Definition 1.1); 'very generic' means: C_k is in addition regular semisimple (Assumption 1.4).

See Remark 3.16 for a discussion when \mathcal{M}_{μ} is only generic.

To prove the main result, we need to improve A. Mellit's cell decomposition [40, §.7], which applies only to a vector bundle over very generic \mathcal{M}_{μ} . More precisely, our second main result answers Mellit's question in [40, §1.4], i.e. we give a honest cell decomposition for \mathcal{M}_{μ} :

Theorem 0.5 (Theorem 2.10). Any very generic \mathcal{M}_{μ} admits a cell decomposition:

$$\mathcal{M}_{\boldsymbol{\mu}} = \sqcup_{(\vec{w}, p) \in \mathcal{W}^*} \mathcal{M}_{\boldsymbol{\mu}}(\vec{w}, p),$$

where each $\mathcal{M}_{\mu}(\vec{w}, p)$ is a locally closed affine subvariety, such that

$$\mathcal{M}_{\mu}(\vec{w},p) \cong (\mathbb{K}^{\times})^{\overline{a}(\vec{w},p)} \times \mathcal{A}_{\mu}(\vec{w},p), \quad \mathcal{A}_{\mu}(\vec{w},p) \times \mathbb{K}^{|U|} \cong \mathbb{K}^{b(\vec{w},p)},$$

where $U \subset G$ is the subgroup of unipotent upper triangular matrices, $\overline{b}(\vec{w}, p) := b(\vec{w}, p) - |U|$, and $\overline{a}(\vec{w}, p) + 2\overline{b}(\vec{w}, p) = d_{\mu} = \dim_{\mathbb{C}} \mathcal{M}_{\mu}$ is a constant.

Moreover, there exists a unique $(\vec{w}_{\max}, p_{\max})$ such that dim $\mathcal{M}_{\mu}(\vec{w}_{\max}, p_{\max})$ is of maximal dimension d_{μ} . Equivalently, $\overline{a}(\vec{w}_{\max}, p_{\max}) = d_{\mu}$ (resp. $\overline{b}(\vec{w}_{\max}, p_{\max}) = 0$). In particular, $\mathcal{M}_{\mu}(\vec{w}_{\max}, p_{\max})$ is an open dense algebraic torus:

$$\mathcal{M}_{\mu}(\vec{w}_{\max}, p_{\max}) \cong (\mathbb{K}^{\times})^{d_{\mu}}, \quad \mathcal{R}_{\mu}(\vec{w}_{\max}, p_{\max}) = \{ \text{pt} \}.$$

Note: $\mathcal{A}_{\mu}(\vec{w}, p)$ is stably isomorphic to $\mathbb{A}^{\overline{b}(\vec{w}, p)}$. We expect that, $\mathcal{A}_{\mu}(\vec{w}, p)$ is in general not isomorphic to $\mathbb{A}^{\overline{b}(\vec{w}, p)}$, hence gives a counterexample to the Zariski cancellation problem for dimension $b = \overline{b}(\vec{w}, p) \ge 3$ in characteristic zero. See Remark 2.12 for a further discussion.

Similar to [40, §.7], Theorem 0.5 is proved via a connection to braid varieties. However, a more careful analysis is required. For the sake of clarity, we will give a self-contained proof. In fact, we use a somewhat different language (diagram calculus of matrices). To prove Theorem 0.4.(2) via Theorem 0.5, the idea is to apply a **remove/reduction lemma** (Lemma 3.7): If X is a connected smooth quasi-projective variety, and $Z \subset X$ is a smooth irreducible closed subvariety with nonempty open complement U, such that $\mathbb{D}\partial Z$ is contractible, then we have a homotopy equivalence $\mathbb{D}\partial X \sim \mathbb{D}\partial U$. We do can apply the remove lemma because of a key property [59, Cor.0.3]: if \mathcal{A}_{μ} is stably isomorphic to \mathbb{A}^{b} for some $b \geq 1$, then $\mathbb{D}\partial \mathcal{A}_{\mu}$ is contractible. This property is proved via a motivic characterization of the integral cohomology of dual boundary complexes (recalled in Proposition 3.6).

As a final remark, we mention that the same strategy in this article can be applied to wild character varieties [4, 6]. This will be pursued elsewhere.

Organization. We have already explained the main ideas above. Now, we sketch the organization.

In Section 1, we set up the basic notions related to character varieties. To clarify the computations in Section 2, we introduce some diagram calculus of matrices in Section 1.3. Morally, it's about braid matrix diagrams generalizing braid matrices associated to positive braids. In Section 1.4, we review braid varieties, complemented by Appendix A.

In Section 2, we prove a strong form of the cell decomposition for \mathcal{M}_{μ} (Theorem 2.10). This involves some routine diagram calculus in Sections 2.1-2.2. In Sections 2.3-2.4, we have borrowed statements on quotients of varieties from Appendix B. In Section 2.5, we illustrate Theorem 2.10 by two examples.

In Section 3, we study dual boundary complexes of character varieties. Section 3.1 reviews the basics on dual boundary complexes. In Section 3.2, we treat the weak geometric P=W conjecture (Theorem 3.10). To end this article, we add a few remarks on some further directions in Section 3.3.

T. SU

1. Setup

Fix the base field \mathbb{K} to be algebraically closed of characteristic 0. For simplicity, $\mathbb{K} = \mathbb{C}$. A \mathbb{K} -variety means a reduced separated scheme of finite type over \mathbb{K} .

Convention 1: Let *H* be a linear algebraic group acting on a variety *X* over \mathbb{K} , then

- A principal *H*-bundle (or a fiber bundle) means so in the étale topology, unless stated otherwise.
- For *H* reductive and *X* affine, $X//H = \text{Spec } O_X(X)^H$ denotes the affine GIT quotient.
- [X/H] denotes the quotient stack of X by H.
- If *H* acts freely on *X*, and $\pi : X \to Y$ is principal *H*-bundle over a \mathbb{K} -variety, so also a geometric quotient (see [30, Def.3.27], Proposition B.9). Then denote X/H := Y.

Equivalently, this says that the quotient stack [X/H] is representable by Y. Thus, we may also use the identification [X/H] = X/H.

We refer to Appendix B for more background on various quotients of varieties.

1.1. **Generic character varieties.** Recall that, $(\Sigma, \sigma = \{q_1, \dots, q_k\})$ is a *k*-punctured genus *g* Riemann surface, $k \ge 1, 2g + k \ge 3$. Let $T \subset G = GL_n(\mathbb{K})$ be the diagonal maximal torus. Let $(C_1, \dots, C_k) \in T^k$ be semisimple elements of type $\mu := (\mu^1, \dots, \mu^k) \in \mathcal{P}_n^k$. That is, the multiplicities of eigenvalues of C_i define a partition of $n: \mu^i = (\mu_1^i, \dots, \mu_{r_i}^i) \in \mathcal{P}_n$.

Let $\mathcal{M}_{\mu} = \mathcal{M}_{B} = \mathcal{M}_{B}(\Sigma, \sigma, G; C_{1}, \dots, C_{k})$ be the character variety of *G*-local systems on Σ whose local monodromy around q_{i} is conjugate to C_{i} . More precisely, define the affine \mathbb{K} -variety

$$M_B = M_B(\Sigma, \sigma, G; C_1, \dots, C_k) := \{ (A_j)_{j=1}^{2g}, x_1, \dots, x_k) \in G^{2g+k} : \prod_{j=1}^g (A_{2j-1}, A_{2j}) \prod_{i=1}^k x_i C_i x_i^{-1} = \mathsf{id} \},$$

where (-, -) stands for the multiplicative commutator. It has an action of the reductive group

$$G_{\text{par}} := G \times \prod_{i=1}^{k} Z(C_i), \quad Z(C_i) = \text{ the centralizer of } C_i,$$

with the action given by:

$$(h_0, h_1, \dots, h_k) \cdot (A_1, \dots, A_{2j}, x_1, \dots, x_k) := (h_0 A_1 h_0^{-1}, \dots, h_0 A_{2j} h_0^{-1}, h_0 x_1 h_1^{-1}, \dots, h_0 x_k h_k^{-1}).$$

Then, the diagonal \mathbb{K}^{\times} acts trivially on M_B , and $\mathcal{M}_{\mu} := M_B / / G_{\text{par}}$ is the affine GIT quotient.

Definition 1.1 ([40, Def.4.6.1]). $(C_1, \dots, C_k) \in T^k$ is generic if: $\prod_{i=1}^k \det C_i = 1$, and for any $1 \le n' < n$, take any n' eigenvalues $\alpha_{i,1}, \dots, \alpha_{i,n'}$ of each C_i , have

$$\prod_{i=1}^{k}\prod_{j=1}^{n'}\alpha_{i,j}\neq 1.$$

In this case, \mathcal{M}_{μ} is called a *generic character variety*.

Lemma 1.2 ([27, Thm.2.1.5], [28, Thm.5.1.1]). If (C_1, \dots, C_k) is generic of type μ , then $\mathcal{M}_{\mu} = M_B/PG_{\text{par}}$ (if nonempty) is a connected smooth affine \mathbb{K} -variety of dimension

$$d_{\mu} := n^2 (2g - 2 + k) - \sum_{i,j} (\mu_j^i)^2 + 2, \qquad (1.1.1)$$

and the quotient map $\pi: M_B \to \mathcal{M}_{\mu} = M_B/PG_{\text{par}}$ is a principal PG_{par} -bundle.

Definition 1.3. $C_i \in T$ is *ordered nicely* if it's of the form $\text{Diag}(\lambda_{i,1}I_{\mu_i^i}, \dots, \lambda_{i,r_i}I_{\mu_{r_i}^i})$.

Without loss of generality, we may assume that each $C_i \in T$ is *ordered nicely*, so that $Z(C_i) \subset G$ is the Levi subgroup of block-diagonal matrices of type μ^i .

1.2. Very generic character varieties. Let $B \subset G$ be the standard Borel subgroup of upper triangular elements, with unipotent radical $U \subset B$. We make the *very generic* assumption:

Assumption 1.4. C_k is regular semisimple. So, $Z(C_k) = T$ and $\mu^k = (1^n) \in \mathcal{P}_n$.

Then, \mathcal{M}_{μ} is called a *very generic character variety*. The nice feature in this case is a cell decomposition [40], and an enhanced version will be proved in Theorem 2.10.

The first step is as follows: Taking the diagonal gives the quotient morphism

$$D: B \twoheadrightarrow B/U \cong T.$$

Define a closed (affine) subvariety of M_B by

$$M'_B := M_B \cap (G^{2g+k-1} \times U), \tag{1.2.1}$$

and a closed subgroup of G_{par} by

$$B_{\text{par}} = B \times \prod_{i=1}^{k-1} Z(C_i) \hookrightarrow G_{\text{par}} : (b, h_1, \dots, h_{k-1}) \mapsto (b, h_1, \dots, h_{k-1}, D(b)).$$
(1.2.2)

Denote $PB_{par} := B_{par}/\mathbb{K}^{\times}$. We have mutually inverse isomorphisms of G_{par} -varieties

$$PG_{\text{par}} \times^{PB_{\text{par}}} M'_B := (PG_{\text{par}} \times M'_B) / PB_{\text{par}} \xrightarrow{\simeq} M_B : [g_{\text{par}}, m'_B = (A_1, \dots, x_k)] \mapsto g_{\text{par}} \cdot m'_B,$$

$$M_B \to PG_{\text{par}} \times^{PB_{\text{par}}} M'_B : (A_1, \dots, x_k) \mapsto [(x_k, \text{id}, \dots, \text{id}), ((x_k^{-1}A_jx_k)_{j=1}^{2g}, (x_k^{-1}x_i)_{i=1}^k)],$$

where $PG_{par} \times^{PB_{par}} M'_B$ is well-defined by Proposition B.9. Then by Proposition B.12 and Proposition B.9, we obtain a natural isomorphism of \mathbb{K} -varieties:

$$M'_B/PB_{\text{par}} \cong (PG_{\text{par}} \times^{PB_{\text{par}}} M'_B)/PG_{\text{par}} \cong M_B/PG_{\text{par}} = \mathcal{M}_{\mu}$$

and the quotient $\pi' : M'_B \to \mathcal{M}_\mu = M'_B / PB_{\text{par}}$ is a principal PB_{par} -bundle. Observe that we have a quotient group $PB_{\text{par}}/U \cong PT_{\text{par}} := (T \times \prod_{i=1}^{k-1} Z(C_i)) / \mathbb{K}^{\times}$.

Take the coordinate change

$$U \to U : x_k \mapsto u_k := x_k C_k x_k^{-1} C_k^{-1}.$$
 (1.2.3)

We can re-write

$$M'_{B} = \{ (A_{1}, \dots, x_{k-1}, u_{k}) \in G^{2g+k-1} \times U : (\prod_{j=1}^{g} (A_{2j-1}, A_{2j})) (\prod_{i=1}^{k-1} x_{i}C_{i}x_{i}^{-1})u_{k}C_{k} = \mathsf{id} \}$$
(1.2.4)

with the action of B_{par} given by:

$$(b, h_1, \dots, h_{k-1}) \cdot (A_1, \dots, x_{k-1}, u_k) \coloneqq (bA_1 b^{-1}, \dots, bx_{k-1} h_{k-1}^{-1}, bu_k (b^{C_k})^{-1}), \quad b^{C_k} \coloneqq C_k b C_k^{-1}.$$
(1.2.5)

Next, for any *nicely ordered* $C \in T$, take the parabolic subgroup $P = BZ(C) \subset G$. Denote the Weyl groups $W \cong S_n \subset G$, W(C) := W(Z(C)). Recall the Bruhat cell decomposition

$$G = \sqcup_{w \in W/W(C)} B \dot{w} P,$$

where $\dot{w} \in W$ denotes the *shortest* representative of $w \in W/W(C)$.

In our setting, $P_i := BZ(C_i) \subset G$. For each sequence

$$\vec{w} = (\tau_1, \dots, \tau_{2g}, w_1, \dots, w_{k-1}) \in W^{2g} \times \prod_{i=1}^{k-1} W/W(C_i),$$
 (1.2.6)

we obtain a locally closed affine B_{par} -subvariety of M'_B :

$$M'_{B}(\vec{w}) := M'_{B} \cap \left(\prod_{j=1}^{2g} B\tau_{j}B \times \prod_{i=1}^{k-1} B\dot{w}_{i}P_{i} \times U\right) = M_{B} \cap \left(\prod_{j=1}^{2g} B\tau_{j}B \times \prod_{i=1}^{k-1} B\dot{w}_{i}P_{i} \times U\right).$$
(1.2.7)

Define

$$\mathcal{M}_{\mu}(\vec{w}) := \pi'(M'_B(\vec{w})) \hookrightarrow \mathcal{M}_{\mu} = M'_B/PB_{\text{par}}.$$
(1.2.8)

By Corollary B.11, $\mathcal{M}_{\mu}(\vec{w}) \subset \mathcal{M}_{\mu}$ is a locally closed K-subvariety, and the quotient map

$$\pi'_{\vec{w}} := \pi'|_{M'_B(\vec{w})} : M'_B(\vec{w}) \to \mathcal{M}_{\mu}(\vec{w}) = M'_B(\vec{w})/PB_{\text{par}}$$

is a principal PB_{par} bundle.

To obtain a cell decomposition for \mathcal{M}_{μ} , the idea is to decompose $M'_B(\vec{w})$ further, via a connection to the so-called *braid varieties*. We will come back to this point after some preparations.

T. SU

1.3. **Diagram calculus of matrices.** To relate character and braid varieties, we introduce some diagram calculus of matrices. Denote

$$FBr_n^+ := \langle \sigma_1, \dots, \sigma_{n-1} \rangle; \text{ (free monoid of } n\text{-strand (positive) braid presentations)} \\Br_n^+ := FBr_n^+ / (\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \forall i; \sigma_i \sigma_j = \sigma_j \sigma_i, \forall |i - j| > 1); \text{ (n-strand positive braids)} \\s : Br_n^+ \to Br_n^+ / (\sigma_i^2 = 1, \forall i) \cong S_n : \sigma_k \mapsto s_k := s(\sigma_k) = (k + 1). \text{ (underlying permuations)}$$

Convention 2: As in Figure 1.3.1, any $\beta = \sigma_{i_{\ell}} \cdots \sigma_{i_1} \in FBr_n^+$ is represented by the *braid diagram* $[\sigma_{i_1}|\cdots|\sigma_{i_{\ell}}]$ going *from left to right*, where $[\beta_1|\beta_2]$ is the concatenation of β_1 with β_2 . Label the left (resp. right) ends *from bottom to top* by 1, 2, ..., *n*. Denote $[n] := \{1, 2, ..., n\}$.



FIGURE 1.3.1. Braid diagram for a positive braid: n = 3.

Definition 1.5. Let $e_{i,j} \in M_{n \times n}(\mathbb{K})$ be the matrix so that $(e_{i,j})_{a,b} = \delta_{a,i}\delta_{b,j}$. Define

$$\begin{split} \mathbf{K}_{k}(\epsilon) &:= \sum_{i \neq k} e_{i,i} + \epsilon e_{k,k} \in G, \ 1 \leq k \leq n, \epsilon \in \mathbb{K}^{\times}; \quad [\mathbf{K}_{k}(\epsilon)] := \text{ Figure 1.3.2 (left).} \\ \mathbf{H}_{i,j}(\epsilon) &:= I_{n} + \epsilon e_{i,j} \in G, \ 1 \leq i < j \leq n, \epsilon \in \mathbb{K}; \quad [\mathbf{H}_{i,j}(\epsilon)] := \text{ Figure 1.3.2 (middle).} \end{split}$$

Denote $H_k(\epsilon) := H_{k,k+1}(\epsilon)$. Define $[s_k] := \sigma_k =$ Figure 1.3.2 (right), for $1 \le k \le n-1$. Each of $[K_k(\epsilon)]$, $[H_{i,j}(\epsilon)]$, $[s_k]$ is called an *elementary braid matrix diagram* of rank *n* over \mathbb{K} .

n n	n n	m - n
$e \stackrel{\epsilon}{\underset{k}{\overset{\ldots}{\overset{\ldots}{\overset{\ldots}{\overset{\ldots}{\overset{\ldots}{\overset{\ldots}{\overset{\ldots}{\overset$	$ \begin{array}{c c} j & \vdots \\ \epsilon & \vdots \\ i & \vdots \\ i & i \end{array} $	k+1 $k+1$ $k+1$ k
1 1	1 1	1 1
$[\mathbf{K}_{k}(\boldsymbol{\epsilon})]$	$[\mathrm{H}_{i,i}(\epsilon)]$	$[s_k] = \sigma_k$

FIGURE 1.3.2. Elementary braid matrix diagrams representing elementary matrices: $[K_k(\epsilon)]$ (*scaling*), $[H_{i,j}(\epsilon)]$ for i < j (*handleslide*), $[s_k]$ (*transposition*).

Definition 1.6 (Braid matrix diagram presentations).

(1) The monoid of *braid matrix diagram presentations* (**bmdp**) of rank *n* over \mathbb{K} is:

 $\operatorname{FBD}_n := \langle \sigma_1, \dots, \sigma_{n-1}, [\mathbf{K}_k(\epsilon)], [\mathbf{H}_{i,j}(\epsilon')] \rangle / ([\mathbf{K}_k(1)] = \operatorname{id}_n = [\mathbf{H}_{i,j}(0)]), \quad \Gamma_2 \circ \Gamma_1 := [\Gamma_1 | \Gamma_2].$

So, a bmdp can be viewd as a finite concatenation of elementary braid matrix diagrams.(2) Define a morphism of monoids

$$g_{-}: \operatorname{FBD}_{n} \to G: \sigma_{k} \mapsto s_{k}, \ [\operatorname{K}_{k}(\epsilon)] \mapsto \operatorname{K}_{k}(\epsilon), \ [\operatorname{H}_{i,j}(\epsilon')] \mapsto \operatorname{H}_{i,j}(\epsilon').$$

Two **bmdp**'s $\Gamma_1, \Gamma_2 \in \text{FBD}_n$ are *weakly equivalent* if $g_{\Gamma_1} = g_{\Gamma_2}$, denoted as $\Gamma_1 \stackrel{\text{w}}{\sim} \Gamma_2$.

Lemma 1.7 (*Elementary moves* of **bmdp**'s). *Denote* $i' := s_k(i)$, then

$$\mathbf{K}_{k}(\epsilon_{1})\circ\mathbf{K}_{\ell}(\epsilon_{2}) = \begin{cases} \mathbf{K}_{k}(\epsilon_{1}\epsilon_{2}) & k = \ell, \\ \mathbf{K}_{\ell}(\epsilon_{2})\circ\mathbf{K}_{k}(\epsilon_{1}) & k \neq \ell. \end{cases} \quad \mathbf{H}_{i,j}(\epsilon_{1})\circ\mathbf{K}_{k}(\epsilon_{2}) = \begin{cases} \mathbf{K}_{i}(\epsilon_{2})\circ\mathbf{H}_{i,j}(\epsilon_{1}^{-1}\epsilon_{1}) & k = i, \\ \mathbf{K}_{i}(\epsilon_{2})\circ\mathbf{H}_{i,j}(\epsilon_{1}\epsilon_{2}) & k = j, \\ \mathbf{K}_{i}(\epsilon_{2})\circ\mathbf{H}_{i,j}(\epsilon_{1}) & k \neq i, j \end{cases}$$

$$s_{k} \circ \mathbf{K}_{i}(\epsilon) = \mathbf{K}_{i'}(\epsilon) \circ s_{k}; \\ s_{k} \circ \mathbf{H}_{i,j}(\epsilon) = \mathbf{H}_{i',j'}(\epsilon) \circ s_{k}. \quad \mathbf{H}_{i,j}(\epsilon_{1}) \circ \mathbf{H}_{k,\ell}(\epsilon_{2}) = \begin{cases} \mathbf{H}_{i,j}(\epsilon_{1}+\epsilon_{2}) & (k,\ell) = (i,j), \\ \mathbf{H}_{j,\ell}(\epsilon_{2}) \circ \mathbf{H}_{i,\ell}(\epsilon_{1}\epsilon_{2}) \circ \mathbf{H}_{i,j}(\epsilon_{1}) & k = j, \\ \mathbf{H}_{k,i}(\epsilon_{2}) \circ \mathbf{H}_{k,j}(-\epsilon_{2}\epsilon_{1}) \circ \mathbf{H}_{i,j}(\epsilon_{1}) & \ell = i, \\ \mathbf{H}_{k,\ell}(\epsilon_{2}) \circ \mathbf{H}_{i,j}(\epsilon_{1}) & \ell = i, \end{cases}$$

Each identity is either trivial (commutative), or represented by an (elementary) move in Figure 1.3.3.

Proof. Let e_1, \dots, e_n be the *standard basis* for \mathbb{K}^n . It's direct to check that, both sides of each identity applied to every e_i , give the same results. See also Figure 1.3.3 for an illustration.



FIGURE 1.3.3. *Elementary moves* for **bmdp**'s: The trivial ones are skipped. Each move is a weak equivalence in FBD_n representing a composition identity in Lemma 1.7, and vice versa. The composition goes from left to right: $\Gamma_2 \circ \Gamma_1 = [\Gamma_1 | \Gamma_2]$.

Definition 1.8 (Braid matrix diagrams).

(1) Let <u>FBD</u>, be the quotient of FBD_n by elementary moves. Then \exists a morphism of monoids

$$\beta_{-}: \underline{\operatorname{FBD}}_{n} \to \operatorname{FBr}_{n}^{+}: \sigma_{k} \mapsto \sigma_{k}, \ [\operatorname{K}_{k}(\epsilon)] \mapsto \operatorname{id}_{n}, \ [\operatorname{H}_{i,j}(\epsilon')] \mapsto \operatorname{id}_{n}$$

(2) The monoid BD_n of rank *n* braid matrix diagrams (**bmd**) is the quotient of <u>FBD</u>_n by usual braid relations/moves. So, BD_n = FBD_n/ $\stackrel{b}{\sim}$, with $\stackrel{b}{\sim}^1$ generated by elementary and braid moves. (3) $\beta_- : \underline{FBD}_n \to FBr_n^+$ and $g_- : FBD_n \to G$ induce morphisms of monoids:

$$\beta_-: \mathrm{BD}_n \to \mathrm{Br}_n^+, \quad g_-: \mathrm{BD}_n \to G$$

Remark 1.9. We have natural morphisms of monoids

$$i: \operatorname{FBr}_n^+ \to \operatorname{\underline{FBD}}_n : \sigma_k \mapsto \sigma_k, \quad \rightsquigarrow \quad i: \operatorname{Br}_n^+ \to \operatorname{BD}_n : \sigma_k \mapsto \sigma_k.$$

 $\beta_- \circ i = id$, so *i* induces embeddings $FBr_n^+ \hookrightarrow \underline{FBD}_n$ and $Br_n^+ \hookrightarrow BD_n$. This morally explains the terminology. Altogether, we get a commutative diagram of monoids:

Next, s : $Br_n^+ \rightarrow S_n$ admits a *canonical section* (as a map of sets) characterized by:

 $[-]: S_n \to \operatorname{Br}_n^+ \subset \operatorname{BD}_n : w \mapsto [w], \text{ with } \operatorname{s}([w]) = w, \ell([w]) = \ell(w).$ (1.3.1)

As we have seen, $[s_k] = \sigma_k$. Up to a choice, we may assume $[w] \in FBr_n^+$. By definition,

$$[w_1] \circ [w_2] = [w_1 w_2] \in \operatorname{Br}_n^+ \Leftrightarrow \ell(w_1 w_2) = \ell(w_1) + \ell(w_2).$$
(1.3.2)

Also, for $B \subset G$, there is a canonical *morphism of monoids* extending Definition 1.5:

$$[-]: B \to \underline{\text{FBD}}_n : b \mapsto [b] \text{ with } g_{[b]} = b.$$
(1.3.3)

¹Term $\stackrel{b}{\sim}$ as braid equivalence. Clearly, $\stackrel{b}{\sim}$ implies $\stackrel{w}{\sim}$. On the other hand, $\sigma_k^2 \stackrel{w}{\sim} id_n$, but $\sigma_k^2 \stackrel{b}{\sim} id_n$.

Proposition 1.10. The map g_- : BD_n \rightarrow G has a canonical section (as a map of sets)

$$[-]: G = \sqcup_{w \in W} B w B \to BD_n : x = b_1 w b_2 \mapsto [x] := [b_1] \circ [w] \circ [b_2]$$

Up to a choice $[w] \in FBr_n^+$, $[x] \in \underline{FBD}_n$. Moreover, for any other $x' = b'_1w'b'_2 \in Bw'B$,

$$[x] \circ [x'] = [xx'] \in \mathrm{BD}_n \Leftrightarrow \ell(ww') = \ell(w) + \ell(w') \Leftrightarrow \mathrm{inv}(ww') = w'^{-1}(\mathrm{inv}(w)) \sqcup \mathrm{inv}(w').$$

In this case, we obtain unique decompositions for $xx' \in Bww'B = Bww'U_{ww'}^-$:

$$Bww'U_{ww'}^{-} = BwU_{w}^{-}w'U_{w'}^{-} = U_{w^{-1}}^{-}wBw'U_{w'}^{-} = U_{w^{-1}}^{-}wU_{w'^{-1}}^{-}w'B = U_{w'^{-1}w^{-1}}^{-}ww'B.$$
(1.3.4)

Convention 3: Often, for $[A] \in BD_n$, we use the same notation for its lift in FBD_n or <u>FBD_n</u>. Before giving the proof, we discuss some linear algebra. Denote

$$I := \{(i, j) : 1 \le i < j \le n\}$$

Then $U = I_n + \sum_{(i,j) \in I} \mathbb{K}e_{i,j} \subset B \subset G$. We say a subset $\mathcal{J} \subset I$ is *multiplicative*, if

$$(i, j), (j, k) \in \mathcal{J} \Rightarrow (i, k) \in \mathcal{J}$$

For each $m \ge 1$, denote $\mathcal{J}_1 := \mathcal{J}$ and in general:

$$\mathcal{J}_m := \{ (j_0, \cdots, j_m) : (j_k, j_{k+1}) \in \mathcal{J}, \forall 0 \le k \le m-1 \}.$$

Lemma 1.11. Let $\mathcal{J} \subset I$ be multiplicative. Denote $U_{\mathcal{J}} := I_n + \sum_{(i,j) \in \mathcal{J}} \mathbb{K}e_{i,j} \subset U$. Then

(1) $U_{\mathcal{J}} \subset U$ is a closed subgroup.

(2) Any fixed total order \leq on \mathcal{J} induces an isomorphism of \mathbb{K} -varieties

$$\phi_{\leq} : \mathbb{A}^{|\mathcal{J}|} \to U_{\mathcal{J}} : (\epsilon_{i,j})_{(i,j) \in \mathcal{I}} \mapsto \prod_{(i,j) \in \mathcal{J}} \mathbf{H}_{i,j}(\epsilon_{i,j})$$

Proof. We prove (2). (1) is similar. Say, $\mathcal{J} = \{(i_1, j_1) < \dots < (i_N, j_N)\}$. Then

$$\prod_{(i,j)\in\mathcal{J}} H_{i,j}(\epsilon_{i,j}) = \prod_{\ell=1}^{N} (I_n + \epsilon_{i_\ell,j_\ell} e_{i_\ell,j_\ell}) = I_n + \sum_{m\geq 1} \sum_{(k_0,k_1)<\cdots<(k_{m-1},k_m) \text{ in }\mathcal{J}} \epsilon_{k_0,k_1}\cdots\epsilon_{k_{m-1},k_m} e_{k_0,k_m}.$$

The equation $I_n + \sum_{(i,j) \in \mathcal{J}} a_{ij} e_{i,j} = \prod_{(i,j) \in \mathcal{J}} H_{i,j}(\epsilon_{i,j})$ becomes

$$a_{i,j} = \epsilon_{i,j} + \sum_{m \ge 2} \sum_{(i,k_1) < \cdots < (k_{m-1},j) \text{ in } \mathcal{J}} \epsilon_{i,k_1} \cdots \epsilon_{k_{m-1},j}, \quad \forall (i,j) \in \mathcal{J}.$$

This uniquely determines the $\epsilon_{i,j}$'s inductively, in the increasing (partial) order on |j - i|.

For any permuation $w \in W = S_n$, denote

$$inv(w) := \{(i, j) \in \mathcal{I} : i < j, w(i) > w(j)\}; \quad noinv(w) := \{(i, j) \in \mathcal{I} : i < j, w(i) < w(j)\}.$$

Then $\ell(w) = |inv(w)|$, and inv(w), noinv(w) are multiplicative subsets of I. So,

$$U_w^+ := U_{\text{noinv}(w)} = \text{id} + \sum_{(i,j) \in \text{noinv}(w)} \mathbb{K}e_{i,j}; \quad U_w^- := U_{\text{inv}(w)} = \text{id} + \sum_{(i,j) \in \text{inv}(w)} \mathbb{K}e_{i,j},$$

are closed subgroups of U, to which Lemma 1.11 apply. Alternatively, we have

$$U_{w}^{+} = U \cap w^{-1}Uw = w^{-1}U_{w^{-1}}^{+}w, \quad U_{w}^{-} = U \cap w^{-1}U^{-}w,$$
(1.3.5)

with $U^- \subset G$ the opposite unipotent subgroup. By Lemma 1.11 (2), we have decompositions:

$$U = U_w^+ U_w^- = U_w^- U_w^+; \quad BwB = U_{w^{-1}}^- wB = BwU_w^-.$$
(1.3.6)

Proof of Proposition 1.10. Clearly, [-] is well-defined, $g_- \circ [-] = id$. To show equivalences. "LHS \Rightarrow Middle": Say, $xx' \in B\widetilde{w}B$, $\widetilde{w} \in S_n$, so $\beta_{[xx']} = [\widetilde{w}]$. If $[x] \circ [x'] = [xx']$, then

$$[\widetilde{w}] = \beta_{[xx']} = \beta_{[x]} \circ \beta_{[x']} = [w] \circ [w'], \quad \widetilde{w} = \mathrm{s}(\beta_{[xx']}) = \mathrm{s}(\beta_{[x]}) \circ \mathrm{s}(\beta_{[x']}) = ww'.$$

Thus, $\ell(ww') = \ell([ww']) = \ell([w] \circ [w']) = \ell([w]) + \ell([w']) = \ell(w) + \ell(w')$, as desired. "Middle \Rightarrow RHS": Denote

$$\begin{aligned} & (a',b') \coloneqq (w'(a),w'(b)), \quad (a'',b'') \coloneqq (w(a'),w(b')), \quad 1 \le a < b \le n; \\ & I_{+-} \coloneqq \{(a,b) \in I : a' < b',a'' > b''\}; \quad I_{-+} \coloneqq \{(a,b) \in I : a' > b',a'' > b''\}. \end{aligned}$$

Observe that $inv(ww') = I_{+-} \sqcup I_{-+}$ and w' induces bijections

$$w': I_{+-} \xrightarrow{\simeq} \operatorname{noinv}(w'^{-1}) \cap \operatorname{inv}(w); \quad R \circ w': I_{-+} \xrightarrow{\simeq} \operatorname{inv}(w'^{-1}) \cap \operatorname{noinv}(w), \quad R(i, j) := (j, i);$$

$$w' \sqcup R \circ w': \operatorname{inv}(ww') = I_{+-} \sqcup I_{-+} \xrightarrow{\simeq} \operatorname{noinv}(w'^{-1}) \cap \operatorname{inv}(w) \sqcup \operatorname{inv}(w'^{-1}) \cap \operatorname{noinv}(w).$$

Hence,

 $\ell(ww') = |\operatorname{noinv}(w'^{-1}) \cap \operatorname{inv}(w)| + |\operatorname{inv}(w'^{-1}) \cap \operatorname{noinv}(w)| \le |\operatorname{inv}(w)| + |\operatorname{inv}(w'^{-1})| = \ell(w) + \ell(w'),$ with equality holds if and only if $\operatorname{inv}(w) \cap \operatorname{inv}(w'^{-1}) = \emptyset$. Then, $I_{+-} = w'^{-1}(\operatorname{inv}(w))$, and $I_{-+} = (R \circ w')^{-1}(\operatorname{inv}(w'^{-1})) = \operatorname{inv}(w')$. Hence, $\operatorname{inv}(ww') = w'^{-1}(\operatorname{inv}(w)) \sqcup \operatorname{inv}(w')$, as desired.

"RHS
$$\Rightarrow$$
 Middle" is clear, so RHS \Leftrightarrow Middle.

"Middle
$$\Rightarrow$$
 LHS": By above, noinv(w) \cup noinv(w'⁻¹) = *I*. By Lemma 1.11, we get a surjection

$$m: U_w^+ \times U_{w'^{-1}}^+ \to U: (u_1, u_2) \mapsto u_1 u_2.$$

By (1.3.6), can assume $b_2 \in U_w^-$, $b_1' \in U_{w'^{-1}}^-$. By above, we may write $b_2b_1' = u_1u_2$, $u_1 \in U_w^+$, $u_2 \in U_{w'^{-1}}^+$. So, $xx' = (b_1wu_1w^{-1})ww'(w'^{-1}u_2w'b_2') \in Bww'B$, and

$$[xx'] = [b_1wu_1w^{-1}] \circ [ww'] \circ [w'^{-1}u_2w'b'_2] \in BD_n \quad \text{(by definition)}$$

= $[b_1] \circ [wu_1w^{-1}] \circ [w] \circ [w'] \circ [w'^{-1}u_2w'] \circ [b'_2] \quad \text{(by (1.3.2), (1.3.3))}$
= $[b_1] \circ ([w] \circ [u_1]) \circ ([u_2] \circ [w']) \circ [b'_2] \quad \text{(by elementary moves as in Figure 1.3.3.(8))}$
= $[b_1] \circ [w] \circ [b_2] \circ [b'_1] \circ [w'] \circ [b'_2] = [x] \circ [x']. \quad \text{(by (1.3.3))}$

It remains to show the decompositions. By Lemma 1.11, we get an isomorphism

$$m: U_{w'^{-1}(inv(w))} \times U_{inv(w')} = w'^{-1}U_w^{-}w' \times U_{w'}^{-} \to U_{inv(ww')} = U_{ww'}^{-}: (u_1, u_2) \mapsto u_1 u_2$$

So is $m: U_{w'}^- \times w'^{-1} U_w^- w' \to U_{ww'}^-$. Then, $U_{ww'}^- = (w'^{-1} U_w^- w') U_{w'}^- = U_{w'}^- (w'^{-1} U_w^- w')$. Also, the same result applies to $(ww')^{-1} = w'^{-1} w^{-1}$. Altogether, we get unique decompositions

$$Bww'B = Bww'U_{ww'}^{-} = BwU_{w}^{-}w'U_{w'}^{-} = U_{w^{-1}}^{-}wBw'U_{w'}^{-} = U_{w^{-1}}^{-}wU_{w'^{-1}}^{-}w'B = U_{w'^{-1}w^{-1}}^{-}ww'B.$$

Back to character varieties. Let's reinterpret $M'_B(\vec{w})$ (see (1.2.7)) via braid matrix diagrams. First, we set up some notations. We have seen (unique) decompositions:

$$U = U_w^+ U_w^- : u = L_w^+(u) L_w^-(u), \quad B = TU = (TU_w^+) U_w^- : b = D(b) b_R = L_w^+(b) L_w^-(b); \quad (1.3.7)$$

$$U = U_w^- U_w^+ : u = R_w^-(u) R_w^+(u), \quad B = UT = (U_w^-) (U_w^+ T) : b = b_L D(b) = R_w^-(b) R_w^+(b).$$

Similarly, each parabolic $P_i \subset G$ has decompositions

$$P_i = N_i Z(C_i) = Z(C_i) N_i : p_i = L_i(p_i) D_i(p_i) = D_i(p_i) R_i(p_i), \quad N_i := P_i \cap U.$$
(1.3.8)
As the shortest representative of $w_i \in W/W(C_i)$, $\dot{w_i}$ gives

$$U^-_{\dot{w}_i} \subset N_i, \quad Z(C_i) \cap U \subset U^+_{\dot{w}_i}.$$

Thus, by Lemma 1.11, we obtain decompositions

$$N_{i} = U_{\dot{w}_{i}}^{-}(U_{\dot{w}_{i}}^{+} \cap N_{i}) = (U_{\dot{w}_{i}}^{+} \cap N_{i})U_{\dot{w}_{i}}^{-}; \ U_{\dot{w}_{i}}^{+} = (U_{\dot{w}_{i}}^{+} \cap N_{i})(Z(C_{i}) \cap U) = (Z(C_{i}) \cap U)(U_{\dot{w}_{i}}^{+} \cap N_{i}). \ (1.3.9)$$
Now, we reinterpret $M'_{i}(\vec{w})$. To begin with we have a capacital isomerphism.

Now, we reinterpret $M'_B(\vec{w})$. To begin with, we have a canonical isomorphism

$$U_{\dot{w}_{i}^{-1}}^{-} \times N_{i} \times Z(C_{i}) \cong U_{\dot{w}_{i}^{-1}}^{-} \dot{w}_{i} P_{i} = B \dot{w}_{i} P_{i} : (\nu_{i}, n_{i}, z_{i}) \mapsto x_{i} = \nu_{i} \dot{w}_{i} n_{i} z_{i}.$$
(1.3.10)

Then, $x_i C_i x_i^{-1} = v_i \dot{w}_i n_i C_i n_i^{-1} \dot{w}_i^{-1} v_i^{-1} = v_i \dot{w}_i n_i' C_i \dot{w}_i^{-1} v_i^{-1}$. Here, we use the isomorphism

$$N_i \xrightarrow{\cong} N_i : n_i \mapsto n'_i = n_i C_i n_i^{-1} C_i^{-1}.$$
(1.3.11)

Recall that for $b \in B$, $[b] \in \underline{FBD}_n$ (see (1.3.3)), $\beta_{[b]} = id_n \in Br_n^+$ and $g_{[b]} = b$. Define

$$[x_i C_i x_i^{-1}]' := [v_i] \circ [\dot{w}_i] \circ [n'_i] \circ [C_i] \circ [\dot{w}_i^{-1}] \circ [v_i^{-1}] = [v_i^{-1} |\dot{w}_i^{-1}| C_i |n'_i |\dot{w}_i |v_i] \in \underline{\text{FBD}}_n.$$
(1.3.12)
Recall that $A_i \in B\tau_i B$, with Proposition 1.10 in mind, define $[M_{\vec{w}}] = [M_{\vec{w}}((A_i)_i, (x_i)_i, u_k)]$ by

$$[\mathbf{M}_{\vec{w}}] := \prod_{j=1}^{g} ([A_{2j-1}] \circ [A_{2j}] \circ [A_{2j-1}^{-1}] \circ [A_{2j}^{-1}]) \circ \prod_{i=1}^{k-1} ([x_i C_i x_i^{-1}]') \circ [u_k C_k] \in \underline{\operatorname{FBD}}_n.$$
(1.3.13)

Then by (1.2.7) and (1.2.4), the defining equation for $M'_B(\vec{w})$ reads

$$[\mathbf{M}_{\vec{w}}] = [\mathbf{M}_{\vec{w}}((A_j)_{j=1}^{2g}, (x_i)_{i=1}^{k-1}, u_k)] \stackrel{\text{w}}{\sim} \mathsf{id}_n \in \underline{\mathrm{FBD}}_n,$$
(1.3.14)

where $[M_{\vec{w}}]$ is a **bmdp** (with varying coefficients) but with fixed *shape*:

$$\beta(\vec{w}) := \beta_{[\mathbf{M}_{\vec{w}}]} = \prod_{j=1}^{8} ([\tau_{2j-1}] \circ [\tau_{2j}] \circ [\tau_{2j-1}^{-1}] \circ [\tau_{2j}]) \circ \prod_{i=1}^{k-1} ([\vec{w}_i] \circ [\vec{w}_i^{-1}]) \in \mathrm{FBr}_n^+.$$
(1.3.15)

The next idea is to canonicalize $[M_{\vec{w}}]$ by diagram calculus: push every handleslide or scaling in $[M_{\vec{w}}] = [C_k |u_k| [x_{k-1}C_{k-1}x_{k-1}^{-1}]' |\cdots |A_2^{-1}|A_1^{-1}|A_2|A_1]$ via elementary moves, to the right as far as possible. Later on, we'll see this leads to braid varieties.

1.4. Braid varieties. As already mentioned above, we now review some basics on braid varieties.

Definition 1.12. The braid matrix (resp. **bmdp**) with coefficient $\epsilon \in \mathbb{K}$ associated to σ_k is

$$B_k(\epsilon) := s_k H_k(\epsilon) \in G; \quad [B_k(\epsilon)] = \sigma_k \circ [H_k(\epsilon)] \in FBD_n \text{ (Figure 1.4.1)}. \tag{1.4.1}$$

For $\beta = \sigma_{i_{\ell}} \cdots \sigma_{i_1} \in FBr_n^+$, and $\vec{\epsilon} = (\epsilon_i)_{i=\ell}^1 \in \mathbb{A}^{\ell}$, define

$$\mathbf{B}_{\beta}(\vec{\epsilon}) \coloneqq \mathbf{B}_{i_{\ell}}(\epsilon_{\ell}) \cdots \mathbf{B}_{i_{1}}(\epsilon_{1}) \in G; \quad [\mathbf{B}_{\beta}(\vec{\epsilon})]' \coloneqq [\mathbf{B}_{i_{\ell}}(\epsilon_{\ell})] \circ \cdots \circ [\mathbf{B}_{i_{1}}(\epsilon_{1})] \in \mathrm{FBD}_{n}. \tag{1.4.2}$$



FIGURE 1.4.1. The braid matrix diagram with coefficient ϵ associated to σ_k .

As an easy application of diagram calculus of matrices, we obtain the following:

Lemma 1.13 (Braid relations for braid matrices). We have

(a) $[B_i(\epsilon_1)] \circ [B_i(\epsilon_2)] = [B_i(\epsilon_2)] \circ [B_i(\epsilon_1)] \in BD_n, |i-j| > 1, \epsilon_{\bullet} \in \mathbb{K};$

 $(b) \qquad [\mathsf{B}_{i}(\epsilon_{1})] \circ [\mathsf{B}_{i+1}(\epsilon_{2})] \circ [\mathsf{B}_{i}(\epsilon_{3})] = [\mathsf{B}_{i+1}(\epsilon_{3})] \circ [\mathsf{B}_{i}(\epsilon_{2} - \epsilon_{3}\epsilon_{1})] \circ [\mathsf{B}_{i+1}(\epsilon_{1})] \in \mathsf{BD}_{n}, \epsilon_{\bullet} \in \mathbb{K}.$

Proof. (*a*) is clear. We prove (*b*) by Figure 1.4.2: (*i*) is a composition of elementary moves Figure 1.3.3 (8) and a trivial move switching ϵ_1, ϵ_2 ; (*ii*) is a composition of an elementary move Figure 1.3.3 (7), a trivial move switching $-\epsilon_3\epsilon_1, \epsilon_3$, and an elementary move Figure 1.3.3 (5); (*iii*) is a braid move; (*iv*) is a composition of elementary moves Figure 1.3.3 (8).



FIGURE 1.4.2. Braid relation for braid matrix diagrams with coefficients.

Definition 1.14 (Braid varieties). Let $\beta = \sigma_{i_{\ell}} \cdots \sigma_{i_1} \in FBr_n^+$. For each $1 \le j \le \ell$, denote $f_j : \mathbb{A}^j \to G : (\epsilon_j, \cdots, \epsilon_1) \mapsto B_{i_j}(\epsilon_j) \cdots B_{i_1}(\epsilon_1).$ (1.4.3) (1) The (*resp. restricted*) braid variety associated to β is a closed subvariety of \mathbb{A}^{ℓ} :

$$X(\beta) := f_{\ell}^{-1}(B); \quad \text{resp. } X(\beta, C) := \mu \mathfrak{mon}^{-1}(C), C \in T, \quad \mu \mathfrak{mon} := D \circ f_{\ell} : X(\beta) \to B \to T. \quad (1.4.4)$$

By Lemma 1.13, $X(\beta)$ (resp. $X(\beta, C)$) depends only on $\beta \in Br_n^+$, up to a canonical isomorphism. (2) $b \in B$ acts on $X(\beta) \in \vec{\epsilon}$ as follows: $\hat{\vec{\epsilon}} = (\hat{\epsilon}_{\ell}, \dots, \hat{\epsilon}_1) := b \cdot \vec{\epsilon}$ is uniquely determined by

$$[\mathbf{B}_{i_{\ell}}(\epsilon_{\ell})] \circ \dots \circ [\mathbf{B}_{i_{1}}(\epsilon_{1})] \circ [b^{-1}] = [\tilde{b}_{\ell}] \circ [\mathbf{B}_{i_{\ell}}(\widehat{\epsilon}_{\ell})] \circ \dots \circ [\mathbf{B}_{i_{1}}(\widehat{\epsilon}_{1})] \in \underline{\mathrm{FBD}}_{n}, \tilde{b}_{\ell} \in B.$$
(1.4.5)

That is, in $[b^{-1}|B_{i_1}(\epsilon_1)|\cdots|B_{i_\ell}(\epsilon_\ell)]$, push $[b^{-1}]$ by elementary moves (Figure 1.3.3) to the right as far as possible, the outcome is $[B_{i_1}(\hat{\epsilon}_1)|\cdots|B_{i_\ell}(\hat{\epsilon}_\ell)|\tilde{b}_\ell]$.

Convention 4: If the context is clear, for a group action of $h \in H$ on any variety $Y \ni y$, denote

$$\widehat{\mathbf{y}} := h \cdot \mathbf{y}. \tag{1.4.6}$$

Next, we recall the cell decomposition of braid varieties. Fix $\beta = \sigma_{i_{\ell}} \cdots \sigma_{i_1} \in FBr_n^+$. Define

$$p: \mathbb{A}^{\ell} \to W^{\ell+1}: \vec{\epsilon} \mapsto p(\vec{\epsilon}) = (p_{\ell}, \dots, p_0), \quad \mathbf{B}_{i_m}(\epsilon_m) \cdots \mathbf{B}_{i_1}(\epsilon_1) \in Bp_m B.$$
(1.4.7)

Alternatively by Proposition 1.10: $[B_{i_m}(\epsilon_m) \cdots B_{i_1}(\epsilon_1)] \in [B] \circ [p_m] \circ [B] \subset BD_n$.

Definition 1.15 ([40, §.5.4]). Let $p = (p_{\ell}, \dots, p_0) \in W^{\ell+1}$. If for any *position* $1 \le m \le \ell$:

$$p_m = \begin{cases} s_{i_m} p_{m-1} \text{ (go-up)} & \text{if } s_{i_m} p_{m-1} > p_{m-1}, \\ s_{i_m} p_{m-1} \text{ (go-down) or } p_{m-1} \text{ (stay)} & \text{if } s_{i_m} p_{m-1} < p_{m-1}. \end{cases}$$

and $p_0 = p_\ell = id$, we say p is a *walk* of β . Denote:

$$U_p := \{\text{go-up's}\}, \ S_p := \{\text{stays}\}, \ D_p := \{\text{go-down's}\}, \Rightarrow [\ell] = \{1, \dots, \ell\} = U_p \sqcup D_p \sqcup S_p.$$

By a length count: $|U_p| - |D_p| = \ell(p_\ell) - \ell(p_0) = 0$. Denote $\mathcal{W}(\beta) := \{ \text{walks of } \beta \}$.

For any $1 \le m \le \ell = \ell(\beta)$, denote

$$\mathbf{s}_{< m}(\beta) := \prod_{q=m-1}^{1} \mathbf{s}_{i_q}, \quad \mathbf{s}_{> m}(\beta) := \prod_{q=\ell}^{m+1} \mathbf{s}_{i_q}.$$
(1.4.8)

We use **Convention** 4. Recall that $x^y = yxy^{-1}$ in *G*, and we write $t = \text{Diag}(t_1, \dots, t_n) \in T$.

Proposition 1.16. We have B-equivariant decompositions into locally closed subvarieties:

$$\begin{split} X(\beta) &= \sqcup_{p \in \mathcal{W}(\beta)} X_p(\beta), \quad \varphi : X_p(\beta) \coloneqq X_p^{\ell}(\beta) \cong (\mathbb{K}^{\times})^{|S_p|} \times \mathbb{K}^{|U_p|} : \vec{\epsilon} \mapsto (\epsilon'_m)_{m \in S_p \sqcup U_p} (1.4.9) \\ X(\beta, C) &= \sqcup_{p \in \mathcal{W}(\beta)} X_p(\beta, C), \quad X_p(\beta, C) \coloneqq X_p(\beta) \cap X(\beta, C), \end{split}$$

such that

(1) The inherited action of $b \in B$ on $(\epsilon'_m)_{m \in S_p \sqcup U_p} \in (\mathbb{K}^{\times})^{|S_p|} \times \mathbb{K}^{|U_p|}$ satisfies:

- (a) If $b = u \in U \subset B$, then $\widehat{\epsilon}'_m = \epsilon'_m, \forall m \in S_p$.
- (b) If $b = t \in T \subset B$, then

$$\widetilde{\epsilon}'_{m} = (t^{p_{m-1}})_{i_{m}}(t^{p_{m-1}})_{i_{m}+1}^{-1}\epsilon'_{m}, \forall m \in S_{p} \sqcup U_{p}.$$
(1.4.10)

(2) $\mu mon : X_p(\beta) \to T$ is identified with

$$\mu \mathfrak{mon}((\epsilon'_m)_{m \in S_p \sqcup U_p}) = \mu \mathfrak{mon}((\epsilon'_m)_{m \in S_p}) = \prod_{m \in S_p} (\mathbf{K}_{i_m}(-\epsilon'_m) \mathbf{K}_{i_m+1}(\epsilon'_m))^{s_{>m}(\beta)}.$$
(1.4.11)

In particular, det($\mu \mathfrak{mon}((\epsilon'_m)_{m \in S_p \sqcup U_p})) = (-1)^{|S_p|}$.

Proof. By diagram calculus, the proof is straightforward. See Appendix A for the details.

2. Cell decomposition of character varieties

In this section, we prove a strong form (Theorem 2.10) of A. Mellit's cell decomposition for very generic character varieties [40]. This will be used to prove our main theorem 3.10.

Recall that we have obtained a decomposition (see (1.2.8), (1.2.7)):

$$\mathcal{M}_{\mu} = \bigsqcup_{\vec{w} \in W^{2g} \times \prod_{i=1}^{k-1} W/W(C_i)} \mathcal{M}_{\mu}(\vec{w}), \quad \mathcal{M}_{\mu}(\vec{w}) = M'_B(\vec{w})/PB_{\text{par}},$$

and the defining equation of $M'_{R}(\vec{w})$ has been interpreted via braid matrix diagrams as (1.3.14):

$$[\mathbf{M}_{\vec{w}}] = \prod_{j=1}^{g} ([A_{2j-1}] \circ [A_{2j}] \circ [A_{2j-1}^{-1}] \circ [A_{2j}^{-1}]) \circ \prod_{i=1}^{k-1} ([x_i C_i x_i^{-1}]') \circ [u_k] \circ [C_k] \stackrel{\text{w}}{\sim} \mathsf{id}_n \in \underline{\mathrm{FBD}}_n$$

To obtain the actual cell decomposition of \mathcal{M}_{μ} , we would like to decompose $\mathcal{M}_{\mu}(\vec{w})$ further. This amounts to decomposing $M'_B(\vec{w})$ equivariantly. For that, as mentioned in the end of Section 1.3, the next step is to canonicalize $[M_{\vec{w}}]$ by diagram calculus. This will be done in the next three subsections: Section 2.1 and Section 2.2 do the puncture and genus calculations respectively; Section 2.3 combines these local calculations to describe $M'_B(\vec{w})$ via braid varieties. Finally, in Section 2.4, we prove a strong form (Theorem 2.10) of the cell decomposition for \mathcal{M}_{μ} .

To simplify our computations, we do a trick as follows.

- Fix an embedding $S_n \hookrightarrow FBr_n^+ \subset FBD_n$ lifting $[-]: S_n \to Br_n^+$ (as a map of sets).
- Via Lemma 1.11, we also fix an embedding [-]: B = U ⋊ T → FBD_n (not as a morphism of monoids) lifting [-]: B → FBD_n.

Definition/Proposition 2.1. Fix $S \subset [N] = \{1, \dots, N\}$. Take Y_1, \dots, Y_N such that: $Y_i \subset B$ is a locally closed \mathbb{K} -subvariety for $i \notin S$, and Y_i is a single permutation² in S_n , for $i \in S$.

(1) For any \mathbb{K} -variety Z, define

$$[Y_1] \circ \dots \circ [Y_N] \times Z := \{ [y_1] \circ \dots \circ [y_N] \in FBD_n : y_i \in Y_i \} \times Z.$$

As a \mathbb{K} -variety, $[Y_1] \circ \cdots \circ [Y_N] \times Z$ is $\prod_{i \notin S} Y_i \times Z$. Define the obvious composition

$$E: [Y_1] \circ \dots \circ [Y_N] \times Z \to [Y_1] \circ \dots \circ [Y_N] \hookrightarrow FBD_n \to \underline{FBD}_n.$$

(2) If $\varphi : [Y_1] \circ \cdots \circ [Y_N] \times Z \to [Y'_1] \circ \cdots \circ [Y'_M] \times Z'$ is an isomorphism of \mathbb{K} -varieties respecting E, we say φ is **elementary**. In this case, we write

$$[Y_1] \circ \dots \circ [Y_N] \times Z \cong [Y'_1] \circ \dots \circ [Y'_M] \times Z'.$$

In particular, any elementary isomorphism respects the maps

$$g_{-}: [Y_{1}] \circ \dots \circ [Y_{N}] \times Z \to G = GL(n, \mathbb{K}) : ([y_{1}] \circ \dots \circ [y_{N}], z) \mapsto y_{1} \dots y_{N},$$

$$\beta_{-}: [Y_{1}] \circ \dots \circ [Y_{N}] \times Z \to FBr_{n}^{+}: ([y_{1}] \circ \dots \circ [y_{N}], z) \mapsto \beta_{[y_{1}]} \circ \dots \circ \beta_{[y_{N}]}.$$

(3) Clearly, we have the following elementary isomorphisms:

(a) Let $Y, Y_1, Y_2 \subset B$. If the multiplication induces an isomorphism $m : Y_1 \times Y_2 \xrightarrow{\cong} Y : (y_1, y_2) \mapsto y_1y_2$, then $[Y_1] \circ [Y_2] \cong [Y]$.

(b) If $Y_1 \subset Y_2 \subset B$ and Y_2 is a closed subgroup, then $[Y_1] \circ [Y_2] \cong [Y_2] \times Y_1 \cong [Y_2] \circ [Y_1]$.

(c) If $w \in S_n$ and $Y \subset U_w^+$, then $[w] \circ [Y] \cong [wYw^{-1}] \circ [w]$.

- (d) For any $C \in T$ and any $Y \subset B$, we have $[C] \circ [Y] \cong [CYC^{-1}] \circ [C]$.
- 2.1. **Diagram calculus for punctures.** Back to the end of Section 1.3. Recall (1.3.10), (1.3.12):

$$U_{\dot{w}_{i}^{-1}}^{-} \times N_{i} \times Z(C_{i}) \cong B\dot{w}_{i}P_{i} : (v_{i}, n_{i}, z_{i}) \mapsto x_{i} = v_{i}\dot{w}_{i}n_{i}z_{i}; \quad N_{i} \cong N_{i} : n_{i} \mapsto n_{i}' = n_{i}C_{i}n_{i}^{-1}C_{i}.$$
$$[x_{i}C_{i}x_{i}^{-1}]' = [v_{i}] \circ [\dot{w}_{i}] \circ [n_{i}'] \circ [C_{i}] \circ [\dot{w}_{i}^{-1}] \circ [v_{i}^{-1}] \in \underline{FBD}_{n}.$$

Using the notations in (1.3.7), define ${}^{+}n'_{i} := R^{+}_{\dot{w}_{i}}(n'_{i}) \in U^{+}_{\dot{w}_{i}} \cap N_{i}$ by (1.3.9).

²By a little abuse of notations, we identify $w \in S_n$ with $\{w\} \subset S_n$. Similarly, we identify $b \in B$ with $\{b\} \subset B$.

Lemma 2.2. For any $D_{i+1} \in T$, we have a natural isomorphism of \mathbb{K} -varieties of the form

$$B\dot{w}_i P_i \times U \xrightarrow{\simeq} U \times U^-_{\dot{w}_i} \times U^-_{\dot{w}_i^{-1}} \times (U^+_{\dot{w}_i} \cap N_i) \times Z(C_i) : (x_i, u_{i+1}) \mapsto (u_i, \xi'_i, \xi_i, {}^+n'_i, z_i)$$
(2.1.1)

such that

 $[x_i C_i x_i^{-1}] \circ [u_{i+1}] \circ [D_{i+1}] = [u_i] \circ [D_i] \circ [\dot{w}_i] \circ [\xi'_i] \circ [\psi_i^{-1}] \circ [\xi_i] \in \underline{FBD}_n, \exists ! D_i = C_i^{\dot{w}_i} D_{i+1} \in T.$ (2.1.2) Moreover, the above equation uniquely determines $(u_i, \xi'_i, \xi_i) \in U \times U^-_{\dot{w}_i} \times U^-_{\dot{w}_i^{-1}}.$

Note: Up to a canonical isomorphism, we have

$$[\dot{w}_i] \circ [\xi'_i] = [\mathbf{B}_{[\dot{w}_i]}(\xi'_i)] \in \underline{\mathrm{FBD}}_n, \quad [\dot{w}_i^{-1}] \circ [\xi_i] = [\mathbf{B}_{[\dot{w}_i^{-1}]}(\xi_i)] \in \underline{\mathrm{FBD}}_n.$$

This will pave the way to braid varieties.

Proof of Lemma 2.2. Observe that we have an isomorphism of K-varieties

$$B\dot{w}_iP_i\times U\xrightarrow{\simeq} U^-_{\dot{w}_i^{-1}}\times N_i\times Z(C_i)\times U: (x_i,u_{i+1})\mapsto (v_i,n_i,z_i,u_{i+1}':=v_i^{-1}u_{i+1}).$$

such that $[x_i C_i x_i^{-1}]' \circ [u_{i+1}] = [v_i] \circ [\dot{w}_i] \circ [C_i] \circ [\dot{w}_i^{-1}] \circ [u'_{i+1}] \in \underline{FBD}_n$. By our trick (Definition/Proposition 2.1), it suffices to compute $[U_{\dot{w}_i^{-1}}] \circ [\dot{w}_i] \circ [N_i] \circ [C_i] \circ [\dot{w}_i^{-1}] \circ [U] \circ [D_{i+1}] \ni [v_i] \circ [\dot{w}_i] \circ [n'_i] \circ [C_i] \circ [\dot{w}_i^{-1}] \circ [U] \circ [D_{i+1}] \equiv [v_i] \circ [\dot{w}_i] \circ [n'_i] \circ [C_i] \circ [\dot{w}_i^{-1}] \circ [U'_{i+1}] \circ [D_{i+1}]$:

$$\begin{split} & [U_{\dot{w}_{i}^{-1}}^{-1}] \circ [\dot{w}_{i}] \circ [N_{i}] \circ [C_{i}] \circ [\dot{w}_{i}^{-1}] \circ [U] \circ [D_{i+1}] \\ &\cong \quad [U_{\dot{w}_{i}^{-1}}^{-1}] \circ [\dot{w}_{i}] \circ [U_{\dot{w}_{i}}^{-1}] \circ [U_{\dot{w}_{i}}^{+} \cap N_{i}] \circ [C_{i}] \circ [\dot{w}_{i}^{-1}] \circ [U] \circ [D_{i+1}] (n_{i}' = {}^{-}n_{i}' \cdot {}^{+}n_{i}') \\ &\cong \quad [U_{\dot{w}_{i}^{-1}}^{-1}] \circ [\dot{w}_{i}] \circ [U_{\dot{w}_{i}}^{-1}] \circ [C_{i}] \circ [\dot{w}_{i}^{-1}] \circ [\dot{w}_{i}C_{i}^{-1}(U_{\dot{w}_{i}}^{+} \cap N_{i})C_{i}\dot{w}_{i}^{-1}] \circ [U] \circ [D_{i+1}] \\ &\cong \quad [U_{\dot{w}_{i}^{-1}}^{-1}] \circ [\dot{w}_{i}] \circ [U_{\dot{w}_{i}}^{-1}] \circ [C_{i}] \circ [\dot{w}_{i}^{-1}] \circ [U] \circ [D_{i+1}] \times (U_{\dot{w}_{i}}^{+} \cap N_{i}) (\text{direct factor} : {}^{+}n_{i}' \in U_{\dot{w}_{i}}^{+} \cap N_{i}) \\ &\cong \quad [U_{\dot{w}_{i}^{-1}}^{-1}] \circ [\dot{w}_{i}] \circ [U_{\dot{w}_{i}}^{-1}] \circ [C_{i}] \circ [\dot{w}_{i}^{-1}] \circ [U_{i}^{+} \circ [U_{\dot{w}_{i}^{-1}}^{-1}] \circ [D_{i+1}] \times (U_{\dot{w}_{i}}^{+} \cap N_{i}) (U = U_{\dot{w}_{i}^{-1}}^{+}U_{\dot{w}_{i}^{-1}}^{-1}) \\ &\cong \quad [U_{\dot{w}_{i}^{-1}}^{-1}] \circ [\dot{w}_{i}] \circ [U_{\dot{w}_{i}}^{-1}] \circ [U_{\dot{w}_{i}}^{+1}] \circ [U_{\dot{w}_{i}^{-1}}^{-1}] \circ [D_{i+1}] \times (U_{\dot{w}_{i}}^{+} \cap N_{i}) (U = U_{\dot{w}_{i}^{-1}}^{+}U_{\dot{w}_{i}^{-1}}^{-1}) \\ &\cong \quad [U_{\dot{w}_{i}^{-1}}^{-1}] \circ [\dot{w}_{i}] \circ [U_{\dot{w}_{i}}^{-1}] \circ [U_{\dot{w}_{i}}^{-1}] \circ [U_{\dot{w}_{i}^{-1}}^{-1}] \circ [D_{i+1}] \times (U_{\dot{w}_{i}}^{+} \cap N_{i}) (U = U_{\dot{w}_{i}^{+1}}^{+}U_{\dot{w}_{i}}^{-1}) \\ &\cong \quad [U_{\dot{w}_{i}^{-1}}^{-1}] \circ [\dot{w}_{i}] \circ [U_{\dot{w}_{i}}^{-1}] \circ [C_{i}] \circ [\dot{w}_{i}^{-1}^{-1}] \circ [U_{\dot{w}_{i}^{-1}}^{-1}] \circ [D_{i+1}] \times (U_{\dot{w}_{i}}^{+} \cap N_{i}) \\ &\cong \quad [U_{\dot{w}_{i}}^{-1}] \circ [U_{\dot{w}_{i}}^{-1}] \circ [U_{\dot{w}_{i}}^{-1}] \circ [U_{\dot{w}_{i}^{-1}}^{-1}] \circ [D_{\dot{w}_{i}^{-1}}^{-1}] \circ [D_{i+1}] \times (U_{\dot{w}_{i}}^{+} \cap N_{i}) \\ &\cong \quad [U] \circ [\dot{w}_{i}] \circ [U_{\dot{w}_{i}}^{-1}] \circ [\dot{w}_{i}^{-1}^{-1}] \circ [U_{\dot{w}_{i}^{-1}}^{-1}] \times [U_{\dot{w}_{i}}^{-1}] \wedge [U_{\dot{w}_{i}}^{-1}] \times [U_{\dot{w}_{i}}^{-} \cap N_{i}) . \\ &\cong \quad [U] \circ [D_{i}] \circ [\dot{w}_{i}] \circ [U_{\dot{w}_{i}}^{-1}] \circ [\dot{w}_{i}^{-1}^{-1}] \circ [U_{\dot{w}_{i}^{-1}}^{-1}] \times (U_{\dot{w}_{i}}^{+} \cap N_{i}) . \\ &= \quad [U] \circ [D_{i}] \circ [\dot{w}_{i}] \circ [U_{\dot{w}_{i}}^{-1}] \circ [\dot{w}_{i}^{-1}^{-1}] \circ [U_{\dot{w}_{i}}^{-1}] \times [U$$

The uniqueness part is clear by definition of FBD_n. This finishes the proof.

The upshot is that, Lemma 2.2 provides the inductive step for the diagram calculus for punctures. In our case, we have seen $u_k \in U$, take $D_k := C_k$, and define D_i 's inductively as above. Thus,

$$D_1 = (\prod_{i=1}^{k-1} C_i^{\dot{w}_i}) C_k \in T.$$
(2.1.3)

Altogether, the defining equation (1.3.14) for $M'_B(\vec{w})$ reduces to:

$$[\mathbf{M}_{\vec{w}}] = \prod_{j=1}^{g} ([A_{2j-1}] \circ [A_{2j}] \circ [A_{2j-1}^{-1}] \circ [A_{2j}]) \circ [u_1] \circ [D_1] \circ \prod_{i=1}^{k-1} ([\dot{w}_i \xi'_i] \circ [\dot{w}_i^{-1} \xi_i]) \stackrel{\mathfrak{w}}{\sim} \mathsf{id}_n.$$
(2.1.4)

For the diagram calculus for genera below, denote $u^g := u_1 \in U$, $D^g := D_1 \in T$.

2.2. **Diagram calculus for genera.** For each $1 \le j \le g$, take the isomorphisms

$$U_{\tau_{2j-1}^{-1}}^{-} \times T \times U \cong B\tau_{2j-1}B : (\mu_{2j-1}, y_{2j-1}, \eta_{2j-1}) \mapsto A_{2j-1} = \mu_{2j-1}\tau_{2j-1}y_{2j-1}\eta_{2j-1}, \quad (2.2.1)$$
$$U \times T \times U_{\tau_{2j}}^{-} \cong B\tau_{2j}B : (\eta_{2j}, y_{2j}, \mu_{2j}) \mapsto A_{2j} = \eta_{2j}y_{2j}\tau_{2j}\mu_{2j}.$$

Here, recall that $A_m \in B\tau_m B$, $\forall 1 \le m \le 2g$. By Proposition 1.10, we have

$$[A_{2j-1}] = [\mu_{2j-1}] \circ [\tau_{2j-1}] \circ [\gamma_{2j-1}] \circ [\eta_{2j-1}], \ [A_{2j}] = [\eta_{2j}] \circ [y_{2j}] \circ [\tau_{2j}] \circ [\mu_{2j}] \in \underline{\text{FBD}}_n, \\ [A_{2j-1}^{-1}] = [\eta_{2j-1}^{-1}] \circ [y_{2j-1}^{-1}] \circ [\tau_{2j-1}^{-1}] \circ [\mu_{2j-1}^{-1}], \ [A_{2j}^{-1}] = [\mu_{2j}^{-1}] \circ [\tau_{2j}^{-1}] \circ [y_{2j}^{-1}] \circ [\eta_{2j}^{-1}] \in \underline{\text{FBD}}_n.$$

T. SU

Lemma 2.3. For any $D^j \in T$, we have a natural isomorphism of \mathbb{K} -varieties

$$B\tau_{2j-1}B \times B\tau_{2j}B \times U \xrightarrow{\simeq} U \times T^2 \times (U^-_{\tau_{2j-1}} \times U^-_{\tau_{2j}} \times U^-_{\tau_{2j-1}} \times U^-_{\tau_{2j}^{-1}}) \times (U^+_{\tau_{2j}^{-1}} \times U^+_{\tau_{2j-1}}), (2.2.2)$$

$$(A_{2j-1}, A_{2j}, u^j) \mapsto (u^{j-1}, y_{2j-1}, y_{2j}, \zeta'_{2j-1}, \zeta'_{2j}, \zeta_{2j-1}, \zeta_{2j}, {}^+n^{2j}, {}^+n^{2j-1}),$$

such that

$$[A_{2j-1}] \circ [A_{2j}] \circ [A_{2j-1}^{-1}] \circ [A_{2j}^{-1}] \circ [D^{j}]$$

$$= [u^{j-1}] \circ [D^{j-1}] \circ [\tau_{2j-1}] \circ [\zeta'_{2j-1}] \circ [\tau_{2j}] \circ [\zeta'_{2j}] \circ [\tau_{2j-1}^{-1}] \circ [\zeta_{2j-1}] \circ [\zeta_{2j}] \in \underline{FBD}_{n},$$

$$(2.2.3)$$

for a unique $D^{j-1} \in T$:

$$D^{j-1} := y_{2j-1}^{\tau_{2j-1}} y_{2j}^{\tau_{2j-1}} (y_{2j-1}^{-1})^{\tau_{2j-1}\tau_{2j}} (y_{2j}^{-1})^{\tau_{2j-1}\tau_{2j}\tau_{2j-1}^{-1}\tau_{2j}^{-1}} (D^{j})^{\tau_{2j-1}\tau_{2j}\tau_{2j-1}^{-1}\tau_{2j}^{-1}} \in T.$$
(2.2.4)

Moreover, (2.2.3) *uniquely determines* $(u^{j-1}, \zeta'_{2j-1}, \zeta'_{2j}, \zeta_{2j-1}, \zeta_{2j}) \in U \times U^{-}_{\tau_{2j-1}} \times U^{-}_{\tau_{2j-1}} \times U^{-}_{\tau_{2j-1}^{-1}} \times U^{-}_{\tau_{2j-1$

This provides the inductive step for the diagram calculus for genera. In our case, we have seen that $u^g \in U$, take $D^g = D_1 \in T$, and define D^j 's inductively as above. Thus,

$$D^{0}((y_{j})_{j=1}^{2g}) = \prod_{j=1}^{g} (y_{2j-1}^{\tau_{2j-1}}(y_{2j-1}^{-1})^{\tau_{2j-1}\tau_{2j}}y_{2j}^{\tau_{2j-1}}(y_{2j}^{-1})^{(\tau_{2j-1},\tau_{2j})}) \prod_{m=1}^{j-1} (\tau_{2m-1},\tau_{2m}) D_{1}^{\prod_{m=1}^{g} (\tau_{2m-1},\tau_{2m})}.$$
 (2.2.5)

Altogether, the defining equation (1.3.14) for $M'_B(\vec{w})$, i.e. $[M_{\vec{w}}] \stackrel{w}{\sim} id_n \in \underline{FBD}_n$, reduces to:

$$[u^{0}D^{0}] \circ \prod_{j=1}^{g} ([\tau_{2j-1}\zeta'_{2j-1}] \circ [\tau_{2j}\zeta'_{2j}] \circ [\tau_{2j-1}^{-1}\zeta_{2j-1}] \circ [\tau_{2j}^{-1}\zeta_{2j}]) \circ \prod_{i=1}^{k-1} ([\dot{w}_{i}\xi'_{i}] \circ [\dot{w}_{i}^{-1}\xi_{i}]) \stackrel{\mathfrak{w}}{\sim} \mathsf{id}_{n}.$$

Proof of Lemma 2.3. As one could expect, the proof is done by diagram calculus. **Step** 1. Denote $u'^j := \eta_{2j}^{-1} u^j \in U, \eta'_{2j} := \eta_{2j-1} \eta_{2j} \in U, \eta'_{2j-1} := \eta_{2j-1} \mu_{2j}^{-1} \in U$. So, $\eta'_{2j-1}^{-1} = \mu_{2j} \eta_{2j-1}^{-1}$. Then $(A_{2j-1}, A_{2j}, u^j) \mapsto (\mu_{2j-1}, y_{2j-1}, \eta'_{2j-1}, \eta'_{2j}, y_{2j}, \mu_{2j}, u'^j)$ defines an isomorphism

$$B\tau_{2j-1}B \times B\tau_{2j}B \times U \xrightarrow{\cong} U^{-}_{\tau^{-1}_{2j-1}} \times T \times U^2 \times T \times U^{-}_{\tau_{2j}} \times U$$

such that we obtain an identity in \underline{FBD}_n :

$$[A_{2j-1}] \circ [A_{2j}] \circ [A_{2j-1}^{-1}] \circ [A_{2j}^{-1}] \circ [u^{j}]$$

$$= [\mu_{2j-1}] \circ [\tau_{2j-1}] \circ [\gamma_{2j-1}] \circ [\eta'_{2j}] \circ [y_{2j}] \circ [\tau_{2j}]$$

$$\circ [\eta'_{2j-1}] \circ [y_{2j-1}^{-1}] \circ [\tau_{2j-1}^{-1}] \circ [\mu_{2j-1}^{-1}] \circ [\mu_{2j}^{-1}] \circ [\tau_{2j}^{-1}] \circ [y_{2j}^{-1}] \circ [u'^{j}].$$

$$(2.2.6)$$

Note: The variable $\mu_{2j}^{-1} \in U_{\tau_{2j}}^{-}$ becomes free (appears only once).

Step 2. We firstly compute $[\mu_{2j-1}^{-1}] \circ [\mu_{2j}^{-1}] \circ [\tau_{2j}^{-1}] \circ [y_{2j}^{-1}] \circ [u'^j]$. By Definition/Proposition 2.1,

$$\begin{split} & [\mu_{2j-1}^{-1}] \circ [U_{\tau_{2j}}^{-1}] \circ [\tau_{2j}^{-1}] \circ [y_{2j}^{-1}] \circ [U] \cong [\mu_{2j-1}^{-1}] \circ [U_{\tau_{2j}}^{-1}] \circ [\tau_{2j}^{-1}] \circ [U] \circ [y_{2j}^{-1}] \\ & \cong \quad [\mu_{2j-1}^{-1}] \circ [U_{\tau_{2j}}^{-1}] \circ [\tau_{2j}^{-1}] \circ [U_{\tau_{2j}^{-1}}^{+1}] \circ [U_{\tau_{2j}^{-1}}^{-1}] \circ [y_{2j}^{-1}] \quad (U = U_{\tau_{2j}^{-1}}^{+}U_{\tau_{2j}^{-1}}^{-}) \\ & \cong \quad [\mu_{2j-1}^{-1}] \circ [U_{\tau_{2j}}^{-1}] \circ [U_{\tau_{2j}^{-1}}^{+1}] \circ [\tau_{2j}^{-1}] \circ [U_{\tau_{2j}^{-1}}^{-1}] \circ [y_{2j}^{-1}] \cong [\mu_{2j-1}^{-1}] \circ [U] \circ [\tau_{2j}^{-1}] \circ [U_{\tau_{2j}^{-1}}^{-1}] \circ [y_{2j}^{-1}] \\ & \cong \quad [U] \circ [\tau_{2j}^{-1}] \circ [U_{\tau_{2j}^{-1}}^{-1}] \circ [y_{2j}^{-1}]. \end{split}$$

This means we obtain an isomorphism of K-varieties

$$U_{\tau_{2j-1}^{-1}}^{-} \times T \times U^{2} \times T \times U_{\tau_{2j}}^{-} \times U \xrightarrow{\cong} U_{\tau_{2j-1}^{-1}}^{-} \times T \times U^{3} \times T \times U_{\tau_{2j}^{-1}}^{-}, \qquad (2.2.7)$$

$$(\mu_{2j-1}, y_{2j-1}, \eta'_{2j-1}, \eta'_{2j}, y_{2j}, \mu_{2j}, u'^{j}) \mapsto (\mu_{2j-1}, y_{2j-1}, \eta'_{2j-1}, \eta'_{2j}, u^{j}_{3}, y_{2j}, L_{1}^{-}),$$

such that
$$[\mu_{2j-1}^{-1}] \circ [\mu_{2j}^{-1}] \circ [\tau_{2j}^{-1}] \circ [y_{2j}^{-1}] \circ [u'^{j}] = [u_{3}^{j}] \circ [\tau_{2j}^{-1}] \circ [L_{1}^{-1}] \circ [y_{2j}^{-1}] \in \underline{FBD}_{n}$$
. Hence,

$$[A_{2j-1}] \circ [A_{2j}] \circ [A_{2j-1}^{-1}] \circ [A_{2j}^{-1}] \circ [u^{j}] \qquad (2.2.8)$$

$$= [\mu_{2j-1}] \circ [\tau_{2j-1}] \circ [y_{2j-1}] \circ [\eta'_{2j}] \circ [\tau_{2j}]$$

$$\circ [\eta'_{2j-1}^{-1}] \circ [y_{2j-1}^{-1}] \circ [\tau_{2j-1}^{-1}] \circ [L_{1}^{-1}] \circ [y_{2j}^{-1}].$$

Note: the non-torus variables $\mu_{2j-1} \in U^-_{\tau^{-1}_{2j-1}}, \eta'_{2j}, \eta'^{-1}_{2j-1}, u^j_3 \in U, L^-_1 \in U^-_{\tau^{-1}_{2j}}$ all become free.

Step 3. By our trick (Definition/Proposition 2.1), the computation of $[A_{2j-1}] \circ [A_{2j}] \circ [A_{2j-1}^{-1}] \circ [A_{2j}^{-1}] \circ [u^j]$ then reduces to that of the following variety in <u>FBD</u>_n:

$$[U_{\tau_{2j-1}^{-1}}] \circ [\tau_{2j-1}] \circ [y_{2j-1}] \circ [U] \circ [y_{2j}] \circ [\tau_{2j}] \circ [U] \circ [y_{2j-1}^{-1}] \circ [\tau_{2j-1}^{-1}] \circ [U] \circ [\tau_{2j}^{-1}] \circ [U_{\tau_{2j}^{-1}}^{-1}] \circ [U_{$$

The idea is 'canonicalize'. According to the expression above, reorder the variables:

$$U_{\tau_{2j-1}^{-1}}^{-} \times T \times U^{3} \times T \times U_{\tau_{2j}^{-1}}^{-} \xrightarrow{\cong} U_{\tau_{2j-1}^{-1}}^{-} \times (T \times U)^{2} \times U \times U_{\tau_{2j}^{-1}}^{-}, \qquad (2.2.10)$$

$$(\mu_{2j-1}, y_{2j-1}, \eta_{2j-1}', \eta_{2j}', u_{3}^{j}, y_{2j}, L_{1}^{-}) \mapsto (\mu_{2j-1}, y_{2j-1}, \eta_{2j}', y_{2j}, \eta_{2j-1}', u_{3}^{j}, L_{1}^{-})$$

Step 3.1. We firstly compute $[U] \circ [y_{2j-1}^{-1}] \circ [\tau_{2j-1}^{-1}] \circ [U] \ni [\eta_{2j-1}^{\prime-1}] \circ [y_{2j-1}^{-1}] \circ [\tau_{2j-1}^{-1}] \circ [u_3^j]$:

$$\begin{split} & [U] \circ [y_{2j-1}^{-1}] \circ [\tau_{2j-1}^{-1}] \circ [U] \cong [U] \circ [y_{2j-1}^{-1}] \circ [\tau_{2j-1}^{-1}] \circ [U_{\tau_{2j-1}^{-1}}^{-1}] \circ [U_{\tau_{2j-1}^{-1}}^{-1}] (U = U_{\tau_{2j-1}^{-1}}^{+}U_{\tau_{2j-1}^{-1}}^{-1}) \\ & \cong \quad [U] \circ [U_{\tau_{2j-1}}^{+}] \circ [y_{2j-1}^{-1}] \circ [\tau_{2j-1}^{-1}] \circ [U_{\tau_{2j-1}^{-1}}^{-1}] \cong [U] \circ [y_{2j-1}^{-1}] \circ [\tau_{2j-1}^{-1}] \circ [U_{\tau_{2j-1}^{-1}}^{-1}] \times U_{\tau_{2j-1}}^{+}, \end{split}$$

with the direct factor $U_{\tau_{2j-1}}^+ \ni {}^+ n^{2j-1}$. This means we obtain an isomorphism of \mathbb{K} -varieties

$$U_{\tau_{2j-1}^{-1}}^{-} \times (T \times U)^{2} \times U \times U_{\tau_{2j}^{-1}}^{-} \xrightarrow{\simeq} U_{\tau_{2j-1}^{-1}}^{-} \times (T \times U)^{2} \times U_{\tau_{2j-1}^{-1}}^{-} \times U_{\tau_{2j}^{-1}}^{-} \times U_{\tau_{2j-1}}^{+},$$
(2.2.11)
$$(\mu_{2j-1}, y_{2j-1}, \eta'_{2j}, y_{2j}, \eta'_{2j-1}^{-1}, u_{3}^{j}, L_{1}^{-}) \mapsto (\mu_{2j-1}, y_{2j-1}, \eta'_{2j}, y_{2j}, u_{4}^{j}, L_{3}^{-}, L_{1}^{-}, {}^{+}n^{2j-1}),$$

such that $[\eta_{2j-1}^{\prime-1}] \circ [y_{2j-1}^{-1}] \circ [\tau_{2j-1}^{-1}] \circ [u_3^j] = [u_4^j] \circ [y_{2j-1}^{-1}] \circ [\tau_{2j-1}^{-1}] \circ [L_3^-] \in \underline{FBD}_n$. **Step** 3.2. We compute $[U] \circ [y_{2j}] \circ [\tau_{2j}] \circ [U] \ni [\eta_{2j}'] \circ [y_{2j}] \circ [\tau_{2j}] \circ [u_4^j]$, which is similar:

$$\begin{split} & [U] \circ [y_{2j}] \circ [\tau_{2j}] \circ [U] \cong [U] \circ [y_{2j}] \circ [\tau_{2j}] \circ [U_{\tau_{2j}}^+] \circ [U_{\tau_{2j}}^-] \quad (U = U_{\tau_{2j}}^+ U_{\tau_{2j}}^-) \\ \cong & [U] \circ [U_{\tau_{2j}}^+] \circ [y_{2j}] \circ [\tau_{2j}] \circ [U_{\tau_{2j}}^-] \cong [U] \circ [y_{2j}] \circ [\tau_{2j}] \circ [U_{\tau_{2j}}^-] \times U_{\tau_{\tau}^{-1}}^+, \end{split}$$

with the direct factor $U^+_{\tau_{2i}^{-1}} \ni {}^+n^{2j}$. This means we obtain an isomorphism of \mathbb{K} -varieties

$$U_{\tau_{2j-1}^{-1}}^{-} \times (T \times U)^{2} \times U_{\tau_{2j-1}^{-1}}^{-} \times U_{\tau_{2j}^{-1}}^{-} \times U_{\tau_{2j-1}}^{+} \xrightarrow{\simeq} U_{\tau_{2j-1}^{-1}}^{-} \times T \times U \times T \times U_{\tau_{2j}}^{-} \times U_{\tau_{2j-1}^{-1}}^{-} \times U_{\tau_{2j}^{-1}}^{-} \times U_{\tau_{2j}^{-1}}^{+} \times U_{\tau_{2j}^{-1}}^{+} \times U_{\tau_{2j-1}^{-1}}^{+},$$

$$(\mu_{2j-1}, y_{2j-1}, \eta'_{2j}, y_{2j}, u_{4}^{j}, L_{3}^{-}, L_{1}^{-}, {}^{+}n^{2j-1}) \mapsto (\mu_{2j-1}, y_{2j-1}, u_{5}^{j}, y_{2j}, L_{5}^{-}, L_{3}^{-}, L_{1}^{-}, {}^{+}n^{2j-1}),$$

such that $[\eta'_{2j}] \circ [y_{2j}] \circ [\tau_{2j}] \circ [u_4^j] = [u_5^j] \circ [y_{2j}] \circ [\tau_{2j}] \circ [L_5^-] \in \underline{FBD}_n.$ **Step** 3.3. Now, we compute $[U_{\tau_{2j-1}}^-] \circ [\tau_{2j-1}] \circ [y_{2j-1}] \circ [U] \ni [\mu_{2j-1}] \circ [\tau_{2j-1}] \circ [y_{2j-1}] \circ [u_5^j]:$ $[U_{\tau_{2j-1}}^-] \circ [\tau_{2j-1}] \circ [y_{2j-1}] \circ [U] \cong [U_{\tau_{2j-1}}^-] \circ [\tau_{2j-1}] \circ [U_{\tau_{2j-1}}^+] \circ [U_{\tau_{2j-1}}^-]$

$$\cong \quad [U_{\tau_{2j-1}^{-1}}^{-1}] \circ [U_{\tau_{2j-1}^{-1}}^{+1}] \circ [\tau_{2j-1}] \circ [y_{2j-1}] \circ [U_{\tau_{2j-1}}^{-1}] \cong [U] \circ [\tau_{2j-1}] \circ [y_{2j-1}] \circ [U_{\tau_{2j-1}}^{-1}]$$

This means we obtain an isomorphism of $\mathbb K\text{-varieties}$

$$\begin{split} & U_{\tau_{2j-1}^{-1}}^{-} \times T \times U \times T \times U_{\tau_{2j}}^{-} \times U_{\tau_{2j-1}^{-1}}^{-} \times U_{\tau_{2j}^{-1}}^{-} \times U_{\tau_{2j}^{-1}}^{+} \times U_{\tau_{2j-1}^{-1}}^{+} \xrightarrow{-} \\ & U \times T \times U_{\tau_{2j-1}}^{-} \times T \times U_{\tau_{2j}}^{-} \times U_{\tau_{2j-1}^{-1}}^{-} \times U_{\tau_{2j}^{-1}}^{-} \times U_{\tau_{2j}^{-1}}^{+} \times U_{\tau_{2j-1}^{+}}^{+}, \\ & (\mu_{2j-1}, y_{2j-1}, u_{5}^{j}, y_{2j}, L_{5}^{-}, L_{3}^{-}, L_{1}^{-}, {}^{+}n^{2j}, {}^{+}n^{2j-1}) \mapsto (u^{j-1}, y_{2j-1}, L_{7}^{-}, y_{2j}, L_{5}^{-}, L_{3}^{-}, L_{1}^{-}, {}^{+}n^{2j-1}) \end{split}$$

such that $[\mu_{2j-1}] \circ [\tau_{2j-1}] \circ [y_{2j-1}] \circ [u_5^j] = [u^{j-1}] \circ [\tau_{2j-1}] \circ [y_{2j-1}] \circ [L_7^-] \in \underline{FBD}_n$. In summary, we have obtained

$$\begin{bmatrix} A_{2j-1} \end{bmatrix} \circ \begin{bmatrix} A_{2j} \end{bmatrix} \circ \begin{bmatrix} A_{2j-1}^{-1} \end{bmatrix} \circ \begin{bmatrix} A_{2j}^{-1} \end{bmatrix} \circ \begin{bmatrix} u^{j} \end{bmatrix}$$
(2.2.12)

$$= \begin{bmatrix} u^{j-1} \end{bmatrix} \circ \begin{bmatrix} \tau_{2j-1} \end{bmatrix} \circ \begin{bmatrix} y_{2j-1}L_{7}^{-}y_{2j} \end{bmatrix} \circ \begin{bmatrix} \tau_{2j} \end{bmatrix} \circ \begin{bmatrix} L_{5}^{-}y_{2j-1}^{-1} \end{bmatrix} \circ \begin{bmatrix} \tau_{2j-1}^{-1} \end{bmatrix} \circ \begin{bmatrix} \tau_{2j}^{-1} \end{bmatrix} \circ \begin{bmatrix} L_{1}^{-}y_{2j}^{-1} \end{bmatrix} .$$
and
$$\begin{bmatrix} U_{\tau_{2j-1}^{-1}} \end{bmatrix} \circ \begin{bmatrix} \tau_{2j-1} \end{bmatrix} \circ \begin{bmatrix} U \end{bmatrix} \circ \begin{bmatrix} y_{2j-1} \end{bmatrix} \circ \begin{bmatrix} U \end{bmatrix} \circ \begin{bmatrix} y_{2j} \end{bmatrix} \circ \begin{bmatrix} \tau_{2j} \end{bmatrix} \circ \begin{bmatrix} U \end{bmatrix} \circ \begin{bmatrix} \tau_{2j-1} \end{bmatrix} \circ \begin{bmatrix} U \end{bmatrix} \circ \begin{bmatrix} \tau_{2j}^{-1} \end{bmatrix} \circ \begin{bmatrix} U_{\tau_{2j}^{-1}} \end{bmatrix} \circ \begin{bmatrix} U_{\tau_{2j}^{-1}} \end{bmatrix} \circ \begin{bmatrix} U_{\tau_{2j}^{-1}} \end{bmatrix} \circ \begin{bmatrix} U_{\tau_{2j}^{-1}} \end{bmatrix} \circ \begin{bmatrix} \tau_{2j} \end{bmatrix} \circ$$

Step 4. Deal with the torus variables. Now, for any $D^j \in T$, it follows that

$$\begin{split} & [A_{2j-1}] \circ [A_{2j}] \circ [A_{2j-1}^{-1}] \circ [A_{2j}^{-1}] \circ [u^{j}] \circ [D^{j}] \\ & = \quad [u^{j-1}] \circ [\tau_{2j-1}] \circ [y_{2j-1}L_{7}^{-}y_{2j}] \circ [\tau_{2j}] \circ [L_{5}^{-}y_{2j-1}^{-1}] \circ [\tau_{2j-1}^{-1}] \circ [\tau_{2j}^{-1}] \circ [L_{1}^{-}y_{2j}^{-1}] \circ [D^{j}] \\ & = \quad [u^{j-1}] \circ [D^{j-1}] \circ [\tau_{2j-1}] \circ [\zeta_{2j-1}] \circ [\tau_{2j}] \circ [\zeta_{2j}'] \circ [\tau_{2j-1}^{-1}] \circ [\zeta_{2j-1}] \circ [\tau_{2j}^{-1}] \circ [\zeta_{2j}] \in \underline{\text{FBD}}_{n}, \end{split}$$

where $(\zeta'_{2j-1}, \zeta'_{2j}, \zeta_{2j-1}, \zeta_{2j}) \in U^-_{\tau_{2j-1}} \times U^-_{\tau_{2j}} \times U^-_{\tau_{2j-1}^{-1}} \times U^-_{\tau_{2j}^{-1}}$, and D^{j-1} is indeed given by (2.2.4).

In other words, we have obtained an isomorphism of the form (2.2.2) in Lemma 2.3 such that (2.2.3) and (2.2.4) hold. By definition of \underline{FBD}_n , the uniqueness part is clear. Done.

Remark 2.4. By a careful check of the proof, there're formulas for ${}^{+}n^{2j-1}$, ${}^{+}n^{2j}$ using (1.3.7):

2.3. Connection to braid varieties. We relate $M'_B(\vec{w})$ to braid varieties. Sum up Section 2.1-2.2, we've obtained an isomorphism

$$(\prod_{j=1}^{2g} B\tau_{j}B) \times (\prod_{i=1}^{k-1} B\dot{w}_{i}P_{i}) \times U \xrightarrow{\cong}$$

$$(2.3.1)$$

$$U \times \prod_{j=1}^{g} (T^{2} \times (U_{\tau_{2j-1}}^{-} \times U_{\tau_{2j}}^{-} \times U_{\tau_{2j-1}}^{-} \times U_{\tau_{2j}}^{-}) \times (U_{\tau_{2j-1}}^{+} \times U_{\tau_{2j-1}}^{+})) \times \prod_{i=1}^{k-1} (U_{\dot{w}_{i}}^{-} \times U_{\dot{w}_{i}}^{-}) \times (U_{\dot{w}_{i}}^{+} \cap N_{i}) \times Z(C_{i})),$$

$$((A_{j})_{j}, (x_{i})_{i}, u_{k}) \mapsto (u^{0}, (y_{2j-1}, y_{2j}, \zeta_{2j-1}', \zeta_{2j}', \zeta_{2j-1}, \zeta_{2j}, {}^{+}n^{2j}, {}^{+}n^{2j-1})_{j=1}^{g}, (\xi_{i}', \xi_{i}, {}^{+}n_{i}', z_{i})_{i=1}^{k-1}),$$

such that we obtain an equality in FBD_n:

$$\begin{split} [\mathbf{M}_{\vec{w}}] &= \prod_{j=1}^{s} ([A_{2j-1}] \circ [A_{2j}] \circ [A_{2j-1}^{-1}] \circ [A_{2j}^{-1}]) \circ \prod_{i=1}^{k-1} [x_i C_i x_i^{-1}]' \circ [u_k] \circ [D_k] \\ &= [u^0 D^0] \circ \prod_{j=1}^{s} ([\tau_{2j-1}] \circ [\zeta'_{2j-1}] \circ [\tau_{2j}] \circ [\zeta'_{2j}] \circ [\tau_{2j-1}^{-1}] \circ [\zeta_{2j-1}] \circ [\tau_{2j}^{-1}] \circ [\zeta_{2j}]) \circ \prod_{i=1}^{k-1} ([\dot{w}_i] \circ [\xi'_i] \circ [\dot{w}_i^{-1}] \circ [\xi_i]). \end{split}$$

Recall by (1.3.15), $\beta(\vec{w}) = \prod_{j=1}^{g} ([\tau_{2j-1}] \circ [\tau_{2j}] \circ [\tau_{2j-1}]^{-1} \circ [\tau_{2j}^{-1}]) \circ \prod_{i=1}^{k-1} ([\dot{w}_i] \circ [\dot{w}_i^{-1}]) \in FBr_n^+$. By Definition 1.12, the defining equation (1.3.14) for $M'_B(\vec{w})$ in <u>FBD</u>_n becomes

$$[\mathbf{B}_{\beta(\vec{w})}(\vec{\epsilon})]' \stackrel{\mathbf{w}}{\sim} [(D^0)^{-1}] \circ [(u^0)^{-1}], \ \vec{\epsilon} := ((\zeta'_{2j-1}, \zeta'_{2j}, \zeta_{2j-1}, \zeta_{2j})^g_{j=1}, (\xi'_i, \xi_i)^{k-1}_{i=1}) \in \mathbb{K}^{\ell(\beta(\vec{w}))}.$$
(2.3.2)

By (2.2.5), (2.1.3), and the generic assumption (Definition 1.1), det $D^0 = \det D_1 = 1$. Define

$$\phi_{\vec{w}}: T^{2g} \times X(\beta(\vec{w})) \to T_1: ((y_j)_{j=1}^{2g}, \vec{\epsilon}) \mapsto D^0 \mu \mathfrak{mon}(\vec{\epsilon}); \quad T_1 := \{t \in T : \det t = 1\}.$$
(2.3.3)

Here, det $\circ \mu \mathfrak{mon} = 1$ by Proposition 1.16: $\forall p \in \mathcal{W}(\beta(\vec{w})) \Rightarrow |S_p| = \ell(\beta(\vec{w})) - 2|U_p| = \sum_{j=1}^g 2\ell(\tau_j) + \sum_{i=1}^{k-1} 2\ell(\vec{w}_i) - 2|U_p|$, which is even. Define a closed K-subvariety

$$\widetilde{X}(\beta(\vec{w})) := \phi_{\vec{w}}^{-1}(I_n) \subset T^{2g} \times X(\beta(\vec{w})).$$
(2.3.4)

Then by (2.3.2), $M'_B(\vec{w})$ is related to the restricted twisted braid variety $\widetilde{X}(\beta(\vec{w}))$:

$$M'_{B}(\vec{w}) \cong \widetilde{X}(\beta(\vec{w})) \times \prod_{j=1}^{g} (U^{+}_{\tau_{2j}^{-1}} \times U^{+}_{\tau_{2j-1}}) \times \prod_{i=1}^{k-1} ((U^{+}_{\dot{w}_{i}} \cap N_{i}) \times Z(C_{i})).$$
(2.3.5)

Now, $(b, (h_i)_{i=1}^{k-1}) \in B_{\text{par}}$ acts on $(u^0, (y_j, \zeta'_j, \zeta_j, {}^+n^j)_{j=1}^{2g}, (\xi'_i, \xi_i, {}^+n'_i, z_i)_{i=1}^{k-1}) \in M'_B(\vec{w})$. Our major interest is that of $T_{\text{par}} := T \times \prod_{i=1}^{k-1} Z(C_i) \subset B_{\text{par}}$. Let's study the action. We use **Convention** 4. Recall by (1.2.5) and (1.3.10) that

$$\widehat{A}_{j} = bA_{j}b^{-1}, \quad \widehat{x}_{i} = bx_{i}h_{i}^{-1}, \quad \widehat{u}_{k} = bu_{k}(b^{C_{k}})^{-1}; \quad x_{i} = v_{i}\dot{w}_{i}n_{i}z_{i}.$$

Notice that $h_i \in Z(C_i)$ acts only on x_i , hence only on $z_i \in Z(C_i)$ via: $z_i \mapsto z_i h_i^{-1}$. Thus, define

$$M_B''(\vec{w}) := \{ (u^0, (y_j)_{j=1}^{2g}, \vec{\epsilon}, ({}^+n^j)_{j=1}^{2g}, ({}^+n'_i)_{i=1}^{k-1}) : u^0 D^0 \mathbf{B}_{\beta(\vec{w})}(\vec{\epsilon}) = \mathsf{id}_n \}.$$
(2.3.6)

Then it admits an induced action of $b \in B$, and the above computation shows that

$$M_B''(\vec{w}) \cong \widetilde{X}(\beta(\vec{w})) \times \prod_{j=1}^{g} (U_{\tau_{2j}^{-1}}^+ \times U_{\tau_{2j-1}}^+) \times \prod_{i=1}^{k-1} (U_{\vec{w}_i}^+ \cap N_i); \quad M_B'(\vec{w}) = M_B''(\vec{w}) \times \prod_{i=1}^{k-1} Z(C_i).$$
(2.3.7)

Recall that $PB_{par} = B_{par}/\mathbb{K}^{\times}$ acts freely on $M'_B(\vec{w})$, and $M'_B(\vec{w}) \to \mathcal{M}_{\mu}(\vec{w}) = M'_B(\vec{w})/PB_{par}$ is a principal PB_{par} -bundle. Observe that $\prod_{i=1}^{k-1} Z(C_i) \cong (\mathbb{K}^{\times} \times \prod_{i=1}^{k-1} Z(C_i))/\mathbb{K}^{\times} \hookrightarrow PB_{par} = (B \times \prod_{i=1}^{k-1} Z(C_i))/\mathbb{K}^{\times}$ is a normal closed subgroup, with quotient $PB_{par}/\prod_{i=1}^{k-1} Z(C_i) \cong PB := B/\mathbb{K}^{\times}$. Clearly, we have $M'_B(\vec{w})/\prod_{i=1}^{k-1} Z(C_i) \cong M''_B(\vec{w})$. Then, by Proposition B.14 and Proposition B.9,

$$\mathcal{M}_{\mu}(\vec{w}) = M'_{B}(\vec{w})/PB_{\text{par}} \cong M''_{B}(\vec{w})/PB,$$
 (2.3.8)

and the induced quotient map

$$\pi''_{\vec{w}}: M''_B(\vec{w}) \to \mathcal{M}_\mu(\vec{w}) = M''_B(\vec{w})/PB$$

is a principal *PB*-bundle. It leads to study the *B*-action on $M_B''(\vec{w})$.

Lemma 2.5. The induced action of $b \in B$ on $(u^0, (y_j)_{j=1}^{2g}, \vec{\epsilon}, (+n^j)_{j=1}^{2g}, (+n'_i)_{i=1}^{k-1}) \in M''_B(\vec{w})$ satisfies: a) The canonical projection $M''_B(\vec{w}) \to \widetilde{X}(\beta(\vec{w})) \subset T^{2g} \times X(\beta(\vec{w}))$ is B-equivariant. Here, $b \in B$ acts on $((y_j)_{j=1}^{2g}, \vec{\epsilon}) \in T^{2g} \times X(\beta(\vec{w}))$ diagonally: $B \sim X(\beta(\vec{w}))$ via Definition 1.14, and

$$\widehat{y}_{2j-1} = D(b)^{\tau_{2j-1}^{-1}} D(b)^{-1} y_{2j-1}, \ \widehat{y}_{2j} = D(b) (D(b)^{\tau_{2j}})^{-1} y_{2j}.$$
 (2.3.9)

b) If $b = t \in T$, then

$$\hat{n}'_{i} = t^{\hat{w}_{i}^{-1} + n'_{i}}(t^{-1})^{\hat{w}_{i}^{-1}}; \hat{n}^{2j-1} = t^{+}n^{2j-1}t^{-1}; \hat{n}^{2j} = t^{+}n^{2j}t^{-1},$$

$$(2.3.10)$$

$$(\hat{y}_{2j-1} = t^{\tau_{2j-1}^{-1}}t^{-1}y_{2j-1}; \hat{y}_{2j} = t(t^{\tau_{2j}})^{-1}y_{2j}; \hat{u}^{0} = tu^{0}t^{-1}; \hat{D}^{0} = tD^{0}(t^{-1})^{\prod_{m=1}^{g}(\tau_{2m-1}, \tau_{2m})})$$

$$[\hat{u}^{0}] \circ [\hat{D}^{0}] \circ [\mathbf{B}_{\beta(\vec{w})}(\hat{\vec{\epsilon}})]' = [t] \circ [u^{0}] \circ [D^{0}] \circ [\mathbf{B}_{\beta(\vec{w})}(\vec{\epsilon})]' \circ [t^{-1}] \in \underline{FBD}_{n}.$$

Here, the last equation uniquely determines $(\hat{u}^0, \hat{D}^0, \hat{\vec{\epsilon}})$.

Proof. Now, $\widehat{A}_j = bA_j b^{-1}$, $\widehat{x}_i = bx_i$. So, $[\widehat{A}_j] = [b] \circ [A_j] \circ [b^{-1}]$, $[\widehat{x}_i C_i \widehat{x}_i^{-1}]' = [b] \circ [x_i C_i x_i^{-1}]' \circ [b^{-1}]$. *a*). By Section 2.2, $(\widehat{\mu}_{2j-1}, \widehat{y}_{2j-1}, \widehat{\eta}_{2j-1}) \in U^-_{\tau_{2j-1}^{-1}} \times T \times U$ is uniquely determined by:

$$\widehat{\mu}_{2j-1}\tau_{2j-1}\widehat{y}_{2j-1}\widehat{\eta}_{2j-1} = \widehat{A}_{2j-1} = bA_{2j-1}b^{-1} = b\mu_{2j-1}\tau_{2j-1}y_{2j-1}\eta_{2j-1}b^{-1}$$

Thus, $\hat{y}_{2j-1} = D(b)^{\tau_{2j-1}^{-1}} y_{2j-1} D(b)^{-1} \in T$, as desired. Similarly, $\hat{y}_{2j} = D(b) y_{2j} (D(b)^{-1})^{\tau_{2j}} \in T$. Recall that $\hat{u}_k = b u_k (b^{C_k})^{-1}$, $D_k = C_k$, then we have equalities in <u>FBD</u>_n:

$$\begin{split} & [\widehat{u}^{0}\widehat{D}^{0}] \circ [\mathbf{B}_{\beta(\vec{w})}(\vec{\hat{\epsilon}})] = [\widehat{\mathbf{M}}_{\vec{w}}] = \prod_{j=1}^{g} ([\widehat{A}_{2j-1}] \circ [\widehat{A}_{2j}] \circ [\widehat{A}_{2j-1}^{-1}] \circ [\widehat{A}_{2j}^{-1}]) \circ \prod_{i=1}^{k-1} [\widehat{x}_{i}C_{i}\widehat{x}_{i}^{-1}]' \circ [\widehat{u}_{k}D_{k}] \mathcal{B}. 11) \\ & = [b] \circ \prod_{j=1}^{g} ([A_{2j-1}] \circ [A_{2j}] \circ [A_{2j-1}^{-1}] \circ [A_{2j}^{-1}]) \circ \prod_{i=1}^{k-1} [x_{i}C_{i}x_{i}^{-1}]' \circ [b^{-1}] \circ [\widehat{u}_{k}D_{k}] \\ & = [b] \circ [u^{0}D^{0}] \circ [\mathbf{B}_{\beta(\vec{w})}(\vec{\epsilon})] \circ [D_{k}^{-1}u_{k}^{-1}b^{-1}\widehat{u}_{k}D_{k}] = [b] \circ [u^{0}D^{0}] \circ [\mathbf{B}_{\beta(\vec{w})}(\vec{\epsilon})] \circ [b^{-1}]. \end{split}$$

This equation uniquely determines $(\hat{u}^0, \hat{D}^0, \hat{\vec{\epsilon}})$, and we see that $\hat{\vec{\epsilon}}$ coincides with the action of $b \in B$ on $\vec{\epsilon} \in X(\beta(\vec{w}))$ in Definition 1.14. This show *a*) and most part of *b*).

b). By above, it remains to check the action of $b = t \in T$ on $((+n^j)_j, (+n'_j)_i)$.

By the equation $\hat{\nu}_i \dot{w}_i \hat{n}_i \hat{z}_i = \hat{x}_i = t x_i = t \nu_i \dot{w}_i n_i z_i$, we see that $\hat{n}_i = t^{\dot{w}_i^{-1}} n_i (t^{\dot{w}_i^{-1}})^{-1}$. It follows that $\hat{n}'_i = \hat{n}_i C_i \hat{n}_i^{-1} C_i^{-1} = t^{\dot{w}_i^{-1}} n'_i (t^{\dot{w}_i^{-1}})^{-1}$. Then $\hat{n}'_i = R^+_{\dot{w}_i} (\hat{n}'_i) = t^{\dot{w}_i^{-1}} n'_i (t^{\dot{w}_i^{-1}})^{-1}$, as desired.

It remains to compute \widehat{n}^{j} . By (2.1.4) and similar to (2.3.11), we get by induction that $\widehat{u}_{i} = tu_{i}t^{-1}$. Similarly, by (2.2.3) and induction, we get $\widehat{u}^{j} = tu^{j}t^{-1}$. By the equation $\widehat{\mu}_{2j-1}\tau_{2j-1}\widehat{y}_{2j-1}\widehat{\eta}_{2j-1} = t\mu_{2j-1}\tau_{2j-1}y_{2j-1}\eta_{2j-1}t^{-1}$, we see $\widehat{\mu}_{2j-1} = t\mu_{2j-1}t^{-1}$, $\widehat{\eta}_{2j-1} = t\eta_{2j-1}t^{-1}$. Similarly, $\widehat{\mu}_{2j} = t\mu_{2j}t^{-1}$, $\widehat{\eta}_{2j} = t\eta_{2j}t^{-1}$. Now, by Remark 2.4, we see $\widehat{n}^{j} = t^{+1}\widehat{n}^{j}t^{-1}$, as desired.

2.4. The cell decomposition. Recall that $\pi''_{\vec{w}} : M''_B(\vec{w}) \to \mathcal{M}_\mu(\vec{w}) = M''_B(\vec{w})/PB$ is a principal *PB*bundle. Moreover, $M''_B(\vec{w}) \cong \widetilde{X}(\beta(\vec{w})) \times_{\prod_{j=1}^g} (U^+_{\tau_{2j}^{-1}} \times U^+_{\tau_{2j-1}}) \times_{\prod_{i=1}^{k-1}} (U^+_{\vec{w}_i} \cap N_i)$ by (2.3.7). Say, $\beta := \beta(\vec{w}) = \sigma_{i_\ell} \cdots \sigma_{i_1} \in FBr_n^+$. Define a *B*-invariant closed subvariety of $T^{2g} \times X(\beta)$ by

$$\widetilde{X}_p(\beta) := \widetilde{X}(\beta) \cap (T^{2g} \times X_p(\beta)) \subset T^{2g} \times X(\beta), \quad p \in \mathcal{W}(\beta).$$
(2.4.1)

Then by Lemma 2.5, the *B*-equivariant decomposition $\widetilde{X}(\beta) = \bigsqcup_{p \in W(\beta)} \widetilde{X}_p(\beta)$ induces a (resp. *B*-equivariant) decomposition of $\mathcal{M}_{\mu}(\vec{w})$ (resp. $\mathcal{M}''_B(\vec{w})$), as desired. It remains to give a more concrete description of each piece in these decompositions. This reduces to describe $\widetilde{X}_p(\beta)$.

We start with the following observation: Remember that *PB* acts freely on $M''_B(\vec{w})$. By (2.3.10), we then see that $PT = T/\mathbb{K}^{\times} \subset PB$ preserves and acts *freely* on $\widetilde{X}(\beta) \cong \widetilde{X}(\beta) \times \{0\} \subset M''_B(\vec{w})$. It turns out that, this free action leads to a more concrete description of $\widetilde{X}(\beta)$, which we now pursue.

By Proposition 1.16, $X_p(\beta) \cong (\mathbb{K}^{\times})^{|S_p|} \times \mathbb{K}^{|U_p|} : (\vec{\epsilon}) = (\epsilon_m)^1_{m=\ell} \mapsto (\epsilon'_m)_{m \in S_p \sqcup U_p}$. Define

$$\overline{\phi}_{\vec{w},p} : T^{2g} \times (\mathbb{K}^{\times})^{|S_p|} \to T_1 : (\vec{y} = (y_j)_{j=1}^{2g}, (\epsilon'_m)_{m \in S_p}) \mapsto D^0(\vec{y}) \mu \mathfrak{mon}((\epsilon'_m)_{m \in S_p}), \quad (2.4.2)$$

$$\widetilde{T}_p(\beta) := \overline{\phi}_{\vec{w},p}^{-1}(I_n) \subset T^{2g} \times (\mathbb{K}^{\times})^{|S_p|}.$$

Here, $D^0(\vec{y})$ is given by (2.2.5), and $\mu \mathfrak{mon}((\epsilon'_m)_{m \in S_p})$ is given by Proposition 1.16 (2). Clearly, $\phi_{\vec{w},p} := \phi_{\vec{w}}|_{T^{2g} \times X_p(\beta)}$ is the composition of $\overline{\phi}_{\vec{w},p}$ with the obvious projection:

$$\phi_{\vec{w},p}: T^{2g} \times X_p(\beta) \cong T^{2g} \times (\mathbb{K}^{\times})^{|S_p|} \times \mathbb{K}^{|U_p|} \twoheadrightarrow T^{2g} \times (\mathbb{K}^{\times})^{|S_p|} \xrightarrow{\phi_{\vec{w},p}} T_1.$$

Hence, we have

$$\widetilde{X}_{p}(\beta) = \phi_{\widetilde{w},p}^{-1}(I_{n}) = \widetilde{T}_{p}(\beta) \times \mathbb{K}^{|U_{p}|}.$$
(2.4.3)

In particular, $\widetilde{X}_p(\beta) \neq \emptyset$ if and only $\widetilde{T}_p(\beta) \neq \emptyset$.

Also, by Proposition 1.16 and (2.3.9), $T^{2g} \times (\mathbb{K}^{\times})^{|S_p|}$ inherits an action of T such that, the projection $T^{2g} \times X_p(\beta) \cong T^{2g} \times (\mathbb{K}^{\times})^{|S_p|} \times \mathbb{K}^{|U_p|} \twoheadrightarrow T^{2g} \times (\mathbb{K}^{\times})^{|S_p|}$ is equivariant with respect to the projection $B \to T : b \mapsto D(b)$. Besides, observe that

PT acts freely on
$$\tilde{X}_p(\beta)$$
 if and only if it does so on $\tilde{T}_p(\beta)$. (2.4.4)

Indeed, this is the case if $\widetilde{X}_p(\beta) \neq \emptyset$ by the observation above.

The key property is the following

Lemma 2.6. The PT-action on $\widetilde{T}_p(\beta)$ is free if and only if $\overline{\phi}_{w,p} : T^{2g} \times (\mathbb{K}^{\times})^{|S_p|} \to T_1$ is surjective.

Proof. Firstly, we consider the PT-action. For that, we make some preparations:

(1) For any $\tau \in S_n$, define a subgroup of *T* by

$${}^{\tau}T := \{t \in T : t^{\tau} = t, \text{ i.e., } t_a = t_{\tau(a)}, \forall a \in [n]\} \subset T.$$
(2.4.5)

So, ${}^{(a \ b)}T = \{t \in T : t_a = t_b\}$ for any transposition $(a \ b) \in S_n$. Define a subgroup of T by

$${}^{I}T := \{t \in T : t^{\tau} = t, \forall \tau \in I\} \subset T, \quad I \subset S_{n}; \quad \Leftrightarrow \quad {}^{I}T = \cap_{\tau \in I}{}^{\tau}T = \cap_{a \in [n], \tau \in I}{}^{(a \ \tau(a))}T.$$
(2.4.6)

(2) By definition, $\tau T = \tau^{-1}T$. Also, $t = t^{\tau_1}, t = t^{\tau_2}$ implies that $t = t^{\tau_1} = (t^{\tau_2})^{\tau_1} = t^{\tau_1\tau_2}$, so by (1),

$${}^{I}T = {}^{\langle I \rangle}T = {}^{\overline{\langle I \rangle}}T.$$
(2.4.7)

Here, we denote by $\langle I \rangle \subset S_n$ the subgroup generated by *I*, and

$$\langle I \rangle := \langle (a \ \tau(a)), \forall a \in [n], \tau \in I \rangle \subset S_n.$$
(2.4.8)

(3) Let $[n] = O_1 \sqcup \cdots \sqcup O_r$ be any partition, denoted by \vec{O} . Define subgroups of T and S_n :

$${}^{O}T := \{t \in T : a, b \in O_m \Rightarrow t_a = t_b, \forall 1 \le m \le r\} \subset T; \quad S_{\vec{O}} := S_{|O_1|} \times \dots \times S_{|O_r|} \subset S_n.$$
(2.4.9)
Then for any subset $I \subset S_n$, we have

$${}^{I}T = {}^{\vec{O}}T \Leftrightarrow \{O_i : 1 \le i \le r\} = \{\langle I \rangle \text{-orbits on } [n]\} \Leftrightarrow \overline{\langle I \rangle} = S_{\vec{O}}.$$
 (2.4.10)

Now, we see that ${}^{I}T = {}^{J}I$ if and only if $\overline{\langle I \rangle} = \overline{\langle J \rangle}$. In particular, ${}^{I}T = \mathbb{K}^{\times}I_{n}$ if and only if $\overline{\langle I \rangle} = S_{n}$.

Come back to the proof. The free *PT*-action on $\widetilde{T}_p(\beta)$ means: $\forall t \in T$, $\forall ((y_j)_{j=1}^{2g}, (\epsilon'_m)_{m \in S_p}) \in \widetilde{T}_p(\beta)$, we have $t \cdot ((y_j)_{j=1}^{2g}, (\epsilon'_m)_{m \in S_p}) = ((y_j)_{j=1}^{2g}, (\epsilon'_m)_{m \in S_p}) \Rightarrow t \in \mathbb{K}^{\times}I_n$. We use **Convention** 4. Recall by (2.3.10) that, $\widehat{y}_j = y_j$ if and only if $t^{\tau_j} = t$. By Proposition 1.16 (1), for any $m \in S_p$, $\widehat{\epsilon}'_m = \epsilon'_m$ if and only if $(t^{p_{m-1}})_{i_m} = (t^{p_{m-1}})_{i_m+1}$. Equivalently, $(t^{p_{m-1}})^{s_{i_m}} = t^{p_{m-1}}$, i.e., $t^{\underline{s}_{i_m}} = t$, where:

$$\underline{\mathbf{s}}_{i_m} := p_{m-1}^{-1} \mathbf{s}_{i_m} p_{m-1} \in S_n.$$
(2.4.11)

Thus, denote

$$J_p := \{\tau_j : 1 \le j \le 2g, \, \underline{s}_{i_m} : m \in S_p\} \subset S_n.$$
(2.4.12)

By the preparation above, we conclude that

$$PT \text{ acts freely on } \widetilde{T}_p(\beta) \iff {}^{J_p}T = \mathbb{K}^{\times}I_n \Leftrightarrow \overline{\langle J_p \rangle} = S_n.$$
(2.4.13)

On the other hand, we consider the image of $\overline{\phi}_{\vec{w},p}$. Again, we begin with some preparations:

(1) For any $\tau \in S_n$, define a subgroup of T_1 by

$$_{T}T := \{y(y^{-1})^{\tau} : y \in T\} \subset T_{1}.$$
 (2.4.14)

So, $(a \ b)T = \{K_a(\lambda)K_b(\lambda^{-1}) : \lambda \in \mathbb{K}^{\times}\}$ for any transposition $(a \ b) \in S_n$. For any $I \subset S_n$, define $_IT \subset T_1$ as the subgroup generated by $_{\tau}T, \tau \in I$, and write $_IT := \langle_{\tau}T : \tau \in I\rangle$. Equivalently,

$$_{I}T = \langle_{(a \ \tau(a))}T : a \in [n], \tau \in I \rangle = \langle \mathbf{K}_{a}(\lambda)\mathbf{K}_{\tau(a)}(\lambda^{-1}) : \lambda \in \mathbb{K}^{\times}, a \in [n], \tau \in I \rangle \subset T_{1}.$$

(2) By definition, $_{\tau}T = _{\tau^{-1}}T$. Also, as $y(y^{-1})^{\tau_1\tau_2} = (y(y^{-1})^{\tau_2})(y^{\tau_2}((y^{-1})^{\tau_2})^{\tau_1})$, we have by (1):

$$_{I}T = {}_{\langle I \rangle}T = {}_{\overline{\langle I \rangle}}T$$

(3) Let $[n] = O_1 \sqcup \cdots \sqcup O_r$ be any partition, denoted by \vec{O} . Define a subgroup

$$_{\vec{O}}T := \{t \in T : \prod_{q \in O_a} t_q = 1, \forall 1 \le a \le r\} \subset T_1.$$

Then for any subset $I \subset S_n$, we have

$${}_{I}T = {}_{\vec{O}}T \Leftrightarrow \{O_i : 1 \le i \le r\} = \{\langle I \rangle \text{-orbits on } [n]\} \Leftrightarrow \overline{\langle I \rangle} = S_{\vec{O}}.$$

Now, we see that $_{I}T = _{J}I$ if and only if $\overline{\langle I \rangle} = \overline{\langle J \rangle}$. In particular, $_{I}T = T_{1}$ if and only if $\overline{\langle I \rangle} = S_{n}$.

Now, come back to our setting. Observe that $s(\beta) = \prod_{j=1}^{g} (\tau_{2j-1}, \tau_{2j})$. Inspired by Proposition 1.16 (2) and equation (2.2.5), we define an isomorphism

$$m_p: T_1 \xrightarrow{\simeq} T_1: t \mapsto t(D_1^{-1})^{s(\beta)} \prod_{m \in S_p} (\mathbf{K}_{i_m}(-1))^{s_{>m}(\beta)}$$

So $\text{Im}(\overline{\phi}_{\vec{w},p}) = T_1$ if and only if $\text{Im}(m_p \circ \overline{\phi}_{\vec{w},p}) = T_1$. It suffices to consider the latter. For any $1 \le j \le g$, denote

$$c_{$$

In (2.2.5), observe that

$$(y_{2j-1}^{\tau_{2j-1}}(y_{2j-1}^{-1})^{\tau_{2j-1}\tau_{2j}})^{\prod_{m=1}^{j-1}(\tau_{2m-1},\tau_{2m})} = \widetilde{y}_{2j-1}(\widetilde{y}_{2j-1}^{-1})^{c_{

$$(y_{2j}^{\tau_{2j-1}}(y_{2j}^{-1})^{(\tau_{2j-1},\tau_{2j})})^{\prod_{m=1}^{j-1}(\tau_{2m-1},\tau_{2m})} = \widetilde{y}_{2j}(\widetilde{y}_{2j}^{-1})^{c_{$$$$

In Proposition 1.16 (2), for each $m \in S_p$, observe that

$$\{(\mathbf{K}_{i_m}(\epsilon'_m)\mathbf{K}_{i_m+1}(\epsilon'_m))^{s_{>m}(\beta)} = \mathbf{K}_{s_{>m}(\beta)(i_m)}(\epsilon'_m)\mathbf{K}_{s_{>m}(\beta)(i_m+1)}(\epsilon'_m) : \epsilon'_m \in \mathbb{K}^{\times}\}$$

$$= (s_{>m}(\beta)(i_m) s_{>m}(\beta)(i_m+1))T = s_{>m}(\beta)s_{i_m}s_{>m}(\beta)^{-1}T \subset T_1.$$

Now, by above, (2.2.5), and Proposition 1.16 (2), we have $\text{Im}(m_p \circ \overline{\phi}_{\vec{w},p}) = \overline{J}_p I$, where

$$\widetilde{J}_{p} := \{ c_{m}(\beta)s_{i_{m}}s_{>m}(\beta)^{-1}, m \in S_{p} \}.$$

$$(2.4.15)$$

Then by the preparation above, we conclude that

$$\operatorname{Im}(m_p \circ \overline{\phi}_{\vec{w},p}) = {}_{\widetilde{J}_p}I = T_1 \Leftrightarrow \langle \widetilde{J}_p \rangle = S_n.$$
(2.4.16)

Combined with (2.4.13), this shows that the lemma is equivalent to the following statement:

$$\overline{\langle J_p \rangle} = S_n \Leftrightarrow \overline{\langle \widetilde{J}_p \rangle} = S_n,$$

which will follow from the claim below.

Claim: we have $\langle J_p \rangle = \langle \widetilde{J}_p \rangle$.

Proof of Claim. Denote

$$J_{\tau} := \{\tau_m : 1 \le m \le 2g\} \subset S_n, \quad \widetilde{J}_{\tau} := \{\widetilde{\tau}_{2j} := c_{
Firstly, we prove by induction that, for each $1 \le i \le a$, we have$$

Firstly, we prove by induction that, for each $1 \le j \le g$, we have

$$\langle \tau_m : 1 \le m \le 2j \rangle = \langle \widetilde{\tau}_m : 1 \le m \le 2j \rangle \subset S_n.$$
(2.4.17)

In particular, j = g gives $\langle J_{\tau} \rangle = \langle \widetilde{J}_{\tau} \rangle$. For j = 1, we have $\widetilde{\tau}_2 = \tau_1 \tau_2 \tau_1^{-1}$, $\widetilde{\tau}_1 = \tau_1 \tau_2 \tau_1^{-1} \tau_2^{-1} \tau_1^{-1}$. So, $\widetilde{\tau}_2^{-1} \widetilde{\tau}_1 \widetilde{\tau}_2 = \tau_1^{-1} \in \langle \widetilde{\tau}_1, \widetilde{\tau}_2 \rangle$. It follows that $\langle \widetilde{\tau}_1, \widetilde{\tau}_2 \rangle = \langle \tau_1^{-1}, \widetilde{\tau}_2 = \tau_1 \tau_2 \tau_1^{-1} \rangle = \langle \tau_1, \tau_2 \rangle$, as desired. Suppose (2.4.17) holds for '< j', so

$$c_{$$

Observe that $\tilde{\tau}_{2j}^{-1}\tilde{\tau}_{2j-1}\tilde{\tau}_{2j} = c_{<j}\tau_{2j}^{-1}\tau_{2j-1}^{-1}(\tau_{2j-1},\tau_{2j})\tau_{2j}c_{<j}^{-1} = c_{<j}\tau_{2j-1}^{-1}c_{<j}^{-1}$. It follows that

$$\begin{aligned} &\langle \widetilde{\tau}_m : 1 \le m \le 2j \rangle = \langle \tau_m : 1 \le m \le 2(j-1), c_{$$

This finishes the induction, and hence proves (2.4.17).

Now, observe that $s(\beta) = \prod_{j=1}^{g} (\tau_{2j-1}, \tau_{2j}) \in \langle \widetilde{J}_{\tau} \rangle = \langle J_{\tau} \rangle$. Thus,

$$\langle \widetilde{J}_p \rangle = \langle J_\tau, \mathbf{s}(\beta)^{-1} \mathbf{s}_{>m}(\beta) \mathbf{s}_{i_m} \mathbf{s}_{>m}(\beta)^{-1} \mathbf{s}(\beta), m \in S_p \rangle = \langle J_\tau, \mathbf{s}_{$$

Say, $S_p = \{m_1 < \dots < m_N\}$. Set $J_s := \{\underline{s}_{i_m} : m \in S_p\} \subset W = S_n, \widetilde{J}_s := \{\widetilde{s}_{i_m} : m \in S_p\} \subset W = S_n$. We will show that

$$\langle \underline{\mathbf{s}}_{i_{m_j}}, 1 \le j \le L \rangle = \langle \widetilde{\mathbf{s}}_{i_{m_j}}, 1 \le j \le L \rangle.$$
(2.4.18)

In particular, L = N gives $\langle J_s \rangle = \langle \widetilde{J_s} \rangle$, and the Claim will follow immediately.

For that, we firstly prove by induction that, for each $1 \le j \le N$, we have

$$\prod_{q=j}^{1} \underline{\mathbf{s}}_{i_{m_q}} = p_{m_j-1}^{-1} \mathbf{s}_{< m_j+1}(\beta).$$
(2.4.19)

For j = 1, as $p_{m_1-1} = s_{<m_1}(\beta)$, we get $\prod_{q=1}^{1} \underline{s}_{i_{m_q}} = p_{m_1-1}^{-1} s_{i_{m_1}} s_{<m_1}(\beta) = p_{m_1-1}^{-1} s_{<m_1+1}(\beta)$. Done. Suppose (2.4.19) holds for ' $\leq j$ ', then

$$\prod_{q=j+1}^{1} \underline{\mathbf{s}}_{i_{m_q}} = \underline{\mathbf{s}}_{i_{m_{j+1}}} \prod_{q=j}^{1} \underline{\mathbf{s}}_{i_{m_q}} = (p_{m_{j+1}-1}^{-1} \mathbf{s}_{i_{m_{j+1}}} p_{m_{j+1}-1}) p_{m_j-1}^{-1} \mathbf{s}_{< m_j+1}(\beta) = p_{m_{j+1}-1}^{-1} \mathbf{s}_{< m_{j+1}+1}(\beta).$$

This finishes the induction, and hence proves (2.4.19).

Now, for any $1 \le j \le N$, we have

$$(\prod_{q=j-1}^{1} \underline{s}_{i_{m_{j}}})^{-1} \underline{s}_{i_{m_{j}}} (\prod_{q=j-1}^{1} \underline{s}_{i_{m_{q}}}) = (p_{m_{j-1}-1}^{-1} s_{< m_{j-1}+1}(\beta))^{-1} p_{m_{j}-1}^{-1} s_{i_{m_{j}}} p_{m_{j}-1}(p_{m_{j-1}-1}^{-1} s_{< m_{j-1}+1}(\beta))$$

= $s_{< m_{j}}(\beta)^{-1} s_{i_{m_{j}}} s_{< m_{j}}(\beta) = \widetilde{s}_{i_{m_{j}}}.$

It follows that for any $1 \le L \le N$, we have

$$\langle \underline{\mathbf{s}}_{i_{m_j}}, 1 \leq j \leq L \rangle = \langle (\prod_{q=j-1}^{1} \underline{\mathbf{s}}_{i_{m_q}})^{-1} \underline{\mathbf{s}}_{i_{m_j}} (\prod_{q=j-1}^{1} \underline{\mathbf{s}}_{i_{m_q}}), 1 \leq j \leq L \rangle = \langle \widetilde{\mathbf{s}}_{i_{m_j}}, 1 \leq j \leq L \rangle.$$

This is exactly what we want, hence the claim follows. Done.

Inspired by the lemma, we make the following

Definition 2.7. For $\beta = \beta(\vec{w})$, define

$$\mathcal{W}^*(\beta) := \{ p \in \mathcal{W}(\beta) : \overline{T}_p(\beta) \neq \emptyset \},$$
(2.4.20)

and any $p \in W^*(\beta)$ is called an *admissible walk*. By the proof of Lemma 2.6,

$$p \in \mathcal{W}^*(\beta) \Leftrightarrow \langle J_p \rangle$$
 acts transitively on $[n] \Leftrightarrow \langle J_p \rangle = S_n; \quad \forall p \in \mathcal{W}(\beta)$,
where $J_p = \{\tau_j : 1 \le j \le 2g, \underline{s}_{i_m} = p_{m-1}^{-1} s_{i_m} p_{m-1} : m \in S_p\} \subset S_n$. Alternatively, denote

$$'_p := \{\tau_j : 1 \le j \le 2g, \ \widetilde{\mathbf{s}}_{i_m} = \mathbf{s}_{< m}(\beta)^{-1} \mathbf{s}_{i_m} \mathbf{s}_{< m}(\beta) : m \in S_p\} \subset S_n.$$

We have seen that $\langle J_p \rangle = \langle J'_p \rangle$, then the same holds if we replace J_p above by J'_p .

As an immediate corollary of Lemma 2.6, we obtain

Corollary 2.8. If $\widetilde{X}_p(\beta) \neq \emptyset$, then we have a (B-equivariant) isomorphism

$$\widetilde{X}_p(\beta) = \widetilde{T}_p(\beta) \times \mathbb{K}^{|U_p|} \cong (\mathbb{K}^{\times})^{|S_p| + 2gn - n + 1} \times \mathbb{K}^{|U_p|}.$$
(2.4.21)

Proof. If $\widetilde{X}_p(\beta) \neq \emptyset$, then we know that the *PT*-action on $\widetilde{X}_p(\beta)$ is free. By Lemma 2.6, the map

$$\overline{\phi}_{\vec{w},p}: T^{2g} \times (\mathbb{K}^{\times})^{|S_p|} \cong (\mathbb{K}^{\times})^{|S_p|+2gn} \to T_1 \cong (\mathbb{K}^{\times})^{n-1}.$$

is surjective. By (2.3.3), Proposition 1.16 (2), and (2.2.5), the composition $m_p \circ \overline{\phi}_{\vec{w},p}$ is a surjective group homomorphism of algebraic tori. It follows that

$$\widetilde{T}_{p}(\beta) = \overline{\phi}_{\vec{w},p}^{-1}(I_{n}) = (m_{p} \circ \overline{\phi}_{\vec{w},p})^{-1}((D_{1}^{-1})^{s(\beta)} \prod_{m \in S_{p}} (\mathbf{K}_{i_{m}}(-1))^{s_{>m}(\beta)}) \cong (\mathbb{K}^{\times})^{|S_{p}|+2gn-n+1}.$$

Thus, $\widetilde{X}_p(\beta) = \widetilde{T}_p(\beta) \times \mathbb{K}^{|U_p|} \cong (\mathbb{K}^{\times})^{|S_p| + 2gn - n + 1} \times \mathbb{K}^{|U_p|}$, as desired.

Now, denote

$$n_{\vec{w}} := \sum_{j=1}^{2g} |U_{\tau_j}^+| + \sum_{i=1}^{k-1} |U_{\dot{w}_i}^+ \cap N_i| = 2g|U| + \sum_{i=1}^{k-1} |N_i| - \frac{1}{2}\ell(\beta).$$
(2.4.22)

By Lemma 2.5 and Corollary 2.8, we have shown the following

Proposition 2.9. We have a B-equivariant decomposition into locally closed \mathbb{K} -subvarieties

$$M''_{B}(\vec{w}) = \sqcup_{p \in \mathcal{W}^{*}(\beta)} M''_{B}(\vec{w}, p), \quad M''_{B}(\vec{w}, p) := \widetilde{X}_{p}(\beta) \times \prod_{j=1}^{s} (U^{+}_{\tau_{2j}^{-1}} \times U^{+}_{\tau_{2j-1}}) \times \prod_{i=1}^{k-1} (U^{+}_{\dot{w}_{i}} \cap N_{i}).$$

In particular,

$$M_B''(\vec{w}, p) \cong (\mathbb{K}^{\times})^{a(\vec{w}, p)} \times \mathbb{K}^{b(\vec{w}, p)} = (\mathbb{K}^{\times})^{|S_p| + 2gn - n + 1} \times \mathbb{K}^{|U_p| + n_{\vec{w}}}$$

and $a(\vec{w}, p) + 2b(\vec{w}, p) = 4g|U| + 2\sum_{i=1}^{k-1} |N_i| + 2gn - n + 1$ is a constant independent of \vec{w}, p .

Recall by (2.3.8) that, *PB* acts freely on $M''_B(\vec{w})$ and $\pi''_{\vec{w}} : M''_B(\vec{w}) \to \mathcal{M}_{\mu}(\vec{w}) = M''_B(\vec{w})/PB$ is a principal *PB*-bundle. Define

$$\mathcal{M}_{\mu}(\vec{w}, p) \coloneqq \pi_{\vec{w}}^{\prime\prime}(M_B^{\prime\prime}(\vec{w}, p)) \hookrightarrow \mathcal{M}_{\mu}(\vec{w}) \tag{2.4.23}$$

as a locally closed K-subvariety. By base change (see Corollary B.11), the restriction of $\pi''_{\vec{w}}$

$$\pi_{\vec{w},p}: M_B''(\vec{w},p) \cong (\mathbb{K}^{\times})^{a(\vec{w},p)} \times \mathbb{K}^{b(\vec{w},p)} \to \mathcal{M}_{\mu}(\vec{w},p) = M_B''(\vec{w},p)/PB_{\mu}(\vec{w},p)$$

is a principal *PB*-bundle as well as a geometric quotient.

Our first main result improves A. Mellit's cell decomposition theorem [40, §7]:

Theorem 2.10. If $(C_1, \dots, C_k) \in T^k$ is very generic (Definition 1.1 and Assumption 1.4) of type μ , and \mathcal{M}_{μ} is nonempty, then there is a decomposition into locally closed affine \mathbb{K} -subvarieties

$$\mathcal{M}_{\mu} = \bigsqcup_{\vec{w} \in W^{2g} \times \prod_{i=1}^{k-1} W/W(C_i)} \mathcal{M}_{\mu}(\vec{w}) = \bigsqcup_{\vec{w} \in W^{2g} \times \prod_{i=1}^{k-1} W/W(C_i)} \bigsqcup_{p \in W^*(\beta(\vec{w}))} \mathcal{M}_{\mu}(\vec{w}, p), \qquad (2.4.24)$$

such that:

(1) We have

$$\mathcal{M}_{\mu}(\vec{w}, p) \cong (\mathbb{K}^{\times})^{\overline{a}(\vec{w}, p)} \times \mathcal{A}_{\mu}(\vec{w}, p), \quad \mathcal{A}_{\mu}(\vec{w}, p) \times \mathbb{K}^{|U|} \cong \mathbb{K}^{b(\vec{w}, p)}.$$
(2.4.25)
$$\overline{a}(\vec{w}, p) = a(\vec{w}, p) - n + 1 = |S_p| + 2gn - 2n + 2, \quad \overline{b}(\vec{w}, p) := b(\vec{w}, p) - |U|.$$

In particular, $\mathcal{M}_{\mu}(\vec{w}, p)$ is of dimension $\overline{a}(\vec{w}, p) + \overline{b}(\vec{w}, p)$, and

$$\overline{a}(\vec{w}, p) + 2\overline{b}(\vec{w}, p) = d_{\mu} = n^2(2g - 2 + k) - \sum_{i,j} (\mu_j^i)^2 + 2k^2 + 2k^2$$

is a constant independent of (\vec{w}, p) .

(2) There exists a unique $(\vec{w}_{\max}, p_{\max})$ such that dim $\mathcal{M}_{\mu}(\vec{w}_{\max}, p_{\max})$ is of maximal dimension d_{μ} . Equivalently, $\overline{a}(\vec{w}_{\max}, p_{\max}) = d_{\mu}$ (resp. $\overline{b}(\vec{w}_{\max}, p_{\max}) = 0$). In particular, $\mathcal{M}_{\mu}(\vec{w}_{\max}, p_{\max})$ is an open dense algebraic torus:

$$\mathcal{M}_{\mu}(\vec{w}_{\max}, p_{\max}) \cong (\mathbb{K}^{\times})^{d_{\mu}}, \quad \mathcal{R}_{\mu}(\vec{w}_{\max}, p_{\max}) = \{pt\}.$$

Proof. (0). By (1.2.8) and Proposition 2.9, the decomposition of \mathcal{M}_{μ} is clear. Next, we show that $\mathcal{M}_{\mu}(\vec{w})$ is affine. Inspired by (1.2.7), we define a closed (affine) subvariety of $M'_B(\vec{w})$:

$$\underline{M}'_{B}(\vec{w}) := M'_{B} \cap (\prod_{j=1}^{2g} B\tau_{j}B \times \prod_{i=1}^{k-1} B\dot{w}_{i}P_{i} \times \{I_{n}\}) \subset M'_{B}(\vec{w}).$$

Observe that we have mutually inverse *U*-equivariant isomorphisms:

$$U \times \underline{M}'_{B}(\vec{w}) \to M'_{B}(\vec{w}) : (u, (A_{1}, \dots, x_{k-1}, I_{n})) \mapsto (uA_{1}u^{-1}, \dots, ux_{k-1}, u(u^{C_{k}})^{-1});$$
(2.4.26)
$$M'_{B}(\vec{w}) \to U \times \underline{M}'_{B}(\vec{w}) : (A_{1}, \dots, x_{k-1}, u_{k} = \widetilde{u}(\widetilde{u}^{C_{k}})^{-1}) \mapsto (\widetilde{u}, (\widetilde{u}^{-1}A_{1}\widetilde{u}, \dots, \widetilde{u}^{-1}x_{k-1}, I_{n})),$$

where $\tilde{u} \in U$ is uniquely determined by the equation $u_k = \tilde{u}(\tilde{u}^{C_k})^{-1}$. Thus, $\underline{M'_B}(\vec{w}) = M'_B(\vec{w})/U$. Recall that $U \subset PB_{\text{par}}$ is a normal closed subgroup with quotient $PB_{\text{par}}/U \cong PT_{\text{par}} = (T \times \prod_{i=1}^{k-1} Z(C_i))$. Then by Proposition B.14 and Proposition B.9, we get an isomorphism

$$\mathcal{M}_{\mu}(\vec{w}) = M'_B(\vec{w})/PB_{\text{par}} \cong \underline{M'_B}(\vec{w})/(PB_{\text{par}}/U) = \underline{M'_B}(\vec{w})/PT_{\text{par}},$$

and the quotient map $\underline{M'_B} \to \mathcal{M}_{\mu}(\vec{w}) \cong \underline{M'_B}(\vec{w})/PT_{\text{par}}$ is a principal PT_{par} -bundle as well as a geometric quotient, which is unique up to unique isomorphism. However, $\underline{M'_B}(\vec{w})$ is an affine \mathbb{K} -variety and PT_{par} is reductive, we must have

$$\mathcal{M}_{\mu}(\vec{w}) \cong \underline{M}'_{B}(\vec{w})/PT_{\text{par}} \cong \underline{M}'_{B}(\vec{w})/PT_{\text{par}} = \text{Spec } O(\underline{M}'_{B}(\vec{w}))^{PT_{\text{par}}}.$$

In particular, $\mathcal{M}_{\mu}(\vec{w})$ is affine, as desired. It suffices to prove (1) and (2).

(1). We firstly show that $\mathcal{M}_{\mu}(\vec{w}, p)$ is affine, the argument is similar to (0).

Recall by (2.3.1), we have a PB_{par} -equvariant isomorphism of \mathbb{K} -varieties

$$\begin{split} &(\prod_{j=1}^{2g} B\tau_{j}B) \times (\prod_{i=1}^{k-1} B\dot{w}_{i}P_{i}) \times U \xrightarrow{\cong} \\ &U \times \prod_{j=1}^{g} (T^{2} \times (U_{\tau_{2j-1}}^{-} \times U_{\tau_{2j}}^{-} \times U_{\tau_{2j-1}}^{-} \times U_{\tau_{2j-1}}^{-}) \times (U_{\tau_{2j}^{-1}}^{+} \times U_{\tau_{2j-1}}^{+})) \times \prod_{i=1}^{k-1} (U_{\dot{w}_{i}}^{-} \times U_{\dot{w}_{i}^{-1}}^{-} \times (U_{\dot{w}_{i}}^{+} \cap N_{i}) \times Z(C_{i})), \\ &((A_{j})_{j=1}^{2g}, (x_{i})_{i=1}^{k-1}, u_{k}) \mapsto (u^{0}, (y_{2j-1}, y_{2j}, \zeta_{2j-1}^{\prime}, \zeta_{2j}^{\prime}, \zeta_{2j-1}, \zeta_{2j}, +n^{2j}, +n^{2j-1})_{j=1}^{g}, (\xi_{i}^{\prime}, \xi_{i}, +n_{i}^{\prime}, z_{i})_{i=1}^{k-1}) \end{split}$$

Then, $M'_B(\vec{w})$ is a closed affine PB_{par} -subvariety of the latter. Also, $M'_B(\vec{w}, p) := M''_B(\vec{w}, p) \times_{\prod_{i=1}^{k-1}} Z(C_i) \subset M'_B(\vec{w}) = M''_B(\vec{w}) \times_{\prod_{i=1}^{k-1}} Z(C_i)$ is a locally closed affine PB_{par} -subvariety. Define

$$\underline{M}'_B(\vec{w}, p) := M'_B(\vec{w}, p) \cap \left(\prod_{j=1}^{2g} B\tau_j B \times \prod_{i=1}^{k-1} B\dot{w}_i P_i \times \{I_n\}\right) \subset M'_B(\vec{w})$$

as a closed subvariety. By the same formula (2.4.26), we obtain an U-equivariant isomorphism:

$$U \times \underline{M'_B}(\vec{w}, p) \cong M'_B(\vec{w}, p).$$

Thus, $M'_B(\vec{w}, p)/U \cong \underline{M}'_B(\vec{w}, p)$ is affine. Recall that $\prod_{i=1}^{k-1} Z(C_i) = (\mathbb{K}^{\times} \times \prod_{i=1}^{k-1} Z(C_i))/\mathbb{K}^{\times} \triangleleft PB_{\text{par}}$ with quotient $PB_{\text{par}}/\prod_{i=1}^{k-1} Z(C_i) = PB$. Similar to (0), by Proposition B.14, we have isomorphisms

$$\mathcal{M}_{\mu}(\vec{w},p) = M''_{B}(\vec{w},p)/PB \cong M'_{B}(\vec{w},p)/PB_{\text{par}} \cong \underline{M'}_{B}(\vec{w},p)/PT_{\text{par}} \cong \text{Spec } O(\underline{M'}_{B}(\vec{w},p))^{PT_{\text{par}}}.$$

In particular, $\mathcal{M}_{\mu}(\vec{w},p)$ is affine, as desired.

Secondly, by definition, the quotient map $M_B''(\vec{w}, p) \to \mathcal{M}_\mu(\vec{w}, p) = M_B''(\vec{w}, p)/PB$ is a principal *PB*bundle. Notice that $U \hookrightarrow PB$ is a normal closed subgroup with quotient $PB/U \cong PT$. Then by Proposition B.14 and Proposition B.9, the quotient map

$$\mathfrak{p}_U: M''_B(\vec{w}, p) \cong (\mathbb{K}^{\times})^{a(\vec{w}, p)} \times \mathbb{K}^{b(\vec{w}, p)} \to \mathcal{P}(\vec{w}, p) \coloneqq M''_B(\vec{w}, p)/U$$

is a principal U-bundle, and the quotient map

$$q_U: \mathcal{P}(\vec{w}, p) \to \mathcal{M}_{\mu}(\vec{w}, p) = M_B''(\vec{w}, p)/PB$$

is a principal *PT*-bundle. By Lemma B.8, q_U is affine, then so is $\mathcal{P}(\vec{w}, p)$, as $\mathcal{M}_{\mu}(\vec{w}, p)$ is.

Recall that U only acts on the factor $\mathbb{K}^{b(\vec{w},p)}$ of $M''_B(\vec{w},p)$ (see also Lemma 2.11 below). Consider the closed U-subvariety $\{1\}\times\mathbb{K}^{b(\vec{w},p)} \subset M''_B(\vec{w},p)$, by Corollary B.11 and Proposition B.9,

$$\mathfrak{p}_U|_{\mathbb{K}^{b(\vec{w},p)}}:\mathbb{K}^{b(\vec{w},p)}\cong\{1\}\times\mathbb{K}^{b(\vec{w},p)}\to\mathcal{A}_{\mu}(\vec{w},p):=\mathfrak{p}_U(\mathbb{K}^{b(\vec{w},p)})\cong\mathbb{K}^{b(\vec{w},p)}/U\subset\mathcal{P}(\vec{w},p)$$

is a principal U-bundle as well as a geometric quotient. Also, $\mathcal{A}_{\mu}(\vec{w}, p) \subset \mathcal{P}(\vec{w}, p)$ is a closed subvariety, hence affine. Clearly, $\mathcal{A}_{\mu}(\vec{w}, p)$ is connected of dimension $\overline{b}(\vec{w}, p) = b(\vec{w}, p) - |U|$. Now, by Proposition B.15, $\mathfrak{p}_{U}|_{\mathbb{K}^{b(\vec{w},p)}}$ is trivial, i.e. we obtain an U-equivariant isomorphism

$$U \times \mathcal{A}_{\mu}(\vec{w}, p) \cong \mathbb{K}^{b(\vec{w}, p)}$$

This proves the second isomorphism in (2.4.25).

Thirdly, let's prove the first isomorphism in (2.4.25). By the uniqueness of geometric quotient, we obtain an isomorphism

$$\mathcal{P}(\vec{w},p) = (\mathbb{K}^{\times})^{a(\vec{w},p)} \times \mathbb{K}^{b(\vec{w},p)})/U \cong (\mathbb{K}^{\times})^{a(\vec{w},p)} \times (\mathbb{K}^{b(\vec{w},p)}/U) = (\mathbb{K}^{\times})^{a(\vec{w},p)} \times \mathcal{A}_{\mu}(\vec{w},p).$$

Recall by (2.4.4) that *PT* acts freely on the torus factor $\widetilde{T}_p(\beta(\vec{w})) \cong (\mathbb{K}^{\times})^{a(\vec{w},p)} = \mathbb{K}^{|S_p|+2gn-n+1}$ of $M''_B(\vec{w},p) \cong (\mathbb{K}^{\times})^{a(\vec{w},p)} \times \mathbb{K}^{b(\vec{w},p)}$. By Lemma 2.11 below, this is equivalent to an injective algebraic group homomorphism $PT \cong (\mathbb{K}^{\times})^{n-1} \hookrightarrow (\mathbb{K}^{\times})^{a(\vec{w},p)}$. So, up to a coordinate change, we may assume $(\mathbb{K}^{\times})^{a(\vec{w},p)} \cong (\mathbb{K}^{\times})^{\overline{a}(\vec{w},p)} \times PT$, where $\overline{a}(\vec{w},p) = a(\vec{w},p) - n + 1$, and *PT* acts via translation on the second factor. Now, we have

$$\mathcal{M}_{\mu}(\vec{w},p) = \mathcal{P}(\vec{w},p)/PT \cong (\mathbb{K}^{\times})^{\overline{a}(\vec{w},p)} \times (PT \times \mathcal{A}_{\mu}(\vec{w},p))/PT \cong (\mathbb{K}^{\times})^{\overline{a}(\vec{w},p)} \times \mathcal{A}_{\mu}(\vec{w},p),$$

as desired. Here, in the last step: By Proposition B.9, the natural map $(PT \times \mathcal{A}_{\mu}(\vec{w}, p))/PT \rightarrow PT/PT =$ Spec K is a fiber bundle with fiber $\mathcal{A}_{\mu}(\vec{w}, p)$, hence $(PT \times \mathcal{A}_{\mu}(\vec{w}, p))/PT \cong \mathcal{A}_{\mu}(\vec{w}, p)$.

Finally, it remains to compute $\overline{a}(\vec{w}, p) + 2\overline{b}(\vec{w}, p)$. By Proposition 2.9, we have

$$\begin{aligned} \overline{a}(\vec{w},p) + 2\overline{b}(\vec{w},p) &= a(\vec{w},p) - n + 1 + 2b(\vec{w},p) - 2|U| \\ &= (4g|U| + 2\sum_{i=1}^{k-1}|N_i| + 2gn - n + 1) - n + 1 - 2|U| = (4g - 4)|U| + 2\sum_{i=1}^{k}|N_i| + 2gn - 2n + 2 \\ &= (2g - 2)(n^2 - n) + \sum_{i=1}^{k}(n^2 - \sum_{j}(\mu_j^i)^2) + 2gn - 2n + 2 = (2g - 2 + k)n^2 - \sum_{i,j}(\mu_j^i)^2 + 2 = d_{\mu}, \end{aligned}$$

which is clearly independent of \vec{w} , p. This completes the proof of (1).

(2). By Lemma 1.2, \mathcal{M}_{μ} (if nonempty) is connected smooth affine of dimension d_{μ} . So, in the decomposition, there exists a unique (\vec{w}, p) such that $\mathcal{M}_{\mu}(\vec{w}, p) \subset \mathcal{M}_{\mu}$ is (open dense) of dimension d_{μ} . Moreover, for all (\vec{w}, p) in the decomposition,

$$\dim \mathcal{M}_{\mu}(\vec{w}, p) = \overline{a}(\vec{w}, p) + \overline{b}(\vec{w}, p) \le \overline{a}(\vec{w}, p) + 2\overline{b}(\vec{w}, p) = d_{\mu}.$$

Thus, dim $\mathcal{M}_{\mu}(\vec{w}, p) = d_{\mu} \Leftrightarrow \overline{b}(\vec{w}, p) = 0 \Leftrightarrow \overline{a}(\vec{w}, p) = 0$. The rest follows immediately. Done.

Lemma 2.11. Let *H* be a connected algebraic group and $T' = (\mathbb{K}^{\times})_{z_1,\dots,z_m}^m$ is an algebraic torus, then any algebraic *H*-action on *T'* is identical to an algebraic group homomorphism

$$\rho: H \to T',$$

where T' acts on itself by translation. Thus, if H is unipotent, then the H-action on T' is trivial.

T. SU

Proof. For any $a \in H$, the *H*-action ρ on *T'* induces an isomorphism

$$\rho_a: T' \xrightarrow{\simeq} T': z_j \mapsto c_i(a) \prod_{i=1}^m z_i^{a_{ij}},$$

for some $(a_{ij}) \in GL(n, \mathbb{Z})$ and $c_i(a) \in \mathbb{K}^{\times}$. As $\rho_1 = id$, by continuity, $a_{ij} = \delta_{ij}$. So, $\rho : H \to T' : a \mapsto \rho_a = \text{Diag}(c_1(a), \dots, c_m(a))$ becomes an algebraic group morphism. This gives the desired identification. In addition, if *H* is unipotent, then any $a \in H$ is unipotent, hence so is $\rho_a \in T'$, which is also semisimple. It follows that $\rho_a = id$, $\forall a \in H$. We're done. \Box

Question: If μ is only generic, does \mathcal{M}_{μ} contain an open algebraic torus?

Remark 2.12. If $(\vec{w}, p) \neq (\vec{w}_m, p_m)$ in Theorem 2.10, then $\mathcal{A}_{\mu}(\vec{w}, p)$ is smooth affine of dimension $\overline{b}(\vec{w}, p) > 0$, and $\mathcal{A}_{\mu}(\vec{w}, p)$ is *stably isomorphic* to $\mathbb{A}_{\mathbb{K}}^{\overline{b}(\vec{w}, p)} : \mathcal{A}_{\mu}(\vec{w}, p) \times \mathbb{A}_{\mathbb{K}}^{\frac{n^2 - n}{2}} \cong \mathbb{A}_{\mathbb{K}}^{\overline{b}(\vec{w}, p) + \frac{n^2 - n}{2}}$. This is closely related to the famous **Zariski cancellation problem (ZCP)**:

If *Y* is an affine \mathbb{K} -variety of dimension *d* such that $Y \times \mathbb{A}^1 \cong \mathbb{A}^{d+1}$, is *Y* always isomorphic to \mathbb{A}^d ? The answer is positive if d = 1, 2 [1, 20, 44, 49], and negative for $d \ge 3$ in positive characteristic [24, 25]; For $d \ge 3$ and char $\mathbb{K} = 0$, the problem is still open.

We tend to believe that $\mathcal{A}_{\mu}(\vec{w}, p)$ is not isomorphic to $\mathbb{A}_{\mathbb{K}}^{\overline{b}(\vec{w}, p)}$ in general, thus providing a systematical way of constructing counterexamples to **ZCP** for $d \ge 3$ and char $\mathbb{K} = 0$. For example, take $(g, k, n, \mu) = (0, 6, 2, ((1, 1)^6))$, by a computation similar to Example 2.13 below, we obtain a cell decomposition of \mathcal{M}_{μ} :

$$\mathcal{M}_{\mu} = (\mathbb{K}^*)^6 \sqcup ((\mathbb{K}^*)^4 \times \mathbb{K})^{\sqcup 12} \sqcup ((\mathbb{K}^*)^2 \times \mathbb{K}^2)^{\sqcup 40} \sqcup \sqcup_{j=1}^{80} \mathcal{A}_j; \quad \mathcal{A}_j \times \mathbb{K} \cong \mathbb{K}^4, \forall 1 \le j \le 80.$$

In other words, the \mathcal{A}_i 's are 80 potential counterexamples to the 3-dim **ZCP** in characteristic 0.

2.5. Examples. We illustrate Theorem 2.10 by two examples.

Example 2.13 $((g, k, n, \mu) = (0, 4, 2, ((1^2), (1^2), (1^2)))$: Fricke-Klein cubic). Let $\Sigma_{0,4} := (\Sigma_0, \sigma = \{q_1, q_2, q_3, q_4\})$ be a four-punctured two-sphere, $G = GL_2(\mathbb{K})$. So, $T \cong (\mathbb{K}^{\times})^2$, and $W = S_2 = \{1, s_1 = (1 2)\}$. Let $\mu = ((1)^2, (1)^2, (1)^2, (1)^2)$, so μ is very generic (Definition 1.1, Assumption 1.4) and

$$C_{i} = \text{Diag}(a_{i,1}, a_{i,2}) \in T, a_{i,1} \neq a_{i,2}, \quad 1 \le i \le 4; \quad \prod_{i,j} a_{i,j} = 1, \quad \prod_{i=1}^{4} a_{i,\psi_{i}(1)} \neq 1, \forall \psi_{i} \in W.$$
(2.5.1)

Clearly, dim $\mathcal{M}_{\mu} = d_{\mu} = n^2(2g - 2 + k) - \sum_{i=1}^{k} \sum_{j} (\mu_j^i)^2 + 2 = 2$. The example goes back to [18]. 1) We firstly compute the cell decomposition of \mathcal{M}_{μ} (Theorem 2.10):

$$\begin{split} \mathcal{M}_{\mu} &= \sqcup_{(\vec{w},p) \in \mathcal{W}^*} \mathcal{M}_{\mu}(\vec{w},p), \quad \mathcal{W}^* := \{ (\vec{w},p) : \vec{w} \in W^{k-1} = W^3, p \in \mathcal{W}^*(\beta(\vec{w})) \}, \\ \mathcal{M}_{\mu}(\vec{w},p) &\cong (\mathbb{K}^{\times})^{\overline{a}(\vec{w},p)} \times \mathcal{A}_{\mu}(\vec{w},p), \quad \mathcal{A}_{\mu}(\vec{w},p) \times \mathbb{A}^{|U|} \cong \mathbb{A}^{\overline{b}(\vec{w},p)+|U|}, \end{split}$$

where |U| = 1, and

$$\overline{a}(\vec{w},p) = |S_p| + 2gn - 2n + 2 = |S_p| - 2 \ge 0, \quad \overline{b}(\vec{w},p) = \frac{1}{2}(d_\mu - \overline{a}(\vec{w},p)) = \frac{1}{2}(4 - |S_p|) \ge 0.$$

Thus, $|S_p| = 2$ or 4. Accordingly, $(\overline{a}(\vec{w}, p), \overline{b}(\vec{w}, p)) = (0, 1)$ or (2, 0). By the affirmative answer [1, 20, 44, 49] to the Zariski cancellation problem in dim ≤ 2 , we have

$$\mathcal{A}_{\mu}(\vec{w},p) \cong \mathbb{A}^{\overline{b}(\vec{w},p)} \implies \mathcal{M}_{\mu}(\vec{w},p) \cong (\mathbb{K}^{\times})^{\overline{a}(\vec{w},p)} \times \mathbb{K}^{\overline{b}(\vec{w},p)} = \mathbb{K} \text{ or } (\mathbb{K}^{\times})^{2}.$$

We would like to compute \mathcal{W}^* . For each $\vec{w} = (\dot{w}_1, \dot{w}_2, \dot{w}_3) = (w_1, w_2, w_3) \in W^{k-1} = W^3 = S_2^3$, denote $\beta := \beta(\vec{w})$ and $\ell := \ell(\beta)$. Recall that, for any $p \in \mathcal{W}(\beta) \subset W^{\ell+1}$, denote

$$J_p = \{\underline{\mathbf{s}}_{i_m} = p_{m-1}^{-1} \mathbf{s}_{i_m} p_{m-1}, m \in S_p\} \subset W,$$

then $p \in \mathcal{W}^*(\beta)$ if and only if the group $\langle J_p \rangle$ acts transitively on $[2] = \{1, 2\}$. **Note:** In this case, it means that $\langle J_p \rangle = W$, equivalently, $S_p \neq \emptyset$. That is,

$$W^*(\beta) = \{ p \in \mathcal{W}(\beta) : S_p \neq \emptyset \} \subset \mathcal{W}(\beta).$$

Recall that, $\beta = \beta(\vec{w}) = [w_1] \circ [w_1^{-1}] \circ [w_2] \circ [w_2^{-1}] \circ [w_3] \circ [w_3^{-1}]$, so $\ell = \ell(\beta) = 2\ell(w_1) + 2\ell(w_2) + 2\ell(w_3)$. If $\exists p \in \mathcal{W}(\beta)$ such that $S_p \neq \emptyset$, then $\ell(\beta) > 0$, so $\ell(\beta) \ge 2$, and $w_i = s_1 = (1 \ 2)$ for at least one $i \in \{1, 2, 3\}$. Recall that p is of the form $(p_\ell = \mathsf{id}, \dots, p_1, p_0 = \mathsf{id}) \in W^{\ell+1}$. By Definition 1.15 of a walk, there's no *i* such that $(p_{i+1}, p_i) = (id, id)$ (if we could go up, then we must go up). In particular, we must have $p_1 = s_1, p_{\ell-1} = s_1$. As $S_p \neq \emptyset$, we must have $\ell \ge 4$. This means that $w_i = s_1 = (1 \ 2)$ for at least two $i \in \{1, 2, 3\}$. We're left with 4 cases:

(a) $\vec{w} = (s_1, s_1, id) \in W^3$. Then, $\beta := \beta(\vec{w}) = \sigma_1^4 \in FBr_2^+$ and $\ell = \ell(\beta) = 4$. Observe that

$$\mathcal{W}(\beta) = \{ (\mathsf{id}, \mathsf{s}_1, \mathsf{id}, \mathsf{s}_1, \mathsf{id}), p^1 = (\mathsf{id}, \mathsf{s}_1, \mathsf{s}_1, \mathsf{s}_1, \mathsf{id}) \} \subset W^{\ell+1} = W^5.$$

Then $\mathcal{W}^*(\beta) = \{p^1\}$, with $S_{p^1} = \{2, 3\} \subset [\ell] = [4] = \{1, 2, 3, 4\}$, $U_{p^1} = \{1\} \subset [4]$, and $D_{p^1} = \{4\} \subset [4]$. Denote $\vec{w}^1 := (s_1, s_1, id)$. By the previous computation, we have

$$\overline{a}(\vec{w}^1, p^1) = |S_{p^1}| - 2 = 0, \ \overline{b}(\vec{w}^1, p^1) = \frac{1}{2}(4 - |S_{p^1}|) = 1, \ \mathcal{M}_{\mu}(\vec{w}^1, p^1) \cong \mathbb{K}.$$

(b) Similarly, for $\vec{w}^2 = (s_1, id, s_1) \in W^3$ (resp. $\vec{w}^3 = (id, s_1, s_1) \in W^3$), we have $\mathcal{W}^*(\beta(\vec{w}^2)) = \{p^2 = (id, s_1, s_1, s_1, id)\}$ (resp. $\mathcal{W}^*(\beta(\vec{w}^3)) = \{p^3 = (id, s_1, s_1, s_1, id)\}$), and

$$\mathcal{M}_{\mu}(\vec{w}^2, p^2) \cong \mathbb{K}, \quad \mathcal{M}_{\mu}(\vec{w}^3, p^3) \cong \mathbb{K}.$$

(c) $\vec{w} = (s_1, s_1, s_1) =: \vec{w}^4 \in W^3$. Then, $\beta := \beta(\vec{w}^4) = \sigma_1^6 \in FBr_2^+$ and $\ell(\beta) = 6$. Observe that

$$(\beta) = \{ (\mathsf{id}, \mathsf{s}_1, \mathsf{id}, \mathsf{s}_1, \mathsf{id}, \mathsf{s}_1, \mathsf{id}), p^4 = (\mathsf{id}, \mathsf{s}_1, \mathsf{s}_1, \mathsf{s}_1, \mathsf{id}, \mathsf{s}_1, \mathsf{id}), p^5 = (\mathsf{id}, \mathsf{s}_1, \mathsf{s}_1, \mathsf{id}, \mathsf{s}_1, \mathsf{s}_1, \mathsf{id}), \\ p^6 = (\mathsf{id}, \mathsf{s}_1, \mathsf{id}, \mathsf{s}_1, \mathsf{s}_1, \mathsf{s}_1, \mathsf{id}), p^7 = (\mathsf{id}, \mathsf{s}_1, \mathsf{s}_1, \mathsf{s}_1, \mathsf{s}_1, \mathsf{s}_1, \mathsf{s}_1, \mathsf{id}) \} \subset W^{\ell+1} = W^7.$$

Thus, $\mathcal{W}^*(\beta) = \{p \in \mathcal{W}(\beta) : S_p \neq \emptyset\} = \{p^j : 4 \le j \le 7\}$. Denote $\vec{w}^j := (s_1, s_1, s_1), 4 \le j \le 7$. So, $\mathcal{M}_{(\vec{w}^j, p^j)} \cong \mathbb{K} \quad \forall A \le i \le 6; \quad \mathcal{M}_{(\vec{w}^j, p^j)} \simeq (\mathbb{K}^{\times})^2$

$$\mathcal{M}_{\mu}(w^{j}, p^{j}) \cong \mathbb{K}, \forall 4 \leq j \leq 6; \quad \mathcal{M}_{\mu}(w^{j}, p^{j}) \cong (\mathbb{K}^{\wedge})^{2}.$$

In summary, $\mathcal{W}^* = \{(\vec{w}^j, p^j), 1 \le j \le 7\}$, and the cell decomposition of \mathcal{M}_{μ} reads:

$$\mathcal{M}_{\boldsymbol{\mu}} = \sqcup_{j=1}^{7} \mathcal{M}_{\boldsymbol{\mu}}(\vec{w}^{j}, p^{j}) = \mathbb{K}^{\sqcup 6} \sqcup (\mathbb{K}^{\times})^{2}.$$

The order on indices is *admissible* (Definition 3.12). Our example matches with [40, §1.4].

2) Let's give a concrete description of the variety \mathcal{M}_{μ} . The defining equation of M_B reads

$$x_1C_1x_1^{-1}x_2C_2x_2^{-1}x_3C_3x_3^{-1}x_4C_4x_4^{-1} = id, \quad x_i \in G.$$

with an action of $G_{par} = G \times T^4$ via conjugation:

$$h \cdot (x_1, x_2, x_3, x_4) = (h_0 x_1 h_1^{-1}, h_0 x_2 h_2^{-1}, h_0 x_3 h_3^{-1}, h_0 x_4 h_4^{-1}), \quad h = (h_0, h_1, h_2, h_3, h_4) \in G \times T^4.$$

Denote

W

$$\underline{M}'_B := M_B \cap (G^3 \times \{I_2\}) \subset M_B,$$

so the defining equation of \underline{M}'_{B} becomes

$$x_1C_1x_1^{-1}x_2C_2x_2^{-1}x_3C_3x_3^{-1}C_4 = id, \quad x_1, x_2, x_3 \in G.$$

Now by Proposition B.9 and Proposition B.12, we have

$$\mathcal{M}_{\mu} = M_B / PG_{\text{par}} \cong \underline{M}'_B / PT_{\text{par}}, \quad PT_{\text{par}} = T^4 / \mathbb{K}^{\times} \hookrightarrow PG_{\text{par}} = (G \times T^4) / \mathbb{K}^{\times},$$

where $PT_{\text{par}} \hookrightarrow PG_{\text{par}} : (h_0, h_1, h_2, h_3) \mapsto (h_0, h_1, h_2, h_3, h_0)$ acts on \underline{M}'_B by:

$$(h_0, h_1, h_2, h_3) \cdot (x_1, x_2, x_3) := (h_0 x_1 h_1^{-1}, h_0 x_2 h_2^{-1}, h_0 x_3 h_3^{-1}).$$

Denote $X^i := x_i C_i x_i^{-1} \in G \cdot C_i \cong G/T$. Clearly, $G \cdot C_i$ is affine. Define

$$\underline{M}'_{B}(\vec{X}) := \{ (X^{i})_{i=1}^{3} \in \prod_{i=1}^{3} G \cdot C_{i} : X^{1} X^{2} X^{3} C_{4} = \mathsf{id} \},\$$

equipped with the action of $T \ni h_0$ via conjugation. Then we obtain a cartesian diagram

$$\underbrace{\begin{array}{ccc} \underline{M}'_B & & & \\ & \downarrow & & \\ & \downarrow & & \\ \underline{M}'_B(\vec{X}) & & & \\ \end{array} \\ \end{array} \xrightarrow{\begin{array}{c} G^3 \\ & \downarrow & \\ \end{array} \\ G \cdot C_i \cong (G/T)^2$$

By base change, $\underline{M'_B} \to \underline{M'_B}(\vec{X})$ is a principal T^3 -bundle. As $T^3 \hookrightarrow PT_{\text{par}} : (h_1, h_2, h_3) \mapsto [\text{id}, h_1, h_2, h_3]$ is a normal subgroup with quotient $PT \ni [h_0]$, by Proposition B.14, we have

$$\mathcal{M}_{\mu} \cong \underline{M}'_{B}/PT_{\text{par}} \cong (\underline{M}'_{B}/T^{3})/PT \cong \underline{M}'_{B}(\vec{X})/PT$$

Clearly, the conjugate action of $h_0 = \text{Diag}(h_{0,1}, h_{0,2}) \in T$ on X^i is:

$$h_0 \cdot X^i = \begin{pmatrix} X_{11}^i & h_{0,1} X_{12}^i h_{0,2}^{-1} \\ h_{0,2} X_{21}^i h_{0,1}^{-1} & X_{22}^i \end{pmatrix}.$$

Denote

$$y_3 := \operatorname{Tr}(X^1 X^2 = (X^3 C_4)^{-1}), y_1 := \operatorname{Tr}(X^2 X^3 = (C_4 X^1)^{-1}), y_2 := \operatorname{Tr}(C_4^{-1} X^3 C_4 X^1 = C_4^{-1} (X^2)^{-1}).$$

Then a direct computation shows that, \mathcal{M}_{μ} is a (smooth) affine cubic surface defined by

$$\sum_{i=1}^{3} \det C_{i} y_{i}^{2} + y_{1} y_{2} y_{3} - \det C_{4}^{-1} \sum_{\text{cyclic rotation on } 1,2,3} (\text{Tr}C_{4}\text{Tr}C_{1} + \text{Tr}C_{2}^{-1}\text{Tr}C_{3}^{-1}) y_{1}$$
(2.5.2)
+
$$\det C_{4}^{-1} (\sum_{i=1}^{4} \text{Tr}C_{i}\text{Tr}C_{i}^{-1} + \text{Tr}C_{1}\text{Tr}C_{2}\text{Tr}C_{3}\text{Tr}C_{4} - 4) = 0.$$

Note: when det $C_i = 1, \forall 1 \le i \le 4$, this recovers the Fricke-Klein cubic [18], [21, §5].

Next, we consider a rank 3 example.

Example 2.14 ((g, k, n, μ) = (0, 3, 3, ((1³), (1³), (1³)))). Let $\Sigma_{0,3} := (\Sigma_0, \sigma = \{q_1, q_2, q_3\})$ be the pair of pants, $G = GL_3(\mathbb{K})$. So, $W = S_3$. Let $\mu = ((1^3), (1^3), (1^3))$. So, μ is very generic and

$$C_i = \text{Diag}(a_{i,1}, a_{i,2}, a_{i,3}) \in T, \ a_{i,1}, a_{i,2}, a_{i,3}$$
: pairwise distinct; $\prod_{i,j} a_{i,j} = 1, \ \prod_{i=1}^3 a_{i,\psi_i(1)} \neq 1, \forall \psi_i \in W.$

Clearly, dim $\mathcal{M}_{\mu} = d_{\mu} = n^2 (2g - 2 + k) - \sum_{i=1}^k \sum_j (\mu_j^i)^2 + 2 = 9 - 3 \times 3 + 2 = 2.$

By a similar computation, the cell decomposition of \mathcal{M}_{μ} reads:

$$\mathcal{M}_{\mu} = \sqcup_{j=1}^{9} \mathcal{M}_{\mu}(\vec{w}^{j}, p^{j}) = \mathbb{K}^{\sqcup 8} \sqcup (\mathbb{K}^{\times})^{2}.$$
(2.5.3)

We have ordered the indices so that we get an admissible total order (Definition 3.12).

3. DUAL BOUNDARY COMPLEXES OF CHARACTER VARIETIES

3.1. Preliminaries on dual boundary complexes.

3.1.1. *Setup.* Let *X* be a smooth quasi-projective \mathbb{K} -variety.

Definition 3.1. A *log compactification* of X is a smooth projective variety \overline{X} with simple normal crossing (snc) boundary divisor $\partial X = \overline{X} \setminus X$. Moreover, we say that ∂X is *very simple normal crossing*, if every nonempty finite intersection of its irreducible components is *connected*.

By Hironaka's resolution of singularities, a log compactification $(\overline{X}, \partial X)$ always exists. By blowing up further if necessary, ∂X will be very simple normal crossing. Say, we're in this case.

Definition 3.2. The *dual boundary complex* $\mathbb{D}\partial X$ is a simplicial complex such that:

• Vertices of $\mathbb{D}\partial X$ are in one-to-one correspondence with the irreducible components of ∂X ;

• k vertices of $\mathbb{D}\partial X$ spans a (k-1)-simplex if and only if the corresponding irreducible components have non-empty intersection.

Proposition 3.3 ([10]). *The homotopy type of* $\mathbb{D}\partial X$ *is an invariant of* X*. i.e. independent of the choice of the log compactification* ($\overline{X}, \partial X$).

Example 3.4. For any two quasi-projective smooth varieties X, Y, take the log compactifications $\overline{X}, \overline{Y}$ separately, then $\overline{X} \times \overline{Y}$ is a log compactification of $X \times Y$ with $\partial(X \times Y) = \partial X \times \overline{Y} \cup_{\partial X \times \partial Y} \overline{X} \times \partial Y$. By a direct calculation, we then have a homotopy equivalence:

$$\mathbb{D}\partial(X \times Y) \sim \mathbb{D}\partial X \star \mathbb{D}\partial Y, \tag{3.1.1}$$

where ' \star ' stands for the *join* of simplicial complexes. Clearly, $\mathbb{D}\partial \mathbb{K} \sim *$ and $\mathbb{D}\partial \mathbb{K}^* \sim S^0$. Thus,

$$\mathbb{D}\partial(\mathbb{A}^1 \times Y) \sim *$$

i.e. the dual boundary complex of $\mathbb{A}^1 \times Y$ is contractible. As another example, we have

$$\mathbb{D}\partial(\mathbb{K}^{\times})^{d} \sim S^{0} \star \mathbb{D}\partial(\mathbb{K}^{\times})^{d-1} \sim \cdots \sim S^{d-1}.$$

The cohomology with rational coefficients of a dual boundary complex is computed by:

Proposition 3.5 (See e.g. [48]). Let X be a connected smooth quasi-projective complex variety of dimension d, then the reduced rational (co)homology of the dual boundary complex corresponds to one piece of the weight filtration:

$$\widetilde{H}_{i-1}(\mathbb{D}\partial X,\mathbb{Q}) \cong \mathrm{Gr}_{2d}^{W} H^{2d-i}(X(\mathbb{K}),\mathbb{Q}), \quad \widetilde{H}^{i-1}(\mathbb{D}\partial X,\mathbb{Q}) \cong \mathrm{Gr}_{0}^{W} H_{c}^{i}(X(\mathbb{K}),\mathbb{Q}).$$

The latter is equivalent to the former by Poincaré duality. More recently, the author has provided a motivic generalization, encoding the integral cohomology of the dual boundary complex:

Proposition 3.6 ([59, Prop.0.2]). For any smooth complex quasi-projective variety X, we have

$$H^{i-1}(\mathbb{D}\partial X;\mathbb{Z}) \cong H^{i,0}_{W,c}(X;\mathbb{Z}), \tag{3.1.2}$$

where $H^{a,b}_{W,c}(X;\mathbb{Z})$ is the integral singular weight cohomology with compact support of X defined via motives.

3.1.2. A remove/reduction lemma. In a case, one can simplify the computation of dual boundary complexes: **Lemma 3.7.** [55, Lem.2.3] Let X be a smooth irreducible quasi-projective \mathbb{K} -variety, and $Z \subset X$ be a smooth irreducible closed subvariety of smaller dimension with complement U. If N_Z is the normal bundle of Z in X, then we have a natural homotopy cofiber sequence

$$\mathbb{D}\partial Z \sim \mathbb{D}\partial \mathbb{P}(N_Z) \to \mathbb{D}\partial X \sim \mathbb{D}\mathrm{Bl}_Z(X) \to \mathbb{D}\partial U.$$

In particular, if $\mathbb{D}\partial Z \sim pt$, then the natural map $\mathbb{D}\partial X \to \mathbb{D}\partial U$ is a homotopy equivalence.

By an inductive procedure, one may obtain a further simplification:

Lemma 3.8. [55, Prop.2.6] Let X be a smooth irreducible quasi-projective \mathbb{K} -variety with a non-empty open subset U. If the complement $Z = X \setminus U$ admits a finite decomposition into locally closed smooth subvarieties Z_j such that: $\mathbb{D}\partial Z_j \sim *$; there is a total order on the indices such that $\cup_{j \leq a} Z_j$ is closed for all a. Then the natural map $\mathbb{D}\partial X \to \mathbb{D}\partial U$ is a homotopy equivalence.

This will be our key tool for computing the dual boundary complex of the character variety \mathcal{M}_{μ} .

3.1.3. A fibration over the dual boundary complex. For completeness, we review "the fibration at infinity" associated to any connected smooth quasi-projective \mathbb{K} -variety, say X. For $X = \mathcal{M}_B$, such a fibration α appeared in the statement of the geometric P=W conjecture 0.1.

Fix any log compactification $(\overline{X}, D = \partial X)$ with D very simple normal crossing. Fix a Riemann metric on \overline{X} . For $0 < \delta \ll 1$, let $N_{\delta}(D) \subset \overline{X}$ be the δ -neighborhood of D. Let $D_i, i \in I = \{1, 2, \dots, m\}$ be the irreducible components of D. For each i, let $N_{\delta}(D_i) \subset \overline{X}$ be the δ -neighborhood of D_i , which is a tubular neighborhood of D_i . Then $\{N_{\delta}(D_i)\}_{i \in I}$ is an open cover of $N_{\delta}(D)$, and

$$\bigcap_{1 \le j \le r} N_{\delta}(D_{i_j}) \neq \emptyset \Leftrightarrow \bigcap_{1 \le j \le r} D_{i_j} \neq \emptyset.$$
(3.1.3)

Now, take any partition of unity $\{\rho_i\}_{i \in I}$ associated to this cover. That is, $\rho_i \in C^{\infty}(N_{\delta}(D))$ s.t.:

(1) $0 \le \rho_i \le 1$ and $\sum_{i \in I} \rho_i = 1$.

(2) For each *i*, the support Supp ρ_i is a compact subset contained in $N_{\delta}(D_i)$.

$$\overline{\alpha} = \overline{\alpha}[\{\rho_i\}] : N_{\delta}(D) \to \mathbb{D}D, \quad \overline{\alpha}(x) := \sum_{i=1}^{m} \rho_i(x) v_i,$$

where v_i is the **vertex** of $\mathbb{D}D$ corresponding to D_i . By (3.1.3), the map is indeed well-defined. **Claim**: the map $\overline{\alpha}$, up to homotopy, is independent of the choice of partition of unity.

Proof of Claim. Indeed, for any other partition of unity $\{\widetilde{\rho}_i\}$, we get a continuous family of partitions of unity $\{\rho_i^t := (1-t)\rho_i + t\widetilde{\rho}_i\}_{i \in I}$, parameterized by $t \in [0, 1]$. Apply the above definition, we then obtain a continuous family of maps $\overline{\alpha}_t = \sum \rho_i^t v_i : N_{\delta}(D) \to \mathbb{D}D$.

Remark 3.9. By the Claim and Proposition 3.3, the map obtained by composition

$$\alpha: N^*_{\delta}(D) = N_{\delta}(D) \setminus D \hookrightarrow N_{\delta}(D) \xrightarrow{\alpha} \mathbb{D}\partial X, \tag{3.1.4}$$

is well-defined up to homotopy.

3.2. Dual boundary complexes of very generic character varieties. Here's our main result:

Theorem 3.10. The weak geometric P=W conjecture 0.2 for very generic character varieties holds: If $(C_1, ..., C_k) \in T^k$ is very generic (Definition 1.1, Assumption 1.4) of type μ , and \mathcal{M}_{μ} is nonempty, then we have a homotopy equivalence

$$\mathbb{D}\partial \mathcal{M}_{\mu} \sim S^{d_{\mu}-1}, \quad d_{\mu} = \dim \mathcal{M}_{\mu}.$$

It relies on the following lemma:

28

Lemma 3.11 ([59, Cor.0.3]). If Y is a \mathbb{K} -variety stably isomorphic to \mathbb{A}^{ℓ} , $\ell \geq 1$, then $\mathbb{D}\partial Y$ is contractible.

Proof of Theorem 3.10. By Theorem 2.10, we get a decomposition into locally closed subvarieties

$$\begin{aligned} \mathcal{M}_{\mu} &= \sqcup_{\vec{w} \in W^{2g} \times \prod_{i=1}^{k-1} W/W(C_i)} \mathcal{M}_{\mu}(\vec{w}) = \sqcup_{\vec{w} \in W^{2g} \times \prod_{i=1}^{k-1} W/W(C_i)} \sqcup_{p \in W^*(\beta(\vec{w}))} \mathcal{M}_{\mu}(\vec{w}, p), \\ \mathcal{M}_{\mu}(\vec{w}, p) &\cong (\mathbb{K}^{\times})^{\overline{a}(\vec{w}, p)} \times \mathcal{A}_{\mu}(\vec{w}, p), \quad \mathcal{A}_{\mu}(\vec{w}, p) \times \mathbb{K}^{|U|} \cong \mathbb{K}^{b(\vec{w}, p)}, \end{aligned}$$

and dim $\mathcal{M}_{\mu}(\vec{w}, p) = d_{\mu}$ if and only if $(\vec{w}, p) = (\vec{w}_{\max}, p_{\max})$, that is, $\overline{a}(\vec{w}, p) = d_{\mu}$, i.e., $\overline{b}(\vec{w}, p) = \dim \mathcal{H}_{\mu}(\vec{w}, p) = 0$. Thus, for any $(\vec{w}, p) \neq (\vec{w}_{\max}, p_{\max})$, by (3.1.1) and Lemma 3.11, we have

$$\mathbb{D}\partial \mathcal{M}_{\mu}(\vec{w},p) \sim \mathbb{D}\partial(\mathbb{K}^{\times})^{\overline{a}(\vec{w},p)} \star \mathbb{D}\partial \mathcal{A}_{\mu}(\vec{w},p) \sim \mathbb{D}\partial(\mathbb{K}^{\times})^{\overline{a}(\vec{w},p)} \star \mathrm{pt} \sim \mathrm{pt}.$$

To apply Lemma 3.8, it remains to produce an *admissible total order* (Definition 3.12) on

$$\mathcal{W}^* := \{ (\vec{w}, p) : \vec{w} \in W^{2g} \times \prod_{i=1}^{k-1} W/W(C_i), p \in \mathcal{W}^*(\beta(\vec{w})) \}.$$
(3.2.1)

This is done in Corollary 3.15 below. Thus, by Lemma 3.8, with $X = \mathcal{M}_{\mu}$ and $U = \mathcal{M}_{\mu}(\vec{w}_m, p_m) = (\mathbb{K}^{\times})^{d_{\mu}}$, we get a homotopy equivalence $\mathbb{D}\partial \mathcal{M}_{\mu} \xrightarrow{\sim} \mathbb{D}\partial \mathcal{M}_{\mu}(\vec{w}_m, p_m) \sim S^{d_{\mu}-1}$, as desired.

To finish the proof of our main theorem, we're left with the question of admissible total orders.

Definition 3.12. Let $X = \bigsqcup_{i \in I} Z_i$ be a finite decomposition of a \mathbb{K} -variety into locally closed subvarieties. We say that a total order \leq on *I* is *admissible*, if

$$Z_{\leq a} := \bigcup_{i \in I: i \leq a} Z_i \subset X$$

is closed, all $a \in I$. In particular, for the maximal index $m \in I$, $U := Z_m \subset X$ is open. In this case, we call $(X = \bigsqcup_{i \in I} Z_i, \leq)$ an *admissible decomposition*.

For simplicity, we also denote

$$I_{\leq a} := \{i \in I : i \leq a\}, \quad I_{< a} := \{i \in I : i < a\}, \quad Z_{< a} := \bigcup_{i \in I_{< a}} Z_i.$$

Observe that $I_{<i} = I_{\le m_{<i}}$, where $m_{<i}$ is the maximal element of $I_{<i}$. It follow by definition that, $Z_{<i} \subset Z_{\le i} \subset X$ are two closed subsets. Hence, the complement $Z_i = Z_{\le i} \setminus Z_{<i} \subset Z_{\le i}$ is open.

Lemma 3.13. Let $(X = \bigsqcup_{i \in I} Z_i, \leq)$ be an admissible decomposition. Suppose that, for each $i \in I$, we have an admissible decomposition $(Z_i = \bigsqcup_{j \in J_i} Z_{i,j}, \leq)$. Denote $\tilde{I} := \{(i, j) : i \in I, j \in I_i\} \cong \bigsqcup_{i \in I} J_i$, and define a total order on \tilde{I} by

 $(i, j) \le (i', j') \Leftrightarrow i < i', \text{ or } i = i' \text{ and } j \le j'.$

Then $(X = \sqcup_{(i,i) \in \widetilde{I}} Z_{i,j}, \leq)$ is an admissible decomposition.

Proof. Clearly, we have a decomposition of X into locally closed subvarieties $X = \bigsqcup_{(i,j) \in I} Z_{i,j}$. It suffices to show that the total order \leq on \widetilde{I} is admissible. Indeed, we have

$$Z_{\leq (i,j)} = Z_{< i} \cup (Z_i)_{\leq j} \hookrightarrow Z_{< i} \cup Z_i = Z_{\leq i} \hookrightarrow X.$$

By assumption, $Z_{\leq i} \subset X$ and $(Z_i)_{\leq j} \subset Z_i$ are closed. By the observation above, the composition

$$Z_{\leq i} \setminus Z_{\leq i,j} = Z_i \setminus (Z_i)_{\leq j} \subset Z_i \subset Z_{\leq i}$$

is then open. So, the complement $Z_{\leq i,j}$ is closed in $Z_{\leq i}$, hence also closed in X. Done.

Next, consider the Bruhat cell decomposition $G = \bigsqcup_{\dot{w} \in W/W(P)} B\dot{w}P$, where W(P) is the Weyl group of a Levi subgroup of *P*. Recall that there is a Bruhat partial order on W/W(p): $\dot{\lambda} \leq \dot{\mu}$ if and only if $B\dot{\lambda}P \subset \overline{B\dot{\mu}P}$. That is, $\overline{B\dot{\mu}P} = \bigsqcup_{\dot{\lambda}\leq \dot{\mu}} B\dot{\lambda}P$. It follows that any total order extending the Bruhat partial order is admissible. From now on, we alway fix such an extension.

Let $\beta \in Br_n^+$ be a *n*-strand positive braid with a braid presentation $\beta = \sigma_{i\ell} \circ \cdots \sigma_{i_1}$ as usual. Recall that the braid variety $X(\beta)$ has a *B*-equivariant decomposition (1.4.9):

$$X(\beta) = \sqcup_{p \in W(\beta)} X_p(\beta).$$

Lemma 3.14. There exists a natural admissible total order on $W(\beta)$.

Proof. For each $p = (p_{\ell} = id, \dots, p_1, p_0 = id) \in \mathcal{W}(\beta)$, define locally closed subvarieties of $X(\beta)$:

$$X_{(p_i,\dots,p_0)}(\beta) = \bigcap_{1 \le j \le i} \overline{f_j}^{-1}(Bp_j B); \quad \overline{f_j} : X(\beta) \to G : \vec{\epsilon} = (\epsilon_i)_{i=\ell}^1 \mapsto B_{i_j}(\epsilon_j) \cdots B_{i_1}(\epsilon_1).$$
(3.2.2)

So, $X_{(p_i,\dots,p_0)}(\beta) = \bigsqcup_{p_{i+1} \in W} X_{(p_{i+1},\dots,p_0)}(\beta)$. Then by the Bruhat cell decomposition, have

$$X_{(\leq p_{i+1},p_i,\cdots,p_0)}(\beta) = \bigcup_{w \leq p_{i+1}} X_{(p_i,\cdots,p_0)}(\beta) \cap \overline{f}_{i+1}^{-1}(\overline{BwB}),$$

hence is closed in $X_{(p_i, \dots, p_0)}(\beta)$. In other words, the Bruhat total order on

 $W_{(p_i,\cdots,p_0)} := \{ p_{i+1} \in W : X_{(p_{i+1},\cdots,p_0)}(\beta) \neq \emptyset \} \subset W$

is admissible. Then by induction, Lemma 3.13 induces an admissible total order \leq on $\mathcal{W}(\beta)$.

Finally, as promised in the proof of Theorem 3.10, we obtain

Corollary 3.15. *There is a natural admissible total order on the cell decomposition:*

$$\mathcal{M}_{\mu} = \sqcup_{(\vec{w},p) \in \mathcal{W}^*} \mathcal{M}_{\mu}(\vec{w},p),$$

such that (\vec{w}_m, p_m) is the maximal index.

Proof. This is more or less a consequence of Lemma 3.14, and the argument is similar.

Firstly, consider the decomposition

$$\mathcal{M}_{\mu} = \sqcup_{\vec{w} \in W^{2g} \times \prod_{i=1}^{k-1} W/W(C_i)} \mathcal{M}_{\mu}(\vec{w})$$

Let's show that it's *admissible*: there exists an admissible total order on $W^{2g} \times \prod_{i=1}^{k-1} W/W(C_i)$. Equivalently by definition (1.2.8) and (1.2.7), it means the equivariant decomposition

$$M'_B = \sqcup_{\vec{w} \in W^{2g} \times \prod_{i=1}^{k-1} W/W(C_i)} M'_B(\vec{w})$$

is admissible. Indeed, similar to the proof of Lemma 3.14, by Lemma 3.13, we obtain an admissible total order on $W^{2g} \times \prod_{i=1}^{k-1} W/W(C_i)$ as the compositions of the total Bruhat orders on $G = \bigsqcup_{w \in W} BwB$ and $G = \bigsqcup_{w_i \in W/W(C_i)} Bw_i P_i$.

Now, again by Lemma 3.13, it suffices to show that the decomposition

$$\mathcal{M}_{\mu}(\vec{w}) = \sqcup_{p \in \mathcal{W}^*(\beta(\vec{w}))} \mathcal{M}_{\mu}(\vec{w}, p)$$

is admissible. By definition (2.4.23), it suffices to show that the equivariant decomposition

$$M_B''(\vec{w}) = \sqcup_{p \in \mathcal{W}^*(\beta(\vec{w}))} M_B''(\vec{w}, p)$$

in Proposition 2.9 is admissible. By (2.3.7), it amounts to show that the equivariant decomposition

$$\widetilde{X}(\beta(\vec{w})) = \sqcup_{p \in W^*(\beta(\vec{w}))} \widetilde{X}_p(\beta(\vec{w}))$$

defined by (2.3.4), (2.4.1) is admissible. By definition, this follows from Lemma 3.14.

Remark 3.16. If \mathcal{M}_{μ} is only generic, a weaker result holds: $\mathbb{D}\partial \mathcal{M}_{\mu}$ is a rational homology $(d_{\mu} - 1)$ -sphere. This can be shown by a completely different argument using the curious hard Lefschetz property [40, Thm.1.5.3]. For example, see [41, Rmk.7.0.7].

T. SU

Lemma 3.17. [59, Lem.3.1], Let X be a smooth connected affine algebraic variety of dimension \geq 3, with any log compactification $(\overline{X}, D = \overline{X} - X)$. Then $\mathbb{D}\partial X$ is connected and we have a natural surjection

$$\pi_1(X) \twoheadrightarrow \pi_1(\overline{X}) \simeq \pi_1(D) \twoheadrightarrow \pi_1(\mathbb{D}\partial X).$$

Assuming (I), observe that (II) is more or less manageable. If dim $\mathcal{M}_{\mu} > 2$, by Remark 3.16, (I), and Lemma 3.17, (II) holds once we know that $\pi_1(\mathcal{M}_{\mu})$ is abelian, which is expected to be the case. If \mathcal{M}_{μ} is very generic, by Theorem 2.10, \mathcal{M}_{μ} contains an open dense algebraic torus $(\mathbb{K}^{\times})^{d_{\mu}}$, which induces a surjection $\pi_1((\mathbb{K}^{\times})^{d_{\mu}}) = \mathbb{Z}^{d_{\mu}} \twoheadrightarrow \pi_1(\mathcal{M}_{\mu})$. This confirms that $\pi_1(\mathcal{M}_{\mu})$ is abelian. If \mathcal{M}_{μ} is only generic, we expect that \mathcal{M}_{μ} still contains an open dense algebraic torus, then the same argument applies.

Alternatively, we consider a case when \mathcal{M}_{μ} is only generic. Let $\mathcal{N}(n, c_1)$ be the moduli space of stable rank *n* holomorphic bundles of degree c_1 on a Riemann surface Σ_g of genus *g*. In [15, Thm.3.1], the Yang-Mills functional was used as a Morse-Bott function to show that $\pi_1(\mathcal{N}(n, c_1))$ is abelian, if *n*, c_1 are coprime and $(g, n) \neq (2, 2)$. In this case, $T^*\mathcal{N}(n, c_1)$ is open dense in $\mathcal{M}_{\text{Dol}}(n, c_1)$, the Dolbeault moduli space of stable rank *n* Higgs bundles of degree c_1 on Σ_g . So, we obtain a surjection $\pi_1(\mathcal{N}(n, c_1)) \approx$ $\pi_1(T^*\mathcal{N}(n, c_1)) \twoheadrightarrow \pi_1(\mathcal{M}_{\text{Dol}}(n, c_1))$, and hence $\pi_1(\mathcal{M}_{\text{Dol}}(n, c_1))$ is also abelian. Under NAH, we get a diffemorphism $\mathcal{M}_{\text{Dol}}(n, c_1) \simeq \mathcal{M}_B(n, c_1)$, with $\mathcal{M}_B(n, c_1) = \mathcal{M}_{\mu}$, k = 1, $\mu = ((n))$, and $C_1 = \exp(-\frac{2\pi i c_1}{n})$. Then, $\pi_1(\mathcal{M}_B(n, c_1))$ and $\pi_1(\overline{\mathcal{M}}_B(n, c_1))$ are abelian. See also [19, Prop.6.30]. We expect that such an argument works more generally.

Now, we speculate on (*I*). By Proposition 3.6, (*I*) amounts to part of the weight cohomology with compact support $H^{a,0}_{W,c}(\mathcal{M}_{\mu};\mathbb{Z})$ being torsion-free. We refer to [59, §1] for a quick review on weight cohomology with compact support. Indeed, examples suggest that the following much more general statement should hold for all generic character varieties:

- (1) The weight cohomology with compact support $H^{a,b}_{W,c}(\mathcal{M}_{\mu};\mathbb{Z}), \forall a, b \geq 0$ is always torsion-free;
- (2) The integral cohomological descent spectral sequence [23, Thm.3] for \mathcal{M}_{μ}

$$E_2^{a,b} = H^{a,b}_{\mathrm{W},c}(X;\mathbb{Z}) \implies H^{a+b}_c(X(\mathbb{C});\mathbb{Z})$$

degenerates at E_2 . In particular, $H_c^j(\mathcal{M}_{\mu};\mathbb{Z}) = \bigoplus_{a+b=j} H_{W,c}^{a,b}(\mathcal{M}_{\mu};\mathbb{Z})$ is torsion-free.

Notice that such a degeneration fails dramatically for general varieties, it's then natural to ask that what is the reason behind the degeneration for character varieties. Due to the motivic nature of $H^{a,b}_{W,c}(X;A)$, we are led to expect that the motives with compact support of generic character varieties take some simple form, in a way compatible with the HLRV conjecture [27, Conj.1.2.1-1.2.2].

Finally, we give a remark on a motivic study of character varieties.

Remark 3.18. Recall that A. Mellit has established the curious hard Lefschetz (CHL) theorem [40] for all generic character varieties M_{μ} :

$$\omega^m \cup -: \operatorname{Gr}^W_{d_{\mu}-2m} H^i_c(\mathcal{M}_{\mu}; \mathbb{C}) \xrightarrow{\simeq} \operatorname{Gr}^W_{d_{\mu}+2m} H^{i+2m}_c(\mathcal{M}_{\mu}: \mathbb{C}),$$

where ω is the canonical holomorphic symplectic form obtained from quasi-Hamiltonian geometry. Adapted to our setting, his proof can be divided into two main steps: First, prove CHL for very generic character varieties using the cell decomposition and a gluing property for CHL; Second, reduce CHL of *generic* \mathcal{M}_{μ} to that of very generic character varieties by a **degeneration argument**:

(1) By varying the monodromy at a virtual puncture, \mathcal{M}_{μ} is embedded into a central fiber of a singular family of character varieties. By taking a resolution of the latter, \mathcal{M}_{μ} fits into a cartesian diagram (left) which is morally a base change of the cartesian diagram (right)

$$\begin{array}{cccc} X_{\mathrm{sm}}^{=1} & \stackrel{\iota_X}{\longrightarrow} & X_{\mathrm{sm}}^1 & \stackrel{\frown}{\longrightarrow} & X_{\mathrm{sm}} & & G/B \stackrel{\iota_i}{\to} & \widetilde{N} \hookrightarrow & \widetilde{G} \\ & \downarrow^{\pi_X} & \downarrow^{\pi_X} & \downarrow^{\pi_X} & & \downarrow^{\pi} & \downarrow^{\pi} \\ \mathcal{M}_{\mu} = X_{\mathrm{sing}}^{=1} & \stackrel{i_X}{\to} & X_{\mathrm{sing}}^1 & \stackrel{\frown}{\hookrightarrow} & X_{\mathrm{sing}} & & \{1\} \stackrel{\iota_i}{\to} & N \hookrightarrow & G \end{array}$$

30

where $\pi : \widetilde{G} \to G = GL_n(\mathbb{C})$ is the Grothendieck-Springer resolution and N is the subvariety of unipotent matrices. Moreover, X_{sm}^1 behaves like a very generic character variety, satisfying CHL with middle weight $d_{\mu} + 2 \dim G/B$;

(2) The BBDG decomposition theorem [3] for the Springer resolution $\pi : \tilde{N} \to N$ implies a natural isomorphism of mixed Hodge complexes of sheaves:

$$(R\pi_*\mathbb{C}_{\widetilde{N}})^- \cong i_*\mathbb{C}(-\dim G/B)[-2\dim G/B],$$

where $(-)^{-}$ stands for the sign component as a S_n -representation;

(3) By base change, there is a natural isomorphism of mixed Hodge complexes of sheaves:

$$(R\pi_{X*}X_{\rm sm}^1)^- \cong i_{X*}\mathbb{C}_{X_{\rm sing}^{-1}}(-\dim G/B)[-2\dim G/B].$$

Passing to cohomology with compact support, this induces an isomorphism of mixed Hodge structures:

$$H_c^*(X_{\operatorname{sm}}^1;\mathbb{C})^- \cong H_c^*(X_{\operatorname{sing}}^{-1};\mathbb{C})(-\dim G/B)[-2\dim G/B].$$

(4) The above isomorphism is compatible with the Lefschetz operator $\omega \cup -$. Then, by the CHL for X_{sm}^1 , $\mathcal{M}_{\mu} = X_{sing}^{=1}$ satisfies CHL with middle weight d_{μ} .

Given the discussion above, it seems natural to pose the following question:

Does the degenerating argument for proving CHL for character varieties admit a motivic improvement?

If so, such a motivic result could be used to detect the integral cohomology of the dual boundary complex of generic but not necessarily very generic character varieties. The latter is the main obstruction for us to prove the full weak geometric P=W conjecture 0.2. There're at least two main challenges in the motivic question: 1. Existence of a motivic lifting/variation of the decomposition theorem for the Springer resolution? For the rational version, see [16]; 2. The six functor formalism for integral motivic sheaves. For related work, see [7, 56].

APPENDIX A. CELL DECOMPOSITION OF BRAID VARIETIES

We'll prove Proposition 1.16. We use the notations of Definition 1.14, 1.15. Let $p \in W(\beta)$, $0 \le m \le \ell$. Define a closed subvariety of \mathbb{A}^m and subsets of $[\ell]$:

$$\begin{split} X_p^m(\beta) &:= \cap_{1 \le j \le m} f_j^{-1}(Bp_j B) \subset \mathbb{A}^m, \quad X_p^0(\beta) = \text{pt.} \\ U_p^m &:= U_p \cap [m], \ S_p^m := S_p \cap [m], \ D_p^m := D_p \cap [m]. \ \Rightarrow \ [m] = U_p^m \sqcup D_p^m \sqcup S_p^m. \end{split}$$
(A.0.1)

Lemma A.1. We have $p(X(\beta)) \subset W(\beta)$. Moreover, for any $p \in W(\beta)$ and $1 \le m \le \ell$,

$$X_{p}^{m}(\beta) \cong \begin{cases} \mathbb{K}_{\epsilon'_{m}} \times X_{p}^{m-1}(\beta) & \text{if } p_{m} = s_{i_{m}}p_{m-1} > p_{m-1} \quad (go-up); \\ X_{p}^{m-1}(\beta) & \text{if } p_{m} = s_{i_{m}}p_{m-1} < p_{m-1} \quad (go-down); \\ \mathbb{K}_{\epsilon'_{m}}^{\times} \times X_{p}^{m-1}(\beta) & \text{if } s_{i_{m}}p_{m-1} < p_{m-1} \text{ and } p_{m} = p_{m-1} \quad (stay). \end{cases}$$

Proof. For any $\vec{\epsilon} \in \mathbb{A}^{\ell}$, denote $p := p(\vec{\epsilon}) \in W^{\ell+1}$.

(1) If $s_{i_m} p_{m-1} > p_{m-1}$, that is, $\ell(s_{i_m} p_{m-1}) = \ell(p_{m-1}) + 1$. By Proposition 1.10, we have

 $[\mathbf{B}_{i_m}(\epsilon_m)\cdots\mathbf{B}_{i_1}(\epsilon_1)] = [\mathbf{B}_{i_m}(\epsilon_m)] \circ [\mathbf{B}_{i_{m-1}}(\epsilon_{m-1})\cdots\mathbf{B}_{i_1}(\epsilon_1)] \in \mathbf{BD}_n; \Rightarrow p_m = \mathbf{s}_{i_m}p_{m-1} \text{ (go-up)}.$

Let's define a parameter $\epsilon'_m \in \mathbb{K}$: By the unique decomposition

$$Bp_{m-1}B = Bp_{m-1}U^{-}_{p_{m-1}} : B_{i_{m-1}}(\epsilon_{m-1}) \cdots B_{i_{1}}(\epsilon_{1}) = b_{m-1}(\epsilon_{m-1}, \cdots, \epsilon_{1})p_{m-1}L^{-}_{p_{m-1}}(\epsilon_{m-1}, \cdots, \epsilon_{1}),$$

we get $b_{m-1} \in B$. Then $\epsilon'_m \in \mathbb{K}$ and $b_m \in B$ are uniquely determined by the equation

$$[\mathbf{B}_{i_m}(\boldsymbol{\epsilon}_m)] \circ [\boldsymbol{b}_{m-1}] = [\boldsymbol{b}_m] \circ [\mathbf{B}_{i_m}(\boldsymbol{\epsilon}'_m)] \in \underline{\mathrm{FBD}}_n. \tag{A.0.2}$$

Or, by Proposition 1.10: $[\mathbf{B}_{i_m}(\epsilon_m)\cdots\mathbf{B}_{i_1}(\epsilon_1)] = [b_m] \circ [\mathbf{B}_{i_m}(\epsilon'_m)] \circ [p_{m-1}] \circ [L^-_{p_{m-1}}] \in \mathbf{BD}_n$. In fact,

$$\epsilon'_{m} = (b_{m-1})_{i_{m},i_{m}}^{-1} (b_{m-1})_{i_{m}+1,i_{m}+1} \epsilon_{m} + (b_{m-1})_{i_{m},i_{m}}^{-1} (b_{m-1})_{i_{m},i_{m}+1}.$$
(A.0.3)

This shows that $X_p^m(\beta) \cong \mathbb{K}_{\epsilon'_m} \times X_p^{m-1}(\beta)$.

T. SU

(2) If
$$p'_{m-1} := s_{i_m} p_{m-1} < p_{m-1}$$
. So, $\ell(p_{m-1} = s_{i_m} p'_{m-1}) = \ell(p'_{m-1}) + 1$. By Proposition 1.10,
 $Bs_{i_m} p'_{m-1} B = U^-_{s^{-1}_{i_m}} s_{i_m} B p'_{m-1} U^-_{p'_{m-1}} : B_{i_{m-1}}(\epsilon_{i_{m-1}}) \cdots B_{i_1}(\epsilon_1) = H_{i_m}(c_{m-1}) s_{i_m} b'_{m-1} p'_{m-1} L^-_{p'_{m-1}}$

So, $[\mathbf{B}_{i_m}(\epsilon_m)\cdots\mathbf{B}_{i_1}(\epsilon_1)] = [\mathbf{s}_{i_m}\mathbf{H}_{i_m}(\epsilon_m + c_{m-1})\mathbf{s}_{i_m}b'_{m-1}p'_{m-1}L^-_{p'_{m-1}}] \in \mathbf{BD}_n$. Define $\epsilon'_m \in \mathbb{K}$ by

$$B = TU : b'_{m-1} = D'_{m-1}u'_{m-1}; \quad s_{i_m}H_{i_m}(\epsilon_m + c_{m-1})s_{i_m}D'_{m-1} = D'_{m-1}s_{i_m}H_{i_m}(\epsilon'_m)s_{i_m}.$$
(A.0.4)
More concretely, write $D'_{m-1} = \text{Diag}((D'_{m-1})_1, \dots, (D'_{m-1})_n) \in T$, then we have

$$\epsilon'_{m} = (D'_{m-1})_{i_{m}+1}^{-1} (\epsilon_{m} + c_{m-1}) (D'_{m-1})_{i_{m}}.$$
(A.0.5)

By the Bruhat cell decomposition for $GL(2, \mathbb{K})$, we have:

$$\mathbf{s}_{i_m} \mathbf{H}_{i_m}(\epsilon'_m) \mathbf{s}_{i_m} = \begin{cases} I_n & \epsilon'_m = 0; \\ a'_m \mathbf{s}_{i_m} d'_m \in B_{\mathbf{s}_{i_m}} U^-_{\mathbf{s}_{i_m}}, \ \epsilon'_m \neq 0. \end{cases} a'_m = \mathbf{K}_{i_m}(-\epsilon'_m) \mathbf{K}_{i_m+1}(\epsilon'_m) \mathbf{H}_{i_m}(-\epsilon'_m), \ d'_m = \mathbf{H}_{i_m}(\epsilon'_m), \quad (A.0.6) \end{cases}$$

according to the following computation:

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & \epsilon'_m \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} -(\epsilon'_m)^{-1} & 0 \\ 0 & \epsilon'_m \end{pmatrix} \begin{pmatrix} 1 & -\epsilon'_m \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & (\epsilon'_m)^{-1} \\ 0 & 1 \end{pmatrix}$$
.
a) If $\epsilon'_m = 0$, then

(2.a) If $\epsilon'_m = 0$,

$$[\mathbf{B}_{i_m}(\epsilon_m)\cdots\mathbf{B}_{i_1}(\epsilon_1)] = [b'_{m-1}p'_{m-1}L^-_{p'_{m-1}}] \Rightarrow p_m = p'_{m-1} = \mathbf{s}_{i_m}p_{m-1} \text{ (go-down)}$$

This shows that $X_p^m(\beta) \cong X_p^{m-1}(\beta)$. (2.b) If $\epsilon'_m \neq 0$, then by Proposition 1.10, we have:

$$\begin{bmatrix} B_{i_m}(\epsilon_m)\cdots B_{i_1}(\epsilon_1) \end{bmatrix} = \begin{bmatrix} D'_{m-1}a'_m s_{i_m}d'_m u'_{m-1}p'_{m-1}L^-_{p'_{m-1}} \end{bmatrix} = \begin{bmatrix} D'_{m-1}a'_m s_{i_m}d'_m \end{bmatrix} \circ \begin{bmatrix} u'_{m-1}p'_{m-1}L^-_{p'_{m-1}} \end{bmatrix} \in BD_n,$$

and $B_{i_m}(\epsilon_m)\cdots B_{i_1}(\epsilon_1) \in Bs_{i_m}p'_{m-1}B$. So, $p_m = p_{m-1}$ (stay), and $X_p^m(\beta) \cong \mathbb{K}_{\epsilon'_m}^{\times} \times X_p^{m-1}(\beta)$.
Now, if $\vec{\epsilon} \in X(\beta)$, then $p_0 = p_\ell = id$ by definition. So, $p \in \mathcal{W}(\beta)$. Done.

Now, we fulfill our promise:

Proof of Proposition 1.16. (0). The action of $b \in B$ on $\vec{\epsilon} \in X(\beta)$ is uniquely determined by

$$[\tilde{b}_m] \circ [B_{i_m}(\widehat{\epsilon}_m)] \circ \dots \circ [B_{i_1}(\widehat{\epsilon}_1)] = [B_{i_m}(\epsilon_m)] \circ \dots \circ [B_{i_1}(\epsilon_1)] \circ [b^{-1}] \in \underline{FBD}_n. \ (\forall 1 \le m \le \ell)$$

Apply g_{-} (Definition 1.6), we see $X_{p}(\beta)$ is *B*-invariant. The rest follows from Lemma A.1.

(1). By above, $\tilde{b}_m \in B$ and $\hat{\epsilon}_m \in \mathbb{K}$ are determined inductively by

$$[\tilde{b}_m] \circ [B_{i_m}(\hat{\epsilon}_m)] = [B_{i_m}(\epsilon_m)] \circ [\tilde{b}_{m-1}] \in \underline{FBD}_n.$$
(A.0.7)

By diagram calculus (Lemma 1.7), $\tilde{b}_m \in TU^+_{s_{i_m}}$, and by induction, $\tilde{b}_m \in U^+_{s_{i_m}}$ if $b = u \in U$. (1.a). If $b = u \in U$, so $\tilde{b}_m = \tilde{u}_m \in U^+_{s_{i_m}}$. Again by diagram calculus, have $U^+_{s_{i_m}} X = XU^+_{s_{i_m}}$, $\forall X \in T, X = s_{i_m}$, or $X = H_{i_m}(c)$. In particular, we can define $\tilde{u}'_m \in U$ by

$$\widetilde{u}_m^{-1}D'_{m-1}\mathbf{s}_{i_m}\mathbf{H}_{i_m}(\epsilon'_m)\mathbf{s}_{i_m} = D'_{m-1}\mathbf{s}_{i_m}\mathbf{H}_{i_m}(\epsilon'_m)\mathbf{s}_{i_m}\widetilde{u}_m^{\prime-1}.$$

If $m \in S_p$, so $p_m = p_{m-1} = s_{i_m} p'_{m-1}$ and $\epsilon'_m \neq 0$. By the decomposition (1.3.7), we compute

$$\begin{split} & [\mathbf{B}_{i_m}(\widehat{\epsilon}_m) \cdots \mathbf{B}_{i_1}(\widehat{\epsilon}_1)] = [\widetilde{u}_m^{-1} \mathbf{B}_{i_m}(\epsilon_m) \cdots \mathbf{B}_{i_1}(\epsilon_1) u^{-1}] \\ & = [\widetilde{u}_m^{-1} D'_{m-1} \mathbf{s}_{i_m} \mathbf{H}_{i_m}(\epsilon'_m) \mathbf{s}_{i_m} u'_{m-1} p'_{m-1} L^{-}_{p'_{m-1}} u^{-1}] \quad (\text{By } (2.b) \text{ in the proof of Lemma A.1}) \\ & = [D'_{m-1} \mathbf{s}_{i_m} \mathbf{H}_{i_m}(\epsilon'_m) \mathbf{s}_{i_m} \widetilde{u}'_m^{-1} u'_{m-1} (p'_{m-1} L^{+}_{p'_{m-1}} (L^{-}_{p'_{m-1}} u^{-1}) p'_{m-1} L^{-}_{p'_{m-1}} (L^{-}_{p'_{m-1}} u^{-1})] \\ & = [\widehat{D}'_{m-1} \mathbf{s}_{i_m} \mathbf{H}_{i_m}(\widehat{\epsilon}'_m) \mathbf{s}_{i_m} \widetilde{u}'_{m-1} p'_{m-1} \widehat{L}^{-}_{p'_{m-1}}] \in \text{BD}_n, \end{split}$$

where the last equality gives: $\widehat{L}_{p'_{m-1}} \in U_{p'_{m-1}}^-$, $\widehat{u}'_{m-1} \in U$, $\widehat{D}'_{m-1} = D'_{m-1}$, and $\widehat{\epsilon}'_m = \epsilon'_m$, as desired. (1.b). By diagram calculus (Lemma 1.7), for any $\lambda = \text{Diag}(\lambda_1, \dots, \lambda_n) \in T$, we have

$$[\mathbf{B}_{j}(\epsilon)] \circ [\lambda^{-1}] = [\mathbf{s}_{j}\mathbf{H}_{j}(\epsilon)] \circ [\lambda^{-1}] = [(\lambda^{\mathbf{s}_{j}})^{-1}] \circ [\mathbf{s}_{j}\mathbf{H}_{j}(\lambda_{j}\lambda_{j+1}^{-1}\epsilon)] = [(\lambda^{\mathbf{s}_{j}})^{-1}] \circ [\mathbf{B}_{j}(\lambda_{j}\lambda_{j+1}^{-1}\epsilon)].$$

Then by (A.0.7) and induction, we obtain

$$\widehat{\epsilon}_m = (t^{s_{i_{m-1}}\cdots s_{i_1}})_{i_m}(t^{s_{i_{m-1}}\cdots s_{i_1}})_{i_m+1}^{-1}\epsilon_m = t_{(s_{i_{m-1}}\cdots s_{i_1})^{-1}(i_m)}t_{(s_{i_{m-1}}\cdots s_{i_1})^{-1}(i_m+1)}^{-1}\epsilon_m, \quad \widetilde{b}_m = (t^{s_{i_m}\cdots s_{i_1}})^{-1}.$$

(1.b.1). If $m \in U_p$, then $p_m = s_{i_m} p_{m-1}$. By (1) in the proof of Lemma A.1, $\widehat{b}_{m-1} p_{m-1} \widehat{L}_{p_{m-1}}^- = \mathbf{B}_{i_{m-1}}(\widehat{\epsilon}_{m-1}) \cdots \mathbf{B}_{i_1}(\widehat{\epsilon}_1) = \widetilde{b}_{m-1}^{-1} \mathbf{B}_{i_{m-1}}(\epsilon_{m-1}) \cdots \mathbf{B}_{i_1}(\epsilon_1) t^{-1} = \widetilde{b}_{m-1}^{-1} b_{m-1} p_{m-1} L_{p_{m-1}}^- t^{-1}$. Then a simple computation gives $\widehat{b}_{m-1} = t^{s_{i_{m-1}} \cdots s_{i_1}} b_{m-1} (t^{p_{m-1}})^{-1}$, $\widehat{L}_{p_{m-1}}^- = tL_{p_{m-1}}^- t^{-1}$. So,

$$[\mathbf{B}_{i_m}(\widehat{\epsilon}_m)] \circ [\widehat{b}_{m-1}] = [t^{\mathbf{s}_{i_m}\cdots\mathbf{s}_{i_1}}\mathbf{B}_{i_m}(\epsilon_m)(t^{\mathbf{s}_{i_{m-1}}\cdots\mathbf{s}_{i_1}})^{-1}] \circ [t^{\mathbf{s}_{i_{m-1}}\cdots\mathbf{s}_{i_1}}b_{m-1}(t^{p_{m-1}})^{-1}]$$

= $[t^{\mathbf{s}_{i_m}\cdots\mathbf{s}_{i_1}}] \circ [b_m] \circ [\mathbf{B}_{i_m}(\epsilon'_m)] \circ [(t^{p_{m-1}})^{-1}] = [\widehat{b}_m] \circ [\mathbf{B}_{i_m}(\widehat{\epsilon}'_m)] \in \underline{\mathrm{FBD}}_n.$

This implies that

$$\hat{\epsilon}'_{m} = (t^{p_{m-1}})_{i_{m}}(t^{p_{m-1}})_{i_{m}+1}^{-1}\epsilon'_{m}, \quad \hat{b}_{m} = t^{s_{i_{m}}\cdots s_{i_{1}}}b_{m}(t^{p_{m}})^{-1}$$

(1.b.2). If $m \in S_p$, so $p_m = p_{m-1} = s_{i_m} p'_{m-1}$ and $\epsilon'_m \neq 0$. By (2.*b*) in the proof of Lemma A.1,

$$\begin{bmatrix} \mathbf{B}_{i_m}(\widehat{\epsilon}_m) \cdots \mathbf{B}_{i_1}(\widehat{\epsilon}_1) \end{bmatrix} = \begin{bmatrix} \widetilde{b}_m^{-1} \mathbf{B}_{i_m}(\epsilon_m) \cdots \mathbf{B}_{i_1}(\epsilon_1) t^{-1} \end{bmatrix}$$

=
$$\begin{bmatrix} \widetilde{b}_m^{-1} D'_{m-1} \mathbf{s}_{i_m} \mathbf{H}_{i_m}(\epsilon'_m) \mathbf{s}_{i_m} u'_{m-1} p'_{m-1} L_{p'_{m-1}}^{-1} t^{-1} \end{bmatrix} = \begin{bmatrix} \widehat{D}'_{m-1} \mathbf{s}_{i_m} \mathbf{H}_{i_m}(\widehat{\epsilon}'_m) \mathbf{s}_{i_m} \widehat{u}'_{m-1} p'_{m-1} \widehat{L}_{p'_{m-1}}^{-1} \end{bmatrix} \in \mathbf{BD}_n$$

Then a simple computation gives

$$\widehat{D}'_{m-1} = t^{s_{i_m}\cdots s_{i_1}} D'_{m-1} (t^{-1})^{p'_{m-1}}, \quad \widehat{\epsilon}'_m = (t^{p_{m-1}})_{i_m} (t^{p_{m-1}})^{-1}_{i_m+1} \epsilon'_m$$

$$\widehat{u}'_{m-1} = t^{p'_{m-1}} u'_{m-1} (t^{p'_{m-1}})^{-1}, \quad \widehat{L}^-_{p'_{m-1}} = tL^-_{p'_{m-1}} t^{-1}.$$

Altogether, we have proved (1).

(2). By the decomposition $Bp_m B = Bp_m U_{p_m}^-$: $B_{i_m}(\epsilon_m) \cdots B_{i_1}(\epsilon_1) = b_m p_m L_{p_m}^-$, we define

$$\mu \mathfrak{mon}_m : X_p^m(\beta) \to T : (\epsilon_m, \dots, \epsilon_1) \mapsto D(b_m) \in T.$$
(A.0.8)

In particular, $\mu mon = \mu mon_{\ell}$, and $\mu mon_0 = I_n$. We will prove by induction. Case 1. If $\ell(s_{i_m}p_{m-1}) = \ell(p_{m-1}) + 1$, i.e. $m \in U_p$ and $p_m = s_{i_m}p_{m-1}$. By (2) in Lemma A.1,

 $\mathbf{B}_{i_{m-1}}(\epsilon_{m-1})\cdots\mathbf{B}_{i_1}(\epsilon_1) = b_{m-1}p_{m-1}L_{p_{m-1}}^-; \quad [\mathbf{B}_{i_m}(\epsilon_m)] \circ [b_{m-1}] = [b_m] \circ [\mathbf{B}_{i_m}(\epsilon_m')] \in \underline{\mathrm{FBD}}_n.$

If follows that

$$\mu\mathfrak{mon}_m(\epsilon_m,\cdots,\epsilon_1) = D(b_m) = D(b_{m-1})^{s_{im}} = (\mu\mathfrak{mon}_{m-1}(\epsilon_{m-1},\cdots,\epsilon_1))^{s_{im}} \in T.$$
(A.0.9)

Case 2. If $p_{m-1} = s_{i_m} p'_{m-1}$ with $\ell(p_{m-1}) = \ell(p'_{m-1}) + 1$. We use (2) in Lemma A.1. So,

$$\mathbf{B}_{i_{m-1}}(\epsilon_{m-1})\cdots\mathbf{B}_{i_1}(\epsilon_1) = \mathbf{H}_{i_m}(c_{m-1})\mathbf{s}_{i_m}b'_{m-1}p'_{m-1}L^-_{p'_{m-1}} = b_{m-1}p_{m-1}L^-_{p_{m-1}}.$$

Observe that $D(b_{m-1}) = (D(b'_{m-1}))^{s_{i_m}} = (D'_{m-1})^{s_{i_m}} \in T$. Again, by (2) in Lemma A.1,

$$\mathbf{B}_{i_m}(\epsilon_m)\cdots\mathbf{B}_{i_1}(\epsilon_1) = D'_{m-1}\mathbf{s}_{i_m}\mathbf{H}_{i_m}(\epsilon'_m)\mathbf{s}_{i_m}u'_{m-1}p'_{m-1}L^-_{p'_{m-1}}.$$

Case 2.1. If $m \in D_p$, then $p_m = p'_{m-1}$ and $\epsilon'_m = 0$. Thus,

$$B_{i_m}(\epsilon_m)\cdots B_{i_1}(\epsilon_1) = b'_{m-1}p'_{m-1}L^-_{p'_{m-1}} = b_m p_m L^-_{p_m},$$

with $b_m = b'_{m-1}$ and $L^-_{p_m} = L^-_{p'_{m-1}}$. By above, it follows that

$$\mu_m(\epsilon_m, \dots, \epsilon_1) = D(b_m) = (D(b_{m-1}))^{s_{i_m}} = (\mu \mathfrak{mon}_{m-1}(\epsilon_{m-1}, \dots, \epsilon_1))^{s_{i_m}} \in T.$$
(A.0.10)

Case 2.2. If $m \in S_p$, then $p_m = p_{m-1}$ and $\epsilon'_m \neq 0$. By (2.*b*) in Lemma A.1,

$$\mathbf{B}_{i_m}(\epsilon_m)\cdots\mathbf{B}_{i_1}(\epsilon_1) = D'_{m-1}a'_m \mathbf{s}_{i_m} d'_m u'_{m-1} p'_{m-1} L^-_{p'_{m-1}} = b_m p_m L^-_{p_m},$$

where a'_{m}, d'_{m} are given by (A.0.6). So $D(b_{m}) = D(a'_{m})D'_{m-1} = D(a'_{m})D(b_{m-1})^{s_{i_{m}}}$. That is,

$$\mu_m(\epsilon_m, \dots, \epsilon_1) = \mathbf{K}_{i_m}(-\epsilon_m'^{-1})\mathbf{K}_{i_m+1}(\epsilon_m')\mu_{m-1}(\epsilon_{m-1}, \dots, \epsilon_1)^{\mathbf{s}_{i_m}} \in T.$$
(A.0.11)

Now, (1.4.11) follows by induction from (A.0.9), (A.0.10), and (A.0.11). This proves (2).

Appendix B. Quotients of varieties

We collect some results on quotients of varieties. Hopefully, it helps to digest the main body of this article. Most statements below are standard, so we skip the proofs whenever possible.

Recall our **Convention** 1: A \mathbb{K} -variety means a reduced separated scheme of finite type over \mathbb{K} .

Convention 7: Fix G, H as linear algebraic groups over \mathbb{K} , unless otherwise stated.

We refer to [30] for the background on various quotients for algebraic group actions. Occasionally, the notation $X/g^{e}G$ is used for geometric quotient.

B.1. Principal bundles. A principal G-bundle means so in the étale topology unless stated otherwise.

Proposition B.1. If $G \curvearrowright X$ freely, and $\pi : X \to Y$ is a flat orbit map into a K-variety, then:

(1) The fiber product $X \times_Y X$ is *G*-isomorphic to $X \times G$, i.e. we get a cartesian diagram:

$$\begin{array}{cccc} X \times G & \xrightarrow{(x,g)\mapsto(x,xg)} & X \times_Y X & \xrightarrow{p_2} & X \\ \downarrow p_1 & & \downarrow p_1 & & \downarrow \pi \\ X & \xrightarrow{\qquad} & X & \xrightarrow{\qquad} & \pi & \to Y \end{array} \tag{B.1.1}$$

(2) $\pi: X \to Y$ is smooth and affine.

(3) $\pi: X \to Y$ is a principal G-bundle (in the étale topology).

Remark B.2. In above, $\pi : X \to Y$ is in fact a geometric quotient. See Proposition B.9.

Proof of Proposition B.1. (1). The proof is covered by the three commutative diagrams:

First, consider the left cartesian diagram. As *Y* is **separated**, Δ_Y is a closed embedding, so is $\widetilde{\Delta}_Y : X \times_Y X \to X \times X$ by base change. A free action means the morphism $a : X \times G \to X \times X : (x, g) \mapsto (x, xg)$ is a closed embedding. π is *G*-invariant means $\pi \circ p_1 = \pi \circ p_1 \circ a = \pi \circ p_2 \circ a : X \times G \to Y$.

This induces the middle commutative diagram. As *a* and $\widetilde{\Delta}_Y$ are closed embeddings, so is $c: X \times G \to X \times_Y X: (x, g) \mapsto (x, xg)$, by the **cancellation property** for closed embeddings.

Now, consider the right commutative diagram. By assumption, each closed fiber of π is isomorphic to *G*. Then so is $p_1 : X \times_Y X \to X$ by base change. Thus, *c* is an isomorphism on the closed fibers over *X*, hence a dominant morphism.

By base change and composition, $X \times_Y X$ is of finite type over \mathbb{K} , and hence $p_1 : X \times_Y X \to X$ is a surjective **flat** morphism of schemes **of finite type** over \mathbb{K} , with *reduced* closed fibers and *reduced* base X. Then, $X \times_Y X$ is reduced by Lemma B.3 below. Now, *c* is a a closed embedding and a dominant morphism between reduced schemes, hence an isomorphism. This proves (1).

(2). Clearly, $\pi : X \to Y$ is **fpqc**. By [57, Lemma 02VL] (resp. [57, Lemma 02L5]), a morphism being smooth (resp. affine) is fpqc local on the target. Then by the cartesian diagram in (1), π is smooth and affine, as $p_1 : X \times G \to X$ is.

(3). By (2), $\pi : X \to Y$ is a smooth covering. By [57, Lemma 055V] (slicing smooth morphisms and refining a smooth covering by an étale covering), there exists a morphism $s : \mathcal{V} \to X$ such that the composition $\pi \circ s : \mathcal{V} \to X \to Y$ is an étale covering. Now, by (1), the base change of $\pi : X \to Y$ along $\pi \circ s : \mathcal{V} \xrightarrow{s} X \xrightarrow{\pi} Y$ is *G*-isomorphic to $\mathcal{V} \times G$ over \mathcal{V} . This gives a local trivialization of $\pi : X \to Y$ in the étale topology. This finishes the proof.

Lemma B.3 (Reducedness). If $f : X \to Y$ be a surjective morphism of schemes of finite type over any field *k*, and all closed fibers (i.e. X_y , \forall closed point $y \in Y$) of *f* are **reduced**, then

(1) Any fiber X_y of f is reduced (the point $y \in Y$ may not be closed).

(2) If in addition Y is reduced and f is flat, then X is also reduced.

Remark B.4 (Jacobson schemes). Let *Y* be a scheme of finite type over any field *k*.

(1) Recall by [57, Definition 02J1] that, a point $y \in Y$ is a **finite type point** if the canonical morphism Spec $k(y) \to Y$ is of finite type. Equivalently by [57, Lemma 01TA], y is a closed point in some affine open subset U = Spec R of Y, and the field extension $k \hookrightarrow k(y)$ is finite.

(2) By [57, Lemma 02J6], *Y* is **Jacobson**: the closed points are dense in every closed subset [57, Definition 01P2]. Equivalently, every nonempty locally closed subset contains a closed point.

(3) Now by [57, Lemma 01TB], the closed points in *Y* are precisely the finite type points.

In particular, if k is an algebraically closed field, then the closed points of Y are the k-points, and every nonempty locally closed subset contains a closed (i.e. k-) point.

Proof of Lemma B.3. (1). Take any point $y \in Y$, define $Y' := \overline{\{y\}} \hookrightarrow Y$ equipped with the reduced close subscheme structure. Then Y is integral and y is a generic point of Y'. Let $f' : X' = X \times_Y Y' \to Y'$ be the base change of f along $Y' \hookrightarrow Y$, which is still a surjective morphism of schemes of finite type over k. By base change and our assumption, all the closed fibers of f' are also reduced. Moreover, $X'_y = (f')^{-1}(y) = X_y$.

If X'_{y} is non-reduced, then by [57, Lemma 0575], there exists a nonempty open subset $V \subset Y'$ such that, for all $v \in V$ the fiber X'_{v} is non-reduced. However, by Remark B.4, Y is Jacobson³, hence V contains a closed point. This is a contradiction.

(2). This follows from (1) and [22, Cor.3.3.5] (alternatively, see [38, Thm.23.9, Cor.]):

Let $f : X \to Y$ be a flat morphism between two locally Noetherian schemes. If Y is reduced at the points of f(X), and $f^{-1}(y)$ is a reduced k(y)-scheme, \forall point $y \in f(X)$, then X is reduced. \Box

Remark B.5. As in the proof of Proposition B.1 (3), a similar argument also shows: If $\pi : X \to Y$ is a morphism of \mathbb{K} -varieties and *F* is a \mathbb{K} -variety, then π is a fiber bundle with fiber *F* in the étale topology if and only if π is a fiber bundle with fiber *F* in the smooth topology.

The flatness condition in Proposition **B**.1 can be relaxed when the base is normal:

Proposition B.6. If G acts freely on a pure dimensional variety X over \mathbb{K} (of char. 0), and $\pi : X \to Y$ is an orbit map, with Y a normal pure dimensional \mathbb{K} -variety, then:

(1) π is flat. So Proposition B.1 applies: π is smooth, affine, and a principal G-bundle, etc. (2) X is normal.

Proof. (1). By Lemma B.7, π is flat, so Proposition B.1 applies.

(2). This follows from (1) and [57, Lemma 034F]: a morphism being normal is local in the smooth topology: If $S' \to S$ is a smooth morphism, then S is normal \Rightarrow so is S'. If $S' \to S$ is smooth surjective, then S' is normal \Rightarrow so is S.

Lemma B.7 (A variant of miracle flatness. [50, Thm.3.3.27]). If $R \to S$ be a local morphism of Noetherian local rings, R is an excellent normal local domain with perfect residue field, and the closed fiber is regular of dimension dim S – dim R, then $R \to S$ is faithfully flat.

As a moral converse to Proposition **B**.1, we have

Lemma B.8. If $\pi : X \to Y$ is a principal *G*-bundle in **fpqc topology**, with X, Y \mathbb{K} -varieties, then:

(1) π is smooth and affine.

(2) We have a natural G-isomorphism $c: X \times G \xrightarrow{\simeq} X \times_Y X : (x,g) \mapsto (x,xg)$, i.e. the cartesian diagram

(B.1.1) holds. In addition, the G-action on X is free.

(3) $\pi: X \to Y$ is an orbit map and a principal *G*-bundle in the étale topology.

Proof. (1). The proof is similar to that of Proposition B.1 (2).

(2). As in the proof of Proposition B.1 (1), we get a G-morphism c in a commutative diagram:

$$\begin{array}{cccc} X \times G & \stackrel{c}{--} & X \times_Y X & \stackrel{p_2}{\longrightarrow} X \\ & \downarrow^{p_1} & \downarrow^{p_1} & \downarrow^{\pi} \\ X & \stackrel{\pi}{\longrightarrow} & X & \stackrel{\pi}{\longrightarrow} Y \end{array}$$

³It's this step that the finite type assumption in the lemma becomes essential.

and a closed embedding $X \times_Y X \hookrightarrow X \times X$ (Y is *separated*). It remains to show c is an isomorphism.

By our assumption and base change, $p_1 : X \times_Y X \to X$ is a principal *G*-bundle in the fpqc topology. So, there exists a fpqc covering $\tau : \mathcal{V} \to X$ and a *G*-isomorphism $\phi_\tau : \mathcal{V} \times_Y X \xrightarrow{\simeq} \mathcal{V} \times G$ over \mathcal{V} . In other words, we obtain the following commutative diagram:

$$\mathcal{V} \times G \xrightarrow{\widetilde{c}_{\tau}} \mathcal{V} \times G \xrightarrow{\tau \times \mathsf{id}_G} X \times G$$

$$\overset{\widetilde{c}_{\tau}}{\xrightarrow{\varphi_{\tau}}} \mathcal{V} \times_Y X \xrightarrow{\downarrow_{\mathcal{C}_{\tau}}} X \times_Y X$$

$$\overset{\simeq}{\xrightarrow{\varphi_{\tau}}} \mathcal{V} \times_Y X \xrightarrow{\downarrow_{\mathcal{C}_{\tau}}} X \times_Y X$$

$$\overset{\varphi_{\tau}}{\xrightarrow{\varphi_{\tau}}} \mathcal{V} \xrightarrow{\varphi_{\tau}} X$$

As a *G*-morphism over $\mathcal{V}, \tilde{c}_{\tau} := \phi_{\tau} \circ c_{\tau} : \mathcal{V} \times G \to \mathcal{V} \times G$ is then a *G*-isomorphism over \mathcal{V} . Thus, so is c_{τ} . Now, by [57, Lemma 02L4], *a morphism being an isomorphism is fpqc local on the target*. It follows that $c : X \times G \to X \times_Y X$ is a *G*-isomorphism over *X*. We're done.

(3). As $\pi : X \to Y$ is smooth surjective, by base change, the closed fibers of $\pi : X \to Y$ are the same as those of $p_1 : X \times_Y X \to X$, which are isomorphic to *G* by (2). Thus, $\pi : X \to Y$ is an orbit map. Now, the result follows from Proposition B.1 (3).

B.2. Associated fiber bundles. A useful reference is [35]. Recall Convention 7. As promised in Remark B.2, we complement Proposition B.1, Proposition B.6, and Lemma B.8.

Proposition B.9 (Associated fiber bundles). Let F be an affine H-variety over \mathbb{K} , and $\mathfrak{p} : P \to B$ be an (étale) principal H-bundle⁴. H acts on $P \times F$ diagonally: $h \cdot (a, z) := (ah^{-1}, hz)$. Then:

(1) The action $H \curvearrowright P \times F$ admits a geometric quotient $\pi : P \times F \to P \times^H F = (P \times F)/H$, with $P \times^H F$ is a \mathbb{K} -variety. In particular, $\mathfrak{p} : P \to B$ is a geometric quotient, with F = Spec k.

(2) The canonical map $\pi : P \times F \to P \times^H F$ is a principal H-bundle.

(3) The induced map $q: P \times^H F \to B: [a, z] \mapsto \mathfrak{p}(a)$ is an (étale) fiber bundle with fiber F.

(4) We have a fiber product diagram l

$$\begin{array}{ccc} P \times F & \xrightarrow{p_1} & P \\ \downarrow^{\pi} & \downarrow^{\mathfrak{p}} & & \downarrow^{\mathfrak{p}} \\ P \times^H F & \xrightarrow{q} & B \end{array} \tag{B.2.1}$$

Here, p_i *always stands for the projection to the i-th factor.*

Proof. This is similar to [35, Lem.3.4.1].

Example B.10 (Homogeneous spaces. [5, II.Thm.6.8]). Let $H \subset G$ be a closed subgroup. Then $q : G \to G/H = G/{^{ge}}H$ is a principal *H*-bundle over a smooth quasi-projective \mathbb{K} -variety.

B.3. Functorial properties and applications. Finally, we give some applications.

Corollary B.11 (base change). If $\mathfrak{p} : P \to B$ is a principal *H*-bundle, with $P' \subset P$ a locally closed *H*-subvariety, then $B' := \mathfrak{p}(P') \subset B$ is a locally closed subvariety, and we get a cartesian square

$$\begin{array}{ccc} P' & \longleftrightarrow & P \\ \mathfrak{p}' = \mathfrak{p}|_{P'} \downarrow & \dashv & \downarrow \mathfrak{p} \\ B' & \longleftrightarrow & B \end{array}$$

In particular, $\mathfrak{p}': P' \to B'$ is a principal H-bundle as well as a geometric quotient.

Proof. By the composition $P' \hookrightarrow \overline{P}' \hookrightarrow P$, it suffices to consider the case when $P' \subset P$ is an open or a closed *H*-subvariety. By Proposition B.9, $\mathfrak{p} : P \to B$ is a geometric quotient. Then as an *H*-invariant subset, $P' \subset P$ is open (resp. closed) if and only if $B' \subset B$ is open (resp. closed), and $P' = \mathfrak{p}^{-1}(B')$. It remains to show that the obviously commutative diagram is cartesian.

The open case is trivial. For the closed case, take $P'' := B' \times_B P$ and $\mathfrak{p}'' := \mathfrak{p}|_{P''} : P'' \to B'$ in the category of schemes. By base change, \mathfrak{p}'' is a principal *H*-bundle. In particular, \mathfrak{p}'' is flat, and its closed

⁴By Proposition B.1, it suffices to assume: \mathfrak{p} is a flat orbit map of \mathbb{K} -varieties for a free action $H \sim P$.

fibers are isomorphic to *H*, hence reduced. Of course, *B'* is reduced, then so is *P''* by Lemma B.3. On the closed subset $P' = \mathfrak{p}^{-1}(B')$, there is a unique reduced close subscheme structure. Thus, the natural morphism $P' \to P''$ is an isomorphism.

Proposition B.12 (Reduction of principal bundles). Let $H \subset G$ be a closed subgroup, X be a G-variety, and $i : Z \hookrightarrow X$ be a closed H-subvariety. So $H \curvearrowright G \times Z : h \cdot (g, z) = (gh^{-1}, h \cdot z)$. If:

- (1) $\pi: X \to B$ be a principal G-bundle over a \mathbb{K} -variety B.
- (2) $\tilde{a}: G \times Z \to X: (g, z) \mapsto gz$ is a principal H-bundle.

Then $\mathfrak{r} := \pi \circ i : Z \to B$ *is a principal H-bundle, and we have a cartesian diagram:*

$$\begin{array}{ccc} H \times Z & \longrightarrow & G \times Z & \xrightarrow{p_2} & Z \\ \downarrow^{a_Z} & & \downarrow^{\widetilde{a}} & \downarrow & \downarrow_{\mathfrak{f}} \\ Z & \stackrel{i}{\longleftrightarrow} & X & \xrightarrow{\pi} & B \end{array} (\Rightarrow Z/H = Z/^{\operatorname{ge}}H \xrightarrow{\widetilde{\to}} X/G = X/^{\operatorname{ge}}G \xrightarrow{\widetilde{\to}} B).$$
 (B.3.1)

Proof. Clearly, $\mathfrak{r}: \mathbb{Z} \to B$ is an *H*-invariant morphism. We have the following cartesian diagram:

$$\begin{array}{cccc} G \times Z & \xrightarrow{p_2} & Z \\ & & & \downarrow^{\operatorname{id}_G \times i} & & & \downarrow^i \\ G \times X & \xrightarrow{(g,x) \mapsto (gx,x)} & X \times_B X & \xrightarrow{p_2} & X \\ & & & \downarrow^a & & \downarrow^{p_1} & \downarrow^{\pi} \\ & X & \xrightarrow{\qquad} & X & \xrightarrow{\qquad} & B \end{array}$$

Here, p_i denotes the projection to the *i*-th factor. $a : G \times X \to X : (g, x) \mapsto gx$ is the action map. So, $\tilde{a} = a \circ (id_G \times i)$. The isomorphism $G \times X \xrightarrow{\approx} X \times_B X$ follows from Lemma B.8. Then by assumption and base change, $\mathfrak{r} : Z \to B$ is a principal *H*-bundle in the smooth topology, hence in the étale topology by Lemma B.8. It remains to show (B.3.1): the right square is cartesian by above; By Lemma B.8, so is the outer square. The rest is due to Proposition B.9.

Remark B.13. In above, by Propositions B.1, B.6, the conditions (1)-(2) can be relaxed to:

• The *G*-action on *X* is free.

• $\pi : X \to B$ and $\tilde{a} : G \times Z \to X$ are either *flat* orbit maps, or orbit maps between *pure dimensional* varieties with *B normal*.

Proposition B.14 (Quotient of a principal bundle by a subgroup). Let $H \subset G$ be a closed subgroup such that G/H is affine. Let $\mathfrak{p} : P \to B = P/G$ be a principal G-bundle, then:

- (1) There exists a geometric quotient $\mathfrak{p}_H : P \to P/H$, and \mathfrak{p}_H is a principal H-bundle.
- (2) The natural map $q_H : P/H \to B$ is a fiber bundle with fiber G/H in the étale topology.
- (3) If in addition, $H \triangleleft G$ is a normal subgroup, then $q_H : P/H \rightarrow B$ is a principal G/H-bundle.

Proof. Let F := G/H and $X := P \times F$. So, G acts diagonally on X: $g \cdot (p, zH) := (pg^{-1}, gzH)$. As F = G/H is **affine**, by Proposition B.9, we conclude that $G \curvearrowright X$ admits a geometric quotient $\pi : X = P \times F \rightarrow Y := P \times^G F$, which is a principal G-bundle, and the induced map $q_H = q : Y = P \times^G F \rightarrow B$ is a fiber bundle with fiber F in the étale topology. Moreover, we obtain the following cartesian diagram

$$\begin{array}{c} P \times G \xrightarrow{|\mathsf{Id}_P \times \pi} & X = P \times G/H \xrightarrow{p_1} P \\ \stackrel{\square}{a}(p,g) \mapsto pg & \downarrow \pi & \downarrow p \\ P = P \times^G G \xrightarrow{\mathfrak{p}_H} & Y = P \times^G G/H \xrightarrow{q_H} B \end{array}$$

(1). By the left cartesian square, $\mathfrak{p}_H : P \to Y = P \times^G G/H$ is a principal *H*-bundle. By Proposition B.9 (1), \mathfrak{p}_H is also a geometric quotient. So we can write $P/H = Y = P \times^G G/H$.

(2). By the right cartesian square above, $q_H : P/H = Y \rightarrow B$ is a fiber bundle with fiber G/H in the smooth topology, hence in the étale topology by Remark B.5.

(3). The action map $a : P \times G \to P$ induces an action $P/H \times G/H \to P/H : (pH, gH) \mapsto pHgH = pgH$. Then by the right cartesian square above $q_H : Y = P/H \to B$ is a principal G/H-bundle in the smooth topology, hence in the étale topology by Lemma B.8. **Proposition B.15.** Let G be a unipotent algebraic group and B be an affine variety over \mathbb{K} . Then any principal G-bundle $\mathfrak{p}: P \to B$ is trivial.

Note: G is connected: $\pi_0(G) = G/G^0$ is a finite unipotent algebraic group, which must be trivial.

Proof. First, assume G is commutative. Then, $G = G_a^m$ for some $m \ge 0$, as its (abelian) Lie algebra is a K-vector space with trivial Lie bracket. Now, the principal G-bundles over B are classified by $H_{\text{ét}}^1(B, G_a^m)$ (see [57, Lemma 03F7]). By [57, Proposition 03DW] and the affine vanishing property, $H_{\text{ét}}^1(B, G_a^m) \cong H^1(B, O_B^m) \cong 0$. Thus, p is a trivial principal G-bundle.

In general, we prove by induction on dim G. Say, $n := \dim G > 0$. Let H := Z(G) be the center of G. Then, H is a commutative unipotent algebraic group of positive dimension, and $\overline{G} := G/H$ is a unipotent algebraic group of dimension < n. By Proposition B.14, there exists a \mathbb{K} -variety $\overline{P} := P/H$ such that, $\mathfrak{p}_H : P \to \overline{P}$ is a principal H-bundle, and $q_H : \overline{P} \to B$ is a principal \overline{G} -bundle. By induction, q_H is trivial, hence admits a section $s : B \to \overline{P}$. We then obtain the following commutative diagram in which the square is cartesian:



By base change, $\tilde{p}_H : s^{-1}P \to B$ is a principal *H*-bundle. As *H* is a commutative unipotent algebraic group, \tilde{p}_H is trivial by above, hence admits a section $s' : B \to s^{-1}P$. Then, $\tilde{s} \circ s' : B \to P$ defines a section of $\mathfrak{p} : P \to B$. This implies that \mathfrak{p} is a trivial principal *G*-bundle. \Box

Acknowledgements. The author would like to thank his Ph.D advisor, Vivek Shende, for the interest and the suggestion to work along the direction of this article. He thanks Chenglong Yu, Penghui Li, Mingchen Xia, Yu Zhao, Mirko Mauri, Junliang Shen, Changjian Su, and Jia-Choon Lee for helpful discussions and comments. Furthermore, the author is very grateful to David Nadler, Lenhard Ng, Shing-Tung Yau, and Shaoyuan Zheng, for their valuable support during his job search. The initial version was done when the author was a postdoc at IMS, CUHK. During the revision, the author benefited a lot from a visit invited by Changjian Su at YMSC, and a visit invited by Song-Yan Xie at AMSS, CAS.

References

- S. S. Abhyankar, W. Heinzer and P. Eakin, On the uniqueness of the coefficient ring in a polynomial ring, J. Alg. 23 (1972), 310–342.
- [2] O. Biquard and P. Boalch, Wild non-abelian Hodge theory on curves, Compos. Math. 140 (2004), 179-204.
- [3] A. Beilinson, J. Bernstein, P. Deligne and O. Gabber, Faisceaux pervers, Société mathématique de France Paris, 2018.
- [4] P. P. Boalch, Geometry and braiding of Stokes data; fission and wild character varieties, Annals of math. (2014), 301-365.
- [5] A. Borel, Linear algebraic groups, Springer Science & Business Media, 2012.
- [6] P. Boalch and D. Yamakawa, Twisted wild character varieties, arXiv:1512.08091.
- [7] D.-C. Cisinski and F. Déglise, Triangulated categories of mixed motives, Springer, 2019.
- [8] M. de Cataldo, D. Maulik and J. Shen, *Hitchin fibrations, abelian surfaces, and the P= W conjecture*, J. Amer. Math. Soc. **35** (2022), 911–953.
- [9] K. Corlette, Flat G-bundles with canonical metrics, J. Diff. Geom. 28 (1988), 361-382.
- [10] V. I. Danilov, Polyhedra of schemes and algebraic varieties, Math. of the USSR-Sbornik 26 (1975), 137.
- [11] B. Davison, Nonabelian Hodge theory for stacks and a stacky P=W conjecture, Adv. Math. 415 (2023), 108889.
- [12] M. de Cataldo, T. Hausel and L. Migliorini, *Topology of Hitchin systems and Hodge theory of character varieties: the case A1*, Annals of Math. (2012), 1329–1407.
- [13] M. de Cataldo and D. Maulik, *The perverse filtration for the Hitchin fibration is locally constant*, Pure Appl. Math. Q. **16** (2020), 1444–1464.
- [14] T. de Fernex, J. Kollár and C. Xu, *The dual complex of singularities*, Higher dimensional algebraic geometry—in honour of Professor Yujiro Kawamata's sixtieth birthday 74 (2017), 103–129.
- [15] G. D. Daskalopoulos and K. K. Uhlenbeck, An application of transversality to the topology of the moduli space of stable bundles, Topology 34 (1995), 203–215.
- [16] J. N. Eberhardt and C. Stroppel, Motivic Springer theory, Indag. Math. 33 (2022), 190-217.
- [17] M. Farajzadeh-Tehrani and C. Frohman, On compactifications of the SL (2, C) character varieties of punctured surfaces, arXiv:2305.12306.

- [18] R. Fricke and F. Klein, Vorlesungen über die Theorie der automorphen Funktionen. Band 1: Die gruppentheoretischen Grundlagen. Band II: Die funktionentheoretischen Ausführungen und die Andwendungen, Bibliotheca Mathematica Teubneriana, Bände 3 (1965).
- [19] C. Felisetti and M. Mauri, P = W conjectures for character varieties with symplectic resolution, J. Ec. Polytech. Math. 9 (2022), 853–905.
- [20] T. Fujita, On Zariski problem, 1979.
- [21] W. M. Goldman and W. D. Neumann, *Homological action of the modular group on some cubic moduli spaces*, Math. Res. Lett. **12** (2005), 575–591.
- [22] A. Grothendieck, Éléments de géométrie algébrique: IV. Étude locale des schémas et des morphismes de schémas, Seconde partie, Publ. Math. IHES 24 (1965), 5–231.
- [23] H. Gillet and C. Soulé, Descent, motives and K-theory, Walter de Gruyter, Berlin/New York (1996).
- [24] N. Gupta, On the cancellation problem for the affine space in characteristic p, Invent. Math. 195 (2014), 279–288.
- [25] N. Gupta, On Zariski's cancellation problem in positive characteristic, Adv. Math. 264 (2014), 296–307.
- [26] P. Huang, G. Kydonakis, H. Sun and L. Zhao, Tame parahoric nonabelian Hodge correspondence on curves, arXiv:2205.15475.
- [27] T. Hausel, E. Letellier and F. Rodriguez-Villegas, Arithmetic harmonic analysis on character and quiver varieties, Duke Math. J. 160 (2011), 323–400.
- [28] T. Hausel, E. Letellier and F. Rodriguez-Villegas, Arithmetic harmonic analysis on character and quiver varieties II, Adv. Math. 234 (2013), 85–128.
- [29] T. Hausel, A. Mellit, A. Minets and O. Schiffmann, $P = W via H_2$, arXiv:2209.05429.
- [30] V. Hoskins, Moduli problems and geometric invariant theory, Lecture notes 2016.
- [31] P. Huang and H. Sun, Meromorphic parahoric Higgs torsors and filtered Stokes G-local systems on curves, arXiv:2212.04939.
- [32] L. Katzarkov, A. Noll, P. Pandit and C. Simpson, *Harmonic maps to buildings and singular perturbation theory*, Commun. Math. Phys. 336 (2015), 853–903.
- [33] A. Komyo, On compactifications of character varieties of n-punctured projective line, Ann. Inst. Fourier 65 (2015), 1493–1523.
- [34] H. Konno, Construction of the moduli space of stable parabolic Higgs bundles on a Riemann surface, J. Math. Soc. Japan 45 (1993), 253–276.
- [35] H. Kraft, Fiber bundles, slice theorem and applications, (online) Lecture notes, 2015.
- [36] M. Kontsevich and Y. Soibelman, Wall-crossing structures in Donaldson–Thomas invariants, integrable systems and mirror symmetry, Homological mirror symmetry and tropical geometry (2014), 197–308.
- [37] J. Kollár and C. Xu, The dual complex of Calabi-Yau pairs, Invent. Math. 205 (2016), 527-557.
- [38] H. Matsumura, Commutative ring theory, Cambridge university press, 1989.
- [39] M. Mauri, Intersection cohomology of rank 2 character varieties of surface groups, J. Inst. Math. Jussieu (2021), 1–40.
- [40] A. Mellit, Cell decompositions of character varieties, arXiv:1905.10685.
- [41] M. Mauri, E. Mazzon and M. Stevenson, On the geometric P=W conjecture, Selecta Math. (N.S.) 28 (2022).
- [42] T. Mochizuki, Wild harmonic bundles and wild pure twistor D-modules, Asterisque (2011).
- [43] T. Mochizuki, Good wild harmonic bundles and good filtered Higgs bundles, SIGMA 17 (2021), 068.
- [44] M. Miyanishi and T. Sugie, Affine surfaces containing cylinderlike open sets, Kyoto J. Math. 20 (1980), 11-42.
- [45] D. Maulik and J. Shen, The P = W conjecture for GL_n , arXiv:2209.02568.
- [46] D. Maulik, J. Shen and Q. Yin, Perverse filtrations and Fourier transforms, arXiv:2308.13160.
- [47] A. Némethi and S. Szabó, The geometric P=W conjecture in the Painlevé cases via plumbing calculus, IMRN 2022 (2022), 3201–3218.
- [48] S. Payne, Boundary complexes and weight filtrations, Mich. Math. J. 62 (2013), 293-322.
- [49] P. Russell, On affine-ruled rational surfaces, Math. Ann. 255 (1981), 287-302.
- [50] H. Schoutens, The use of ultraproducts in commutative algebra, (Springer-Verlag, 2010).
- [51] C. T. Simpson, Harmonic bundles on noncompact curves, J. Am. Math. Soc. 3 (1990), 713–770.
- [52] C. T. Simpson, Higgs bundles and local systems, Publ. Math. IHES 75 (1992), 5-95.
- [53] C. T. Simpson, Moduli of representations of the fundamental group of a smooth projective variety I, Publ. Math. IHES 79 (1994), 47–129.
- [54] C. T. Simpson, Moduli of representations of the fundamental group of a smooth projective variety II, Publ. Math. IHES 80 (1995), 5–79.
- [55] C. T. Simpson, *The dual boundary complex of the SL*₂ character variety of a punctured sphere, Ann. Fac. Sci. Toulouse **25** (2016), 317–361.
- [56] M. Spitzweck, A commutative P¹-spectrum representing motivic cohomology over Dedekind domains, Mém. Soc. Math. Fr. 157 (2018), 1–114.
- [57] The Stacks project authors, *The Stacks project*, https://stacks.math.columbia.edu, 2023.
- [58] T. Su, Dual boundary complexes of Betti moduli spaces over the two sphere with one irregular singularity, arXiv:2109.01645.
- [59] T. Su, Integral cohomology of dual boundary complexes is motivic, arXiv:???.
- [60] S. Szabo, Simpson's geometric P=W conjecture in the Painlevé VI case via abelianization, arXiv:1906.01856.
- [61] S. Szabo, Perversity equals weight for Painlevé spaces, Adv. Math. 383 (2021), 107667.
- [62] J. P. Whang, Global geometry on moduli of local systems for surfaces with boundary, Compos. Math. 156 (2020), 1517–1559. Email address: sutao@bimsa.cn

BEIJING INSTITUTE OF MATHEMATICAL SCIENCES AND APPLICATIONS