

CONVERGENCE TO EQUILIBRIUM FOR A DEGENERATE MCKEAN-VLASOV EQUATION

MANH HONG DUONG¹ & AMIT EINA²

ABSTRACT. In this work we study the convergence to equilibrium for a (potentially) degenerate nonlinear and nonlocal McKean-Vlasov equation. We show that the solution to this equation is related to the solution of a linear degenerate and/or defective Fokker-Planck equation and employ recent sharp convergence results to obtain an easily computable (and many times sharp) rates of convergence to equilibrium for the equation in question.

KEYWORDS. McKean-Vlasov equation, Fokker-Planck equation, entropy, long time behaviour.

MSC. 35Q84 (Fokker-Planck equations), 35Q82 (PDEs in connection with statistical mechanics), 35H10 (Hypoelliptic equations), 35K10 (second order parabolic equations), 35B40 (Asymptotic behavior of solutions to PDEs).

CONTENTS

1. Introduction	1
1.1. Background	1
1.2. The setting of the problem	2
1.3. Organisation of the paper	6
2. Solving the McKean-Vlasov equation	6
3. About the convergence to equilibrium	9
4. Final remarks	13
Appendix A. The verification of our solution	13
References	14

1. INTRODUCTION

1.1. **Background.** In this work we consider the following McKean-Vlasov type equation

$$(1.1) \quad \partial_t \rho(x, t) = \operatorname{div} \left[\mathbf{C}x\rho(x, t) + \left(\int_{\mathbb{R}^d} \mathbf{K}(x-y)\rho(y, t)dy \right) \rho(x, t) + \mathbf{D}\nabla \rho(x, t) \right],$$

where $t \in (0, \infty)$, $x \in \mathbb{R}^d$, and $\mathbf{C}, \mathbf{K}, \mathbf{D}$ are $d \times d$ constant matrices that correspond to the drift, interaction and diffusion phenomena of the system, respectively.

The research of MHD was supported by EPSRC Grants EP/V038516/1 and EP/W008041/1.

Equation (1.1) is nonlinear and nonlocal in a typical way of McKean-Vlasov type of equations - that is, via an interaction term which is presented as a convolution with a kernel \mathbf{K} . McKean-Vlasov PDEs, originally proposed by McKean in his 1966 work [14] where he explored connections between a wider class of Markov processes and non-linear parabolic equations, have been shown to have a strong connection to systems of interacting particle/agents by means of the so-called mean field limit approach. In particular, it can be shown that (1.1) arises naturally as the mean-field limit of the following interacting particle system

$$(1.2) \quad dX_i(t) = -\mathbf{C}X_i(t) dt - \frac{1}{N} \sum_{j=1}^N \mathbf{K}(X_i(t) - X_j(t)) dt + \sqrt{2\mathbf{D}} dW_i(t)$$

with $i = 1, \dots, N$ and where $\{W_i(t)\}_{i=1, \dots, N}$ are d -dimensional standard independent Wiener processes. Indeed, as can be seen in the classical works [14, 15, 18], the particle system described by (1.2) satisfies the *propagation of chaos* property, i.e. as N goes to infinity any fixed number of particles in the system become independent, and the limiting SDE equation that described an average particle in (1.2) is given by

$$(1.3) \quad d\bar{X}(t) = -\mathbf{C}\bar{X}(t) dt - \mathbf{K}(\bar{X}(t) - \mathbb{E}(\bar{X}(t))) dt + \sqrt{2\mathbf{D}} dW(t).$$

The law of this limiting particle is a weak solution to our McKean-Vlasov equation (1.1).

Systems such as (1.2), describing the evolution of N interacting particles where every particle moves under the influence of an external force, an interacting force (Curie-Weiss type interaction) and a stochastic noise, have many applications in physics, biology and social sciences. Such systems have been investigated intensively in the literature and their study remains a very active field. In particular, the McKean-Vlasov equation (1.1) has been considered recently in [8, 10] in the context of interacting diffusions driven by (non-Markovian) coloured noise. For additional information on the topic we refer the reader to recent review works [9, 13].

The goal of our presented work is to consider degenerate and defective McKean-Vlasov type of equations and utilise recent study of similar Fokker-Planck equation to conclude easily computable (and many times sharp) rates of convergence to equilibrium for (1.1).

1.2. The setting of the problem. We would like to start and emphasise that our main focus in this paper is the long time behaviour of the solution to (1.1) and *not* its existence, uniqueness and regularity. The well-posedness and smoothness of the solution to general McKean-Vlasov equations under certain restricted Hörmander conditions (which are slightly stronger than the assumptions of this work), has been shown in [1] in the one dimensional case. For additional information and settings we refer the interested reader to [5, 6, 7]. We will assume from this point onwards that the solution to our equation exists, is unique, and is smooth with finite first moment. We will, in fact, find an explicit and smooth solution to the problem under relatively mild conditions on the initial datum.

Thus, in our setting, the uniqueness of solutions to (1.1) will be enough (see Remark 2.2 in §2).

The study of the long time behaviour for (1.1) which we will present here is heavily motivated by (and relies on) recent works on the Fokker-Planck equation (which in our setting can be viewed as (1.1) with no interaction term, i.e. $\mathbf{K} = 0$) such as [3, 16] and [2, 4]. Due to the particular structure and type of McKean-Vlasov equation we consider here, we are able to allow for more degeneracies and find an explicit rate of convergence to the equilibrium of the system under our underline “distance” - the Boltzmann entropy. This will not only provide a different approach to this study than those found in recent work on more general (over-damped) McKean-Vlasov equations and Vlasov-Fokker-Planck equations (such as [6, 12, 11, 17]), but will also explicitly show the impact of \mathbf{C} , \mathbf{K} , and \mathbf{D} on the convergence rate.

Following on the above, we present some important notions from the recent studies of Fokker-Planck equations which will allow us to state our main theorem.

Definition 1.1. The (relative) Boltzmann entropy of a non-negative measurable function $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ with respect to a positive measurable function $g : \mathbb{R}^d \rightarrow \mathbb{R}$, both of which of unit mass, i.e.

$$\int_{\mathbb{R}^d} f(x) dx = \int_{\mathbb{R}^d} g(x) dx = 1,$$

is defined as

$$\mathcal{H}(f|g) = \int_{\mathbb{R}^d} f(x) \log\left(\frac{f(x)}{g(x)}\right) dx.$$

Remark 1.2. The Boltzmann entropy plays a fundamental role in many of the studies of convergence to equilibrium for various physically, biologically, and chemically motivated equations. While it is nonlinear and not a distance in nature, the Csiszár-Kullback-Pinsker inequality states that for any unit mass f and g as above we have that

$$\|f - g\|_{L^1(\mathbb{R}^d)} \leq \sqrt{2\mathcal{H}(f|g)}$$

which reassures us that the entropy is enough to give us “reasonable” convergence. The requirement that both f and g will be of unit mass is very natural in numerous equations which describe the evolution of a probability density of a physical, biological, or chemical phenomenon. The McKean-Vlasov equation is slightly different due to the (quadratic) interaction term but, as we will mention shortly, we are always able to assume that our solution has a unit mass by scaling the interaction matrix \mathbf{K} and using the fact that mass is conserved under (1.1) (see Remark 1.7).

Definition 1.3. Let \mathcal{A} and \mathcal{B} be $d \times d$ matrices. We say that the pair $(\mathcal{A}, \mathcal{B})$ is an admissible pair, or admissible in short, if:

(A) \mathcal{B} is positive semi-definite with

$$1 \leq \text{rank}(\mathcal{B}) \leq d.$$

- (B) \mathcal{A} is positively stable, i.e. all its eigenvalues have positive real part.
 (C) There exists no \mathcal{A}^T -invariant subspace of $\ker(\mathcal{B})$.

It is worth to mention that condition (C) in the above definition has been shown to be equivalent to Hörmander condition for the following operator [3]

$$L_{\mathcal{A}, \mathcal{B}} f = \operatorname{div}(\mathcal{A} x f + \mathcal{B} \nabla f),$$

associated to the Fokker-Planck equation

$$\partial_t f(x, t) = \operatorname{div}(\mathcal{A} x f + \mathcal{B} \nabla f).$$

In the subsequent analysis, we will relate the solution of the McKean-Vlasov equation (1.1) to that of the above Fokker-Planck equation with \mathcal{A} being a linear combination of \mathbf{C} and \mathbf{K} and $\mathcal{B} = \mathbf{D}$.

Lastly, we introduce a new notion for matrices that will prove to be very useful in our current study

Definition 1.4. We say that a square matrix \mathcal{A} is *almost-positively stable* if each of its eigenvalue satisfies

- (i) Its real part is positive, or
- (ii) It is zero. In this case, the algebraic and geometric multiplicity of the zero eigenvalue should equal.

Remark 1.5. From the definition it is simple to see that if \mathcal{A} is a $d \times d$ almost-positively stable matrix then its associated Jordan matrix, \mathbf{J} , must be of the form

$$(1.4) \quad \mathbf{J} = \begin{pmatrix} \mathbf{0}_{r \times r} & \mathbf{0}_{r \times (d-r)} \\ \mathbf{0}_{(d-r) \times r} & \mathbf{J}'_{(d-r) \times (d-r)} \end{pmatrix}$$

where \mathbf{J}' is, in general, a Jordan matrix with no zero entries on its diagonal. Consequently, denoting by $\{v_1, \dots, v_d\}$ the basis of generalised eigenvectors of \mathcal{A} such that $\ker \mathcal{A} = \operatorname{span}\{v_1, \dots, v_r\}$ (which can be inferred from \mathbf{J} and the appropriate similarity matrix) we see that the linear operator

$$(1.5) \quad P_{\ker \mathcal{A}} \left(\sum_{i=1}^d \alpha_i v_i \right) = \sum_{i=1}^r \alpha_i v_i$$

defines a projection on $\ker \mathcal{A}^1$. Moreover, the space

$$(1.6) \quad V := \operatorname{span}(\mathbf{I} - P_{\ker \mathcal{A}}) = \operatorname{span}\{v_{r+1}, \dots, v_d\}$$

is an \mathcal{A} -invariant subspace and the Jordan form associated to $\mathcal{A}|_V$ is given by \mathbf{J}' . We can conclude from this that $\mathcal{A}|_V$ is, in fact, positively stable.

With these definitions and remark at hand, we are ready to state our main theorem.

Theorem 1.6. *Consider the McKean-Vlasov equation (1.1) with drift, interaction, and diffusion matrices \mathbf{C} , \mathbf{K} , and \mathbf{D} respectively. Assume that the pair $(\mathbf{C} + \mathbf{K}, \mathbf{D})$ is admissible.*

¹Indeed, $P_{\ker \mathcal{A}}^2 = P_{\ker \mathcal{A}}$ and $P_{\ker \mathcal{A}}|_{\ker \mathcal{A}} = \mathbf{I}_{\ker \mathcal{A}}$.

Let $\rho(t)$ be the solution to the McKean-Vlasov equation with a unit mass initial datum $\rho_0 \in L^1_+(\mathbb{R}^d)^2$ that has a finite first moment and which guarantees the existence of a unique smooth solution to the problem. Denote by

$$(1.7) \quad \mathbf{m}_1 := \int_{\mathbb{R}^d} x \rho_0(x) dx,$$

$$(1.8) \quad s(t) := (e^{-\mathbf{C}t} - e^{-(\mathbf{C}+\mathbf{K})t}) \mathbf{m}_1$$

and

$$(1.9) \quad \rho_\infty(x) := \frac{1}{(2\pi \det \mathcal{K})^{\frac{d}{2}}} e^{-\frac{1}{2}x^T \mathcal{K} x},$$

with \mathcal{K} being the unique positive definite solution to the continuous Lyapunov equation

$$2\mathbf{D} = (\mathbf{C} + \mathbf{K}) \mathcal{K} + \mathcal{K} (\mathbf{C} + \mathbf{K})^T.$$

Let

$$(1.10) \quad \mu := \min \{ \operatorname{Re} \lambda \mid \lambda \text{ is an eigenvalue of } \mathbf{C} + \mathbf{K} \},$$

and

$$(1.11) \quad \nu := \begin{cases} \min \{ \operatorname{Re} \lambda \mid \lambda \neq 0 \text{ is an eigenvalue of } \mathbf{C} \}, & \mathbf{C} \neq 0, \\ 0, & \mathbf{C} = 0. \end{cases}$$

Then $s(t)$ converges to $s_\infty := P_{\ker \mathbf{C}} \mathbf{m}_1$ and there exists an explicit constant $c > 0$ that depends only on \mathbf{C} , \mathbf{K} , and \mathbf{D} , such that

$$(1.12) \quad \begin{aligned} \mathcal{H}(\rho(t) | \rho_\infty(\cdot - s_\infty)) &\leq c \mathcal{H}(\rho_0 | \rho_\infty) (1 + t^{2n_1}) e^{-2\mu t} \\ &+ c \left[(1 + t^{n_2}) e^{-\nu t} |(\mathbf{I} - P_{\ker \mathbf{C}}) \mathbf{m}_1| + (1 + t^{n_1}) e^{-\mu t} |\mathbf{m}_1| \right] \\ &\left[(1 + (1 + t^{n_1}) e^{-\mu t}) |P_{\ker \mathbf{C}} \mathbf{m}_1| + (1 + t^{n_2}) e^{-\nu t} |(\mathbf{I} - P_{\ker \mathbf{C}}) \mathbf{m}_1| \right] \end{aligned}$$

where $P_{\ker \mathbf{C}}$ is the projection on the kernel of \mathbf{C} , and n_1 and n_2 are the maximal defect of all eigenvalues associated to μ and ν respectively (n_2 is defined to be zero when $\mathbf{C} = 0$).

Remark 1.7. The keen reader would notice that we have required a unit mass initial datum ρ_0 . It is immediate to see that under the assumption of the existence of a smooth solution, $\rho(t)$, the mass is a conserved quantity³, i.e.

$$m_0(t) := \int_{\mathbb{R}^d} \rho(x, t) dx = \int_{\mathbb{R}^d} \rho_0(x) dx = m_0.$$

² $L^1_+(\mathbb{R}^d)$ is the space of all non-negative $L^1(\mathbb{R}^d)$ functions.

³Indeed

$$\frac{d}{dt} m_0(t) = \int_{\mathbb{R}^d} \operatorname{div} \left[\mathbf{C} x \rho(x, t) + \left(\int_{\mathbb{R}^d} \mathbf{K}(x-y) \rho(y, t) dy \right) \rho(x, t) + \mathbf{D} \nabla \rho(x, t) \right] dx = 0.$$

The McKean-Vlasov equation, (1.1), is not invariant under mass scaling. This is due to the interaction term. Replacing $\rho(t)$ by $\tilde{\rho}(t) = \rho(t)/m_0$, however, yields a *unit mass* solution to the McKean-Vlasov equation

$$\partial_t \tilde{\rho}(x, t) = \operatorname{div} \left[\mathbf{C}x \tilde{\rho}(x, t) + \left(\int_{\mathbb{R}^d} (m_0 \mathbf{K})(x - y) \tilde{\rho}(y, t) dy \right) \tilde{\rho}(x, t) + \mathbf{D} \nabla \tilde{\rho}(x, t) \right],$$

which is governed by the matrices \mathbf{C} , $m_0 \mathbf{K}$, and \mathbf{D} . We can thus apply our main theorem in this case as long as the associated matrices satisfy the required conditions.

Remark 1.8. The reader might notice that our expression for the rate of convergence of the solution under the Boltzmann entropy functional, inequality (1.12), seems quite cumbersome. The reason we kept it as such and didn't estimate it more crudely is to not only get a better explicit bound, but also to showcase the impact of the drift matrix \mathbf{C} on the result. In particular, if $\ker \mathbf{C} = \{0\}$ we get a much improved rate of convergence and we find that $s_\infty = P_{\ker \mathbf{C}} \mathbf{m}_1 = 0$, i.e. we find that the equilibrium to (1.1) is the same as that for the Fokker-Planck equation with drift matrix $(\mathbf{C} + \mathbf{K})$ and diffusion matrix \mathbf{D} .

1.3. Organisation of the paper. We start the body of our work finding an explicit solution to our equation (1.1) in §2, relying on known results from the study of Fokker-Planck equations. §3 will be dedicated to showing the main result of our paper followed by a short discussion about our results and future research in §4.

2. SOLVING THE MCKEAN-VLASOV EQUATION

The goal of this section is to provide an explicit solution to our McKean-Vlasov equation (1.1).

Our main result for this section is:

Theorem 2.1. *Consider the McKean-Vlasov equation (1.1) with drift, interaction, and diffusion matrices \mathbf{C} , \mathbf{K} , and \mathbf{D} respectively. Assume that the pair $(\mathbf{C} + \mathbf{K}, \mathbf{D})$ satisfies condition (C) of Definition 1.3, i.e. that there is no $(\mathbf{C} + \mathbf{K})^T$ -invariant subspace of $\ker \mathbf{D}$. Then, a unit mass solution to the equation with initial datum $\rho_0 \in L^1_+(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$, where $p \in (1, \infty]$, is given by*

$$(2.1) \quad \rho(x, t) = f_{FP}^{\mathbf{C}+\mathbf{K}, \mathbf{D}}(x - s(t), t),$$

with $f_{FP}^{\mathcal{A}, \mathcal{B}}(x, t)$ being the (unique) smooth solution to the Fokker-Planck equation

$$(2.2) \quad \partial_t f(x, t) = \operatorname{div} [\mathcal{A}x f(x, t) + \mathcal{B} \nabla f(x, t)]$$

where $t \in (0, \infty)$, $x \in \mathbb{R}^d$, $f(x, 0) = \rho_0(x)$, and $s(t)$ is defined as (1.8).

Remark 2.2. Theorem 2.1 will guarantee that we have an explicit expression to our solution to (1.1) as long as we know that $\rho_0 \in L^1_+(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$ and that the solution is unique.

Remark 2.3. Much like we've noted in Remark 1.7, the above theorem can easily be modified for the case we are considering a general solution to the problem by replacing the matrix \mathbf{K} with $m_0\mathbf{K}$, where m_0 is the mass of the solution.

Proof of Theorem 2.1. To emphasise the point that the mass doesn't matter, we will provide the general proof of this theorem, i.e. we assume that $m_0 > 0$ is arbitrary.

We start by assuming the our solution is smooth and decays fast enough and rewrite (1.1) as

$$(2.3) \quad \begin{aligned} \partial_t \rho(x, t) = & \operatorname{div} \left[\mathbf{C}x\rho(x, t) + \mathbf{K}x \underbrace{\left(\int_{\mathbb{R}^d} \rho(y, t) dy \right)}_{=m_0} \right] \rho(x, t) \\ & - \mathbf{K} \left(\int_{\mathbb{R}^d} y\rho(y, t) dy \right) \rho(x, t) + \mathbf{D}\nabla\rho(x, t). \end{aligned}$$

This motivates us to explore the momentum of our solution

$$\mathbf{m}_1(t) := \int_{\mathbb{R}^d} x\rho(x, t) dx.$$

Using (2.3) together with the conservation of mass, we see that for any $i = 1, \dots, d$

$$\begin{aligned} \frac{d}{dt} m_{1,i}(t) &= \int_{\mathbb{R}^d} x_i \partial_t \rho(x, t) dx \\ &= \int_{\mathbb{R}^d} x_i \operatorname{div} \left[(\mathbf{C} + m_0\mathbf{K})x\rho(x, t) - \left(\int_{\mathbb{R}^d} (\mathbf{K}y)\rho(y, t) dy \right) \rho(x, t) + \mathbf{D}\nabla\rho(x, t) \right] dx \\ &= - \int_{\mathbb{R}^d} ((\mathbf{C} + m_0\mathbf{K})x)_i \rho(x, t) dx + \left(\int_{\mathbb{R}^d} (\mathbf{K}y)_i \rho(y, t) dy \right) \underbrace{\int_{\mathbb{R}^d} \rho(x, t) dx}_{=m_0} - \sum_{j=1}^d d_{i,j} \int_{\mathbb{R}^d} \partial_{x_j} \rho(x, t) dx \\ &= - \int_{\mathbb{R}^d} (\mathbf{C}x)_i \rho(x, t) dx = - \sum_{j=1}^d c_{i,j} \int_{\mathbb{R}^d} x_j \rho(x, t) dx = -(\mathbf{C}\mathbf{m}_1(t))_i. \end{aligned}$$

We conclude that

$$\frac{d}{dt} \mathbf{m}_1(t) = -\mathbf{C}\mathbf{m}_1(t)$$

and consequently,

$$(2.4) \quad \mathbf{m}_1(t) = e^{-\mathbf{C}t} \mathbf{m}_1(0) := e^{-\mathbf{C}t} \mathbf{m}_1.$$

The above allows us to rewrite (2.3), and as a result (1.1), as

$$(2.5) \quad \partial_t \rho(x, t) = \operatorname{div} \left[(\mathbf{C} + m_0\mathbf{K})x\rho(x, t) - \mathbf{K}e^{-\mathbf{C}t} \mathbf{m}_1 \rho(x, t) + \mathbf{D}\nabla\rho(x, t) \right],$$

which could be recast as a Fokker-Planck equation with a "transport part"

$$(2.6) \quad \partial_t \rho(x, t) + \mathbf{K}e^{-\mathbf{C}t} \mathbf{m}_1 \cdot \nabla \rho(x, t) = \operatorname{div} \left[(\mathbf{C} + m_0\mathbf{K})x\rho(x, t) + \mathbf{D}\nabla\rho(x, t) \right].$$

Defining

$$\tilde{\rho}(x, t) := \rho(x + s(t), t)$$

for a function $s(t)$ which we will soon find, and denoting $y(x, t) = x + s(t)$ we find that

$$\begin{aligned}
\partial_t \tilde{\rho}(x, t) &= \partial_t \rho(y(x, t), t) + \nabla_y \rho(y(x, t), t) \cdot \frac{ds}{dt}(t) \\
&= \partial_t \rho(y(x, t), t) + \mathbf{K} e^{-\mathbf{C}t} \mathbf{m}_1 \cdot \nabla_y \rho(y(x, t), t) + \left(\frac{ds}{dt}(t) - \mathbf{K} e^{-\mathbf{C}t} \mathbf{m}_1 \right) \nabla_y \rho(y(x, t), t) \\
&= \operatorname{div}_y \left[(\mathbf{C} + m_0 \mathbf{K}) y(x, t) \rho(y(x, t), t) + \mathbf{D} \nabla_y \rho(y(x, t), t) \right] + \left(\frac{ds}{dt}(t) - \mathbf{K} e^{-\mathbf{C}t} \mathbf{m}_1 \right) \cdot \nabla_y \rho(y(x, t), t) \\
&= \operatorname{div}_x \left[(\mathbf{C} + m_0 \mathbf{K}) x \tilde{\rho}(x, t) + \mathbf{D} \nabla_x \tilde{\rho}(x, t) \right] + \left(\frac{ds}{dt}(t) - \mathbf{K} e^{-\mathbf{C}t} \mathbf{m}_1 + (\mathbf{C} + m_0 \mathbf{K}) s(t) \right) \cdot \nabla_y \rho(y(x, t), t).
\end{aligned}$$

Thus, if

$$(2.7) \quad \frac{ds}{dt}(t) = \mathbf{K} e^{-\mathbf{C}t} \mathbf{m}_1 - (\mathbf{C} + m_0 \mathbf{K}) s(t)$$

we conclude that $\tilde{\rho}(x, t)$ solved the Fokker-Planck equation (2.2) with drift matrix $\mathcal{A} = \mathbf{C} + m_0 \mathbf{K}$ and diffusion matrix $\mathcal{B} = \mathbf{D}$.

It is easy to verify that a general solution to (2.7) is given by

$$s_G(t) = \frac{1}{m_0} \left(e^{-\mathbf{C}t} - e^{-(\mathbf{C} + m_0 \mathbf{K})t} \right) \mathbf{m}_1 + e^{-(\mathbf{C} + m_0 \mathbf{K})t} s_G(0)$$

As was discussed in [3], the conditions on the pair $(\mathbf{C} + m_0 \mathbf{K}, \mathbf{D})$ guarantee the existence of a unique smooth solution to the Fokker-Planck equation

$$(2.8) \quad \partial_t \tilde{\rho}(x, t) = \operatorname{div}_x \left[(\mathbf{C} + m_0 \mathbf{K}) x \tilde{\rho}(x, t) + \mathbf{D} \nabla_x \tilde{\rho}(x, t) \right]$$

as long as the initial datum is in $L_+^1(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$ for some $p \in (1, \infty]$. We conclude that

$$(2.9) \quad \rho_G(x, t) = \tilde{\rho}(x - s_G(t), t) = f_{FP}^{\mathbf{C} + m_0 \mathbf{K}, \mathbf{D}}(x - s_G(t), t).$$

is a candidate for a smooth and strongly decaying solution to (1.1). By choosing $s_G(0) = 0$, i.e. considering the function

$$\rho(x, t) = f_{FP}^{\mathbf{C} + m_0 \mathbf{K}, \mathbf{D}}(x - s(t), t)$$

where

$$(2.10) \quad s(t) = \frac{1}{m_0} \left(e^{-\mathbf{C}t} - e^{-(\mathbf{C} + m_0 \mathbf{K})t} \right) \mathbf{m}_1,$$

we see that

$$\tilde{\rho}(x, 0) = \rho(x + s(0), 0) = \rho(x, 0) = \rho_0(x) \in L_+^1(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$$

and consequently have managed to find a potential solution. Plugging this function back in (1.1) completes the proof. \square

Remark 2.4. The verification that (2.9) is indeed a solution to (1.1) relies on the explicit form of $f_{FP}^{\mathbf{C} + m_0 \mathbf{K}, \mathbf{D}}$. It was shown in [3], the solution of the Fokker Planck equation (2.2) when the pair $(\mathcal{A}, \mathcal{B})$ is admissible is given explicitly by

$$(2.11) \quad f_{FP}^{\mathcal{A}, \mathcal{B}}(x, t) = \frac{1}{(2\pi \det \mathcal{Q}(t))^{\frac{d}{2}}} \int_{\mathbb{R}^d} e^{-\frac{1}{2}(x - e^{-\mathcal{A}t} y)^T \mathcal{Q}(t)^{-1} (x - e^{-\mathcal{A}t} y)} \rho_0(y) dy,$$

where the convolution kernel $\mathcal{Q}(t)$ is given by

$$\mathcal{Q}(t) := 2 \int_0^t e^{-\mathcal{A}s} \mathcal{B} e^{-\mathcal{A}^T s} ds.$$

We leave the proof to the Appendix as it is a straightforward calculation.

With an explicit expression of the solution to the McKean-Vlasov equation, one that is intimately connected to a Fokker-Planck equation, we can start considering the question of convergence to equilibrium.

3. ABOUT THE CONVERGENCE TO EQUILIBRIUM

The main result shown in the previous section - namely the fact that a solution to our McKean-Vlasov equation (1.1) is nothing but a “transported” solution to an appropriate Fokker-Planck equation - allows us to utilise recent sharp results about convergence to equilibrium in the setting of Fokker-Planck equations, in particular those achieved in the work of Monmarché [16], to conclude explicit and quantitative convergence to equilibrium in (1.1)

We begin our section by quoting the main convergence to equilibrium result from [16]:

Theorem 3.1. *Consider the Fokker-Planck equation*

$$\partial_t f(x, t) = \operatorname{div} [\mathcal{A}x f(x, t) + \mathcal{B} \nabla f(x, t)]$$

where the pair $(\mathcal{A}, \mathcal{B})$ is admissible. Let $f(\cdot, t)$ be the solution to the above equation with a unit mass initial datum $f_0 \in L^1_+(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$ for some $p \in (1, \infty]$ and let

$$\eta := \min \{ \operatorname{Re} \lambda \mid \lambda \text{ is an eigenvalue of } \mathcal{A} \}.$$

Then, there exists an explicit $c > 0$ that only depends on \mathcal{A} and \mathcal{B} such that

$$\mathcal{H}(f(t) | f_\infty) \leq c \mathcal{H}(f_0 | f_\infty) (1 + t^{2n}) e^{-2\eta t}$$

where n is the maximal defect of all eigenvalues associated to η and where

$$f_\infty(x) := \frac{1}{(2\pi \det \mathcal{Q})^{\frac{d}{2}}} e^{-\frac{1}{2} x^T \mathcal{Q} x},$$

with \mathcal{Q} being the unique positive definite solution to

$$2\mathcal{B} = \mathcal{A}\mathcal{Q} + \mathcal{Q}\mathcal{A}^T.$$

The above theorem and Theorem 2.1 provide us with the following result:

Theorem 3.2. *Consider the McKean-Vlasov equation (1.1) with drift, interaction, and diffusion matrices \mathbf{C} , \mathbf{K} , and \mathbf{D} respectively. Assume in addition that the pair $(\mathbf{C} + \mathbf{K}, \mathbf{D})$ is admissible and that the equation admits a unique solution. Let $\rho(\cdot, t)$ be the solution to the equation with initial unit mass datum $\rho_0 \in L^1_+(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$ for some $p \in (1, \infty]$. Then, following on the notations for $s(t)$, ρ_∞ , μ , and n_1 from Theorem 1.6 we have that*

$$(3.1) \quad \mathcal{H}(\rho(t) | \rho_\infty(\cdot - s(t))) \leq c \mathcal{H}(\rho_0 | \rho_\infty) (1 + t^{2n_1}) e^{-2\mu t}$$

for an explicit $c > 0$ that only depends on \mathbf{C} , \mathbf{K} , and \mathbf{D} .

Proof. Following on Theorem 2.1 we have that

$$\begin{aligned} \mathcal{H}(\rho(t)|\rho_\infty(\cdot - s(t))) &= \mathcal{H}\left(f_{FP}^{\mathbf{C}+\mathbf{K},\mathbf{D}}(\cdot - s(t), t), |\rho_\infty(\cdot - s(t))\right) \\ &= \mathcal{H}\left(f_{FP}^{\mathbf{C}+\mathbf{K},\mathbf{D}}(t)|\rho_\infty\right) \leq c\mathcal{H}(f_0|\rho_\infty)(1 + t^{2m})e^{-2\mu t}, \end{aligned}$$

where we have used the fact that a time translation in the spatial variable doesn't change the value of the relative entropy and Theorem 3.1. \square

Theorem 3.2 assures us that the solution to (1.1) will “stabilise”, in a sense, as long as $(\mathbf{C} + \mathbf{K}, \mathbf{D})$ (or $(\mathbf{C} + m_0\mathbf{K}, \mathbf{D})$ to be more general) is an admissible pair. The profile function, however, given by $\rho_\infty(\cdot - s(t))$ is not time independent in general. It is not surprising, however, that if $s(t)$ converges as time goes to infinity then a true equilibrium (which is by definition stationary) emerges.

We present the following technical lemma to show that under the conditions of Theorem 1.6 this is always the case.

Lemma 3.3. *Let \mathcal{A} and \mathcal{B} be $d \times d$ matrices such that $(\mathcal{A} + \mathcal{B})$ is positively stable. Define the operator*

$$\xi(t) := e^{-\mathcal{A}t} - e^{-(\mathcal{A}+\mathcal{B})t}.$$

Then $\xi(t)$ has a limit as t goes to infinity if and only if \mathcal{A} is almost-positively stable and in that case

$$\lim_{t \rightarrow \infty} \xi(t) = P_{\ker \mathcal{A}}.$$

Moreover, if we denote by

$$\alpha := \min \{ \operatorname{Re} \lambda \mid \lambda \text{ is an eigenvalue of } \mathcal{A} + \mathcal{B} \},$$

and

$$\beta := \begin{cases} \min \{ \operatorname{Re} \lambda \mid \lambda \neq 0 \text{ is an eigenvalue of } \mathcal{A} \}, & \mathcal{A} \neq 0, \\ 0, & \mathcal{A} = 0. \end{cases}$$

Then for any $x \in \mathbb{R}^d$

$$(3.2) \quad \begin{aligned} |\xi(t)x - P_{\ker \mathcal{A}}x| &\leq c_{\mathcal{A}}(1 + t^{n_2})e^{-\beta t} |(\mathbf{I} - P_{\ker \mathcal{A}})x| \\ &\quad + c_{\mathcal{A}+\mathcal{B}}(1 + t^{n_1})e^{-\alpha t} |x|, \end{aligned}$$

where $c_{\mathcal{A}} > 0$ and $c_{\mathcal{A}+\mathcal{B}}$ are explicit constants that depend only on \mathcal{A} and \mathcal{B} , and n_1 and n_2 are the maximal defect of all eigenvalues associated to α and β respectively (n_2 is defined to be zero when $\mathcal{A} = 0$).

Proof. To show the desired result, we start by considering single $r \times r$ Jordan block matrix of the form

$$\mathfrak{J} = \begin{pmatrix} \lambda & 1 & 0 & \dots & 0 & 0 \\ 0 & \lambda & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \lambda & 1 \\ 0 & 0 & 0 & \dots & 0 & \lambda \end{pmatrix}$$

A well known result from the theory of ODEs states that we can find an explicit constant $c_{\mathfrak{J}}$ such that⁴

$$\|e^{-\mathfrak{J}t}\| \leq c_{\mathfrak{J}}(1+t^r)e^{-\lambda t}.$$

The above implies that for any matrix \mathcal{C} with a Jordan form

$$\mathcal{J} = \begin{pmatrix} \mathcal{J}_1 & \mathbf{0} & \dots & \mathbf{0} \\ \vdots & \vdots & \vdots & \vdots \\ \mathbf{0} & \mathbf{0} & \dots & \mathcal{J}_l \end{pmatrix}$$

we have that

$$(3.3) \quad \|e^{-\mathcal{C}t}\| \leq c_S \|e^{-\mathcal{J}t}\| \leq c_S \max_{i=1,\dots,l} c_{\mathcal{J}_i} (1+t^{r_i}) e^{-\lambda_i t}$$

where c_S is a constant that depends on the similarity matrix taking \mathcal{C} to \mathcal{J} .

We can thus conclude that, with the notation of the lemma, if $\mathcal{A} + \mathcal{B}$ is a positively stable matrix then

$$(3.4) \quad \|e^{-(\mathcal{A}+\mathcal{B})t}\| \leq c_{\mathcal{A}+\mathcal{B}} (1+t^{n_1}) e^{-\alpha t}.$$

Next we notice that the definition of $P_{\ker \mathcal{A}}$, (1.5), for an almost-positively stable matrix imply that

$$e^{-\mathcal{A}t} - P_{\ker \mathcal{A}} = e^{-\mathcal{A}t} (\mathbf{I} - P_{\ker \mathcal{A}}).$$

Consequently, if we consider the \mathcal{A} -invariant space $V := \text{span}(\mathbf{I} - P_{\ker \mathcal{A}})$ on which \mathcal{A} is positively stable (see Remark 1.5), we find that (3.3) implies that for any $x \in \mathbb{R}^d$

$$(3.5) \quad \begin{aligned} \|e^{-\mathcal{A}t} x - P_{\ker \mathcal{A}} x\| &= \|e^{-\mathcal{A}t} (\mathbf{I} - P_{\ker \mathcal{A}}) x\| \\ &\leq \|e^{-\mathcal{A}t}|_V\| |(\mathbf{I} - P_{\ker \mathcal{A}}) x| \leq c_{\mathcal{A}} (1+t^{n_2}) e^{-\beta t} |(\mathbf{I} - P_{\ker \mathcal{A}}) x|. \end{aligned}$$

Since

$$|\xi(t)x - P_{\ker \mathcal{A}} x| \leq \left| e^{-\mathcal{A}t} x - P_{\ker \mathcal{A}} x \right| + \left| e^{-(\mathcal{A}+\mathcal{B})t} x \right|$$

the desired inequality follows directly from (3.4) and (3.5). \square

Remark 3.4. Lemma 3.3 and the conditions for Theorem 1.6 guarantee that $s(t)$, which is given by (1.8), will converge to $s_{\infty} = P_{\ker \mathbf{C}} \mathbf{m}_1$ as time goes to infinity. Moreover, using the notations of Theorem 1.6

$$(3.6) \quad \begin{aligned} |s(t) - s_{\infty}| &\leq c(1+t^{n_2}) e^{-\nu t} |(\mathbf{I} - P_{\ker \mathbf{C}}) \mathbf{m}_1| \\ &\quad + c(1+t^{n_1}) e^{-\mu t} |\mathbf{m}_1|, \end{aligned}$$

for any appropriate explicit constant $c > 0$.

It is worth to mention that $s(t)$ may converge under less restrictive conditions than the ones imposed in Theorem 1.6. For instance, if $\mathbf{m}_1 = 0$ then $s(t) = 0$ for all time regardless of \mathbf{C} and \mathbf{K} .

We are now ready to prove our main theorem.

⁴the interested reader may find a proof to this statement in the Proof of Lemma 4.7 in [2].

Proof of Theorem 1.6. We start by noticing that

$$\begin{aligned}
(3.7) \quad \mathcal{H}(\rho(t)|\rho_\infty(\cdot - s_\infty)) &= \int_{\mathbb{R}^d} \rho(x, t) \log\left(\frac{\rho(x, t)}{\rho_\infty(x - s_\infty)}\right) dx \\
&= \mathcal{H}(\rho(t)|\rho_\infty(\cdot - s(t))) + \int_{\mathbb{R}^d} \rho(x, t) \log\left(\frac{\rho_\infty(x - s(t))}{\rho_\infty(x - s_\infty)}\right) dx.
\end{aligned}$$

Since

$$\begin{aligned}
\frac{\rho_\infty(x - s(t))}{\rho_\infty(x - s_\infty)} &= e^{-\frac{1}{2}[(x-s(t))^T \mathcal{K}(x-s(t)) - (x-s_\infty)^T \mathcal{K}(x-s_\infty)]} \\
&= e^{(s(t)-s_\infty)^T \mathcal{K} x} e^{-\frac{1}{2}[s(t)^T \mathcal{K} s(t) - s_\infty^T \mathcal{K} s_\infty]} = e^{(s(t)-s_\infty)^T \mathcal{K} x} e^{-\frac{1}{2}(s(t)-s_\infty)^T \mathcal{K} (s(t)+s_\infty)},
\end{aligned}$$

where we have used the fact that for any symmetric matrix \mathcal{A}

$$(x - y)^T \mathcal{A} (x + y) = x^T \mathcal{A} x - y^T \mathcal{A} y,$$

we conclude that

$$\begin{aligned}
(3.8) \quad \mathcal{H}(\rho(t)|\rho_\infty(\cdot - s_\infty)) &= \mathcal{H}(\rho(t)|\rho_\infty(\cdot - s(t))) \\
&+ \int_{\mathbb{R}^d} (s(t) - s_\infty)^T \mathcal{K} x \rho(x, t) dx \\
&- \frac{1}{2} (s(t) - s_\infty)^T \mathcal{K} (s(t) + s_\infty) \int_{\mathbb{R}^d} \rho(x, t) dx \\
&= \mathcal{H}(\rho(t)|\rho_\infty(\cdot - s(t))) + (s(t) - s_\infty)^T \mathcal{K} e^{-\mathbf{C}t} \mathbf{m}_1 \\
&- \frac{1}{2} (s(t) - s_\infty)^T \mathcal{K} (s(t) + s_\infty)
\end{aligned}$$

where we have used the conservation of (unit) mass and (2.4).

Consequently, using the fact that \mathcal{K} is positive definite and as such there exists $\kappa > 0$ such that $\mathcal{K} \leq \kappa \mathbf{I}$, we find that

$$\begin{aligned}
(3.9) \quad \mathcal{H}(\rho(t)|\rho_\infty(\cdot - s_\infty)) &\leq \mathcal{H}(\rho(t)|\rho_\infty(\cdot - s(t))) \\
&+ \frac{\kappa}{2} |s(t) - s_\infty| (|P_{\ker \mathbf{C}} \mathbf{m}_1| + |(e^{-\mathbf{C}t} - P_{\ker \mathbf{C}}) \mathbf{m}_1|) \\
&+ \frac{\kappa}{2} |s(t) - s_\infty|^2 + \kappa |s_\infty| |s(t) - s_\infty|.
\end{aligned}$$

Using Theorem 3.2, inequality (3.6), the facts that $s_\infty = P_{\ker \mathbf{C}} \mathbf{m}_1$ and

$$|(e^{-\mathbf{C}t} - P_{\ker \mathbf{C}}) \mathbf{m}_1| \leq c_{\mathbf{C}} (1 + t^{n_2}) e^{-\nu t} |(\mathbf{I} - P_{\ker \mathbf{C}}) \mathbf{m}_1|$$

we find that

$$\begin{aligned}
\mathcal{H}(\rho(t)|\rho_\infty(\cdot - s_\infty)) &\leq c \mathcal{H}(\rho_0|\rho_\infty) (1 + t^{2n_1}) e^{-2\mu t} \\
&+ c [(1 + t^{n_2}) e^{-\nu t} |(\mathbf{I} - P_{\ker \mathbf{C}}) \mathbf{m}_1| + (1 + t^{n_1}) e^{-\mu t} |\mathbf{m}_1|] \\
&\left[(1 + (1 + t^{n_1}) e^{-\mu t}) |P_{\ker \mathbf{C}} \mathbf{m}_1| + (1 + t^{n_2}) e^{-\nu t} |(\mathbf{I} - P_{\ker \mathbf{C}}) \mathbf{m}_1| \right]
\end{aligned}$$

for an appropriate constant $c > 0$, which is the desired result. The proof is now complete. \square

4. FINAL REMARKS

While our main estimate, inequality (1.12), seems complicated, it gives us an easily computable and explicit rate of convergence to equilibrium. It is clear from the proof of our theorems, that various additional conditions on the drift, interaction, and diffusion matrices will allow us to refine our proofs and get better rates of convergence. For example, in the case $\mathbf{K} = 0$ we find that $s(t) = s_\infty = 0$ (as expected). Consequently, the rate of convergence to equilibrium, according to Theorem 3.2, is given by $(1 + t^{2n_1}) e^{-2\mu t}$, which is a better rate than the one we'll get from (1.12).

We would like to emphasise that the method we presented here can deal with many degenerate cases, such as the case $\mathbf{C} = 0$. In that case, as long as the pair (\mathbf{K}, \mathbf{D}) is admissible, we find that

$$\mathcal{H}(\rho(t)|\rho_\infty(\cdot - s_\infty)) \leq c \mathcal{H}(\rho_0|\rho_\infty) (1 + t^{2n_1}) e^{-2\mu t} + c(1 + t^{n_1}) e^{-\mu t} |\mathbf{m}_1|.$$

This showcase the importance of the interaction term, governed by \mathbf{K} , as in this instance a part of it takes the role of a drift which allows us to attain an equilibrium.

Another point we would like to mention is that the identity

$$\begin{aligned} \mathcal{H}(\rho(t)|\rho_\infty(\cdot - s_\infty)) &= \mathcal{H}(\rho(t)|\rho_\infty(\cdot - s(t))) \\ &+ (s(t) - s_\infty)^T \mathcal{K} e^{-\mathbf{C}t} \mathbf{m}_1 - \frac{1}{2} (s(t) - s_\infty)^T \mathcal{K} (s(t) + s_\infty), \end{aligned}$$

which was shown in the proof of our main theorem, Theorem 1.6, shows that the optimal rate of convergence to equilibrium is determined by the profile convergence, $\mathcal{H}(\rho_0|\rho_\infty) (1 + t^{2n_1}) e^{-2\mu t}$, and the optimal rate of convergence of $s(t)$ to s_∞ . As

$$s(t) = (e^{-\mathbf{C}t} - e^{-(\mathbf{C}+\mathbf{K})t}) \mathbf{m}_1,$$

we immediately see the significant impact of \mathbf{C} and \mathbf{K} , as well as the interplay between them, on the convergence. As an example we notice that since

$$\begin{aligned} |s(t) - s_\infty| &\leq c(1 + t^{n_2}) e^{-\nu t} |(\mathbf{I} - P_{\ker \mathbf{C}}) \mathbf{m}_1| \\ &+ c(1 + t^{n_1}) e^{-\mu t} |\mathbf{m}_1|, \end{aligned}$$

when \mathbf{C} is not positively stable (but is almost positively stable), the above will dominate the convergence to equilibrium.

In the near future, we would be interested to investigate if the result attained in this work can be attained via a process of mean field limit from the particle system associated to the McKean-Vlasov equation (1.1), as well as whether or not our result can be extended to non-quadratic potentials.

APPENDIX A. THE VERIFICATION OF OUR SOLUTION

In this short appendix we will show that the function

$$\rho(x, t) = f_{FP}^{\mathbf{C}+m_0\mathbf{K}, \mathbf{D}}(x - s(t), t),$$

with $f_{FP}^{\mathcal{A}, \mathcal{B}}$ defined as in (2.11) and $s(t)$ defined in (1.8), is a solution to our McKean-Vlasov equation (1.1).

Since

$$\frac{1}{(2\pi \det \mathcal{Q}(t))^{\frac{d}{2}}} \int_{\mathbb{R}^d} x e^{-\frac{1}{2}(x-e^{-\mathcal{A}t}y)^T \mathcal{Q}(t)^{-1}(x-e^{-\mathcal{A}t}y)} dx = e^{-\mathcal{A}t} y$$

we have that

$$\begin{aligned} & \mathbf{K} \int_{\mathbb{R}^d} x f_{FP}^{\mathbf{C}+m_0\mathbf{K}, \mathbf{D}}(x-s(t), t) dx = \mathbf{K}s(t) \int_{\mathbb{R}^d} f_{FP}^{\mathbf{C}+m_0\mathbf{K}, \mathbf{D}}(x, t) dx \\ & + \mathbf{K} \int_{\mathbb{R}^d} x f_{FP}^{\mathbf{C}+m_0\mathbf{K}, \mathbf{D}}(x, t) dx = \mathbf{K}s(t) \int_{\mathbb{R}^d} \rho_0(y) dy + \mathbf{K} \int_{\mathbb{R}^d} e^{-(\mathbf{C}+m_0\mathbf{K})t} y \rho_0(y) dy \\ & = m_0 \mathbf{K}s(t) + \mathbf{K} e^{-(\mathbf{C}+m_0\mathbf{K})t} \mathbf{m}_1 = \mathbf{K} (e^{-\mathbf{C}t} - e^{-(\mathbf{C}+m_0\mathbf{K})t}) \mathbf{m}_1 \\ & \quad + \mathbf{K} e^{-(\mathbf{C}+m_0\mathbf{K})t} \mathbf{m}_1 = \mathbf{K} e^{-\mathbf{C}t} \mathbf{m}_1. \end{aligned}$$

Consequently

$$\mathbf{K} \int_{\mathbb{R}^d} (x-y) f_{FP}^{\mathbf{C}+m_0\mathbf{K}, \mathbf{D}}(y-s(t), t) dy = m_0 \mathbf{K}x - \mathbf{K} e^{-\mathbf{C}t} \mathbf{m}_1$$

and we find that $\rho(x, t)$ satisfies

$$\begin{aligned} \partial_t \rho(x, t) &= \partial_t f_{FP}^{\mathbf{C}+m_0\mathbf{K}, \mathbf{D}}(y(x, t), t) - \nabla_y f_{FP}^{\mathbf{C}+m_0\mathbf{K}, \mathbf{D}}(y(x, t), t) \cdot \frac{ds}{dt}(t) \\ \operatorname{div}_y \left((\mathbf{C} + m_0\mathbf{K}) y(x, t) \rho(x, t) + \mathbf{D} \nabla_x \rho(x, t) \right) &- (\mathbf{K} e^{-\mathbf{C}t} \mathbf{m}_1 - (\mathbf{C} + m_0\mathbf{K}) s(t)) \nabla_y f_{FP}^{\mathbf{C}+m_0\mathbf{K}, \mathbf{D}}(y(x, t), t) \\ &= \operatorname{div}_y \left(((\mathbf{C} + m_0\mathbf{K}) y(x, t) - (\mathbf{K} e^{-\mathbf{C}t} \mathbf{m}_1 - (\mathbf{C} + m_0\mathbf{K}) s(t))) \rho(x, t) + \mathbf{D} \nabla_x \rho(x, t) \right) \\ &= \operatorname{div}_y \left(((\mathbf{C} + m_0\mathbf{K}) (y(x, t) + s(t)) - \mathbf{K} e^{-\mathbf{C}t} \mathbf{m}_1) \rho(x, t) + \mathbf{D} \nabla_x \rho(x, t) \right) \\ &= \operatorname{div} \left(((\mathbf{C} + m_0\mathbf{K}) x - \mathbf{K} e^{-\mathbf{C}t} \mathbf{m}_1) \rho(x, t) + \mathbf{D} \nabla_x \rho(x, t) \right) \\ &= \operatorname{div} \left(\mathbf{C}x \rho(x, t) + (m_0 \mathbf{K}x - \mathbf{K} e^{-\mathbf{C}t} \mathbf{m}_1) \rho(x, t) + \mathbf{D} \nabla_x \rho(x, t) \right) \\ &= \operatorname{div} \left(\mathbf{C}x \rho(x, t) + \left(\int_{\mathbb{R}^d} \mathbf{K}(x-y) \rho(y, t) dy \right) \rho(x, t) + \mathbf{D} \nabla \rho(x, t) \right), \end{aligned}$$

where here $y(x, y) = x - s(t)$. This shows the desired result.

REFERENCES

- [1] F. Antonelli, and A. Kohatsu-Higa, *Rate of convergence of a particle method to the solution of the McKean–Vlasov equation*, Ann. Appl. Probab. (2002), 423–476.
- [2] A. Arnold, A. Einav, and T. Wöhner, *On the rates of decay to equilibrium in degenerate and defective Fokker–Planck equations*, J. Differential Equations. (2018), 6843–6872.
- [3] A. Arnold, and J. Erb, *Sharp entropy decay for hypocoercive and non-symmetric Fokker–Planck equations with linear drift*, arXiv preprint (arXiv:1409.5425). (2014).
- [4] A. Arnold, A. Einav, and T. Wöhner, *Generalised Fisher Information approach to defective Fokker–Planck equations*, arXiv preprint (arXiv:2210.14033). (2022).
- [5] R. Buckdahn, J. Li, S. Peng, and C. Rainer, *Mean-field stochastic differential equations and associated PDEs*, Ann. Probab. (2017), 824 – 878.
- [6] J. A. Carrillo, E. J. McCann, and C. Villani, *Kinetic equilibration rates for granular media and related equations: entropy dissipation and mass transportation estimates*, Rev. Mat. Iberoamericana (2003), 971–1018.
- [7] D. Crisan, and E. McMurray, *Smoothing properties of McKean–Vlasov SDEs*, Probab. Theory Related Fields. (2018), 97–148.

- [8] H. Duong and G. A. Pavliotis, *Mean field limits for non-Markovian interacting particles: convergence to equilibrium, GENERIC formalism, asymptotic limits and phase transitions*, Comm. Math. Sci. **16** (2018), 2199–2230.
- [9] F. Golse, *On the dynamics of large particle systems in the mean field limit*, Macroscopic and large scale phenomena: coarse graining, mean field limits and ergodicity. (2016), 1–144.
- [10] S. N. Gomes and G. A. Pavliotis, *Mean field limits for interacting diffusions in a two-scale potential*, J. Nonlinear Sci. **28** (2018), 905–941.
- [11] A. Guillin, W. Liu, L. Wu, and C. Zhang, *Uniform Poincaré and logarithmic Sobolev inequalities for mean field particle systems*, Ann. App. Prob. **32** (3), 1590 – 1614, 2022.
- [12] A. Guillin, and P. Monmarché, *Uniform long-time and propagation of chaos estimates for mean field kinetic particles in non-convex landscapes*, J. Stat. Phys. (2021), 1–20.
- [13] P-E. Jabin and Z. Wang, *Mean field limit for stochastic particle systems*, Active Particles, Volume 1: Advances in Theory, Models, and Applications. (2017), 379–402.
- [14] H. P. Jr. McKean, *A class of Markov processes associated with nonlinear parabolic equations*, Proc. Nat. Acad. Sci. U.S.A., **56** (1966), 1907–1911.
- [15] H. P. Jr. McKean, *Propagation of chaos for a class of non-linear parabolic equations*, Stochastic Differential Equations (Lecture Series in Differential Equations, Session 7, Catholic Univ. (1967), 41–57.
- [16] P. Monmarché, *Generalized Γ calculus and application to interacting particles on a graph*, Potential Anal. **3** (2019), 439–466.
- [17] P. Ren, and F-Y. Wang, *Exponential convergence in entropy and Wasserstein for McKean-Vlasov SDEs*, Nonlinear Anal. (2021).
- [18] A-S. Sznitman, *Topics in propagation of chaos*, Ecole d’été de probabilités de Saint-Flour XIX 1989. (1991), 165–251.

¹UNIVERSITY OF BIRMINGHAM, SCHOOL OF MATHEMATICS, EDGBASTON, BIRMINGHAM, B15 2TT, UNITED KINGDOM

Email address: h.duong@bham.ac.uk

²DURHAM UNIVERSITY, SCHOOL OF MATHEMATICAL SCIENCES, UPPER MOUNTJOY CAMPUS, STOCKTON ROAD, DH1 3LE, DURHAM, UNITED KINGDOM

Email address: amit.einav@durham.ac.uk