

Contractive coupling rates and curvature lower bounds for Markov chains

Francesco Pedrotti

August 2, 2023

Abstract

Contractive coupling rates have been recently introduced by Conforti as a tool to establish convex Sobolev inequalities (including modified log-Sobolev and Poincaré inequality) for some classes of Markov chains. In this work, we show how contractive coupling rates can also be used to prove stronger inequalities, in the form of curvature lower bounds for Markov chains and geodesic convexity of entropic functionals. We illustrate this in several examples discussed by Conforti, where in particular, after appropriately choosing a parameter function, we establish positive curvature in the entropic and (discrete) Bakry–Émery sense. In addition, we recall and give straightforward generalizations of some notions of coarse Ricci curvature, and we discuss some of their properties and relations with the concepts of couplings and coupling rates: as an application, we show exponential contraction of the p -Wasserstein distance for the heat flow in the aforementioned examples.

Contents

1	Introduction	2
2	Preliminaries and main inequality	4
2.1	Logarithmic mean and entropic curvature	5
2.2	Weight functions and convex Sobolev inequalities	6
2.3	Arithmetic mean and discrete Bakry–Émery curvature	8
3	Coupling rates and curvature lower bound	9
4	Applications	11
4.1	Glauber dynamics	11
4.1.1	Examples of Glauber dynamics	14
4.2	Bernoulli-Laplace model	16
4.2.1	Proof of Theorem 4.8	16
4.3	Hardcore model	18
4.3.1	Proof of Theorem 4.12	19
4.4	Interacting random walks	22
4.4.1	Proof of Theorem 4.17	24
4.4.2	Examples of interacting random walks	26
5	Couplings and coarse Ricci curvature	28
5.1	Discrete time	28
5.2	Continuous time	31
5.3	Comparison discrete and continuous time	37
5.4	Applications and related problems	38
A	Proof of Proposition 2.2	40
B	Proof of Proposition 2.4	40
C	Grönwall’s lemma with Dini derivative	41

1 Introduction

In this work, we are mostly concerned with finite state space continuous time Markov chains and we assume that they are irreducible and reversible: we use the letter Ω for the state space, L for the generator and m for the invariant measure. A fundamental problem in the theory of Markov chains consists in estimating the speed of convergence to the stationary distribution and giving upper bound for its mixing time. Convex Sobolev inequalities are a particular useful tool to address this task. Given a convex function $\phi: \mathbb{R}_{>0} \rightarrow \mathbb{R}$ such that $\phi \in C^1((0, \infty))$, we define the ϕ -entropy of a function $\rho: \Omega \rightarrow \mathbb{R}_{>0}$ by

$$\mathcal{H}^\phi(\rho) := \mathbb{E}_m[\phi \circ \rho] - \phi(\mathbb{E}_m[\rho]). \quad (1)$$

Notice that by Jensen's inequality $\mathcal{H}^\phi(\rho) \geq 0$ and $\mathcal{H}^\phi(C) = 0$ for any constant $C > 0$. Therefore, if we denote by $\rho = \frac{d\mu}{dm}$ the density of a probability measure μ with respect to m , we can think of $\mathcal{H}^\phi(\rho)$ as a (non-symmetric) measure of distance of μ from m . We also recall the definition of the Dirichlet form $\mathcal{E}: \mathbb{R}^\Omega \times \mathbb{R}^\Omega \rightarrow \mathbb{R}$ via

$$\mathcal{E}(f, g) := -\mathbb{E}_m[f(Lg)],$$

and of the ϕ -Fisher information

$$\mathcal{I}^\phi(\rho) := \mathcal{E}(\rho, \phi' \circ \rho). \quad (2)$$

We then say that a ϕ -convex Sobolev inequality holds with constant $K > 0$ (notation: $\text{CSI}_\phi(K)$) if for all positive functions $\rho: \Omega \rightarrow \mathbb{R}_{>0}$ we have that

$$K \mathcal{H}^\phi(\rho) \leq \mathcal{I}^\phi(\rho). \quad (3)$$

The interest behind this inequality lies in the fact that, denoting by $P_t = e^{tL}$ the semigroup associated to the generator L , we have the well known identity

$$\frac{d}{dt} \mathcal{H}^\phi(P_t \rho) = -\mathcal{I}^\phi(P_t \rho)$$

for any $\rho: \Omega \rightarrow \mathbb{R}_{\geq 0}$. Thus, by Grönwall Lemma, (3) is equivalent to the exponential decay of the entropy along the heat flow

$$\mathcal{H}^\phi(P_t \rho) \leq e^{-Kt} \mathcal{H}^\phi(\rho),$$

and therefore quantifies the speed of convergence to equilibrium of the Markov chain. Classical choices of the function ϕ include the function ϕ_α for $\alpha \in [1, 2]$ defined by

$$\phi_\alpha(t) = \begin{cases} t \log t - t + 1 & \text{if } \alpha = 1, \\ \frac{t^\alpha - t}{\alpha - 1} - t + 1 & \text{if } \alpha \in (1, 2]. \end{cases} \quad (4)$$

When $\phi = \phi_1$, we get the relative entropy and inequality (3) is the celebrated modified log-Sobolev inequality [2] (notation: $\text{MLSI}(K)$); when $\phi = \phi_2$, we find the variance and (3) is the Poincaré inequality (notation: $\text{PO}(K)$). For $\alpha \in (1, 2)$, inequalities (3) are known as Beckner inequalities, which interpolate between modified log-Sobolev and Poincaré [2, 16].

Curvature of Markov chains In the setting of Riemannian manifolds, positive Ricci curvature lower bounds have been linked to many functional inequalities: this has motivated the seminal independent works of Sturm [27] and Lott and Villani [18], who extended the notion of curvature lower bound and many of its consequences (including some logarithmic Sobolev inequalities) to a large class of geodesic metric measure spaces. In spite of its generality, this theory does not apply to Markov chains on discrete spaces; for this reason, several adapted notions of curvature have been proposed, based on different equivalent characterisations of Ricci curvature for Riemannian manifolds. Among these, we recall in particular the entropic curvature by Erbar and Maas [11], which is based on displacement convexity of the relative entropy with respect to an adapted Wasserstein-like metric \mathscr{W} introduced in [20] (see also the work of Mielke [21]). This theory shares many similarities with the classical Lott–Sturm–Villani one, and among its merits it is such that many of the desired functional inequalities follow from positive lower bounds on the Ricci curvature, including in particular the modified log-Sobolev inequality. Moreover, as shown in [10], the role of the classical relative entropy (with respect to m) can be taken over by other ϕ -entropy functionals as defined in (1), provided one changes accordingly a parameter function in the definition of \mathscr{W} : once again, from positive geodesic convexity one can derive many consequences, including the convex Sobolev inequality (3). Unfortunately, establishing positive lower bound for the entropic Ricci curvature (or more generally K -geodesic convexity of an entropic functional \mathcal{H}^ϕ) of a Markov chain can be challenging, and in many interesting examples good estimates are not available.

Coupling rates and Conforti’s results While studying the entropic curvature of a Markov chain is a difficult task, in general even finding good estimates on the best constant for the modified log-Sobolev inequality (or other convex Sobolev inequalities) can be difficult. For this reason, in the recent paper [5], Conforti introduced a method based on the new notion of *coupling rates* to study general convex Sobolev inequalities, and applied it to some interesting classes of Markov chains. Coupling rates are a modification of the familiar notion of couplings: roughly speaking, they are used to “couple” the action of the generator L from two different states. While couplings have been extensively used to establish fast convergence of Markov chains (“probabilistic” approach to fast mixing), their use to establish convex Sobolev inequalities (which belong to the “analytic” approach to fast mixing) is less common, and the results of [5] give an interesting connection between these two families of methods.

Our contribution and organization of the paper In this work, we show that couplings rates are also a powerful tool to establish entropic curvature lower bounds for Markov chains and other related inequalities. By adapting Conforti’s arguments, we illustrate this in most of the examples discussed in [5]. As an illustration of the applications of these methods, we state below one particular instance of our results. We refer to Sections 2–4 below for precise definitions.

Theorem 1.1 (Cf. Sections 4.1.1, 4.3). *Denote by Ric_e the entropic curvature of a Markov chain, as introduced in [11].*

- *For the Curie–Weiss model with size N and parameter $\beta > 0$, in the limit $N \rightarrow \infty$ we have*

$$\text{Ric}_e \geq (1 - \beta) + (1 - 2\beta)e^{-\beta}$$

for $\beta \leq \frac{1}{2}$.

- *For the Ising model in dimension d with parameter $\beta > 0$, we have*

$$\text{Ric}_e \geq 1 + e^{-2\beta d} - 3d(1 - e^{-2\beta})e^{2\beta d}$$

if $2d(1 - e^{-2\beta})e^{4d\beta} \leq 1$.

- *For the hardcore model on a graph with maximum degree Δ and parameter $\beta > 0$, set $\kappa_* = 1 - \beta(\Delta - 1)$ and $\overline{\kappa_*} = \min\{\beta, 1 - \beta\Delta\}$. Then*

$$\text{Ric}_e \geq \frac{\kappa_*}{2} + \overline{\kappa_*}$$

provided that $\beta\Delta \geq 1$.

We remark that in all the examples above we find new estimates for the entropic curvature of those Markov chains: these estimates imply in particular MLSI with the same constant obtained in [5], but also other interesting functional inequalities, such as exponential contractivity along the heat flow of the popular Wasserstein-like metric \mathscr{W} of [20] (see Section 2.1 and references therein for more details). Therefore, in this sense, we provide a strengthening of the results of [5].

To conclude this section, we briefly present below the organization of this paper while giving a more complete overview of our contributions.

- In Section 2 we give some preliminary definitions and introduce the general abstract inequality that we consider: it reads $\mathscr{B}(\rho, \psi) \geq K\mathscr{A}(\rho, \psi)$ for a constant K and all $\rho: \Omega \rightarrow \mathbb{R}_{>0}$ and $\psi: \Omega \rightarrow \mathbb{R}$ and depends on an additional weight function $\theta: \mathbb{R}_{>0} \times \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$. In the following subsections, we motivate the interest in studying this inequality. In particular:
 - In Section 2.1 we explain that when θ is the logarithmic mean then the inequality corresponds to an entropic curvature lower bound for the Markov chain (see [11]). We recall some necessary definitions and consequences of such curvature lower bound, including MLSI($2K$).
 - In Section 2.2 we extend the considerations of the previous section to the case where the relative entropy is replaced by some other ϕ -entropy functionals, and in particular explain how we recover a family of convex Sobolev inequalities as in (3) (consistently with [5]), together with other functional inequalities, by applying the results of [10].
 - In Section 2.3 we recall that if θ is the arithmetic mean then the inequality of interest corresponds to a lower bound for another notion of discrete curvature, namely the discrete Bakry–Émery one (see [26]).

- In Section 3 we recall the definition of coupling rates. Moreover, we show how (under some basic assumptions on the weight function θ) they can be useful for establishing our main inequality, especially when they are “contractive” as in [5]: we explain this by providing a technical lemma and some heuristic considerations, which can also guide future applications of this tool.
- In Section 4 we illustrate the considerations of the previous section by considering most of the examples discussed in [5] and correspondingly establishing the general abstract inequality $\mathcal{B}(\rho, \psi) \geq K\mathcal{A}(\rho, \psi)$ introduced in Section 2. More specifically:
 - In Section 4.1 we consider Glauber dynamics, which includes in particular the Ising model and the Curie–Weiss model.
 - In Section 4.2 we consider a simplified version of the Bernoulli–Laplace model.
 - In Section 4.3 we consider the classical hardcore model.
 - In Section 4.4 we consider the case of interacting random walks on the discrete grid \mathbb{N}^d .

In particular, by choosing θ to be the logarithmic mean, we find new estimates for the entropic curvature of the Ising model, the Curie–Weiss model and the hardcore model, and we recover the best known lower bound for the entropic curvature of the Bernoulli–Laplace model.

- Finally, in Section 5 we make some considerations about the notion of contractive coupling rates and the connection to the notion of coarse Ricci curvature [24] and its variants, reviewing some well known definitions and properties and providing some natural generalizations. In particular, as done by Conforti for the specific case of the interacting random walks of Section 4.4, we show that in all the other examples discussed in Section 4 we have an exponential contraction of the Wasserstein distance (with respect to the natural graph distance d on Ω) of the form

$$W_p(P_t\rho, P_t\sigma) \leq e^{-\frac{Kt}{p}} W_p(\rho, \sigma)$$

for all starting densities ρ, σ and $p \geq 1$, when the assumptions of the respective main theorems are satisfied.

We emphasize how, in some sense, this section shows that the connections between probabilistic and analytic methods emerging in [5] carry over at the level of curvature: contractive couplings are indeed naturally linked to coarse Ricci curvature, and yet we use them to establish in particular entropic curvature lower bounds, a rather analytic notion of curvature.

Acknowledgment

The author gratefully acknowledges support by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No. 716117) and by the Austrian Science Fund (FWF), Project SFB F65. The author also warmly thanks Jan Maas for suggesting the project and for his guidance, and Melchior Wirth and Haonan Zhang for useful discussions.

2 Preliminaries and main inequality

Following [5], we work with a so called “mapping representation” of the Markov chain, which we briefly recall. We are given a set of moves G (where a move $\sigma \in G$ is a function $\sigma: \Omega \rightarrow \Omega$) together with a transition rate function $c: \Omega \times G \rightarrow \mathbb{R}_{\geq 0}$, so that $c(\eta, \sigma)$ represents the rate of using move σ starting from state η . Such a mapping representation has already proved useful before in establishing functional inequalities and curvature lower bounds for Markov chains [25, 4, 11]. Typically (and if not otherwise specified) we use the letter η for a state and $\sigma, \gamma, \bar{\gamma}$ for moves. We make the assumption that for each move σ there exists a unique inverse $\sigma^{-1} \in G$ such that $\sigma^{-1}\sigma\eta = \eta$ whenever $m(\eta)c(\eta, \sigma) > 0$ and, without loss of generality, we also assume that a null move e (the identity map) belongs to G . With this notation, we can write explicitly the action of the generator L of the continuous time Markov chain in the form

$$L\psi(\eta) = \sum_{\sigma} c(\eta, \sigma)(\psi(\sigma\eta) - \psi(\eta)) \tag{5}$$

for any $\psi: \Omega \rightarrow \mathbb{R}$. We also use the notation

$$\nabla_{\sigma}\psi(\eta) := \psi(\sigma\eta) - \psi(\eta)$$

for the discrete gradient and

$$S := \{(\eta, \sigma) \in \Omega \times G \mid c(\eta, \sigma) > 0\}.$$

The reversibility condition implies that for all bounded functions $F: \Omega \times G \rightarrow \mathbb{R}$ we have

$$\sum_{\eta \in \Omega, \sigma \in G} m(\eta)c(\eta, \sigma)F(\eta, \sigma) = \sum_{\eta \in \Omega, \sigma \in G} m(\eta)c(\eta, \sigma)F(\sigma\eta, \sigma^{-1}). \quad (6)$$

To introduce our main inequality of interest in an abstract way, we first need an additional ingredient, i.e. a weight function $\theta: \mathbb{R}_{>0} \times \mathbb{R}_{>0} \rightarrow \mathbb{R}_{\geq 0}$. In this paper, we always work under the following basic

Assumption 1. *The weight function θ is such that:*

1. θ is not identically 0;
2. $\theta(s, t) = \theta(t, s)$;
3. θ is differentiable;
4. θ is concave.

For given θ , it will be convenient to also introduce the additional quantity

$$M_\theta := \inf_{\substack{s, t > 0: \\ \theta(s, t) > 0}} \frac{\theta(s, s) + \theta(t, t)}{2\theta(s, t)} \in [0, 1], \quad (7)$$

where we see that $M_\theta \leq 1$ by choosing $s = t$. From the definition it follows immediately that $\forall s, t \geq 0$

$$2M_\theta\theta(s, t) \leq \theta(s, s) + \theta(t, t).$$

We are now ready to state the inequality which is the main object of study of this paper: given a weight function θ , we aim to prove that

$$\mathcal{B}(\rho, \psi) \geq K\mathcal{A}(\rho, \psi) \quad (8)$$

for all positive functions $\rho: \Omega \rightarrow \mathbb{R}_{>0}$, functions $\psi: \Omega \rightarrow \mathbb{R}$ and for a constant $K \in \mathbb{R}$ independent of ρ, ψ , where we have

$$\begin{aligned} \mathcal{A}(\rho, \psi) &= \frac{1}{2} \sum_{\rho, \sigma} m(\eta)c(\eta, \sigma)\theta(\rho(\eta), \rho(\sigma\eta))[\psi(\eta) - \psi(\sigma\eta)]^2, \\ \mathcal{B}(\rho, \psi) &= \mathcal{C}(\rho, \psi) - \mathcal{D}(\rho, \psi), \end{aligned}$$

with

$$\begin{aligned} \mathcal{C}(\rho, \psi) &= \frac{1}{4} \sum_{\eta, \sigma} m(\eta)c(\eta, \sigma) \left\{ \nabla\theta(\rho(\eta), \rho(\sigma\eta)) \cdot \begin{pmatrix} L\rho(\eta) \\ L\rho(\sigma\eta) \end{pmatrix} \right\} [\psi(\eta) - \psi(\sigma\eta)]^2, \\ \mathcal{D}(\rho, \psi) &= \frac{1}{2} \sum_{\eta, \sigma} m(\eta)c(\eta, \sigma)\theta(\rho(\eta), \rho(\sigma\eta))(\psi(\eta) - \psi(\sigma\eta))(L\psi(\eta) - L\psi(\sigma\eta)). \end{aligned}$$

Depending on the choice of θ , inequality (8) has different interpretations and consequences, which we explore in the next subsections.

2.1 Logarithmic mean and entropic curvature

The main reason for studying inequality (8) comes from the work [11] and for choosing as θ the logarithmic mean

$$\theta_1(s, t) := \int_0^1 s^{1-pt} dp = \begin{cases} \frac{s-t}{\log s - \log t} & \text{if } s \neq t, \\ s & \text{if } s = t. \end{cases}$$

Let us now denote by $\mathcal{P}(\Omega)$ the set of probability densities on Ω with respect to m , i.e. functions $\rho: \Omega \rightarrow \mathbb{R}_{\geq 0}$ such that $\mathbb{E}_m(\rho) = 1$, and by $\mathcal{P}_*(\Omega)$ the set of strictly positive densities. In [20], Maas introduced a Wasserstein-like

metric \mathscr{W} on $\mathcal{P}(\Omega)$ via a discrete variant of the Benamou–Brenier formula, and showed that, as in the classical setting, for any $\rho \in \mathcal{P}(\Omega)$ the heat flow $t \rightarrow P_t \rho = e^{tL} \rho$ is the gradient flow of the relative entropy functional in $(\mathcal{P}(\Omega), \mathscr{W})$ started at ρ , where the relative entropy is the restriction of the functional \mathcal{H}^{ϕ_1} (as in (1)) to $\mathcal{P}(\Omega)$. Writing it explicitly, for $\rho \in \mathcal{P}(\Omega)$ we have that

$$\mathcal{H}^{\phi_1}(\rho) = \sum_{\eta \in \Omega} m(\eta) \rho(\eta) \log(\rho(\eta)),$$

with the convention that $0 \log 0 = 0$ in the above sum. In this setting, inequality (8) can be interpreted as a lower bound of the Hessian of \mathcal{H}^{ϕ_1} with respect to \mathscr{W} (see [11]), or equivalently as a statement of K -geodesic convexity (see also the independent work of Mielke [21]). As gradient flows of geodesically K -convex functionals enjoy many useful properties, functional inequalities can be subsequently derived for the Markov chains. In the next proposition, we collect some of them and we refer the reader to [11, 14, 8] for the details and additional interesting results.

Proposition 2.1. *Assume that θ is the logarithmic mean and that inequality (8) holds for some $K \in \mathbb{R}$ and for all $\psi: \Omega \rightarrow \mathbb{R}$, $\rho \in \mathcal{P}_*(\Omega)$. Then:*

- the HWI(K) inequality

$$H^{\phi_1}(\rho) \leq \mathscr{W}(\rho, \mathbf{1}) \sqrt{I^{\phi_1}(\rho)} - \frac{K}{2} \mathscr{W}(\rho, \mathbf{1})^2$$

holds for all $\rho \in \mathcal{P}(\Omega)$;

- for any $\rho, \sigma \in \mathcal{P}(\Omega)$

$$\mathscr{W}(P_t \rho, P_t \sigma) \leq e^{-Kt} \mathscr{W}(\rho, \sigma);$$

- if $K > 0$ then the modified log-Sobolev inequality MLSI($2K$)

$$2K \mathcal{H}^{\phi_1}(\rho) \leq \mathcal{I}^{\phi_1}(\rho)$$

holds for all $\rho \in \mathcal{P}_*(\Omega)$.

When (8) holds for the logarithmic mean, we say that the *entropic curvature* of the Markov chain is bounded from below by K , and we use the notation

$$\text{Ric}_e \geq K.$$

For the logarithmic mean, we have $M_{\theta_1} = 1$ (cf. equation (7) and Proposition 2.4).

2.2 Weight functions and convex Sobolev inequalities

In some cases, it is possible to let other ϕ -entropies take over the role of the relative entropy in the previous section, following [10] and by choosing an appropriate weight function θ . First, we consider a function $\phi: \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ and correspondingly we make the following

Assumption 2. *The function $\phi: \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$ is such that*

1. ϕ is continuous and $\phi \in C^2(\mathbb{R}_{>0})$;
2. ϕ is strictly convex.

Moreover the weight function $\theta = \theta_\phi$ defined by

$$\theta(s, t) := \begin{cases} \frac{s-t}{\phi'(s) - \phi'(t)} & \text{if } s \neq t, \\ \frac{1}{\phi''(s)} & \text{if } s = t \end{cases} \quad (9)$$

satisfies Assumption 1.

A first motivation for defining θ as in (9) comes from the following

Proposition 2.2. *Suppose Assumption 2 is satisfied for some $K > 0$ and that inequality (8) holds. Then the convex Sobolev inequality (3) holds with constant $2K$ (notation: $\text{CSI}_\phi(2K)$).*

The proof of this proposition is given in Section A: the idea is that, when restricting to the specific choice $\psi := \phi' \circ \rho$, inequality (8) is equivalent to the second order differential inequality

$$\left. \frac{d^2}{dt^2} \right|_{t=0} \mathcal{H}^\phi(P_t \rho) \geq 2K \mathcal{I}^\phi(\rho). \quad (10)$$

From this, it is standard to deduce the convex Sobolev inequality (3) with constant $2K$, essentially by integration, following what is known as the ‘‘Bakry–Émery argument’’. Actually, this is exactly the approach used by Conforti in [5], that is, he uses couplings rates to establish (10) and subsequently deduces the convex Sobolev inequality $\text{CSI}_\phi(2K)$. Since (10) is a particular case of (8) for a specific choice of ψ , it is clear that proving (8) under Assumption 2 gives a stronger result as compared to (10) and is in general more challenging to achieve as one has to deal with two unknown functions (ρ and ψ) as opposed to just one (i.e. ρ). Another difference with the work of Conforti lies in the assumptions on the convex function ϕ : indeed, our Assumption 2 requires that θ is concave. This assumption was present also in [16] and (as already observed there) implies in particular that $\frac{1}{\phi''}$ is concave, which is a classical assumption for the continuous setting. On the other hand, Conforti does not assume concavity of θ , but requires instead that the function

$$(s, t) \rightarrow (s - t) \cdot (\phi'(s) - \phi'(t)) \quad (11)$$

is convex.

While both assumptions are enough to deduce convex Sobolev inequalities, it is possible to make a further more demanding one and deduce stronger consequences from inequality (8).

Assumption 3. *Assumption 2 is satisfied. Moreover, with θ as in (9), we have that:*

- $\phi \in C^\infty(\mathbb{R}_{>0})$;
- $\theta \in C^\infty(\mathbb{R}_{>0} \times \mathbb{R}_{>0})$;
- θ extends to a continuous function defined on $\mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}$;
- $\theta(r, s) \leq \theta(r, t)$ for all $0 \leq s \leq t$ and $0 \leq r$.

If the above is satisfied, then in [10] the authors showed it is possible to adapt some of the results of Section 2.1. Replacing the \mathcal{W} metric with a suitable modified metric \mathcal{W}_ϕ (where the new weight function θ replaces the logarithmic mean), it holds that for any starting density $\rho \in \mathcal{P}(\Omega)$ the heat flow $t \rightarrow P_t \rho$ is the gradient flow of the ϕ -entropy \mathcal{H}^ϕ in $(\mathcal{P}(\Omega), \mathcal{W}_\phi)$. Moreover, as in the previous section, inequality 8 is equivalent to K -geodesic convexity of \mathcal{H}^ϕ in $(\mathcal{P}(\Omega), \mathcal{W}_\phi)$ and the following holds

Proposition 2.3. *Under Assumption 3, suppose that inequality (8) holds for some $K \in \mathbb{R}$ and for all $\psi: \Omega \rightarrow \mathbb{R}$, $\rho \in \mathcal{P}_*(\Omega)$. Then:*

- the inequality

$$H^\phi(\rho) \leq \mathcal{W}_\phi(\rho, \mathbf{1}) \sqrt{I^\phi(\rho)} - \frac{K}{2} \mathcal{W}_\phi(\rho, \mathbf{1})^2$$

holds for all $\rho \in \mathcal{P}(\Omega)$;

- for any $\rho, \sigma \in \mathcal{P}(\Omega)$

$$\mathcal{W}_\phi(P_t \rho, P_t \sigma) \leq e^{-Kt} \mathcal{W}_\phi(\rho, \sigma);$$

- if $K > 0$ then the ϕ -convex Sobolev inequality $\text{CSI}_\phi(2K)$

$$2K \mathcal{H}^\phi(\rho) \leq \mathcal{I}^\phi(\rho)$$

holds for all $\rho \in \mathcal{P}_*(\Omega)$.

All the functions ϕ_α defined in (4) satisfy Assumption 3 (see also [16, Lemma 16]): for the corresponding weight function, we use the notation θ_α . Notice in particular that for $\alpha = 1$ we recover the logarithmic mean of the previous subsection, while for $1 < \alpha < 2$ we have

$$\theta_\alpha(s, t) = \begin{cases} \frac{\alpha-1}{\alpha} \frac{s-t}{s^{\alpha-1}-t^{\alpha-1}} & \text{if } s \neq t, \\ \frac{1}{\alpha} s^{2-\alpha} & \text{if } s = t. \end{cases} \quad (12)$$

Proposition 2.4. For $\alpha \in [1, 2]$ we have

$$M_{\theta_\alpha} = \begin{cases} 1 & \text{if } \alpha \in [1, \frac{3}{2}] \text{ or } \alpha = 2; \\ \frac{1}{2(\alpha-1)} & \text{if } \alpha \in (\frac{3}{2}, 2). \end{cases}$$

The proof of this proposition is given in Appendix B.

Remark 2.5 (Case $\alpha = 2$). The case $\alpha = 2$ is particular and should be studied separately. In this case, indeed, the weight function satisfies $\theta_2 \equiv \frac{1}{2}$. Therefore, the quantities $\mathcal{B}(\rho, \psi)$ and $\mathcal{A}(\rho, \psi)$ become independent of ρ , which makes establishing inequality (8) significantly simpler. Actually, for $\theta = \theta_2$ establishing (8) for all ρ, ψ is equivalent to establishing (10) for all ρ , as done by Conforti. Therefore, in this case it is usually possible to establish inequality (8) with a better constant than what would happen just under Assumption 1. In all the examples of Section 4, this can be achieved by a simple modification of the arguments after substituting $\theta \equiv \frac{1}{2}$, or alternatively, given the equivalence of (8) and (10), by just applying the results of [5]. For this reasons, for the results of Section 4 applied to $\theta = \theta_\alpha$ we focus on $\alpha \in [1, 2)$ when comparing to [5].

2.3 Arithmetic mean and discrete Bakry–Émery curvature

If θ is the arithmetic mean (for which $M_\theta = 1$), it is known that inequality (8) is equivalent to a lower bound for the discrete Bakry–Émery curvature (see for example [19]). For completeness, we recall the definitions and include a proof of this fact. In analogy with the classical setting, for $f, g: \Omega \rightarrow \mathbb{R}$ define

$$\Gamma(f, g)(\eta) := \frac{1}{2} \sum_{\sigma} c(\eta, \sigma) (f(\sigma\eta) - f(\eta))(g(\sigma\eta) - g(\eta)),$$

$\Gamma(f) := \Gamma(f, f)$ and

$$\Gamma_2(f) := \frac{1}{2} L\Gamma(f) - \Gamma(f, Lf).$$

Definition 2.6 ([26]). We say that the curvature condition $\text{CD}(K, \infty)$ is satisfied if for all $f: \Omega \rightarrow \mathbb{R}$

$$\Gamma_2(f) \geq K\Gamma(f). \tag{13}$$

Proposition 2.7. Suppose that θ is the arithmetic mean. Then for any $K \in \mathbb{R}$ inequality (8) holds if and only if $\text{CD}(K, \infty)$ holds.

Proof. Notice that, using reversibility,

$$\begin{aligned} \mathcal{A}(\rho, \psi) &= \frac{1}{4} \sum_{\eta, \sigma} m(\eta) c(\eta, \sigma) (\rho(\eta) + \rho(\sigma\eta)) [\psi(\eta) - \psi(\sigma\eta)]^2 \\ &= \frac{1}{2} \sum_{\eta} m(\eta) \rho(\eta) \left[\sum_{\sigma} c(\eta, \sigma) [\psi(\eta) - \psi(\sigma\eta)]^2 \right] \\ &= \sum_{\eta} m(\eta) \rho(\eta) \Gamma(\psi)(\eta). \end{aligned}$$

Moreover, using reversibility multiple times (cf. also (6)),

$$\begin{aligned}
\mathcal{E}(\rho, \psi) &= \frac{1}{8} \sum_{\eta} m(\eta) c(\eta, \sigma) (L\rho(\eta) + L\rho(\sigma\eta)) [\psi(\eta) - \psi(\sigma\eta)]^2 \\
&= \frac{1}{4} \sum_{\eta, \sigma} m(\eta) c(\eta, \sigma) (L\rho(\eta)) [\psi(\eta) - \psi(\sigma\eta)]^2 \\
&= \frac{1}{4} \sum_{\eta, \sigma, \gamma} m(\eta) c(\eta, \sigma) c(\eta, \gamma) (\rho(\gamma\eta) - \rho(\eta)) [\psi(\eta) - \psi(\sigma\eta)]^2 \\
&= \frac{1}{4} \sum_{\eta, \sigma, \gamma} m(\eta) c(\gamma\eta, \sigma) c(\eta, \gamma) \rho(\eta) [\psi(\gamma\eta) - \psi(\sigma\gamma\eta)]^2 \\
&\quad - \frac{1}{4} \sum_{\eta, \sigma, \gamma} m(\eta) c(\eta, \sigma) c(\eta, \gamma) \rho(\eta) [\psi(\eta) - \psi(\sigma\eta)]^2 \\
&= \frac{1}{4} \sum_{\eta} m(\eta) \rho(\eta) \sum_{\gamma} c(\eta, \gamma) \sum_{\sigma} \left\{ c(\gamma\eta, \sigma) [\psi(\gamma\eta) - \psi(\sigma\gamma\eta)]^2 - c(\eta, \sigma) [\psi(\eta) - \psi(\sigma\eta)]^2 \right\} \\
&= \sum_{\eta} m(\eta) \rho(\eta) \frac{1}{2} L\Gamma\psi(\eta),
\end{aligned}$$

and

$$\begin{aligned}
\mathcal{D}(\rho, \psi) &= \frac{1}{4} \sum_{\eta, \sigma} m(\eta) c(\eta, \sigma) (\rho(\eta) + \rho(\sigma\eta)) (\psi(\eta) - \psi(\sigma\eta)) (L\psi(\eta) - L\psi(\sigma\eta)) \\
&= \frac{1}{2} \sum_{\eta} m(\eta) \rho(\eta) \sum_{\sigma} c(\eta, \sigma) (\psi(\eta) - \psi(\sigma\eta)) (L\psi(\eta) - L\psi(\sigma\eta)) \\
&= \sum_{\eta} m(\eta) \rho(\eta) \Gamma(\psi, L\psi)(\eta).
\end{aligned}$$

Therefore, (8) is equivalent to

$$\sum_{\eta} m(\eta) \rho(\eta) \Gamma_2\psi(\eta) \geq K \sum_{\eta} m(\eta) \rho(\eta) \Gamma\psi(\eta). \tag{14}$$

From this, it is clear that $\text{CD}(K, \infty)$ implies (8) by choosing $f = \psi$. Conversely, choosing $\rho = \frac{d\delta_{\eta}}{dm}$ to be the density of a Dirac and $\psi = f$ in (14) gives the converse implication. \square

For more details about and consequences of the discrete Bakry–Émery curvature we refer the reader to [15] and the references therein.

3 Coupling rates and curvature lower bound

Coupling rates were introduced by Conforti in [5] as a tool to establish convex Sobolev inequalities. They are a modifications of the usual notion of coupling, and they apply to continuous time Markov chains. Roughly speaking, they are a way of letting the generator L act at the same time on two different states in a way that is consistent with equation (5) when one looks separately at the two states. More precisely, for any $\eta \in \Omega$ and $\sigma \in G$, we consider coupling rates between the neighbouring state $\eta, \sigma\eta$ in the form of a function $c^{\text{cpl}}(\eta, \sigma\eta, \cdot, \cdot): G \times G \rightarrow \mathbb{R}_{\geq 0}$ such that

$$\begin{aligned}
\sum_{\bar{\gamma}} c^{\text{cpl}}(\eta, \sigma\eta, \gamma, \bar{\gamma}) &= c(\eta, \gamma), \\
\sum_{\gamma} c^{\text{cpl}}(\eta, \sigma\eta, \gamma, \bar{\gamma}) &= c(\sigma\eta, \bar{\gamma}).
\end{aligned}$$

It can be seen easily seen that coupling rates always exist (up to changing without loss of generality the values $c(\sigma, e)$ and $c(\sigma\eta, e)$, where we recall that $e \in G$ is the identity map/null move. In [5], Conforti showed that coupling

rates are useful for organizing the terms appearing in the inequality (10) and thus (if one manages to establish it), in proving convex Sobolev inequalities via the Bakry–Émery argument. Heuristically, it turns out it is convenient to consider not arbitrary coupling rates, but rather “contractive” ones, where contractive informally means that “as often as possible” $\gamma\eta = \bar{\gamma}\sigma\eta$ (and, in particular, a fruitful choice is $(\gamma, \bar{\gamma}) = (\sigma, e)$ or (e, σ^{-1})). Indeed, when this is achieved, some terms cancellations going into the right direction occur when studying inequality (10).

In the rest of this section, we show that similar considerations also hold when studying the stronger inequality (8). Below, we derive a lower bound for $\mathcal{B}(\rho, \psi)$ using coupling rates under only Assumption 1 on θ : this gives a sufficient condition for establishing the inequality $\mathcal{B}(\rho, \psi) \geq K\mathcal{A}(\rho, \psi)$. In general, this is a more challenging situation compared to (10), since now we are dealing with two unknowns (ρ, ψ) as opposed to just ρ , and we also have an additional non linear weight function θ to deal with. Moreover, as discussed in Section 2.2, inequality (10) corresponds to a particular case of (8) with the choice $\psi = \phi' \circ \rho$ and θ as in Assumption 2.

We now proceed to show how arbitrary coupling rates can help rewrite the main inequality (8) in a convenient way, by organizing the involved terms. Notice first that we can write, using coupling rates,

$$\begin{aligned}
& \nabla\theta(\rho(\eta), \rho(\sigma\eta)) \cdot \begin{pmatrix} L\rho(\eta) \\ L\rho(\sigma\eta) \end{pmatrix} \\
&= \nabla\theta(\rho(\eta), \rho(\sigma\eta)) \cdot \left[\sum_{\gamma, \bar{\gamma}} c^{\text{cpl}}(\eta, \sigma\eta, \gamma, \bar{\gamma}) \begin{pmatrix} \rho(\gamma\eta) - \rho(\eta) \\ \rho(\bar{\gamma}\sigma\eta) - \rho(\sigma\eta) \end{pmatrix} \right] \\
&= \sum_{\gamma, \bar{\gamma}} c^{\text{cpl}}(\eta, \sigma\eta, \gamma, \bar{\gamma}) \nabla\theta(\rho(\eta), \rho(\sigma\eta)) \cdot \left[\begin{pmatrix} \rho(\gamma\eta) \\ \rho(\bar{\gamma}\sigma\eta) \end{pmatrix} - \begin{pmatrix} \rho(\eta) \\ \rho(\sigma\eta) \end{pmatrix} \right] \\
&\geq \sum_{\gamma, \bar{\gamma}} c^{\text{cpl}}(\eta, \sigma\eta, \gamma, \bar{\gamma}) [\theta(\rho(\gamma\eta), \rho(\bar{\gamma}\sigma\eta)) - \theta(\rho(\eta), \rho(\sigma\eta))],
\end{aligned} \tag{15}$$

where we used concavity of θ in the last line.

Therefore for $\mathcal{C}(\rho, \psi)$ we have the lower bound

$$\begin{aligned}
\mathcal{C}(\rho, \psi) &\geq \frac{1}{4} \sum_{\eta, \sigma} \sum_{\gamma, \bar{\gamma}} m(\eta) c(\eta, \sigma) c^{\text{cpl}}(\eta, \sigma\eta, \gamma, \bar{\gamma}) \theta(\rho(\gamma\eta), \rho(\bar{\gamma}\sigma\eta)) [\psi(\eta) - \psi(\sigma\eta)]^2 \\
&\quad - \frac{1}{4} \sum_{\eta, \sigma} \sum_{\gamma, \bar{\gamma}} m(\eta) c(\eta, \sigma) c^{\text{cpl}}(\eta, \sigma\eta, \gamma, \bar{\gamma}) \theta(\rho(\eta), \rho(\sigma\eta)) [\psi(\eta) - \psi(\sigma\eta)]^2.
\end{aligned}$$

As for the term $\mathcal{D}(\rho, \psi)$, we can write

$$\begin{aligned}
\mathcal{D}(\rho, \psi) &= \frac{1}{2} \sum_{\eta, \sigma} \sum_{\gamma, \bar{\gamma}} m(\eta) c(\eta, \sigma) c^{\text{cpl}}(\eta, \sigma\eta, \gamma, \bar{\gamma}) \theta(\rho(\eta), \rho(\sigma\eta)) (\psi(\eta) - \psi(\sigma\eta)) (\psi(\gamma\eta) - \psi(\bar{\gamma}\sigma\eta)) \\
&\quad - \frac{1}{2} \sum_{\eta, \sigma} \sum_{\gamma, \bar{\gamma}} m(\eta) c(\eta, \sigma) c^{\text{cpl}}(\eta, \sigma\eta, \gamma, \bar{\gamma}) \theta(\rho(\eta), \rho(\sigma\eta)) (\psi(\eta) - \psi(\sigma\eta))^2.
\end{aligned}$$

Combining the bound for \mathcal{C} and the expression for \mathcal{D} we derive the following:

Lemma 3.1. *We have*

$$\mathcal{B}(\rho, \psi) \geq \frac{1}{4} \sum_{\eta, \sigma} \sum_{\gamma, \bar{\gamma}} m(\eta) c(\eta, \sigma) c^{\text{cpl}}(\eta, \sigma\eta, \gamma, \bar{\gamma}) J(\eta, \sigma, \gamma, \bar{\gamma}) \tag{16}$$

for all $\rho: \Omega \rightarrow \mathbb{R}_{>0}$ and $\psi: \Omega \rightarrow \mathbb{R}$, where we define the function $J: \Omega \times G \times G \times G \rightarrow \mathbb{R}$ by

$$\begin{aligned}
J(\eta, \sigma, \gamma, \bar{\gamma}) &:= \{\theta(\rho(\gamma\eta), \rho(\bar{\gamma}\sigma\eta)) + \theta(\rho(\eta), \rho(\sigma\eta))\} [\psi(\eta) - \psi(\sigma\eta)]^2 \\
&\quad - 2\theta(\rho(\eta), \rho(\sigma\eta)) (\psi(\eta) - \psi(\sigma\eta)) (\psi(\gamma\eta) - \psi(\bar{\gamma}\sigma\eta)).
\end{aligned}$$

It is also convenient to define the function $I: \Omega \times G \times G \times G \rightarrow \mathbb{R}$ by

$$\begin{aligned}
I(\eta, \sigma, \gamma, \bar{\gamma}) &= I_1(\eta, \sigma, \gamma, \bar{\gamma}) - I_2(\eta, \sigma, \gamma, \bar{\gamma}), \\
I_1(\eta, \sigma, \gamma, \bar{\gamma}) &= \theta(\rho(\gamma\eta), \rho(\bar{\gamma}\sigma\eta)) [\psi(\eta) - \psi(\sigma\eta)]^2 \geq 0, \\
I_2(\eta, \sigma, \gamma, \bar{\gamma}) &= \theta(\rho(\eta), \rho(\sigma\eta)) [\psi(\gamma\eta) - \psi(\bar{\gamma}\sigma\eta)]^2 \geq 0.
\end{aligned}$$

Notice we have

$$\begin{aligned} J(\eta, \sigma, \gamma, \bar{\gamma}) &= I(\eta, \sigma, \gamma, \bar{\gamma}) + \theta(\rho(\eta), \rho(\sigma\eta))[\psi(\eta) - \psi(\sigma\eta) - \psi(\gamma\eta) + \psi(\bar{\gamma}\sigma\eta)]^2 \\ &\geq I(\eta, \sigma, \gamma, \bar{\gamma}). \end{aligned} \tag{17}$$

At this point, we can see at least heuristically why it is useful to consider especially *contractive* coupling rates. In view of Lemma 3.1, to establish inequality (8) it suffices to prove that for some coupling rates

$$\begin{aligned} &\frac{1}{2} \sum_{\eta, \sigma} \sum_{\gamma, \bar{\gamma}} m(\eta) c(\eta, \sigma) c^{\text{cpl}}(\eta, \sigma\eta, \gamma, \bar{\gamma}) J(\eta, \sigma, \gamma, \bar{\gamma}) \\ &\geq K \sum_{\eta, \sigma} m(\eta) c(\eta, \sigma) \theta(\rho(\eta), \rho(\sigma\eta)) [\psi(\sigma\eta) - \psi(\eta)]^2. \end{aligned} \tag{18}$$

Notice now that when $\gamma\eta = \bar{\gamma}\sigma\eta$ the second term in the definition of $J(\eta, \sigma\eta, \gamma, \bar{\gamma})$ is 0, so J is non-negative. In particular this is true when $(\gamma, \bar{\gamma})$ is equal to either (σ, e) or (e, σ^{-1}) , and in this case we find some terms appearing in the sum defining $\mathcal{A}(\rho, \psi)$ (i.e. in the right hand side of the previous inequality (18)).

We conclude this section by emphasizing that, while this method potentially applies to a wide variety of settings, in general it seems that some extra assumptions on the Markov chains are helpful to get the desired conclusions. In particular, reversibility of the model and an underlying symmetry of the structure of the Markov chain can help obtain useful terms cancellations.

4 Applications

In this section, we apply Lemma 3.1 to establish the general inequality of interest (8) for most of the examples considered in [5], under just Assumption 1. In particular, Section 4.1, 4.2, 4.3 and 4.4 corresponds to Section 4, 5.1, 5.2 and 3 of [5] respectively. Not surprisingly, the proofs are similar to the ones of Conforti, and in all these examples the considered contractive coupling rates are the ones constructed in [5]. The case of the interacting random walks of Section 4.4 is the only one where an additional assumption is present as compared to [5]: moreover, as done by Conforti, in that section a localization procedure is used to deal with the infinite cardinality of the state space.

4.1 Glauber dynamics

We work in the setting of Section 4 of [5] (i.e. Glauber dynamics) and we use the same notation, which we briefly recall. The state space is a finite set Ω . We assume that $\sigma = \sigma^{-1}$ and $\sigma\gamma\eta = \gamma\sigma\eta$ for all moves $\sigma, \gamma \in G$ and states $\eta \in \Omega$. Given an inverse temperature parameter $\beta > 0$ and an Hamiltonian function $H: \Omega \rightarrow \mathbb{R}$, the rates are defined by

$$c(\eta, \sigma) = \exp\left(-\frac{\beta}{2} \nabla_{\sigma} H(\eta)\right),$$

where we recall the notation $\nabla_{\sigma} H(\eta) = H(\sigma\eta) - H(\eta)$ for the discrete gradient. The reversible measure is then the Gibbs measure

$$m(\eta) = \frac{1}{Z_{\beta}} \exp(-\beta H(\eta))$$

where $Z_{\beta} > 0$ is the appropriate normalization constant. Finally we make the technical assumption that $\kappa(\eta, \sigma) \geq 0$ for all states η and moves σ , where we define

$$\kappa(\eta, \sigma) := c(\sigma\eta, \sigma) - \sum_{\gamma: \gamma \neq \sigma} \max\{-\nabla_{\sigma} c(\eta, \gamma), 0\}.$$

This assumption is crucial for the construction of appropriate contractive coupling rates, for which we will apply Lemma 3.1. We also define the quantities

$$\kappa_* := \inf_{\eta, \sigma} \kappa(\eta, \sigma) + \kappa(\sigma\eta, \sigma), \quad \overline{\kappa}_* := \inf_{\eta, \sigma} \kappa(\eta, \sigma).$$

Notice that $2\overline{\kappa}_* \leq \kappa_*$.

Theorem 4.1. *With the previous notation, suppose that for all $\eta \in \Omega$ and $\sigma, \gamma \in G$ we have $\sigma\gamma = \gamma\sigma$, $\sigma = \sigma^{-1}$ and $\kappa(\eta, \sigma) \geq 0$. Let θ be a weight functions θ satisfying Assumption 1. Then*

$$\mathcal{B}(\rho, \psi) \geq \left(M_\theta \overline{\kappa_*} + \frac{\kappa_*}{2} \right) \mathcal{A}(\rho, \psi)$$

for all functions $\rho: \Omega \rightarrow \mathbb{R}_{>0}$ and $\psi: \Omega \rightarrow \mathbb{R}$.

Remark 4.2 (Comparison with [5]). *In [5, Thm. 4.1], under the same assumptions on the model, Conforti establishes inequality (10) and thus CSI $\phi(2K)$ with constant K equal to*

- $\frac{\kappa_*}{2}$ for general convex ϕ satisfying convexity of (11);
- $\frac{\kappa_*}{2} + \overline{\kappa_*}$ for $\phi = \phi_1$ (thus $\text{MLSI}(\kappa_* + 2\overline{\kappa_*})$);
- $\frac{\alpha}{2}\kappa_*$ for $\phi = \phi_\alpha$ with $\alpha \in (1, 2]$.

Thus, by the discussion in Section 2 (recalling Proposition 2.4), we obtain a stronger results for the case $\theta = \theta_1$ and complementary results for other choices of θ .

Proof of Theorem 4.1

As done in [5], we define

$$\begin{aligned} \Upsilon^<(\eta) &= \{(\sigma, \gamma) \mid \sigma \neq \gamma, \nabla_\sigma c(\eta, \gamma) < 0\}, \\ \Upsilon^>(\eta) &= \{(\sigma, \gamma) \mid \sigma \neq \gamma, \nabla_\sigma c(\eta, \gamma) > 0\}, \\ \Upsilon^=(\eta) &= \{(\sigma, \gamma) \mid \sigma \neq \gamma, \nabla_\sigma c(\eta, \gamma) = 0\}. \end{aligned}$$

We then define the same coupling rates:

$$c^{\text{cpl}}(\eta, \sigma\eta, \gamma, \bar{\gamma}) = \begin{cases} \min\{c(\sigma\eta, \gamma), c(\eta, \gamma)\} & \text{if } \gamma = \bar{\gamma} \text{ and } \sigma \neq \gamma, \\ -\nabla_\sigma c(\eta, \gamma) & \text{if } \bar{\gamma} = \sigma \text{ and } (\sigma, \gamma) \in \Upsilon^<(\eta), \\ \nabla_\sigma c(\eta, \bar{\gamma}) & \text{if } \gamma = \sigma \text{ and } (\sigma, \bar{\gamma}) \in \Upsilon^>(\eta), \\ \kappa(\sigma\eta, \sigma) & \text{if } \gamma = \sigma, \bar{\gamma} = e, \\ \kappa(\eta, \sigma) & \text{if } \gamma = e, \bar{\gamma} = \sigma, \\ 0 & \text{otherwise.} \end{cases}$$

Notice that these are admissible, thanks to the assumptions on the model. With these coupling rates and using Lemma 3.1 and (17) we can write $\mathcal{B}(\rho, \psi) \geq \frac{1}{4}(A + B + C + D)$ with

$$\begin{aligned} A &= \sum_{\eta, \sigma \neq \gamma} m(\eta) c(\eta, \sigma) \min\{c(\eta, \gamma), c(\sigma\eta, \gamma)\} I(\eta, \sigma, \gamma, \gamma), \\ B &= - \sum_{\eta, (\sigma, \gamma) \in \Upsilon^<(\eta)} m(\eta) c(\eta, \sigma) \nabla_\sigma c(\eta, \gamma) I(\eta, \sigma, \gamma, \sigma), \\ C &= \sum_{\eta, (\sigma, \bar{\gamma}) \in \Upsilon^>(\eta)} m(\eta) c(\eta, \sigma) \nabla_\sigma c(\eta, \bar{\gamma}) I(\eta, \sigma, \sigma, \bar{\gamma}), \\ D &= \sum_{\eta, \sigma} m(\eta) c(\eta, \sigma) (\kappa(\sigma\eta, \sigma) J(\eta, \sigma, \sigma, e) + \kappa(\eta, \sigma) J(\eta, \sigma, e, \sigma)). \end{aligned}$$

We show below that $A = B = C = 0$ and that $D \geq (4M_\theta \overline{\kappa_*} + 2\kappa_*) \mathcal{A}(\rho, \psi)$, which concludes the proof of the theorem. It is useful to have an auxiliary lemma:

Lemma 4.3. *For all $\eta \in \Omega$ and $\sigma, \gamma \in G$ with $\sigma \neq \gamma$ the following hold:*

1. $c(\eta, \sigma) \nabla_\sigma c(\eta, \gamma) = c(\eta, \gamma) \nabla_\gamma c(\eta, \sigma)$.
2. $c(\eta, \sigma) c(\sigma\eta, \gamma) = c(\eta, \gamma) c(\gamma\eta, \sigma)$.
3. $\nabla_\sigma c(\sigma\eta, \gamma) = -\nabla_\sigma c(\eta, \gamma)$.
4. $(\sigma, \gamma) \in \Upsilon^<(\eta) \iff (\gamma, \sigma) \in \Upsilon^<(\eta)$.

5. $(\sigma, \gamma) \in \Upsilon^>(\gamma\eta) \iff (\sigma, \gamma) \in \Upsilon^<(\eta)$.
6. $(\sigma, \gamma) \in \Upsilon^>(\sigma\eta) \iff (\sigma, \gamma) \in \Upsilon^<(\eta)$.
7. $(\sigma, \gamma) \in \Upsilon^=(\gamma\eta) \iff (\sigma, \gamma) \in \Upsilon^=(\eta)$
8. $I(\eta, \sigma, \gamma, \sigma) = -I(\eta, \gamma, \sigma, \gamma)$.
9. $I(\sigma\eta, \sigma, \sigma, \gamma) = -I(\gamma\eta, \gamma, \gamma, \sigma)$.
10. $I(\eta, \sigma, \gamma, \gamma) = -I(\gamma\eta, \sigma, \gamma, \gamma)$.

Proof of Lemma. Statements 1–7 were already observed in the proof of [5, Thm. 4.1] and are easy to check, while statements 8–10 are immediate from the definitions. \square

Term D We have

$$\begin{aligned} J(\eta, \sigma, \sigma, e) &= \{\theta(\rho(\sigma\eta), \rho(\sigma\eta)) + \theta(\rho(\eta), \rho(\sigma\eta))\}[\psi(\eta) - \psi(\sigma\eta)]^2 \\ J(\eta, \sigma, e, \sigma) &= \{\theta(\rho(\eta), \rho(\eta)) + \theta(\rho(\eta), \rho(\sigma\eta))\}[\psi(\eta) - \psi(\sigma\eta)]^2 \end{aligned}$$

and so

$$\begin{aligned} &\kappa(\sigma\eta, \sigma)J(\eta, \sigma, \sigma, e) + \kappa(\eta, \sigma)J(\eta, \sigma, e, \sigma) \\ &\geq \{\overline{\kappa_*}[\theta(\rho(\sigma\eta), \rho(\sigma\eta)) + \theta(\rho(\eta), \rho(\eta))] + \kappa_*\theta(\rho(\eta), \rho(\sigma\eta))\}[\psi(\eta) - \psi(\sigma\eta)]^2 \\ &\geq (2M_\theta\overline{\kappa_*} + \kappa_*)\theta(\rho(\eta), \rho(\sigma\eta))[\psi(\eta) - \psi(\sigma\eta)]^2. \end{aligned}$$

Therefore we get

$$D \geq (4M_\theta\overline{\kappa_*} + 2\kappa_*)\mathcal{A}(\rho, \psi).$$

Term B We have using Lemma 4.3

$$\begin{aligned} B &= - \sum_{\eta, (\sigma, \gamma) \in \Upsilon^<(\eta)} m(\eta)c(\eta, \sigma)\nabla_\sigma c(\eta, \gamma)I(\eta, \sigma, \gamma, \sigma) \\ &= - \sum_{\eta, (\gamma, \sigma) \in \Upsilon^<(\eta)} m(\eta)c(\eta, \sigma)\nabla_\sigma c(\eta, \gamma)I(\eta, \sigma, \gamma, \sigma) \\ &= - \sum_{\eta, (\sigma, \gamma) \in \Upsilon^<(\eta)} m(\eta)c(\eta, \gamma)\nabla_\gamma c(\eta, \sigma)I(\eta, \gamma, \sigma, \gamma) \\ &= \sum_{\eta, (\sigma, \gamma) \in \Upsilon^<(\eta)} m(\eta)c(\eta, \sigma)\nabla_\sigma c(\eta, \gamma)I(\eta, \sigma, \gamma, \sigma) \\ &= -B \end{aligned}$$

and so $B = 0$.

Term C This is similar to term B using reversibility. We have

$$C = \sum_{\eta, (\sigma, \bar{\gamma}) \in \Upsilon^>(\eta)} m(\eta)c(\eta, \sigma)\nabla_\sigma c(\eta, \bar{\gamma})I(\eta, \sigma, \sigma, \bar{\gamma}).$$

Using reversibility with $F(\eta, \sigma) = \sum_{\bar{\gamma}: (\sigma, \bar{\gamma}) \in \Upsilon^>(\eta)} \nabla_\sigma c(\eta, \bar{\gamma})I(\eta, \sigma, \sigma, \bar{\gamma})$, the other assumptions ($\sigma = \sigma^{-1}$) and Lemma 4.3 we get

$$C = - \sum_{\eta, (\sigma, \bar{\gamma}) \in \Upsilon^<(\eta)} m(\eta)c(\eta, \sigma)\nabla_\sigma c(\eta, \bar{\gamma})I(\sigma\eta, \sigma, \sigma, \bar{\gamma}).$$

We want to show this expression is 0 analogously to what done for B . Notice that

$$\begin{aligned}
C &= - \sum_{\eta, (\sigma, \bar{\gamma}) \in \Upsilon^<(\eta)} m(\eta) c(\eta, \sigma) \nabla_{\sigma} c(\eta, \bar{\gamma}) I(\sigma \eta, \sigma, \sigma, \bar{\gamma}) \\
&= - \sum_{\eta, (\bar{\gamma}, \sigma) \in \Upsilon^<(\eta)} m(\eta) c(\eta, \sigma) \nabla_{\sigma} c(\eta, \bar{\gamma}) I(\sigma \eta, \sigma, \sigma, \bar{\gamma}) \\
&= - \sum_{\eta, (\sigma, \bar{\gamma}) \in \Upsilon^<(\eta)} m(\eta) c(\eta, \bar{\gamma}) \nabla_{\bar{\gamma}} c(\eta, \sigma) I(\bar{\gamma} \eta, \bar{\gamma}, \bar{\gamma}, \sigma) \\
&= \sum_{\eta, (\sigma, \bar{\gamma}) \in \Upsilon^<(\eta)} m(\eta) c(\eta, \sigma) \nabla_{\sigma} c(\eta, \bar{\gamma}) I(\sigma \eta, \sigma, \sigma, \bar{\gamma}) \\
&= -C,
\end{aligned}$$

which implies $C = 0$.

Term A We split A into three different terms $A = A_1 + A_2 + A_3$, where

$$\begin{aligned}
A_1 &= \sum_{\eta, (\sigma, \gamma) \in \Upsilon^<(\eta)} m(\eta) c(\eta, \sigma) c(\sigma \eta, \gamma) I(\eta, \sigma, \gamma, \gamma) \\
A_2 &= \sum_{\eta, (\sigma, \gamma) \in \Upsilon^>(\eta)} m(\eta) c(\eta, \sigma) c(\eta, \gamma) I(\eta, \sigma, \gamma, \gamma) \\
A_3 &= \frac{1}{2} \sum_{\eta, (\sigma, \gamma) \in \Upsilon^=(\eta)} m(\eta) c(\eta, \sigma) (c(\sigma \eta, \gamma) + c(\eta, \gamma)) I(\eta, \sigma, \gamma, \gamma)
\end{aligned}$$

We want to show $A_1 + A_2 = 0$ and $A_3 = 0$. Notice that using reversibility with the function $F(\eta, \gamma) = \sum_{\sigma: (\sigma, \gamma) \in \Upsilon^>(\eta)} c(\eta, \sigma) I(\eta, \sigma, \gamma, \gamma)$ we get, using Lemma 4.3 and the assumptions of the model,

$$\begin{aligned}
A_2 &= \sum_{\eta, (\sigma, \gamma) \in \Upsilon^>(\eta)} m(\eta) c(\eta, \sigma) c(\eta, \gamma) I(\eta, \sigma, \gamma, \gamma) \\
&= \sum_{\eta, (\sigma, \gamma) \in \Upsilon^>(\gamma \eta)} m(\eta) c(\eta, \gamma) c(\gamma \eta, \sigma) I(\gamma \eta, \sigma, \gamma, \gamma) \\
&= - \sum_{\eta, (\sigma, \gamma) \in \Upsilon^<(\eta)} m(\eta) c(\eta, \sigma) c(\sigma \eta, \gamma) I(\eta, \sigma, \gamma, \gamma) \\
&= -A_1.
\end{aligned}$$

Hence the contribution of the first two terms is 0. It remains to show that $A_3 = 0$. This is done in a similar way by using reversibility with the function $F(\eta, \gamma) = \sum_{\sigma: (\sigma, \gamma) \in \Upsilon^=(\eta)} c(\eta, \sigma) I(\eta, \sigma, \gamma, \gamma)$ to see that (with the help of Lemma 4.3)

$$\begin{aligned}
&\sum_{\eta, (\sigma, \gamma) \in \Upsilon^=(\eta)} m(\eta) c(\eta, \sigma) c(\eta, \gamma) I(\eta, \sigma, \gamma, \gamma) \\
&= \sum_{\eta, (\sigma, \gamma) \in \Upsilon^=(\gamma \eta)} m(\eta) c(\eta, \gamma) c(\gamma \eta, \sigma) I(\gamma \eta, \sigma, \gamma, \gamma) \\
&= - \sum_{\eta, (\sigma, \gamma) \in \Upsilon^=(\eta)} m(\eta) c(\eta, \sigma) c(\sigma \eta, \gamma) I(\eta, \sigma, \gamma, \gamma).
\end{aligned}$$

Therefore $A_3 = 0$. This concludes the proof of the theorem.

4.1.1 Examples of Glauber dynamics

Below, following [5], we present two examples of Glauber dynamics models satisfying the assumptions of Theorem 4.1.

Curie Weiss model For the Curie Weiss model, the state space is the discrete hypercube $\Omega = \{-1, 1\}^N$ for some integer $N > 0$. The set of moves G is given by $G = \{\sigma_1, \dots, \sigma_N\} \cup \{e\}$ where $\sigma_i: \Omega \rightarrow \Omega$ corresponds to flipping the i -th bit. Finally, the Hamiltonian function is

$$H(\eta) := -\frac{1}{2N} \sum_{i,j} \eta_i \eta_j.$$

In this setting, the assumptions of Theorem 4.1 and the explicit values of κ_* , $\overline{\kappa}_*$ were checked by Conforti, who proves the following

Theorem 4.4 (Thm. 4.2 of [5]). *Assume that*

$$(N-1) \left(e^{\frac{2\beta}{N}} - 1 \right) \leq 1.$$

Then the assumptions of 4.1 are satisfied with

$$\begin{aligned} \kappa_* &= f_{CW,\beta,N} \left(\left\lfloor \frac{N-1}{2} \right\rfloor \right), \\ \overline{\kappa}_* &= e^{-\frac{\beta}{N}(N-1)} \left[1 - (N-1) \left(1 - e^{\frac{2\beta}{N}} \right) \right], \end{aligned}$$

where $f_{CW,\beta,N}: \mathbb{N} \rightarrow \mathbb{R}$ is defined by

$$\begin{aligned} f_{CW,\beta,N}(m) &:= e^{-\frac{\beta}{N}(N-1-2m)} \left[1 - (N-1-m) \left(e^{\frac{2\beta}{N}} - 1 \right) \right] \\ &\quad + e^{\frac{\beta}{N}(N-1-2m)} \left[1 - m \left(e^{\frac{2\beta}{N}} - 1 \right) \right]. \end{aligned}$$

Remark 4.5 (Comparison with [9]). *In particular, in the limit $N \rightarrow \infty$, the condition above reads $\beta \leq \frac{1}{2}$. Thus by choosing $\theta = \theta_1$ and combining Theorems 4.1 and 4.4 we have that as $N \rightarrow \infty$ the entropic curvature of the Curie–Weiss model satisfies*

$$\text{Ric}_e \geq (1-\beta) + (1-2\beta)e^{-\beta}$$

for $\beta \leq \frac{1}{2}$. This improves both in the estimate and in the range of admissible β over [9, Cor. 4.5], where it was proved that, as $N \rightarrow \infty$, $\text{Ric}_e \geq 2(1-2\beta e^{2\beta})e^{-\beta}$ for $\beta \lesssim 0.284$.

Ising model The second example of Glauber dynamics that we consider is the Ising model. For the Ising model, we let $\Lambda \subset \mathbb{Z}^d$ be a connected subset of \mathbb{Z}^d , endowed with the inherited graph structure \sim of the discrete grid, and consider state space $\Omega = \{-1, 1\}^\Lambda$. The set of moves is $G = \{\sigma_x\}_{x \in \Lambda} \cup \{e\}$ where $\sigma_x: \Omega \rightarrow \Omega$ acts on a state η by flipping the spin η_x at site x . Finally, the Hamiltonian is defined by

$$H(\eta) := -\frac{1}{2} \sum_{x \sim y} \eta_x \eta_y.$$

Again, the assumptions and values of κ_* , $\overline{\kappa}_*$ in Theorem 4.1 were checked in [5], where the following result is proved.

Theorem 4.6 (Thm. 4.3 of [5]). *Assume that*

$$2d(1 - e^{-2\beta})e^{4d\beta} \leq 1. \tag{19}$$

Then the assumptions of Theorem 4.1 are satisfied and we have

$$\begin{aligned} \kappa_* &= 2 - 2d(1 - e^{-2\beta})e^{2\beta d} \\ \overline{\kappa}_* &= e^{-2\beta d} - 2d(1 - e^{-2\beta})e^{2\beta d}. \end{aligned}$$

Remark 4.7 (Comparison with Cor. 4.4. of [9]). *By combining Theorems 4.1 and 4.4 and by choosing the logarithmic mean $\theta = \theta_1$, it follows that*

$$\text{Ric}_e \geq 1 + e^{-2\beta d} - 3d(1 - e^{-2\beta})e^{2\beta d} \quad \text{if} \quad 2d(1 - e^{-2\beta})e^{4d\beta} \leq 1. \tag{20}$$

On the other hand, in [9, Cor. 4.4], it was proved for the Ising model that

$$\text{Ric}_e \geq 2 \left[1 - (2d-1)(1 - e^{-2\beta})e^{4\beta d} \right] e^{-2\beta d} \quad \text{if} \quad (2d-1)(1 - e^{-2\beta})e^{4\beta d} \leq 1. \tag{21}$$

As observed by Conforti, the condition in (20) is a bit more demanding than the one in (21), but when it is satisfied then the corresponding lower bound for the entropic Ricci curvature is better (for $d \geq 2$).

4.2 Bernoulli-Laplace model

In this subsection we analyze a simplified version of the Bernoulli Laplace model, following Section 5.1 of [5]. Given integers $L > N \in \mathbb{N}$, where L represents the number of sites and N the number of particles, the state space is

$$\Omega = \left\{ \eta \in \{0, 1\}^{[L]} \mid \sum_{i=1}^L \eta_i = N \right\},$$

where $[L] = \{1, \dots, L\}$. Let $\delta_i \in \{0, 1\}^{[L]}$ be defined by $\delta_i(k) = \begin{cases} 1 & \text{if } i = k, \\ 0 & \text{otherwise.} \end{cases}$ Then the set of moves is $G = \{\sigma_{ik} \mid i, k \in [L]\} \cup \{e\}$ where σ_{ik} moves a particle from site i to site k if possible, i.e.

$$\sigma_{ij}(\eta) = \begin{cases} \eta - \delta_i + \delta_j & \text{if } \eta_i(1 - \eta_j) > 0, \\ \eta & \text{otherwise.} \end{cases}$$

The transition rates are given by

$$c(\eta, \sigma_{ij}) = \eta_i(1 - \eta_j)$$

and the reversible measure m is the uniform one on Ω . Finally, notice that we have $\sigma_{ij}^{-1} = \sigma_{ji}$.

Theorem 4.8. *For the Bernoulli Laplace model we have*

$$\mathcal{B}(\rho, \psi) \geq \left(M_\theta + \frac{L}{2} \right) \mathcal{A}(\rho, \psi)$$

for all functions $\rho: \Omega \rightarrow \mathbb{R}_{>0}$ and $\psi: \Omega \rightarrow \mathbb{R}$ and weight functions θ satisfying Assumption 1.

Remark 4.9 (Comparison with [5]). *In [5, Thm. 5.1], under the same assumptions on the model, Conforti establishes inequality (10) and thus CSI $\phi(2K)$ with constant K equal to*

- $\frac{L}{2}$ for general convex ϕ satisfying convexity of (11);
- $\frac{L}{2} + 1$ for $\phi = \phi_1$, corresponding to MLSI($L + 2$);
- $\frac{\alpha L}{2}$ for $\phi = \phi_\alpha$ with $\alpha \in (1, 2]$.

Thus, by the discussion in Section 2, we obtain a stronger results for the case $\theta = \theta_1$ and complementary results for other choices of θ .

Remark 4.10. *The entropic curvature of the Bernoulli Laplace model has been studied before (see [12]), also in the more general case of non homogeneous rates (see [14]). In the homogenous setting, our result for $\theta = \theta_1$ recovers the same (best known) lower bound of [12].*

4.2.1 Proof of Theorem 4.8

Again, the proof is based on Lemma 3.1 and adapts the arguments of [5], from which we use the same coupling rates: for $(\eta, \sigma_{ij}) \in S$ (i.e. $\eta_i = 1, \eta_j = 0$) set

$$c^{\text{pl}}(\eta, \sigma_{ij}\eta, \gamma, \bar{\gamma}) = \begin{cases} \min\{c(\eta, \gamma), c(\sigma_{ij}\eta, \gamma)\} & \text{if } \gamma = \bar{\gamma}, \\ 1 & \text{if } \gamma = \sigma_{ij}, \bar{\gamma} = e \text{ or if } \gamma = e, \bar{\gamma} = \sigma_{ji}, \\ 1 - \eta_l & \text{if } \gamma = \sigma_{il}, \bar{\gamma} = \sigma_{jl} \text{ and } l \notin \{i, j\}, \\ \eta_k & \text{if } \gamma = \sigma_{kj}, \bar{\gamma} = \sigma_{ki} \text{ and } k \notin \{i, j\}, \\ 0 & \text{otherwise.} \end{cases}$$

With these coupling rates, the right hand side of equation (16) from Lemma 3.1 is lower bounded by $\frac{1}{4}(A + B + C + D)$, where we define

$$\begin{aligned}
A &= \sum_{\eta, i, j, k, l} m(\eta) c(\eta, \sigma_{ij}) \min\{c(\eta, \sigma_{kl}), c(\sigma_{ij}\eta, \sigma_{kl})\} I(\eta, \sigma_{ij}, \sigma_{kl}, \sigma_{kl}), \\
B &= \sum_{\eta, i, j} m(\eta) c(\eta, \sigma_{ij}) [J(\eta, \sigma_{ij}, \sigma_{ij}, e) + J(\eta, \sigma_{ij}, e, \sigma_{ji})], \\
C &= \sum_{\eta, i, j} m(\eta) c(\eta, \sigma_{ij}) \left[\sum_{l \neq i, j} (1 - \eta_l) J(\eta, \sigma_{ij}, \sigma_{il}, \sigma_{jl}) \right], \\
D &= \sum_{\eta, i, j} m(\eta) c(\eta, \sigma_{ij}) \left[\sum_{k \neq i, j} \eta_k J(\eta, \sigma_{ij}, \sigma_{kj}, \sigma_{ki}) \right].
\end{aligned}$$

We show that

- $A = 0$,
- $B \geq (4 + 4M_\theta) \mathcal{A}(\rho, \psi)$,
- $C \geq 2(L - N - 1) \mathcal{A}(\rho, \psi)$,
- $D \geq 2(N - 1) \mathcal{A}(\rho, \psi)$,

from which the theorem follows by Lemma 3.1. It is convenient to prove first the following

Lemma 4.11. *For all $\eta \in \Omega$ and $i, j \in [L]$ the following hold:*

1. $c(\eta, \sigma_{ij}) \min\{c(\eta, \sigma_{kl}), c(\sigma_{ij}\eta, \sigma_{kl})\} = \begin{cases} 1 & \text{if } i \neq k, j \neq l, \eta_i = \eta_k = 1, \eta_j = \eta_l = 0, \\ 0 & \text{otherwise.} \end{cases}$
2. $c(\eta, \sigma_{ij}) \min\{c(\eta, \sigma_{kl}), c(\sigma_{ij}\eta, \sigma_{kl})\} = c(\eta, \sigma_{kl}) \min\{c(\eta, \sigma_{ij}), c(\sigma_{kl}\eta, \sigma_{ij})\}$.
3. $I(\eta, \sigma_{ij}, \sigma_{kl}, \sigma_{kl}) = -I(\sigma_{kl}\eta, \sigma_{ij}, \sigma_{lk}, \sigma_{lk})$ whenever $c(\eta, \sigma_{ij}) \min\{c(\eta, \sigma_{kl}), c(\sigma_{ij}\eta, \sigma_{kl})\} > 0$.

Proof of Lemma. Statements 1–2 were already observed by Conforti in the proof of [5, Thm 5.1] and are easy to check, while statement 3 is immediate from the definitions. \square

Term A We have, using Lemma 4.11,

$$\begin{aligned}
A &= \sum_{\eta, i, j, k, l} m(\eta) c(\eta, \sigma_{ij}) \min\{c(\eta, \sigma_{kl}), c(\sigma_{ij}\eta, \sigma_{kl})\} I(\eta, \sigma_{ij}, \sigma_{kl}, \sigma_{kl}) \\
&= \sum_{\eta, i, j, k, l} m(\eta) c(\eta, \sigma_{kl}) \min\{c(\eta, \sigma_{ij}), c(\sigma_{kl}\eta, \sigma_{ij})\} I(\eta, \sigma_{ij}, \sigma_{kl}, \sigma_{kl}) \\
&= \sum_{\eta, k, l} m(\eta) c(\eta, \sigma_{kl}) F(\eta, \sigma_{kl})
\end{aligned}$$

with

$$F(\eta, \sigma_{kl}) = \sum_{i, j} \min\{c(\eta, \sigma_{ij}), c(\sigma_{kl}\eta, \sigma_{ij})\} I(\eta, \sigma_{ij}, \sigma_{kl}, \sigma_{kl}).$$

Using reversibility (6) we deduce

$$\begin{aligned}
A &= \sum_{\eta, k, l} m(\eta) c(\eta, \sigma_{kl}) F(\sigma_{kl}\eta, \sigma_{lk}) \\
&= \sum_{\eta, i, j, k, l} m(\eta) c(\eta, \sigma_{kl}) \min\{c(\eta, \sigma_{ij}), c(\sigma_{kl}\eta, \sigma_{ij})\} I(\sigma_{kl}\eta, \sigma_{ij}, \sigma_{lk}, \sigma_{lk}) \\
&= \sum_{\eta, i, j, k, l} m(\eta) c(\eta, \sigma_{ij}) \min\{c(\eta, \sigma_{kl}), c(\sigma_{ij}\eta, \sigma_{kl})\} I(\sigma_{kl}\eta, \sigma_{ij}, \sigma_{lk}, \sigma_{lk}) \\
&= - \sum_{\eta, i, j, k, l} m(\eta) c(\eta, \sigma_{ij}) \min\{c(\eta, \sigma_{kl}), c(\sigma_{ij}\eta, \sigma_{kl})\} I(\eta, \sigma_{ij}, \sigma_{kl}, \sigma_{kl}) \\
&= -A
\end{aligned}$$

where we used again Lemma 4.11. It follows that $A = 0$.

Term B Notice that for $(\eta, \sigma_{ij}) \in S$

$$\begin{aligned} J(\eta, \sigma_{ij}, \sigma_{ij}, e) &= \{\theta(\rho(\sigma_{ij}\eta), \rho(\sigma_{ij}\eta)) + \theta(\rho(\eta), \rho(\sigma_{ij}\eta))\}[\psi(\eta) - \psi(\sigma_{ij}\eta)]^2 \\ J(\eta, \sigma_{ij}, e, \sigma_{ji}) &= \{\theta(\rho(\eta), \rho(\eta)) + \theta(\rho(\eta), \rho(\sigma_{ij}\eta))\}[\psi(\eta) - \psi(\sigma_{ij}\eta)]^2 \end{aligned}$$

and so

$$J(\eta, \sigma, \sigma, e) + J(\eta, \sigma, e, \sigma) \geq \{2M_\theta + 2\}\theta(\rho(\eta), \rho(\sigma\eta))[\psi(\eta) - \psi(\sigma\eta)]^2.$$

Therefore we get

$$B \geq 4(M_\theta + 1)\mathcal{A}(\rho, \psi).$$

Term C Notice that for $(\eta, \sigma_{ij}) \in S$ we have $\eta_i = 1, \eta_j = 0$ and there are $L - N - 1$ empty sites left. Moreover when $l \neq i, j$ and $\eta_l = 0$ we have

$$J(\eta, \sigma_{ij}, \sigma_{il}, \sigma_{jl}) \geq \theta(\rho(\eta), \rho(\sigma_{ij}\eta))[\psi(\eta) - \psi(\sigma_{ij}\eta)]^2.$$

Therefore we get

$$C \geq 2(L - N - 1)\mathcal{A}(\rho, \psi).$$

Term D Notice that for $(\eta, \sigma_{ij}) \in S$ we have $\eta_i = 1, \eta_j = 0$ and there are $N - 1$ other occupied sites. Moreover when $k \neq i, j$ and $\eta_k = 1$ we have

$$J(\eta, \sigma_{ij}, \sigma_{kj}, \sigma_{ki}) \geq \theta(\rho(\eta), \rho(\sigma_{ij}\eta))[\psi(\eta) - \psi(\sigma_{ij}\eta)]^2.$$

Therefore we get

$$D \geq 2(N - 1)\mathcal{A}(\rho, \psi).$$

Combining these results, an application of Lemma 3.1 gives

$$\mathcal{B}(\rho, \psi) \geq \left(2 + \frac{L}{2} - 1\right)\mathcal{A}(\rho, \psi) = \left(1 + \frac{L}{2}\right)\mathcal{A}(\rho, \psi).$$

This concludes the proof of the theorem.

4.3 Hardcore model

Following Section 5.2 of [5], we consider the classical hardcore model. Let (V, E) be a simple, finite, connected graph and write $x \sim y$ if the vertices x, y are connected (in the rest of this subsection, x, y always denote general elements of V). The state space is

$$\Omega = \left\{ \eta \in \{0, 1\}^V \mid \eta_x \eta_y = 0 \text{ if } x \sim y \right\}.$$

In other words, each vertex can either be empty or be occupied by a particle, with the rule that if a site is occupied then its neighbors are all free. Let $N_x = \{y \in V \mid x \sim y\}$ be the set of neighbors of vertex x , $\bar{N}_x = N_x \cup \{x\}$

and as before let $\delta_x \in \{0, 1\}^V$ be defined by $\delta_x(y) = \begin{cases} 1 & \text{if } x = y, \\ 0 & \text{otherwise.} \end{cases}$ Then the set of moves is given by

$G = \{\gamma_x^+, \gamma_x^- \mid i, k \in [L]\} \cup \{e\}$ where γ_x^+ adds a particle to site x if possible and γ_x^- removes it if possible, i.e.

$$\begin{aligned} \gamma_x^+(\eta) &= \begin{cases} \eta + \delta_x & \text{if } \eta + \delta_x \in \Omega, \\ \eta & \text{otherwise,} \end{cases} \\ \gamma_x^-(\eta) &= \begin{cases} \eta - \delta_x & \text{if } \eta - \delta_x \in \Omega, \\ \eta & \text{otherwise.} \end{cases} \end{aligned}$$

We also denote $G^+ = \{\gamma_x^+ \mid x \in V\}$ and $G^- = \{\gamma_x^- \mid x \in V\}$. For a given parameter $\beta \in (0, 1)$ the transition rates are given by

$$\begin{aligned} c(\eta, \gamma_x^+) &= \beta \prod_{y \in \bar{N}_x} (1 - \eta_y), \\ c(\eta, \gamma_x^-) &= \eta_x \end{aligned}$$

for all $\eta \in \Omega, x \in V$. With these choices, we have $(\gamma_x^+)^{-1} = \gamma_x^-$ and the reversible measure is given by

$$m(\eta) = \frac{1}{Z} \mathbb{1}_{\eta \in \Omega} \prod_{x \in V} \beta^{\eta_x},$$

where $Z > 0$ is the normalization constant.

Theorem 4.12. *Let Δ be the maximum degree of (V, E) and assume that*

$$\beta \Delta \leq 1. \tag{22}$$

Set

$$\kappa_* = 1 - \beta(\Delta - 1), \quad \bar{\kappa}_* = \min\{\beta, 1 - \beta\Delta\}.$$

Then

$$\mathcal{B}(\rho, \psi) \geq \left(\frac{\kappa_*}{2} + M_\theta \bar{\kappa}_* \right) \mathcal{A}(\rho, \psi)$$

for all functions $\rho: \Omega \rightarrow \mathbb{R}_{>0}$ and $\psi: \Omega \rightarrow \mathbb{R}$ and weight functions θ satisfying Assumption 1.

Remark 4.13 (Comparison with [5]). *In [5, Thm. 5.2], under the same assumptions on the model, Conforti establishes inequality (10) and thus CSI $\phi(2K)$ with constant K equal to*

- $\frac{\kappa_*}{2}$ for general convex ϕ satisfying convexity of (11);
- $\frac{\kappa_*}{2} + \bar{\kappa}_*$ for $\phi = \phi_1$, corresponding to $\text{MLSI}(\kappa_* + 2\bar{\kappa}_*)$;
- $\frac{\alpha\kappa_*}{2}$ for $\phi = \phi_\alpha$ with $\alpha \in (1, 2]$.

Thus, by the discussion in Section 2, we obtain a stronger results for the case $\theta = \theta_1$ and complementary results for other choices of θ .

Remark 4.14. *The entropic curvature of the hardcore model has been studied before in [9], where a more general version of the model is considered. When restricting to the classical version discussed in this section, it was proved in [9, Cor. 4.8] that*

$$\text{Ric}_e \geq \frac{\kappa_*}{2}$$

under condition (22). Therefore, in this setting, by choosing $\theta = \theta_1$ in Theorem 4.12 we find a better lower bound for the entropic curvature.

4.3.1 Proof of Theorem 4.12

The proof is based on Lemma 3.1 and on the arguments in [5], from which we use the same coupling rates. For $(\eta, \gamma_x^+) \in S$ (i.e. $\eta|_{\bar{N}_x} = 0$) we set

$$c^{\text{cpl}}(\eta, \gamma_x^+ \eta, \gamma, \bar{\gamma}) = \begin{cases} \min\{c(\eta, \gamma), c(\gamma_x^+ \eta, \gamma)\} & \text{if } \gamma = \bar{\gamma}, \\ \beta & \text{if } \gamma = \gamma_y^+, \bar{\gamma} = \gamma_x^- \text{ with } y \sim x, \eta|_{\bar{N}_y} = 0, \\ \beta & \text{if } \gamma = \gamma_x^+, \bar{\gamma} = e, \\ 1 - \beta \left| \left\{ y : y \sim x, \eta|_{\bar{N}_y} = 0 \right\} \right| & \text{if } \gamma = e, \bar{\gamma} = \gamma_x^-, \\ 0 & \text{otherwise.} \end{cases}$$

If $(\eta, \gamma_x^-) \in S$ then also $(\gamma_x^- \eta, \gamma_x^+) \in S$, and so we can set

$$c^{\text{cpl}}(\eta, \gamma_x^- \eta, \gamma, \bar{\gamma}) = c^{\text{cpl}}(\gamma_x^- \eta, \gamma_x^+ \gamma_x^- \eta, \bar{\gamma}, \gamma).$$

With these coupling rates the right hand side of equation (16) from Lemma 3.1 reads

$$\begin{aligned} & \frac{1}{4} \sum_{(\eta, \gamma_x^+) \in S} \sum_{\gamma, \bar{\gamma}} m(\eta) c(\eta, \gamma_x^+) c^{\text{cpl}}(\eta, \gamma_x^+ \eta, \gamma, \bar{\gamma}) J(\eta, \gamma_x^+, \gamma, \bar{\gamma}) \\ & + \frac{1}{4} \sum_{(\eta, \gamma_x^-) \in S} \sum_{\gamma, \bar{\gamma}} m(\eta) c(\eta, \gamma_x^-) c^{\text{cpl}}(\gamma_x^- \eta, \eta, \bar{\gamma}, \gamma) J(\eta, \gamma_x^-, \gamma, \bar{\gamma}). \end{aligned}$$

Using reversibility (6) with

$$F(\eta, \sigma) = \mathbb{1}_{G^-}(\sigma) \sum_{\gamma, \bar{\gamma}} c^{\text{cpl}}(\sigma \eta, \eta, \bar{\gamma}, \gamma) J(\eta, \sigma, \gamma, \bar{\gamma})$$

and that $J(\eta, \sigma \eta, \gamma, \bar{\gamma}) = J(\sigma \eta, \sigma^{-1}, \bar{\gamma}, \gamma)$ (when $\sigma^{-1} \sigma \eta = \eta$) the second term is equal to the first, so we can rewrite the previous quantity as

$$\frac{1}{2} \sum_{(\eta, \gamma_x^+) \in S} \sum_{\gamma, \bar{\gamma}} m(\eta) c(\eta, \gamma_x^+) c^{\text{cpl}}(\eta, \gamma_x^+ \eta, \gamma, \bar{\gamma}) J(\eta, \gamma_x^+, \gamma, \bar{\gamma}). \quad (23)$$

Similarly we have

$$\begin{aligned} \mathcal{A}(\rho, \psi) &= \frac{1}{2} \sum_{(\eta, \gamma_x^+) \in S} m(\eta) c(\eta, \gamma_x^+) [\psi(\eta) - \psi(\gamma_x^+ \eta)]^2 \\ &+ \frac{1}{2} \sum_{(\eta, \gamma_x^-) \in S} m(\eta) c(\eta, \gamma_x^-) [\psi(\eta) - \psi(\gamma_x^- \eta)]^2 \end{aligned}$$

and using again reversibility the second term is equal to the first, so we can write

$$\mathcal{A}(\rho, \psi) = \sum_{(\eta, \gamma_x^+) \in S} m(\eta) c(\eta, \gamma_x^+) [\psi(\eta) - \psi(\gamma_x^+ \eta)]^2.$$

We then have that (using that $\forall \eta, x, y$ $c(\eta, \gamma_y^-) \leq c(\gamma_x^+ \eta, \gamma_y^-)$ and $c(\eta, \gamma_y^+) \geq c(\gamma_x^+ \eta, \gamma_y^+)$) the quantity (23) (and in particular $B(\rho, \psi)$ too) is lower bounded by $\frac{1}{2}(A + B + C)$ with

$$\begin{aligned} A &= \sum_{\eta, x, y} m(\eta) c(\eta, \gamma_x^+) c(\eta, \gamma_y^-) I(\eta, \gamma_x^+, \gamma_y^-, \gamma_y^-) \\ &+ \sum_{\eta, x, y} m(\eta) c(\eta, \gamma_x^+) c(\gamma_x^+ \eta, \gamma_y^+) I(\eta, \gamma_x^+, \gamma_y^+, \gamma_y^+), \\ B &= \sum_{\substack{\eta, x, y: \\ x \sim y, \eta|_{\bar{N}_y} = 0}} m(\eta) c(\eta, \gamma_x^+) I(\eta, \gamma_x^+, \gamma_y^+, \gamma_x^-), \\ C &= \sum_{\eta, x} m(\eta) c(\eta, \gamma_x^+) \left[\left(1 - \beta \left| \left\{ y : y \sim x, \eta|_{\bar{N}_y} = 0 \right\} \right| \right) J(\eta, \gamma_x^+, e, \gamma_x^-) + \beta J(\eta, \gamma_x^+, \gamma_x^+, e) \right]. \end{aligned}$$

We will show that

- $A = 0$,
- $B = 0$,
- $C \geq (\kappa_* + 2M_\theta \bar{\kappa}_*) \mathcal{A}(\rho, \psi)$.

An application of Lemma 3.1 then concludes the proof of the theorem. To do so, we will use the following:

Lemma 4.15. *For all $\eta \in \Omega$ and $x, y \in V$ the following hold:*

1. $c(\eta, \gamma_x^+) c(\gamma_x^+ \eta, \gamma_y^+) = c(\eta, \gamma_x^+) c(\gamma_x^+ \eta, \gamma_y^+) = \begin{cases} \beta^2 & \text{if } x \asymp y, \eta|_{\bar{N}_x \cup \bar{N}_y} = 0, \\ 0 & \text{otherwise.} \end{cases}$

2. $x \approx y, \eta|_{\bar{N}_x \cup \bar{N}_y} = 0 \implies \gamma_x^+ \gamma_y^+ \eta = \gamma_y^+ \gamma_x^+ \eta$.
3. If $\eta|_{\bar{N}_x \cup \bar{N}_y} = 0$ then $c(\eta, \gamma_x^+) = c(\eta, \gamma_y^+) = \beta$.
4. $I(\gamma_y^+ \eta, \gamma_x^+, \gamma_y^-, \gamma_x^-) = -I(\eta, \gamma_x^+, \gamma_y^+, \gamma_x^-)$ if $x \approx y, \eta|_{\bar{N}_x \cup \bar{N}_y} = 0$.
5. $I(\eta, \gamma_x^+, \gamma_y^+, \gamma_x^-) = -I(\eta, \gamma_y^+, \gamma_x^+, \gamma_y^-)$ if $\eta|_{\bar{N}_x \cup \bar{N}_y} = 0$.

Proof of Lemma. Statements 1–3 were already in the proof of [5, Thm. 5.2] and are easy to check, while statement 4–5 are immediate from the definitions. \square

Term A We look at the first term in the sum defining A : we can write it as

$$\sum_{\eta, \gamma} m(\eta) c(\eta, \gamma) F(\eta, \gamma)$$

with

$$F(\eta, \gamma) = \mathbb{1}_{G^-}(\gamma) \sum_x c(\eta, \gamma_x^+) I(\eta, \gamma_x^+, \gamma, \gamma).$$

Using reversibility (6) we can rewrite it as

$$\sum_{\eta, x, y} m(\eta) c(\eta, \gamma_y^+) c(\gamma_y^+ \eta, \gamma_x^+) I(\gamma_y^+ \eta, \gamma_x^+, \gamma_y^-, \gamma_x^-).$$

Then we have

$$\begin{aligned} & \sum_{\eta, x, y} m(\eta) c(\eta, \gamma_y^+) c(\gamma_y^+ \eta, \gamma_x^+) I(\gamma_y^+ \eta, \gamma_x^+, \gamma_y^-, \gamma_x^-) \\ &= \sum_{\eta, x, y} m(\eta) c(\eta, \gamma_x^+) c(\gamma_x^+ \eta, \gamma_y^+) I(\gamma_y^+ \eta, \gamma_x^+, \gamma_y^-, \gamma_x^-) \\ &= - \sum_{\eta, x, y} m(\eta) c(\eta, \gamma_x^+) c(\gamma_x^+ \eta, \gamma_y^+) I(\eta, \gamma_x^+, \gamma_y^+, \gamma_x^-) \end{aligned}$$

using Lemma 4.15. Therefore, the first term in the sum of A is the opposite of the second, which implies $A = 0$.

Term B We have

$$B = \sum_{\substack{\eta, x, y: \\ x \sim y, \eta|_{\bar{N}_y} = 0}} m(\eta) c(\eta, \gamma_x^+) I(\eta, \gamma_x^+, \gamma_y^+, \gamma_x^-).$$

Noticing as in [5] that $(\eta, \gamma_x^+) \in S$ if and only if $\eta|_{\bar{N}_x} = 0$ we can write

$$B = \sum_{\substack{\eta, x, y: \\ x \sim y, \eta|_{\bar{N}_y} = 0, \eta|_{\bar{N}_x} = 0}} m(\eta) c(\eta, \gamma_x^+) I(\eta, \gamma_x^+, \gamma_y^+, \gamma_x^-). \quad (24)$$

By exchanging x, y we therefore also have

$$B = \sum_{\substack{\eta, x, y: \\ x \sim y, \eta|_{\bar{N}_y} = 0, \eta|_{\bar{N}_x} = 0}} m(\eta) c(\eta, \gamma_y^+) I(\eta, \gamma_y^+, \gamma_x^+, \gamma_y^-). \quad (25)$$

Summing these two expressions we get

$$\begin{aligned} 2B &= \sum_{\substack{\eta, x, y: \\ x \sim y, \eta|_{\bar{N}_y} = 0, \eta|_{\bar{N}_x} = 0}} m(\eta) [c(\eta, \gamma_x^+) I(\eta, \gamma_x^+, \gamma_y^+, \gamma_x^-) + c(\eta, \gamma_y^+) I(\eta, \gamma_y^+, \gamma_x^+, \gamma_y^-)] \\ &= \sum_{\substack{\eta, x, y: \\ x \sim y, \eta|_{\bar{N}_y} = 0, \eta|_{\bar{N}_x} = 0}} m(\eta) [c(\eta, \gamma_x^+) - c(\eta, \gamma_y^+)] I(\eta, \gamma_x^+, \gamma_y^+, \gamma_x^-) \\ &= 0, \end{aligned}$$

where we used Lemma 4.15.

Term C We have

$$\begin{aligned}
C &= \sum_{\eta, x} m(\eta) c(\eta, \gamma_x^+) \left[\left(1 - \beta \left| \left\{ y : y \sim x, \eta|_{\bar{N}_y} = 0 \right\} \right| \right) J(\eta, \gamma_x^+, e, \gamma_x^-) + \beta J(\eta, \gamma_x^+, \gamma_x^+, e) \right] \\
&= \sum_{\eta, x} m(\eta) c(\eta, \gamma_x^+) [\psi(\eta) - \psi(\gamma_x^+ \eta)]^2 \\
&\quad \cdot \left[\left(1 - \beta \left| \left\{ y : y \sim x, \eta|_{\bar{N}_y} = 0 \right\} \right| \right) (\theta(\rho(\eta), \rho(\eta)) + \theta(\rho(\eta), \rho(\gamma_x^+ \eta))) + \beta (\theta(\rho(\gamma_x^+ \eta), \rho(\gamma_x^+ \eta)) + \theta(\rho(\eta), \rho(\gamma_x^+ \eta))) \right] \\
&\geq \sum_{\eta, x} m(\eta) c(\eta, \gamma_x^+) [\psi(\eta) - \psi(\gamma_x^+ \eta)]^2 (\kappa_* + 2M_\theta \bar{\kappa}_*) \\
&= (\kappa_* + 2M_\theta \bar{\kappa}_*) \mathcal{A}(\rho, \psi).
\end{aligned}$$

This concludes the proof of the theorem.

4.4 Interacting random walks

Following Section 3 of [5], we consider now the case of interacting random walks. One motivation in this subsection is to find a discrete analogue of the following classical result.

Proposition 4.16. *Consider \mathbb{R}^d equipped with the standard Euclidean distance d . Let $V : \mathbb{R}^d \rightarrow \mathbb{R}$ be **convex**, γ_d be the law of a standard Gaussian in \mathbb{R}^d and $Z > 0$ be a normalization constant so that $\frac{1}{Z} e^{-V} d\gamma_d$ is a probability measure. Then the metric measure space $(\mathbb{R}, d, \frac{1}{Z} e^{-V} d\gamma_d)$ has Ricci curvature $\text{Ric} \geq 1$ in the sense of the Lott–Sturm–Villani theory.*

To find a discrete analogue, here the role of \mathbb{R}^d is taken over by the discrete state space $\Omega = \mathbb{N}^d$, while the Gaussian measure γ_d is replaced by the multivariate Poisson distribution μ_λ given by the product measure of d one-dimensional Poisson distribution of intensity $\frac{1}{\lambda}$, i.e. for $\eta \in \Omega$ we have

$$\mu_\lambda(\eta) = \prod_{i=1}^d e^{-\frac{1}{\lambda}} \frac{\lambda^{-\eta_i}}{\eta_i!}.$$

It remains to define a Markov chain dynamic on this state space. We consider the set of moves G containing $e, \gamma_i^+, \gamma_i^-$ for $i \in [d]$, where

$$\begin{aligned}
\gamma_i^+ \eta &= \eta + \mathbf{e}_i, \\
\gamma_i^- \eta &= \eta - \mathbf{e}_i \mathbb{1}_{\eta_i > 0},
\end{aligned}$$

and e is the null move as usual. Consider two potentials $V^+, V^- : \mathbb{N}^d \rightarrow \mathbb{R}$ and correspondingly define transition rates

$$\begin{aligned}
c(\eta, \gamma_i^+) &= \exp(-\nabla_i^+ V^+(\eta)), \\
c(\eta, \gamma_i^-) &= \begin{cases} \exp(-\nabla_i^- V^-(\eta)) & \text{if } \eta_i > 0, \\ 0 & \text{if } \eta_i = 0, \end{cases}
\end{aligned}$$

where $\nabla_i^\pm = \nabla_{\gamma_i^\pm}$. Then the reversible measure takes the form

$$m = \frac{1}{Z} \exp(-V^+ - V^-) d\mu_\lambda,$$

where Z is normalization constant. An interesting choice is

$$V^-(\eta) = \sum_{i=1}^d \log(\lambda) \eta_i + \log(\eta_i!), \quad (26)$$

which corresponds to $c(\eta, \gamma_i^-) = \lambda \eta_i$. In this case, we write $V = V^+$ and the reversible measure becomes

$$\frac{1}{Z} e^{-V} d\mu_\lambda,$$

which resembles the setting of Proposition 4.16. Therefore, to find an analogue discrete result, we are left to look for conditions on the potential V that resemble convexity and yield positive entropic curvature of the corresponding Markov chain; we will do this in Section 4.4.2, as an application of the main theorem of this section below.

At this point, we make the technical assumption that for all $\eta \in \Omega, i \in [d]$ the following quantities are non-negative

$$\begin{aligned}\kappa^+(\eta, i) &:= -\nabla_i^+ c(\eta, \gamma_i^+) - \sum_{\bar{\gamma} \in G, \bar{\gamma} \neq \gamma_i^+, \gamma_i^-} \max\{\nabla_i^+ c(\eta, \bar{\gamma}), 0\} \geq 0, \\ \kappa^-(\eta, i) &:= \nabla_i^+ c(\eta, \gamma_i^-) - \sum_{\gamma \in G, \gamma \neq \gamma_i^+, \gamma_i^-} \max\{-\nabla_i^+ c(\eta, \gamma), 0\} \geq 0.\end{aligned}$$

Correspondingly we set

$$\kappa_* = \inf_{\eta \in \mathbb{N}^d, i \in [d]} \kappa^+(\eta, i) + \kappa^-(\eta, i).$$

It is natural to introduce the additional quantity

$$\overline{\kappa_*} = \min\left\{\inf_{\eta, i} \kappa^+(\eta, i), \inf_{\eta, i} \kappa^-(\eta, i)\right\}.$$

Finally, compared to [5], we make the *additional assumption* that for all η, i, j we have

$$\nabla_i^+ c(\eta, \gamma_j^+) \leq 0; \tag{27}$$

$$\nabla_i^+ c(\eta, \gamma_j^-) \geq 0. \tag{28}$$

Notice in the case $c(\eta, \gamma_j^-) = \lambda \eta_j$ (corresponding to V^- as in (26)) the second additional assumption (28) is always satisfied.

It is important to note that this is the only example that we discuss where the cardinality of the state space is not finite and that, because of this, there are some additional technical difficulties and not all the considerations of Section 2 can be directly applied here. In this paper, to deal with the infinite cardinality of the state space, we proceed as in [5] and make use of a localization argument, which we now briefly describe; more precisely, with this procedure and with ϕ and θ satisfying Assumption 2, we explain how to derive inequality (3) from establishing (8) for a localizing sequence of finite state space Markov chains. Let $\Omega_N = \{\eta \in \Omega \mid \eta_i \leq N \forall i \in [d]\}$, and on Ω_N consider the Markov chain with moves $\gamma_i^{+,N}(\eta) = \eta + e_i \mathbb{1}_{\eta_i < N}$ and $\gamma_i^{-,N}(\eta) = \eta - e_i \mathbb{1}_{\eta_i > 0}$, rates $c^N(\eta, \gamma_i^{\pm, N}) = c(\eta, \gamma_i^{\pm})$ and reversible measure $m_N = \frac{m}{m(\Omega_N)}$. Finally, denote by \mathcal{B}_N and \mathcal{A}_N the corresponding quantities \mathcal{B} and \mathcal{A} for this Markov chain. The main theorem of this section reads as follows.

Theorem 4.17. *With the previous notation, suppose that for all $\eta \in \Omega$ and $i, j \in [d]$ we have*

$$\begin{aligned}\kappa^+(\eta, i) &\geq 0, \\ \kappa^-(\eta, i) &\geq 0, \\ \nabla_i^+ c(\eta, \gamma_j^+) &\leq 0, \\ \nabla_i^+ c(\eta, \gamma_j^-) &\geq 0.\end{aligned}$$

Let also θ be a weight function satisfying Assumption 1. Then we have that

$$\mathcal{B}_N(\rho, \psi) \geq \left(\frac{\kappa_*}{2} + M_\theta \overline{\kappa_*}\right) \mathcal{A}_N(\rho, \psi)$$

for all $N \geq 2$, for all functions $\rho: \Omega_N \rightarrow \mathbb{R}_{>0}$ and $\psi: \Omega_N \rightarrow \mathbb{R}$.

Remark 4.18. *With θ as in Assumption 2, by the discussion of Section 2 the previous theorem allows to deduce the convex Sobolev inequality (3) for all Markov chains (Ω_n, L_n, m_n) with uniform constant. As observed in [5], this allows to deduce the same convex Sobolev inequality for the original Markov chain on by taking limits, when ϕ is lower bounded (as it is the case for $\phi = \phi_\alpha$ with $\alpha \in [1, 2]$ in particular), cf. Corollary 2.1 of [5].*

Remark 4.19 (Comparison with [5]). *In [5, Thm. 3.1], Conforti establishes inequality (10) for the localizing sequence of Markov chain on Ω_N with a uniform constant K , and thus also $\text{CSI}\phi(2K)$ for the original Markov chain, with constant K equal to*

- $\frac{\kappa_*}{2}$ for general convex ϕ satisfying convexity of (11);
- $\frac{\alpha\kappa_*}{2}$ for $\phi = \phi_\alpha$ with $\alpha \in (1, 2]$.

Compared to our assumptions on the model, he does not assume non negativity in (27), (28); however, these additional assumptions are satisfied in the examples of Section 4.4.2.

By the discussion in Section 2, we therefore obtain a complementary result to [5, Thm. 3.1].

4.4.1 Proof of Theorem 4.17

Fix N : since there is no ambiguity, we drop the sub(/sup)scripts N . We apply Lemma 3.1 and so to prove the theorem it is enough to show that

$$\frac{1}{4} \sum_{\eta, \sigma} \sum_{\gamma, \bar{\gamma}} m(\eta) c(\eta, \sigma) c^{\text{cpl}}(\eta, \sigma \eta, \gamma, \bar{\gamma}) J(\eta, \sigma, \gamma, \bar{\gamma}) \geq K \mathcal{A}(\rho, \psi), \quad (29)$$

where the sum is for $\eta \in \Omega^N$, $\sigma, \gamma, \bar{\gamma} \in G$. We define the same coupling rates as in [5], namely for $(\eta, \gamma_i^+) \in S$

$$c^{\text{cpl}}(\eta, \gamma_i^+ \eta, \gamma, \bar{\gamma}) = \begin{cases} \min\{c(\eta, \gamma), c(\gamma_i^+ \eta, \gamma)\} & \text{if } \gamma = \bar{\gamma} \\ \max\{\nabla_i^+ c(\eta, \bar{\gamma}), 0\} & \text{if } \gamma = \gamma_i^+ \text{ and } \bar{\gamma} \neq \gamma_i^+, \gamma_i^- \\ \max\{-\nabla_i^+ c(\eta, \gamma), 0\} & \text{if } \gamma \neq \gamma_i^+, \gamma_i^- \text{ and } \bar{\gamma} = \gamma_i^- \\ \kappa^+(\eta, i) & \text{if } \gamma = \gamma_i^+, \bar{\gamma} = e, \\ \kappa^-(\eta, i) & \text{if } \gamma = e, \bar{\gamma} = \gamma_i^-, \\ 0 & \text{otherwise.} \end{cases}$$

Correspondingly, for $(\eta, \gamma_i^-) \in S$ we set also $c^{\text{cpl}}(\eta, \gamma_i^-, \gamma, \bar{\gamma}) = c^{\text{cpl}}(\gamma_i^- \eta, \gamma_i^+ \gamma_i^- \eta, \bar{\gamma}, \gamma)$.

The left hand side of equation (29) reads (summing for $i \in [d]$)

$$\begin{aligned} & \frac{1}{4} \sum_{\eta, i} \sum_{\gamma, \bar{\gamma}} m(\eta) c(\eta, \gamma_i^+) c^{\text{cpl}}(\eta, \gamma_i^+ \eta, \gamma, \bar{\gamma}) J(\eta, \gamma_i^+ \eta, \gamma, \bar{\gamma}) \\ & + \frac{1}{4} \sum_{\eta, i} \sum_{\gamma, \bar{\gamma}} m(\eta) c(\eta, \gamma_i^-) c^{\text{cpl}}(\gamma_i^- \eta, \eta, \bar{\gamma}, \gamma) J(\eta, \gamma_i^- \eta, \gamma, \bar{\gamma}). \end{aligned}$$

Using reversibility (6) and that $J(\eta, \sigma, \gamma, \bar{\gamma}) = J(\sigma \eta, \sigma^{-1}, \bar{\gamma}, \gamma)$ (when $\sigma^{-1} \sigma \eta = \eta$) we get that the second summand is equal to the first and so we can rewrite our quantity as

$$\frac{1}{2} \sum_{\eta, i} \sum_{\gamma, \bar{\gamma}} m(\eta) c(\eta, \gamma_i^+) c^{\text{cpl}}(\eta, \gamma_i^+ \eta, \gamma, \bar{\gamma}) J(\eta, \gamma_i^+ \eta, \gamma, \bar{\gamma}).$$

With this explicit choice of coupling rates the left hand side of (29) becomes $\frac{1}{2}(\tilde{A} + \tilde{B} + \tilde{C} + \tilde{D})$, where

$$\begin{aligned} \tilde{A} &= \sum_{\eta, i, \gamma} m(\eta) c(\eta, \gamma_i^+) \min\{c(\eta, \gamma), c(\gamma_i^+ \eta, \gamma)\} J(\eta, \gamma_i^+, \gamma, \gamma), \\ \tilde{B} &= \sum_{\eta, i} \sum_{\gamma \neq \gamma_i^\pm} m(\eta) c(\eta, \gamma_i^+) \max\{\nabla_i^+ c(\eta, \gamma), 0\} J(\eta, \gamma_i^+, \gamma_i^+, \gamma), \\ \tilde{C} &= \sum_{\eta, i} \sum_{\gamma \neq \gamma_i^\pm} m(\eta) c(\eta, \gamma_i^+) \max\{-\nabla_i^+ c(\eta, \gamma), 0\} J(\eta, \gamma_i^+, \gamma, \gamma_i^-), \\ \tilde{D} &= \sum_{\eta, i} m(\eta) c(\eta, \gamma_i^+) [\kappa^+(\eta, i) J(\eta, \gamma_i^+, \gamma_i^+, e) + \kappa^-(\eta, i) J(\eta, \gamma_i^+, e, \gamma_i^-)]. \end{aligned}$$

This is then lower bounded (since $J \geq I$ by (17)) by $\frac{1}{2}(A + B + C + D)$ where

$$\begin{aligned}
A &= \sum_{\eta, i, \gamma} m(\eta) c(\eta, \gamma_i^+) \min\{c(\eta, \gamma), c(\gamma_i^+ \eta, \gamma)\} I(\eta, \gamma_i^+, \gamma, \gamma), \\
B &= \sum_{\eta, i} \sum_{\gamma \neq \gamma_i^\pm} m(\eta) c(\eta, \gamma_i^+) \max\{\nabla_i^+ c(\eta, \gamma), 0\} I(\eta, \gamma_i^+, \gamma_i^+, \gamma), \\
C &= \sum_{\eta, i} \sum_{\gamma \neq \gamma_i^\pm} m(\eta) c(\eta, \gamma_i^+) \max\{-\nabla_i^+ c(\eta, \gamma), 0\} I(\eta, \gamma_i^+, \gamma, \gamma_i^-), \\
D &= \sum_{\eta, i} m(\eta) c(\eta, \gamma_i^+) [\kappa^+(\eta, i) J(\eta, \gamma_i^+, \gamma_i^+, e) + \kappa^-(\eta, i) J(\eta, \gamma_i^+, e, \gamma_i^-)].
\end{aligned}$$

We also have

$$\begin{aligned}
\mathcal{A}(\rho, \psi) &= \frac{1}{2} \sum_{\eta, i} m(\eta) c(\eta, \gamma_i^+) \theta(\rho(\eta), \rho(\gamma_i^+ \eta)) [\psi(\eta) - \psi(\gamma_i^+ \eta)]^2 \\
&\quad + \frac{1}{2} \sum_{\eta, i} m(\eta) c(\eta, \gamma_i^-) \theta(\rho(\eta), \rho(\gamma_i^- \eta)) [\psi(\eta) - \psi(\gamma_i^- \eta)]^2 \\
&= \sum_{\eta, i} m(\eta) c(\eta, \gamma_i^+) \theta(\rho(\eta), \rho(\gamma_i^+ \eta)) [\psi(\eta) - \psi(\gamma_i^+ \eta)]^2
\end{aligned}$$

using reversibility again.

Term D We have

$$J(\eta, \gamma_i^+, \gamma_i^+, e) = (\theta(\rho(\gamma_i^+ \eta), \rho(\gamma_i^+ \eta)) + \theta(\rho(\eta), \rho(\gamma_i^+ \eta))) [\psi(\eta) - \psi(\gamma_i^+ \eta)]^2$$

and

$$J(\eta, \gamma_i^+, e, \gamma_i^-) = (\theta(\rho(\eta), \rho(\eta)) + \theta(\rho(\eta), \rho(\gamma_i^+ \eta))) [\psi(\eta) - \psi(\gamma_i^+ \eta)]^2$$

and so

$$\begin{aligned}
&\kappa^+(\eta, i) J(\eta, \gamma_i^+, \gamma_i^+, e) + \kappa^-(\eta, i) J(\eta, \gamma_i^+, e, \gamma_i^-) \\
&\geq \{\overline{\kappa_*} [\theta(\rho(\gamma_i^+ \eta), \rho(\gamma_i^+ \eta)) + \theta(\rho(\eta), \rho(\eta))] + \kappa_* \theta(\rho(\eta), \rho(\gamma_i^+ \eta))\} [\psi(\eta) - \psi(\gamma_i^+ \eta)]^2 \\
&\geq (2M_\theta \overline{\kappa_*} + \kappa_*) \theta(\rho(\eta), \rho(\gamma_i^+ \eta)) [\psi(\eta) - \psi(\gamma_i^+ \eta)]^2.
\end{aligned}$$

Therefore we get

$$D \geq (2M_\theta \overline{\kappa_*} + \kappa_*) \mathcal{A}(\rho, \psi).$$

Other terms We now show that each one of the other terms is 0, concluding the proof of the theorem. Using our additional assumptions we have

$$\begin{aligned}
A &= \sum_{\eta, i, j} m(\eta) c(\eta, \gamma_i^+) c(\gamma_i^+ \eta, \gamma_j^+) I(\eta, \gamma_i^+, \gamma_j^+, \gamma_j^+) \\
&\quad + \sum_{\eta, i, j} m(\eta) c(\eta, \gamma_i^+) c(\eta, \gamma_j^-) I(\eta, \gamma_i^+, \gamma_j^-, \gamma_j^-), \\
B &= \sum_{\eta, i, j \neq i} m(\eta) c(\eta, \gamma_i^+) \nabla_i^+ c(\eta, \gamma_j^-) I(\eta, \gamma_i^+, \gamma_i^+, \gamma_j^-), \\
C &= \sum_{\eta, i, j \neq i} m(\eta) c(\eta, \gamma_i^+) \{-\nabla_i^+ c(\eta, \gamma_j^+)\} I(\eta, \gamma_i^+, \gamma_j^+, \gamma_i^-).
\end{aligned}$$

To show $A = B = C = 0$ we proceed similarly to [5]. Again it is useful to have an auxiliary lemma.

Lemma 4.20. *For all $\eta \in \mathbb{N}^d$ and $i, j \in [d]$*

1. $c(\eta, \gamma_j^+) c(\gamma_j^+ \eta, \gamma_i^+) = c(\eta, \gamma_i^+) c(\gamma_i^+ \eta, \gamma_j^+)$.

2. $c(\eta, \gamma_i^-) \nabla_i^- c(\eta, \gamma_j^-) = c(\eta, \gamma_j^-) \nabla_j^- c(\eta, \gamma_i^-)$.
3. $c(\eta, \gamma_i^+) \nabla_i^+ c(\eta, \gamma_j^+) = c(\eta, \gamma_j^+) \nabla_j^+ c(\eta, \gamma_i^+)$.
4. $I(\gamma_j^+ \eta, \gamma_i^+, \gamma_j^-, \gamma_i^-) = -I(\eta, \gamma_i^+, \gamma_j^+, \gamma_i^-)$.
5. $I(\gamma_i^- \eta, \gamma_i^+, \gamma_j^+, \gamma_i^-) = -I(\gamma_j^- \eta, \gamma_j^+, \gamma_i^+, \gamma_i^-)$ if $\eta_i, \eta_j > 0$.
6. $I(\eta, \gamma_i^+, \gamma_j^+, \gamma_i^-) = -I(\eta, \gamma_j^+, \gamma_i^+, \gamma_j^-)$.

Remark 4.21. In general the quantities

- $I(\eta, \gamma_i^+, \gamma_i^+, \gamma_j^+) - I(\eta, \gamma_i^+, \gamma_j^+, \gamma_i^+)$,
- $I(\gamma_i^- \eta, \gamma_i^+, \gamma_j^-, \gamma_i^-) - I(\gamma_i^- \eta, \gamma_i^+, \gamma_j^-, \gamma_i^-)$

are not antisymmetric in i and j and so we could not get all the desired cancellations in the more general setting of [5]: to overcome this issue, we introduced the additional assumptions (27) and (28).

Proof of Lemma. Statements 1–3 were already observed in the proof of [5, Thm. 3.1] and are easy to check, while statements 4–6 are immediate from the definitions and the assumptions. \square

Term A Use reversibility in the second summand with

$$F(\eta, \sigma) = 1_{G^-}(\sigma) \sum_i c(\eta, \gamma_i^+) I(\eta, \gamma_i^+, \sigma, \sigma).$$

This way we get

$$\begin{aligned} & \sum_{\eta, i, j} m(\eta) c(\eta, \gamma_i^+) c(\eta, \gamma_j^-) I(\eta, \gamma_i^+, \gamma_j^-, \gamma_j^-) \\ &= \sum_{\eta, i, j} m(\eta) c(\eta, \gamma_j^+) c(\gamma_j^+ \eta, \gamma_i^+) I(\gamma_j^+ \eta, \gamma_i^+, \gamma_j^-, \gamma_j^-) \\ &= - \sum_{\eta, i, j} m(\eta) c(\eta, \gamma_i^+) c(\gamma_i^+ \eta, \gamma_j^+) I(\eta, \gamma_i^+, \gamma_j^+, \gamma_j^+). \end{aligned}$$

Therefore $A = 0$ as desired.

Term B First, using reversibility with

$$F(\eta, \sigma) = 1_{G^+}(\sigma) \sum_{\gamma \in G^-, \gamma \neq \sigma, \sigma^{-1}} \nabla_\sigma c(\eta, \gamma) I(\eta, \sigma, \sigma, \gamma)$$

and Lemma 4.20 we get the equivalent expression

$$B = \sum_{\eta, i \neq j} m(\eta) c(\eta, \gamma_i^-) \{-\nabla_i^- c(\eta, \gamma_j^-)\} I(\gamma_i^- \eta, \gamma_i^+, \gamma_i^+, \gamma_j^-).$$

By exchanging i, j and using the lemma we see that $B = 0$.

Term C We have

$$C = \sum_{\eta, i, j \neq i} m(\eta) c(\eta, \gamma_i^+) \{-\nabla_i^+ c(\eta, \gamma_j^+), 0\} I(\eta, \gamma_i^+, \gamma_j^+, \gamma_i^-).$$

Similarly to the term B , by exchanging i, j and using Lemma 4.20 we see that $C = 0$.

This concludes the proof of the theorem.

4.4.2 Examples of interacting random walks

As anticipated before and following [5], as an application of Theorem 4.17 and looking for a discrete analogue of Proposition 4.16 we now consider two particular examples of interacting random walks. In this subsection, we stick to the particular choice of V^- given in (26) and write $V = V^+$.

Example 1 For a function $h: \mathbb{R} \rightarrow \mathbb{R}$ write $\nabla^+ h(m) := h(m+1) - h(m)$ and for $\eta \in \Omega$ set $|\eta| = \sum_i |\eta_i|$. We consider a potential V of the form

$$V(\eta) := \beta h(|\eta|)$$

for some function $h: \mathbb{R} \rightarrow \mathbb{R}$ and $\beta > 0$.

Corollary 4.22 (Cor 3.1 of [5]). *With the notation of this section, suppose that h is **convex** and that*

$$\inf_{m \in \mathbb{N}} \lambda - (d-1) \left[e^{-\beta \nabla^+ h(m)} - e^{-\beta \nabla^+ h(m+1)} \right] \geq 0.$$

Then the assumptions of Theorem 4.17 are satisfied and

$$\kappa_* = \inf_{m \in \mathbb{N}} \lambda - (d-2) \left[e^{-\beta \nabla^+ h(m)} - e^{-\beta \nabla^+ h(m+1)} \right].$$

In particular, if h is strictly increasing and

$$\beta \geq \frac{\log(d-1) - \log(\lambda)}{h(1) - h(0)}$$

then the assumptions of Theorem 4.17 are satisfied with

$$\kappa_* = \lambda - (d-2)e^{-\beta \nabla^+ h(0)}.$$

Note that

- similarly to the classical Proposition 4.16, a convexity assumption on the potential comes into play;
- the necessary condition of a lower bound on β means this is a non perturbative criterion, since the resulting reversible measure is far from being a product measure. It is known that product measures behave well with the entropic curvature, i.e. they tensorise (see Theorem 6.2 of [11]). Therefore, it is particularly interesting to have conditions implying positive entropic curvature for a Markov chain whose stationary measure is not close to a product measure.

Example 2 Here, we see another example where a convexity condition on V is linked to inequality (8): in this case, convexity takes the form of non-negativity of the entries of a discrete Hessian for the potential V .

Corollary 4.23 (Cor 3.2 of [5]). *With the notation of this section, suppose that*

$$\nabla_i^+ \nabla_j^+ V(\eta) \geq 0 \tag{30}$$

for all $\eta \in \Omega$, $i, j \in [d]$ and that

$$\lambda - \sum_{\substack{j=1, \\ j \neq i}}^d \left[e^{-\nabla_j^+ V(\eta)} - e^{-\nabla_j^+ V(\gamma_i^+ \eta)} \right] \geq 0$$

for all $\eta \in \omega$, $i \in [d]$. Then the assumptions of Theorem 4.17 are satisfied and

$$\kappa_* = \inf_{\substack{\eta \in \Omega \\ i \in [d]}} \lambda + \left[e^{-\nabla_i^+ V(\eta)} - e^{-\nabla_i^+ V(\gamma_i^+ \eta)} \right] - \sum_{\substack{j=1, \\ j \neq i}}^d \left[e^{-\nabla_j^+ V(\eta)} - e^{-\nabla_j^+ V(\gamma_i^+ \eta)} \right].$$

If in addition

$$\min_{i \in [d]} \lambda - \sum_{\substack{j=1 \\ j \neq i}} e^{-\nabla_j^+ V(\mathbf{0})} \geq 0 \tag{31}$$

then κ_ is given by the previous equation (31).*

5 Couplings and coarse Ricci curvature

In this section, we recall some well known related definitions of curvature for discrete and continuous time Markov chains (generally referred to as “coarse Ricci curvature”), give some natural generalizations and discuss the relations among them and with the concept of couplings. As an application, we show that in all the examples of Section 4 (except for Section 4.4 - which was however covered in [5, Sec. 3.3] - since we restrict to the case of finite state space Markov chains), when the assumptions of the respective main theorems are satisfied then the coarse curvature is positive, and for all starting probability densities we have exponential contraction of the p -Wasserstein distances along the heat flow P_t (see the precise details later). The coarse Ricci curvature was first introduced by Ollivier for discrete time Markov chains (see [24, 23]) and later modified to apply to continuous time models (see [17, 28]). Compared to the entropic Ricci curvature, it is often easier to establish positive curvature for this notion. However, it is not known whether positive coarse Ricci curvature implies some functional inequalities, and in particular the modified log-Sobolev inequality (see Section 5.4 for a discussion).

Throughout this section, we switch to a more standard notation, and we don't employ the description of the Markov chain in terms of its allowed moves. As anticipated, we always assume in this section that we are working with finite state space, irreducible and reversible Markov chains. We use the letter Ω for the state space, x, y, w, z for elements of Ω , P for a stochastic transition matrix and L, Q for a generator L with transition rates Q , so that the action of L on a function $\psi: \Omega \rightarrow \mathbb{R}$ is given by

$$L\psi(x) = \sum_y Q(x, y)(\psi(y) - \psi(x)).$$

Here we have $Q(x, y) \geq 0$ and we do not assume $Q(x, x) = 0$, and in fact we often change the value of $Q(x, x)$ (without loss of generality) depending on convenience; on the other hand, we sometimes identify L , as a linear operator, with a matrix, in which case by construction $L(x, x) = -\sum_{y \neq x} Q(x, y)$. We typically identify measures with row vectors: in particular, let δ_x be the row vector with entry 1 corresponding to x and 0 everywhere else, which is identified with the Dirac measure at x ; on the other hand, densities with respect to π and other functions on the state space are identified with column vectors. We also introduce a simple graph structure, where $x \sim y$ if and only if $x \neq y$ and $P(x, y) > 0$ or $Q(x, y) > 0$ respectively, and correspondingly consider the unweighted graph distance d . With respect to this graph distance d , we will consider the p -Wasserstein distances W_p . Couplings for the transition measures/rates from starting points x, y will be described by non-negative functions $\Pi(x, y, \cdot, \cdot), C(x, y, \cdot, \cdot): \Omega \times \Omega \rightarrow \mathbb{R}_{\geq 0}$ respectively in discrete and continuous time, so that

$$\begin{cases} \sum_w \Pi(x, y, w, z) &= P(y, z) \text{ for all } z, \\ \sum_z \Pi(x, y, w, z) &= P(x, w) \text{ for all } w, \end{cases} \quad (32)$$

$$\begin{cases} \sum_w C(x, y, w, z) &= Q(y, z) \text{ for all } z, \\ \sum_z C(x, y, w, z) &= Q(x, w) \text{ for all } w. \end{cases} \quad (33)$$

The set of such admissible couplings is non empty, provided that in continuous time one redefines $Q(x, x)$ appropriately without loss of generality: notice indeed that the existence of the coupling rates implies (by summing over w, z) that $\sum_w Q(x, w) = \sum_z Q(y, z) =: Z$; if this holds, the “product” coupling rates $C(x, y, w, z) = \frac{1}{Z}Q(x, w)Q(y, z)$ are admissible. Notice that, to compare with the notation of the previous sections of this work in terms of allowed moves, we could write $c^{\text{pl}}(\eta, \sigma, \gamma, \bar{\gamma}) = C(\eta, \sigma\eta, \gamma\eta, \bar{\gamma}\sigma\eta)$ for a state η and moves $\sigma, \gamma, \bar{\gamma}$.

5.1 Discrete time

An important notion of curvature for Markov chains is the *coarse Ricci curvature* introduced by Ollivier (see [24] and [23]).

Definition 5.1. *Given $p \geq 1$ and $x \neq y$, we say that the Markov chain has (discrete time) p -coarse Ricci curvature $K_{dc,p}(x, y)$ in direction (x, y) if*

$$W_p(\delta_x P, \delta_y P) = (1 - K_{dc,p}(x, y))d(x, y).$$

Remark 5.2. *Ollivier focused in particular on the case $p = 1$; in these paper, however, it will be useful to consider also other values of p sometimes.*

We also give the following definition, inspired by the properties of the couplings constructed in the previous sections.

Definition 5.3. For $x \neq y$, we define the ∞ -coarse Ricci curvature $K_{dc,\infty}(x, y)$ in direction (x, y) to be the supremum of all $K \in \mathbb{R}$ such that there exists a coupling $\Pi(x, y, \cdot, \cdot)$ of $(\delta_x P, \delta_y P)$ satisfying:

- $\sum_{w, z \in \Omega} \Pi(x, y, w, z) \mathbb{1}_{d(w, z) > d(x, y)} = 0$;
- $\sum_{w, z \in \Omega} \Pi(x, y, w, z) d(w, z) \leq (1 - K) d(x, y)$.

We use the convention that $\sup \emptyset = -\infty$.

Remark 5.4. We have that $K_{dc,\infty}(x, y) \in \mathbb{R}_{\geq 0} \cup \{-\infty\}$; if $K_{dc,\infty}(x, y) \geq 0$ then the supremum in the definition is attained.

Notice that, equivalently, $K_{dc,\infty}(x, y) \geq 0$ means that there exists a coupling (X, Y) of the one step probability distribution $(\delta_x P, \delta_y P)$ which proves $K_{dc,1}(x, y) \geq K_{dc,\infty}(x, y)$ and at the same time satisfies $d(X, Y) \leq d(x, y)$ almost surely.

For $p \in [1, \infty]$ we write

$$\text{Ric}_{dc,p} \geq K$$

if $K_{dc,p}(x, y) \geq K$ for all $x \neq y \in \Omega$. The next propositions collect some useful results.

Proposition 5.5. The following hold:

1. For $p \in [1, \infty]$, if $K_{dc,p}(x, y) \geq K$ for all $x \sim y$ then $\text{Ric}_{dc,p} \geq K$.
2. For $1 \leq p \leq q < \infty$ we have $K_{dc,p}(x, y) \geq K_{dc,q}(x, y)$. Moreover if $x \sim y$ we have

$$K_{dc,p}(x, y) \geq \frac{K_{dc,\infty}(x, y)}{p}.$$

3. If $x \sim y$ and $\lim_{p \rightarrow \infty} K_{dc,p}(x, y) \geq 0$ we have

$$K_{dc,\infty}(x, y) \geq 1 - e^{-\limsup_{p \rightarrow \infty} p \cdot K_{dc,p}(x, y)}.$$

4. For $p \in [1, \infty)$, if $\text{Ric}_{dc,p} \geq K$ then for any starting probability measures μ, ν and $n \geq 0$ we have

$$W_p(\mu P^n, \nu P^n) \leq (1 - K)^n W_p(\mu, \nu).$$

Proof. 1. If $p < \infty$, this is done as in [24, Prop. 19]: suppose $d(x, y) = n$ and let $x = z_0 \sim z_1 \sim \dots \sim z_n = y$. Then

$$W_p(\delta_x P, \delta_y P) \leq \sum_{i=0}^{n-1} W_p(\delta_{z_{i-1}} P, \delta_{z_i} P) \leq (1 - K)n = (1 - K)d(x, y).$$

Suppose now that $p = \infty$: if $K = -\infty$ the conclusion is trivial, hence assume that $K \geq 0$. Let again $n = d(x, y)$ and $x = z_0 \sim z_1 \sim \dots \sim z_n = y$: we prove the claim by induction over n . The base case $n = 1$ follows directly by the assumption. Now suppose $n > 1$ and that the inductive hypothesis holds. Let $\Pi(x, z_{n-1}, \cdot, \cdot)$ and $\Pi(z_{n-1}, y, \cdot, \cdot)$ be such that

$$\begin{aligned} \sum_{v, w} \Pi(x, z_{n-1}, v, w) \mathbb{1}_{d(v, w) > d(x, z_{n-1})} &= 0, \\ \sum_{v, w} \Pi(x, z_{n-1}, v, w) d(v, w) &\leq (1 - K) d(x, z_{n-1}), \\ \sum_{v, w} \Pi(z_{n-1}, y, v, w) \mathbb{1}_{d(v, w) > 1} &= 0, \\ \sum_{v, w} \Pi(z_{n-1}, y, v, w) d(v, w) &\leq (1 - K). \end{aligned}$$

By the Gluing lemma (see [29]), there exists $\hat{\Pi}(x, z_{n-1}, y, \cdot, \cdot, \cdot) \in \mathcal{P}(\Omega \times \Omega \times \Omega)$ such that $p_{1,2} \# \hat{\Pi} = \Pi(x, z_{n-1}, \cdot, \cdot)$ and $p_{2,3} \# \hat{\Pi} = \Pi(z_{n-1}, y, \cdot, \cdot)$, where $p_{i,j}$ is the projection on coordinates i, j and $\#$ denotes

the pushforward of a measure via a map. The measure $\Pi(x, y, \cdot, \cdot) := p_{1,3} \# \hat{\Pi}$ then realizes a coupling with the desired properties, since, given that $d(v, w) \leq d(v, s) + d(s, w)$, we have

$$\begin{aligned}
\sum_{v,w} \Pi(x, y, v, w) d(v, w) &= \sum_{v,w,s} \hat{\Pi}(x, z_{n-1}, y, v, s, w) d(v, w) \\
&\leq \sum_{v,w,s} \hat{\Pi}(x, z_{n-1}, y, v, s, w) (d(v, s) + d(s, w)) \\
&= \sum_{v,s} \Pi(x, z_{n-1}, v, s) d(v, s) + \sum_{s,w} \Pi(z_{n-1}, y, s, w) d(s, w) \\
&\leq (1 - K)(d(x, z_{n-1}) + d(z_{n-1}, y)) \\
&= (1 - K)d(x, y),
\end{aligned}$$

and similarly

$$\begin{aligned}
&\sum_{v,w} \Pi(x, y, v, w) \mathbb{1}_{d(v,w) > d(x,y)} \\
&= \sum_{v,w,s} \hat{\Pi}(x, z_{n-1}, y, v, s, w) \mathbb{1}_{d(v,w) > d(x,y)} \\
&\leq \sum_{v,w,s} \hat{\Pi}(x, z_{n-1}, y, v, s, w) \mathbb{1}_{d(v,s) + d(s,w) \geq d(x, z_{n-1}) + d(z_{n-1}, y)} \\
&\leq \sum_{v,w,s} \hat{\Pi}(x, z_{n-1}, y, v, s, w) (\mathbb{1}_{d(v,s) \geq d(x, z_{n-1})} + \mathbb{1}_{d(s,w) \geq d(z_{n-1}, y)}) \\
&= \sum_{v,s} \Pi(x, z_{n-1}, v, s) \mathbb{1}_{d(v,s) \geq d(x, z_{n-1})} + \sum_{s,w} \Pi(z_{n-1}, y, s, w) \mathbb{1}_{d(s,w) \geq d(z_{n-1}, y)} \\
&= 0.
\end{aligned}$$

2. The first statement follows by the inequality $W_p(\mu, \nu) \leq W_q(\mu, \nu)$ for $1 \leq p \leq q < \infty$. For the second statement, suppose $K \geq 0$ and $x \sim y$. Let Π be the optimal coupling in the definition of $K_{dc, \infty}(x, y)$. Then notice that

$$W_p(\delta_x P, \delta_y P) \leq [1 - K]^{\frac{1}{p}} \leq 1 - \frac{K}{p},$$

from which the conclusion follows.

3. Let $K = \limsup_{p \rightarrow \infty} p \cdot K_{dc, p}(x, y) \geq 0$. By restricting to appropriate subsequences, we can find sequences $(p_n)_n \subset [1, \infty)$, $(K_n)_n \subset (-\infty, 1]$ and couplings $(\Pi_n(x, y, \cdot, \cdot))_n, \Pi$ of $(\delta_x P, \delta_y P)$ such that $p_n \rightarrow \infty$ and $\Pi_n \rightarrow \Pi$ as $n \rightarrow \infty$, $W_{p_n}(\delta_x P, \delta_y P) \leq (1 - K_n)$ when using coupling Π_n , that is

$$\left(\sum_{w,z} \Pi_n(x, y, w, z) d(w, z)^{p_n} \right)^{\frac{1}{p_n}} \leq 1 - K_n.$$

We claim that the coupling Π has the desired properties. Notice indeed that

$$\begin{aligned}
&2^{p_n} \sum_{w,z} \Pi_n(x, y, w, z) \mathbb{1}_{d(w,x) \geq 2} + \sum_{w,z} \Pi_n(x, y, w, z) \mathbb{1}_{d(w,x) = 1} \\
&\leq W_{p_n}^{p_n}(\delta_x P, \delta_y P) \\
&\leq (1 - K_n)^{p_n} \\
&\leq e^{-K_n p_n}.
\end{aligned}$$

Letting $n \rightarrow \infty$ in the previous we deduce

$$\begin{cases} \sum_{w,z} \Pi(x, y, w, z) \mathbb{1}_{d(w,x) \geq 2} = 0, \\ \sum_{w,z} \Pi(x, y, w, z) \mathbb{1}_{d(w,x) = 1} \leq e^{-K}, \end{cases}$$

which yields the conclusion.

4. Let $\Pi_{\mu,\nu}$ and $\Pi_{x,y}$ be the optimal couplings in the definitions of $W_p(\mu,\nu)$ and $K_{dc,p}(x,y)$ respectively. Then

$$\begin{aligned} W_p^p(\mu P, \nu P) &\leq \sum_{x,y} d(x,y)^p \sum_{w,z} \Pi_{\mu,\nu}(w,z) \Pi_{w,z}(x,y) \\ &\leq (1-K)^p \sum_{w,z} \Pi_{\mu,\nu}(w,z) d(w,z)^p \\ &\leq (1-K)^p W_p^p(\mu,\nu). \end{aligned}$$

□

5.2 Continuous time

In this subsection, we describe the analogue notions of coarse Ricci curvature for continuous time Markov chains with generator L (see [28, 17, 22]). Recalling we identify L with a matrix with zero row sums, we use the notation

$$\tilde{P}_t = I + tL,$$

so that for $t > 0$ small enough \tilde{P}_t is a stochastic matrix too and we expect it to approximate P_t (given it corresponds to the first order Taylor expansion of $P_t = e^{tL}$). The next definition is motivated by the study of the idleness function in [3].

Definition 5.6. Let $T = (\max_{x \in \Omega} -L(x,x))^{-1} = \left(\max_{x \in \Omega} \sum_{y \neq x} Q(x,y) \right)^{-1}$. Fix $x \neq y \in \Omega$ and let $p \in [1, \infty)$. Then we define the function $I_{p,x,y}: [0, T] \rightarrow \mathbb{R}$ by

$$I_{p,x,y}(t) := W_p^p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t).$$

For simplicity, we drop the subscripts x, y when there is no confusion. Notice that $I_p(0) = d(x,y)^p$. The next propositions and lemma are straightforward adaptations of the results in [3], where the case $p = 1$ was considered. Our motivation here is to establish linearity of $t \rightarrow I_p(t)$ for small values of t , hence the differentiability at $t = 0$ (cf. Proposition 5.9): this allows us to consider a first definition of coarse Ricci curvature in continuous time, cf. Definition 5.11. This definition was first given for $p = 1$ in [17] for combinatorial graphs (i.e. simple random walks on graphs), and later studied in a more general setting in [22].

Proposition 5.7. The function $t \rightarrow I_p(t)$ is convex.

Proof. We use Kantorovich duality, where for $\psi: \Omega \rightarrow \mathbb{R}$ we denote by $\psi^{c,p}$ its d^p transform, i.e.

$$\psi^{c,p}(x) := \inf_y d(x,y)^p - \psi(y).$$

Then we have for $t \in [0, T]$

$$\begin{aligned} &W_p^p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t) \\ &= \sup_{\psi} \left\{ \sum_{z \in \Omega} \psi(z) \cdot (\delta_x \tilde{P}_t)(z) + \psi^{c,p}(z) \cdot (\delta_y \tilde{P}_t)(z) \right\} \\ &= \sup_{\psi} \left\{ \psi(x) \left(1 - t \sum_z Q(x,z) \right) + \psi^{c,p}(y) \left(1 - t \sum_z Q(y,z) \right) + t \left[\sum_z Q(x,z) \psi(z) + Q(y,z) \psi^{c,p}(z) \right] \right\}. \end{aligned}$$

This is the supremum of affine functions of t , hence $t \rightarrow W_p^p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t) = I_{p,x,y}(t)$ is convex. □

Lemma 5.8. Let $0 \leq t_1 < t_2 < T$ and suppose that $(\psi, \psi^{c,p})$ is a pair of optimal Kantorovich potentials in the definition of both $W_p(\delta_x \tilde{P}_{t_1}, \delta_y \tilde{P}_{t_1})$ and $W_p(\delta_x \tilde{P}_{t_2}, \delta_y \tilde{P}_{t_2})$. Then $t \rightarrow I_p(t)$ is linear over $[t_1, t_2]$.

Proof. We already know the function $t \rightarrow W_p^p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t)$ is convex, hence it suffices to show it is also concave over $[t_1, t_2]$. This follows by the assumption using Kantorovich duality, since for $\alpha \in [0, 1]$

$$\begin{aligned}
& W_p^p(\delta_x \tilde{P}_{\alpha t_1 + (1-\alpha)t_2}, \delta_y \tilde{P}_{\alpha t_1 + (1-\alpha)t_2}) \\
& \geq \sum_{z \in \Omega} \psi(z) \cdot (\delta_x \tilde{P}_{\alpha t_1 + (1-\alpha)t_2})(z) + \psi^{c,p}(z) \cdot (\delta_y \tilde{P}_{\alpha t_1 + (1-\alpha)t_2})(z) \\
& = \psi(x) \left(1 - (\alpha t_1 + (1-\alpha)t_2) \sum_z Q(x, z) \right) + \psi^{c,p}(y) \left(1 - (\alpha t_1 + (1-\alpha)t_2) \sum_z Q(y, z) \right) \\
& \quad + (\alpha t_1 + (1-\alpha)t_2) \left[\sum_z Q(x, z) \psi(z) + \sum_z Q(y, z) \psi^{c,p}(z) \right] \\
& = \alpha W_p^p(\delta_x \tilde{P}_{t_1}, \delta_y \tilde{P}_{t_1}) + (1-\alpha) W_p^p(\delta_x \tilde{P}_{t_2}, \delta_y \tilde{P}_{t_2}).
\end{aligned}$$

□

Proposition 5.9. *The function I_p is linear over $[0, \frac{T}{2}]$.*

Proof. Fix $t < \frac{T}{2}$ and let ψ and Γ be a couple of optimal Kantorovich potential and plan for $W_p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t)$. Then it is easy to see that $\Gamma(x, y) > 0$ (since the laziness of \tilde{P}_t is at least $1 - t/T > \frac{1}{2}$) and so $\psi(x) + \psi^c(y) = d(x, y)^p$ (cf. [29, Thm. 5.10] for example). This implies that (ψ, ψ^c) is optimal also for $W_p(\delta_x \tilde{P}_0, \delta_y \tilde{P}_0) = W_p(\delta_x, \delta_y) = d(x, y)$. The conclusion follows by Lemma 5.8 and by letting $t \rightarrow \frac{1}{2T}$ (by continuity at $\frac{T}{2}$, which follows from the convexity of I_p). □

Since linear functions are easily differentiated, the previous proposition immediately implies the following corollary, which gives an expression for the derivative of $I_p(t)$ at $t = 0$.

Corollary 5.10. *For $x \neq y$ there exists*

$$\left. \frac{d}{dt} \right|_{t=0} W_p^p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t) = -\frac{1}{s} \left(d(x, y)^p - W_p^p(\delta_x \tilde{P}_s, \delta_y \tilde{P}_s) \right) \quad (34)$$

for all $0 < s \leq \frac{T}{2}$.

Definition 5.11. *The continuous time p -coarse Ricci curvature in direction (x, y) is defined via*

$$\begin{aligned}
K_{cc,p}(x, y) & := -\frac{1}{d(x, y)} \left. \frac{d}{dt} \right|_{t=0} W_p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t) \\
& = -\frac{1}{p \cdot d(x, y)^p} \left. \frac{d}{dt} \right|_{t=0} W_p^p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t) \\
& = \frac{1}{sp} \left(1 - \frac{W_p^p(\delta_x \tilde{P}_s, \delta_y \tilde{P}_s)}{d(x, y)^p} \right)
\end{aligned} \quad (35)$$

for all $0 < s \leq \frac{T}{2}$.

Remark 5.12. *The original definition in [17] (see also [22]) focused on the case $p = 1$, but in this paper it will be useful to consider also other values of p .*

Next, we prove a few preliminary results needed for the proof of Lemma 5.16, which will show that we can define $K_{cc,p}$ also by involving P_t instead of \tilde{P}_t . For $p = 1$, this shows the equivalence with the notion of continuous time coarse Ricci curvature defined in [28], which was first proved in [22, Thm. 5.8].

Lemma 5.13. *Let X, Y be finite sets and $\mu, \tilde{\mu} \in \mathcal{P}(X), \nu \in \mathcal{P}(Y)$ probability measures. Suppose that $\Gamma \in \Pi(\mu, \nu)$ is a coupling. Then there exists a coupling $\tilde{\Gamma} \in \Pi(\tilde{\mu}, \nu)$ that satisfies*

$$\|\tilde{\Gamma} - \Gamma\|_{TV} = \|\tilde{\mu} - \mu\|_{TV}$$

where $\|\mu - \tilde{\mu}\|_{TV} \in [0, 1]$ is the total variation distance.

Proof. We will construct a coupling $\tilde{\Gamma} \in \Pi(\tilde{\mu}, \nu)$ such that

$$\tilde{\Gamma}(x, y) \geq \Gamma(x, y) \text{ if and only if } \tilde{\mu}(x) \geq \mu(x). \quad (36)$$

Notice that this implies the thesis since we would have

$$\begin{aligned} \|\tilde{\Gamma} - \Gamma\|_{\text{TV}} &= \frac{1}{2} \sum_{x \in X, y \in Y} \left| \tilde{\Gamma}(x, y) - \Gamma(x, y) \right| \\ &= \frac{1}{2} \sum_{x, y: \tilde{\mu}(x) \geq \mu(x)} \left[\tilde{\Gamma}(x, y) - \Gamma(x, y) \right] + \frac{1}{2} \sum_{x, y: \tilde{\mu}(x) < \mu(x)} \left[\Gamma(x, y) - \tilde{\Gamma}(x, y) \right] \\ &= \frac{1}{2} \sum_{x: \tilde{\mu}(x) \geq \mu(x)} [\tilde{\mu}(x) - \mu(x)] + \frac{1}{2} \sum_{x: \tilde{\mu}(x) < \mu(x)} [\mu(x) - \tilde{\mu}(x)] \\ &= \frac{1}{2} \sum_{x \in X} |\tilde{\mu}(x) - \mu(x)| \\ &= \|\tilde{\mu} - \mu\|_{\text{TV}}. \end{aligned}$$

Let us thus construct $\tilde{\Gamma}$ that satisfies (36). To do that, for every $x \in X, y \in Y$ set

$$\begin{aligned} \alpha(x) &= \tilde{\mu}(x) - (\tilde{\mu} \wedge \mu)(x), \\ \beta(y) &= \nu(y) - \sum_{x \in X: \tilde{\mu}(x) < \mu(x)} \Gamma(x, y) \frac{\tilde{\mu}(x)}{\mu(x)} - \sum_{x \in X: \tilde{\mu}(x) \geq \mu(x)} \Gamma(x, y). \end{aligned}$$

It is easy to see that $\sum_{x \in X} \alpha(x) = \sum_{y \in Y} \beta(y) = \|\mu - \tilde{\mu}\|_{\text{TV}}$. Then we define

$$\tilde{\Gamma}(x, y) := \mathbb{1}_{\tilde{\mu}(x) < \mu(x)} \frac{\tilde{\mu}(x)}{\mu(x)} \Gamma(x, y) + \mathbb{1}_{\tilde{\mu}(x) \geq \mu(x)} \Gamma(x, y) + \frac{1}{\|\mu - \tilde{\mu}\|_{\text{TV}}} \alpha(x) \beta(y).$$

It is easy to check that $\tilde{\Gamma} \in \Pi(\tilde{\mu}, \nu)$ and that $\tilde{\Gamma}$ satisfies (36). \square

Corollary 5.14. *Let X, Y be finite sets, $\mu_1, \mu_2 \in \mathcal{P}(X)$, $\nu_1, \nu_2 \in \mathcal{P}(Y)$ and $c: X \times Y \rightarrow \mathbb{R}_{\geq 0}$ be a cost function. For any probability measures $\mu \in \mathcal{P}(X), \nu \in \mathcal{P}(Y)$, let $\mathcal{T}(\mu, \nu)$ be the optimal transport cost associated to the cost c . Then*

$$|\mathcal{T}(\mu_1, \nu_1) - \mathcal{T}(\mu_2, \nu_2)| \leq 2 \max_{x \in X, y \in Y} c(x, y) \cdot (\|\mu_1 - \mu_2\|_{\text{TV}} + \|\nu_1 - \nu_2\|_{\text{TV}}).$$

Proof. Let $M := \max_{x \in X, y \in Y} c(x, y) \geq 0$. Notice that

$$|\mathcal{T}(\mu_1, \nu_1) - \mathcal{T}(\mu_2, \nu_2)| \leq |\mathcal{T}(\mu_1, \nu_1) - \mathcal{T}(\mu_2, \nu_1)| + |\mathcal{T}(\mu_2, \nu_1) - \mathcal{T}(\mu_2, \nu_2)|,$$

hence it suffices to show that

$$\begin{aligned} |\mathcal{T}(\mu_1, \nu_1) - \mathcal{T}(\mu_2, \nu_1)| &\leq 2M \cdot \|\mu_1 - \mu_2\|_{\text{TV}}, \\ |\mathcal{T}(\mu_2, \nu_1) - \mathcal{T}(\mu_2, \nu_2)| &\leq 2M \cdot \|\nu_1 - \nu_2\|_{\text{TV}}. \end{aligned}$$

We prove only the first inequality, since the proof of the second one is similar. Let Γ be an optimal coupling in the definition of $\mathcal{T}(\mu_1, \nu_1)$ and $\tilde{\Gamma}$ be the coupling for (μ_2, ν_1) given by Lemma 5.13. Then it follows that

$$\begin{aligned} \mathcal{T}(\mu_2, \nu_1) &\leq \sum_{x, y} c(x, y) \tilde{\Gamma}(x, y) \\ &\leq \sum_{x, y} c(x, y) \left(\Gamma(x, y) + \left| \tilde{\Gamma} - \Gamma \right|(x, y) \right) \\ &\leq \mathcal{T}(\mu_1, \nu_1) + 2M \|\tilde{\Gamma} - \Gamma\|_{\text{TV}} \\ &\leq \mathcal{T}(\mu_1, \nu_1) + 2M \|\mu_1 - \mu_2\|_{\text{TV}}. \end{aligned}$$

Similarly one shows that

$$\mathcal{T}(\mu_1, \nu_1) \leq \mathcal{T}(\mu_2, \nu_1) + 2M \|\mu_1 - \mu_2\|_{\text{TV}},$$

which concludes the proof of the corollary. \square

Corollary 5.15. For any probability measures $\mu, \nu \in \mathcal{P}(\Omega)$ and $t > 0$ small enough we have

$$\left| W_p^p(\mu P_t, \nu P_t) - W_p^p(\mu \tilde{P}_t, \nu \tilde{P}_t) \right| = O(t^2)$$

as $t \rightarrow 0$.

Proof. Fix $\mu, \nu \in \mathcal{P}(X)$ and $t > 0$ small and let $D := \text{diam}(\Omega)$. Since Ω is finite, we have that $D < \infty$ and $\|\mu \tilde{P}_t - \mu P_t\|_{\text{TV}}, \|\nu \tilde{P}_t - \nu P_t\|_{\text{TV}} = O(t^2)$. By applying Corollary 5.14 for $t > 0$ small enough with $\mu_1 = \mu P_t, \mu_2 = \mu \tilde{P}_t, \nu_1 = \nu P_t, \nu_2 = \nu \tilde{P}_t$, we find

$$\left| W_p^p(\mu P_t, \nu P_t) - W_p^p(\mu \tilde{P}_t, \nu \tilde{P}_t) \right| \leq 2D^p \cdot \left(\|\mu \tilde{P}_t - \mu P_t\|_{\text{TV}} + \|\nu \tilde{P}_t - \nu P_t\|_{\text{TV}} \right) = O(t^2),$$

as desired. □

We can finally give an equivalent definition of $K_{cc,p}$ using P_t instead of \tilde{P}_t .

Corollary 5.16. For $x \neq y \in \Omega$ we have

$$K_{cc,p}(x, y) = -\frac{1}{d(x, y)} \frac{d}{dt} \Big|_{t=0} W_p(\delta_x P_t, \delta_y P_t). \quad (37)$$

Proof. Recalling Definition 5.11, it suffices to show that

$$\frac{d}{dt} \Big|_{t=0} W_p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t) = \frac{d}{dt} \Big|_{t=0} W_p(\delta_x P_t, \delta_y P_t).$$

Equivalently, since $W_p(\delta_x P_0, \delta_y P_0) = W_p(\delta_x \tilde{P}_0, \delta_y \tilde{P}_0) = d(x, y) > 0$, it is enough to show

$$\frac{d}{dt} \Big|_{t=0} W_p^p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t) = \frac{d}{dt} \Big|_{t=0} W_p^p(\delta_x P_t, \delta_y P_t),$$

but this follows from Corollary 5.15. □

Remark 5.17. As anticipated, for $p = 1$ the right hand side in (37) corresponds to the definition of curvature for continuous time Markov chain introduced by Veysseire (see [28]). For $p = 1$, the identity in the corollary was established in [22] (in a more general setting).

As in discrete time, we define also the curvature for $p = \infty$.

Definition 5.18. For $x \neq y$, we define the continuous time ∞ -coarse Ricci curvature $K_{cc,\infty}(x, y)$ in direction (x, y) to be the supremum of all $K \in \mathbb{R}$ such that there exist coupling rates $C(x, y, \cdot, \cdot)$ satisfying:

- $\sum_{v,z \in \Omega} C(x, y, v, z) \mathbb{1}_{d(v,z) > d(x,y)} = 0$;
- $\sum_{v,z \in \Omega} C(x, y, v, z) d(v, z) \leq \left(\sum_{v,z} C(x, y, v, z) - K \right) d(x, y)$.

Remark 5.19. We have that $K_{cc,\infty}(x, y) \in \mathbb{R}_{\geq 0} \cup \{-\infty\}$; if $K_{cc,\infty}(x, y) \geq 0$, the supremum in the definition is attained. Notice that if $C(x, y, \cdot, \cdot)$ gives optimal coupling rates, then we can change the value $C(x, y, x, y)$ arbitrarily obtaining other optimal coupling rates.

As in discrete time, for $p \in [1, \infty]$ we write

$$\text{Ric}_{cc,p} \geq K$$

if $K_{cc,p}(x, y) \geq K$ for all $x \neq y \in \Omega$.

The next results shows that, given coupling rates from $x \neq y$, we can construct a coupling for the probability measures $(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t)$ for any $t > 0$ small enough. This will be useful to connect the notions of coarse Ricci curvature for discrete and continuous time Markov chains, when we consider a natural correspondence between stochastic transition matrices P and generators L , see Section 5.3.

Lemma 5.20. Let $x \neq y \in \Omega$ and $C(x, y, \cdot, \cdot)$ denote coupling rates from x, y and set

$$M := \sum_{v,w} C(x, y, v, w) = \sum_v Q(x, v) = \sum_w Q(y, w).$$

Then for any $0 < t \leq \frac{1}{M - C(x, y, x, y)}$ we have that Γ_t is a coupling for $(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t)$, where we define

$$\Gamma_t(v, w) = \begin{cases} C(x, y, v, w) \cdot t & \text{if } (v, w) \neq (x, y), \\ 1 - M \cdot t + C(x, y, x, y) \cdot t & \text{if } (v, w) = (x, y). \end{cases}$$

Proof. Clearly for $0 < t \leq \frac{1}{M - C(x, y, x, y)}$ we have $\Gamma_t(v, w) \geq 0$. Now if $v \neq x$ we have

$$\sum_w \Gamma_t(v, w) = t \sum_w C(x, y, v, w) = tQ(x, v) = (\delta_x \tilde{P}_t)(v).$$

If $v = x$ then

$$\sum_w \Gamma_t(x, w) = t \sum_{w \neq y} C(x, y, x, w) + 1 - M \cdot t + C(x, y, x, y) \cdot t = 1 + Q(x, x) \cdot t - M \cdot t = (\delta_x \tilde{P}_t)(x).$$

Similarly one checks that the second marginal of Γ_t is $\delta_y \tilde{P}_t$. \square

The next propositions collect some useful results, and it is a continuous time analogue of Proposition 5.5.

Proposition 5.21. *The following hold:*

1. For $p \in [1, \infty]$, if $K_{cc,p}(x, y) \geq K$ for all $x \sim y$ then $\text{Ric}_{cc,p} \geq K$.
2. For $1 \leq p \leq q < \infty$ we have $K_{cc,p}(x, y) \geq K_{cc,q}(x, y)$. Moreover if $x \sim y$ we have

$$K_{cc,p}(x, y) \geq \frac{K_{cc,\infty}(x, y)}{p}.$$

3. For $p \in [1, \infty)$, if $\text{Ric}_{dc,p} \geq K$ then for any starting measures μ, ν and $t \geq 0$ we have

$$W_p(\mu P_t, \nu P_t) \leq e^{-Kt} W_p(\mu, \nu).$$

Proof. 1. Suppose first that $p < \infty$ and let $n = d(x, y)$ with $x = z_0 \sim z_1 \sim \dots \sim z_n = y$. Then for $t > 0$ small enough we have that

$$-W_p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t) \geq \sum_{i=0}^{n-1} -W_p(\delta_{z_i} \tilde{P}_t, \delta_{z_{i+1}} \tilde{P}_t)$$

by the triangle inequality. Therefore adding n and dividing by t

$$\frac{d(x, y) - W_p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t)}{t} \geq \sum_{i=0}^{n-1} \frac{1 - W_p(\delta_{z_i} \tilde{P}_t, \delta_{z_{i+1}} \tilde{P}_t)}{t}.$$

Letting $t \rightarrow 0$ and using the assumption gives the conclusion.

Suppose now that $p = \infty$: if $K = -\infty$ the conclusion is trivial, hence assume that $K \geq 0$. Let again $n = d(x, y)$ and $x = z_0 \sim z_1 \sim \dots \sim z_n = y$: we prove the claim by induction over n . The base case $n = 1$ follows directly by the assumption. Now suppose $n > 1$ and that the inductive hypothesis holds. Let $C(x, z_{n-1}, \cdot, \cdot)$ and $C(z_{n-1}, y, \cdot, \cdot)$ be such that

$$\begin{aligned} \sum_{v,w} C(x, z_{n-1}, v, w) \mathbb{1}_{d(v,w) > d(x, z_{n-1})} &= 0, \\ \sum_{v,w} C(x, z_{n-1}, v, w) d(v, w) &\leq \left(\sum_{v,w} C(x, z_{n-1}, v, w) - K \right) d(x, z_{n-1}), \\ \sum_{v,w} C(z_{n-1}, y, v, w) \mathbb{1}_{d(v,w) > 1} &= 0, \\ \sum_{v,w} C(z_{n-1}, y, v, w) d(v, w) &\leq \left(\sum_{v,w} C(z_{n-1}, y, v, w) - K \right). \end{aligned}$$

Without loss of generality, by changing if needed the values of $C(x, z_{n-1}, x, z_{n-1})$, $C(z_{n-1}, y, z_{n-1}, y)$, $Q(x, x)$, $Q(z_{n-1}, z_{n-1})$, $Q(y, y)$, we can assume that

$$\sum_v Q(x, v) = \sum_z Q(z_{n-1}, z) = \sum_w Q(y, w) = \sum_{v,w} C(x, z_{n-1}, v, w) = \sum_{v,w} C(z_{n-1}, y, v, w) =: M > 0.$$

Therefore, we can apply the Gluing lemma (which easily extends to measures having the same total mass) to conclude that there exists $\hat{C}(x, z_{n-1}, y, \cdot, \cdot, \cdot): \Omega \times \Omega \times \Omega \rightarrow \mathbb{R}_{\geq 0}$ such that $p_{1,2}\#\hat{C} = C(x, z_{n-1}, \cdot, \cdot)$ and $p_{2,3}\#\hat{C} = C(z_{n-1}, y, \cdot, \cdot)$, where $p_{i,j}$ is the projection on coordinates i, j and $\#$ is the pushforward, so that

$$\begin{aligned} \sum_w \hat{C}(x, z_{n-1}, y, v, s, w) &= C(x, z_{n-1}, v, s), \\ \sum_v \hat{C}(x, z_{n-1}, y, v, s, w) &= C(z_{n-1}, y, s, w). \end{aligned}$$

Defining $C(x, y, \cdot, \cdot) := p_{1,3}\#\hat{C}$ gives coupling rates with the desired properties, analogously to the proof of Proposition 5.5. Indeed, we have that

$$\begin{aligned} \sum_{v,w} C(x, y, v, w) d(v, w) &= \sum_{v,w,s} \hat{C}(x, z_{n-1}, y, v, s, w) d(v, w) \\ &\leq \sum_{v,w,s} \hat{C}(x, z_{n-1}, y, v, s, w) (d(v, s) + d(s, w)) \\ &= \sum_{v,s} C(x, z_{n-1}, v, s) d(v, s) + \sum_{s,w} C(z_{n-1}, y, s, w) d(s, w) \\ &\leq (M - K)(d(x, z_{n-1}) + d(z_{n-1}, y)) \\ &= \left(\sum_{v,w} C(x, y, v, w) - K \right) d(x, y), \end{aligned}$$

and similarly

$$\begin{aligned} &\sum_{v,w} C(x, y, v, w) \mathbb{1}_{d(v,w) > d(x,y)} \\ &= \sum_{v,w,s} \hat{C}(x, z_{n-1}, y, v, s, w) \mathbb{1}_{d(v,w) > d(x,y)} \\ &\leq \sum_{v,w,s} \hat{C}(x, z_{n-1}, y, v, s, w) \mathbb{1}_{d(v,s) + d(s,w) \geq d(x, z_{n-1}) + d(z_{n-1}, y)} \\ &\leq \sum_{v,w,s} \hat{C}(x, z_{n-1}, y, v, s, w) (\mathbb{1}_{d(v,s) \geq d(x, z_{n-1})} + \mathbb{1}_{d(s,w) \geq d(z_{n-1}, y)}) \\ &= \sum_{v,s} C(x, z_{n-1}, v, s) \mathbb{1}_{d(v,s) \geq d(x, z_{n-1})} + \sum_{s,w} C(z_{n-1}, y, s, w) \mathbb{1}_{d(s,w) \geq d(z_{n-1}, y)} \\ &= 0. \end{aligned}$$

2. The first statement follows by the inequality $W_p(\mu, \nu) \leq W_q(\mu, \nu)$: indeed it implies

$$\frac{d(x, y) - W_p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t)}{t d(x, y)} \geq \frac{d(x, y) - W_q(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t)}{t d(x, y)},$$

from which the conclusion follows by letting $t \rightarrow 0$. For the second statement, suppose $K_{cc,\infty}(x, y) \geq 0$ for $x \sim y$. Let $C(x, y, \cdot, \cdot)$ be optimal coupling rates in the definition of $K_{cc,\infty}(x, y)$. Then for $t > 0$ small enough consider the coupling Γ_t given by Lemma 5.20. It is easy to see that this coupling is such that

$$\begin{aligned} \sum_{v,w} \Gamma_t(v, w) \mathbb{1}_{d(v,w) > 1} &= 0; \\ \sum_{v,w} \Gamma_t(v, w) \mathbb{1}_{d(v,w) = 0} &\geq K_{cc,\infty}(x, y) \cdot t. \end{aligned}$$

Therefore, it shows that

$$W_p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t) \leq (1 - K_{cc,\infty}(x, y) \cdot t)^{\frac{1}{p}} \leq 1 - \frac{K_{cc,\infty}(x, y) \cdot t}{p}.$$

Hence we have

$$\frac{1 - W_p(\delta_x \tilde{P}_t, \delta_y \tilde{P}_t)}{t} \geq \frac{K_{cc,\infty}(x, y)}{p},$$

from which the conclusion follows by letting $t \rightarrow 0$.

3. Let $\Pi_{\mu,\nu}$ and $\Pi_{x,y,t}$ be optimal couplings in the definitions of $W_p(\mu, \nu)$ and $W_p(\delta_x P_t, \delta_y P_t)$ respectively. Then notice that

$$\begin{aligned} W_p^p(\mu P_t, \nu P_t) &\leq \sum_{x,y} d(x, y)^p \sum_{w,z} \Pi_{\mu,\nu}(w, z) \Pi_{w,z,t}(x, y) \\ &= \sum_{w,z} \Pi_{\mu,\nu}(w, z) W_p^p(\delta_w P_t, \delta_z P_t), \end{aligned}$$

and so

$$\frac{W_p^p(\mu P_t, \nu P_t) - W_p^p(\mu, \nu)}{t} \leq \sum_{w,z} \Pi_{\mu,\nu}(w, z) \frac{W_p^p(\delta_w P_t, \delta_z P_t) - d(w, z)^p}{t}.$$

Taking the $\limsup_{t \rightarrow 0^+}$ and denoting by $\frac{d^+}{dt}$ the upper Dini derivative (cf. Section C) we find that

$$\begin{aligned} &\frac{d^+}{dt} \Big|_{t=0} W_p^p(\mu P_t, \nu P_t) \\ &\leq \sum_{w,z} \Pi_{\mu,\nu}(w, z) \frac{d}{dt} \Big|_{t=0} W_p^p(\delta_w P_t, \delta_z P_t) \\ &= \sum_{w,z} \Pi_{\mu,\nu}(w, z) p d(w, z)^{p-1} \frac{d}{dt} \Big|_{t=0} W_p(\delta_w P_t, \delta_z P_t) \\ &\leq -K p \sum_{w,z} \Pi_{\mu,\nu}(w, z) d(w, z)^p \\ &= -K p W_p^p(\mu, \nu), \end{aligned}$$

where we also used Corollary 5.16. Therefore,

$$\frac{d^+}{dt} \Big|_{t=0} W_p^p(\mu P_t, \nu P_t) \leq -K p W_p^p(\mu, \nu).$$

and Markovianity this extends to every $\bar{t} > 0$, i.e.

$$\frac{d^+}{dt} \Big|_{t=\bar{t}} W_p^p(\mu P_t, \nu P_t) \leq -K p W_p^p(\mu P_{\bar{t}}, \nu P_{\bar{t}}).$$

Noticing also that $t \rightarrow W_p^p(\mu P_t, \nu P_t)$ is continuous by Corollary 5.14, we can apply Lemma C.1 to conclude that

$$W_p^p(\mu P_t, \nu P_t) \leq e^{-K p t} W_p^p(\mu, \nu),$$

as desired. □

5.3 Comparison discrete and continuous time

There is a natural way to construct a continuous time Markov chain from a discrete time one: namely, for $\lambda > 0$ and a stochastic matrix P , the generator is defined by $L = \lambda(P - I)$. On the other hand, it is readily seen that, given a generator L , for any $\lambda > 0$ big enough there exists a corresponding stochastic matrix $P = I + \frac{1}{\lambda} L$ (recall we are assuming finiteness of the state space, so the entries of L are bounded).

Remark 5.22. Let P be a stochastic matrix and $\lambda > 0$. In view of Definition 5.11, we see that for $p = 1$ the $\text{Ric}_{\text{dc},1}$ and $\text{Ric}_{\text{cc},1}$ notions of curvature are essentially equivalent for the continuous time Markov chain with generator $L = \lambda(P - I)$ (and transition semigroup P_t) and the **lazy** discrete time Markov chain with transition matrix $\tilde{P} := \frac{I+P}{2}$. Indeed, it follows from the identity in (35) with $s = \frac{1}{2\lambda}$ that

$$K_{\text{dc},1}^{(\tilde{P})}(x, y) = \frac{1}{2\lambda} K_{\text{cc},1}(x, y), \quad (38)$$

where above $K_{\text{dc},1}^{(\tilde{P})}(x, y)$ is the curvature $K_{\text{dc},p}$ for the Markov chain with transition matrix \tilde{P} , not P . Notice that here it is fundamental to consider the lazy version with transition matrix \tilde{P} . To see why, consider the following simple example: the state space is the two-point space $\Omega = \{x, y\}$, the stochastic matrix is

$$P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

and $\lambda = 1$. Then we have $K_{\text{dc},1}^{(P)}(x, y) = 0$ for P , $K_{\text{dc},1}^{(\tilde{P})}(x, y) = 1$ for \tilde{P} and $K_{\text{cc},1}(x, y) = 2$ for $L = P - I$. Hence (38) is satisfied only when $K_{\text{dc},p}(x, y)$ is defined for \tilde{P} and not for P .

The next proposition shows that an analogue relation as the one described in the above remark holds true also for $\text{Ric}_{\text{dc},\infty}$ and $\text{Ric}_{\text{cc},\infty}$.

Proposition 5.23. Suppose $L = \lambda(P - I)$ for a stochastic matrix P and $\lambda > 0$. Then the following hold for any $x \neq y$:

1. $K_{\text{cc},\infty}(x, y) \geq \lambda K_{\text{dc},\infty}^{(P)}(x, y)$.
2. For the lazy Markov chain with transition $\tilde{P} = \frac{P+I}{2}$, we have $K_{\text{dc},\infty}^{(\tilde{P})}(x, y) \geq \frac{1}{2\lambda} K_{\text{cc},\infty}(x, y)$.

Proof. 1. Assume $K_{\text{dc},\infty}^{(P)}(x, y) \geq 0$, otherwise the claim is trivial. Let $\Pi(x, y, \cdot, \cdot)$ be an optimal coupling in the definition of $K_{\text{dc},\infty}^{(P)}(x, y)$ and define then the coupling rates $C(x, y, w, z) = \lambda \cdot \Pi(x, y, w, z)$. It is easy to check that these coupling rates are admissible and yield the first conclusion.

2. Again, assume $K_{\text{cc},\infty}(x, y) \geq 0$, otherwise the conclusion is trivial. Let $C(x, y, \cdot, \cdot)$ be optimal coupling rates in the definition of $K_{\text{cc},\infty}(x, y)$. Notice that for $L = \lambda(P - I)$ and $t = \frac{1}{2\lambda}$ we have

$$\tilde{P}_t = \tilde{P}_{\frac{1}{2\lambda}} = I + \frac{1}{2\lambda} \cdot \lambda(P - I) = \tilde{P}.$$

Therefore, since

$$\begin{aligned} \sum_{w,z} C(x, y, w, z) &= C(x, y, x, y) + \sum_{w,z:(w,z) \neq (x,y)} C(x, y, w, z) \\ &\leq C(x, y, x, y) + \sum_{w,z:w \neq x} C(x, y, w, z) + \sum_{w,z:z \neq y} C(x, y, w, z) \\ &= C(x, y, x, y) + \sum_{w \neq x} Q(x, w) + \sum_{z \neq y} Q(y, z) \\ &\leq C(x, y, x, y) + 2\lambda, \end{aligned}$$

we can apply Lemma 5.20 to obtain coupling $\Pi(x, y, \cdot, \cdot)$ for $(\delta_x \tilde{P}, \delta_y \tilde{P})$. It is easy to check that this coupling proves that $K_{\text{dc},\infty}^{(\tilde{P})}(x, y) \geq \frac{1}{2\lambda} K_{\text{cc},\infty}(x, y)$ for the lazy Markov chain \tilde{P} . \square

5.4 Applications and related problems

Recalling Definition 5.18 and Proposition 5.21, we see that for a continuous time Markov chain on a finite state space Ω we have that $\text{Ric}_{\text{cc},\infty} \geq K > 0$ if and only if for all *neighbouring* states $x \sim y \in \Omega$ there exist coupling rates $C(x, y, \cdot, \cdot)$ satisfying

- $\sum_{v,z \in \Omega} C(x,y,v,z) \mathbb{1}_{d(v,z) > 1} = 0$;
- $\sum_{v,z \in \Omega} C(x,y,v,z) \mathbb{1}_{d(v,z) = 0} \geq K$.

Bearing this mind, we can see that in the examples of Section 4 the constructed coupling rates immediately yield positive $\text{Ric}_{\text{cc},\infty}$ curvature: more precisely, we have the following result.

Theorem 5.24. *The following hold:*

- Under the assumptions of Theorem 4.1, we have $\text{Ric}_{\text{cc},\infty} \geq \kappa_*$ for Glauber dynamics.
- Under the assumptions of Theorem 4.8, we have $\text{Ric}_{\text{cc},\infty} \geq L$ for the Bernoulli–Laplace model.
- Under the assumptions of Theorem 4.12, we have $\text{Ric}_{\text{cc},\infty} \geq \kappa_*$ for the hardcore model.

In particular, by Proposition 5.21, under the respective theorems’ assumptions for all probability measures $\mu, \nu \in \mathcal{P}(\Omega)$, $p \geq 1$ and $t \geq 0$ we have

$$W_p(\mu P_t, \nu P_t) \leq e^{-\frac{\kappa_*}{p} t} W_p(\mu, \nu), \quad (39)$$

with $K = \begin{cases} \kappa_* & \text{for Glauber dynamics,} \\ L & \text{for the Bernoulli–Laplace model,} \\ \kappa_* & \text{for the hardcore model.} \end{cases}$

We remark here that the estimate (39) was already established in [5] only for the specific case of interacting random walks on the grid of Section 4.4 (see Theorem 3.2 of [5]), which doesn’t follow directly from the arguments of this section since we restricted our discussion to finite state space Markov chains.

We also remark that Theorem 5.24 shows that, for some Markov chains, assumptions that are strictly connected to positive $\text{Ric}_{\text{cc},\infty}$ curvature are useful for establishing the modified log-Sobolev inequality, positive entropic curvature, positive discrete Bakry–Émery curvature and other related inequalities (cf. the discussion in Section 2). This suggests interesting connections with some other open problems in the theory of functional inequalities and discrete curvature for Markov chains, as we discuss next.

Peres–Tetali Conjecture One notable example is the following important unpublished conjecture by Peres and Tetali, which links coarse Ricci curvature to the modified log-Sobolev inequality in the setting of lazy simple random walks on finite graphs (see also [7, Con. 3.1], [13, Con. 4], [1, Rmk. 1.1]).

Conjecture 5.25. *There exists a universal constant $\alpha > 0$ such that the following holds. Let Ω be a finite unweighted graph and consider the stochastic matrix P associated with the simple random walk on this graph, $\tilde{P} = \frac{P+I}{2}$ associated to the lazy simple random walk and the generator $L = P - I$. If $\text{Ric}_{\text{dc},1} \geq K > 0$ for the lazy stochastic matrix \tilde{P} (or, equivalently, if $\text{Ric}_{\text{cc},1} \geq 2K$ for L), then $\text{MLSI}(\alpha K)$ holds.*

In all the examples of Theorem 5.24 we have positive $\text{Ric}_{\text{cc},\infty}$ curvature, which implies in particular positive $\text{Ric}_{\text{cc},1}$ curvature by Proposition 5.21. Therefore, it is natural to study the following problem related to the above conjecture: assuming a strictly positive lower bound of $\text{Ric}_{\text{cc},\infty}$ (and under some additional assumptions), is it possible to deduce a lower bound of the same order for the MLSI constant? In particular, if the additional assumptions are that we are in the setting of simple random walks on finite graphs, this problem constitutes a weaker form of the Peres–Tetali Conjecture.

Coarse and entropic curvature Another important open problem consists in comparing the different notions of discrete curvature. For example, it is not known when a positive lower bound for the coarse Ricci curvature implies a positive lower bound of the same order for the entropic curvature, or vice versa (and similarly for the discrete Bakry–Émery curvature). In light of the results of this paper, the following is also a natural question: assuming a strictly positive lower bound of $\text{Ric}_{\text{cc},\infty}$ (and under some additional assumption), is it possible to deduce a lower bound of the same order for the entropic curvature? Interestingly, we remark that positive lower bounds for the entropic curvature are linked to exponential contraction with respect to the metric \mathscr{W} . Precisely, if $\text{Ric}_e \geq K$ then

$$\mathscr{W}(\mu P_t, \nu P_t) \leq e^{-Kt} \mathscr{W}(\mu, \nu)$$

for all $t \geq 0$ and starting probability measures μ, ν (see [11, Prop 4.7] and cf. Proposition 2.1). This is of course reminiscent of the exponential decay of (39), which follows from $\text{Ric}_{\text{cc},\infty} \geq K$. However, not much is known about the relationship between \mathscr{W} and W_p (see [11, Prop. 2.12, 2.14] for a lower(/upper) bound of \mathscr{W} in terms of W_1 (/ W_2)), so it is not clear how these exponential decay estimates are connected.

A Proof of Proposition 2.2

Here, we work in the setting of Section 5 and with that notation; recall in particular that the state space is finite and the Markov chain is irreducible and reversible.

Proof. Suppose Ω is finite and Assumption 2 is satisfied and that (8) holds for all $\rho: \Omega \rightarrow \mathbb{R}_{>0}$ and $\psi: \Omega \rightarrow \mathbb{R}$. Fix now $\rho: \Omega \rightarrow \mathbb{R}_{>0}$ and choose $\psi = \phi' \circ \rho$. Notice then that

$$\mathcal{A}(\rho, \phi' \circ \rho) = \frac{1}{2} \sum_{x,y:\psi(x) \neq \psi(y)} \pi(x)Q(x,y) \frac{\rho(x) - \rho(y)}{(\phi' \circ \rho)(x) - (\phi' \circ \rho)(y)} [(\phi' \circ \rho)(x) - (\phi' \circ \rho)(y)]^2 = \mathcal{E}(\rho, \phi' \circ \rho). \quad (40)$$

Moreover if $\psi(x) \neq \psi(y)$ we have that

$$\nabla\theta(\rho(x), \rho(y)) = \frac{1}{\psi(x) - \psi(y)} [1 - \phi''(\rho(x)) \theta(\rho(x), \rho(y)), -1 + \phi''(\rho(y)) \theta(\rho(x), \rho(y))].$$

It follows that

$$\begin{aligned} \mathcal{E}(\rho, \phi' \circ \rho) &= \frac{1}{4} \sum_{x,y} \pi(x)Q(x,y) \left\{ [1 - \phi''(\rho(x)) \theta(\rho(x), \rho(y)), -1 + \phi''(\rho(y)) \theta(\rho(x), \rho(y))] \cdot \begin{pmatrix} L\rho(x) \\ L\rho(y) \end{pmatrix} \right\} \\ &\quad \cdot [(\phi' \circ \rho)(x) - (\phi' \circ \rho)(y)] \\ &= \frac{1}{2} \mathcal{E}(L\rho, \phi' \circ \rho) - \frac{1}{2} \mathcal{E}(\rho, (\phi'' \circ \rho) \cdot L\rho) \\ &= \frac{1}{2} \mathcal{E}(\rho, L(\phi' \circ \rho)) - \frac{1}{2} \mathcal{E}(\rho, (\phi'' \circ \rho) \cdot L\rho) \\ \mathcal{D}(\rho, \phi' \circ \rho) &= \frac{1}{2} \sum_{x,y} \pi(x)Q(x,y) (\rho(x) - \rho(y)) (L(\phi' \circ \rho)(x) - L(\phi' \circ \rho)(y)) = \mathcal{E}(\rho, L(\phi' \circ \rho)). \end{aligned}$$

Therefore

$$\mathcal{B}(\rho, (\phi' \circ \rho)) = -\frac{1}{2} \mathcal{E}[\rho, (\phi'' \circ \rho) \cdot L\rho + L(\phi' \circ \rho)].$$

From this, letting $\rho_t = P_t \rho$, we see that

$$\left. \frac{d}{dt} \right|_{t=0} \mathcal{E}(\rho_t, \phi' \circ \rho_t) = -2\mathcal{B}(\rho, \phi' \circ \rho).$$

Therefore the inequality $\mathcal{B}(\rho, \phi' \circ \rho) \geq K\mathcal{A}(\rho, \phi' \circ \rho)$ is equivalent to inequality (10), from which $\text{CSI}_\phi(2K)$ follows. \square

Remark A.1. In the particular case when θ is the logarithmic mean, for any constant $Z > 0$ we have $\mathcal{B}(Z \cdot \rho, \psi) = Z\mathcal{B}(\rho, \psi)$ and $\mathcal{A}(Z \cdot \rho, \psi) = Z\mathcal{A}(\rho, \psi)$, so it suffices to consider the case where ρ is a density with respect to π when proving inequality (8).

B Proof of Proposition 2.4

In this section we compute the parameter M_θ defined in (7) for the weight function θ_α , which was defined in (9) with $\phi = \phi_\alpha$ as in (4), corresponding to Beckner functionals. In other words, we have

$$\theta(s, t) = \frac{\alpha - 1}{\alpha} \frac{s - t}{s^{\alpha-1} - t^{\alpha-1}}$$

for $s \neq t > 0$ and $\theta(s, s) = \frac{1}{\alpha} s^{2-\alpha}$.

Proof. Recall the definition of

$$M_\theta := \inf_{\substack{s,t \geq 0: \\ \theta(s,t) > 0}} \frac{\theta(s,s) + \theta(t,t)}{2\theta(s,t)} \in [0, 1].$$

For $\alpha = 2$ the result is trivial, while for $\alpha = 1$ it follows from equation (2.1) in [11]. Suppose hence that $\alpha \in (1, 2)$: without loss of generality we can minimize over $s > t > 0$, so that substituting the expression of θ_α from (12) the problem reduces to computing

$$\begin{aligned} M_{\theta_\alpha} &= \inf_{s>t>0} \frac{1}{2(\alpha-1)} \frac{(s^{2-\alpha} + t^{2-\alpha}) \cdot (s^{\alpha-1} - t^{\alpha-1})}{s-t} \\ &= \frac{1}{2(\alpha-1)} \left\{ 1 + \inf_{s>t>0} \frac{s^{\alpha-1}t^{2-\alpha} - s^{2-\alpha}t^{\alpha-1}}{s-t} \right\} \\ &= \frac{1}{2(\alpha-1)} \left\{ 1 + \inf_{\lambda>1} \frac{\lambda^{\alpha-1} - \lambda^{2-\alpha}}{\lambda-1} \right\} \end{aligned} \quad (41)$$

where we set $\lambda := \frac{s}{t} > 1$. If $\alpha \geq \frac{3}{2}$, the conclusion follows by noticing that $\frac{\lambda^{\alpha-1} - \lambda^{2-\alpha}}{\lambda-1} \geq 0$ for $\lambda > 1$ and by letting $\lambda \rightarrow \infty$. Suppose hence now that $\alpha \in (1, \frac{3}{2})$: to conclude that in this case $M_{\theta_\alpha} = 1$ it is enough to show that for $\lambda > 1$

$$\frac{\lambda^{\alpha-1} - \lambda^{2-\alpha}}{\lambda-1} \geq 2\alpha - 3. \quad (42)$$

Notice that equality holds as $\lambda \rightarrow 1^+$. By density of \mathbb{Q} in \mathbb{R} and by a continuity argument, it suffices to show that for all $\lambda > 1$ and all *even* integers $p, s \in \mathbb{N}$ with $p > s$ we have that

$$\frac{\lambda^{\frac{s}{p+s}} - \lambda^{\frac{p}{p+s}}}{\lambda-1} \geq -\frac{p-s}{p+s},$$

where we used the substitution $\alpha = 1 + \frac{s}{p+s}$. Rearranging this and renaming $\lambda \leftarrow \lambda^{\frac{1}{p+s}}$ we need to prove equivalently that for all $\lambda > 1$

$$\frac{\lambda^{p+s} - 1}{p+s} \geq \frac{\lambda^p - \lambda^s}{p-s}.$$

Equivalently, dividing both sides by $\lambda - 1$ and denoting by AM the arithmetic mean, we need to prove that

$$\text{AM}(1, \dots, \lambda^{p+s-1}) \geq \text{AM}(\lambda^s, \dots, \lambda^{p-1}).$$

This last inequality holds true, since for $\lambda > 1$ and integers $0 < i < j$ we have that

$$\lambda^{i-1} + \lambda^{j+1} \geq \lambda^i + \lambda^j$$

by the classical rearrangement inequality. □

C Grönwall's lemma with Dini derivative

For a continuous function $f: I \rightarrow \mathbb{R}$, where $I \subset \mathbb{R}$ is an interval, we consider the *upper/lower Dini derivative*, which are defined respectively by

$$\begin{aligned} \frac{d^+}{dt} \Big|_{t_0} f(t) &= \limsup_{h \rightarrow 0^+} \frac{f(t+h) - f(t)}{h}, \\ \frac{d^-}{dt} \Big|_{t_0} f(t) &= \liminf_{h \rightarrow 0^+} \frac{f(t+h) - f(t)}{h}. \end{aligned}$$

Clearly, $\frac{d^-}{dt} f(t) \leq \frac{d^+}{dt} f(t)$. It may be useful to apply Grönwall's lemma in absence of differentiability, by considering instead the Dini derivatives. In particular, the following variant holds.

Lemma C.1 (Grönwall's lemma). *Let $I = [a, b) \subset \mathbb{R}$ be an interval, with $a < b \leq \infty$, and consider real valued continuous functions $u, \beta: I \rightarrow \mathbb{R}$. Suppose that for all $t \in I$*

$$\frac{d^-}{dt} u(t) \leq \beta(t)u(t). \quad (43)$$

Then for all $t \in I$

$$u(t) \leq \exp\left(\int_0^t \beta(s)ds\right)u(a).$$

Proof. Set $v(t) = \exp\left(-\int_0^t \beta(s)ds\right)$, which satisfies $v(t) > 0$, $v'(t) = -\beta(t)v(t)$. Notice that for all $t \in I$

$$\begin{aligned} \frac{d^-}{dt}[u(t)v(t)] &= \liminf_{h \rightarrow 0^+} \frac{u(t+h)v(t+h) - u(t)v(t)}{h} + \frac{u(t)[v(t+h) - v(t)]}{h} \\ &= \left[\frac{d^-}{dt}u(t) \right] v(t) + u(t)v'(t) \\ &\leq \beta(t)u(t)v(t) - u(t)\beta(t)v(t) \\ &= 0. \end{aligned}$$

Next, we notice as in [6, Eqn. (3.8)] that this implies that the continuous function $u(t)v(t)$ is non-increasing on I . In particular, we have $u(t)v(t) \leq u(a)$, which implies the thesis by substituting the expression for $v(t)$ and rearranging. \square

References

- [1] Antonio Blanca, Pietro Caputo, Zongchen Chen, Daniel Parisi, Daniel Štefankovič, and Eric Vigoda. “On mixing of Markov chains: coupling, spectral independence, and entropy factorization”. English. In: *Electron. J. Probab.* 27 (2022). Id/No 142, p. 42. ISSN: 1083-6489. DOI: 10.1214/22-EJP867.
- [2] Sergey G. Bobkov and Prasad Tetali. “Modified logarithmic Sobolev inequalities in discrete settings”. English. In: *J. Theor. Probab.* 19.2 (2006), pp. 289–336. ISSN: 0894-9840. DOI: 10.1007/s10959-006-0016-3. URL: citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.124.6483.
- [3] D. P. Bourne, D. Cushing, S. Liu, F. Münch, and N. Peyerimhoff. “Ollivier–Ricci Idleness Functions of Graphs”. In: *SIAM Journal on Discrete Mathematics* 32.2 (2018), pp. 1408–1424. DOI: 10.1137/17M1134469. eprint: <https://doi.org/10.1137/17M1134469>. URL: <https://doi.org/10.1137/17M1134469>.
- [4] Pietro Caputo, Paolo Dai Pra, and Gustavo Posta. “Convex entropy decay via the Bochner–Bakry–Emery approach”. English. In: *Ann. Inst. Henri Poincaré, Probab. Stat.* 45.3 (2009), pp. 734–753. ISSN: 0246-0203. DOI: 10.1214/08-AIHP183.
- [5] Giovanni Conforti. “A probabilistic approach to convex (ϕ) -entropy decay for Markov chains”. English. In: *Ann. Appl. Probab.* 32.2 (2022), pp. 932–973. ISSN: 1050-5164. DOI: 10.1214/21-AAP1700.
- [6] Sara Daneri and Giuseppe Savaré. “Eulerian Calculus for the Displacement Convexity in the Wasserstein Distance”. In: *SIAM Journal on Mathematical Analysis* 40.3 (2008), pp. 1104–1122. DOI: 10.1137/08071346X. eprint: <https://doi.org/10.1137/08071346X>. URL: <https://doi.org/10.1137/08071346X>.
- [7] Ronen Eldan, James R. Lee, and Joseph Lehec. “Transport-entropy inequalities and curvature in discrete-space Markov chains”. In: *A journey through discrete mathematics*. Springer, 2017, pp. 391–406.
- [8] Matthias Erbar and Max Fathi. “Poincaré, modified logarithmic Sobolev and isoperimetric inequalities for Markov chains with non-negative Ricci curvature”. In: *Journal of Functional Analysis* 274.11 (2018), pp. 3056–3089. ISSN: 0022-1236. DOI: <https://doi.org/10.1016/j.jfa.2018.03.011>.
- [9] Matthias Erbar, Christopher Henderson, Georg Menz, and Prasad Tetali. “Ricci curvature bounds for weakly interacting Markov chains”. In: *Electronic Journal of Probability* 22.none (2017), pp. 1–23. DOI: 10.1214/17-EJP49. URL: <https://doi.org/10.1214/17-EJP49>.
- [10] Matthias Erbar and Jan Maas. “Gradient flow structures for discrete porous medium equations”. In: *Discrete and Continuous Dynamical Systems* 34.4 (2014), pp. 1355–1374.
- [11] Matthias Erbar and Jan Maas. “Ricci curvature of finite Markov chains via convexity of the entropy”. English. In: *Arch. Ration. Mech. Anal.* 206.3 (2012), pp. 997–1038. ISSN: 0003-9527. DOI: 10.1007/s00205-012-0554-z.
- [12] Matthias Erbar, Jan Maas, and Prasad Tetali. “Discrete Ricci curvature bounds for Bernoulli–Laplace and random transposition models”. English. In: *Ann. Fac. Sci. Toulouse, Math. (6)* 24.4 (2015), pp. 781–800. ISSN: 0240-2963. DOI: 10.5802/afst.1464. URL: citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.740.3480.
- [13] Max Fathi. “Quelques applications du transport optimal en analyse et en probabilités”. Habilitation à diriger des recherches. Université Paul Sabatier (Toulouse 3), Apr. 2019. URL: <https://hal.science/tel-02105311>.
- [14] Max Fathi and Jan Maas. “Entropic Ricci curvature bounds for discrete interacting systems”. English. In: *Ann. Appl. Probab.* 26.3 (2016), pp. 1774–1806. ISSN: 1050-5164. DOI: 10.1214/15-AAP1133.

- [15] Max Fathi and Yan Shu. “Curvature and transport inequalities for Markov chains in discrete spaces”. In: *Bernoulli* 24.1 (2018), pp. 672–698. DOI: 10.3150/16-BEJ892. URL: <https://doi.org/10.3150/16-BEJ892>.
- [16] Ansgar Jüngel and Wen Yue. “DISCRETE BECKNER INEQUALITIES VIA THE BOCHNER–BAKRY–EMERY APPROACH FOR MARKOV CHAINS”. In: *The Annals of Applied Probability* 27.4 (2017), pp. 2238–2269. ISSN: 10505164. URL: <http://www.jstor.org/stable/26361546>.
- [17] Yong Lin, Linyuan Lu, and Shing-Tung Yau. “Ricci curvature of graphs”. In: *Tohoku Mathematical Journal* 63.4 (2011), pp. 605–627. DOI: 10.2748/tmj/1325886283. URL: <https://doi.org/10.2748/tmj/1325886283>.
- [18] John Lott and Cedric Villani. “Ricci curvature for metric-measure spaces via optimal transport”. English. In: *Ann. Math. (2)* 169.3 (2009), pp. 903–991. ISSN: 0003-486X. DOI: 10.4007/annals.2009.169.903. URL: annals.math.princeton.edu/annals/2009/169-3/p04.xhtml.
- [19] Jan Maas. “Entropic Ricci curvature for discrete spaces”. In: *Modern approaches to discrete curvature*. Vol. 2184. Lecture Notes in Math. Springer, Cham, 2017, pp. 159–174. ISBN: 978-3-319-58001-2.
- [20] Jan Maas. “Gradient flows of the entropy for finite Markov chains”. English. In: *J. Funct. Anal.* 261.8 (2011), pp. 2250–2292. ISSN: 0022-1236. DOI: 10.1016/j.jfa.2011.06.009.
- [21] Alexander Mielke. “Geodesic convexity of the relative entropy in reversible Markov chains”. English. In: *Calc. Var. Partial Differ. Equ.* 48.1-2 (2013), pp. 1–31. ISSN: 0944-2669. DOI: 10.1007/s00526-012-0538-8.
- [22] Florentin Münch and Radosław K. Wojciechowski. “Ollivier Ricci curvature for general graph Laplacians: heat equation, Laplacian comparison, non-explosion and diameter bounds”. English. In: *Adv. Math.* 356 (2019). Id/No 106759, p. 45. ISSN: 0001-8708. DOI: 10.1016/j.aim.2019.106759.
- [23] Yann Ollivier. “A survey of Ricci curvature for metric spaces and Markov chains”. English. In: *Probabilistic approach to geometry. Proceedings of the 1st international conference, Kyoto, Japan, 28th July – 8th August, 2008*. Tokyo: Mathematical Society of Japan (MSJ), 2010, pp. 343–381. ISBN: 978-4-931469-58-7.
- [24] Yann Ollivier. “Ricci curvature of Markov chains on metric spaces”. English. In: *J. Funct. Anal.* 256.3 (2009), pp. 810–864. ISSN: 0022-1236. DOI: 10.1016/j.jfa.2008.11.001.
- [25] Paolo Dai Pra and Gustavo Posta. “Entropy decay for interacting systems via the Bochner-Bakry-Émery approach”. English. In: *Electron. J. Probab.* 18 (2013). Id/No 52, p. 21. ISSN: 1083-6489. DOI: 10.1214/EJP.v18-2041.
- [26] Michael Schmuckenschläger. “Curvature of nonlocal Markov generators”. English. In: *Convex geometric analysis*. Cambridge: Cambridge University Press, 1999, pp. 189–197. ISBN: 0-521-64259-0.
- [27] Karl-Theodor Sturm. “On the geometry of metric measure spaces”. In: *Acta Mathematica* 196.1 (2006), pp. 65–131. DOI: 10.1007/s11511-006-0002-8. URL: <https://doi.org/10.1007/s11511-006-0002-8>.
- [28] Laurent Veysseire. *Coarse Ricci curvature for continuous-time Markov processes*. 2012. arXiv: 1202.0420 [math.PR].
- [29] Cédric Villani. *Optimal transport. Old and new*. English. Vol. 338. Grundlehren Math. Wiss. Berlin: Springer, 2009. ISBN: 978-3-540-71049-3. DOI: 10.1007/978-3-540-71050-9.