

Absolute continuity, supports and idempotent splitting in categorical probability

Tobias Fritz¹, Tomáš Gonda², Antonio Lorenzin¹,
Paolo Perrone³, and Dario Stein⁴

¹*Department of Mathematics, University of Innsbruck, Austria*

²*Institute for Theoretical Physics, University of Innsbruck, Austria*

³*Department of Computer Science, University of Oxford, United Kingdom*

⁴*iHub, Radboud University Nijmegen, The Netherlands*

September 7, 2023

Abstract

Markov categories have recently turned out to be a powerful high-level framework for probability and statistics. They accommodate purely categorical definitions of notions like conditional probability and almost sure equality, as well as proofs of fundamental results such as the Hewitt–Savage 0/1 Law, the de Finetti Theorem and the Ergodic Decomposition Theorem.

In this work, we develop additional relevant notions from probability theory in the setting of Markov categories. This comprises improved versions of previously introduced definitions of absolute continuity and supports, as well as a detailed study of idempotents and idempotent splitting in Markov categories. Our main result on idempotent splitting is that every idempotent measurable Markov kernel between standard Borel spaces splits through another standard Borel space, and we derive this as an instance of a general categorical criterion for idempotent splitting in Markov categories.

Contents

1	Introduction	2
2	Absolute continuity	7
2.1	An extended definition of almost sure equality	7
2.2	Definition and examples of absolute continuity	10
2.3	Properties of absolute continuity	14
2.4	Absolute bicontinuity	16
3	Supports	17
3.1	Definition and basic properties of supports	18
3.2	Support completeness	24
3.3	The input-output relation functor	31
3.4	Categories of statistical models	35
3.5	The equalizer principle	37
3.6	Split supports	39


4	Idempotents and their splitting in Markov categories	45
4.1	Static, strong and balanced idempotents	46
4.2	Split idempotents in Markov categories	52
4.3	Supports and splittings of static idempotents	55
4.4	General conditions for the splitting of idempotents	57
4.5	The Blackwell envelope	63
A	Background on Markov categories	65
A.1	Some basics	65
A.2	Parametric Markov categories	67
A.3	Some auxiliary results on almost sure equality	70
A.4	Observationally representable Markov categories	74
A.5	The Cauchy–Schwarz Axiom	78
B	An isomorphism of measure spaces which does not preserve supports	80

1 Introduction

In recent years, there has been growing interest in categorical methods for probability theory and related fields. Structures such as *garbage-share monoidal categories* [20] (also called *copy-discard categories* [8]) and *Markov categories* [11] have appeared in contexts as diverse as mathematical statistics [18], graphical models [12], ergodic theory [29], machine learning [40], and information theory [36].


The common theme in all these works is the aim distill the most conceptual aspects of reasoning in the face of uncertainty, such as stochastic independence or the composition of Markov kernels, and to prove theorems in probability theory in terms of these prime principles. The hope is to achieve both greater conceptual clarity and generality. There are several aspects of probability theory that lend themselves very well to this point of view. The idea of forming *conditional distributions* and *Bayesian inverses* can be expressed categorically and even graphically [8], and proofs in probability theory involving conditioning can be simplified by resorting to this abstract definition. This facilitated for example the categorical proof of de Finetti’s theorem [15], as well as a generalization of Blackwell’s theorem on the comparison of statistical experiments to the case of nondiscrete random variables [18]. Similarly, a categorical notion of *determinism* has allowed us to formulate and prove certain *zero-one laws* [14], and can be seen to be the main idea behind the concept of *ergodicity* [29]. Moreover, measuring the deviation from determinism can lead to the notion of *entropy* [36]. Last but not least, the aforementioned idea of stochastic dependence and independence can be refined to a rich theory of conditional independence and Markov separation properties that reduces stochastic dependencies to topological path-connectedness in a graph [12].

Supports. One of the conceptual aspects of probabilistic processes that we address in this work is the notion of *support*. Roughly speaking, one could define the support of a probability measure as the smallest set of probability 1, if it exists. For example, the probability distribution p on the set $\{a, b, c\}$ given by




has as support the subset $\{a, b\}$. For distributions on discrete spaces, this definition is appropriate, and the support consists exactly of the points with non-zero probability.

In the case of continuous sample spaces, points may have measure zero while still being in the support — this is the case, for example, for the normal distribution:



In this case, one can say that a point $x_0 \in X$ lies in the support of a probability measure if every open neighborhood U of x_0 has non-zero measure. This is equivalent to saying that the support is the smallest *closed* subset of full measure. Such an approach, which we call the *topological support* (Example 3.1.5), works well for (τ -smooth) probability measures on topological spaces [16].

On a general measurable space, there are no closed sets, and so this approach cannot work. Moreover, even in those measurable spaces which do come from a topological space, such as real numbers, using a topological notion in a measure-theoretic context is not going to be invariant under isomorphism. For example, we can equip the unit interval with the following two probability measures:



Topologically, it is clear that the first measure has full support, and the second measure is supported on $[0, 1/2]$. However, as measure spaces, the two are isomorphic, as we show in Theorem B.0.1. In particular, no invariant of measure spaces can tell the two situations apart. Intuitively, the reason is that measurable functions alone cannot tell apart a region of zero measure which is outside the (topological) support, such as the half-interval $(1/2, 1]$ on the right-hand side, from a region within the topological support, but which does not fill the space enough to have positive measure (see Appendix B for details). On this note, we may wonder whether such probability measures should even admit a support. In fact, the existence of a support is deeply connected to the notion of atomicity (Corollary 3.2.15 and Example 3.2.16). In particular, non-atomic measures like the two described in (1.1) will never admit a support.

In this work, we provide a categorical perspective on supports. Both the topological and measure-theoretic perspectives on supports are reflected in the general categorical notion. It is not obvious how to define a notion of support in categorical terms, since abstracting away from the concrete category of measurable spaces and Markov kernels deprives one of the possibility to talk about sets of measure zero. The way we deal with such situations abstractly is via the categorical notion of *almost sure equality* [11, Section 13], which we discuss in more detail in Section 2.1 of the present paper. This concept reduces to the usual almost sure equality of functions and kernels when instantiated in the category $\mathbf{BorelStoch}$ of standard Borel spaces and Markov kernels, and can be used conveniently in statements and proofs. For example, conditionals and Bayesian inverses, when they exist, are unique up to almost sure equality in any Markov category (Lemma A.3.4), and this is an immediate consequence of the definition.

What could then be a categorical definition of support? In discrete probability theory, two functions on a probability space are almost surely equal if and only if they agree on the support of the probability measure. One can turn this idea into a categorical definition of support via the universal property of *representing* almost sure equality in the sense of representable functors. This was suggested in [11, Definition 13.20], and then used in [6] and [7]. Similar ideas were given in [10] as well.

Here we expand on that idea, but we also give an alternative and weaker definition of support. The notion from the previous paragraph—which we rename to *split support* in this paper—is a “mapping-out” universal property, i.e. it is about representing a *covariant* functor to \mathbf{Set} . We can also define supports via a “mapping-in” universal property, representing a *contravariant* functor to \mathbf{Set} . The latter notion is called a *support* in the present article (Definition 3.1.2), and we show that it is strictly weaker than that of a split support (Proposition 3.6.14). Concrete instantiations of the resulting theory are the following:

- In the category $\mathbf{FinStoch}$ of finite sets and stochastic matrices, all supports exist (Example 3.1.4), and the support of every morphism with non-empty domain is split (Example 3.6.4).
- In the category $\mathbf{TychStoch}$ of Tychonoff topological spaces and continuous Markov kernels from Example A.1.4, all supports exist and they correspond exactly to the traditional topological supports (Example 3.1.5). Note that Tychonoff topological spaces include all metric spaces and all compact Hausdorff spaces. However, these supports do not split in general (Proposition 3.6.14).
- For measure-theoretic probability theory, i.e. in $\mathbf{BorelStoch}$, a probability measure has a support if and only if it is atomic, and similarly for Markov kernels (Theorem 3.2.7). In particular, the Lebesgue measure on the unit interval has no support in our sense.

The second item in particular shows that our newly developed mapping-in universal property of support seems to be the right notion to use in general. The third item is what one should expect in light of isomorphisms like the one of (1.1) above.

Another weaker property that has a similar flavour is a condition that we call the *equalizer principle* (Section 3.5). It states that if two deterministic morphisms agree p -almost surely, then p factors through their equalizer. If supports exist, this simply means that the support of p is included in the equalizer, but it makes sense and is useful even when p has no support. This property always holds in $\mathbf{BorelStoch}$, and it is crucial for our results about idempotents in Section 4.

Absolute continuity. The mapping-in universal property of supports discussed above is based on the idea that a Markov kernel lands on the support of a measure if and only if it is absolutely continuous with respect to that measure. In light of this conceptualization, Section 2 is devoted to the notion of *absolute continuity* (Definition 2.2.1). This definition improves upon our earlier definition given in [18, Definition 2.8] by allowing for extra input wires. As we show, in the general categorical setting this new definition ensures the compatibility with tensor products (Proposition 2.3.6), which the previous definition does not (Remark 2.3.7). Moreover, this abstract absolute continuity reduces to the usual measure-theoretic notion when instantiated in \mathbf{Stoch} (Example 2.2.6), and it corresponds to the inclusion of supports whenever they exist.

Idempotents. The other main theme of this paper is that of *idempotents*. While Section 3 on supports makes use of concepts and results developed in Section 2, Section 4 on idempotents can be read independently of the other two for the most part.

To say that an endomorphism $e: X \rightarrow X$ is an idempotent means that it satisfies $ee = e$. Idempotent Markov kernels secretly come up in several technical aspects of probability theory and ergodic theory. For example, conditional expectations can be thought of as idempotent kernels acting on bounded measurable functions by precomposition. In the study of ergodic theorems, forming the long-term average of a process produces an idempotent kernel [3, p. 560].

Given an idempotent $e: X \rightarrow X$, it is typically of interest to know whether e *splits* [5]. Such a splitting is given by another object T together with morphisms

$$\iota: T \hookrightarrow X, \quad \pi: X \twoheadrightarrow T$$

satisfying

$$\pi \iota = \text{id}_T, \quad \iota \pi = e.$$

When this is the case, we may think of e as projecting via π onto the subobject given by ι . Conversely, whenever we have morphisms ι and π such that $\pi \iota = \text{id}_T$, then the other composition $\iota \pi$ is an idempotent, which we think of as the projection onto T . For example, split idempotents in this form arise from the split supports mentioned above: A split support is a subobject onto which one can project.

As it turns out, there are different ways in which an idempotent can interact with the Markov category structure (see Figure 1 in Section 4), and this is related to the problem of splitting. A particularly important role is played by *balanced idempotents*, which are defined as satisfying a form of detailed balance (Remark 4.1.11). About these, we prove that:

- Every split idempotent in a positive Markov category, such as `BorelStoch`, is balanced (Section 4.2).
- Under suitable extra conditions on a positive Markov category, every idempotent is balanced and splits (Section 4.4).
- The free completion under balanced idempotents of a Markov category is again a Markov category (Section 4.5).

The extra conditions for the second item are the aforementioned equalizer principle, and the fact that the Markov category is *observationally representable*. “Representable” roughly means that we are in the Kleisli category of a probability monad (see [18] for more details). “Observational” means that morphisms into a space of distributions PX can be distinguished by taking repeated independent samples, as it is in ordinary probability (and as it reflects common practice in statistics and in all of science). The mathematical details of this idea are outlined in Appendix A.4. Since `BorelStoch` is positive and satisfies these extra conditions, we obtain our main result about measure-theoretic probability:

Theorem. *Every idempotent Markov kernel between standard Borel spaces is balanced and splits.*

We prove this as Corollary 4.4.5 and note that it strengthens a classical result of Blackwell on idempotent Markov kernels (Theorem 4.4.6). As far as we know, this strengthening is new, and in particular no measure-theoretic proof exists to date. We also note the analogous result that idempotents in the category of compact Hausdorff spaces and continuous Markov kernels are balanced and split (Corollary 4.4.10).

Outline

We start in Section 2 with an extended notion of almost sure equality (Definition 2.1.1) that appears subsequently in our new definition of absolute continuity between morphisms in a Markov category (Definition 2.2.1). We then show that tensor product and post-composition respect absolute continuity (Lemma 2.3.2 and Proposition 2.3.6).

In Section 3, we define the support of a morphism p via a mapping-in universal property as the universal object classifying morphisms absolutely continuous with respect to p . We study supports in this sense in some detail, showing in particular that this recovers the traditional notion in $\mathbf{FinStoch}$ (Example 3.1.4), as well as in categories of continuous Markov kernels between topological spaces (Example 3.1.5). We also show that a Markov category with all supports is causal if and only if its supports assemble into a functor (Theorem 3.1.17).

In Section 3.2, we establish which morphisms stand a chance to have a support at all (Lemma 3.2.1). We call morphisms satisfying this condition *atomic* (Definition 3.2.2). For $\mathbf{BorelStoch}$, we show that a Markov kernel is atomic in this sense if and only if it is atomic in the standard measure-theoretic sense. However, not even all atomic morphisms have a support, as Theorem 3.2.7 proves. We also show how to extend every causal Markov category to a *free support completion* — another causal Markov category in which every atomic morphism has a support (Theorem 3.2.14). Finding a good 2-categorical universal property for this construction remains an open question (Remark 3.2.18).

We give a first application of supports in Section 3.3. Under causality, supports and an additional assumption, we construct a functor from a Markov category to $\mathbf{SetMulti}$, the category of sets and multivalued functions in Section 3.3. Intuitively, the functor assigns to every morphisms its “possibilistic shadow” in the form of the *input-output relation*. In $\mathbf{FinStoch}$, every object is mapped to the set of deterministic states on it every stochastic matrix is sent to the relation of all input-output pairs with non-zero probability (Example 3.3.7).

As another application of supports, we show in Section 3.4 how they allow one to prove an equivalence between *parametric* statistical models and *non-parametric* statistical models. In particular, we define a category in which the objects, thought of as parametric statistical models, are morphisms in a suitable Markov category (Definition 3.4.1). We then show an abstract version of the statement that the behavior of a statistical model p can be identified with that of the collection of probability distributions that p parametrizes, provided that p has a support (Proposition 3.4.3).

In Section 3.5, we argue that even in the absence of supports, the weaker *equalizer principle* can be used as a substitute in certain situations. For example, it allows one to talk about the largest subobject on which two deterministic morphisms agree. This principle holds in Markov categories with supports (Remark 3.5.2), but also in categories such as $\mathbf{BorelStoch}$ (Proposition 3.5.4) in which generic morphisms do not have a support. Equalizer principle also plays an important role in our study of idempotents.

We finally turn to *split supports* in Section 3.6, which extends the notion of supports originally proposed in [11, Definition 13.20]. We introduce them as supports for which the support inclusion has a left inverse (Definition 3.6.2) and then describe their universal property in Proposition 3.6.9. The support of every stochastic matrix is in fact a split support (Example 3.6.4), and likewise all supports that exist in $\mathbf{BorelStoch}$ are split (Example 3.6.5).

We then turn to idempotents in Section 4. First, we introduce several types of interaction between idempotents and the structure of Markov categories (Definition 4.1.1). These give rise to *static*, *strong* and *balanced* idempotents respectively. We give examples of each kind and provide several reformulations of their definitions (Proposition 4.1.10 and Lemma 4.1.12). A simple but important observation is that in Markov categories satisfying the *Cauchy-Schwarz property* (Definition A.5.1), every idempotent is balanced (Theorem 4.1.8). This holds in particular for

any of the categories describing discrete or measure-theoretic probability (Corollary 4.1.9).

In Section 4.2, we study split idempotents and give an illuminating characterization above types of idempotents in terms of their splittings (Theorem 4.2.1). We also present a number of examples of split idempotents.

The interplay between the two main themes of the paper, supports and idempotents, is explored in Section 4.3. In particular, we prove that split supports are equivalently splittings of certain static idempotents (Theorem 4.3.5).

With the goal of showing that all idempotents of `BorelStoch` split, we study sufficient conditions for the splitting of balanced idempotents in Section 4.4. Our main result is Theorem 4.4.3 that establishes conditions under which all (balanced) idempotents in a Markov category split. It follows that every idempotent in `BorelStoch` splits (Corollary 4.4.5). This strengthens an existing result of Blackwell (Theorem 4.4.6).

Finally, in Section 4.5, we define the *Blackwell envelope* as the balanced idempotent completion that mimics the Karoubi envelope from ordinary category theory. We show that the Blackwell envelope of a Markov category is also a Markov category (Proposition 4.5.4), while the Karoubi envelope need not be.

In Appendix A, we provide relevant background material on Markov categories used in the rest of this work. We start with basic definitions and examples in Appendix A.1, and turn to the construction of parametric Markov categories (where morphisms have additional input wires) in Appendix A.2. The following two sections, Appendix A.3 and Appendix A.4, present novel material. In the former, we derive basic results on almost sure equality, which are used in Section 2. In the latter, we give some results on *observationally representable* Markov categories. In particular, every observationally representable Markov category is shown to be a.s.-compatibly representable (Proposition A.4.2). Finally, inspired by work of Parzygnat [33], we introduce a new information flow axiom named the *Cauchy-Schwarz Property* in Appendix A.5.

In Appendix B, we give an example of an isomorphism of measure spaces which does not preserve topological supports, making mathematically precise some of the ideas given in the introduction.

Acknowledgments

We thank Tommaso Russo for suggestions and references on the proof of Proposition 3.6.14, and Sean Moss for the pointers on observability. TF and AL acknowledge support from the Austrian Science Fund (FWF P 35992-N). PP acknowledges support from the European Research Council (grant “BLaSt – A Better Language for Statistics”). TG acknowledges support from the Austrian Science Fund (FWF) via the START Prize Y1261-N.

2 Absolute continuity

The goal of this short section is to introduce and study absolute continuity in an improved version and in more detail than was done in our earlier work [18], including the derivation of some properties relevant for later in the paper. Throughout, all our considerations take place in a Markov category \mathcal{C} (for the basics of the theory, see Appendix A.1 and the references therein).

2.1 An extended definition of almost sure equality

In the context of Markov categories, almost sure equality has been defined in [11, Definition 13.1]. In our improved notion of absolute continuity coming up as Definition 2.2.1, we use the

following notion of almost sure equality extended to morphisms with an extra input W .

Definition 2.1.1. Let $p: A \rightarrow X$ and $f, g: W \otimes X \rightarrow Y$. We say that f and g are **p -a.s. equal**, and write $f =_p g$, if we have

$$(2.1)$$

As in [11], we write A for the domain of p in order to indicate that we may think of p as a statistical model with parameter space A .

Remark 2.1.2. The extended notion of almost sure equality can be repackaged neatly in terms of parametric Markov categories (Appendix A.2). Indeed the above f and g can be interpreted as morphisms f and g of type $X \rightarrow Y$ in the parametric Markov category \mathbf{C}_W . It is easy to see that $f =_p g$ in \mathbf{C}_W is equivalent to Equation (2.1) in \mathbf{C} , and the equation $f =_p g$ is indeed the “old” definition of almost sure equality (without the extra wire W) in \mathbf{C}_W .

Alternatively, the extended notion of almost sure equality can also be interpreted using the one without the extra wire upon replacing p by $\text{id}_W \otimes p$:

Lemma 2.1.3. Let $p: A \rightarrow X$ and $f, g: W \otimes X \rightarrow Y$. Then we have

$$f =_p g \iff f =_{\text{id}_W \otimes p} g \quad (2.2)$$

Because of this, $f =_p g$ can also be read as a short-hand notation for $f =_{\text{id}_W \otimes p} g$, and we may think of the object W as left implicit and clear from context.

Proof. The equality $f =_{\text{id}_W \otimes p} g$ expressed in string diagrams as follows:

$$(2.3)$$

By marginalizing the middle output W , we conclude $f =_p g$ in the sense of Equation (2.1). For the converse, we merely copy the W input in Equation (2.1) to get $f =_{\text{id}_W \otimes p} g$ by the standard invariance of string diagrams in a symmetric monoidal category under swapping wires. \square

Example 2.1.4. Suppose that \mathbf{C} is a cartesian monoidal category, or equivalently a Markov category in which all morphisms are deterministic. Then the universal property of the product $X \times Y$ shows that $f =_p g$ is equivalent to $f(\text{id}_W \times p) = g(\text{id}_W \times p)$.

Example 2.1.5. For the category \mathbf{Stoch} , we have that Markov kernels $f, g: W \otimes X \rightarrow Y$ are p -almost surely equal for a probability measure $p: I \rightarrow X$ if and only if for every $w \in W$ and for p -almost all $x \in X$ we have $f(_ | w, x) = g(_ | w, x)$ (as measures on Y). In other words, we recover exactly almost sure equality in one of the two variables (and equality in the other variable).

The next result establishes the link between almost sure equality in Markov categories of topological spaces and the topological notion of support.¹ It can serve as a motivation for several of the upcoming constructions in this manuscript. As detailed in Example A.1.4, we consider $\mathbf{TopStoch}$, the category of topological spaces and continuous τ -smooth Markov kernels. A morphism $p: A \rightarrow X$ in $\mathbf{TopStoch}$ has a **support** $S_p \subseteq X$ defined as

$$S_p := \bigcap \{C \subseteq X \text{ closed} \mid p(C|a) = 1 \text{ for all } a \in A\}. \quad (2.4)$$

The τ -smoothness implies $p(S_p|a) = 1$ for all $a \in A$, and therefore S_p can also be characterized as the smallest closed set of full measure under all $p(_ | a)$.²

In the following result, we restrict to Tychonoff spaces, which plays an important role in the proof.

Proposition 2.1.6. *In the category $\mathbf{TychStoch}$ of Tychonoff spaces and continuous Markov kernels (see Example A.1.4 for more details), almost sure equality is equality on the support: For $p: A \rightarrow X$ and $f, g: W \otimes X \rightarrow Y$, we have*

$$f =_p g \quad \iff \quad f(_ | w, x) = g(_ | w, x) \quad \forall w \in W, x \in S_p, \quad (2.5)$$

where S_p is given by Equation (2.4).

Proof. Two τ -smooth probability measures on a topological space are equal if and only if they coincide on a basis of opens [16, Proposition 3.8]. In particular, we can test equality of morphisms on a product space by checking equality on all product opens. Therefore, the almost sure equality $f =_p g$ can be explicitly written out as

$$\int_{x \in U} f(V|w, x) p(dx|a) = \int_{x \in U} g(V|w, x) p(dx|a) \quad (2.6)$$

for all $a \in A$ and $w \in W$ and all opens $U \subseteq X$ and $V \subseteq Y$. Suppose now that equality on the support in the sense of Equation (2.5) holds. Then also Equation (2.6) holds, since by $p(X \setminus S_p|a) = 0$ the integrands coincide almost everywhere.

The converse direction is more difficult. Assuming Equation (2.6), we want to show that $f(_ | w, x) = g(_ | w, x)$ for all $w \in W$ and $x \in S_p$. Since w can be fixed throughout, let us suppress it from the notation to improve readability. Then since Y is Tychonoff, the two probability measures $f(_ | x)$ and $g(_ | x)$ on Y are equal if and only if the expectation values of all bounded continuous functions coincide (see for example [16, Proposition 4.7]). Therefore it is enough to show that for every $x \in S_p$ and every bounded continuous $h: Y \rightarrow \mathbb{R}$, we have

$$\int_{y \in Y} h(y) f(dy|x) = \int_{y \in Y} h(y) g(dy|x). \quad (2.7)$$

What we know from Equation (2.6) is that

$$\int_{x \in U} \int_{y \in Y} h(y) f(dy|x) p(dx|a) = \int_{x \in U} \int_{y \in Y} h(y) g(dy|x) p(dx|a),$$

where we have switched the order of integration by Fubini's theorem. Since this holds for arbitrary open $U \subseteq X$, it likewise holds for every Borel set in place of U , and therefore the

¹See [16, Section 5] for a treatment of the support as a morphism of monads.

²See e.g. [16, Proposition 5.2], where this is spelled out for $A = I$, i.e. for probability measures without an additional parameter.

integrands $\int_{y \in Y} h(y) f(dy|x)$ and $\int_{y \in Y} h(y) g(dy|x)$ must be almost surely equal: there is a Borel set $N_a \subseteq X$ such that $p(N_a|a) = 0$ and such that Equation (2.7) holds for all $x \in X \setminus N_a$. The crucial point is now that both integrands are continuous functions of x , since integrating against h is itself a continuous function on the space of probability measures [16, Proposition 4.7]. Therefore Equation (2.7) holds even on the closure $\overline{X \setminus N_a}$. Since this closure still has full measure by $p(N_a|a) = 0$, we can conclude $S_p \subseteq \overline{X \setminus N_a}$, and therefore Equation (2.7) holds for all $x \in S_p$, as was to be shown. \square

An important application of almost sure equality is that weakening equality to a.s. equality ensures the uniqueness of conditionals, whenever these exist (Lemma A.3.4). Some additional properties of almost sure equality used throughout the paper can be also found in Appendix A.3.

2.2 Definition and examples of absolute continuity

A definition of absolute continuity for morphisms in Markov categories was first given in [18, Definition 2.8]. Here we introduce and study an improved version of that definition. One important improvement is that it is automatically compatible with \otimes , as we show in Proposition 2.3.6, while this fails for the earlier one (Remark 2.3.7). We also show that the two definitions coincide in the main examples of Markov categories.

Definition 2.2.1 (Absolute continuity). *Let $p: A \rightarrow X$ and $q: B \rightarrow X$ be morphisms in a Markov category \mathcal{C} . We say that p is **absolutely continuous with respect to q** , denoted³*

$$p \ll q \quad \text{or} \quad q \gg p,$$

if for all objects W and Y and any two parallel morphisms $f, g: W \otimes X \rightarrow Y$, we have

More concisely, $q \gg p$ says that q -a.s. equality implies p -a.s. equality. Since the relation \gg is clearly reflexive and transitive, it equips the class of morphisms having the same codomain X with a preorder relation.

Example 2.2.2. Since $f =_{\text{id}_X} g$ is equivalent to $f = g$, every morphism $p: A \rightarrow X$ satisfies $\text{id}_X \gg p$. In other words, id_X is a greatest element with respect to \gg .

One may wonder whether it is enough to consider the case $W = I$ only in Definition 2.2.1. That is, can one detect asymptotic continuity by merely comparing almost sure equality of morphisms without an extra input W ? This turns out to be related to the following notions.

Definition 2.2.3. *A Markov category \mathcal{C} is:*

³While the notation $p \ll q$ is arguably more commonly used in measure theory, we write $q \gg p$ in this section as it matches the direction of implication (2.8).

(i) **locally state-separable** if for any parallel morphisms $f, g: W \otimes X \rightarrow Y$, we have

$$\begin{array}{c} Y \\ | \\ \boxed{f} \\ | \\ \nabla_w \\ | \\ X \end{array} = \begin{array}{c} Y \\ | \\ \boxed{g} \\ | \\ \nabla_w \\ | \\ X \end{array} \quad \forall w \in \mathbf{C}(I, W) \quad \Longrightarrow \quad f = g. \quad (2.9)$$

(ii) **point-separable** if for all parallel morphisms $f, g: X \rightarrow Y$, we have

$$\begin{array}{c} Y \\ | \\ \boxed{f} \\ | \\ \nabla_a \end{array} = \begin{array}{c} Y \\ | \\ \boxed{g} \\ | \\ \nabla_a \end{array} \quad \forall a \in \mathbf{C}_{\text{det}}(I, X) \quad \Longrightarrow \quad f = g. \quad (2.10)$$

For example, it is easy to see that all of the usual Markov categories of interest in categorical probability (like **FinStoch**, **BorelStoch** and **Stoch**) are point-separable. We introduce the point-separation condition mainly since it provides a convenient way to verify local state-separation:

Lemma 2.2.4. *If \mathbf{C} is point-separable, then it is locally state-separable.*

Proof. To show that point-separability implies local state-separability, assume that the antecedent of Implication (2.9) holds. Then we have

$$\begin{array}{c} Y \\ | \\ \boxed{f} \\ | \\ \nabla_w \quad \nabla_x \end{array} = \begin{array}{c} Y \\ | \\ \boxed{g} \\ | \\ \nabla_w \quad \nabla_x \end{array} \quad (2.11)$$

for all deterministic states $x: I \rightarrow X$ and $w: I \rightarrow W$. But since every deterministic state on $W \otimes X$ is of the form $w \otimes x$ (because \mathbf{C}_{det} is cartesian monoidal), we obtain the desired $f = g$ from the assumed point-separability. \square

This gives a criterion for when absolute continuity can be detected without the extra wire W .

Proposition 2.2.5. *Consider a locally state-separable Markov category and morphisms $p: A \rightarrow X$ and $q: B \rightarrow X$. Then $q \gg p$ holds if and only if we have*

$$\begin{array}{c} Y \quad X \\ | \quad | \\ \boxed{m} \quad \boxed{k} \\ | \quad | \\ \bullet \\ | \\ \boxed{q} \\ | \\ A \end{array} = \begin{array}{c} Y \quad X \\ | \quad | \\ \boxed{k} \quad \boxed{m} \\ | \quad | \\ \bullet \\ | \\ \boxed{q} \\ | \\ A \end{array} \quad \Longrightarrow \quad \begin{array}{c} Y \quad X \\ | \quad | \\ \boxed{m} \quad \boxed{k} \\ | \quad | \\ \bullet \\ | \\ \boxed{p} \\ | \\ B \end{array} = \begin{array}{c} Y \quad X \\ | \quad | \\ \boxed{k} \quad \boxed{m} \\ | \quad | \\ \bullet \\ | \\ \boxed{p} \\ | \\ B \end{array} \quad (2.12)$$

for all $m, k: X \rightarrow Y$.

Proof. The “only if” direction is trivial, since each Implication (2.12) is a special case of Implication (2.8). For the converse direction, assume that all Implications (2.12) hold. Further note that, by local state-separability, the antecedent of Implication (2.8)—the defining property of the relation $q \gg p$ —is equivalent to

$$\begin{array}{c}
 Y \quad X \\
 \downarrow \quad \downarrow \\
 \boxed{f} \quad \downarrow \\
 \downarrow \quad \downarrow \\
 \nabla_w \quad \bullet \\
 \downarrow \quad \downarrow \\
 \boxed{q} \\
 \downarrow \\
 A
 \end{array}
 =
 \begin{array}{c}
 Y \quad X \\
 \downarrow \quad \downarrow \\
 \boxed{g} \quad \downarrow \\
 \downarrow \quad \downarrow \\
 \nabla_w \quad \bullet \\
 \downarrow \quad \downarrow \\
 \boxed{q} \\
 \downarrow \\
 A
 \end{array}
 \quad (2.13)$$

for all $w: I \rightarrow W$, and similarly for the consequent with p instead of q . In this way, each Implication (2.8) becomes of the form of the assumed (2.12), thus proving the result. \square

Example 2.2.6. For two Markov kernels $p: A \rightarrow X$ and $q: B \rightarrow X$ (i.e. morphisms in the Markov category Stoch), we can use Proposition 2.2.5 to characterize $q \gg p$. In particular, it says that for all measurable sets $S \in \Sigma_X$ we have

$$q(S|b) = 0 \quad \forall b \in B \quad \implies \quad p(S|a) = 0 \quad \forall a \in A. \quad (2.14)$$

Indeed it is known that $m =_q k$ holds if and only if for every $T \in \Sigma_Y$, the measurable functions

$$m(T|_-, k(T|_-) : X \rightarrow [0, 1] \quad (2.15)$$

are almost surely equal with respect to the probability measure $q(_||b)$ for every $b \in B$ [11, Example 13.3], and this directly shows Implication (2.14) by choosing the indicator functions. Conversely, if Implication (2.14) holds, then the description of almost sure equality expressed in terms of symmetric difference given in [11, Example 13.3] is sufficient to prove that $q \gg p$. In conclusion, our categorical notion of absolute continuity is equivalent to the standard one.

Let us consider two special cases.

- (i) For states $p, q: I \rightarrow X$, we recover the standard notion of absolute continuity of probability measures, where (2.14) becomes

$$q(S) = 0 \quad \implies \quad p(S) = 0. \quad (2.16)$$

- (ii) Suppose that p and q are deterministic morphisms in BorelStoch , i.e. measurable maps (which we also denote by p and q by abuse of notation) between standard Borel spaces. Then (2.14) becomes

$$S \cap \text{im}(q) = \emptyset \quad \implies \quad S \cap \text{im}(p) = \emptyset, \quad (2.17)$$

or equivalently simply $\text{im}(q) \subseteq \text{im}(p)$ since every singleton is measurable.

Example 2.2.7. Recall the Markov category SetMulti of multi-valued functions from Example A.1.3, and consider two of its morphisms $p: A \rightarrow X$ and $q: B \rightarrow X$. Then we have

$$q \gg p \quad \iff \quad \bigcup_{a \in A} p(a) \subseteq \bigcup_{b \in B} q(b), \quad (2.18)$$

where $p(a)$ denotes the image of a under p and similarly for $q(b)$.

Example 2.2.8. In the cartesian case (Example 2.1.4), absolute continuity $q \gg p$ simply means

$$f(\text{id}_W \times q) = g(\text{id}_W \times q) \quad \implies \quad f(\text{id}_W \times p) = g(\text{id}_W \times p). \quad (2.19)$$

For example, for $p = \text{id}_X$ this condition states that $\text{id}_W \times q$ must be an epimorphism for every W .

The next example shows that the notion of absolute continuity may depend heavily on the choice of morphisms of the category. In particular, restricting to continuous kernels can result in something quite different than allowing all measurable kernels.

Example 2.2.9. Just as **Stoch**, it is easy to see that the Markov category **TychStoch** is also point-separable, and so Proposition 2.2.5 can be applied. By Proposition 2.1.6 we have $q \gg p$ if and only if the topological support of q contains the topological support of p . In other words, the absolute continuity preorder is the one induced by the inclusion order of topological supports.

Note that this is in stark contrast to the measure-theoretic case: Consider the space Y given by the unit interval $[0, 1]$ with its usual topology and Borel σ -algebra. Further consider the Dirac measure $\delta_{1/2}$ and the Lebesgue measure λ . We can see both as morphisms of type $I \rightarrow Y$. In the Markov category **Stoch** as well as in **BorelStoch**, we have the familiar facts $\delta_{1/2} \not\gg \lambda$ and $\delta_{1/2} \not\ll \lambda$. That is, in measure-theoretic terminology, the two measures are mutually singular. However, $\delta_{1/2} \ll \lambda$ holds in **TychStoch**, since the topological support of $\delta_{1/2}$, the singleton $\{1/2\}$, is contained in the topological support of λ , which is the whole of $[0, 1]$.

For example, a pair of measurable maps $m, k: Y \rightarrow \mathbb{R}$ witnessing $\delta_{1/2} \ll \lambda$ in **Stoch** is

$$m(y) := 0 \quad \forall y, \quad k(y) := \begin{cases} 1 & \text{for } y = 1/2, \\ 0 & \text{for } y \neq 1/2, \end{cases} \quad (2.20)$$

where we have $m =_\lambda k$ but $m \neq_{\delta_{1/2}} k$. The same argument does not work in **TychStoch** because k is not continuous. In fact by Proposition 2.1.6, any two continuous maps or kernels that are λ -almost surely equal must also be $\delta_{1/2}$ -almost surely equal.

While local state-separability allows one to leave out the extra input W in Implication (2.8), this is not true for all Markov categories, as we show next. Here is a general criterion for when this does happen.

Lemma 2.2.10. *Let $q: A \rightarrow X$ be an epimorphism and W an object such that $W \otimes A$ is initial but $W \otimes X$ is not. Then we have*

$$m =_q k \quad \implies \quad m =_{\text{id}_X} k, \quad (2.21)$$

for all $m, k: X \rightarrow Y$, but $q \not\gg \text{id}_X$.

Proof. Implication (2.21) follows directly from the assumption that q is an epimorphism.

For the second statement, consider arbitrary morphisms $f, g: W \otimes X \rightarrow Y$. We trivially have $f =_q g$ since $W \otimes A$ is initial. If $q \gg \text{id}_X$ was true, then we would obtain $f = g$. But this would mean that also $W \otimes X$ is initial, in contradiction to our assumptions. Therefore, $q \gg \text{id}_X$ cannot hold. \square

Example 2.2.11. As in [14, Example 2.3], let $\mathbf{CRing}_+^{\text{op}}$ be the opposite of the category of commutative rings and unital additive maps, considered as a symmetric monoidal category with respect to the usual tensor product. Using the ring multiplication for the comonoid structure on every object turns $\mathbf{CRing}_+^{\text{op}}$ into a Markov category.

Then the assumptions of Lemma 2.2.10 can be satisfied if we choose $q: \mathbb{R} \leftarrow \mathbb{Z}^2$ to be any unital additive inclusion, where the arrow direction indicates the direction as a function. To this end, we can take $W = \mathbb{Z}/2$ to be the field with two elements, since $\mathbb{Z}/2 \otimes \mathbb{R}$ is the zero ring while $\mathbb{Z}/2 \otimes \mathbb{Z}^2$ is not. Hence Lemma 2.2.10 applies, and therefore Implication (2.12) is not equivalent to $q \gg p$ in $\text{CRing}_+^{\text{op}}$.

By Proposition 2.2.5, we can also conclude that $\text{CRing}_+^{\text{op}}$ is not locally state-separable.

2.3 Properties of absolute continuity

Lemma 2.3.1. *For any $h: X \rightarrow Y$ and $p: A \rightarrow X$, we have*

$$h \gg h p. \quad (2.22)$$

In other words, if $h: X \rightarrow Y$ and $g: A \rightarrow Y$ are such that g factors across h , then $h \gg g$ holds.

Proof. To obtain $=_{hp}$ from $=_h$, we simply pre-compose the relevant instances of Equation (2.1) with p . \square

Since $\text{id}_X \gg p$ is always true, the above Lemma 2.3.1 says that in this case, post-composition with h preserves the absolute continuity relation. General post-composition is also \gg -monotone, but only under the additional assumption of causality.

Lemma 2.3.2. *If C is a causal Markov category, then post-composition is \gg -monotone. That is, for all $p: A \rightarrow X$, $q: B \rightarrow X$, and $h: X \rightarrow Y$, we have*

$$q \gg p \implies h q \gg h p. \quad (2.23)$$

Proof. We begin with the first statement. Assuming $f =_{hq} g$ amounts to

$$(2.24)$$

An application of the causality axiom in the form of Implication (A.14) then gives

$$(2.25)$$

Using the assumption $q \gg p$ lets us replace q by p in this equation, and the required $f =_{hp} g$ then follows upon discarding the X output. \square

Under stronger assumptions on the morphism that we post-compose with, we can do away with the causality assumption and even obtain an equivalence.

Lemma 2.3.3. *In an arbitrary Markov category, consider morphisms $p: A \rightarrow T$ and $q: B \rightarrow T$. Furthermore, let $\iota: T \rightarrow X$ be a deterministic split monomorphism. Then we have*

$$q \gg p \iff \iota q \gg \iota p. \quad (2.26)$$

Proof. This is a direct consequence of Lemma A.3.3. \square

Another property of such a left-invertible ι is that morphisms that are absolutely continuous with respect to it factor through ι .

Lemma 2.3.4. *Consider a deterministic split monomorphism $\iota: T \rightarrow X$ with left inverse given by $\pi: X \rightarrow T$, and an arbitrary morphism $p: A \rightarrow X$. Then we have*

$$\iota \gg p \implies p = \iota \pi p. \quad (2.27)$$

Proof. The assumptions imply that we have $\iota \pi = \text{id}_X$. This can be shown by direct calculation, but it also follows from equivalence (A.24) in Lemma A.3.3 by taking $p = \text{id}_X$, $f_2 = \iota \pi$, and $g_2 = \text{id}_X$. Recalling that relation $\iota \gg p$ means, by definition, that ι -a.s. equality implies p -a.s. equality, we get $\iota \pi =_p \text{id}_X$. The desired equation $\iota \pi p = p$ then follows by marginalization. \square

Remark 2.3.5. Unlike post-composition, pre-composition generally does not preserve absolute continuity, not even in FinStoch. For example, if we take $A = B = X = \{0, 1\}$ with

$$q := \begin{pmatrix} 1 & 1/2 \\ 0 & 1/2 \end{pmatrix} \quad p := \begin{pmatrix} 1/2 & 1/2 \\ 1/2 & 1/2 \end{pmatrix}, \quad (2.28)$$

then $q \gg p$ holds. For the deterministic state $\delta_0: I \rightarrow A$, however, we also have $q \delta_0 \not\gg p \delta_0$, since the latter has strictly larger support:

$$q \delta_0 = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad p \delta_0 = \begin{pmatrix} 1/2 \\ 1/2 \end{pmatrix}. \quad (2.29)$$

However, our definition of absolute continuity interacts very well with the monoidal structure.

Proposition 2.3.6. *The monoidal product \otimes is \gg -monotone. That is, for any choice of morphisms $q: B \rightarrow X$, $q': B' \rightarrow X'$, $p: A \rightarrow X$, and $p': A' \rightarrow X'$ we have*

$$q \gg p \text{ and } q' \gg p' \implies q \otimes q' \gg p \otimes p'. \quad (2.30)$$

Proof. Consider generic morphisms $f, g: W \otimes X \otimes X' \rightarrow Y$ and assume that

$$\begin{array}{c} \begin{array}{c} Y & X & X' \\ \downarrow & \downarrow & \downarrow \\ \boxed{f} \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ \boxed{q} \quad \boxed{q'} \\ \downarrow \quad \downarrow \\ W \quad B \quad B' \end{array} \\ = \\ \begin{array}{c} \begin{array}{c} Y & X & X' \\ \downarrow & \downarrow & \downarrow \\ \boxed{g} \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ \boxed{q} \quad \boxed{q'} \\ \downarrow \quad \downarrow \\ W \quad B \quad B' \end{array} \end{array} \quad (2.31)$$

holds. We can rearrange these string diagrams to get

$$(2.32)$$

Using $q' \gg p'$, we can replace q' with p' in this equation and rearrange back to obtain

$$(2.33)$$

By swapping wires, we can similarly use $q \gg p$ to replace q by p and get the desired conclusion of the implication that defines $q \otimes q' \gg p \otimes p'$. \square

Remark 2.3.7. Example 2.2.11 shows that our old definition of absolute continuity without the extra input W [18, Definition 2.8] does not satisfy Equation (2.30): with that definition, we would get $q \gg \text{id}_X$ and $q \otimes \text{id}_W \not\gg \text{id}_X \otimes \text{id}_W$.

2.4 Absolute bicontinuity

The following symmetrized version of absolute continuity comes up frequently in the paper, so it is useful to introduce notation for it.

Definition 2.4.1. We say that two morphisms $p: A \rightarrow X$ and $q: B \rightarrow X$ are **absolutely bicontinuous**, and write $p \approx q$, if both

$$p \gg q \quad \text{and} \quad q \gg p$$

are satisfied. In this case, we also say that p is **absolutely bicontinuous with respect to** q .

Lemma 2.4.2. If $\pi: X \rightarrow T$ is such that $\text{id}_W \otimes \pi$ is an epimorphism for every W , then we have $\pi \approx \text{id}_T$. More generally, $f \pi \approx f$ holds for any $f: T \rightarrow Y$.

For example, this applies if π is a split epimorphism.

Proof. The absolute continuity direction $f \gg f \pi$ is an instance of Lemma 2.3.1, while $f \pi \gg f$ is straightforward from the definition of absolute continuity and the assumption that $\text{id}_W \otimes \pi$ is an epimorphism for every W . \square

Thanks to Proposition 2.3.6, it is also clear that absolute bicontinuity is preserved by the monoidal product.

3 Supports

The support of a probability measure on a finite set or discrete measurable space is a classical concept. Here, we investigate an abstract categorical generalization of this notion in terms of a universal property. Besides the intrinsic interest and utility of having a universal property, one of the features of this development is that it clarifies the significance of topology in the context of supports: A continuous Markov kernel between Polish spaces has a support in `TychStoch`—which turns out to be a Markov category in which *every* morphism has a support—but the same Markov kernel considered as a morphism in the purely measure-theoretic context of `BorelStoch` will typically not have a support.⁴

We start in Section 3.1 by giving our new definition of a support and developing some of its basic properties. We make frequent use of the characterization of supports from Lemma 3.1.3. Using it, Example 3.1.5 shows that all morphisms in `TychStoch` have a support, and these are given by the usual topological supports. This demonstrates that our definition of support is a natural generalization of the classical notion. In Example 3.1.6, we prove that `SetMulti` has supports as well. Theorem 3.1.17 shows that the functoriality of supports is closely related to the causality axiom [11, Definition 11.31].

Section 3.2 is devoted to a more detailed study of the question: “Which morphisms can be expected to have a support?” This results in the definition of an *atomic* morphism in Definition 3.2.2, which is intimately related to the traditional notion of an atomic measure. We use these insights to characterize which morphisms in `BorelStoch` have a support in Theorem 3.2.7. In Proposition 3.2.11, we construct a *free support completion* for every causal Markov category by freely adjoining a support for every atomic morphism. A good universal property of this construction remains to be found, as discussed in Remark 3.2.18.

The following Section 3.3 on the input-output relations functor and Section 3.4 on categories of statistical models provide applications of absolute continuity and supports. These parts can safely be skipped without impeding understanding of the rest of the paper.

The lack of supports in categories like `BorelStoch` can be seen as an (expected) shortcoming. We partially remedy this in Section 3.5 by introducing the *equalizer principle*. This is an axiom for a Markov category \mathbb{C} that can serve as a substitute for supports to some extent. Remark 3.5.2 shows that the equalizer principle is implied by the existence of supports together with the existence of equalizers in the deterministic subcategory \mathbb{C}_{det} . We use this principle in Section 4 as a crucial assumption in our main result on the splitting of idempotents. We also introduce the *relative equalizer principle* in Definition 3.5.5, which is a stronger version of the equalizer principle that still holds in `BorelStoch`. We do not otherwise use it, but merely note that, together with an additional assumption, it implies causality. This results in a statement (Proposition 3.5.7) that is vaguely reminiscent of the earlier Theorem 3.1.17 where causality was derived from the functoriality of supports.

In the final Section 3.6, we introduce a stronger notion of supports that is more in line with our original definition of supports given in [11, Definition 13.20]. In order to distinguish it from Section 3.1, we call them *split supports*. Per Definition 3.6.2, the idea behind these is that not only is the support of a morphism included in its codomain, but this inclusion also has a *splitting* which projects back onto the support. This definition is motivated by the fact that split supports, when they do exist, are better-behaved than plain supports. In particular, we show in Proposition 3.6.12 that split supports are stable under tensoring, which may not be the case for plain supports in general (Conjecture 3.1.16). Proposition 3.6.9 shows that they also enjoy a *second* universal property in addition to the mapping-in universal property of plain


⁴See Example 3.1.5 and Theorem 3.2.7.

supports, namely the *mapping-out* universal property considered originally in [11, Definition 13.20]. As we show in Remark 3.6.15 and Section 4, they are also closely related to idempotents. Unfortunately, not all supports are split supports: Proposition 3.6.14 gives an example of a probability measure on a compact Hausdorff space whose support is not split. In summary, the notion of split support results in a theory that is more appealing than that of plain supports, but at the same time has more limited applicability.

3.1 Definition and basic properties of supports

A first definition of support for morphisms in Markov categories was given in [11, Definition 13.20]. Here we introduce a more intuitive and more broadly applicable definition, while a close variant of the previous notion will be developed in Section 3.6. Let us start with a motivating example.

Example 3.1.1. In $\mathbf{FinStoch}$, let X be the three element set $\{a, b, c\}$, and consider the probability distribution p on X from the introduction, which is



The support of p in the standard sense is the subset $\{a, b\} \subseteq X$. This subset is characterized by the universal property that a stochastic matrix $f: Z \rightarrow X$ factors across the inclusion $\{a, b\} \hookrightarrow X$ if and only if f is absolutely continuous with respect to p . Indeed by Example 2.2.6, the condition $f \ll p$ is equivalent to $f(c|z) = 0$ for all $z \in Z$, and this in turn is equivalent to f factoring across $\{a, b\} \hookrightarrow X$.

In a general Markov category \mathbf{C} , we can define the support of a morphism by means of a universal property generalizing the observation from this example. For any morphism $p: A \rightarrow X$ in \mathbf{C} , consider the functor $\mathbf{C}(_, X)_{\ll p}: \mathbf{C}^{\text{op}} \rightarrow \mathbf{Set}$ defined as the subfunctor of $\mathbf{C}(_, X)$ on all morphisms absolutely continuous with respect to p , i.e. as

$$\mathbf{C}(Z, X)_{\ll p} := \{f: Z \rightarrow X \mid f \ll p\}. \quad (3.1)$$

By definition, having a support amounts to representing this functor.

Definition 3.1.2. Let $p: A \rightarrow X$ be a morphism in a Markov category \mathbf{C} . A **support** for p is an object S together with a natural isomorphism

$$\mathbf{C}(_, S) \cong \mathbf{C}(_, X)_{\ll p} \quad (3.2)$$

which restricts to a bijection between deterministic morphisms on the two sides. \mathbf{C} has **supports** if there exists a support for every morphism p in \mathbf{C} .

Applying the Yoneda lemma shows that a support is given by a deterministic **support inclusion** $\text{si}: S \rightarrow X$ which must be such that composing with si establishes a bijection

$$\mathbf{C}(Z, S) \cong \mathbf{C}(Z, X)_{\ll p} \quad (3.3)$$

for every $Z \in \mathbf{C}$. In other words, every $f: Z \rightarrow X$ satisfying $f \ll p$ must factor uniquely through si . The requirement that si should be deterministic is precisely what guarantees that

the bijection restricts to a bijection at the level of deterministic morphisms. Applying this to the reflexivity inequality $p \ll p$ produces a unique factorization

$$p = \text{si} \widehat{p}, \quad (3.4)$$

where we generally use the hat notation $\widehat{p}: A \rightarrow S$ for this factorization of a morphism across its support. Whenever we want to emphasize the morphism p to which the support object S and the support inclusion si refer to, we denote them instead by S_p and si_p , respectively.

Let us emphasize again that Definition 3.1.2 differs from our earlier one [11, Definition 13.20]. There, the mapping-out property used to define the notion of a support in Markov categories is instead similar to the stronger notion of split supports studied in Section 3.6. A non-trivial example of a morphism for which the two definitions are inequivalent is provided in Proposition 3.6.14.

For many purposes, it is helpful to reformulate the definition of support as follows.

Lemma 3.1.3. *A deterministic monomorphism $\text{si}: S \rightarrow X$ is a support inclusion for a morphism $p: A \rightarrow X$ if and only if it satisfies $\text{si} \ll p$ and, for every $f: Z \rightarrow X$, we have*

$$f \ll p \quad \implies \quad f \text{ factors across } \text{si}. \quad (3.5)$$

Proof. We gave an argument for the forward implication above. Let us prove the reverse implication here. Since si is a monomorphism, composing with it embeds $\mathbf{C}(Z, S)$ into $\mathbf{C}(Z, X)$ injectively. All morphisms in the image of this inclusion factor through si , so that applying Lemma 2.3.1 shows that they are all absolutely continuous with respect to si . As $\text{si} \ll p$ holds by assumption, this allows us to restrict the codomain of the injection to the subset $\mathbf{C}(Z, X)_{\ll p}$ of $\mathbf{C}(Z, X)$. Moreover, this embedding is surjective by the assumed condition (3.5). \square

In all of the following examples, it is obvious that the purported support inclusion is indeed a deterministic monomorphism, and so it only remains to check $\text{si} \ll p$ as well as the factorization condition. Verifying both of these properties is straightforward based on the characterization of the absolute continuity relation in each category.

Example 3.1.4. In FinStoch , every morphism $p: A \rightarrow X$ has a support, namely the set

$$S = \{x \in X \mid \exists a \in A : p(x|a) > 0\}, \quad (3.6)$$

with si given by the inclusion map into X . Using Example 2.2.6, we indeed have $\text{si} \ll p$, and also that an arbitrary $f: Z \rightarrow X$ factors across si if and only if we have $f(x|z) = 0$ for all $x \in X \setminus S$.

Example 3.1.5. In the category TychStoch , the support is given by the topological support, i.e. the smallest closed set of full measure. In more detail, let $p: A \rightarrow X$ be any morphism and define the set $S \subseteq X$ as in Equation (2.4) as

$$S := \bigcap \{C \subseteq X \text{ closed} \mid p(C|a) = 1 \text{ for all } a \in A\}. \quad (3.7)$$

Consider the deterministic inclusion $\text{si}: S \rightarrow X$, and let us apply Lemma 3.1.3 to show that it really is a support inclusion.

- We have $\text{si} \ll p$, and even $\text{si} \approx p$ (see Lemma 3.1.7 below), as both si -almost sure equality and p -almost sure equality are equivalent to equality of kernels restricted to S by Proposition 2.1.6.

- Let now $f: Z \rightarrow X$ be any morphism absolutely continuous with respect to p . By Proposition 2.1.6 again, this means that for all $z \in Z$ we have $f(X \setminus S|z) = 0$. That is, f can be considered as a morphism of type $Z \rightarrow S$, and this provides the desired factorization.

By Lemma 3.1.3, the topological support of p thus coincides with the support of p in our sense.

Example 3.1.6. In SetMulti , every morphism $p: A \rightarrow X$ has a support. In particular, the support inclusion si is the inclusion of the set

$$S := \bigcup_{a \in A} p(a) \quad (3.8)$$

into X , where $p(a)$ is the image of a under p . The property $\text{si} \ll p$ and the factorization condition hold by Example 2.2.7.

Let us return to the general theory for the moment.

Lemma 3.1.7. *For every morphism $p: A \rightarrow X$ with a support, we have*

$$\text{si} \approx p. \quad (3.9)$$

Proof. Relation $\text{si} \gg p$ follows directly from the factorization (3.4) in view of Lemma 2.3.1. The opposite one, $\text{si} \ll p$, holds because si is defined as an element of the set $\mathcal{C}(S, X)_{\ll p}$ from the right-hand side of bijection (3.3). \square

This fact yields in the following characteristic connection between supports and almost sure equality, which may elucidate the definition of supports further.

Lemma 3.1.8 (Almost surely is restriction to the support). *With p and si as above, we have*

$$\begin{array}{c}
 \begin{array}{ccc}
 Y & X & \\
 | & | & \\
 \boxed{f} & \text{---} & \\
 | & \text{---} & \\
 W & A & \\
 \end{array}
 & = &
 \begin{array}{ccc}
 Y & X & \\
 | & | & \\
 \boxed{g} & \text{---} & \\
 | & \text{---} & \\
 W & A & \\
 \end{array}
 & \iff &
 \begin{array}{ccc}
 Y & & \\
 | & & \\
 \boxed{f} & & \\
 | & \boxed{\text{si}} & \\
 W & S & \\
 \end{array}
 & = &
 \begin{array}{ccc}
 Y & & \\
 | & & \\
 \boxed{g} & & \\
 | & \boxed{\text{si}} & \\
 W & S & \\
 \end{array}
 \end{array} \quad (3.10)$$

for all $f, g: W \otimes X \rightarrow X$ with arbitrary W and X .

Proof. The left-hand side of (3.10) is just $f =_p g$. If this holds, then we also get $f =_{\text{si}} g$ by Lemma 3.1.7, and the right-hand side of equivalence (3.10) follows by marginalization. Conversely, if the right-hand side of (3.10) holds, then we have $f =_{\text{si}} g$ by the fact that si is deterministic via Proposition A.3.1 (v). We then conclude by another application of Lemma 3.1.7. \square

Proposition 3.1.9 (Absolute bicontinuity and supports). *Let $p: A \rightarrow X$ and $q: B \rightarrow Y$ be arbitrary. Then:*

- (i) *If p and q have supports with the same support inclusion, then we have $p \approx q$.*
- (ii) *If $p \approx q$ holds and p has a support, then q has the same support.*

Proof.

(i) Since $\text{si}_p = \text{si}_q$, we just use Lemma 3.1.7.

(ii) This holds by the definition of a support. \square

Example 3.1.10. If $p: A \rightarrow X$ is a split epimorphism, then $p \approx \text{id}_X$ holds by Lemma 2.4.2, and therefore the support inclusion of p is just id_X .

Proposition 3.1.11 (Supports of composite morphism). *Consider an arbitrary morphism $p: A \rightarrow T$ and a deterministic split monomorphism $\iota: T \rightarrow X$ with left inverse $\pi: X \rightarrow T$. Then we have the following:*

(i) If p has a support, then so does ιp , and its support inclusion is

$$\text{si}_{\iota p} = \iota \text{si}_p. \quad (3.11)$$

(ii) If ιp has a support, then so does p , and its support inclusion is

$$\text{si}_p = \pi \text{si}_{\iota p}. \quad (3.12)$$

Proof. We employ Lemma 3.1.3 in both directions.

(i) We show that the composite ιsi_p is indeed a support inclusion $\text{si}_{\iota p}$. First, ιsi_p is a deterministic monomorphism as a composite of two deterministic monomorphisms. The required inequality $\iota \text{si}_p \ll \iota p$ holds as a consequence of $\text{si}_p \ll p$ and Lemma 2.3.3. Second, if $f: Z \rightarrow X$ is absolutely continuous with respect to ιp , then we obtain $\iota \pi =_f \text{id}_X$ from $\iota \pi =_{\iota p} \text{id}_X$. This implies $\iota \pi f = f$. By Lemma 2.3.3, we therefore obtain $\pi f \ll p$ from the assumed $f \ll \iota p$. The assumption that si_p is a support inclusion implies now that πf factors across si_p . But then of course also $f = \iota \pi f$ factors across ιsi_p , and we are done.

(ii) We show that the composite $\pi \text{si}_{\iota p}$ is indeed a support inclusion si_p . Using Lemma 2.3.3 again, we obtain $\iota \pi \text{si}_{\iota p} = \text{si}_{\iota p}$ from $\text{si}_{\iota p} \ll \iota p$. Therefore our putative support inclusion for p satisfies $\iota \text{si}_p = \text{si}_{\iota p}$. In particular, si_p is a deterministic monomorphism.⁵ The required inequality $\text{si}_p \ll p$ holds as a consequence of $\iota \text{si}_p \ll \iota p$ and Lemma 2.3.3 once again.

Finally, let $q: Z \rightarrow T$ be absolutely continuous with respect to p . Then $\iota q \ll \iota p$ by Lemma 2.3.3, and hence ιq factors across $\text{si}_{\iota p} = \iota \text{si}_p$. The claim now follows as ι is a monomorphism. \square

Example 3.1.12 (Copying does not change the support). As a special case of Proposition 3.1.11, consider $X = T \otimes T$ and let

$$q := \text{copy}_T p \quad (3.13)$$

be the output-copy version of a morphism $p: A \rightarrow T$. Since copy_T is a deterministic split monomorphism with left inverse $\text{id}_T \otimes \text{del}_T$, we can apply Proposition 3.1.11 (i) to obtain that q has

$$\begin{array}{c} T \quad T \\ \curvearrowright \\ \bullet \\ \boxed{\text{si}} \\ \downarrow \\ S_p \end{array} \quad (3.14)$$

as the support inclusion, where si is the support inclusion of p .

⁵To see that it is deterministic, note that it is enough to prove the determinism equation composed with $\iota \otimes \iota$, and use the assumption that both ι and ιsi_p are deterministic.

In light of Lemma 3.1.3, we do not know which deterministic monomorphisms precisely can arise as support inclusions. However, at least the following is an obvious consequence of Lemma 3.1.3 that also arises as a special case of Proposition 3.1.11 by taking p to be the identity morphism.

Corollary 3.1.13. *Every deterministic split monomorphism is its own support inclusion.*

Example 3.1.14. If an object X has a distribution object PX [18], then we have a deterministic split monomorphism $\delta_X: X \rightarrow PX$ with left inverse given by the sampling map $\text{samp}_X: PX \rightarrow X$. Thus, δ_X is its own support inclusion. The support inclusion of samp_X is given by id_X thanks to Example 3.1.10.

Although properties of cartesian monoidal categories viewed as Markov categories are usually trivial, the existence of supports is a non-trivial requirement.

Proposition 3.1.15. *Let \mathbf{C} be a point-separable cartesian monoidal category with pushouts. Then for a morphism $p: A \rightarrow X$, the following are equivalent:*

- (i) p has a support.
- (ii) The canonical parallel pair⁶ $X \rightrightarrows X \sqcup_A X$ has an equalizer.

If these conditions hold, then the equalizer from (ii) is the support inclusion of p .

Note that a cartesian monoidal category is point-separable if and only if $\mathbf{C}(I, _)$ is a faithful functor, and this is also equivalent to being locally state-separable. Furthermore, the equalizer condition is reminiscent of the notion of *regular category* [4] considered in dual form.

Proof. By the point separability assumption, we have $f \ll p$ if and only if f equalizes all parallel pairs out of X which are equalized by p . But the latter pairs are precisely those which factor across the pushout $X \sqcup_A X$ (by its universal property). Therefore we have $f \ll p$ if and only if f equalizes the parallel pair of (ii).

Now if the support exists, then it follows that si must be the equalizer. The converse is likewise straightforward. \square

Another question is whether supports interact well with the monoidal structure. This remains open, but we believe that this is arguably not the case.

Conjecture 3.1.16. *There exists a Markov category \mathbf{C} with supports and two morphisms p and q such that $S_{p \otimes q} \not\cong S_p \otimes S_q$.*

In particular, if Conjecture 3.1.16 holds, then the canonical comparison morphism

$$S_p \otimes S_q \longrightarrow S_{p \otimes q}$$

induced by the universal property of $S_{p \otimes q}$ and $\text{si}_p \otimes \text{si}_q$ is not an isomorphism. Indeed, since there does not seem to be any obvious way to prove that it is an isomorphism, it seems natural to expect that it is not in general.

We finally turn to the functoriality of supports. Surprisingly, this turns out to be closely related to the causality axiom.

⁶Here, we write $X \sqcup_A X$ for the object of the pushout of $p: A \rightarrow X$ along itself. The parallel pair of morphisms of type $X \rightarrow X \sqcup_A X$ from the colimit cone is also known as the *cokernel pair* of p , a concept better known in the dual form as kernel pair.

Theorem 3.1.17 (Functoriality of supports). *For a Markov category \mathcal{C} with supports, the following are equivalent:*

(i) \mathcal{C} is causal.

(ii) Supports are functorial: Every commutative square

$$\begin{array}{ccc} A & \xrightarrow{p} & X \\ f \downarrow & & \downarrow g \\ B & \xrightarrow{q} & Y \end{array} \quad (3.15)$$

factors into

$$\begin{array}{ccccc} A & \xrightarrow{\hat{p}} & S_p & \xrightarrow{\text{si}_p} & X \\ f \downarrow & & \vdots \downarrow & & \downarrow g \\ B & \xrightarrow{\hat{q}} & S_q & \xrightarrow{\text{si}_q} & Y \end{array} \quad (3.16)$$

for a (necessarily unique) dashed arrow $S_p \rightarrow S_q$.

Note that it is enough for the right square in Equation (3.16) to commute, since then the left square commutes automatically by commutativity of the composite square and the fact that si_q is a monomorphism.

Proof. (i) \Rightarrow (ii): We construct the dashed arrow from the universal property of S_q by showing that $g \text{si}_p \ll q$ holds. Using Lemma 3.1.7 gives $\text{si}_p \approx p$, and applying Lemma 2.3.2 then yields

$$g \text{si}_p \approx g p. \quad (3.17)$$

Moreover, the assumed commutative diagram and Lemma 2.3.1 imply

$$g p = q f \ll q, \quad (3.18)$$

as was to be shown. Therefore, we obtain a unique morphism of type $S_p \rightarrow S_q$ which makes the right square commute. The left square then commutes automatically as noted above.

(ii) \Rightarrow (i): We will show the causality criterion as expressed in Lemma A.3.2. Let $p: A \rightarrow X$, $g: X \rightarrow Y$, and $h_1, h_2: Y \rightarrow Z$ be such that $h_1 =_{gp} h_2$ holds. Then the assumed functoriality of supports applied to the square

$$\begin{array}{ccc} A & \xrightarrow{p} & X \\ \text{id}_A \downarrow & & \downarrow g \\ A & \xrightarrow{gp} & Y \end{array} \quad (3.19)$$

gives us a commutative diagram

$$\begin{array}{ccc} S_p & \xrightarrow{\text{si}_p} & X \\ \vdots \downarrow & & \downarrow g \\ S_{gp} & \xrightarrow{\text{si}_{gp}} & Y \end{array} \quad (3.20)$$

By Lemma 3.1.8, the assumption $h_1 =_{gp} h_2$ is equivalent to $h_1 \text{si}_{gp} = h_2 \text{si}_{gp}$. But then the above diagram implies that we also have $h_1 g \text{si}_p = h_2 g \text{si}_p$. Thus we obtain $h_1 g =_p h_2 g$ by Lemma 3.1.8 again, which completes the proof. \square

The following example shows that supports in **TychStoch** are functorial. By Theorem 3.1.17, this proves that **TychStoch** is a causal Markov category.

Example 3.1.18. **TychStoch** has supports by Example 3.1.5. To show that they are indeed functorial, let us consider the topological support of the composite gp of two stochastic maps $p: A \rightarrow X$ and $g: X \rightarrow Y$. We claim that the support object is given by⁷

$$S_{gp} = \overline{\bigcup_{x \in S_p} S_{g(_ | x)}}. \quad (3.21)$$

Indeed, if we tentatively write S_{gp} as shorthand for the right-hand side, then we have

$$\int_{x \in X} g(S_{gp}|x) p(dx|a) \geq \int_{x \in S_p} g(S_{gp}|x) p(dx|a) = \int_{x \in S_p} p(dx|a) = p(S_p|a) = 1, \quad (3.22)$$

where the first equality holds by $S_{gp} \supseteq S_{g(_ | x)}$ for all $x \in S_p$. Conversely, if $C \subseteq Y$ is a closed set of full measure, meaning

$$\int_{x \in X} g(C|x) p(dx|a) = 1, \quad (3.23)$$

then we need to show $S_{gp} \subseteq C$, or equivalently $S_{g(_ | x)} \subseteq C$ for all $x \in S_p$. But this in turn amounts to showing $g(C|x) = 1$ for all $x \in S_p$. By assumption, we know that the integrand $g(C|x)$ is $p(_ | a)$ -almost everywhere equal to 1. But since $g(C|_)$ is an upper semi-continuous function on X , the set where it is equal to 1 is closed, and hence contains S_p . This shows the required $g(C|x) = 1$ for all $x \in S_p$. Hence the above S_{gp} is indeed the topological support of gp .

Equipped with this description of the topological support of a composite morphism, we can show that supports in **TychStoch** are functorial. To show the desired factorization, we need to prove $g(S_q|x) = 1$ for all $x \in S_p$ given a commutative square as in (3.15). This is an immediate consequence of

$$S_q \supseteq S_{qf} = S_{gp} \supseteq S_{g(_ | x)} \quad \forall x \in S_p, \quad (3.24)$$

where the first containment also follows by the formula for supports of a composite derived above, but now applied to the composite qf .

In conclusion, supports in **TychStoch** are functorial, and Theorem 3.1.17 implies that **TychStoch** is causal.

3.2 Support completeness

As we have seen, the Markov categories **FinStoch**, **TychStoch** and **SetMulti** all have supports. Can we expect supports in measure-theoretic probability? How common are supports in general? In this subsection, we consider these questions and show that supports in measure-theoretic probability (in **BorelStoch** specifically) are quite rare. First, we start with a necessary condition for any morphism to have a support.

Lemma 3.2.1. *If a morphism $p: A \rightarrow X$ has a support, then it satisfies $\text{copy } p \ll p \otimes p$.*

Proof. Whenever p has a support, $p \approx \text{si}$ holds by Lemma 3.1.7. Therefore, we have

$$\text{copy } p \approx \text{copy } \text{si} = (\text{si} \otimes \text{si}) \text{copy} \ll \text{si} \otimes \text{si} \approx p \otimes p, \quad (3.25)$$

where we use first

⁷This equation can also be seen as a manifestation of the fact that the topological support is a morphism of monads [16].

- Lemma 2.3.3 and the fact that copy is a deterministic split monomorphism, then
- the fact that si is deterministic, followed by
- Lemma 2.3.1 with h and p from the lemma given by $\text{si} \otimes \text{si}$ and copy respectively, and finally
- Proposition 2.3.6 in the last step. □

This property of morphisms with a support is important enough to give it its own name, the choice of which we justify in Example 3.2.3.

Definition 3.2.2. *A morphism p in a Markov category is **atomic** if it satisfies $\text{copy } p \ll p \otimes p$.*

Example 3.2.3. The Lebesgue measure $\lambda: I \rightarrow [0, 1]$ in **BorelStoch** is not atomic, since $\text{copy } \lambda$ is the Lebesgue measure on the diagonal of the unit square $[0, 1]^2$, while $\lambda \otimes \lambda$ is the Lebesgue measure on all of $[0, 1]^2$, and the former is not absolutely continuous with respect to the latter (by Example 2.2.6). Hence by Lemma 3.2.1, the Lebesgue measure on $[0, 1]$ does not have a support.

The same argument works generally, and shows that a state $I \rightarrow X$ in **BorelStoch** is atomic if and only if it is atomic in the usual sense as a probability measure on X (see the forthcoming Theorem 3.2.7).

Remark 3.2.4. If $p \approx q$ holds for any deterministic morphism q , then p is an atomic morphism. The proof is the same as that of Lemma 3.2.1, but with q in place of si .

On the other hand, since **FinStoch**, **TychStoch** and **SetMulti** have all supports, it follows a posteriori that every morphism in these categories is atomic. Since not every morphism is atomic in general, the following more restricted existence of supports is the most that one can expect in general.

Definition 3.2.5. *A Markov category \mathcal{C} is **support complete** if every atomic morphism has a support.*

We now turn to a more detailed investigation of supports in measure-theoretic probability, which in particular justifies the term “atomic” further. For a generic morphism $p: A \rightarrow X$ in **BorelStoch**, let us say that an element $x \in X$ is an **atom** if there is $a \in A$ such that the conditional probability $p(\{x\}|a)$ is non-zero. We owe the following auxiliary result to a MathOverflow user.⁸

Lemma 3.2.6 (Packo). *For every morphism p in **BorelStoch**, its set of atoms*

$$\mathcal{A} := \{x \in X \mid \exists a \in A \text{ such that } p(\{x\}|a) > 0\} \tag{3.26}$$

is universally measurable.

Proof. The statement is trivial if X is at most countable, since then every subset is measurable. By Kuratowski’s theorem, we can assume we have $X = \mathbb{R}$ without loss of generality.

The function

$$\begin{aligned} A \times \mathbb{R} &\longrightarrow [0, 1] \\ (a, x) &\longmapsto p([0, x]|a) \end{aligned} \tag{3.27}$$

⁸See <https://mathoverflow.net/questions/448485/atoms-for-markov-kernels/448496> for the relevant answer by Packo.

is measurable in the first argument and right-continuous in the second, which implies that it is jointly measurable.⁹ Similarly, also the function

$$\begin{aligned} A \times \mathbb{R} &\longrightarrow [0, 1] \\ (a, x) &\longmapsto p([0, x] | a) \end{aligned} \tag{3.28}$$

is jointly measurable. Their difference is precisely

$$\begin{aligned} A \times \mathbb{R} &\longrightarrow [0, 1] \\ (a, x) &\longmapsto p(\{x\} | a) \end{aligned} \tag{3.29}$$

which is thus jointly measurable as well. The set of atoms \mathcal{A} is then the projection to \mathbb{R} of the preimage of the set $(0, 1]$ under the function from (3.29). As a projection of a measurable set, the set of atoms is therefore universally measurable. \square

Lemma 3.2.6 allows us to consider the conditional probability $p(\mathcal{A} | a)$ for every $a \in A$. Generalizing the notion of an atomic measure, we may then call a Markov kernel p *atomic* if it satisfies $p(\mathcal{A} | a) = 1$ for all $a \in A$. The following theorem establishes that this is consistent with the notion of atomicity from Definition 3.2.2.

Theorem 3.2.7. *For a morphism $p: A \rightarrow X$ in $\mathbf{BorelStoch}$, consider the following conditions:*

- (i) *p has a support.*
- (ii) *There is a smallest measurable set $S \in \Sigma_X$ with $p(S | a) = 1$ for all $a \in A$.*
- (iii) *p is atomic in the sense of Definition 3.2.2.*
- (iv) *$p(\mathcal{A} | a) = 1$ for all $a \in A$.*

Then we have:¹⁰

$$(i) \iff (ii) \implies (iii) \iff (iv).$$

The converse implication $(iii) \implies (ii)$ holds if A is at most countable, but not in general.

Proof. If p satisfies property (i), then we can take $S \in \Sigma_X$ to be the image of the support inclusion. This set is measurable by [26, Corollary 15.2]. Applying the universal property of the support of p with respect to the inclusion of any other measurable subset establishes the minimality in (ii).

Conversely, suppose that (ii) holds, and take $\mathbf{si}: S \rightarrow X$ to be the inclusion map. Clearly \mathbf{si} is a deterministic monomorphism satisfying $\mathbf{si} \ll p$. By Lemma 3.1.3, it remains to be shown that if $f: Z \rightarrow X$ satisfies $f \ll p$, then it factors across \mathbf{si} . Indeed $f \ll p$ gives $f(X \setminus S | z) = 0$ for all $z \in Z$, and hence f factors across \mathbf{si} . Therefore p has a support.

The implication from (i) to (iii) is exactly Lemma 3.2.1 applied to $\mathbf{BorelStoch}$.

Next, assume that (iii) holds. Following the proof of [36, Theorem 4.6], we write $\Delta \subseteq A \times A$ for the diagonal and \mathcal{A} for the set of atoms of p as in Lemma 3.2.6. Let us argue that the conditional probability

$$(p \otimes p)(\Delta \setminus (\mathcal{A} \times \mathcal{A}) | a_1, a_2) \tag{3.30}$$

⁹This is a standard observation in stochastic process theory, see e.g. [25, Proposition 1.13] applied to the constant filtration.

¹⁰For the equivalence of (iii) and (iv) in the case of states, see also [36, Theorem 4.6].

of $\Delta \setminus (\mathcal{A} \times \mathcal{A})$ given any $a_1, a_2 \in A$ vanishes. Indeed, this follows upon decomposing every $p(_ | a)$ into its atomic and diffuse parts, noting that the atomic part vanishes outside of \mathcal{A} , and that the product of two diffuse finite measures vanishes on the diagonal.¹¹ By Equation (2.14) and the assumed property (iii), we therefore obtain

$$(\text{copy } p)(\Delta \setminus (\mathcal{A} \times \mathcal{A}) | a) = 0, \quad (3.31)$$

or equivalently $p(A \setminus \mathcal{A} | a) = 0$, for all a , which implies (iv).

Conversely, suppose that (iv) holds, so that we have $p(\mathcal{A} | a) = 1$ for all $a \in A$. To show $\text{copy } p \ll p \otimes p$, suppose that we have a measurable set $T \in \Sigma_{X \times X}$ satisfying $(p \otimes p)(T | a_1, a_2) = 0$ for all $a_1, a_2 \in A$. Then taking $a_1 = a_2$ shows that T is disjoint from $\mathcal{A} \times \mathcal{A}$. But then also the intersection of T with the diagonal does not contain any atom, and hence $(\text{copy } p)(T | a) = 0$ holds for all $a \in A$, which gives the desired property (iv).

For the final statement, we note that there is an implication from (iv) to (ii) if the set of atoms is measurable. In such case, we can take S from (ii) to be the set \mathcal{A} itself. This is guaranteed to be the case if A is countable, since then \mathcal{A} is countable as well and hence trivially measurable.

To see that this implication does not hold for general A , consider any measurable function $\mathbb{R} \rightarrow \mathbb{R}$ with non-measurable image. Considered as a deterministic morphism $p: \mathbb{R} \rightarrow \mathbb{R}$ in BorelStoch , it is clearly atomic with the set of atoms \mathcal{A} being its image. If p had a support, then by $\text{si} \approx p$ and Example 2.2.6 (ii), such support would have to be given by the inclusion map of a non-measurable subset, which is not possible in BorelStoch [26, Corollary 15.2]. \square

Corollary 3.2.8 (Supports of measurable maps). *A deterministic morphism p in BorelStoch has a support if and only if its image (as a measurable map) is measurable, and in this case the support is given by the image.*

Proof. If p has measurable image, then this image serves as the set S in Theorem 3.2.7 (ii). Conversely, suppose that p has a support. Then we have $\text{si} \approx p$ by Lemma 3.1.7, which implies $\text{im}(\text{si}) = \text{im}(p)$ by Example 2.2.6 (ii). In particular $\text{im}(p)$ is measurable, since si is an injective measurable map and therefore has a measurable image [26, Corollary 15.2]. \square

The final statement of Theorem 3.2.7—or in other words the existence of measurable maps between standard Borel spaces with non-measurable image—has the following unfortunate consequence.

Corollary 3.2.9. *BorelStoch is not support complete.*

In order to address this shortcoming, one may wonder whether it is possible to construct a support complete Markov category that contains BorelStoch as a full Markov subcategory. This is true, and it is an instance of a more general construction which we call *free support completion*. This term is by analogy with the free cocompletion of a category, which is its presheaf category [4, Exercise 6.7.4].

Definition 3.2.10. *Let \mathcal{C} be a causal Markov category. Its **free support completion** $\text{Supp}(\mathcal{C})$ is the causal Markov category described as follows:*

- (i) *Objects are pairs (X, p) , where $X \in \mathcal{C}$ is an object and p is any atomic morphism landing in X .*

¹¹This can be shown by decomposing A into pieces of mass below ε for both measures.

- (ii) Morphisms $(X, p) \rightarrow (Y, q)$ are p -a.s. classes of morphisms $f: X \rightarrow Y$ satisfying $f p \ll q$.
- (iii) The Markov category structure is induced from \mathbf{C} .

Let us first demonstrate that this definition actually defines a causal Markov category.

Proposition 3.2.11. *Supp(C) is indeed a well-defined causal Markov category, and C is naturally identified with the full Markov subcategory whose objects are of the form (X, id_X) .*

Proof. We first show that composition in Supp(C) is well-defined. If $[f]: p \rightarrow q$ and $[g]: q \rightarrow r$ are morphisms represented by $f: X \rightarrow Y$ and $g: Y \rightarrow Z$, then also gf represents a morphism $p \rightarrow r$, since we have

$$(gf)p = g(fp) \ll gq \ll r, \quad (3.32)$$

where the second step is by Lemma 2.3.2. We also need to show that this composition is well-defined with respect to the equivalence classes. To this end, consider a morphism $f': X \rightarrow Y$ satisfying $f' =_p f$. Then, by Proposition A.3.1 (i), we also have $gf =_p gf'$, which amounts to the desired $[gf] = [gf']$. Similarly, if $g': Y \rightarrow Z$ satisfies $g' =_q g$, then we obtain $g =_{fp} g'$ by $fp \ll q$. Hence the required $gf =_p g'f$ follows by Lemma A.3.2.

It is clear that the symmetric monoidal structure morphisms of \mathbf{C} also belong to Supp(C), where (I, id_I) is the monoidal unit of Supp(C). Moreover, the tensor product of two morphisms respects the absolute continuity condition: For $[f]: (X, p) \rightarrow (Y, q)$ and $[f']: (X', p') \rightarrow (Y', q')$, we have

$$(f \otimes f')(p \otimes p') = (fp) \otimes (f'p') \ll q \otimes q' \quad (3.33)$$

by Proposition 2.3.6 and the assumed $fp \ll q$ and $f'p' \ll q'$, as was to be shown. Hence by what has been proven so far, Supp(C) is a symmetric monoidal category. It is obvious that it is again semicartesian.

Concerning the Markov category structure, we define the copy morphisms as

$$\text{copy}_{(X,p)} := [\text{copy}_X]: (X, p) \rightarrow (X, p) \otimes (X, p). \quad (3.34)$$

We emphasize that such a $\text{copy}_{(X,p)}$ is indeed a morphism in Supp(C) because p is atomic. Direct checks show that these morphisms indeed equip Supp(C) with a Markov category structure.

From this description, it is immediate that we can identify \mathbf{C} with the full subcategory whose objects are of the form (X, id_X) , and the inclusion functor is a strict Markov functor by construction.

It remains to be shown that Supp(C) is causal. This holds as a special case of the following implication in \mathbf{C} :

This is exactly *relative causality* in the sense of [17, Corollary 2.28], which is shown there to be equivalent to causality. \square

Before showing that $\text{Supp}(\mathbf{C})$ is support complete, we give an explicit description of absolute continuity in this category.

Lemma 3.2.12. *Let $[f]: (X, p) \rightarrow (Y, q)$ and $[g]: (Z, r) \rightarrow (Y, q)$ be two generic morphisms in $\text{Supp}(\mathbf{C})$. Then $[f] \ll [g]$ holds in $\text{Supp}(\mathbf{C})$ if and only if we have $f p \ll g r$ in \mathbf{C} .*

Proof. The absolute continuity relation $[f] \ll [g]$ in $\text{Supp}(\mathbf{C})$ reads as

$$\begin{array}{c}
 \begin{array}{c} A \quad Y \\ \boxed{h} \quad \text{---} \\ | \quad | \\ \bullet \\ | \quad | \\ W \quad Z \\ \boxed{g} \end{array} \\
 =_r \\
 \begin{array}{c} A \quad Y \\ \boxed{k} \quad \text{---} \\ | \quad | \\ \bullet \\ | \quad | \\ W \quad Z \\ \boxed{g} \end{array} \\
 \Rightarrow \\
 \begin{array}{c} A \quad Y \\ \boxed{h} \quad \text{---} \\ | \quad | \\ \bullet \\ | \quad | \\ W \quad X \\ \boxed{f} \end{array} \\
 =_p \\
 \begin{array}{c} A \quad Y \\ \boxed{k} \quad \text{---} \\ | \quad | \\ \bullet \\ | \quad | \\ W \quad X \\ \boxed{f} \end{array}
 \end{array} \tag{3.36}$$

in \mathbf{C} , where the h and k both have domain $(W, \text{id}) \otimes (Y, q)$ in $\text{Supp}(\mathbf{C})$. Note that we can set their left input to have id_W as the atomic morphism without loss of generality. Any other choice can be subsumed by adjusting the morphisms themselves, but this is fine as they are arbitrary.

Using the causality of \mathbf{C} , the antecedent of Implication (3.36) can be equivalently written as

$$\begin{array}{c}
 \begin{array}{c} A \quad Y \\ \boxed{h} \quad \text{---} \\ | \quad | \\ \bullet \\ | \quad | \\ W \quad Z \\ \boxed{g} \\ \boxed{r} \end{array} \\
 = \\
 \begin{array}{c} A \quad Y \\ \boxed{k} \quad \text{---} \\ | \quad | \\ \bullet \\ | \quad | \\ W \quad Z \\ \boxed{g} \\ \boxed{r} \end{array}
 \end{array} \tag{3.37}$$

where we use Implication (A.14). Since the same is true for the consequent of Implication (3.36) with f and p in place of g and r , we conclude that $f p \ll g r$ is equivalent to Implication (3.36). \square

Lemma 3.2.13. *Let $[f]: (X, p) \rightarrow (Y, q)$ be an atomic morphism in $\text{Supp}(\mathbf{C})$. Then $(Y, f p)$ is a support of f with support inclusion $[\text{id}_Y]: (Y, f p) \rightarrow (Y, q)$,*

Proof. We use the conditions of Lemma 3.1.3. First of all, to verify that $(Y, f p)$ is an object of $\text{Supp}(\mathbf{C})$, we need to ensure that $f p$ is atomic. This holds by Lemma 3.2.12 because $[f]$ is atomic in $\text{Supp}(\mathbf{C})$.

Since id_Y is a deterministic monomorphism, it is easy to verify that the proposed support inclusion si given by $[\text{id}_Y]: (Y, f p) \rightarrow (Y, q)$ is a deterministic monomorphism as well.¹²

By Lemma 3.2.12, the condition $\text{si} \ll [f]$ from Lemma 3.1.3 is immediate because it corresponds to $f p \ll f p$ in \mathbf{C} . Similarly, if any $[g]: (Z, r) \rightarrow (Y, q)$ satisfies $[g] \ll [f]$, then we have $g r \ll f p$ in \mathbf{C} , so that g defines a morphism $(Z, r) \rightarrow (Y, f p)$ in $\text{Supp}(\mathbf{C})$. This is the desired factorization of $[g]$ across si . \square

¹²The reader may wonder whether this morphism is also an epimorphism. From the definition, one can check that this is true if and only if $f p \approx q$ holds.

We therefore have the following main result on the free support completion.

Theorem 3.2.14. *For every causal Markov category \mathbf{C} :*

- (i) *The free support completion $\mathbf{Supp}(\mathbf{C})$ is support complete.*
- (ii) *Any atomic morphism $p: A \rightarrow X$ in \mathbf{C} has support (X, p) in the larger category $\mathbf{Supp}(\mathbf{C})$.*

Proof. (i) This follows from Lemma 3.2.13.

- (ii) In light of Lemma 3.2.12, $[p]: (A, \text{id}_A) \rightarrow (X, \text{id}_X)$ is still atomic in $\mathbf{Supp}(\mathbf{C})$. Hence this follows from Lemma 3.2.13 as well. \square

Corollary 3.2.15. *Let \mathbf{C} be a causal Markov category. A morphism admits a support in a Markov category containing \mathbf{C} as a Markov subcategory if and only if it is atomic.*

Proof. If p is atomic, then Theorem 3.2.14 shows that p has a support in $\mathbf{Supp}(\mathbf{C})$. Conversely, if we assume that p is not atomic, then $\text{copy } p \ll p \otimes p$ fails already in \mathbf{C} , i.e. there exist morphisms f and g satisfying both $f =_{p \otimes p} g$ and $f \neq_{\text{copy } p} g$. This remains the case in any Markov category containing \mathbf{C} as a Markov subcategory. The contrapositive of Lemma 3.2.1 says that such p cannot have a support. \square

Example 3.2.16. By applying Corollary 3.2.15 to $\mathbf{BorelStoch}$, we conclude that the Lebesgue measure (as well as any other non-atomic morphism) will never admit a support, even if we enlarge our category.

Remark 3.2.17. The inclusion functor $\mathbf{C} \rightarrow \mathbf{Supp}(\mathbf{C})$ does not preserve supports in general. If $p: A \rightarrow X$ in \mathbf{C} admits a support, then there at least exists a canonical comparison morphism

$$[\text{si}]: (S, \text{id}_S) \rightarrow (X, p)$$

from the image of its support in \mathbf{C} to its support in $\mathbf{Supp}(\mathbf{C})$. This morphism is clearly deterministic, and it can be seen to be an epimorphism by Lemma 3.1.8. Hence it is an isomorphism if and only if it has a left inverse $[\text{sp}]: (X, p) \rightarrow (S, \text{id}_S)$. However, this would require $\text{sp si} = \text{id}_S$ in \mathbf{C} , which means that the support S would have to be a *split support* in the sense of the upcoming Section 3.6. This is not the case in general (Proposition 3.6.14).

In fact, the example from Proposition 3.6.14 can be considered as a counterexample in $\mathbf{TychStoch}$. This suggests that the inclusion functor of $\mathbf{TychStoch}$ into its free support completion may not be an equivalence of categories, although $\mathbf{TychStoch}$ already has all supports.

These phenomena underline again why we speak of the *free* support completion: By adding new supports, $\mathbf{Supp}(\mathbf{C})$ can break the existing ones. All of this is analogous to the free cocompletion of a category, where the analogous inclusion functor does not preserve colimits in general, and it is not an equivalence of categories.

Remark 3.2.18 (Universal property of the support completion?). A first problem regarding the free support completion is that we have not found a good universal property for it so far. By analogy with the free cocompletion, we would expect a universal property of the following kind: Given a support complete Markov category \mathbf{D} and a Markov functor $F: \mathbf{C} \rightarrow \mathbf{D}$ preserving absolute continuity, this functor can be extended to a support-preserving Markov functor $\mathbf{Supp}(\mathbf{C}) \rightarrow \mathbf{D}$ by mapping every object (X, p) in $\mathbf{Supp}(\mathbf{C})$ to the support $S_{F(p)}$ in \mathbf{D} . For a morphism $[f]: (X, p) \rightarrow (Y, q)$ in $\mathbf{Supp}(\mathbf{C})$, we have $F(f) F(p) \ll F(q)$ since F preserves absolute continuity by assumption, and hence $F(f) \text{si}_{F(p)} \ll F(q)$ holds by Lemmas 2.3.2 and 3.1.7, so that we obtain the desired induced morphism $S_{F(p)} \rightarrow S_{F(q)}$. This is well-defined with respect to the equivalence relation $=_p$ thanks to Lemma 3.1.8 and clearly functorial.

A first problem is now that we do not know whether or not monoidal products of supports in \mathbf{D} are supports again (Conjecture 3.1.16). If this indeed was the case, then we should indeed get a strong monoidal functor $\text{Supp}(\mathbf{C}) \rightarrow \mathbf{D}$, and in fact a Markov functor, which should facilitate the formulation of a well-behaved 2-categorical universal property. If this is not the case in general—as we expect—then we could at most hope to obtain a lax Markov functor $\text{Supp}(\mathbf{C}) \rightarrow \mathbf{D}$.

3.3 The input-output relation functor

In the following two sections, we present several applications of supports and absolute continuity. These are not otherwise used in the rest of the paper.

For a morphism $p: A \rightarrow X$ in FinStoch , it is sometimes of interest to consider the set of all pairs $(a, x) \in A \times X$ satisfying $p(x|a) > 0$. This set is a relation from A to X which we call the *input-output relation*, since it assigns to each input a the set of possible outputs x . Mapping every stochastic matrix p to its input-output relation defines an identity-on-objects functor $\text{FinStoch} \rightarrow \text{SetMulti}$. As we show below, one can associate an input-output relation to morphisms in any suitably well-behaved Markov category \mathbf{C} , and this results in a Markov functor $\mathbf{C} \rightarrow \text{SetMulti}$. Whenever this is possible, one can say that every morphism has a consistent “possibilistic shadow” [21].

Concretely, we map each object A of \mathbf{C} to the set of its deterministic states

$$\overline{A} := C_{\text{det}}(I, A), \quad (3.38)$$

which one may also think of as “points” of A . Further, we map a morphism $p: A \rightarrow X$ to the **input-output relation** $p \gg$ given by

$$p \gg := \{(a, x) \in \overline{A} \times \overline{X} \mid pa \gg x\}. \quad (3.39)$$

In other words, this relation is defined by

$$a \overset{p}{\gg} x \quad :\iff \quad pa \gg x, \quad (3.40)$$

and we think of this as stating that the output x is possible for p under input a . In order to ensure the functoriality of the assignment $p \mapsto p \gg$, we need an additional assumption.

Definition 3.3.1. *A Markov category \mathbf{C} has **point liftings** if for all $p: A \rightarrow X$ and all $x \in \overline{X}$ satisfying $p \gg x$, there exists an $a \in \overline{A}$ satisfying $pa \gg x$.*

Intuitively, this says that every output that can occur under p must be able to arise from *some* input.

Remark 3.3.2. Definition 3.3.1 can be easier to check if supports exist, which is going to be the case in the assumptions of Theorem 3.3.6 below. For in this case, Lemma 3.1.3 implies that for every $a \in \overline{A}$, we have a canonical bijection

$$\{x \in \overline{X} \mid a \overset{p}{\gg} x\} \cong \overline{S_{pa}} \quad (3.41)$$

between points absolutely continuous with respect to pa on the left and deterministic states “within the support of pa ” on the right. The point lifting condition then says that each point $x \in \overline{X}$ in the support of p is also in the support of the image of some point $a \in \overline{A}$ under p .

Example 3.3.3. FinStoch has point liftings because the support of a stochastic matrix is the union of its “pointwise” supports (i.e. supports of its columns).

Example 3.3.4. BorelStoch also has point liftings. Indeed, by the characterization of absolute continuity of Example 2.2.6 used in the contrapositive, $p \gg x$ means that if x is an element of a measurable set $S \in \Sigma_X$, then $p(S|a) > 0$ holds for some $a \in A$. Taking $S = \{x\}$ shows that there is $a \in A$ such that x is an atom of the probability measure $p(_ | a)$. This is why we also have the required $pa \gg x$.

For an object $A \in \mathbf{C}$, we have $\text{del}_A \gg \text{id}_I$ as soon as $\mathbf{C}(I, A) \neq \emptyset$, since then one can witness the relevant instance of the absolute continuity implication (2.8) by pre-composing with any state on A . In this case, the point lifting property implies that A also must have a deterministic state.

Example 3.3.5. Let G be a finite group, and let FinStoch^G be the Markov category of finite G -sets together with G -equivariant Markov kernels. Then FinStoch^G does not have point liftings, unless G is the trivial group. Indeed the set $A := G$ itself with its canonical G -action has a state given by the uniform distribution, but it has no deterministic state unless G is trivial.

Theorem 3.3.6. *Let \mathbf{C} be a causal Markov category with supports and point liftings. Then there is an **input-output relation functor** $\Upsilon: \mathbf{C} \rightarrow \text{SetMulti}$ given on objects and morphisms by*

$$\Upsilon(A) := \overline{A}, \quad \Upsilon(p) := {}^p \gg,$$

as defined in (3.38) and (3.39), and this Υ is a Markov functor.

Proof. We show first that $\Upsilon(p)$ is actually a morphism in SetMulti for every $p: A \rightarrow X$. This amounts to proving that for every $a \in \overline{A}$, there is $x \in \overline{X}$ with $a {}^p \gg x$, or equivalently $pa \gg x$. Indeed the support object S_{pa} has a state given by the support factorization \widehat{pa} , and by what was noted above this implies that S_{pa} also has a deterministic state. Composing this state with the support inclusion si_{pa} produces $x \in \overline{X}$. The desired $pa \gg x$ then follows by $pa \approx \text{si}_{pa}$ and Lemma 2.3.4.

Next, we show that Υ is a functor. Proving that Υ preserves identities amounts to showing that if $a \gg b$ for $a, b \in \overline{A}$, then $a = b$. Since every deterministic state is its own support by Corollary 3.1.13, the assumption $a \gg b$ implies that b factors across a , and is thus equal to a .

Consider now arbitrary morphisms $p: A \rightarrow X$ and $f: X \rightarrow Y$. We need to show that for every deterministic $a: I \rightarrow A$ and $y: I \rightarrow Y$, we have

$$a {}^f p \gg y \iff a ({}^f \gg \circ {}^p \gg) y, \tag{3.42}$$

where ${}^f \gg \circ {}^p \gg$ is the composite relation. This amounts to showing

$$f p a \gg y \iff \exists x \in \overline{X} : pa \gg x \text{ and } f x \gg y. \tag{3.43}$$

We consider both implications.

(\Leftarrow): Using the causality assumption and Lemma 2.3.2, we have

$$pa \gg x \implies f p a \gg f x. \tag{3.44}$$

Together with the transitivity of \gg and the assumed $f x \gg y$, this gives the desired $f p a \gg y$.

(\Rightarrow): We now assume that $f p a \gg y$ holds. Since $\text{si}_{pa} \gg pa$ holds by Lemma 3.1.7, we also have $f \text{si}_{pa} \gg f p a$ by Lemma 2.3.2, and therefore

$$f \text{si}_{pa} \gg y \tag{3.45}$$

holds by the transitivity of \gg . Hence, by point liftings, there is a deterministic $\tilde{x}: I \rightarrow S_{pa}$ that satisfies

$$f \text{ si}_{pa} \tilde{x} \gg y. \quad (3.46)$$

If we define $x := \text{si}_{pa} \tilde{x}$, then $fx \gg y$ is necessarily true. The required $pa \gg x$ then holds by $pa \gg \text{si}_{pa} \gg \text{si}_{pa} \tilde{x}$, where the first relation is by Lemma 3.1.7 again and the second by Lemma 2.3.1. Since x is deterministic, this concludes the proof of the forward implication in (3.43).

Having shown that Υ is a functor, let us provide the details of its strong monoidal structure. First, the monoidal structure morphisms are the relations given by the graphs of the functions

$$\begin{aligned} \psi_0 : I &\longrightarrow \bar{I} \\ * &\longmapsto \text{id}_I \end{aligned} \quad (3.47)$$

where the monoidal unit $I \in \text{SetMulti}$ is a singleton set with element $*$, and

$$\begin{aligned} \psi_{A,B} : \bar{A} \otimes \bar{B} &\longrightarrow \overline{A \otimes B} \\ (a, b) &\longmapsto a \otimes b. \end{aligned} \quad (3.48)$$

The only non-trivial condition to check in order for these to equip Υ with the structure of a strong symmetric monoidal functor is the naturality of ψ . This amounts to showing that, for all morphisms $p: A \rightarrow X$ and $q: B \rightarrow Y$ and all deterministic states

$$a \in \bar{A}, \quad b \in \bar{B}, \quad x \in \bar{X}, \quad y \in \bar{Y},$$

we have

$$(a \otimes b)^{p \otimes q} \gg (x \otimes y) \iff a^p \gg x \wedge b^q \gg y, \quad (3.49)$$

or equivalently

$$pa \otimes qb \gg x \otimes y \iff pa \gg x \wedge qb \gg y. \quad (3.50)$$

The right-to-left implication is clear by Proposition 2.3.6. In the other direction, we can post-compose with marginalization maps in order to obtain the claim from Lemma 2.3.2.

To conclude the proof that Υ is a Markov functor, it remains to be shown that we have $\Upsilon(\text{copy}_A) = \psi_{A,A} \text{ copy}_{\Upsilon(A)}$ for every $A \in \mathcal{C}$. This amounts to proving that for all $a, b, c \in \bar{A}$, we have

$$\text{copy}_A a \gg b \otimes c \iff a = b = c. \quad (3.51)$$

Using $\text{copy}_A a = a \otimes a$, this follows by the same arguments as in the previous paragraph. \square

Example 3.3.7. In FinStoch , the Markov functor $\Upsilon: \text{FinStoch} \rightarrow \text{SetMulti}$ is precisely the input-output relation functor discussed at the beginning of this subsection.

Example 3.3.8. SetMulti also satisfies the assumptions of Theorem 3.3.6, and the associated input-output relation functor $\Upsilon: \text{SetMulti} \rightarrow \text{SetMulti}$ is the identity functor.

Example 3.3.9. One may wonder how relevant the assumption of supports in Theorem 3.3.6 really is, given that its statement does not refer to supports in any way. To see that merely assuming point liftings is not sufficient, consider BorelStoch and let $\lambda: I \rightarrow [0, 1]$ be the Lebesgue measure, or similarly any other non-atomic measure on a standard Borel space. Then $\Upsilon(\lambda) = \lambda \gg$ is the empty relation since λ has no atoms. We could then still hope to get a functor to Rel rather than to SetMulti . If functoriality held, then also $\Upsilon(\text{del}_{[0,1]} \lambda)$ would have to be the empty

relation. But this is clearly equal to $\Upsilon(\text{del}_I) = \Upsilon(\text{id}_I) = \text{id}_{\{*\}}$, which is not the empty relation. Therefore functoriality does not hold.

Incidentally, looking at the proof around (3.45) shows that the proof would go through if λ had a support. We therefore obtain another proof of the fact that λ does not have a support, as in Theorem 3.2.7.

In [37, Definition IV.4], the property of a Markov category with “precise supports” was given in the context of conditioning. Note that this notion refers only to the absolute continuity relation \gg rather than to supports in the sense of this paper.¹³ More concretely, a Markov category \mathbf{C} is said to have **precise supports** if, for all $x \in \overline{X}$ and $y \in \overline{Y}$ and arbitrary morphisms $p: I \rightarrow X$ and $f: X \rightarrow Y$, the relation

$$\begin{array}{c} X \quad Y \\ \downarrow \quad \downarrow \\ \text{---} \quad \boxed{f} \\ \downarrow \quad \downarrow \\ \bullet \\ \downarrow \\ \triangle p \end{array} \gg \begin{array}{c} X \quad Y \\ \downarrow \quad \downarrow \\ \triangle x \quad \triangle y \end{array} \quad (3.52)$$

is equivalent to

$$\begin{array}{c} X \\ \downarrow \\ \triangle p \end{array} \gg \begin{array}{c} X \\ \downarrow \\ \triangle x \end{array} \quad \text{and} \quad \begin{array}{c} Y \\ \downarrow \\ \boxed{f} \\ \downarrow \\ \triangle x \end{array} \gg \begin{array}{c} Y \\ \downarrow \\ \triangle y \end{array} \quad (3.53)$$

This property follows from point liftings in a causal Markov category.

Proposition 3.3.10. *If a causal Markov category \mathbf{C} has point liftings, then it also has precise supports.*

Proof. First, note that we can use Lemma 2.3.2 to marginalize Y in relation (3.52) to get $p \gg x$. Furthermore, relation (3.52) also implies

$$\begin{array}{c} X \quad Y \\ \downarrow \quad \downarrow \\ \text{---} \quad \boxed{f} \\ \downarrow \quad \downarrow \\ \bullet \\ \downarrow \\ X \end{array} \gg \begin{array}{c} X \quad Y \\ \downarrow \quad \downarrow \\ \triangle x \quad \triangle y \end{array} \quad (3.54)$$

by Lemma 2.3.1. Using point liftings, we get that there exists a deterministic $a: I \rightarrow X$ such that we have

$$\begin{array}{c} X \quad Y \\ \downarrow \quad \downarrow \\ \text{---} \quad \boxed{f} \\ \downarrow \quad \downarrow \\ \triangle a \quad \triangle a \end{array} \gg \begin{array}{c} X \quad Y \\ \downarrow \quad \downarrow \\ \triangle x \quad \triangle y \end{array} \quad (3.55)$$

By Lemma 2.3.2 again, we get $a \gg x$ and $fa \gg y$ by marginalization. Note that a is its own support inclusion (Corollary 3.1.13), so that x factors through a . In other words, we have $x = a$, so that condition (3.53) follows.

¹³Also, the definition of absolute continuity used in [37] was still the “old” one without the extra input wire in Definition 2.2.1.

Conversely, using Lemma 2.3.2, we can post-compose the relation $p \gg x$ appearing in (3.53) to get the left-most relation in

$$\begin{array}{c} X \quad Y \\ \downarrow \quad \downarrow \\ \text{---} \quad \text{---} \\ \downarrow \quad \downarrow \\ \bullet \quad \downarrow \\ \downarrow \quad \downarrow \\ \triangleleft \quad \downarrow \\ p \quad f \end{array} \ggg \begin{array}{c} X \quad Y \\ \downarrow \quad \downarrow \\ \text{---} \quad \text{---} \\ \downarrow \quad \downarrow \\ \bullet \quad \downarrow \\ \downarrow \quad \downarrow \\ \triangleleft \quad \downarrow \\ x \quad f \end{array} = \begin{array}{c} X \quad Y \\ \downarrow \quad \downarrow \\ \triangleleft \quad \downarrow \\ x \quad f \end{array} \ggg \begin{array}{c} X \quad Y \\ \downarrow \quad \downarrow \\ \triangleleft \quad \downarrow \\ x \quad y \end{array} \quad (3.56)$$

where the right-most relation follows from the assumed $fx \gg y$ by Proposition 2.3.6. In conclusion, condition (3.53) implies condition (3.52). \square

3.4 Categories of statistical models

As another application of absolute continuity and supports, we now consider categories of statistical models. Concrete measure-theoretic versions of such categories have been studied since the pioneering work of Čencov [38, 39]. Recent works in this direction include a paper by Jost, Lê and Tran [24] and the thesis of Patterson [34].

Here, we introduce the category of parametric statistical models internal to any causal and representable Markov category. Subsequently, in Proposition 3.4.3 we use supports to show that in the measure-theoretic setting of standard Borel spaces, a parametric statistical model consisting of a family of distributions $p = (p_a)_{a \in A}$ is isomorphic to the model given by its image in PX , provided that this image is measurable.

We assume familiarity with representable Markov categories [18]; see Appendix A.4 for a brief recap.

Definition 3.4.1. *Let \mathcal{C} be a Markov category that is both causal and representable. Then its **category of statistical models** $\text{Stat}(\mathcal{C})$ is the symmetric monoidal category where:*

- *Objects are pairs (X, p) where $X \in \mathcal{C}$ and $p: A \rightarrow X$ is any morphism in \mathcal{C} .*
- *Morphisms $(X, p) \rightarrow (Y, q)$ are p -a.s. equivalence classes of morphisms $f: X \rightarrow Y$ such that $P(f)p^\# \ll q^\#$, or equivalently $(fp)^\# \ll q^\#$, holds.*
- *Composition and symmetric monoidal structure are inherited from \mathcal{C} .*

The idea is that a pair $(X, p: A \rightarrow X)$ can be interpreted as a statistical model with A as parameter space [11, Definition 14.1]. The absolute continuity condition on $f: X \rightarrow Y$ intuitively encodes the requirement that f must map distributions on X compatible with the model p to distributions on Y compatible with the model q (Example 3.4.4). If $f, g: X \rightarrow Y$ satisfy $f =_p g$, then they intuitively behave the same on all distributions that are compatible with p , and hence we consider them as representatives of the same morphism in $\text{Stat}(\mathcal{C})$. The monoidal structure on objects of $\text{Stat}(\mathcal{C})$ is given by the monoidal product of morphisms in \mathcal{C} , the idea being that if $(X, p: A \rightarrow X)$ and $(Y, q: B \rightarrow Y)$ are statistical models, then $(X \otimes Y, p \otimes q)$ is a statistical model with parameter space $A \otimes B$ such that the factors X and Y are modelled as independent.

For the following proof, note that in a causal Markov category, we have

$$P(f)p^\# = (fp)^\# \ll q^\# \implies fp \ll q \quad (3.57)$$

by using post-composition with samp_Y and Lemma 2.3.2.

Proposition 3.4.2. *Stat(C) is a symmetric monoidal category.*

The proof bears similarity to the analogous statement for the free support completion, Proposition 3.2.11.

Proof. We first consider composition. If $[f]: (X, p) \rightarrow (Y, q)$ and $[g]: (Y, q) \rightarrow (Z, r)$ are morphisms, then also $[g f]: (X, p) \rightarrow (Z, r)$ is, since

$$P(g f) p^\sharp = P(g) P(f) p^\sharp \ll P(g) q^\sharp \ll r^\sharp, \quad (3.58)$$

where the middle step follows by Lemma 2.3.2 upon composing the assumed relation $P(f) p^\sharp \ll q^\sharp$ with $P(g)$. We also need to show that this composition is well-defined with respect to the equivalence classes. So consider a morphism $f': X \rightarrow Y$ with $f' =_p f$. Then, by Proposition A.3.1 (i), we also have $g f =_p g f'$, which amounts to the desired $[g f] = [g f']$. Similarly, if $g': Y \rightarrow Z$ satisfies $g' =_q g$, then we obtain $g =_{f p} g'$ by $f p \ll q$, which holds by Implication (3.57). Hence the required $g f =_p g' f$ follows by Lemma A.3.2.

It is clear that the symmetric monoidal structure morphisms of \mathbf{C} also belong to $\text{Stat}(\mathbf{C})$. Moreover, Proposition 2.3.6 shows that the tensor product of two morphisms respects the absolute continuity condition: For $[f]: (X, p) \rightarrow (Y, q)$ and $[f']: (X', p') \rightarrow (Y', q')$, we have

$$(f p)^\sharp \otimes (f' p')^\sharp \ll q^\sharp \otimes q'^\sharp \quad (3.59)$$

by Proposition 2.3.6 and the assumed $(f p)^\sharp \ll q^\sharp$ and $(f' p')^\sharp \ll q'^\sharp$. This is an absolute continuity relation between morphisms with codomain $PY \otimes PY'$. Now recall that in terms of the symmetric monoidal structure of P given by

$$\nabla : PY \otimes PY' \longrightarrow P(Y \otimes Y'), \quad (3.60)$$

we have

$$(q \otimes q')^\sharp = \nabla (q^\sharp \otimes q'^\sharp), \quad (3.61)$$

and similarly for the left-hand side of Equation (3.59). Therefore composing this absolute continuity relation with ∇ yields the desired

$$(f p \otimes f' p')^\sharp \ll (q \otimes q')^\sharp \quad (3.62)$$

by Lemma 2.3.2 and the proof is complete. Finally, the well-definedness of the monoidal product with respect to $=_p$ -classes is straightforward: For any f_2 satisfying $f_2 =_p f$, then we also have $f_2 \otimes f' =_{p \otimes p'} f \otimes f'$ by Proposition A.3.1 (iii). \square

Proposition 3.4.3. *Let $p: A \rightarrow X$ be such that $p^\sharp: A \rightarrow PX$ has a support. Then there is an isomorphism*

$$(X, p) \cong (X, \text{samp}_X \text{si}_{p^\sharp}) \quad (3.63)$$

in $\text{Stat}(\mathbf{C})$.

Intuitively, the support of p^\sharp coincides with its image of PX , which is the set of distributions compatible with the statistical model (X, p) . Therefore the proposition states that (X, p) is equivalent as a statistical model to this image, as one would expect.

Proof. Let us write $q: S_{p^\sharp} \rightarrow PX$ as shorthand for the morphism $\text{samp}_X \text{si}_{p^\sharp}$ in \mathbf{C} . We show that id_X gives both a morphism $(X, p) \rightarrow (X, q)$ and a morphism $(X, q) \rightarrow (X, p)$ in $\text{Stat}(\mathbf{C})$. Since $P(\text{id}_X) = \text{id}_{PX}$, we have to show that $p^\sharp \approx q^\sharp$ holds. Indeed,

$$q^\sharp = (\text{samp}_X \text{si}_{p^\sharp})^\sharp = \text{si}_{p^\sharp} \quad (3.64)$$

follows because si_{p^\sharp} is deterministic. Then the required $p^\sharp \approx \text{si}_{p^\sharp}$ holds by Lemma 3.1.7. \square

Example 3.4.4. Consider any statistical model $(X, p: A \rightarrow X)$ in BorelStoch , which we can think of as an A -indexed family of distributions $(p_a)_{a \in A}$ on X . The measurable map $p^\sharp: A \rightarrow PX$ is precisely the one that assigns to every parameter value $a \in A$ the associated distribution $p_a \in PX$.

If the image of p^\sharp , namely the set of distributions

$$\{p_a \mid a \in A\} \subseteq PX, \quad (3.65)$$

is measurable, then its support is given by this image (Corollary 3.2.8). In this case, Proposition 3.4.3 applies, and we find that the parametric statistical model $(X, p: A \rightarrow X)$ is isomorphic to a model given simply by a subset of PX , namely the set of distributions compatible with the model.

3.5 The equalizer principle

In this subsection, we introduce an axiom for Markov categories that can sometimes be used as a substitute for supports when these do not exist. This axiom plays an important role in Section 4.

Definition 3.5.1. A Markov category \mathcal{C} satisfies the *equalizer principle* if:

- (i) Equalizers in \mathcal{C}_{det} exist.
- (ii) For every equalizer diagram

$$E \xrightarrow{\text{eq}} X \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} Y \quad (3.66)$$

in \mathcal{C}_{det} , every $p: A \rightarrow X$ in \mathcal{C} satisfying $f =_p g$ factors uniquely across eq .

Since finite products exist in \mathcal{C}_{det} by virtue of it being cartesian monoidal, the existence of equalizers implies that \mathcal{C}_{det} is even finitely complete.

The relation between the equalizer principle and supports is as follows.

Remark 3.5.2. Property (ii) holds automatically for every p which has a support. In such case, the assumption $f =_p g$ is equivalent to $f \text{ si} =_p g \text{ si}$ by Lemma 3.1.8, and therefore si factors across eq by the universal property. Furthermore, since we have $p = \text{si} \hat{p}$, it follows that p itself factors across eq .

In particular, if \mathcal{C}_{det} has equalizers and \mathcal{C} has supports, then the equalizer principle holds. For example:

- (i) FinStoch satisfies the equalizer principle, since its deterministic subcategory is FinSet (which is finitely complete) and every morphism has a support by Example 3.1.4.
- (ii) SetMulti satisfies the equalizer principle, since its deterministic subcategory is Set (which is complete), and every morphism has a support by Example 3.1.6.
- (iii) TychStoch satisfies the equalizer principle, since its deterministic subcategory is the category of Tychonoff spaces and continuous maps (which has equalizers since every subspace of a Tychonoff space is Tychonoff) and every morphism has a support by Example 3.1.5.

In the other direction, we apply the equalizer principle in the proof of Theorem 4.4.3 to *construct* supports for certain morphisms. Such relations to supports illustrate why we believe that the equalizer principle can sometimes be used as a substitute for supports.

Remark 3.5.3. Even when the equalizer principle holds, an equalizer in \mathbf{C}_{det} is usually *not* an equalizer in \mathbf{C} . In other words, merely having $fp = gp$ instead of $f =_p g$ is not enough to guarantee that p factors across eq . For example in $\mathbf{FinStoch}$, consider any finite set Y with $|Y| \geq 2$, and let f and g be given by the two deterministic projections $Y \otimes Y \rightarrow Y$. Then their equalizer in $\mathbf{FinStoch}_{\text{det}} = \mathbf{FinSet}$ is given by the inclusion of the diagonal. Taking p to be the uniform probability measure on $Y \otimes Y$, we have $fp = gp$, but p does not factor across the inclusion of the diagonal.

We cannot derive the equalizer principle from supports in measure-theoretic probability, since supports do typically not exist there (Theorem 3.2.7). Nevertheless, we have the following positive result.

Proposition 3.5.4. *BorelStoch satisfies the equalizer principle.*

Proof. Equalizers in $\mathbf{BorelStoch}_{\text{det}} = \mathbf{BorelMeas}$ clearly exist, since for a given parallel pair as in Equation (3.66), we simply consider the subset

$$E := \{x \in X \mid f(x) = g(x)\}, \quad (3.67)$$

which is measurable in X since f and g are measurable maps and the diagonal of their output is measurable. Letting $\text{eq}: E \hookrightarrow X$ be the inclusion, the universal property of an equalizer is obvious.

Property (ii) is more interesting. Let $p: A \rightarrow X$ be any Markov kernel satisfying $f =_p g$. By Kuratowski's theorem, we know that Y is either discrete and (at most) countable, or isomorphic to \mathbb{R} . In each of the cases, we can embed Y into \mathbb{R} while preserving the equalizer as well as the equation $f =_p g$. Therefore, we can assume $Y = \mathbb{R}$ without loss of generality.

In short, f and g are assumed to be real-valued measurable functions which are $p(_|_a)$ -almost surely equal for every $a \in A$. That is, we have $p(X \setminus E|a) = 0$ for all $a \in A$, and hence p factors across the inclusion $\text{eq}: E \hookrightarrow X$. The uniqueness of the factorization is obvious. \square

The following stronger version of the equalizer principle may also be of interest, although it does not appear in this paper besides the present Section 3.5. We prove that it is also satisfied for Markov kernels in $\mathbf{BorelStoch}$ and illustrate its power by showing how it can be used to obtain the causality axiom.

Definition 3.5.5. *A Markov category \mathbf{C} satisfies the relative equalizer principle if:*

- (i) *Equalizers in \mathbf{C}_{det} exist.*
- (ii) *For every equalizer diagram*

$$E \xrightarrow{\text{eq}} X \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} Y \quad (3.68)$$

in \mathbf{C}_{det} , every $p: A \rightarrow X$ satisfying $f =_{pq} g$ for some $q: Z \rightarrow A$ factors uniquely across eq up to q -a.s. equality.

Proposition 3.5.6. *BorelStoch satisfies the relative equalizer principle.*

Proof. Proceeding as in the previous proof, we have that $f, g: X \rightarrow \mathbb{R}$ are almost surely equal with respect to the probability measure

$$\int_{a \in A} p(_|_a) q(da|z) \quad (3.69)$$

for every $z \in Z$, so that we get

$$\int_{a \in A} p(X \setminus E|a) q(da|z) = 0. \quad (3.70)$$

This implies that the integrand vanishes almost surely, meaning that with the notation

$$N := \{a \in A \mid p(X \setminus E|a) > 0\} \quad (3.71)$$

we have $q(N|z) = 0$ for all $z \in Z$. This implies the unique factorization as desired by constructing a Markov kernel $A \rightarrow E$ to be given by p on $A \setminus N$ and arbitrarily¹⁴ on N . \square

Proposition 3.5.7. *Let \mathcal{C} be an a.s.-compatibly representable¹⁵ Markov category satisfying the relative equalizer principle. Then \mathcal{C} is causal.*

Proof. We use the reformulation of the causality axiom from Lemma A.3.2. Let $h_1 =_{gf} h_2$ be given, where $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ and $h_1, h_2: Z \rightarrow A$ are morphisms of the appropriate type. By a.s.-compatible representability, we obtain $h_1^\# =_{gf} h_2^\#$, and it is enough to show $h_1^\# g =_f h_2^\# g$. To simplify the notation, we therefore just assume that h_1 and h_2 are deterministic without loss of generality.

With eq denoting the equalizer of h_1 and h_2 in \mathcal{C}_{det} , the relative equalizer principle applied to $h_1 =_{gf} h_2$ gives us the dashed morphism in

$$\begin{array}{ccccc} E & \xleftarrow{\text{eq}} & Z & \begin{array}{c} \xrightarrow{h_1} \\ \xrightarrow{h_2} \end{array} & A \\ & \swarrow \ell & \uparrow g & & \\ X & \xrightarrow{f} & Y & & \end{array} \quad (3.72)$$

where the triangle commutes up to $=_f$. Therefore we have

$$h_1 g =_f h_1 \text{eq} \ell = h_2 \text{eq} \ell =_f h_2 g, \quad (3.73)$$

which amounts to the desired $h_1 g =_f h_2 g$. \square


3.6 Split supports

In this subsection, we study a stronger notion of support, which we call “split support”. This has several independent motivations, starting with another universal property, which in contrast to the mapping-in universal property of supports given in Definition 3.1.2 is a mapping-out universal property, extending the original one [11, Definition 13.20]. Another point is that split supports are more well-behaved than “plain” supports when they do exist. For example, it is easy to see that split supports are stable under tensor product, while this is probably not the case for plain supports (Conjecture 3.1.16).


¹⁴If N is non-empty, this is possible only if E is non-empty. However, we can assume this without loss of generality. If Z is the empty set, the overall claim is trivial. On the other hand, if Z is non-empty, then f and g are almost surely equal with respect to some probability measure, and thus E is also non-empty.

¹⁵See Appendix A.4 for the definition.

Example 3.6.1 (Split support for a stochastic map). Let us illustrate the basic idea by considering Example 3.1.1 again. For another set $Y = \{x, y, z\}$, let functions $f, g: X \rightarrow Y$ be given by



$$f(a) = x, \quad f(b) = y, \quad f(c) = z;$$



$$g(a) = x, \quad g(b) = y, \quad g(c) = y.$$

Note that f and g are equal p -almost surely. The intuitive reason behind this is that restricting them to the support $S = \{a, b\}$ of p gives the *same* function of type $S \rightarrow Y$. This is true also for general supports as given by Definition 3.1.2 (see Lemma 3.1.8). However, notice an additional property that is *not captured* by our definition of “plain” supports in Markov categories. Namely, every morphism $h: S \rightarrow Y$ arises from a restriction as above. One can always extend such h to a morphism $X \rightarrow Y$ (necessarily p -a.s. uniquely), so that restricting to S produces h itself again.

Of course, to be able to extend every h , it is sufficient for the identity map $S \rightarrow S$ to be extendable, or equivalently for the support inclusion $\text{si}: S \rightarrow X$ to have a left inverse $\text{sp}: X \rightarrow S$. In our example, we can take it to be any map satisfying

$$\text{sp}(a) = a, \quad \text{sp}(b) = b. \tag{3.74}$$

The value of $\text{sp}(c)$ is arbitrary.

We can express an analogous property in any Markov category.

Definition 3.6.2. Let $p: A \rightarrow X$ be a morphism in a Markov category \mathbb{C} . Then a **split support** for p is a support such that the support inclusion $\text{si}: S \rightarrow X$ has a left inverse $\text{sp}: X \rightarrow S$.

We also call such a left inverse a **support projection** for p . We assume a fixed choice of support projection sp throughout, while noting that the particular choice plays no role due to the following result.

Lemma 3.6.3. The support inclusion and projection of a split support satisfy

$$\text{sp si} = \text{id}_S, \quad \text{si sp} =_p \text{id}_X. \tag{3.75}$$

Moreover, sp is p -a.s. deterministic and uniquely determined by the first equation up to p -a.s. equality.

Proof. The first equation is by definition, while the second is a consequence of Lemma 3.1.8

and $\text{si sp si} = \text{si}$. The uniqueness up to p -a.s. equality is now immediate. Finally, we compute

$$\begin{array}{c} S \quad S \\ | \quad | \\ \text{---} \quad \text{---} \\ | \quad | \\ \bullet \\ | \\ \boxed{\text{sp}} \\ | \\ X \end{array} = \begin{array}{c} S \quad S \\ | \quad | \\ \boxed{\text{sp}} \quad \boxed{\text{sp}} \\ | \quad | \\ \boxed{\text{si}} \quad \boxed{\text{si}} \\ | \quad | \\ \bullet \\ | \\ \boxed{\text{sp}} \\ | \\ X \end{array} = \begin{array}{c} S \quad S \\ | \quad | \\ \boxed{\text{sp}} \quad \boxed{\text{sp}} \\ | \quad | \\ \bullet \\ | \\ \boxed{\text{si}} \\ | \\ \boxed{\text{sp}} \\ | \\ X \end{array} \stackrel{=p}{=} \begin{array}{c} S \quad S \\ | \quad | \\ \boxed{\text{sp}} \quad \boxed{\text{sp}} \\ | \quad | \\ \bullet \\ | \\ X \end{array} \quad (3.76)$$

to conclude that sp is p -a.s. deterministic. \square

Let us turn to some examples of split supports. A nontrivial example of a support that is not split will be given later in Proposition 3.6.14.

Example 3.6.4. The support of any morphism $p: A \rightarrow X$ in $\mathbf{FinStoch}$ is a split support, provided that A is non-empty. The reason is that its support inclusion $\text{si}: S \rightarrow X$ is a split monomorphism with a deterministic left inverse as illustrated in Example 3.6.1.

Example 3.6.5. Similar statements apply to $\mathbf{BorelStoch}$. In fact, every deterministic monomorphism with non-empty domain in $\mathbf{BorelStoch}$ is also a split monomorphism. By a classical result, such a map is a measurable isomorphism onto its image [26, Corollary 15.2], and therefore we can obtain a (deterministic) left inverse by mapping the complement of the image to any element of its domain. It follows that any Markov kernel $p: A \rightarrow X$ in $\mathbf{BorelStoch}$ with a support also has a split support (unless A is empty).

The following result is an analogue of Lemma 3.1.8 for split supports, and it can be helpful for showing that a given morphism has a split support.

Lemma 3.6.6 (Concrete characterization of split supports). *To give a split support for a morphism $p: A \rightarrow X$ is to give an object S and morphism $\text{si}: S \rightarrow X$ such that the following conditions hold:*

- (i) si is a deterministic split monomorphism.
- (ii) for all $f, g: W \otimes X \rightarrow Y$ with arbitrary W and Y , we have

$$\begin{array}{c} Y \quad X \\ | \quad | \\ \boxed{f} \quad \bullet \\ | \quad | \\ W \quad A \\ | \quad | \\ \boxed{p} \end{array} = \begin{array}{c} Y \quad X \\ | \quad | \\ \boxed{g} \quad \bullet \\ | \quad | \\ W \quad A \\ | \quad | \\ \boxed{p} \end{array} \iff \begin{array}{c} Y \\ | \\ \boxed{f} \\ | \\ W \quad S \\ | \\ \boxed{\text{si}} \end{array} = \begin{array}{c} Y \\ | \\ \boxed{g} \\ | \\ W \quad S \\ | \\ \boxed{\text{si}} \end{array} \quad (3.77)$$

Proof. Given a split support as in Definition 3.1.2, the first property holds by definition, and the second is precisely Lemma 3.1.8.

Conversely, suppose that we have a morphism $\text{si}: S \rightarrow X$ satisfying the two properties, and choose any left inverse sp . Then it is enough to apply Lemma 3.1.3 to show that si is a support inclusion. Indeed the condition $\text{si} \ll p$ is straightforward to check directly from (3.77) and an

application of Proposition A.3.1 (iv) to si . It remains to be shown that if $q: Z \rightarrow X$ satisfies $q \ll p$, then it factors across si . To this end, note that by (3.75) and $q \ll p$ we get

$$\text{si sp} =_q \text{id}_X, \quad (3.78)$$

which gives the explicit factorization $q = \text{si sp}$. \square

Example 3.6.7. Every deterministic split monomorphism si is its own split support inclusion, since Equation (3.77) holds with $p = \text{si}$ by Proposition A.3.1 (v). This strengthens Corollary 3.1.13.

For example, every deterministic state is trivially a split monomorphism by the terminality of the unit object I . Hence, deterministic states are their own split support inclusions.

Remark 3.6.8. In a locally state-separable Markov category, a deterministic split monomorphism $\text{si}: S \rightarrow X$ is a split support for p already if it satisfies condition (ii) from Lemma 3.6.6 without an additional input W : For all $f, g: X \rightarrow Y$, we have

$$\begin{array}{c} Y \quad A \\ \boxed{f} \quad \boxed{p} \\ \text{---} \quad \text{---} \\ \bullet \\ \text{---} \\ A \end{array} = \begin{array}{c} Y \quad A \\ \boxed{g} \quad \boxed{p} \\ \text{---} \quad \text{---} \\ \bullet \\ \text{---} \\ A \end{array} \iff \begin{array}{c} Y \\ \boxed{f} \\ \text{---} \\ \boxed{\text{si}} \\ \text{---} \\ S \end{array} = \begin{array}{c} Y \\ \boxed{g} \\ \text{---} \\ \boxed{\text{si}} \\ \text{---} \\ S \end{array} \quad (3.79)$$

This follows by the same argument as the proof of Proposition 2.2.5.

Lemma 3.6.6 also lets us prove the announced mapping-out universal property of split supports. For any morphism $p: A \rightarrow X$ and arbitrary objects W and Y , let us write

$$\mathbb{C}(W \otimes X, Y)_p := \mathbb{C}(W \otimes X, Y) / \equiv_p \quad (3.80)$$

for the hom-set modulo p -a.s. equality.

Proposition 3.6.9. *Let $p: A \rightarrow X$ be any morphism in a Markov category \mathbb{C} . Then a morphism $\text{si}: S \rightarrow X$ is a split support inclusion for p if and only if for all objects Y and W in \mathbb{C} , the map*

$$\begin{aligned} \mathbb{C}(W \otimes X, Y)_p &\longrightarrow \mathbb{C}(W \otimes S, Y) \\ f &\longmapsto f(\text{id}_W \otimes \text{si}) \end{aligned} \quad (3.81)$$

is a bijection, and it restricts to a bijection between equivalence classes of p -a.s. deterministic morphisms on the left and deterministic morphisms on the right. When this is the case, then the inverse map is given by $g \mapsto g(\text{id}_W \otimes \text{sp})$ for any support projection sp .

Proof. Suppose that si is a split support inclusion with support projection sp . Then the fact that the two composition maps $f \mapsto f(\text{id}_W \otimes \text{si})$ and $g \mapsto g(\text{id}_W \otimes \text{sp})$ are well-defined and mutually inverse bijections follows from Lemma 3.6.6 and Equation (3.75). That this restricts to a bijection at the level of (p -a.s.) deterministic morphisms follows from the fact that si is deterministic and sp is p -a.s. deterministic (Lemma 3.6.3).

Conversely, suppose that the composition map is a bijection restricting to a bijection at the (p -a.s.) deterministic level. Then define si as the counterpart on the right of the identity on the left, which we know to be deterministic. Furthermore, we have an $\text{sp}: X \rightarrow S$ which is the

counterpart on the left of the identity on the right, and Yoneda lemma shows that for $W = I$, the map from right to left is given by $g \mapsto g \mathbf{sp}$, and in particular \mathbf{sp} is a left inverse for \mathbf{si} . It is now clear that both properties of Lemma 3.6.6 are satisfied, and hence \mathbf{si} is a split support inclusion. \square

Remark 3.6.10. In particular, taking $W = I$ in Proposition 3.6.9 produces a bijection

$$\mathbf{C}(X, Y)_p \cong \mathbf{C}(S, Y),$$

which is natural in Y . This is the promised mapping-out universal property, which also was our original definition of support in [11, Definition 13.20]. If we consider the characterization of Proposition 3.6.9 as the definition of split supports, then the present notion strengthens the earlier one in two ways:

- As in our new definition of absolute continuity (Definition 2.2.1), an extra input W is allowed.
- We now have an additional correspondence at the level of deterministic morphisms.

The latter point is analogous to the definition of Kolmogorov products [14, Definition 4.1], where both general morphisms and deterministic morphisms into a Kolmogorov product are fixed via its universal property.

As yet another possible characterization of split supports, we have the following elegant statement.

Corollary 3.6.11. *A morphism p has a split support if and only if there is a deterministic split monomorphism \mathbf{si} satisfying $p \approx \mathbf{si}$, in which case \mathbf{si} provides the split support.*

Proof. The “if” direction follows by Proposition 3.1.9 (ii) and the fact that every deterministic split monomorphism is its own split support inclusion (Example 3.6.7). The “only if” direction is by Lemma 3.1.7. \square

One of the distinguishing features of split supports is that they are preserved by monoidal products. Recall that this is an open problem for “plain” supports (Conjecture 3.1.16).

Proposition 3.6.12 (Split supports of monoidal products). *If morphisms*

$$p: A \rightarrow X, \quad q: B \rightarrow Y$$

have split supports, then also $p \otimes q$ has a split support with support inclusion given by $\mathbf{si}_p \otimes \mathbf{si}_q$.

Proof. We apply Corollary 3.6.11. Indeed deterministic split monomorphisms are stable under the monoidal product: One can take $\mathbf{sp}_p \otimes \mathbf{sp}_q$ as a left inverse for $\mathbf{si}_p \otimes \mathbf{si}_q$. Furthermore, $\mathbf{si}_p \otimes \mathbf{si}_q \approx p \otimes q$ follows from $\mathbf{si}_p \approx p$ and $\mathbf{si}_q \approx q$ by Proposition 2.3.6. \square

We also have an analogue of Proposition 3.1.11 on how split supports transport along split monomorphisms.

Proposition 3.6.13. *Consider an arbitrary morphism $p: A \rightarrow T$ and a deterministic split monomorphism $\iota: T \rightarrow X$ with left inverse $\pi: X \rightarrow T$. Then we have the following:*

- (i) *If p has a split support, then so does ιp , and its support inclusion is*

$$\mathbf{si}_{\iota p} = \iota \mathbf{si}_p. \tag{3.82}$$

(ii) If ιp has a split support, then so does p , and its support inclusion is

$$\text{si}_p = \pi \text{si}_{\iota p}. \quad (3.83)$$

Proof. By Proposition 3.1.11, it is enough to establish the existence of a support projection in each case.

(i) $\text{si}_{\iota p}$ is a split monomorphism as a composite of two split monomorphisms, since

$$\text{sp}_{\iota p} := \text{sp}_p \pi \quad (3.84)$$

serves as a support projection for ιp .

(ii) The support projection

$$\text{sp}_p := \text{sp}_{\iota p} \iota \quad (3.85)$$

is indeed a left inverse of si_p , since $\iota \pi \text{si}_{\iota p} = \text{si}_{\iota p}$ was noted in the proof of Proposition 3.1.11. \square

Finally, we note that not all supports are split, even in the very well-behaved case of supports in CHausStoch , which always exist (Example 3.1.5). The relevant idea that an embedding $\beta\mathbb{N} \hookrightarrow [0, 1]^{\mathbb{R}}$ cannot have a continuous stochastic retraction was suggested to us by Tommaso Russo, who also noted the stronger statement that there is no Banach space embedding of $C(\beta\mathbb{N})$ into $C([0, 1]^{\mathbb{R}})$ due to known results in renorming theory [9].

Proposition 3.6.14. *There is a probability measure $\nu: I \rightarrow [0, 1]^{\mathbb{R}}$ in CHausStoch such that ν has no split support.*

Surprisingly, the ν that we construct is even atomic (in the measure-theoretic sense of having atoms).

Proof. Let $\beta\mathbb{N}$ be the Stone-Ćech compactification of a countably infinite set. Then the hom-set

$$\text{CHaus}(\beta\mathbb{N}, [0, 1]) \cong \text{Top}(\mathbb{N}, [0, 1]) \quad (3.86)$$

has the cardinality of \mathbb{R} . Using this together with a standard double dualization argument gives us a homeomorphic embedding $\beta\mathbb{N} \hookrightarrow [0, 1]^{\mathbb{R}}$. We identify $\beta\mathbb{N}$ with the image of this embedding and let ν be any atomic probability measure whose set of atoms is given by $\mathbb{N} \subseteq \beta\mathbb{N} \subseteq [0, 1]^{\mathbb{R}}$.

Then since $\mathbb{N} \subseteq \beta\mathbb{N}$ is dense and $\beta\mathbb{N} \subseteq [0, 1]^{\mathbb{R}}$ is closed, the topological support of ν is $\beta\mathbb{N}$. Therefore, by Example 3.1.5, the support inclusion in our sense is $\text{si}: \beta\mathbb{N} \hookrightarrow [0, 1]^{\mathbb{R}}$. Under Gelfand duality, this inclusion corresponds to an algebra homomorphism $C(\text{si}): C([0, 1]^{\mathbb{R}}) \rightarrow C(\beta\mathbb{N})$ given by the restriction of functions to the subspace. Now, if a support projection existed, then by probabilistic Gelfand duality [19], the homomorphism $C(\text{si})$ would have a positive unital linear section $C(\beta\mathbb{N}) \rightarrow C([0, 1]^{\mathbb{R}})$, assigning to every continuous function on $\beta\mathbb{N}$ a continuous extension to $[0, 1]^{\mathbb{R}}$ of the same norm. By a result of Pełczyński [35, Corollary 8.14], such a section does not exist.¹⁶ \square

Let us finish with a teaser for the next section.

¹⁶Pełczyński's result applies to non-metrizable compact spaces which are extremally disconnected, and it is well-known that $\beta\mathbb{N}$ has these properties. It follows that $\beta\mathbb{N}$ is not an *almost Dugundji space*, and in particular not a *Dugundji space*, and this implies our claim by his [35, Proposition 6.2].

Remark 3.6.15. For every split support, the composite sp si is an idempotent. As a consequence of Lemma 3.6.6, we also have

$$(3.87)$$

This is an instance of the *relative positivity* axiom [17, Section 2.5] which automatically holds in this case. Since we have $\text{si} \approx p$, we can also write this as

$$(3.88)$$

In the parlance of Definition 4.1.1, this implies that the idempotent sp si is of a special type, namely a *static* idempotent. We study these in more in detail in Section 4.3.

4 Idempotents and their splitting in Markov categories

In applications of probability, one often deals with certain operations—such as averaging over a group action or applying a conditional expectation—which are idempotent: Applying such an operation twice is the same as applying it only once.

The main goal of this section, which turns out to be closely related to supports, is to prove that all idempotents split in Markov categories satisfying suitable basic axioms. We achieve this in Theorem 4.4.3, and obtain as a straightforward consequence that idempotents in BorelStoch split. As we explain with Theorem 4.4.6, this strengthens an existing theorem of Blackwell on idempotent Markov kernels.

Being able to state and prove these statements requires a good deal of preparation. First of all, Section 4.1 introduces various ways in which an idempotent can interact with a Markov category structure, resulting in the definition of *static*, *strong* and *balanced* idempotents. We provide various examples, such as Reynolds operators, as well as reformulations of the definitions involving notions like detailed balance. In the auxiliary Section 4.3, we study splittings of static idempotents and show that the splitting of a static idempotent is the same as a (split) support. This is then put to use in Section 4.4, which contains our main results. Our construction of splittings of the more general balanced idempotents in the proof of Theorem 4.4.3 is reduced to the splitting of static idempotents from the previous subsection.

The final Section 4.5 offers a general formal construction for the splitting of balanced idempotents which we call the *Blackwell envelope*. It is analogous to the Karoubi envelope from ordinary category theory, which embeds every category in a larger category in which all idempotents split.

4.1 Static, strong and balanced idempotents

In a generic category, an **idempotent** is any endomorphism $e: X \rightarrow X$ satisfying $e^2 = e$, where e^2 stands for the sequential composite of two instances of e . As one would expect in a more specific setting like that of Markov categories, relevant notions of idempotents satisfy additional compatibility conditions with the extra structure that is present.

Here, we present three types of idempotents, namely *static*, *strong* and *balanced*. These have the following significance:

- Static idempotents are closely related to supports, as investigated in Section 4.3. Moreover, they play a crucial role in Theorem 4.4.3 on splitting of idempotents, which is the main result of this section. Indeed, a preliminary result (Lemma 4.4.2) reduces the proof of the splitting of all idempotents to those that are static.
- Strong idempotents have connections with conditional expectations, as we point out in Example 4.1.7, but they appear only in our present general theory.
- The weakest compatibility condition is that of balanced idempotents. As shown in Figure 1, these contain both static and strong idempotents as special cases. In a positive Markov category, also all split idempotents are balanced (Theorem 4.2.1). Moreover, assuming the Cauchy–Schwarz property (Definition A.5.1), *every* idempotent is balanced (Theorem 4.1.8).

Definition 4.1.1. *An idempotent $e: X \rightarrow X$ in a Markov category \mathcal{C} is:*

- *static if it satisfies*

$$(4.1)$$

- *strong if it satisfies*

$$(4.2)$$

- *balanced if it satisfies*

$$(4.3)$$

To understand these equations intuitively, note that the left-hand side is the same for all, and it amounts to considering the Markov chain with transition kernel e for two time steps and recording the intermediate value. Each of these equations implies that e is an idempotent, by marginalizing either on the right (static, strong) or left (balanced).

The reasons behind our choice of names for these idempotents are as follows:




Figure 1: Venn diagram of types of idempotents in a Markov category.

- The defining Equation (4.1) of a static idempotent can also be written as $e =_e \text{id}_X$. This is static in the sense that the Markov chain already stabilizes after one step.
- A balanced idempotent satisfies a version of *detailed balance*. This is discussed in the upcoming Remark 4.1.11 and reflected in the more symmetrical characterization from Proposition 4.1.10 (ii).
- The definition of strong idempotents arises as a strengthening of the characterization of balanced idempotents via Proposition 4.1.10 (iii).

These interpretations are also bolstered by Theorem 4.2.1, which classifies split idempotents by their types. For now, let us discuss the relation between the three types of idempotents just introduced and give some examples.

Remark 4.1.2. We can infer from the definition that both static and strong idempotents are balanced. Indeed, in the static case, we can compose Equation (4.1) with e on the right output to get the mirror image of Equation (4.3), which is an equivalent equation by the commutativity of copy. In the strong case, we simply pre-compose Equation (4.2) with e and use $e^2 = e$ to obtain Equation (4.3).

Furthermore, any idempotent that is both static and strong is deterministic, since that is an obvious consequence of chaining Equations (4.1) and (4.2). The Venn diagram of Figure 1 depicts all of these implications.

Example 4.1.3. Let us consider *group averaging* as a general motivating example in FinStoch . To this end, let G be a finite group acting on a finite set X . Then the G -averaging is the stochastic matrix $X \rightarrow X$ with entries

$$e(y|x) = \frac{1}{|G|} \sum_{g \in G} \delta_{gx,y}, \quad (4.4)$$

where $\delta_{a,b}$ stands for $\text{id}(a|b)$. Intuitively, this amounts to choosing a uniformly random element $g \in G$ and applying it to x , or equivalently choosing a uniformly random element in the orbit Gx . A direct calculation shows that this e is a strong idempotent: Looking at the probability for input x and output (y, z) , both the left-hand side and the right-hand side of Equation (4.2) take the value 0 if these three points are not all in the same orbit, and the value $|G_x|^2 / |G|^2 = 1 / |Gx|^2$ if they are, where $G_x = \{g \in G \mid gx = x\}$ denotes the stabilizer of x .

Example 4.1.4. We show, via examples, that in `FinStoch` all three types of idempotents actually differ. More precisely, all regions of the Venn diagram of Figure 1, besides the outer “general idempotents” one, are inhabited by some stochastic matrices. We also give a **splitting** for each of the examples, i.e. a pair π, ι of morphisms such that both $\pi \iota = \text{id}$ and $\iota \pi = e$ hold. Computations are omitted for the sake of brevity.

- A strong but not static idempotent is

$$e = \begin{pmatrix} 1/2 & 1/2 \\ 1/2 & 1/2 \end{pmatrix}, \quad \iota = \begin{pmatrix} 1/2 \\ 1/2 \end{pmatrix}, \quad \pi = (1 \quad 1). \quad (4.5)$$

This amounts to a stochastic map which ignores its input and outputs a fair coin flip instead.

- A static but not strong idempotent is

$$e = \begin{pmatrix} 1 & 0 & 1/2 \\ 0 & 1 & 1/2 \\ 0 & 0 & 0 \end{pmatrix}, \quad \iota = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \pi = \begin{pmatrix} 1 & 0 & 1/2 \\ 0 & 1 & 1/2 \end{pmatrix}. \quad (4.6)$$

This stochastic map acts as the identity on the first two states and sends the third state to a fair coin flip on the first two.

- A balanced idempotent that is neither static nor strong is

$$e = \begin{pmatrix} 1/2 & 1/2 & 0 & 1/4 \\ 1/2 & 1/2 & 0 & 1/4 \\ 0 & 0 & 1 & 1/2 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \iota = \begin{pmatrix} 1/2 & 0 \\ 1/2 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \pi = \begin{pmatrix} 1 & 1 & 0 & 1/2 \\ 0 & 0 & 1 & 1/2 \end{pmatrix}. \quad (4.7)$$

The structure of this map is easier to understand based on Remark 4.4.8 presented further on.

Note that π is deterministic for the strong idempotent and ι is deterministic for the static idempotent, while neither of these statements is true for the balanced idempotent. These are general facts, see Theorem 4.2.1.

Example 4.1.5. In order to populate the remaining region in Figure 1, let us present two examples of idempotents in `SetMulti` which are not balanced:

- On a two-element set $X = \{0, 1\}$, consider the multivalued function $e: X \rightarrow X$ given by $e(0) = \{0, 1\}$ and $e(1) = \{1\}$. Then e is an idempotent, but it is not balanced, as one can see by direct calculation: The right-hand side of Equation (4.3) relates $0 \in X$ to $(0, 1) \in X \times X$, but the left-hand side does not.
- Consider $X = \mathbb{R}$ and the strict ordering relation $>$ on real numbers. Define e to be the morphism mapping each number to the upset it generates:

$$e(x) = \{y \mid y > x\}. \quad (4.8)$$

This is another idempotent that is not balanced: For any ordered triple $z > y > x$, the right-hand side of Equation (4.3) relates x to (y, z) , but the left-hand side does not. This example works with any nontrivial dense¹⁷ partially ordered set in place of $(\mathbb{R}, >)$.

¹⁷Density means that for every ordered pair $z > x$ there is an element y satisfying $z > y > x$ [23, Definition 10.5]. This is relevant in order to get $e^2 = e$. We need the partial order to be nontrivial only in that there should be *some* nontrivial order relation $y > x$.

Example 4.1.6. Also FinStoch_\pm has idempotents which are not balanced, such as

$$e = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 1 \end{pmatrix}, \quad (4.9)$$

Indeed upon indexing the rows and columns as $X = \{a, b, c\}$, the right-hand side of Equation (4.3) takes a to (a, b) with probability 1, while this transition has probability 0 on the left-hand side.

Example 4.1.7. It is instructive to spell out the defining equations of the three types of idempotents in the Markov category $\text{CRing}_+^{\text{op}}$ from Example 2.2.11. This means that we consider a commutative ring R and an additive map $e: R \rightarrow R$ satisfying $e(1) = 1$. Then we have:

- e is a static idempotent if and only if

$$e(e(a)b) = e(ab) \quad (4.10)$$

holds for all $a, b \in R$.

- e is a strong idempotent if and only if

$$e(e(a)b) = e(a)e(b) \quad (4.11)$$

holds for all $a, b \in R$. This is precisely the defining equation of an *averaging operator* (also known as a *Reynolds operator*)¹⁸ on R [2, 31]. Moreover, Equation (4.11), together with linearity and continuity conditions, characterizes *conditional expectations* on the algebra of bounded measurable functions on a probability space [30].

- e is a balanced idempotent if and only if

$$e(e(a)b) = e(e(a)e(b)) \quad (4.12)$$

holds for all $a, b \in R$.

More general examples of idempotents arising in categorical probability can be found after Theorem 4.2.1.

We now turn to a conceptually important result stating that all idempotents are balanced in Markov categories formalizing probability theory, either in the discrete setting or via measure theory.

Theorem 4.1.8. *Let \mathcal{C} be a Markov category. If \mathcal{C} satisfies the Cauchy–Schwarz Property (Definition A.5.1), then every idempotent in \mathcal{C} is balanced.*

The proof is essentially trivial.

Proof. Let e be any idempotent in \mathcal{C} . In the Cauchy–Schwarz Property from Equation (A.41), take $f = g = h = e$. Then the antecedent of Implication (A.41) holds by the assumed $e^2 = e$. Its consequent tells us that e is a balanced idempotent. \square

¹⁸Reynolds operators are frequently defined by a more general equation which does not imply they are idempotent, but which is equivalent to Equation (4.11) in the idempotent case [2].

Corollary 4.1.9. *Let \mathcal{C} be FinStoch , BorelStoch , Stoch , TopStoch , TychStoch , or CHausStoch . Then every idempotent in \mathcal{C} is balanced.*

Proof. This is a direct application of Theorem 4.1.8, since all these Markov categories have the Cauchy–Schwarz Property (Proposition A.5.2 and Corollary A.5.3). \square

We conclude this subsection by giving alternative descriptions of balanced and static idempotents.

Proposition 4.1.10 (Alternative description of balanced idempotents). *For an idempotent e , the following are equivalent:*

- (i) e is balanced.
- (ii) We have

$$(4.13)$$

- (iii) Equation (4.2) holds e -almost surely, i.e. we have

$$(4.14)$$

- (iv) For every morphism $p: A \rightarrow X$ that is invariant, i.e. satisfies $ep = p$, we have

$$(4.15)$$

Proof.

- (i) \Rightarrow (ii): As noted in Remark 4.1.2, applying Equation (4.3) twice yields the required

$$(4.16)$$

- (ii) \Rightarrow (i): This follows by applying e to the left output in Equation (4.13).

(i) \Rightarrow (iii): Using the associativity of copying as well as Equations (4.2) and (4.16), we prove Equation (4.14) via the following equalities:

$$(4.17)$$

This is the desired statement.

(iii) \Rightarrow (i): Whenever e satisfies Equation (4.14), Equation (4.3) defining balanced idempotents follows simply by marginalizing the extra output of that equation and using $e^2 = e$.

(ii) \Rightarrow (iv): We can use the invariance of p followed by Equation (4.13) and the invariance of p again:

$$(4.18)$$

(iv) \Rightarrow (ii): Since e is assumed to be an idempotent, it is itself invariant, so that we can set $p = e$ in Equation (4.15) to get Equation (4.13). \square

Remark 4.1.11. In point-separable categories such as **Stoch**, condition (iv) can be equivalently checked just on invariant states, in which case it looks like

$$(4.19)$$

which says that e satisfies *detailed balance* with respect to p in the language of Markov processes [27, Section 4.10]. This is also equivalent to saying that the morphism e is its own Bayesian inverse with respect to p . By Proposition 4.1.10, balanced idempotents in **Stoch** are thus those

idempotents satisfying *detailed balance* for every invariant state, i.e. ones for which the dynamics is *microscopically reversible*.

Alternatively, Equation (4.19) can also be interpreted as saying that e is *self-adjoint*. Indeed, if p is an invariant measure for a balanced idempotent Markov kernel $e: X \rightarrow X$, then the operator $L^2(X, p) \rightarrow L^2(X, p)$ induced by pre-composition with e is idempotent and self-adjoint, making it an orthogonal projection on the Hilbert space $L^2(X, p)$.

We have the following weaker formulation of the definition of static idempotent.

Lemma 4.1.12 (Alternative description of static idempotents). *Let e be an idempotent. Then e is static if and only if it is e -a.s. deterministic.*

Proof. Let us consider an idempotent e which is e -a.s. deterministic. We can then use this assumption together with $e^2 = e$ to get

$$(4.20)$$

The desired Equation (4.1) can then be derived as follows.

$$(4.21)$$

Conversely, a static idempotent e is balanced, as noted in Remark 4.1.2. Therefore, we have

$$(4.22)$$

by Proposition 4.1.10, which says exactly that e is e -a.s. deterministic. \square

4.2 Split idempotents in Markov categories

The goal of this and the following subsection is mainly to study **splitting** of idempotents in Markov categories and to find criteria which guarantee that all (balanced) idempotents split. Recall first that a splitting of a generic morphism e in a category is a factorization $e = \iota \pi$ such that $\pi \iota$ is the identity morphism [4, Section 6.5]. Clearly, if e has such a splitting, then it is an idempotent, and the splitting is unique up to unique isomorphism. In a Markov category, the types of idempotents from Definition 4.1.1 are reflected in properties of a splitting as follows.

Theorem 4.2.1. *In a positive Markov category \mathbf{C} , let $e: X \rightarrow X$ be an idempotent that splits as $e = \iota \pi$ for $\pi: X \rightarrow T$ and $\iota: T \rightarrow X$.*

(i) *e is static if and only if ι is deterministic.*¹⁹

¹⁹The “if” part holds even without assuming that \mathbf{C} is positive (as our proof shows).

(ii) e is strong if and only if π is deterministic.

(iii) e is balanced (without further assumptions).

Moreover, π is necessarily ι -a.s. deterministic.

Proof. Since $\pi \iota$ is the identity and thus deterministic, the positivity assumption on C gives

$$(4.23)$$

(i) Using the assumption that ι is deterministic twice, we get

$$(4.24)$$

and thus obtain that e is static.

Conversely, using Equation (4.23) and the assumption that e is static yields

$$(4.25)$$

and the claim follows by cancelling the epimorphism π at the bottom.

(ii) If π is deterministic, we calculate

$$(4.26)$$

where the first step is by Equation (4.23) and the second is by assumption. Thus, e is a strong idempotent.

Conversely, if e is strong, then we similarly get

$$(4.27)$$

and the claim follows by post-composing with π on both outputs, which cancels both the monomorphisms ι at the top.

(iii) We apply Equation (4.23) twice to obtain

$$(4.28)$$

so that the claim follows by Proposition 4.1.10.

In order to show that π is ι -a.s. deterministic, we use that e is balanced together with Proposition 4.1.10 to obtain

$$(4.29)$$

Post-composing with $\pi \otimes \pi$ cancels the two ι morphisms on top, and therefore π is e -a.s. deterministic. Since we have $\iota = e\iota$ by assumption and $e\iota \ll e$ by Lemma 2.3.1, we conclude that π is also ι -a.s. deterministic. \square

Example 4.2.2. Since the idempotents of Example 4.1.5 are not balanced although SetMulti is positive, it follows that these idempotents do not split. Note that the first one of these already lives in FinSetMulti .

Example 4.2.3. In any Markov category, a morphism of the form

$$(4.30)$$

is a strong idempotent by direct calculations. Under positivity, this idempotent is immediately strong because it is defined in terms of a splitting across X (dashed line) with a deterministic projection given by $\text{id}_X \otimes \text{dely}_Y$.

Example 4.2.4. Let \mathbf{C} be a representable positive Markov category. Then the composite morphism $\delta \text{samp}: PX \rightarrow PX$ is a static idempotent for every object X because δ is deterministic and samp is its left inverse.

Example 4.2.5. Let \mathbf{C} be a positive²⁰ Markov category and p a morphism with a split support. Then the composition of its support projection and injection

$$e := \text{si}_p \text{sp}_p \tag{4.31}$$

is a static idempotent, since $\text{sp}_p \text{si}_p = \text{id}_S$ and si_p is deterministic.

4.3 Supports and splittings of static idempotents

We now focus further on the case of static idempotents and show that for them, splitting is closely related to supports. In the following subsection, we will put these results to use in order to construct splittings of more general idempotents under suitable additional assumptions.

Lemma 4.3.1. *In any Markov category, let $e: X \rightarrow X$ be a static idempotent and let $p: A \rightarrow X$ be a morphism with $p \ll e$. Then $ep = p$.*

Proof. e being static means $e =_e \text{id}_X$, so that the assumption $p \ll e$ gives $e =_p \text{id}_X$. By marginalizing the output of p in this equation, we get $ep = p$. \square

Proposition 4.3.2 (Support and splitting of static idempotents). *Let e be a static idempotent in a positive Markov category. Then the following are equivalent:*

- (i) e is a split idempotent.
- (ii) e has a split support.
- (iii) e has a support.

Moreover, this equivalence is such that the split support gives the splitting via $e = \text{si}_e \text{sp}_e$.

Proof. We first show that (i) implies (ii). Suppose that e splits as $e = \iota \pi$ for $\pi: X \rightarrow T$ and $\iota: T \rightarrow X$ satisfying $\pi \iota = \text{id}_T$. We know from Theorem 4.2.1 that ι is a deterministic split monomorphism, and we have $\iota \ll e$ by Lemma 2.3.1 and $\iota = e \iota$. Thus, by the support criterion of Lemma 3.1.3, it is enough to show that any $p: A \rightarrow X$ satisfying $p \ll e$ also factors across ι . Since we have $p = ep$ by Lemma 4.3.1, we can indeed write $p = \iota \pi p$.

The implication from (ii) to (iii) is trivial.

Finally, assume that the static idempotent e has a support with inclusion $\text{si}: S \rightarrow X$ and let us prove that e splits. Since $e =_e \text{id}$ holds by definition of static idempotents, we obtain $e \text{si} = \text{si}$ by Lemma 3.1.8. The support factorization of Equation (3.4) reads as $e = \text{si} \widehat{e}$, and therefore also

$$\text{si} \widehat{e} \text{si} = e \text{si} = \text{si} \tag{4.32}$$

holds. Since si is a monomorphism, we get $\widehat{e} \text{si} = \text{id}$, so that $e = \text{si} \widehat{e}$ is indeed a split idempotent. \square

²⁰The attentive reader may notice that the positivity assumption is redundant here by Footnote 19.

Example 4.3.3. If a morphism p has a split support, then $\text{si}_p \text{sp}_p$ is a static idempotent by Example 4.2.5. Proposition 4.3.2 now allows us to conclude that this idempotent has the same split support as p itself.

Corollary 4.3.4. *Let \mathcal{C} be a positive Markov category and let $e: X \rightarrow X$ be an arbitrary morphism. Then the following are equivalent:*

- (i) e is a static idempotent that splits.
- (ii) We have $e = \text{si}_p \text{sp}_p$ for some morphism p with a split support.

Proof. If e is a static idempotent that splits, then by Proposition 4.3.2 we have $e = \text{si}_e \text{sp}_e$. The other direction is Example 4.2.5. \square

We can apply this to obtain another characterization of split supports similar to Lemma 3.6.6, but where the second condition only requires one direction of implication.

Theorem 4.3.5 (Split supports via static idempotents). *Let \mathcal{C} be a positive Markov category and $p: A \rightarrow X$ a morphism in \mathcal{C} . Then the following are equivalent:*

- (i) p admits a split support.
- (ii) There exists a split static idempotent $e: X \rightarrow X$ satisfying $ep = p$, and such that for all $f, g: W \otimes X \rightarrow Y$ with arbitrary W and Y , we have

$$\begin{array}{c}
 \begin{array}{ccc}
 Y & & X \\
 | & & | \\
 \boxed{f} & & \text{---} \\
 | & \text{---} & \bullet \\
 | & & | \\
 W & & A \\
 \boxed{p} & & \\
 | & & \\
 & &
 \end{array}
 =
 \begin{array}{ccc}
 Y & & X \\
 | & & | \\
 \boxed{g} & & \text{---} \\
 | & \text{---} & \bullet \\
 | & & | \\
 W & & A \\
 \boxed{p} & & \\
 | & & \\
 & &
 \end{array}
 \implies
 \begin{array}{ccc}
 Y \\
 | \\
 \boxed{f} \\
 | \\
 W \quad X \\
 \boxed{e} \\
 | \\
 W \quad X
 \end{array}
 =
 \begin{array}{ccc}
 Y \\
 | \\
 \boxed{g} \\
 | \\
 W \quad X \\
 \boxed{e} \\
 | \\
 W \quad X
 \end{array}
 \quad (4.33)
 \end{array}$$

Proof. Assuming (i), take $e := \text{si}_p \text{sp}_p$, which is a split static idempotent as in Example 4.2.5. Then $ep = p$ holds by the second equality in (3.75) and Implication (4.33) is a direct consequence of Lemma 3.6.6.

Conversely, suppose that (ii) holds with a splitting given by $e = \iota\pi$. By Lemma 2.4.2, we have the absolute bicontinuity relation $e = \iota\pi \approx \iota$. So if we can show that its left-hand side e is absolutely bicontinuous to p , then p has a split support by Corollary 3.6.11, since ι is indeed a deterministic split monomorphism (see Theorem 4.2.1).

To this end, note that $ep \ll e$ holds by Lemma 2.3.1, so that we get $p \ll e$ by the assumed $ep = p$. To prove absolute continuity in the other direction, suppose that we have $f =_p g$ and let us show that necessarily also $f =_e g$ holds. Using Implication (4.33) and the fact that $\text{id}_W \otimes \pi$ is an epimorphism, we get

$$\begin{array}{ccc}
 Y \\
 | \\
 \boxed{f} \\
 | \\
 W \quad T \\
 \boxed{\iota} \\
 | \\
 W \quad T
 \end{array}
 =
 \begin{array}{ccc}
 Y \\
 | \\
 \boxed{g} \\
 | \\
 W \quad T \\
 \boxed{\iota} \\
 | \\
 W \quad T
 \end{array}
 \quad (4.34)$$

By the second part of Lemma A.3.3, Equation (4.34) is equivalent to $f =_\iota g$, which directly implies the desired $f =_e g$ by pre-composition with π . In conclusion, we thus have both $p \ll e$ and $e \ll p$, so that p has a split support by the above argument. \square

4.4 General conditions for the splitting of idempotents

We now turn to the question of when *general* idempotents split in a Markov category, and we will derive sufficient conditions in order for this to be the case. The resulting Theorem 4.4.3 is the main result of this section. It implies in particular that all idempotents in **BorelStoch** split (Corollary 4.4.5). As far as we know, this is an entirely new result which has not been proven by measure-theoretic methods before, and we note in Theorem 4.4.6 that it strengthens an existing result of Blackwell on idempotent Markov kernels.

To this end, we assume familiarity with the *equalizer principle* from Section 3.5 as well as with the notion of *observationally representable* Markov category from Appendix A.4. Roughly speaking, the latter postulates that one can distinguish between morphisms by iterated sampling.

Lemma 4.4.1. *Let e be a balanced idempotent in an observationally representable Markov category \mathcal{C} . Then $e^\sharp e =_e e^\sharp$.*

Proof. By the definition of observationally representable Markov categories and Proposition A.4.2, it suffices to show the desired equation for n independent samples of both sides, i.e. post-composed with $\text{samp}^{(n)}$ defined in (A.30) for every natural number n . Since e^\sharp is deterministic and satisfies $\text{samp} e^\sharp = e$, we only need to show that

$$\begin{array}{c} \overbrace{\begin{array}{ccc} \boxed{e} & \cdots & \boxed{e} \end{array}}^{n \text{ outputs}} \\ \downarrow \\ \boxed{e} \end{array} =_e \begin{array}{c} \overbrace{\begin{array}{ccc} \boxed{e} & \cdots & \boxed{e} \end{array}}^{n \text{ outputs}} \\ \downarrow \end{array} \quad (4.35)$$

holds for any number of outputs n . This can be proven by induction. The case $n = 1$ just expresses that e is idempotent. For the induction step, we compute

$$\begin{array}{c} \overbrace{\begin{array}{ccc} \boxed{e} & \cdots & \boxed{e} \end{array}}^{n+1 \text{ outputs}} \\ \downarrow \\ \boxed{e} \end{array} = \begin{array}{c} \begin{array}{ccc} & \boxed{e} & \\ & \downarrow & \\ \boxed{e} & \cdots & \boxed{e} \end{array} \\ \downarrow \\ \boxed{e} \end{array} =_e \begin{array}{c} \begin{array}{ccc} & \boxed{e} & \\ & \downarrow & \\ \boxed{e} & \cdots & \boxed{e} \end{array} \\ \downarrow \end{array} =_e \begin{array}{c} \overbrace{\begin{array}{ccc} \boxed{e} & \cdots & \boxed{e} \end{array}}^{n+1 \text{ outputs}} \\ \downarrow \end{array}$$

where the second equation uses the induction assumption and the other steps follow from the characterization of balanced idempotents in Proposition 4.1.10 (iii). \square

Splittings of balanced idempotents can be constructed as in the following lemma, which plays a crucial role in the proof of the forthcoming Theorem 4.4.3.

Lemma 4.4.2. *Let \mathcal{C} be a positive and observationally representable Markov category. If every static idempotent in \mathcal{C} splits, then also every balanced idempotent in \mathcal{C} splits.*

Proof. Consider an arbitrary balanced idempotent $e: X \rightarrow X$. Applying the distribution functor to it gives $Pe: PX \rightarrow PX$, which is also an idempotent by the functoriality of P and idempotency of e :

$$(Pe)(Pe) = P(ee) = Pe. \quad (4.36)$$

In particular, since Pe is a deterministic morphism by definition, it is both a static and a strong idempotent. By assumption, we thus have a splitting $Pe = \iota\pi$ with $\pi\iota = \text{id}_T$. Since Pe is static and strong, $\iota: T \rightarrow PX$ and $\pi: PX \rightarrow T$ are both deterministic by Theorem 4.2.1.

Let us now prove the following auxiliary equations:

$$\text{samp } \iota\pi\delta = e, \quad (4.37)$$

$$\text{samp } \iota = e \text{ samp } \iota, \quad (4.38)$$

$$\pi\delta = \pi e^\sharp. \quad (4.39)$$

The first one can be verified as

$$\text{samp } \iota\pi\delta = \text{samp } (Pe)\delta = e \text{ samp } \delta = e, \quad (4.40)$$

where the first equation is by definition of ι and π , the second by naturality of samp and the third by $\text{samp } \delta = \text{id}_X$. Equation (4.38) follows via

$$\text{samp } \iota = \text{samp } (Pe)\iota = e \text{ samp } \iota, \quad (4.41)$$

where we first use $\iota = (Pe)\iota$ and then the naturality of samp . Lastly, Equation (4.39) can be derived similarly via

$$\pi\delta = \pi(Pe)\delta = \pi e^\sharp, \quad (4.42)$$

where we use $\pi = \pi(Pe)$ and then $(Pe)\delta = e^\sharp$.

We now employ these facts to show that the composite morphism

$$T \xrightarrow{\iota} PX \xrightarrow{\text{samp}} X \xrightarrow{\delta} PX \xrightarrow{\pi} T \quad (4.43)$$

is also a static idempotent. To this end, we compute

where the first equality uses that both π and δ are deterministic as well as Equation (4.37), and the second equality uses Equations (4.38) and (4.39). We can further reduce the right-hand

side to

$$= \quad = \quad = \quad (4.45)$$

where the first step is by Lemma 4.4.1, the second by Equations (4.38) and (4.39) again, and the third by π and δ being deterministic. This proves that the composite morphism is indeed a static idempotent.

Therefore, by assumption, we have another splitting consisting of morphisms κ and ρ with

$$\pi \delta \text{samp } \iota = \kappa \rho, \quad \rho \kappa = \text{id}. \quad (4.46)$$

Using Equation (4.37), we construct a splitting of the original idempotent e as

$$e = e e = \text{samp } \iota \pi \delta \text{samp } \iota \pi \delta = (\text{samp } \iota \kappa) (\rho \pi \delta). \quad (4.47)$$

This is indeed a splitting since we have

$$(\rho \pi \delta) (\text{samp } \iota \kappa) = \rho (\pi \delta \text{samp } \iota) \kappa = \rho \kappa \rho \kappa = \text{id}, \quad (4.48)$$

which concludes the proof. \square

Lemma 4.4.2 reduces the problem of splitting balanced idempotents to the problem of splitting static idempotents. Fortunately, the equalizer principle from Section 3.5 is an additional condition which guarantees that all static idempotents split. We therefore obtain the following main result on idempotent splitting in Markov categories.

Theorem 4.4.3. *Let \mathbf{C} be a Markov category such that*

- (i) \mathbf{C} is positive,
- (ii) \mathbf{C} is observationally representable, and
- (iii) \mathbf{C} satisfies the equalizer principle (Definition 3.5.1).

Then an idempotent in \mathbf{C} splits if and only if it is balanced. In particular, if \mathbf{C} also has the Cauchy–Schwarz Property, then all idempotents split.

Proof. Every split idempotent in a positive Markov category is balanced by Theorem 4.2.1, so we focus on showing that all balanced idempotents split.

By Lemma 4.4.2, it is enough to show that every static idempotent splits. To this end, consider a static idempotent $e: X \rightarrow X$ and the equalizer in \mathbf{C}_{det} given by

$$T \xrightarrow{\iota} X \xrightarrow[\delta]{e^\#} PX \quad (4.49)$$

As our notation already suggests, the deterministic morphism ι will eventually be the monomorphism part of a splitting of e .

Since e is a static idempotent, we have $e =_e \text{id}$ by definition. Because \mathbf{C} is a.s.-compatibly representable by Proposition A.4.2 and $\text{id}^\# = \delta$, we can infer

$$e^\# =_e \delta. \quad (4.50)$$

Using the equalizer principle, the a.s. equality of the parallel pair from diagram (4.49) means that we have a factorization $e = \iota \pi$ for some $\pi: X \rightarrow T$.

To show that this provides the requisite splitting, we still need to prove $\pi \iota = \text{id}_T$. First note that we have

$$\iota \pi \iota = e \iota = \text{samp } e^\# \iota = \text{samp } \delta \iota = \iota. \quad (4.51)$$

In order to conclude the proof, it is enough to argue that ι is a monomorphism in \mathbf{C} .²¹ For this purpose, consider two arbitrary morphisms f and g satisfying $\iota f = \iota g$. By Equation (4.51) and Lemma 2.3.1, we have $\iota f = e \iota f \ll e$. Combining this absolute continuity relation with Equation (4.50) yields $e^\# =_{\iota f} \delta$. The equalizer principle thus says that the factorization of ιf through ι is unique. Therefore $\iota f = \iota g$ implies $f = g$. Consequently, ι is a monomorphism and thus Equation (4.51) implies the desired $\pi \iota = \text{id}_T$.

The last sentence of the theorem statement is now an immediate consequence of Theorem 4.1.8. \square

Our main application of Theorem 4.4.3 is to **BorelStoch**, which we consider shortly. Before that, it is instructive to note an example application in which *not* all idempotents split due to the failure of the Cauchy-Schwarz property.

Corollary 4.4.4. *Every balanced idempotent in **FinSetMulti** splits, but not all idempotents split.*

Proof. **FinSetMulti** satisfies the assumptions of Theorem 4.4.3:

- (i) It has conditionals [12, Proposition 16], and this implies positivity [11, Lemma 11.24].
- (ii) It is observationally representable by Example A.4.6.
- (iii) It satisfies the equalizer principle by Remark 3.5.2.

Hence the split idempotents in **FinSetMulti** are exactly the balanced idempotents. However, by Example 4.2.2, there are non-balanced idempotents in **FinSetMulti**, so not all idempotents split by Theorem 4.2.1. \square

We also already know that **BorelStoch** satisfies all of the relevant assumptions, so here is our previously announced key result.

Corollary 4.4.5. *Every idempotent in **BorelStoch** is balanced and splits.*

Proof. **BorelStoch** has the Cauchy–Schwarz Property (Proposition A.5.2) and also satisfies the other hypotheses of Theorem 4.4.3, since:

- (i) It is positive ([11, Example 11.25]).
- (ii) It is observationally representable (Example A.4.5).
- (iii) It satisfies the equalizer principle (Proposition 3.5.4). \square

²¹It is clear that ι is a monomorphism in \mathbf{C}_{det} by virtue of being an equalizer, but this does not directly imply that it is a monomorphism in \mathbf{C} as well.

In order to put this result into context, let us note that it strengthens the following theorem due to Blackwell.

Theorem 4.4.6 ([3, Theorem 7]). *Let $e: X \rightarrow X$ be an idempotent Markov kernel on a standard Borel space X . Then there exists a partition*

$$X = N \sqcup \bigsqcup_{t \in T} X_t \tag{4.52}$$

of X into measurable sets²² such that:

- (i) $e(X_t|x) = 1$ for all $x \in X_t$ and all $t \in T$.
- (ii) The probability measure $e(_||x)$ is independent of x within each X_t .
- (iii) $e(N|x) = 0$ for all $x \in X$.

The following reduction of this result to Corollary 4.4.5 amounts to translating the existence of a splitting into measure-theoretic language.

Proof. By Corollary 4.4.5, we know that e can be split as $e = \iota \pi$ for some $\pi: X \rightarrow T$ and $\iota: T \rightarrow X$ with $\pi \iota = \text{id}$. Consider the set of all $x \in X$ for which $\pi(_||x)$ is a Dirac delta measure, or equivalently for which

$$\pi(U|x)^2 = \pi(U|x) \tag{4.53}$$

holds for all $U \in \Sigma_X$. Since Σ_X is countably generated, this set is itself measurable, so that we can define N to be its complement. Then because π is ι -a.s. deterministic by Theorem 4.2.1, we have $\iota(N|t) = 0$ for all $t \in T$.

If we now define

$$X_t := \{x \in X \setminus N \mid \pi(_||x) = \delta_t\} \tag{4.54}$$

for each $t \in T$, it is straightforward to check that the required properties hold. Indeed, for (i), we have for every $x \in X_t$,

$$e(X_t|x) = \int_{t' \in T} \iota(X_t|t') \pi(dt'|x) = \int_{t' \in T} \iota(X_t|t') \delta_t(dt') = \iota(X_t|t) = 1, \tag{4.55}$$

where the final step is due to $\pi \iota = \text{id}_T$. Property (ii) holds simply by the definition of X_t . Finally, (iii) is immediate from $\iota(N|t) = 0$ for all t . \square

Remark 4.4.7. It is worth emphasizing that Blackwell's Theorem 4.4.6 does not immediately imply that every idempotent splits, since T is merely an index set rather than a standard Borel measurable space itself. It is in this sense that our Corollary 4.4.5 is stronger than Blackwell's theorem.

Remark 4.4.8 (Blackwell's decomposition for stochastic maps). To gain better intuition for the decomposition of Equation (4.52), as well as for the type classification of idempotents in general, it may help to illustrate it for FinStoch using Markov chain terminology. In these terms, the set T can be understood as the set of *recurrent communication classes* of the transition

²²Note that "partition" here only means decomposition into a disjoint union of measurable sets, not decomposition into a coproduct of measurable spaces. For example for $e = \text{id}_{[0,1]}$, this partition (unique in this case) consists of all the singleton sets in $[0, 1]$.

kernel e . These classes are the non-empty sets $C \subseteq X$ satisfying $e(C|x) = 1$ for all $x \in C$, and such that C is minimal with respect to this property. The recurrent communication classes are mutually disjoint, and the complement of their union corresponds to N in Equation (4.52), which is exactly the set of transient states.

Then the projection π of a splitting $e = \iota \pi$ acts on a state x as follows:

- If $x \notin N$ — meaning that x is recurrent — then π maps x deterministically to the class containing x .
- If $x \in N$ — meaning that x is transient — then $\pi(_ | x)$ is arbitrary.

Given a class C as input, the inclusion ι samples an element from C according to a full-support distribution which only depends on C itself. In particular, the restriction of e to C is a stochastic matrix of rank one and full support.

These observations elucidate the first two items of Theorem 4.2.1 yet again differently:

- e is static if and only if ι is deterministic, which means that all recurrent communication classes are singletons.
- e is strong if and only if π is deterministic, which means that all states are recurrent (or equivalently $N = \emptyset$).

Remark 4.4.9. Given that we have proven idempotent splitting for **BorelStoch**, the reader may wonder whether idempotents split in **Stoch** in general. Here is what we know about this at present.

First of all, it is unlikely that **Stoch** is representable: **Stoch_{det}** does *not* correspond to **Meas** since e.g. the deterministic states $I \rightarrow X$ on a measurable space X are exactly the $\{0, 1\}$ -valued probability measures, and these do not coincide with the points of X in general. However, **Stoch** is still the Kleisli category of the Giry monad on **Meas**. This is a weaker form of representability which has been studied in [28], where it was also shown that a suitable notion of observability still applies in this case [28, Theorem 9.1]. The question is thus whether the proof of Theorem 4.4.3 still goes through in such weaker setting.

For Lemma 4.4.2, this does indeed seem to be the case. However, we do not know whether a suitable formulation of the equalizer principle still holds, and therefore we do not currently know whether the proof of Theorem 4.4.3 can be adapted accordingly.

In order to show how our Theorem 4.4.3 can also be applied fruitfully to other Markov categories, let us derive a similar conclusion for another Markov category of interest.

Corollary 4.4.10. *Every idempotent in **CHausStoch** is balanced and splits.*

Proof. This follows by an application of Theorem 4.4.3, since:

- (i) **CHausStoch** is positive, since even **TychStoch** is positive, of which **CHausStoch** is a full Markov subcategory. The positivity of **TychStoch** follows because it is causal (Example 3.1.18) and causality implies positivity [17, Theorem 2.24].
- (ii) **CHausStoch** is observationally representable (Proposition A.4.8).
- (iii) It satisfies the equalizer principle since even **TychStoch** does (Remark 3.5.2), of which **CHausStoch** is a full Markov subcategory.

Finally, **CHausStoch** also has the Cauchy-Schwarz property (Corollary A.5.3). □

It is worth noting that Corollary 4.4.10 is essentially known: Hamana proved in [22, Theorem 2.3] that every idempotent and Schwarz-positive unital map between unital C^* -algebras splits across another unital C^* -algebra. In the special case where the C^* -algebra on which the idempotent is defined is commutative, this recovers Corollary 4.4.10 thanks to probabilistic Gelfand duality [19].

Question 4.4.11. *Do idempotents still split in TychStoch?*

A quick look at the previous proof shows that the only missing ingredient here is to prove the observational representability of TychStoch.

4.5 The Blackwell envelope

When not all idempotents split in a generic category \mathcal{C} , it is natural to consider the *Karoubi envelope* $\mathcal{C}^{\mathcal{K}}$, which is the new category obtained by formally splitting all idempotents. The Karoubi envelope of a symmetric monoidal category can also be made symmetric monoidal in a canonical way. However, if \mathcal{C} is a Markov category, there does *not* seem to be any obvious way to turn $\mathcal{C}^{\mathcal{K}}$ into a Markov category as well. This works only if one restricts idempotents used to construct $\mathcal{C}^{\mathcal{K}}$ to the balanced idempotents. In this way, we indeed get a new Markov category which we call the *Blackwell envelope* of \mathcal{C} .

Let us recall the definition of the Karoubi envelope first, also known as *idempotent completion*.

Definition 4.5.1 (e.g. [4, Section 6.5]). *For any category \mathcal{C} , its **Karoubi envelope** $\mathcal{C}^{\mathcal{K}}$ is the category where:*

- *Objects are pairs (X, e_X) with $X \in \mathcal{C}$ and $e_X: X \rightarrow X$ an idempotent.*
- *Morphisms $f: (X, e_X) \rightarrow (Y, e_Y)$ are morphisms $X \rightarrow Y$ in \mathcal{C} satisfying $f e_X = f = e_Y f$.*

In order to move closer to Markov categories, let us consider symmetric monoidal structure next.

Proposition 4.5.2 (e.g. [4, Proposition 6.5.9]). *Let \mathcal{C} be a symmetric monoidal category. Then its Karoubi envelope $\mathcal{C}^{\mathcal{K}}$ inherits a canonical symmetric monoidal structure.*

Sketch of proof. Every functor between categories $F: \mathcal{C} \rightarrow \mathcal{D}$ has a natural extension to the Karoubi envelopes $F^{\mathcal{K}}: \mathcal{C}^{\mathcal{K}} \rightarrow \mathcal{D}^{\mathcal{K}}$ given by

$$F^{\mathcal{K}}(X, e_X) := (F(X), F(e_X)), \quad F^{\mathcal{K}}(f) := F(f). \quad (4.56)$$

Likewise, any natural transformation $\eta: F \Rightarrow G$ induces a natural transformation $\eta^{\mathcal{K}}: F^{\mathcal{K}} \Rightarrow G^{\mathcal{K}}$ with components²³

$$\eta_{(X, e_X)}^{\mathcal{K}} := G(e_X) \eta_X F(e_X). \quad (4.57)$$

Therefore, via the obvious definition $(X, e_X) \otimes (Y, e_Y) := (X \otimes Y, e_X \otimes e_Y)$, we can choose the expected associators, unitors and swap morphisms, and get the desired result. \square

Interestingly, there does not seem to be a canonical way to make the Karoubi envelope of a Markov category into a Markov category. However, the following modified construction works.

²³Note that, since η is natural, we have $\eta_{(X, e)}^{\mathcal{K}} = \eta_X F(e) = G(e) \eta_X$.

Definition 4.5.3. If \mathcal{C} is a Markov category, its **Blackwell envelope** $\mathcal{C}^{\mathcal{B}}$ is the full subcategory of $\mathcal{C}^{\mathcal{K}}$ whose objects are given by pairs (X, e_X) where e_X is a balanced idempotent.

We name this construction after Blackwell in honor of his Theorem 4.4.6, which was arguably the first result in the direction of idempotent splitting in measure-theoretic probability.

Proposition 4.5.4. Let \mathcal{C} be a Markov category. Then its Blackwell envelope $\mathcal{C}^{\mathcal{B}}$ inherits a canonical Markov category structure.

Proof. It is straightforward to see that if e_X and e_Y are balanced idempotents, then so is $e_X \otimes e_Y$. Also id_I is trivially a balanced idempotent. Through Proposition 4.5.2, $\mathcal{C}^{\mathcal{B}}$ therefore becomes a symmetric monoidal category. It is obvious that is again semicartesian.

Define now the copy morphism on an object (X, e_X) to be given by

$$\text{copy}_{(X, e_X)} := \begin{array}{c} \begin{array}{cc} X & X \\ \downarrow & \downarrow \\ \boxed{e_X} & \boxed{e_X} \end{array} \\ \downarrow \bullet \\ \boxed{e_X} \\ \downarrow \\ X \end{array} \quad (4.58)$$

Then all relevant properties are trivial except for the associativity of copy, which follows from the assumption that e_X is balanced. \square

In particular, if every idempotent in \mathcal{C} is balanced, then the Karoubi envelope coincides with the Blackwell envelope and is thus a Markov category. For example, this happens by Theorem 4.1.8 as soon as \mathcal{C} has the Cauchy-Schwarz property, which we already noted to be the case in standard probabilistic settings.

Remark 4.5.5. It is difficult to argue that $\mathcal{C}^{\mathcal{K}}$ cannot be a Markov category in general, since there could be many different ways of defining copy morphisms in general, even if one additionally requires the canonical inclusion $\mathcal{C} \rightarrow \mathcal{C}^{\mathcal{K}}$ to be a Markov functor. However, what we can say is that there is no way to achieve this for $\mathcal{C} = \text{SetMulti}$ if one also wants $\mathcal{C}^{\mathcal{K}}$ to be a *positive* Markov category.

The reason is that if there was a positive Markov category structure on $\mathcal{C}^{\mathcal{K}}$, then by Theorem 4.2.1 every idempotent in $\mathcal{C}^{\mathcal{K}}$ would be balanced. But with the inclusion $\mathcal{C} \rightarrow \mathcal{C}^{\mathcal{K}}$ being assumed a Markov functor, this would imply that every idempotent in \mathcal{C} is balanced. But this is not the case for $\mathcal{C} = \text{SetMulti}$ by Example 4.1.5.

Finally, we study the relationship between the Blackwell envelope and the causality axiom.

Lemma 4.5.6. Let \mathcal{C} be a Markov category, and consider morphisms

$$p: (X, e_X) \rightarrow (Y, e_Y), \quad f, g: (Y, e_Y) \rightarrow (Z, e_Z)$$

in its Blackwell envelope $\mathcal{C}^{\mathcal{B}}$. Then $f =_p g$ holds if and only if the same is true for the corresponding morphisms in \mathcal{C} .

Proof. In \mathcal{C} , we have

$$\begin{array}{c}
 \begin{array}{c}
 Z \quad Y \\
 | \quad | \\
 \boxed{f} \quad \boxed{f} \\
 | \quad | \\
 \bullet \\
 | \\
 \boxed{p} \\
 | \\
 X
 \end{array}
 \quad = \quad
 \begin{array}{c}
 Z \quad Y \\
 | \quad | \\
 \boxed{f} \quad \boxed{f} \\
 | \quad | \\
 \boxed{e_Y} \quad \boxed{e_Y} \\
 | \quad | \\
 \bullet \\
 | \\
 \boxed{e_Y} \\
 | \\
 \boxed{p} \\
 | \\
 X
 \end{array}
 \quad = \quad
 \begin{array}{c}
 Z \quad Y \\
 | \quad | \\
 \boxed{f} \quad \boxed{f} \\
 | \quad | \\
 \boxed{e_Y} \quad \boxed{e_Y} \\
 | \quad | \\
 \bullet \\
 | \\
 \boxed{e_Y} \\
 | \\
 \boxed{p} \\
 | \\
 X
 \end{array}
 \end{array}
 \tag{4.59}$$

because e_Y is a balanced idempotent by assumption. The analogous equations hold for g in place of f . Recalling the definition of the copy morphisms in $\mathcal{C}^{\mathcal{B}}$ as given in Equation (4.58), the result follows. \square

Proposition 4.5.7. *A Markov category \mathcal{C} is causal if and only if its Blackwell envelope $\mathcal{C}^{\mathcal{B}}$ is causal.*

Proof. Since \mathcal{C} is a Markov subcategory of $\mathcal{C}^{\mathcal{B}}$, it is obvious that whenever $\mathcal{C}^{\mathcal{B}}$ is causal, then the same is true for \mathcal{C} . To prove the converse, we use the characterization of causality from Lemma A.3.2.

Consider morphisms

$$\begin{aligned}
 f &: (A, e_A) \rightarrow (X, e_X), \\
 g &: (X, e_X) \rightarrow (Y, e_Y), \\
 h_1, h_2 &: (Y, e_Y) \rightarrow (Z, e_Z),
 \end{aligned}
 \tag{4.60}$$

all in the Markov category $\mathcal{C}^{\mathcal{B}}$. By Lemma 4.5.6, if $h_1 =_{gf} h_2$ holds, then the same is true in \mathcal{C} , so that we obtain $h_1 g =_f h_2 g$ in \mathcal{C} by Lemma A.3.2. Another application of Lemma 4.5.6 ensures that this holds also in $\mathcal{C}^{\mathcal{B}}$. In conclusion, $\mathcal{C}^{\mathcal{B}}$ is thus causal by Lemma A.3.2. \square

A Background on Markov categories

A.1 Some basics

Definition A.1.1. *Consider a symmetric monoidal category \mathcal{C} in which every object $X \in \mathcal{C}$ comes equipped with a **copy morphism** $\text{copy}_X: X \rightarrow X \otimes X$ and a **discard morphism** $\text{del}_X: X \rightarrow I$, where I is the monoidal unit, written in string diagram form as*

$$\begin{array}{c}
 \text{copy}_X = \begin{array}{c}
 X \quad X \\
 | \quad | \\
 \bullet \\
 | \\
 X
 \end{array}
 \quad \text{del}_X = \begin{array}{c}
 \bullet \\
 | \\
 X
 \end{array}
 \end{array}
 \tag{A.1}$$

Then \mathcal{C} is a **Markov category** if:

- (i) The triple $(X, \text{copy}_X, \text{del}_X)$ is a commutative comonoid for every object X .
- (ii) Every morphism $f: X \rightarrow Y$ satisfies $\text{del}_Y f = \text{del}_X$.

(iii) The copying of $X \otimes Y$ is obtained by the copying of X and Y in a natural way:

$$\begin{array}{c} X \otimes Y \quad X \otimes Y \\ \cup \\ \bullet \\ | \\ X \otimes Y \end{array} = \begin{array}{c} X \quad Y \quad X \quad Y \\ \cup \quad \cup \\ \bullet \quad \bullet \\ | \quad | \\ X \quad Y \end{array} \quad (\text{A.2})$$

It follows that the monoidal unit I is terminal, meaning that \mathbf{C} is a **semicartesian** monoidal category [11, Remark 2.3].

Definition A.1.2. Let \mathbf{C} be a Markov category. A **Markov subcategory** $\mathbf{D} \subseteq \mathbf{C}$ is a symmetric monoidal subcategory that is a Markov category with respect to the same copy and discard morphisms as \mathbf{C} .

Example A.1.3. Here we list some important examples of Markov categories. More details can be found in [11].

- (i) **FinStoch**, the category with finite sets as objects and stochastic matrices as morphisms. Its variant **FinStoch**_± is defined the same way, but allows negative entries in the matrices.
- (ii) **Stoch**, the category of all measurable spaces with measurable Markov kernels.
- (iii) **BorelStoch**, the category of standard Borel measurable spaces with measurable Markov kernels.
- (iv) **SetMulti**, the category of sets and multivalued functions, where a morphism $X \rightarrow Y$ is a function from X to the power set 2^Y such that each $f(x)$ is non-empty. We write **FinSetMulti** for the full subcategory whose objects are finite sets.
- (v) For a cartesian category \mathbf{D} and an affine symmetric monoidal monad P on \mathbf{D} , the associated Kleisli category $\mathbf{Kl}(P)$ is a Markov category.

For instance, the Giry monad on the category of measurable spaces **Meas** gives rise to **Stoch**, the Giry monad on the category of standard Borel spaces **BorelMeas** to **BorelStoch**, and the non-empty power set monad on **Set** produces **SetMulti**. See [18, Section 3] for more details.

Example A.1.4. Expanding on (v), another example of an affine commutative monad on a cartesian category is the probability monad on **Top**. It is given by taking τ -smooth Borel measures on topological spaces [16, Section 4]. We denote the resulting Kleisli category by **TopStoch**, which is equivalently the category of topological spaces and *continuous* Markov kernels, where the continuity is with respect to the *A-topology* on measures. In particular, a Markov kernel $p: A \rightarrow X$ is continuous if, for every open $U \subseteq X$, the map $a \mapsto p(U|a)$ is lower semi-continuous.

The resulting monad can be restricted to Tychonoff (meaning $T_{31/2}$) spaces, which include all metric spaces and compact Hausdorff spaces. In this case, the topology on each space of probability measures can be equivalently taken to be the weak topology induced by the integration maps against continuous functions [1]. We call the resulting Kleisli category **TychStoch**.

If we restrict the monad further to compact Hausdorff spaces, we obtain the so-called *Radon monad* [41]. Its Kleisli category, which we denote by **CHausStoch**, can equivalently be seen as the opposite of the category of commutative unital C^* -algebras and positive unital maps, a generalization of the usual Gelfand duality to the stochastic case (see [19] and [32] for details).

Another central definition is the one of deterministic morphism.

Definition A.1.5. A morphism $f: X \rightarrow Y$ is **deterministic** if it satisfies

$$\begin{array}{c} Y \quad Y \\ \cup \\ \bullet \\ \downarrow \\ \boxed{f} \\ \downarrow \\ X \end{array} = \begin{array}{c} Y \quad Y \\ \downarrow \quad \downarrow \\ \boxed{f} \quad \boxed{f} \\ \cup \\ \bullet \\ \downarrow \\ X \end{array} \tag{A.3}$$

The deterministic morphisms form a cartesian monoidal subcategory $\mathcal{C}_{\text{det}} \subseteq \mathcal{C}$ [11, Remark 10.13].

Example A.1.6. The deterministic morphisms in TychStoch are precisely the ones coming from continuous functions. They are continuous Markov kernels taking values in $\{0, 1\}$ -valued τ -smooth probability measures, and these correspond to the points by sobriety (see [29, Appendix A] and [28, Section 6] for additional context).

We will also use the following notions that are used in the theory of Markov categories, for which we refer to previous works.

- Positive Markov categories [11, Definition 11.22];
- Causal Markov categories [11, Definition 11.31];
- Representable Markov categories [18, Definition 3.10].

It is also helpful to have terminology for functors between Markov categories.

Definition A.1.7 ([11, Definition 10.14]). Let \mathcal{C} and \mathcal{D} be Markov categories. Then a **Markov functor** $F: \mathcal{C} \rightarrow \mathcal{D}$ is a strong symmetric monoidal functor making the diagram

$$\begin{array}{ccc} & FX & \\ \text{copy}_{FX} \swarrow & & \searrow F(\text{copy}_X) \\ FX \otimes_{\mathcal{D}} FX & \xrightarrow{\cong} & F(X \otimes_{\mathcal{C}} X) \end{array} \tag{A.4}$$

commute for every $X \in \mathcal{C}$, where the horizontal arrow is the coherence isomorphism of F . A **strict Markov functor** is a Markov functor that is strict as a monoidal functor.

Hence a strict Markov functor preserves the copy morphisms on the nose, while a general Markov functor need not. Note also that the preservation of the discard morphisms is automatic as the monoidal units are terminal.

A.2 Parametric Markov categories

An interesting aspect concerning synthetic probability via Markov categories is the fact that we can study parametrization in such a way that many results automatically generalize to the case where everything depends on an additional parameter. Indeed, let us recall the definition of parametric Markov category from [18, Section 2.2], which is such that morphisms in it are morphisms in the original category with a possible additional dependence on a parameter object.

Definition A.2.1. Let \mathcal{C} be a Markov category and $W \in \mathcal{C}$ any object. The **Markov category parametrized by W** , or simply **parametric Markov category**, is the Markov category \mathcal{C}_W defined as follows:

- Objects are those of \mathcal{C} ;
- Morphisms are parametrized by W , i.e.

$$\mathcal{C}_W(A, X) := \mathcal{C}(W \otimes A, X). \quad (\text{A.5})$$

- Composition and tensor are obtained by distributing the parameters via copy morphisms:

$$\begin{array}{c} Y \\ | \\ \boxed{g} \\ | \\ \boxed{f} \\ | \\ A \end{array} := \begin{array}{c} Y \\ | \\ \boxed{g} \\ | \\ \bullet \\ | \\ W \end{array} \begin{array}{c} X \\ | \\ \boxed{f} \\ | \\ A \end{array} \quad \begin{array}{c} X \\ | \\ \boxed{f} \\ | \\ A \end{array} \otimes \begin{array}{c} Y \\ | \\ \boxed{g} \\ | \\ B \end{array} := \begin{array}{c} X \quad Y \\ | \quad | \\ \boxed{f} \quad \boxed{g} \\ | \quad | \\ \bullet \quad | \\ | \quad W \quad A \quad B \end{array} \quad (\text{A.6})$$

- Structure morphisms discard the parameter:

$$\begin{array}{c} X \quad X \\ \cup \\ \bullet \\ | \\ X \end{array} := \begin{array}{c} X \quad X \\ | \quad | \\ \bullet \quad \bullet \\ | \quad | \\ W \quad X \end{array} \quad \begin{array}{c} \bullet \\ | \\ X \end{array} := \begin{array}{c} \bullet \quad \bullet \\ | \quad | \\ W \quad X \end{array} \quad (\text{A.7})$$

We use the same names for arrows and objects of both \mathcal{C} and \mathcal{C}_W . In order to distinguish the two settings, we use blue font for everything in \mathcal{C}_W , as opposed to the standard black used for \mathcal{C} . For example, the identity morphism on an object $X \in \mathcal{C}_W$ is given by $\text{id}_X = \text{del}_W \otimes \text{id}_X$.

Lemma A.2.2. The inclusion functor $\text{incl}: \mathcal{C} \rightarrow \mathcal{C}_W$, given by the identity on objects and by $\text{incl}(p) := \text{del}_W \otimes p$ on morphisms, is a strict Markov functor.

Proof. The proof is a direct check. The preservation of identities is trivial, while for the composition of $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ we have

$$\begin{array}{c} Z \\ | \\ \boxed{\text{incl}(g)} \\ | \\ \boxed{\text{incl}(f)} \\ | \\ X \end{array} = \begin{array}{c} Z \\ | \\ \bullet \\ | \\ W \end{array} \begin{array}{c} X \\ | \\ \boxed{f} \\ | \\ X \end{array} = \begin{array}{c} Z \\ | \\ \boxed{g} \\ | \\ \bullet \\ | \\ W \end{array} \begin{array}{c} X \\ | \\ \boxed{f} \\ | \\ X \end{array} = \begin{array}{c} Z \\ | \\ \boxed{\text{incl}(gf)} \\ | \\ X \end{array} \quad (\text{A.8})$$

For the monoidal product of $f: A \rightarrow X$ and $g: B \rightarrow Y$, we have

$$\begin{array}{c} X \\ | \\ \boxed{\text{incl}(f)} \\ | \\ A \end{array} \otimes \begin{array}{c} Y \\ | \\ \boxed{\text{incl}(g)} \\ | \\ B \end{array} = \begin{array}{c} X \quad Y \\ | \quad | \\ \boxed{f} \quad \boxed{g} \\ | \quad | \\ \bullet \quad | \\ | \quad W \quad A \quad B \end{array} = \begin{array}{c} X \quad Y \\ | \quad | \\ \boxed{f} \quad \boxed{g} \\ | \quad | \\ \bullet \quad | \\ | \quad W \quad A \quad B \end{array} = \begin{array}{c} X \quad Y \\ | \quad | \\ \boxed{\text{incl}(f \otimes g)} \\ | \quad | \\ A \quad B \end{array} \quad (\text{A.9})$$

The fact that copy and discard morphisms are preserved by incl follows immediately from the definitions. \square

We omit explicit mention of incl from most expressions besides this lemma, e.g. in the following proof.

Proposition A.2.3. *Let \mathcal{C} be a causal Markov category and $W \in \mathcal{C}$ be an object. Then \mathcal{C}_W is causal as well.*

Proof. We use the equivalent characterization of causality as parametrized equality strengthening [17, Proposition 2.17]. To this end, consider arbitrary morphisms $h_1, h_2: X \rightarrow Y$ and $p: A \rightarrow X$ in the parametric Markov category \mathcal{C}_W . Let us assume that h_1 is p -a.s. equal to h_2 , which takes the form

$$\begin{array}{c}
 Y \quad X \\
 | \quad | \\
 \boxed{h_1} \\
 | \quad | \\
 \bullet \\
 | \quad | \\
 \boxed{p} \\
 | \quad | \\
 \bullet \\
 | \quad | \\
 W \quad A
 \end{array}
 =
 \begin{array}{c}
 Y \quad X \\
 | \quad | \\
 \boxed{h_2} \\
 | \quad | \\
 \bullet \\
 | \quad | \\
 \boxed{p} \\
 | \quad | \\
 \bullet \\
 | \quad | \\
 W \quad A
 \end{array}
 \tag{A.10}$$

in \mathcal{C} . An arbitrary dilation²⁴ $\pi: X \rightarrow Y \otimes E$ of p is a morphism that satisfies

$$\begin{array}{c}
 X \\
 | \\
 \bullet \\
 | \\
 \boxed{\pi} \\
 | \quad | \\
 W \quad A
 \end{array}
 =
 \begin{array}{c}
 X \\
 | \\
 \boxed{p} \\
 | \quad | \\
 W \quad A
 \end{array}
 \tag{A.11}$$

in \mathcal{C} , so that the morphism

$$\begin{array}{c}
 W \quad X \quad E \\
 | \quad | \quad | \\
 \boxed{\pi} \\
 | \quad | \\
 \bullet \\
 | \quad | \\
 W \quad A
 \end{array}
 \text{ is a dilation of }
 \begin{array}{c}
 W \quad X \\
 | \quad | \\
 \boxed{p} \\
 | \quad | \\
 \bullet \\
 | \quad | \\
 W \quad A
 \end{array}
 \tag{A.12}$$

in \mathcal{C} .

Thus, applying parametrized equality strengthening of \mathcal{C} to Equation (A.10), with additionally the input W copied, yields

$$\begin{array}{c}
 Y \quad E \\
 | \quad | \\
 \boxed{h_1} \\
 | \quad | \\
 \boxed{\pi} \\
 | \quad | \\
 \bullet \\
 | \quad | \\
 W \quad A
 \end{array}
 =
 \begin{array}{c}
 Y \quad E \\
 | \quad | \\
 \boxed{h_2} \\
 | \quad | \\
 \boxed{\pi} \\
 | \quad | \\
 \bullet \\
 | \quad | \\
 W \quad A
 \end{array}
 \tag{A.13}$$

Since this applies for any dilation of p , we get that h_1 is p -dilationally equal to h_2 [17, Definition 4.1] and as a result that \mathcal{C}_W also satisfies parametrized equality strengthening. \square

²⁴For more details on dilations, see [17, Section 4].

Remark A.2.4. In particular, if \mathbf{C}_W is causal, then in \mathbf{C} the implication

$$(A.14)$$

holds for any $f, g, h_1,$ and h_2 . Therefore causality in a Markov category also implies causality with extra inputs in the sense of Implication (A.14).

A.3 Some auxiliary results on almost sure equality

Here, we prove several basic properties of almost sure equality as defined in Definition 2.1.1. One should keep in mind that this definition applies more broadly than the one from previous papers [11, Definition 13.1] due to the extra input wire.

Proposition A.3.1. *With p, f and g as in Definition 2.1.1, we have the following properties of almost sure equality:*

(i) *If $f =_p g$ holds, then also $h f =_p h g$ holds for any h , i.e. we have*

$$(A.15)$$

(ii) *If $h f =_p k f$ and $f =_p g$ hold, then also $h g =_p k g$ holds.*

(iii) *If $f =_p f'$ and $g =_q g'$ hold, then also $f \otimes g =_{p \otimes q} f' \otimes g'$ holds.*

(iv) *$f =_p g$ holds for $p: A \rightarrow X$ if and only if*

$$(A.16)$$

does.

(v) If p is deterministic, then $f =_p g$ is equivalent to

$$\begin{array}{c} Y \\ \downarrow \\ \boxed{f} \\ \downarrow \\ W \end{array} \quad \begin{array}{c} \downarrow \\ \boxed{p} \\ \downarrow \\ A \end{array} = \begin{array}{c} Y \\ \downarrow \\ \boxed{g} \\ \downarrow \\ W \end{array} \quad \begin{array}{c} \downarrow \\ \boxed{p} \\ \downarrow \\ A \end{array} \quad (\text{A.17})$$

Proof. (i): This follows directly from the definition.

(ii): This follows by two applications of property (i): $h g =_p h f =_p k f =_p k g$.

(iii): By Lemma 2.1.3, we can restrict to morphisms f and g with W equal to the unit I . Then the statement follows via:

$$\begin{array}{c} Y \quad Z \quad X \quad V \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ \boxed{f} \quad \boxed{g} \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ \boxed{p} \quad \boxed{q} \\ \downarrow \quad \downarrow \\ A \quad B \end{array} = \begin{array}{c} Y \quad Z \quad X \quad V \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ \boxed{f} \quad \quad \quad \boxed{g} \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ \boxed{p} \quad \boxed{q} \\ \downarrow \quad \downarrow \\ A \quad B \end{array} = \begin{array}{c} Y \quad Z \quad X \quad V \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ \boxed{f'} \quad \quad \quad \boxed{g'} \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ \boxed{p} \quad \boxed{q} \\ \downarrow \quad \downarrow \\ A \quad B \end{array} = \begin{array}{c} Y \quad Z \quad X \quad V \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ \boxed{f'} \quad \boxed{g'} \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ \boxed{p} \quad \boxed{q} \\ \downarrow \quad \downarrow \\ A \quad B \end{array} \quad (\text{A.18})$$

(iv): Once again by Lemma 2.1.3, we can restrict to morphisms without extra inputs. In such case we can use the manipulations

$$\begin{array}{c} Y \quad X \quad X \\ \downarrow \quad \downarrow \quad \downarrow \\ \boxed{f} \\ \downarrow \\ \bullet \\ \downarrow \\ \bullet \\ \downarrow \\ \boxed{p} \\ \downarrow \\ X \end{array} = \begin{array}{c} Y \quad X \quad X \\ \downarrow \quad \downarrow \quad \downarrow \\ \boxed{f} \\ \downarrow \\ \bullet \\ \downarrow \\ \bullet \\ \downarrow \\ \boxed{p} \\ \downarrow \\ X \end{array} = \begin{array}{c} Y \quad X \quad X \\ \downarrow \quad \downarrow \quad \downarrow \\ \boxed{g} \\ \downarrow \\ \bullet \\ \downarrow \\ \bullet \\ \downarrow \\ \boxed{p} \\ \downarrow \\ X \end{array} = \begin{array}{c} Y \quad X \quad X \\ \downarrow \quad \downarrow \quad \downarrow \\ \boxed{g} \\ \downarrow \\ \bullet \\ \downarrow \\ \bullet \\ \downarrow \\ \boxed{p} \\ \downarrow \\ X \end{array} \quad (\text{A.19})$$

to prove the claim.

(v): This is analogous to [11, Lemma 15.5]. \square

Lemma A.3.2. *Let \mathcal{C} be a Markov category. Then \mathcal{C} is causal if and only if for all $f: A \rightarrow X$ and $g: X \rightarrow Y$ and $h_1, h_2: Y \rightarrow Z$, we have*

$$\begin{array}{c} Z \quad Y \\ \downarrow \quad \downarrow \\ \boxed{h_1} \\ \downarrow \\ \bullet \\ \downarrow \\ \boxed{g} \\ \downarrow \\ \boxed{f} \\ \downarrow \\ A \end{array} = \begin{array}{c} Z \quad Y \\ \downarrow \quad \downarrow \\ \boxed{h_2} \\ \downarrow \\ \bullet \\ \downarrow \\ \boxed{g} \\ \downarrow \\ \boxed{f} \\ \downarrow \\ A \end{array} \quad \Rightarrow \quad \begin{array}{c} Z \quad X \\ \downarrow \quad \downarrow \\ \boxed{h_1} \\ \downarrow \\ \boxed{g} \\ \downarrow \\ \bullet \\ \downarrow \\ \boxed{f} \\ \downarrow \\ A \end{array} = \begin{array}{c} Z \quad X \\ \downarrow \quad \downarrow \\ \boxed{h_2} \\ \downarrow \\ \boxed{g} \\ \downarrow \\ \bullet \\ \downarrow \\ \boxed{f} \\ \downarrow \\ A \end{array} \quad (\text{A.20})$$

Proof. If \mathbf{C} is causal, we conclude by discarding the middle output in the consequent of causality, as on the right of (A.14) with $W = I$.

Conversely, assume that Implication (A.20) holds. The antecedent of the defining implication of causality, $h_1 =_{gf} h_2$, is equivalent to

$$\begin{array}{ccc}
 \begin{array}{c} Z \quad Y \\ | \quad | \\ \boxed{h_1} \\ | \quad | \\ \bullet \\ | \\ Y \end{array} & =_{gf} & \begin{array}{c} Z \quad Y \\ | \quad | \\ \boxed{h_2} \\ | \quad | \\ \bullet \\ | \\ Y \end{array} \\
 \end{array} \tag{A.21}$$

by Proposition A.3.1(iv), so that applying Implication (A.20) yields the required

$$\begin{array}{ccc}
 \begin{array}{c} Z \quad Y \\ | \quad | \\ \boxed{h_1} \\ | \quad | \\ \bullet \\ | \\ \boxed{g} \\ | \\ X \end{array} & =_f & \begin{array}{c} Z \quad Y \\ | \quad | \\ \boxed{h_2} \\ | \quad | \\ \bullet \\ | \\ \boxed{g} \\ | \\ X \end{array} \\
 \end{array} \tag{A.22}$$

which entails causality of \mathbf{C} . □

Lemma A.3.3. *In an arbitrary Markov category, let us consider a morphism $p: A \rightarrow T$ and a deterministic split monomorphism $\iota: T \rightarrow X$ with left inverse π . Then we have*

$$\begin{array}{ccccc}
 \begin{array}{c} Y \\ | \\ \boxed{f_1} \\ | \quad | \\ W \quad T \end{array} & =_p & \begin{array}{c} Y \\ | \\ \boxed{g_1} \\ | \quad | \\ W \quad T \end{array} & \iff & \begin{array}{c} Y \\ | \\ \boxed{f_1} \\ | \quad | \\ \pi \\ | \quad | \\ W \quad X \end{array} & =_{\iota p} & \begin{array}{c} Y \\ | \\ \boxed{g_1} \\ | \quad | \\ \pi \\ | \quad | \\ W \quad X \end{array} \\
 \end{array} \tag{A.23}$$

and

$$\begin{array}{ccccc}
 \begin{array}{c} Y \\ | \\ \boxed{f_2} \\ | \quad | \\ \iota \\ | \quad | \\ W \quad T \end{array} & =_p & \begin{array}{c} Y \\ | \\ \boxed{g_2} \\ | \quad | \\ \iota \\ | \quad | \\ W \quad T \end{array} & \iff & \begin{array}{c} Y \\ | \\ \boxed{f_2} \\ | \quad | \\ W \quad X \end{array} & =_{\iota p} & \begin{array}{c} Y \\ | \\ \boxed{g_2} \\ | \quad | \\ W \quad X \end{array} \\
 \end{array} \tag{A.24}$$

for any choice of morphisms $f_1, g_1: W \otimes T \rightarrow Y$ and $f_2, g_2: W \otimes X \rightarrow Y$.

Proof. We prove the equivalence (A.23) by rewriting the right-hand side. Specifically, by using the fact that ι is deterministic and satisfies $\pi \iota = \text{id}$, it is equivalent to

$$\begin{array}{ccc}
 \begin{array}{c} Y \quad X \\ | \quad | \\ \boxed{f_1} \quad \boxed{\iota} \\ | \quad | \\ \bullet \\ | \\ \boxed{p} \\ | \\ W \quad A \end{array} & = & \begin{array}{c} Y \quad X \\ | \quad | \\ \boxed{g_1} \quad \boxed{\iota} \\ | \quad | \\ \bullet \\ | \\ \boxed{p} \\ | \\ W \quad A \end{array} \\
 \end{array} \tag{A.25}$$

This equality follows from $f_1 =_p g_1$ by post-composing with ι on the right. For the converse implication, we post-compose with π on the right.

Showing the equivalence (A.24) works similarly. Post-composing the left-hand side with ι on the right gives the right-hand side since ι is deterministic. The converse implication is obtained by post-composing with π on the right. \square

Let $f: A \rightarrow X \otimes Y$ be a morphism in a Markov category \mathcal{C} . Then a *conditional of f given X* is a morphism $f_{|X}: X \otimes A \rightarrow Y$ which satisfies [11, Definiton 11.5]

(A.26)

The following observation generalizes [11, Proposition 13.7].

Lemma A.3.4 (Almost sure uniqueness of conditionals). *If $f_{|X}^{(1)}$ and $f_{|X}^{(2)}$ are two morphisms satisfying Equation (A.26), then we have the almost sure equality*

$$f_{|X}^{(1)} =_b f_{|X}^{(2)}, \tag{A.27}$$

where $b: A \rightarrow X \otimes A$ denotes the dashed box in Equation (A.26).

Thus we can think of the conditional $f_{|X}$ as being defined uniquely up to b -a.s. equality. In particular, if A is the monoidal unit I , then the lemma says that the conditional distribution $f_{|X}$ is f_X -almost surely unique.

Proof. Let us use first the definition of b , then properties of copying, and finally the definition of conditionals to write

(A.28)

which notably only involves f and not the choice of its conditional $f_{|X}^{(1)}$. Since we can do the same for $f_{|X}^{(2)}$, the desired Equation (A.27) follows. \square

A.4 Observationally representable Markov categories

In a representable Markov category \mathbf{C} [18], every object X comes equipped with a sampling morphism $\mathbf{samp}_X: PX \rightarrow X$, and for each morphism $p: A \rightarrow X$ there is a deterministic morphism $p^\sharp: A \rightarrow PX$ such that we have $\mathbf{samp} p^\sharp = p$. These establish a natural bijection

$$\mathbf{C}(A, X) \cong \mathbf{C}_{\text{det}}(A, PX) \quad (\text{A.29})$$

given by $p \mapsto p^\sharp$ from left to right and by post-composition with \mathbf{samp}_X (often denoted by \mathbf{samp} further on) from right to left.

In particular, the sampling morphism is a split epi with right inverse given by $(\text{id}_X)^\sharp$, which is nothing but $\delta_X: X \rightarrow PX$. Moreover, \mathbf{samp} is not monic, unless \mathbf{C} is cartesian, in which case P is isomorphic to the identity functor. Nevertheless, in standard probability theory it is the case that any two distinct probability measures can be distinguished by *iterated* sampling.²⁵ In a Markov category, collecting n independent samples amounts to composition with the morphism

$$\begin{array}{c} X^n \\ \downarrow \\ \boxed{\mathbf{samp}^{(n)}} \\ \downarrow \\ PX \end{array} := \begin{array}{c} \overbrace{}^{n \text{ outputs}} \\ X \quad \dots \quad X \\ \downarrow \quad \dots \quad \downarrow \\ \boxed{\mathbf{samp}} \quad \dots \quad \boxed{\mathbf{samp}} \\ \downarrow \quad \dots \quad \downarrow \\ \bullet \\ \downarrow \\ PX \end{array} \quad (\text{A.30})$$

where X^n denotes the n -fold tensor product of objects identical to X . In this section, we investigate what happens when these morphisms are jointly monic. This means that if two morphisms f, g satisfy $\mathbf{samp}^{(n)} f = \mathbf{samp}^{(n)} g$ for all $n \in \mathbb{N}$, then $f = g$ holds. Our claim is that this cancellation property extends automatically to almost sure equality classes. This is a consequence of the following observation.

Lemma A.4.1. *Let \mathbf{C} be a representable Markov category. Then the multiplication map*

$$\nabla_{X,Y}: PX \otimes PY \xrightarrow{\delta_{PX \otimes PY}} P(PX \otimes PY) \xrightarrow{P(\mathbf{samp} \otimes \mathbf{samp}_Y)} P(X \otimes Y)$$

is a split monomorphism, with left inverse given by $(P(\pi_2) \otimes P(\pi_1)) \text{copy}$, where

$$\pi_1 := \text{del}_X \otimes \text{id}_Y, \quad \pi_2 := \text{id}_X \otimes \text{del}_Y$$

are the two projections.

Proof. This is a general fact about monoidal monads on cartesian monoidal categories [13, Proposition 3.4], which we only need to instantiate on the monad $(P, P\mathbf{samp}, \delta)$ on \mathbf{C}_{det} . \square

Note that the left inverse $P(X \otimes Y) \rightarrow PX \otimes PY$ acts on Kleisli morphisms out of I as the map which takes a joint state to the product of its marginals:

$$\begin{array}{c} X \quad Y \\ \downarrow \quad \downarrow \\ \triangle \\ \downarrow \\ p \end{array} \mapsto \begin{array}{c} X \quad \bullet \quad Y \\ \downarrow \quad \downarrow \quad \downarrow \\ \triangle \quad \bullet \quad \triangle \\ \downarrow \quad \downarrow \quad \downarrow \\ p \quad \bullet \quad p \end{array} \quad (\text{A.31})$$

²⁵For the intuition behind this statement, note that by the strong law of large numbers, the relative frequency of an event in a sequence of samples approaches its actual probability almost surely.

Proposition A.4.2. *Let \mathcal{C} be a representable Markov category. Then the following are equivalent:*

- (i) *For every object, the morphisms $(\text{samp}^{(n)})_{n \in \mathbb{N}}$ are jointly monic.*
- (ii) *For every object, the morphisms $(\text{samp}^{(n)})_{n \in \mathbb{N}}$ are jointly monic modulo a.s. equality: for any morphisms $p: A \rightarrow X$ and $f, g: X \rightarrow PY$, we have*

$$\text{samp}^{(n)} f =_p \text{samp}^{(n)} g \quad \forall n \in \mathbb{N} \quad \implies \quad f =_p g. \quad (\text{A.32})$$

Proof. The implication from (ii) to (i) is trivial, since we can just consider the case $p = \text{id}_X$.

For the converse, we drop the subscript X from samp to simplify the notation. We thus assume that the $(\text{samp}^{(n)})_{n \in \mathbb{N}}$ are jointly monic, and that we have $\text{samp}^{(n)} f =_p \text{samp}^{(n)} g$ for some f, g and p as in the statement. Then we have

$$\text{Diagrammatic proof of Proposition A.4.2.} \quad (\text{A.33})$$

The first step holds because the multiplication maps $\nabla_{Y,X}$ are deterministic. The second step uses an instance of

$$h = \text{samp } h^\# \quad \text{where} \quad h^\# := (Ph) \delta_V, \quad (\text{A.34})$$

which holds for any morphism $h: V \rightarrow W$ in a representable Markov category. In particular, we use it for h given by $\text{samp}_Y \otimes \text{samp}_X: PY \otimes PX \rightarrow Y \otimes X$, since $\nabla_{Y,X}$ is equal to $(\text{samp}_Y \otimes \text{samp}_X)^\#$ by definition.

Since δ_X is deterministic and has samp_X as left inverse, the right-hand side is equal to

$$\text{Diagrammatic proof of Proposition A.4.2.} \quad (\text{A.35})$$

Note that using the assumed $\text{samp}^{(n)} f =_p \text{samp}^{(n)} g$, we can replace f by g in Diagram (A.35). Following the same steps backwards then shows that the left-hand side of Equation (A.33) coincides with the same expression with g instead of f . By assumption, the morphisms in $(\text{samp}^{(n)})_{n \in \mathbb{N}}$ are jointly monic, and by Lemma A.4.1, $\nabla_{Y,X}$ is also a monomorphism, so that we get

$$\begin{array}{c}
 \begin{array}{cc}
 PY & PX \\
 \downarrow & \downarrow \\
 \boxed{f} & \boxed{\delta} \\
 \downarrow & \downarrow \\
 \bullet & \\
 \downarrow & \\
 \boxed{p} & \\
 \downarrow & \\
 A &
 \end{array}
 & = &
 \begin{array}{c}
 \begin{array}{cc}
 PY & PX \\
 \downarrow & \downarrow \\
 \boxed{g} & \boxed{\delta} \\
 \downarrow & \downarrow \\
 \bullet & \\
 \downarrow & \\
 \boxed{p} & \\
 \downarrow & \\
 A &
 \end{array}
 \end{array}
 \end{array}
 \tag{A.36}$$

and applying samp_X on the right then gives $f =_p g$. This concludes the proof of Implication (A.32) from property (i). \square

Definition A.4.3. A representable Markov category is **observationally representable** if it satisfies the equivalent conditions of Proposition A.4.2.

The terminology follows that of [28, Definition 6.1], which is based on the idea that repeated sampling amounts to a sequence of observations. From [18], recall that we call a representable Markov category **a.s.-compatibly representable** if the bijections (A.29) respect almost sure equality.

Proposition A.4.4. Every observationally representable Markov category is a.s.-compatibly representable.

Proof. Consider $f, g: X \rightarrow Y$ with $f =_p g$. Then we have

$$\begin{array}{c}
 \begin{array}{cc}
 Y & Y \\
 \downarrow & \downarrow \\
 \boxed{\text{samp}} & \dots & \boxed{\text{samp}} \\
 \downarrow & & \downarrow \\
 \bullet & & \\
 \downarrow & & \\
 \boxed{f^\sharp} & & \\
 \downarrow & & \\
 X & &
 \end{array}
 & = &
 \begin{array}{c}
 \begin{array}{cc}
 Y & Y \\
 \downarrow & \downarrow \\
 \boxed{f} & \dots & \boxed{f} \\
 \downarrow & & \downarrow \\
 \bullet & & \\
 \downarrow & & \\
 X & &
 \end{array}
 \\
 & =_p & \\
 \begin{array}{c}
 \begin{array}{cc}
 Y & Y \\
 \downarrow & \downarrow \\
 \boxed{g} & \dots & \boxed{g} \\
 \downarrow & & \downarrow \\
 \bullet & & \\
 \downarrow & & \\
 X & &
 \end{array}
 \\
 & = & \\
 \begin{array}{c}
 \begin{array}{cc}
 Y & Y \\
 \downarrow & \downarrow \\
 \boxed{\text{samp}} & \dots & \boxed{\text{samp}} \\
 \downarrow & & \downarrow \\
 \bullet & & \\
 \downarrow & & \\
 \boxed{g^\sharp} & & \\
 \downarrow & & \\
 X & &
 \end{array}
 \end{array}
 \end{array}
 \tag{A.37}$$

where the middle equality follows directly from the assumed $f =_p g$ via [11, Lemma 13.5]. The other two equations hold because f^\sharp is deterministic and satisfies Equation (A.34) by representability. By observational representability, we obtain $f^\sharp =_p g^\sharp$, which concludes the proof of a.s.-compatible representability. \square

Example A.4.5. BorelStoch is observationally representable. Indeed, [28, Theorem 9.1] shows that (even on all of Stoch) the family $(\text{samp}^{(n)})_{n \in \mathbb{N}}$ is jointly monic.

Example A.4.6. FinSetMulti is observationally representable. Indeed, FinSetMulti is representable since it is the Kleisli category of the non-empty powerset monad on FinSet, which is its deterministic subcategory. The sampling morphism $\text{samp}: PX \rightarrow X$ is given by the multi-valued function mapping every non-empty subset of X to its elements. The observability is now the special case of [28, Theorem 9.4] for finite discrete spaces.

Example A.4.7. `SetMulti` is not observationally representable. The functor $X \mapsto PX$ forms the set of all nonempty subsets of X , finite and infinite. A `SetMulti` state $I \rightarrow PX$ is then a set of subsets of X . We now define two distinct states $p, q: I \rightarrow PX$ which will be the same after composing with $\mathbf{samp}^{(n)}$ for any $n \in \mathbb{N}$. To this end, let $p := PX$ be the set of *all* nonempty subsets of X itself, and let $q \subseteq PX$ be the set of *all finite* nonempty subsets of X . Assuming that X is infinite, these two states are trivially distinct. Composing any state with $\mathbf{samp}^{(n)}: PX \rightarrow X^n$ produces the set of sequences (x_1, \dots, x_n) for which there is a set contains all the x_i and belongs to the state. Therefore both $\mathbf{samp}^{(n)} p$ and $\mathbf{samp}^{(n)} q$ are the set of all n -element sequences from X . It follows that the maps $\mathbf{samp}^{(n)}$ cannot distinguish between p and q . Intuitively, the part where they differ — namely the infinite subsets of X — is not detectable by finitary sampling.

Proposition A.4.8. `CHausStoch` is observationally representable.

Proof. By point separability, it is enough to test the conditions of Proposition A.4.2 on states. We will apply [28, Lemma 6.6] (see also the proof of Theorem 9.1 therein for context). For X a compact Hausdorff space, PX is the space of Radon probability measures on X equipped with the topology of weak convergence, which is the weakest topology making the integration map

$$\begin{array}{ccc} \varepsilon_f : PX & \xrightarrow{\varepsilon_f} & [0, 1] \\ p & \longmapsto & \int f dp \end{array}$$

continuous for every continuous map $f: X \rightarrow [0, 1]$. In order to prove observability, by the mentioned [28, Lemma 6.6] it suffices to show that two probability measures μ and ν on PX (i.e. in PPX) are equal whenever they give the same integral on finite products of the integration maps,

$$\int_{PX} \varepsilon_{f_1} \cdots \varepsilon_{f_n} d\mu = \int_{PX} \varepsilon_{f_1} \cdots \varepsilon_{f_n} d\nu, \quad (\text{A.38})$$

for $f_1, \dots, f_n: X \rightarrow [0, 1]$ continuous. Denote now by $C(PX, \mathbb{R})$ the algebra of bounded continuous functions $PX \rightarrow \mathbb{R}$ and by \mathcal{S} the unital subalgebra generated by the functions ε_f for all continuous $f: X \rightarrow [0, 1]$. Given that the integrals (A.38) are equal for all continuous $f_i: X \rightarrow [0, 1]$, and that the finite products of integration maps ε_f span \mathcal{S} , we can conclude that

$$\int_{PX} h d\mu = \int_{PX} h d\nu \quad (\text{A.39})$$

for all $h \in \mathcal{S}$.

With $C(PX, \mathbb{R})$ carrying the usual topology of uniform convergence, we can apply the Stone–Weierstrass theorem (note that PX is compact Hausdorff too) to obtain that $\mathcal{S} \subseteq C(PX, \mathbb{R})$ is dense. Indeed the subalgebra \mathcal{S} separates the points of PX because the ε_f generate the topology of PX , which is Hausdorff.

Finally, let $g \in C(PX, \mathbb{R})$ and $\varepsilon > 0$ be arbitrary. Then by the density of \mathcal{S} there is $h \in \mathcal{S}$ with $\|g - h\|_\infty < \varepsilon$, which implies

$$\begin{aligned} \left| \int g d\mu - \int g d\nu \right| &= \left| \int g d\mu + \int h d\mu - \int h d\nu - \int g d\nu \right| \\ &\leq \int |g(x) - h(x)| \mu(dx) + \int |g(x) - h(x)| \nu(dx) \\ &\leq 2\varepsilon. \end{aligned} \quad (\text{A.40})$$

Therefore μ and ν assign the same integral to each continuous function $g: PX \rightarrow \mathbb{R}$, and so $\mu = \nu$ by the Riesz–Markov representation theorem. \square

A.5 The Cauchy–Schwarz Axiom

Here, we introduce a new information flow axiom for Markov categories, which is a version of the *multiplication lemma* of Parzygnat [33, Lemma 8.10]. As we will see, the proof that it holds in Stoch is based on the Cauchy–Schwarz Inequality as the essential ingredient, hence the name.

Definition A.5.1. A Markov category \mathcal{C} has the **Cauchy–Schwarz Property** if the implication

$$(A.41)$$

holds in \mathcal{C} .

We will show that Stoch (and therefore also BorelStoch and FinStoch) satisfy the Cauchy–Schwarz Property, while FinSetMulti and FinStoch_\pm do not. But first, let us motivate the name. For simplicity, take $A = I$ and $f = \text{id}$ in (A.41), so that g is a state and the implication becomes:

$$(A.42)$$

Instantiating this statement in Stoch , the left-hand side of the implication means that for all measurable sets $U, V \subseteq Y$, we have

$$\int_{x \in X} h(U|x) g(dx) \int_{x' \in X} h(V|x') g(dx') = \int_{x \in X} h(U|x) h(V|x) g(dx). \quad (A.43)$$

In particular, if U and V coincide, and denoting $h(U|x)$ by $H(x)$, we get

$$\left(\int_{x \in X} H(x) g(dx) \right)^2 = \int_{x \in X} H(x)^2 g(dx). \quad (A.44)$$

We know that, by the Cauchy–Schwarz Inequality (or by strict convexity of the function $y \mapsto y^2$), the equality above holds if and only if the function H is g -almost surely constant. This is exactly the consequent of Implication (A.42), which says

$$(A.45)$$

In other words, g -almost surely, the output of h does not depend on X . The Cauchy–Schwarz Property for Markov categories thus axiomatizes that part of the Cauchy–Schwarz Theorem which stipulates under what condition the inequality becomes an equality.

This illustration was given by setting $f = \text{id}$ and $A = I$. As the next proposition shows, the traditional Cauchy–Schwarz result (or at least the part where the inequality is an equality) also implies the axiom (A.5.1) for **Stoch** in full generality. When seen in contrast with the upcoming Example A.5.4, this result establishes an important feature of probabilistic uncertainty which is not shared by possibilistic uncertainty.

Proposition A.5.2. *The Markov category **Stoch** has the Cauchy–Schwarz Property.*

Proof. For $f: A \rightarrow B$, $g: B \rightarrow X$, and $h: X \rightarrow Y$, suppose that the antecedent of Implication (A.41) holds. Let us assume $A = I$, meaning that f is a state. By point-separability of **Stoch** (Definition 2.2.3), this assumption comes without loss of generality.

Fixing $U \in \Sigma_Y$, let us write $H(x) := h(U|x)$ and consider the expectation value

$$E(b) := \int_{x \in X} H(x) g(dx|b) \quad (\text{A.46})$$

for every $b \in B$. Then as in the usual proof of the Cauchy–Schwarz Inequality, the nonnegativity of the function $x \mapsto (H(x) - E(b))^2$ implies

$$\int_{x \in X} H(x)^2 g(dx|b) \geq E(b)^2 \quad (\text{A.47})$$

with equality if and only if $H(x) = E(b)$ for $g(_ | b)$ -almost all x . Evaluating the antecedent of Implication (A.41) on the measurable set $U \times U$ gives

$$\int_{b \in B} \int_{x_1 \in X} \int_{x_2 \in X} H(x_1) H(x_2) g(dx_1|b) g(dx_2|b) f(db) = \int_{b \in B} \int_{x \in X} H(x)^2 g(dx|b) f(db), \quad (\text{A.48})$$

or equivalently

$$\int_{b \in B} \left(\int_{x \in X} H(x)^2 g(dx|b) - E(b)^2 \right) f(db) = 0. \quad (\text{A.49})$$

Together with the established nonnegativity of the integrand, it follows that this integrand must vanish almost surely. By the above tightness condition, we hence conclude $H(x) = E(b)$ for $g(_ | b)$ -almost all x and f -almost all b .

Equipped with this intermediate result, let us now turn towards proving the claim. Since it is enough to prove the equality of two kernels with values in a product space on products of measurable sets [11, Lemma 4.2], it suffices to show that for all $W \in \Sigma_X$,

$$\int_{b \in B} \int_{x \in X} H(x) g(dx|b) g(W|b) f(db) = \int_{b \in B} \int_{x \in W} H(x) g(dx|b) f(db), \quad (\text{A.50})$$

or equivalently

$$\int_{b \in B} E(b) g(W|b) f(db) = \int_{b \in B} \int_{x \in W} H(x) g(dx|b) f(db). \quad (\text{A.51})$$

Since $H(x)$ equals $E(b)$ for $g(_ | b)$ -almost all x and f -almost all b , we can evaluate the right-hand side to

$$\int_{b \in B} E(b) \int_{x \in W} g(dx|b) f(db), \quad (\text{A.52})$$

and this proves Equation (A.51). \square


Corollary A.5.3. *FinStoch, BorelStoch, TopStoch, TychStoch and CHausStoch have the Cauchy–Schwarz property.*

Proof. As all the categories considered are Markov subcategories of Stoch, this is an immediate consequence of Proposition A.5.2. \square

Example A.5.4. In SetMulti and FinSetMulti, letting e be one of the idempotents from Example 4.1.5 violates the Cauchy–Schwarz Property with $f = g = h = e$. The same holds for the idempotent e from Example 4.1.6 in FinStoch $_{\pm}$.

B An isomorphism of measure spaces which does not preserve supports

Theorem B.0.1. *Let us denote the Lebesgue measure on the unit interval $[0, 1]$ by $\lambda_{[0,1]}$ and the normalized Lebesgue measure on the interval $[0, 1/2]$ by $2\lambda_{[0,1/2]}$.*



Then there exists an isomorphism of measure spaces of type $([0, 1], \lambda_{[0,1]}) \rightarrow ([0, 1], 2\lambda_{[0,1/2]})$.

In order to prove the theorem, we make use of the following lemma.

Lemma B.0.2. *There is an isomorphism of measure spaces $[0, 1] \rightarrow [0, 1)$, where both measurable spaces are equipped with the Lebesgue measure.*

Proof of Lemma B.0.2. Denote by $K \subseteq [0, 1]$ the countable subset

$$\left\{ \frac{1}{n} \mid n \in \mathbb{N}_{>0} \right\} = \left\{ 1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \right\}.$$

Consider now the bijection $b: K \rightarrow K \setminus \{1\}$ given by

$$b\left(\frac{1}{n}\right) := \frac{1}{n+1}. \tag{B.1}$$

The map

$$x \mapsto \begin{cases} b(x) & \text{if } x \in K, \\ x & \text{if } x \notin K \end{cases} \tag{B.2}$$

is now a measure-preserving isomorphism $[0, 1] \rightarrow [0, 1)$, since both measurability and measure preservation follow from the fact that it differs from the identity at only countably many points. \square

Proof of Theorem B.0.1. Let $C \subset [0, 1]$ be the Cantor set, which is a closed but uncountable subset of Lebesgue measure zero. Since every measurable subset of a standard Borel space is itself standard Borel, and since by Kuratowski's theorem all uncountable standard Borel spaces are isomorphic, there exists a measurable isomorphism $\Phi_0: C \rightarrow (1/2, 1]$. This becomes

an isomorphism of measure spaces if we equip both C and $(1/2, 1]$ with the zero measure, or equivalently, if we consider C as a null set in $([0, 1], \lambda_{[0,1]})$ and $(1/2, 1]$ as a null set in $([0, 1], 2\lambda_{[0,1/2]})$.

Consider now the complement $[0, 1] \setminus C$, which has Lebesgue measure 1. By construction of the Cantor set, its complement is a countable disjoint union of open intervals $\coprod_{n=0}^{\infty} (a_n, a_n + \ell_n)$, with $0 \leq a_n < a_n + \ell_n \leq 1$, and such that $\sum_{n=0}^{\infty} \ell_n = 1$. Now by Lemma B.0.2, there is a measure-preserving isomorphism $\psi_0: (a_0, a_0 + \ell_0) \rightarrow [0, \ell_0]$, as well as a measure-preserving isomorphism

$$\psi_n : (a_n, a_n + \ell_n) \longrightarrow (\ell_0 + \cdots + \ell_{n-1}, \ell_0 + \cdots + \ell_n]$$

for each $n > 0$. Since we have

$$[0, 1] = [0, \ell_0] \sqcup \coprod_{n=1}^{\infty} (\ell_0 + \cdots + \ell_{n-1}, \ell_0 + \cdots + \ell_n], \quad (\text{B.3})$$

we obtain a measure-preserving isomorphism $\Phi_1: [0, 1] \setminus C \rightarrow [0, 1]$ given by $x \mapsto \psi_n(x)$ for all $x \in (a_n, a_n + \ell_n)$ and $n \in \mathbb{N}$.

The map

$$\Phi_0 \sqcup \Phi_1 : [0, 1] = (X \setminus C) \sqcup C \longrightarrow [0, 1/2] \sqcup (1/2, 1] = [0, 1] \quad (\text{B.4})$$

given by

$$x \mapsto \begin{cases} \Phi_0(x) & \text{if } x \in C, \\ \frac{1}{2} \Phi_1(x) & \text{if } x \notin C \end{cases} \quad (\text{B.5})$$

is now an isomorphism between measure spaces $([0, 1], \lambda_{[0,1]})$ and $([0, 1], 2\lambda_{[0,1/2]})$. \square

References

- [1] T. Banach. Topology of spaces of probability measures. I: The functors P_τ and \hat{P} . *Mat. Stud.*, 5(1-2):65–87, 1995. ISSN 1027-4634. arXiv:1112.6161. ↑66
- [2] M. Billik. Idempotent Reynolds operators. *J. Math. Anal. Appl.*, 18:135–144, 1967. doi:10.1016/0022-247X(67)90188-6. ↑49
- [3] D. Blackwell. Idempotent Markoff chains. *Ann. of Math. (2)*, 43:560–567, 1942. doi:10.2307/1968811. ↑5, 61
- [4] F. Borceux. *Handbook of Categorical Algebra I*. Cambridge University Press, 1994. doi:10.1017/CBO9780511525858. ↑22, 27, 52, 63
- [5] F. Borceux and D. Dejean. Cauchy completion in category theory. *Cahiers de Topologie et Géométrie Différentielle Catégoriques*, 27(2):133–146, 1986. ↑5
- [6] D. Braithwaite and J. Hedges. Dependent Bayesian lenses: Categories of bidirectional Markov kernels with canonical Bayesian inversion, 2022. arXiv:2209.14728. ↑4
- [7] D. Braithwaite, J. Hedges, and T. S. C. Smithe. The compositional structure of Bayesian inference, 2023. arXiv:2305.06112. ↑4
- [8] K. Cho and B. Jacobs. Disintegration and Bayesian inversion via string diagrams. *Math. Structures Comput. Sci.*, 29:938–971, 2019. doi:10.1017/s0960129518000488. arXiv:1709.00322. ↑2
- [9] R. Deville, G. Godefroy, and V. Zizler. *Smoothness and renormings in Banach spaces*, volume 64 of *Pitman Monogr. Surv. Pure Appl. Math.* Longman Publishing Group, 1993. ↑44
- [10] E. Di Lavore, G. de Felice, and M. Román. Monoidal streams for dataflow programming. In *Proceedings of the 37th Annual ACM/IEEE Symposium on Logic in Computer Science*, pages 1–14, 2022. doi:10.1145/3531130.3533365. ↑4
- [11] T. Fritz. A synthetic approach to Markov kernels, conditional independence and theorems on sufficient statistics. *Adv. Math.*, 370:107239, 2020. doi:10.1016/j.aim.2020.107239. arXiv:1908.07021. ↑2, 3, 4, 6, 7, 8, 12, 17, 18, 19, 35, 39, 43, 60, 66, 67, 70, 71, 73, 76, 79
- [12] T. Fritz and A. Klingler. The d -separation criterion in categorical probability. *J. Mach. Learn. Res.*, 24(46):1–49, 2023. arXiv:2207.05740. ↑2, 60
- [13] T. Fritz and P. Perrone. Bimonoidal structure of probability monads. In *Proceedings of the Thirty-Fourth Conference on the Mathematical Foundations of Programming Semantics (MFPS XXXIV)*, volume 341 of *Electron. Notes Theor. Comput. Sci.*, pages 121–149. Elsevier Sci. B. V., Amsterdam, 2018. doi:10.1016/j.entcs.2018.11.007. arXiv:1804.03527. ↑74
- [14] T. Fritz and E. F. Rischel. Infinite products and zero-one laws in categorical probability. *Compositionality*, 2:3, 2020. doi:10.32408/compositionality-2-3. arXiv:1912.02769. ↑2, 13, 43
- [15] T. Fritz, T. Gonda, and P. Perrone. de Finetti’s theorem in categorical probability. *J. Stoch. Anal.*, 2(4), 2021. doi:10.31390.josa.2.4.06. arXiv:2105.02639. ↑2

- [16] T. Fritz, P. Perrone, and S. Rezagholi. Probability, valuations, hyperspace: Three monads on Top and the support as a morphism. *Math. Structures Comput. Sci.*, 31(8):850–897, 2021. doi:10.1017/S0960129521000414. arXiv:1910.03752. ↑3, 9, 10, 24, 66
- [17] T. Fritz, T. Gonda, N. G. Houghton-Larsen, A. Lorenzin, P. Perrone, and D. Stein. Dilations and information flow axioms in categorical probability, 2022. arXiv:2211.02507. ↑28, 45, 62, 69
- [18] T. Fritz, T. Gonda, P. Perrone, and E. F. Rischel. Representable Markov categories and comparison of statistical experiments in categorical probability. *Theoretical Computer Science*, 961:113896, 2023. doi:10.1016/j.tcs.2023.113896. arXiv:2010.07416. ↑2, 4, 5, 7, 10, 16, 22, 35, 66, 67, 74, 76
- [19] R. W. J. Furber and B. P. F. Jacobs. From Kleisli categories to commutative C*-algebras: Probabilistic Gelfand duality. *Logical Methods in Computer Science*, 11, 2015. doi:10.2168/LMCS-11(2:5)2015. arXiv:1303.1115. ↑44, 63, 66
- [20] F. Gadducci. *On The Algebraic Approach To Concurrent Term Rewriting*. PhD thesis, University of Pisa, 1996. ↑2
- [21] T. Gonda, T. Reinhart, S. Stengele, and G. D. I. Coves. A framework for universality in physics, computer science, and beyond, 2023. arXiv:2307.06851. ↑31
- [22] M. Hamana. Injective envelopes of c^* -algebras. *J. Math. Soc. Japan*, 31(1):181–197, 1979. ↑63
- [23] E. Harzheim. *Ordered sets*, volume 7 of *Advances in Mathematics (Springer)*. Springer, New York, 2005. ↑48
- [24] J. Jost, H. V. Lê, and T. D. Tran. Probabilistic morphisms and Bayesian nonparametrics. *Eur. Phys. J. Plus*, 136:441, 2021. doi:10.1140/epjp/s13360-021-01427-7. arXiv:1905.11448. ↑35
- [25] I. Karatzas and S. E. Shreve. *Brownian motion and stochastic calculus.*, volume 113 of *Grad. Texts Math.* Springer, 2nd ed. edition, 1991. ↑26
- [26] A. S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. ↑26, 27, 41
- [27] K. J. Laidler. *Chemical Kinetics*. Pearson, 3rd edition, 1987. ↑51
- [28] S. Moss and P. Perrone. Probability monads with submonads of deterministic states. In *Proceedings of the 37th Annual ACM/IEEE Symposium on Logic in Computer Science*, pages 1–13, 2022. doi:10.1145/3531130.3533355. arXiv:2204.07003. ↑62, 67, 76, 77
- [29] S. Moss and P. Perrone. A category-theoretic proof of the ergodic decomposition theorem. *Ergodic Theory Dynam. Systems*, pages 1–27, 2023. doi:10.1017/etds.2023.6. arXiv:2207.07353. ↑2, 67
- [30] S.-T. C. Moy. Characterizations of conditional expectation as a transformation on function spaces. *Pacific J. Math.*, 4:47–63, 1954. ↑49
- [31] D. Mumford, J. Fogarty, and F. Kirwan. *Geometric invariant theory*, volume 34 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)]*. Springer-Verlag, Berlin, third edition, 1994. ↑49

- [32] A. J. Parzygnat. Discrete probabilistic and algebraic dynamics: a stochastic commutative Gelfand-Naimark theorem, 2017. [arXiv:1708.00091](#). ↑66
- [33] A. J. Parzygnat. Inverses, disintegrations, and Bayesian inversion in quantum Markov categories, 2020. [arXiv:2001.08375](#). ↑7, 78
- [34] E. Patterson. *The algebra and machine representation of statistical models*. PhD thesis, Stanford University, 2020. [arXiv:2006.08945](#). ↑35
- [35] A. Pełczyński. Linear extensions, linear averagings, and their applications to linear topological classification of spaces of continuous functions. *Dissertationes Math. (Rozprawy Mat.)*, 58:92, 1968. [eudml.org/doc/268522](#). ↑44
- [36] P. Perrone. Markov categories and entropy, 2022. [arXiv:2212.11719](#). ↑2, 26
- [37] D. Stein and S. Staton. Compositional semantics for probabilistic programs with exact conditioning. In *Logic in Computer Science*. IEEE, 2021. doi:10.1109/LICS52264.2021.9470552. [arXiv:2101.11351](#). ↑34
- [38] N. N. Čencov. The categories of mathematical statistics. *Dokl. Akad. Nauk SSSR*, 164: 511–514, 1965. [mathnet.ru/rus/dan31602](#). ↑35
- [39] N. N. Čencov. *Statistical decision rules and optimal inference*, volume 53 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, R.I., 1982. Translation from the Russian edited by Lev J. Leifman. ↑35
- [40] Y. Zhang and M. Sugiyama. A category-theoretical meta-analysis of definitions of disentanglement, 2023. [arXiv:2305.06886](#). ↑2
- [41] T. Świrszcz. Monadic functors and convexity. *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.*, 22, 1974. ↑66