

# SUPERSYMMETRIC EXTENSION OF UNIVERSAL ENVELOPING VERTEX ALGEBRAS

UHI RINN SUH<sup>1</sup> AND SANGWON YOON<sup>1</sup>

**ABSTRACT.** In this paper, we study the construction of the supersymmetric extensions of vertex algebras. In particular, for  $N = n \in \mathbb{Z}_+$ , we show the universal enveloping  $N = n$  supersymmetric (SUSY) vertex algebra of an  $N = n$  SUSY Lie conformal algebra can be extended to an  $N = n' > n$  SUSY vertex algebra. We also show the  $N = 2$  SUSY affine vertex algebra of level 0 associated with a Lie superalgebra, which is an  $N = 2$  SUSY extension of the affine vertex algebra of level 0, can be embedded as an  $N = 2$  SUSY vertex subalgebra into an  $N = 2$  superconformal vertex algebra.

## 1. INTRODUCTION

Vertex algebras were introduced by R. Borcherds in the 1980's (see [3]) to describe the chiral part of two-dimensional conformal field theory (see [4]). A vertex algebra consists of a vector superspace  $V$ ,  $\text{End}(V)$ -valued quantum fields  $a(z)$  for  $a \in V$  and a translation operator  $\partial$  with certain axioms. One of the axiom called locality tells that, for a given pair of fields  $(a, b)$  in  $V$ , there is a positive integer  $t \in \mathbb{N}$  such that  $(z - w)^t [a(z), b(w)] = 0$ . Considering the formal delta distribution  $\delta(z, w) := \sum_{j \in \mathbb{Z}} z^j w^{-j-1}$ , the elements  $a_{(m)}b \in V$  for  $m = 0, 1, \dots, t$  are determined by the supercommutator formula:

$$(1.1) \quad [a(z), b(w)] = \sum_{m=0}^t \frac{1}{m!} a_{(m)}b(w) \partial_w^m \delta(z, w).$$

The existence of such formula (1.1) is guaranteed by so-called the decomposition theorem. Additionally, (1.1) and the state-field correspondence yield another well-known definition of a vertex algebra using the  $\lambda$ -bracket formalism (see Section 2.1). Speaking briefly, a vertex algebra  $V$  is a differential quasi-commutative quasi-associative superalgebra endowed with a  $\lambda$ -bracket which is compatible with the differential algebra structure. Here, the derivation on  $V$  for the differential algebra structure arises from the translation operator  $\partial$ .

To describe two-dimensional conformal field theory with supersymmetries in mathematical language, it is good to use the notion of supersymmetric vertex algebra, which is a generalization of the superconformal vertex algebras (see Section 5.9 of [19]). A supersymmetric vertex algebra consists of a vector superspace  $V$  and superfields  $a(z, \Theta) := a(z, \theta_1, \theta_2, \dots, \theta_n)$  for  $a \in V$  and odd variables  $\theta_i$ . Roughly speaking, the superfield

$$a(z, \Theta) = \sum_I \pm \theta^I D^I a(z),$$

where the summation is over the ordered subset  $I = \{i_1, i_2, \dots, i_s\} \in \{1, 2, \dots, n\}$  satisfying  $i_1 < i_2 < \dots < i_s$ ,  $D^I a(z) := D^{i_1} D^{i_2} \dots, D^{i_s} a(z)$  is a field of parity  $p(a) + \bar{s}$  and  $\theta_I = \theta_{i_1} \theta_{i_2} \dots \theta_{i_s}$ . Here  $D^i$ 's can be understood as odd endomorphisms of fields such that  $D^i D^j + D^j D^i = \delta_{i,j} 2\partial$  for  $i, j \in I$  and, indeed, the odd endomorphisms determine the structures of  $V$  coming from the supersymmetry. In particular, in  $N = 1$  (resp.  $N = 2$ ) case, a superfield  $a(z, \Theta)$  consists of a couple (resp. quadruple) of fields  $(a(z), Da(z))$

---

<sup>1</sup>This work was supported by NRF Grant, #2022R1C1C1008698 and Creative-Pioneering Researchers Program through Seoul National University.

(resp.  $(a(z), D^1a(z), D^2a(z), D^1D^2(z))$ ). Analogous to the  $\lambda$ -bracket formalism in vertex algebras theory, an  $N = n$  supersymmetric vertex algebra can be introduced via so-called  $\Lambda = (\lambda, \chi^1, \chi^2, \dots, \chi^n)$ -bracket formalism (see Section 2.2).

As one can expect, a SUSY vertex algebra is a vertex algebra by itself but the converse cannot be true. However, it is well-known that if a vertex algebra has a superconformal vector then the vector induces a supersymmetry in the vertex algebra (see, for example [13, 19]). Here a superconformal vector  $G$  in a vertex algebra  $V$  is a primary of conformal weight  $\frac{3}{2}$  element satisfying

$$[G \lambda G] = 2L + \frac{c}{3}\lambda^2$$

for a conformal vector  $L$  of  $V$ . Indeed, in this case, we can define an odd operator  $D$  by letting  $D(a)$  to be the coefficient of  $\lambda^0$  in  $[G \lambda a]$  and the vertex algebra  $V$  together with  $D$  becomes an  $N = 1$  SUSY vertex algebra. We remark that constructions of  $N = 1$  and  $N = 2$  superconformal structures in vertex algebras have been studied in various contexts (see [16, 18]).

If a vertex algebra does not have a superconformal vector, we cannot guarantee the existence of supersymmetries. For example, the Virasoro vertex algebra and affine vertex algebras do not have neither superconformal vectors nor known supersymmetries. Nevertheless, we can impose supersymmetries on these vertex algebras by embedding them into bigger supersymmetric vertex algebras. For the Virasoro vertex algebra, consider the  $N = 1$  SUSY vertex algebra generated by the odd element  $G$  and endowed with the  $\Lambda$ -bracket:

$$[G \Lambda G] = (2\partial + 3\lambda + \chi D)G + \frac{c}{3}\lambda^2\chi.$$

This SUSY vertex algebra is called the  $N = 1$  super-Virasoro vertex algebra and the vertex subalgebra generated by the even element  $D(G)$  is the Virasoro vertex algebra. Hence the  $N = 1$  super-Virasoro vertex algebra can be considered as a supersymmetric extension of the Virasoro vertex algebra.

Similarly, the affine vertex algebra  $V^k(\mathfrak{g})$  of level  $k$  associated with a Lie superalgebra  $\mathfrak{g}$  can be embedded into an  $N = 1$  SUSY vertex algebra  $V_{N=1}^k(\mathfrak{g})$  called the  $N = 1$  SUSY affine vertex algebra of level  $k$ . As a SUSY vertex algebra,  $V_{N=1}^k(\mathfrak{g})$  is generated by the parity reversed vector superspace  $\bar{\mathfrak{g}}$  of  $\mathfrak{g}$  and the  $\Lambda$ -bracket is given by:

$$[\bar{a} \Lambda \bar{b}] = (-1)^{p(a)} \left( \overline{[a, b]} + k\chi(a|b) \right)$$

for  $a, b \in \mathfrak{g}$ . Now, the vertex subalgebra generated by  $D\bar{\mathfrak{g}}$  is isomorphic to the affine vertex algebra. In addition, in various articles [14, 22], supersymmetric extensions of well-known vertex algebras such as lattice vertex algebras and  $W$ -algebras have been discussed.

In this paper, we investigate where supersymmetries of vertex algebras come from. In short, if a vertex algebra  $V$  has an odd derivation  $D$  which induces the sesquilinearity of the  $\Lambda$ -bracket and satisfies  $[D, D] = 2\partial$  then the operator  $D$  gives a supersymmetry in  $V$ . In terms of  $\lambda$ -bracket, the sesquilinearity of  $\Lambda$ -bracket can be rewritten as (SEF1) and (SEF2) in Section 3 and they are necessary and sufficient conditions for a supersymmetry of a vertex algebra (see Proposition 3.4). Using these observation, we show the main theorem in Section 3:


**Theorem 1.1** (Theorem 3.6). *Let  $R = \mathbb{C}[\partial] \otimes \mathcal{U}$  for a vector superspace  $\mathcal{U}$  be a Lie conformal algebra and  $V(R)$  be the universal enveloping vertex algebra of the Lie conformal algebra  $R$ . The differential algebra  $V(\bar{R} \oplus R)$ , where  $\bar{R}$  is the parity reversed  $\mathbb{C}[\partial]$ -module of  $R$ , endowed with the  $\Lambda$ -bracket*

$$(1.2) \quad [\bar{a} \Lambda \bar{b}] := (-1)^{p(a)} \overline{[a \lambda b]}$$

and the odd derivation  $D$  such that

$$D(\bar{a}) = a \quad \text{and} \quad D(a) = \partial \bar{a}$$

for  $a, b \in \mathcal{U}$  is an  $N = 1$  SUSY vertex algebra having  $V(R)$  as a vertex subalgebra. (See the picture below.)



Construction of  $N = 1$  supersymmetry in Theorem 3.6

The conclusion of Theorem 1.1 is, for the universal enveloping vertex algebra of a Lie conformal algebra, one can always find a supersymmetric extension of the vertex algebra. Note that the  $\Lambda$ -bracket (1.2) is inspired from the bracket of  $V_{N=1}^0(\mathfrak{g})$ , the SUSY extension of  $V^0(\mathfrak{g})$ . However, it is not the only supersymmetric extension of such vertex algebras. For example, the  $bc$ - $\beta\gamma$  system is a well-known supersymmetric extension of the  $\beta\gamma$  system but the SUSY vertex algebra obtained by Theorem 1.1 is not the  $bc$ - $\beta\gamma$  system (see Example 3.7). For the Virasoro vertex algebra, the supersymmetric extension via Theorem 1.1 is not the  $N = 1$  super-Virasoro vertex algebra (see Example 3.8).

Moreover, by using the SUSY extension formulas (SEF1) and (SEF2) in Section 3, it is possible to impose desired constraints and thereby construct SUSY extensions of a given vertex algebra. As an example (see Remark 3.9), we can reproduce the operator product expansion of the superconformal current algebra, which was defined in [20]. In other words, the  $N = 1$  SUSY affine vertex algebra  $V_{N=1}^0(\mathfrak{g})$  of level 0 can be embedded into an  $N = 1$  superconformal vertex algebra  $\text{SVir}^c \otimes V_{N=1}^0(\mathfrak{g})$ , where  $\text{SVir}^c$  is the  $N = 1$  super-Virasoro vertex algebra of central charge  $c \in \mathbb{C}$ . Note that  $V_{N=1}^k(\mathfrak{g})$  for a basic Lie superalgebra  $\mathfrak{g}$  and  $k \neq 0$  is known to be an  $N = 1$  superconformal vertex algebra by itself, since it has the Kac-Todorov superconformal vector (see [19, 20]). Meanwhile, when  $\mathfrak{g}$  does not have a non-degenerate even supersymmetric bilinear form or the level  $k = 0$ , the Kac-Todorov construction collapses and no superconformal vector has been discovered.

In Section 4, we begin with the characterization of the  $N = n$  SUSY vertex algebras using  $n$  odd derivations, as mentioned in Section 5.9 of [19]. With this characterization, we observe an action of the orthogonal group on the collection of  $N = n$  SUSY structures of the given vertex algebra, which can be viewed as a generalization of the relationships between SUSY structures derived from the superconformal structures (see Proposition 4.1). In particular, we deal with some examples of the SUSY structures associated with superconformal structures and, in Remark 4.8, we discuss comments related to the shift of superconformal vectors and its application to the superconformal structures of SUSY  $W$ -algebras.

In Theorem 4.9, we generalize the result in Section 3 to obtain an  $N = 2$  SUSY extension of a universal enveloping  $N = 1$  SUSY vertex algebra. Moreover, in Remark 4.14, we show the same argument can be applied to find an  $N = n + 1$  SUSY extension of a universal enveloping  $N = n$  SUSY vertex algebra.

Inductively, a universal enveloping vertex algebra can be extended to an  $N = n$  SUSY vertex algebra for  $n \geq 1$ . In particular, applying the above process to centerless current Lie conformal algebra yields the super-loop algebra in [6].

Furthermore, in Proposition 4.13, we show the  $N = 2$  SUSY affine vertex algebra  $V_{N=2}^0(\mathfrak{g})$  of level 0, which is an  $N = 2$  SUSY extension of the affine vertex algebra of level 0, can be embedded into an  $N = 2$  superconformal vertex algebra. More precisely, there is an  $N = 2$  superconformal vertex algebra  $\tilde{V}_{N=2}^0(\mathfrak{g})$  such that

$$\tilde{V}_{N=2}^0(\mathfrak{g}) \simeq \text{SVir}_{N=2}^c \otimes V_{N=2}^0(\mathfrak{g}),$$

where  $\text{SVir}_{N=2}^c$  is the  $N = 2$  super-Virasoro vertex algebra, and the two odd derivations giving  $N = 2$  SUSY structure of  $V_{N=2}^0(\mathfrak{g})$  can be realized by the  $N = 2$  superconformal structure of  $\tilde{V}_{N=2}^0(\mathfrak{g})$ .

## 2. PRELIMINARIES

**2.1. Vertex algebras.** In this section, we review basic notions and properties used throughout this article. For more details, we refer to [8, 13, 19].

In this paper, the base field is  $\mathbb{C}$ . A vector superspace  $V = V_{\bar{0}} \oplus V_{\bar{1}}$  is a  $\mathbb{Z}/2\mathbb{Z}$ -graded vector space. An element  $a \in V$  is called even (resp. odd) if  $a \in V_{\bar{0}}$  (resp.  $a \in V_{\bar{1}}$ ). When  $a \in V$  is even (resp. odd), the parity of  $a$  is given by  $p(a) = 0$  (resp.  $p(a) = 1$ ). A vector superspace  $V$  is called a *vertex algebra* if it is endowed with  $m$ -th products  $a_{(m)}b$  for  $a, b \in V$  and  $m \in \mathbb{Z}$  with the properties described below. The  $(-1)$ -th product  $a_{(-1)}b$  is usually denoted by  $:ab:$  and called the *normally ordered product* of  $a$  and  $b$ . In addition,  $V$  has an even derivation  $\partial$  such that

$$a_{(-m-1)}b = \frac{1}{m!} : (\partial^m a)b :$$

for  $m \in \mathbb{N}$ . Therefore the structure of  $V$  is determined by (i) the  $m$ -th product for  $m \in \mathbb{Z}_+$ , (ii) the normally ordered product  $: :$ , and (iii) the derivation  $\partial$ . In order to deal with all the information from (i) together, we define a bracket  $V \otimes V \rightarrow \mathbb{C}[\lambda] \otimes V$  by

$$(2.1) \quad (a, b) \mapsto [a \ \lambda \ b] = \sum_{m \in \mathbb{Z}_+} \frac{\lambda^m}{m!} a_{(m)}b,$$

where  $\lambda$  is an even indeterminate. The linear bracket (2.1) satisfies the following property:

$$(\text{Sesquilinearity}) \quad [\partial a \ \lambda \ b] = -\lambda [a \ \lambda \ b], \quad [a \ \lambda \ \partial b] = (\partial + \lambda)[a \ \lambda \ b].$$

The RHS of the second equality in sesquilinearity can be computed by the relation  $\partial\lambda = \lambda\partial$ . More generally, for a  $\mathbb{C}[\partial]$ -module  $R$ , a bracket  $R \times R \rightarrow R[\lambda]$  with the sesquilinearity is called a  $\lambda$ -*bracket* on  $R$ . In other words, the bracket (2.1) is a  $\lambda$ -bracket on the vertex algebra  $V$ . Moreover, a vertex algebra is a Lie conformal algebra, which is defined as below.

**Definition 2.1** ([8]). A *Lie conformal algebra* is a  $\mathbb{Z}/2\mathbb{Z}$ -graded  $\mathbb{C}[\partial]$ -module  $R$  with a  $\lambda$ -bracket which is a parity preserving  $\mathbb{C}$ -linear map

$$[\ \lambda \ ] : R \otimes R \rightarrow \mathbb{C}[\lambda] \otimes R, \quad a \otimes b \mapsto [a \ \lambda \ b],$$

satisfying the following conditions:

$$\begin{aligned} (\text{Skew-symmetry}) \quad & [b \ \lambda \ a] = (-1)^{p(a)p(b)+1} [a \ -\partial-\lambda \ b], \\ (\text{Jacobi identity}) \quad & [a \ \lambda \ [b \ \gamma \ c]] = [[a \ \lambda \ b] \ \lambda+\gamma \ c] + (-1)^{p(a)p(b)} [b \ \gamma \ [a \ \lambda \ c]], \end{aligned}$$

where  $a, b, c$  are homogeneous elements in  $R$ . More precisely, the RHS of skew-symmetry can be rewritten as

$$[a \text{ }_{-\partial-\lambda} b] = \sum_{m \in \mathbb{Z}_+} \frac{(-\partial - \lambda)^m}{m!} a_{(m)} b,$$

where we denote the  $\lambda$ -bracket as in (2.1), and the Jacobi identity is defined in  $\mathbb{C}[\lambda, \gamma] \otimes R$ . In addition, we assume  $\lambda\gamma = \gamma\lambda$ .

Furthermore, there are more properties required to the normally ordered product on a vertex algebra  $V$  and for the compatibility between the normally ordered product and the  $\lambda$ -bracket or derivation. In the following definition, one can find all axioms for vertex algebras.

**Definition 2.2** ([8]). A *vertex algebra* is a tuple  $(V, \partial, [\ \lambda ], |0\rangle, \ : \ :)$  such that:

- $(V, \partial, [\ \lambda ])$  is a Lie conformal algebra,
- $(V, \partial, |0\rangle, \ : \ :)$  is a unital differential superalgebra with a derivation  $\partial$ , satisfying the following properties:

$$\text{(Quasi-commutativity)} \quad : ab : - (-1)^{p(a)p(b)} : ba : = \int_{-\partial}^0 [a \ \lambda b] d\lambda,$$

(Quasi-associativity)

$$: ab : c : - : a : bc :: = : \left( \int_0^\partial d\lambda a \right) [b \ \lambda c] : + (-1)^{p(a)p(b)} : \left( \int_0^\partial d\lambda b \right) [a \ \lambda c] :,$$

- the  $\lambda$ -bracket and the product  $\ : \ :$  are related by

(Non-commutative Wick formula)

$$[a \ \lambda : bc :] = : [a \ \lambda b] c : + (-1)^{p(a)p(b)} : b [a \ \lambda c] : + \int_0^\lambda [[a \ \lambda b] \ \gamma c] d\gamma.$$

In the quasi-associativity, we note that  $: \left( \int_0^\partial d\lambda a \right) [b \ \lambda c] : = \sum_{n \in \mathbb{Z}_+} \sum_{m \in \mathbb{Z}_+} (a_{(-n-2)} (b_{(m)} c))$ .

**Remark 2.3.** From now on, we denote the normally ordered product  $\ : ab :$  simply by  $ab$  for  $a, b \in V$ . For  $a_1, a_2, \dots, a_l \in V$ , the element  $a_1 a_2 \cdots a_{l-1} a_l$  is the element obtained by performing normally ordered product from right to left. For example:

$$a_1 a_2 a_3 = (a_1 (a_2 a_3)).$$

As we introduced in Definition 2.2, a vertex algebra is a Lie conformal algebra. On the other hand, once we have a Lie conformal algebra, one can extend it to a vertex algebra by the following proposition.

**Proposition 2.4** ([7]). *Let  $R$  be a Lie conformal algebra and consider the Lie bracket on  $R$  defined by*

$$[a, b] = \int_{-\partial}^0 d\lambda [a \ \lambda b].$$

*Then the universal enveloping algebra  $U(R)$  induces a canonical vertex algebra  $V(R)$ . More precisely, the associative product on  $U(R)$  induces the normally ordered product on  $V(R)$  and the unique  $\lambda$ -bracket on  $V(R)$  whose restriction to  $R \otimes R$  is the  $\lambda$ -bracket on  $R$  can be obtained by the non-commutative Wick formula.*

The vertex algebra  $V(R)$  in Proposition 2.4 is called the *universal enveloping vertex algebra* of  $R$ . By the construction of  $V(R)$ , one can see that  $V(R)$  is strongly generated by  $R$ . Here, the phrase *strongly generated* means that  $V(R)$  is spanned by the normally ordered products of a  $\mathbb{C}[\partial]$ -module  $R$ . Moreover, since  $V(R)$

is defined via the universal enveloping algebra of  $R$ , one can realize  $V(R)$  is *freely generated* by  $R$ , that is, if  $\{v_i | i \in I\}$  is an ordered  $\mathbb{C}$ -basis of  $R$  then

$$\{v_{i_1} v_{i_2} \cdots v_{i_l} \mid i_1 \leq i_2 \leq \cdots \leq i_l, i_t < i_{t+1} \text{ if } p(v_{i_t}) = 1\}$$

is a basis of  $V(R)$ .

**Example 2.5** (Affine vertex algebra). Let  $\mathfrak{g}$  be a Lie superalgebra with an even invariant supersymmetric bilinear form  $(\cdot | \cdot)$ . The  $\mathbb{C}[\partial]$ -module  $\text{Cur}\mathfrak{g} = \mathbb{C}[\partial] \otimes \mathfrak{g} \oplus \mathbb{C}K$  with the  $\lambda$ -brackets

$$[a \lambda b] = [a, b] + K\lambda(ab), \quad [K \lambda a] = [K \lambda K] = 0$$

for  $a, b \in \mathfrak{g}$ , is called the *current Lie conformal algebra*. For  $k \in \mathbb{C}$ , the quotient of the universal enveloping vertex algebra  $V^k(\mathfrak{g}) := V(\text{Cur}\mathfrak{g}) / \langle K - k \rangle$  is called the *affine vertex algebra of level  $k$* .

**Example 2.6** (Virasoro vertex algebra). The Lie conformal algebra  $\text{Vir} = \mathbb{C}[\partial] \otimes \mathbb{C}L \oplus \mathbb{C}C$  endowed with the  $\lambda$ -brackets

$$(2.2) \quad [L \lambda L] = (\partial + 2\lambda)L + \frac{C}{12}\lambda^3, \quad [C \lambda L] = [C \lambda C] = 0$$

is called the *Virasoro Lie conformal algebra*. For  $c \in \mathbb{C}$ , the *Virasoro vertex algebra of central charge  $c$*  is the quotient vertex algebra  $\text{Vir}^c := V(\text{Vir}) / \langle C - c \rangle$ .

In a vertex algebra  $V$ , an even vector  $L \in V$  is called a *conformal vector* if  $L_{(0)} = \partial$ ,  $L_{(1)}$  is diagonalizable on  $V$  and it satisfies the Virasoro relation (2.2) with central charge  $c$ . If a vertex algebra has a conformal vector then it is called a *conformal vertex algebra*. For a conformal vector  $L$  of a conformal vertex algebra  $V$ , if  $a \in V$  satisfies  $[L \lambda a] = (\partial + \Delta_a)a + O(\lambda^2)$  for  $\Delta_a \in \mathbb{C}$ , then  $\Delta_a$  is called the *conformal weight* of  $a$ . If  $O(\lambda^2) = 0$ , then we call  $a$  a *primary of conformal weight  $\Delta_a$* .

**Example 2.7** ([13, 19, 21]). The following vertex algebras are the most well-known examples of the extensions of Virasoro vertex algebras. The  $N = 1$  *super-Virasoro vertex algebra*  $\text{SVir}^c$  is generated by an even vector  $L$  and an odd vector  $G$  satisfying the following super-Virasoro relation

$$(2.3) \quad [L \lambda L] = (\partial + 2\lambda)L + \frac{c}{12}\lambda^3, \quad [L \lambda G] = (\partial + \frac{3}{2}\lambda)G, \quad [G \lambda G] = 2L + \frac{c}{3}\lambda^2.$$

The  $N = 2$  *super-Virasoro vertex algebra*  $\text{SVir}_{N=2}^c$  is generated by two even vectors  $L, J$ , and two odd vectors  $G^+, G^-$ , satisfying the following  $N = 2$  super-Virasoro relation

$$(2.4) \quad [L \lambda L] = (\partial + 2\lambda)L + \frac{c}{12}\lambda^3, \quad [L \lambda G^\pm] = (\partial + \frac{3}{2}\lambda)G^\pm, \quad [G^\pm \lambda G^\pm] = 0,$$

$$[G^+ \lambda G^-] = L + (\frac{1}{2}\partial + \lambda)J + \frac{c}{6}\lambda^2, \quad [L \lambda J] = (\partial + \lambda)J, \quad [G^\pm \lambda J] = \mp G^\pm, \quad [J \lambda J] = \frac{c}{3}\lambda.$$

The  $N = 3$  *super-Virasoro vertex algebra* is generated by a conformal vector  $L$  with central charge  $c$ , three even primary vectors  $J^+, J^-, J^0$  of conformal weight 1, three odd primary vectors  $G^+, G^-, G^0$  of conformal weight  $\frac{3}{2}$ , and an odd primary vector  $\Phi$  of conformal weight  $\frac{1}{2}$ , satisfying the following  $N = 3$  super-Virasoro relation where the non-zero  $\lambda$ -brackets are:

$$(2.5) \quad [G^+ \lambda G^-] = L + (\frac{1}{2}\partial + \lambda)J^0 + \frac{c}{6}\lambda^2, \quad [G^\pm \lambda G^0] = \pm(\frac{1}{2}\partial + \lambda)J^\pm, \quad [G^0 \lambda G^0] = L + \frac{c}{6}\lambda^2,$$

$$[J^+ \lambda J^-] = J^0 + \frac{c}{3}\lambda, \quad [J^\pm \lambda J^0] = \mp J^\pm, \quad [J^0 \lambda J^0] = \frac{c}{3}\lambda,$$

$$[G^\pm \lambda J^\mp] = \mp G^0 + (\partial + \lambda)\Phi, \quad [G^\pm \lambda J^0] = [G^0 \lambda J^\pm] = \mp G^\pm, \quad [G^0 \lambda J^0] = -(\partial + \lambda)\Phi,$$

$$[G^\pm \lambda \Phi] = \frac{1}{2}J^\pm, \quad [G^0 \lambda \Phi] = -\frac{1}{2}J^0, \quad [\Phi \lambda \Phi] = \frac{c}{6}.$$

Note that, for the  $N = 3$  super-Virasoro vertex algebra, there is an automorphism of the form

$$G^\pm \mapsto G^\mp, \quad G^0 \mapsto -G^0, \quad J^\pm \mapsto J^\mp, \quad J^0 \mapsto -J^0.$$

For  $\mu \in \mathbb{C}^*$ , the following map defined by  $G^\pm \mapsto \mu^\pm G^\pm$ ,  $J^\pm \mapsto \mu^\pm J^\pm$  is also an automorphism of the  $N = 3$  super-Virasoro vertex algebra. For more details, the reader is referred to [23].

An odd vector  $G$  of a vertex algebra  $V$  is called a *superconformal vector* if it satisfies the super-Virasoro relation (2.3) with a conformal vector  $L = \frac{1}{2}G_{(0)}G$  of central charge  $c$ . In this case, we say that  $V$  is an  $N = 1$  *superconformal vertex algebra*.

**2.2. Supersymmetric vertex algebras.** In this section, we introduce supersymmetric (SUSY) vertex algebras using so-called  $\Lambda$ -bracket. It can be understood as an analogous definition to that of vertex algebras in Definition 2.2. We remark, all the SUSY vertex algebras considered in this paper are the  $N_K = n$  SUSY vertex algebras in [13]. For the simplicity of notations, let us denote by  $N = n$  SUSY vertex algebras, instead of  $N_K = n$  SUSY vertex algebras. The main reference of this section is [13].

Let  $R$  be a  $\mathbb{C}[\nabla]$ -module, where  $\nabla = (\partial, D^1, D^2, \dots, D^n)$  consists of an even operator  $\partial$  and odd operators  $D^i$  such that

$$(2.6) \quad [D^i, D^j] = \delta_{i,j} 2\partial$$

for  $i, j = 1, 2, \dots, n$ . Note that the bracket in (2.6) is a supercommutator. In addition, consider a tuple  $\Lambda := (\lambda, \chi^1, \chi^2, \dots, \chi^n)$  of an even formal variable  $\lambda$  and odd formal variables  $\chi^i$  which are subject to the relations

$$[\chi^i, \chi^j] = -\delta_{i,j} 2\lambda,$$

where  $i, j = 1, 2, \dots, n$ , and the bracket is again a supercommutator. Now, a  $\Lambda$ -bracket  $R \otimes R \rightarrow \mathbb{C}[\Lambda] \otimes R$  is a degree  $\bar{n}$  linear map with the following property:

$$(\text{Sesquilinearity}) \quad [D^i a \underset{\Lambda}{\smile} b] = (-1)^{n+1} \chi^i [a \underset{\Lambda}{\smile} b], \quad [a \underset{\Lambda}{\smile} D^i b] = (-1)^{p(a)+n} (D^i + \chi^i) [a \underset{\Lambda}{\smile} b].$$

The RHS of the second equality in sesquilinearity can be computed by the relation

$$(2.7) \quad [D^i, \chi^j] = \delta_{i,j} 2\lambda, \quad [D^i, \lambda] = 0.$$

**Definition 2.8** ([13]). An  $N = n$  *supersymmetric (SUSY) Lie conformal algebra* is a  $\mathbb{Z}/2\mathbb{Z}$ -graded  $\mathbb{C}[\nabla]$ -module  $R$  with a  $\Lambda$ -bracket which is a  $\mathbb{C}$ -linear map of degree  $\bar{n}$ :

$$[\underset{\Lambda}{\smile} ] : R \otimes R \rightarrow \mathbb{C}[\Lambda] \otimes R, \quad a \otimes b \mapsto [a \underset{\Lambda}{\smile} b],$$

satisfying the following conditions:

$$(\text{Skew-symmetry}) \quad [b \underset{\Lambda}{\smile} a] = (-1)^{p(a)p(b)+n+1} [a \underset{-\nabla-\Lambda}{\smile} b],$$

$$(\text{Jacobi identity}) \quad [a \underset{\Lambda}{\smile} [b \underset{\Gamma}{\smile} c]] = (-1)^{(p(a)+1)n} [[a \underset{\Lambda}{\smile} b] \underset{\Lambda+\Gamma}{\smile} c] + (-1)^{(p(a)+n)(p(b)+n)} [b \underset{\Gamma}{\smile} [a \underset{\Lambda}{\smile} c]],$$

for  $a, b, c \in R$ . We remark that the Jacobi identity holds in  $\mathbb{C}[\Lambda, \Gamma]$ , where  $\Gamma = (\gamma, \eta^1, \eta^2, \dots, \eta^n)$  consists of an even formal variable  $\gamma$  and odd formal variables  $\eta^i$  with the relations  $[\eta^i, \eta^j] = -\delta_{i,j} 2\gamma$  and  $\Lambda$  and  $\Gamma$  are supercommute. In addition,  $\mathbb{C}[\Lambda, \Gamma] \otimes R$  is a  $\mathbb{C}[\nabla]$ -module via the relation (2.7) and

$$[D^i, \eta^j] = \delta_{i,j} 2\gamma, \quad [D^i, \gamma] = 0.$$

In addition, the  $\Lambda$ -bracket supercommutes with  $\mathbb{C}[\Lambda, \Gamma]$ . For example,  $[a \underset{\Lambda}{\smile} \eta^i b] = (-1)^{p(a)+n} \eta^i [a \underset{\Lambda}{\smile} b]$  for any  $a, b \in R$ .

In particular, we denote the  $\Lambda$ -bracket for  $N = 1$  SUSY Lie conformal algebras by

$$[a \ \Lambda \ b] = \sum_{m \in \mathbb{Z}_+} \frac{\lambda^m}{m!} a_{(m|0)} b + \chi \sum_{m \in \mathbb{Z}_+} \frac{\lambda^m}{m!} a_{(m|1)} b,$$

and the  $\Lambda$ -bracket for  $N = 2$  SUSY Lie conformal algebras by

$$[a \ \Lambda \ b] = \sum_{m \in \mathbb{Z}_+} \frac{\lambda^m}{m!} a_{(m|00)} b - \chi^1 \sum_{m \in \mathbb{Z}_+} \frac{\lambda^m}{m!} a_{(m|10)} b - \chi^2 \sum_{m \in \mathbb{Z}_+} \frac{\lambda^m}{m!} a_{(m|01)} b - \chi^1 \chi^2 \sum_{m \in \mathbb{Z}_+} \frac{\lambda^m}{m!} a_{(m|11)} b.$$

**Definition 2.9** ([13]). An  $N = n$  supersymmetric (SUSY) vertex algebra is a tuple  $(V, \nabla, [ \ \Lambda \ ], |0\rangle, : :)$  such that:

- $(V, \nabla, [ \ \Lambda \ ])$  is an  $N = n$  SUSY Lie conformal algebra,
- $(V, \nabla, |0\rangle, : :)$  is a unital differential superalgebra satisfying the following properties:

$$\text{(Quasi-commutativity)} \quad : ab : - (-1)^{p(a)p(b)} : ba : = \int_{-\nabla}^0 [a \ \Lambda \ b] \, d\Lambda,$$

(Quasi-associativity)

$$:: ab : c : - : a : bc :: = \left( \int_0^\nabla d\Lambda a \right) [b \ \Lambda \ c] : + (-1)^{p(a)p(b)} : \left( \int_0^\nabla d\Lambda b \right) [a \ \Lambda \ c] :,$$

- the  $\Lambda$ -bracket and the product  $: :$  are related by

(Non-commutative Wick formula)

$$[a \ \Lambda \ : bc :] = : [a \ \Lambda \ b] c : + (-1)^{(p(a)+n)p(b)} : b [a \ \Lambda \ c] : + \int_0^\Lambda [[a \ \Lambda \ b] \ \Gamma \ c] \, d\Gamma,$$

where the integral  $\int_0^\Lambda d\Gamma$  is computed as  $\partial_{\eta^1} \partial_{\eta^2} \cdots \partial_{\eta^n} \int_0^\Lambda d\gamma$ .

As an analogue of the universal enveloping vertex algebras, one can construct the universal enveloping  $N = n$  SUSY vertex algebra  $V(R)$  of an  $N = n$  SUSY Lie conformal algebra  $R$ . More precisely,  $V(R)$  is freely generated by  $R$  with respect to the  $N = n$  SUSY normally ordered product and the  $\Lambda$ -bracket on  $V(R)$  is induced from the  $\Lambda$ -bracket on  $R$  and the  $N = n$  SUSY Wick formula (see [13]). We also note that, throughout this article, the normally ordered product  $: ab :$  is denoted by  $ab$  for  $a, b \in V$  as in the non-SUSY case (see Remark 2.3) to simplify notations.

**Example 2.10** ([13]). Let  $\mathfrak{g}$  be a Lie superalgebra with an even invariant supersymmetric bilinear form  $(|)$ . For an even  $n$ , the  $\mathbb{C}[\nabla]$ -module  $R = \mathbb{C}[\nabla] \otimes \mathfrak{g} \oplus \mathbb{C}K$  with the  $\Lambda$ -brackets

$$[a \ \Lambda \ b] = [a, b] + K\lambda(a|b)$$

for  $a, b \in \mathfrak{g}$ , is an  $N = n$  SUSY Lie conformal algebra. For an odd  $n$ , the  $\mathbb{C}[\nabla]$ -module  $R = \mathbb{C}[\nabla] \otimes \bar{\mathfrak{g}} \oplus \mathbb{C}K$  with the  $\Lambda$ -brackets

$$[\bar{a} \ \Lambda \ \bar{b}] = (-1)^{p(a)} (\overline{[a, b]}) + K \left( \sum_{i=1}^n \chi^i \right) (a|b)$$

for  $\bar{a}, \bar{b} \in \bar{\mathfrak{g}}$ , is an  $N = n$  SUSY Lie conformal algebra. Here  $\bar{\mathfrak{g}}$  denotes  $\mathfrak{g}$  with reversed parity. The quotient of the universal enveloping  $N = n$  SUSY vertex algebra  $V_{N=n}^k(\mathfrak{g}) := V(R)/\langle K - k \rangle$  is called the  $N = n$  SUSY affine vertex algebra of level  $k$ .

We remark that every  $N = n$  SUSY vertex algebra  $V$  is an  $N = n'$  SUSY vertex algebra for  $n' < n$ . In particular, if  $V$  is an  $N = 1$  SUSY vertex algebra, the  $m$ -th product  $a_{(m)} b := a_{(m|1)} b$  for  $a, b \in V$



and  $m \in \mathbb{Z}_+$  induces a vertex algebra structure on  $V$  together with the normally ordered product and the derivation  $\partial = \frac{1}{2}[D, D]$ . For instance, the  $N = 1$  SUSY affine vertex algebra  $V_{N=1}^k(\mathfrak{g})$  in Example 2.10 is a vertex algebra strongly generated by  $\mathbb{C}[\partial] \otimes (\bar{\mathfrak{g}} \oplus D\bar{\mathfrak{g}})$ . In addition, the differential algebra homomorphism  $V^k(\mathfrak{g}) \rightarrow V_{N=1}^k(\mathfrak{g})$  for the affine vertex algebra  $V^k(\mathfrak{g})$  (see Example 2.5) defined by  $a \mapsto D\bar{a}$  is an injective vertex algebra homomorphism. In other words,  $V^k(\mathfrak{g})$  is a proper vertex subalgebra of  $V_{N=1}^k(\mathfrak{g})$ . If  $V$  is an  $N = 2$  SUSY vertex algebra, the  $m$ -th product  $a_{(m)}b := a_{(m|11)}b$  for  $m \in \mathbb{Z}_+$  induces a vertex algebra structure on  $V$ .

**Example 2.11** ([10, 13]). As an  $N = 1$  SUSY vertex algebra, the  $N = 1$  super-Virasoro vertex algebra in Example 2.7 is generated by an odd vector  $G$  with the  $N = 1$   $\Lambda$ -bracket relation (see Example 5.5 of [13])

$$[G \wedge G] = (2\partial + 3\lambda + \chi D)G + \frac{c}{3}\lambda^2\chi,$$

and the  $N = 2$  super-Virasoro vertex algebra is generated by an even vector  $J$  as an  $N = 2$  SUSY vertex algebra (see Section 3.2 of [10]), with the  $N = 2$   $\Lambda$ -bracket relation

$$[J \wedge J] = (2\partial + 2\lambda + \chi^1 D^1 + \chi^2 D^2)J + \frac{c}{3}\lambda\chi^1\chi^2.$$

As in Section 1.2 of [13], an  $N = 1$  superconformal vertex algebra  $V$  with a superconformal vector  $G \in V$  has an  $N = 1$  SUSY vertex algebra structure by taking the odd derivation  $D := G_{(0)}$ . Then, as an  $N = 1$  SUSY vertex algebra,  $V$  is a conformal  $N = 1$  SUSY vertex algebra with a conformal vector  $G$  (see Definition 5.6 of [13]). Conversely, by using the sesquilinearity of  $N = 1$  SUSY Lie conformal algebras, we can check that a conformal  $N = 1$  SUSY vertex algebra  $V$  with a conformal vector  $G$  is just an  $N = 1$  superconformal vertex algebra with a superconformal vector  $G$  as a vertex algebra.

### 3. SUPERSYMMETRIC EXTENSION THEOREM

Recall that an  $N = 1$  (resp.  $N = 2$ ) SUSY vertex algebra  $V$  is a vertex algebra with the  $m$ -th product for  $m \in \mathbb{Z}_+$  defined by  $a_{(m)}b := a_{(m|1)}b$  (resp.  $a_{(m)}b := a_{(m|11)}b$ ). Conversely, in this section, we investigate conditions to embed a given vertex algebra into an  $N = 1$  SUSY vertex algebra. In Theorem 3.6, we also present the construction of an  $N = 1$  SUSY extension of  $V(R)$  for any Lie conformal algebra  $R$ , by considering some proper ghost vectors which consist of the SUSY counterpart of vectors in the given Lie conformal algebra.

In order to impose an  $N = 1$  SUSY structure on a vertex algebra  $V$ , we need an odd derivation  $D$  on  $V$  which satisfies  $[D, D] = 2\partial$  and

$$(3.1) \quad D[a \wedge b] = [Da \wedge b] + (-1)^{p(a)}[a \wedge Db],$$

where  $[D, \lambda] = 0$  (see Section 5.9 of [19]). In other words, the operator  $D$  should also be an odd derivation with respect to the  $\lambda$ -bracket. Indeed, by assuming the above properties, the  $\Lambda$ -bracket

$$(3.2) \quad [a \wedge b] := [Da \wedge b] + \chi[a \wedge b],$$

defined on  $V$  satisfies all the axioms for  $N = 1$  SUSY vertex algebras. The sesquilinearity, skew-symmetry and Jacobi identity of  $N = 1$  SUSY Lie conformal algebras follow from those properties of non-SUSY Lie conformal algebras, (3.1) and the property  $[D, D] = 2\partial$ . The quasi-commutativity, quasi-associativity and non-commutative Wick formula of  $N = 1$  SUSY vertex algebras also follow from those properties of vertex algebras. For example, the non-commutative Wick formula of  $N = 1$  SUSY vertex algebras holds since we

can check the following computations:

$$\begin{aligned}
[a \wedge bc] &= [Da \wedge bc] + \chi[a \wedge bc] \\
&= ([Da \wedge b]c + (-1)^{(p(a)+1)p(b)}b[Da \wedge c] + \int_0^\lambda [[Da \wedge b] \wedge_\gamma c] d\gamma) \\
&\quad + \chi([a \wedge b]c + (-1)^{p(a)p(b)}b[a \wedge c] + \int_0^\lambda [[a \wedge b] \wedge_\gamma c] d\gamma) \\
&= ([Da \wedge b]c + \chi[a \wedge b]c) + (-1)^{(p(a)+1)p(b)}(b[Da \wedge c] + (-1)^{p(b)}\chi b[a \wedge c]) \\
&\quad + \partial_\eta \int_0^\lambda [D[Da \wedge b] \wedge_\gamma c] + \eta[[Da \wedge b] \wedge_\gamma c] - \chi[D[a \wedge b] \wedge_\gamma c] - \chi\eta[[a \wedge b] \wedge_\gamma c] d\gamma \\
&= [a \wedge b]c + (-1)^{(p(a)+1)p(b)}b[a \wedge c] + \int_0^\lambda [[a \wedge b] \wedge_\Gamma c] d\Gamma,
\end{aligned}$$

for the  $\wedge$ -bracket in (3.2) and  $a, b, c \in V$ . Hence if there is an odd derivation  $D$  of the normally ordered product satisfying  $[D, D] = 2\partial$  and (3.1), then  $V$  itself can be considered as an  $N = 1$  SUSY vertex algebra.

Now, let us discuss the case when we cannot find such an odd derivation  $D$ . In this case, we aim to see if  $V$  can be embedded into an  $N = 1$  SUSY vertex algebra. More concretely, the following question is the main topic we will deal with in the rest of Section 3.

**Question 3.1.** Consider a Lie conformal algebra

$$(3.3) \quad R = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{u_i\}_{i \in I} \oplus E_C,$$

where the set  $\{u_i\}_{i \in I}$  is linearly independent and  $E_C$  is the central extension part of  $R$  with  $\partial E_C = 0$ . Can we find a Lie conformal algebra  $\tilde{R}$  containing  $R$  such that the corresponding universal enveloping vertex algebra  $V(\tilde{R})$  is an  $N = 1$  SUSY vertex algebra?

If there is an  $N = 1$  SUSY vertex algebra  $V(\tilde{R})$  containing the original vertex algebra  $V(R)$  as a vertex subalgebra, then we say  $V(\tilde{R})$  is an  $N = 1$  SUSY extension of  $V(R)$ . Suppose  $R$  is the Lie conformal algebra and  $I$  is the index set in (3.3). For subindex sets  $J, \bar{J} \subset I$  such that  $I = J \sqcup \bar{J}$ , we consider new data  $\bar{R}, D$  and  $V(\bar{R})$  for  $\tilde{R} = R \oplus \bar{R}$  which are given as follows:

(D-1)  $\bar{R} = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{v_i\}_{i \in I}$  is a  $\mathbb{C}[\partial]$ -module and  $p(v_i) = 1 - p(u_i)$  for any  $i \in I$ .

(D-2)  $D$  is an odd endomorphism on  $\tilde{R} = R \oplus \bar{R}$ , such that  $[D, D] = 2\partial$  and  $D(u_j) = v_j$ ,  $D(v_{\bar{j}}) = u_{\bar{j}}$  for  $j \in J$ ,  $\bar{j} \in \bar{J}$ , and  $D(E_C) = 0$ . For the simplicity of notations, let us denote  $\bar{v}_j := u_j$  and  $\bar{u}_{\bar{j}} := v_{\bar{j}}$ . Then we have

$$D(\bar{u}) = u \quad \text{for } u \in \{v_j \mid j \in J\} \cup \{u_{\bar{j}} \mid \bar{j} \in \bar{J}\}.$$

(D-3) Take an ordered  $\mathbb{C}$ -basis of  $R$  and extend it to an ordered basis  $\{r_i\}_{i \in \mathcal{I}}$  of  $\tilde{R}$ . Let  $V(\tilde{R})$  be a  $\mathbb{C}$ -algebra endowed with a basis

$$\{r_{i_1} r_{i_2} \cdots r_{i_l} \mid l \in \mathbb{Z}_+, i_1 \leq i_2 \leq \cdots \leq i_l, i_t < i_{t+1} \text{ if } p(r_{i_t}) = 1\},$$

where if  $l = 0$  then  $r_{i_1} r_{i_2} \cdots r_{i_l} := 1$ . Moreover, we assume  $V(\tilde{R})$  is the differential algebra with the odd derivation  $D$  induced from  $D$  in (D-2).

In the rest of this section, we will look for assumptions to give an  $N = 1$  SUSY vertex algebra structure on  $V(\tilde{R})$ . The following example shows  $bc$ - $\beta\gamma$  system is an  $N = 1$  SUSY extension of  $\beta\gamma$  system (see Example 5.12 of [13]).

**Example 3.2** ( $\beta\gamma$  system vs  $bc$ - $\beta\gamma$  system). Let  $R = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{\beta, \gamma\} \oplus \mathbb{C}C$  be an even Lie conformal algebra with the  $\lambda$ -bracket

$$[\beta \lambda \gamma] = -[\gamma \lambda \beta] = C,$$

where  $C$  is a central element and satisfies  $\partial C = 0$ . Then the universal enveloping vertex algebra  $V(R)$  is called the  $\beta\gamma$  system. Let  $\bar{R} = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{b, c\}$  and  $\tilde{R} = R \oplus \bar{R} = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{\beta, \gamma, b, c\} \oplus \mathbb{C}C$ , where  $b$  and  $c$  are odd elements. If the non-zero  $\lambda$ -brackets between the generators of  $\tilde{R}$  are given by

$$[\beta \lambda \gamma] = C, \quad [\gamma \lambda \beta] = -C, \quad [b \lambda c] = C, \quad [c \lambda b] = C,$$

then the universal enveloping vertex algebra  $V(\tilde{R})$  is called the  $bc$ - $\beta\gamma$  system and it is an  $N = 1$  SUSY extension of  $V(R)$ . Indeed, for the odd derivation  $D$  on  $\tilde{R}$  defined by

$$D(b) = \beta, \quad D(c) = \partial\gamma, \quad D(\beta) = \partial b, \quad D(\gamma) = c,$$

i.e.,  $b = \bar{\beta}$  and  $\gamma = \bar{c}$ , the  $\lambda$ -bracket on  $V(\tilde{R})$  can be rewritten via a  $\Lambda$ -bracket as follows:

$$[b \Lambda \gamma] = [\gamma \Lambda b] = C.$$

To deal with general cases, observe the sesquilinearity of an  $N = 1$  SUSY vertex algebra  $V$  in terms of the  $\lambda$ -bracket

$$[a \lambda b] = \sum_{m \in \mathbb{Z}_+} \frac{\lambda^m}{m!} a_{(m|1)} b \quad \text{for } a, b \in V.$$

Note that  $V$  is a vertex algebra with respect to the above  $\lambda$ -bracket. Consider  $\bar{a}$  and  $\bar{b}$  in  $V$  such that  $D\bar{a} = a$  and  $D\bar{b} = b$ . Then the sesquilinearity of  $N = 1$  SUSY Lie conformal algebras for the pair  $(\bar{a}, \bar{b})$  tells that  $[a \lambda b] = (-1)^{p(a)}(D[a \lambda \bar{b}] + \lambda[\bar{a} \lambda \bar{b}])$ , and  $[\bar{a} \lambda b] = (-1)^{p(a)}([a \lambda \bar{b}] - D[\bar{a} \lambda \bar{b}])$ . Equivalently, we have

$$\text{(SEF1)} \quad [D\bar{a} \lambda D\bar{b}] = (-1)^{p(\bar{a})+1}(D[D\bar{a} \lambda \bar{b}] + \lambda[\bar{a} \lambda \bar{b}]),$$

$$\text{(SEF2)} \quad [\bar{a} \lambda D\bar{b}] = (-1)^{p(\bar{a})+1}([D\bar{a} \lambda \bar{b}] - D[\bar{a} \lambda \bar{b}]),$$

which are called *supersymmetric extension formulas*. These formulas are crucially needed to see if the differential algebra  $V(\tilde{R})$  in (D-3) is indeed an  $N = 1$  SUSY vertex algebra.

**Definition 3.3.** Let  $R$  be a  $\mathbb{C}[\partial]$ -module with a  $\lambda$ -bracket. For a subset  $S \subseteq R$  and an odd endomorphism  $D \in \text{End}(R)$  such that  $[D, D] = 2\partial$ , we say the pair  $(S, D)$  satisfies the supersymmetric extension formulas, if (SEF1) and (SEF2) hold for all  $\bar{a}, \bar{b} \in S$ . In this case, we also call the subset  $S$  a *supersymmetric generator in the  $D$ -direction*.

Under the assumptions in Definition 3.3, note that if an ordered pair  $(\bar{a}, \bar{b})$  satisfies (SEF1) and (SEF2), then the odd endomorphism  $D$  acts as an odd derivation on the following  $\lambda$ -brackets:

$$[\bar{a} \lambda \bar{b}], \quad [D\bar{a} \lambda \bar{b}], \quad [\bar{a} \lambda D\bar{b}], \quad [D\bar{a} \lambda D\bar{b}],$$

since we know that the following equalities hold:

$$\begin{aligned} D[\bar{a} \lambda D\bar{b}] &= (-1)^{p(\bar{a})}(D[D\bar{a} \lambda \bar{b}] - \partial[\bar{a} \lambda \bar{b}]) \\ &= [D\bar{a} \lambda D\bar{b}] + (-1)^{p(\bar{a})+1}(\partial + \lambda)[\bar{a} \lambda \bar{b}] = [D\bar{a} \lambda D\bar{b}] + (-1)^{p(\bar{a})}[\bar{a} \lambda \partial\bar{b}], \\ \text{(3.4)} \quad D[D\bar{a} \lambda D\bar{b}] &= (-1)^{p(\bar{a})}(\partial[D\bar{a} \lambda \bar{b}] + \lambda D[\bar{a} \lambda \bar{b}]) \\ &= -\lambda[\bar{a} \lambda D\bar{b}] + (-1)^{p(\bar{a})}(\partial + \lambda)[D\bar{a} \lambda \bar{b}] = [\partial\bar{a} \lambda D\bar{b}] + (-1)^{p(\bar{a})}[D\bar{a} \lambda \partial\bar{b}]. \end{aligned}$$

Furthermore, Proposition 3.4 shows the two formulas (SEF1) and (SEF2) are sufficient to give solutions to Question 3.1.

**Proposition 3.4.** *Let  $R = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{u_i\}_{i \in I} \oplus E_C$  be a Lie conformal algebra in (3.3), and let  $J$  and  $\bar{J}$  be subsets of  $I$  such that  $I = J \sqcup \bar{J}$ . Suppose  $\tilde{R}$  and  $D$  are given by the data in (D-1) and (D-2), and the  $\lambda$ -brackets between any two elements in  $\{u_i, v_i\}_{i \in I}$  are given so that  $\tilde{R}$  is a Lie conformal algebra. If  $S = \{u_i = \bar{v}_i \mid i \in J\} \cup \{v_i = \bar{u}_i \mid i \in \bar{J}\}$  is a supersymmetric generator with the  $D$ -direction, then  $V(\tilde{R})$  is an  $N = 1$  SUSY vertex algebra with the odd derivation  $D$ .*

*Proof.* From (3.4), the odd endomorphism  $D$  is indeed an odd derivation of the  $\lambda$ -brackets of  $\tilde{R}$ . Moreover, from the non-commutative Wick formula, the following equality holds:

$$\begin{aligned}
D[a \lambda bc] &= D([a \lambda b]c + (-1)^{p(a)p(b)}b[a \lambda c] + \int_0^\lambda [[a \lambda b] \gamma c] d\gamma) \\
&= (D[a \lambda b])c + (-1)^{p(a)+p(b)}[a \lambda b]Dc \\
&\quad + (-1)^{p(a)p(b)}(Db)[a \lambda c] + (-1)^{(p(a)+1)p(b)}bD[a \lambda c] + \int_0^\lambda D[[a \lambda b] \gamma c] d\gamma \\
&= [Da \lambda b]c + (-1)^{p(a)}[a \lambda Db]c + (-1)^{p(a)+p(b)}[a \lambda b]Dc \\
&\quad + (-1)^{p(a)p(b)}(Db)[a \lambda c] + (-1)^{(p(a)+1)p(b)}b[Da \lambda c] + (-1)^{p(a)p(b)+p(a)+p(b)}b[a \lambda Dc] \\
&\quad + \int_0^\lambda [[Da \lambda b] \gamma c] d\gamma + (-1)^{p(a)} \int_0^\lambda [[a \lambda Db] \gamma c] d\gamma + (-1)^{p(a)+p(b)} \int_0^\lambda [[a \lambda b] \gamma Dc] d\gamma \\
&= [Da \lambda bc] + (-1)^{p(a)}[a \lambda (Db)c] + (-1)^{p(a)+p(b)}[a \lambda bDc] \\
&= [Da \lambda bc] + (-1)^{p(a)}[a \lambda D(bc)],
\end{aligned}$$

for  $a, b, c \in \tilde{R}$ . Hence,  $D$  is an odd derivation of the  $\lambda$ -brackets on  $V(\tilde{R})$  so that  $V(\tilde{R})$  is an  $N = 1$  SUSY vertex algebra with the  $\Lambda$ -brackets in (3.2).  $\square$

In Proposition 3.4, we find an  $N = 1$  SUSY Lie conformal algebra structure on  $\tilde{R}$  properly concatenating the  $\mathbb{C}[\partial]$ -module  $\tilde{R}$  to the Lie conformal algebra  $R$  which yields the  $N = 1$  SUSY extension  $V(\tilde{R})$  of  $V(R)$ . In this process, (SEF1) and (SEF2) are essential in the sense that they guarantee the  $N = 1$  supersymmetry of  $V(\tilde{R})$ . In the following example, we present another fundamental example for Question 3.1 when  $V(R)$  is the affine vertex algebra (see also Example 5.9 of [13]).

**Example 3.5** (Affine vertex algebra vs  $N = 1$  SUSY affine vertex algebra). Let  $\mathfrak{g}$  be a Lie superalgebra with an even invariant supersymmetric bilinear form  $(\mid)$ . Consider the current Lie conformal algebra  $\text{Curg} = \mathbb{C}[\partial] \otimes \mathfrak{g} \oplus \mathbb{C}K$  introduced in Example 2.5 and let  $\widetilde{\text{Curg}} = \mathbb{C}[\partial] \otimes (\mathfrak{g} \oplus \bar{\mathfrak{g}}) \oplus \mathbb{C}K$  be a  $\mathbb{C}[\partial]$ -module, where  $\bar{\mathfrak{g}}$  is the parity reversed vector superspace of  $\mathfrak{g}$ , i.e.  $\bar{\mathfrak{g}} = \{\bar{a} \mid a \in \mathfrak{g}, p(\bar{a}) = 1 - p(a)\}$ . Define the  $\lambda$ -bracket on  $\widetilde{\text{Curg}}$  by

$$(3.5) \quad [a \lambda b] = [a, b] + K\lambda(a|b), \quad [a \lambda \bar{b}] = (-1)^{p(a)}\overline{[a, \bar{b}]}, \quad [\bar{a} \lambda \bar{b}] = (-1)^{p(a)}K(a|b),$$

and the odd derivation  $D$  on  $\widetilde{\text{Curg}}$  by  $D\bar{a} = a$  and  $Da = \partial\bar{a}$  for  $a, b \in \mathfrak{g}$  (see Example 2.6 of [15]). Then the  $\lambda$ -brackets in (3.5) satisfy (SEF1) and (SEF2) and thus  $\bar{\mathfrak{g}}$  is a supersymmetric generator in the  $D$ -direction. In terms of  $\Lambda$ -bracket, the  $\lambda$ -bracket (3.5) on  $V(\widetilde{\text{Curg}})$  can be rewritten as

$$[\bar{a} \Lambda \bar{b}] = (-1)^{p(a)}(\overline{[a, \bar{b}]} + K\chi(a|b)),$$

and the quotient vertex algebra  $V(\widetilde{\text{Curg}})/\langle K - k \rangle$  is the  $N = 1$  SUSY affine vertex algebra  $V_{N=1}^k(\mathfrak{g})$  of level  $k$  introduced in Example 2.10.

Recall, in Example 3.2, that  $R$  is the Lie conformal algebra generated by  $\beta, \gamma$  and the central element  $C$  whose universal enveloping vertex algebra  $V(R)$  is the  $\beta\gamma$  system and the  $bc$ - $\beta\gamma$  system  $V(\tilde{R})$  is an  $N = 1$

SUSY extension of  $V(R)$ . In this case, the choice of  $\mathbb{C}[\partial]$ -module  $\bar{R}$  and derivation  $D$  is depicted by the diagram (Case 1 :  $bc$ - $\beta\gamma$  system). On the other hand, suppose we consider another  $\mathbb{C}[\partial]$ -module  $\bar{R}$  generated by  $\bar{\beta}$  and  $\bar{\gamma}$ , as in the diagram (Case 2 : No supersymmetry). Then there is no supersymmetry in  $V(\bar{R})$ .

$$\begin{array}{ccc} \bar{u}_1 = b & \xrightarrow{D} & \beta = u_1 \\ & & \\ u_2 = \gamma & \xrightarrow{D} & c = D(u_2) \end{array} \quad \begin{array}{ccc} \bar{u}_1 = \bar{\beta} & \xrightarrow{D} & \beta = u_1 \\ & & \\ \bar{u}_2 = \bar{\gamma} & \xrightarrow{D} & \gamma = u_2 \end{array}$$

(Case 1 :  $bc$ - $\beta\gamma$  system) (Case 2 : No supersymmetry)

That is because, in this case, (SEF1) does not have a solution. For example, for  $u_1 = \beta$ ,  $u_2 = \gamma$ , we cannot find the  $\lambda$ -brackets  $[u_1 \lambda \bar{u}_2]$  and  $[\bar{u}_1 \lambda \bar{u}_2]$  satisfying the equation below:

$$C = D[u_1 \lambda \bar{u}_2] + \lambda[\bar{u}_1 \lambda \bar{u}_2].$$

From the observation above, for a given Lie conformal algebra  $R$  in (3.3), the choice of  $\bar{R}$  in which  $\tilde{R} = R \oplus \bar{R}$  and  $D$  is a crucial part to give an answer to Question 3.1. The subsequent theorem (Theorem 3.6) shows the universal enveloping vertex algebra  $V(R)$  can be extended to an  $N = 1$  SUSY vertex algebra using  $\bar{R}$  and  $D$  given as follows:

(D-2')  $\bar{R} = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{\bar{u}_i\}_{i \in I} \oplus \bar{E}_C$  is the parity reversed  $\mathbb{C}[\partial]$ -module of  $R$ , where  $\bar{E}_C = \{\bar{C} \mid C \in E_C\}$ . The odd derivation  $D$  is defined by  $D(\bar{u}_i) = u_i$  for  $i \in I$ , and  $D(\bar{C}) = C$  for  $C \in E_C$ . In addition,  $\partial \bar{E}_C = 0$ .

**Theorem 3.6.** *Let  $R = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{u_i\}_{i \in I} \oplus E_C$  be a Lie conformal algebra in (3.3). Then  $\tilde{R} = R \oplus \bar{R}$  for the  $\mathbb{C}[\partial]$ -module  $\bar{R}$  given by (D-2') is a Lie conformal algebra endowed with the  $\lambda$ -bracket defined by*

$$(3.6) \quad [\bar{u}_i \lambda u_j] := \overline{[u_i \lambda u_j]}, \quad [u_i \lambda \bar{u}_j] := (-1)^{p(u_i)} \overline{[u_i \lambda u_j]}, \quad [\bar{u}_i \lambda \bar{u}_j] := 0,$$

and  $[C \lambda \tilde{R}] = [\bar{C} \lambda \tilde{R}] = 0$  for  $C \in E_C$ . Moreover,  $V(\tilde{R})$  is an  $N = 1$  SUSY extension of  $V(R)$ .

*Proof.* For simplicity, we use the notation  $\overline{[a \lambda b]} = \sum_{m \in \mathbb{Z}_+} \frac{\lambda^m}{m!} \overline{a^{(m)} b}$  for  $a, b \in R$ , where  $\overline{\partial^s u_i} = \partial^s \bar{u}_i$ . Then the skew-symmetry and the Jacobi identity of the  $\lambda$ -bracket can be verified by direct computations. For example, the Jacobi identity for the triple  $\{(u_i, u_j, \bar{u}_k) \mid i, j, k \in I\}$  can be checked as follows:

$$\begin{aligned} [u_i \lambda [u_j \gamma \bar{u}_k]] &= (-1)^{p(u_j)} [u_i \lambda \overline{[u_j \gamma u_k]}] \\ &= (-1)^{p(u_i) + p(u_j)} \overline{[u_i \lambda [u_j \gamma u_k]]} \\ &= (-1)^{p(u_i) + p(u_j)} \overline{[[u_i \lambda u_j]_{\lambda + \gamma} u_k]} + (-1)^{p(u_i)p(u_j) + p(u_i) + p(u_j)} \overline{[u_j \gamma [u_i \lambda u_k]]} \\ &= [[u_i \lambda u_j]_{\lambda + \gamma} \bar{u}_k] + (-1)^{p(u_i)p(u_j)} [u_j \gamma [u_i \lambda \bar{u}_k]]. \end{aligned}$$

Moreover, one can check the pair  $(\{\bar{u}_i\}_{i \in I}, D)$  satisfies the SUSY extension formulas. Hence, by Proposition 3.4, the vertex algebra  $V(\tilde{R}) = V(R) \otimes V(\bar{R})$  has an  $N = 1$  supersymmetry.  $\square$

The  $\lambda$ -bracket (3.6) is inspired from the  $\Lambda$ -bracket of the  $N = 1$  SUSY affine vertex algebra (see also Section 3.2.1 of [6]). In terms of  $\Lambda$ -bracket, the bracket (3.6) can be written as follows:

$$[\bar{a} \Lambda \bar{b}] = (-1)^{p(a)} \overline{[a \lambda b]}$$

for  $a, b \in R$ . Indeed, if  $R$  is a centerless current Lie conformal algebra, the  $N = 1$  SUSY extension obtained from Theorem 3.6 is the  $N = 1$  SUSY affine vertex algebra of level  $k = 0$  (see Example 2.10). In the

following examples, we find an  $N = 1$  SUSY extension of the  $\beta\gamma$  system and the Virasoro vertex algebra using Theorem 3.6.

**Example 3.7** ( $N = 1$  SUSY extension of  $\beta\gamma$  system). Let  $R = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{\beta, \gamma\} \oplus \mathbb{C}C$  be the Lie conformal algebra for the  $\beta\gamma$  system (see Example 3.2). If we apply Theorem 3.6 to  $R$  by taking  $u_1 = \beta$  and  $u_2 = \gamma$ , we know that  $\widetilde{R} = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{\beta, \gamma, \bar{\beta}, \bar{\gamma}\} \oplus \mathbb{C}C \oplus \mathbb{C}\bar{C}$  with the remaining non-zero  $\lambda$ -brackets

$$[\beta \ \lambda \ \bar{\gamma}] = [\bar{\beta} \ \lambda \ \gamma] = \bar{C},$$

is an  $N = 1$  SUSY extension of the  $\beta\gamma$  system. Note that  $V(\widetilde{R})$  is a distinct vertex algebra from the  $bc$ - $\beta\gamma$  system, which is another SUSY extension introduced in Example 3.2.

**Example 3.8** ( $N = 1$  SUSY extension of Virasoro vertex algebra). Consider the *centerless Virasoro Lie conformal algebra*  $\text{Vir} = \mathbb{C}[\partial] \otimes \mathbb{C}L$  with the  $\lambda$ -bracket  $[L \ \lambda \ L] = (\partial + 2\lambda)L$ , and its universal enveloping vertex algebra  $\text{Vir}^0$  (see Example 2.6). First of all, for the  $N = 1$  super-Virasoro Lie conformal algebra  $\text{SVir} = \mathbb{C}[\partial] \otimes (\mathbb{C}L \oplus \mathbb{C}G)$  in which the  $\lambda$ -brackets are defined as

$$[L \ \lambda \ G] = (\partial + \frac{3}{2}\lambda)G, \quad [G \ \lambda \ G] = 2L,$$

if we define the odd derivation  $D$  by  $DG = 2L$  then  $(\{G\}, D)$  satisfies the SUSY extension formulas. Hence the  $N = 1$  super-Virasoro vertex algebra  $\text{SVir}^0$  is an  $N = 1$  SUSY extension of  $\text{Vir}^0$ . On the other hand, by using the Theorem 3.6, the universal enveloping vertex algebra  $V(\widetilde{\text{Vir}})$  for  $\widetilde{\text{Vir}} = \mathbb{C}[\partial] \otimes (\mathbb{C}L \oplus \mathbb{C}\bar{L})$  is also an  $N = 1$  SUSY extension of  $\text{Vir}^0$ , where the  $\lambda$ -brackets of  $\widetilde{\text{Vir}}$  are given by

$$[L \ \lambda \ \bar{L}] = (\partial + 2\lambda)\bar{L}, \quad [\bar{L} \ \lambda \ \bar{L}] = 0.$$

Note that, as in Example 2.7 of [13], if we take an another generating set for the  $N = 2$  super-Virasoro vertex algebra by defining  $L' := L - \frac{1}{2}\partial J$ , then we know that especially the  $\lambda$ -brackets for  $L'$  and  $G^+$  are

$$[L' \ \lambda \ L'] = (\partial + 2\lambda)L', \quad [L' \ \lambda \ G^+] = (\partial + 2\lambda)G^+, \quad [G^+ \ \lambda \ G^+] = 0.$$

Hence, the vertex algebra  $V(\widetilde{\text{Vir}})$  can be viewed as a vertex subalgebra of the  $N = 2$  super-Virasoro vertex algebra of any central charge  $c \in \mathbb{C}$ .

In Example 3.8, we introduced two distinct  $N = 1$  SUSY vertex algebras  $\text{SVir}^0$  and  $V(\widetilde{\text{Vir}})$  which are strongly generated by two elements  $L$  and  $D^{-1}(L)$ . Furthermore, an  $N = 1$  SUSY extension of  $\text{Vir}^0$  strongly generated by  $L$  and a primary field  $\bar{L}$  such that  $D(\bar{L}) = L$  is one of the two vertex algebras mentioned above. To check the detail, suppose the conformal weight of  $\bar{L}$  is  $\Delta \in \mathbb{C}$ , i.e.  $[L \ \lambda \ \bar{L}] = (\partial + \Delta\lambda)\bar{L}$ . Then, from the equation (SEF1), we have

$$[\bar{L} \ \lambda \ \bar{L}] = \lambda^{-1}([L \ \lambda \ L] - D[L \ \lambda \ \bar{L}]) = (2 - \Delta)L.$$

Moreover, by using the Jacobi identity, the possible values of  $\Delta$  is  $\frac{3}{2}$  or 2 since

$$\begin{aligned} (2 - \Delta)(\partial + 2\lambda)L &= [L \ \lambda \ [\bar{L} \ \gamma \ \bar{L}]] \\ &= [[L \ \lambda \ \bar{L}] \ \lambda + \gamma \ \bar{L}] + [\bar{L} \ \gamma \ [L \ \lambda \ \bar{L}]] \\ &= ((\Delta - 1)\lambda - \gamma)(2 - \Delta)L + (\partial + \Delta\lambda + \gamma)(2 - \Delta)L \\ &= (2 - \Delta)(\partial + (2\Delta - 1)\lambda)L. \end{aligned}$$

As the Virasoro vertex algebra  $\text{Vir}^c$  can be extended to the  $N = 1$  super-Virasoro vertex algebra  $\text{SVir}^c$ , it is natural to ask if a vertex algebra can be extended to an  $N = 1$  superconformal vertex algebra. In the following Remark 3.9, using the SUSY extension formulas, one can reproduce the  $N = 1$  superconformal

extension of the affine vertex algebra of level 0, which is called the *superconformal current algebra* in [20]. Note that the  $N = 1$  SUSY affine vertex algebra of level  $k \neq 0$  associated with a basic Lie superalgebra is an  $N = 1$  superconformal vertex algebra with the *Kac-Todorov superconformal vector* (see Section 5.9 of [19]). For the  $N = 2$  superconformal structures of the  $N = 1$  SUSY affine vertex algebra of level  $k \neq 0$  associated with the double of a Lie bialgebra, refer to [15].

**Remark 3.9** ([20]). Let  $\mathfrak{g} = \bigoplus_{\omega \in \Omega} \mathfrak{g}_\omega$  be an  $\Omega$ -graded Lie superalgebra. Then, the  $N = 1$  SUSY affine vertex algebra  $V_{N=1}^0(\mathfrak{g})$  can be embedded into an  $N = 1$  superconformal vertex algebra with the conformal weight  $\Delta_a = 1 - \omega$ , where  $a \in \mathfrak{g}_\omega$  is a homogeneous element for  $\omega \in \Omega$ . More precisely, consider the Lie conformal algebra  $\text{Curg}_{N=1} = \mathbb{C}[\partial] \otimes (\mathfrak{g} \oplus \bar{\mathfrak{g}})$  with the  $\lambda$ -brackets

$$[a \ \lambda \ b] = [a, b], \quad [a \ \lambda \ \bar{b}] = (-1)^{p(a)} \overline{[a, b]}, \quad [\bar{a} \ \lambda \ \bar{b}] = 0,$$

for  $a, b \in \mathfrak{g}$ , and let  $\text{SVir} = \mathbb{C}[\partial] \otimes (\mathbb{C}L \oplus \mathbb{C}G) \oplus \mathbb{C}C$  be the  $N = 1$  super-Virasoro Lie conformal algebra with the  $\lambda$ -brackets in (2.3), where  $C$  is central in the  $\mathbb{C}[\partial]$ -module  $\text{SVir} \oplus \text{Curg}_{N=1}$  and  $\partial C = 0$ . For the  $\lambda$ -brackets given by:

$$[L \ \lambda \ a] = (\partial + \Delta_a \lambda) a, \quad [L \ \lambda \ \bar{a}] = (\partial + (\Delta_a - \frac{1}{2}) \lambda) \bar{a},$$

where  $\Delta_a = 1 - \omega$  for  $a \in \mathfrak{g}_\omega$ , one can reproduce the remaining  $\lambda$ -brackets of the superconformal current algebra as follows

$$[G \ \lambda \ a] = (\partial + (2\Delta_a - 1) \lambda) \bar{a}, \quad [G \ \lambda \ \bar{a}] = a,$$

from the SUSY extension formulas in (SEF1) and (SEF2). Note that  $\Delta_{[a,b]} = \Delta_a + \Delta_b - 1$  for all  $a, b \in \mathfrak{g}$ , since  $\omega_{[a,b]} = \omega_a + \omega_b$ . Then, one can also check that  $\text{SVir} \oplus \text{Curg}_{N=1}$  is indeed a Lie conformal algebra. For example, the following equality holds:

$$\begin{aligned} [G \ \lambda \ [a \ \gamma \ b]] &= [G \ \lambda \ [a, b]] = (\partial + (2\Delta_{[a,b]} - 1) \lambda) \overline{[a, b]} = (\partial + (2\Delta_a + 2\Delta_b - 3) \lambda) \overline{[a, b]} \\ &= ((2\Delta_a - 2) \lambda - \gamma) \overline{[a, b]} + (\partial + (2\Delta_b - 1) \lambda + \gamma) \overline{[a, b]} \\ &= ((2\Delta_a - 2) \lambda - \gamma) [\bar{a} \ \lambda + \gamma \ b] + (-1)^{p(a)} (\partial + (2\Delta_b - 1) \lambda + \gamma) [a \ \gamma \ \bar{b}] \\ &= [(\partial + (2\Delta_a - 1) \lambda) \bar{a} \ \lambda + \gamma \ b] + (-1)^{p(a)} [a \ \gamma \ (\partial + (2\Delta_b - 1) \lambda) \bar{b}] \\ &= [[G \ \lambda \ a] \ \lambda + \gamma \ b] + (-1)^{p(a)} [a \ \gamma \ [G \ \lambda \ b]]. \end{aligned}$$

Hence,  $V(\text{SVir} \oplus \text{Curg}_{N=1}) / \langle C - c \rangle \simeq \text{SVir}^c \otimes V_{N=1}^0(\mathfrak{g})$  is an  $N = 1$  superconformal vertex algebra where the odd derivation  $D$  is given by  $DG = 2L$  and  $D\bar{a} = a$ .

Let  $\mathfrak{g}$  be a basic Lie superalgebra. For the  $N = 1$  SUSY affine vertex algebra  $V_{N=1}^k(\mathfrak{g})$  with  $k \neq 0$ , any element  $\bar{a} \in \bar{\mathfrak{g}}$  is primary of conformal weight  $\frac{1}{2}$  with respect to the Kac-Todorov superconformal vector  $\tau_{\mathfrak{g}}$  (see Section 3.4 of [8]). Hence  $a := D\bar{a} \in D(\bar{\mathfrak{g}})$  has the conformal weight 1. In addition, the element  $\tau_{\mathfrak{g},h} = \tau_{\mathfrak{g}} + \partial h \in V_{N=1}^k(\mathfrak{g})$  is a superconformal vector (see Example 2.13 of [15]) for  $h$  in the Cartan subalgebra of  $\mathfrak{g}$  and  $\tau_{\mathfrak{g},h}$  yields another conformal structure induced from the eigenspace of  $\text{adh}$ . Similarly, in Remark 3.9, one can construct an  $N = 1$  SUSY extension of  $V^0(\mathfrak{g})$  such that every element  $a \in \mathfrak{g}$  has conformal weight 1 by taking  $\Omega = \{0\}$ . Furthermore, by letting  $\Omega$  to be the grading of  $\mathfrak{g}$  defined by  $\text{adh}$ ,  $V^0(\mathfrak{g})$  can be extended to an  $N = 1$  superconformal vertex algebra with another conformal weight decomposition. In this case, the conformal weight of  $a \in \mathfrak{g}$  is the same as the conformal weight of  $a$  in  $V_{N=1}^k(\mathfrak{g})$  via the superconformal vector  $\tau_{\mathfrak{g},h}$ .

## 4. GENERALIZATIONS

The goal of this section is to generalize the discussions in the Section 3. Similar to the  $N = 1$  case, we say that an  $N = n$  SUSY vertex algebra  $V$  has an  $N = n' > n$  *supersymmetric extension* if there is a differential algebra  $W$  such that  $V \otimes W$  has an  $N = n'$  SUSY vertex algebra structure, containing the original vertex algebra  $V$  as an  $N = n$  SUSY vertex subalgebra.

Note that we can also impose an  $N = 2$  SUSY structure on a vertex algebra  $V$ , if it has two odd derivations  $D^1, D^2$  such that  $[D^i, D^j] = 2\delta_{ij}\partial$ , and the equation (3.1) holds for each  $D^i$ . Then the  $\Lambda$ -bracket:

$$(4.1) \quad [a \ \Lambda \ b] = [D^2 D^1 a \ \lambda \ b] - \chi^1 [D^2 a \ \lambda \ b] + \chi^2 [D^1 a \ \lambda \ b] - \chi^1 \chi^2 [a \ \lambda \ b]$$

gives an  $N = 2$  SUSY vertex algebra structure on  $V$ . The proof is similar to the proof of the  $N = 1$  case. For example, the non-commutative Wick formula of  $N = 2$  SUSY vertex algebras holds since:

$$\begin{aligned} [a \ \Lambda \ bc] &= [D^2 D^1 a \ \lambda \ bc] - \chi^1 [D^2 a \ \lambda \ bc] + \chi^2 [D^1 a \ \lambda \ bc] - \chi^1 \chi^2 [a \ \lambda \ bc] \\ &= ([D^2 D^1 a \ \lambda \ b]c + (-1)^{p(a)p(b)} b [D^2 D^1 a \ \lambda \ c] + \int_0^\lambda [[D^2 D^1 a \ \lambda \ b] \ \gamma \ c] \ d\gamma) \\ &\quad - \chi^1 ([D^2 a \ \lambda \ b]c + (-1)^{(p(a)+1)p(b)} b [D^2 a \ \lambda \ c] + \int_0^\lambda [[D^2 a \ \lambda \ b] \ \gamma \ c] \ d\gamma) \\ &\quad + \chi^2 ([D^1 a \ \lambda \ b]c + (-1)^{(p(a)+1)p(b)} b [D^1 a \ \lambda \ c] + \int_0^\lambda [[D^1 a \ \lambda \ b] \ \gamma \ c] \ d\gamma) \\ &\quad - \chi^1 \chi^2 ([a \ \lambda \ b]c + (-1)^{p(a)p(b)} b [a \ \lambda \ c] + \int_0^\lambda [[a \ \lambda \ b] \ \gamma \ c] \ d\gamma) \\ &= [a \ \Lambda \ b]c + (-1)^{p(a)p(b)} b [a \ \Lambda \ c] + \int_0^\Lambda [[a \ \Lambda \ b] \ \Gamma \ c] \ d\Gamma. \end{aligned}$$

Similarly, in the case of  $N = n$ , we know that  $n$  odd derivations of the normally ordered product and the  $\lambda$ -brackets, denoted as  $D^1, \dots, D^n$ , and satisfying  $[D^i, D^j] = 2\delta_{ij}\partial$ , also provide an  $N = n$  SUSY vertex algebra structure. For these  $n$  odd derivations satisfying the conditions described above, we denote by  $\text{Der}_{N=n}(V)$  the set of  $n$ -tuples of the form  $(D^1, \dots, D^n)$ , i.e. the set of all  $N = n$  SUSY structures of  $V$ . Then, in the following proposition, we can consider the complex orthogonal group action on the set of  $N = n$  SUSY structures of a vertex algebra. Indeed, the orthogonal groups were studied as the automorphism groups of the  $N = 2, 3, 4$  superconformal algebras in [23].

**Proposition 4.1.** *Let  $V$  be a vertex algebra. Then  $\text{Der}_{N=n}(V)$  admits an action of the orthogonal group in dimension  $n$ .*

*Proof.* Let  $\mathbf{O}_n(\mathbb{C})$  be the complex orthogonal group in dimension  $n$ . If we define an  $\mathbf{O}_n(\mathbb{C})$ -action by

$$A \cdot (D^1, \dots, D^n) := \left( \sum_{i=1}^n A_{1i} D^i, \dots, \sum_{i=1}^n A_{ni} D^i \right),$$

for  $A \in \mathbf{O}_n(\mathbb{C})$  and  $(D^1, \dots, D^n) \in \text{Der}_{N=n}(V)$ , then the result follows from direct calculations.  $\square$

Note that a vertex algebra is called an  $N = 2$  *superconformal vertex algebra* if it has two odd vectors  $G^\pm$  satisfying the  $N = 2$  super-Virasoro relation (2.4) with a conformal vector  $L = \frac{1}{2}(G^+_{(0)}G^- + G^-_{(0)}G^+)$  and  $J = G^+_{(1)}G^-$ . Then, from Remark 2.8 of [13] (see also Remark 1.3 of [2]), an  $N = 2$  superconformal vertex algebra has an  $N = 2$  SUSY vertex algebra structure by taking  $D^1 = (\mu G^+ + \mu^{-1} G^-)_{(0)}$  and  $D^2 = \pm\sqrt{-1}(\mu G^+ - \mu^{-1} G^-)_{(0)}$  for  $\mu \in \mathbb{C}^*$ . This  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{C}^* \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbf{SO}_2(\mathbb{C})$  family of  $N = 2$  SUSY structures is derived from the automorphism group of the  $N = 2$  super-Virasoro vertex algebra, which preserve the



conformal vector. Therefore, Proposition 4.1 can be viewed as a generalization of the relationships between SUSY vertex algebra structures induced from the superconformal structures.

We will now generalize the result from Section 3. Recall that we have shown that there exists an  $N = 1$  SUSY extension of any universal enveloping vertex algebras. Along with the result of Theorem 3.6, we can also consider the following question, which will be proven in Theorem 4.9.

**Question 4.2.** For a given  $N = 1$  supersymmetric Lie conformal algebra  $R_{N=1}$ , can we find an  $N = 2$  supersymmetric extension of the  $N = 1$  supersymmetric vertex algebra  $V(R_{N=1})$ ?

In order to answer to Question 4.2, we aim to find conditions to assign an  $N = 2$  supersymmetry on the  $\mathbb{C}[\partial]$ -module:

$$(4.2) \quad \tilde{R} = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{u_i, \bar{u}_i, u_i^\circ, \bar{u}_i^\circ\}_{i \in I} \oplus E_C,$$

where the set  $\{u_i, \bar{u}_i, u_i^\circ, \bar{u}_i^\circ\}_{i \in I}$  is linearly independent and  $E_C$  is the central extension part of  $\tilde{R}$  such that  $\partial E_C = 0$ . Additionally, we assume the parity condition in (D-4) and the existence of two odd endomorphisms with properties in (D-5).

(D-4) For the generators of  $\tilde{R}$ , their parities satisfy  $p(u_i) = p(\bar{u}_i^\circ) = 1 - p(\bar{u}_i) = 1 - p(u_i^\circ)$  for any  $i \in I$ .

(D-5)  $D^1, D^2$  are odd endomorphisms on  $\tilde{R}$  such that  $[D^i, D^j] = 2\delta_{ij}\partial$ , and satisfy the following diagram:

$$\begin{array}{ccc} & \bar{u}_i & \xrightarrow{D^1} & u_i \\ D^2 \nearrow & & & \nearrow D^2 \\ \bar{u}_i^\circ & \xrightarrow{-D^1} & u_i^\circ & \end{array}$$

for each  $i \in I$ , i.e.  $D^1(\bar{u}_i) = u_i$ ,  $D^1(\bar{u}_i^\circ) = -u_i^\circ$ ,  $D^2(u_i^\circ) = u_i$ ,  $D^2(\bar{u}_i^\circ) = \bar{u}_i$ , and  $D^1(E_C) = D^2(E_C) = 0$ .

**Proposition 4.3.** Let  $\tilde{R}$  be the  $\mathbb{C}[\partial]$ -module in (4.2) satisfying (D-4), and let  $D^1$  and  $D^2$  be given by the data in (D-5). Suppose the  $\lambda$ -brackets between the elements in  $\{u_i, \bar{u}_i, u_i^\circ, \bar{u}_i^\circ\}_{i \in I}$  are given so that  $\tilde{R}$  is a Lie conformal algebra, and each of the following pairs

$$(4.3) \quad (\{\bar{u}_i, \bar{u}_i^\circ\}_{i \in I}, D^1), \quad (\{u_i^\circ, \bar{u}_i^\circ\}_{i \in I}, D^2),$$

satisfies the SUSY extension formulas. Then  $V(\tilde{R})$  is an  $N = 2$  SUSY vertex algebra with the odd derivations  $D^1$  and  $D^2$ .

*Proof.* By the proof of Proposition 3.4, we know that each of  $D^1$  and  $D^2$  acts as an odd derivation of the  $\lambda$ -brackets on  $V(\tilde{R})$ . Hence the universal enveloping vertex algebra  $V(\tilde{R})$  is an  $N = 2$  SUSY vertex algebra via the  $\Lambda$ -brackets in (4.1).  $\square$

Note that the SUSY extension formulas for the pair  $(\{\bar{u}_i, \bar{u}_i^\circ\}_{i \in I}, D^1)$  in (4.3) can be written explicitly as the following equalities:

$$\begin{aligned} [u_i^\circ \lambda u_j^\circ] &= (-1)^{p(u_i)}(D^1[u_i^\circ \lambda \bar{u}_j^\circ] - \lambda[\bar{u}_i^\circ \lambda \bar{u}_j^\circ]), & [\bar{u}_i^\circ \lambda u_j^\circ] &= (-1)^{p(u_i)+1}([u_i^\circ \lambda \bar{u}_j^\circ] + D^1[\bar{u}_i^\circ \lambda \bar{u}_j^\circ]), \\ [u_i \lambda u_j^\circ] &= (-1)^{p(u_i)+1}(D^1[u_i \lambda \bar{u}_j^\circ] + \lambda[\bar{u}_i \lambda \bar{u}_j^\circ]), & [\bar{u}_i \lambda u_j^\circ] &= (-1)^{p(u_i)+1}([u_i \lambda \bar{u}_j^\circ] - D^1[\bar{u}_i \lambda \bar{u}_j^\circ]), \\ [u_i \lambda u_j] &= (-1)^{p(u_i)}(D^1[u_i \lambda \bar{u}_j] + \lambda[\bar{u}_i \lambda \bar{u}_j]), & [\bar{u}_i \lambda u_j] &= (-1)^{p(u_i)}([u_i \lambda \bar{u}_j] - D^1[\bar{u}_i \lambda \bar{u}_j]), \end{aligned}$$

and the SUSY extension formulas for the second pair  $(\{u_i^\circ, \bar{u}_i^\circ\}_{i \in I}, D^2)$  are given by the following equalities:

$$\begin{aligned} [\bar{u}_i \lambda \bar{u}_j] &= (-1)^{p(u_i)+1}(D^2[\bar{u}_i \lambda \bar{u}_j^\circ] + \lambda[\bar{u}_i^\circ \lambda \bar{u}_j^\circ]), & [\bar{u}_i^\circ \lambda \bar{u}_j] &= (-1)^{p(u_i)+1}([\bar{u}_i \lambda \bar{u}_j^\circ] - D^2[\bar{u}_i^\circ \lambda \bar{u}_j^\circ]), \\ [u_i \lambda \bar{u}_j] &= (-1)^{p(u_i)}(D^2[u_i \lambda \bar{u}_j^\circ] + \lambda[u_i^\circ \lambda \bar{u}_j^\circ]), & [u_i^\circ \lambda \bar{u}_j] &= (-1)^{p(u_i)}([u_i \lambda \bar{u}_j^\circ] - D^2[u_i^\circ \lambda \bar{u}_j^\circ]), \end{aligned}$$

$$[u_i \lambda u_j] = (-1)^{p(u_i)}(D^2[u_i \lambda u_j^\circ] + \lambda[u_i^\circ \lambda u_j^\circ]), \quad [u_i^\circ \lambda u_j] = (-1)^{p(u_i)}([u_i \lambda u_j^\circ] - D^2[u_i^\circ \lambda u_j^\circ]).$$

Proposition 4.3 tells that, for a given Lie conformal algebra  $R$ , if we can find a Lie conformal algebra  $\tilde{R}$  that contains  $R$  as a Lie conformal subalgebra and satisfies the conditions of the proposition, then we can conclude that  $V(R)$  has an  $N = 2$  SUSY extension.

**Example 4.4** ( $N = 2$  SUSY structure of  $N = 2$  super-Virasoro vertex algebra). From Example 2.9 of [13], we already know that the  $N = 2$  super-Virasoro vertex algebra (see also Examples 2.7 and 2.11) has an  $N = 2$  SUSY vertex algebra structure as in the diagram:

$$\begin{array}{ccc} & \sqrt{-1}(G^+ + G^-) & \xrightarrow{D^1} & 2\sqrt{-1}L \\ & \nearrow^{D^2} & & \nearrow^{D^2} \\ J & \xrightarrow{-D^1} & G^+ - G^- & \end{array}$$

Indeed, if we choose a generating set of the  $N = 2$  super-Virasoro vertex algebra  $V(\tilde{R})/\langle C - c \rangle$  as

$$\bar{u}_1^\circ = J, \quad u_1^\circ = G^+ - G^-, \quad \bar{u}_1 = \sqrt{-1}(G^+ + G^-), \quad u_1 = 2\sqrt{-1}L,$$

where  $\tilde{R} = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{u_1, \bar{u}_1, u_1^\circ, \bar{u}_1^\circ\} \oplus \mathbb{C}C$ , then the SUSY extension formulas for the pairs in (4.3) can be checked by direct computations.

In the consecutive examples (Example 4.5, 4.6 and 4.7), we investigate several  $N = 2$  SUSY vertex algebra structures of the  $N = 2$  SUSY extension of the  $bc$ - $\beta\gamma$  system. The subsequent example is a construction of an  $N = 2$  SUSY extension of the  $bc$ - $\beta\gamma$  system via Proposition 4.3.

**Example 4.5** ( $N = 2$  SUSY extension of  $bc$ - $\beta\gamma$  system). Let  $\tilde{R} = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{u_i, \bar{u}_i, u_i^\circ, \bar{u}_i^\circ\}_{i \in \{1,2\}} \oplus \mathbb{C}C$  be a  $\mathbb{C}[\partial]$ -module where  $C$  is a central element satisfying  $\partial C = 0$  and the generators are given by

$$u_1 = \beta, \quad \bar{u}_1 = b, \quad u_1^\circ = -a, \quad \bar{u}_1^\circ = \alpha, \quad u_2 = \delta, \quad \bar{u}_2 = d, \quad u_2^\circ = -c, \quad \bar{u}_2^\circ = \gamma,$$

where  $\alpha, \beta, \gamma, \delta$  are even elements and  $a, b, c, d$  are odd elements. Define the odd endomorphisms  $D^1$  and  $D^2$  on  $\tilde{R}$  as in the diagram below

$$\begin{array}{ccc} & b & \xrightarrow{D^1} & \beta \\ & \nearrow^{D^2} & & \nearrow^{D^2} \\ \alpha & \xrightarrow{-D^1} & -a & \end{array} \quad \begin{array}{ccc} & d & \xrightarrow{D^1} & \delta \\ & \nearrow^{D^2} & & \nearrow^{D^2} \\ \gamma & \xrightarrow{-D^1} & -c & \end{array}$$

with  $[D^i, D^j] = 2\delta_{ij}\partial$  and  $D^1 C = D^2 C = 0$ . For the Lie conformal algebra structure of  $\tilde{R}$ , define the non-zero  $\lambda$ -brackets of  $\tilde{R}$  as

$$\begin{aligned} [\beta \lambda \gamma] &= C, & [\gamma \lambda \beta] &= -C, & [b \lambda c] &= C, & [c \lambda b] &= C, \\ [\delta \lambda \alpha] &= -C, & [\alpha \lambda \delta] &= C, & [d \lambda a] &= -C, & [a \lambda d] &= -C. \end{aligned}$$

Then, we can check each of the pairs  $(\{\bar{u}_i, \bar{u}_i^\circ\}_{i \in \{1,2\}}, D^1)$  and  $(\{u_i^\circ, \bar{u}_i^\circ\}_{i \in \{1,2\}}, D^2)$  satisfies the SUSY extension formulas. Hence, by Proposition 4.3,  $V(\tilde{R})$  is an  $N = 2$  SUSY extension of the  $bc$ - $\beta\gamma$  system where the non-zero  $\Lambda$ -bracket of the generators as an  $N = 2$  SUSY vertex algebra is

$$[\alpha \Lambda \gamma] = -C,$$

and, we call the vertex algebra  $V(\tilde{R})$  or its quotient vertex algebra  $V(\tilde{R})/\langle C - 1 \rangle$  the *extended  $bc$ - $\beta\gamma$  system* in the rest of the paper.

In the following examples, we describe other  $N = 2$  SUSY structures derived from  $N = 2$  superconformal structures in the extended  $bc$ - $\beta\gamma$  system (see also [9, 12]).

**Example 4.6** ( $N = 2$  superconformal structures of the extended  $bc$ - $\beta\gamma$  system (1)). The extended  $bc$ - $\beta\gamma$  system has an  $N = 2$  superconformal structure associated to the following vectors

$$L_h = \frac{1}{2}(-c\partial b + \partial cb + 2\partial\gamma\beta + a\partial d - \partial ad - 2\partial\alpha\delta),$$

$$G_h^+ = \sqrt{-1}(c\beta - a\delta), \quad G_h^- = \sqrt{-1}(-\partial\gamma b + \partial\alpha d), \quad J_h = cb - ad,$$

with central charge 6, and the  $N = 2$  SUSY vertex algebra structure induced from the above  $N = 2$  superconformal structure is depicted in the following picture

$$\begin{array}{ccc} & -\beta & \xrightarrow{D_h^1} \sqrt{-1}\partial b \\ D_h^2 \nearrow & & \searrow D_h^2 \\ b & \xrightarrow{-D_h^1} -\sqrt{-1}\beta & \\ & -a & \xrightarrow{D_h^1} \sqrt{-1}\partial\alpha \\ D_h^2 \nearrow & & \searrow D_h^2 \\ \alpha & \xrightarrow{-D_h^1} -\sqrt{-1}a & \end{array} \quad \begin{array}{ccc} & -c & \xrightarrow{D_h^1} \sqrt{-1}\partial\gamma \\ D_h^2 \nearrow & & \searrow D_h^2 \\ \gamma & \xrightarrow{-D_h^1} -\sqrt{-1}c & \\ & -\delta & \xrightarrow{D_h^1} \sqrt{-1}\partial d \\ D_h^2 \nearrow & & \searrow D_h^2 \\ d & \xrightarrow{-D_h^1} -\sqrt{-1}\delta & \end{array}$$

where  $D_h^1 = (G_h^+ + G_h^-)_{(0)}$  and  $D_h^2 = \sqrt{-1}(G_h^+ - G_h^-)_{(0)}$ . Observe that the non-zero  $\Lambda$ -brackets of generators are given by

$$[b \wedge \gamma] = \chi^1 + \sqrt{-1}\chi^2, \quad [\alpha \wedge d] = -\chi^1 - \sqrt{-1}\chi^2,$$

and hence each of the  $bc$ - $\beta\gamma$  system and the  $ad$ - $\alpha\delta$  system (which refers to the vertex algebra generated by the four elements  $a, d, \alpha, \delta$ ) is already closed under the  $N = 2$  SUSY vertex algebra structure, respectively. In other words, this  $N = 2$  SUSY vertex algebra structure which is derived from the  $N = 2$  superconformal structure associated to  $G_h^\pm$  provides an *internal*  $N = 2$  SUSY structure for both the  $bc$ - $\beta\gamma$  system and the  $ad$ - $\alpha\delta$  system. Therefore, the extended  $bc$ - $\beta\gamma$  system with  $D_h^1$  and  $D_h^2$  can be understood as the product of two independent copies of the  $bc$ - $\beta\gamma$  system as  $N = 2$  SUSY vertex algebras.

**Example 4.7** ( $N = 2$  superconformal structures of the extended  $bc$ - $\beta\gamma$  system (2)). We have another  $N = 2$  superconformal structure of the extended  $bc$ - $\beta\gamma$  system induced from the following vectors

$$L_v = L_h, \quad G_v^+ = d\beta - b\delta, \quad G_v^- = -\partial\gamma a + \partial\alpha c, \quad J_v = -cb + ad.$$

The  $N = 2$  SUSY structure associated to  $G_v^\pm$  is presented in the diagram below:

$$\begin{array}{ccc} & \sqrt{-1}b & \xrightarrow{D_v^1} \sqrt{-1}\partial\alpha \\ D_v^2 \nearrow & & \searrow D_v^2 \\ \alpha & \xrightarrow{-D_v^1} -b & \\ & -\sqrt{-1}\beta & \xrightarrow{D_v^1} \sqrt{-1}\partial a \\ D_v^2 \nearrow & & \searrow D_v^2 \\ a & \xrightarrow{-D_v^1} \beta & \end{array} \quad \begin{array}{ccc} & -\sqrt{-1}\delta & \xrightarrow{D_v^1} \sqrt{-1}\partial c \\ D_v^2 \nearrow & & \searrow D_v^2 \\ c & \xrightarrow{-D_v^1} \delta & \\ & \sqrt{-1}d & \xrightarrow{D_v^1} \sqrt{-1}\partial\gamma \\ D_v^2 \nearrow & & \searrow D_v^2 \\ \gamma & \xrightarrow{-D_v^1} -d & \end{array}$$

and the non-zero  $\Lambda$ -brackets of generators are

$$[\alpha \wedge c] = -\sqrt{-1}\chi^1 + \chi^2, \quad [a \wedge \gamma] = \sqrt{-1}\chi^1 - \chi^2.$$

Hence the extended  $bc$ - $\beta\gamma$  system with  $D_v^1$  and  $D_v^2$  is the product of the  $bc$ - $\alpha\delta$  system and the  $ad$ - $\beta\gamma$  system, each having its own internal  $N = 2$  SUSY structure.

We remark that each of the odd derivations  $D^1, D^2$  in Example 4.5 can be realized as an odd derivation derived from the independent superconformal structures, with  $D^1 = -\sqrt{-1}D_h^1 = -D_h^2$  for  $D_h^1$  and  $D_h^2$  in Example 4.6, and  $D^2 = D_v^1 = -\sqrt{-1}D_v^2$  for  $D_v^1$  and  $D_v^2$  in Example 4.7. Furthermore, there is an  $N = 2$  superconformal structure of the extended  $bc$ - $\beta\gamma$  system that simultaneously induces  $D^1$  and  $D^2$  (see Section 6.4 of [9]). Precisely, the  $N = 2$  superconformal structure induced from the following vectors

$$\begin{aligned} L &= L_h, & G^+ &= -\frac{\sqrt{-1}}{2}(G_h^+ - G_h^- + G_v^+ + G_v^-), \\ G^- &= -\frac{\sqrt{-1}}{2}(G_h^+ - G_h^- - G_v^+ - G_v^-), & J &= \sqrt{-1}(ac + bd), \end{aligned}$$

give rise to the  $N = 2$  SUSY vertex algebra structure of the extended  $bc$ - $\beta\gamma$  system defined in Example 4.5.

**Remark 4.8.** Note that, there is a 1-parameter deformation of the superconformal vector of the  $bc$ - $\beta\gamma$  system (see, for example [5, 11]). For an  $N = 1$  superconformal vertex algebra  $V$  with a superconformal vector  $G$  with central charge  $c$ , in general, we can shift the superconformal vector  $G$  to another superconformal vector, giving rise to various conformal weights. More precisely, suppose that  $v \in V$  is an odd vector satisfying the following relations:

$$[L \lambda v] = (\partial + \frac{\lambda}{2})v, \quad [G \lambda v] = G_{(0)}v, \quad [v \lambda v] = c',$$

for some  $c' \in \mathbb{C}$  and  $G_{(0)}v$  is diagonalizable on  $V$ , where  $L$  is the conformal vector. Then, the vector  $G + \partial v$  is also a superconformal vector and the corresponding shifted conformal vector is  $L + \frac{1}{2}\partial G_{(0)}v$  with central charge  $c - 3c'$ . At the level of SUSY vertex algebra structures, the shifting of superconformal vectors does not change the  $N = 1$  SUSY structures induced from the superconformal vectors.

In the context of conformal vertex algebras, in Section 5 of [1], the effect of the shifted conformal vectors to the Zhu's algebra and Frenkel-Zhu's bimodules was studied. Furthermore, there is an application of the shifted conformal vectors to the construction of the conformal structures of  $W$ -algebras (see Section 7 of [1] and Section 5.1 of [8] for more details).

For the superconformal cases, by combining the shifted superconformal vector of the  $bc$ - $\beta\gamma$  system with the shifted Kac-Todorov superconformal vector (see Example 2.13 of [15]), we can obtain the realization of the  $N = 1$  supersymmetry of SUSY  $W$ -algebras via superconformal structures, as in Remark 4.10 of [24].

Now, returning to the Question 4.2, let us construct an  $N = 2$  SUSY extension of a universal enveloping  $N = 1$  SUSY vertex algebra  $V(R_{N=1})$ . For the purpose of finding  $N = 2$  SUSY extensions, we use the construction in Theorem 3.6 and give an additional  $N = 1$  supersymmetry in an independent direction from the given  $N = 1$  supersymmetry of  $V(R_{N=1})$ .

**Theorem 4.9.** *Let  $R_{N=1} = \mathbb{C}[D] \otimes \text{span}_{\mathbb{C}}\{\bar{u}_i\}_{i \in I} \oplus E_C$  be an  $N = 1$  SUSY Lie conformal algebra, where the set  $\{D(\bar{u}_i), \bar{u}_i\}_{i \in I}$  is linearly independent and  $E_C$  is the central extension part of  $R_{N=1}$  with  $DE_C = 0$ . Then  $V(R_{N=1})$  has an  $N = 2$  SUSY extension.*

*Proof.* If we set  $u_i := D\bar{u}_i$ , then we can consider the  $N = 1$  SUSY Lie conformal algebra  $R_{N=1}$  as a non-SUSY Lie conformal algebra  $R_{N=1} = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{u_i, \bar{u}_i\}_{i \in I} \oplus E_C$  with a SUSY generator  $\{\bar{u}_i\}_{i \in I}$  in the  $D$ -direction.

By using the construction in Theorem 3.6, we have a SUSY generator  $\{u_i^\circ, \bar{u}_i^\circ\}_{i \in I}$  in the  $D^\circ$ -direction of the  $\mathbb{C}[\partial]$ -module  $\tilde{R}_{N=1} := R_{N=1} \oplus R_{N=1}^\circ$ , where  $R_{N=1}^\circ = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{u_i^\circ, \bar{u}_i^\circ\}_{i \in I} \oplus E_C^\circ$ . The remaining

non-zero  $\lambda$ -brackets on  $\widetilde{R}_{N=1}$  are given by the following relations

$$\begin{aligned} [u_i^\circ \lambda u_j] &= [u_i \lambda u_j]^\circ, & [u_i \lambda u_j^\circ] &= (-1)^{p(u_i)} [u_i \lambda u_j]^\circ, \\ [u_i^\circ \lambda \bar{u}_j] &= [u_i \lambda \bar{u}_j]^\circ, & [u_i \lambda \bar{u}_j^\circ] &= (-1)^{p(u_i)} [u_i \lambda \bar{u}_j]^\circ, \\ [\bar{u}_i^\circ \lambda u_j] &= [\bar{u}_i \lambda u_j]^\circ, & [\bar{u}_i \lambda u_j^\circ] &= (-1)^{p(u_i)+1} [\bar{u}_i \lambda u_j]^\circ, \\ [\bar{u}_i^\circ \lambda \bar{u}_j] &= [\bar{u}_i \lambda \bar{u}_j]^\circ, & [\bar{u}_i \lambda \bar{u}_j^\circ] &= (-1)^{p(u_i)+1} [\bar{u}_i \lambda \bar{u}_j]^\circ. \end{aligned}$$

Now, to give an  $N = 2$  SUSY structure on  $\widetilde{R}_{N=1}$ , let us define odd derivations  $D^1$  and  $D^2$  as follows:

$$D^1(u_i) := D(u_i), \quad D^1(\bar{u}_i) := D(\bar{u}_i), \quad D^1(u_i^\circ) := (-D(u_i))^\circ, \quad D^1(\bar{u}_i^\circ) := (-D(\bar{u}_i))^\circ, \quad D^2 := D^\circ.$$

Then it can be verified that the pair  $(\{\bar{u}_i, \bar{u}_i^\circ\}_{i \in I}, D^1)$  also satisfies the SUSY extension formulas, especially the non-trivial parts can be verified through the following calculations:

$$\begin{aligned} [u_i \lambda u_j^\circ] &= (-1)^{p(u_i)} [u_i \lambda u_j]^\circ = (D[u_i \lambda \bar{u}_j] + \lambda[\bar{u}_i \lambda \bar{u}_j])^\circ \\ &= (D[u_i \lambda \bar{u}_j])^\circ + \lambda[\bar{u}_i \lambda \bar{u}_j]^\circ = -D^1([u_i \lambda \bar{u}_j]^\circ) + \lambda[\bar{u}_i \lambda \bar{u}_j]^\circ \\ &= (-1)^{p(u_i)+1} (D^1[u_i \lambda \bar{u}_j] + \lambda[\bar{u}_i \lambda \bar{u}_j^\circ]), \\ [\bar{u}_i \lambda u_j^\circ] &= (-1)^{p(u_i)+1} [\bar{u}_i \lambda u_j]^\circ = -([u_i \lambda \bar{u}_j] - D[\bar{u}_i \lambda \bar{u}_j])^\circ \\ &= -[u_i \lambda \bar{u}_j]^\circ + (D[\bar{u}_i \lambda \bar{u}_j])^\circ = -[u_i \lambda \bar{u}_j]^\circ - D^1([\bar{u}_i \lambda \bar{u}_j]^\circ) \\ &= (-1)^{p(u_i)+1} ([u_i \lambda \bar{u}_j^\circ] - D^1[\bar{u}_i \lambda \bar{u}_j^\circ]), \end{aligned}$$

which follow from the fact that  $\{\bar{u}_i\}_{i \in I}$  is a SUSY generator in the  $D$ -direction. Therefore, by Proposition 4.3,  $V(\widetilde{R}_{N=1}) = V(R_{N=1}) \otimes V(R_{N=1}^\circ)$  is an  $N = 2$  SUSY vertex algebra.  $\square$

Combining the results in Section 3 and Theorem 4.9, we conclude that any non-SUSY universal enveloping vertex algebra can be embedded into an  $N = 2$  SUSY vertex algebra.

**Corollary 4.10.** *Let  $R = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{u_i\}_{i \in I} \oplus E_C$  be a Lie conformal algebra in (3.3). Then  $V(R)$  has an  $N = 2$  SUSY extension.*

*Proof.* It follows from Theorem 3.6 and 4.9.  $\square$

Recall that the  $N = 2$  super-Virasoro vertex algebra is an  $N = 2$  SUSY extension of the  $N = 1$  super-Virasoro vertex algebra, which is an  $N = 1$  SUSY extension of the Virasoro vertex algebra. In the following examples, by applying Theorem 4.9 (resp. Corollary 4.10) to the  $N = 1$  super-Virasoro (resp. Virasoro) vertex algebra, we obtain  $N = 2$  SUSY extensions of the Virasoro vertex algebra, which are distinct from the  $N = 2$  super-Virasoro vertex algebra.

**Example 4.11** ( $N = 2$  SUSY extension of Virasoro and  $N = 1$  super-Virasoro vertex algebra). In this example, we consider the centerless Virasoro and  $N = 1$  super-Virasoro Lie conformal algebras (see Example 3.8). Recall that the  $N = 1$  super-Virasoro Lie conformal algebra  $\text{SVir} = \mathbb{C}[\partial] \otimes (\mathbb{C}L \oplus \mathbb{C}G)$  admits an  $N = 1$  supersymmetry defined by  $DG = 2L$ . If we apply the construction of Theorem 4.9 to the  $N = 1$  super-Virasoro vertex algebra, we obtain the  $N = 2$  SUSY vertex algebra  $V(\widetilde{\text{SVir}})$ , where  $\widetilde{\text{SVir}} = \mathbb{C}[\partial] \otimes (\mathbb{C}L \oplus \mathbb{C}G \oplus \mathbb{C}L^\circ \oplus \mathbb{C}G^\circ)$  is the Lie conformal algebra endowed with the following  $\lambda$ -brackets:

$$\begin{aligned} [L \lambda L] &= (\partial + 2\lambda)L, & [L \lambda L^\circ] &= (\partial + 2\lambda)L^\circ, & [L \lambda G] &= (\partial + \frac{3}{2}\lambda)G, & [L \lambda G^\circ] &= (\partial + \frac{3}{2}\lambda)G^\circ, \\ [G \lambda L^\circ] &= -(\frac{1}{2}\partial + \frac{3}{2}\lambda)G^\circ, & [G \lambda G] &= 2L, & [G \lambda G^\circ] &= -2L^\circ, & [L^\circ \lambda L^\circ] &= [L^\circ \lambda G^\circ] = [G^\circ \lambda G^\circ] = 0. \end{aligned}$$

In other words, as an  $N = 2$  SUSY vertex algebra,  $V(\widetilde{\text{SVir}})$  is generated by one element  $G^\circ$  and the  $N = 2$   $\Lambda$ -bracket is given by

$$[G^\circ \ \Lambda \ G^\circ] = -(2\partial + 3\lambda)G^\circ + 2\chi^1 L^\circ.$$

**Example 4.12** (Another  $N = 2$  SUSY extension of Virasoro vertex algebra). If we apply Corollary 4.10, i.e. apply Theorem 3.6 and Theorem 4.9 subsequently, to the Virasoro vertex algebra, we have an  $N = 2$  SUSY extension of the Virasoro vertex algebra which is generated by one element  $\bar{L}^\circ$  endowed with the following  $N = 2$   $\Lambda$ -bracket:

$$[\bar{L}^\circ \ \Lambda \ \bar{L}^\circ] = -(\partial + 2\lambda)\bar{L}^\circ.$$

Note that this extended vertex algebra is distinct from the extended vertex algebra  $V(\widetilde{\text{SVir}})$  described in Example 4.11.

By Corollary 4.10, the non-zero  $\lambda$ -brackets of the  $N = 2$  SUSY extension of the affine vertex algebra of level 0 are given by

$$(4.4) \quad \begin{aligned} [a \ \lambda \ b] &= [a, b], \quad [\bar{a} \ \lambda \ b] = (-1)^{p(a)}[a \ \lambda \ \bar{b}] = \overline{[a, b]}, \quad [a^\circ \ \lambda \ b] = (-1)^{p(a)}[a \ \lambda \ b^\circ] = [a, b]^\circ, \\ [\bar{a}^\circ \ \lambda \ b] &= [a \ \lambda \ \bar{b}^\circ] = (-1)^{p(a)}[a^\circ \ \lambda \ \bar{b}] = (-1)^{p(a)+1}[\bar{a} \ \lambda \ b^\circ] = \overline{[a, b]^\circ}, \end{aligned}$$

and this vertex algebra is isomorphic to  $V_{N=2}^0(\mathfrak{g})$  in Example 2.10. As an  $N = 2$  analogue of the superconformal current algebra in Remark 3.9, the following proposition shows that a natural  $N = 2$  superconformal extension of  $V_{N=2}^0(\mathfrak{g})$  can be obtained by considering it as an  $N = 2$  SUSY vertex subalgebra of an  $N = 2$  superconformal vertex algebra.

**Proposition 4.13.** *Let  $\mathfrak{g} = \bigoplus_{\omega \in \Omega} \mathfrak{g}_\omega$  be an  $\Omega$ -graded Lie superalgebra. Then the  $N = 2$  SUSY affine vertex algebra  $V_{N=2}^0(\mathfrak{g})$  can be embedded into an  $N = 2$  superconformal vertex algebra, where the conformal weight of  $a \in \mathfrak{g}_\omega$  is given by  $\Delta_a = 1 - \omega$ .*

*Proof.* Let  $\text{Curg}_{N=2} = \mathbb{C}[\partial] \otimes (\mathfrak{g} \oplus \bar{\mathfrak{g}} \oplus \mathfrak{g}^\circ \oplus \bar{\mathfrak{g}}^\circ)$  be the Lie conformal algebra with the  $\lambda$ -brackets in (4.4) and consider the following  $N = 2$  super-Virasoro Lie conformal algebra

$$\text{SVir}_{N=2} = \mathbb{C}[\partial] \otimes (\mathbb{C}L \oplus \mathbb{C}G^+ \oplus \mathbb{C}G^- \oplus \mathbb{C}J) \oplus \mathbb{C}C$$

with the  $N = 2$  SUSY structure in Example 4.4, where  $C$  is central and  $\partial C = 0$ . First of all, define the following  $\lambda$ -brackets

$$(4.5) \quad \begin{aligned} [L \ \lambda \ a] &= (\partial + \Delta_a \lambda)a, \quad [L \ \lambda \ \bar{a}] = (\partial + (\Delta_a - \frac{1}{2})\lambda)\bar{a}, \\ [L \ \lambda \ a^\circ] &= (\partial + (\Delta_a - \frac{1}{2})\lambda)a^\circ, \quad [L \ \lambda \ \bar{a}^\circ] = (\partial + (\Delta_a - 1)\lambda)\bar{a}^\circ, \end{aligned}$$

where  $\Delta_a = 1 - \omega$ , for  $a \in \mathfrak{g}_\omega$ . Then, by solving the SUSY extension formulas with the constraints (4.5), we can define the remaining  $\lambda$ -brackets as follows

$$\begin{aligned} [G^+ + G^- \ \lambda \ a] &= (\partial + (2\Delta_a - 1)\lambda)\bar{a}, \quad [G^+ + G^- \ \lambda \ \bar{a}] = a, \\ [G^+ + G^- \ \lambda \ a^\circ] &= -(\partial + (2\Delta_a - 2)\lambda)\bar{a}^\circ, \quad [G^+ + G^- \ \lambda \ \bar{a}^\circ] = -a^\circ, \\ [G^+ - G^- \ \lambda \ a] &= \sqrt{-1}(\partial + (2\Delta_a - 1)\lambda)a^\circ, \quad [G^+ - G^- \ \lambda \ a^\circ] = \sqrt{-1}\bar{a}, \\ [G^+ - G^- \ \lambda \ \bar{a}] &= \sqrt{-1}(\partial + (2\Delta_a - 2)\lambda)\bar{a}^\circ, \quad [G^+ - G^- \ \lambda \ \bar{a}^\circ] = \sqrt{-1}\bar{a}, \\ [J \ \lambda \ a] &= \sqrt{-1}(2\Delta_a - 2)\lambda\bar{a}^\circ, \quad [J \ \lambda \ \bar{a}] = \sqrt{-1}a^\circ, \quad [J \ \lambda \ a^\circ] = -\sqrt{-1}\bar{a}, \quad [J \ \lambda \ \bar{a}^\circ] = 0. \end{aligned}$$

The Jacobi identity of the  $\mathbb{C}[\partial]$ -module  $\text{SVir}_{N=2} \oplus \text{Curg}_{N=2}$  follows from direct computations. Hence the product of the  $N = 2$  super-Virasoro vertex algebra and the  $N = 2$  SUSY affine vertex algebra of level 0

$$V(\text{SVir}_{N=2} \oplus \text{Curg}_{N=2}) / \langle C - c \rangle \simeq \text{SVir}_{N=2}^c \otimes V_{N=2}^0(\mathfrak{g})$$

is an  $N = 2$  superconformal vertex algebra. Moreover, the two odd derivations

$$D^1 = (G^+ + G^-)_{(0)}, \quad D^2 = -\sqrt{-1}(G^+ - G^-)_{(0)},$$

induce the  $N = 2$  SUSY vertex algebra structures of both  $\text{SVir}_{N=2}^c$  and  $V_{N=2}^0(\mathfrak{g})$ .  $\square$

Now, we generalize the construction of Theorem 4.9 for any  $N = n$  SUSY cases. For example, in Remark 4.14, we present the construction of  $N = 3$  SUSY extensions of a given universal enveloping  $N = 2$  SUSY vertex algebra. The proof for the arbitrary  $N = n$  case is identical to the proofs for  $N = 2$  (Theorem 4.9) and  $N = 3$  (Remark 4.14) cases in essence.

**Remark 4.14.** For a given  $N = 2$  SUSY vertex algebra  $V(R_{N=2})$ , where the  $N = 2$  SUSY Lie conformal algebra is given by  $R_{N=2} = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{u_i, \bar{u}_i, u_i^\circ, \bar{u}_i^\circ\}_{i \in I}$  with the  $N = 2$  supersymmetry  $D^1$  and  $D^2$ , we can also construct an  $N = 3$  SUSY extension of  $V(R_{N=2})$  as in the proof of Theorem 4.9. More precisely, we construct an  $N = 1$  supersymmetry on the  $\mathbb{C}[\partial]$ -module  $\tilde{R}_{N=2} := R_{N=2} \oplus R_{N=2}^\dagger$  in the  $D^\dagger$ -direction, where the  $\mathbb{C}[\partial]$ -module  $R_{N=2}^\dagger = \mathbb{C}[\partial] \otimes \text{span}_{\mathbb{C}}\{u_i^\dagger, \bar{u}_i^\dagger, u_i^{\circ,\dagger}, \bar{u}_i^{\circ,\dagger}\}_{i \in I}$  is given by Theorem 3.6. If we define the three odd endomorphisms  $D^1$ ,  $D^2$  and  $D^3 = D^\dagger$  as shown in the diagram below:

$$\begin{array}{ccccc}
 & & \bar{u}_i & \xrightarrow{D^1} & u_i \\
 & \nearrow^{D^2} & \uparrow & & \nearrow^{D^2} \\
 \bar{u}_i^\circ & \xrightarrow{D^3} & u_i^\circ & & \uparrow \\
 & \searrow^{D^3} & \downarrow & & \searrow^{D^3} \\
 & & \bar{u}_i^\dagger & \xrightarrow{D^3} & u_i^\dagger \\
 & \nearrow^{-D^2} & \uparrow & & \nearrow^{-D^2} \\
 \bar{u}_i^{\circ,\dagger} & \xrightarrow{D^1} & u_i^{\circ,\dagger} & & \uparrow
 \end{array}$$

then each of the following pairs satisfies the SUSY extension formulas:

$$(\{\bar{u}_i, \bar{u}_i^\circ, \bar{u}_i^\dagger, \bar{u}_i^{\circ,\dagger}\}_{i \in I}, D^1), \quad (\{u_i^\circ, \bar{u}_i^\circ, u_i^{\circ,\dagger}, \bar{u}_i^{\circ,\dagger}\}_{i \in I}, D^2), \quad (\{u_i^\dagger, \bar{u}_i^\dagger, u_i^{\circ,\dagger}, \bar{u}_i^{\circ,\dagger}\}_{i \in I}, D^3).$$

Indeed, the non-trivial part of the proof is the verification of the SUSY extension formulas for the pairs:

$$(\{\bar{u}_i, \bar{u}_i^{\circ,\dagger}\}_{i \in I}, D^1), \quad (\{\bar{u}_i^\dagger, \bar{u}_i^\circ\}_{i \in I}, D^1), \quad (\{u_i^\circ, \bar{u}_i^{\circ,\dagger}\}_{i \in I}, D^2), \quad (\{u_i^{\circ,\dagger}, \bar{u}_i^\circ\}_{i \in I}, D^2),$$

which follows from the fact that  $(\{\bar{u}_i, \bar{u}_i^\circ\}_{i \in I}, D^1)$  and  $(\{u_i^\circ, \bar{u}_i^\circ\}_{i \in I}, D^2)$  satisfy the SUSY extension formulas. For example, we have the (SEF1) of the pair  $(\{\bar{u}_i\}_{i \in I}, \{\bar{u}_i^{\circ,\dagger}\}_{i \in I})$  in the  $D^1$ -direction from the following computations:

$$\begin{aligned}
 [D^1 \bar{u}_i \lambda D^1 \bar{u}_j^{\circ,\dagger}] &= [u_i \lambda u_j^{\circ,\dagger}] = (-1)^{p(u_i)} [u_i \lambda u_j^{\circ,\dagger}]^\dagger = -(D^1 [u_i \lambda \bar{u}_j^\circ])^\dagger - \lambda [\bar{u}_i \lambda \bar{u}_j^{\circ,\dagger}]^\dagger \\
 &= D^1 ([u_i \lambda \bar{u}_j^\circ]^\dagger) - \lambda [\bar{u}_i \lambda \bar{u}_j^{\circ,\dagger}]^\dagger = (-1)^{p(u_i)} (D^1 [u_i \lambda \bar{u}_j^\circ]^\dagger + \lambda [\bar{u}_i \lambda \bar{u}_j^{\circ,\dagger}]^\dagger).
 \end{aligned}$$

Before the closing of this paper, we suggest additional systematic methods to find SUSY structures of vertex algebras (see Remark 4.15, 4.16 and 4.17). These methods can be used for a vertex algebra which is not a universal enveloping vertex algebra.

**Remark 4.15.** Let  $V$  be a vertex algebra with an  $N = 1$  superconformal vector  $G$ . Suppose there is a decomposition  $G = G^+ + G^-$  satisfying  $(G^\pm_{(0)}G^\pm)_{(0)} = 0$ . Then  $V$  has an  $N = 2$  SUSY vertex algebra structure. Indeed, as in the case of  $N = 2$  superconformal vertex algebra, define the odd derivations as

$$D^1 = (\mu G^+ + \mu^{-1}G^-)_{(0)}, \quad D^2 = \sqrt{-1}(\mu G^+ - \mu^{-1}G^-)_{(0)},$$

for  $\mu \in \mathbb{C}^*$ . Then, by the Jacobi identity, we can check  $[D^i, D^j] = 2\delta_{ij}\partial$  for  $i, j = 1, 2$ . Also, it follows from Proposition 1.12 of [8] that  $D^1$  and  $D^2$  are odd derivations of the normally ordered product and the  $\lambda$ -brackets of  $V$ .

**Remark 4.16.** A vertex algebra is called an  $N = 3$  *superconformal vertex algebra* if it has a conformal vector  $L$  with central charge  $c$ , three even primary vectors  $J^+, J^-, J^0$  of conformal weight 1, three odd primary vectors  $G^+, G^-, G^0$  of conformal weight  $\frac{3}{2}$ , and an odd primary vector  $\Phi$  of conformal weight  $\frac{1}{2}$ , satisfying the  $N = 3$  super-Virasoro relation (2.5). For an  $N = 3$  superconformal vertex algebra, there is an  $N = 3$  SUSY vertex algebra structure with the following odd derivations

$$D^1 = (\mu G^+ + \mu^{-1}G^-)_{(0)}, \quad D^2 = \sqrt{-1}(\mu G^+ - \mu^{-1}G^-)_{(0)}, \quad D^3 = \sqrt{2}G^0_{(0)}.$$

For an  $N = 4$  superconformal vertex algebra, if we adapt the definition and notation in Section 8.4 of [21], the following odd derivations

$$\begin{aligned} D^1 &= (\mu G^+ + \mu^{-1}\bar{G}^-)_{(0)}, & D^2 &= \sqrt{-1}(\mu G^+ - \mu^{-1}\bar{G}^-)_{(0)}, \\ D^3 &= (\nu G^- + \nu^{-1}\bar{G}^+)_{(0)}, & D^4 &= \sqrt{-1}(\nu G^- - \nu^{-1}\bar{G}^+)_{(0)}, \end{aligned}$$

give rise to an  $N = 4$  SUSY vertex algebra structure, for  $\mu, \nu \in \mathbb{C}^*$ .

**Remark 4.17.** For any vertex algebra  $V$ , it can be embedded into SUSY vertex algebras by considering the universal enveloping vertex algebra of the vertex algebra  $V$ . Therefore, we know that there is an  $N = n'$  SUSY vertex algebra containing a given  $N = n$  SUSY vertex algebra, for any  $n' \geq n$ .



## REFERENCES

- [1] T. Arakawa, *Rationality of  $W$ -algebras: Principal Nilpotent Cases*, Ann. Math. 182, 565–604 (2015)
- [2] K. Barron, *Axiomatic Aspects of  $N = 2$  Vertex Superalgebras with Odd Formal Variables*, Commun. Alg. 38, 1199–1268 (2010)
- [3] R. Borcherds, *Vertex Algebras, Kac–Moody Algebras, and the Monster*, Proc. Nat. Acad. Sci. USA 83, 10, 3068–3071 (1986)
- [4] A. A. Belavin, A. M. Polyakov, A. B. Zamolodchikov, *Infinite Conformal Symmetry in Two-Dimensional Quantum Field Theory*, Nucl. Phys. B 241, 333–380 (1984)
- [5] S. -J. Cheng, *Construction of  $N = 2$  Superconformal Algebra from Affine Algebras with Extended Symmetry: I*, Lett. Math. Phys. 33, 23–31 (1995)
- [6] R. Coquereaux, L. Frappat, E. Ragoucy, P. Sorba, *Extended Super–Kac–Moody Algebras and Their Super–Derivation Algebras*, Commun. Math. Phys. 133, 1–35 (1990)
- [7] A. De Sole, V. G. Kac, *V. Freely Generated Vertex Algebras and Non-Linear Lie Conformal Algebras*, Commun. Math. Phys. 254, 659–694 (2005)
- [8] A. De Sole, V. G. Kac, *Finite vs Affine  $W$ -Algebras*, Jpn. J. Math. 1, 137–261 (2006)
- [9] J. Ekstrand, R. Heluani, J. Källén, M. Zabzine, *Non-Linear Sigma Models via the Chiral de Rham Complex*, Adv. Theor. Math. Phys. 13, 1221–1254 (2009)
- [10] J. Ekstrand, R. Heluani, M. Zabzine, *Sheaves of  $N = 2$  Supersymmetric Vertex Algebras on Poisson Manifolds*, J. Geom. Phys. 62, 11, 2259–2278 (2012)
- [11] P. Frè, L. Girardello, A. Lerda, P. Soriani, *Topological First-Order Systems with Landau–Ginzburg Interactions*, Nucl. Phys. B 387, 333–372 (1992)
- [12] R. Heluani, *Supersymmetry of the Chiral de Rham Complex 2: Commuting Sectors*, Int. Math. Res. Not. 6, 953–987 (2009)
- [13] R. Heluani, V. G. Kac, *Supersymmetric Vertex Algebras*, Commun. Math. Phys. 271, 103–178 (2007)
- [14] R. Heluani, V. G. Kac, *SUSY Lattice Vertex Algebras*, Bulg. J. Phys. 35, S1, 3–24 (2008)
- [15] R. Heluani, M. Zabzine, *Generalized Calabi–Yau Manifolds and the Chiral de Rham Complex*, Adv. Math. 223, 1815–1844 (2010)
- [16] R. Heluani, M. Zabzine, *Superconformal Structures on Generalized Calabi–Yau Metric Manifolds*, Commun. Math. Phys. 306, 333–364 (2011)
- [17] J. M. Isidro, A. V. Ramallo, *Superconformal Current Algebras and Topological Field Theories*, Phys. Lett. B 340, 48–56 (1994)
- [18] T. Johnson–Freyd, *Supersymmetry and the Suzuki Chain*, Tunis. J. Math. 3, 309–359 (2021)
- [19] V. G. Kac, *Vertex Algebras for Beginners*, University Lecture Series, AMS, Vol. 10, 1996. Second edition, 1998.
- [20] V. G. Kac, I. T. Todorov, *Superconformal Current Algebras and Their Unitary Representations*, Commun. Math. Phys. 102, 337–347 (1985)
- [21] V. G. Kac, M. Wakimoto, *Quantum Reduction and Representation Theory of Superconformal Algebras*, Adv. Math. 185, 400–458 (2004)
- [22] A. Molev, E. Ragoucy, U. R. Suh, *Supersymmetric  $W$ -Algebras*, Lett. Math. Phys. 111, 6 (2021)
- [23] A. Schwimmer, N. Seiberg, *Comments on the  $N = 2, 3, 4$  Superconformal Algebras in Two Dimensions*, Phys. Lett. B 184, 191–196 (1987)
- [24] S. Yoon, *Superconformal Structures of Supersymmetric Free Fermion Vertex Algebras*, MSc Thesis, Seoul National University, Department of Mathematical Sciences, February 2023

DEPARTMENT OF MATHEMATICAL SCIENCES AND RESEARCH INSTITUTE OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY,  
 GWANAK-RO 1, GWANAK-GU, SEOUL 08826, KOREA

*Email address:* uhrisu1@snu.ac.kr

DEPARTMENT OF MATHEMATICAL SCIENCES, SEOUL NATIONAL UNIVERSITY, GWANAK-RO 1, GWANAK-GU, SEOUL 08826,  
 KOREA

*Email address:* ysw317@snu.ac.kr