Sourceless Maxwell and Dirac equations via Clifford Analysis

Calum Robson c.j.robson@lse.ac.uk CPNSS, London School of Economics

August 4, 2023

Abstract

The study of complex functions is based around the study of holomorphic functions, satisfying the Cauchy-Riemann equations. The complex numbers are a subset [the even subalgebra] of Cl(2), and therefore we can ask whether there are analogues for the Cauchy-Riemann equations for other Clifford algebras. This has been extensively explored under the name of Clifford Analysis. Here I explicitly decompose the Cauchy-Riemann equations for a general Clifford algebra into grades using the Geometric Algebra formalism,, and show that for the Spacetime Algebra Cl(3, 1) these equations are the equations for a self-dual source free Electromagnetic field, and for a massless uncharged Spinor.

1 Introduction and Background

This paper is about the application of Clifford (or Geometric) algebras to physics. In particular, it is about Clifford Analysis, which is an attempt to extend the results of Complex Analysis first to the Quaterions, and then to a general Clifford algebra. The Clifford approach to geometry is based upon the ideas of W.K.Clifford and H.G. Grassmann [25], and involves giving a direct geometric representation of the n- dimensional subspaces of a Clifford algebra in terms of the n-dimensional linear subspaces of a geometric space, for example lines, planes, volumes; and so on.

Dieudonne clarifies the technical distincion between Clifford geometry and the more standard approach as follows [9]. In standard approaches, we consider both a vector space \mathcal{V} , and its dual \mathcal{V}^* . We cannot identify these two spaces, but we can associate elements in \mathcal{V} to elements in \mathcal{V}^* if we define an inner product between these spaces. Then a *p*-dimensional subspace $V_p \in \mathcal{V}$ will be dual to an (n-p)-dimensional subspace $W_{n-p} \in \mathcal{V}^*$ iff (v, w) = 0 for all $v \in V_p$ and $w \in W_{n-p}$, where *n* is the total dimension of \mathcal{V} and \mathcal{V}^* , and (,) denotes the inner product.

Conversely, in the Clifford approach to Geometry, we only have one space, \mathcal{V} , which is equipped with a product to turn it into an algebra; in exact analogy to

the transition between \mathbb{R}^2 and \mathbb{C} . More details are given below, but the essential point is that the dual of a *p*-dimensional subspace V_p is given by its (n-p) dimensional complement in \mathcal{V} itself. This has advantages for physics; first of all by cutting down on the mathematical entities required, and secondly because most objects in physics can be given a direct geometric interpretation in this construction (See [10] for examples).

This fits well with approaches to theoretical physics which seeks to describe fundamental processes as an algebra. As explicitly described by Bohm [6], since the elements of the algebra act on each other, they can model both the physical objects in the theory and the interactions between them which make up the process. Common examples of theories in this approach are Twistor theories [2], Octonion models [16][12] and related strategies, e.g. [5]. The paradigmatic example is the complex plane, where an element $Re^{i\theta}$ both represents a point, and the action of rotating anticlockwise by θ , and dilating by scale factor R.

I begin the paper with some definitions and notation. First, I will review the Geometric Algebra formalism for Clifford algebras. Then, I will discuss Clifford Analysis as a generalisation of Complex analysis. Finally, I will give some results from Hodge Theory which will be needed for this paper.

1.1 Geometric Algebras

First, we need to give an overview of the Geometric Algebra formalism for Clifford algebra calculations. This section is mainly based on [1] and [10]. Other excellent references are [7] and [11].

An (orthogonal) Clifford Algebra Cl(p,q) with a non- degenerate metric is described by d = p + q generators e_i , satisfying

$$e_i e_j - e_j e_i = \eta_{ij} \tag{1}$$

Where η_{ii} is 1 for *i* from 1 to p + 1, $\eta_{ii} = -1$ for *i* from p + 1 to *d*, and $\eta_{ij} = 0$ otherwise.

Clifford algebras are a graded algebra, with the grading given by the number of generators which are multipled together. For example, e_i is grade 1, $e_i e_j e_k$ is grade 3, and so on.

A Clifford algebra element which is contains only objects of the same grade is called a blade, and is denoted by $\langle E \rangle_i$, where E is the label of the element, and i is the grade of the blade. We can also use this notation to describe taking the component of a general Clifford algebra element which has grade i.

A Clifford algebra can be split into even and odd parts, corresponding to collecting the blades with even and odd grade. The even part is called the Spin algebra, denoted Spin(p,q). [24]. The Clifford algebra for a space are also isomorphic as vector spaces to the Differential Forms on the same space. This is what allows us to use techniques from the cohomology theory of forms in the Clifford Algebra context [29].

In the Geometric Algebra (GA) representation of Clifford Algebras, we take each grade to describe an k- dimensional subspace of \mathbb{R}^d . The highest grade object is denoted I, and describes a d-volume. So in 3 dimensions, the scalar describes a point, e_i describes a line, $e_i e_j$ describes a plane, and the pseudoscalar $I = e_1 e_2 e_3$ describes a 3-volume.

When we multiply two blades $\langle A \rangle_i$ and $\langle B \rangle_j$, the lowest grade object we can form has grade |i - j|, and is called the dot product, denoted by $A \cdot B$ or $\langle A \cdot B \rangle_{i-j}$. The highest grade element has grade i + j, and is called the wedge product, denoted $A \wedge B$ or $\langle A \wedge B \rangle_{i+j}$. Note that these may be identically zero (for example if i + j > d), and that in general one or both will be. For vectors a, b these definitions become

$$a \cdot b = ab - ba$$
$$a \wedge b = ab - ba \tag{2}$$

Here, $a \cdot b$ is the usual dot product, and in 3d we have

$$a \wedge b = (a \times b)I^{-1} \tag{3}$$

Where $a \times b$ is the cross product. This formula just means that $a \wedge b$ is the area between a and b, and $a \times b$ is the length orthogonal to that area. Note, however, that we can define $a \wedge b$ in any dimension, not just in 3d.

We call an element made up of different grades a Multivector. Then the most general function we can write from $Cl(p,q) \rightarrow Cl(p,q)$ is

$$\mathbf{z} = f_0 + f_i e_i + f_{ij} e_i e_j + \dots + f_{1\dots d} e_1 \dots e_d \tag{4}$$

Where the f are scalar functions. In this paper I shall only consider functions of \mathbf{x} , where $\mathbf{x} = x_i e_i$ is the usual position vector.

Duality relations exist between blades of grades k and d - k. These are defined in various ways in the literature, which differ up to a sign. I shall use the convention

$$A^{\star} = AI^{-1} \tag{5}$$

Where A^* is the dual of A. If A has grade k, then A^* has grade d - k. The following property of the dual will be important later

$$A \wedge BI^{-1} = (A \wedge B)I^{-1}$$
$$A \cdot BI^{-1} = (A \cdot B)I^{-1}$$
(6)

Next, we can define a vector derivative

$$\partial = \frac{\partial}{\partial \mathbf{x}} \equiv e_i \frac{\partial}{\partial x} \tag{7}$$

Once we have this, we can define the wave/laplacian operator as ∂^2 . Applying the vector derivative to a blade will either raise or lower the grades of the components of multivector by 1. In general it will do both. We therefore write

$$\partial \mathbf{z} = \partial \cdot \mathbf{z} + \partial \wedge \mathbf{z} \tag{8}$$

where $\partial \cdot \mathbf{z}$ and $\partial \wedge \mathbf{z}$ are the grade lowering and grade raising parts respectively.

1.2 Clifford Analysis

The introduction of the vector derivative brings us back to Clifford Analysis. Clifford Analysis is focused around finding and understanding solutions of the Dirac eqation $\partial \phi = 0$. Indeed, a key motivator for extending Complex Analysis to Clifford algebras is the fact that this equation is a higher dimensional analogue of the Cauchy- Reimann equations. This means that Clifford Analysis intersects with other areas of mathematics, for example Spin Geometry, in paricular index theory [24]

The motivation for a Clifford algebra version of complex analysis comes from the fact that we can use Clifford algebras to take the square root of the wave equation, as we saw below equation (7). Then we can call ∂ a Dirac operator, as it has exactly the same algebraic properties as the matrix operator ∂ from the standard Dirac theory. Note, however, that we can define different Dirac operators as the square roots of different second order PDEs.

Clifford Analysis has its beginnings in the work of Moisil, Théodoresco and Riesz [26] [28] in the 1930s, which successfully extended the key results of Complex Analysis to the quaternionic case. R. Fueter then extended this to higher dimensions via Clifford Algebras[14][15]. In the late 20th century, this was developed by figures like Hestenes, Sobcyck and Delenghe [20][8]with results like the Cauchy Intergral theorem and the Cauchy- Kowalseka extension being extended from the complex numbers to general Clifford Algebras.

In two dimensions, we have the Geometric Algebra Cl(2). This consists of a scalar, two vectors e_1 and e_2 , and a bivector pseudoscalar, $I = e_1e_2$. The even subalgebra, Spin(2), is spanned by $\{1, I\}$, with $I^2 = -1$. Therefore $Spin(2) \cong \mathbb{C}$. A generic Spin(2)-valued function is given by u(z) + Iv(z), where z = x + Iy. Equivalently, we can consider the variable $\tilde{z} = e_1 z = xe_1 + ye_2$. Then

$$\partial (u(\tilde{z})) + Iv(\tilde{z})) = e_1 \frac{\partial u}{\partial x} + e_2 \frac{\partial u}{\partial y} + e_1 I \frac{\partial v}{\partial x} + e_2 I \frac{\partial v}{\partial y}$$
$$= e_1 \left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) + e_2 \left(\frac{\partial u}{\partial y} + \frac{\partial v}{\partial x} \right)$$
(9)

Therefore the condition $\partial(u + Iv) = 0$ is equivalent to the pair of equations

$$u_x = v_y; \quad u_y = v_x \tag{10}$$

We can recognise these as the Cauchy-Riemann equations. The natural question is whether there is an analogue to these for a general Clifford Algebra. It turns out that the answer is Yes- see the lecture notes [22], the book [8] and the references cited within for more details. Solutions to the equation $\partial \mathbf{f} = 0$ are usually called Monogenic functions, though they are occasionally known as Clifford Holomorphic functions. or Regular functions in older literature. I shall refer to them as Monogenic functions in this paper.

1.3 Hodge Theory

We will need to use some elements of Hodge Theory in this paper. Whilst there is not the space here for a full discussion of that important and complicated branch of mathematics, I will review the main results we will be using. It is important to bear in mind that Hodge Theory is true only for compact (i.e. closed and bounded) Reimannian Manifolds. There are strong links between Hodge theory and Clifford analysis which are only beginning to be explored. See [29] for an excellent and very recent treatment. As explained in [29] we can use the fact that Clifford Algebras are isomorphic as Vector Spaces to the space of differential forms to use Hodge decomposition on the Clifford algebra itself. The main result of Hodge Theory is the Hodge Decomposition Theorem, which states that the space of grade-k differential forms Ω_k decomposes as

$$\Omega_k = Har_k + d\Omega_{k-1} + \delta\Omega_{k+1} \tag{11}$$

This then lifts to a decomposition of the whole space of differential forms. In this expression, Har_k is the space of harmonic forms, i.e. those satisfying $\partial^2 \omega = 0$. A consequence of this is that the condition for a form ϕ to be harmonic is that $\delta \phi = d\phi = 0$. The expressions $d\Omega_{k-1} + \delta\Omega_{k+1}$ refer to the spaces of exact and co-exact k-forms respectively. Here, d refers to the usual exterior derivative, and δ refers to the coderivative, which is the adjoint of the exterior derivative under the \mathcal{L}_2 norm

$$\langle \omega, \xi \rangle := \int \omega \wedge \star \xi \, dx^4 \tag{12}$$

Here, \star refers to the Hodge dual. It is defined by the relation

$$A \wedge \star A = |A|^2 I \tag{13}$$

This interchanges k and n-k forms, and in Geometric Algebra notation is given by

$$\star \omega = (\omega I^{-1})^{\dagger} = (-1)^{\frac{k(k-1)}{2}}$$
(14)

where the factor $(-1)^{\frac{k(k-1)}{2}}$ comes from the fact that

$$|A|^{2} = A^{\dagger} \cdot A = (-1)^{\frac{k(k-1)}{2}} A \cdot A$$
(15)

In terms of the Hodge dual, we have $\delta = \star d \star$, however, in geometric algebra notation, as discussed in [29], we have

$$d\omega \equiv \partial \wedge \omega \tag{16}$$
$$\delta \omega \equiv -\partial \cdot \omega$$

Where k is the degree of ω . The operators d and δ have the important property that they are adjoint with respect to the inner product (12)– that is

$$\langle d\omega, \xi \rangle = \langle \omega, \delta \xi \rangle \tag{17}$$

Now, in Geometric Algebra language this translates into an adjoint property between $\partial \cdot$ and $\partial \wedge$

$$\langle \boldsymbol{\partial} \wedge \boldsymbol{\omega}, \boldsymbol{\xi} \rangle = -\langle \boldsymbol{\omega}, \boldsymbol{\partial} \cdot \boldsymbol{\xi} \rangle \tag{18}$$

We will use this property frequently during this paper

1.4 Outline of the Paper

After introducing the main concepts, I begin the paper by decomposing the equation $\partial \mathbf{z} = 0$ grade by grade in order give to give a closed system of equations defining a monogenic function. I call these the, 'Clifford-Cauchy-Reimann' equations, or CCR equations for short. I then discuss some properties of these equations, showing how they require each grade of $\mathbf{z}(x)$ to be harmonic.

I then show that the CCR equations applied to a general multivector in the, 'Spacetime Algebra' Cl(3, 1), the Clifford algebra representing Minkowski Space, give the Dirac and Maxwell equations. This is not totally surprising— the Dirac equation is a monogenic function by definition, and it has long been realised that Maxwell's equations can be written as Cauchy- Reimann equations in quaternionic analysis (called regularity conditions) [32] and also via complexified quaterions (or biquaternions) [21], including in the context of Clifford Analysis [30]. However this paper unites both equations into a single framework, and shows that the (source-free) Maxwell and Dirac equations are the CCR equations for a general multivector in Cl(3, 1), implying that they are mathematically fundamental in a way linked to the geometry of Minkowski space. Having done this, I finish with a short discussion of future work which could be done in this area.

2 Clifford-Cauchy-Riemann equations

I am not aware of an explicit grade-by-grade decomposition of the Cauchy-Reimann equations for a Clifford algebra in the literature, though they must have been derived before in the course of calculations. I present here such a decomposition, which I will use for the main results in the next section. We write a general x-valued multivector function in Cl(p,q) as

$$\mathbf{z}(\mathbf{x}) = \sum_{i} \mathbf{f}_{i} = f_{0}(\mathbf{x}) + f_{1i}(\mathbf{x})e_{i} + f_{2ij}(\mathbf{x})e_{i}e_{j} + f_{3ijk}(\mathbf{x})e_{i}e_{j}e_{k} + \dots + f_{d}e_{1}\dots e_{d}$$
(19)

where \mathbf{f}_i is a blade of grade i, $f(\mathbf{x})$ is a scalar function of \mathbf{x} and i runs from 1 to d = p + q. We are interested in Monogenic functions, for which $\partial \mathbf{z} = 0$ This implies that

$$\sum_{i} \partial \mathbf{f}_{i} = \sum_{i} \left(\partial \cdot \mathbf{f}_{i+1} + \partial \wedge \mathbf{f}_{i-1} \right) = 0$$
(20)

equating all terms with the same grades, we find that

$$\partial \cdot \mathbf{f}_{i+1} = -\partial \wedge \mathbf{f}_{i-1}, i \neq \{1, d-1\}$$
(21)

$$\boldsymbol{\partial} \cdot \mathbf{f}_1 = 0 \tag{22}$$

$$\partial \wedge \mathbf{f}_{d-1} = 0 \tag{23}$$

with $\partial \cdot f_0$ and $\partial \wedge \mathbf{f}_d \equiv 0$ identically (you can't take the exterior derivative of a top form, or the divergence of a scalar function). These are the generalisation of the Cauchy-Riemann equations to general Clifford Algebras for the Dirac operator ∂ . I shall refer to them as the Clifford-Cauchy-Riemann, or CCR, equations. Note also that the condition $\partial \cdot \mathbf{f}_{i+1} = -\partial \wedge \mathbf{f}_{i-1}$ becomes $\delta f_{i-1} = df_{i+1}$ in the language of differential forms.

2.1 Properties of the CCR Equations

The first hing to note about these equations is that the equations for the odd and even parts separate. Equation (21) links \mathbf{f}_{n-1} and \mathbf{f}_{n+1} , whose grades are always both odd, or both even. Then, equation (22) always involves only \mathbf{f}_1 , which is odd; and (23) only involves \mathbf{f}_{d-1} , which is either odd or even depending upon *d*. Therefore the CCR equations split into parts involving only the odd or even components. This simplifies the analysis of these equations, and we shall make use of this below in section 3 when analysing the spacetime algebra. We can rewrite the CCR equations using various dualities. If we define $g_i = f_i I^{-1}$, then they become

$$\boldsymbol{\partial} \wedge \mathbf{g}_{d-(i+1)} I^{-1} = -\boldsymbol{\partial} \cdot \mathbf{g}_{d-(i-1)} I^{-1}, i \neq \{1, d-1\}$$
(24)

$$\partial \wedge \mathbf{g}_{d-1} I^{-1} = 0 \tag{25}$$

$$\partial \cdot \mathbf{g}_1 I^{-1} = 0 \tag{26}$$

Alternatively, we can only take the dual of one side of equation (21) to get

$$\partial \wedge \mathbf{f}_{i-1} = \partial \wedge \mathbf{g}_{d-(i+1)} I^{-1}$$
(27)

Note that whilst in general \mathbf{f}_{i-1} and $\mathbf{g}_{d-(i+1)}$ have different grades, when d is even, $\mathbf{f}_{d/2-1}$ and $\mathbf{g}_{d/2-1}$ both have the same grade in the equation, giving

$$\partial \wedge \mathbf{f}_{d/2-1} = \partial \wedge \mathbf{g}_{d/2-1} I^{-1}$$
(28)

This suggests that mathematically it could be interesting to investigate functions satisfying $\mathbf{z} = \pm \mathbf{z}I^{-1}$, which could be another avenue to explore to better understand the properties of solutions to these equations.

An important property of the Cauchy-Riemann equations is that any solution to them is also a harmonic function, i.e. $\partial^2 \mathbf{z} = 0$. This also applies to Clifford analysis [8]

Theorem 2.1. $Mon_p(\mathcal{M})$, the vector space of monogenic functions of degree p over a space \mathcal{M} , is identical to $Harm_p(\mathcal{M})$, the vector space of harmonic forms of degree p.

Since the spaces of harmonic forms are isomorphic to the de Rahm homology groups, this has as its corollary the well known result [31]

Theorem 2.2. $Mon_p(\mathcal{M})$ is isomorphic to $\mathcal{H}^p_{dR}(\mathcal{M})$, the *p*th de Rham cohomology group of \mathcal{M}

Although it is a long established result, I will give a proof of theorem 2.1 using the CCR equations, as it illuminates the structure of the equations, and shows how they are linked to Hodge Theory.

Proof. We begin with a Hodge decomposition of the vector spaces of p-blades:

$$\mathbf{z}(\mathbf{x}) = \sum_{i} \mathbf{f}_{i} = \sum_{i} \partial \wedge \mathbf{X}_{i-1} + \partial \cdot \mathbf{Y}_{i+1} + \mathbf{Z}_{i}$$
(29)

Now, for $p \neq 1, d-1$ imposing the condition that $\partial \mathbf{z} = 0$ implies that at degree p,

$$\partial \wedge \partial \cdot \mathbf{Y}_p = -\partial \cdot \wedge \mathbf{X}_p \equiv \mathbf{P}_i \tag{30}$$

But then, using the L^2 - norm (12), we have

$$\left\langle \mathbf{P}_{i}, \mathbf{P}_{i} \right\rangle = \left\langle \boldsymbol{\partial} \cdot \mathbf{Y}_{p}, -\boldsymbol{\partial} \wedge \mathbf{X}_{p} \right\rangle = \left\langle \boldsymbol{\partial} \cdot \boldsymbol{\partial} \cdot \mathbf{Y}_{p}, \mathbf{X}_{p} \right\rangle \equiv 0$$
 (31)

where the second inequality uses the skew- adjoint relation between $\partial \wedge$ and $\partial \cdot$. This implies that $P_i = 0$, and therefore separately that $\partial \wedge \partial \cdot \mathbf{Y}_p = 0$ and $\partial \cdot \partial \wedge \mathbf{X}_p = 0$. This implies that $\partial \wedge X_{i-1}$ and $\partial \cdot Y_{i+1}$ are harmonic, since $\partial \wedge \partial \wedge X_{i-1}$ and $\partial \cdot \partial \cdot Y_{i+1}$ are zero by definition, and harmonic functions satisfy $\partial \wedge \phi = \partial \cdot \phi = 0$. However, we assumed that $\partial \cdot \mathbf{Y}_p$ and $\partial \wedge \mathbf{X}_p$ were not harmonic functions, (and are therefore orthogonal to them under the L^2 -norm) and therefore they must be the zero- function.

For p = 1, $\partial \wedge X_1$ is covered by the above argument, and $\partial \cdot Y_1 \equiv 0$ by the CCR equations. Similarly for p = d - 1, $\partial \cdot \mathbf{Y}_{d-1}$ is covered by the above whereas $\partial \wedge \mathbf{X}_{d-1} \equiv 0$ by the CCR equations.

This shows that the only nonzero parts of \mathbf{z} are the harmonic blades \mathbf{Z}_p , proving that \mathbf{z} monogenic implies that \mathbf{z} is harmonic. Since harmonic forms are trivially monogenic, this establishes that the set of harmonic forms on a space is identical to the set of monogenic functions on that space.

This proof allows us to see how satisfying the CCR equations implies that a function is harmonic– the CCR equations imply that the exact and coexact parts at each grade of the function are equal to zero. This analysis also allows us to see what happens if the function fails to satisfy the CCR equations at grade p. In this case

$$\partial \wedge f_{i-1} \neq -\partial \cdot f_{i+1} \tag{32}$$

which means that neither $\partial \cdot \mathbf{X}_p \in \Lambda_{p-1}$ or $\partial \wedge \mathbf{Y}_p \in \Lambda_{p+1}$ are equal to zero. Therefore

$$f_{p-1} = \mathbf{Z}_{p-1} + \partial \cdot \mathbf{X}_p$$
 and $f_{p+1} = \mathbf{Z}_{p+1} + \partial \wedge \mathbf{Y}_p$ (33)

This means that if a multivector function \mathbf{z} does not satify the CCR equations at grade p, then neither the grade p-1 nor the grade p+1 component of \mathbf{z} is

harmonic.

Finally, in complex analysis, we can show that every real harmonic function on \mathbb{R}^n is the real part of a holomorphic function. For Clifford analysis, we have the following theorem (which I believe is original)

Theorem 2.3. Every harmonic function f(x) on \mathbb{R}^n is the 0-grade component of a monogenic function $\mathbf{z}(x) \in Cl(n)$.. Conversely, every monogenic function $\mathbf{z}(x)$ defines a harmonic function $f(x) = \langle \mathbf{z}(x) \rangle_0$

Proof. For the first direction, we start with a harmonic function f(x). For a scalar function, we can write the condition that f(x) is harmonic as $\partial \cdot \partial f = 0$. Expanding out, this implies that

$$\partial \cdot \partial \wedge f(x) = 0 \tag{34}$$

Now we consider

$$|\boldsymbol{\partial} \wedge f(x)| = \langle \boldsymbol{\partial} \wedge f(x), \boldsymbol{\partial} \wedge f(x) \rangle = \langle f(x), \boldsymbol{\partial} \cdot \boldsymbol{\partial} \wedge f(x) \rangle = 0$$
(35)

where we have used the adjoint property in the second equality, and the harmonic property of f(x) in the final one. But now since we are in a Reimannian space, $|\partial \wedge f(x)| = 0$ implies that $\partial \wedge f(x) \equiv 0$. From the above dicussion, this is the condition for f(x) to be the grade 0 part of a monogenic function.

Now consider the other direction. Suppose $f_0(x)$ is the grade-0 part of some monogenic function $\mathbf{z}(x)$. Then, by the CCR equations, $\partial \wedge f_0 = 0$. Then $\partial \cdot \partial f_0 = \partial \cdot \partial \wedge f_0 = 0$. This means that f_0 is a harmonic function on \mathbb{R}^n

3 CCR and the Spacetime Algebra

We now examine the CCR equations for the Spacetime Algebra. This is the Clifford Algebra Cl(3, 1), whose Geometric Algebra representation is associated to Minkowski Spacetime. It consists of 1 scalar, 4 vectors e_a , six bivectors e_ae_b , 4 trivectors $e_ae_be_c$, and 1 Pseudoscalar $I = e_0e_1e_2e_3$.

Here we use the index 0 to refer to the timelike direction, for which $e_0^2 = -1$, and $i, j, k = \{1, 2, 3\}$ for spacelike directions, for which $e_i^2 = 1$. These correspond to the gamma matrices γ_a in the usual Dirac theory, and the pseudoscalar I corresponds to $\gamma_5 = \gamma_0 \gamma_1 \gamma_2 \gamma_3$.

It is also worth noting that the bivector part of the algebra corresponds to Lorentz rotations [10]. The Timelike rotations correspond to terms of the form e_0e_i , and spacelike rotations correspond to e_ie_j . Finally, the vector derivative $\partial = e_0 \frac{\partial}{\partial t} + e_i \frac{\partial}{\partial x_i}$ Now, a general multivector in the spacetime algebra can be written in the form

This leads to the CCR equations

$$\partial \cdot \mathbf{f}_1 = 0$$

$$\partial f_0 = -\partial \cdot \mathbf{f}_2$$

$$\partial \wedge \mathbf{f}_1 = -\partial \cdot \mathbf{f}_3$$

$$\partial \wedge \mathbf{f}_3 = 0$$

$$\partial \cdot \mathbf{f}_4 = -\partial \wedge \mathbf{f}_2$$
(37)

As discussed in section 2.1, these split into odd and even sectors.

3.1 The Odd Sector

We start with the odd sector

$$\boldsymbol{\partial} \cdot \mathbf{f}_1 = 0 \tag{38}$$

$$\boldsymbol{\partial} \wedge \mathbf{f}_1 = -\boldsymbol{\partial} \cdot \mathbf{f}_3 \tag{39}$$

$$\boldsymbol{\partial} \wedge \mathbf{f}_3 = 0 \tag{40}$$

Since 4 is an even number, we can use the discussion of duality in equation (28) to write the second of these equations as

$$\partial \wedge \mathbf{f}_1 = -\partial \wedge \mathbf{g}_1 I^{-1} \tag{41}$$

where $g_3 = f_1 I^{-1}$. To solve this equation, we first of all note that the field strength for an electromagnetic potential is always anti-selfdual [13]. To see this, we can write $\partial \wedge \mathbf{A} = F_{ab} = E_{0i} + B_{jk}$, where E, B are the electric and magnetic fields respectively. The Electric field is a timelike bivector, and the magnetic field is a spacelike one. Then we can directly calculate

$$-\left(\partial_{0}A_{1}e_{0}e_{1}-\partial_{1}A_{0}e_{1}e_{0}\right)I^{-1} = \partial_{2}A_{3}e_{2}e_{3}-\partial_{3}A_{2}e_{3}e_{2}$$
$$-\left(\partial_{0}A_{2}e_{0}e_{2}-\partial_{2}A_{0}e_{2}e_{0}\right)I^{-1} = \partial_{3}A_{1}e_{3}e_{1}-\partial_{1}A_{3}e_{1}e_{3}$$
$$-\left(\partial_{0}A_{3}e_{0}e_{3}-\partial_{3}A_{0}e_{3}e_{0}\right)I^{-1} = \partial_{1}A_{2}e_{1}e_{2}-\partial_{2}A_{1}e_{2}e_{1}$$
(42)

This shows that $-\mathbf{E}I^{-1} = \mathbf{B}$, and therefore that $\mathbf{F} = -\mathbf{F}I^{-1}$. Hence $f_1 = g_1 = \mathbf{A}$ is always a solution to equation (39) since $\partial \wedge \mathbf{A} = -(\partial \wedge \mathbf{A})I^{-1}$. We now check for other solutions. Suppose that $\partial \wedge \mathbf{A} = -(\partial \wedge \mathbf{C})I^{-1}$ for $\mathbf{C} \neq \mathbf{A}$. Then $-(\partial \wedge \mathbf{C})I^{-1} - (\partial \wedge \mathbf{A})I^{-1} = 0$, which implies that $\partial \wedge (\mathbf{C} - \mathbf{A}) = 0$, and therefore \mathbf{C} and \mathbf{A} differ by the gradient of a scalar function, since $\partial \wedge \partial \lambda = 0$ automatically. Then the most general solution we can write is

$$\mathbf{f}_1 = \mathbf{A} + \partial \lambda_1, f_3 = g_1 I^{-1} = \left(\mathbf{A} + \partial \lambda_2\right) I^{-1}$$
(43)

for Scalar functions λ_1 and λ_2 . If we set $\lambda_1 = \lambda_2 \equiv \lambda$, and write $\tilde{\mathbf{A}} = \mathbf{A} + \partial \lambda$ then the CCR equations become the equation for a single anti-selfdual Field Strength.

$$\partial \wedge \tilde{\mathbf{A}} = (\partial \wedge \tilde{\mathbf{A}})I^{-1}$$
(44)

I will discuss the situation where $\lambda_1 = \lambda_2$ at the end of this subsection. Finally, if we write $\mathbf{F} = \partial \wedge \tilde{\mathbf{A}}$ then

$$oldsymbol{\partial} \mathbf{F} = oldsymbol{\partial} \cdot oldsymbol{\hat{\mathbf{A}}} + oldsymbol{\partial} \wedge oldsymbol{\hat{\mathbf{A}}} + oldsymbol{\partial} \wedge oldsymbol{\hat{\mathbf{A}}} = oldsymbol{\partial} \cdot oldsymbol{\hat{\mathbf{A}}} - oldsymbol{\partial} \cdot oldsymbol{\left(\mathbf{A}I^{-1}
ight)} = 0$$

which is simply the equation for a source-free Maxwell field written in Geometric Algebra notation [10]. I have used the fact that $\partial \cdot \partial \cdot \mathbf{f} = \partial \wedge \partial \wedge \mathbf{f} = 0$ for any \mathbf{f} in the second and fourth inequalities, and equation (44) in the second. We can now look at the equation $\partial \cdot \mathbf{f}_1 = 0$. This is Gauss' law

$$\boldsymbol{\partial} \cdot \tilde{\mathbf{A}} = 0 \tag{45}$$

Note that if we extract the gauge function λ we get

$$\partial \cdot \mathbf{A} + \partial^2 \lambda = 0 \tag{46}$$

which allows us to use λ to set a gauge exactly as in standard presentations of electromagnetism.

What about the remaining equation $\partial \wedge \mathbf{f}_3 = 0$? By the solution in equation (44) this is equivalent to $\partial \wedge (\tilde{A}I^{-1}) = 0$ which implies that $(\partial \cdot \tilde{\mathbf{A}})I^{-1} = 0$, which implies that $\partial \cdot \mathbf{A} + \partial^2 \lambda = 0$, just the same as equation (38) Putting it all together, we see that the odd sector of the CCR equations is

$$\boldsymbol{\partial} \wedge \tilde{\mathbf{A}} = \left(\boldsymbol{\partial} \wedge \tilde{\mathbf{A}}\right) I^{-1} \tag{47}$$

$$\boldsymbol{\partial} \cdot \tilde{\mathbf{A}} = 0 \tag{48}$$

(49)

This describes an anti-selfdual field arising from a vector potential \tilde{A} (equation (47), which obeys Gauss' Law (equation (48). We also have a gauge freedom to rescale \tilde{A} by a scalar function λ , via $\tilde{A} \to \tilde{A} + \partial \lambda$. What about when $\lambda_1 \neq \lambda_2$? In this case we have

$$f_1 = \mathbf{A} + \partial \lambda_1, \qquad f_3 = ig(\mathbf{A} + \partial \lambda_2ig) I^{-1}$$

Putting this into equations (38) and (40) we get that

$$\boldsymbol{\partial} \cdot \mathbf{A} = -\boldsymbol{\partial}^2 \lambda_1, \quad \boldsymbol{\partial} \cdot \mathbf{A} = -\boldsymbol{\partial}^2 \lambda_2 \tag{50}$$

This implies that $\partial^2 (\lambda_1 - \lambda_2) = 0$, and therefore that $\lambda_1 - \lambda_2 = a + b\mathbf{x}$, for $a, b \in \mathbb{R}$. I am unsure of the physical significance of this, and so I have focussed on the solutions where $\lambda_1 = \lambda_2$; however this would be a good topic for future work.

3.2 The Even Sector

The CCR equations for the even subalgebra are given by

$$\partial f_0 = -\partial \cdot \mathbf{f}_2 \tag{51}$$

$$\partial \wedge \mathbf{f}_2 = -\partial \cdot \mathbf{f}_4 \tag{52}$$

The physical meaning of these equations can be seen by considering the Dirac equation for a massless, uncharged spinor. Written in the Geometric Algebra representation, this is

$$\partial \phi = 0 \tag{53}$$

where $\phi = \rho^{1/2} e^{IB} e^{\theta/2}$. Here, $\rho(\mathbf{x})$ and $B(\mathbf{x})$ are scalar functions, and $\theta(\mathbf{x})$ is a bivector function. This form of the solution to Dirac's equation is due to David Hestenes [18]. Mathematically this corresponds to a polar decomposition of ϕ . There are eight degrees of freedom, just as we would expect – one each for ρ and B, and six contained in θ , which geneates a Lorentz rotation. Physically, following Hestenes, we can interpret the multivector function ϕ as a physical wave in Minkowski space, with $\rho^{1/2}$ as the amplitude, and $e^{\theta/2}$ being the spinor generator of a rotation into the rest frame of the particle. The physical interpretation of B is more ambigous, but Hestenes has suggested that for the full Dirac equation it corresponds to a hypothetical rapid oscillation of the electron called Zitterbewegung [19]. I will not addess these interpretational issues here. We can evaluate

$$\partial \phi = \left(\frac{\partial \rho}{\rho} + \partial (IB) + \partial \theta\right) \rho^{1/2} e^{IB} e^{\theta/2} = 0$$
(54)

Which implies

$$\frac{\partial \rho}{\rho} + \partial (IB) + \partial \theta/2 = 0 \tag{55}$$

collecting terms of the same grade, we find

$$\partial \ln(\rho) = -\partial \cdot \theta/2 \tag{56}$$

for grade 1, and

$$\partial \wedge \theta / 2 = -\partial \cdot (IB) \tag{57}$$

for grade 3, with $\partial \wedge (IB) \equiv 0$ identically. But these are just the CCR equations (51) and (52), with $f_0 = \ln\rho$, $\mathbf{f}_2 = \theta/2$ and $\mathbf{f}_4 = IB$. Therefore, $f_0, \mathbf{f}_2, \mathbf{f}_4$ satisfying the CCR equations automatically define a free Dirac field via (53). A final note: We could have written the scalar part of ϕ as $e^{a(\mathbf{x})/2}$ for some scalar function $a(\mathbf{x})$, rather than using $\rho^{1/2}$. This would have given us $f_0 = a/2$, rather than $\ln(\rho)$. I chose the notation $\rho^{1/2}$ partly to fit with the notation of Hestenes [18], and partly because of the similarity of the term $\frac{\partial \rho}{\rho}$ to the quantum potential of Bohm [6], which is derived from a similar polar decomposition of the wavefunction.

4 Conclusion and Further Work

Putting it all together, we find that a multivector

$$\mathbf{z} = \ln(\rho) + \tilde{\mathbf{A}} + \boldsymbol{\theta}/2 + \tilde{\mathbf{A}}I^{-1} + IB$$
(58)

satisfying the CCR equations in the Spacetime Algebra Cl(3,1) defines both a free Dirac field $\phi = \rho^{1/2} e^{IB} e^{\theta/2}$, and an electromagnetic field strength $\mathbf{F} = \boldsymbol{\partial} \wedge \tilde{\mathbf{A}}$, where $\tilde{\mathbf{A}}$ satisfies Gauss' law, and is defined up to the addition of a gradient $\boldsymbol{\partial}\lambda$.

Aside from the points mentioned earlier in the paper (For example, below equation (50)), there are many things to explore. Chief among these for me is the question whether we can obtain the equations for a massive or charged Dirac field, or for an electromagnetic field with a source $\mathbf{J} = \partial \mathbf{F}$? I suspect that this would involve considering Dirac operators of the form $\partial + \mathbf{A}$, where \mathbf{A} is a gauge potential.

Once this had been done, it would be interesting to look at the geometric meaning of the cohomology groups of this operator. We could perform a Hodge decomposition of the space of differential forms by looking for forms which are harmonic with respect to this gauged Dirac operator. We could then compare the resulting gauge theory cohomologies with more standard methods such as the BRST or BV cohomologies [17][27], making use of the fact that Geometric Algebra allows us to give a direct geometric interpretation of all our expressions to determine the physical meaning of the resulting cohomology groups.

Another avenue would be to investigate Hodge theory on Lorentzian manifolds. The motivation for this comes from the fact that in the proof of theorem 2.1, the key argument was that the condition $\partial \wedge X_{i-1} = -\partial \cdot Y_{i+1} \equiv P_i$ implied that $\langle P_i, P_i \rangle = 0$, and hence that $P_i = 0$.

In the Lorentzian case, the condition $\langle P_i, P_i \rangle = 0$ no longer implies that $P_i = 0$, but that P_i is null. Therefore the condition for a multivector function $\mathbf{z}(x)$ to be monogenic is no longer that it is made up of harmonic forms, but that it is made up of forms whose boundaries and coboundaries are null under the L_2 norm, such that $\partial \wedge X_{i-1} = -\partial \cdot Y_{i+1}$. Investigating these solutions could be a valuable tool for understanding cohomology on Lorentzian manifolds, which so far has focused on the timelike and spacelike parts of the spacetime [23][4][3] . I hope to explore this in future work.

References

- [1] A.MACDONALD, Linear and Geometric Algebra, Luther College, 2010.
- [2] M. ATIYAH, M. DUNAJSKI, AND L. MASON, Twistor theory at fifty, Proc. Royal. Soc. A, (2017).
- [3] C. BAER, Green-hyperbolic operators on globally hyperbolic spacetimes, Commun. Math. Phys., 333 (2015).
- M. BENINI, Optimal space of linear classical observables for maxwell kforms via spacelike and timelike compact de rham cohomologies, J. Math. Phys., 57 (2016).

- [5] E. BINZ, M. DE GOSSON, AND B. HILEY, Clifford algebras in symplectic geometry and quantum mechanics, Foundations of Physics, 43 (2013), pp. 424–439.
- [6] D. BOHM, Wholeness and the Implicate Order, Routledge, 1980.
- [7] E. CHISHOLM, Geometric algebra. arXiv preprint 1205.5935, 2012.
- [8] R. DELANGHE, F. SOMMEN, AND V. SOČEK, Clifford Algebra and Spinor Valued Functions, Springer, 1992.
- [9] J. DIEUDONNE, The tragedy of grassman, Linear and Multilinear Algebra, 8 (1979).
- [10] C. DORAN AND G. LASENBY, *Geometric Algebra for Physicists*, CUP, 2003.
- [11] L. DORST, D. FONTIJNE, AND S. MANN, *Geometric Algebra for Computer Science*, Morgan Kaufmann, 2009.
- [12] M. DUBOIS-VIOLETTE, Exceptional quantum geometry and particle physics, Nucl. Phys. B, 912 (2016).
- [13] J. FIGUEROA-O'FARRILL, Electromagnetic duality for children.
- [14] R. FUETER, Die funktionentheorie der differentialgleichungen $\delta u = 0$ und $\delta \delta u = 0$ mit vier reellen variablen, Comment. Math. Helv., 7 (1935), pp. 307–330.
- [15] R. FUETER, über die funktionentheorie in einer hyperkomplexen algebra, Elemente der Mathematik, Band III/5 (1948), pp. 89–94.
- [16] C. FUREY, Standard model physics from an algebra?, PhD thesis, Waterloo U., 2015.
- [17] M. HENNEAUX AND C. TEITELBOIM, Quantization of Gauge Systems, Princeton, 1994.
- [18] D. HESTENES, Mysteries and insights of dirac theory, Annales de la Fondation Louis de Broglie, 28 (2003), pp. 390–408.
- [19] D. HESTENES, Zitterbewegung in quantum mechanics, Foundations of Physics, 40 (2010), pp. 1–54.
- [20] D. HESTENES AND G. SOBCZYK, Clifford Algebra to Geometric Calculus, Springer, 1987.
- [21] K. IMAEDA, A new formulation of classical electrodynamics, Nuevo Cimento, Sezione B, 32B (1976), pp. 138–162.
- [22] J.RYAN, Foundations of clifford analysis.

- [23] I. KHAVKINE, Cohomology with causally restricted supports, Annales Henri Poincaré, 17 (2016).
- [24] H. LAWSON AND M.-L. MICHELSOHN, Spin Geometry, vol. PMS, Princeton, 1990.
- [25] P. LOUNESTO, Clifford Algebras and Spinors, CUP, 2009.
- [26] G. MOISIL AND N. THÉODORESCO, Functions holomorphes dans l'espace, Mathematica, Cluj, 3 (1931), pp. 142–159.
- [27] K. REJZNER, Bv quantization in perturbative algebraic qft: Fundamental concepts and perspectives, in The Philosophy and Physics of Noether's Theorems, 2018.
- [28] M. RIESZ, Clifford Numbers and Spinors, Springer, 1993 (Facsimile of 1958 original).
- [29] C. ROBERTS, Hodge and Gelfand theory in Clifford analysis and tomography, PhD thesis, Colorado State University, 2022.
- [30] J. RYAN, Duality in complex clifford analysis, Journal of Functional Analysis, 61 (1985), pp. 117–135.
- [31] F. SOMMEN, Monogenic differential forms and homology theory, Proc. Royal. Irish Academy, (1984), pp. 87–109.
- [32] Z. TYPALDOS AND R. POGORZELSKI, Quaternion calculus and the solution of maxwell's equations. scanned typewritten notes, ece-research.umn.edu/summa/notes/Mathematics/0043.pdf, 1975.