

SOME MULTIPLICATION FORMULAS IN QUEER q -SCHUR SUPERALGEBRAS

JIE DU^{*}, HAIXIA GU^{**}, ZHENHUA LI^{*}, AND JINKUI WAN^{**}

ABSTRACT. Building on the work [18], where some standard basis for the queer q -Schur superalgebra $\mathcal{Q}_q(n, r; R)$ is defined by a labelling set of matrices and their associated double coset representatives, we investigate the matrix representation of the regular module of $\mathcal{Q}_q(n, r; R)$ with respect to this basis. More precisely, we derive explicitly (resp., partial explicitly) the multiplication formulas of the basis elements by certain even (resp., odd) generators of a queer q -Schur superalgebra.

These multiplication formulas are highly technical to derive, especially in the odd case. It requires to discover many multiplication (or commutation) formulas in the Hecke–Clifford algebra $\mathcal{H}_{r,R}^c$ associated with the labelling matrices. For example, for a given such a labelling matrix A^* , there are several matrices $w(A)$, $\sigma(A)$, \tilde{A} , and \hat{A} associated with the base matrix A of A^* , where $w(A)$ is used to compute a reduced expression of the distinguished double coset representatives d_A , and the other matrices are used to describe the permutation d_A and the SDP (commutation) condition between T_{d_A} and generators of the Clifford subsuperalgebra.

With these multiplication formulas, we will construct a new realisation of the quantum queer supergroup in a forthcoming paper [13], and to give new applications to the integral Schur–Olshanski duality and its associated representation theory at roots of unity.

CONTENTS

1.	Introduction	1
2.	The queer q -Schur superalgebra	3
3.	Some commutation relations in the Hecke–Clifford superalgebra	7
4.	Distinguished double coset representatives and the SDP condition	11
5.	Multiplication formulas in $\mathcal{Q}_q(n, r; R)$: the even case	15
6.	Multiplication formulas in $\mathcal{Q}_q(n, r; R)$: the odd case	21
7.	Some special cases	30
8.	An outline of some applications	34
	Appendix A. Some multiplication formulas in $\mathcal{H}_{r,R}^c$	35
	References	39

1. INTRODUCTION

For a given finitely generated algebra \mathcal{A} over a field, the regular representation of ${}_{\mathcal{A}}\mathcal{A}$ can be understood as the matrix representations of generators with respect to a certain given basis. If the basis contains the generators as a subset, computing such matrix representations amounts to the determination of the structure constants associated with the generators or to the explicit multiplication formulas of the basis by the generators. When \mathcal{A} is a member of the family of Schur and q -Schur algebras, which are fundamental objects in representation theory, such multiplication formulas are first established by Beilinson–Lusztig–MacPherson (BLM) in [3]

Date: August 7, 2023.

2010 Mathematics Subject Classification. 20B30, 20G43, 17A70, 20C08.

Key words and phrases. quantum supergroup, q -Schur superalgebra, Hecke–Clifford superalgebra, Schur–Olshanski duality.

^{*}Partially supported by the Science FRG; ^{**}Partially supported by NSFC-12071129.

^{*}Partially supported by NSFC-11871404; ^{**}Partially supported by NSFC-12122101 & NSFC-12071026.

(see also [8, §13.5] for more details), and used to determine the regular representation (or a new realisation) of quantum linear groups.

The method in [3] is geometric, using the convolution algebra interpretation of q -Schur algebras associated with certain partial flag varieties, and the multiplication formulas are derived directly from the convolution products. This geometric method has also been generalised to the q -Schur algebra of affine type A [22, 25] and, more recently, to q -Schur algebras arising from i -quantum groups and quantum symmetric pairs [2, 1] of type AIII, whose Satake diagrams have no black nodes (see also [23] for a coordinate algebra approach to these q -Schur algebras), and to affine type C [21]. However, for quantum linear supergroups and q -Schur superalgebras, the geometric method cannot be applied. Thus, an algebraic method is developed in [12, 14]. In this approach, the multiplication formulas for q -Schur superalgebras, regarded as endomorphism algebras of q -permutation supermodules over Hecke algebras, are derived using certain multiplication (or commutation) formulas in the Hecke algebras involving distinguished double coset representatives.

The aim of this paper is to establish similar fundamental multiplication formulas for queer q -Schur superalgebras arising from the Schur-Olshanski duality [26]. We will use the definition of queer q -Schur superalgebras $\mathcal{Q}_q(n, r; R)$ introduced by the first and last authors [18]. As endomorphism algebras of a direct sum of certain cyclic modules (or queer q -permutation supermodules) over the Hecke-Clifford superalgebra $\mathcal{H}_{r,R}^c$, such a queer q -Schur superalgebra admits a standard basis $\{\phi_{A^*}\}$ labelled by certain $n \times 2n$ matrices $A^* = (A^{\bar{0}}|A^{\bar{1}})$. Here each such a matrix defines a distinguished double coset representative d_A associated with the “base matrix” $A := A^{\bar{0}} + A^{\bar{1}}$ in the symmetric group \mathfrak{S}_r and an element c_{A^*} in the Clifford superalgebra, and the standard basis element ϕ_{A^*} is defined by an element T_{A^*} in $\mathcal{H}_{r,R}^c$ associated with d_A and the element c_{A^*} . Continuing with this construction, we now compute the structure constants occurring in the multiplication $\phi_{X^*}\phi_{A^*}$, for an arbitrary A^* , where X^* is one of the following six matrices

$$\begin{aligned} &(\lambda|O), \quad (\lambda + E_{h,h+1} - E_{h+1,h+1}|O), \quad (\lambda - E_{h,h} + E_{h+1,h}|O) \quad (\text{of parity } 0); \\ &(\lambda - E_{h,h}|E_{h,h}), \quad (\lambda - E_{h+1,h+1}|E_{h,h+1}), \quad (\lambda - E_{h,h}|E_{h+1,h}) \quad (\text{of parity } 1). \end{aligned}$$

Here $\lambda \in \mathbb{N}^n$ represent diagonal matrices. These matrices correspond to the Chevalley type generators for the queer Lie superalgebra and its universal enveloping algebra.

However, unlike the quantum \mathfrak{gl}_n or $\mathfrak{gl}_{m|n}$ case, there are no complete multiplication formulas for X^* being the odd matrices above. This subtlety was observed over 10 years ago! In order to seek new ideas and methods, we tested two experiments. First, can the universal enveloping algebra of the queer Lie superalgebra be approached by queer Schur superalgebras? Second, can the quantum queer supergroups be approached by constructing its regular representation directly? Both questions had affirmative answers; see [20] and [16]. The two works convinced us that a construction using queer q -Schur superalgebras must exist and so does the integral Schur-Olshanski duality.

Here is the new approach to tackle the problem. Instead of seeking complete multiplication formulas in the odd case, we focus on the “head part”, which is determined by a so-called SDP condition, leaving out the big “tail part”; see Theorems 6.2–6.4. The SDP condition is about commuting certain generators in the Clifford superalgebra with a basis element T_{d_A} in the Hecke algebra associated with the distinguished double coset representative d_A defined by a matrix A ; see Definition 4.2. If a matrix satisfies this condition, then certain multiplication formulas in the odd case can be completely computed. Fortunately, these head part formulas do provide a path to solving this decade long problem.

In the forthcoming paper [13], we will use the multiplication formulas (and partial formulas in the odd case) in this paper to give a new construction (a BLM type realisation) for the quantum queer supergroup $\mathbf{U}(\mathfrak{q}_n)$. Furthermore, this breakthrough will allow us to establish the integral Schur-Olshanski duality and to study the polynomial representation theory at roots of unity [19]. We also remark that, as a by product, all above given ϕ_{X^*} generate the queer q -Schur

algebra $\mathcal{Q}_q(n, r; R)$ and the multiplication formulas in §§5–6 provide the regular representation on the standard basis. See Remark 6.5.

We organise the paper as follows. We start with the definitions of Hecke-Clifford superalgebras $\mathcal{H}_{r,R}^c$ and queer q -Schur superalgebras $\mathcal{Q}_q(n, r; R)$ in §2, and discuss the construction of their standard bases. In §3, we establish several commutation formulas in $\mathcal{H}_{r,R}^c$ which involve distinguished double coset representatives and the c -elements aforementioned. The SDP condition is introduced and investigated in §4. The main result of this section is Theorem 4.6 which characterises the SDP condition by the shape of the matrix A . The fundamental multiplication formulas for the even (resp., odd) case are established in §5 (resp. in §6). In §7, we present some special cases for the multiplication formulas and relations relative to the generators. Finally, in the last section, we outline some ongoing work on applications of these multiplication formulas, including the aforementioned new realisation of the quantum queer supergroup $\mathbf{U}(\mathfrak{q}_n)$.

Some Notations. Let \mathbb{Z} (resp., \mathbb{N}) be the set of integers (resp. non-negative integers). Let $\mathbb{Z}_2 = \{\bar{0}, \bar{1}\}$ denote the abelian group of order 2 and $\mathbb{N}_2 = \{0, 1\} \subset \mathbb{N}$.

For any positive integer n , let $[1, n] = \{1, 2, \dots, n\}$, and define $[i+1, i+n]$ similarly for any integer i . Let $[[n]] = [[n]]_q = \frac{q^n - 1}{q - 1}$ denote the Gaussian integers and let $[[n]]_{x,y} = [[n]]_x - [[n]]_y$. The free \mathbb{Z} -module \mathbb{Z}^n has standard basis $\epsilon_1, \epsilon_2, \dots, \epsilon_n$ and “positive roots” $\alpha_1, \dots, \alpha_{n-1}$, where $\alpha_h = \epsilon_h - \epsilon_{h+1}$.

Let $M_n(\mathbb{N})$ be the set of $n \times n$ -matrices over \mathbb{N} . For any $m \times n$ matrix A over \mathbb{N} , let $|A|$ denote the sum of entries of A .

2. THE QUEER q -SCHUR SUPERALGEBRA

Let R be a commutative ring with 1. The Clifford superalgebra \mathcal{C}_r is an associative superalgebra over R defined by odd generators c_1, \dots, c_r and relations: for $1 \leq i, j \leq r$,

$$c_i^2 = -1, \quad c_i c_j = -c_j c_i \quad (i \neq j). \quad (2.0.1)$$

Let \mathfrak{S}_r be the symmetric group on r letters, with Coxeter generators (or simple reflections) $s_i = (i, i+1)$, for all $1 \leq i < r$ and length function ℓ . Let $y < w$ denote the Bruhat partial order relation on \mathfrak{S}_r . The Hecke algebra $\mathcal{H}_{r,R} = \mathcal{H}(\mathfrak{S}_r)_R$ associated with \mathfrak{S}_r and $1 \neq q \in R^\times$ is an algebra over R generated by $T_i = T_{s_i}$ for all $1 \leq i < r$, subject to the relations: for $1 \leq i, j \leq r-1$,

$$\begin{aligned} (T_i - q)(T_i + 1) &= 0, \\ T_i T_j &= T_j T_i \quad (|i - j| > 1), \\ T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1} \quad (i \neq r-1). \end{aligned} \quad (2.0.2)$$

The *Hecke-Clifford superalgebra* $\mathcal{H}_{r,R}^c$ is a superalgebra over R generated by even generators T_1, \dots, T_{r-1} and odd generators c_1, \dots, c_r , subject to the relations (2.0.1)-(2.0.2) and the extra relations: for $1 \leq i \leq r-1, 1 \leq j \leq r$,

$$\begin{aligned} c_j T_i &= T_i c_j \quad (j \neq i, i+1), \\ c_{i+1} T_i &= T_i c_i, \\ c_i T_i &= T_i c_{i+1} + (q-1)(c_i - c_{i+1}). \end{aligned} \quad (2.0.3)$$

For any homogeneous element $h \in \mathcal{H}_{r,R}^c$, let $p(h)$ be its parity, more precisely, $p(h) = \bar{0}$ if h is even, or $p(h) = \bar{1}$ if h is odd.

The following relations can be deduced directly from (2.0.2)-(2.0.3) and will be used in the sequel: for any $1 \leq k \leq r-1$,

$$\begin{aligned} (1) \quad qT_k^{-1} &= T_k - (q-1), \\ (2) \quad T_k c_{k+1} &= q c_k T_k^{-1} + (q-1)c_{k+1}, \\ (3) \quad c_k T_k &= q T_k^{-1} c_{k+1} + (q-1)c_k. \end{aligned} \quad (2.0.4)$$

For $w \in \mathfrak{S}_r$ with reduced expression $w = s_{i_1} s_{i_2} \cdots s_{i_k}$ and $\alpha \in \mathbb{N}_2^r$, let

$$T_w = T_{i_1} \cdots T_{i_k}, \quad c^\alpha = c_1^{\alpha_1} c_2^{\alpha_2} \cdots c_r^{\alpha_r}.$$

Then it is known (cf. [17, Lemma 2.2]) that both

$$\mathcal{B} = \{c^\alpha T_w \mid w \in \mathfrak{S}_r, \alpha \in \mathbb{N}_2^r\} \quad \text{and} \quad \mathcal{B}' = \{T_w c^\alpha \mid w \in \mathfrak{S}_r, \alpha \in \mathbb{N}_2^r\}$$

form (standard) R -bases for $\mathcal{H}_{r,R}^c$, consisting of homogeneous elements.

Lemma 2.1. *The transition matrix from \mathcal{B} to \mathcal{B}' has the lower triangular form*

$$\begin{pmatrix} P_1 & O & \cdots & O \\ * & P_2 & \cdots & O \\ \vdots & \vdots & \ddots & \vdots \\ * & * & \cdots & P_r! \end{pmatrix}, \quad (2.1.1)$$

where each block P_k ($k \in [1, r!]$) on the diagonal is a signed permutation matrix of size $2^r \times 2^r$. Moreover, each P_k is a block diagonal of blocks of sizes $\binom{r}{s}$, for all $s \in [1, r]$.

Proof. Define the place permutation of \mathfrak{S}_r on \mathbb{N}_2^r by setting $(\alpha_1, \dots, \alpha_r) \cdot w = (\alpha_{w(1)}, \dots, \alpha_{w(r)})$. We have, by [18, (2.4)],

$$c^\alpha T_w = \pm T_w c^{\alpha \cdot w} + \sum_{y < w, \beta \in \mathbb{N}_2^r} f_{\beta, y}^{\alpha, w} T_y c^\beta$$

for some $f_{\beta, y}^{\alpha, w} \in R$. For $1 \leq s \leq r$, let $\mathbb{N}_2^{r, s} = \{\alpha \in \mathbb{N}_2^r \mid s = |\alpha|\}$. If $\alpha \in \mathbb{N}_2^{r, s}$, the formula above can be further refined to

$$c^\alpha T_w = \pm T_w c^{\alpha \cdot w} + \sum_{y < w, \beta \in \mathbb{N}_2^{r, s}} f_{\beta, y}^{\alpha, w} T_y c^\beta$$

(For a proof, use induction on $|\alpha|$.) Thus, if we order the basis \mathcal{B} by linearly refining the partial order defined by setting

$$T_w c^\alpha < T_y c^\beta \iff w > y \text{ or } y = w, |\alpha| \leq |\beta| \quad (\text{and in lexicographic order if } |\alpha| = |\beta|)$$

(and define $c^\alpha T_w < c^\beta T_y$ similarly), then the transition matrix has the desired form. \square

For later use, we record the following special case:

$$c_j T_w = T_w c_{w^{-1}(j)} + \sum_{y < w, i \in [1, r]} f_{i, y}^{j, w} T_y c_i. \quad (2.1.2)$$

Define the set of compositions of r into n parts:

$$\Lambda(n, r) = \{\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n) \in \mathbb{N}^n \mid r = |\lambda| := \lambda_1 + \cdots + \lambda_n\}. \quad (2.1.3)$$

For $\lambda, \mu \in \Lambda(n, r)$, let \mathfrak{S}_λ be the standard Young subgroup of \mathfrak{S}_r corresponding to λ , and let \mathcal{D}_λ be the distinguished (i.e. shortest) representatives of right cosets of \mathfrak{S}_λ in \mathfrak{S}_r . Then $\mathcal{D}_{\lambda, \mu} := \mathcal{D}_\lambda \cap \mathcal{D}_\mu^{-1}$ is the distinguished representatives of the \mathfrak{S}_λ - \mathfrak{S}_μ double cosets in \mathfrak{S}_r .

Associated with \mathfrak{S}_λ , define the even element

$$x_\lambda = \Sigma_{\mathfrak{S}_\lambda} := \sum_{w \in \mathfrak{S}_\lambda} T_w. \quad (2.1.4)$$

It is known that, for $s_i \in \mathfrak{S}_\lambda$, we have

$$T_i x_\lambda = x_\lambda T_i = q x_\lambda, \quad (2.1.5)$$

and $x_\lambda \mathcal{H}_{r,R}^c$ is an R -free supermodule¹, as well as a right $\mathcal{H}_{r,R}^c$ -supermodule.

Given R -free supermodules V, W , let $\text{Hom}_R(V, W)$ be the set of all R -linear maps from V to W . We make $\text{Hom}_R(V, W)$ into an R -free supermodule by declaring $\text{Hom}_R(V, W)_i$ to be the set of homogeneous maps of parity $i \in \mathbb{Z}_2$, that is, the linear maps $\theta : V \rightarrow W$ with $\theta(V_j) \subset W_{i+j}$ for $j \in \mathbb{Z}_2$. Note that it is typical in a superalgebra to write the expressions which only make

¹By an R -free supermodule, we mean an R -module of the form $V_0 \oplus V_1$ such that both V_0 and V_1 are R -free.

sense for homogeneous elements, and the expected meaning for arbitrary elements is obtained by extending linearly from the homogeneous case.

Following [18], the *queer q -Schur superalgebra* over R is defined to be

$$\mathcal{Q}_q(n, r; R) := \text{End}_{\mathcal{H}_{r,R}^c} \left(\bigoplus_{\lambda \in \Lambda(n,r)} x_\lambda \mathcal{H}_{r,R}^c \right). \quad (2.1.6)$$

Here we view $T_R(n, r) := \bigoplus_{\lambda \in \Lambda(n,r)} x_\lambda \mathcal{H}_{r,R}^c$ as a right $\mathcal{H}_{r,R}^c$ -module and homomorphisms are $\mathcal{H}_{r,R}^c$ -module homomorphisms, i.e., R -linear maps $\phi : T_R(n, r) \rightarrow T_R(n, r)$ satisfying $\phi(mh) = \phi(m)h$, for all $m \in T_R(n, r)$ and $h \in \mathcal{H}_{r,R}^c$.

Remark 2.2. Notice that $\text{End}_R(T_R(n, r))$ is an R -free supermodule. Thus, we may also define the endomorphism superalgebra $\text{End}_{\mathcal{H}_{r,R}^c}^s(T_R(n, r))$ which consists of $\mathcal{H}_{r,R}^c$ -supermodule homomorphisms defined by the rule

$$\Phi(mh) = (-1)^{p(\Phi)p(h)} \Phi(m)h, \quad (2.2.1)$$

for all $m \in T_R(n, r)$ and (homogeneous) $h \in \mathcal{H}_{r,R}^c$, $\Phi \in \text{End}_R(T_R(n, r))$ (see [7, Def. 1.1]). These superalgebras over $R = \mathbb{Q}(v)$ ($q = v^2$) will be used in [13] for a new construction of the quantum queer supergroup. It can be proved that if $\sqrt{-1} \in R$ then

$$\text{End}_{\mathcal{H}_{r,R}^c}^s(T_R(n, r)) \cong \text{End}_{\mathcal{H}_{r,R}^c}(T_R(n, r)) = \mathcal{Q}_q(n, r; R).$$

Let $\tilde{\mathcal{Q}}(n, r) := \text{End}_{\mathcal{H}_{r,\mathbb{Q}(v)}^c}^s(T_{\mathbb{Q}(v)}(n, r))$. Note that $T_{\mathbb{Q}(v)}(n, r)$ can be identified as the tensor space considered in [26]; see [28, Lem. 3.1].

In order to describe a natural (standard) basis for $\mathcal{Q}_q(n, r; R)$, we need a few more notations. Associated with $\lambda = (\lambda_1, \dots, \lambda_n) \in \Lambda(n, r)$, define its *partial sum sequence*,

$$\tilde{\lambda} := (\tilde{\lambda}_1, \dots, \tilde{\lambda}_n), \text{ where } \tilde{\lambda}_i = \sum_{k=1}^i \lambda_k \quad (1 \leq i \leq n), \text{ and } \tilde{\lambda}_0 = 0. \quad (2.2.2)$$

Let $M_n(\mathbb{N})$ be the set of $n \times n$ -matrices over \mathbb{N} . For any $A = (a_{i,j}) \in M_n(\mathbb{N})$, let

$$\nu_A := (a_{1,1}, \dots, a_{n,1}, a_{1,2}, \dots, a_{n,2}, \dots, a_{1,n}, \dots, a_{n,n}) = (\nu_1, \nu_2, \dots, \nu_{n^2}). \quad (2.2.3)$$

Define $\tilde{\nu}_A = \nu_{\tilde{A}}$ similarly, where

$$\tilde{A} = (\tilde{a}_{i,j}), \text{ with } \tilde{a}_{i,j} := \sum_{p=1}^{j-1} \sum_{u=1}^n a_{u,p} + \sum_{u=1}^i a_{u,j}, \text{ and set } \tilde{a}_{0,1} = 0, \tilde{a}_{0,j} = \tilde{a}_{n,j-1} \quad (j \geq 2). \quad (2.2.4)$$

Following [18, Section 4], we introduce the following odd elements in $\mathcal{H}_{r,R}^c$: for $1 \leq i \leq j \leq r$,

$$c_{q,i,j} = q^{j-i} c_i + \dots + q c_{j-1} + c_j, \quad c'_{q,i,j} = c_i + q c_{i+1} + \dots + q^{j-i} c_j. \quad (2.2.5)$$

For $\lambda = (\lambda_1, \dots, \lambda_m) \in \mathbb{N}^m$ and $\alpha \in \mathbb{N}^m$ such that $\alpha \leq \lambda$ (meaning $\alpha_i \leq \lambda_i$, for all $1 \leq i \leq m$), recall the following elements introduced in [18, §4]

$$\begin{aligned} c_\lambda^\alpha &= (c_{q,1,\tilde{\lambda}_1})^{\alpha_1} (c_{q,\tilde{\lambda}_1+1,\tilde{\lambda}_2})^{\alpha_2} \dots (c_{q,\tilde{\lambda}_{m-1}+1,\tilde{\lambda}_m})^{\alpha_m}, \\ (c_\lambda^\alpha)' &= (c'_{q,1,\tilde{\lambda}_1})^{\alpha_1} (c'_{q,\tilde{\lambda}_1+1,\tilde{\lambda}_2})^{\alpha_2} \dots (c'_{q,\tilde{\lambda}_{m-1}+1,\tilde{\lambda}_m})^{\alpha_m}. \end{aligned} \quad (2.2.6)$$

Here we used the convention $(c_{q,\tilde{\lambda}_{i-1}+1,\tilde{\lambda}_i})^{\alpha_i} = 1$ if $\lambda_i = 0$ (so, $\alpha_i = 0$). By [18, Lem. 4.1, Cor. 4.3], we have, for all $\alpha \leq \lambda$,

$$x_\lambda c_\lambda^\alpha = (c_\lambda^\alpha)' x_\lambda. \quad (2.2.7)$$

Let $M_n(\mathbb{N})_r = \{A = (a_{i,j}) \in M_n(\mathbb{N}) \mid r = |A|\}$, where $|A| := \sum_{i,j} a_{i,j}$. It is well known (see, e.g., [8, Section 4.2]) that there is a bijection

$$\begin{aligned} j : M_n(\mathbb{N})_r &\rightarrow \{(\lambda, d, \mu) \mid \lambda, \mu \in \Lambda(n, r), d \in \mathcal{D}_{\lambda,\mu}\} \\ A &\mapsto (\text{ro}(A), d_A, \text{co}(A)), \end{aligned} \quad (2.2.8)$$

where, for $A = (a_{i,j})$,

$$\text{ro}(A) = \left(\sum_{j=1}^n a_{1,j}, \dots, \sum_{j=1}^n a_{n,j} \right) \text{ and } \text{co}(A) = \left(\sum_{i=1}^n a_{i,1}, \dots, \sum_{i=1}^n a_{i,n} \right). \quad (2.2.9)$$

Call $\text{ro}(A)$ (resp., $\text{co}(A)$) the *row sum* (resp., *column sum*) vector of A . For the distinguished double coset representative d_A , see (3.1.4) below for its reduced expression and (4.0.1) for the precise description as a permutation.

We are now ready to describe a basis for $\mathcal{Q}_q(n, r; R)$.

Recall $\mathbb{N}_2 = \{0, 1\} \subset \mathbb{N}$ and the following matrix sets introduced in [18]:²

$$\begin{aligned} M_n(\mathbb{N}|\mathbb{N}_2) &= \{A^* = (A^{\bar{0}}|A^{\bar{1}}) \mid A^{\bar{0}} \in M_n(\mathbb{N}), A^{\bar{1}} \in M_n(\mathbb{N}_2)\}, \\ M_n(\mathbb{N}|\mathbb{N}_2)_r &= \{A^* = (A^{\bar{0}}|A^{\bar{1}}) \in M_n(\mathbb{N}|\mathbb{N}_2) \mid A^{\bar{0}} + A^{\bar{1}} \in M_n(\mathbb{N})_r\}. \end{aligned} \quad (2.2.10)$$

If $A^* = (A^{\bar{0}}|A^{\bar{1}}) \in M_n(\mathbb{N}|\mathbb{N}_2)_r$ with $A^{\bar{0}} = (a_{i,j}^{\bar{0}})$, $A^{\bar{1}} = (a_{i,j}^{\bar{1}})$, then we often write $A^* = (a_{i,j}^{\bar{0}}|a_{i,j}^{\bar{1}})$. We call the matrix

$$A = \lfloor A^* \rfloor := A^{\bar{0}} + A^{\bar{1}} = (a_{i,j}) \in M_n(\mathbb{N})_r, \text{ where } a_{i,j} = a_{i,j}^{\bar{0}} + a_{i,j}^{\bar{1}}, \quad (2.2.11)$$

the *base* (matrix) of A^* . For later use in Section 6, we also let, for $\lambda, \mu \in \Lambda(n, r)$,

$$M_n(\mathbb{N}|\mathbb{N}_2)_{\lambda, \mu} = \{A^* \in M_n(\mathbb{N}|\mathbb{N}_2) \mid \text{ro}(A) = \lambda, \text{co}(A) = \mu\}.$$

With the compositions $\nu = \nu_A$ and $\alpha = \nu_{A^{\bar{1}}}$ as defined in (2.2.3), we define

$$c_{A^*} := c_\nu^\alpha \quad \text{and} \quad T_{A^*} := x_{\text{ro}(A)} T_{d_A} c_{A^*} \sum_{\sigma \in \mathcal{D}_\nu \cap \mathfrak{S}_{\text{co}(A)}} T_\sigma. \quad (2.2.12)$$

For the simplicity of later use, we abbreviate the ‘‘tail’’ of the element T_{A^*} by setting

$$\Sigma_A := \Sigma_{\mathcal{D}_{\nu_A} \cap \mathfrak{S}_{\text{co}(A)}} = \sum_{\sigma \in \mathcal{D}_{\nu_A} \cap \mathfrak{S}_{\text{co}(A)}} T_\sigma. \quad (2.2.13)$$

We also set

$$c'_{A^*} = (c_\nu^\alpha)', \quad p(A^*) := p(T_{A^*}) \equiv |A^{\bar{1}}| \pmod{2}. \quad (2.2.14)$$

Proposition 2.3 ([18, Prop. 5.2, Th. 5.3]). (1) For any $\lambda, \mu \in \Lambda(n, r)$, the intersection $x_\lambda \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$ is a free R -module with basis $\{T_{A^*} \mid A^* \in M_n(\mathbb{N}|\mathbb{N}_2)_r, \lambda = \text{ro}(A), \mu = \text{co}(A)\}$.

(2) The queer q -Schur superalgebra $\mathcal{Q}_q(n, r; R)$ is a free R -supermodule with basis $\{\phi_{A^*} \mid A^* \in M_n(\mathbb{N}|\mathbb{N}_2)_r\}$, where ϕ_{A^*} is defined by

$$\phi_{A^*} : \bigoplus_{\lambda \in \Lambda(n, r)} x_\lambda \mathcal{H}_{r,R}^c \longrightarrow \bigoplus_{\lambda \in \Lambda(n, r)} x_\lambda \mathcal{H}_{r,R}^c, \quad x_\mu h \longmapsto \delta_{\mu, \text{co}(A)} T_{A^*} h, \quad \forall \mu \in \Lambda(n, r), h \in \mathcal{H}_{r,R}^c,$$

where A is the base of A^* . Moreover, ϕ_{A^*} has the parity $p(A^*)$.

We have, for $A^*, B^* \in M_n(\mathbb{N}|\mathbb{N}_2)_r$ with base A, B , respectively,

$$\phi_{B^*} \phi_{A^*} \neq 0 \implies \text{co}(B) = \text{ro}(A). \quad (2.3.1)$$

By (2.2.12), if we write $T_{A^*} = x_{\text{ro}(A)} h'$, where $h' = T_{d_A} c_{A^*} \Sigma_A$, then h' is a homogeneous element in the superalgebra $\mathcal{H}_{r,R}^c$ with parity $p(h') = p(A^*)$ and the structure constants $\gamma_{B^*, A^*}^{M^*}$ appearing in $\phi_{B^*} \phi_{A^*} = \sum_{M^*} \gamma_{B^*, A^*}^{M^*} \phi_{M^*}$ are determined by writing $T_{B^*} h' \in x_{\text{ro}(B)} \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_{\text{co}(A)}$ as a linear combination of T_{M^*} 's. In other words,

$$z = \phi_{B^*} \phi_{A^*} (x_{\text{co}(A)}) = T_{B^*} h' = \sum_{M^* \in M_n(\mathbb{N}|\mathbb{N}_2)_r} \gamma_{B^*, A^*}^{M^*} T_{M^*}. \quad (2.3.2)$$

Moreover, we have $p(M^*) = p(A^*) + p(B^*)$, for every M^* with $\gamma_{B^*, A^*}^{M^*} \neq 0$.

²The set $M_n(\mathbb{N}|\mathbb{N}_2)$ is denoted $M_n(\mathbb{N}|\mathbb{Z}_2)$ in [18].

Corollary 2.4. *Let $\mathcal{S}_q(n, r; R)$ be the R -submodule spanned by $\{\phi_{(A|O)} \mid A \in M_n(\mathbb{N})_r\}$. Then $\mathcal{S}_q(n, r; R)$ is a subalgebra of $\mathcal{Q}_q(n, r; R)$ (with the same identity). Moreover, $\mathcal{S}_q(n, r; R)$ is isomorphic to the q -Schur algebra (of type A).*

3. SOME COMMUTATION RELATIONS IN THE HECKE-CLIFFORD SUPERALGEBRA

In this section, we establish a number of commutation relations in $\mathcal{H}_{r,R}^c$ which will be useful to compute the multiplication formulas in the queer q -Schur superalgebra.

Let $E_{i,j} \in M_n(\mathbb{N})$ denote the matrix unit with entry 1 at (i, j) position and 0 elsewhere. For any $A = (a_{i,j}) \in M_n(\mathbb{N})_r$ and $h, k \in [1, n]$, $h < n$, define matrices in $M_n(\mathbb{N})_r$:

$$\begin{aligned} A_{h,k}^+ &:= A + E_{h,k} - E_{h+1,k}, \text{ if } a_{h+1,k} > 0; \\ A_{h,k}^- &:= A - E_{h,k} + E_{h+1,k}, \text{ if } a_{h,k} > 0. \end{aligned} \quad (3.0.1)$$

As shown in [14, Prop. 3.6], explicit relations between the distinguished double coset representatives d_A and $d_{A_{h,k}^+}$, $d_{A_{h,k}^-}$ are given via their reduced expressions. In fact, such matrices occur naturally in the product $\phi_{X^*} \phi_{A^*}$ for the X^* mentioned in the introduction.

We further define the partial sums of row h of A by setting $\overleftarrow{\mathbf{r}}_h^1(A) = 0$, $\overrightarrow{\mathbf{r}}_h^n(A) = 0$, and

$$\begin{aligned} \overleftarrow{\mathbf{r}}_h^k(A) &:= \sum_{u=1}^{k-1} a_{h,u}, \text{ for } 2 \leq k \leq n+1; \\ \overrightarrow{\mathbf{r}}_h^k(A) &:= \sum_{j=k+1}^n a_{h,j}, \text{ for } 0 \leq k \leq n-1. \end{aligned} \quad (3.0.2)$$

Note that $\overleftarrow{\mathbf{r}}_h^{n+1}(A) = \text{ro}(A)_h$, the h -th component of $\text{ro}(A)$, and $\overrightarrow{\mathbf{r}}_h^0(A) = \text{ro}(A)_h$.

For $A = (a_{i,j}) \in M_n(\mathbb{N})$ with $\lambda = \text{ro}(A)$, $r = |A|$, $\overleftarrow{\mathbf{r}}_h^k = \overleftarrow{\mathbf{r}}_h^k(A)$, let

$$\widehat{A} = (\widehat{a}_{i,j}), \text{ where } \widehat{a}_{i,j} = \widetilde{\lambda}_{i-1} + \overleftarrow{\mathbf{r}}_i^{j+1}, \text{ and set } \widehat{a}_{1,0} = 0, \widehat{a}_{i,0} = \widehat{a}_{i-1,n} (i > 1). \quad (3.0.3)$$

Observe that the matrix \widehat{A} and the matrix \widetilde{A} defined in (2.2.4) are related by ${}^t\widetilde{A} = {}^t(\widehat{A})$, where tB is the transpose of B .

The following corner sub-matrices will be useful later on.

Definition 3.1. For $A = (a_{i,j}) \in M_n(\mathbb{N})_r$ and $h, k \in [1, n]$, deleting h th row and k th column divides the matrix into four submatrices in general. We will call them, respectively, *the upper left/right corner matrix* and *the lower left/right corner matrix* at (h, k) . We denote the upper left/right corner matrix at (h, k) by $A_{h,k}^\downarrow/A_{h,k}^\uparrow$, and the lower left/right corner matrix at (h, k) by $A_{\downarrow}^{h,k}/A_{\uparrow}^{h,k}$.

We also set $A_{1,k}^\downarrow = 0$, $A_{h,n}^\uparrow = 0$, $A_{\downarrow}^{h,1} = 0$, and $A_{\uparrow}^{n,k} = 0$, etc.

We will use the lower left corner matrices to characterise a SDP condition in Section 4.

We first describe a reduced expression of d_A ; see [14, Lemma 3.2]. For $\lambda = \text{ro}(A)$, $\mu = \text{co}(A)$, define the ‘‘hook sums’’ matrix $\sigma(A) = (\sigma_{i,j})$ of A , where, for $i, j \in [1, n]$,

$$\sigma_{i,j} := \widetilde{\mu}_{j-1} + \sum_{u \leq i, t \geq j} a_{u,t} = \widetilde{\lambda}_i + \sum_{u > i, t < j} a_{u,t}. \quad (3.1.1)$$

(Recall the $\widetilde{\cdot}$ notation in (2.2.2) and (2.2.4) and see Figure 1 below for a depiction of $\sigma_{h-1,k}$.)

For $i \in [2, n]$, $j \in [1, n-1]$, define $w_{i,j} \in \mathfrak{S}_r$ by


$$w_{i,j} = \begin{cases} 1, & \text{if } a_{i,j} = 0, \text{ or } a_{i,j} > 0 \text{ and } \sigma_{i-1,j} = \widetilde{a}_{i-1,j}, \\ w_{i,j}^{(1)} w_{i,j}^{(2)} \cdots w_{i,j}^{(a_{i,j})}, & \text{if } a_{i,j} > 0 \text{ and } \sigma_{i-1,j} > \widetilde{a}_{i-1,j}, \end{cases} \quad (3.1.2)$$

where $w_{i,j}^{(k)} = s_{\sigma_{i-1,j+k-1}} s_{\sigma_{i-1,j+k-2}} \cdots s_{\widetilde{a}_{i-1,j+k}}$ for $k \in [1, a_{i,j}]$.

Note that, for $h \in [2, n], k \in [1, n-1]$, the two corner matrices $A_{\uparrow}^{h,k}$ and $A_{h,k}^{\downarrow}$ play interesting roles in linking the entries of $\sigma(A)$, \tilde{A} , and \hat{A} in the following way:

$$\begin{aligned}\sigma_{h-1,k} &= \tilde{a}_{h-1,k} + \sum_{s < h, t > k} a_{s,t} = \tilde{a}_{h-1,k} + |A_{h,k}^{\downarrow}| \\ &= \tilde{\lambda}_{h-1} + \overleftarrow{\mathbf{r}}_h^k + \sum_{i > h, j < k} a_{i,j} = \hat{a}_{h,k-1} + |A_{\uparrow}^{h,k}|.\end{aligned}\quad (3.1.3)$$

We use a picture (Figure 1) to indicate the various entry positions $(h-1, k) = \bullet$, $(h, k-1) = *$ and $(h, k) = \circ$. The two grey areas represent the submatrices $A_{h,k}^{\downarrow}$ and $A_{\uparrow}^{h,k}$. (Recall Definition 3.1 for corner matrices.)



(Figure 1)

Thus, each $w_{i,j}^{(k)}$ has length $\ell(w_{i,j}^{(k)}) = |A_{i,j}^{\downarrow}|$. By [14, Lem. 3.2], we have

$$d_A = (w_{2,1}w_{3,1} \cdots w_{n,1})(w_{2,2}w_{3,2} \cdots w_{n,2}) \cdots (w_{2,n-1}w_{3,n-1} \cdots w_{n,n-1}), \quad (3.1.4)$$

and (cf. [8, Exer. 8.2])

$$\ell(d_A) = \sum_{1 \leq i, j \leq n} a_{i,j} |A_{i,j}^{\downarrow}|.$$

Remark 3.2. (1) We remark that the condition $\sigma_{i-1,j} = \tilde{a}_{i-1,j}$ (i.e., $A_{i,j}^{\downarrow} = 0$) in (3.1.2) is trivially true for $i = 1$ or $j = n$. Thus, we may extend the definition of $w_{i,j}$ to include $w_{1,j} = 1 = w_{i,n}$, for all $i, j \in [1, n]$. Thus, we may depict d_A as the $n \times n$ matrix $w(A) = (w_{i,j})_{1 \leq i, j \leq n}$ and regard d_A as the product down column 1 of $w(A)$, then column 2 of $w(A)$, and so on. We may identify d_A as $w(A)$.

(2) Observe the following:

$$\begin{aligned}d_A = 1 &\iff \sigma_{i-1,j} - \tilde{a}_{i-1,j} = 0, \text{ for all } 2 \leq i \leq n, 1 \leq j \leq n-1 \text{ with } a_{i,j} > 0; \\ &\iff A_{i,j}^{\downarrow} = 0, \text{ for all } 2 \leq i \leq n, 1 \leq j \leq n-1 \text{ with } a_{i,j} > 0.\end{aligned}$$

(3) For convenience of later use, if $w_{i,j} \neq 1$, we may display all the simple reflections s_k in $w_{i,j}$ as follows:

$$\begin{aligned}s_{\sigma_{i-1,j}} s_{\sigma_{i-1,j-1}} \cdots s_{\tilde{a}_{i-1,j}+1} \\ s_{\sigma_{i-1,j}+1} s_{\sigma_{i-1,j}} \cdots s_{\tilde{a}_{i-1,j}+2} \\ \dots \dots \dots \\ s_{\sigma_{i-1,j}+a_{i,j}-1} s_{\sigma_{i-1,j}+a_{i,j}-2} \cdots s_{\tilde{a}_{i-1,j}+a_{i,j}}.\end{aligned}\quad (3.2.1)$$

If s_k appears in the display, then we write $s_k|w_{i,j}$. If w is a product of some $w_{i,j}$ and $w \neq 1$, then $s_k|w$ is well defined. For such a w , define

$$\max(w) := \max\{k \mid k \in [1, r], s_k|w\}, \quad \min(w) := \min\{k \mid k \in [1, r], s_k|w\}. \quad (3.2.2)$$

We see easily from (3.2.1) that if $w_{i,j} \neq 1$, then

$$\max(w_{i,j}) = \sigma_{i-1,j} + a_{i,j} - 1 \quad \text{and} \quad \min(w_{i,j}) = \tilde{a}_{i-1,j} + 1. \quad (3.2.3)$$

Call w a *segment* of d_A if w is a product from $w_{i,j}$ to $w_{i',j'}$ in (3.1.4). In the sequel, we often write $d_A = \overleftarrow{w} \cdot w \cdot \overrightarrow{w}$, where \overleftarrow{w} (resp., \overrightarrow{w}) is the segment of d_A before (resp., after) w .

Lemma 3.3. (1) If $w = w_{i,j} \cdots w_{h,k}$ is a segment of d_A and $w \neq 1$, then $\min(w) \geq \tilde{a}_{i-1,j} + 1$.
(2) If $y = w(A_{h+1,k+1}^{\downarrow}) := (w_{1,1}w_{2,1}w_{3,1} \cdots w_{h,1}) \cdots (w_{1,k}w_{2,k}w_{3,k} \cdots w_{h,k})$ and $w \neq 1$, then $\max(y) \leq \sigma_{h-1,k} + a_{h,k} - 1$.

Proof. (1) is clear since $\min(w) = \min(w_{i',j'}) = \tilde{a}_{i'-1,j'} + 1$, where $w_{i',j'}$ is the first non-identity $w_{s,t}$ appearing in w , and $\tilde{a}_{i'-1,j'} \geq \tilde{a}_{i-1,j}$ whenever $j' > j$ or $j' = j$ but $i' > i$.

(2) follows from the fact that $\sigma_{i-1,j} + a_{i,j} - 1 \leq \sigma_{h-1,k} + a_{h,k} - 1$ whenever $i \leq h$ and $j \leq k$. \square

We are now ready to look at some commutation relations in $\mathcal{H}_{r,R}^c$. The first involves the distinguished representatives d_A , $d_{A_{h,k}^+}$, and $d_{A_{h,k}^-}$. It tells how to turn an expression involving T_{d_A} to an expression involving $T_{d_{A_{h,k}^+}}$ or $T_{d_{A_{h,k}^-}}$.

The following elements in $\mathcal{H}_{r,R}^c$ will be frequently used later on in this paper. For any $1 \leq i, j \leq r-1$, define

$$\begin{aligned} T_{(i,j)}^{\triangleleft} &= \begin{cases} 1 + T_i + T_i T_{i+1} + \cdots + T_i T_{i+1} \cdots T_j, & \text{if } i \leq j; \\ 1, & \text{otherwise,} \end{cases} \\ T_{(j,i)}^{\triangleright} &= \begin{cases} 1 + T_j + T_j T_{j-1} + \cdots + T_j T_{j-1} \cdots T_i, & \text{if } j \geq i; \\ 1, & \text{otherwise.} \end{cases} \end{aligned} \quad (3.3.1)$$

For example, in $\mathcal{H}(\mathfrak{S}_{r+1})_R$, $\Sigma_{\mathcal{D}(1,r)} = T_{(1,r)}^{\triangleleft}$ and $\Sigma_{\mathcal{D}(r,1)} = T_{(r,1)}^{\triangleright}$.

Lemma 3.4. For $A = (a_{i,j}) \in M_n(\mathbb{N})$, let $\lambda = \text{ro}(A)$, $\overleftarrow{\mathbf{r}}_h^k = \overleftarrow{\mathbf{r}}_h^k(A)$, and $\overrightarrow{\mathbf{r}}_h^k = \overrightarrow{\mathbf{r}}_h^k(A)$.

(1) For each $1 \leq h \leq n-1$ and $1 \leq k \leq n$ such that $a_{h+1,k} > 0$, we have

$$\sum_{j=\overleftarrow{\mathbf{r}}_{h+1}^k}^{\overleftarrow{\mathbf{r}}_{h+1}^{k+1}-1} T_{\tilde{\lambda}_{h+1}}^- T_{\tilde{\lambda}_{h+2}}^- \cdots T_{\tilde{\lambda}_{h+j}}^- T_{d_A} = T_{\tilde{\lambda}_h}^- T_{\tilde{\lambda}_{h-1}}^- \cdots T_{\tilde{\lambda}_h - \overrightarrow{\mathbf{r}}_{h+1}^k}^- T_{d_{A_{h,k}^+}} T_{(\tilde{a}_{h,k}+1, \tilde{a}_{h,k}+a_{h+1,k}-1)}^{\triangleleft}.$$

(2) For each $1 \leq h \leq n$ and $1 \leq k \leq n$ such that $a_{h,k} > 0$, we have

$$\sum_{j=\overrightarrow{\mathbf{r}}_h^k}^{\overrightarrow{\mathbf{r}}_h^{k-1}-1} T_{\tilde{\lambda}_{h-1}}^- \cdots T_{\tilde{\lambda}_{h-j}}^- T_{d_A} = T_{\tilde{\lambda}_h}^- T_{\tilde{\lambda}_{h+1}}^- \cdots T_{\tilde{\lambda}_h + \overleftarrow{\mathbf{r}}_{h+1}^k}^- T_{d_{A_{h,k}^-}} T_{(\tilde{a}_{h,k}-1, \tilde{a}_{h,k}-a_{h,k}+1)}^{\triangleright}.$$

Proof. We only prove (1), and the proof of (2) is similar. For each $j \in [\overleftarrow{\mathbf{r}}_{h+1}^k, \overleftarrow{\mathbf{r}}_{h+1}^{k+1} - 1]$ and $p_j = j - \overleftarrow{\mathbf{r}}_{h+1}^k$, according to [14, Proposition 3.6(1)], we have

$$s_{\tilde{\lambda}_h+1}^- s_{\tilde{\lambda}_{h+2}}^- \cdots s_{\tilde{\lambda}_{h+j}}^- d_A = s_{\tilde{\lambda}_h}^- s_{\tilde{\lambda}_{h-1}}^- \cdots s_{\tilde{\lambda}_h - \overrightarrow{\mathbf{r}}_{h+1}^k}^- d_{A_{h,k}^+} s_{\tilde{a}_{h,k}+1}^- \cdots s_{\tilde{a}_{h,k}+p_j}^-.$$

Since both sides are reduced expressions, it follows that

$$T_{\tilde{\lambda}_{h+1}}^- T_{\tilde{\lambda}_{h+2}}^- \cdots T_{\tilde{\lambda}_{h+j}}^- T_{d_A} = T_{\tilde{\lambda}_h}^- T_{\tilde{\lambda}_{h-1}}^- \cdots T_{\tilde{\lambda}_h - \overrightarrow{\mathbf{r}}_{h+1}^k}^- T_{d_{A_{h,k}^+}} T_{\tilde{a}_{h,k}+1}^- \cdots T_{\tilde{a}_{h,k}+p_j}^-.$$

Hence, (1) follows. \square

The next relations show how to interchange the tails Σ_A and $\Sigma_{A_{h,k}^+}$ or Σ_A and $\Sigma_{A_{h,k}^-}$, where $A_{h,k}^+$ and $A_{h,k}^-$ are defined in (3.0.1) and their tails in (2.2.13).

Lemma 3.5. Let $h \in [1, n-1]$, $k \in [1, n]$, and $A = (a_{i,j}) \in M_n(\mathbb{N})_r$. Then we have

$$(1) \quad T_{(\tilde{a}_{h,k}+1, \tilde{a}_{h,k}+a_{h+1,k}-1)}^{\triangleleft} \Sigma_A = T_{(\tilde{a}_{h,k}, \tilde{a}_{h,k}-a_{h,k}+1)}^{\triangleright} \Sigma_{A_{h,k}^+}, \quad \text{if } a_{h+1,k} \geq 1;$$

$$(2) \quad T_{(\tilde{a}_{h,k}-1, \tilde{a}_{h,k}-a_{h,k}+1)}^{\triangleright} \Sigma_A = T_{(\tilde{a}_{h,k}, \tilde{a}_{h,k}+a_{h+1,k}-1)}^{\triangleleft} \Sigma_{A_{h,k}^-}, \quad \text{if } a_{h,k} \geq 1.$$

Proof. Let $\mu = \text{co}(A) = \text{co}(A_{h,k}^+)$, $\nu = \nu_A$, $\nu^+ = \nu_{A_{h,k}^+}$, and $\mathfrak{S}_{\delta^+} = \mathfrak{S}_{\nu} \cap \mathfrak{S}_{\nu^+}$ with δ^+ as given in (5.2.2). Then, for $a_{h+1,k} \geq 1$,

$$\mathcal{D}_{\delta^+} \cap \mathfrak{S}_{\mu} = (\mathcal{D}_{\delta^+} \cap \mathfrak{S}_{\nu})(\mathcal{D}_{\nu} \cap \mathfrak{S}_{\mu}) = (\mathcal{D}_{\delta^+} \cap \mathfrak{S}_{\nu^+})(\mathcal{D}_{\nu^+} \cap \mathfrak{S}_{\mu}),$$

where

$$\begin{aligned} \mathcal{D}_{\delta^+} \cap \mathfrak{S}_{\nu} &= \{1, s_{\tilde{a}_{h,k}+1}^-, s_{\tilde{a}_{h,k}+1}^- s_{\tilde{a}_{h,k}+2}^-, \cdots, s_{\tilde{a}_{h,k}+1}^- \cdots s_{\tilde{a}_{h,k}+a_{h+1,k}-1}^-\} \\ \mathcal{D}_{\delta^+} \cap \mathfrak{S}_{\nu^+} &= \{1, s_{\tilde{a}_{h,k}}^-, s_{\tilde{a}_{h,k}}^- s_{\tilde{a}_{h,k}-1}^-, \cdots, s_{\tilde{a}_{h,k}}^- \cdots s_{\tilde{a}_{h,k}-a_{h,k}+1}^-\}. \end{aligned}$$

Hence, by length additivity,

$$T_{(\tilde{a}_{h,k+1}, \tilde{a}_{h,k+a_{h+1,k-1}})}^{\triangleleft} \Sigma_A = \Sigma_{\mathcal{D}_{\delta+n} \mathfrak{S}_{\nu}} \Sigma_{\mathcal{D}_{\nu} \mathfrak{S}_{\mu}} = \Sigma_{\mathcal{D}_{\delta+n} \mathfrak{S}_{\nu+}} \Sigma_{\mathcal{D}_{\nu+} \mathfrak{S}_{\mu}} = T_{(\tilde{a}_{h,k}, \tilde{a}_{h,k-a_{h,k+1}})}^{\triangleright} \Sigma_{A_{h,k}^+},$$

proving assertion (1). If $B = A_{h,k}^+$, then $b_{h,k} > 0$ and $B_{h,k}^- = A$. Hence, assertion (2) follows from (1). \square

Lemma 3.6. *For $\alpha \in \Lambda(n, r)$, $1 \leq u < r$, and $j \geq 0$, assume $s_{u+1}, \dots, s_{u+j} \in \mathfrak{S}_{\alpha}$. Then $x_{\alpha} T_{(u, u+j)}^{\triangleleft} = x_{\alpha} T_{(u+j, u)}^{\triangleright} = \llbracket j+2 \rrbracket_q x_{\alpha}$ if, in addition, $s_u \in \mathfrak{S}_{\alpha}$. Moreover, the following holds:*

$$x_{\alpha} T_u^{-1} T_{u+1}^{-1} \cdots T_{u+j}^{-1} = q^{-1-j} x_{\alpha} T_u T_{u+1} \cdots T_{u+j} - q^{-j-1} (q-1) x_{\alpha} T_{(u, u+j-1)}^{\triangleleft}.$$

Proof. Assertion (1) is clear by (2.1.5) and (3.3.1), while assertion (2) can be proved by induction on j , noting (2.0.4)(1). \square

Finally, we look at some commutation relations involving odd generators c_i . Recall from (2.2.5) the elements $c_{q,i,j}$.

Lemma 3.7. *For $\alpha \in \Lambda(n, r)$, $1 \leq u < r$, and $j \geq 0$, assume $s_{u+1}, \dots, s_{u+j} \in \mathfrak{S}_{\alpha}$ if $j \geq 1$.*

(1) *For $j \geq 0$, we have in $\mathcal{H}_{r,R}^c$*

$$x_{\alpha} c_u T_u T_{u+1} \cdots T_{u+j} = q^{j+1} x_{\alpha} T^{-1}(u, j+1) c_{u+j+1} + (q-1) q^j x_{\alpha} \sum_{k=0}^j T^{-1}(u, k) c_{u+k}$$

$$\text{where } T^{-1}(u, k) = \begin{cases} 1, & \text{if } k = 0; \\ T_u^{-1} T_{u+1}^{-1} T_{u+2}^{-1} \cdots T_{u+k-1}^{-1}, & \text{if } k \in [1, j]. \end{cases}$$

(2) *Further, we have*

$$x_{\alpha} c_u T_{(u, u+j)}^{\triangleleft} = q^{j+1} x_{\alpha} (c_u + T_u^{-1} c_{u+1} + T_u^{-1} T_{u+1}^{-1} c_{u+2} + \cdots + T_u^{-1} T_{u+1}^{-1} T_{u+2}^{-1} \cdots T_{u+j}^{-1} c_{u+j+1}).$$

If, in addition, $s_u \in \mathfrak{S}_{\alpha}$, then $x_{\alpha} c_u T_{(u, u+j)}^{\triangleleft} = x_{\alpha} c_{q, u, u+j+1}$.

(3) *If $s_u, s_{u-1}, \dots, s_{u-j} \in \mathfrak{S}_{\alpha}$, then $x_{\alpha} c_{u+1} T_{(u, u-j)}^{\triangleright} = x_{\alpha} c_{q, u-j, u+1}$.*

(4) *For $1 \leq i \leq j \leq r$, $(c_{q,i,j})^2 = -\llbracket j-i+1 \rrbracket_{q^2}$.*

Proof. We first prove (1). Repeatedly applying (2.0.4)(3) gives

$$\begin{aligned} & x_{\alpha} c_u T_u T_{u+1} \cdots T_{u+j} \\ &= x_{\alpha} (q T_u^{-1} c_{u+1} + (q-1) c_u) T_{u+1} \cdots T_{u+j} \\ &= q x_{\alpha} T_u^{-1} c_{u+1} T_{u+1} \cdots T_{u+j} + (q-1) q^j x_{\alpha} c_u \\ &= q x_{\alpha} T_u^{-1} (q T_{u+1}^{-1} c_{u+2} + (q-1) c_{u+1}) T_{u+2} \cdots T_{u+j} + (q-1) q^j x_{\alpha} c_u \\ &= q^2 x_{\alpha} T_u^{-1} T_{u+1}^{-1} c_{u+2} T_{u+2} \cdots T_{u+j} + (q-1) q^j x_{\alpha} (c_u + T_u^{-1} c_{u+1}) \\ &= \cdots \\ &= q^{j+1} x_{\alpha} T_u^{-1} T_{u+1}^{-1} T_{u+2}^{-1} \cdots T_{u+j}^{-1} c_{u+j+1} \\ &\quad + (q-1) q^j x_{\alpha} (c_u + T_u^{-1} c_{u+1} + T_u^{-1} T_{u+1}^{-1} c_{u+2} + \cdots + T_u^{-1} T_{u+1}^{-1} \cdots T_{u+j-1}^{-1} c_{u+j}), \end{aligned}$$

proving (1). Now, by (1), we have

$$\begin{aligned} x_{\alpha} c_u T_{(u, u+j)}^{\triangleleft} &= x_{\alpha} c_u + \sum_{i=0}^j x_{\alpha} c_u T_u \cdots T_{u+i} \\ &= x_{\alpha} c_u + \sum_{i=0}^j ((q-1) q^i x_{\alpha} \sum_{k=0}^i T^{-1}(u, k) c_{u+k} + q^{i+1} x_{\alpha} T^{-1}(u, i+1) c_{u+i+1}) \\ &= \sum_{k=0}^j (q^k + \sum_{i=k}^j (q-1) q^i) x_{\alpha} T^{-1}(u, k) c_{u+k} + q^{j+1} x_{\alpha} T^{-1}(u, j+1) c_{u+j+1} \\ &= q^{j+1} x_{\alpha} (c_u + T_u^{-1} c_{u+1} + T_u^{-1} T_{u+1}^{-1} c_{u+2} + \cdots + T_u^{-1} T_{u+1}^{-1} T_{u+2}^{-1} \cdots T_{u+j}^{-1} c_{u+j+1}), \end{aligned}$$

since the telescoping sum $q^k + \sum_{i=k}^j (q-1)q^i = q^{j+1}$. This proves the first assertion of (2). The second assertion in (2) is clear.

For assertion (3), $s_{u-i} \in \mathfrak{S}_\alpha$ implies $x_\alpha T_{u-i} = qx_\alpha$ for $0 \leq i \leq j$. Thus, by (2.0.3), we obtain

$$\begin{aligned} x_\alpha c_{u+1} T_{(u,u-j)}^\triangleright &= x_\alpha c_{u+1} (1 + T_u + T_u T_{u-1} + \cdots + T_u T_{u-1} \cdots T_{u-j}) \\ &= x_\alpha (c_{u+1} + T_u c_u + T_u T_{u-1} c_{u-1} + \cdots + T_u T_{u-1} \cdots T_{u-j} c_{u-j}) \\ &= x_\alpha (c_{u+1} + qc_u + \cdots + q^{j+1} c_{u-j}) = x_\alpha c_{q,u-j,u+1}, \end{aligned}$$

proving (3).

Finally, (4) is straightforward by an induction. \square

4. DISTINGUISHED DOUBLE COSET REPRESENTATIVES AND THE SDP CONDITION

Recall from (2.2.8) the distinguished double coset representative $d_A \in \mathfrak{S}_r$ associated with $A \in M_n(\mathbb{N})_r$. To compute some structure constants in (2.2.12), especially in the odd case, we need to impose some commutation relations between T_{d_A} and certain c -elements. The SDP condition introduced in this section is convenient for the purpose.

This element d_A has a reduced expression as given in (3.1.4). As a permutation, d_A is given in [9, p. 424] (or [8, Ex. 8.2]). More precisely, for $A = (a_{i,j}) \in M_n(\mathbb{N})$ with $\tilde{A} = (\tilde{a}_{i,j})$ and $\hat{A} = (\hat{a}_{i,j})$ as in (2.2.4) and (3.0.3), the description of the pseudo-matrix \ddot{A} associated with A in loc. cit. (see Example 4.1 below) can be easily interpreted as follow:

$$d_A(\tilde{a}_{h-1,k} + p) = \hat{a}_{h,k-1} + p, \quad (4.0.1)$$

for all $h, k \in [1, n]$ with $a_{h,k} > 0$ and $p \in [1, a_{h,k}]$. In other words, d_A is the permutation obtained by concatenating the permutations

$$\left(\begin{array}{cccc} \tilde{a}_{i-1,j} + 1 & \tilde{a}_{i-1,j} + 2 & \cdots & \tilde{a}_{i-1,j} + a_{i,j} \\ \hat{a}_{i,j-1} + 1 & \hat{a}_{i,j-1} + 2 & \cdots & \hat{a}_{i,j-1} + a_{i,j} \end{array} \right) \quad (4.0.2)$$

for all $a_{i,j} > 0$ via the ordering from top to bottom of column 1, then top to bottom of column 2, and so on.

Example 4.1. Let $A = \begin{pmatrix} 0 & 3 & 2 \\ 1 & 1 & 0 \\ 2 & 0 & 0 \end{pmatrix} \in M_3(\mathbb{N})_9$. Then

$$d_A = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 6 & 8 & 9 & 1 & 2 & 3 & 7 & 4 & 5 \end{pmatrix}.$$

The images $d_A(1), d_A(2), \dots, d_A(9)$ of d_A (i.e., the second row in d_A) is obtained as follows:

Form the pseudo-matrix $\ddot{A} = \begin{pmatrix} - & (1,2,3) & (4,5) \\ 6 & 7 & - \\ (8,9) & - & - \end{pmatrix}$ obtained by replacing each nonzero entry

$a_{i,j}$ by the second row of (4.0.2) — a sequence of length $a_{i,j}$. In other words, \ddot{A} is obtained by filling $1, 2, \dots, 9$ along each row from left to right (using the sequence length $a_{i,j}$), and then down successive rows. Now, the sequence $d_A(1), \dots, d_A(9)$ is obtained by reading the numbers from left to right inside each sequence associated with $a_{i,j}$ down column 1, then column 2, and so on.

By specialising q to 1, $\mathfrak{H}_r^c = \mathcal{H}_{r,R}^c|_{q=1}$ is the Sergeev algebra (see [27]), which is the “semi-direct (or smash) product” $\mathcal{C}_r \rtimes R\mathfrak{S}_r$. (See [4] for another such a kind of example.) Then, we have $wc_i = c_{w(i)}w$ in \mathfrak{H}_r^c and, in particular,

$$d_A c_{\tilde{a}_{h-1,k}+p} = c_{\hat{a}_{h,k-1}+p} d_A$$

for all $1 \leq h, k \leq n$ with $a_{h,k} > 0$ and $1 \leq p \leq a_{h,k}$. However, if we replace d_A by T_{d_A} , this commutation formula may not hold in the Hecke-Clifford superalgebra $\mathcal{H}_{r,R}^c$. So we introduce the following definition.

Definition 4.2. Let $A = (a_{i,j}) \in M_n(\mathbb{N})$. If $a_{h,k} > 0$ and

$$c_{\widehat{a}_{h,k-1}+p} T_{d_A} = T_{d_A} c_{\widetilde{a}_{h-1,k}+p} \quad (\text{in } \mathcal{H}_{r,R}^c)$$

for each $p \in [1, a_{h,k}]$, then A is said to satisfy the *semi-direct product (SDP) condition* at (h, k) .

If A satisfies the SDP condition at (h, k) for every $k \in [1, n]$ (resp., $h \in [1, n]$) with $a_{h,k} > 0$, then A is said to satisfy the SDP condition on the h th row (resp., k th column).

Using the element defined in (2.2.5), A satisfied the SDP condition at (h, k) implies

$$c_{q, \widehat{a}_{h,k-1}+1, \widehat{a}_{h,k-1}+a_{h,k}} T_{d_A} = T_{d_A} c_{q, \widetilde{a}_{h-1,k}+1, \widetilde{a}_{h-1,k}+a_{h,k}}. \quad (4.2.1)$$


Example 4.3. For the given A as above, if $d_A = 1$, then, for any $h, k \in [1, n]$ satisfying $a_{h,k} > 0$, the equation (4.0.1) implies $\widehat{a}_{h,k-1} + p = \widetilde{a}_{h-1,k} + p$. Hence, A satisfies the SDP condition everywhere (i.e., at any (h, k) with $a_{h,k} > 0$).

In particular, if $A = (a_{i,j})$ is one of the following matrices in $M_n(\mathbb{N})_r$:

$$\text{diag}(\lambda) + \sum_{i=1}^{n-1} u_i E_{i,i+1}, \quad \text{diag}(\lambda) + \sum_{i=1}^{n-1} u_i E_{i+1,i},$$

where $\lambda \in \mathbb{N}^n$ and $\mathbf{u} = (u_1, \dots, u_{n-1}) \in \mathbb{N}^{n-1}$, then $d_A = 1$, and A satisfies the SDP condition everywhere.

In general, one checks easily by Remark 3.2(2) (cf. Example 4.1) that $d_A = 1$ if and only if the nonzero entries of A are on a zig-zag line of the shape



Remark 4.4. The SDP condition on the base $A = A^{\bar{0}} + A^{\bar{1}}$ of $A^* = (A^{\bar{0}} | A^{\bar{1}})$ makes the computation for some of the multiplication formulas in Sections 5 and 6 possible. Interestingly, these multiplication formulas associated with matrices that satisfy some SDP conditions play a key role for the establishment of the BLM type realisation for quantum queer supergroups.

We now derive an equivalent condition for a matrix to satisfy the SDP condition at (h, k) .

Lemma 4.5. (1) If $1 \leq i \leq t \leq j < n$, then we have

$$c_t T_j T_{j-1} \cdots T_i = T_j T_{j-1} \cdots T_i c_{t+1} + (q-1) T_j \cdots T_{t+1} T_{t-1} \cdots T_i (c_i - c_{t+1}).$$

(2) Let $A = (a_{i,j}) \in M_n(\mathbb{N})$. If $a_{h,k} > 0$, then there are some $f_{w,i}^{\cdots} \in R$ such that, for all $p \in [1, a_{h,k}]$, we have in $\mathcal{H}_{r,R}^c$

$$c_{\widehat{a}_{h,k-1}+p} T_{d_A} = T_{d_A} c_{\widetilde{a}_{h-1,k}+p} + \sum_{w < d_A, i \in [1, r]} f_{w,i}^{\cdots} T_w c_i.$$

Proof. According to (2.0.3), we have

$$\begin{aligned} c_t T_j T_{j-1} \cdots T_i &= T_j \cdots T_{t+1} c_t T_t T_{t-1} \cdots T_i \\ &= T_j \cdots T_{t+1} (T_t c_{t+1} + (q-1)(c_t - c_{t+1})) T_{t-1} \cdots T_i \\ &= T_j \cdots T_{t+1} (T_t c_{t+1}) T_{t-1} \cdots T_i + T_j \cdots T_{t+1} ((q-1)(c_t - c_{t+1})) T_{t-1} \cdots T_i \\ &= T_j T_{j-1} \cdots T_i c_{t+1} + (q-1) T_j \cdots T_{t+1} T_{t-1} \cdots T_i (c_i - c_{t+1}). \end{aligned}$$

Assertion (2) follows from (2.1.2). \square

Using the notation in (2.2.5) and (3.0.3), Lemma 4.5(2) implies

$$c_{q, \widehat{a}_{h,k-1}+1, \widehat{a}_{h,k-1}+a_{h,k}} T_{d_A} = T_{d_A} c_{q, \widetilde{a}_{h-1,k}+1, \widetilde{a}_{h-1,k}+a_{h,k}} + \sum_{w < d_A, i \in [1, r]} g_{w,i}^{\cdots} T_w c_i. \quad (4.5.1)$$

Recall the lower left corner matrix $A_{\overline{1}}^{h,k}$ defined in Definition 3.1.

Theorem 4.6. *Let $A \in M_n(\mathbb{N})$ and $h, k \in [1, n]$. Then A satisfies the SDP condition at (h, k) if and only if $a_{h,k} > 0$ and $a_{i,j} = 0$, for $i > h$ and $j < k$ (equivalently, the lower left corner matrix $A_{\overleftarrow{1}}^{h,k}$ at (h, k) is 0).*

Proof. We first recall from (3.1.4) the reduced expression of d_A :

$$d_A = (w_{2,1}w_{3,1} \cdots w_{n,1})(w_{2,2}w_{3,2} \cdots w_{n,2}) \cdots (w_{2,n-1}w_{3,n-1} \cdots w_{n,n-1}),$$

where $w_{i,j}$, for $i \in [2, n]$, $j \in [1, n-1]$, are defined in (3.1.2). Thus, if $w_{i,j} \neq 1$, then $a_{i,j} > 0$, $|A_{i,j}^{\overleftarrow{1}}| \neq 0$, and $w_{i,j} = w_{i,j}^{(1)}w_{i,j}^{(2)} \cdots w_{i,j}^{(a_{i,j})}$ has length $a_{i,j}|A_{i,j}^{\overleftarrow{1}}|$, where, for $k \in [1, a_{i,j}]$,

$$w_{i,j}^{(k)} = (s_{\sigma_{i-1,j+k-1}}s_{\sigma_{i-1,j+k-2}} \cdots s_{\tilde{a}_{i-1,j+k}}). \quad (4.6.1)$$

Moreover, we also adopt the following notational scheme when breaking d_A as a product of “segments”:

$$d_A = \overleftarrow{(d_A)}_{i,j} \cdot w_{i,j} \cdot \overrightarrow{(d_A)}_{i,j} = \overleftarrow{(d_A)}_{i,j}^{(k)} \cdot w_{i,j}^{(k)} \cdot \overrightarrow{(d_A)}_{i,j}^{(k)} \quad (4.6.2)$$

We first prove that “if” part.

Suppose $a_{h,k} > 0$ and $a_{i,j} = 0$ for all $i > h$ and $j < k$, i.e., $A_{\overleftarrow{1}}^{h,k} = 0$. We prove that A satisfies SDP condition at (h, k) : $c_{\tilde{a}_{h,k-1+p}}T_{d_A} = T_{d_A}c_{\tilde{a}_{h-1,k+p}}$, for all $p \in [1, a_{h,k}]$.

By the hypothesis, we have $w_{i,j} = 1$ for all $i > h$ and $j < k$. Thus, d_A has the form

$$d_A = (w_{2,1}w_{3,1} \cdots w_{h,1}) \cdots (w_{2,k-1}w_{3,k-1} \cdots w_{h,k-1})(w_{2,k}w_{3,k} \cdots w_{h-1,k} \cdot w_{h,k} \cdots w_{n,k}) \\ (w_{2,k+1}w_{3,k+1} \cdots w_{n,k+1}) \cdots (w_{2,n-1}w_{3,n-1} \cdots w_{n,n-1}) \quad (4.6.3)$$

Consider the maximal index defined in (3.2.2) for

$$w := (w_{2,1}w_{3,1} \cdots w_{h,1}) \cdots (w_{2,k-1}w_{3,k-1} \cdots w_{h,k-1}).$$

By Lemma 3.3(2), $m_1 = \max(w) < \sigma_{h-1,k-1} + a_{h,k-1}$. Similarly, $m_2 = \max(w_{2,k}w_{3,k} \cdots w_{h-1,k}) < \sigma_{h-2,k} + a_{h-1,k}$. Now, $A_{\overleftarrow{1}}^{h,k} = 0$ implies

$$\sigma_{h-2,k} + a_{h-1,k} = \tilde{\lambda}_{h-2} + \sum_{j \leq k} a_{h-1,j} + \sum_{j < k} a_{h,j} = \tilde{\lambda}_{h-2} + \sum_{j \leq k} a_{h-1,j} + \overleftarrow{\mathbf{r}}_h^k \\ \leq \tilde{\lambda}_{h-1} + \overleftarrow{\mathbf{r}}_h^k (= \sigma_{h-1,k-1} + a_{h,k-1}) < \tilde{\lambda}_{h-1} + \overleftarrow{\mathbf{r}}_h^k + p = \hat{a}_{h,k-1} + p,$$

for all $p \in [1, a_{h,k}]$. Hence, $\hat{a}_{h,k-1} + p > \max(m_1, m_2) + 1$. This implies, by the notation in (4.6.2), that $c_{\tilde{a}_{h,k-1+p}}T_{\overleftarrow{(d_A)}_{h,k}} = T_{\overrightarrow{(d_A)}_{h,k}}c_{\tilde{a}_{h,k-1+p}}$ and so

$$c_{\tilde{a}_{h,k-1+p}}T_{d_A} = T_{\overleftarrow{(d_A)}_{h,k}}c_{\tilde{a}_{h,k-1+p}}T_{w_{h,k}}T_{\overrightarrow{(d_A)}_{h,k}}. \quad (4.6.4)$$

We now commute $c_{\tilde{a}_{h,k-1+p}}T_{w_{h,k}}$ for a fixed $p \in [1, a_{h,k}]$. The hypothesis $A_{\overleftarrow{1}}^{h,k} = 0$ together with (3.1.3) gives

$$\sigma_{h-1,k} = \tilde{\lambda}_{h-1} + \overleftarrow{\mathbf{r}}_h^k = \hat{a}_{h,k-1}.$$

Recall that the j th factor of $w_{h,k} = w_{h,k}^{(1)}w_{h,k}^{(2)} \cdots w_{h,k}^{(a_{h,k})}$ has the form

$$w_{h,k}^{(j)} = s_{\tilde{a}_{h,k-1+j-1}}s_{\tilde{a}_{h,k-1+j-2}} \cdots s_{\tilde{a}_{h-1,k+j}}.$$

If $j < p$, the index of s_i in the expression of $w_{h,k}^{(j)}$ is smaller than $\hat{a}_{h,k-1} + p - 1$. So, $c_{\tilde{a}_{h,k-1+p}}$ commutes with these $T_{w_{h,k}^{(j)}}$. If $j = p$, repeatedly applying (2.0.3) yields

$$c_{\tilde{a}_{h,k-1+p}}T_{w_{h,k}^{(p)}} = c_{\tilde{a}_{h,k-1+p}}T_{\tilde{a}_{h,k-1+p-1}} \cdots T_{\tilde{a}_{h-1,k+p}} = T_{w_{h,k}^{(p)}}c_{\tilde{a}_{h-1,k+p}}. \quad (4.6.5)$$

Finally, if $j > p$, then $\tilde{a}_{h-1,k} + p < \tilde{a}_{h-1,k} + j$. Hence, $c_{\tilde{a}_{h-1,k+p}}$ commutes with $T_{w_{h,k}^{(j)}}$ and (4.6.4)

becomes $c_{\tilde{a}_{h,k-1+p}}T_{d_A} = T_{\overleftarrow{(d_A)}_{h,k}}T_{w_{h,k}}c_{\tilde{a}_{h-1,k+p}}T_{\overrightarrow{(d_A)}_{h,k}}$.

It remains to prove that $c_{\tilde{a}_{h-1,k+p}}T_{\overrightarrow{(d_A)}_{h,k}} = T_{\overrightarrow{(d_A)}_{h,k}}c_{\tilde{a}_{h-1,k+p}}$. This is clear, since

$$\overrightarrow{(d_A)}_{h,k} = w_{h+1,k} \cdots w_{n,k}(w_{2,k+1}w_{3,k+1} \cdots w_{n,k+1}) \cdots (w_{2,n-1}w_{3,n-1} \cdots w_{n,n-1}),$$

and Lemma 3.3(1) implies

$$\min(\overrightarrow{(d_A)}_{h,k}) \geq \tilde{a}_{h,k} + 1 > \tilde{a}_{h-1,k} + p.$$

We now prove the “only if” part. This is equivalent to proving that if $a_{h,k} > 0$ and $A_{\uparrow}^{h,k} \neq 0$, then A does not satisfy the SDP condition at (h, k) .

Suppose that there exists $a_{i,j} > 0$ for some i, j with $i > h$ and $j < k$. Then we must have $k > 1$ (and $h < n$). Set

$$t = \min\{j \mid a_{i,j} > 0, i > h, j < k\}, \quad l = \min\{i \mid a_{i,t} > 0, i > h\}.$$

Then $a_{l,t}$ is the top left most nonzero entry in the corner matrix $A_{\uparrow}^{h,k}$. Thus, $a_{i,j} = 0$ for $i > h$ and $j < t$ or $h < i < l$ and $j = t$, and so, $w_{i,j} = 1$ in these cases. Hence, d_A takes the form

$$\begin{aligned} d_A = & (w_{2,1}w_{3,1} \cdots w_{h,1}) \cdots (w_{2,t-1}w_{3,t-1} \cdots w_{h,t-1})(w_{2,t}w_{3,t} \cdots w_{h,t} \cdot w_{l,t}^{(1)} \cdots w_{n,t}) \\ & (w_{2,t+1}w_{3,t+1} \cdots w_{n,t+1}) \cdots (w_{2,n-1}w_{3,n-1} \cdots w_{n,n-1}) = \overleftarrow{(d_A)}_{l,t}^{(1)} \cdot w_{l,t}^{(1)} \cdot \overrightarrow{(d_A)}_{l,t}^{(1)} \end{aligned} \quad (4.6.6)$$

Now we derive the commutation formula for $c_{\widehat{a}_{h,k-1}+p}T_{d_A}$.

First, we have

$$c_{\widehat{a}_{h,k-1}+p}T_{\overleftarrow{(d_A)}_{l,t}^{(1)}} = T_{\overleftarrow{(d_A)}_{l,t}^{(1)}}c_{\widehat{a}_{h,k-1}+p}, \quad (4.6.7)$$

since, by Lemma 3.3(2) and noting $A_{\uparrow}^{h,t} = 0$,

$$\max(\overleftarrow{(d_A)}_{l,t}^{(1)}) \leq \sigma_{h-1,t} + a_{h,t} - 1 = \widehat{a}_{h,t-1} + \sum_{i>h, j<t} a_{i,j} + a_{h,t} - 1 = \widehat{a}_{h,t-1} - 1 < \widehat{a}_{h,k-1}.$$

Second, we consider the commutation relation for $c_{\widehat{a}_{h,k-1}+p}T_{w_{l,t}^{(1)}}$. The hypothesis $a_{h,k} > 0$ together with $a_{i,j} = 0$, for $i > h$ and $j < t$ (and noting $l > h$) imply

$$\sigma_{l-1,t} = \tilde{\lambda}_{l-1} + \sum_{i>l-1, j<t} a_{i,j} = \tilde{\lambda}_{l-1} \geq \tilde{\lambda}_h \geq \tilde{\lambda}_{h-1} + \overleftarrow{\Upsilon}_h^k + p = \widehat{a}_{h,k-1} + p =: p'.$$

On the other hand, $a_{l,t} > 0$ together with $a_{i,t} = 0$, for $h < i < l$, and $A_{\uparrow}^{h,t} = 0$ implies $\tilde{a}_{l-1,t} = \tilde{a}_{h,t}$ and

$$\tilde{a}_{l-1,t} + 1 = \tilde{a}_{h,t} + 1 \leq \tilde{\lambda}_{h-1} + \overleftarrow{\Upsilon}_h^k + p = p'.$$

Hence, $\sigma_{l-1,t} \geq p' \geq \tilde{a}_{l-1,t} + 1$ and so, $w_{l,t}^{(1)} = s_{\sigma_{l-1,t}} \cdots s_{p'} \cdots s_{\tilde{a}_{l-1,t}+1}$ (i.e., $s_{p'} | w_{l,t}^{(1)}$). Applying Lemma 4.5(1) yields

$$\begin{aligned} c_{p'}T_{w_{l,t}^{(1)}} &= T_{w_{l,t}^{(1)}}c_{p'+1} + (q-1)T_{w_{l,t}^{(1)} \setminus s_{p'}}(c_{\tilde{a}_{l-1,t}+1} - c_{p'+1}) \\ &= (q-1)T_{w_{l,t}^{(1)} \setminus s_{p'}}c_{\tilde{a}_{l-1,t}+1} + (T_{w_{l,t}^{(1)}} - (q-1)T_{w_{l,t}^{(1)} \setminus s_{p'}})c_{p'+1}. \end{aligned} \quad (4.6.8)$$

where $w_{l,t}^{(1)} \setminus s_{p'}$ ($p' = \widehat{a}_{h,k-1} + p$) is the element obtained by deleting $s_{p'}$ from $w_{l,t}^{(1)}$.

Finally, consider the multiplication (4.6.8) $\times T_{\overleftarrow{(d_A)}_{l,t}^{(1)}}$. We must analyse the commutation relations for

$$(1) \quad c_{\tilde{a}_{l-1,t}+1}T_{\overleftarrow{(d_A)}_{l,t}^{(1)}}, \quad (2) \quad c_{p'+1}T_{\overleftarrow{(d_A)}_{l,t}^{(1)}}.$$

For (1), observe that the first factor of $\overleftarrow{(d_A)}_{l,t}^{(1)}$ is $w_{l,t}^{(2)}$. By Lemma 3.3(1),

$$\min(\overleftarrow{(d_A)}_{l,t}^{(1)}) \geq \tilde{a}_{l-1,t} + 2 > \tilde{a}_{l-1,t} + 1.$$

Thus, by (2.0.3), we have

$$c_{\tilde{a}_{l-1,t}+1}T_{\overleftarrow{(d_A)}_{l,t}^{(1)}} = T_{\overleftarrow{(d_A)}_{l,t}^{(1)}}c_{\tilde{a}_{l-1,t}+1}. \quad (4.6.9)$$

For (2), since

$$\overleftarrow{(d_A)}_{l,t}^{(1)} = w_{l,t}^{(2)} \cdots w_{l,t}^{(a_{l,t})} w_{l+1,t} \cdots w_{n,t} (w_{2,t+1}w_{3,t+1} \cdots w_{n,t+1}) \cdots (w_{2,n-1}w_{3,n-1} \cdots w_{n,n-1}),$$

and every nonidentity factor $w_{i,j}$ of $\overrightarrow{(d_A)}_{l,t}^{(1)}$ is a product of nonidentity $w_{i,j}^{(k)}$ ($k \in [1, a_{i,j}]$) satisfying $i > l$, if $j = t$, or $j > t$ and the inequality $\min(w_{i,j}^{(k)}) = \tilde{a}_{i-1,j} + k \geq \tilde{a}_{i-1,j} + 1 > \tilde{a}_{l-1,t} + 1$ by (3.2.3). By deleting all the identity $w_{i,j}$ from $\overrightarrow{(d_A)}_{l,t}^{(1)}$, we may assume that $\overrightarrow{(d_A)}_{l,t}^{(1)} = y_1 y_2 \cdots y_m$ is a product of those nonidentity $w_{i,j}^{(k)}$. Let $m_1 = \min(y_1)$ and $m = \min(m_1, p' + 1)$. Then $\min(y_i) \geq \min(y_1)$ for all $i \in [1, m]$ and $m > \tilde{a}_{l-1,t} + 1$. Moreover, each y_i has the form

$$y_i = s_b s_{b-1} \cdots s_a, \quad \text{where } b = \sigma_{i-1,j} + k - 1 \geq \tilde{a}_{i-1,j} + k = a \geq m. \quad (4.6.10)$$

Let $\mathcal{H}_{\geq m}$ be the subalgebra of $\mathcal{H}_{r,R}^c$ generated by T_i, c_j with $m \leq i \leq r-1$ and $m \leq j \leq r$. Then $c_{p'+1} \in \mathcal{H}_{\geq m}$ and $T_{\overrightarrow{(d_A)}_{l,t}^{(1)}} \in \mathcal{H}_{\geq m}$. Hence, $c_{p'+1} T_{\overrightarrow{(d_A)}_{l,t}^{(1)}} \in \mathcal{H}_{\geq m}$. Consequently, by (2.1.2),

$$c_{p'+1} T_{\overrightarrow{(d_A)}_{l,t}^{(1)}} = T_{\overrightarrow{(d_A)}_{l,t}^{(1)}} c_{\tilde{a}_{h-1,k+p}} + \sum_{d' < \overrightarrow{(d_A)}_{l,t}^{(1)}, u \geq m} f_{d',u} T_{d'} c_u \quad (f_{d',u} \in R). \quad (4.6.11)$$

Combining (4.6.7)–(4.6.11) gives

$$c_{\tilde{a}_{h,k-1+p}} T_{d_A} = T_{d_A} c_{\tilde{a}_{h-1,k+p}} + (q-1) T_{d_A \setminus s_{\tilde{\lambda}_{h-1} + \tilde{\tau}_h^{k+p}}} c_{\tilde{a}_{l-1,t+1}} + \sum_{\substack{d < d_A, \\ u' > \tilde{a}_{l-1,t+1}}} g_{d,u'} T_d c_{u'}. \quad (4.6.12)$$

Since the “tail” $(q-1) T_{d_A \setminus s_{\tilde{\lambda}_{h-1} + \tilde{\tau}_h^{k+p}}} c_{\tilde{a}_{l-1,t+1}} + \sum_{u' > \tilde{a}_{l-1,t+1}} g_{d,u'} T_d c_{u'}$ is nonzero (as $q \neq 1$), it follows that A does not satisfy the SDP condition at (h, k) . \square

Corollary 4.7. (1) Every $A = (a_{i,j}) \in M_n(\mathbb{N})$ satisfies the SDP condition on the 1st column or n th row.

(2) If $l \in [1, n]$ and $A = (a_{i,j}) \in M_n(\mathbb{N})$ satisfy $a_{i,j} = 0$, for $i > l$ and $j < n$, then A satisfies SDP condition on the l -th row.

5. MULTIPLICATION FORMULAS IN $\mathcal{Q}_q(n, r; R)$: THE EVEN CASE

We are now ready to compute, for any $A^* = (a_{i,j}^{\bar{0}} | a_{i,j}^{\bar{1}}) \in M_n(\mathbb{N} | \mathbb{N}_2)_r$, the structure constants that appear in $\phi_{X^*} \phi_{A^*}$, where X^* is one of the following six matrices:

$$\begin{aligned} \text{the even case: } & (\lambda | O), \quad (\lambda + E_{h,h+1} - E_{h+1,h+1} | O), \quad (\lambda - E_{h,h} + E_{h+1,h} | O); \\ \text{the odd case: } & (\lambda - E_{h,h} | E_{h,h}), \quad (\lambda - E_{h+1,h+1} | E_{h,h+1}), \quad (\lambda - E_{h,h} | E_{h+1,h}). \end{aligned} \quad (5.0.1)$$

Here $1 \leq h \leq n-1$, and we regard any $\lambda \in \Lambda(n, r)$ as a diagonal matrix $\lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$ in $M_n(\mathbb{N})$ and write $\lambda + B := \text{diag}(\lambda_1, \dots, \lambda_n) + B$, for $B \in M_n(\mathbb{N})$.

We first deal with the even case in this section.

Convention 5.1. The terms ϕ_{M^*} appearing in $\phi_{X^*} \phi_{A^*}$ have the form $M^* = (A^{\bar{0}} \pm E | A^{\bar{1}} \pm E')$ for some $E \in M_n(\mathbb{N})$, $E' \in M_n(\mathbb{N}_2)$. If $M^* \notin M_n(\mathbb{N} | \mathbb{N}_2)$ (i.e., M^* has a negative entry or $A^{\bar{1}} \pm E'$ has an entry > 1), then we set ϕ_{M^*} to be 0. We make a similar convention for the elements T_{M^*} in (2.2.12) that defines ϕ_{M^*} .

The following notations will be used in the proofs of the multiplication formulas in this and next sections.

Notation 5.2 (for the proofs in Sections 5 and 6). Let $A^* = (a_{i,j}^{\bar{0}} | a_{i,j}^{\bar{1}}) \in M_n(\mathbb{N} | \mathbb{N}_2)$ and $A = [A^*]$ its base matrix. For fixed $1 \leq h, k \leq n$, we decompose the elements c_{A^*}, c'_{A^*} defined in (2.2.12) as a product of three segments:

$$c_{A^*} = \overleftarrow{c}_{A^*}^{h,k} \cdot c_{A^*;k}^{h,h+1} \cdot \overrightarrow{c}_{A^*}^{h+1,k}, \quad c'_{A^*} = (\overleftarrow{c}_{A^*}^{h,k})' \cdot (c_{A^*;k}^{h,h+1})' \cdot (\overrightarrow{c}_{A^*}^{h+1,k})' \quad (5.2.1)$$

where $c_{A^*;k}^{h,h+1} := (c_{q, \tilde{a}_{h-1,k+1}, \tilde{a}_{h,k}})^{\bar{1}} \cdot (c_{q, \tilde{a}_{h,k+1}, \tilde{a}_{h+1,k}})^{\bar{1}}$, and $\overleftarrow{c}_{A^*}^{h,k}$ (resp., $\overrightarrow{c}_{A^*}^{h+1,k}$) are the segment of the product before (resp., after) $c_{A^*;k}^{h,h+1}$. The decomposition for c'_{A^*} is similar.

Recall the notations in (3.0.1) and (3.0.2) with $\lambda = \text{ro}(A)$, and let $\overleftarrow{\mathbf{r}}_{h+1}^k = \overleftarrow{\mathbf{r}}_{h+1}^k(A)$, $\overrightarrow{\mathbf{r}}_h^k = \overrightarrow{\mathbf{r}}_h^k(A)$. Then

$$\lambda_{(h)}^+ := \text{ro}(A_{h,k}^+) = \lambda + \epsilon_h - \epsilon_{h+1} = \lambda + \alpha_h, \quad \lambda_{(h)}^- := \text{co}(A_{h,k}^-) = \lambda - \epsilon_h + \epsilon_{h+1} = \lambda - \alpha_h,$$

and $\text{co}(A_{h,k}^+) = \text{co}(A_{h,k}^-) = \text{co}(A)$. Let

$$\begin{aligned} \nu &= \nu_A = (\cdots, a_{h,k}, a_{h+1,k}, \cdots) \quad (\text{as in (2.2.3)}), \\ \nu^+ &= \nu_{A_{h,k}^+} = (\cdots, a_{h,k} + 1, a_{h+1,k} - 1, \cdots), \quad \delta^+ = (\cdots, a_{h,k}, 1, a_{h+1,k} - 1, \cdots), \\ \nu^- &= \nu_{A_{h,k}^-} = (\cdots, a_{h,k} - 1, a_{h+1,k} + 1, \cdots), \quad \delta^- = (\cdots, a_{h,k} - 1, 1, a_{h+1,k}, \cdots). \end{aligned} \quad (5.2.2)$$

Here the omitted components are identical. Clearly, $\mathfrak{S}_{\delta^\pm} \subset \mathfrak{S}_{\nu^\pm}$. If \mathcal{D}_\pm denote the sets of the shortest representatives of the left cosets of $\mathfrak{S}_{\delta^\pm}$ in \mathfrak{S}_{ν^\pm} , i.e.,

$$\mathcal{D}_+ = \mathcal{D}_{\delta^+}^{-1} \cap \mathfrak{S}_{\nu^+}, \quad \mathcal{D}_- = \mathcal{D}_{\delta^-}^{-1} \cap \mathfrak{S}_{\nu^-},$$

and set $\tilde{a} = \tilde{a}_{h,k}$, $a = a_{h,k}$, $b = a_{h+1,k}$, and $x_{\mathcal{D}_\pm} = \Sigma_{\mathcal{D}_\pm}$, then we have

$$\begin{aligned} x_{\nu^+} &= x_{\mathcal{D}_+} x_{\delta^+} = x_{\delta^+} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright}, & x_{\nu^-} &= x_{\delta^-} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} = x_{\delta^-} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright}. \end{aligned} \quad (5.2.3)$$

Let $\nu^+ = \nu_{A_{h,k}^+}$ and $\nu^- = \nu_{A_{h,k}^-}$ be defined as in (2.2.3) by using the transposes ${}^t A_{h,k}^+$, ${}^t A_{h,k}^-$ of $A_{h,k}^+$, $A_{h,k}^-$, respectively. Then $\mathfrak{S}_{\nu^\pm} \subseteq \mathfrak{S}_{\lambda_{(h)}^\pm}$ and we have (cf. [18, Corollary 3.5])

$$\begin{aligned} x_{\lambda_{(h)}^+} T_{d_{A_{h,k}^+}} &= x_{\lambda_{(h)}^+} \setminus \nu^+ T_{d_{A_{h,k}^+}} x_{\nu^+}, \quad \text{where} \quad x_{\lambda_{(h)}^+} \setminus \nu^+ := \sum_{u' \in \mathcal{D}_{\nu^+}^{-1} \cap \mathfrak{S}_{\lambda_{(h)}^+}} T_{u'}; \\ x_{\lambda_{(h)}^-} T_{d_{A_{h,k}^-}} &= x_{\lambda_{(h)}^-} \setminus \nu^- T_{d_{A_{h,k}^-}} x_{\nu^-}, \quad \text{where} \quad x_{\lambda_{(h)}^-} \setminus \nu^- := \sum_{u' \in \mathcal{D}_{\nu^-}^{-1} \cap \mathfrak{S}_{\lambda_{(h)}^-}} T_{u'}. \end{aligned} \quad (5.2.4)$$

The following displays the multiplication formulas for $\phi_{X^*} \phi_{A^*}$ in the queer q -Schur superalgebra $\mathcal{Q}_q(n, r; R)$, where X is one of the following matrices

$$D_\mu^* := (\mu|O), \quad E_{h,\lambda}^* := (\lambda + E_{h,h+1} - E_{h+1,h+1}|O), \quad F_{h,\lambda}^* := (\lambda - E_{h,h} + E_{h+1,h}|O). \quad (5.2.5)$$

Theorem 5.3. *Let $h \in [1, n-1]$ and $A^* = (a_{i,j}^{\bar{0}} | a_{i,j}^{\bar{1}}) \in M_n(\mathbb{N}|\mathbb{N}_2)_r$. Assume $A = (a_{i,j})$ is the base of A^* with $a_{i,j} = a_{i,j}^{\bar{0}} + a_{i,j}^{\bar{1}}$, and $\overrightarrow{\mathbf{r}}_h^k = \overrightarrow{\mathbf{r}}_h^k(A)$. Then, for $\lambda, \mu \in \Lambda(n, r)$ and $\varepsilon = \delta_{\lambda, \text{ro}(A)}$, the following multiplication formulas hold in $\mathcal{Q}_q(n, r; R)$:*

- (1) $\phi_{D_\mu^*} \phi_{A^*} = \delta_{\mu, \text{ro}(A)} \phi_{A^*}, \quad \phi_{A^*} \phi_{D_\mu^*} = \delta_{\mu, \text{co}(A)} \phi_{A^*}.$
- (2) $\phi_{E_{h,\lambda}^*} \phi_{A^*} = \varepsilon \sum_{k=1}^n \left\{ q^{\overrightarrow{\mathbf{r}}_h^k + a_{h+1,k}^{\bar{1}}} \llbracket a_{h,k}^{\bar{0}} + 1 \rrbracket_q \phi_{(A^{\bar{0}} + E_{h,k} - E_{h+1,k} | A^{\bar{1}})} + q^{\overrightarrow{\mathbf{r}}_h^k} \phi_{(A^{\bar{0}} | A^{\bar{1}} + E_{h,k} - E_{h+1,k})} \right. \\ \left. + q^{\overrightarrow{\mathbf{r}}_h^k - 1} \llbracket a_{h,k} + 1 \rrbracket_{q,q^2} \phi_{(A^{\bar{0}} + 2E_{h,k} | A^{\bar{1}} - E_{h,k} - E_{h+1,k})} \right\}.$
- (3) $\phi_{F_{h,\lambda}^*} \phi_{A^*} = \varepsilon \sum_{k=1}^n \left\{ q^{\overleftarrow{\mathbf{r}}_{h+1}^k} \llbracket a_{h+1,k}^{\bar{0}} + 1 \rrbracket_q \phi_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} | A^{\bar{1}})} + q^{\overleftarrow{\mathbf{r}}_{h+1}^k + a_{h,k} - 1} \phi_{(A^{\bar{0}} | A^{\bar{1}} - E_{h,k} + E_{h+1,k})} \right. \\ \left. + q^{\overleftarrow{\mathbf{r}}_{h+1}^k + a_{h,k} - 2} \llbracket a_{h+1,k} + 1 \rrbracket_{q,q^2} \phi_{(A^{\bar{0}} + 2E_{h+1,k} | A^{\bar{1}} - E_{h,k} - E_{h+1,k})} \right\}.$

Proof. Formula (1) is a special case of (2.3.1), noting that $\phi_{(\lambda|O)}$ is the identity map on $x_\lambda \mathcal{H}_{r,R}^c$. We now prove (2).

Let $X^* = E_{h,\lambda}^* = (\lambda + E_{h,h+1} - E_{h+1,h+1}|O)$ whose base matrix $X = \lambda + E_{h,h+1} - E_{h+1,h+1}$. Then $\text{ro}(X) = \lambda + \alpha_h = \lambda_{(h)}^+$, $\text{co}(X) = \lambda$, $d_X = 1$, $T_{d_X} = 1$, and $c_{X^*} = 1$. Further, we have

$\mathfrak{S}_{\nu_X} = \mathfrak{S}_{\lambda_{(h)}^+} \cap \mathfrak{S}_\lambda$ and $\mathcal{D}_{\nu_X} \cap \mathfrak{S}_\lambda = \{1, s_{\tilde{\lambda}_{h+1}}, s_{\tilde{\lambda}_{h+1}} s_{\tilde{\lambda}_{h+2}}, \dots, s_{\tilde{\lambda}_{h+1}} s_{\tilde{\lambda}_{h+2}} \dots s_{\tilde{\lambda}_{h+1-1}}\}$. Thus,

$$T_{X^*} = x_{\lambda_{(h)}^+} \Sigma_{\mathcal{D}_{\nu_X} \cap \mathfrak{S}_\lambda} = x_{\lambda_{(h)}^+} T_{(\tilde{\lambda}_{h+1}, \tilde{\lambda}_{h+1-1})}^{\triangleleft} = \sum_{\substack{1 \leq k \leq n \\ a_{h+1,k} > 0}} x_{\lambda_{(h)}^+} \cdot \sum_{j=\overleftarrow{\mathfrak{F}}_{h+1}^k}^{\overleftarrow{\mathfrak{F}}_{h+1}^{k+1}-1} T_{\tilde{\lambda}_{h+1}} T_{\tilde{\lambda}_{h+2}} \dots T_{\tilde{\lambda}_{h+j}}.$$

Hence, for $\lambda = \text{ro}(A)$ and $\mu = \text{co}(A)$, (2.3.2) with $B^* = X^*$ becomes

$$z := x_{\lambda_{(h)}^+} T_{(\tilde{\lambda}_{h+1}, \tilde{\lambda}_{h+1-1})}^{\triangleleft} T_{d_A} c_{A^*} \sum_{\sigma \in \mathcal{D}_{\nu_A} \cap \mathfrak{S}_\mu} T_\sigma = \sum_{\substack{1 \leq k \leq n \\ a_{h+1,k} > 0}} z_k,$$

where, for each $k \in [1, n]$ with $a_{h+1,k} > 0$ (and $\Sigma_A := \sum_{\sigma \in \mathcal{D}_{\nu_A} \cap \mathfrak{S}_\mu} T_\sigma$),

$$z_k = x_{\lambda_{(h)}^+} \left(\sum_{j=\overleftarrow{\mathfrak{F}}_{h+1}^k}^{\overleftarrow{\mathfrak{F}}_{h+1}^{k+1}-1} T_{\tilde{\lambda}_{h+1}} T_{\tilde{\lambda}_{h+2}} \dots T_{\tilde{\lambda}_{h+j}} T_{d_A} \right) c_{A^*} \Sigma_A. \quad (5.3.1)$$

Then, with $\tilde{a} = \tilde{a}_{h,k}$, $a = a_{h,k}$, and $b = a_{h+1,k}$, (5.3.1) becomes, by Lemma 3.4,

$$\begin{aligned} z_k &= x_{\lambda_{(h)}^+} T_{\tilde{\lambda}_h} T_{\tilde{\lambda}_{h-1}} \dots T_{\tilde{\lambda}_h - \overrightarrow{\mathfrak{F}}_{h+1}^k} T_{d_{A_{h,k}^+}} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} c_{A^*} \Sigma_A \\ &= q^{\overrightarrow{\mathfrak{F}}_{h+1}^k} x_{\lambda_{(h)}^+} T_{d_{A_{h,k}^+}} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} c_{A^*} \Sigma_A \quad (\text{since } s_{\tilde{\lambda}_h}, s_{\tilde{\lambda}_{h-1}}, \dots, s_{\tilde{\lambda}_h - \overrightarrow{\mathfrak{F}}_{h+1}^k} \in \mathfrak{S}_{\lambda_{(h)}^+}) \\ &= q^{\overrightarrow{\mathfrak{F}}_{h+1}^k} x_{\lambda_{(h)}^+} \setminus_{\nu^+} T_{d_{A_{h,k}^+}} x_{\mathcal{D}_+} x_{\delta^+} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} c_{A^*} \Sigma_A \quad (x_{\nu^+} = x_{\mathcal{D}_+} x_{\delta^+} \text{ by (5.2.4) and (5.2.3)}) \\ &= q^{\overrightarrow{\mathfrak{F}}_{h+1}^k} x_{\lambda_{(h)}^+} \setminus_{\nu^+} T_{d_{A_{h,k}^+}} x_{\mathcal{D}_+} (c_{A^*})' x_{\nu} \Sigma_A \quad (x_{\delta^+} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} = x_{\nu} \text{ by (5.2.3) and by (2.2.7)}). \end{aligned}$$

By rewriting $x_{\nu} = x_{\delta^+} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft}$ again and applying Lemma 3.5(1), we have

$$\begin{aligned} z_k &= q^{\overrightarrow{\mathfrak{F}}_{h+1}^k} x_{\lambda_{(h)}^+} \setminus_{\nu^+} T_{d_{A_{h,k}^+}} x_{\mathcal{D}_+} (c_{A^*})' x_{\delta^+} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} \Sigma_{A_{h,k}^+}, \quad \text{where } \Sigma_{A_{h,k}^+} = \sum_{\sigma \in \mathcal{D}_{\nu^+} \cap \mathfrak{S}_\mu} T_\sigma, \\ &= q^{\overrightarrow{\mathfrak{F}}_{h+1}^k} x_{\lambda_{(h)}^+} \setminus_{\nu^+} T_{d_{A_{h,k}^+}} x_{\mathcal{D}_+} (\overleftarrow{c}_{A^*}^{h,k})' (c_{A^*}^{h,h+1})' (\overrightarrow{c}_{A^*}^{h+1,k})' x_{\delta^+} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} \Sigma_{A_{h,k}^+} \\ &= (-1)^{(a_{h,k}^{\bar{1}} + a_{h+1,k}^{\bar{1}}) \tilde{a}_{h-1,k}^{\bar{1}}} q^{\overrightarrow{\mathfrak{F}}_{h+1}^k} x_{\lambda_{(h)}^+} \setminus_{\nu^+} T_{d_{A_{h,k}^+}} \cdot \underbrace{x_{\mathcal{D}_+} (c_{A^*}^{h,h+1})' x_{\delta^+} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright}}_{=\mathfrak{m}^+} \cdot \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+}. \end{aligned} \quad (5.3.2)$$

Here, for the last equality, we used the following relations:

- (a) $(\overleftarrow{c}_{A^*}^{h,k})' (c_{A^*}^{h,h+1})' (\overrightarrow{c}_{A^*}^{h+1,k})' = (-1)^{(a_{h,k}^{\bar{1}} + a_{h+1,k}^{\bar{1}}) \tilde{a}_{h-1,k}^{\bar{1}}} (c_{A^*}^{h,h+1})' (\overleftarrow{c}_{A^*}^{h,k})' (\overrightarrow{c}_{A^*}^{h+1,k})'$ (noting that $\tilde{a}_{h-1,k}^{\bar{1}}$ is the partial sum at the $(h-1, k)$ position associated with $\nu_{A^{\bar{1}}}$);
- (b) $(\overleftarrow{c}_{A^*}^{h,k})' (\overrightarrow{c}_{A^*}^{h+1,k})' = (c_{\delta^+}^\alpha)'$ for some $\alpha \leq \delta^+$ with $\alpha_i = 0$, $i \in [\tilde{a}_{h-1,k} + 1, \tilde{a}_{h+1,k}]$, and hence $(\overleftarrow{c}_{A^*}^{h,k})' (\overrightarrow{c}_{A^*}^{h+1,k})' x_{\delta^+} = x_{\delta^+} (\overleftarrow{c}_{A^*}^{h,k}) (\overrightarrow{c}_{A^*}^{h+1,k})$ by (2.2.7);
- (c) $(\overleftarrow{c}_{A^*}^{h,k}) (\overrightarrow{c}_{A^*}^{h+1,k})$ commutes with $T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright}$ since $T_i c_j = c_j T_i$ for all $j \neq i, i+1$.

To complete the computation of z_k , we need to manipulate the ‘‘middle part’’ \mathfrak{m}^+ . This requires to consider the following four cases. We record the following fact

$$x_{\nu^+} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} = \llbracket a+1 \rrbracket_q x_{\nu^+}, \quad \text{since } s_{\tilde{a}}, s_{\tilde{a}-1}, \dots, s_{\tilde{a}-a+1} \in \mathfrak{S}_{\nu^+}. \quad (5.3.3)$$

The first case is simple.

Case 1 ($a_{h,k}^{\bar{1}} = 0, a_{h+1,k}^{\bar{1}} = 0$). In this case, $(c_{A^*;k}^{h,h+1})' = 1, c_{A^*} = \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} = c_{A_{h,k}^+}$ and

$$\begin{aligned} z_k &= q^{\vec{r}_h^k} x_{\lambda_{(h)}^+} \vee_{\nu^+} T_{d_{A_{h,k}^+}} x_{\nu^+} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+} \quad (\text{as } x_{\mathcal{D}_+} x_{\delta^+} = x_{\nu^+}) \\ &= q^{\vec{r}_h^k} \llbracket a+1 \rrbracket_q x_{\lambda_{(h)}^+} T_{d_{A_{h,k}^+}} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+} \\ &= q^{\vec{r}_h^k} \llbracket a_{h,k}^{\bar{0}} + 1 \rrbracket_q T_{(A^{\bar{0}} + E_{h,k} - E_{h+1,k} | A^{\bar{1}})}. \end{aligned}$$

Case 2 ($a_{h,k}^{\bar{1}} = 0, a_{h+1,k}^{\bar{1}} = 1$). Then $(c_{A^*;k}^{h,h+1})' = c'_{q,\tilde{a}+1,\tilde{a}+b}$ and

$$z_k = (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\vec{r}_h^k} x_{\lambda_{(h)}^+} \vee_{\nu^+} T_{d_{A_{h,k}^+}} \cdot \underbrace{x_{\mathcal{D}_+} (c_{q,\tilde{a}+1,\tilde{a}+b})' x_{\delta^+} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright}}_{=\mathbf{m}^+} \cdot \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+}.$$

We now rewrite the middle part \mathbf{m}^+ using (5.2.3), (2.2.7), and (2.2.5), etc. as follows:

$$\begin{aligned} \mathbf{m}^+ &= x_{\mathcal{D}_+} (q(c_{q,\tilde{a}+2,\tilde{a}+b})' + c_{\tilde{a}+1}) x_{\delta^+} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} \quad (\text{by (2.2.5)}) \\ &= q x_{\mathcal{D}_+} x_{\delta^+} c_{q,\tilde{a}+2,\tilde{a}+b} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} + x_{\mathcal{D}_+} x_{\delta^+} c_{\tilde{a}+1} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} \quad (\text{by (2.2.7)}) \\ &= q(c_{q,\tilde{a}+2,\tilde{a}+b})' x_{\nu^+} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} + x_{\nu^+} c_{q,\tilde{a}-a+1,\tilde{a}+1} \quad (\text{by (5.2.3), Lemma 3.7(3)}) \\ &= q \llbracket a+1 \rrbracket_q x_{\nu^+} c_{q,\tilde{a}+2,\tilde{a}+b} + x_{\nu^+} c_{q,\tilde{a}-a+1,\tilde{a}+1} \quad (\text{by (5.3.3)}). \end{aligned}$$

Notice that, if $a_{h+1,k}^{\bar{0}} = 0$ (thus, $b = 1$), then the term $(c_{q,\tilde{a}+2,\tilde{a}+b})' = 0$ (hence, $c_{q,\tilde{a}+2,\tilde{a}+b} = 0$). Hence, by noting $\overleftarrow{c}_{A^*}^{h,k} c_{q,\tilde{a}+2,\tilde{a}+b} \overrightarrow{c}_{A^*}^{h+1,k} = c_{\nu^+}^{\alpha}$ with $\alpha = \nu_{A^{\bar{1}}}$ and $\overleftarrow{c}_{A^*}^{h,k} c_{q,\tilde{a}-a+1,\tilde{a}+1} \overrightarrow{c}_{A^*}^{h+1,k} = c_{\nu^+}^{\beta}$ with $\beta = \nu_{A^{\bar{1}} - E_{h+1,k} + E_{h,k}}$,

$$\begin{aligned} z_k &= q^{\vec{r}_h^k + 1} \llbracket a+1 \rrbracket_q x_{\lambda_{(h)}^+} T_{d_{A_{h,k}^+}} \overleftarrow{c}_{A^*}^{h,k} c_{q,\tilde{a}+2,\tilde{a}+b} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+} \\ &\quad + q^{\vec{r}_h^k} x_{\lambda_{(h)}^+} T_{d_{A_{h,k}^+}} \overleftarrow{c}_{A^*}^{h,k} c_{q,\tilde{a}-a+1,\tilde{a}+1} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+} \\ &= q^{\vec{r}_h^k + 1} \llbracket a_{h,k}^{\bar{0}} + 1 \rrbracket_q T_{(A^{\bar{0}} + E_{h,k} - E_{h+1,k} | A^{\bar{1}})} + q^{\vec{r}_h^k} T_{(A^{\bar{0}} | A^{\bar{1}} + E_{h,k} - E_{h+1,k})}. \end{aligned}$$

Here the last step follows from the facts that $(A^{\bar{0}} + E_{h,k} - E_{h+1,k} | A^{\bar{1}})$ and $(A^{\bar{0}} | A^{\bar{1}} + E_{h,k} - E_{h+1,k})$ have the same base $A_{h,k}^+$, and the sign in z_k is cancelled after swapping $c_{q,\tilde{a}+2,\tilde{a}+b}$ (resp., $c_{q,\tilde{a}-a+1,\tilde{a}+1}$) with $\overleftarrow{c}_{A^*}^{h,k}$. (Note that, by Convention 5.1, $T_{(A^{\bar{0}} + E_{h,k} - E_{h+1,k} | A^{\bar{1}})} = 0$ if $a_{h+1,k}^{\bar{0}} = 0$.)

Case 3 ($a_{h,k}^{\bar{1}} = 1, a_{h+1,k}^{\bar{1}} = 0$). Then we have $c_{A^*;k}^{h,h+1} = c_{q,\tilde{a}-a+1,\tilde{a}}$ and

$$z_k = (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\vec{r}_h^k} x_{\lambda_{(h)}^+} \vee_{\nu^+} T_{d_{A_{h,k}^+}} \cdot \underbrace{x_{\mathcal{D}_+} (c_{q,\tilde{a}-a+1,\tilde{a}})' x_{\delta^+} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright}}_{=\mathbf{m}^+} \cdot \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+}.$$

For the middle part \mathbf{m}^+ , again using (5.2.3), (2.2.7), and (2.2.5), one can deduce

$$\begin{aligned} q\mathbf{m}^+ &= x_{\mathcal{D}_+} x_{\delta^+} \cdot q c_{q,\tilde{a}-a+1,\tilde{a}} \cdot T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} \\ &= x_{\nu^+} c_{q,\tilde{a}-a+1,\tilde{a}+1} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} - x_{\nu^+} c_{\tilde{a}+1} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} \\ &= (c_{q,\tilde{a}-a+1,\tilde{a}+1})' x_{\nu^+} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} - x_{\nu^+} c_{q,\tilde{a}-a+1,\tilde{a}+1} \quad (\text{by Lemma 3.7(3)}) \\ &= (\llbracket a+1 \rrbracket_q - 1) x_{\nu^+} c_{q,\tilde{a}-a+1,\tilde{a}+1} \quad (\text{by (5.3.3)}). \end{aligned}$$

Substituting leads to

$$\begin{aligned} z_k &= q^{\vec{r}_h^k - 1} (\llbracket a+1 \rrbracket_q - 1) x_{\lambda_{(h)}^+} T_{d_{A_{h,k}^+}} \overleftarrow{c}_{A^*}^{h,k} c_{q,\tilde{a}-a+1,\tilde{a}+1} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+} \\ &= q^{\vec{r}_h^k} \llbracket a_{h,k}^{\bar{0}} + 1 \rrbracket_q T_{(A^{\bar{0}} + E_{h,k} - E_{h+1,k} | A^{\bar{1}})}. \end{aligned}$$

Case 4 ($a_{h,k}^{\bar{1}} = 1, a_{h+1,k}^{\bar{1}} = 1$). Then $(c_{A^*;k}^{h,h+1})' = c'_{q,\tilde{a}-a+1,\tilde{a}} c'_{q,\tilde{a}+1,\tilde{a}+b}$ and

$$z_k = q^{\vec{r}_h^k} x_{\lambda_{(h)}^+} \vee_{\nu^+} T_{d_{A_{h,k}^+}} \cdot \underbrace{x_{\mathcal{D}^+} c'_{q,\tilde{a}-a+1,\tilde{a}} c'_{q,\tilde{a}+1,\tilde{a}+b} x_{\delta^+} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright}}_{=\mathfrak{m}^+} \cdot \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+}.$$

This time, we handle the middle part \mathfrak{m}^+ in a more tricky way: first swapping the two c' factors, then commuting one with x_{δ^+} , then separating $c_{\tilde{a}+1}$ from the c' factor and adding the tail $c_{\tilde{a}+1}$ to the q times the c -factor, and finally expanding the product. In other words, we have

$$\begin{aligned} qc'_{q,\tilde{a}-a+1,\tilde{a}} c'_{q,\tilde{a}+1,\tilde{a}+b} x_{\delta^+} &= -qc'_{q,\tilde{a}+1,\tilde{a}+b} c'_{q,\tilde{a}-a+1,\tilde{a}} x_{\delta^+} = -c'_{q,\tilde{a}+1,\tilde{a}+b} x_{\delta^+} (qc_{q,\tilde{a}-a+1,\tilde{a}}) \\ &= -(c_{\tilde{a}+1} + qc'_{q,\tilde{a}+2,\tilde{a}+b}) x_{\delta^+} (c_{q,\tilde{a}-a+1,\tilde{a}+1} - c_{\tilde{a}+1}) \\ &= -x_{\delta^+} (qc_{q,\tilde{a}+2,\tilde{a}+b} + c_{\tilde{a}+1}) (c_{q,\tilde{a}-a+1,\tilde{a}+1} - c_{\tilde{a}+1}) \\ &= -x_{\delta^+} (qc_{q,\tilde{a}+2,\tilde{a}+b} c_{q,\tilde{a}-a+1,\tilde{a}+1} - qc_{q,\tilde{a}+2,\tilde{a}+b} c_{\tilde{a}+1} + c_{\tilde{a}+1} c_{q,\tilde{a}-a+1,\tilde{a}+1} - c_{\tilde{a}+1} c_{\tilde{a}+1}). \end{aligned} \quad (5.3.4)$$

Now, substituting into \mathfrak{m}^+ , noting $c_{\tilde{a}+1} c_{q,\tilde{a}-a+1,\tilde{a}+1} = -(c_{q,\tilde{a}-a+1,\tilde{a}+1} c_{\tilde{a}+1} + 2)$, $x_{\nu^+} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} = \llbracket a+1 \rrbracket_q x_{\nu^+}$, and repeatedly applying (5.2.3), (2.2.7) yield

$$\begin{aligned} q\mathfrak{m}^+ &= -qx_{\nu^+} c_{q,\tilde{a}+2,\tilde{a}+b} c_{q,\tilde{a}-a+1,\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} + qx_{\nu^+} c_{q,\tilde{a}+2,\tilde{a}+b} c_{\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} \\ &\quad - x_{\nu^+} c_{\tilde{a}+1} c_{q,\tilde{a}-a+1,\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} + x_{\nu^+} c_{\tilde{a}+1} c_{\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} \\ &= -q(c_{q,\tilde{a}+2,\tilde{a}+b})' (c_{q,\tilde{a}-a+1,\tilde{a}+1})' x_{\nu^+} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} + q(c_{q,\tilde{a}+2,\tilde{a}+b})' x_{\nu^+} c_{\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} \\ &\quad + x_{\nu^+} (c_{q,\tilde{a}-a+1,\tilde{a}+1} c_{\tilde{a}+1} + 2) T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} - x_{\nu^+} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} \\ &= -q\llbracket a+1 \rrbracket_q x_{\nu^+} c_{q,\tilde{a}+2,\tilde{a}+b} c_{q,\tilde{a}-a+1,\tilde{a}+1} + qx_{\nu^+} c_{q,\tilde{a}+2,\tilde{a}+b} c_{q,\tilde{a}-a+1,\tilde{a}+1} \\ &\quad + (c_{q,\tilde{a}-a+1,\tilde{a}+1})' x_{\nu^+} c_{\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} + \llbracket a+1 \rrbracket_q x_{\nu^+} \quad (2\text{nd term by Lemma 3.7(3)}) \\ &= q(\llbracket a+1 \rrbracket_q - 1) x_{\nu^+} c_{q,\tilde{a}-a+1,\tilde{a}+1} c_{q,\tilde{a}+2,\tilde{a}+b} + (\llbracket a+1 \rrbracket_q - \llbracket a+1 \rrbracket_{q^2}) x_{\nu^+}, \end{aligned}$$

where the third term $= x_{\nu^+} (c_{q,\tilde{a}-a+1,\tilde{a}+1})^2$ by Lemma 3.7(3) and then apply Lemma 3.7(4). Since $q(\llbracket a+1 \rrbracket_q - 1) = q^2 \llbracket a \rrbracket_q$, we have then, in this case,

$$\begin{aligned} z_k &= q^{\vec{r}_h^k+1} \llbracket a \rrbracket_q x_{\lambda_{(h)}^+} T_{d_{A_{h,k}^+}} \overleftarrow{c}_{A^*}^{h,k} c_{q,\tilde{a}-a+1,\tilde{a}+1} c_{q,\tilde{a}+2,\tilde{a}+b} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+} \\ &\quad + q^{\vec{r}_h^k-1} (\llbracket a+1 \rrbracket_q - \llbracket a+1 \rrbracket_{q^2}) x_{\lambda_{(h)}^+} T_{d_{A_{h,k}^+}} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+} \\ &= q^{\vec{r}_h^k+1} \llbracket a_{h,k}^{\bar{0}} + 1 \rrbracket_q T_{(A^{\bar{0}}+E_{h,k}-E_{h+1,k}|A^{\bar{1}})} + q^{\vec{r}_h^k-1} \llbracket a_{h,k} + 1 \rrbracket_{q,q^2} T_{(A^{\bar{0}}+2E_{h,k}|A^{\bar{1}}-E_{h,k}-E_{h+1,k})}. \end{aligned}$$

Let

$$\begin{aligned} z'_k &= q^{\vec{r}_h^k+a_{h+1,k}^{\bar{1}}} \llbracket a_{h,k}^{\bar{0}} + 1 \rrbracket_q T_{(A^{\bar{0}}+E_{h,k}-E_{h+1,k}|A^{\bar{1}})} + q^{\vec{r}_h^k} T_{(A^{\bar{0}}|A^{\bar{1}}+E_{h,k}-E_{h+1,k})} \\ &\quad + q^{\vec{r}_h^k-1} \llbracket a_{h,k} + 1 \rrbracket_{q,q^2} T_{(A^{\bar{0}}+2E_{h,k}|A^{\bar{1}}-E_{h,k}-E_{h+1,k})}. \end{aligned}$$

Then, by the notational Convention 5.1, $z'_k = 0$ if $a_{h+1,k} = 0$. A case-by-case argument following the four cases above shows that $z'_k = z_k$ if $a_{h+1,k} > 0$. This proves (2).

Finally, we prove the formula in (3). Observe that, for $X^* = F_{h,\lambda}^* = (\lambda - E_{h,h} + E_{h+1,h}|O)$ with $\lambda = \text{ro}(A)$, we have

$$\begin{aligned} X &= \lfloor X^* \rfloor = \text{diag}(\lambda) - E_{h,h} + E_{h+1,h}, \quad \text{and} \\ \mathcal{D}_{\nu_X} \cap \mathfrak{S}_\lambda &= \{1, s_{\tilde{\lambda}_{h-1}}, s_{\tilde{\lambda}_{h-1}} s_{\tilde{\lambda}_{h-2}}, \dots, s_{\tilde{\lambda}_{h-1}} s_{\tilde{\lambda}_{h-2}} \dots s_{\tilde{\lambda}_{h-1+1}}\}. \end{aligned} \quad (5.3.5)$$

By replacing the (+) notations by the (-) notations in various places in the proof of (2), (2.3.2) with $B^* = X^*$ becomes

$$z = x_{\lambda_{(h)}^-} T_{(\tilde{\lambda}_{h-1}, \tilde{\lambda}_{h-1+1})}^{\triangleright} T_{d_A} c_{A^*} \Sigma_A = \sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} z_k,$$

where, for each $k \in [1, n]$ with $a_{h,k} > 0$ and by Lemma 3.4(2),

$$\begin{aligned}
z_k &= \sum_{j=\overleftarrow{\mathfrak{r}}_h^{k-1}}^{\overleftarrow{\mathfrak{r}}_h^{k-1}-1} x_{\lambda_{(h)}^-} T_{\tilde{\lambda}_{h-1}} T_{\tilde{\lambda}_{h-2}} \cdots T_{\tilde{\lambda}_{h-j}} T_{d_{A_{h,k}^-}} c_{A^*} \Sigma_A \\
&= x_{\lambda_{(h)}^-} T_{\tilde{\lambda}_h} T_{\tilde{\lambda}_{h+1}} \cdots T_{\tilde{\lambda}_h + \overleftarrow{\mathfrak{r}}_{h+1}^k - 1} T_{d_{A_{h,k}^-}} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A \\
&= q^{\overleftarrow{\mathfrak{r}}_{h+1}^k} x_{\lambda_{(h)}^- \setminus \nu^-} T_{d_{A_{h,k}^-}} x_{\nu^-} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A,
\end{aligned} \tag{5.3.6}$$

since $s_{\tilde{\lambda}_h}, s_{\tilde{\lambda}_{h+1}}, \dots, s_{\tilde{\lambda}_h + \overleftarrow{\mathfrak{r}}_{h+1}^k - 1}$ are all in $\mathfrak{S}_{\lambda_{(h)}^-}$. (Recall from Notation 5.2 that $\tilde{a} = \tilde{a}_{h,k}$, $a = a_{h,k}$, and $b = a_{h+1,k}$.) Thus, by (5.2.3),

$$\begin{aligned}
z_k &= q^{\overleftarrow{\mathfrak{r}}_{h+1}^k} x_{\lambda_{(h)}^- \setminus \nu^-} T_{d_{A_{h,k}^-}} x_{\mathcal{D}_-} x_{\delta^-} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A \\
&= q^{\overleftarrow{\mathfrak{r}}_{h+1}^k} x_{\lambda_{(h)}^- \setminus \nu^-} T_{d_{A_{h,k}^-}} x_{\mathcal{D}_-} (c_{A^*})' x_{\delta^-} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} \Sigma_A \quad (x_{\delta^-} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} = x_{\nu^-}, (2.2.7)) \\
&= q^{\overleftarrow{\mathfrak{r}}_{h+1}^k} x_{\lambda_{(h)}^- \setminus \nu^-} T_{d_{A_{h,k}^-}} x_{\mathcal{D}_-} (c_{A^*})' x_{\delta^-} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \Sigma_{A_{h,k}^-} \quad (\text{by Lemma 3.5(2)}) \\
&= (-1)^{(a_{h,k}^{\bar{1}} + a_{h+1,k}^{\bar{1}}) \tilde{a}_{h-1,k}^{\bar{1}}} q^{\overleftarrow{\mathfrak{r}}_{h+1}^k} x_{\lambda_{(h)}^- \setminus \nu^-} T_{d_{A_{h,k}^-}} \cdot x_{\mathcal{D}_-} (c_{A^*,k}^{h,h+1})' x_{\delta^-} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \cdot \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^-}.
\end{aligned} \tag{5.3.7}$$

For later use, we extract from (5.3.6) and (5.3.7) the element

$$\mathfrak{M}^- := x_{\nu^-} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A = (-1)^{(a_{h,k}^{\bar{1}} + a_{h+1,k}^{\bar{1}}) \tilde{a}_{h-1,k}^{\bar{1}}} \underbrace{x_{\mathcal{D}_-} (c_{A^*,k}^{h,h+1})' x_{\delta^-} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft}}_{=\mathfrak{m}^-} \cdot \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^-}.$$

Note that $z_k = q^{\overleftarrow{\mathfrak{r}}_{h+1}^k} x_{\lambda_{(h)}^- \setminus \nu^-} T_{d_{A_{h,k}^-}} \mathfrak{M}^-$.

We summarise the computation of \mathfrak{m}^- , \mathfrak{M}^- and z_k corresponding to the four values of $(x, y) := (a_{h,k}^{\bar{1}}, a_{h+1,k}^{\bar{1}})$ in Table 1.

(Table 1)

(x, y)	\mathfrak{m}^-	\mathfrak{M}^-	z_k
$(0, 0)$	$\mathfrak{m}^- = \llbracket b+1 \rrbracket_q x_{\nu^-}$	$\mathfrak{M}^- = \llbracket b+1 \rrbracket_q x_{\nu^-} c_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} A^{\bar{1}})} \Sigma_{A_{h,k}^-}$	$z_k = q^{\overleftarrow{\mathfrak{r}}_{h+1}^k} \llbracket b+1 \rrbracket_q T_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} A^{\bar{1}})}$
$(1, 0)$	$\mathfrak{m}^- = \llbracket b+1 \rrbracket_q x_{\nu^-} c_{q, \tilde{a}-a+1, \tilde{a}-1} + q^{a-1} x_{\nu^-} c_{q, \tilde{a}, \tilde{a}+b}$	$\mathfrak{M}^- = \llbracket b+1 \rrbracket_q x_{\nu^-} c_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} A^{\bar{1}})} \Sigma_{A_{h,k}^-} + q^{a-1} x_{\nu^-} c_{(A^{\bar{0}} A^{\bar{1}} - E_{h,k} + E_{h+1,k})} \Sigma_{A_{h,k}^-}$	$z_k = q^{\overleftarrow{\mathfrak{r}}_{h+1}^k} \llbracket b+1 \rrbracket_q T_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} A^{\bar{1}})} + q^{\overleftarrow{\mathfrak{r}}_{h+1}^k + a - 1} T_{(A^{\bar{0}} A^{\bar{1}} - E_{h,k} + E_{h+1,k})}$
$(0, 1)$	$\mathfrak{m}^- = \llbracket b \rrbracket_q x_{\nu^-} (c_{q, \tilde{a}, \tilde{a}+b})$	$\mathfrak{M}^- = \llbracket b \rrbracket_q x_{\nu^-} c_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} A^{\bar{1}})} \Sigma_{A_{h,k}^-}$	$z_k = q^{\overleftarrow{\mathfrak{r}}_{h+1}^k} \llbracket b \rrbracket_q T_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} A^{\bar{1}})}$
$(1, 1)$	$\mathfrak{m}^- = \llbracket b \rrbracket_q x_{\nu^-} (c_{q, \tilde{a}-a+1, \tilde{a}-1}) (c_{q, \tilde{a}, \tilde{a}+b}) - q^{a-2} \llbracket b+1 \rrbracket_{q^2} x_{\nu^-}$	$\mathfrak{M}^- = \llbracket b \rrbracket_q x_{\nu^-} c_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} A^{\bar{1}})} \Sigma_{A_{h,k}^-} - q^{a-2} \llbracket b+1 \rrbracket_{q^2} x_{\nu^-} c_{(A^{\bar{0}} + 2E_{h+1,k} A^{\bar{1}} - E_{h,k} - E_{h+1,k})} \Sigma_{A_{h,k}^-}$	$z_k = q^{\overleftarrow{\mathfrak{r}}_{h+1}^k} \llbracket b \rrbracket_q T_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} A^{\bar{1}})} - q^{\overleftarrow{\mathfrak{r}}_{h+1}^k + a - 2} \llbracket b+1 \rrbracket_{q^2} T_{(A^{\bar{0}} + 2E_{h+1,k} A^{\bar{1}} - E_{h,k} - E_{h+1,k})}$

Let

$$\begin{aligned}
z'_k &= q^{\overleftarrow{\mathfrak{r}}_{h+1}^k} \llbracket a_{h+1,k}^{\bar{0}} + 1 \rrbracket_q T_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} | A^{\bar{1}})} + q^{\overleftarrow{\mathfrak{r}}_{h+1}^k + a_{h,k} - 1} T_{(A^{\bar{0}} | A^{\bar{1}} - E_{h,k} + E_{h+1,k})} \\
&\quad - q^{\overleftarrow{\mathfrak{r}}_{h+1}^k + a_{h,k} - 2} \llbracket a_{h+1,k} + 1 \rrbracket_{q^2} T_{(A^{\bar{0}} + 2E_{h+1,k} | A^{\bar{1}} - E_{h,k} - E_{h+1,k})}.
\end{aligned}$$

Then, by the notational Convention 5.1, if $(a_{h,k}^{\bar{1}}, a_{h+1,k}^{\bar{1}}) = (0, 0)$, then $b = a_{h+1,k}^{\bar{0}}$ and the second and third summands in z'_k are 0. Hence, z'_k is the z_k of line $(0, 0)$ in Table 1. The other three cases can be checked easily. This completes the proof of (3). \square

Using the notational Convention 5.1, the element \mathfrak{M}^- can be unified as follows. This multiplication formulas in $\mathcal{H}_{r,R}^c$ will be used in the proof of Theorem 6.4(2).

Corollary 5.4. *Maintain the notation in Theorem 5.3. We have in $\mathcal{H}_{r,R}^c$*

$$\begin{aligned} x_{\nu_{A_{h,k}^-}} T_{(\tilde{a}_{h,k}-1, \tilde{a}_{h,k}-a_{h,k}+1)}^{\triangleright} c_{A^*} \Sigma_A &= \llbracket a_{h+1,k}^{\bar{0}} + 1 \rrbracket_q x_{\nu_{A_{h,k}^-}} c_{(A^{\bar{0}}-E_{h,k}+E_{h+1,k}|A^{\bar{1}})} \Sigma_{A_{h,k}^-} \\ &+ q^{a_{h,k}-1} x_{\nu_{A_{h,k}^-}} c_{(A^{\bar{0}}|A^{\bar{1}}-E_{h,k}+E_{h+1,k})} \Sigma_{A_{h,k}^-} \\ &- q^{a_{h,k}-2} \llbracket a_{h+1,k} + 1 \rrbracket_{q^2, q} x_{\nu_{A_{h,k}^-}} c_{(A^{\bar{0}}+2E_{h+1,k}|A^{\bar{1}}-E_{h,k}-E_{h+1,k})} \Sigma_{A_{h,k}^-}. \end{aligned} \quad (5.4.1)$$

Remark 5.5. If $A^{\bar{1}} = 0$, then the multiplication formulas coincide with those for quantum \mathfrak{gl}_n (cf. [3, Lem. 3.4]).

6. MULTIPLICATION FORMULAS IN $\mathcal{Q}_q(n, r; R)$: THE ODD CASE

In this section, we derive the multiplication formulas for $\phi_{X^*} \phi_{A^*}$ under the SDP condition, where X^* is one of the matrices

$$(\lambda - E_{h,h}|E_{h,h}), \quad (\lambda - E_{h+1,h+1}|E_{h,h+1}), \quad (\lambda - E_{h,h}|E_{h+1,h}). \quad (6.0.1)$$

We also describe the general case using the following order relation.

Recall from [3] the partial order ' \preceq ' defined over $M_n(\mathbb{N})$. For $A, B \in M_n(\mathbb{N})$, we say $B \preceq A$ if the following conditions hold:

- (1) $\sum_{i \leq s, j \geq t} b_{i,j} \leq \sum_{i \leq s, j \geq t} a_{i,j}$ for all $1 \leq s < t \leq n$;
- (2) $\sum_{i \geq s, j \leq t} b_{i,j} \leq \sum_{i \geq s, j \leq t} a_{i,j}$ for all $1 \leq t < s \leq n$.

In particular, $B \prec A$ if one of the inequalities is strict.

For $A, B \in M_n(\mathbb{N})_r$, if d_A, d_B are the associated double coset representatives, then we have, by [8, Lem. 13.20],

$$d_A < d_B \implies A \prec B. \quad (6.0.2)$$

For any $\mu \in \Lambda(n, r)$, denote by $\mathcal{H}_{\mu,R}^c$ the subsuperalgebra of $\mathcal{H}_{r,R}^c$ associated with the Young subgroup \mathfrak{S}_μ and the full Clifford superalgebra \mathcal{C}_r .

Lemma 6.1 ([18, Cors 3.5 & 3.8]). *For any given $\lambda, \mu \in \Lambda(n, r)$, the right (resp. left) $\mathcal{H}_{r,R}^c$ -module $x_\lambda \mathcal{H}_{r,R}^c$ (resp. $\mathcal{H}_{r,R}^c x_\mu$) admits a basis $\{x_\lambda T_d c^\alpha T_v \mid d \in \mathcal{D}_{\lambda,\mu}, \alpha \in \mathbb{N}_2^r, v \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu\}$ (resp. $\{T_u c^\alpha T_d x_\mu \mid u \in \mathcal{D}_{\nu(d-1)}^{-1} \cap \mathfrak{S}_\lambda, \alpha \in \mathbb{N}_2^r, d \in \mathcal{D}_{\lambda,\mu}\}$), where $\mathfrak{S}_{\nu(d)} = d^{-1} \mathfrak{S}_\lambda d \cap \mathfrak{S}_\mu$ and $\mathfrak{S}_{\nu(d-1)} = \mathfrak{S}_\lambda \cap d \mathfrak{S}_\mu d^{-1}$. Moreover*

$$x_\lambda T_d = \sum_{u' \in \mathcal{D}_{\nu(d-1)}^{-1} \cap \mathfrak{S}_\lambda} T_{u'} T_d x_{\nu(d)} \quad (6.1.1)$$

$$\mathcal{H}_{\mu,R}^c = \bigoplus_{\sigma \in \mathcal{D}_{\nu(d)} \cap \mathfrak{S}_\mu} \mathcal{H}_{\nu(d),R}^c T_\sigma \quad (6.1.2)$$

for each $d \in \mathcal{D}_{\lambda,\mu}$.

Recall the notations in (3.0.2), and Notation 5.2. We also apply (2.2.4) to define the partial sum matrix $\tilde{A}^{\bar{1}} = (\tilde{a}_{i,j}^{\bar{1}})$. Recall also the SDP condition in Definition 4.2.

Theorem 6.2. *For $h \in [1, n]$ and $A^* = (A^{\bar{0}}|A^{\bar{1}}) = (a_{i,j}^{\bar{0}}|a_{i,j}^{\bar{1}}) \in M_n(\mathbb{N}|\mathbb{N}_2)_r$, let $A = A^{\bar{0}} + A^{\bar{1}}$, $\lambda = \text{ro}(A)$, and $\vec{\mathbf{r}}_h^k = \vec{\mathbf{r}}_h^k(A)$. Let $D_h^* = (\lambda - E_{h,h}|E_{h,h})$.*

(1) Assume that A satisfies the SDP condition on the h -th row. Then we have in $\mathcal{Q}_q(n, r; R)$

$$\phi_{D_h^*} \phi_{A^*} = \sum_{k=1}^n (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\tilde{\mathbf{r}}_h^k} \left\{ \phi_{(A^{\bar{0}} - E_{h,k} | A^{\bar{1}} + E_{h,k})} - \llbracket a_{h,k} \rrbracket_{q^2} \phi_{(A^{\bar{0}} + E_{h,k} | A^{\bar{1}} - E_{h,k})} \right\} =: {}_{\text{SDP}} \mathbf{HK}.$$

(2) In general, we have

$$\phi_{D_h^*} \phi_{A^*} = {}_{\text{SDP}} \mathbf{HK} + \sum_{\substack{B^* \in M_n(\mathbb{N} | \mathbb{N}_2)_r \\ \llbracket B^* \rrbracket < A}} f_{B^*}^{D_h^*, A^*} \phi_{B^*} \quad (f_{B^*}^{D_h^*, A^*} \in R).$$

Proof. Let $X^* = D_h^* = (\lambda - E_{h,h} | E_{h,h})$. Then $X = \llbracket D_h^* \rrbracket = \text{diag}(\lambda)$. Clearly $\text{ro}(X) = \lambda$, $d_X = 1$, $\mathcal{D}_{\nu_X} \cap \mathfrak{S}_\lambda = \{1\}$, and $c_{X^*} = c_{q, \tilde{\lambda}_{h-1} + 1, \tilde{\lambda}_h}$. Then, for $B^* = X^*$ and $\mu = \text{co}(A)$, (2.3.2) becomes

$$\begin{aligned} z &:= \phi_{X^*} \phi_{A^*}(x_\mu) = x_\lambda c_{q, \tilde{\lambda}_{h-1} + 1, \tilde{\lambda}_h} T_{d_A} c_{A^*} \Sigma_A \\ &= x_\lambda (q^{\lambda_h - 1} c_{\tilde{\lambda}_{h-1} + 1} + q^{\lambda_h - 2} c_{\tilde{\lambda}_{h-1} + 2} + \cdots + q c_{\tilde{\lambda}_{h-1}} + c_{\tilde{\lambda}_h}) T_{d_A} c_{A^*} \Sigma_A \\ &= \sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} x_\lambda \left(\sum_{j=\tilde{\mathbf{r}}_h^k + 1}^{\tilde{\mathbf{r}}_h^{k+1}} q^{\lambda_h - j} c_{\tilde{\lambda}_{h-1} + j} \right) T_{d_A} c_{A^*} \Sigma_A \\ &= \sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} x_\lambda q^{\lambda_h - \tilde{\mathbf{r}}_h^{k+1}} (c_{q, \hat{a}_{h,k-1} + 1, \hat{a}_{h,k-1} + a_{h,k}} T_{d_A}) c_{A^*} \Sigma_A \quad (\text{as } \tilde{\lambda}_{h-1} + \tilde{\mathbf{r}}_h^k = \hat{a}_{h,k-1}), \end{aligned} \quad (6.2.1)$$

where the tail Σ_A of T_{A^*} is defined in (2.2.13).) Applying the SDP condition (4.2.1) may turn z into a summation:

$$z = \sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} q^{\lambda_h - \tilde{\mathbf{r}}_h^{k+1}} x_\lambda (T_{d_A} c_{q, \tilde{a}_{h-1,k} + 1, \tilde{a}_{h-1,k} + a_{h,k}}) c_{A^*} \Sigma_A = \sum_{1 \leq k \leq n, a_{h,k} > 0} z_k,$$

where, for each $1 \leq k \leq n$ with $a_{h,k} > 0$,

$$z_k = q^{\lambda_h - \tilde{\mathbf{r}}_h^{k+1}} x_\lambda T_{d_A} \cdot \underbrace{c_{q, \tilde{a}_{h-1,k} + 1, \tilde{a}_{h-1,k} + a_{h,k}}}_{=\mathbf{m}} c_{A^*} \cdot \Sigma_A. \quad (6.2.2)$$

If we decompose $c_{A^*} = \overleftarrow{c}_{A^*}^{h,k} c_{q, \tilde{a}_{h-1,k} + 1, \tilde{a}_{h-1,k} + a_{h,k}}^{a_{h,k}^{\bar{1}}} \overrightarrow{c}_{A^*}^{h,k}$, similar to (5.2.1), then the middle part

$$\mathbf{m} = (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} \left(\overleftarrow{c}_{A^*}^{h,k} c_{q, \tilde{a}_{h-1,k} + 1, \tilde{a}_{h-1,k} + a_{h,k}}^{a_{h,k}^{\bar{1}}} \overrightarrow{c}_{A^*}^{h,k} \right).$$

If $a_{h,k}^{\bar{1}} = 0$, then $\mathbf{m} = (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} c_{M^*}$, where $M^* = (A^{\bar{0}} - E_{h,k} | A^{\bar{1}} + E_{h,k})$. Since A^* and M^* have the same base so that $d_A = d_M$, $\Sigma_A = \Sigma_M$, it follows that

$$z_k = (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\lambda_h - \tilde{\mathbf{r}}_h^{k+1}} x_\lambda T_{d_M} c_{M^*} \Sigma_M = (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\lambda_h - \tilde{\mathbf{r}}_h^{k+1}} T_{(A^{\bar{0}} - E_{h,k} | A^{\bar{1}} + E_{h,k})}.$$

If $a_{h,k}^{\bar{1}} = 1$, then

$$\begin{aligned} \mathbf{m} &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} \left(\overleftarrow{c}_{A^*}^{h,k} (c_{q, \tilde{a}_{h-1,k} + 1, \tilde{a}_{h-1,k} + a_{h,k}})^2 \overrightarrow{c}_{A^*}^{h,k} \right) \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} \left(-\llbracket a_{h,k} \rrbracket_{q^2} \right) \left(\overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h,k} \right) \quad (\text{Lemma 3.7(4)}) \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + 1} \llbracket a_{h,k} \rrbracket_{q^2} c_{N^*}, \quad \text{where } N^* = (A^{\bar{0}} + E_{h,k} | A^{\bar{1}} - E_{h,k}). \end{aligned}$$

Now, A^* and N^* have the same base so that $d_A = d_N$, $\Sigma_A = \Sigma_N$. Hence, (6.2.2) becomes

$$z_k = (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + 1} q^{\lambda_h - \tilde{\mathbf{r}}_h^{k+1}} \llbracket a_{h,k} \rrbracket_{q^2} x_\lambda T_{d_N} c_{N^*} \Sigma_N = (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + 1} q^{\lambda_h - \tilde{\mathbf{r}}_h^{k+1}} \llbracket a_{h,k} \rrbracket_{q^2} T_{(A^{\bar{0}} + E_{h,k} | A^{\bar{1}} - E_{h,k})}.$$

Finally, putting $z'_k = (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\tilde{\mathbf{r}}_h^k} (T_{(A^{\bar{0}} - E_{h,k} | A^{\bar{1}} + E_{h,k})} - \llbracket a_{h,k} \rrbracket_{q^2} T_{(A^{\bar{0}} + E_{h,k} | A^{\bar{1}} - E_{h,k})})$, one checks easily $z'_k = z_k$, by Convention 5.1. This proves (1).

We now prove the general case (2). Applying Lemma 4.5(2) to (6.2.1) yields

$$\begin{aligned} z &= \sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} x_\lambda \left(\sum_{p=1}^{a_{h,k}} q^{\lambda_h - \overleftarrow{\mathbf{r}}_h^k - p} (T_{d_A} c_{\bar{a}_{h-1,k}+p} + \sum_{w < d_A, j \in [1,r]} f_{w,j}^{\cdots} T_w c_j) \right) c_{A^*} \Sigma_A \\ &=_{\text{SDP}} \text{HK} + \text{TK} \end{aligned}$$

where ${}_{\text{SDP}}\text{HK} = {}_{\text{SDP}}\text{HK}(x_\mu)$, $\Sigma_A = \sum_{v \in \mathcal{D}_{\nu_A} \cap \mathfrak{S}_\mu}$ and

$$\text{TK} = \sum_{1 \leq k \leq n, a_{h,k} > 0} \sum_{p=1}^{a_{h,k}} \sum_{w < d_A, j \in [1,r]} f_{w,j}^{\cdots} q^{\lambda_h - \overleftarrow{\mathbf{r}}_h^k - p} x_\lambda T_w c_j c_{A^*} \Sigma_A.$$

Since every w in the expression of TK satisfies $w < d_A$ and may be written as $w = xdy$ with $w \in \mathfrak{S}_\lambda$, $d \in \mathcal{D}_{\lambda,\mu}$, and $y \in \mathfrak{S}_\mu$, it follows that $d < d_A$. Also, $c_j c_{A^*} \Sigma_A \in \mathcal{H}_{\mu,R}^c$. By [18, Cor. 3.8] (or Lemma 6.1), TK is a linear combination of basis elements of the form $x_\lambda T_d c^\alpha T_v$ for $x_\lambda \mathcal{H}_{r,R}^c$, where $d \in \mathcal{D}_{\lambda,\mu}$ with $d < d_A$, $\alpha \in \mathbb{N}_2^r$ and $v \in \mathcal{D}_{\nu_A} \cap \mathfrak{S}_\mu$.

On the other hand, since both z and ${}_{\text{SDP}}\text{HK}$ are in $x_\lambda \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$, it follows that

$$\text{TK} = z - {}_{\text{SDP}}\text{HK} \in x_\lambda \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu.$$

Thus, by Proposition 2.3(1), $\text{TK} = \sum_{M^* \in M_n(\mathbb{N}|\mathbb{N}_2)_{\lambda,\mu}} f_{M^*}^{D_h^*, A^*} T_{M^*} (f_{M^*}^{D_h^*, A^*} \in R)$, which is also a linear combination of the same basis for $x_\lambda \mathcal{H}_{r,R}^c$.

Equating coefficients and using [8, Lem. 13.20] or (6.0.2) conclude that

$$\text{TK} = \sum_{B^* \in M_n(\mathbb{N}|\mathbb{N}_2)_r, d_{[B^*]} < d_{[A^*]}} f_{B^*}^{D_h^*, A^*} T_{B^*} = \sum_{\substack{B^* \in M_n(\mathbb{N}|\mathbb{N}_2)_r \\ [B^*] \prec A}} f_{B^*}^{D_h^*, A^*} \phi_{B^*}(x_{\text{co}(B^*)}),$$

proving (2). \square

We now tackle the remaining odd cases. The proof is parallel to the even cases in Section 5 under the SDP condition, but with extra care of the c -elements. The general case are even more complicated.

Theorem 6.3. *Let $h \in [1, n-1]$ and $A^* = (A^{\bar{0}}|A^{\bar{1}}) = (a_{i,j}^{\bar{0}}|a_{i,j}^{\bar{1}}) \in M_n(\mathbb{N}|\mathbb{N}_2)_r$ with base $A = A^{\bar{0}} + A^{\bar{1}}$, $\lambda = \text{ro}(A)$, and $\overrightarrow{\mathbf{r}}_h^k = \overrightarrow{\mathbf{r}}_h^k(A)$. Let $E_h^* = (\lambda - E_{h+1,h+1}|E_{h,h+1})$.*

(1) *Suppose that, for every $k \in [1, n]$ such that $a_{h+1,k} > 0$, A satisfies the SDP condition at (h, k) if $a_{h,k} > 0$ and satisfies $A_{\bar{1}}^{h,k} = 0$ if $a_{h,k} = 0$.³ Then we have in $\mathcal{Q}_q(n, r; R)$*

$$\begin{aligned} \phi_{E_h^*} \phi_{A^*} &= \sum_{k=1}^n \left\{ (-1)^{\bar{a}_{h-1,k}^{\bar{1}}} q^{\overrightarrow{\mathbf{r}}_h^k + a_{h+1,k}^{\bar{1}}} \phi_{(A^{\bar{0}} - E_{h+1,k}|A^{\bar{1}} + E_{h,k})} \right. \\ &\quad + (-1)^{\bar{a}_{h-1,k}^{\bar{1}} + 1 - a_{h,k}^{\bar{1}}} q^{\overrightarrow{\mathbf{r}}_h^k} \llbracket a_{h,k}^{\bar{0}} + 1 \rrbracket_q \phi_{(A^{\bar{0}} + E_{h,k}|A^{\bar{1}} - E_{h+1,k})} \\ &\quad \left. + (-1)^{\bar{a}_{h-1,k}^{\bar{1}}} q^{\overrightarrow{\mathbf{r}}_h^k - 1 + a_{h+1,k}^{\bar{1}}} \llbracket a_{h,k} + 1 \rrbracket_{q^2} \phi_{(A^{\bar{0}} + 2E_{h,k} - E_{h+1,k}|A^{\bar{1}} - E_{h,k})} \right\} \\ &=: {}_{\text{SDP}}\text{HE}. \end{aligned} \tag{6.3.1}$$

(2) *In general, we have*

$$\phi_{E_h^*} \phi_{A^*} = {}_{\text{SDP}}\text{HE} + \sum_{\substack{B^* \in M_n(\mathbb{N}|\mathbb{N}_2)_r \\ \exists k, [B^*] \prec A_{h,k}^+}} f_{B^*}^{E_h^*, A^*} \phi_{B^*}.$$

³This hypothesis is equivalent to that, for every $k \in [1, n]$, $A_{h,k}^+$ satisfies the SDP condition at (h, k) .

Proof. Let $X^* = E_h^* = (\lambda - E_{h+1,h+1} | E_{h,h+1})$. Then its base $X = \lambda + E_{h,h+1} - E_{h+1,h+1}$ which is the same base of the matrix X^* in Theorem 5.3(2). Thus, we have $\text{ro}(X) = \lambda_{(h)}^+$, $c_{X^*} = c_{\lambda_{h+1}^+}$. Similar to Theorem 5.3(2), the equation (2.3.2) with $B^* = X^*$ (and $a = a_{h,k}$, $b = a_{h+1,k}$) becomes

$$\begin{aligned} z &= \sum_{\substack{1 \leq k \leq n \\ a_{h+1,k} > 0}} x_{\lambda_{(h)}^+} c_{\lambda_{h+1}^+} \left(\sum_{j=\overleftarrow{\mathbf{r}}_{h+1}^k}^{\overleftarrow{\mathbf{r}}_{h+1}^{k+1}-1} T_{\lambda_{h+1}}^- T_{\lambda_{h+2}}^- \cdots T_{\lambda_{h+j}}^- T_{d_A} \right) c_{A^*} \Sigma_A \\ &= \sum_{\substack{1 \leq k \leq n \\ a_{h+1,k} > 0}} x_{\lambda_{(h)}^+} c_{\lambda_{h+1}^+} \left(T_{\lambda_h}^- T_{\lambda_{h-1}}^- \cdots T_{\lambda_{h-\overrightarrow{\mathbf{r}}_h^k+1}}^- T_{d_{A_{h,k}^+}} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} \right) c_{A^*} \Sigma_A \quad (\text{Lemma 3.4}) \quad (6.3.2) \\ &= \sum_{\substack{1 \leq k \leq n \\ a_{h+1,k} > 0}} x_{\lambda_{(h)}^+} T_{\lambda_h}^- T_{\lambda_{h-1}}^- \cdots T_{\lambda_{h-\overrightarrow{\mathbf{r}}_h^k+1}}^- (c_{\lambda_{h-\overrightarrow{\mathbf{r}}_h^k+1}}^- T_{d_{A_{h,k}^+}}) T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} c_{A^*} \Sigma_A. \end{aligned}$$

Observe that $\tilde{\lambda}_h - \overrightarrow{\mathbf{r}}_h^k + 1 = \tilde{\lambda}_{h-1} + \overleftarrow{\mathbf{r}}_h^{k+1} + 1 = \hat{a}_{h,k-1} + a_{h,k} + 1 = \hat{a}_{h,k-1} + a_{h,k}^+$, where $a_{i,j}^+$ is the (i,j) -entry of $A_{h,k}^+$ (thus, $a_{h,k-1}^+ = a_{h,k-1}$ and $a_{h,k}^+ = a_{h,k} + 1$). Since the hypothesis in (1) together with Theorem 4.6 implies that $A_{h,k}^+$ satisfies the SDP condition at (h,k) , for all $k \in [1, n]$ with $a_{h,k}^+ > 0$, it follows that

$$c_{\lambda_{h-\overrightarrow{\mathbf{r}}_h^k+1}}^- T_{d_{A_{h,k}^+}} = T_{d_{A_{h,k}^+}} c_{\tilde{a}_{h-1,k}^+ + a_{h,k}^+} = T_{d_{A_{h,k}^+}} c_{\tilde{a}+1}.$$

Hence, since $x_{\lambda_{(h)}^+} T_{\lambda_h}^- T_{\lambda_{h-1}}^- \cdots T_{\lambda_{h-\overrightarrow{\mathbf{r}}_h^k+1}}^- = q^{\overrightarrow{\mathbf{r}}_h^k} x_{\lambda_{(h)}^+}$, we have

$$z = \sum_{\substack{1 \leq k \leq n \\ a_{h+1,k} > 0}} z_k, \quad \text{where } z_k = q^{\overrightarrow{\mathbf{r}}_h^k} x_{\lambda_{(h)}^+} T_{d_{A_{h,k}^+}} c_{\tilde{a}+1} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} c_{A^*} \Sigma_A,$$

and, similar to the even case, using an argument for turning Σ_A to $\Sigma_{A_{h,k}^+}$ right above (5.3.2) together with the fact $x_{\delta} + c_{\tilde{a}+1} = c_{\tilde{a}+1} x_{\delta}$ gives rise to

$$z_k = (-1)^{(a_{h,k}^{\bar{1}} + a_{h+1,k}^{\bar{1}}) \tilde{a}_{h-1,k}^{\bar{1}}} q^{\overrightarrow{\mathbf{r}}_h^k} x_{\lambda_{(h)}^+} \setminus_{\nu^+} T_{d_{A_{h,k}^+}} \cdot \underbrace{x_{\mathcal{D}_+} c_{\tilde{a}+1} (c_{A^*;k}^{h,h+1})' x_{\delta} + T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright}}_{=\underline{\mathbf{m}}^+} \cdot \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+}. \quad (6.3.3)$$

Similar to the even situation, we need to analyse the middle part $\underline{\mathbf{m}}^+$ in four cases.

Case 1. $a_{h,k}^{\bar{1}} = 0$, $a_{h+1,k}^{\bar{1}} = 0$. In this case we have $(c_{A^*;k}^{h,h+1})' = 1$ and, by Lemma 3.7(3),

$$\underline{\mathbf{m}}^+ = x_{\mathcal{D}_+} c_{\tilde{a}+1} x_{\delta} + T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} = x_{\nu^+} c_{\tilde{a}+1} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} = x_{\nu^+} c_{q, \tilde{a}-a+1, \tilde{a}+1}.$$

Thus, if $M^* = (A^{\bar{0}} - E_{h+1,k} | A^{\bar{1}} + E_{h,k})$, then M^* has the base $M = A_{h,k}^+$ and $\text{ro}(M) = \lambda_{(h)}^+$. Moreover, we have $c_{M^*} = \overleftarrow{c}_{A^*}^{h,k} c_{q, \tilde{a}-a+1, \tilde{a}+1} \overrightarrow{c}_{A^*}^{h+1,k}$. Hence,

$$\begin{aligned} z_k &= q^{\overrightarrow{\mathbf{r}}_h^k} x_{\lambda_{(h)}^+} \setminus_{\nu^+} T_{d_{A_{h,k}^+}} x_{\nu^+} c_{q, \tilde{a}-a+1, \tilde{a}+1} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+} \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\overrightarrow{\mathbf{r}}_h^k} x_{\lambda_{(h)}^+} d_M \overleftarrow{c}_{A^*}^{h,k} c_{q, \tilde{a}-a+1, \tilde{a}+1} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_M \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\overrightarrow{\mathbf{r}}_h^k} T_{(A^{\bar{0}} - E_{h+1,k} | A^{\bar{1}} + E_{h,k})}. \end{aligned}$$

Case 2. $a_{h,k}^{\bar{1}} = 0$, $a_{h+1,k}^{\bar{1}} = 1$. Then $(c_{A^*;k}^{h,h+1})' = (c_{q, \tilde{a}+1, \tilde{a}+b})' = c_{\tilde{a}+1} + q(c_{q, \tilde{a}+2, \tilde{a}+b})'$ and

$$\begin{aligned} \underline{\mathbf{m}}^+ &= q x_{\mathcal{D}_+} c_{\tilde{a}+1} (c_{q, \tilde{a}+2, \tilde{a}+b})' x_{\delta} + T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} + x_{\mathcal{D}_+} c_{\tilde{a}+1}^2 x_{\delta} + T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} \\ &= q x_{\nu^+} c_{\tilde{a}+1} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} c_{q, \tilde{a}+2, \tilde{a}+b} - x_{\nu^+} T_{(\tilde{a}, \tilde{a}-a+1)}^{\triangleright} \\ &= q x_{\nu^+} c_{q, \tilde{a}-a+1, \tilde{a}+1} c_{q, \tilde{a}+2, \tilde{a}+b} - \llbracket a+1 \rrbracket_q x_{\nu^+}. \end{aligned}$$

Substituting into (6.3.3) and noting that the above M^* and $N^* = (A^{\bar{0}} + E_{h,k} | A^{\bar{1}} - E_{h+1,k})$ have the same base $A_{h,k}^+$ and

$$c_{q,\tilde{a}-a+1,\tilde{a}+1} c_{q,\tilde{a}+2,\tilde{a}+b} \cdot \overleftarrow{c_{A^*}^{h,k}} \overrightarrow{c_{A^*}^{h+1,k}} = \overleftarrow{c_{A^*}^{h,k}} c_{q,\tilde{a}-a+1,\tilde{a}+1} c_{q,\tilde{a}+2,\tilde{a}+b} \overrightarrow{c_{A^*}^{h+1,k}} = c_{M^*},$$

we obtain

$$\begin{aligned} z_k &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\vec{r}_h^k+1} x_{\lambda_{(h)}^+ \setminus \nu^+} T_{d_{A_{h,k}^+}} x_{\nu^+} c_{q,\tilde{a}-a+1,\tilde{a}+1} c_{q,\tilde{a}+2,\tilde{a}+b} \cdot \overleftarrow{c_{A^*}^{h,k}} \overrightarrow{c_{A^*}^{h+1,k}} \Sigma_{A_{h,k}^+} \\ &\quad - (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\vec{r}_h^k} \llbracket a+1 \rrbracket_q x_{\lambda_{(h)}^+ \setminus \nu^+} T_{d_{A_{h,k}^+}} x_{\nu^+} \overleftarrow{c_{A^*}^{h,k}} \overrightarrow{c_{A^*}^{h+1,k}} \Sigma_{A_{h,k}^+} \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\vec{r}_h^k+1} T_{(A^{\bar{0}}-E_{h+1,k} | A^{\bar{1}}+E_{h,k})} + (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\vec{r}_h^k} \llbracket a_{h,k}^{\bar{0}} + 1 \rrbracket_q T_{(A^{\bar{0}}+E_{h,k} | A^{\bar{1}}-E_{h+1,k})}. \end{aligned}$$

Case 3. $a_{h,k}^{\bar{1}} = 1$, $a_{h+1,k}^{\bar{1}} = 0$. It follows $c_{A^*}^{h,h+1} = c_{q,\tilde{a}-a+1,\tilde{a}}$ and the middle part has the form

$$z_k = (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\vec{r}_h^k} x_{\lambda_{(h)}^+ \setminus \nu^+} T_{d_{A_{h,k}^+}} \underbrace{x_{\mathcal{D}_+} c_{\tilde{a}+1}(c_{q,\tilde{a}-a+1,\tilde{a}})' x_{\delta} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright}}_{\underline{\mathbf{m}}^+} \overleftarrow{c_{A^*}^{h,k}} \overrightarrow{c_{A^*}^{h+1,k}} \Sigma_{A_{h,k}^+}.$$

Using (2.2.5), (5.2.3), Lemma 3.7(2)&(3), we have

$$\begin{aligned} \underline{\mathbf{m}}^+ &= q x_{\mathcal{D}_+} c_{\tilde{a}+1}(c_{q,\tilde{a}-a+1,\tilde{a}})' x_{\delta} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} \\ &= -x_{\mathcal{D}_+} x_{\delta} + q c_{q,\tilde{a}-a+1,\tilde{a}} c_{\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} \\ &= -x_{\nu^+} c_{q,\tilde{a}-a+1,\tilde{a}+1} c_{\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} + x_{\nu^+} (c_{\tilde{a}+1})^2 T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} \\ &= -(c_{q,\tilde{a}-a+1,\tilde{a}+1})' x_{\nu^+} c_{\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} - x_{\nu^+} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} \\ &= -x_{\nu^+} (c_{q,\tilde{a}-a+1,\tilde{a}+1})^2 - \llbracket a+1 \rrbracket_q x_{\nu^+} \quad (\text{by Lemma 3.7(3)}) \\ &= (\llbracket a+1 \rrbracket_{q^2} - \llbracket a+1 \rrbracket_q) x_{\nu^+}, \quad (\text{by Lemma 3.7(4)}). \end{aligned}$$

Hence, taking $N^* = (A^{\bar{0}} + 2E_{h,k} - E_{h+1,k} | A^{\bar{1}} - E_{h,k})$ which has base $N = A_{h,k}^+$ and $c_{N^*} = \overleftarrow{c_{A^*}^{h,k}} \overrightarrow{c_{A^*}^{h+1,k}}$, we have

$$\begin{aligned} z_k &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\vec{r}_h^k-1} x_{\lambda_{(h)}^+ \setminus \nu^+} T_{d_{A_{h,k}^+}} ((\llbracket a+1 \rrbracket_{q^2} - \llbracket a+1 \rrbracket_q) x_{\nu^+}) \overleftarrow{c_{A^*}^{h,k}} \overrightarrow{c_{A^*}^{h+1,k}} \Sigma_{A_{h,k}^+} \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\vec{r}_h^k-1} (\llbracket a_{h,k} + 1 \rrbracket_{q^2} - \llbracket a_{h,k} + 1 \rrbracket_q) T_{(A^{\bar{0}}+2E_{h,k}-E_{h+1,k} | A^{\bar{1}}-E_{h,k})}. \end{aligned}$$

Case 4. $a_{h,k}^{\bar{1}} = 1$, $a_{h+1,k}^{\bar{1}} = 1$. Then $(c_{A^*}^{h,h+1})' = c'_{q,\tilde{a}-a+1,\tilde{a}} c'_{q,\tilde{a}+1,\tilde{a}+b}$ and (6.3.3) takes the form

$$z_k = q^{\vec{r}_h^k-1} x_{\lambda_{(h)}^+ \setminus \nu^+} T_{d_{A_{h,k}^+}} (\underline{\mathbf{q}}\underline{\mathbf{m}}^+) \overleftarrow{c_{A^*}^{h,k}} \overrightarrow{c_{A^*}^{h+1,k}} \Sigma_{A_{h,k}^+}, \quad (6.3.4)$$

where $\underline{\mathbf{q}}\underline{\mathbf{m}}^+ = x_{\mathcal{D}_+} c_{\tilde{a}+1} c'_{q,\tilde{a}-a+1,\tilde{a}} c'_{q,\tilde{a}+1,\tilde{a}+b} x_{\delta} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright}$. Multiplying (5.3.4) by $c_{\tilde{a}+1}$ to the left yields

$$q c_{\tilde{a}+1} c'_{q,\tilde{a}-a+1,\tilde{a}} c'_{q,\tilde{a}+1,\tilde{a}+b} x_{\delta} = x_{\delta} (q c_{q,\tilde{a}+2,\tilde{a}+b} c_{\tilde{a}+1} c_{q,\tilde{a}-a+1,\tilde{a}+1} + q c_{q,\tilde{a}+2,\tilde{a}+b} + c_{q,\tilde{a}-a+1,\tilde{a}+1} - c_{\tilde{a}+1})$$

Substituting into $\underline{\mathbf{q}}\underline{\mathbf{m}}^+$ with a similar argument right after (5.3.4), we have

$$\begin{aligned} \underline{\mathbf{q}}\underline{\mathbf{m}}^+ &= q x_{\nu^+} c_{q,\tilde{a}+2,\tilde{a}+b} (-c_{q,\tilde{a}-a+1,\tilde{a}+1} c_{\tilde{a}+1} - 2) T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} + q x_{\nu^+} c_{q,\tilde{a}+2,\tilde{a}+b} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} \\ &\quad + x_{\nu^+} c_{q,\tilde{a}-a+1,\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} - x_{\nu^+} c_{\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} \\ &= -q x_{\nu^+} c_{q,\tilde{a}+2,\tilde{a}+b} c_{q,\tilde{a}-a+1,\tilde{a}+1} c_{\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} - q x_{\nu^+} c_{q,\tilde{a}+2,\tilde{a}+b} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} \\ &\quad + x_{\nu^+} c_{q,\tilde{a}-a+1,\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} - x_{\nu^+} c_{\tilde{a}+1} T_{(\tilde{a},\tilde{a}-a+1)}^{\triangleright} \\ &= -q c'_{q,\tilde{a}+2,\tilde{a}+b} c'_{q,\tilde{a}-a+1,\tilde{a}+1} x_{\nu^+} c_{q,\tilde{a}-a+1,\tilde{a}+1} - q c'_{q,\tilde{a}+2,\tilde{a}+b} \llbracket a+1 \rrbracket_q x_{\nu^+} \\ &\quad + c'_{q,\tilde{a}-a+1,\tilde{a}+1} \llbracket a+1 \rrbracket_q x_{\nu^+} - x_{\nu^+} c_{q,\tilde{a}-a+1,\tilde{a}+1} \\ &= q (\llbracket a+1 \rrbracket_{q^2} - \llbracket a+1 \rrbracket_q) x_{\nu^+} c_{q,\tilde{a}+2,\tilde{a}+b} + (\llbracket a+1 \rrbracket_q - 1) x_{\nu^+} c_{q,\tilde{a}-a+1,\tilde{a}+1}. \end{aligned}$$

Then (6.3.4) becomes after substitution

$$\begin{aligned} z_k &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\vec{r}_h^k} (\llbracket a+1 \rrbracket_{q^2} - \llbracket a+1 \rrbracket_q) x_{\lambda_{(h)}^+} T_{d_{A_{h,k}^+}} \overleftarrow{c}_{A^*}^{h,k} c_{q,\tilde{a}+2,\tilde{a}+b} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+} \\ &\quad + (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\vec{r}_h^k-1} (\llbracket a+1 \rrbracket_q - 1) x_{\lambda_{(h)}^+} T_{d_{A_{h,k}^+}} \overleftarrow{c}_{A^*}^{h,k} c_{q,\tilde{a}-a+1,\tilde{a}+1} \overrightarrow{c}_{A^*}^{h+1,k} \Sigma_{A_{h,k}^+}. \end{aligned}$$

Let $B^* = (b_{i,j}^{\bar{0}} | b_{i,j}^{\bar{1}}) = (A^{\bar{0}} + 2E_{h,k} - E_{h+1,k} | A^{\bar{1}} - E_{h,k})$. Then its base $B = (b_{i,j}) = A_{h,k}^+$ and $b_{h,k}^{\bar{1}} = 0, b_{h+1,k}^{\bar{1}} = 1$, and $b_{h+1,k} = a_{h+1,k} - 1 = b - 1$. Thus, $\tilde{b}_{h,k} + 1 = \tilde{a} + 2$, and $\tilde{b}_{h+1,k} = \tilde{a} + b$. Hence, $c_{B^*} = \overleftarrow{c}_{A^*}^{h,k} c_{q,\tilde{a}+2,\tilde{a}+b} \overrightarrow{c}_{A^*}^{h+1,k}$. Similarly, putting $C^* = (A^{\bar{0}} + E_{h,k} | A^{\bar{1}} - E_{h+1,k})$, we have $c_{C^*} = \overleftarrow{c}_{A^*}^{h,k} c_{q,\tilde{a}-a+1,\tilde{a}+1} \overrightarrow{c}_{A^*}^{h+1,k}$. Consequently,

$$\begin{aligned} z_k &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\vec{r}_h^k} (\llbracket a_{h,k} + 1 \rrbracket_{q^2} - \llbracket a_{h,k} + 1 \rrbracket_q) T_{(A^{\bar{0}}+2E_{h,k}-E_{h+1,k}|A^{\bar{1}}-E_{h,k})} \\ &\quad + (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{\vec{r}_h^k} \llbracket a_{h,k}^{\bar{0}} + 1 \rrbracket_q T_{(A^{\bar{0}}+E_{h,k}|A^{\bar{1}}-E_{h+1,k})} \quad (\text{noting } \llbracket a+1 \rrbracket_q - 1 = q \llbracket a_{h,k} \rrbracket_q). \end{aligned}$$

Finally, by using Convention 5.1, one checks easily $\sum_{\substack{1 \leq k \leq n \\ a_{h+1,k} > 0}} z_k$ is the image of the right hand side of (6.3.1) at $x_{\text{co}(A)}$. This proves (1).

In general, by Lemma 4.5(2), $c_{\tilde{\lambda}_h - \vec{r}_h^k + 1} T_{d_{A_{h,k}^+}} = T_{d_{A_{h,k}^+}} c_{\tilde{a}_{h,k} + 1} + \sum_{w < d_{A_{h,k}^+}, j \in [1,r]} f_{w,j}^{\dots} T_w c_j$. Thus, (6.3.2) takes the form

$$\begin{aligned} z &= \sum_{\substack{1 \leq k \leq n \\ a_{h+1,k} > 0}} q^{\vec{r}_h^k} x_{\lambda_{(h)}^+} \left(T_{d_{A_{h,k}^+}} c_{\tilde{a}_{h,k} + 1} + \sum_{w < d_{A_{h,k}^+}, j \in [1,r]} f_{w,j}^{\dots} T_w c_j \right) T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} c_{A^*} \Sigma_A \\ &= {}_{\text{SDP}} \text{HE} + \text{T}\overline{\text{E}}, \end{aligned}$$

where ${}_{\text{SDP}} \text{HE} = {}_{\text{SDP}} \text{HE}(x_\mu)$ ($\mu = \text{co}(A)$)

$$\text{T}\overline{\text{E}} = \sum_{\substack{1 \leq k \leq n \\ a_{h+1,k} > 0}} \sum_{w < d_{A_{h,k}^+}, j \in [1,r]} f_{w,j}^{\dots} q^{\vec{r}_h^k} x_{\lambda_{(h)}^+} T_w c_j T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} c_{A^*} \Sigma_A. \quad (6.3.5)$$

Since $T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} c_{A^*} \Sigma_A \in \mathcal{H}_{\mu,R}^c$ and every $w < d_{A_{h,k}^+}$ has the form $w = xdy$ with $x \in \mathfrak{S}_{\lambda_{(h)}^+}$, $d \in \mathcal{D}_{\lambda_{(h)}^+, \mu}$ and $y \in \mathfrak{S}_\mu$, it follows that $\text{T}\overline{\text{E}}$ is a linear combination of the basis elements $x_{\lambda_{(h)}^+} T_d c^\alpha T_v$ for $x_{\lambda_{(h)}^+} \mathcal{H}_{r,R}^c$ (see [18, Cor. 3.8] or Lemma 6.1) where $d < d_{A_{h,k}^+}$, for some k .

On the other hand, since both z and ${}_{\text{SDP}} \text{HE}$ are in $x_{\lambda_{(h)}^+} \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$, it follows that $\text{T}\overline{\text{E}} \in x_{\lambda_{(h)}^+} \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$. Hence, by Proposition 2.3(1),

$$\text{T}\overline{\text{E}} = \sum_{M^* \in M_n(\mathbb{N}|\mathbb{N}_2)_{\lambda_{(h)}^+, \mu}} f_{M^*}^{E_h^*, A^*} T_{M^*}. \quad (6.3.6)$$

Writing this sum as a linear combination of the same basis for $x_{\lambda_{(h)}^+} \mathcal{H}_{r,R}^c$ and equating coefficients show that every $d_{[M^*]} < d_{A_{h,k}^+}$, for some k . Now assertion (2) follows from the fact $d_A < d_B \implies A \prec B$; see [8, Lem. 13.20] or (6.0.2). \square

We now deal with the most complicated case.

Theorem 6.4. *Let $h \in [1, n-1]$ and $A^* = (A^{\bar{0}} | A^{\bar{1}}) = (a_{i,j}^{\bar{0}} | a_{i,j}^{\bar{1}}) \in M_n(\mathbb{N}|\mathbb{N}_2)_r$ with $A = A^{\bar{0}} + A^{\bar{1}}$, $\lambda = \text{ro}(A)$, and $\vec{r}_{h+1}^k = \vec{r}_{h+1}^k(A)$. Let $F_h^* = (\lambda - E_{h,h} | E_{h+1,h})$.*

(1) If A satisfies the SDP condition at (h, k) , for all $k \in [1, n]$ with $a_{h,k} > 0$, then we have

$$\begin{aligned} \phi_{F_h^*} \phi_{A^*} &= \sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} \left\{ (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + a_{h,k}^{\bar{1}}} \phi_{(A^{\bar{0}} - E_{h,k} | A^{\bar{1}} + E_{h+1,k})} \right. \\ &\quad + (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + a_{h,k}^{\bar{1}} + 1} q^{-1} \llbracket a_{h+1,k} + 1 \rrbracket_{q^2, q} \phi_{(A^{\bar{0}} - E_{h,k} + 2E_{h+1,k} | A^{\bar{1}} - E_{h+1,k})} \\ &\quad \left. + (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + 1} q^{a_{h,k} - 1} \llbracket a_{h+1,k}^{\bar{0}} + 1 \rrbracket_q \phi_{(A^{\bar{0}} + E_{h+1,k} | A^{\bar{1}} - E_{h,k})} \right\} =: {}_{\text{SDP}} \text{HF}. \end{aligned}$$

(2) In general, let

$$\begin{aligned} \text{HHF} &= \sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} \sum_{l=1}^{k-1} (-1)^{\tilde{a}_{h,l}^{\bar{1}}} \cdot q^{\vec{r}_{h+1}^l - \vec{r}_{h+1}^{k-1}} \left\{ \llbracket a_{h+1,k}^{\bar{0}} + 1 \rrbracket_q \cdot \right. \\ &\quad \left(\phi_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} - E_{h+1,l} | A^{\bar{1}} + E_{h+1,l})} - \llbracket a_{h+1,l} \rrbracket_{q^2} \phi_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} + E_{h+1,l} | A^{\bar{1}} - E_{h+1,l})} \right) \\ &\quad + q^{a_{h,k} - 1} \left(\phi_{(A^{\bar{0}} - E_{h+1,l} | A^{\bar{1}} - E_{h,k} + E_{h+1,k} + E_{h+1,l})} - \llbracket a_{h+1,l} \rrbracket_{q^2} \phi_{(A^{\bar{0}} + E_{h+1,l} | A^{\bar{1}} - E_{h,k} + E_{h+1,k} - E_{h+1,l})} \right) \\ &\quad \left. + q^{a_{h,k} - 2} \llbracket a_{h+1,k} + 1 \rrbracket_{q^2, q} \left(\phi_{(A^{\bar{0}} + 2E_{h+1,k} - E_{h+1,l} | A^{\bar{1}} - E_{h,k} - E_{h+1,k} + E_{h+1,l})} \right. \right. \\ &\quad \left. \left. - \llbracket a_{h+1,l} \rrbracket_{q^2} \phi_{(A^{\bar{0}} + 2E_{h+1,k} + E_{h+1,l} | A^{\bar{1}} - E_{h,k} - E_{h+1,k} - E_{h+1,l})} \right) \right\}. \end{aligned}$$

Then

$$\phi_{F_h^*} \phi_{A^*} = {}_{\text{SDP}} \text{HF} + (q-1) \text{HHF} + \sum_{\substack{B^* \in \mathcal{M}_n(\mathbb{N} | \mathbb{N}_2)_r \\ \exists k, [B^*] \prec A_{h,k}^-}} f_{B^*}^{F_h^*, A^*} \phi_{B^*}.$$

Proof. We want to compute $z := \phi_{X^*} \phi_A(x_\mu)$ with $X^* = F_h^* = (\lambda - E_{h,h} | E_{h+1,h})$ and $\mu = \text{co}(A)$. Since $X = [F_h^*]$ is the same matrix X as given in (5.3.5), and $c_{X^*} = c_{\tilde{\lambda}_h}$, it follows from (5.3.5) that $\sum_{\sigma \in \mathcal{D}_{\nu_X} \cap \mathfrak{S}_\lambda} T_\sigma = T_{(\tilde{\lambda}_{h-1}, \tilde{\lambda}_{h-1} + 1)}^\triangleright$, and thus,

$$\begin{aligned} z &= x_{\lambda_{(h)}^-} T_{d_X} c_{X^*} T_{(\tilde{\lambda}_{h-1}, \tilde{\lambda}_{h-1} + 1)}^\triangleright \cdot (T_{d_A} c_{A^*} \Sigma_A) \\ &= \sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} x_{\lambda_{(h)}^-} c_{\tilde{\lambda}_h} \left(\sum_{j=\vec{r}_h^{k-1} - 1}^{\vec{r}_h^{k-1} - 1} T_{\tilde{\lambda}_{h-1}} \cdots T_{\tilde{\lambda}_{h-j}} T_{d_A} \right) c_{A^*} \Sigma_A \\ &= \sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} x_{\lambda_{(h)}^-} c_{\tilde{\lambda}_h} (T_{\tilde{\lambda}_h} T_{\tilde{\lambda}_{h+1}} \cdots T_{\tilde{\lambda}_h + \overleftarrow{r}_{h+1}^k - 1} T_{d_{A_{h,k}^-}} T_{(\tilde{a}-1, \tilde{a}-a+1)}^\triangleright) c_{A^*} \Sigma_A \quad (a = a_{h,k}), \end{aligned} \tag{6.4.1}$$

by Lemma 3.4(2). For each $k \in [1, n]$ with $a_{h,k} > 0$, let

$$z_k = x_{\lambda_{(h)}^-} c_{\tilde{\lambda}_h} T_{\tilde{\lambda}_h} T_{\tilde{\lambda}_{h+1}} \cdots T_{\tilde{\lambda}_h + \overleftarrow{r}_{h+1}^k - 1} T_{d_{A_{h,k}^-}} T_{(\tilde{a}-1, \tilde{a}-a+1)}^\triangleright c_{A^*} \Sigma_A. \tag{6.4.2}$$

The hypothesis that A satisfies the SDP condition at (h, k) implies by Theorem 4.6 that $A_{\tilde{1}}^{h,k} = 0$. Thus, the lower left corner matrix of $A_{h,k}^- = (a_{i,j}^-)$ at (h, k) (and hence, at $(h+1, k)$) is zero. This implies, on the one hand, $a_{h+1,j} = 0$ for $j < k$, forcing $\overleftarrow{r}_{h+1}^k = 0$ and so, $T_{\tilde{\lambda}_h} \cdots T_{\tilde{\lambda}_h + \overleftarrow{r}_{h+1}^k - 1} = 1$. On the other hand, since $a_{h+1,k}^- = a_{h+1,k} + 1 > 0$, it follows that $A_{h,k}^-$ satisfies the SDP condition at $(h+1, k)$, by Theorem 4.6 again. Thus, as $\tilde{\lambda}_h = \hat{a}_{h,n} = \hat{a}_{h+1,k-1} = \hat{a}_{h+1,k-1}^- + 1$, we have

$$\begin{aligned} c_{\tilde{\lambda}_h} T_{d_{A_{h,k}^-}} &= c_{\hat{a}_{h+1,k-1}^- + 1} T_{d_{A_{h,k}^-}} = T_{d_{A_{h,k}^-}} c_{\hat{a}_{h,k}^- + 1} = T_{d_{A_{h,k}^-}} c_{\hat{a}_{h,k}}, \quad \text{and} \\ z_k &= x_{\lambda_{(h)}^-} c_{\tilde{\lambda}_h} T_{d_{A_{h,k}^-}} T_{(\tilde{a}-1, \tilde{a}-a+1)}^\triangleright c_{A^*} \Sigma_A = x_{\lambda_{(h)}^-} T_{d_{A_{h,k}^-}} c_{\hat{a}_{h,k}} T_{(\tilde{a}-1, \tilde{a}-a+1)}^\triangleright c_{A^*} \Sigma_A. \end{aligned} \tag{6.4.3}$$

Hence, by Lemma A.1,

$$\begin{aligned}
z &= \sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} z_k = \sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} \left\{ (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + a_{h,k}^{\bar{1}}} T_{(A^{\bar{0}} - E_{h,k} | A^{\bar{1}} + E_{h+1,k})} \right. \\
&\quad + (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + a_{h,k}^{\bar{1}} + 1} q^{-1} \llbracket a_{h+1,k} + 1 \rrbracket_{q^2, q} T_{(A^{\bar{0}} - E_{h,k} + 2E_{h+1,k} | A^{\bar{1}} - E_{h+1,k})} \\
&\quad \left. + (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + 1} q^{a_{h,k} - 1} \llbracket a_{h+1,k}^{\bar{0}} + 1 \rrbracket_q T_{(A^{\bar{0}} + E_{h+1,k} | A^{\bar{1}} - E_{h,k})} \right\} =_{\text{SDP}} \text{HF}(x_\mu),
\end{aligned} \tag{6.4.4}$$

proving (1).

We now prove the general case (2).

By applying Lemma 3.7 (1) to $x_{\lambda_{(h)}^-} c_{\tilde{\lambda}_h} T_{\tilde{\lambda}_h} T_{\tilde{\lambda}_{h+1}} \cdots T_{\tilde{\lambda}_h + \tilde{\mathfrak{F}}_{h+1}^k - 1}$, (6.4.2) becomes

$$\begin{aligned}
z_k &= \left(q^{\tilde{\mathfrak{F}}_{h+1}^k} x_{\lambda_{(h)}^-} T_{\tilde{\lambda}_h}^{-1} T_{\tilde{\lambda}_{h+1}}^{-1} T_{\tilde{\lambda}_{h+2}}^{-1} \cdots T_{\tilde{\lambda}_h + \tilde{\mathfrak{F}}_{h+1}^k - 1}^{-1} c_{\tilde{\lambda}_h + \tilde{\mathfrak{F}}_{h+1}^k} \right. \\
&\quad + (q-1) q^{\tilde{\mathfrak{F}}_{h+1}^k - 1} x_{\lambda_{(h)}^-} \left(c_{\tilde{\lambda}_h} + T_{\tilde{\lambda}_h}^{-1} c_{\tilde{\lambda}_{h+1}} + T_{\tilde{\lambda}_h}^{-1} T_{\tilde{\lambda}_{h+1}}^{-1} c_{\tilde{\lambda}_{h+2}} \right. \\
&\quad \left. + \cdots + T_{\tilde{\lambda}_h}^{-1} T_{\tilde{\lambda}_{h+1}}^{-1} \cdots T_{\tilde{\lambda}_h + \tilde{\mathfrak{F}}_{h+1}^k - 2}^{-1} c_{\tilde{\lambda}_h + \tilde{\mathfrak{F}}_{h+1}^k - 1} \right) \cdot T_{d_{A_{h,k}^-}} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A \\
&= \left(x_{\lambda_{(h)}^-} c_{\tilde{\lambda}_h + \tilde{\mathfrak{F}}_{h+1}^k} + (q-1) x_{\lambda_{(h)}^-} \left(q^{\tilde{\mathfrak{F}}_{h+1}^k - 1} c_{\tilde{\lambda}_h} + q^{\tilde{\mathfrak{F}}_{h+1}^k - 2} c_{\tilde{\lambda}_{h+1}} + q^{\tilde{\mathfrak{F}}_{h+1}^k - 3} c_{\tilde{\lambda}_{h+2}} \right. \right. \\
&\quad \left. \left. + \cdots + c_{\tilde{\lambda}_h + \tilde{\mathfrak{F}}_{h+1}^k - 1} \right) \right) \cdot T_{d_{A_{h,k}^-}} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A \\
&= \left(x_{\lambda_{(h)}^-} c_{\tilde{a}_{h+1,k-1}} + (q-1) x_{\lambda_{(h)}^-} c_{q, \tilde{\lambda}_h, \tilde{a}_{h+1,k-1} - 1} \right) \cdot T_{d_{A_{h,k}^-}} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A = Z_k + N_k,
\end{aligned} \tag{6.4.5}$$

where

$$\begin{aligned}
Z_k &= x_{\lambda_{(h)}^-} c_{\tilde{a}_{h+1,k-1}} T_{d_{A_{h,k}^-}} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A \\
N_k &= (q-1) x_{\lambda_{(h)}^-} c_{q, \tilde{\lambda}_h, \tilde{a}_{h+1,k-1} - 1} T_{d_{A_{h,k}^-}} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A.
\end{aligned} \tag{6.4.6}$$

Consider Z_k first. Since $\tilde{a}_{h+1,k-1} = \tilde{a}_{h+1,k-1}^- + 1$ and $\tilde{a}_{h,k} = \tilde{a}_{h,k}^- + 1$, by Lemma 4.5(2),

$$c_{\tilde{a}_{h+1,k-1}} T_{d_{A_{h,k}^-}} = T_{d_{A_{h,k}^-}} c_{\tilde{a}_{h,k}} + \sum_{w < d_{A_{h,k}^-}, j \in [1, r]} f_{w,j}^{\cdots} T_w c_j. \tag{6.4.7}$$

Thus, $Z_k = (\diamond)_k + (\clubsuit)_k$, where

$$\begin{aligned}
(\diamond)_k &:= x_{\lambda_{(h)}^-} T_{d_{A_{h,k}^-}} c_{\tilde{a}_{h,k}} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A, \\
(\clubsuit)_k &:= x_{\lambda_{(h)}^-} \left(\sum_{w < d_{A_{h,k}^-}, j \in [1, r]} f_{w,j}^{\cdots} T_w c_j \right) T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A.
\end{aligned}$$

Note that, according to (6.4.3), $(\diamond)_k$ is the k th summand of the head term $_{\text{SDP}} \text{HF}(x_{\text{co}(A)})$ in (1).

Suppose $\tilde{\mathfrak{F}}_{h+1}^k = a_{h+1,l_1} + a_{h+1,l_2} + \cdots + a_{h+1,l_s}$, where $a_{h+1,l_i} > 0$ and $1 \leq l_1 < l_2 < \cdots < l_s < k$. Then

$$\begin{aligned}
c_{q, \tilde{\lambda}_h, \tilde{a}_{h+1,k-1} - 1} &= q^{\tilde{\mathfrak{F}}_{h+1}^k - 1} c_{\tilde{\lambda}_h} + q^{\tilde{\mathfrak{F}}_{h+1}^k - 2} c_{\tilde{\lambda}_{h+1}} + q^{\tilde{\mathfrak{F}}_{h+1}^k - 3} c_{\tilde{\lambda}_{h+2}} + \cdots + c_{\tilde{\lambda}_h + \tilde{\mathfrak{F}}_{h+1}^k - 1} \\
&= \sum_{i=1}^s q^{\tilde{\mathfrak{F}}_{h+1}^k - \tilde{\mathfrak{F}}_{h+1}^{l_i} + 1} c_{q, \tilde{a}_{h+1,l_i}^- - 1, \tilde{a}_{h+1,l_i}^-}.
\end{aligned}$$

Thus, by (4.5.1)

$$\begin{aligned}
N_k &= (q-1)x_{\lambda_{(h)}^-} \left(\sum_{i=1}^s q^{\overleftarrow{\mathbb{F}}_{h+1}^k - \overleftarrow{\mathbb{F}}_{h+1}^{l_i+1}} c_{q, \tilde{a}_{h+1, l_i-1}+1, \tilde{a}_{h+1, l_i}^-} \right) T_{d_{A_{h,k}^-}} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A \\
&= (q-1) \sum_{i=1}^s q^{\overleftarrow{\mathbb{F}}_{h+1}^k - \overleftarrow{\mathbb{F}}_{h+1}^{l_i+1}} x_{\lambda_{(h)}^-} \left(T_{d_{A_{h,k}^-}} c_{(q, \tilde{a}_{h, l_i-1}+1, \tilde{a}_{h+1, l_i}^-)} + \sum_{w < d_{A_{h,k}^-}, j \in [1, r]} g_{w,j}^{\dots} T_w c_j \right) T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A \\
&= (q-1)((\diamond\diamond)_k + (\clubsuit\clubsuit)_k),
\end{aligned}$$

where (noting $\tilde{a}_{h, l_i-1}^- = \tilde{a}_{h, l_i-1}$ and $\tilde{a}_{h, l_i}^- = \tilde{a}_{h, l_i}$)

$$\begin{aligned}
(\diamond\diamond)_k &:= \sum_{\substack{1 \leq l \leq k-1 \\ a_{h+1, l} > 0}} q^{\overleftarrow{\mathbb{F}}_{h+1}^k - \overleftarrow{\mathbb{F}}_{h+1}^{l+1}} x_{\lambda_{(h)}^-} T_{d_{A_{h,k}^-}} c_{(q, \tilde{a}_{h, l}+1, \tilde{a}_{h+1, l})} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A \\
(\clubsuit\clubsuit)_k &:= \sum_{\substack{1 \leq l \leq k-1 \\ a_{h+1, l} > 0}} \sum_{w < d_{A_{h,k}^-}, j \in [1, r]} q^{\overleftarrow{\mathbb{F}}_{h+1}^k - \overleftarrow{\mathbb{F}}_{h+1}^{l+1}} g_{w,j}^{\dots} x_{\lambda_{(h)}^-} T_w c_j T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A \\
&= \sum_{w < d_{A_{h,k}^-}, j \in [1, r]} f_{w,j}^{\dots} x_{\lambda_{(h)}^-} T_w c_j T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A.
\end{aligned}$$

Similar to the computations in (5.3.6) and (5.3.7), the element $(\diamond\diamond)_k$ has the form

$$\begin{aligned}
(\diamond\diamond)_k &= \sum_{\substack{1 \leq l \leq k-1 \\ a_{h+1, l} > 0}} q^{\overleftarrow{\mathbb{F}}_{h+1}^k - \overleftarrow{\mathbb{F}}_{h+1}^{l+1}} x_{\lambda_{(h)}^-} T_{d_{A_{h,k}^-}} c_{(q, \tilde{a}_{h, l}+1, \tilde{a}_{h+1, l})} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A \\
&= \sum_{\substack{1 \leq l \leq k-1 \\ a_{h+1, l} > 0}} q^{\overleftarrow{\mathbb{F}}_{h+1}^k - \overleftarrow{\mathbb{F}}_{h+1}^{l+1}} x_{\lambda_{(h)}^-} \setminus \nu^- T_{d_{A_{h,k}^-}} x_{\nu^-} c_{(q, \tilde{a}_{h, l}+1, \tilde{a}_{h+1, l})} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A \\
&= \sum_{\substack{1 \leq l \leq k-1 \\ a_{h+1, l} > 0}} q^{\overleftarrow{\mathbb{F}}_{h+1}^k - \overleftarrow{\mathbb{F}}_{h+1}^{l+1}} x_{\lambda_{(h)}^-} \setminus \nu^- T_{d_{A_{h,k}^-}} c'_{(q, \tilde{a}_{h, l}+1, \tilde{a}_{h+1, l})} x_{\nu^-} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A,
\end{aligned} \tag{6.4.8}$$

where $\nu^- = \nu_{A_{h,k}^-}$; see (5.2.2). By Corollary 5.4,

$$\begin{aligned}
x_{\nu^-} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A &= \llbracket a_{h+1, k}^{\bar{0}} + 1 \rrbracket_q x_{\nu^-} c_{(A^{\bar{0}} - E_{h, k} + E_{h+1, k} | A^{\bar{1}})} \Sigma_{A_{h, k}^-} \\
&\quad + q^{a_{h, k} - 1} x_{\nu^-} c_{(A^{\bar{0}} | A^{\bar{1}} - E_{h, k} + E_{h+1, k})} \Sigma_{A_{h, k}^-} \\
&\quad + q^{a_{h, k} - 2} \llbracket a_{h+1, k} + 1 \rrbracket_{q^2} x_{\nu^-} c_{(A^{\bar{0}} + 2E_{h+1, k} | A^{\bar{1}} - E_{h, k} - E_{h+1, k})} \Sigma_{A_{h, k}^-}.
\end{aligned} \tag{6.4.9}$$

Let B^* be one of the following three matrices occurring in (6.4.9)

$$(A^{\bar{0}} - E_{h, k} + E_{h+1, k} | A^{\bar{1}}), (A^{\bar{0}} | A^{\bar{1}} - E_{h, k} + E_{h+1, k}), (A^{\bar{0}} + 2E_{h+1, k} | A^{\bar{1}} - E_{h, k} - E_{h+1, k}),$$

and write $B^* = (B^{\bar{0}} | B^{\bar{1}}) = (b_{i,j}^{\bar{0}} | b_{i,j}^{\bar{1}})$. Then we have $[B^*] = A_{h, k}^-$, $\nu_{[B^*]} = \nu_{A_{h, k}^-} = \nu^-$, and $c_{B^*} = c_{\nu_{[B^*]}^-}$. Note that, for $1 \leq l < k$,

$$b_{h+1, l}^{\bar{0}} = a_{h+1, l}^{\bar{0}}, b_{h+1, l}^{\bar{1}} = a_{h+1, l}^{\bar{1}}, \tilde{b}_{h, l} = \tilde{a}_{h, l}, \tilde{b}_{h+1, l} = \tilde{a}_{h+1, l}.$$

Thus,

$$\begin{aligned}
&c'_{(q, \tilde{a}_{h, l}+1, \tilde{a}_{h+1, l})} x_{\nu^-} c_{B^*} = x_{\nu^-} c_{(q, \tilde{a}_{h, l}+1, \tilde{a}_{h+1, l})} c_{B^*} = x_{\nu_{[B^*]}^-} c_{(q, \tilde{b}_{h, l}+1, \tilde{b}_{h+1, l})} c_{B^*} \\
&= \begin{cases} (-1)^{\tilde{a}_{h, l}^{\bar{1}}} x_{\nu_{[B^*]}^-} c_{(B^{\bar{0}} - E_{h+1, l} | B^{\bar{1}} + E_{h+1, l})}, & \text{if } b_{h+1, l}^{\bar{1}} = a_{h+1, l}^{\bar{1}} = 0; \\ -(-1)^{\tilde{a}_{h, l}^{\bar{1}}} \llbracket a_{h+1, l} \rrbracket_{q^2} x_{\nu_{[B^*]}^-} c_{(B^{\bar{0}} + E_{h+1, l} | B^{\bar{1}} - E_{h+1, l})}, & \text{if } b_{h+1, l}^{\bar{1}} = a_{h+1, l}^{\bar{1}} = 1. \end{cases} \tag{6.4.10}
\end{aligned}$$

Substituting (6.4.9) and (6.4.10) into (6.4.8) yields

$$\begin{aligned}
(\diamond\diamond)_k &= \sum_{l=1}^{k-1} (-1)^{\tilde{a}_{h,l}^-} \cdot q^{\overleftarrow{\mathbb{F}}_{h+1}^k - \overleftarrow{\mathbb{F}}_{h+1}^{l+1}} \\
&\left\{ \llbracket a_{h+1,k}^{\bar{0}} + 1 \rrbracket_q (T_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} - E_{h+1,l} | A^{\bar{1}} + E_{h+1,l})} - \llbracket a_{h+1,l} \rrbracket_{q^2} T_{(A^{\bar{0}} - E_{h,k} + E_{h+1,k} + E_{h+1,l} | A^{\bar{1}} - E_{h+1,l})}) \right. \\
&+ q^{a_{h,k}^{-1}} (T_{(A^{\bar{0}} - E_{h+1,l} | A^{\bar{1}} - E_{h,k} + E_{h+1,k} + E_{h+1,l})} - \llbracket a_{h+1,l} \rrbracket_{q^2} T_{(A^{\bar{0}} + E_{h+1,l} | A^{\bar{1}} - E_{h,k} + E_{h+1,k} - E_{h+1,l})}) \\
&+ q^{a_{h,k}^{-2}} \llbracket a_{h+1,k} + 1 \rrbracket_{q^2, q} (T_{(A^{\bar{0}} + 2E_{h+1,k} - E_{h+1,l} | A^{\bar{1}} - E_{h,k} - E_{h+1,k} + E_{h+1,l})} \\
&\quad \left. - \llbracket a_{h+1,l} \rrbracket_{q^2} T_{(A^{\bar{0}} + 2E_{h+1,k} + E_{h+1,l} | A^{\bar{1}} - E_{h,k} - E_{h+1,k} - E_{h+1,l})}) \right\}, \tag{6.4.11}
\end{aligned}$$

which is the k th term of $\text{HH}\overline{\mathbb{F}}(x_{\text{co}(A)})$.

Finally, combining above gives

$$z = \sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} \{(\diamond)_k + (q-1)(\diamond\diamond)_k\} + \sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} \{(\clubsuit)_k + (q-1)(\clubsuit\clubsuit)_k\}.$$

Since $z \in x_{\lambda_{(h)}^-} \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$ and $(\diamond)_k$ and $(\diamond\diamond)_k$ are in $x_{\lambda_{(h)}^-} \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$, it follows that

$$\sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} \{(\clubsuit)_k + (q-1)(\clubsuit\clubsuit)_k\} \in x_{\lambda_h^-} \mathcal{H}_{r,R}^c \cap \mathcal{H}_{r,R}^c x_\mu$$

Since $T_{(\tilde{a}-1, \tilde{a}-a+1)}^\triangleright \in \mathcal{H}_{\mu,R}$, and so, $\sum_{\sigma \in \mathcal{D}_{\nu_A} \cap \mathfrak{S}_\mu} T_{(\tilde{a}-1, \tilde{a}-a+1)}^\triangleright c_{A^*} T_\sigma \in \mathcal{H}_{\mu,R}^c$, a similar argument at the end of the proofs of Theorems 6.2 and 6.3 shows that

$$\sum_{\substack{1 \leq k \leq n \\ a_{h,k} > 0}} \{(\clubsuit)_k + (q-1)(\clubsuit\clubsuit)_k\} = \sum_{\substack{B^* \in M_n(\mathbb{N}|\mathbb{N}_2)_r \\ \exists k, \lfloor B^* \rfloor \prec A_{h,k}^-}} f_{B^*}^{F_h^*, A^*} T_{B^*},$$

for some $f_{B^*}^{F_h^*, A^*} \in R$. This completes the proof of (2). \square

Remark 6.5. Let $\mathcal{Q}(n, r)_{\mathbb{C}(v)} = \mathcal{Q}_q(n, r; \mathbb{C}(v))$, where $q = v^2$ and let $\mathbf{U}(q_n)$ be the quantum queer supergroup over $\mathbb{C}(v)$ with generators $E_h, F_h, K_j, K_{\bar{n}}$ ($h \in [1, n-1], j \in [1, n]$). Then, by [26], there is a superalgebra homomorphism $\varphi : \mathbf{U}(q_n) \rightarrow \mathcal{Q}(n, r)_{\mathbb{C}(v)}$. Let $e_h = \varphi(E_h)$, $f_h = \varphi(F_h)$, $k_j = \varphi(K_j)$, and $k_{\bar{n}} = \varphi(K_{\bar{n}})$. Then, by [18, Th.9.5], $\mathcal{Q}(n, r) = \mathcal{Q}_q(n, r; \mathbb{Q}(v))$ is generated by $e_h \phi_{(\lambda|O)}$, $f_h \phi_{(\lambda|O)}$, $k_i \phi_{(\lambda|O)}$, and $k_{\bar{n}} \phi_{(\lambda|O)}$, for all $h \in [1, h-1]$, $\lambda \in \Lambda(n, r)$. We will see in [13] that these generators are exactly those ϕ_{X^*} , where X^* is one of the matrices

$$(\lambda|O), \quad (\lambda + E_{h,h+1} - E_{h+1,h+1}|O), \quad (\lambda - E_{h,h} + E_{h+1,h}|O), \quad (\lambda - E_{n,n}|E_{n,n}).$$

Hence, the multiplication formulas given in Theorem 5.3 and Theorem 6.2 give the matrix form of the regular representation of $\mathcal{Q}(n, r)$.

7. SOME SPECIAL CASES

In this section, we establish several special multiplication formulas related to certain generators. We single out them for the convenience when checking the defining relations in [13]. We see below that the tail parts in Theorems 6.3 and 6.4 are computable if A is simple enough.

Proposition 7.1. *For any given $\mu \in \Lambda(n, r-1)$, $h \in [1, n-1]$, the following formulas hold in $\mathcal{Q}_q(n, r; R)$.*

- (1) $\phi_{(\mu|E_{h,h+1})}\phi_{(\mu+E_{h+1,h}|O)} = \phi_{(\mu-E_{h+1,h+1}+E_{h+1,h}|E_{h,h+1})} + q^{\mu_{h+1}}\phi_{(\mu|E_{h,h})} - (q-1)\llbracket\mu_h+1\rrbracket_q\phi_{(\mu-E_{h+1,h+1}+E_{h,h}|E_{h+1,h+1})},$
- (2) $\phi_{(\mu|E_{h,h+1})}\phi_{(\mu|E_{h+1,h})} = -\llbracket\mu_h+1\rrbracket_q q^{\mu_{h+1}}\phi_{(\mu+E_{h,h}|O)} - \phi_{(\mu-E_{h+1,h+1}|E_{h,h+1}+E_{h+1,h})} + (q-1)\phi_{(\mu-E_{h+1,h+1}|E_{h,h}+E_{h+1,h+1})}$
- (3) $\phi_{(\mu|E_{h+1,h})}\phi_{(\mu+E_{h,h+1}|O)} = \phi_{(\mu-E_{h,h}+E_{h,h+1}|E_{h+1,h})} + \phi_{(\mu|E_{h+1,h+1})},$
- (4) $\phi_{(\mu|E_{h+1,h})}\phi_{(\mu|E_{h,h+1})} = \phi_{(\mu-E_{h,h}|E_{h,h+1}+E_{h+1,h})} - \llbracket\mu_{h+1}+1\rrbracket_q\phi_{(\mu+E_{h+1,h+1}|O)}.$

Proof. Parts (3) and (4) clearly follow from Theorem 6.4(1) as the matrix $\text{diag}(\mu) + E_{h,h+1}$ satisfies SDP condition on the h th row by Theorem 4.6. We now prove parts (1) and (2).

Let $A^* = (a_{i,j}^{\bar{0}}|a_{i,j}^{\bar{1}}) = (\mu + E_{h+1,h}|O)$ or $(\mu|E_{h+1,h})$ which have the same base matrix

$$A = [A^*] = \text{diag}(\mu) + E_{h+1,h} = (a_{i,j}).$$

Thus, $\lambda := \text{ro}(A) = \mu + \epsilon_{h+1}$. Hence, $X^* = (\mu|E_{h,h+1}) = (\lambda - E_{h+1,h+1}|E_{h,h+1}) = E_h^*$ in both cases. This means that cases (1) and (2) are special cases of Theorem 6.3(2) and $\text{ro}(X) = \text{ro}([X^*]) = \lambda_{(h)}^+$, and $c_{X^*} = c_{\lambda_{h+1}}^-$. Since $A = \text{diag}(\lambda) + E_{h+1,h} - E_{h+1,h+1}$, it follows that

$$\begin{aligned} \nu_A &= (\lambda_1, 0, \dots, 0, \lambda_h, 0, \dots, 0, 1, \lambda_{h+1} - 1, 0, \dots), \\ \text{co}(A) &= (\lambda_1, \dots, \lambda_h + 1, \lambda_{h+1} - 1, \dots, \lambda_n), \\ \mathcal{D}_{\nu_A} \cap \mathfrak{S}_{\text{co}(A)} &= \{1, s_{\lambda_h}^-, s_{\lambda_h}^- s_{\lambda_{h-1}}^- \cdots, s_{\lambda_h}^- s_{\lambda_{h-1}}^- \cdots s_{\lambda_{h-1}+1}^-\}, \\ \Sigma_A(\Sigma_{v \in \mathcal{D}_{\nu_A} \cap \mathfrak{S}_{\text{co}(A)}} T_v) &= T_{(\lambda_h, \tilde{\lambda}_{h-1}+1)}^{\triangleright}. \end{aligned} \tag{7.1.1}$$

By Example 4.3 and (2.2.12), we also have

$$d_A = 1, \quad c_{A^*} = \begin{cases} 1, & \text{if } A^* = (\mu + E_{h+1,h}|O); \\ c_{\lambda_{h+1}}^-, & \text{if } A^* = (\mu|E_{h+1,h}). \end{cases} \tag{7.1.2}$$

Since, by Theorem 4.6, $A_{h,h+1}^+$ does not satisfy the SDP condition at $(h, h+1)$ when $\mu_{h+1} > 0$ (see footnote 3), we must compute the tail term explicitly in Theorem 6.3(2).

By (6.3.2), we have, for $\overleftarrow{\mathbf{r}}_h^k = \overleftarrow{\mathbf{r}}_h^k(A)$,

$$z = \sum_{\substack{1 \leq k \leq n \\ a_{h+1,k} > 0}} z_k, \quad \text{where } z_k = \sum_{j=\overleftarrow{\mathbf{r}}_{h+1}^k}^{\overleftarrow{\mathbf{r}}_{h+1}^{k+1}-1} x_{\lambda_{(h)}^+} c_{\tilde{\lambda}_{h+1}}^- T_{\tilde{\lambda}_{h+1}}^- T_{\tilde{\lambda}_{h+2}}^- \cdots T_{\tilde{\lambda}_{h+j}}^- T_{d_A} c_{A^*} \Sigma_A. \tag{7.1.3}$$

Further, since $a_{h+1,k} = 0$, for all $k \neq h, h+1$, it follows that $z = z_h + z_{h+1}$, where

$$z_h = x_{\lambda_{(h)}^+} c_{\tilde{\lambda}_{h+1}}^- c_{A^*} \Sigma_A = x_{\lambda_{(h)}^+} c_{\tilde{\lambda}_{h+1}}^- c_{A^*} T_{(\lambda_h, \tilde{\lambda}_h - (\lambda_h - 1))}^{\triangleright}, \quad \text{as } d_A = 1, a_{h+1,h} = 1,$$

and, as $\overleftarrow{\mathbf{r}}_{h+1}^{h+1} = 1$, $\overleftarrow{\mathbf{r}}_{h+1}^{h+2} = \overleftarrow{\mathbf{r}}_{h+1}^{h+1} + a_{h+1,h+1} = \lambda_{h+1}$, and $d_A = 1$,

$$\begin{aligned} z_{h+1} &= x_{\lambda_{(h)}^+} c_{\tilde{\lambda}_{h+1}}^- \left(\sum_{j=\overleftarrow{\mathbf{r}}_{h+1}^{h+1}}^{\overleftarrow{\mathbf{r}}_{h+1}^{h+2}-1} T_{\tilde{\lambda}_{h+1}}^- T_{\tilde{\lambda}_{h+2}}^- \cdots T_{\tilde{\lambda}_{h+j}}^- \right) T_{d_A} c_{A^*} \Sigma_A \\ &= x_{\lambda_{(h)}^+} c_{\tilde{\lambda}_{h+1}}^- T_{\tilde{\lambda}_{h+1}}^- T_{(\tilde{\lambda}_h+2, \tilde{\lambda}_h+\lambda_{h+1}-1)}^{\triangleleft} c_{A^*} \Sigma_A. \end{aligned}$$

Now, depending on the choice of c_{A^*} in (7.1.2), Lemma 3.7(3) together with (2.2.12) implies

$$z_h = \begin{cases} (1) x_{\lambda_{(h)}^+} c_{q, \tilde{\lambda}_{h-1}+1, \tilde{\lambda}_{h+1}} = T_{(\lambda - E_{h+1,h+1}|E_{h,h})}, & \text{if } A^* = (\mu + E_{h+1,h}|O), \\ (2) -x_{\lambda_{(h)}^+} T_{(\tilde{\lambda}_h, \tilde{\lambda}_h - (\lambda_h - 1))}^{\triangleright} = -\llbracket\lambda_h+1\rrbracket_q T_{(\lambda - E_{h+1,h+1} + E_{h,h}|O)}, & \text{if } A^* = (\mu|E_{h+1,h}). \end{cases} \tag{7.1.4}$$

It remains to compute z_{h+1} . Note that $c_{\tilde{\lambda}_{h+1}}T_{\tilde{\lambda}_{h+1}} = T_{\tilde{\lambda}_{h+1}}c_{\tilde{\lambda}_{h+2}} + (q-1)(c_{\tilde{\lambda}_{h+1}} - c_{\tilde{\lambda}_{h+2}})$, by (2.0.3).

$$\begin{aligned}
z_{h+1} &= x_{\lambda_{(h)}^+} (T_{\tilde{\lambda}_{h+1}}c_{\tilde{\lambda}_{h+2}} + (q-1)(c_{\tilde{\lambda}_{h+1}} - c_{\tilde{\lambda}_{h+2}}))T_{(\tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1})}^{\triangleleft} c_{A^*} \Sigma_A \\
&= x_{\lambda_{(h)}^+} T_{\tilde{\lambda}_{h+1}}c_{\tilde{\lambda}_{h+2}}T_{(\tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1})}^{\triangleleft} c_{A^*} \Sigma_A \\
&\quad + (q-1)x_{\lambda_{(h)}^+} T_{(\tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1})}^{\triangleleft} c_{\tilde{\lambda}_{h+1}} c_{A^*} \Sigma_A \\
&\quad - (q-1)x_{\lambda_{(h)}^+} c_{\tilde{\lambda}_{h+2}} T_{(\tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1})}^{\triangleleft} c_{A^*} \Sigma_A.
\end{aligned} \tag{7.1.5}$$

If $A^* = (\mu + E_{h+1, h} | O)$, then $c_{A^*} = 1$ and

$$T_{(\tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1})}^{\triangleleft} \Sigma_A = T_{(\tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1})}^{\triangleleft} T_{(\tilde{\lambda}_h, \tilde{\lambda}_h - (\lambda_h - 1))}^{\triangleright} = \sum_{\sigma \in \mathcal{D}_{\nu_C} \cap \mathfrak{S}_{\text{co}(C)}} T_{\sigma} = \Sigma_C, \tag{7.1.6}$$

where $C = \lambda - 2E_{h+1, h+1} + E_{h, h+1} + E_{h+1, h}$ with $\nu_C = (\dots, \lambda_h, 1, 0, \dots, 0, 1, \lambda_{h+1} - 2, \dots)$. (Compare ν_C with $\text{co}(A)$ in (7.1.1).) Hence, by Lemmas 3.6 and 3.7(2)&(3), we obtain

$$\begin{aligned}
z_{h+1} &= x_{\lambda_{(h)}^+} T_{\tilde{\lambda}_{h+1}}c_{\tilde{\lambda}_{h+2}} \Sigma_C + (q-1) \llbracket \lambda_{h+1} - 1 \rrbracket_q x_{\lambda_{(h)}^+} c_{\tilde{\lambda}_{h+1}} T_{(\tilde{\lambda}_h, \tilde{\lambda}_h - (\lambda_h - 1))}^{\triangleright} \\
&\quad - (q-1)x_{\lambda_{(h)}^+} c_{q, \tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1}} T_{(\tilde{\lambda}_h, \tilde{\lambda}_h - (\lambda_h - 1))}^{\triangleright} \\
&= x_{\lambda_{(h)}^+} T_{\tilde{\lambda}_{h+1}}c_{\tilde{\lambda}_{h+2}} \Sigma_C + (q-1) \llbracket \lambda_{h+1} - 1 \rrbracket_q x_{\lambda_{(h)}^+} c_{q, \tilde{\lambda}_{h-1} + 1, \tilde{\lambda}_{h+1}} \\
&\quad - (q-1) \llbracket \lambda_h + 1 \rrbracket_q x_{\lambda_{(h)}^+} c_{q, \tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1}} \\
&= T_{(\lambda - 2E_{h+1, h+1} + E_{h+1, h} | E_{h, h+1})} + (q-1) \llbracket \lambda_{h+1} - 1 \rrbracket_q T_{(\lambda - E_{h+1, h+1} | E_{h, h})} \\
&\quad - (q-1) \llbracket \lambda_h + 1 \rrbracket_q T_{(\lambda - 2E_{h+1, h+1} + E_{h, h} | E_{h+1, h+1})}.
\end{aligned}$$

This together with (7.1.4)(1), after replacing $\lambda - E_{h+1, h+1}$ by μ , proves (1).

If $A = (\mu | E_{h+1, h})$, then $c_{A^*} = c_{\tilde{\lambda}_{h+1}}$ and, hence, $T_{(\tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1})}^{\triangleleft} c_{A^*} = c_{A^*} T_{(\tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1})}^{\triangleleft}$. Then, with a similar argument, (7.1.5) becomes

$$\begin{aligned}
z_{h+1} &= -x_{\lambda_{(h)}^+} T_{\tilde{\lambda}_{h+1}}c_{\tilde{\lambda}_{h+1}}c_{\tilde{\lambda}_{h+2}}T_{(\tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1})}^{\triangleleft} \Sigma_A - (q-1)x_{\lambda_{(h)}^+} T_{(\tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1})}^{\triangleleft} T_{(\tilde{\lambda}_h, \tilde{\lambda}_h - (\lambda_h - 1))}^{\triangleright} \\
&\quad - (q-1)x_{\lambda_{(h)}^+} c_{\tilde{\lambda}_{h+2}} T_{(\tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1})}^{\triangleleft} c_{\tilde{\lambda}_{h+1}} T_{(\tilde{\lambda}_h, \tilde{\lambda}_h - (\lambda_h - 1))}^{\triangleright} \\
&= -x_{\lambda_{(h)}^+} T_{\tilde{\lambda}_{h+1}}c_{\tilde{\lambda}_{h+1}}c_{\tilde{\lambda}_{h+2}} \Sigma_C - (q-1)x_{\lambda_{(h)}^+} \llbracket \lambda_{h+1} - 1 \rrbracket_q \llbracket \lambda_h + 1 \rrbracket_q \\
&\quad + (q-1)x_{\lambda_{(h)}^+} c_{q, \tilde{\lambda}_{h-1} + 1, \tilde{\lambda}_{h+1}} c_{q, \tilde{\lambda}_{h+2}, \tilde{\lambda}_{h+1}} \\
&= -T_{(\lambda - 2E_{h+1, h+1} | E_{h, h+1} + E_{h+1, h})} - (q-1) \llbracket \lambda_{h+1} - 1 \rrbracket_q \llbracket \lambda_h + 1 \rrbracket_q T_{(\lambda - E_{h+1, h+1} | E_{h, h} | O)} \\
&\quad + (q-1) T_{(\lambda - 2E_{h+1, h+1} | E_{h, h} + E_{h+1, h+1})}.
\end{aligned}$$

This together with (7.1.4)(2), after replacing $\lambda - E_{h+1, h+1}$ by μ , proves (2). \square

The following relation gives an example of the fact that a product of two odd elements in the superalgebra $\mathcal{Q}_q(n, r; R)$ is an even element.

Lemma 7.2. *Let $h \in [1, n-1]$ and $\lambda \in \Lambda(n, r-1)$. Then the following relation holds in $\mathcal{Q}_q(n, r; R)$:*

$$\begin{aligned}
&\phi(\lambda | E_{h, h+1}) \phi(\lambda | E_{h+1, h+2}) + q \phi(\lambda + \alpha_h | E_{h+1, h+2}) \phi(\lambda - \alpha_{h+1} | E_{h, h+1}) = \\
&- \phi(\lambda + E_{h, h+1} | O) \phi(\lambda + E_{h+1, h+2} | O) + q \phi(\lambda + \alpha_h + E_{h+1, h+2} | O) \phi(\lambda - \alpha_{h+1} + E_{h, h+1} | O).
\end{aligned} \tag{7.2.1}$$

Proof. Observe that each term in (7.2.1) is a product of the form $\phi_{A^*} \phi_{B^*}$. For all choices of B^* , the base matrix B have the same $\text{co}(B) = \lambda + \epsilon_{h+2}$. We prove the result by applying both sides to $x_{\lambda + \epsilon_{h+2}}$ and check the identity in $\mathcal{H}_{r, R}^c$.

Let $A_1^* = (\lambda | E_{h, h+1})$ and $B_1^* = (\lambda | E_{h+1, h+2})$ with base A and B , respectively. Then

$$\begin{aligned}
\mathcal{Y}_1 &= \phi(\lambda | E_{h, h+1}) (\phi(\lambda | E_{h+1, h+2}) (x_{\lambda + \epsilon_{h+2}})) = \phi(\lambda | E_{h, h+1}) (T_{(\lambda | E_{h+1, h+2})}) \\
&= \phi(\lambda | E_{h, h+1}) (x_{\text{ro}(B)} T_{d_B} c_{B_1^*} \Sigma_B) = x_{\text{ro}(A)} T_{d_A} c_{A_1^*} \Sigma_A \cdot T_{d_B} c_{B_1^*} \Sigma_B.
\end{aligned}$$

By Example 4.3 and various definitions, it is easy to verify $d_A = d_B = 1$, $c_{A_1^*} = c_{\tilde{\lambda}_{h+1}}$, $c_{B_1^*} = c_{\tilde{\lambda}_{h+1+1}}$, $\text{ro}(A) = \lambda + \epsilon_h$, and

$$\Sigma_A = T_{(\tilde{\lambda}_{h+1}, \tilde{\lambda}_h + \lambda_{h+1})}^{\triangleleft}, \quad \Sigma_B = T_{(\tilde{\lambda}_{h+1+1}, \tilde{\lambda}_{h+1} + \lambda_{h+2})}^{\triangleleft}.$$

Hence,

$$\mathcal{Y}_1 = x_{\lambda + \epsilon_h} c_{\tilde{\lambda}_{h+1}} T_{(\tilde{\lambda}_{h+1}, \tilde{\lambda}_h + \lambda_{h+1})}^{\triangleleft} c_{\tilde{\lambda}_{h+1+1}} T_{(\tilde{\lambda}_{h+1+1}, \tilde{\lambda}_{h+1} + \lambda_{h+2})}^{\triangleleft}.$$

Similarly, set $M_1^* = (\lambda + \alpha_h | E_{h+1, h+2})$, $N_1^* = (\lambda - \alpha_{h+1} | E_{h, h+1})$, having base matrices, M , N , respectively, then we have

$$\Sigma_M = T_{(\tilde{\lambda}_{h+1+1}, \tilde{\lambda}_{h+1} + \lambda_{h+2})}^{\triangleleft}, \quad \Sigma_N = T_{(\tilde{\lambda}_h + 1, \tilde{\lambda}_h + \lambda_{h+1} - 1)}^{\triangleleft},$$

and direct calculation shows

$$\begin{aligned} \mathcal{Y}_2 &= \phi_{(\lambda + \epsilon_h - \epsilon_{h+1} | E_{h+1, h+2})} \cdot \phi_{(\lambda + \epsilon_{h+2} - \epsilon_{h+1} | E_{h, h+1})}(x_{\lambda + \epsilon_{h+2}}) \\ &= x_{\lambda + \epsilon_h} c_{\tilde{\lambda}_{h+1+1}} \Sigma_M \cdot c_{\tilde{\lambda}_h + 1} \Sigma_N \\ &= x_{\lambda + \epsilon_h} c_{\tilde{\lambda}_{h+1+1}} T_{(\tilde{\lambda}_{h+1+1}, \tilde{\lambda}_{h+1} + \lambda_{h+2})}^{\triangleleft} c_{\tilde{\lambda}_h + 1} T_{(\tilde{\lambda}_h + 1, \tilde{\lambda}_h + \lambda_{h+1} - 1)}^{\triangleleft}. \end{aligned}$$

On the other hand, let $A_2^* = (\lambda + E_{h, h+1} | O)$, $B_2^* = (\lambda + E_{h+1, h+2} | O)$ which have the same base matrices A and B as A_1^* and B_1^* . Using the proof of Proposition 5.3(2), a straightforward calculation implies

$$\begin{aligned} \mathcal{Y}_3 &= \phi_{(\lambda + E_{h, h+1} | O)}(\phi_{(\lambda + E_{h+1, h+2} | O)}(x_{\lambda + \epsilon_{h+2}})) = \phi_{(\lambda + E_{h, h+1} | O)}(T_{(\lambda + E_{h+1, h+2} | O)}) \\ &= \phi_{(\lambda + E_{h, h+1} | O)}(x_{\text{ro}(B)} T_{d_B} c_{B_2^*} \Sigma_B) = x_{\text{ro}(A)} T_{d_A} c_{A_2^*} \Sigma_A \cdot T_{d_B} c_{B_2^*} \Sigma_B \\ &= x_{\lambda + \epsilon_h} T_{(\tilde{\lambda}_h + 1, \tilde{\lambda}_h + \lambda_{h+1})}^{\triangleleft} T_{(\tilde{\lambda}_{h+1} + 1, \tilde{\lambda}_{h+1} + \lambda_{h+2})}^{\triangleleft}. \end{aligned}$$

Similarly, let $M_2^* = (\lambda + \epsilon_h - \epsilon_{h+1} + E_{h+1, h+2} | O)$, $N_2^* = (\lambda + \epsilon_{h+2} - \epsilon_{h+1} + E_{h, h+1} | O)$, and we have

$$\begin{aligned} \mathcal{Y}_4 &= \phi_{(\lambda + \epsilon_h - \epsilon_{h+1} + E_{h+1, h+2} | O)}(\phi_{(\lambda + \epsilon_{h+2} - \epsilon_{h+1} + E_{h, h+1} | O)}(x_{\lambda + \epsilon_{h+2}})) \\ &= x_{\lambda + \epsilon_h} \Sigma_M \cdot \Sigma_N = x_{\lambda + \epsilon_h} T_{(\tilde{\lambda}_{h+1} + 1, \tilde{\lambda}_{h+1} + \lambda_{h+2})}^{\triangleleft} T_{(\tilde{\lambda}_h + 1, \tilde{\lambda}_h + \lambda_{h+1} - 1)}^{\triangleleft}. \end{aligned}$$

Let $\tilde{a} = \tilde{\lambda}_h$, $b = \lambda_{h+1}$, $\tilde{c} = \tilde{\lambda}_{h+2}$. Then, by Lemma 3.7(2),

$$\begin{aligned} \mathcal{Y}_1 &= x_{\lambda + \epsilon_h} c_{\tilde{a}+1} T_{(\tilde{a}+1, \tilde{a}+b)}^{\triangleleft} c_{\tilde{a}+b+1} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft} \\ &= q^b x_{\lambda + \epsilon_h} (c_{\tilde{a}+1} + T_{\tilde{a}+1}^{-1} c_{\tilde{a}+2} + T_{\tilde{a}+1}^{-1} T_{\tilde{a}+2}^{-1} c_{\tilde{a}+3} + \cdots + T_{\tilde{a}+1}^{-1} \cdots T_{\tilde{a}+b}^{-1} c_{\tilde{a}+b+1}) c_{\tilde{a}+b+1} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft}, \\ q\mathcal{Y}_2 &= qx_{\lambda + \epsilon_h} c_{\tilde{a}+b+1} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft} c_{\tilde{a}+1} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} \\ &= -q^b x_{\lambda + \epsilon_h} (c_{\tilde{a}+1} + T_{\tilde{a}+1}^{-1} c_{\tilde{a}+2} + T_{\tilde{a}+1}^{-1} T_{\tilde{a}+2}^{-1} c_{\tilde{a}+3} + \cdots + T_{\tilde{a}+1}^{-1} \cdots T_{\tilde{a}+b-1}^{-1} c_{\tilde{a}+b}) c_{\tilde{a}+b+1} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft}. \end{aligned}$$

Consequently,

$$\begin{aligned} \mathcal{Y}_1 + q\mathcal{Y}_2 &= q^b x_{\lambda + \epsilon_h} (c_{\tilde{a}+1} + T_{\tilde{a}+1}^{-1} c_{\tilde{a}+2} + T_{\tilde{a}+1}^{-1} T_{\tilde{a}+2}^{-1} c_{\tilde{a}+3} + \cdots + T_{\tilde{a}+1}^{-1} \cdots T_{\tilde{a}+b}^{-1} c_{\tilde{a}+b+1}) c_{\tilde{a}+b+1} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft} \\ &\quad - q^b x_{\lambda + \epsilon_h} (c_{\tilde{a}+1} + T_{\tilde{a}+1}^{-1} c_{\tilde{a}+2} + T_{\tilde{a}+1}^{-1} T_{\tilde{a}+2}^{-1} c_{\tilde{a}+3} + \cdots + T_{\tilde{a}+1}^{-1} \cdots T_{\tilde{a}+b-1}^{-1} c_{\tilde{a}+b}) c_{\tilde{a}+b+1} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft} \\ &= -q^b x_{\lambda + \epsilon_h} T_{\tilde{a}+1}^{-1} T_{\tilde{a}+2}^{-1} T_{\tilde{a}+3}^{-1} \cdots T_{\tilde{a}+b}^{-1} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft} \\ &= -q^b (q^{-b} x_{\lambda + \epsilon_h} T_{\tilde{a}+1} T_{\tilde{a}+2} \cdots T_{\tilde{a}+b} - q^{-b} (q-1) x_{\lambda + \epsilon_h} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft}) T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft} \quad (\text{by Lemma 3.6}) \\ &= -q^b q^{-b} x_{\lambda + \epsilon_h} T_{\tilde{a}+1} T_{\tilde{a}+2} \cdots T_{\tilde{a}+b} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft} + q^b q^{-b} (q-1) x_{\lambda + \epsilon_h} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft} \\ &= -x_{\lambda + \epsilon_h} T_{\tilde{a}+1} T_{\tilde{a}+2} \cdots T_{\tilde{a}+b} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft} + (q-1) x_{\lambda + \epsilon_h} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft}. \end{aligned}$$

On the other hand, we have

$$\begin{aligned} \mathcal{Y}_3 &= x_{\lambda + \epsilon_h} T_{(\tilde{a}+1, \tilde{a}+b)}^{\triangleleft} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft}, \\ q\mathcal{Y}_4 &= qx_{\lambda + \epsilon_h} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} = qx_{\lambda + \epsilon_h} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft}, \end{aligned}$$

and

$$-\mathcal{Y}_3 + q\mathcal{Y}_4 = -x_{\lambda+\epsilon_n} T_{\tilde{a}+1} T_{\tilde{a}+2} \cdots T_{\tilde{a}+b} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft} + (q-1)x_{\lambda+\epsilon_n} T_{(\tilde{a}+1, \tilde{a}+b-1)}^{\triangleleft} T_{(\tilde{a}+b+1, \tilde{c})}^{\triangleleft}.$$

Hence, $\mathcal{Y}_1 + q\mathcal{Y}_2 = -\mathcal{Y}_3 + q\mathcal{Y}_4$, as desired. \square

Remarks 7.3. The multiplication formulas in Theorems 5.3, 6.2, 6.3 and 6.4 are the “raw” formulas which are the counterpart of [3, Lemma 3.2] or [12, Lemma 3.1]. By normalising the ϕ -basis, a normalised version of these formulas using symmetric Gaussian polynomials in [3, Lemma 3.4 and §4.6] or [12, Proposition 4.4 and 4.5] are used in the construction of the BLM type realisations. However, an inspection on the coefficients of these multiplication formulas in Sections 5 and 6 shows that, for the candidate basis $\{[A^*] \mid A^* \in M_n(\mathbb{N}|\mathbb{N}_2)_r\}$, setting $[A^*] := v^{-l(w_A^+) + l(w_{0,ro(A)})} \phi_{A^*}$ is not enough to obtain a normalised version of these formulas.⁴ In fact, the factor c_{A^*} in defining ϕ_{A^*} needs to be normalised too. For more details, see [13].

8. AN OUTLINE OF SOME APPLICATIONS

We end this paper with an outline of some works in progress. The first main application is to use the multiplication formulas discovered in §§5–7 to give a BLM type realisation for the quantum queer supergroup.

Using a modified approach [10], the work in [3] can be interpreted as an algebra monomorphism $f_{\mathcal{S}}$ from the quantum linear group $\mathbf{U}(\mathfrak{gl}_n)$ to the direct product of q -Schur algebras $\mathcal{S}(n, r) := \mathcal{S}_q(n, r; \mathbb{Q}(v))$ ($q = v^2$), the top horizontal map in the diagram below (see Corollary 2.4).

$$\begin{array}{ccc} \mathbf{U}(\mathfrak{gl}_n) & \xrightarrow{f_{\mathcal{S}}} & \prod_{r \geq 0} \mathcal{S}(n, r) \\ \downarrow \cap & & \downarrow \cap \\ \mathbf{U}(\mathfrak{q}_n) & \xrightarrow{\exists f_{\mathcal{Q}}?} & \prod_{r \geq 0} \tilde{\mathcal{Q}}(n, r) \end{array}$$

The image of $f_{\mathcal{S}}$ can be presented in terms of a basis and explicit multiplication formulas as relations. Since $\mathbf{U}(\mathfrak{gl}_n)$ is a subalgebra of the quantum queer supergroup $\mathbf{U}(\mathfrak{q}_n)$ and, by Corollary 2.4, each q -Schur algebra is a subalgebra of the “twisted” queer q -Schur superalgebra $\tilde{\mathcal{Q}}(n, r)$ (see Remark 2.2), there are two vertical inclusion maps.⁵ Thus, it is natural to ask if there exists a superalgebra homomorphism $f_{\mathcal{Q}}$ such that the above diagram commutes. We will answer this question in the forthcoming paper [13].

Here we outline the method used in earlier works and point out the degree of difficulty in the queer case. The fundamental multiplication formulas established in §§5–7 are related to the standard basis for queer q -Schur superalgebras. In the non-super case, stabilisation properties of these formulas are established to construct the modified quantum linear groups, i.e., the algebra $\dot{\mathbf{U}}$ in [24] associated with quantum \mathfrak{gl}_n ; see [3] and [8, Part 5] for more details. For the algebra $\dot{\mathbf{U}}$ (without 1) of $\mathbf{U} = \mathbf{U}(\mathfrak{gl}_{m|n})$, a similar work has been done; see [12, 15]. Then, by introducing certain “long elements” from the standard basis, the “short” multiplication formulas can be extended to these long elements. One then uses these long element multiplication formulas to define a subalgebra inside the direct product of q -Schur algebra/superalgebras. By further checking the defining relations, one proves that this subalgebra is isomorphic to a quantum linear group/supergroup.

⁴Here w_A^+ is the longest elements of the double coset associated with A ; see [8, Lemma 13.10] for a geometric interpretation of $-w_A^+ + w_{0,ro(A)}$.

⁵In the diagram, a twisted version $\tilde{\mathcal{Q}}(n, r)$ is used in order to make it true over \mathbb{Q} . If we use the complex number field \mathbb{C} , then the twisted version is isomorphic to $\mathcal{Q}(n, r)$; see [13].

It has been expected for years that such a construction should be generalised to quantum queer supergroup $\mathbf{U}(\mathfrak{q}_n)$. However, deriving the multiplication formulas by the odd generators became the main hurdle. Just as indicated in the theorems in Section 6, there are no such explicit formulas! However, the SDP condition introduced in the paper provides a path for us to overcome the hurdle. The work [13] will complete the task above for quantum queer supergroups. In this way, we also reestablish the Schur–Olshanski duality given by truncating an infinite sum to a finite one as surjective algebra homomorphisms from $\mathbf{U}(\mathfrak{q}_n)$ to queer q -Schur algebras.

We then can give another important application which establishes the Schur–Olshanski duality at the integral level. Thus, representations of the hyperalgebra of $\mathbf{U}(\mathfrak{q}_n)$, especially those polynomial ones, in positive quantum characteristic can be investigated with a connection to the modular representations of Hecke–Clifford superalgebras, generalising some of the classical theories in [5, 6] to the quantum case. For more details, see the forthcoming paper [19].

APPENDIX A. SOME MULTIPLICATION FORMULAS IN $\mathcal{H}_{r,R}^c$

In this appendix, we prove (6.4.4). Let $(\diamond)_k := x_{\lambda_{(h)}^-} \left(T_{d_{A_{h,k}^-}} c_{\tilde{a}_{h,k}} \right) T_{(\tilde{a}_{h,k}-1, \tilde{a}_{h,k}-a_{h,k}+1)}^{\triangleright} c_{A^*} \Sigma_A$.

Lemma A.1. *Let $h \in [1, n-1]$ and $A^* = (a_{i,j}^{\bar{0}} | a_{i,j}^{\bar{1}}) \in M_n(\mathbb{N} | \mathbb{N}_2)_r$. Assume $A = \lfloor A^* \rfloor$ and $\lambda = \text{ro}(A)$. The following formula holds in $\mathcal{H}_{r,R}^c$.*

$$\begin{aligned} (\diamond)_k &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + a_{h,k}^{\bar{1}}} T_{(A^{\bar{0}} - E_{h,k} | A^{\bar{1}} + E_{h+1,k})} \\ &\quad + (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + a_{h,k}^{\bar{1}} + 1} q^{-1} \llbracket a_{h+1,k} + 1 \rrbracket_{q^2, q} T_{(A^{\bar{0}} - E_{h,k} + 2E_{h+1,k} | A^{\bar{1}} - E_{h+1,k})} \\ &\quad + (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + 1} q^{a_{h,k}} q^{-1} \llbracket a_{h+1,k}^{\bar{0}} + 1 \rrbracket_q T_{(A^{\bar{0}} + E_{h+1,k} | A^{\bar{1}} - E_{h,k})}. \end{aligned} \quad (\text{A.1.1})$$

Proof. Throughout the proof, we keep the notations in the proof of Theorem 6.4 as well as Notation 5.2. In particular, $\nu = \nu_A$, $\nu^- = \nu_{A_{h,k}^-}$ and δ^- are as in (5.2.2) and $\tilde{a} = \tilde{a}_{h,k}$, $a = a_{h,k}$, $b = a_{h+1,k}$. We also record the following some of which have been (implicitly) used before.

$$\begin{aligned} (1) \quad &x_{\nu^-} = x_{\mathcal{D}_-} x_{\delta^-} = x_{\delta^-} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft}, \quad x_{\nu} = x_{\delta^-} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright}; \\ (2) \quad &x_{\nu} c_{q, \tilde{a}-a+1, \tilde{a}} = (c_{q, \tilde{a}-a+1, \tilde{a}})' x_{\nu}, \quad x_{\nu} c_{q, \tilde{a}+1, \tilde{a}+b} = (c_{q, \tilde{a}+1, \tilde{a}+b})' x_{\nu}; \\ (3) \quad &x_{\nu^-} c_{q, \tilde{a}-a+1, \tilde{a}-1} = (c_{q, \tilde{a}-a+1, \tilde{a}-1})' x_{\nu^-}, \quad x_{\nu^-} c_{q, \tilde{a}, \tilde{a}+b} = (c_{q, \tilde{a}, \tilde{a}+b})' x_{\nu^-}; \\ (4) \quad &x_{\delta^-} c_{q, \tilde{a}-a+1, \tilde{a}-1} = (c_{q, \tilde{a}-a+1, \tilde{a}-1})' x_{\delta^-}, \quad x_{\delta^-} c_{\tilde{a}} = c_{\tilde{a}} x_{\delta^-}, \quad x_{\delta^-} c_{q, \tilde{a}+1, \tilde{a}+b} = (c_{q, \tilde{a}+1, \tilde{a}+b})' x_{\delta^-}. \end{aligned} \quad (\text{A.1.2})$$

Because it can not be able to commute x_{ν^-} with $(c_{q, \tilde{a}-a+1, \tilde{a}})'$ and $c_{q, \tilde{a}+1, \tilde{a}+b}$ as above formulas, in the following proof, we usually rewrite

$$(c_{q, \tilde{a}-a+1, \tilde{a}})' = (c_{q, \tilde{a}-a+1, \tilde{a}-1})' + q^{a-1} c_{\tilde{a}}, \quad c_{q, \tilde{a}+1, \tilde{a}+b} = c_{q, \tilde{a}, \tilde{a}+b} - q^b c_{\tilde{a}}. \quad (\text{A.1.3})$$

Since $s_{\tilde{a}}, s_{\tilde{a}+1}, \dots, s_{\tilde{a}+b-1} \in \mathfrak{S}_{\nu^-}$,

$$x_{\nu^-} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} = \llbracket b+1 \rrbracket_q x_{\nu^-}, \quad (\text{A.1.4})$$

and from Lemma 3.7,

$$x_{\nu^-} c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} = x_{\nu^-} c_{q, \tilde{a}, \tilde{a}+b}. \quad (\text{A.1.5})$$

By (5.2.4) and (A.1.2)(1),

$$x_{\lambda_{(h)}^-} T_{d_{A_{h,k}^-}} = x_{\lambda_{(h)}^-} \setminus \nu^- T_{d_{A_{h,k}^-}} x_{\mathcal{D}_-} x_{\delta^-}.$$

Thus, $(\diamond)_k$ can be rewritten as

$$(\diamond)_k = x_{\lambda_{(h)}^-} \setminus \nu^- T_{d_{A_{h,k}^-}} x_{\mathcal{D}_-} x_{\delta^-} c_{\tilde{a}} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A.$$

Moreover, by (A.1.2)(2)&(4), $x_{\delta^-}c_{\tilde{a}} = c_{\tilde{a}}x_{\delta^-}$ and $x_{\nu^-}c_{A^*} = c'_{A^*}x_{\nu^-}$. Combining $x_{\nu^-} = x_{\delta^-}T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright}$ and $x_{\nu^-}\Sigma_A = x_{\mu^-}\Sigma_{A_{h,k}^-}$ gives

$$(\diamondsuit)_k = x_{\lambda_{(h)}^-} \setminus \nu^- T_{d_{A_{h,k}^-}} x_{\mathcal{D}_-} c_{\tilde{a}} x_{\delta^-} T_{(\tilde{a}-1, \tilde{a}-a+1)}^{\triangleright} c_{A^*} \Sigma_A = x_{\lambda_{(h)}^-} \setminus \nu^- T_{d_{A_{h,k}^-}} \underbrace{x_{\mathcal{D}_-} c_{\tilde{a}} c'_{A^*} x_{\nu^-}}_{=\underline{\mathbf{m}}^-} \Sigma_{A_{h,k}^-}. \quad (\text{A.1.6})$$

Since $c'_{A^*} = (\overleftarrow{c}_{A^*}^{h,k})' (c_{A^*;k}^{h,h+1})' (\overrightarrow{c}_{A^*}^{h+1,k})'$ (see (5.2.1)), moving $(c_{A^*;k}^{h,h+1})'$ to the left yields

$$c'_{A^*} = (-1)^{(a_{h,k}^{\bar{1}} + a_{h+1,k}^{\bar{1}}) \tilde{a}_{h-1,k}^{\bar{1}}} (c_{A^*;k}^{h,h+1})' (\overleftarrow{c}_{A^*}^{h,k})' (\overrightarrow{c}_{A^*}^{h+1,k})'.$$

On the other hand, $x_{\nu^-} = x_{\delta^-} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft}$, $(\overleftarrow{c}_{A^*}^{h,k})' (\overrightarrow{c}_{A^*}^{h+1,k})' x_{\delta^-} = x_{\delta^-} (\overleftarrow{c}_{A^*}^{h,k}) (\overrightarrow{c}_{A^*}^{h+1,k})$, and $T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft}$ commutes with $\overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k}$. The middle part $\underline{\mathbf{m}}^-$ becomes

$$\underline{\mathbf{m}}^- = (-1)^{(a_{h,k}^{\bar{1}} + a_{h+1,k}^{\bar{1}}) \tilde{a}_{h-1,k}^{\bar{1}}} x_{\mathcal{D}_-} c_{\tilde{a}} (c_{A^*;k}^{h,h+1})' x_{\delta^-} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k}.$$

The following discussion is parallel to the cases in the proof of Theorem 6.3 according to the four values of $(a_{h,k}^{\bar{1}}, a_{h+1,k}^{\bar{1}})$ in $c_{A^*;k}^{h,h+1} = (c_{q, \tilde{a}_{h-1,k}+1, \tilde{a}_{h,k}})^{a_{h,k}^{\bar{1}}} \cdot (c_{q, \tilde{a}_{h,k}+1, \tilde{a}_{h+1,k}})^{a_{h+1,k}^{\bar{1}}}$ to determine $\underline{\mathbf{m}}^-$ and then $(\diamondsuit)_k$.

Case 1 ($a_{h,k}^{\bar{1}} = 0, a_{h+1,k}^{\bar{1}} = 0$). In this case, $c_{A^*;k}^{h,h+1} = 1$ and so,

$$\underline{\mathbf{m}}^- = x_{\mathcal{D}_-} c_{\tilde{a}} x_{\delta^-} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} = x_{\mathcal{D}_-} x_{\delta^-} c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k}$$

Applying (A.1.2)(2) and (A.1.5), we have $x_{\mathcal{D}_-} x_{\delta^-} c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} = x_{\nu^-} c_{q, \tilde{a}, \tilde{a}+b}$. So,

$$\begin{aligned} \underline{\mathbf{m}}^- &= x_{\nu^-} c_{q, \tilde{a}, \tilde{a}+b} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} = (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} x_{\nu^-} \overleftarrow{c}_{A^*}^{h,k} c_{q, \tilde{a}, \tilde{a}+b} \overrightarrow{c}_{A^*}^{h+1,k} \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} x_{\nu^-} c_{(A^{\bar{0}} - E_{h,k} | A^{\bar{1}} + E_{h+1,k})} \end{aligned}$$

Hence, (A.1.6) becomes

$$\begin{aligned} (\diamondsuit)_k &= x_{\lambda_{(h)}^-} \setminus \nu^- T_{d_{A_{h,k}^-}} ((-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} x_{\nu^-} c_{(A^{\bar{0}} - E_{h,k} | A^{\bar{1}} + E_{h+1,k})}) \Sigma_{A_{h,k}^-} \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} x_{\lambda_{(h)}^-} T_{d_{A_{h,k}^-}} c_{(A^{\bar{0}} - E_{h,k} | A^{\bar{1}} + E_{h+1,k})} \Sigma_{A_{h,k}^-} = (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} T_{(A^{\bar{0}} - E_{h,k} | A^{\bar{1}} + E_{h+1,k})}. \end{aligned} \quad (\text{A.1.7})$$

Case 2 ($a_{h,k}^{\bar{1}} = 0, a_{h+1,k}^{\bar{1}} = 1$). Here $c_{A^*;k}^{h,h+1} = c_{q, \tilde{a}+1, \tilde{a}+b}$. By (A.1.2)(4),

$$c_{\tilde{a}} (c_{q, \tilde{a}+1, \tilde{a}+b})' x_{\delta^-} = x_{\delta^-} c_{\tilde{a}} (c_{q, \tilde{a}+1, \tilde{a}+b})$$

Thus,

$$\begin{aligned} \underline{\mathbf{m}}^- &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} x_{\mathcal{D}_-} c_{\tilde{a}} (c_{q, \tilde{a}+1, \tilde{a}+b})' x_{\delta^-} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} x_{\nu^-} c_{\tilde{a}} (c_{q, \tilde{a}+1, \tilde{a}+b}) T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k}. \end{aligned}$$

Since $c_{\tilde{a}} c_{q, \tilde{a}+1, \tilde{a}+b} = -c_{q, \tilde{a}+1, \tilde{a}+b} c_{\tilde{a}}$, (A.1.3) and (A.1.2)(3) imply

$$\begin{aligned} x_{\nu^-} c_{\tilde{a}} c_{q, \tilde{a}+1, \tilde{a}+b} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} &= -x_{\nu^-} c_{q, \tilde{a}+1, \tilde{a}+b} c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \\ &= -x_{\nu^-} (c_{q, \tilde{a}, \tilde{a}+b} - q^b c_{\tilde{a}}) c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} = -x_{\nu^-} c_{q, \tilde{a}, \tilde{a}+b} c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} + x_{\nu^-} q^b c_{\tilde{a}} c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft}. \end{aligned}$$

By (A.1.2)(3) again and (A.1.5),

$$\begin{aligned} -x_{\nu^-} c_{q, \tilde{a}, \tilde{a}+b} c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} &= -c'_{q, \tilde{a}, \tilde{a}+b} x_{\nu^-} c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \\ &= -c'_{q, \tilde{a}, \tilde{a}+b} x_{\nu^-} c_{q, \tilde{a}, \tilde{a}+b} = -x_{\nu^-} c_{q, \tilde{a}, \tilde{a}+b}^2 = \llbracket b+1 \rrbracket_{q^2} x_{\nu^-} \end{aligned}$$

and, by applying (A.1.4), $x_{\nu^-} q^b c_{\tilde{a}} c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} = -q^b \llbracket b+1 \rrbracket_q x_{\nu^-}$. Hence,

$$\begin{aligned} \underline{\mathbf{m}}^- &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} x_{\nu^-} (\llbracket b+1 \rrbracket_{q^2} - q^b \llbracket b+1 \rrbracket_q) \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{-1} \llbracket b+1 \rrbracket_{q^2} x_{\nu^-} c_{(A^{\bar{0}} - E_{h,k} + 2E_{h+1,k} | A^{\bar{1}} - E_{h+1,k})}, \end{aligned}$$

and

$$\begin{aligned} (\diamond)_k &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}+1} q^{-1} \llbracket b+1 \rrbracket_{q^2, q} x_{\nu^-} c_{(A^{\bar{0}}-E_{h,k}+2E_{h+1,k}|A^{\bar{1}}-E_{h+1,k})} \Sigma_{A_{h,k}^-} \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}+1} q^{-1} \llbracket b+1 \rrbracket_{q^2, q} T_{(A^{\bar{0}}-E_{h,k}+2E_{h+1,k}|A^{\bar{1}}-E_{h+1,k})}. \end{aligned} \quad (\text{A.1.8})$$

Case 3 ($a_{h,k}^{\bar{1}} = 1, a_{h+1,k}^{\bar{1}} = 0$). We have $c_{A^*;k}^{h,h+1} = c_{q,\tilde{a}-a+1,\tilde{a}}$ and

$$\underline{\mathbf{m}}^- = (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} x_{\mathcal{D}_-} c_{\tilde{a}} (c_{q,\tilde{a}-a+1,\tilde{a}})' x_{\delta^-} T_{(\tilde{a},\tilde{a}+b-1)}^{\triangleleft} \overleftarrow{C}_{A^*}^{h,k} \overrightarrow{C}_{A^*}^{h+1,k}.$$

Applying (A.1.2)(4),(A.1.3),(A.1.4) and (A.1.5) yields

$$\begin{aligned} & x_{\mathcal{D}_-} c_{\tilde{a}} (c_{q,\tilde{a}-a+1,\tilde{a}})' x_{\delta^-} T_{(\tilde{a},\tilde{a}+b-1)}^{\triangleleft} \\ &= x_{\mathcal{D}_-} c_{\tilde{a}} ((c_{q,\tilde{a}-a+1,\tilde{a}-1})' + q^{a-1} c_{\tilde{a}}) x_{\delta^-} T_{(\tilde{a},\tilde{a}+b-1)}^{\triangleleft} \\ &= x_{\mathcal{D}_-} c_{\tilde{a}} (c_{q,\tilde{a}-a+1,\tilde{a}-1})' x_{\delta^-} T_{(\tilde{a},\tilde{a}+b-1)}^{\triangleleft} + q^{a-1} x_{\mathcal{D}_-} c_{\tilde{a}} c_{\tilde{a}} x_{\delta^-} T_{(\tilde{a},\tilde{a}+b-1)}^{\triangleleft} \\ &= -x_{\mathcal{D}_-} (c_{q,\tilde{a}-a+1,\tilde{a}-1})' x_{\delta^-} c_{\tilde{a}} T_{(\tilde{a},\tilde{a}+b-1)}^{\triangleleft} - q^{a-1} x_{\mathcal{D}_-} x_{\delta^-} T_{(\tilde{a},\tilde{a}+b-1)}^{\triangleleft} \\ &= -x_{\nu^-} (c_{q,\tilde{a}-a+1,\tilde{a}-1}) (c_{q,\tilde{a},\tilde{a}+b}) - q^{a-1} \llbracket b+1 \rrbracket_q x_{\nu^-}, \end{aligned}$$

and

$$\begin{aligned} \underline{\mathbf{m}}^- &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} (-x_{\nu^-} (c_{q,\tilde{a}-a+1,\tilde{a}-1}) (c_{q,\tilde{a},\tilde{a}+b}) - q^{a-1} \llbracket b+1 \rrbracket_q x_{\nu^-}) \overleftarrow{C}_{A^*}^{h,k} \overrightarrow{C}_{A^*}^{h+1,k} \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}+1} x_{\nu^-} \overleftarrow{C}_{A^*}^{h,k} (c_{q,\tilde{a}-a+1,\tilde{a}-1}) (c_{q,\tilde{a},\tilde{a}+b}) \overrightarrow{C}_{A^*}^{h+1,k} + (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}} q^{a-1} \llbracket b+1 \rrbracket_q x_{\nu^-} \overleftarrow{C}_{A^*}^{h,k} \overrightarrow{C}_{A^*}^{h+1,k} \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}+1} (x_{\nu^-} c_{(A^{\bar{0}}-E_{h,k}|A^{\bar{1}}+E_{h+1,k})} + q^{a-1} \llbracket b+1 \rrbracket_q x_{\nu^-} c_{(A^{\bar{0}}+E_{h+1,k}|A^{\bar{1}}-E_{h,k})}). \end{aligned}$$

Consequently, (A.1.6) becomes

$$\begin{aligned} (\diamond)_k &= x_{\lambda_{(h)}^-} \nu^- T_{d_{A_{h,k}^-}} \{ (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}+1} (x_{\nu^-} c_{(A^{\bar{0}}-E_{h,k}|A^{\bar{1}}+E_{h+1,k})} \\ &\quad + q^{a-1} \llbracket b+1 \rrbracket_q x_{\nu^-} c_{(A^{\bar{0}}+E_{h+1,k}|A^{\bar{1}}-E_{h,k})}) \} \Sigma_{A_{h,k}^-} \\ &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}}+1} (T_{(A^{\bar{0}}-E_{h,k}|A^{\bar{1}}+E_{h+1,k})} + q^{a-1} \llbracket b+1 \rrbracket_q T_{(A^{\bar{0}}+E_{h+1,k}|A^{\bar{1}}-E_{h,k})}). \end{aligned} \quad (\text{A.1.9})$$

Case 4 ($a_{h,k}^{\bar{1}} = 1, a_{h+1,k}^{\bar{1}} = 1$). Here we have $c_{A^*;k}^{h,h+1} = c_{q,\tilde{a}-a+1,\tilde{a}} c_{q,\tilde{a}+1,\tilde{a}+b}$ and

$$\underline{\mathbf{m}}^- = x_{\mathcal{D}_-} \underbrace{c_{\tilde{a}} (c_{q,\tilde{a}-a+1,\tilde{a}} c_{q,\tilde{a}+1,\tilde{a}+b})}'_{=\mathcal{N}} x_{\delta^-} T_{(\tilde{a},\tilde{a}+b-1)}^{\triangleleft} \overleftarrow{C}_{A^*}^{h,k} \overrightarrow{C}_{A^*}^{h+1,k}.$$

Since

$$\begin{aligned} c_{\tilde{a}} c'_{q,\tilde{a}-a+1,\tilde{a}} &= c_{\tilde{a}} (c'_{q,\tilde{a}-a+1,\tilde{a}-1} + q^{a-1} c_{\tilde{a}}) = -c'_{q,\tilde{a}-a+1,\tilde{a}-1} c_{\tilde{a}} - q^{a-1}, \\ c_{\tilde{a}} c_{q,\tilde{a},\tilde{a}+b} &= c_{\tilde{a}} (c_{q,\tilde{a}+1,\tilde{a}+b} + q^b c_{\tilde{a}}) = -c_{q,\tilde{a}+1,\tilde{a}+b} c_{\tilde{a}} - q^b = -c_{q,\tilde{a},\tilde{a}+b} c_{\tilde{a}} - 2q^b, \end{aligned}$$

by (A.1.2)(4) and (A.1.3), we have

$$\begin{aligned} \mathcal{N} &= c_{\tilde{a}} (c_{q,\tilde{a}-a+1,\tilde{a}})' (c_{q,\tilde{a}+1,\tilde{a}+b})' x_{\delta^-} \\ &= (-c'_{q,\tilde{a}-a+1,\tilde{a}-1} c_{\tilde{a}} - q^{a-1}) x_{\delta^-} (c_{q,\tilde{a}+1,\tilde{a}+b}) \\ &= x_{\delta^-} (-c_{q,\tilde{a}-a+1,\tilde{a}-1} c_{\tilde{a}} - q^{a-1}) (c_{q,\tilde{a}+1,\tilde{a}+b}) \\ &= -x_{\delta^-} c_{q,\tilde{a}-a+1,\tilde{a}-1} c_{\tilde{a}} (c_{q,\tilde{a}+1,\tilde{a}+b}) - q^{a-1} x_{\delta^-} (c_{q,\tilde{a}+1,\tilde{a}+b}). \end{aligned}$$

Thus,

$$\begin{aligned} \underline{\mathbf{m}}^- &= x_{\mathcal{D}_-} \mathcal{N} T_{(\tilde{a},\tilde{a}+b-1)}^{\triangleleft} \overleftarrow{C}_{A^*}^{h,k} \overrightarrow{C}_{A^*}^{h+1,k} \\ &= x_{\mathcal{D}_-} \{ -x_{\delta^-} c_{q,\tilde{a}-a+1,\tilde{a}-1} c_{\tilde{a}} (c_{q,\tilde{a}+1,\tilde{a}+b}) - q^{a-1} x_{\delta^-} (c_{q,\tilde{a}+1,\tilde{a}+b}) \} T_{(\tilde{a},\tilde{a}+b-1)}^{\triangleleft} \overleftarrow{C}_{A^*}^{h,k} \overrightarrow{C}_{A^*}^{h+1,k} \\ &= \{ -x_{\nu^-} c_{q,\tilde{a}-a+1,\tilde{a}-1} c_{\tilde{a}} (c_{q,\tilde{a}+1,\tilde{a}+b}) \} T_{(\tilde{a},\tilde{a}+b-1)}^{\triangleleft} \overleftarrow{C}_{A^*}^{h,k} \overrightarrow{C}_{A^*}^{h+1,k} \\ &\quad + \{ -q^{a-1} x_{\nu^-} (c_{q,\tilde{a}+1,\tilde{a}+b}) \} T_{(\tilde{a},\tilde{a}+b-1)}^{\triangleleft} \overleftarrow{C}_{A^*}^{h,k} \overrightarrow{C}_{A^*}^{h+1,k} \\ &= \mathcal{M}_1 + \mathcal{M}_2, \end{aligned} \quad (\text{A.1.10})$$

where

$$\begin{aligned}\mathcal{M}_1 &= \{-x_{\nu^-} c_{q, \tilde{a}-a+1, \tilde{a}-1} c_{\tilde{a}}(c_{q, \tilde{a}+1, \tilde{a}+b})\} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k}, \\ \mathcal{M}_2 &= \{-q^{a-1} x_{\nu^-}(c_{q, \tilde{a}+1, \tilde{a}+b})\} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k}.\end{aligned}$$

We now modify \mathcal{M}_1 and \mathcal{M}_2 into a desired form. Since

$$\begin{aligned}& x_{\nu^-}(c_{q, \tilde{a}-a+1, \tilde{a}-1}) c_{\tilde{a}}(c_{q, \tilde{a}+1, \tilde{a}+b}) T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \\ &= x_{\nu^-}(c_{q, \tilde{a}-a+1, \tilde{a}-1}) c_{\tilde{a}}(c_{q, \tilde{a}, \tilde{a}+b} - q^b c_{\tilde{a}}) T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \\ &= x_{\nu^-}(c_{q, \tilde{a}-a+1, \tilde{a}-1}) (-c_{q, \tilde{a}, \tilde{a}+b} c_{\tilde{a}} + q^b c_{\tilde{a}}^2) T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \\ &= (-x_{\nu^-}(c_{q, \tilde{a}-a+1, \tilde{a}-1}) (c_{q, \tilde{a}, \tilde{a}+b}) c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft}) + (-q^b x_{\nu^-}(c_{q, \tilde{a}-a+1, \tilde{a}-1}) T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft}),\end{aligned}$$

applying (A.1.2)(3) and (A.1.4) yields

$$\begin{aligned}& -x_{\nu^-}(c_{q, \tilde{a}-a+1, \tilde{a}-1}) (c_{q, \tilde{a}, \tilde{a}+b}) c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \\ &= -(c_{q, \tilde{a}-a+1, \tilde{a}-1})' (c_{q, \tilde{a}, \tilde{a}+b})' x_{\nu^-} c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} = -(c_{q, \tilde{a}-a+1, \tilde{a}-1})' (c_{q, \tilde{a}, \tilde{a}+b})' x_{\nu^-} (c_{q, \tilde{a}, \tilde{a}+b}) \\ &= -x_{\nu^-}(c_{q, \tilde{a}-a+1, \tilde{a}-1}) (c_{q, \tilde{a}, \tilde{a}+b})^2 = \llbracket b+1 \rrbracket_{q^2} x_{\nu^-}(c_{q, \tilde{a}-a+1, \tilde{a}-1}),\end{aligned}$$

and, by (A.1.5),

$$-q^b x_{\nu^-}(c_{q, \tilde{a}-a+1, \tilde{a}-1}) T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} = -q^b \llbracket b+1 \rrbracket_q x_{\nu^-}(c_{q, \tilde{a}-a+1, \tilde{a}-1}).$$

Because of $\llbracket b+1 \rrbracket_{q^2} - q^b \llbracket b+1 \rrbracket_q = -q^{-1} \llbracket b+1 \rrbracket_{q^2, q}$,

$$\begin{aligned}\mathcal{M}_1 &= (-x_{\nu^-} c_{q, \tilde{a}-a+1, \tilde{a}-1} c_{\tilde{a}}(c_{q, \tilde{a}+1, \tilde{a}+b})) T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \\ &= q^{-1} \llbracket b+1 \rrbracket_{q^2, q} x_{\nu^-}(c_{q, \tilde{a}-a+1, \tilde{a}-1}) \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \\ &= (-1)^{\tilde{a}_{h-1, k}^{\bar{1}}} q^{-1} \llbracket b+1 \rrbracket_{q^2, q} x_{\nu^-} \overleftarrow{c}_{A^*}^{h,k} (c_{q, \tilde{a}-a+1, \tilde{a}-1}) \overrightarrow{c}_{A^*}^{h+1,k} \\ &= (-1)^{\tilde{a}_{h-1, k}^{\bar{1}}} q^{-1} \llbracket b+1 \rrbracket_{q^2, q} x_{\nu^-} c_{(A^{\bar{0}} - E_{h, k} + 2E_{h+1, k} | A^{\bar{1}} - E_{h+1, k})}.\end{aligned}\tag{A.1.11}$$

Now, we calculate \mathcal{M}_2 . Since

$$\begin{aligned}& -q^{a-1} x_{\nu^-}(c_{q, \tilde{a}+1, \tilde{a}+b}) T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \\ &= -q^{a-1} x_{\nu^-}(c_{q, \tilde{a}, \tilde{a}+b} - q^b c_{\tilde{a}}) T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \quad (\text{by (A.1.3)}) \\ &= -q^{a-1} x_{\nu^-} c_{q, \tilde{a}, \tilde{a}+b} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} + q^{a+b-1} x_{\nu^-} c_{\tilde{a}} T_{(\tilde{a}, \tilde{a}+b-1)}^{\triangleleft} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \\ &= \left(-q^{a-1} \llbracket b+1 \rrbracket_q x_{\nu^-} c_{q, \tilde{a}, \tilde{a}+b} + q^{a+b-1} x_{\nu^-} c_{q, \tilde{a}, \tilde{a}+b}\right) \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \quad (\text{by (A.1.4) and (A.1.5)}) \\ &= (-q^{a-1} \llbracket b+1 \rrbracket_q + q^{a+b-1}) x_{\nu^-} c_{q, \tilde{a}, \tilde{a}+b} \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k},\end{aligned}$$

and $(-q^{a-1} \llbracket b+1 \rrbracket_q + q^{a+b-1}) = -q^{a-1} \llbracket b \rrbracket_q$, we have

$$\begin{aligned}\mathcal{M}_2 &= -q^{a-1} \llbracket b \rrbracket_q x_{\nu^-}(c_{q, \tilde{a}, \tilde{a}+b}) \overleftarrow{c}_{A^*}^{h,k} \overrightarrow{c}_{A^*}^{h+1,k} \\ &= (-1)^{\tilde{a}_{h-1, k}^{\bar{1}}} q^{a-1} \llbracket b \rrbracket_q x_{\nu^-} \overleftarrow{c}_{A^*}^{h,k} (c_{q, \tilde{a}, \tilde{a}+b}) \overrightarrow{c}_{A^*}^{h+1,k} \\ &= (-1)^{\tilde{a}_{h-1, k}^{\bar{1}}} q^{a-1} \llbracket b \rrbracket_q x_{\nu^-} c_{(A^{\bar{0}} + E_{h+1, k} | A^{\bar{1}} - E_{h, k})}.\end{aligned}\tag{A.1.12}$$

Now, combining the formulas (A.1.10), (A.1.11) and (A.1.12), we obtain in this case

$$\begin{aligned}(\diamondsuit)_k &= x_{\lambda_{(h)}^-} \nu^- T_{d_{A_{h, k}^-}} (\underline{\mathbf{m}}^-) \Sigma_{A_{h, k}^-} \\ &= (-1)^{\tilde{a}_{h-1, k}^{\bar{1}} + a_{h, k}^{\bar{1}}} q^{-1} \llbracket b+1 \rrbracket_{q^2, q} T_{(A^{\bar{0}} - E_{h, k} + 2E_{h+1, k} | A^{\bar{1}} - E_{h+1, k})} \\ &\quad + (-1)^{\tilde{a}_{h-1, k}^{\bar{1}}} q^{a-1} \llbracket b \rrbracket_q T_{(A^{\bar{0}} + E_{h+1, k} | A^{\bar{1}} - E_{h, k})}.\end{aligned}\tag{A.1.13}$$

Finally, let

$$\begin{aligned} (\diamond)'_k &= (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + a_{h,k}^{\bar{1}}} T_{(A^{\bar{0}} - E_{h,k} | A^{\bar{1}} + E_{h+1,k})} \\ &\quad + (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + a_{h,k}^{\bar{1}} + 1} q^{-1} \llbracket a_{h+1,k} + 1 \rrbracket_{q^2, q} T_{(A^{\bar{0}} - E_{h,k} + 2E_{h+1,k} | A^{\bar{1}} - E_{h+1,k})} \\ &\quad + (-1)^{\tilde{a}_{h-1,k}^{\bar{1}} + 1} q^{a_{h,k} - 1} \llbracket a_{h+1,k}^{\bar{0}} + 1 \rrbracket_q T_{(A^{\bar{0}} + E_{h+1,k} | A^{\bar{1}} - E_{h,k})}. \end{aligned}$$

A case by case discussion, using (A.1.7), (A.1.8), (A.1.9), (A.1.13), and Convention 5.1, proves $(\diamond)'_k = (\diamond)_k$. \square

REFERENCES

- [1] H. Bao, J. Kujawa, Y. Li, W. Wang. *Geometric Schur duality of classical type*. Transf. Groups, **23** (2014), 329–389.
- [2] H. Bao, W. Wang. *A new approach to Kazhdan-Lusztig theory of type B via quantum symmetric pairs*. Asterisque **402**, Soc. Math. de France, Paris, 2018.
- [3] A. Beilinson, G. Lusztig, R. MacPherson. *A geometric setting for the quantum deformation of \mathfrak{gl}_n* . Duke Math. J., **61** (1990), 655–677.
- [4] G. Benkart, N. Guay, J. Jung, S. Kang, S. Wilcox. *Quantum walled Brauer-Clifford superalgebras*. J. Algebra, **454** (2016), 433–474.
- [5] J. Brundan, A. Kleshchev. *Projective representations of symmetric groups via Sergeev duality*. Math. Z. **239** (2002), 27–68.
- [6] J. Brundan, A. Kleshchev. *Modular representations of the supergroup $Q(n)$, I*. J. Algebra, **260** (2003), 64–98.
- [7] S. Cheng, W. Wang. *Dualities and Representations of Lie Superalgebras*. Graduate Studies in Mathematics **144**, Amer. Math. Soc., Providence, 2009.
- [8] B. Deng, J. Du, B. Parshall, J. Wang. *Finite Dimensional Algebras and Quantum Groups*. Math. Surveys and Monographs **150**. Amer. Math. Soc., Providence, 2008.
- [9] J. Du. *Cells in certain sets of matrices*. Tôhoku Math. J. **48** (1996), 417–427.
- [10] J. Du, Q. Fu. *Quantum affine \mathfrak{gl}_n via Hecke algebras*. Adv. Math. **282** (2015), 23–46.
- [11] J. Du, Q. Fu. *A modified BLM approach to quantum affine \mathfrak{gl}_n* . Math. Z. **266** (2010), 747–781.
- [12] J. Du, H. Gu. *A realisation of the quantum supergroup $U(\mathfrak{gl}_{m|n})$* . J. Algebra **404** (2014), 60–99.
- [13] J. Du, H. Gu, Z. Li, J. Wan, *Constructing the quantum queer supergroup using Hecke-Clifford superalgebras*, in preparation.
- [14] J. Du, H. Gu, Z. Zhou. *Multiplication formulas and semisimplicity for q -Schur superalgebras*. Nagoya Math. J. **237** (2020), 98–126.
- [15] J. Du, H. Gu, Z. Zhou, *Canonical bases for the modified quantum $\mathfrak{gl}_{m|n}$* , Proc. Symp. Pure Math., to appear.
- [16] J. Du, Y. Lin, Z. Zhou. *Quantum queer supergroups via v -differential operators*. J. Algebra **599** (2022), 48–103.
- [17] J. Du, J. Wan. *Presenting queer Schur superalgebras*. Internat. Math. Res. Notices (IMRN) Vol. 2015:8 (2015), 2210–2272.
- [18] J. Du, J. Wan. *The queer q -Schur superalgebra*. J. Aust. Math. Soc. **105** (2018), 316–346.
- [19] J. Du, J. Wan, Z. Zhou, *Polynomial superrepresentations of $\mathbf{U}(\mathfrak{q}_n)$ at roots of unity*, in preparation.
- [20] H. Gu, Z. Li, Y. Lin. *The integral Schur-Weyl-Sergeev duality*. J. Pure Appl. Algebra **226** (2022), 107044.
- [21] Z. Fan, C. Lai, Y. Li, L. Luo, W. Wang. *Affine flag varieties and quantum symmetric pairs*, Mem. Amer. Math. Soc. **265** (2020), no.1285, v+123pp.
- [22] V. Ginzburg, E. Vasserot. *Langlands reciprocity for affine quantum groups of type A_n* . Internat. Math. Res. Notices (IMRN) Vol. 1993:3 (1993), 67–85.
- [23] C.-J. Lai, D. K. Nakano, Z. Xiang. *q -Schur algebras corresponding to Hecke algebras of type B*. Transf. Groups **27** (2020), 983–1024.
- [24] G. Lusztig, *Introduction to Quantum Groups*, Progress in Math. **110**, Birkhäuser, 1993.
- [25] G. Lusztig. *Aperiodicity in quantum affine \mathfrak{gl}_n* . Asian J. Math. **3** (1999), 147–177.
- [26] G. I. Olshanski. *Quantized universal enveloping superalgebra of type Q and a super-extension of the Hecke algebra*. Letters Math. Phys. **24** (1992), 93–102.
- [27] A. N. Sergeev. *The tensor algebra of the identity representation as a module over the Lie superalgebras $GL(n, m)$ and $Q(n)$* . Math. USSR Sbornik **51** (1985) 419–427.
- [28] J. Wan, W. Wang. *Frobenius character formula and spin generic degrees for Hecke-Clifford algebra*, Proc. London Math. Soc. **106** (2013), 287–317.

J. D., SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW SOUTH WALES, SYDNEY NSW 2052, AUSTRALIA

Email address: j.du@unsw.edu.au

H. G., SCHOOL OF SCIENCE, HUZHOU UNIVERSITY, HUZHOU, CHINA

Email address: ghx@zjhu.edu.cn

Z. L., SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN 361005, CHINA

Email address: zhen-hua.li@qq.com

J. W., SCHOOL OF MATHEMATICS AND STATISTICS, BEIJING INSTITUTE OF TECHNOLOGY, BEIJING 100081, CHINA.

Email address: wjk302@hotmail.com