

Re-imagining the Future of Forest Management — An Age-Dependent Approach towards Harvesting

Shuyang Bian, Yuanyuan Xie, Flora Zhang

College of Arts and Sciences, Emory University

Primary Contacts: simon.bian@emory.edu, yuanyuan.xie@emory.edu

1 Abstract

Facing the drastic climate changes, current strategies for enhancing carbon dioxide stocks need to be thoroughly honed. While technological developments improve the quality of human life, they also severely damage the global carbon cycle with CO_2 emissions. Sole preservation is not a panacea, for it may well pack the woods, resulting in fire and ultimately carbon emissions. We attempt to search for a well-designed model of forests' carbon storage management, such that harvested carbon may yield economic return without compromising the delicate balance with nature.

To address the problem, we first built a carbon sequestration growth model driven by **growth rate dependency (GRDM)**. We abstracted the carbon cycling system into the process of photosynthesis, the humidity fluctuation, and the original storage of carbon in the trees. In the photosynthesis model, we considered various factors, including transition rate of absorption and organic matter production. In the humidity influence model, we defined the range of humidity that assisted plant growth and approximated the volume of carbon storage. The carbon storage in each tree is denoted based on the summation of per unit volume of carbon in each branch. Further, noticing the age-based differences in species ability to store carbon, we simulated the carbon harvest based on natural selection.

We also designed an **Economic Return Evaluation Model (EREM)** to estimate the optimal distribution of trees in the forest based on the utility function. Maximizing the util-

ity brought by the amount of carbon storage, we derived the equation for profit optimization with the constraints of total economic expenses allowed.

To assess its performance, we took an **object-oriented approach**, simulated an 8×8 ideal forest by placing instances of trees and plotted a time-dependent forest composition graph. With a time interval of 100 years, we recognize that the forest management strategies should be adjusted with a combination of three approaches: age-dependent, height-dependent, and density-dependent approach. Model-wise speaking, we find that selecting for trees' age would typically yield higher trees, a subsequent reduction in density, and smaller wildfires. Parameters adjustment show that when cutting trees whose ages are within 10% of the age at which wood loses its carbon sequestration ability, we will not compromise environmental protection for the sake of economic benefits. After proper normalization of climate and economic data, we also make predictions for 169 worldwide forest-covered countries. Our model further suggests high sensitivity and robustness with a similar trend of overall utility when environmental aridity or proportion of harvested woods are varied.

Finally, we apply the model to Georgia temperate deciduous forest, and we evaluate the carbon storage ability to adjust the Red Spruce based on available biological literature research. We recognize that while the model is preliminary in its failure to identify a diverse array of variables, it has encapsulated key features of idealized forests.

Keywords: Carbon Cycle, Utility Model,

2 Introduction

2.1 Problem Background

As organisms of the primary tropic level, green plants - with trees in forests contributing the most - fixate carbon dioxide by photosynthesis, thereby contributing to the circulation of global carbon flow and acting as a carbon sink. ^[1] Human activities and logging, nevertheless, serve as a carbon source that contributes to the increase of global CO_2 . ^[2]

While it seems natural that minimizing forest use to be feasible in preserving natural environments, studies reveal the possibility of striking a balance between economic benefits of carbon sequestration and maintaining forest vivacity. ^[3]

Furthermore, inter-species competition may arise when succession occurs, meaning that without proper forest management plans, prior fundamental species are replaced and converted into detritus, acting as a carbon source. ^[1] Environmental conditions including forest fires may also occur under high species density ^[4] more frequently, hence releasing carbon dioxide.

2.2 Problem Restatement

We need to create a carbon sequestration model to estimate the amount of carbon dioxide a forest will store and absorb over time. The model identifies the carbon sequestering management strategy to maximize forest's competence. After that, we develop a decision model to assist forest managers in determining the most efficient usage of a forest. The model develops a forest management strategy that considers many ways forests are valued including carbon sequestration, biodiversity aspects, and recreational uses, etc.

Considering the spectrum of management strategies, we need to design the decision-making model concerning tree types, geography, and lifespan, etc. We will test the model with various real-life forests to adjust the error

term of the model. With the advanced model, we predict the carbon dioxide a forest produces and absorbs over the next 100 years. With the optimum management plan and the prediction model, we suggest possible transitioning strategies for all forest users to add on to existing practices. Finally, as the forest is more and more open to the public, we support the statement of limiting the trees to optimal amount with the model.

2.3 Scenario-based Methodologies

1. Regarding net carbon dioxide absorption, we use ordinary differential equations to model the growth rate dependency (GRDM). As known ^[1], growth rate positively correlate with carbon retention. We consider three chronology-dependent aspects, including environmental temperature, overall humidity, and biological characteristics.
2. In light of the gain of net carbon dioxide absorption from the previous model, we then model rival factors, including human use of plants. Using the limitation of budget model (LBM), we consider aspects including human resource investment in forest management and commodity values of carbon sequestration.

Finally, a utility function (UF) is established to balance the gains and losses of both models. And different approaches were examined using the mathematical model.

3 Carbon Sequestration Models (CSM)

3.1 Assumptions

1. Concerning the chemicals involved in this model, we assume that no atoms other than Carbon are discussed for this mathematical model.
2. Concerning the biological systems involved in global carbon cycles, it is suggested that
(1) Unless otherwise amended, growth of trees are affected by no other factors than

age, volume, temperature and moisture availability.

(2) No other biomass other than trees in forests should be discussed. Further, trees in the forest rigidly do not expire or their expiration rates are also less than their inspiration of CO_2 .

(3) Human activities involve only logging and bringing carbon dioxide to the atmosphere, and its only economically favorable outcomes are the sequestered carbon from woods.

Other specific assumption, if necessary, will be mentioned and illustrated while we're building the models.

3. Concerning the mathematical model we use, we consider that:

(1) The tree can be modeled in a cylinder and whose volume would change in a predictable way;

(2) Once the trees were cut for carbon sequestration, the harvested carbon would no longer emit or absorb any other chemical, and also that once cut, the trees were considered dead. Seedlings would go from the exact location at which sawing happen.

3.2 Notations

The primary notations used in this paper are listed in the table in the last few pages.

If additional notations are necessary, they will be explained and illustrated while building models.

3.3 Carbon Sequestration Model: Growth Rate Dependency (GRDM)

The variation of carbon in forests over time is written as the total of the carbon stored and the carbon used for volume growth:

$$\frac{dC_{tot}}{dt} = F(H(t, T), h, rad, s, A) + P(C_a, T, h, rad, \rho) \quad (1)$$

Here, F is the gross carbon storage rate due to Humidity(H), height of the tree(h), radius to

the tree trunk(r), number of trees harvested(s), and the age of the tree(A). P denote the carbon assimilation due to gross photosynthesis eliminating the usage of foliage respiration. However, compare to the total carbon storage, the respiration of plants is almost negligible under our assumption.

$$H(t, T) = H(\sin(\omega t)), T(t) = \kappa \cos(B\omega t) + D \quad (2)$$

Among all species, there are non-resistant β species which takes $\frac{dC_{tot}}{dt} \propto H$, the rest $1 - \beta$ drought resistant thrives under low humidity such that $\frac{dC_{tot}}{dt} \propto \frac{1}{H}$. Notice that such periodic relationship only works when $T \leq 451^\circ F$, that is, when the temperature has not exceeded the point to set wood on fire. As shown below, temperature is also a function of time t. m represents the mass of the tree at time t, while χ , σ , γ , κ are constant parameters, and ϵ is the total error term.

$$H(t, T) = e^{-\frac{\chi}{2m}t}(\gamma \sin(\omega t)) + e^{-\frac{\chi}{2m}t} B \cos(\omega t) + \epsilon \quad (3)$$

$$\text{with } \omega = \frac{\sqrt{\chi^2 - 4m\kappa}}{2m} \quad (4)$$

$$\text{and } m = \rho V = \rho \left(\pi rad_1^2 h + \frac{2}{3} \pi rad_2^3 \right) \quad (5)$$

The above eq.3 would give a graph with oscillation behavior that tends to a set humidity H_0 asymptotically with the setting of constant ω (eq.4) and mass at time t m (eq.5).




Figure 1: Three-dimensional visualization of formula (3)




Figure 2: Three dimension of formula (3) asymptotic behavior

$$S_c = S \frac{\lambda}{\left(1 + \frac{dV}{dt}\right)}, \quad (6)$$

where $V \approx \pi \text{ rad}^2 h$.

Approximating the volume of tree trunks with the shape of cylinder, and the volume of canopy with a half of a sphere, we have the storage of carbon to the relation of the volume as eq.6 above.

$$\frac{dC_{tot}}{dt} = \frac{1}{s} \quad (7)$$

Harvesting trees would bring the carbon storage within each tree directly out while remain nearly neglectable damage to the environment. Such linear relationship is denoted by eq.7 above.

P is modeling photosynthesis considering the influential factors of atmospheric CO_2 , the growth rate of the tree $\frac{h'-h}{h'}$, the temperature of the growing environment $(T - T_{opt})$, and water potential.

$$P\alpha(1 - \exp C_a) \quad (8)$$

Photosynthesis is positively related to 1 minus the exponential of CO_2 concentration in the atmosphere C_a .

$$P\alpha \left(\Psi + \frac{2}{3} \pi \text{rad}^3 h \right) \quad (9)$$

The solar radiation absorbed is positively influenced by the depth of each canopy of tree,

photosynthesis tends to decrease with declining water potential due to the height of the trees.

$$P\alpha(1 - (T - T_{opt})^2) \quad (10)$$

Finally, in the assumed test environment of rain forests, photosynthesis's dependence on temperature is approximately a concave down parabola. Hence, P increase with the water potential minus the temperature difference to the optimal temperature (T_{opt}) squared.

Combining relations, we have from (8), (9), and (10), we have:

$$P = (1 - \exp(C_a))(\psi + 2/3\pi \text{rad}^3 h)(1 - (T - T_{opt})^2) \quad (11)$$

There are four defined stages for the age of trees: birth to sprout, sprout to youth, youth to mature, and mature to death. For all four stages, we have $\frac{dC_{tot}}{dt} \propto -\ln(A)$ for each tree. μ represents the constant that compensates for the errors.

Thus, we get

$$\frac{dC_{tot}}{dt} = -\ln(A) + \mu \quad (12)$$

3.4 Utility function based - Economic Return Evaluation Model (EREM)

Utility function $U(C_{tot}, E)$ is denoted by:

$$U(C_{tot}, E) = \theta \ln \left(\frac{1}{C_{tot}} \right) + (1 - \theta) \ln E + \tau \text{Simp} \ln(452 - T) \quad (13)$$

Where θ represents the number of drought resistant Keystone Species and $(1 - \theta)$ denotes the rest of the species which are non-resistant. The constant τ weights the influence of drought resistant species to utility.

With the restrain of $P_1 C_{tot} + P_2 E = I$, we apply the Lagrangian Method:

$$\begin{aligned} \max_{C_{tot}, E, \lambda} U(C_{tot}, E) &= \theta \ln \left(\frac{1}{C_{tot}} \right) \\ &+ (1 - \theta) \ln(E) + \tau \ln(T - 452) \\ &- \lambda(P_1 \cdot C_{tot} + P_2 \cdot E - I) \end{aligned}$$

Here, the inverse Simpson index has no effect on utility when temperature T goes over $451^\circ F$, the burning point of wood, and does not make sense when the temperature goes higher than the limit. That is, we also need the condition of $T \geq 451^\circ F$.

We take the First Order Condition (F.O.C) as below:

$$\frac{\partial U}{\partial C_{tot}} = \theta C_{tot} - \lambda P_1 = 0$$

$$\frac{\partial U}{\partial E} = \frac{1 - \theta}{E} - \lambda P_2 = 0$$

$$\frac{\partial U}{\partial \lambda} = P_1 C_{tot} + P_2 E - I = 0$$

from the first two equation, we have:

$$P_1 = \frac{\theta C_{tot} P_2 E}{1 - \theta}$$

then we rewrite our equation of $\frac{dU}{d\lambda}$ as:

$$\frac{\theta C_{tot} P_2 E}{1 - \theta} + P_2 E = I$$

Therefore,

$$P_1 = \frac{I}{(1 - \theta) C_{tot} E}$$

and

$$P_2 = \frac{I}{\theta C_{tot}^2 E^2}$$

4 Decision Model

4.1 Baseline Approach - A one-hundred year simulation

Before taking on any forest management plans, we attempt to determine the projected forest growth without human intervention. We devise a Java algorithm (see Attachment) that takes into account different characteristics of trees. As shown in the UML diagram below, trees were considered as a class and were inherited into the forest class. Different properties of trees, including age, water-sensitivity, growth rate, and aging, are considered. These tree instances are then placed in an 8×8 2-D array that would mimic a typical forest distribution.

Based on available literature and assumption, the dryness of the environment would occur in a time-dependent, trigonometric way; and under a given set threshold, this would result in all trees decimated in a 4×4 grid. Data is obtained with randomized initial status. Carbon dioxide absorption was calculated grid by grid. Colorization of data is achieved through normalized scaling as follows.



The below represents a time-based change of carbon dioxide with random forest fire, tree death, and tree growth over a period of 100 years. Two colored grid cells at the start of the simulated experiment and the end of it were included above the trendline of total carbon absorption. It is clear that after the disturbances by fire, forests need restoration. Evidently, with zero human intervention, forest fires would occur and yield no particular benefits.

87	67	78	99	90	81	121	115
67	99	81	115	110	102	96	94
78	81	115	102	77	26	86	98
99	115	102	94	104	98	74	115
90	110	77	104	105	74	110	96
81	102	26	98	74	112	58	27
121	96	86	74	110	58	113	96
115	94	98	115	96	27	96	107
115	26	78	112	68	87	61	112

(a) Carbon Sequestered in Randomized Initial-ized Forest

2	162	133	98	77	44	82	3
162	98	44	3	52	9	64	101
133	44	57	9	46	59	48	3
98	3	9	101	15	3	16	96
77	52	46	15	4	34	111	45
44	9	59	3	34	113	35	32
82	64	48	16	111	35	81	61
3	101	3	96	45	32	61	143
57	59	71	113	80	51	38	66

(b) Carbon Sequestered in Ten years with no Forest Management




Figure 3: Change curve of CO_2 and Economic Benefits with time

4.2 Age-dependent Approach

Literature has revealed that with increase in time, trees are increasingly susceptible to various environmental cues, including pathogen infection, interspecies competition, and environmental stress. Based on the model, we devise an age based tree logging approach to harvest wood while maintaining forest vitality. Notably, in the Java algorithm, we extend the trees in the aforementioned model to be a *Comparable* one, and modeled tree growth and carbon sequestered from each year by pushing the resulting ArrayList object onto the Stack. This approach ensured the constancy of comparison for different methods, and further ensured the randomness of initialized status.

We assume that consistently, every year regardless of the environmental status, the tree whose age is close to the imagined death age (1 year from the death expectancy) would be clearly identified and cut completely, thereby providing equal amounts of economic benefits as that of the CO_2 .

Matplotlib was then utilized to provide a plot of Benefits from both environmental CO_2 and wood collection. In comparison with the baseline approach, the scatter plot suggests that due to non-consideration of environmental conditions, continuous wood logging shows temporary decreased overall benefits based on utility function under arid conditions. However, in other times, not only was the forest still a net CO_2 sink, but also there had been continuous economic benefits.

We also confirm the regeneration ability of forests under appropriate logging under the age-selection approach. Color scale follows the same

methodology as described in the aforementioned baseline approach. As shown in the page that continues, we see a clear pattern in which a 55-year prediction suggests that tree growth are unencumbered by human logging.




Figure 4: Economic Benefits and Carbon Sequestered versus Baseline. In this figure, dots represent economic benefits, dashed red lines benefits obtained from age-specified tree logging activities, and solid line represents net carbon benefits with no cation.

4.3 Indications and Contraindications for Forest Management Plan - Sensitivity and Robustness Analysis

We adjust the following variables used in our algorithm, and obtain the following data for understanding of proper use of timber cutting as management strategies. Four different forest were evaluated based on existing literature on environmental science, and expected benefits as compared to baseline approach was summarized.

4.3.1 Effects of Harvest Age concerning Humidity and Harvest Percentage

We modeled three different hypothetical situations that relates to harvest age: (i) We want to understand whether this logging approach would be effective in different drought possibility, with step wise progression of 0.3. This

has its contemporary implications in that with the advent of global warming, extreme weathers have become more likely and would therefore impact the robustness of prior climate data's use on timber harvesting strategies.

The above trend lines yield several important conclusions which visualizes the following mathematical trends:

- (A) The higher the cutoff age for harvesting, the more likely it is going to yield more benefits than no clear cutting of the trees.
- (B) The humidity does not necessarily guarantee a lower cutoff age to be beneficial. We see that at wetness equivalent as 0.6, meaning that when environmental humidity is 0.6 out of a total of 1, the trees in a 3×3 grid would go on fire, which is when the threshold of maximum age types is actually the largest.
- (C) The model effectively pictures a similar trend line, albeit providing the critical information as to the non-linear dependency of age groups versus the aridity of the environment.

We also are interested in investigating variable factors of harvest percentage. Given the algorithm, our approach is to determine a random number and compare it to 1 – harvest percentage. Data was run on matplotlib and obtained as below. This test primarily serves as a sensitivity approach towards the values, and we recognize a decrease in the intersection between age of trees before its death and the percentage we count. This complements and corroborates the methodology depicted in part (i) of this section.

4.3.2 Effects of Wood Price and Carbon Dioxide Emission Price

As modeled, Net Primary Production (NPP) is essential to the cycling of the carbon, which is defined as the Gross Primary Production (GPP), that is, the overall amount of organic matter that all photosynthetic organisms can

consolidate from environmental CO_2 minus the respired CO_2 from them when daylight available. Historically, to increase the amount of NPP and presumably to decrease the value of NEE - defined as the negative of NPP - multiple nations have established a Kyoto Protocol, in which a global price for carbon is established based on country characteristics. We are interested in, therefore, implementing the utility function model by taking into account the different countries' forest coverage rate and past CO_2 emission. We consider annual data of Country's Forest coverage area, Country's forest-contributing GDP, alongside Climate Factors including temperature and humidity from open data sources. We normalize these data by ranking these values through Z score distribution, with the formula being the following, and μ being the average and σ being the standard deviation.

$$Z = (x - \mu) / \sigma$$




Figure 7: Geographical mapping of carbon sequestration model




Figure 8: Geographical mapping of carbon loss

5 Case Analysis

We choose Georgia state to be our region of interest. As a Southeastern state, Georgia boasts itself with the most forest coverage in the United

States^[7]. With the majority of the state in humid subtropical climate, Georgia becomes excellent in providing growth of forest region. Furthermore, Georgia's climate has been inevitably warming due to anthropogenic activities. What the region may act to save the continuous increase of temperature while not exceeding the budget on managing the at-risk situation has become crucial.

Further, one of its key cities, the metropolitan downtown area, Atlanta, needs economic incentive for supplying the demand of local people. Therefore, the state becomes our exemplary model for analyzing the effectiveness of modified forest management strategies. We attempt to demonstrate the model's performance of carbon storage model and economic return model.

In the following sections, we zoom into Red Spruce as the sample species and use our model to help decide the status quo of forest conditions in Appalachian area and strategies of how to improve the forest management. We visualize the carbon storage along with the economic returns to illustrate the cost and benefit for further actions suggested by our modifications.

5.1 Carbon Storage Benefits of Red Spruce in Appalachian Region

Picea rubens, commonly known as Red Spruce, is a native species to eastern North America. More specifically, its distinct adaptation to the ecosystem of southern Appalachian spruce-fir forest. It is specifically adapted to high altitude, medium amount of moisture and grows on well-drained sand land. Its peculiarity of being perennial provides a good fit to the assumption of our model to predict the actions in the near hundred years.

Such species has the average height of 5 to 54 meters and grow 10.6cm on a yearly basis. Its radius would therefore change accordingly. The average longevity range from 250 to more than 450 years. ^[7] This long time of growth results in a long term of sprout to mature stage when Red Spruce absorb the most carbon. We take data from literature, and estimate the function of Age and Carbon Dioxide Absorption to be

(illustrated by the figure below):

$$-0.008A^2 + \ln(A) + 8$$




Figure 9: Red Spruce Tree Age (in decades) vs. CO_2 absorbed

in which the visualization shows a prime age of carbon absorption at 19.75% of the total living period before death. At the point shown, the ability of carbon storage peak for Red Spruce for a rather short time and soon decrease until it reach the stage of late maturity and death.

While the carbon storage dumped with aging, the direct logging of Red Spruce may be subject to insect parasitism, acid rains, and regional warming. Instead of cutting down directly the aged Red Spruce, we should approximate with the model to find the ones that are already severely damaged by insects and can barely benefit the surrounding natural environment.

5.2 Economic Benefit of Red Spruce

With mature technology of wood processing, Red Spruce plays a crucial role in the production of paper pulp, acoustic instrument, and even food productions. Its wide varieties of usage promise a logged Red Spruce just after its death would directly decrease the forest density, raise the related local economic income, and remove the carbon storage to allow more species in their prime age to attribute to carbon storage.

With the current Carbon Volume and Price Ratio of 500\$ per tree, as well as the 20\$ per kilogram CO_2 of emitted based on Kyoto Protocol on Annex I countries, we recognize that carefully cutting down the aged Red Spruce which are impotent is critical in contributing in the natural may give more in the economic markets.

5.3 Conclusion of Red Spruce Region

Current Georgia management practices involve the use of controlled fire as a strategy. However, in such regard, carbon dioxide would be emitted from the burned trees, thereby creating a greater carbon source rather than flux. The Georgia Forestry Commission announces the probability of fire from a scale of 1 to 5, with the values closely mimick our simulation. Our strategy, in comparison with previous practices, aims at more specifically target at trees that need to be managed, while this may bring about an additional cost of human resources, the overall benefits are comparably huge. We take the above values into the mathematical equations and obtain the following strategies to stratify the forest environment. Additional data are obtained from the open data file from US Forest Service Department of Agriculture (2022), in which tree density is estimated based on available Growth-Removal-Mortality Model, which closely aligns with our computer simulation. Pandas module was imported in aiding the analysis of the problem.

Based on FIA data, our rough estimate concerning Georgia tree occupancy of the entire state is 15.4 billion. With a absorption rate of 21 kg / year, we have the total absorption without consideration of forest fire to be:

$$21 \times 15.4 \times 10^9 \times 10^{-3} = 3.3 \times 10^8 \text{ton}$$

We also obtain average forest fire affected acres from Fire and Smoke provided by the Augusta Chronicle, and assume all the carbon dioxide that the trees have absorbed over a lifetime was returned to the environment, and obtain the following two parts:

$$\begin{aligned} \text{Total } CO_2 &= \text{Wild} + \text{Human Induced} \\ &= 10 \times \frac{458861}{60} \times 3.3 \times (10^8) \\ &\quad \times \frac{1}{15.4 \times 10^9} \times 21 \times 100 \\ &= 344145 + 5601729 \\ &= 5.9 \times 10^6 \text{ton} \end{aligned}$$

This amount of CO_2 loss is equivalent to, as in

USD:

$$(5.9 \times 10^6) \times 20 = 1.2 \times 10^7 \text{USD}$$

Meanwhile, based on commodity price of carbon sequestered product, we estimate that with clear cut forests, the obtained carbon otherwise burned would yield a **gain** of the following economic benefits, equivalent as:

$$\begin{aligned} E &= \frac{458861}{60} \times 3.3 \times (10^8) \\ &\quad \times \frac{1}{15.4 \times 10^9} \times 21 \times 100 \times 20 \\ &= 6.9 \times 10^6 \text{USD} \end{aligned}$$

Given these data above, we have therefore quantitatively demonstrated an increase in the overall benefits as far as current tree management strategies are concerned.

6 Discussion: Strengths and Weaknesses

6.1 Strengths

- Considering various indicators in environmental aspects, we also combined possible factors of economic returns to make a more comprehensive and objective model.
- The simulation process may be replicated while changing the key conditions of average longevity of a specific forest, making our model effective and valuable under different environmental settings.
- The model of the carbon storage considers the influence of humidity, tree volume, tree growth rate, and the process of photosynthesis separately to fully identify the key linkages of the carbon cycling system.

6.2 Weaknesses

- With the more drastic climate changes which result in rise of sea level, acid rains, and extreme weathers, the benefit of our model won't last forever. Upgrades are needed for the new dominating factors to adjust continuously for even longer term of development.

- Some data is difficult to obtain and is dependent on rules of thumb encountered in biology, which may therefore cause sample errors.

7 Conclusion and Results

Logging as a Part of Long-term Forest Management

To address the ongoing discussion on the forest condition related to climate change, we conclude that it is unacceptable to pursue zero-logging for the so-called "eco-consciousness", as it may easily turn into extremism. Logging, when done appropriately, can prevent unfavorable situations, such as wildfires, which are caused by high density of trees in the forest. Hence, we investigated a considerable number of variables and found the relationship between proper logging and carbon dioxide.

Recent environmental protection strategies tend to favor the diminishing human activities on the natural environment as much as possible. In terms of active pollution and over-harvesting, such activities are not commendable, but we should also realize that human intervention can sometimes play a positive role. When done correctly, human intervention can help stabilize ecosystems. Providing underdeveloped areas with economic support to harvest carbon products—through creating tourism opportunities for rural areas—can also benefit communities and the environment at large. All things considered, an arbitrary closure of all-natural forests toward human beings is an impractical approach, and it is not economically feasible in the long run.

Species density plays a substantial role in advocating for proper forest management strategies. As known in previous literature, drought occurs cyclically. When using computations to model tree growth in arid areas within a simulated environment, data reveals several potential factors of harm: a high likelihood of forest fires, consequently loss of carbon storage, an imbalance between carbon release and absorption, and further diminishing of

keystone species with high volume of carbon storage (even in droughts). As a result, the anticipated species diversity would remain low despite the efforts of maintaining the secluded environment. In contrast, when implementing an age-based approach towards logging of trees, though the area suffers from a lack of water in arid times, this method effectively protects the trees against the possibility of forest fire. Furthermore, the obtained logs help ensure a decent income for the local economy and uplifts the community with natural resources, thereby balancing the loss of carbon in the process of sequestration.


In most situations, the area does not suffer from year-round drought, the surrounding industry does not depend actively on harvesting wood, nor does local water loss and soil erosion greatly rely on the number of trees. Thus, an appropriate amount of tree logging would protect the carbon storage in forest while yielding sustainable economic return.

Humans, by our presence upon this Earth, is already impacting every inch of the environment, be it directly or indirectly. And it is time for us to collectively be responsible for the fact that we owe it both to the Earth and to ourselves to draw up our next steps with sustainable development. Instead of blindly taking extreme approaches due to shallow look-ups, we should rather remain a humble and careful approach with thorough researches, such that humans can maintain harmony with the Earth and the countless species who also call this planet "home."


References

- [1] Stiling, Peter D. *Ecology: Global Insights & Investigations*. No. 577 St53e Ej. 1 024994. McGraw-Hill, 2012.
- [2] Alexiadis, Alessio. "Global Warming and Human Activity: A Model for Studying the Potential instability of the Carbon Dioxide/Temperature Feedback Mech-

- anism." *Ecological modelling*, vol.203, no.3, 2007, pp. 243-256.
- [3] Kishor, Nalin, et al. "Economic and Environmental Benefits of Eliminating Log Export Bans—The Case of Costa Rica." *World Economy*, VOL.27, no.4, 2004, PP. 609-624.
- [4] Song, Hark-Soo, and Sang-Hee Lee. "Effects of Wind and Tree Density on Forest Fire Patterns in a Mixed-Tree Species Forest." *Forest Science and Technology*, vol. 13, no.1,2017,pp. 9-16.
- [5] Hayat, Amaury, et al. "Modeling Tree Growth Taking into Account Carbon Source and Sink Limitations." *Frontiers in Plant Science*, vol. 8, 2017, p. 182.
- [6] Peng, Changhui, et al. "TRIPLEX: a Generic Hybrid Model for Predicting Forest Growth and Carbon and Nitrogen Dynamics." *Ecological Modelling*, vol. 153, no. 1, 2002, pp. 109–130.
- [7] Scott, D. "Comparison between Lodgepole Pine and Mountain Beech in Establishment and CO₂ Exchange." *New Zealand Journal of Botany*, vol. 8, no. 3, 1970, pp. 357–360.
- [8] Taylor, G. E. Jr, et al. "Carbon Dioxide Assimilation and Growth of Red Spruce (*Picea Rubens* Sarg.) Seedlings in Response to Ozone, Precipitation Chemistry, and Soil Type." *Oecologia*, vol. 70, no. 2, 1986, pp. 163–171.
- [9] Daigneault, Adam J, et al. "Optimal Forest Management with Carbon Sequestration Credits and Endogenous Fire Risk." *Land Economics*, vol. 86, no. 1, 2010, pp. 155–172.
- [10] Wagner, C. E. Van. "Age-Class Distribution and the Forest Fire Cycle." *Canadian Journal of Forest Research*, vol. 8, no. 2, 1978, pp. 220–227.
- [11] Sat Feb 19 17:00:12 CST 2022. Forest Inventory and Analysis Database, St. Paul, MN: U.S. Department of Agriculture, Forest Service, Northern Research Station. [Available only on internet: <http://apps.fs.fed.us/fiadb-downloads/datamart.html>]
- [12] Sat Feb 19 17:00:12 CST 2022. Key Fire Wildfire and Smoke Map - Data.augustachronicle .com. Fire and Smoke, <https://data.augustachronicle.com/fires/incident/7972/key-fire/>.



The structure of carbon simulation model TRIPLEX1.0 in forest (modified from Peng et al., 2002). The round cornered rectangles represent state variables while ovals represent core simulation processes, point lines represent controls, and solid lines represent flows of carbon (C), water, and the fluxes between the forest ecosystem and external environment.



Simulated Tree Growth Experiment. Squares are arranged based on time, in which those to the upper and left are ideal forests in earliest time, and those to the lower and right are ideal are forests in the latest time. It can be observed that while logging lowers the trees' density and lowers carbon dioxide absorption, they provide economic value and do not necessarily perform much inferior than the baseline approach.

Table 1: Notations

Symbol	Definition
C_{tot}	represents the overall carbon storage
t	represents the time passed since the origin, set the original time at $t(0)$
T	represents a fluctuating temperature of the given environment
$L(t, T)$	represents the overall humidity of the modeled natural due to time and temperature of surrounding environment
h	represents the height of the photosynthetic organism
rad	represents the radius of the tree
s	represents the amount of chopped down trees
S	represents carbon stored per unit volume of stem
A	represents the age of the tree
λ	represents the geometric factor
m	represents the mass of the tree at time t
ρ	represents the average density of carbon storage in each tree
ψ	represents the water potential
E	represents the economic return due to carbon storage
P_1	represents the cost of physical and human capital
P_2	represents the economic expenses on getting economic return
τ	represents the ratio of the number of drought resistant Keystone Species over the number of non-resistant common species
I	represents the total budget may be put in controlling carbon storage
$Simp$	represents the Inverse Simpson index which indicate higher species diversity and consequently a higher possibility of carbon fixation