EXACT SEPARATION OF EIGENVALUES OF LARGE DIMENSIONAL NONCENTRAL SAMPLE COVARIANCE MATRICES

By Zhidong Bai^{1,a} Jiang Hu^{1,b} Jack W. Silverstein^{2,d} , and Huanchao Zhou^{1,c}

¹KLASMOE and School of Mathematics and Statistics, Northeast Normal University, China, ^abaizd@nenu.edu.cn; ^bhuj156@nenu.edu.cn; ^czhouhc782@nenu.edu.cn

²Department of Mathematics, North Carolina State University, USA, ^djack@ncsu.edu

Let $\mathbf{B}_n = (\mathbf{R}_n + \frac{1}{\sqrt{n}}\mathbf{T}_n^{1/2}\mathbf{X}_n)(\mathbf{R}_n + \frac{1}{\sqrt{n}}\mathbf{T}_n^{1/2}\mathbf{X}_n)^*$ where \mathbf{X}_n is a $p \times n$ matrix with independent standardized random variables, \mathbf{R}_n is a $p \times n$ non-random matrix, representing the information, and \mathbf{T}_n is a $p \times p$ non-random nonnegative definite Hermitian matrix. Under some conditions on $\mathbf{R}_n \mathbf{R}_n^*$ and \mathbf{T}_n , it has been proved that for any closed interval outside the support of the limit spectral distribution, with probability one there will be no eigenvalues falling in this interval for all p sufficiently large. The purpose of this paper is to carry on with the study of the support of the limit spectral distribution, and we show that there is an exact separation phenomenon: with probability one, the proper number of eigenvalues lie on either side of these intervals.

1. Introduction. Let \mathbf{B}_n be a $p \times p$ matrix defined by

(1.1)
$$\mathbf{B}_n = \frac{1}{n} (\mathbf{R}_n + \mathbf{T}_n^{1/2} \mathbf{X}_n) (\mathbf{R}_n + \mathbf{T}_n^{1/2} \mathbf{X}_n)^*,$$

where $\mathbf{X}_n = (x_{ij})$ is a $p \times n$ matrix of independent and standardized random variables $(\mathbf{E}x_{ij} = 0, \mathbf{E}|x_{ij}|^2 = 1)$, \mathbf{R}_n is a $p \times n$ non-random matrix and \mathbf{T}_n is a $p \times p$ non-random non-negative definite Hermitian matrix. The model \mathbf{B}_n can be viewed as the non-central sample covariance matrix and it is also referred in the literature (see [1]) to as the general information-plus-noise type matrix where the information is contained in the matrix $(1/n)\mathbf{R}_n\mathbf{R}_n^*$ and the matrix $\mathbf{T}_n^{1/2}\mathbf{X}_n$ is the additive noise. In this paper, we carry on with the study of the location of eigenvalues of the matrix \mathbf{B}_n

In this paper, we carry on with the study of the location of eigenvalues of the matrix \mathbf{B}_n when p and n are large with the ratio $y_n = \frac{p}{n} \rightarrow y \in (0, 1)$. Before that, there have been some results on the non-central matrix \mathbf{B}_n . Under certain conditions, especially assuming \mathbf{T}_n and $\mathbf{R}_n \mathbf{R}_n^*$ are commutative, the result of the limiting spectral distribution (LSD) of \mathbf{B}_n has been studied in terms of the empirical spectral distribution (ESD) function $F^{\mathbf{B}_n}$ in [1]. And the analytic behavior of the LSD of the matrix \mathbf{B}_n have been studied in [2]. It is shown that the LSD has a continuous derivative away from zero, the derivative being analytic wherever it is positive, and the determination criterion is available for its support. The support of the probability distribution plays a fundamental role in the study of the spectrum of \mathbf{B}_n . In [3], it proved that for any closed interval outside the support of the LSD of \mathbf{B}_n , with probability one there will be no eigenvalues falling in this interval for all n sufficiently large. The aim of this paper is to prove that the proper number of eigenvalues lie on either side of these intervals. The analog of he property was first proposed in [5] called "exact separation".

Our work has been originally motivated by the popular sample covariance matrix $\mathbf{C}_n = \frac{1}{n} \mathbf{T}_n^{1/2} \mathbf{X}_n \mathbf{X}_n^* \mathbf{T}_n^{1/2}$ of *n* samples of the random vector $\mathbf{T}_n^{1/2} X_{\cdot 1}(X_{\cdot 1})$ denoting the *j*-th column

MSC2020 subject classifications: Primary 60E99, 26A46; secondary 62H99.

Keywords and phrases: Random matrix, LSD, Stieltjes transform, Information-plus-noise matrix.

of \mathbf{X}_n). Bai and Silverstein established the complete results about the almost sure localization of the eigenvalues of \mathbf{C}_n in the non-Gaussian case in [4, 5]. It is shown that for a closed interval [a, b] contained in an open interval (c, d) out the supports of LSD's F^{y_n, H_n} for almost all large n, the numbers of eigenvalues of \mathbf{C}_n falling on either side of [a, b] are exactly the same of eigenvalues of \mathbf{T}_n falling on the corresponding sides of the interval $J \subset \mathbb{R}^+$ which determines the interval [a, b].

Similar ideas were also developed for the information-plus-noise matrix

$$\mathbf{D}_n = \frac{1}{n} (\mathbf{R}_n + \sigma \mathbf{X}_n) (\mathbf{R}_n + \sigma \mathbf{X}_n)^*.$$

Using the same technique of [6, 7], Dozier and Silverstein established the LSD of D_n in [8] and derived the analytical behavior of the LSD of D_n in [9]. Such exact separation phenomenon was also exhibited for information-plus-noise matrix in [16], they proved that a gap in the spectrum of D_n which corresponds a gap in the spectrum of $\mathbf{R}_n \mathbf{R}_n^*$ splits the spectrum of $\mathbf{R}_n \mathbf{R}_n^*$ exactly as that of D_n dealing with independent Gaussian variables X_{ij} , and later, Capitaine extended the results to the framework of non-Gaussian information-plus-noise type matrices investigated in [11]. In addition, they addressed the behaviour of the largest eigenvalues of the information-plus-noise spiked model. And both results were obtained when the information matrix $\mathbf{R}_n \mathbf{R}_n^*$ is low rank.

The majority of known results are under the central condition, that is, the entries of X_n are zero mean (which can be extended to allowing the entries to have a common mean). In fact, the large non-central random matrices also have significant implications that may be considered as an extension of non-central Wishart matrices, an important random matrix in multivariate linear regression under a non-null hypothesis. We turn now to the aim of this paper, that is, to extend the above results of exact separation for the non-central random matrices B_n when $R_n R_n^*$ is full rank. Note that these results may also hold in the full rank case, not treated in this paper.

The rest sections of this paper are organized as follows. In section 2, we review some important results of the general information-plus-noise type matrix \mathbf{B}_n as well as some useful mathematical tools. Our main result and detailed proof are present in Section 3.

2. Preliminary Results. In this section, we recall some known results related to B_n . For the convenience of readers, we will expand here to illustrate. To establish exact separation, it is necessary to review properties of the LSD of of B_n . The best way in understanding both the LSD and analytic properties of the LSD is investigating the equations of the Stieltjes transforms of the LSD. Let F be any probability distribution, its Stieltjes transform is defined as

$$s_F(z) = \int \frac{1}{\lambda - z} \mathrm{d}F(\lambda), z \in \mathbb{C}^+ \equiv \{ z \in \mathbb{C} : \Im z > 0 \},\$$

and F can be obtained by the inversion formula

(2.2)
$$F(b) - F(a) = \frac{1}{\pi} \lim_{v \to 0^+} \int_a^b \Im s_F(x+iv) \mathrm{d}x,$$

where a, b are continuity points of F.

2.1. Convergence of the ESD of \mathbf{B}_n . The main result of the LSD can be summarized in the following lemma (see [1] for more details).

LEMMA 2.1 (Theorem 2.1 of [1]). Under the assumptions:

- (a) $\frac{p}{n} = y_n \rightarrow y$, as $\min\{p, n\} \rightarrow \infty$, and $y \in (0, 1]$.
- (b) The entries of $\mathbf{X}_n = (X_{ij})$ are independent and normalized, i.e., with zero mean and unit variance. Also, satisfy the Lindeberg condition: for any $\eta > 0$,

$$\frac{1}{\eta^2 pn} \sum_{i \le p; j \le n} E|X_{ij}|^2 I_{\{|X_{ij}| > \eta\sqrt{n}\}} \to 0.$$

- (c) $\mathbf{R}_n \mathbf{R}_n^*$ and \mathbf{T}_n are commutative.
- (d) As $\min\{p, n\} \to \infty$, the two-dimensional distribution function $H_n(u, t) = p^{-1} \sum_{i=1}^p I(u_i \le u, t_i \le t)$ converges weakly to a nonrandom limit distribution H(u, t), where u_i, t_i are the paired eigenvalues of $\frac{1}{n} \mathbf{R}_n \mathbf{R}_n^*$ and \mathbf{T}_n , respectively.

Then, with probability one, $F^{\mathbf{B}_n}$ converges in distribution to F, a nonrandom probability distribution function, whose Stieltjes transform $s = s_F(z)$ satisfies the equation system

(2.3)
$$\begin{cases} s = \int \frac{\mathrm{d}H(u,t)}{\frac{u}{1+yg} - (1+yst)z + t(1-y)}, \\ g = \int \frac{\mathrm{t}\mathrm{d}H(u,t)}{\frac{u}{1+yg} - (1+yst)z + t(1-y)}. \end{cases}$$

Moreover, for each $z \in \mathbb{C}^+$ *,* (s, g) *is the unique solution to* (2.3) *in* \mathbb{C}^+ *.*

For convenience, we also consider the convergence of the ESD of

$$\underline{\mathbf{B}}_n = \frac{1}{n} (\mathbf{R}_n + \mathbf{T}_n^{\frac{1}{2}} \mathbf{X}_n)^* (\mathbf{R}_n + \mathbf{T}_n^{\frac{1}{2}} \mathbf{X}_n)$$

The eigenvalues of the matrix $\underline{\mathbf{B}}_n$ are the same as those of the matrix \mathbf{B}_n except |n - p| zero eigenvalues. Let $F^{\underline{\mathbf{B}}_n}$ denote the almost sure limiting distribution function of the eigenvalues of \underline{B}_n . Therefore, it is easy to verify the following relations

$$F^{\underline{\mathbf{B}}_n} = \left(1 - \frac{p}{n}\right) I_{[0,\infty]} + \frac{p}{n} F^{\mathbf{B}_n},$$

where I_A denoting the indicator function of the set A. Then making a variable transformation

$$\underline{s}(z) = -\frac{1-y}{z} + ys(z),$$
$$\underline{g}(z) = -\frac{1}{z(1+yg(z))},$$

where $\underline{s}(z)$ is the Stieltjes transform of the LSD $F^{\underline{\mathbf{B}}_n}$ and $\underline{s}_n(z) = s_{F^{\underline{\mathbf{B}}_n}}(z)$ and $\underline{g}(z)$ is the limit of $\underline{g}_n = \frac{1}{p} \operatorname{tr} \mathbf{T}_n (\underline{\mathbf{B}}_n - z\mathbf{I})^{-1}$. Because Stieltjes transform is invertible, then the equations in (2.3) become

(2.4)
$$z = -\frac{1-y}{\underline{s}} - \frac{y}{\underline{s}} \int \frac{\mathrm{d}H(u,t)}{1+u\underline{g}(z)+t\underline{s}(z)},$$
$$z = -\frac{1}{\underline{g}} + y \int \frac{t\mathrm{d}H(u,t)}{1+u\underline{g}(z)+t\underline{s}(z)}.$$

Equating the two equations in (2.4), we get

$$-\frac{1-y}{\underline{s}} - \frac{y}{\underline{s}} \int \frac{\mathrm{d}H(u,t)}{1+u\underline{g}+t\underline{s}} = -\frac{1}{\underline{g}} + y \int \frac{t\mathrm{d}H(u,t)}{1+u\underline{g}+t\underline{s}}.$$

4

This is equivalent to

(2.5)
$$y\underline{g}^2 \int \frac{u\mathrm{d}H(u,t)}{1+u\underline{g}+t\underline{s}} + \underline{s} - \underline{g} = 0.$$

In addition, we consider several important quantities for the subsequent proof. Write $\underline{s} = \underline{s}_1 + i\underline{s}_2$, $\underline{g} = \underline{g}_1 + i\underline{g}_2$, $\underline{z}\underline{s} = (\underline{z}\underline{s})_1 + i(\underline{z}\underline{s})_2$, and $\underline{z}\underline{g} = (\underline{z}\underline{g})_1 + i(\underline{z}\underline{g})_2$. Fix $z = x + iv \in \mathbb{C}^+$. Multiplying by \underline{s} on both sides to (2.4) and comparing the imaginary part of the resulting equation, we obtain

$$(z\underline{s})_2 = cA_1\underline{g}_2 + cB_1\underline{s}_2,$$
$$v = \frac{\underline{g}_2}{|g|^2} - (cA_2\underline{g}_2 + cB_2\underline{s}_2)$$

where

(2.6)

$$A_{j} = \int \frac{ut^{j-1} dH(u,t)}{|1+u\underline{g}+t\underline{s}|^{2}}, j = 1, 2,$$
$$B_{j} = \int \frac{t^{j} dH(u,t)}{|1+u\underline{g}+t\underline{s}|^{2}}, j = 0, 1, 2.$$

Lemma 1 of [2] has been proved that all the four quantities A_j , B_j , j = 1, 2 are bounded.

2.2. Properties of the LSD of \mathbf{B}_n and characterization of its support. The behaviour of the Stieltjes transform of the LSD of \mathbf{B}_n around the real axis is fundamental to evaluate its support of F. It is necessary at this point to review some of the properties of F and s_F . From (2.3), much of the analytic behavior of F can be inferred in [2]. And we recall the main properties in the following lemma.

LEMMA 2.2 (Theorem 3 of [2]). Assume $c \leq 1$ and the conditions imposed in the limiting H in Lemma 2.1. Suppose $\underline{s}(x)$ is the solution to (2.4) for $x \neq 0$. Then $x \in \mathbb{R} \setminus \{0\}$, $\lim_{z \in \mathbb{C}^+ \to x} s_{\underline{F}}(z) \equiv \underline{s}(x)$ exists. The function \underline{s} is continuous on $\mathbb{R} \setminus \{0\}$, and F has a continuous derivative f on $\mathbb{R} \setminus \{0\}$ given by $f(x) = \frac{1}{\pi} \Im \underline{s}(x)$. Furthermore, if $\Im \underline{s}(x) > 0$ (f(x) > 0) for $x \in \mathbb{R}^+$, then the density f is analytic about x.

Moreover, most important for this paper is to understanding the support of F. For any probability distribution function G, let S_G and S_G^c denote the support of G and the complement of its support, respectively. By definition of support, we have $S_F \subset [0, \infty)$ and $S_H \subset [0, \infty) \times (0, \infty)$. It is shown in [2] that on any closed intervals outside the support of F, $s_F(x)$ exists and is increasing. Therefore, the inverse (2.2) can be used to identify these intervals. The following lemma has been proved in [2].

LEMMA 2.3 (Theorem 5 of [2]). Assume $c \leq 1$ and the conditions imposed on H in Lemma 2.1 hold. Let $x_0 \in S_F^c \cap \mathbb{R}^+$,

- (a) Then $\underline{s}(z) = \int (t-z)^{-1} d\underline{F}(t)$ is analytic in a neighborhood D_{x_0} of x_0 and there exists a co-solution $\underline{g}(z)$ which is also analytic in D_{x_0} . The triple $(x, \underline{s}(x), \underline{g}(x)), x \in D_{x_0} \cap \mathbb{R}^+$ is an extended solution to (2.4) with V < x.
- (b) For any support point (u, t) of H, $u\underline{g}(x_0) + t\underline{s}(x_0) \neq -1$.

On the other hand, if $x_0, \underline{s}_0, \underline{g}_0$, with $x_0 > 0$, form a real extended solution to (2.4) satisfying (b), then from (2.5), there exists a real analytic function $x = x(\underline{g})$, defined in an interval containing \underline{g}_0 which satisfy (2.4), and if $x'(\underline{g}_0) \neq 0$, then $x_0 \in S_F^c$.

As for whether F places any mass at 0, it is also shown in [2] that when $y \le 1$, the LSD F determined by (2.3) has no mass at zero.

2.3. No eigenvalues outside the support of the LSD of \mathbf{B}_n . The focus of [3] is on intervals $[a, b] \subset (c, d)$ lying outside the union of supports of $F^{c,H}$ and F^{c_n,H_n} , for all large n, it proves that there is no eigenvalues of \mathbf{B}_n falling in a, b], where $F^{c,H}$ is the LSD of \mathbf{B}_n and F^{c_n,H_n} is an $F^{c,H}$ with c and H replaced by $c_n = p/n$ and H_n .

Roughly speaking, when n is large, with probability one, there are no sample eigenvalues of \mathbf{B}_n falling into the limiting spectral gaps.

More exactly, the following result for \mathbf{B}_n is established in[3].

LEMMA 2.4 (Theorem 1.1 of [3]). Assume that

- (a) $[a,b] \subset (c,d) \subset S^c_{F^{y_n,H_n}}$, with c > 0 for all large n;
- (b) The matrix X_n is the p × n upper-left conner of the double array of random variables x_{ij} having means zero, variances one, second moments zero if complex and there is a random variable X with finite fourth moment such that for a constant K and for all x > 0

$$\frac{1}{p}\sum_{i=1}^{p}\mathbf{P}(|x_{ij}| > x) \le K\mathbf{P}(|X|) > x),$$

and

$$\frac{1}{n}\sum_{j=1}^{n} \mathbf{P}(|x_{ij}| > x) \le K\mathbf{P}(|X|) > x);$$

(c) There exists a positive function $\psi(x) \uparrow \infty$ as $x \to \infty$, and M > 0 such that

$$\max_{ij} \mathbf{E} |x_{ij}^2| \psi(|x_{ij}|) \le M;$$

- (d) n = n(p) with $y_n = p/n \rightarrow y > 0$ as $n \rightarrow \infty$;
- (e) For $n = 1, 2, ..., \mathbf{R}_n$ is a $p \times n$ nonrandom matrix with $\frac{1}{\sqrt{n}} \mathbf{R}_n$ uniformly bounded in special norm for all n;
- (f) The matrix \mathbf{T}_n is uniformly bounded in spectral norm and $\lambda_{-1} \leq K$ for some constant K, and is also commutative with $(1/n)\mathbf{R}_n\mathbf{R}_n^*$ and their joint spectral distribution $H_n(u,t)$ tends to a proper distribution H(u,t), where $\lambda_{-1} = \int t^{-1} dH(u,t)$.

Then, we have that

(2.7) $\mathbf{P}(no \ eigenvalues \ of \ \mathbf{B}_n \ appear \ in \ [a, b] \ for \ all \ large \ n) = 1.$

As mentioned in [3], assumptions (b)-(c) allow for the x_{ij} to depart from merely being i.i.d.. After suitable truncation, centralization, and scaling of the x_{ij} 's one can assume these variables to be uniformly bounded.

2.4. Other useful lemmas.

LEMMA 2.5. Assume that the entries of $\{x_{ij}\}$ are a double array of independent complex random variables with mean zero, variance σ^2 , and satisfy the assumptions (b) - (e) of Lemma 2.4. Let $\mathbf{X}_n = (x_{ij}; i \leq p, j \leq n)$ be the $p \times n$ matrix of the upper-left corner of the double array. Then, with probability one, we have

$$-2\sqrt{y}\sigma^{2} \leq \liminf_{n \to \infty} \lambda_{\min}(\mathbf{S}_{n} - \sigma^{2}(1+y)\mathbf{I}_{n}) \leq \liminf_{n \to \infty} \lambda_{\max}(\mathbf{S}_{n} - \sigma^{2}(1+y)\mathbf{I}_{n}) \leq 2\sqrt{y}\sigma^{2}$$

where $\mathbf{S}_{n} = n^{-1}\mathbf{X}_{n}\mathbf{X}_{n}^{*}$.

The proof of Lemma 2.5 is proved in [3] under the truncation and centralization, so the detailed proof of the lemma is omitted. For details, the readers are referred to [3].

LEMMA 2.6 (Theorem A.46. of [15]). Let A and B be two $n \times p$ complex matrices. Then,

$$\max_{k} |s_k(\mathbf{A}) - s_k(\mathbf{B})| \le \|\mathbf{A} - \mathbf{B}\|,$$

where $s_k(\mathbf{A})$ is the kth largest singular value of the matrix \mathbf{A} . If \mathbf{A} and \mathbf{B} are Hermitian, then the singular values can be replaced by eigenvalues; i.e.,

$$\max_{k} |\lambda_k(\mathbf{A}) - \lambda_k(\mathbf{B})| \le \|\mathbf{A} - \mathbf{B}\|$$

3. Main Result. In this section, we will study the exact separation problem of \mathbf{B}_n . For better characterize the exact separation, we equivalently reformulate the model \mathbf{B}_n as

(3.8)
$$\mathbf{B}_n = (\mathbf{R}_n + \frac{1}{\sqrt{n}} \mathbf{T}_n^{1/2} \mathbf{X}_n) (\mathbf{R}_n + \frac{1}{\sqrt{n}} \mathbf{T}_n^{1/2} \mathbf{X}_n)^*.$$

The change is only redefined $\frac{1}{\sqrt{n}}\mathbf{R}_n$ as \mathbf{R}_n . It is not difficult to see that the LSD of \mathbf{B}_n exists with probability one and its Stieltjes transform is given by (2.3) or equivalently by (2.4).

When $\mathbf{T}_n = \sigma^2 \mathbf{I}_n$, we recall the results of [12, 11] that with probability one, the number of eigenvalues of \mathbf{B}_n and $\mathbf{R}_n \mathbf{R}_n^*$ lying on one side of their respective intervals are identical for all large *n*. Then considering the model (3.8), we suppose that [a, b] is the interval defined in Lemma 2.4, and based on the conditions about \mathbf{R}_n , \mathbf{T}_n , define

$$h_j(x) = u_j g(x) + t_j \underline{s}(x), x \in [a, b],$$

where u_j , t_j are the paired eigenvalues of $\mathbf{R}_n \mathbf{R}_n^*$ and \mathbf{T}_n , respectively.

Then, we have the following theorem.

THEOREM 3.1. Under the assumptions of Lemma 2.4, with probability one, when n is large, for each $j \leq n$, the function $h_j(x)$ is either > -1 for all $x \in [a,b]$ or < -1 for all $x \in [a,b]$. Also, the number of eigenvalues of \mathbf{B}_n that are below a is exactly equal the number of $h_j(x)$ is below -1; and the number of eigenvalues of \mathbf{B}_n that are above b is exactly equal the number of $h_j(x)$ is above -1.

Before starting the proof of the theorem, we first truncate and renormalize the random variables of \mathbf{X}_n . We define $\hat{\mathbf{X}} = (\hat{X}_{ij})_{p \times n}$ and

$$\hat{X}_{ij} = X_{ij}I(|X_{ij}| < C) - \mathbf{E}X_{ij}I(|X_{ij}| < C)$$

for some constant C, and define

$$\hat{\mathbf{B}}_n = (\mathbf{R}_n + \frac{1}{\sqrt{n}} \mathbf{T}_n^{1/2} \hat{\mathbf{X}}) (\mathbf{R}_n + \frac{1}{\sqrt{n}} \mathbf{T}_n^{1/2} \hat{\mathbf{X}})^*.$$

By Lemma 2.5 and Corollary 7.3.8 of [17], we have

$$\max_{i \le p} |\lambda_i(\mathbf{B}_n) - \lambda_i(\hat{\mathbf{B}}_n)| \le \frac{1}{\sqrt{n}} \|\mathbf{T}_n^{1/2}(\mathbf{X} - \hat{\mathbf{X}})\| \le \sqrt{\|\mathbf{T}_n\|} (1 + \sqrt{y}) \sqrt{\mathbf{E}|X_{ij}^2|I(|X_{ij}| \ge C)},$$

which can be arbitrarily small when C is large.

Another truncations of \mathbf{R}_n and \mathbf{T}_n can be found in [2], which ensured that $\mathbf{R}_n \mathbf{R}_n^*$ and \mathbf{T}_n are of bounded norm for all n, and the details are not covered here.

3.1. Spectral Gap Dependence on y as Tending to 0. Now, using the same approach as in [5], we set up a series of new models such that $y = y_{\ell} = p/n_{\ell} \to 0$ as $\ell \to \infty$. To this end, we have to consider the model as

$$\mathbf{B}_{n,\ell} = (\mathbf{R}_{n_{\ell}} + n_{\ell}^{-1/2} \mathbf{T}_{n}^{1/2} \mathbf{X}_{n_{\ell}}) (\mathbf{R}_{n_{\ell}} + n_{\ell}^{-1/2} \mathbf{T}_{n}^{1/2} \mathbf{X}_{n_{\ell}})^{*},$$

where $n_{\ell} = n + \ell M$, and $M/n \to \tau > 0$ a small positive number. That means, for each model $\mathbf{B}_{n,\ell}$, we extend the $\mathbf{X}_{n_{\ell}}$ with M columns to obtain $\mathbf{X}_{n_{\ell+1}}$ and construct $\mathbf{B}_{n,\ell+1}$ accordingly. Also, we add 0 columns to $\mathbf{R}_{n_{\ell}}$ to match the order of $\mathbf{B}_{n_{\ell}}$, enabling feasible matrix addition.

Suppose that [a, b] represents the spectral gap for $\mathbf{B}_n = \mathbf{B}_{n,0}$. We will denote a series $[a_{\ell}, b_{\ell}]$ as the spectral gaps for $\mathbf{B}_{n,\ell}$ at each ℓ and demonstrate that the number of eigenvalues of $\mathbf{B}_{n,\ell}$ greater than b_{ℓ} (or smaller than a_{ℓ}) is the same as the number of eigenvalues of $\mathbf{B}_{n,\ell+1}$ greater than $b_{\ell+1}$ (or smaller than $a_{\ell+1}$, correspondingly). To achieve this, we need to define $[a_{\ell}, b_{\ell}]$ and prove that $b_{\ell} - a_{\ell}$ increases as $\ell \to \infty$.

We will consider solutions for (2.4) in the following way: for each $\underline{g} \in \mathbb{C}^+$ and $y \leq 1$, define \underline{s} by (2.5) which exists uniquely and belongs to \mathbb{C}^+ . Further, by the second equation in (2.4) we may find $z = z_{y,H(\underline{g},\underline{s},y)} \in \mathbb{C}^+$ and get a set of solutions to (2.4). According to Lemma 2.2, we can extend the solutions to (2.4) to the real axis. We remind the readers that for any real \underline{g} and y, there is a unique real \underline{s} solves (2.5) and consequently, there is a unique real z which solves the second equation of (2.4). The triple $(\underline{g}, \underline{s}, x)$ is a set of solutions to (2.4) but may not be a set of extended solutions to it, that means, the limit of $\underline{s}(z)$ when $z \xrightarrow{\mathbb{C}^+} x + 0i$ may not be $\underline{s}(x)$. According to Lemma 2.3, this limit holds if and only if $x \in S^c(F^{y,H})$. Therefore, we may use the positiveness of the derivative of x with respect to g to examine x is outside the support of $F^{y,H}$.

When the boundary point x of the support of $F^{y,H}$ is infinity, then by (2.4), the corresponding <u>s</u> and <u>g</u> are both 0. Hence, for smaller y, $(0, 0, \infty)$ is still a set of solution. Hence, ∞ is still a boundary point of the support of $F^{y,H}$. Consequently, when y decreases, the upper boundary of the spectral gap doesn't become smaller. Therefore, we only need to consider the case where the upper boundary of the spectral gap is finite.

Recall the definition of (2.6),

$$A_{j} = \int \frac{ut^{j-1} dH(u,t)}{|1+u\underline{g}+t\underline{s}|^{2}}, j = 1, 2,$$

$$B_{j} = \int \frac{t^{j} dH(u,t)}{|1+u\underline{g}+t\underline{s}|^{2}}, j = 0, 1, 2.$$

by conclusion (b) of Lemma 1 of [2], it has been proved that all the five quantities A_i , B_j are all bounded for all $z \in \mathbb{C}^+$. Hence, all x are in the complement of the support of $F^{y,H}$ by Fatou Lemma. That means $(1 + u\underline{g} + t\underline{s})^{-2}$ and its product with t, t^2, ut are integrable with respect to H, and thus the derivative of x with respect to \underline{g} exists and is continuous. Hence, the boundary points of spectral gaps are zero.

A spectral gap is an open interval (c, d) that is an interval of the complement of support of $F^{y,H}$. Suppose the end points of the spectral gap are both finite and denoted by x_j , j = 1, 2. Then, according to (2.5), we have

(3.9)
$$\begin{aligned} x_j &= -\frac{1}{\underline{g}_j} + y \int \frac{t \mathrm{d}H(u,t)}{1 + u\underline{g}_j + t\underline{s}_j},\\ 1 &= \frac{\underline{s}_j}{\underline{g}_j} + y \int \frac{u\underline{g}_j \mathrm{d}H(u,t)}{1 + u\underline{g}_j + t\underline{s}_j}. \end{aligned}$$

Note that

$$\frac{\partial x_j}{\partial \underline{g}_j} = \frac{1}{\underline{g}_j^2} - yA_2 - yB_2\underline{s}_j' = 0,$$

Hence

(3.10)
$$\frac{\mathrm{d}x_j}{\mathrm{d}y} = \frac{\partial x_j}{\partial y} + \frac{\partial x_j}{\partial \underline{g}} \frac{\mathrm{d}\underline{g}}{\mathrm{d}y} = \int \frac{t\mathrm{d}H(u,t)}{1 + u\underline{g}_j + t\underline{s}_j}.$$

Because $(\underline{g}_1, \underline{s}_1)$ and $(\underline{g}_2, \underline{s}_2)$ are in the same spectral gap, thus by the second conclusion of Lemma 2.3, for any support point (u, t) of H,

$$(1+u\underline{g}_1+t\underline{s}_1)(1+u\underline{g}_2+t\underline{s}_2)>0.$$

Therefore, we have

$$\frac{\mathrm{d}}{\mathrm{d}y}(x_1-x_2) = \int \frac{t[u(\underline{g}_2-\underline{g}_1)+t(\underline{s}_2-\underline{s}_1)]\mathrm{d}(u,t)}{(1+u\underline{g}_1+t\underline{s}_1)(1+u\underline{g}_2+t\underline{s}_2)}$$

By Lemma 2.3, both x and <u>s</u> are increasing functions of <u>g</u> outside the support of $F^{y,H}$ and hence $(x_1 - x_2)$ has the opposite signs as $(\underline{g}_2 - \underline{g}_1)$ and $(\underline{s}_2 - \underline{s}_1)$, hence

$$\frac{\mathrm{d}}{\mathrm{d}y}|x_1 - x_2| < 0,$$

thus, the size of spectral gap increases when y decreases.

Finally, we consider the case where $x_1 = \infty$ and x_2 finite. By (3.10), we have

$$\frac{\mathrm{d}x_2}{\mathrm{d}y} = \int \frac{t\mathrm{d}H(u,t)}{1+u\underline{g}_j + t\underline{s}_j}.$$

Since $(\underline{g}, \underline{s})$ is in the same spectral gap as (0, 0), we know that $1 + u\underline{g} + t\underline{s} > 0$ and hence $dx_2/dy < 0$ which implies that x_2 decreases as y decreases. Therefore, in all cases, when y decreases, the size of the spectral gap increases.

By Lemma 2.3, there are \underline{g}_a and \underline{g}_b in $\mathbb{R} \setminus \{0\}$ such that $a = x_{y,H}(\underline{g}_a)$ and $b = x_{y,H}(\underline{g}_b)$. Then, for any ℓ , we may select $[a_\ell, b_\ell]$ such that $b_\ell - a_\ell$ increases as ℓ increases. By Lemma 2.6, we have

(3.11)
$$\max_{k \le p} |\lambda_k^{1/2}(\mathbf{B}_{n,\ell+1}) - \lambda_k^{1/2}(\mathbf{B}_{n,\ell})| \le ||n_\ell^{-1/2} \mathbf{T}^{1/2} \mathbf{X}_{n_\ell} - n_{\ell+1}^{-1/2} \mathbf{T}^{1/2} \mathbf{X}_{n_{\ell+1}}||$$
$$\le (n_\ell^{-1/2} - n_{\ell+1}^{-1/2}) ||\mathbf{T}_n^{1/2} \mathbf{X}_{n_\ell}|| + n_{\ell+1} ||\mathbf{X}_\ell - \mathbf{X}_{n_{\ell+1}}|| \le K\tau$$

which can be made arbitrarily small and hence we may assume that $a_{\ell+1} \in (a_{\ell} - \frac{1}{2}(b_{\ell} - a_{\ell}), \frac{1}{2}(b_{\ell} + a_{\ell}))$ and $b_{\ell+1} \in (\frac{1}{2}(b_{\ell} + a_{\ell}), b_{\ell} + \frac{1}{2}(b_{\ell} - a_{\ell}),)$.

3.2. Proof of Theorem 3.1. In Lemma 2.4, it has proved that with probability one, when n is large, there are no eigenvalues of \mathbf{B}_n falling into the interval [a, b]. By the discussion in last section, there are intervals $[a_\ell, b_\ell]$ satisfies the conditions of Lemma 2.4 such that when n is large, there are no eigenvalues of $\mathbf{B}_{n,\ell}$ falling into the intervals $[a_\ell, b_\ell]$.

We claim that for each $j \leq p$, $\lambda_j(\mathbf{B}_{n,\ell})$ and $\lambda_j(\mathbf{B}_{n,\ell+1})$ are on the same sides of $[a_\ell, b_\ell]$ and $[a_{\ell+1}, b_{\ell+1}]$, respectively. If it is not the case, then we should have for some j and ℓ ,

$$|\lambda_j(\mathbf{B}_{n,\ell+1}) - \lambda_j(\mathbf{B}_{n,\ell})| \ge b_{\ell+1} \wedge b_\ell - a_{\ell+1} \vee a_\ell \ge \frac{1}{2}(b-a)$$

which contradicts (3.11).

Therefore, the numbers of eigenvalues of \mathbf{B}_n on each side of the interval [a, b] are exactly the same as those of eigenvalues of $\mathbf{B}_{n,\ell}$ on the corresponding sides of $[a_\ell, b_\ell]$ for any finitely ℓ . When $\ell \to \infty$, the matrix $\mathbf{B}_{n,\ell}$ tends to the non-random $\mathbf{B}_{n,\infty} = \mathbf{R}_n \mathbf{R}_n^* + \mathbf{T}_n$ whose paired eigenvalues are $\{u_{n,i} + t_{n,i}\}$.

The spectral gaps of the non-random $\mathbf{B}_{n,\infty}$ are the spacings of eigenvalues $u_{n,i} + t_{n,i}$. And the limiting spectral gap must be a subinterval of some limiting spacings. Suppose that the limiting spectral gap (a_{∞}, b_{∞}) is a subinterval of $(u_{i+1,n} + t_{i+1,n}, u_{i,n} + t_{i,n})$ which satisfies $d - c < u_{i,n} + t_{i,n} - (u_{i+1,n} + t_{i+1,n})$. By Lemma2.6, we have

(3.12)
$$\max_{k \le p} |\lambda_k^{1/2}(\mathbf{B}_{n,\ell}) - \lambda_k^{1/2}(\mathbf{B}_{n,\infty})| \le ||n_\ell^{-1/2} \mathbf{T}_n^{1/2} \mathbf{X}_{n_\ell} - \mathbf{T}_n^{1/2}||$$
$$\le K \max_{k \le p} ||s_k(n_{n_\ell}^{-1} \mathbf{X}_{n_\ell} \mathbf{B}_{n,\ell}^* - \mathbf{I}_p)|| \le 3K\sqrt{y_\ell},$$

with probability $o(n_{\ell}^{-\mu})$ for any fixed $\mu > 0$. Therefore, when ℓ is large so that $3K\sqrt{y_{\ell}} < \max(d-b,c-a)$, the numbers of eigenvalues of $\mathbf{B}_{n,\ell}$ larger than b_{ℓ} (smaller than a_{ℓ}) are the same of eigenvalues of $\mathbf{B}_{n,\infty}$ larger than $u_{i,n} + t_{i,n}$ (smaller than $u_{i+1,n} + t_{i+1,n}$).

Now, we complete the proof of Theorem 3.1. Note that for the non-random matrix $\mathbf{B}_{n,\infty}$, its eigenvalues are $\{u_{i,n} + t_{i,n}\}$ and $y_{\infty} = 0$ which implies that $\underline{s}(x) = \underline{g}(x) = -1/x$. Therefore, $h_j(x) = u_{j,n}\underline{g} + t_{j,n}\underline{s} = -(u_{j,n} + t_{j,n})/x$. Then, that $h_j(x)$ is less than -1 is equivalent to $u_{i,n} + t_{i,n}$ is larger than x. Hence, by the continuity and monotonicity, the number eigenvalues $u_{i,n} + t_{i,n}$ are larger than b_{∞} , is the same that $h_j(x)$ less than -1 and that the eigenvalues are smaller than a_{∞} is the same that $h_j(x) > -1$.

In viewing (3.11), we may select τ small enough so that

$$\max_{k \le p} |\lambda_k(\mathbf{B}_{n,\ell+1}) - \lambda_k(\mathbf{B}_{n,\ell})| < (b-a)/2.$$

Also, when $x \in [a_{\ell}, b_{\ell}] \cap [a_{\ell+1}, b_{\ell+1}]$,

$$|\underline{g}^{n,\ell+1}(x) - \underline{g}^{n,\ell}(x)| < \max_{y-x > (b-a)/2} (\underline{g}^{n,\ell}(y) - \underline{g}^{n,\ell}(x))$$

and

$$|\underline{s}^{n,\ell+1}(x) - \underline{s}^{n,\ell}(x)| < \max_{y-x > (b-a)/2} (\underline{s}^{n,\ell}(y) - \underline{s}^{n,\ell}(x)),$$

where $\underline{g}^{n,\ell}$ and $\underline{s}^{n,\ell}$ are the Stieltjes transforms \underline{g} and \underline{s} for the LSD of $\mathbf{B}_{n,\ell}$ respectively. Hence, if $h_i^{\ell}(x) > -1$ for all $x \in [a_{\ell}, b_{\ell}]$, then $h_j(a_{\ell}) > -1$ and thus

$$h_{j}^{\ell+1}(\frac{1}{2}(b_{\ell}+a_{\ell})) = u_{jn}\underline{g}^{\ell}(\frac{1}{2}(b_{\ell}+a_{\ell})) + t_{jn}\underline{s}_{\ell}(\frac{1}{2}(b_{\ell}+a_{\ell}))$$
$$\leq u_{jn}\underline{g}^{\ell}(a_{\ell}) + t_{jn}\underline{s}_{\ell}(a_{\ell}) = h_{j}^{\ell}(a_{\ell}) > -1.$$

Then, by the continuity and monotonicity of $h_j^{\ell+1}$, we conclude that $h_j^{\ell+1}(x) > -1$ for all $x \in [a_{\ell+1}, b_{\ell+1}]$.

Similarly, we can prove that $h_j^{\ell+1}(x) < -1$ for all $x \in [a_{\ell+1}, b_{\ell+1}]$ if $h_j^{\ell}(x) < -1$ for $x \in [a_{\ell}, b_{\ell}]$.

Finally, by (3.12), we can prove that for the last ℓ , $h_j^{\ell}(x)$ is larger (smaller) than -1 are the same as $\mathbf{B}_{n,\infty}$. Thus, Theorem 3.1 is proved.

Acknowledgments. The authors would like to thank the anonymous referee, the Associate Editor and the Editor for their invaluable and constructive comments. J. Hu was supported by NSFC (No. 12171078, 11971097) and National Key R & D Program of China (No. 2020YFA0714102). Z. D. Bai was partially supported by NSFC Grant 12171198 and 12271536 and Team Project of Jilin Provincial Department of Science and Technology (No.20210101147JC).

REFERENCES

- ZHOU, H. C., BAI, Z. D. and HU, J. (2022). The limiting spectral distribution of large dimensional general information-plus-noise type matrices. *Journal of Theoretical Probability*. 36 1203–1226.
- [2] ZHOU, H. C., BAI, Z. D., HU, J. and SILVERSTEIN, JACK. W. (2022). Analysis of the limiting spectral distribution of large dimensional general information-plus-noise type matrices. *Journal of Theoretical Probability*.
- [3] BAI, Z. D., HU, J. and SILVERSTEIN, JACK. W. ZHOU, H. C., (2023). No eigenvalues outside the support of the limiting spectral distribution of large dimensional noncentral sample covariance matrices. arXiv preprint arXiv:2303.12478.
- [4] BAI, Z. D. and SILVERSTEIN, J. W. (1998). No eigenvalues outside the support of the limiting spectral distribution of large-dimensional sample covariance matrices. *The Annals of Probability* 26 316–345.
- [5] BAI, Z. D. and SILVERSTEIN, J. W. (1999). Exact Separation of Eigenvalues of Large Dimensional Sample Covariance Matrices. *The Annals of Probability* 27 1536–1555.
- [6] SILVERSTEIN, J. W. (1995). Strong convergence of the empirical distribution of eigenvalues of large dimensional random matrices. *Journal of Multivariate Analysis* 55 331–339.
- [7] SILVERSTEIN, J. W. and CHOI (1995). Analysis of the limiting spectral distribution of large dimensional random matrices. *Journal of Multivariate Analysis* 54 295–309.
- [8] DOZIER, R. B. and SILVERSTEIN, J. W. (2007). On the empirical distribution of eigenvalues of large dimensional information-plus-noise-type matrices. *Journal of Multivariate Analysis* 98 678–694.
- [9] DOZIER, R. B. and SILVERSTEIN, J. W. (2007). Analysis of the limiting spectral distribution of large dimensional information-plus-noise type matrices. *Journal of Multivariate Analysis* 98 1099–1122.
- [10] LOUBATON, P. and VALLET, P. (2011). Almost sure localization of the eigenvalues in a Gaussian information plus noise model. Application to the spiked models. *Electronic Journal of Probability* 16 1934-1959.
- [11] CAPITAINE, M. (2014). Exact separation phenomenon for the eigenvalues of large information-plus-noise type matrices, and an application to spiked models. *Indiana University Mathematics Journal* 63 1875– 1910.
- [12] COUILLET, R., DEBBAH, M. and SILVERSTEIN, J. W. (2011). A deterministic equivalent for the analysis of correlated MIMO multiple access channels. *IEEE Transactions on Information Theory* 57 3493–3514.
- BURKHOLDER, D. L. (1973). Distribution function inequalities for martingales. *the Annals of Probability* 1 19–42.
- [14] BAI, Z. D. and SILVERSTEIN, J. W. (2012). No eigenvalues outside the support of the limiting spectral distribution of information-plus-noise type matrices. *Random Matrices: Theory and Applications* 1 1150004.
- [15] BAI, Z. D and SILVERSTEIN, J. W. (2010). Spectral analysis of large dimensional random matrices 20. Springer.
- [16] LOUBATON, P. and VALLET, P. (2011). Almost sure localization of the eigenvalues in a Gaussian information plus noise model. Application to the spiked models. *Electronic Journal of Probability* 16 1934– 1959.
- [17] HOM, R. A. and JOHNSON, C. R. (1985). Matrix analysis. Cambridge University Express 455.