

# FORMAL CATEGORY THEORY IN $\infty$ -EQUIPMENTS I

JACO RUIT

**ABSTRACT.** We introduce a generalization of proarrow equipments to the  $\infty$ -categorical setting, termed  $\infty$ -equipments. These equipments take the form of particular double  $\infty$ -categories that support a notion of internal formal higher category theory. We will explore several examples of  $\infty$ -equipments. In particular, we will study the prototypical example given by the  $\infty$ -equipment of  $\infty$ -categories, and construct  $\infty$ -equipments of  $\infty$ -categories internal to general  $\infty$ -toposes as well. The final objective of this article is to study the fundamental notions of formal category theory within an arbitrary  $\infty$ -equipment. To this end, we not only follow the classical approach but also expand upon it.

## CONTENTS

1. Introduction	1
2. Two-dimensional $\infty$ -category theory	6
3. Companions, conjoinants and $\infty$ -equipments	16
4. Example: the $\infty$ -equipment of $\infty$ -categories	26
5. Example: the $\infty$ -equipment of internal categories	34
6. First notions of formal category theory	44
References	55

## 1. INTRODUCTION

In the field of *formal category theory*, the goal is to distill the fundamental concepts that enable one to make sense of category theory internal to a 2-category; the archetype being the ambient 2-category of (small) categories. Ideally, one would then obtain an umbrella framework that encompasses enriched category theory, indexed category theory, internal category theory, and fibered category theory. In this paper, we will aim to extend the methods from this field to the higher categorical setting.

Partly, this is already being pursued by Riehl and Verity in their successful *synthetic or formal* treatise of the foundations of  $\infty$ -category theory. Nowadays, one can find a complete introduction to  $\infty$ -category theory from this perspective [RV22]. The formal (higher) category theory one will encounter in this paper, is, in contrast, not designed to study  $\infty$ -categories *ex nihilo*. Instead, we will use the strength of existing  $\infty$ - and  $(\infty, 2)$ -categorical methods to set up formal category theories internal to an  $(\infty, 2)$ -category. This can then be specialized to enriched, internal, indexed, and fibered  $\infty$ -category theory, and combined flavors of these.

One of the motivations for pursuing this project lies in giving new (conceptual) foundations of the theory of indexed (or parametrized)  $\infty$ -categories, particularly, equivariant  $\infty$ -categories. Indexed  $\infty$ -category theory was introduced by Barwick, Dotto, Glasman, Nardin and Shah in [BDG<sup>+</sup>16]. Its fundamental notions were further studied by Shah [Sha18]. Another far-developed approach to indexed  $\infty$ -category theory, more generally, to internal  $\infty$ -category theory, is due to Martini and Wolf [MW21]. In the sequel to this paper [Rui], we will develop machinery to formally retrieve the fundamental results of [Sha18] using the theory of this article.

The starting insight in the development of formal category theory is the following: merely having the structure of an ambient 2-category or more generally, an ambient  $(\infty, 2)$ -category, is inadequate to define all desirable (internal) categorical notions. Such a structure would give a good notion of (internal) adjunctions. However, the ambient  $(\infty, 2)$ -category is incapable of expressing the expected defining properties of fully faithful arrows or pointwise Kan extensions in general, to name just two examples.

Let us illustrate this with a first naive approach. If  $\mathcal{X}$  is a  $(\infty, 2)$ -category<sup>1</sup> (or, for illustrative purposes, a 2-category), one could attempt to define a notion of fully faithful arrows as follows. Namely, say that an arrow  $f : x \rightarrow y$  in  $\mathcal{X}$  is fully faithful if and only if for every object  $z \in \mathcal{X}$ , the induced pushforward functor

$$f_* : \mathcal{X}(z, x) \rightarrow \mathcal{X}(z, y)$$

is fully faithful on mapping  $\infty$ -categories. This recovers the notion of fully faithful functors if  $\mathcal{X}$  is the  $(\infty, 2)$ -category of (small)  $\infty$ -categories. However, one can check that, in general, this definition does not recover the correct classical notion of enriched fully faithful functors if  $\mathcal{X}$  is the ambient 2-category of enriched categories over an arbitrary monoidal category.

One could also attempt to define a notion of Kan extensions in  $\mathcal{X}$ . Namely, one could say that the datum of a 2-cell

$$\begin{array}{ccc} i & \xrightarrow{f} & x \\ w \downarrow & \Downarrow & \nearrow g \\ & & j \end{array}$$

in  $\mathcal{X}$  exhibits  $g$  as the (naive) left Kan extension of  $f$  if and only if the 2-cell witnesses the presheaf

$$\mathrm{Map}_{\mathcal{X}(i,x)}(f, w^*(-)) : \mathcal{X}(j, x) \rightarrow \mathcal{S}$$

to be representable by  $g$ . Here  $\mathcal{S}$  denotes the  $\infty$ -category of *spaces* or  $\infty$ -*groupoids*. This definition already fails to capture the notion of pointwise left Kan extensions if  $\mathcal{X}$  is the ambient 2-category of categories. In this case, every pointwise left Kan extension is a naive left Kan extension, but the converse is not necessarily true (see [Cam15] for some counterexamples). Moreover, this naive definition does not come with a method to compute such extensions.

These illustrative examples bear witness to the fact that having just an ambient 2-category or  $(\infty, 2)$ -category, does not suffice to define basic categorical notions. Consequently, one will need structures that are better suited for this

<sup>1</sup>We will make precise which model we use in Section 2.

purpose. A first successful approach to formal category theory is due to the Australian school of category theorists. In [SW78], Street and Walters axiomatize so-called *Yoneda structures*. These are structures on 2-categories that allow one to do category theory in these 2-categories. The name of these structures already reflects what obstructs the development of formal category theory in a plain ambient 2-category. Namely, a 2-category does not come with natural notions of presheaves and representable presheaves. These notions are crucial to encode desirable universal properties of pointwise Kan extensions.

Since the work of Street and Walters, there have been more approaches to formal category theory that saw the light of day. The famous approach of Wood is of importance to us. In [Woo82], Wood builds on the ideas of Street and Walters and introduces *proarrow equipments* (sometimes, simply called *equipments*). We are interested in Verity’s equivalent *double categorical* perspective on these equipments [Ver92] (for the relation to Wood’s original formulation, see [Ver92, Definition 1.2.4] and [Shu08, Appendix C]).

To a great extent, this paper is concerned with generalizing the theory of proarrow equipments to the  $\infty$ -categorical context. We make use of the formalism of *double  $\infty$ -categories*. A double  $\infty$ -category is a 2-dimensional  $\infty$ -categorical construct that has (a space of) objects and two types of (spaces of) arrows (1-cells) between objects: *vertical arrows* and *horizontal arrows*. Moreover, a double  $\infty$ -category contains (a space of) 2-cells. A 2-cell may be pictured as a square

$$\begin{array}{ccc} a & \longrightarrow & b \\ \downarrow & \Downarrow & \downarrow \\ c & \longrightarrow & d, \end{array}$$

where the arrows pointing vertically (resp. horizontally) are vertical (resp. horizontal) arrows of the double  $\infty$ -category. Informally, a double  $\infty$ -category may be viewed as the data of these spaces of objects, 1- and 2-cells, together with compatible and coherent composition laws for these cells. Double  $\infty$ -categories were first considered by Haugseng [Hau13] and have also been studied by Moser in [Mos20].

The  $\infty$ -categorical generalizations of proarrow equipments we will introduce, take the form of double  $\infty$ -categories which satisfy some additional properties. The definition of these  $\infty$ -equipments is more or less analogous to Verity’s formulation of classical proarrow equipments. The horizontal arrows of an  $\infty$ -equipment are called *proarrows*, which are used to encode desirable universal properties. Proarrows can be viewed as an abstraction of presheaves and they may be represented by vertical arrows. The vertical arrows often play the role of generalized functors. Imprecisely, an  $\infty$ -equipment is required to admit two specific ways in which each vertical arrow can represent a proarrow: either as a *companion* or *conjoint*. Among other things, this formalism of proarrows and their representability ultimately allows one to express the desired universal property of pointwise Kan extensions in [Subsection 6.3](#).

**Layout of this paper.** Our first substantive section, [Section 2](#), contains the basic theory of (2-fold, 2-uple) Segal spaces, double  $\infty$ -categories,  $(\infty, 2)$ -categories, and introduces terminology that is used throughout the article.

In [Section 3](#), we will define  $\infty$ -equipments, the  $\infty$ -categorical analog of proarrow equipments. Ultimately, we see that the classical definition may be (more or less) repeated in the higher context. This crucially builds on joint work in progress with Ozornova and Rovelli [[ORR](#)] where companions and conjoiners in double  $\infty$ -categories are studied. Because this is still unfinished, we will refer to the material in this future work as hypotheses.

The prototypical example of an  $\infty$ -equipment will be given by the equipment

$$\mathbb{C}at_{\infty}$$

of  $\infty$ -categories ([Construction 4.9](#)). Its objects are given by  $\infty$ -categories, the (vertical) arrows by usual functors, and the (horizontal) proarrows by *profunctors*. A profunctor  $\mathcal{C} \rightarrow \mathcal{D}$  between  $\infty$ -categories is given by a functor  $\mathcal{D}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{S}$ . Equivalently, such a profunctor may be viewed as a *correspondence* or *two-sided discrete fibration*, similar to the classical case. As one might expect, the construction of this  $\infty$ -equipment is more involved than the construction of its classical counterpart. One possible construction using *Conduché fibrations* (or *exponentiable fibrations*) of this equipment is implicit in the work of Ayala and Francis [[AF20](#)]. We will recall this construction in [Section 4](#). Moreover, we will demonstrate that the underlying vertical  $(\infty, 2)$ -category of  $\mathbb{C}at_{\infty}$  is equivalent to a known model for the  $(\infty, 2)$ -category of  $\infty$ -categories.

The construction of  $\mathbb{C}at_{\infty}$  can be generalized. Let  $\mathcal{E}$  be an  $\infty$ -topos. Using closely related ideas of Haugseng [[Hau17](#)], we will construct an  $\infty$ -equipment

$$\mathbb{C}at_{\infty}(\mathcal{E})$$

of  *$\infty$ -categories internal to  $\mathcal{E}$*  in [Section 5](#). We expect that its associated formal category theory recovers the internal  $\infty$ -category theory developed by Rasekh [[Ras18](#)], and by Martini and Wolf [[MW21](#)].

These formal category theories associated with  $\infty$ -equipments are studied in [Section 6](#), and it closely resembles the classical story. Most of the material that we will cover here, is inspired, or, is often completely analogous to the classical counterparts. Albeit in a different set-up, some of the theory dates back to Street and Walters [[SW78](#)], Guitart [[Gui80](#)], and Wood [[Woo82](#)]. In the language of classical double categorical equipments, parts of the formal category theory introduced here can be found in the work of Shulman [[Shu13](#)]. Classical *virtual* equipments and their associated formal category play a role in the synthetic approach to  $\infty$ -category theory by Riehl and Verity as well [[RV22](#), Chapter 9]. This work has also been a source of inspiration. In [Section 6](#), we will define notions of *fully faithful arrows*, *weighted (co)limits*, *pointwise Kan extensions* and *exact squares* in an  $\infty$ -equipment, and study their properties. Under an extra *pointedness* assumption on the equipments, one can also define notions of *conical (co)limits*, and *initial* and *final* arrows. Under yet another extra (strong) assumption on the equipment, one may even retrieve an internal form of Quillen's theorem A, purely formally.

**Preview of part II.** We will further shape the theory of  $\infty$ -equipments in the sequel of this paper [[Rui](#)]. It is a general principle that one may express more aspects of formal category theory when an  $\infty$ -equipment has more properties or is endowed with more structure. In the next article, we will consider *closed monoidal  $\infty$ -equipments*. This extra structure allows one to study the existence

of internal arrows of Kan extension (i.e. adjoints to restriction arrows). To this end, we will also need to set up a theory of *lax double functors* and *(lax) double adjunctions*. We will exhibit a supply of interesting examples. For instance, a geometric morphism between  $\infty$ -toposes will induce a lax adjunction between their associated  $\infty$ -equipments of internal  $\infty$ -categories (constructed in [Section 5](#)). We study properties of (lax) double functors and adjunctions between  $\infty$ -equipments that allow one to relate the formal category theories associated with the corresponding  $\infty$ -equipments to each other. The highlight of [\[Rui\]](#) is the study and construction of *cotensor  $\infty$ -equipments*. This construction allows one to construct new  $\infty$ -equipments out of existing ones by cotensoring with  $\infty$ -categories. Applying this to the archetypical  $\infty$ -equipment  $\text{Cat}_\infty$  from [Section 4](#), one obtains  $\infty$ -equipments for indexed  $\infty$ -category theories that coincide with the  $\infty$ -equipments for category theory internal to corresponding presheaf  $\infty$ -toposes of [Section 5](#). We study abstract properties of cotensor  $\infty$ -equipments that allow one to obtain the basic theory of indexed  $\infty$ -category theory established by Shah [\[Sha18\]](#). The expectation is that this approach is fruitful in the enriched setting as well, yielding similar results for indexed enriched  $\infty$ -category theory.

**Food for thought.** This article invites for further research. We list here some directions that would be worth pursuing:

- (1) We may apply the material in this article to the  $\infty$ -equipment of  $\infty$ -categories enriched over a (suitably) cocomplete monoidal  $\infty$ -category. A double  $\infty$ -category of enriched  $\infty$ -categories, functors and bimodules over a compatibly cocomplete monoidal  $\infty$ -category  $\mathcal{V}$  has already been constructed by Haugseng [\[Hau16\]](#). One can show that this is an  $\infty$ -equipment. In case that  $\mathcal{V}$  is given by the  $\infty$ -category  $\mathcal{S}$  of spaces, one should be able to show that this recovers  $\text{Cat}_\infty$  using [Proposition 5.18](#). Now, interpreting the formal category theory of [Section 6](#) inside the  $\infty$ -equipment of enriched  $\infty$ -categories, one obtains the basic notions for enriched  $\infty$ -category theory. It would be worthwhile to flesh out the details. Moreover, one could compare the theory of weighted colimits that this yields to the one of Hinich [\[Hin21\]](#).
- (2) One may try to compare the different models of enriched  $\infty$ -categories by not only setting up equivalences between their ambient  $\infty$ -categories, but also setting up equivalences between their  $\infty$ -equipments.
- (3) It would be good to understand how far the formalization of category theory may be pushed. For instance, would one be able to set up a formal theory of *doctrines*? Or may one formalize *Yoneda embeddings*? Probably, more structure or properties are needed of the  $\infty$ -equipments one deals with. An approach for the classical case can be found in [\[MT08\]](#).
- (4) One should be able to repeat Shulman's theory of *monoids in equipments* and *monoidal fibrations* [\[Shu08, Section 11, 12\]](#) in the higher setting. How would one go about this?
- (5) The  $\infty$ -cosmology developed by Riehl and Verity [\[RV22\]](#) is related to the theory of  $\infty$ -equipments. One should be able to associate an  $\infty$ -equipment to an  $\infty$ -cosmos so that its 'homotopy equipment' recovers

the equipment studied in [RV22, Chapter 9]. One could try to make this precise.

**Acknowledgements.** I am indebted to Lennart Meier, my PhD supervisor. His invaluable comments, insightful discussions, review of drafts, and above all, his support by granting me the freedom to explore my research interests in the realm of higher formal category theory, have been of great importance in shaping this article. During my visit to the Johns Hopkins University in the spring of 2023, I had many fruitful conversations with Emily Riehl regarding the contents of this paper. I would like to thank her once again for these sessions, her sharing of expertise, and the hospitality extended to me during my stay in Baltimore. During the writing of this paper, the author was funded by the Dutch Research Council (NWO) through the grant “The interplay of orientations and symmetry”, grant no. OCENW.KLEIN.364.

## 2. TWO-DIMENSIONAL $\infty$ -CATEGORY THEORY

We commence by recalling the theory of (*2-uple, 2-fold*) *Segal spaces* and *double  $\infty$ -categories*. Moreover, we will discuss Barwick’s model of  $(\infty, 2)$ -categories via *complete 2-fold Segal spaces* [Bar05].

**2.1. Double Segal spaces.** To efficiently introduce these concepts, we use the notion of *categorical objects*:

**Definition 2.1.** Let  $\mathcal{C}$  be an  $\infty$ -category with finite limits. Then a *categorical object*  $X$  in  $\mathcal{C}$  is a simplicial object  $X : \Delta^{\text{op}} \rightarrow \mathcal{C}$  such that the so-called *Segal map*

$$X([n]) \rightarrow X(\{0 \leq 1\}) \times_{X(\{1\})} \cdots \times_{X(\{n-1\})} X(\{n-1 \leq n\})$$

is an equivalence for all  $n \geq 0$ . The full subcategory of  $\text{Fun}(\Delta^{\text{op}}, \mathcal{C})$  spanned by the categorical objects in  $\mathcal{C}$  is denoted by  $\text{Cat}(\mathcal{C})$ .

**Definition 2.2.** The following types of categorical objects have a special name:

- (1) A *Segal space* (resp. *Segal set*) is a categorical object in the  $\infty$ -category  $\mathcal{S}$  of spaces (resp. the category  $\text{Set}$  of sets). We write  $\text{Seg}(\mathcal{S}) := \text{Cat}(\mathcal{S})$  and  $\text{Seg}(\text{Set}) := \text{Cat}(\text{Set})$  for, respectively, the  $\infty$ -category of Segal spaces and the category of Segal sets.
- (2) In turn, a *2-uple Segal space* or *double Segal space* is categorical object in  $\text{Cat}(\mathcal{S})$ . We will write  $\text{Cat}^2(\mathcal{S}) := \text{Cat}(\text{Cat}(\mathcal{S})) \subset \text{Fun}(\Delta^{\text{op}, \times 2}, \mathcal{S})$  for the  $\infty$ -category of 2-uple Segal spaces.
- (3) A *double  $\infty$ -category* is a categorical object in  $\text{Cat}_{\infty}$ . We write  $\text{DblCat}_{\infty} := \text{Cat}(\text{Cat}_{\infty})$  for the  $\infty$ -category of double  $\infty$ -categories.

Unwinding the definition, we see that a 2-uple Segal space  $\mathcal{P}$  contains a space  $\mathcal{P}_{0,0}$  of objects (0-cells), a space  $\mathcal{P}_{0,1}$  of *vertical arrows* (1-cells), a space  $\mathcal{P}_{1,0}$  of *horizontal arrows* (1-cells), and a space  $\mathcal{P}_{1,1}$  of 2-cells. The 2-cells and the compatible arrows of  $\mathcal{P}$  that have the same direction may be composed in a mutually compatible and coherently associative fashion. A 2-cell may be pictured as a square

$$\begin{array}{ccc} a & \longrightarrow & b \\ \downarrow & \Downarrow & \downarrow \\ c & \longrightarrow & d, \end{array}$$

where the arrows pointing vertically (resp. horizontally) are vertical (resp. horizontal) morphisms of  $\mathcal{P}$ . When dealing with 2-cells in a 2-uple Segal space, we will usually leave out the notation of the arrow ( $\Rightarrow$ ) that fills the square.

We may view double  $\infty$ -categories as 2-uple Segal spaces with extra completeness conditions. Namely, the famous result of Joyal and Tierney [JT07] asserts that the restriction of the Yoneda embedding

$$\mathrm{Cat}_\infty \rightarrow \mathrm{PSh}(\Delta) : \mathcal{C} \mapsto ([n] \mapsto \mathrm{Map}_{\mathrm{Cat}_\infty}([n], \mathcal{C}))$$

is fully faithful, and its essential image consists of those Segal spaces which are *complete*:

**Definition 2.3.** A Segal space  $X$  is called *complete* if the degeneracy map

$$\mathrm{Map}_{\mathrm{Seg}(\mathcal{S})}([0], X) \rightarrow \mathrm{Map}_{\mathrm{Seg}(\mathcal{S})}(J, X),$$

where  $J$  is the Segal set that sits in the pushout square

$$\begin{array}{ccc} [1]^{\sqcup 2} & \xrightarrow{\{0 \leq 2\} \sqcup \{1 \leq 3\}} & [3] \\ \downarrow & & \downarrow \\ [0]^{\sqcup 2} & \longrightarrow & J \end{array}$$

of Segal sets (this is a pushout square in  $\mathrm{Seg}(\mathcal{S})$  as well).

**Remark 2.4.** Viewed as a (discrete) Segal space,  $J$  is equivalent to the image of the free isomorphism under the composite

$$\mathrm{Cat} \rightarrow \mathrm{Seg}(\mathrm{Set}) \rightarrow \mathrm{Seg}(\mathcal{S}),$$

where the first arrow is the inclusion of categories inside Segal sets, and the second arrow is the inclusion of discrete Segal spaces, i.e. it is induced by the inclusion  $\mathrm{Set} \rightarrow \mathcal{S}$ . This fact was shown by Rezk [Rez01, Proposition 11.1].

In light of this result of Joyal and Tierney, we obtain an induced fully faithful functor

$$\mathrm{DblCat}_\infty \rightarrow \mathrm{Cat}^2(\mathcal{S}) : \mathcal{P} \mapsto (([n], [m]) \mapsto \mathrm{Map}_{\mathrm{Cat}_\infty}([m], \mathcal{P}_n))$$

whose essential image consists of all those 2-uple Segal spaces  $X$  so that  $X_{n, \bullet}$  is complete for all  $n$ . This is readily checked to be an inclusion of a reflective subcategory.

**Remark 2.5.** The perspective of double Segal spaces turns out to be useful to prove certain results of combinatorial nature, see for instance [Rui22]. In this article, we introduced a category of *2-dimensional pasting shapes*, denoted by  $\mathrm{Shape}^2$ . These pasting shapes encode particular arrangements in which cells of 2-uple Segal spaces may be composed. This is done using the language of *2-boxes*. A 2-box is simply a pair  $(x, y) \in \mathbb{N}^{\times 2} \times \mathbb{N}^{\times 2}$  of a top-left and bottom-right corner coordinate, so that  $x_a \leq y_a$  for all  $a$ . A 2-dimensional pasting shape is then simply a collection of 2-boxes subject to some conditions (see [Rui22, Definition 3.2] for a precise definition). An illustrative example can be found in [Rui22, Example 3.4]. The category  $\mathrm{Shape}^2$  comes, as expected, with a fully faithful nerve functor

$$\mathrm{Shape}^2 \rightarrow \mathrm{Cat}^2(\mathcal{S}),$$

which factors through  $\text{DblCat}_\infty$  (Construction 3.10 of loc. cit.). The main result [Rui22, Theorem 3.42] stated that this nerve functor preserved particular kinds of colimits.

We will use the terminology of (2-dimensional) pasting shapes in a few instances throughout this article, and, motivated by the fact that the nerve functor is fully faithful, we will make no distinction between the notation of pasting shapes and their images under the nerve functor. Moreover, we will write

$$[a, b] \in \text{Shape}^2$$

for the pasting shape that was previously denoted by  $\square[a, b]$  in [Rui22]. As a double Segal space, it is precisely the image of  $([a], [b])$  under the Yoneda embedding  $\Delta^{\times 2} \rightarrow \text{Cat}^2(\mathcal{S})$ .

In some proofs, we will also use the following notation. If  $S, T$  are subsets of  $\mathbb{N}$ , we will write

$$[S, T] \in \text{Shape}^2$$

for the largest 2-dimensional pasting shape whose vertices are in  $S \times T$ , so that  $[a, b] = [\{0, \dots, a\}, \{0, \dots, b\}]$ . Moreover, if  $\mathcal{P}$  is a 2-uple Segal space and  $I$  is a pasting shape, then we write

$$\mathcal{P}(I) := \text{Map}_{\text{Cat}^2(\mathcal{S})}(I, \mathcal{P}).$$

There are few involutive operations on  $\text{Cat}^2(\mathcal{S})$ :

**Definition 2.6.** Let  $\mathcal{P}$  be a 2-uple Segal space. Then the *horizontal opposite*  $\mathcal{P}^{\text{hop}}$  is given by the composite

$$\mathcal{P}^{\text{hop}} : \Delta^{\text{op}} \times \Delta^{\text{op}} \xrightarrow{(-)^{\text{op}} \times \text{id}} \Delta^{\text{op}} \times \Delta^{\text{op}} \xrightarrow{\mathcal{P}} \mathcal{S},$$

similarly, its *vertical opposite* is given by

$$\mathcal{P}^{\text{vop}} : \Delta^{\text{op}} \times \Delta^{\text{op}} \xrightarrow{\text{id} \times (-)^{\text{op}}} \Delta^{\text{op}} \times \Delta^{\text{op}} \xrightarrow{\mathcal{P}} \mathcal{S}.$$

Note that if  $\mathcal{P}$  is a double  $\infty$ -category, then the vertical and horizontal opposites are again double  $\infty$ -categories. Finally, we also have the *transpose*  $\mathcal{P}^t$  of  $\mathcal{P}$ , given level-wise by

$$\mathcal{P}_{n,m}^t := \mathcal{P}_{m,n}.$$

**Example 2.7.** In Section 4, we will recall the construction of the double  $\infty$ -category

$$\mathbb{C}\text{at}_\infty$$

of  $\infty$ -categories. Its objects are given by  $\infty$ -categories and the vertical arrows correspond to functors between  $\infty$ -categories. A horizontal arrow  $\mathcal{C} \rightarrow \mathcal{D}$  between  $\infty$ -categories is given by a functor  $\mathcal{D}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{S}$ . This is also called a *profunctor*. A 2-cell in  $\mathbb{C}\text{at}_\infty$

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{F} & \mathcal{B} \\ f \downarrow & & \downarrow g \\ \mathcal{C} & \xrightarrow{G} & \mathcal{D} \end{array}$$



corresponds to a natural transformation

$$\begin{array}{ccc} \mathcal{B}^{\text{op}} \times \mathcal{A} & \xrightarrow{F} & \mathcal{S}. \\ g^{\text{op}} \times f \downarrow & \Downarrow & \nearrow G \\ \mathcal{D}^{\text{op}} \times \mathcal{C} & & \end{array}$$

The double  $\infty$ -category  $\text{Cat}_\infty$  is a special example of a double  $\infty$ -category

$$\text{Cat}_\infty(\mathcal{E}),$$

which one can associate to any  $\infty$ -topos  $\mathcal{E}$  (see [Section 5](#)), and whose objects are given by *complete categorical objects* in  $\mathcal{E}$ .

**Example 2.8.** Let  $\mathcal{C}$  be a monoidal  $\infty$ -category. We further assume that  $\mathcal{C}$  admits all geometric realizations (i.e.  $\mathcal{C}$  has all colimits of  $\Delta^{\text{op}}$ -shaped diagrams) and that the tensor product  $\otimes : \mathcal{C}^{\times 2} \rightarrow \mathcal{C}$  of  $\mathcal{C}$  preserves these geometric realizations in each variable. Then Lurie constructs in [[Lur17](#), Remark 4.4.3.11] an associated double  $\infty$ -category,

$$\text{Alg}(\mathcal{C})$$

called the *Morita double  $\infty$ -category*. Its objects are given by algebras in  $\mathcal{C}$ , the vertical arrows are algebra morphisms, and a horizontal arrow  $F : A \rightarrow B$  between algebras corresponds to an  $(A, B)$ -bimodule  $F$ . A 2-cell

$$\begin{array}{ccc} A & \xrightarrow{F} & B \\ f \downarrow & & \downarrow g \\ C & \xrightarrow{G} & D \end{array}$$

in this double  $\infty$ -category corresponds to a map of  $(A, B)$ -bimodules  $F \rightarrow A \otimes_C G \otimes_D B$ .

**Example 2.9.** Let  $\mathcal{C}$  be an  $\infty$ -category with finite products. In [[Hau18](#)], Haugseng constructs the double  $\infty$ -category

$$\text{Span}(\mathcal{C})$$

of spans in  $\mathcal{C}$ . The objects of this double  $\infty$ -category are given by the objects of  $\mathcal{C}$ , and the vertical arrows are the arrows in  $\mathcal{C}$ . A horizontal arrow  $a \rightarrow b$  in this double  $\infty$ -category is given by a span

$$a \leftarrow x \rightarrow b.$$

A 2-cell in  $\text{Span}(\mathcal{C})$

$$\begin{array}{ccc} a & \longrightarrow & b \\ \downarrow & & \downarrow \\ c & \longrightarrow & d \end{array}$$

corresponds to a commutative diagram

$$\begin{array}{ccccc} a & \longleftarrow & x & \longrightarrow & b \\ \downarrow & & \downarrow & & \downarrow \\ c & \longleftarrow & y & \longrightarrow & d \end{array}$$

in  $\mathcal{C}$ .

**2.2. 2-fold Segal spaces.** Recall that 2-uple Segal spaces may be used as a model for  $(\infty, 2)$ -categories. This is a generalization of Rezk's model of complete Segal spaces for  $\infty$ -categories that is due to Barwick [Bar05]. Let us first discuss the notion of *2-fold Segal spaces* and its relation to 2-uple Segal spaces:

**Definition 2.10.** A 2-uple Segal space  $\mathcal{X}$  is called a *2-fold Segal space* if the restriction  $\mathcal{X}_{0,\bullet} : \Delta^{\text{op}} \rightarrow \mathcal{S}$  is essentially constant. We write  $\text{Seg}^2(\mathcal{S})$  for the full subcategory of  $\text{Cat}^2(\mathcal{S})$  spanned by those 2-fold Segal spaces.

The canonical inclusion

$$(-)_h : \text{Seg}^2(\mathcal{S}) \rightarrow \text{Cat}^2(\mathcal{S})$$

of the full subcategory of 2-fold Segal spaces is called the *horizontal inclusion*. By [Hau18, Proposition 4.12], this admits a right adjoint, called the *horizontal fragment functor*

$$\text{Hor}(-) : \text{Cat}^2(\mathcal{S}) \rightarrow \text{Seg}^2(\mathcal{S}).$$

If  $\mathcal{P}$  is a 2-fold Segal space, then the horizontal fragment of  $\mathcal{P}$  is given level-wise by the pullback squares

$$\begin{array}{ccc} \text{Hor}(\mathcal{P})_{n,m} & \longrightarrow & \mathcal{P}_{n,m} \\ \downarrow & & \downarrow \\ \mathcal{P}_{0,0}^{\times(n+1)} & \longrightarrow & \mathcal{P}_{0,m}^{\times(n+1)}, \end{array}$$

see also [Hau18, Remark 4.13]. Similarly, we have an adjunction between the *vertical inclusion* and *fragment functor*

$$(-)_v : \text{Seg}^2(\mathcal{S}) \rightleftarrows \text{Cat}^2(\mathcal{S}) : \text{Vert}(-),$$

where

$$\mathcal{X}_v := (\mathcal{X}_h^t)^{\text{hop}}, \quad \text{and} \quad \text{Vert}(\mathcal{P}) := \text{Hor}((\mathcal{P}^{\text{hop}})^t).$$

The fragment functors are compatible with opposites. We have the following special names for vertical and horizontal opposites of 2-fold Segal spaces:

**Definition 2.11.** If  $\mathcal{X}$  is a 2-fold Segal space, then its *1-opposite* and *2-opposite* are defined by respectively  $\mathcal{X}^{1\text{-op}} := \mathcal{X}^{\text{hop}}$  and  $\mathcal{X}^{2\text{-op}} := \mathcal{X}^{\text{vop}}$ .

It readily follows that  $\text{Hor}(-)$  and  $\text{Vert}(-)$  are compatible with taking opposites as follows: there are equivalences  $\text{Hor}(\mathcal{P}^{\text{hop}}) \simeq \text{Hor}(\mathcal{P})^{1\text{-op}}$ ,  $\text{Hor}(\mathcal{P}^{\text{vop}}) \simeq \text{Hor}(\mathcal{P})^{2\text{-op}}$ ,  $\text{Vert}(\mathcal{P}^{\text{hop}}) \simeq \text{Vert}(\mathcal{P})^{2\text{-op}}$  and  $\text{Vert}(\mathcal{P}^{\text{vop}}) \simeq \text{Vert}(\mathcal{P})^{1\text{-op}}$  that are natural in  $\mathcal{P}$ .

**Remark 2.12.** With these observations in place, note that  $\text{Vert}(\mathcal{P}) \simeq \text{Hor}(\mathcal{P}^t)^{2\text{-op}}$  and  $\mathcal{X}_v \simeq (\mathcal{X}^{2\text{-op}})_h^t$ . One could also have defined the vertical inclusion and vertical fragments so that there do not appear 2-opposites in these descriptions. However, the convention that we have introduced here, seems more natural regarding the material that will follow.

To model  $(\infty, 2)$ -categories via 2-fold Segal spaces, we need to invert the 2-categorical Dwyer-Kan equivalences in  $\text{Seg}^2(\mathcal{S})$ . This requires the introduction of a completeness condition:

**Definition 2.13.** A 2-fold Segal space  $\mathcal{X}$  is said to be *complete* if it has the property that  $\mathcal{X}_{1,\bullet}$  and  $\mathcal{X}_{\bullet,0}$  are complete Segal spaces.

The full subcategory  $\text{Seg}^2(\mathcal{S})$  spanned by the complete 2-fold Segal spaces satisfies the axioms of [BSP21] for an  $\infty$ -category of  $(\infty, 2)$ -categories. Consequently, we will write

$$\text{Cat}_{(\infty,2)} \subset \text{Seg}^2(\mathcal{S})$$


for this full subcategory, and also refer to complete 2-fold Segal spaces as  $(\infty, 2)$ -categories. As expected, Moser showed [Mos20, Theorem B] that there is a fully faithful (or *nerve*) functor

$$\text{Cat}_2 \rightarrow \text{Cat}_{(\infty,2)},$$

where  $\text{Cat}_2$  is the underlying  $\infty$ -category (actually, weak  $(2, 1)$ -category) of the *canonical model structure on 2-categories* [Lac02]. We will make no distinction between 2-categories and the notation of images under this nerve functor. In practice, we will just be interested in the restriction of the nerve to *gaunt 2-categories*, i.e. 2-categories without isomorphisms. If  $G$  is a *gaunt 2-category*, i.e. a 2-category without isomorphisms, then, it is discrete as a complete 2-fold Segal space so that

$$G_{n,m} = \text{Hom}([n; m, \dots, m], G),$$

the *set* of functors from the *globular 2-category*



(ranging  $n, m$ , this naturally forms a double cosimplicial object) to  $G$ .

A few comparisons of the model of complete 2-fold Segal spaces for  $(\infty, 2)$ -categories and some other known models for  $(\infty, 2)$ -categories may be found in [Lur09b], [BR13] and [BR20].

**Definition 2.14.** Let  $\mathcal{X}$  be a 2-fold Segal space. Then the underlying Segal space  $\mathcal{X}^{(1)}$  of  $\mathcal{X}$  is given by  $\mathcal{X}_{\bullet,0}$ . If  $x$  and  $y$  are objects of  $\mathcal{X}$ , then the *mapping Segal space*  $\mathcal{X}(x, y)$  is defined by the pullback square

$$\begin{array}{ccc} \mathcal{X}(x, y) & \longrightarrow & \mathcal{X}_{1,\bullet} \\ \downarrow & & \downarrow \\ \{(x, y)\} & \longrightarrow & \mathcal{X}_{0,\bullet}^{\times 2}. \end{array}$$

A few of the important examples of 2-fold Segal spaces that we will encounter in this paper are not complete, but only locally in the following sense:

**Definition 2.15.** A 2-fold Segal space  $\mathcal{X}$  is called *locally complete* if the mapping Segal spaces  $\mathcal{X}(x, y)$  are complete for all  $x, y \in \mathcal{X}$ .

We will recall some results from [Hau18]:

**Proposition 2.16.** *Suppose that  $\mathcal{X}$  is a 2-fold Segal space. Then  $\mathcal{X}$  is locally complete if and only if  $\mathcal{X}_{n,\bullet}$  is complete for all  $n$ .*

*Proof.* Note that complete Segal spaces are closed under limits. In light of the Segal conditions, we thus have to show that  $\mathcal{X}$  is locally complete if and only if  $\mathcal{X}_{1,\bullet}$  is complete. If  $\mathcal{X}_{1,\bullet}$  is complete, then the mapping Segal spaces of  $\mathcal{X}$  fit in pullback squares of complete Segal spaces, hence they are again complete. The converse assertion follows from applying [Hau18, Lemma 7.25] to the source-target map  $\mathcal{X}_{1,\bullet} \rightarrow \mathcal{X}_{0,\bullet}^2$ .  $\square$

**Corollary 2.17.** *A 2-fold Segal space  $\mathcal{X}$  is complete if and only if it is locally complete and  $\mathcal{X}^{(1)}$  is complete.*

*Proof.* This follows from [Hau18, Lemma 7.21].  $\square$

**Definition 2.18.** We say that a 2-uple Segal space is (locally) horizontally complete or (locally) vertically complete if  $\text{Hor}(\mathcal{P})$ , respectively,  $\text{Vert}(\mathcal{P})$ , is (locally) complete.

Throughout this article, vertically complete double  $\infty$ -categories will play an important role. We will write

$$\text{DblCat}_\infty^v \subset \text{DblCat}_\infty \subset \text{Cat}^2(\mathcal{S})$$

for the full subcategory of  $\text{DblCat}_\infty$  spanned by the vertically complete double  $\infty$ -categories. Since the vertical fragment functor  $\text{Vert}(-)$  is a right adjoint, this is a reflective subcategory.

**Remark 2.19.** If  $\mathcal{P}$  is a double  $\infty$ -category, then  $\mathcal{P}$  is always locally horizontally complete. Namely, we have a pullback square of Segal spaces

$$\begin{array}{ccc} \text{Hor}(\mathcal{P})(x, y) & \longrightarrow & \mathcal{P}_1 \\ \downarrow & & \downarrow \\ \{(x, y)\} & \longrightarrow & \mathcal{P}_0^{\times 2}, \end{array}$$

for  $x, y \in \mathcal{P}$ , where the right arrow is the source-target projection. However,  $\mathcal{P}$  is not necessarily horizontally complete. Similarly, the vertical fragment of  $\mathcal{P}$  is not complete in general. If  $x, y \in \mathcal{P}$ , then the mapping Segal space is given level-wise by

$$\text{Vert}(\mathcal{P})(x, y)_n \simeq \text{Map}_{\mathcal{P}_n}([n] \rightarrow [0]^*x, [n] \rightarrow [0]^*y).$$

Note that  $\text{Vert}(\mathcal{P})^{(1)} \simeq \mathcal{P}_0$  as Segal spaces, hence  $\mathcal{P}$  is vertically complete if and only if  $\mathcal{P}$  is locally vertically complete.

Finally, there is also a local notion for fully faithfulness:

**Definition 2.20.** A map  $F : \mathcal{X} \rightarrow \mathcal{Y}$  between 2-fold Segal spaces is called *locally fully faithful* if and only if the induced map  $\mathcal{X}(x, y) \rightarrow \mathcal{Y}(Fx, Fy)$  is fully faithful for all  $x, y \in \mathcal{X}$ .

Using the language of pasting shapes (see Remark 2.5), we can give a geometric characterization of locally fully faithful maps. In what follows, we will write  $\text{tr}_{\leq 1} I$  for the 1-truncation of a 2-dimensional pasting shape  $I$  as defined in [Rui22, Definition 3.13].

**Proposition 2.21.** *Let  $F : \mathcal{X} \rightarrow \mathcal{Y}$  be a map between 2-fold Segal spaces. Then the following statements are equivalent:*

- (1)  $F$  is locally fully faithful,
- (2) for any 2-dimensional locally composable pasting shape  $I$ , the induced map

$$\mathcal{X}(I) \rightarrow \mathcal{Y}(I) \times_{\mathcal{Y}(\mathrm{tr}_{\leq 1} I)} \mathcal{X}(\mathrm{tr}_{\leq 1} I)$$

is an equivalence.

*Proof.* It follows from the pasting theorem of [Rui22] that (2) holds if and only if (2) holds for  $I = [1, 1]$ . The latter assertion is equivalent to the statement that the commutative square

$$\begin{array}{ccc} \mathcal{X}_{1,1} & \longrightarrow & \mathcal{Y}_{1,1} \\ \downarrow & & \downarrow \\ \mathcal{X}_{1,0}^{\times 2} \times_{\mathcal{X}_{0,0}^{\times 4}} \mathcal{X}_{0,0}^{\times 2} & \longrightarrow & \mathcal{Y}_{1,0}^{\times 2} \times_{\mathcal{Y}_{0,0}^{\times 4}} \mathcal{Y}_{0,0}^{\times 2} \end{array}$$

is a pullback square (here we used that  $\mathcal{X}_{0,\bullet}$  and  $\mathcal{Y}_{0,\bullet}$  are essentially constant). To this end, it suffices to show that for all  $x, y \in \mathcal{X}$  the commutative square

$$\begin{array}{ccc} \mathcal{X}_{1,1} \times_{\mathcal{X}_{0,0}^{\times 4}} \{(x, y)\}^{\times 2} & \longrightarrow & \mathcal{Y}_{1,1} \times_{\mathcal{Y}_{0,0}^{\times 4}} \{(Fx, Fy)\}^{\times 2} \\ \downarrow & & \downarrow \\ \mathcal{X}_{1,0}^{\times 2} \times_{\mathcal{X}_{0,0}^{\times 4}} \{(x, y)\}^{\times 2} & \longrightarrow & \mathcal{Y}_{1,0}^{\times 2} \times_{\mathcal{Y}_{0,0}^{\times 4}} \{(Fx, Fy)\}^{\times 2} \end{array}$$

is a pullback square. But this may be identified with the commutative square

$$\begin{array}{ccc} \mathcal{X}(x, y)_1 & \longrightarrow & \mathcal{Y}(Fx, Fy)_1 \\ \downarrow & & \downarrow \\ \mathcal{X}(x, y)_0^{\times 2} & \longrightarrow & \mathcal{Y}(Fx, Fy)_0^{\times 2} \end{array}$$

which is a pullback square for all  $x, y \in \mathcal{X}$  if and only if  $F$  is locally fully faithful.  $\square$

**2.3. Adjunctions in 2-fold Segal spaces.** Let us recall the notion of adjunctions in terms of pastings:

**Definition 2.22.** Let  $\mathcal{X}$  be a 2-fold Segal space. Then a pair  $(f, g)$  of arrows  $f : x \rightarrow y$  and  $g : y \rightarrow x$  in  $\mathcal{X}$  is called an *adjoint pair*, if there exists two 2-cells, a unit and counit respectively:

$$\begin{array}{ccc} x & \xlongequal{\quad} & x \\ \parallel & & \parallel \\ x & \xrightarrow{f} y \xrightarrow{g} x & \\ \parallel & & \parallel \\ y & \xlongequal{\quad} & y \end{array} \quad \begin{array}{ccc} y & \xrightarrow{g} x \xrightarrow{f} y \\ \parallel & & \parallel \\ y & \xlongequal{\quad} & y \end{array}$$

So that these 2-cells satisfy the triangle identities:

$$\begin{array}{c}
 x \xrightarrow{\quad} x \xrightarrow{f} y \\
 \parallel \qquad \qquad \parallel \\
 x \xrightarrow{f} y \xrightarrow{g} x \xrightarrow{f} y \\
 \parallel \qquad \qquad \parallel \\
 x \xrightarrow{f} y \xrightarrow{\quad} y
 \end{array}
 \simeq
 \begin{array}{c}
 x \xrightarrow{f} y \\
 \parallel \qquad \parallel \\
 x \xrightarrow{f} y
 \end{array},$$

$$\begin{array}{c}
 y \xrightarrow{g} x \xrightarrow{\quad} x \\
 \parallel \qquad \qquad \parallel \\
 y \xrightarrow{g} x \xrightarrow{f} y \xrightarrow{g} x \\
 \parallel \qquad \qquad \parallel \\
 y \xrightarrow{\quad} y \xrightarrow{g} x
 \end{array}
 \simeq
 \begin{array}{c}
 y \xrightarrow{g} x \\
 \parallel \qquad \parallel \\
 y \xrightarrow{g} x
 \end{array}.$$

In this case,  $f$  is the *left adjoint* and  $g$  is the *right adjoint*.

**Proposition 2.23.** *Suppose that  $F : \mathcal{X} \rightarrow \mathcal{Y}$  is a locally full faithful map between 2-fold Segal spaces. Let  $f : x \rightarrow y$  and  $g : y \rightarrow x$  be arrows in  $\mathcal{X}$ . Then  $(f, g)$  is an adjoint pair in  $\mathcal{X}$  if and only if  $(Ff, Fg)$  is an adjoint pair in  $\mathcal{Y}$ .*

*Proof.* Since  $F$  is locally faithful, we obtain (essentially unique) liftings of the unit and counit of the adjunction  $(Ff, Fg)$ . These liftings will again satisfy the triangle identities. This can readily be verified by applying characterization (2) of [Proposition 2.21](#) to the (composable) pastings shapes that govern the pastings of the triangle identities.  $\square$

Note that there are no completeness assumptions in the definition of the adjunctions. We will however be interested in the case that the 2-fold Segal spaces are at least locally complete. In this case, we already have the following two properties:

**Proposition 2.24.** *Suppose that*

$$f : x \rightleftarrows y : g$$

*is an adjunction in a locally complete 2-fold Segal space  $\mathcal{X}$ .<sup>2</sup> Then any arrow  $f' : x \rightarrow y$  is a left adjoint to  $g$  if and only if  $f \simeq f'$  in  $\mathcal{X}(x, y)$ . Dually, any arrow  $g' : y \rightarrow x$  is a right adjoint to  $f$  if and only if  $g \simeq g'$  in  $\mathcal{X}(y, x)$ .*

*Proof.* The proof goes precisely the same as in the classical 2-categorical case using pastings. To make this work, the local completeness of  $\mathcal{X}$  has to come into play in the use of the fact that the commutative square

$$\begin{array}{ccc}
 [1, \{0, 2\}] \sqcup [1, \{0, 3\}] & \longrightarrow & [1, 3] \\
 \downarrow & & \downarrow \\
 [1, 0] \sqcup [1, 0] & \longrightarrow & [1, 0]
 \end{array}$$

is a pushout square of *locally complete* 2-fold Segal spaces (this follows from [Proposition 2.16](#) and [Definition 2.3](#)).  $\square$

<sup>2</sup>We adhere to the convention that the left adjoint arrow appears on top.

**Remark 2.25.** Let  $\text{adj}$  be the *free adjunction* 2-category (see [RV16] for a construction). There are maps  $l : [1] \rightarrow \text{adj}$  and  $r : [1] \rightarrow \text{adj}$  that select, respectively, the left and right adjoint arrow. Suppose that  $\mathcal{X}$  is an  $(\infty, 2)$ -category. Then Riehl and Verity demonstrate [RV16, Theorem 4.3.9] that if  $(f, g)$  is an adjunction in  $\mathcal{X}$ , then there exists a functor

$$a : \text{adj} \rightarrow \mathcal{X}$$

so that  $al \simeq f$  and  $ar \simeq g$  (and this extension is essentially unique if one fixes either a unit or counit, see Theorem 4.4.11 of loc. cit.). Moreover, they prove in [RV16, Theorem 4.4.18] that the restriction maps

$$l^*, r^* : \text{Map}_{\text{Cat}_{(\infty, 2)}}(\text{adj}, \mathcal{X}) \rightarrow \text{Map}_{\text{Cat}_{(\infty, 2)}}([1], \mathcal{X})$$

are equivalences onto the spaces spanned by, respectively, the left and right adjoint arrows in  $\mathcal{X}$ . From this, the preceding two propositions may also be deduced (in the special case that  $\mathcal{X}$  is complete). Here, we implicitly used the equivalence between  $\text{Cat}_{(\infty, 2)}$  and the underlying  $\infty$ -category of the model category of (Joyal) sSet-enriched categories (cf. Subsection 4.3), constructed by Lurie [Lur09b].

**Proposition 2.26.** *Suppose that*

$$f : x \rightleftarrows y : g$$

*is an adjunction in a locally complete 2-fold Segal space  $\mathcal{X}$ . Then for every  $z \in \mathcal{X}$ , we obtain induced adjunctions*

$$\begin{aligned} f_* : \mathcal{X}(z, x) &\rightleftarrows \mathcal{X}(z, y) : g_* \\ g^* : \mathcal{X}(y, z) &\rightleftarrows \mathcal{X}(x, z) : f^* \end{aligned}$$

*of  $\infty$ -categories.*

*Proof.* Let  $\eta$  and  $\epsilon$  be respectively the unit and counit for the adjunction  $(f, g)$ . These are classified by maps  $[1] \rightarrow \mathcal{X}(x, x)$  and  $[1] \rightarrow \mathcal{X}(y, y)$  respectively. They give rise to natural transformations

$$\eta_* : [1] \times \mathcal{X}(z, x) \rightarrow \mathcal{X}(x, x) \times \mathcal{X}(z, x) \rightarrow \mathcal{X}(z, x)$$

and

$$\epsilon_* : [1] \times \mathcal{X}(z, y) \rightarrow \mathcal{X}(y, y) \times \mathcal{X}(z, y) \rightarrow \mathcal{X}(z, y)$$

which are (resp.) our candidates for the unit and counit of the desired adjunction  $(f_*, g_*)$ . The triangle identities for  $\eta$  and  $\epsilon$  now readily imply that the triangle identities hold for  $(\eta_*, \epsilon_*)$ . The other case is handled similarly.  $\square$

**Definition 2.27.** We say that a locally complete 2-fold Segal space  $\mathcal{X}$  is *closed* if for any arrow  $f : x \rightarrow y$  in  $\mathcal{X}$ , the induced functors

$$f_* : \mathcal{X}(z, x) \rightarrow \mathcal{X}(z, y)$$

and

$$f^* : \mathcal{X}(y, z) \rightarrow \mathcal{X}(x, z)$$

admit right adjoints for every  $z \in \mathcal{X}$ .

A locally horizontally (resp. vertically) complete 2-uple Segal space  $\mathcal{P}$  is said to be *horizontally* (resp. *vertically*) *closed* if  $\text{Hor}(\mathcal{P})$  (resp.  $\text{Vert}(\mathcal{P})$ ) is closed.

### 3. COMPANIONS, CONJOINTS AND $\infty$ -EQUIPMENTS

In this section, we will define  $\infty$ -equipments. Essentially, this section shows that the definition of classical equipments can be repeated in the higher setting.

**3.1. Companions and conjoints.** Like in the classical setting of proarrow equipments, the notion of *companions* and *conjoints* will be central to the theory of  $\infty$ -equipments. These notions can be used to interpolate between the vertical and horizontal fragments of a double  $\infty$ -category.

Informally, the process of constructing an  $\infty$ -equipment with vertical fragment given by an  $(\infty, 2)$ -category  $\mathcal{X}$  should be viewed as equipping  $\mathcal{X}$  with *proarrows*. As sketched in the introduction, these proarrows are essential to formulate an internal category theory for  $\mathcal{X}$ . In the case that  $\mathcal{X}$  is the  $(\infty, 2)$ -category of  $\infty$ -categories, the proarrows are given by *profunctors* (see [Example 2.7](#)). Given a functor  $f : \mathcal{C} \rightarrow \mathcal{D}$ , we can associate two canonical ‘representable’ profunctors to it. Namely, there is a *companion* profunctor  $\mathcal{C} \rightarrow \mathcal{D}$  given by

$$\mathcal{D}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{S} : (d, c) \mapsto \text{Map}_{\mathcal{D}}(d, fc),$$

and a *conjoint* profunctor  $\mathcal{D} \rightarrow \mathcal{C}$  given by

$$\mathcal{C}^{\text{op}} \times \mathcal{D} \rightarrow \mathcal{S} : (c, d) \mapsto \text{Map}_{\mathcal{D}}(fc, d).$$

This phenomenon is formalized in the following definition (cf. [\[GP04, Subsection 1.1\]](#) and [\[Shu08, Definition A.1\]](#)):

**Definition 3.1.** Let  $f : x \rightarrow y$  be a vertical arrow in a 2-uple Segal space  $\mathcal{P}$ . Then a *companion for  $f$*  is a horizontal arrow  $y_f : x \rightarrow y$  in  $\mathcal{P}$  which admits two 2-cells in  $\mathcal{P}$

$$\begin{array}{ccc} x & \xrightarrow{y_f} & y \\ f \downarrow & & \parallel \\ y & \xlongequal{\quad} & y \end{array} \quad \text{and} \quad \begin{array}{ccc} x & \xlongequal{\quad} & x \\ \parallel & & \downarrow f \\ x & \xrightarrow{y_f} & y \end{array}$$

which paste together as follows:

$$\begin{array}{ccc} \begin{array}{ccc} x & \xlongequal{\quad} & x \\ \parallel & & \downarrow \\ x & \longrightarrow & y \\ \downarrow & & \parallel \\ y & \xlongequal{\quad} & y \end{array} & \simeq & \begin{array}{ccc} x & \xlongequal{\quad} & x \\ \downarrow & \xlongequal{\quad} & \downarrow \\ y & \xlongequal{\quad} & y, \end{array} \quad \begin{array}{ccc} x & \xlongequal{\quad} & x & \longrightarrow & y \\ \parallel & & \downarrow & & \parallel \\ x & \longrightarrow & y & \xlongequal{\quad} & y \end{array} & \simeq & \begin{array}{ccc} x & \longrightarrow & y \\ \parallel & & \parallel \\ x & \longrightarrow & y. \end{array} \end{array}$$

Dually, a *conjoint for  $f$*  is a horizontal arrow  $f_y : y \rightarrow x$  in  $\mathcal{P}$  so that there exist two 2-cells in  $\mathcal{P}$

$$\begin{array}{ccc} y & \xrightarrow{f_y} & x \\ \parallel & & \downarrow f \\ y & \xlongequal{\quad} & y \end{array} \quad \text{and} \quad \begin{array}{ccc} x & \xlongequal{\quad} & x \\ f \downarrow & & \parallel \\ y & \xrightarrow{f_y} & x \end{array}$$



which compose as follows:

$$\begin{array}{c}
 x \xlongequal{\quad} x \\
 \downarrow \quad \quad \parallel \\
 y \longrightarrow x \\
 \parallel \quad \quad \downarrow \\
 y \xlongequal{\quad} y
 \end{array}
 \approx
 \begin{array}{c}
 x \xlongequal{\quad} x \\
 \downarrow \xlongequal{\quad} \downarrow \\
 y \xlongequal{\quad} y,
 \end{array}
 \quad
 \begin{array}{c}
 y \longrightarrow x \xlongequal{\quad} x \\
 \parallel \quad \quad \downarrow \quad \quad \parallel \\
 y \xlongequal{\quad} y \longrightarrow x
 \end{array}
 \approx
 \begin{array}{c}
 y \longrightarrow x \\
 \parallel \quad \parallel \quad \parallel \\
 y \longrightarrow x
 \end{array}.$$

**Remark 3.2.** A horizontal arrow  $F$  is a conjoint for a vertical arrow  $f$  if and only if  $F$  is a companion for  $f^{\text{op}}$  in  $\mathcal{P}^{\text{vop}}$ . Or similarly,  $F$  is a conjoint for  $f$  if and only if  $F^{\text{op}}$  is a companion for  $f$  in  $\mathcal{P}^{\text{hop}}$ .

**Example 3.3.** Let  $\mathcal{C}$  be an  $\infty$ -category with finite products and consider the associated double  $\infty$ -category of spans  $\text{Span}(\mathcal{C})$  (see [Example 2.9](#)). A vertical arrow  $f : x \rightarrow y$  of  $\text{Span}(\mathcal{C})$  is precisely an arrow in  $\mathcal{C}$ , and it admits both a companion and a conjoint. Namely, the companion  $y_f$  is given by the span

$$x = x \xrightarrow{f} y.$$

and the conjoint  ${}_f y$  is given by span

$$y \xleftarrow{f} x = x.$$

**Proposition 3.4.** Let  $f$  be a vertical arrow in a double Segal space  $\mathcal{P}$  that admits both a conjoint and companion. Then the companion  $y_f$  is left adjoint to the conjoint  ${}_f y$  in  $\text{Hor}(\mathcal{P})$ .

*Proof.* It is proven in exactly the same manner as in the classical case [[Shu08](#), Proposition 5.3]. For completeness, we repeat it here. For the unit, we take the pasting  $\eta$

$$\begin{array}{c}
 x \xlongequal{\quad} x \xlongequal{\quad} x \\
 \parallel \quad \quad \downarrow f \quad \quad \parallel \\
 x \xrightarrow{y_f} y \xrightarrow{{}_f y} x
 \end{array}$$

For the counit, we take the pasting  $\epsilon$

$$\begin{array}{c}
 y \xrightarrow{{}_f y} x \xrightarrow{y_f} y \\
 \parallel \quad \quad \downarrow f \quad \quad \parallel \\
 y \xlongequal{\quad} y \xlongequal{\quad} y.
 \end{array}$$

It is readily verified that this choice of unit and counit satisfies the triangle identity. We will demonstrate just one of them:

$$\begin{array}{c}
 x \xlongequal{\quad} x \xlongequal{\quad} x \xrightarrow{y_f} y \\
 \parallel \quad \quad \downarrow f \quad \quad \parallel \quad \quad \parallel \\
 x \xrightarrow{y_f} y \xrightarrow{{}_f y} x \xrightarrow{y_f} y \\
 \parallel \quad \quad \parallel \quad \quad \downarrow f \quad \quad \parallel \\
 x \xrightarrow{y_f} y \xlongequal{\quad} y \xlongequal{\quad} y
 \end{array}
 \approx
 \begin{array}{c}
 x \xlongequal{\quad} x \xlongequal{\quad} x \xrightarrow{y_f} y \\
 \parallel \quad \quad \downarrow f \quad \quad \downarrow f \quad \quad \parallel \\
 x \xrightarrow{y_f} y \xlongequal{\quad} y \xlongequal{\quad} y.
 \end{array}$$

Here the left hand-side is the whiskered composite  $(\epsilon y_f)(y_f \eta)$ . Pasting further, we obtain

$$(\epsilon y_f)(y_f \eta) \simeq \begin{array}{ccc} x & \xlongequal{\quad} & x \xrightarrow{y_f} y \\ \parallel & & \downarrow f \\ x & \xrightarrow{y_f} & y \xlongequal{\quad} y \end{array} \simeq \begin{array}{ccc} x & \xrightarrow{y_f} & y \\ \parallel & \parallel & \parallel \\ x & \xrightarrow{y_f} & y \end{array},$$

as desired.  $\square$

In [ORR], we will thoroughly study the notion of companions. We will briefly summarize the main prospected results of this work here, which will henceforth be labeled as hypotheses. An important tool in understanding companions which is central to loc. cit., is the *square* or *quintet functor*. In the classical setting, this construction goes back to Ehresmann [Ehr63]. In the (classical) context of companions and conjoinths, this functor has been considered by Grandis and Paré in [GP04].

**Definition 3.5** ([ORR]). The *square functor*

$$\text{Sq} : \text{Cat}_{(\infty,2)} \rightarrow \text{DblCat}_{\infty}^v$$

is given by

$$\text{Sq}(\mathcal{X})_{n,m} := \text{Map}_{\text{Cat}_{(\infty,2)}}([n] \otimes [m], \mathcal{X}).$$

**Remark 3.6.** Here the tensor product

$$- \otimes - : \text{Cat}_2^{\times 2} \rightarrow \text{Cat}_{(\infty,2)}$$

denotes the *oplax Gray tensor product* for strict 2-categories (see [Gra74] for the original definition, and [Lac02] for compatibility with the canonical model structure), followed by the inclusion  $\text{Cat}_2 \rightarrow \text{Cat}_{(\infty,2)}$ . We will just make use of the double cosimplicial gaunt 2-category

$$\Delta \times \Delta \rightarrow \text{Cat}_{(\infty,2)} : ([n], [m]) \mapsto [n] \otimes [m].$$

In this paper, we will not be needing an  $(\infty, 2)$ -categorical (enhanced) version of the Gray tensor product, which is still an active subject of research (e.g. see the introduction of [CM23] for a summary of the current state of affairs). However, in the sequel of this paper [Rui], we will make use of such an enhancement, and be careful about the properties we use.

In [ORR], we aim to show the following:

**Hypothesis 3.7** ([ORR]). *The square functor admits a right adjoint*

$$\text{Comp} : \text{DblCat}_{\infty}^v \rightarrow \text{Cat}_{(\infty,2)}.$$

**Construction 3.8.** Note that there exists (canonical) *tautological double functors*

$$[0, 1] = [1]_v \rightarrow \text{Sq}([1]), \quad \text{and}, \quad [1, 0] = [1]_h \rightarrow \text{Sq}([1]),$$

which select the identity  $[1] = [0] \otimes [1] \rightarrow [1]$  and  $[1] = [1] \otimes [0] \rightarrow [1]$ , respectively. In fact, if

$$[1; 1] = 0 \begin{array}{c} \xrightarrow{0} \\ \downarrow \\ \xrightarrow{1} \end{array} 1$$

denotes the 2-category of the *free 2-cell*, then there are tautological maps

$$[1; 1]_v \rightarrow \mathrm{Sq}([1; 1]), \quad \text{and,} \quad [1; 1]_h \rightarrow \mathrm{Sq}([1; 1]),$$

determined by sending their unique non-degenerate 2-cell to the lax commutative squares

$$\begin{array}{ccc} 0 & \xlongequal{\quad} & 0 \\ 1 \downarrow & \swarrow & \downarrow 0 \\ 1 & \xlongequal{\quad} & 1, \end{array} \quad \text{and,} \quad \begin{array}{ccc} 0 & \xrightarrow{0} & 1 \\ \parallel & \swarrow & \parallel \\ 0 & \xrightarrow{1} & 1, \end{array}$$

respectively. The defined tautological maps are compatible with the face maps  $[1] \rightrightarrows [1; 1]$  between the free 1- and 2-cell. Since  $\mathrm{Cat}_{(\infty, 2)}$  is generated by these free cells, and  $(-)_v$ ,  $(-)_h$  and  $\mathrm{Sq}(-)$  preserve colimits, one can extend the above tautological maps to obtain natural tautological maps

$$\mathcal{X}_v \rightarrow \mathrm{Sq}(\mathcal{X}) \leftarrow \mathcal{X}_h,$$

for any  $(\infty, 2)$ -category  $\mathcal{X}$ .

**Definition 3.9.** The *free companion* is the double category given by

$$\mathbf{comp} := \mathrm{Sq}([1]).$$

**Hypothesis 3.10** ([[ORR](#)]). *Let  $\mathcal{P}$  be a vertically complete double  $\infty$ -category. Then the tautological inclusion  $[1]_v = [0, 1] \rightarrow \mathbf{comp}$  induces an equivalence*

$$\mathrm{Map}_{\mathrm{DblCat}_{\infty}}(\mathbf{comp}, \mathcal{P}) \rightarrow \mathrm{Map}_{\mathrm{DblCat}_{\infty}}([0, 1], \mathcal{P}) = \mathcal{P}_{0,1}$$

*onto the space of vertical arrows of  $\mathcal{P}$  that admit a companion. Similarly, the tautological inclusion  $[1]_h = [1, 0] \rightarrow \mathbf{comp}$  gives rise to an equivalence*

$$\mathrm{Map}_{\mathrm{DblCat}_{\infty}}(\mathbf{comp}, \mathcal{P}) \rightarrow \mathrm{Map}_{\mathrm{DblCat}_{\infty}}([1, 0], \mathcal{P}) = \mathcal{P}_{1,0}$$

*onto the space of horizontal arrows of  $\mathcal{P}$  that are a companion.*

**Remark 3.11.** More generally, the theorem holds for any 2-uple Segal space  $\mathcal{P}$  which has the property that  $\mathrm{Vert}(\mathcal{P})_{1, \bullet}$  and  $\mathrm{Hor}(\mathcal{P})_{1, \bullet}$  are both complete.

Note that if  $\mathcal{P}$  is a vertically complete double  $\infty$ -category, then the tautological maps of [Construction 3.8](#) induce (natural) projections  $\mathrm{Comp}(\mathcal{P}) \rightarrow \mathrm{Vert}(\mathcal{P})$  and  $\mathrm{Comp}(\mathcal{P}) \rightarrow \mathrm{Hor}(\mathcal{P})$ .

**Hypothesis 3.12** ([[ORR](#)]). *Let  $\mathcal{P}$  be a vertically complete double  $\infty$ -category. Then  $\mathcal{P}$  admits all companions if and only if the functor*

$$\mathrm{Comp}(\mathcal{P}) \rightarrow \mathrm{Vert}(\mathcal{P})$$

*is an equivalence between  $(\infty, 2)$ -categories.*

**Corollary 3.13.** *Suppose that  $\mathcal{P}$  is a vertically complete double  $\infty$ -category that admits all companions. Then there exists a map, the companion embedding,*

$$\mathrm{Vert}(\mathcal{P}) \rightarrow \mathrm{Hor}(\mathcal{P})$$

*between locally complete 2-fold Segal spaces, which is locally fully faithful, acts as the identity on objects, and carries vertical arrows to companions.*

*Proof.* It follows from [Hypothesis 3.12](#) that the functor  $\mathrm{Comp}(\mathcal{P}) \rightarrow \mathrm{Vert}(\mathcal{P})$  admits an inverse. Composing this inverse with the projection  $\mathrm{Comp}(\mathcal{P}) \rightarrow \mathrm{Hor}(\mathcal{P})$ , we obtain the desired map.  $\square$

All these results admit the appropriate dual versions for conjoinants, which we will also use in the rest of this paper. These versions may be obtained from the previous results by introducing horizontal opposites. For completeness, we record these here:

**Definition 3.14.** The *free conjoinant* is given by

$$\mathfrak{conj} := \mathfrak{comp}^{\text{hop}}.$$

**Corollary 3.15.** Let  $\mathcal{P}$  be a vertically complete double  $\infty$ -category. Then the tautological inclusion  $[0, 1] \rightarrow \mathfrak{conj}$  induces an equivalence

$$\text{Map}_{\text{DblCat}_\infty}(\mathfrak{conj}, \mathcal{P}) \rightarrow \text{Map}_{\text{DblCat}_\infty}([0, 1], \mathcal{P}) = \mathcal{P}_{0,1}$$

onto the space of vertical arrows of  $\mathcal{P}$  that admit a conjoinant. Similarly, the tautological inclusion  $[1, 0] \rightarrow \mathfrak{conj}$  gives rise to an equivalence

$$\text{Map}_{\text{DblCat}_\infty}(\mathfrak{conj}, \mathcal{P}) \rightarrow \text{Map}_{\text{DblCat}_\infty}([1, 0], \mathcal{P}) = \mathcal{P}_{1,0}$$

onto the space of horizontal arrows of  $\mathcal{P}$  that are a conjoinant.

**Corollary 3.16.** Suppose that  $\mathcal{P}$  is a vertically complete double  $\infty$ -category that admits all conjoinants. Then there exists a map, the conjoinant embedding,

$$\text{Vert}(\mathcal{P}) \rightarrow \text{Hor}(\mathcal{P})^{1,2\text{-op}}$$

between locally complete 2-fold Segal spaces, which is locally fully faithful, acts as the identity on objects, and carries vertical arrows to conjoinants.

**3.2. Higher proarrow equipments.** Our next goal is to introduce the notion of  $\infty$ -equipments. Classically, there are several equivalent ways of defining these. In the higher setting, this is still true, albeit it is more complicated to demonstrate that the characterizations agree. We will show this here. Let us start with a few preparatory definitions. Here, we use the language of pasting shapes of [Rui22] (see Remark 2.5 for a brief and rough summary).

**Definition 3.17.** The *niche*

$$N \subset [1, 1]$$

is the largest subshape of  $[1, 1]$  that does not contain the 1-box  $(0, 0) \rightarrow (1, 0)$ . Pictorially,  $N$  is given by

$$N = \begin{array}{ccc} (0, 0) & & (1, 0) \\ \downarrow & & \downarrow \\ (0, 1) & \longrightarrow & (1, 1). \end{array}$$

A *niche* in a double Segal space  $\mathcal{P}$  is a map  $n : N \rightarrow \mathcal{P}$ . An extension  $[1, 1] \rightarrow \mathcal{P}$  of such a niche is called a *filler of  $n$* .

Throughout this subsection, we will consider the subshape

$$\square \subset [1, 2]$$

which is the largest subshape of  $[1, 2]$  that does not contain the 1-box  $(0, 1) \rightarrow (1, 1)$ . Note that there is a canonical inclusion

$$N \rightarrow \square,$$

which is restricted from the inclusion  $[1, 1] \rightarrow [1, 2]$  that selects the bottom box  $[1, \{1, 2\}]$ .

**Proposition 3.18.** *Let  $\mathcal{P}$  be a double  $\infty$ -category and suppose that  $n$  is a niche in  $\mathcal{P}$  with filler  $c$ . Then the following are equivalent:*

(1) *the restriction map*

$$\mathcal{P}([1, 2]) \times_{\mathcal{P}([1, 1])} \{c\} \rightarrow \mathcal{P}(\square) \times_{\mathcal{P}(N)} \{n\}$$

*is an equivalence,*

(2)  *$c$  classifies a cartesian edge of the source-target projection  $\mathcal{P}_1 \rightarrow \mathcal{P}_0^{\times 2}$ .*

*Proof.* Note that the map of item (1) lies over the space of horizontal arrows of  $\mathcal{P}$  via the tautological inclusion  $[1, 0] \rightarrow \square \rightarrow [1, 2]$  that selects the horizontal 1-box  $(0, 0) \rightarrow (1, 0)$ , so that the map fits in a diagram of spaces

$$\begin{array}{ccc} L := \mathcal{P}([1, 2]) \times_{\mathcal{P}([1, 1])} \{c\} & \longrightarrow & \mathcal{P}(\square) \times_{\mathcal{P}(N)} \{n\} =: R, \\ & \searrow & \swarrow \\ & \mathcal{P}([1, 0]) & \end{array}$$

Hence, (1) holds if and only if the map

$$L_F := \mathcal{P}([1, 2]) \times_{\mathcal{P}([1, 0] \sqcup [1, \{1, 2\}]}) \{(F, c)\} \rightarrow \mathcal{P}(\square) \times_{\mathcal{P}([1, 0] \sqcup N)} \{(F, n)\} =: R_F$$

on fibers is an equivalence for every  $F \in \mathcal{P}([1, 0])$ . Since

$$[1, 2] \simeq ([1, \{1, 2\}] \sqcup \{[0, 1], \{0\}\}) \cup_{[1, 0] \sqcup [1, \{1\}]} [1, \{0, 1\}],$$

we can identify the left hand side with

$$L_F \simeq (\mathcal{P}([1, \{0, 1\}]) \times_{\mathcal{P}([1, 0] \sqcup [1, \{1\}])} \{(F, c|[1, \{0\}])\}) \simeq \text{Map}_{\mathcal{P}_1}(F, c|[1, \{0\}]).$$

We will similarly identify the fiber  $R_F$ . We consider the space  $R'_F$  that sits in a diagram of pullback squares

$$\begin{array}{ccc} R_F & \longrightarrow & \mathcal{P}(\square) \\ \downarrow & & \downarrow \\ R'_F & \longrightarrow & \mathcal{P}(\partial[1, 2]) \\ \downarrow & & \downarrow \\ \{(F, n)\} & \longrightarrow & \mathcal{P}([1, 0] \sqcup N). \end{array}$$

Note that we have a decomposition,

$$\square \simeq \partial[1, 2] \cup_{\partial[1, \{0, 2\}]} [1, \{0, 2\}],$$

so that we obtain a commutative diagram where all ticked squares are pullback squares,

$$\begin{array}{ccccc} R_F & \longrightarrow & \mathcal{P}(\square) & \longrightarrow & \mathcal{P}([1, \{0, 2\}]) \\ \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow \\ R'_F & \longrightarrow & \mathcal{P}(\partial[1, 2]) & \longrightarrow & \mathcal{P}(\partial[1, \{0, 2\}]) \\ \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow \\ \{(F, n)\} & \longrightarrow & \mathcal{P}([1, 0] \sqcup N) & \longrightarrow & \mathcal{P}([1, \{0\}] \sqcup [1, \{2\}]). \end{array}$$

From this, one deduces that  $R_F$  and  $R'_F$  sit in a pullback square

$$\begin{array}{ccc} R_F & \xrightarrow{\quad \lrcorner \quad} & \mathcal{P}([1, \{0, 2\}]) \times_{\mathcal{P}([1, \{0\}] \sqcup [1, \{2\}])} \{(F, c|[1, \{1\}])\} =: S_F \\ \downarrow & & \downarrow \\ R'_F & \longrightarrow & \mathcal{P}(\partial[1, \{0, 2\}]) \times_{\mathcal{P}([1, \{0\}] \sqcup [1, \{2\}])} \{(c|[1, \{0\}], c|[1, \{1\}])\} =: S'_F, \end{array}$$

Hence, we deduce that

$$S_F \simeq \text{Map}_{\mathcal{P}_1}(F, c|[1, \{1\}]).$$

Using the decompositions

$$\begin{aligned} \partial[1, 2] &\simeq ([1, \{0\}] \sqcup N) \cup_{[\{0, \{0\}\} \sqcup \{0, \{1\}\} \sqcup \{1, \{0\}\} \sqcup \{1, \{1\}\}] } (\{0, \{0, 1\}\} \sqcup \{1, \{0, 1\}\}), \\ \partial[1, \{0, 2\}] &\simeq ([1, \{0\}] \sqcup [1, \{2\}]) \cup_{[\{0, \{0\}\} \sqcup \{0, \{2\}\} \sqcup \{1, \{0\}\} \sqcup \{1, \{2\}\}] } (\{0, \{0, 2\}\} \sqcup \{1, \{0, 2\}\}), \end{aligned}$$

one sees that

$$\begin{aligned} R'_F &\simeq \text{Map}_{\mathcal{P}_0}(F|[\{0\}], c|[\{0, \{0\}]) \times \text{Map}_{\mathcal{P}_0}(F|[\{1\}], c|[\{1, \{0\}]), \\ S'_F &\simeq \text{Map}_{\mathcal{P}_0}(F|[\{0\}], c|[\{0, \{1\}]) \times \text{Map}_{\mathcal{P}_0}(F|[\{1\}], c|[\{1, \{1\}]). \end{aligned}$$

Tracing through the identifications above, one sees that the map

$$L_F \rightarrow R_F$$

can be identified with the canonical comparison map that is an equivalence for all  $F \in \mathcal{P}([1, 0])$  if and only if  $c$  is cartesian arrow with respect to the functor  $\mathcal{P}_1 \rightarrow \mathcal{P}_0^{\times 2}$ .  $\square$

**Definition 3.19.** We call a 2-cell  $c : [1, 1] \rightarrow \mathcal{P}$  *cartesian* if the equivalent conditions of [Proposition 3.18](#) are met.

There is also a dual notion of *cocartesian fillers*.

**Definition 3.20.** We define the *roof*

$$R \subset [1, 1]$$

to be the largest subshape of  $[1, 1]$  which does not contain the 1-box  $(0, 1) \rightarrow (1, 1)$ . A *roof* in a double Segal space  $\mathcal{P}$  is a map  $R \rightarrow \mathcal{P}$ . A *filler* for such a roof is an extension to a map  $[1, 1] \rightarrow \mathcal{P}$ .

A 2-cell  $c : [1, 1] \rightarrow \mathcal{P}$  is called *cocartesian* if the 2-cell  $c^{\text{vop}} : [1, 1] \simeq [1, 1]^{\text{vop}} \rightarrow \mathcal{P}^{\text{vop}}$  is cartesian in  $\mathcal{P}^{\text{vop}}$ .

With the definitions, we then obtain directly from [Proposition 3.18](#):

**Corollary 3.21.** *Let  $\mathcal{P}$  be a double  $\infty$ -category and suppose that  $r$  is a roof in  $\mathcal{P}$  with filler  $c$ . Then the following are equivalent:*

- (1)  $c$  is a cocartesian filler for  $r$ ,
- (2) the canonical map

$$\mathcal{P}([1, 2]) \times_{\mathcal{P}([1, 1])} \{c\} \rightarrow \mathcal{P}(\square) \times_{\mathcal{P}(R)} \{r\}$$

is an equivalence,

- (3)  $c$  classifies a cocartesian edge of the source-target projection  $\mathcal{P}_1 \rightarrow \mathcal{P}_0^{\times 2}$ .

The general theory of (co)cartesian fibrations immediately yields the following results:

**Corollary 3.22.** *Suppose  $\mathcal{P}$  is a double  $\infty$ -category and that we have a diagram  $p : [1, 2] \rightarrow \mathcal{P}$  in  $\mathcal{P}$ , that witnesses  $e := p[[1, \{0, 2\}]]$  as the vertical composition of two 2-cells  $c := p[[1, \{0, 1\}]]$ ,  $d := p[[1, \{1, 2\}]]$ . Furthermore, assume that  $c$  is cartesian (resp. cocartesian). Then  $d$  is cartesian (resp. cocartesian) if and only if  $e$  is cartesian (resp. cocartesian).*

**Corollary 3.23.** *Any horizontal equivalence is (co)cartesian.*

The notions of (co)cartesian fillers, companions and conjoints are closely related:

**Proposition 3.24.** *Let  $\mathcal{P}$  be a vertically complete double  $\infty$ -category. Suppose that we have a niche  $n$*

$$\begin{array}{ccc} a & & b \\ f \downarrow & & \downarrow g \\ x & \xrightarrow{F} & y \end{array}$$

in  $\mathcal{P}$ . Then if  $f$  has a companion and  $g$  has a conjoint, the pasting

$$\begin{array}{ccccc} a & \xrightarrow{x_f} & x & \xrightarrow{F} & y & \xrightarrow{g_y} & b \\ f \downarrow & & \parallel & & \parallel & & \downarrow g \\ x & \xlongequal{\quad} & x & \xrightarrow{F} & y & \xlongequal{\quad} & y, \end{array}$$

is a cartesian filler for  $n$ .

Dually, suppose that we have a roof  $r$

$$\begin{array}{ccc} a & \xrightarrow{G} & b \\ f \downarrow & & \downarrow g \\ x & & y \end{array}$$

in  $\mathcal{P}$ . Then if  $f$  has a conjoint and  $g$  has a companion, the pasting

$$\begin{array}{ccccc} a & \xlongequal{\quad} & a & \xrightarrow{G} & b & \xlongequal{\quad} & b \\ f \downarrow & & \parallel & & \parallel & & \downarrow g \\ x & \xrightarrow{f_x} & a & \xrightarrow{G} & b & \xrightarrow{y_g} & y, \end{array}$$

is a cocartesian filler for  $r$ .

We will give a proof of this later. The following choice of notation should now lead to no confusion:

**Notation 3.25.** In the first situation of [Proposition 3.24](#), we will denote the upper horizontal arrow that completes the niche by

$${}_g F_f := g_y \circ F \circ x_f.$$

In the second, dual case in [Proposition 3.24](#), we will write

$${}^g G^f := y_g \circ G \circ f_x.$$

In particular, we have  $a^f = f_x$  and  ${}^g b = y_g$  so that equivalently,

$${}^g G^f = {}^g b \circ G \circ a^f.$$

**Corollary 3.26.** *Suppose that  $\mathcal{P}$  is a vertically complete double  $\infty$ -category. Let  $f : x \rightarrow y$  be a vertical arrow in  $\mathcal{P}$ . Then  $\mathcal{P}$  admits a cartesian filler of the niches*

$$\begin{array}{ccc} x & & y \\ f \downarrow & & \parallel \\ y & \xlongequal{\quad} & y, \end{array} \quad \text{and} \quad \begin{array}{ccc} y & & x \\ \parallel & & \downarrow f \\ y & \xlongequal{\quad} & y, \end{array}$$

*if and only if  $f$  admits a companion, respectively, a conjoint.*

*Dually,  $\mathcal{P}$  admits a cocartesian filler of the roofs*

$$\begin{array}{ccc} x & \xlongequal{\quad} & x \\ \parallel & & \downarrow f \\ x & & y, \end{array} \quad \text{and} \quad \begin{array}{ccc} x & \xlongequal{\quad} & x \\ f \downarrow & & \parallel \\ y & & x, \end{array}$$

*if and only if  $f$  admits a companion, respectively, a conjoint.*

*Proof.* One direction follows from [Proposition 3.24](#). The other direction can be deduced by factoring the identity 2-cell on  $f$  (suitably extended to  $\square$ ) via the universal properties [Proposition 3.18\(1\)](#) and [Corollary 3.21\(2\)](#) of (co)cartesian cells. To check that the associated horizontal pastings are identities, one needs to use these universal properties once again.  $\square$

To prove [Proposition 3.24](#), we need the following observation:

**Lemma 3.27.** *Suppose that  $\mathcal{P}$  is a double  $\infty$ -category, and  $c : [1, 1] \rightarrow \mathcal{P}$  is a cell that factors through a double  $\infty$ -category  $C$  such that the canonical map*

$$C \cup_N \square \rightarrow C \cup_{[1,1]} [1, 2]$$

*is an equivalence. Then  $c$  is cartesian.*

*Proof.* By assumption,  $c$  factors as  $[1, 1] \rightarrow C \xrightarrow{c'} \mathcal{P}$ . This gives rise to a commutative diagram

$$\begin{array}{ccc} \text{Map}_{\text{DblCat}_\infty}(C \cup_{[1,1]} [1, 2], \mathcal{P}) \times_{\text{Map}_{\text{DblCat}_\infty}(C, \mathcal{P})} \{c'\} & \longrightarrow & \mathcal{P}([1, 2]) \times_{\mathcal{P}([1,1])} \{c\} \\ \downarrow & & \downarrow \\ \text{Map}_{\text{DblCat}_\infty}(C \cup_N \square, \mathcal{P}) \times_{\text{Map}_{\text{DblCat}_\infty}(C, \mathcal{P})} \{c'\} & \longrightarrow & \mathcal{P}(\square) \times_{\mathcal{P}(N)} \{c|N\}, \end{array}$$

and by the pasting lemma for pullback squares, both horizontal arrows are equivalences. The left vertical arrow is an equivalence by assumption, hence the right arrow must be an equivalence as well.  $\square$

*Proof of Proposition 3.24.* It suffices to show the first assertion since the dual assertion can be obtained by applying  $(-)^{\text{vop}}$ .

Consider the thickened niche  $N'$  defined by the following pushout square

$$\begin{array}{ccc} [0, 1] \sqcup [0, 1] & \longrightarrow & \text{comp} \sqcup \text{conj} \\ \downarrow & & \downarrow \\ N & \longrightarrow & N'. \end{array}$$

Then the cell  $c : [1, 1] \rightarrow \mathcal{P}$  we are interested in, factors through the cell

$$[1, 1] \rightarrow N'$$



obtained by joining the appropriate cells  $[1, 1] \rightarrow \text{comp} \rightarrow N'$ ,  $[1, 1] \rightarrow [1, 0] \rightarrow N \rightarrow N'$ ,  $[1, 1] \rightarrow \text{conj} \rightarrow N'$ . In view of [Lemma 3.27](#), it suffices to show that the inclusion

$$i : \square' := N' \cup_N \square \rightarrow N' \cup_{[1,1]} [1, 2] =: [1, 2]'$$

is an equivalence of double  $\infty$ -categories.

To this end, we will exhibit an explicit inverse to  $i$ . We will make use of the following 2-dimensional pasting shape:

$$P := \begin{array}{ccccccc} (0,0) & \longrightarrow & (1,0) & \longrightarrow & (2,0) & \longrightarrow & (3,0) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ (0,1) & \longrightarrow & (1,1) & & (2,1) & \longrightarrow & (3,1) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ (0,2) & \longrightarrow & (1,2) & \longrightarrow & (2,2) & \longrightarrow & (3,2) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ (0,3) & \longrightarrow & (1,3) & \longrightarrow & (2,3) & \longrightarrow & (3,3). \end{array}$$

There is a map  $p : P \rightarrow \square'$  represented by the diagram

$$\begin{array}{ccccccc} (0,0) & \xlongequal{\quad} & (0,0) & \longrightarrow & (1,0) & \xlongequal{\quad} & (1,0) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ (0,1) & \xlongequal{\quad} & (0,1) & & (1,1) & \xlongequal{\quad} & (1,1) \\ \parallel & & \downarrow & & \downarrow & & \parallel \\ (0,1) & \longrightarrow & (0,2) & \longrightarrow & (1,2) & \longrightarrow & (1,1) \\ \downarrow & & \parallel & & \parallel & & \downarrow \\ (0,2) & \xlongequal{\quad} & (0,2) & \longrightarrow & (1,2) & \xlongequal{\quad} & (1,2). \end{array}$$

Note that  $P$  admits an inclusion  $[1, 2] \rightarrow P$  that factors the inclusion  $[1, 1] \rightarrow P$  of the outer cell. Restricting  $p$  along this inclusion, we obtain a double functor

$$q : [1, 2] \rightarrow \square',$$

with the following properties:

- the restriction  $q|_{\square}$  is equivalent to the canonical inclusion  $\square \rightarrow \square'$ ,
- the composite  $i q$  is equivalent to the canonical inclusion  $[1, 2] \rightarrow [1, 2]'$ .

These properties may be readily verified by the reader, using the companion and conjunction identities (cf. the proof of [\[Shu08, Theorem 4.1\]](#)). Consequently, the map  $q$  gives rise to a double functor

$$r : [1, 2]' \rightarrow \square'$$

which is an inverse to  $i$ . □

We may combine the preceding few established results, to obtain the following consequence:

**Corollary 3.28.** *Let  $\mathcal{P}$  be a vertically complete double  $\infty$ -category. Then the following statements are equivalent:*

- (1) every vertical arrow in  $\mathcal{P}$  admits both a conjoint and a companion,
- (2) every niche in  $\mathcal{P}$  admits a cartesian filler,
- (3) every roof in  $\mathcal{P}$  admits a cocartesian filler,
- (4) the source-target projection  $\mathcal{P}_1 \rightarrow \mathcal{P}_0^{\times 2}$  is a cartesian fibration,
- (5) the source-target projection  $\mathcal{P}_1 \rightarrow \mathcal{P}_0^{\times 2}$  is a cocartesian fibration.

**Definition 3.29.** A double  $\infty$ -category  $\mathcal{P}$  is called an  $\infty$ -equipment if it is vertically complete and the equivalent conditions of [Corollary 3.28](#) are met. We write

$$\text{Equip}_\infty \subset \text{DblCat}_\infty$$

for the full subcategory spanned by the  $\infty$ -equipments.

**Proposition 3.30.** A double functor between  $\infty$ -equipments preserves (co)cartesian cells.

*Proof.* This follows directly from the fact that double functors preserve companions and conjoints and the description of (co)cartesian cells of [Proposition 3.24](#).  $\square$

**Proposition 3.31.** The subcategory  $\text{Equip}_\infty \subset \text{DblCat}_\infty$  is a reflective subcategory. In particular,  $\infty$ -equipments are closed under limits.

*Proof.* Since  $\text{DblCat}_\infty^v$  is a reflective subcategory of  $\text{DblCat}_\infty$ , it suffices to show that  $\text{Equip}_\infty$  is a reflective subcategory of  $\text{DblCat}_\infty^v$ . This follows from [Hypothesis 3.10](#) which asserts that  $\infty$ -equipments are those vertically complete double  $\infty$ -categories which are local with respect to the inclusions  $[0, 1] \rightarrow \text{comp}$  and  $[0, 1] \rightarrow \text{conj}$ .  $\square$

**Example 3.32.** The double  $\infty$ -categories of spans of [Example 2.9](#) and [Example 3.3](#) are examples of  $\infty$ -equipments. Although we do not prove this here, one can also show that the Morita double  $\infty$ -categories of [Example 2.8](#) are  $\infty$ -equipments.

**Convention 3.33.** Throughout this article, we will adhere to the following terminology. If  $\mathcal{P}$  is an  $\infty$ -equipment, then vertical arrows will plainly be called *arrows*, and its horizontal arrows will be called *proarrows*.

**Remark 3.34.** Suppose that  $\mathcal{P}$  is an  $\infty$ -equipment, and suppose that we have arrows  $f : a \rightarrow c$ ,  $g : b \rightarrow d$  and proarrows  $F : a \rightarrow b$ ,  $G : c \rightarrow d$  in  $\mathcal{P}$ . Then it can be deduced from the universal property of (co)cartesian cells, or from combining [Proposition 2.26](#) and [Proposition 3.4](#), that we have equivalences

$$\text{Map}_{\text{Hor}(\mathcal{P})(c,d)}({}^g F^f, G) \simeq \text{Map}_{\text{Hor}(\mathcal{P})(a,b)}(F, {}_g G_f)$$

which are natural in  $F$  and  $G$ . We will use this interchanging property often when carrying out computations with proarrows.

#### 4. EXAMPLE: THE $\infty$ -EQUIPMENT OF $\infty$ -CATEGORIES

Let us now consider the prototypical example of an  $\infty$ -equipment. Our goal of this section is to recall and closely follow the construction given by Ayala and Francis in [\[AF20\]](#), of a double  $\infty$ -category

$$\text{Cat}_\infty$$

whose objects are (small)  $\infty$ -categories, whose vertical arrows are given by functors and whose horizontal arrows given by *profunctors*, or, from an equivalent perspective, given by *two-sided discrete fibrations* (these are called *bifibrations* by Lurie in [Lur09a]) or *correspondences*.

**4.1. Correspondences, two-sided discrete fibrations and profunctors.** As preparation, we will give a quick recollection of these (equivalent) notions.

**Definition 4.1.** Let  $\mathcal{C}$  and  $\mathcal{D}$  be two (small)  $\infty$ -categories. Then:

- a *correspondence*  $F : \mathcal{C} \rightarrow \mathcal{D}$  is a functor

$$E \rightarrow [1]$$

together with equivalences  $E_1 \simeq \mathcal{C}$  and  $E_0 \simeq \mathcal{D}$ .

- a *two-sided discrete fibration*  $F : \mathcal{C} \rightarrow \mathcal{D}$  is a span

$$\mathcal{C} \leftarrow E \rightarrow \mathcal{D}$$

so that the left leg is a cocartesian fibration, and the right leg is cartesian fibration, and moreover, for any  $d \in \mathcal{D}$ , the pullback  $E_d = E \times_{\mathcal{D}} \{d\} \rightarrow \mathcal{C}$  is a left fibration, and for any  $c \in \mathcal{C}$ , the pullback  $E_c = \{c\} \times_{\mathcal{C}} E \rightarrow \mathcal{D}$  is a right fibration,

- a *profunctor*  $F : \mathcal{C} \rightarrow \mathcal{D}$  is a functor

$$\mathcal{D}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{S}.$$

These objects fit in ambient  $\infty$ -categories as follows. The  $\infty$ -category of correspondences is given by  $\text{Corr} := \text{Cat}_{\infty}/[1]$ , and the  $\infty$ -category of two-sided discrete fibrations is given by the full subcategory

$$\text{DFib} \subset \text{Fun}([1] \cup_{\{0\}} [1], \text{Cat}_{\infty})$$

spanned by those spans that are two-sided discrete fibrations. Note that both these  $\infty$ -categories are canonically fibered over  $\text{Cat}_{\infty}^{\times 2}$ . The  $\infty$ -category of profunctors is obtained by unstraightening the functor

$$\text{Cat}_{\infty}^{\text{op}, \times 2} \rightarrow \widehat{\text{Cat}}_{\infty} : (\mathcal{C}, \mathcal{D}) \mapsto \text{Fun}(\mathcal{D}^{\text{op}} \times \mathcal{C}, \mathcal{S})$$

to a (large) cartesian fibration

$$\text{Prof} \rightarrow \text{Cat}_{\infty}^{\times 2}.$$

If  $\mathcal{C}$  and  $\mathcal{D}$  are  $\infty$ -categories, we will write  $\text{Corr}(\mathcal{C}, \mathcal{D})$ ,  $\text{DFib}(\mathcal{C}, \mathcal{D})$  and  $\text{Prof}(\mathcal{C}, \mathcal{D})$  for the fibers above  $(\mathcal{C}, \mathcal{D})$  of the similarly named  $\infty$ -categories, so that they are the ambient  $\infty$ -categories for correspondences, two-sided discrete fibrations and profunctors from  $\mathcal{C}$  to  $\mathcal{D}$ , respectively.

Like in the classical case, profunctors, two-sided discrete fibrations and correspondences encode the same data. Precisely, in [AF20, Subsection 4.1], the following is shown:

**Proposition 4.2** ([AF20]). *There are equivalences*

$$\text{DFib} \rightarrow \text{Corr} \rightarrow \text{Prof}$$

of  $\infty$ -categories fibered over  $\text{Cat}_{\infty}^{\times 2}$  so that a two-sided discrete fibration  $\mathcal{C} \leftarrow E \rightarrow \mathcal{D}$  is carried to the correspondence

$$\mathcal{C} \cup_{E \times \{1\}} E \times [1] \cup_{E \times \{0\}} \mathcal{D} \rightarrow [1],$$

and a correspondence  $E \rightarrow [1]$  is in turn carried to the presheaf

$$E_0^{\text{op}} \times E_1 \rightarrow E^{\text{op}} \times E \xrightarrow{\text{Map}_E} \mathcal{S}.$$

**Remark 4.3.** Another proof of the fiberwise equivalence between DFib and Corr (in the language of model categories) was first given by Stevenson in [Ste18].

**4.2. Construction via Conduché fibrations.** Central to the construction of the double  $\infty$ -category of  $\infty$ -categories by Ayala and Francis, is the notion of *Conduché fibrations*, also known as *exponentiable fibrations*. These types of fibrations are direct generalizations of the similarly named 1-categorical notions.

**Definition 4.4.** A functor  $p : E \rightarrow \mathcal{C}$  between  $\infty$ -categories is called a *Conduché fibration* if the pullback functor

$$p^* : \text{Cat}_\infty/\mathcal{C} \rightarrow \text{Cat}_\infty/E$$

preserves all colimits or, equivalently, is a left adjoint. The full subcategory of  $\text{Cat}_\infty/\mathcal{C}$  spanned by the Conduché fibrations is denoted by  $\text{Con}(\mathcal{C})$ .

**Remark 4.5.** In the model of quasi-categories, these fibrations are also known as *flat inner fibrations* and they were studied by Lurie in [Lur17, Appendix B.3].

In the double  $\infty$ -category  $\text{Cat}_\infty$  that we aim to construct, there is a natural candidate for the horizontal composition of two correspondences  $E \rightarrow [1]$  and  $E' \rightarrow [1]$  so that  $E'_1 \simeq E_0 \simeq \mathcal{D}$ . Namely, we could form the pushout  $E \cup_{\mathcal{D}} E'$ , which is canonically fibered over  $[2]$ . Pulling back this pushout along the inner face map  $d_1 : [1] \rightarrow [2]$ , we obtain a candidate  $E' \circ E$  for the composite of  $E$  and  $E'$ . The following characterization of Conduché fibrations asserts that Conduché fibrations over  $[2]$  are precisely those functors that are obtained by combining two correspondences via a pushout in the manner above:

**Proposition 4.6** ([AFR18, Lemma 5.16(3)]). *A functor  $p : E \rightarrow \mathcal{C}$  is a Conduché fibration if and only if for every map  $[2] \rightarrow \mathcal{C}$ , the commutative square*

$$\begin{array}{ccc} E \times_{\mathcal{C}} \{1\} & \longrightarrow & E \times_{\mathcal{C}} \{0 \leq 1\} \\ \downarrow & & \downarrow \\ E \times_{\mathcal{C}} \{1 \leq 2\} & \longrightarrow & E \times_{\mathcal{C}} [2] \end{array}$$

is a pushout square of  $\infty$ -categories.

**Corollary 4.7.** *Conduché fibrations are closed under base change.*

Moreover, using characterization (6) of [AFR18, Lemma 5.16], Ayala and Francis immediately deduce the following, which we state without proof (cf. [Lur17, Example B.3.10]):

**Proposition 4.8** ([AF20, Corollary 1.13]). *Every correspondence is a Conduché fibration.*

**Construction 4.9.** Since Conduché fibrations are closed under base change, the functor

$$\text{Cat}_\infty^{\text{op}} \rightarrow \widehat{\text{Cat}}_\infty : \mathcal{C} \mapsto \text{Cat}_\infty/\mathcal{C}$$

obtained by straightening the (large) cartesian fibration  $\text{ev}_1 : \text{Fun}([1], \text{Cat}_\infty) \rightarrow \text{Cat}_\infty$ , gives rise to a functor

$$\text{Cat}_\infty^{\text{op}} \rightarrow \widehat{\text{Cat}}_\infty : \mathcal{C} \mapsto \text{Con}(\mathcal{C}).$$

Restricting this functor along the *twisted inclusion*

$$\text{tw} : \Delta \xrightarrow{(-)^{\text{op}}} \Delta \rightarrow \text{Cat}_\infty,$$

where the last functor is given by the usual inclusion, we obtain a simplicial  $\infty$ -category

$$\mathbb{C}\text{at}_\infty.$$

Ayala and Francis establish the following in the proof of [AF20, Corollary 1.25]:

**Proposition 4.10** ([AF20]). *The simplicial  $\infty$ -category  $\mathbb{C}\text{at}_\infty$  is a (large) double  $\infty$ -category.*

**Proposition 4.11.** *Let  $\mathcal{C}$  and  $\mathcal{D}$  be  $\infty$ -categories. Then there is an equivalence*

$$\text{Vert}(\mathbb{C}\text{at}_\infty)(\mathcal{C}, \mathcal{D}) \simeq \text{Fun}(\mathcal{C}, \mathcal{D})$$

*between  $\infty$ -categories. In particular,  $\mathbb{C}\text{at}_\infty$  is vertically complete.*

*Proof.* In light of the discussion in Remark 2.19, we obtain natural identifications

$$\begin{aligned} \text{Vert}(\mathbb{C}\text{at}_\infty)(\mathcal{C}, \mathcal{D})_n &\simeq \text{Map}_{\text{Con}([n])}([n] \times \mathcal{C}, [n] \times \mathcal{D}) \\ &\simeq \text{Map}_{\text{Cat}_\infty}([n] \times \mathcal{C}, \mathcal{D}) \\ &\simeq \text{Map}_{\text{Cat}_\infty}([n], \text{Fun}(\mathcal{C}, \mathcal{D})), \end{aligned}$$

so that  $\text{Vert}(\mathbb{C}\text{at}_\infty)(\mathcal{C}, \mathcal{D})$  is given by the complete Segal space associated to  $\text{Fun}(\mathcal{C}, \mathcal{D})$ . Here, the contribution of the twist  $(-)^{\text{op}}$  in the definition of  $\mathbb{C}\text{at}_\infty$  is cancelled by our definition of the vertical fragment (see Remark 2.12). The latter assertion follows from the same remark as well.  $\square$

**Definition 4.12.** We define the  $(\infty, 2)$ -category of (small)  $\infty$ -categories to be

$$\text{CAT}_\infty := \text{Vert}(\mathbb{C}\text{at}_\infty).$$

**Remark 4.13.** This will be our preferred model. Shortly, in Subsection 4.3, we will compare this definition to Lurie's model of the  $(\infty, 2)$ -category of  $\infty$ -categories.

We want to demonstrate that  $\mathbb{C}\text{at}_\infty$  is an  $\infty$ -equipment. Since we have shown that it is vertically complete, it remains to verify that  $\mathbb{C}\text{at}_\infty$  admits all companions and conjoinants.

**Proposition 4.14.** *Let  $f : \mathcal{C} \rightarrow \mathcal{D}$  be a functor between  $\infty$ -categories. As a vertical arrow in  $\mathbb{C}\text{at}_\infty$ ,  $f$  admits both a companion and conjoint which may be described as follows:*

- (1) *as correspondences, they are given by respectively the cartesian fibration that classifies*

$$[1]^{\text{op}} \simeq [1] \xrightarrow{f} \text{Cat}_\infty,$$

*and the cocartesian fibration that classifies  $f : [1] \rightarrow \text{Cat}_\infty$ ,*

(2) as two-sided discrete fibrations, they are given by respectively the spans

$$\mathcal{C} \leftarrow \mathcal{C} \times_{\text{Fun}(\{1\}, \mathcal{D})} \text{Fun}([1], \mathcal{D}) \rightarrow \mathcal{D}$$

and

$$\mathcal{D} \leftarrow \text{Fun}([1], \mathcal{D}) \times_{\text{Fun}(\{0\}, \mathcal{D})} \mathcal{C} \rightarrow \mathcal{C},$$

(3) as profunctors, they are given by respectively

$$\mathcal{D}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{S} : (d, c) \mapsto \text{Map}_{\mathcal{D}}(d, fc)$$

and

$$\mathcal{C}^{\text{op}} \times \mathcal{D} \rightarrow \mathcal{S} : (c, d) \mapsto \text{Map}_{\mathcal{D}}(fc, d).$$

Consequently,  $\mathbb{C}\text{at}_{\infty}$  is an  $\infty$ -equipment.

*Proof.* Since  $\text{Prof} \rightarrow \text{Cat}_{\infty}^{\times 2}$  is a cartesian fibration, it directly follows from characterization (4) of [Corollary 3.28](#) and [Proposition 4.2](#) that  $\mathbb{C}\text{at}_{\infty}$  is an  $\infty$ -equipment. Moreover, (3) can readily be deduced from the fact that this fibration  $\text{Prof} \rightarrow \text{Cat}_{\infty}^{\times 2}$  straightens to the functor

$$\text{Cat}_{\infty}^{\times 2} \rightarrow \widehat{\text{Cat}}_{\infty} : (\mathcal{C}, \mathcal{D}) \mapsto \text{PSh}(\mathcal{D}^{\text{op}} \times \mathcal{C}).$$

Then (1) and (2) are verified by tracing through the identifications of [Proposition 4.2](#).  $\square$

Identifying correspondences with profunctors and two-sided discrete fibrations, the horizontal composition in  $\mathbb{C}\text{at}_{\infty}$  can be described as follows:

**Proposition 4.15** ([\[AF20, Lemma 4.5\]](#)). *Let  $F : \mathcal{C} \rightarrow \mathcal{D}$  and  $G : \mathcal{D} \rightarrow \mathcal{E}$  be proarrows of  $\mathbb{C}\text{at}_{\infty}$ . Then the composite  $G \circ F$  can be represented as follows:*

(1) *If  $F$  and  $G$  are two-sided discrete fibrations given by spans  $\mathcal{C} \leftarrow E \rightarrow \mathcal{D}$  and  $\mathcal{D} \leftarrow E' \rightarrow \mathcal{E}$ , then  $G \circ F$  is the two-sided discrete fibration*

$$\mathcal{C} \leftarrow (E \times_{\mathcal{D}} E')[W^{-1}] \rightarrow \mathcal{E},$$

*whose legs are induced by projection, and where  $W$  is the subcategory spanned by the morphisms that are carried to equivalences under the projection  $E \times_{\mathcal{D}} E' \rightarrow \mathcal{C} \times \mathcal{E}$ .*

(2) *If  $F$  and  $G$  are profunctors, then the composite profunctor  $G \circ F$  is given by the coend formula*

$$\mathcal{E}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{S} : (e, c) \mapsto \int^{d \in \mathcal{D}} G(e, d) \times F(d, c).$$

Using some (co)end calculus and the coend formula for the composite of profunctors, we may obtain the following result:

**Corollary 4.16.** *The double  $\infty$ -category  $\mathbb{C}\text{at}_{\infty}$  is horizontally closed. Suppose that  $\mathcal{E}$  is an  $\infty$ -category, and let  $W : \mathcal{C} \rightarrow \mathcal{D}$  be a proarrow in  $\mathbb{C}\text{at}_{\infty}$ . From the perspective of profunctors, we have an adjunction*

$$W_* : \text{Prof}(\mathcal{E}, \mathcal{C}) \rightleftarrows \text{Prof}(\mathcal{E}, \mathcal{D}) : {}^W(-),$$

*where the right adjoint carries  $F \in \text{Prof}(\mathcal{E}, \mathcal{D})$  to the profunctor defined by the end*

$${}^W F(c, e) = \int_{d \in \mathcal{D}} \text{Map}_{\mathcal{S}}(W(d, c), F(d, e)).$$

Similarly, we have an adjunction

$$W^* : \text{Prof}(\mathcal{D}, \mathcal{E}) \rightleftarrows \text{Prof}(\mathcal{C}, \mathcal{E}) : (-)^W,$$

where the right adjoint is given by

$$F^W(e, d) = \int_{c \in \mathcal{C}} \text{Map}_{\mathfrak{S}}(W(d, c), F(e, c))$$

for  $F \in \text{Prof}(\mathcal{C}, \mathcal{E})$ .

*Proof.* This result can readily be obtained from [Proposition 4.15\(2\)](#) by using Fubini's theorem for ends [[Lor18](#)] (an alternative proof can be found in [[Hau22](#)]), and the fact that mapping spaces in functor categories can be computed via ends (see for instance Section 5 of [[GHN17](#)]).  $\square$

**Remark 4.17.** Let  $\mathcal{C}$  and  $\mathcal{E}$  be  $\infty$ -categories. Taking  $\mathcal{D} = [0]$ , [Corollary 4.16](#) has also been proven using the perspective of two-sided discrete fibrations by Stevenson in [[Ste18](#), Proposition 4.24]. Suppose that  $W : X \rightarrow \mathcal{E}$  is a right fibration, i.e. a two-sided discrete fibration from  $[0]$  to  $\mathcal{E}$ . If  $F : \mathcal{C} \rightarrow \mathcal{E}$  is a two-sided discrete fibration given by the span

$$\mathcal{C} \leftarrow E \rightarrow \mathcal{E},$$

then Stevenson proves that the two-sided discrete fibration  ${}^W F : \mathcal{C} \rightarrow [0]$  is the left fibration that sits in the following pullback square

$$\begin{array}{ccc} E' & \longrightarrow & \text{Fun}(X, E) \\ {}^{w_F} \downarrow & & \downarrow \text{Fun}(X, F) \\ \mathcal{C} & \longrightarrow & \text{Fun}(X, \mathcal{C}) \times \text{Fun}(X, \mathcal{E}), \end{array}$$

of  $\infty$ -categories, where the bottom arrow is the product of the diagonal and the constant functor that selects  $W$ .

Riehl and Verity have also given a proof of [Corollary 4.16](#), now in full generality, from the perspective of two-sided discrete fibrations [[RV22](#), Theorem 12.3.6].

**Remark 4.18.** Passing to larger universes, one can repeat the same constructions in this section to obtain larger  $\infty$ -equipments for larger  $\infty$ -categories. For instance, we have a (very large)  $\infty$ -equipment

$$\widehat{\text{Cat}}_{\infty}$$

of large  $\infty$ -categories. The proarrows of this  $\infty$ -equipment are again given by correspondences, two-sided discrete fibrations, and profunctors (but the latter are now valued in  $\hat{\mathfrak{S}}$ , the  $\infty$ -category of large spaces). Note that there is a canonical double functor  $\text{Cat}_{\infty} \rightarrow \widehat{\text{Cat}}_{\infty}$  that is levelwise fully faithful. Consequently, throughout this article, we will mostly be implicit about enlarging universes, and hence leave out the decorations with hat symbols.

**4.3. Comparison of the vertical fragment.** As stated in [Definition 4.12](#), the vertical fragment  $\text{CAT}_\infty$  of  $\text{Cat}_\infty$  will be our preferred model of the  $(\infty, 2)$ -category of  $\infty$ -categories. We will show that this is justified by comparing this vertical fragment to another model for the  $(\infty, 2)$ -category of  $\infty$ -categories due to Lurie, constructed in [\[Lur09b\]](#).

Lurie has defined a model structure on the category of *marked simplicial sets*

$$\text{sSet}^+,$$

which is equivalent to the Joyal model structure on simplicial sets ([\[Lur09a, Chapter 3\]](#)). We may then consider the category

$$\text{Cat}^{\text{sSet}^+}$$

of *marked simplicial categories*: categories enriched over  $\text{sSet}^+$  (with respect to the cartesian monoidal structure). This carries an induced model structure [\[Lur09a, Section A.3\]](#). In [\[Lur09b\]](#), Lurie (essentially) shows that there exists a *nerve functor*

$$N : \text{Cat}^{\text{sSet}^+} \rightarrow \text{Cat}_{(\infty, 2)},$$

which carries the weak equivalences  $W$  between fibrants to equivalences (by first passing through the category of *scaled simplicial sets*). He demonstrates that the induced derived functor

$$\text{Cat}^{\text{sSet}^+, \circ}[W^{-1}] \rightarrow \text{Cat}_{(\infty, 2)}$$

is an equivalence (cf. the discussion in [\[HHLN21, Subsection 5.1\]](#)). Here, we wrote  $\text{Cat}^{\text{sSet}^+, \circ} \subset \text{Cat}^{\text{sSet}^+}$  for the full subcategory spanned by the fibrant marked simplicial categories. Similarly, we will write  $\text{sSet}^{+, \circ} \subset \text{sSet}^+$  for the full (marked simplicial) subcategory of  $\text{sSet}^+$  spanned by the fibrants (i.e. quasi-categories with their equivalences marked). Note that this yields a fibrant marked simplicial category. The goal is now to show the following:

**Proposition 4.19.** *There exists an equivalence of  $(\infty, 2)$ -categories*

$$\text{CAT}_\infty \simeq N(\text{sSet}^{+, \circ}).$$

To prove this, we will need a generalization of Lurie's straightening theorem. Recall that this theorem sets up an equivalence

$$\text{Fun}(\mathcal{C}, \text{Cat}_\infty) \rightarrow \text{Cocart}(\mathcal{C}),$$

given by the unstraightening, for an  $\infty$ -category  $\mathcal{C}$ . Here

$$\text{Cocart}(\mathcal{C}) \subset \text{Cat}_\infty/\mathcal{C}$$

is the subcategory spanned by the cocartesian fibrations and morphisms that preserve cocartesian edges. On account of [\[GHN17, Proposition A.31\]](#), the unstraightening functor is (coherently) natural in  $\mathcal{C}$ .

Following [\[HHLN21\]](#), let us write

$$\text{Cocart}^{\text{lax}}(\mathcal{C}) \subset \text{Cat}_\infty/\mathcal{C}$$

for the *full* subcategory spanned by the cocartesian fibrations. Using Lurie's  $(\infty, 2)$ -categorical straightening theorem [\[Lur09b, Theorem 3.8.1\]](#), Haugseng, Hebestreit, Linskens and Nuiten show a lax variant of the  $(\infty, 1)$ -categorical straightening theorem [\[HHLN21, Theorem 5.3.1\]](#). This takes the form of an equivalence on level of  $(\infty, 2)$ -categories between  $(\infty, 2)$ -categories of functors



and lax natural transformations, and  $(\infty, 2)$ -categorical enhancements of the lax variants of the  $\infty$ -category of cocartesian fibrations. In this article, we will not need the full strength of this equivalence, but a less general, specialized form of the lax straightening theorem:

**Theorem 4.20** ([HHLN21, Theorem 5.3.1]). *There is an equivalence*

$$\mathrm{Map}_{\mathrm{Cat}_{(\infty,2)}}([n] \otimes [m], N(\mathrm{sSet}^{+, \circ})) \simeq \mathrm{Map}_{\mathrm{Cat}_{\infty}}([n], \mathrm{Cocart}^{\mathrm{lax}}([m])).$$

natural in  $[n], [m] \in \Delta$ .

**Remark 4.21.** In [HHLN21], the authors use an  $(\infty, 2)$ -categorical oplax Gray tensor product. Tensoring the 1-categories  $[n]$  and  $[m]$  via this refinement, they show in [HHLN21, Proposition 5.1.9] that the resulting  $(\infty, 2)$ -category coincides with the classical Gray tensor product  $[n] \otimes [m]$  (see Remark 3.6).

There is one more ingredient needed to prove the comparison. This is a well-known fact, and we refer to [HHLN21, Lemma 5.1.11] for a proof:

**Lemma 4.22.** *Let  $n, m \geq 0$ . Then there is a pushout square*

$$\begin{array}{ccc} \coprod_{0 \leq i \leq n} \{i\} \otimes [m] & \longrightarrow & [n] \otimes [m] \\ \downarrow & & \downarrow \\ \coprod_{0 \leq i \leq n} \{i\} & \longrightarrow & [n; m, \dots, m] \end{array}$$

of  $(\infty, 2)$ -categories, where the map on the right is natural in  $\Delta \times \Delta$ .

*Proof of Proposition 4.19.* Combining Theorem 4.20 and Lemma 4.22, we deduce that there is a natural pullback square

$$\begin{array}{ccc} (N(\mathrm{sSet}^{+, \circ}))_{n,m} & \longrightarrow & \mathrm{Map}_{\mathrm{Cat}_{\infty}}([n], \mathrm{Cocart}^{\mathrm{lax}}([m])) \\ \downarrow & & \downarrow \\ \mathrm{Map}_{\mathrm{Cat}_{\infty}}([0], \mathrm{Cat}_{\infty})^{\times(n+1)} & \longrightarrow & \mathrm{Map}_{\mathrm{Cat}_{\infty}}([0], \mathrm{Cocart}^{\mathrm{lax}}([m]))^{\times(n+1)} \end{array}$$

for  $n, m \geq 0$ . Here the components of the right map are given by restricting to the objects of  $[n]$ . The bottom map is induced by the restriction of the degeneracy (pullback) functor

$$[m] \times - : \mathrm{Cat}_{\infty} \rightarrow \mathrm{Cat}_{\infty}/[m].$$

Since both  $\mathrm{Cocart}^{\mathrm{lax}}([m])$  and  $\mathrm{Con}([m])$  are full subcategories of  $\mathrm{Cat}_{\infty}/[m]$  that contain the images of these degeneracies, we deduce that, as a 2-fold Segal space,  $\mathrm{sSet}^{+, \circ}$  sits in levelwise pullback squares

$$\begin{array}{ccc} (N(\mathrm{sSet}^{+, \circ}))_{n,m} & \longrightarrow & \mathrm{Map}_{\mathrm{Cat}_{\infty}}([n], \mathrm{Con}([m])) \\ \downarrow & & \downarrow \\ \mathrm{Map}_{\mathrm{Cat}_{\infty}}([0], \mathrm{Cat}_{\infty})^{\times(n+1)} & \longrightarrow & \mathrm{Map}_{\mathrm{Cat}_{\infty}}([0], \mathrm{Con}([m]))^{\times(n+1)}. \end{array}$$

This is also the defining feature of the vertical fragment of  $\mathrm{Cat}_{\infty}$ .  $\square$

5. EXAMPLE: THE  $\infty$ -EQUIPMENT OF INTERNAL CATEGORIES

Given an  $\infty$ -topos  $\mathcal{E}$ , there is a notion of complete categorical objects, i.e. *internal ( $\infty$ -)categories*, in  $\mathcal{E}$ . Moreover, one may develop a category theory for these internal categories, called *internal category theory*. For presheaf  $\infty$ -toposes, this was firstly done in [BDG<sup>+</sup>16] via an ‘external’ approach, where it is named *parametrized higher category theory*. In a slightly more general setting than  $\infty$ -toposes, a first approach to internal  $\infty$ -category theory is due to Rasekh [Ras18]. In the setting of  $\infty$ -toposes, it is intensively studied by Martini and Wolf in (among others) [MW21]. Classically, an introduction to category theory internal to a 1-topos can be found in [Joh02].

As  $\infty$ -equipments have formal category theories (Section 6) associated to them, one might hope that there exists an  $\infty$ -equipment

$$\mathbb{C}\text{at}_\infty(\mathcal{E})$$

that recovers the category theory internal to the  $\infty$ -topos  $\mathcal{E}$ . We will give a construction of such an  $\infty$ -equipment here, which is a suitable generalization of the construction given in Section 4, so that we recover  $\mathbb{C}\text{at}_\infty$  if  $\mathcal{E} = \mathcal{S}$ .

**5.1. The double  $\infty$ -category of categorical objects.** We commence by introducing an internal variant of Conduché fibrations. This is essentially a modified adaptation of the material in [Hau17] (applied to the cartesian case). Throughout, we fix an  $\infty$ -category  $\mathcal{C}$  and assume the following three things about  $\mathcal{C}$ :

- (1)  $\mathcal{C}$  has all finite limits,
- (2)  $\mathcal{C}$  admits all geometric realizations (i.e. all colimits of  $\Delta^{\text{op}}$ -shaped diagrams),
- (3) fibered products and geometric realizations in  $\mathcal{C}$  are compatible: for any pullback square

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ C & \longrightarrow & D \end{array}$$

in  $\text{Fun}(\Delta^{\text{op}}, \mathcal{C})$  so that  $D$  is essentially constant, the induced square

$$\begin{array}{ccc} \text{colim}_{\Delta^{\text{op}}} A & \longrightarrow & \text{colim}_{\Delta^{\text{op}}} B \\ \downarrow & & \downarrow \\ \text{colim}_{\Delta^{\text{op}}} C & \longrightarrow & \text{colim}_{\Delta^{\text{op}}} D \simeq D_0 \end{array}$$

on geometric realizations is a pullback square as well.

By using that  $\Delta^{\text{op}}$  is sifted, we see that it suffices to check assumption (3) for those pullback squares where  $B$  is essentially constant as well. To wit, suppose that (3) holds in these special cases. Then for any general pullback square as in (3), we have a commutative triangle

$$\begin{array}{ccc} \text{colim}_{\Delta^{\text{op}}} A & \xrightarrow{\simeq} & \text{colim}_{([n],[m]) \in \Delta^{\text{op}} \times \Delta^{\text{op}}} (B_n \times_{D_0} C_m) \\ & \searrow & \downarrow \\ & & \text{colim}_{\Delta^{\text{op}}} B \times_{D_0} \text{colim}_{\Delta^{\text{op}}} C, \end{array}$$

and the horizontal arrow is an equivalence since the diagonal  $\Delta^{\text{op}} \rightarrow \Delta^{\text{op}} \times \Delta^{\text{op}}$  is final. Note that the top right colimit can be written as an iterated colimit. With these observations, one can now verify that the right leg is an equivalence under the assumptions (applying (3) twice for the special cases).

In particular, since  $\infty$ -toposes have all small (co)limits and universality of colimits, we see that conditions (1)-(3) are met if  $\mathcal{C}$  is an  $\infty$ -topos.

**Definition 5.1.** A *categorical  $(n+1)$ -module in  $\mathcal{C}$*  is a functor

$$M : (\Delta/[n])^{\text{op}} \rightarrow \mathcal{C}$$

which satisfies the *Segal condition*: for any map  $f : [k] \rightarrow [n]$  the Segal map

$$M(f) \rightarrow M(f|_{\{0,1\}}) \times_{M(f|_{\{1\}})} \cdots \times_{M(f|_{\{k-1\}})} M(f|_{\{k-1,k\}})$$

is an equivalence. We will write

$$\text{Mod}_{n+1}(\mathcal{C}) \subset \text{Fun}((\Delta/[n])^{\text{op}}, \mathcal{C})$$

for the full subcategory spanned by these categorical  $(n+1)$ -modules.

Consider the cosimplicial object  $C_\bullet$  given by the composite

$$\Delta \xrightarrow{(-)^{\text{op}}} \Delta \xrightarrow{\Delta/[\bullet]} \text{Cat}_\infty \xrightarrow{(-)^{\text{op}}} \text{Cat}_\infty$$

where the middle functor acts by pushing forward. Then we obtain a simplicial  $\infty$ -category

$$\text{Fun}(C_\bullet, \mathcal{C}) : \Delta^{\text{op}} \rightarrow \widehat{\text{Cat}}_\infty : [n] \mapsto \text{Fun}((\Delta/[n])^{\text{op}}, \mathcal{C}).$$

One may readily verify that sees that if  $\alpha : [m] \rightarrow [n]$  is a structure map, and  $M$  is a categorical  $(n+1)$ -module in  $\mathcal{C}$  then the *restriction of  $M$  along  $\alpha$*

$$\alpha^* M : (\Delta/[m])^{\text{op}} \xrightarrow{(\alpha_i^{\text{op}})^{\text{op}}} (\Delta/[n])^{\text{op}} \rightarrow \mathcal{C},$$

is a categorical  $(m+1)$ -module. Hence, we obtain a subobject  $\text{Mod}_{\bullet+1}(\mathcal{C})$  of  $\text{Fun}(C_\bullet, \mathcal{C})$  that is spanned by the categorical modules. However, this fails to be a double  $\infty$ -category in general. The problem is that there are too many trimodules: not all are obtained by composing two bimodules. This can be remedied by restricting to the trimodules that are obtained by composing two bimodules. To formulate this, we will need the following auxiliary shape category, defined by Haugseng:

**Definition 5.2** ([Hau17, Definition 4.13]). A map  $f : [m] \rightarrow [n]$  is called *cellular* if  $f(i+1) \leq f(i)+1$  for all  $i \in [m]$ , i.e. if  $f$  has convex image. The full subcategory of  $\Delta/[n]$  spanned by these cellular maps is denoted by  $\Lambda/[n]$ .

We then have the following analogue of [Proposition 4.6](#):

**Proposition 5.3.** *Let  $M$  be a categorical  $(n+1)$ -module in  $\mathcal{C}$ . Then, the following statements are equivalent:*

(1)  *$M$  is left Kan extended along the inclusion*

$$(\Lambda/[n])^{\text{op}} \rightarrow (\Delta/[n])^{\text{op}},$$

(2) *for any map  $\alpha : [2] \rightarrow [n]$ , the restriction  $\alpha^* M$  is left Kan extended along the inclusion*

$$(\Lambda/[2])^{\text{op}} \rightarrow (\Delta/[2])^{\text{op}},$$

(3) for any map  $\alpha : [2] \rightarrow [n]$ , the canonical map

$$\operatorname{colim}_{[k] \in \Delta^{\text{op}}} M([k+2] \xrightarrow{0 \leq 1 \dots \leq 1 \leq 2} [2] \xrightarrow{\alpha} [n]) \rightarrow M([1] \xrightarrow{0 \leq 2} [2] \xrightarrow{\alpha} [n])$$

is an equivalence.

**Remark 5.4.** Note that condition (3) of [Proposition 5.3](#) essentially asserts that if  $M$  is a trimodule in  $\mathcal{C}$ , then  $M$  is left Kan extended along  $(\Lambda/[2])^{\text{op}} \rightarrow (\Lambda/[2])^{\text{op}}$  if and only if  $M(\{0 \leq 2\})$  is the two-sided bar construction of  $M(\{0 \leq 1\})$  and  $M(\{1 \leq 2\})$  over  $M(\{1\})$ . For  $\mathcal{C} = \mathcal{S}$ , the same bar construction formula makes an appearance in [\[AFR18, Lemma 5.16\(5\)\]](#).

In order to prove the above proposition, we will make use of the following construction and observation (cf. [Proposition 4.16](#) and [Lemma 4.17](#) of [\[Hau17\]](#)):

**Construction 5.5.** Let  $f : [m] \rightarrow [n]$  be a map of posets. Then we will write  $D(f)$  for the set that measures the non-cellularity of  $f$ , i.e. it is given by the difference of the convex hull of  $\operatorname{im}(f)$  and  $\operatorname{im}(f)$ :

$$D(f) := \{f(0), f(0) + 1, \dots, f(m) - 1, f(m)\} \setminus \operatorname{im}(f).$$

We define a functor

$$T_f : \prod_{D(f)} \Delta \rightarrow f/(\Lambda/[n]) := f/(\Delta/[n]) \times_{\Delta/[n]} \Lambda/[n]$$

that carries an indexed poset  $[k_{\bullet}]$  to the triangle

$$\begin{array}{ccc} [m] & \xrightarrow{t_{[k_{\bullet}]}} & [m + \sum_{i \in D(f)} (k_i + 1)] \\ & \searrow f & \swarrow f_{[k_{\bullet}]} \\ & [n] & \end{array}$$

where  $f_{[k_{\bullet}]}$  is the map

$$f_{[k_{\bullet}]} : [1 + \sum_{i \in D(f) \setminus \{0,1\}} (k_i + 1)] \cup_{[0]} \dots \cup_{[0]} [1 + \sum_{i \in D(f) \setminus \{m-1,m\}} (k_i + 1)] \rightarrow [n]$$

so that  $f_{[k_{\bullet}]}^{j,j+1} := f_{[k_{\bullet}]}[1 + \sum_{i \in D(f) \setminus \{j,j+1\}} (k_i + 1)]$  is given by

$$f_{[k_{\bullet}]}^{j,j+1}(x) = \begin{cases} f(j), & x = 0 \\ t, & \sum_{i=f(j)+1}^{t-1} (k_i + 1) < x \leq \sum_{i=f(j)+1}^t (k_i + 1) \\ f(j+1), & \text{if } x = 1 + \sum_{i \in D(f) \setminus \{j,j+1\}} (k_i + 1), \end{cases}$$

and the top map  $t_{[k_{\bullet}]}$  is the unique injective map that fits in the triangle. It is readily verified that this construction is functorial, with the functoriality of  $T_f$  being uniquely determined by pushing forward structure maps along the canonical inclusions

$$\rho_{[k_{\bullet}]}^i : [k_i] \rightarrow [1 + \sum_{i \in D(f) \setminus \{j,j+1\}} (k_i + 1)] \rightarrow [m + \sum_{i \in D(f)} (k_i + 1)].$$

**Lemma 5.6.** Let  $f : [m] \rightarrow [n]$  be a map of posets. Then the functor  $T_f$  of [Construction 5.5](#) is final.

*Proof.* In light of Quillen's theorem A [\[Lur09a, Theorem 4.1.3.1\]](#), it suffices to show that the category

$$T_f/X := (f/(\Lambda/[n]))/X \times_{f/(\Lambda/[n])} \prod_{D(f)} \Delta$$

is weakly contractible for every  $X \in f/(\Lambda/[n])$ . Note that such  $X$  is given by a triangle

$$\begin{array}{ccc} [m] & \xrightarrow{h} & [l] \\ & \searrow f & \swarrow g \\ & & [n], \end{array}$$

so that  $g$  is cellular. It is readily verified that maps  $\phi : T_f([k_\bullet]) \rightarrow X$  in  $f/(\Lambda/[n])$  are uniquely determined by the restrictions  $\phi \rho_{[k_\bullet]}^i : [k_i] \rightarrow [l]$ . Since each restriction should have image in the (non-empty) fiber  $g^{-1}(i) \subset [l]$ , we see that the choice of  $\phi$  so that

$$\phi \rho_{[k_\bullet]}^i : [k_i] \rightarrow [l]$$

is the unique injective map with image  $g^{-1}(i)$ , constitutes a final object of the lax slice  $T_f/X$ .  $\square$

Additionally, we will need the following two results:

**Lemma 5.7.** *Suppose that  $N : (\Lambda/[n])^{\text{op}} \rightarrow \mathcal{C}$  satisfies the Segal condition as in [Definition 5.1](#). Then the left Kan extension  $M : (\Delta/[n])^{\text{op}} \rightarrow \mathcal{C}$  of  $N$  along  $(\Lambda/[n])^{\text{op}} \rightarrow (\Delta/[n])^{\text{op}}$  satisfies the Segal condition as well.*

*Proof.* Let  $f : [m] \rightarrow [n]$  be a map. Then, in view of [Lemma 5.6](#) and the Segal condition for  $N$ , we compute that

$$\begin{aligned} M(f) &= \operatorname{colim}_{[k_\bullet] \in \prod_{D(f)} \Delta^{\text{op}}} N(f_{[k_\bullet]}) \\ &\simeq \operatorname{colim}_{[k_\bullet] \in \prod_{D(f)} \Delta^{\text{op}}} N(f_{[k_\bullet]}^{0,1}) \times_{N(f|\{1\})} \cdots \times_{N(f|\{m-1\})} N(f_{[k_\bullet]}^{m-1,m}) \end{aligned}$$

Using the fact that we have a decomposition  $D(f) = D(f|\{0,1\}) \amalg \cdots \amalg D(f|\{m-1,m\})$  and compatibility of the fibered product with geometric realizations, we deduce that

$$M(f) \simeq M(f|\{0,1\}) \times_{M(f|\{1\})} \cdots \times_{M(f|\{m-1\})} M(f|\{m-1,m\}),$$

as desired.  $\square$

**Lemma 5.8.** *Suppose that  $M$  is a categorical  $(n+1)$ -module in  $\mathcal{C}$ . Let  $f : [1] \rightarrow [n]$  be a map of posets that factors as*

$$[1] \xrightarrow{g} [m] \xrightarrow{\alpha} [n]$$

so that  $\alpha$  is injective. Then there is a canonical commutative triangle

$$\begin{array}{ccc} \operatorname{colim}_{[k_\bullet] \in \prod_{D(g)} \Delta^{\text{op}}} M(\alpha g_{[k_\bullet]}) & \longrightarrow & \operatorname{colim}_{[k_\bullet] \in \prod_{D(f)} \Delta^{\text{op}}} M(f_{[k_\bullet]}), \\ & \searrow & \swarrow \\ & & M(f) \end{array}$$

and the top map is an equivalence as soon as for all  $0 \leq j < m$ , the comparison map

$$\operatorname{colim}_{[k_\bullet] \in \prod_{D(\alpha|\{j,j+1\})} \Delta^{\text{op}}} M(\alpha_{[k_\bullet]}^{j,j+1}) \rightarrow M(\alpha|\{j,j+1\})$$

is an equivalence.

*Proof.* Note that  $\alpha$  restricts to an inclusion  $D(g) \rightarrow D(f)$  and gives rise to a canonical natural transformation

$$\begin{array}{ccccc}
& & & & \\
& & \alpha^* & \longrightarrow & \prod_{D(g)} \Delta^{\text{op}} & \xrightarrow{T_g} & (\Delta/[m])^{\text{op}} & \xrightarrow{(\alpha_1)^{\text{op}}} & (\Delta/[n])^{\text{op}} \\
& \nearrow & & & & \Downarrow & & & \\
\prod_{D(f)} \Delta^{\text{op}} & & & & & & & & \\
& \searrow & & & & & & & \\
& & & \xrightarrow{T_f} & & & & & 
\end{array}$$

Hence, we obtain maps

$$\text{colim}_{[k_\bullet] \in \prod_{D(g)} \Delta^{\text{op}}} M(\alpha g_{[k_{\alpha(\bullet)}]}) \xleftarrow{\cong} \text{colim}_{[k_\bullet] \in \prod_{D(f)} \Delta^{\text{op}}} M(\alpha g_{[k_{\alpha(\bullet)}]}) \rightarrow \text{colim}_{[k_\bullet] \in \prod_{D(f)} \Delta^{\text{op}}} M(f_{[k_\bullet]}),$$

which are all compatible with the canonical projections to  $M(f)$ . The first map is an equivalence as  $\alpha^*$  is final, so that we may obtain the desired commutative triangle by picking an inverse. To show the latter statement, we suppose that the comparison maps for  $\alpha$  as stated in the lemma, are equivalences. By 2-out-of-3, it suffices to show that the right arrow in the span displayed above, is an equivalence, which readily follows from a similar computation as in [Lemma 5.7](#).  $\square$

*Proof of Proposition 5.3.* Let us first show that (2) and (3) are equivalent. In light of [Lemma 5.7](#), a categorical module  $M : (\Delta/[2])^{\text{op}} \rightarrow \mathcal{C}$  is Conduché if and only if for any map  $f : [1] \rightarrow [2]$ , the canonical map

$$\text{colim}_{g \rightarrow f \in (\Delta/[2])^{\text{op}}/f} M(g) \rightarrow M(f)$$

is an equivalence. We may assume that  $f$  is non-cellular so that  $f = d_1$ . Hence, rewriting the above colimit using [Lemma 5.6](#) directly yields the necessary and sufficient condition (3) for  $M$  to be Conduché.

Next, we show that (1) implies (2). If  $M$  is a Conduché module, then we have to show that for every map  $\alpha : [2] \rightarrow [n]$ , the restriction  $\alpha^* M$  is again Conduché. In the trivial case that  $\alpha$  is not injective, this is readily verified, and if  $\alpha$  is injective then it follows from [Lemma 5.8](#) (applied to  $\alpha^{\text{op}}$ ).

Finally, suppose that  $M$  is a categorical module so that (3) holds. Then we will show that  $M$  is Conduché. To this end, we proceed by induction as follows. For  $f : [m] \rightarrow [n]$ , let us define its *non-cellularity index*  $I(f)$  by

$$I(f) := \max_{0 \leq j < m} |D(f|_{\{j, j+1\}})|.$$

We will show that for every map  $f$ , the comparison map

$$c(f) : \text{colim}_{[k_\bullet] \in \prod_{D(f)} \Delta^{\text{op}}} M(f_{[k_\bullet]}) \rightarrow M(f)$$

is an equivalence by induction on  $i = I(f)$ . If  $i = 0$ , the statement is trivial. Suppose now that the statement holds for  $i$ . In order to show that the statement holds for  $i + 1$ , we may use [Lemma 5.7](#) to reduce to showing the statement for any map  $f : [1] \rightarrow [n]$  with  $I(f) = i + 1$ . We may factor such a map  $f$  as

$$[1] \xrightarrow{d_1} [2] \xrightarrow{\alpha} [n]$$

so that  $\alpha$  is injective with  $I(\alpha) = i$ . It follows now directly from the assumptions and [Lemma 5.8](#) that  $c(f)$  is an equivalence as well.  $\square$

**Definition 5.9.** If  $M$  is a categorical  $(n+1)$ -module in  $\mathcal{C}$  that satisfies the equivalent *Conduché conditions* (1)-(3) of [Proposition 5.3](#), then we say that  $M$  is a *Conduché module*. We write

$$\text{Con}_{n+1}(\mathcal{C}) \subset \text{Mod}_{n+1}(\mathcal{C})$$

for the full subcategory spanned by the Conduché modules.

**Corollary 5.10.** *Every categorical bimodule in  $\mathcal{C}$  satisfies the Conduché condition.*

**Construction 5.11.** In light of [Proposition 5.3](#), the Conduché property for categorical modules is preserved under restriction along maps in  $\Delta$ . Consequently, we obtain a further subobject

$$\text{Cat}(\mathcal{C}) \subset \text{Mod}_{\bullet, n+1}(\mathcal{C}) \subset \text{Fun}(C_{\bullet}, \mathcal{C})$$

that is spanned by the Conduché modules in each level.

Now, a standard argument shows the following:

**Proposition 5.12.** *The simplicial  $\infty$ -category  $\text{Cat}(\mathcal{C})$  is a double  $\infty$ -category.*

*Proof.* Let  $\text{Con}'_{n+1}(\mathcal{C}) \subset \text{Fun}((\Lambda/[n])^{\text{op}}, \mathcal{C})$  be the full subcategory spanned by those functors that satisfy the Segal condition as in [Definition 5.1](#). Then it readily follows from characterization (1) of [Proposition 5.3](#) and [Lemma 5.7](#) that the fully faithful left Kan extension functor

$$i_! : \text{Fun}((\Lambda/[n])^{\text{op}}, \mathcal{C}) \rightarrow \text{Fun}((\Delta/[n])^{\text{op}}, \mathcal{C})$$

along  $i : (\Lambda/[n])^{\text{op}} \rightarrow (\Delta/[n])^{\text{op}}$  restricts to an equivalence between  $\text{Con}'_{n+1}(\mathcal{C})$  and  $\text{Con}_{n+1}(\mathcal{C})$ . Consequently, we see that it suffices to check that the canonical functor

$$p : \text{Con}'_{n+1}(\mathcal{C}) \rightarrow \text{Con}_1(\mathcal{C}) \times_{\text{Cat}(\mathcal{C})} \cdots \times_{\text{Cat}(\mathcal{C})} \text{Con}_1(\mathcal{C})$$

is an equivalence.

To this end, we note that this functor may be identified with the restriction of the left adjoint that sits in the adjunction

$$j^* : \text{Fun}((\Lambda/[n])^{\text{op}}, \mathcal{C}) \rightleftarrows \text{Fun}((\Lambda/[n])^{\text{cell,op}}, \mathcal{C}) : j_*$$

associated to *right* Kan extending along the inclusion

$$j : (\Lambda/[n])^{\text{cell,op}} \simeq (\Delta/[1])^{\text{op}} \cup_{\Delta^{\text{op}}} \cdots \cup_{\Delta^{\text{op}}} (\Delta/[1])^{\text{op}} \rightarrow (\Lambda/[n])^{\text{op}}$$

of the full subcategory of those cellular maps  $f : [m] \rightarrow [n]$  so that  $\text{im}(f) \subset \{i, i+1\}$  for some  $i$ . Let us write  $\text{Con}''_{n+1}(\mathcal{C}) \subset \text{Fun}((\Lambda/[n])^{\text{cell,op}}, \mathcal{C})$  for the full subcategory of those functors that satisfy the Segal condition. Then  $p$  may be identified with the restriction of  $j^*$  to  $\text{Con}'_{n+1}(\mathcal{C})$  and  $\text{Con}''_{n+1}(\mathcal{C})$ . Since  $j_*$  is fully faithful, we thus see that  $p$  is an equivalence if the following holds:

- (1) the left adjoint  $j_*$  carries  $\text{Con}''_{n+1}(\mathcal{C})$  into  $\text{Con}'_{n+1}(\mathcal{C})$ ,
- (2) if  $M \in \text{Con}'_{n+1}(\mathcal{C})$ , then  $M$  is right Kan extended along  $j$ , i.e. the unit  $M \rightarrow j_* j^* M$  is an equivalence,

in which case  $j_*$  restricts to an inverse of  $p$ .

Both (1) and (2) follow from the following computation. Let  $M : (\Lambda/[n])^{\text{cell,op}} \rightarrow \mathcal{C}$  be a functor and  $f : [m] \rightarrow [n]$  be a cellular map, then

$$j_* M(f) \simeq M(f|f^{-1}(\{\alpha, \alpha+1\})) \times_{M(f|f^{-1}(\{\alpha+1\}))} \cdots \times_{M(f|f^{-1}(\{\omega-1\}))} M(f|f^{-1}(\{\omega-1, \omega\})),$$

where  $\alpha := \min(f)$  and  $\omega := \max(f)$ . Consequently, if  $M$  satisfies the Segal condition then we get that

$$j_*M(f) \simeq M(f|\{0, 1\}) \times_{M(f|\{1\})} \cdots \times_{M(f|\{m-1\})} M(f|\{m-1, m\}). \quad \square$$

Finally, we record the following functoriality of the construction above. It follows immediately from [Definition 5.1](#) and [Proposition 5.3](#).

**Proposition 5.13.** *Suppose that  $u : \mathcal{C} \rightarrow \mathcal{D}$  is a functor to an  $\infty$ -category  $\mathcal{D}$  that also has properties (1)-(3) stated at the start of [Subsection 5.1](#). If  $u$  preserves pull-backs and geometric realizations, then post-composition with  $u$  gives rise to a double functor  $\mathbb{C}at(u) : \mathbb{C}at(\mathcal{C}) \rightarrow \mathbb{C}at(\mathcal{D})$ .*

**5.2. Completeness conditions in  $\infty$ -toposes.** Lurie has developed a theory of complete categorical objects with respect to a *distributor* in [\[Lur09b\]](#). In particular, if  $\mathcal{E}$  is an  $\infty$ -topos, this theory can be specialized to the trivial distributor  $\mathcal{E} \subset \mathcal{E}$ , yielding the notion of *complete categorical objects* in  $\mathcal{E}$ . Since we do not want to recall the general theory of distributors, we give a simpler definition of these objects. It is shown by Haugseng in [\[Hau18, Proposition 7.13\]](#) that this is equivalent to Lurie's definition. Throughout, we fix an  $\infty$ -topos  $\mathcal{E}$ .

**Definition 5.14.** A categorical object in  $\mathcal{C}$  in  $\mathcal{E}$  is called *complete* if the composition

$$\Delta^{\text{op}} \xrightarrow{\mathcal{C}} \mathcal{E} \xrightarrow{\text{Map}_{\mathcal{E}}(X, -)} \mathcal{S}$$

defines a complete Segal space for every object  $X \in \mathcal{E}$ . In this case, we will also call  $\mathcal{C}$  an *internal ( $\infty$ -)category in  $\mathcal{E}$* . We write

$$\text{Cat}_{\infty}(\mathcal{E}) \subset \text{Cat}(\mathcal{E})$$

for the full subcategory spanned by the complete objects.

More generally, a  $(n+1)$ -categorical module  $M$  in  $\mathcal{E}$  is said to be *complete* if for any vertex  $0 \leq i \leq n$ , the restriction  $(\{i\} \rightarrow [n])^*M$  is a complete categorical object in  $\mathcal{E}$ . We write

$$\text{CMod}_{n+1}(\mathcal{E}) \subset \text{Mod}_{n+1}(\mathcal{E}), \quad \text{CCon}_{n+1}(\mathcal{E}) \subset \text{Con}_{n+1}(\mathcal{E}),$$

for the full subcategories of, respectively, complete categorical and Conduché modules in  $\mathcal{E}$ .

**Construction 5.15.** By definition, the property of completeness for categorical modules is preserved under restricting along maps in  $\Delta$ . Moreover, a categorical  $(n+1)$ -module is complete if and only if its restriction along each inert map is complete. Consequently, the double  $\infty$ -category  $\mathbb{C}at(\mathcal{E})$  that we constructed in [Subsection 5.1](#), admits a sub double  $\infty$ -category

$$\text{Cat}_{\infty}(\mathcal{E})$$

that is levelwise spanned by the complete Conduché modules so that  $\mathbb{C}at_{\infty}(\mathcal{E})_n = \text{CCon}_{n+1}(\mathcal{E})$ .

**Proposition 5.16.** *The double  $\infty$ -category  $\mathbb{C}at_{\infty}(\mathcal{E})$  is an  $\infty$ -equipment.*

*Proof.* One may demonstrate in a way that is similar to the computation in the proof of [Proposition 4.11](#), that  $\text{Cat}_{\infty}(\mathcal{E})$  is vertically complete. If  $\mathcal{C}$  and  $\mathcal{D}$  are internal categories in  $\mathcal{E}$ , the vertical  $\infty$ -category  $\text{Vert}(\mathbb{C}at_{\infty}(\mathcal{E}))(\mathcal{C}, \mathcal{D})$  is given by the object that is denoted by  $\text{MAP}(\mathcal{C}, \mathcal{D})$  in [\[Hau18, Definition 7.12\]](#).



We will show that  $\mathbb{C}at_\infty(\mathcal{E})$  is an  $\infty$ -equipment by demonstrating that its source-target projection is a cartesian fibration. Note that we have a commutative square

$$\begin{array}{ccc} \mathbb{C}Mod_2(\mathcal{E}) & \longrightarrow & Mod_2(\mathcal{E}) \\ \downarrow & & \downarrow \\ \mathbb{C}at_\infty(\mathcal{E})^{\times 2} & \longrightarrow & Cat(\mathcal{E})^{\times 2}, \end{array}$$

where the vertical arrows are the source-target projections of the double  $\infty$ -categories  $\mathbb{C}at_\infty(\mathcal{E})$  and  $Cat(\mathcal{E})$ . By definition, this square is a pullback square. Thus, it suffices to show that source-target projection for  $Cat(\mathcal{E})$  is a cartesian fibration.

Since  $Cat(\mathcal{E}) \subset Fun(\Delta^{op}, \mathcal{E})$  is a reflective subcategory, it has all pullbacks. Hence, it is enough to show that  $Mod_2(\mathcal{E}) \rightarrow Cat(\mathcal{E})^{\times 2}$  admits a fully faithful right adjoint.<sup>3</sup> By construction, this projection is the restriction of the functor

$$(d_0^*, d_1^*) : Fun((\Delta/[1])^{op}, \mathcal{E}) \rightarrow Fun(\Delta^{op}, \mathcal{E})^{\times 2},$$

which admits a fully faithful right adjoint  $(-)\star(-)$  given by right Kan extension along  $(d_0)_! \sqcup (d_1)_! : \Delta^{op} \sqcup \Delta^{op} \rightarrow (\Delta/[1])^{op}$ . One may readily check that  $X \star Y$  is a categorical bimodule if  $X$  and  $Y$  are categorical objects in  $\mathcal{E}$ . Hence,  $(-)\star(-)$  restricts to a fully faithful right adjoint for the source-target projection of  $Cat(\mathcal{E})$ .  $\square$

The  $\infty$ -equipments of internal  $\infty$ -categories are functorial with respect to the following class of maps:

**Proposition 5.17.** *Suppose that  $u : \mathcal{E} \rightarrow \mathcal{F}$  is a functor to an  $\infty$ -topos  $\mathcal{F}$ . If  $u$  preserves pullbacks and geometric realizations, and has the property that for every arrow  $f : x \rightarrow y$ , the commutative square*

$$\begin{array}{ccc} u(x \times x) & \xrightarrow{u(f \times f)} & u(y \times y) \\ \downarrow & & \downarrow \\ u(x) \times u(x) & \xrightarrow{u(f) \times u(f)} & u(y) \times u(y) \end{array}$$

where the vertical arrows are given by the limit comparison maps, is a pullback square, then post-composition with  $u$  gives rise to a double functor

$$\mathbb{C}at_\infty(u) : \mathbb{C}at_\infty(\mathcal{E}) \rightarrow \mathbb{C}at_\infty(\mathcal{F}).$$

*Proof.* It follows from [Proposition 5.13](#) that  $u$  induces a double functor  $Cat(\mathcal{E}) \rightarrow Cat(\mathcal{F})$ . To check that it restricts to the desired double functor  $\mathbb{C}at_\infty(\mathcal{E}) \rightarrow \mathbb{C}at_\infty(\mathcal{F})$ , it suffices to check that post-composition with  $u$  preserves the completeness condition for categorical objects, and this readily follows from the same reasoning as in the proof of [\[Mar21, Lemma 3.3.1\]](#).  $\square$

We will use this observation in [\[Rui\]](#), to demonstrate that a geometric morphism  $f : \mathcal{E} \rightarrow \mathcal{F}$  between  $\infty$ -toposes gives rise to *lax right adjoint*  $\mathbb{C}at_\infty(\mathcal{E}) \rightarrow \mathbb{C}at_\infty(\mathcal{F})$  between their associated  $\infty$ -equipments.

<sup>3</sup>It is a general fact that any functor  $p : E \rightarrow \mathcal{C}$  whose target  $\mathcal{C}$  has all pullbacks, and that admits a fully faithful right adjoint, is a cartesian fibration.

Finally, we will show that the construction above recovers the  $\infty$ -equipment of  $\infty$ -categories when  $\mathcal{E}$  is given by the  $\infty$ -topos of spaces, as promised.

**Proposition 5.18.** *There is an equivalence*

$$\mathbb{C}\text{at}_\infty \xrightarrow{\simeq} \mathbb{C}\text{at}_\infty(\mathcal{S})$$

between  $\infty$ -equipments.

To construct a double functor between the two equipments, we will need the following technicalities. Let

$$\text{PSh} \rightarrow \text{Cat}_\infty$$

be the large cartesian fibration that classifies the functor  $\mathcal{C} \mapsto \text{PSh}(\mathcal{C})$ . If we write

$$\text{RFib} \subset \text{Cat}_\infty/[1]$$

for the full subcategory spanned by the right fibrations, then the functor

$$\text{ev}_1 : \text{RFib} \rightarrow \text{Cat}_\infty.$$

given by evaluating in 1, is a cartesian fibration since right fibrations are stable under pullback. Gepner, Haugseng and Nikolaus show that the unstraightening is (coherently) natural:

**Proposition 5.19** ([GHN17, Proposition A.31]). *There is a map between cartesian fibrations*

$$\begin{array}{ccc} \text{PSh} & \longrightarrow & \text{RFib}, \\ & \searrow & \swarrow \\ & \text{Cat}_\infty & \end{array}$$

which may be identified with the unstraightening functor fiberwise. Consequently, the top functor is an equivalence of  $\infty$ -categories.

We will write

$$\text{RFib}(\mathcal{C}) \subset \text{Cat}_\infty/\mathcal{C}$$

for the full subcategory spanned by the right fibrations, or equivalently, the fiber of  $\text{ev}_1$  above  $\mathcal{C}$ . Note that we have a forgetful functor  $\text{fgt} : \text{RFib} \rightarrow \text{Cat}_\infty$  induced by evaluating at 0, which restricts to a forgetful functor  $\text{fgt} : \text{RFib}(\mathcal{C}) \rightarrow \text{Cat}_\infty$ . The following argument also appears in [GHN17]:

**Lemma 5.20** ([GHN17, Corollary 9.8]). *Let  $\mathcal{C}$  be a (small)  $\infty$ -category. Then there is a pullback square of  $\infty$ -categories*

$$\begin{array}{ccc} \text{Fun}([1], \text{PSh}(\mathcal{C})) & \longrightarrow & \text{PSh} \\ \text{ev}_1 \downarrow & & \downarrow \\ \text{PSh}(\mathcal{C}) & \longrightarrow & \text{Cat}_\infty, \end{array}$$

where the bottom arrow is the composition

$$\text{PSh}(\mathcal{C}) \rightarrow \text{RFib}(\mathcal{C}) \xrightarrow{\text{fgt}} \text{Cat}_\infty,$$

so that the first arrow is given by the Grothendieck construction.

*Proof.* By the straightening theorem and its naturality, it suffices to show that we have a pullback square

$$\begin{array}{ccc} \text{Fun}([1], \text{RFib}(\mathcal{C})) & \longrightarrow & \text{RFib} \\ \text{ev}_1 \downarrow & & \downarrow \\ \text{RFib}(\mathcal{C}) & \xrightarrow{\text{fgt}} & \text{Cat}_\infty. \end{array}$$

One readily checks that we have a pullback square

$$\begin{array}{ccc} \text{Fun}([1], \text{Cat}_\infty/\mathcal{C}) & \xrightarrow{\text{Fun}([1], \text{fgt})} & \text{Fun}([1], \text{Cat}_\infty) \\ \text{ev}_1 \downarrow & & \downarrow \text{ev}_1 \\ \text{Cat}_\infty/\mathcal{C} & \xrightarrow{\text{fgt}} & \text{Cat}_\infty, \end{array}$$

which essentially follows from the equivalent presentation  $\text{Fun}([1], \text{Cat}_\infty/\mathcal{C}) \simeq \text{Fun}([2], \text{Cat}_\infty) \times_{\text{Fun}(\{2\}, \text{Cat}_\infty)} \{\mathcal{C}\}$ . From this, one may deduce that there is a pullback square

$$\begin{array}{ccc} \text{Fun}([1], \text{RFib}(\mathcal{C})) & \xrightarrow{\text{Fun}([1], \text{fgt})} & \text{RFib} \\ \text{ev}_1 \downarrow & & \downarrow \\ \text{RFib}(\mathcal{C}) & \xrightarrow{\text{fgt}} & \text{Cat}_\infty. \end{array}$$

Here we used that if we have a commutative triangle

$$\begin{array}{ccc} X & \longrightarrow & Y \\ & \searrow & \swarrow \\ & \mathcal{C} & \end{array}$$

where the slanted arrows are right fibrations, then  $X \rightarrow Y$  is a right fibration as well (working in the model of quasi-categories, this may be verified by taking suitable replacements, cf. [Lur09a, Corollary 2.2.3.14]).  $\square$

**Remark 5.21.** Let  $\mathcal{C}$  be a (small)  $\infty$ -category, and  $F \in \text{PSh}(\mathcal{C})$  a presheaf classified by a right fibration  $p : Y \rightarrow X$ . Then [Lemma 5.20](#) states that there is an equivalence

$$\text{PSh}(Y) \simeq \text{PSh}(\mathcal{C})/F$$

which is moreover natural in  $F$ . In particular, if  $F$  is represented by an object  $x \in \mathcal{C}$ , then we get that

$$\text{PSh}(\mathcal{C}/x) \simeq \text{PSh}(\mathcal{C})/F.$$

This has also been shown by Lurie using a different method [Lur09a, Corollary 5.1.6.12].

*Proof of Proposition 5.18.* The restriction of the Yoneda embedding  $\text{Cat}_\infty \rightarrow \text{PSh}(\Delta)$  gives rise to a map between cartesian fibrations

$$\begin{array}{ccc} \text{Fun}([1], \text{Cat}_\infty) & \longrightarrow & \text{Fun}([1], \text{PSh}(\Delta)) \times_{\text{PSh}(\Delta)} \text{Cat}_\infty \\ \text{ev}_1 \searrow & & \swarrow \text{ev}_1 \\ & \text{Cat}_\infty & \end{array}$$

since it preserves (in particular) pullbacks. We may now pull back along the twisted inclusion  $\text{tw} : \Delta \rightarrow \text{Cat}_\infty$  (see [Construction 4.9](#)), and pass to their straightenings, so that we obtain a natural transformation between the simplicial  $\infty$ -categories that the resulting cartesian fibrations classify. In light of [Lemma 5.20](#), this is a natural transformation

$$y : S_\bullet \rightarrow \text{PSh}(C_\bullet).$$

where  $S_\bullet$  is the simplicial  $\infty$ -category given by the composite

$$\Delta^{\text{op}} \xrightarrow{\text{tw}^{\text{op}}} \text{Cat}_\infty^{\text{op}} \xrightarrow{\text{Cat}_\infty/(-)} \widehat{\text{Cat}}_\infty$$

Level-wise,  $y_n$  may be identified with the functor

$$\text{Cat}_\infty/[n] \rightarrow \text{PSh}(\Delta/[n]) \simeq \text{PSh}(\Delta)/[n]$$

given by restricting the Yoneda embedding along the inclusion  $\Delta/[n] \rightarrow \text{Cat}_\infty/[n]$ , or equivalently, given by slicing the restricted Yoneda embedding along  $\Delta \rightarrow \text{Cat}_\infty$  over  $[n]$ . In light of [\[JT07\]](#), this factors through the equivalence

$$\text{Cat}_\infty/[n] \xrightarrow{\simeq} \text{CMod}_{n+1}(\mathcal{S}) \simeq \text{Cat}_\infty(\mathcal{S})/[n].$$

Thus  $y$  restricts to an equivalence

$$y : S_\bullet \rightarrow \text{CMod}_{\bullet+1}(\mathcal{S})$$

between simplicial  $\infty$ -categories. Now, we claim that  $y$  carries the Conduché fibrations to complete categorical modules that satisfy the Conduché condition. It suffices to check this in the case that  $n = 2$ , where the result follows from characterization (5) of [\[AFR18, Lemma 5.16\]](#). Note that there are no Conduché conditions in the levels  $n = 0, 1$ . Combining these observations and using the Segal conditions, we see that  $y$  restricts to the desired equivalence

$$y : \text{Cat}_\infty \rightarrow \text{Cat}_\infty(\mathcal{S}). \quad \square$$

## 6. FIRST NOTIONS OF FORMAL CATEGORY THEORY

In this section, we introduce and study the basic notions of category theory internal to an  $\infty$ -equipment. The insight that one may encode desirable universal properties via proarrows is central to developing formal category theories using the language of equipments. It is then the question of whether these proarrows are representable in the sense that they are companions or conjoints. Throughout this section, we will fix an  $\infty$ -equipment  $\mathcal{P}$ .

**6.1. Fully faithfulness and adjoints.** We will start by defining fully faithful arrows in an  $\infty$ -equipment, and give a characterization of adjoint pairs in vertical fragments of  $\infty$ -equipments via proarrows.

**Definition 6.1.** An arrow  $f : x \rightarrow y$  in  $\mathcal{P}$  is *fully faithful* if the horizontal identity 2-cell

$$\begin{array}{ccc} x & \xlongequal{\quad} & x \\ f \downarrow & \xlongequal{\quad} & \downarrow f \\ y & \xlongequal{\quad} & y \end{array}$$

is cartesian.

**Proposition 6.2.** *Any equivalence  $f : x \rightarrow y$  in  $\mathcal{P}$  is fully faithful.*

*Proof.* Note that  $f$  is classified by a map  $[1] \rightarrow \mathcal{P}_0$ , and we have to check that

$$\mathrm{id}_f : [1] \rightarrow \mathcal{P}_0 \xrightarrow{s_0} \mathcal{P}_1$$

is cartesian. This follows from the fact that the first arrow factors over the point.  $\square$

**Example 6.3.** If  $\mathcal{P} = \mathbf{Cat}_\infty$ , the above definition recovers the usual notion of fully faithful functors. To wit, suppose that  $f : \mathcal{C} \rightarrow \mathcal{D}$  is a functor between  $\infty$ -categories. Then the identity 2-cell  $\mathrm{id}_f$  factors (essentially) uniquely as

$$\begin{array}{ccc} \mathcal{C} & \xlongequal{\quad} & \mathcal{C} \\ \parallel & & \parallel \\ \mathcal{C} & \xrightarrow{f \mathcal{D}_f} & \mathcal{C} \\ f \downarrow & & \downarrow f \\ \mathcal{D} & \xlongequal{\quad} & \mathcal{D} \end{array}$$

so that the bottom 2-cell is cartesian. From the perspective of profunctors, the top comparison 2-cell is given by the canonical map  $\mathrm{Map}_{\mathcal{C}}(-, -) \rightarrow \mathrm{Map}_{\mathcal{D}}(f-, f-)$  in  $\mathrm{Prof}(\mathcal{C}, \mathcal{C})$ . By the definition above,  $f$  is fully faithful in  $\mathbf{Cat}_\infty$  if and only if this map is an equivalence in  $\mathrm{Prof}(\mathcal{C}, \mathcal{C})$ , hence the result follows.

**Proposition 6.4.** *Let  $f : x \rightarrow y$  be an arrow in  $\mathcal{P}$ . If  $f$  is fully faithful, then the pushforward map*

$$f_* : \mathrm{Vert}(\mathcal{P})(z, x) \rightarrow \mathrm{Vert}(\mathcal{P})(z, y)$$

*is fully faithful for all  $z \in \mathcal{P}$ .*

*Proof.* Using the companion embedding for  $\mathcal{P}$ , we obtain a commutative square

$$\begin{array}{ccc} \mathrm{Vert}(\mathcal{P})(z, x) & \xrightarrow{f_*} & \mathrm{Vert}(\mathcal{P})(z, y) \\ \downarrow & & \downarrow \\ \mathrm{Hor}(\mathcal{P})(z, x) & \xrightarrow{(y_f)_*} & \mathrm{Hor}(\mathcal{P})(z, y). \end{array}$$

Since the vertical arrows are fully faithful, it suffices to show that the bottom functor is fully faithful. In view of [Proposition 2.26](#) and [Proposition 3.4](#), this bottom functor is left adjoint to  $(f y)_*$ . Thus, this functor is fully faithful if (and only if) the unit for this adjunction is an equivalence. But this unit is induced by the pasting  $\eta$  of the proof of [Proposition 3.4](#). This pasting is an equivalence in  $\mathrm{Hor}(\mathcal{P})(x, x)$  since  $f$  is fully faithful, hence  $(y_f)_*$  must be fully faithful.  $\square$

**Proposition 6.5.** *Suppose that  $f : x \rightarrow y$  and  $g : y \rightarrow x$  are arrows in  $\mathcal{P}$ . Then  $(f, g)$  is an adjunction pair in  $\mathrm{Vert}(\mathcal{P})$  if and only if there exists an equivalence  $f y \simeq x_g$  in  $\mathrm{Hor}(\mathcal{P})(y, x)$ .*

*Proof.* The companion embedding  $\mathrm{Vert}(\mathcal{P}) \rightarrow \mathrm{Hor}(\mathcal{P})$  is a locally fully faithful map between 2-fold Segal spaces, thus  $(f, g)$  is an adjunction pair in  $\mathrm{Vert}(\mathcal{P})$  if and only if  $(y_f, x_g)$  is an adjunction pair in  $\mathrm{Hor}(\mathcal{P})$ . In light of [Proposition 3.4](#), the pair  $(y_f, f y)$  is always an adjunction pair in  $\mathrm{Hor}(\mathcal{P})$ . Thus by the essential uniqueness of adjoints [Proposition 2.24](#), we deduce that  $f$  and  $g$  are adjoint if and only if  $f y \simeq x_g$  in  $\mathrm{Hor}(\mathcal{P})(y, x)$ .  $\square$

**Corollary 6.6.** *An arrow  $f : x \rightarrow y$  in  $\mathcal{P}$  admits a right adjoint in  $\text{Vert}(\mathcal{P})$  if and only if  $f_y$  is a companion. Similarly, an arrow  $g : y \rightarrow x$  in  $\mathcal{P}$  admits a left adjoint in  $\text{Vert}(\mathcal{P})$  if and only if  $x_g$  is a conjoint.*

**Remark 6.7.** If we consider  $\mathcal{P} = \mathbb{C}\text{at}_\infty$  and view proarrows as correspondences, this recovers the fact that a functor  $f : \mathcal{C} \rightarrow \mathcal{D}$  is a left adjoint if and only if the cocartesian fibration that classifies  $f$  is also a cartesian fibration.

**Proposition 6.8.** *Suppose that  $f : x \rightarrow y$  and  $g : y \rightarrow x$  are arrows in  $\mathcal{P}$  so that  $(f, g)$  is an adjunction in  $\text{Vert}(\mathcal{P})$ . Then  $f$  is fully faithful if and only if the unit of this adjunction is an equivalence. Similarly,  $g$  is fully faithful if the counit is an equivalence.*

*Proof.* We only demonstrate the first assertion as the statement for counits is shown similarly. It follows from [Proposition 6.5](#) that  $x_g$  is the conjoint of  $f$ . Moreover, the unit for the adjunction  $(f, g)$  corresponds to the pasting

$$\begin{array}{ccccc} x & \xlongequal{\quad} & x & \xlongequal{\quad} & x \\ \parallel & & \downarrow f & & \parallel \\ x & \xrightarrow{y_f} & y & \xrightarrow{x_g} & x \end{array}$$

under the companion embedding (see the proof of [Proposition 3.4](#)). This unit fits in the diagram

$$\begin{array}{ccccc} x & \xlongequal{\quad} & x & \xlongequal{\quad} & x \\ \parallel & & \downarrow f & & \parallel \\ x & \xrightarrow{y_f} & y & \xrightarrow{x_g} & x \\ f \downarrow & & \parallel & & \downarrow f \\ y & \xlongequal{\quad} & y & \xlongequal{\quad} & y, \end{array}$$

which must paste to the horizontal identity 2-cell of  $f$  on account of the companion and conjoint identities. Also, the pasting of the bottom row is cartesian. Consequently, this shows that the unit  $x \rightarrow x_g f$  may be identified with the (essentially) unique factorization  $x \rightarrow_f y_f$  induced by the identity 2-cell  $\text{id}_f$ . Combining this observation with the fact that the companion embedding is locally fully faithful, we obtain the desired result.  $\square$

**6.2. Weighted colimits and limits.** Equipments support an internal notion of *weighted* colimits and limits. This concept finds its roots in enriched category theory as the suitable generalization of ordinary (co)limits and was introduced by Borceux and Kelly [[BK75](#)]. In the setting of  $\infty$ -categories, weighted (co)limits were introduced by Gepner, Haugseng and Nikolaus [[GHN17](#)], and Rovelli [[Rov21](#)], both using a different approach. It was later shown by Haugseng [[Hau22](#), Corollary 4.5] that the two definitions agree. In [[Hin21](#)], Hinich introduced a notion of weighted (co)limits for enriched  $\infty$ -categories.

We will need the following terminology:

**Definition 6.9.** Suppose that  $f : i \rightarrow x$  is an arrow in  $\mathcal{P}$ . Then there are the following notions of *proarrows of cones*:

- If  $W : j \rightarrow i$  is a proarrow of  $\mathcal{P}$ , then a proarrow  $C : x \rightarrow j$  that represents the presheaf

$$\mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(x,i)}(W_*(-), fx) : \mathrm{Hor}(\mathcal{P})(x, j)^{\mathrm{op}} \rightarrow \mathcal{S}$$

is called a *proarrow of  $W$ -cones under  $f$* .

- If  $W : i \rightarrow j$  is a proarrow of  $\mathcal{P}$ , then a proarrow  $C : j \rightarrow x$  that represents the presheaf

$$\mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(i,x)}(W^*(-), x_f) : \mathrm{Hor}(\mathcal{P})(j, x)^{\mathrm{op}} \rightarrow \mathcal{S}$$

is called a *proarrow of  $W$ -cones over  $f$* .

In this context, the proarrows  $W$  that appear above are also called *weights*.

**Remark 6.10.** Note that the choice of conjoint and companion of  $f$  in the definition above do not matter, as they are unique up to suitable equivalence. This also follows from the following observation. If we have a cartesian 2-cell

$$\begin{array}{ccc} a & \xrightarrow{G} & b \\ f \downarrow & & \downarrow g \\ c & \xrightarrow{F} & d, \end{array}$$

in  $\mathcal{P}$ , then we obtain a pullback square

$$\begin{array}{ccc} \mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(a,b)}(-, G) & \longrightarrow & \mathrm{Map}_{\mathcal{P}_1}(-, F) \\ \downarrow & & \downarrow \\ \{(f, g)\} & \longrightarrow & \mathrm{Map}_{\mathcal{P}_0}(c, d) \times \mathrm{Map}_{\mathcal{P}_0}(a, b) \end{array}$$

of presheaves  $(\mathcal{P}_1 \times_{\mathcal{P}_0^2} \{(a, b)\})^{\mathrm{op}} \simeq \mathrm{Hor}(\mathcal{P})(a, b)^{\mathrm{op}} \rightarrow \mathcal{S}$ .

**Remark 6.11.** If  $\mathcal{P}$  is horizontally closed then it admits all proarrows of cones.

**Example 6.12.** Suppose that  $\mathcal{P} = \mathcal{C}\mathrm{at}_\infty$  and let  $f : I \rightarrow \mathcal{C}$  be a functor. We may consider the *coconical weight*  $W : [0] \rightarrow I$ , which has the defining property that it is a conjoint. From the viewpoint of profunctors,  $W$  is given by the constant presheaf

$$I^{\mathrm{op}} \rightarrow \mathcal{S} : i \mapsto *.$$

In light of [Corollary 4.16](#), the proarrow of  $W$ -cones may be computed pointwise by an end:

$$\mathcal{C} \rightarrow \mathcal{S} : c \mapsto \int_{i \in I} \mathrm{Map}_{\mathcal{C}}(f(i), c) \simeq \mathrm{Map}_{\mathrm{Fun}(I, \mathcal{C})}(f, \mathrm{const}_c).$$

This retrieves the usual presheaf that carries  $c \in \mathcal{C}$  to the space of cones under  $f$  with apex  $c$ .

We may also verify this from the viewpoint of two-sided discrete fibrations. In light of Stevenson's result discussed in [Remark 4.17](#), the two-sided discrete fibration  ${}^W(f)\mathcal{C}$  of  $W$ -cones under  $f$  fits in the diagram

$$\begin{array}{ccccc} E & \longrightarrow & \mathrm{Fun}(I, f\mathcal{C}) & \longrightarrow & \mathrm{Fun}(I \times [1], \mathcal{C}) \\ {}^W(f)\mathcal{C} \downarrow & & \downarrow & & \downarrow (ev_1, ev_0) \\ \mathcal{C} & \longrightarrow & \mathrm{Fun}(I, \mathcal{C}) \times \mathrm{Fun}(I, I) & \xrightarrow{\mathrm{id} \times f_*} & \mathrm{Fun}(I, \mathcal{C}) \times \mathrm{Fun}(I, \mathcal{C}), \end{array}$$

where both squares are pullback squares. The left bottom functor is given by the product of the diagonal functor and the constant map that selects the identity functor on  $I$ . Since the outer square is again a pullback square, we deduce that the proarrow of  $W$ -cones under  $f$  is given by the usual projection  $f/\mathcal{C} \rightarrow \mathcal{C}$ .

With the terminology of proarrows of cones in place, may now define a notion of (co)limits:

**Definition 6.13.** Let  $f : i \rightarrow x$  be an arrow in  $\mathcal{P}$ . Then the *colimit of  $f$  weighted by a proarrow  $W : j \rightarrow i$*  is an arrow  $g : j \rightarrow x$  such that its conjoint  ${}_g x$  is a proarrow of  $W$ -cones under  $f$ .

Similarly, the *limit of  $f$  weighted by a proarrow  $W : i \rightarrow j$*  is an arrow  $g : j \rightarrow x$  such that the companion  $x_g$  is a proarrow of  $W$ -cones over  $f$ .

**Example 6.14.** Suppose that  $\mathcal{P} = \text{Cat}_\infty$ . Let  $f : I \rightarrow \mathcal{C}$  be a functor between  $\infty$ -categories. In light of [Example 6.12](#), we see that the notion of colimits of  $f$  weighted by the conical weight  $W : [0] \rightarrow I$ , coincides with the ordinary notion of colimits of  $f$ .

More generally, suppose that  $W : [0] \rightarrow I$  is a weight, not necessarily the conical weight. Viewed as a profunctor, such a weight is given by a presheaf  $W : I^{\text{op}} \rightarrow \mathcal{S}$ . From this perspective, the proarrow  $C$  of  $W$ -cones under  $f$  is then given by the end formula

$$C : \mathcal{C} \rightarrow \mathcal{S} : c \mapsto \int_{i \in I} \text{Map}_{\mathcal{S}}(W(i), \text{Map}_{\mathcal{C}}(f(i), c)).$$

An object  $x \in \mathcal{C}$ , equivalently, a functor  $x : [0] \rightarrow \mathcal{C}$ , is the  $W$ -weighted colimit of  $f$  if and only if  $C$  is represented by  $x$ . This coincides with the notion of weighted colimits of [\[GHN17\]](#) (precisely, the definition given in [\[Hau22, Section 4\]](#)).

From the perspective of two-sided discrete fibrations,  $W$  is given by a right fibration  $p : I' \rightarrow I$ . A similar computation as in [Example 6.12](#), shows that the proarrow  $C$  (as a two-sided discrete fibration) is given by the left fibration

$$fp/\mathcal{C} \rightarrow \mathcal{C}.$$

Hence, an object  $x \in \mathcal{C}$  is the  $W$ -weighted colimit of  $f$  if and only if it is the (ordinary) colimit of the diagram  $fp : I' \rightarrow \mathcal{C}$ . In view of [\[Rov21, Theorem 2.37\]](#), this coincides with Rovelli's definition of weighted colimits. In particular, we recover [\[Hau22, Theorem 1.2\]](#) and the fact that the definitions of weighted colimits by Rovelli and Gepner-Haugsgeng-Nikolaus agree [\[Hau22, Corollary 4.5\]](#).

The analogous statements for weighted limits are true as well.

**Proposition 6.15.** *Suppose that we have an adjunction*

$$u : x \rightleftarrows y : v$$

*in  $\text{Vert}(\mathcal{P})$ . Let  $f : i \rightarrow x$  be an arrow of  $\mathcal{P}$ . Then we have the following preservations of proarrows of cones:*

- (1) *If  $C : x \rightarrow j$  is a proarrow of cones under  $f$  weighted by  $W : j \rightarrow i$ , then  $C^u$  is a proarrow of  $W$ -cones under  $uf$ .*
- (2) *Similarly, if  $C : j \rightarrow x$  is a proarrow of cones over  $f$  weighted by  $W : i \rightarrow j$ , then  ${}_v C$  is a proarrow of  $W$ -cones over  $vf$ .*



*Proof.* We prove the first statement since the proof of the latter is similar. On account of [Proposition 6.5](#), we may identify  $C^u \simeq C \circ_u y \simeq C \circ x_v \simeq C_v$ . Hence, we have an equivalence

$$\mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(y,j)}(F, C^u) \simeq \mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(x,j)}(F^v, C).$$

natural in proarrows  $F : y \rightarrow j$  (see [Remark 3.34](#)). Since  $C$  is a proarrow of  $W$ -cones under  $f$ , we deduce that

$$\mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(y,j)}(F, C^u) \simeq \mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(x,i)}(W \circ F^v, f x) \simeq \mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(y,i)}(W \circ F, f x_v).$$

Note that  $f x_v \simeq f x x_v \simeq f x_u y \simeq_u f x$ . Thus we obtain the desired equivalence

$$\mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(y,j)}(F, C^u) \simeq \mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(y,i)}(W \circ F, u f x). \quad \square$$

**Corollary 6.16.** *Suppose that  $(u, v)$  is an adjunction in  $\mathrm{Vert}(\mathcal{P})$ . Then  $u$  preserves all weighted colimits and  $v$  preserves all weighted limits.*

**Proposition 6.17.** *Let  $x$  be an object of  $\mathcal{P}$  and suppose that  $W : j \rightarrow i$  is a weight. Consider the full subcategory*

$$A \subset \mathrm{Vert}(\mathcal{P})(i, x)$$

*spanned by those arrows that admit a  $W$ -weighted colimit. Then there exists a functor*

$$\mathrm{colim}^W : A \rightarrow \mathrm{Vert}(\mathcal{P})(j, x)$$

*that carries an arrow to its  $W$ -weighted colimit.*

*Proof.* Let us consider the composite functor

$$C : \mathrm{Vert}(\mathcal{P})(i, x)^{\mathrm{op}} \rightarrow \mathrm{Hor}(\mathcal{P})(x, i) \rightarrow \mathrm{PSh}(\mathrm{Hor}(\mathcal{P})(x, i)) \xrightarrow{(W_*)^*} \mathrm{PSh}(\mathrm{Hor}(\mathcal{P})(x, j)),$$

where the first arrow is the conjoint embedding and the second arrow is given by the Yoneda embedding, so that

$$C(f) = \mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(x,i)}(W_*(-), f x).$$

Note that there is a similarly defined fully faithful functor

$$i : \mathrm{Vert}(\mathcal{P})(j, x)^{\mathrm{op}} \rightarrow \mathrm{Hor}(\mathcal{P})(x, j) \rightarrow \mathrm{PSh}(\mathrm{Hor}(\mathcal{P})(x, j)),$$

given by the conjoint embedding followed by the Yoneda embedding, so that

$$i(g) = \mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(x,j)}(-, g x).$$

The defining feature of  $A$  is that it is the largest subcategory so the essential image of  $R|A^{\mathrm{op}}$  coincides with the essential image of  $i$ . Consequently,  $R$  factors through  $i$  so that we obtain the desired functor  $\mathrm{colim}^W$  (after applying the involution  $(-)^{\mathrm{op}}$ ).  $\square$

**Example 6.18.** Consider the  $\infty$ -equipment  $\widehat{\mathrm{Cat}}_\infty$  of large  $\infty$ -categories (see [Remark 4.18](#)). The proarrows that lie in  $\mathrm{Cat}_\infty \subset \widehat{\mathrm{Cat}}_\infty$ , will be called *small proarrows*. If  $\mathcal{C}$  is a cocomplete  $\infty$ -category, then it admits all colimits weighted by these small proarrows. To wit, suppose that  $f : I \rightarrow \mathcal{C}$  is a functor, and let  $W : J \rightarrow I$  be a small proarrow. Then both  $I$  and  $J$  are small  $\infty$ -categories, and as a profunctor,  $W$  is given by a functor  $I^{\mathrm{op}} \times J \rightarrow \mathcal{S}$ . Since  $\mathcal{C}$  is cocomplete, it is tensored over spaces, i.e. there exists a functor

$$- \otimes - : \mathcal{S} \times \mathcal{C} \rightarrow \mathcal{C}$$

so that

$$\mathrm{Map}_{\mathcal{C}}(K \otimes x, y) \simeq \mathrm{Map}_{\mathcal{S}}(K, \mathrm{Map}_{\mathcal{C}}(x, y)),$$

naturally for all  $x, y \in \mathcal{C}$  and (small) spaces  $K$ . Now, one readily verifies that the  $W$ -weighted colimit  $g : J \rightarrow \mathcal{C}$  of  $f$  exists, and is given by the coend

$$g(j) = \int^{i \in I} W(i, j) \otimes f(i).$$

Conversely, [Example 6.14](#) implies that any  $\infty$ -category that admits all colimit weighted by small proarrows, is cocomplete. Again, there are analogous statements for complete  $\infty$ -categories and weighted limits.

**6.3. Pointwise Kan extensions.** We have the following familiar terminology for (co)limits weighted by conjoinants and companions:

**Definition 6.19.** Let  $f : i \rightarrow x$  and  $w : i \rightarrow j$  be arrows of  $\mathcal{P}$ . Then a colimit of  $f$  weighted by the conjoint  ${}_w j$  is called the (*pointwise*) *left Kan extension* of the arrow  $f$  along  $w$ .

Similarly, a limit of  $f$  weighted by the companion  ${}_w j$  is called a (*pointwise*) *right Kan extension* of  $f$  along  $w$ .

**Construction 6.20.** Suppose that  $g : j \rightarrow x$  is the left Kan extension of  $f : i \rightarrow x$  along an arrow  $w : i \rightarrow j$  of  $\mathcal{P}$ . Then there exists a canonical 2-cell

$$\begin{array}{ccc} i & \xrightarrow{f} & x \\ w \downarrow & \Downarrow & \nearrow g \\ j & & \end{array}$$

in  $\mathrm{Vert}(\mathcal{P})$ . Namely, the conjoint  ${}_w j$  represents the presheaf

$$\mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(x, i)}(w j_w \circ (-), f x) : \mathrm{Hor}(\mathcal{P})(x, i)^{\mathrm{op}} \rightarrow \mathcal{S},$$

thus, the canonical comparison 2-cell  $\mathrm{id}_i \rightarrow w j_w$  gives rise to 2-cell

$${}_w j_w x \rightarrow f x$$

and by locally fully faithfulness of the conjoint embedding, this yields a 2-cell  $f \rightarrow g w$ .

There is a similar story for right Kan extensions. Namely, if  $g$  is now the right Kan extension of  $f$  along  $w$ , then there exists a canonical 2-cell

$$\begin{array}{ccc} i & \xrightarrow{f} & x \\ w \downarrow & \Leftarrow & \nearrow g \\ j & & \end{array}$$

in  $\mathrm{Vert}(\mathcal{P})$  which is constructed in the same manner.

**Corollary 6.21.** *Suppose that  $w : i \rightarrow j$  is a fully faithful arrow in  $\mathcal{P}$  and let  $f : i \rightarrow x$  be an arrow of  $\mathcal{P}$ . If  $g$  is the left Kan extension of  $f$  along  $w$ , then the canonical 2-cell  $f \rightarrow g w$  is invertible. Similarly, if  $g$  is the right Kan extension of  $f$  along  $w$ , then the canonical 2-cell  $g w \rightarrow f$  is invertible.*

The following may readily be deduced from [Proposition 6.17](#):

**Proposition 6.22.** *Suppose that  $w : i \rightarrow j$  is an arrow of  $\mathcal{P}$ . Let  $x$  be an object of  $\mathcal{P}$  and write  $A \subset \text{Vert}(\mathcal{P})(i, x)$  the full subcategory spanned by those arrows that admit a left Kan extension along  $w$ . Then there exists a partial left adjoint*

$$w_! : A \rightarrow \text{Vert}(\mathcal{P})(j, x)$$

to the restriction functor

$$w^* : \text{Vert}(\mathcal{P})(j, x) \rightarrow \text{Vert}(\mathcal{P})(i, x)$$

that carries each arrow in  $A$  to its left Kan extension. Moreover, the associated unit of this (partial) adjunction is the one constructed in [Construction 6.20](#). There is a similar statement for right Kan extensions.

**Remark 6.23.** Any  $(\infty, 2)$ -category  $\mathcal{X}$  supports a local notion of Kan extensions. Namely, if  $f : i \rightarrow x$  and  $w : i \rightarrow j$  are arrows of  $\mathcal{X}$ , then the *weak left Kan extension* of  $f$  along  $w$  is any arrow that represents the presheaf

$$\text{Map}_{\mathcal{X}(i, x)}(f, w^*(-)) : \mathcal{X}(j, x) \rightarrow \mathcal{S}.$$

This notion was explored earlier in the introduction of this article. In light of [Proposition 6.22](#), any (pointwise) Kan extension is a weak Kan extension. However, the reverse implication fails in general.

We conclude this subsection by studying *Beck-Chevalley conditions* for these pointwise Kan extensions. Central to this, is the following definition (cf. [\[RV22, Definition 9.2.2\]](#)):

**Definition 6.24.** A lax commutative square

$$\begin{array}{ccc} i & \xrightarrow{p} & k \\ w \downarrow & \swarrow & \downarrow v \\ j & \xrightarrow{q} & l \end{array}$$

in  $\text{Vert}(\mathcal{P})$  is said to be *exact* if the induced comparison 2-cell

$$p_! w \rightarrow v_! q$$

is an equivalence in  $\text{Hor}(\mathcal{P})(j, k)$ .

**Remark 6.25.** This terminology dates back to Hilton [\[Hil66\]](#), and was originally introduced in the context of abelian categories. It was later further categorified by Guitart [\[Gui80\]](#). In Section 4 of loc. cit., Guitart studies similar phenomena in the Yoneda structures of Street and Walters.

**Proposition 6.26.** *Suppose that we have a lax commutative square as in [Definition 6.24](#). Let  $f : k \rightarrow x$  be an arrow of  $\mathcal{P}$  that admits a left Kan extension  $v_! f : l \rightarrow x$  along  $v$ . Then the following is true:*

- (1) *if  $w_!(fp) : j \rightarrow x$  is a left Kan extension along of  $fp$  along  $w$ , then there exists a canonical comparison 2-cell*

$$w_!(fp) \rightarrow (v_! f)q,$$

- (2) *and if the square is exact, then the left Kan extension  $w_!(fp)$  exists and this comparison 2-cell is an equivalence.*

*Proof.* The existence of the comparison 2-cell of (1) follows from the universal property of weak Kan extensions (cf. [Remark 6.23](#)). This can also be constructed as follows. Note that the conjoint of  $(v_!f)q$  represents the presheaf

$$\mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(x,k)}(v_! \circ^q(-), fx) \simeq \mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(x,k)}(v_!q \circ(-), fx).$$

Hence, the canonical 2-cell  ${}^P i^w \rightarrow v_!q$  gives rise to a map between presheaves

$$\begin{aligned} \mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(x,j)}(-, (v_!f)q_x) &\rightarrow \mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(x,k)}({}^P i^w \circ(-), fx) \\ &\xrightarrow{\simeq} \mathrm{Map}_{\mathrm{Hor}(\mathcal{P})(x,i)}(w_!j \circ(-), f_p x). \end{aligned}$$

In the situation of (1), the presheaf on the right is represented by the conjoint of  $w_!(fp)$ . Hence, under the Yoneda and conjoint embeddings, this would give the comparison 2-cell. Moreover, (2) follows from these observations as well.  $\square$

**Corollary 6.27.** *Suppose that  $\mathcal{P}$  is an  $\infty$ -equipment with an exact square as in [Definition 6.24](#). Let  $x$  be an object of  $\mathcal{P}$ , and suppose that each arrow  $i \rightarrow x$  and  $j \rightarrow x$  in  $\mathcal{P}$  admit left Kan extensions along  $w$  and  $v$  respectively so that the Kan extension functors  $w_!$  and  $v_!$  of [Proposition 6.22](#) exist. Then the mate transformation*

$$w_!p^* \rightarrow w_!p^*v^*v_! \rightarrow w_!w^*q^*v_! \rightarrow q^*v_!$$

is an equivalence in  $\mathrm{Vert}(\mathcal{P})(k, x)$ .

**Example 6.28.** Consider a pullback square

$$\begin{array}{ccc} I & \xrightarrow{p} & K \\ w \downarrow & & \downarrow v \\ J & \xrightarrow{q} & L \end{array}$$

of  $\infty$ -categories. If  $q$  is smooth (see [[Lur09a](#), Definition 4.1.2.9]) or  $v$  is proper (see [[Cis19](#), Definition 4.4.1]), then this square is exact in  $\mathrm{Cat}_\infty$ . This may be deduced from [[Cis19](#), Proposition 6.4.3]

**Example 6.29.** A map  $f : x \rightarrow y$  in  $\mathcal{P}$  is fully faithful if and only if the commutative square

$$\begin{array}{ccc} x & \xlongequal{\quad} & x \\ \parallel & & \downarrow f \\ x & \xrightarrow{f} & y \end{array}$$

is exact.

**6.4. Initiality and finality.** Under an extra assumption on the  $\infty$ -equipment  $\mathcal{P}$ , we may define a notion of initial and final arrows in  $\mathcal{P}$ .

**Definition 6.30.** We call  $\mathcal{P}$  *pointed* if its vertical fragment  $\mathrm{Vert}(\mathcal{P})$  has a final object  $*$ , in the sense that the (unique) map

$$\mathrm{Vert}(\mathcal{P})(x, *) \rightarrow [0]$$

is an equivalence for all  $x \in \mathcal{P}$ .

Suppose that  $\mathcal{P}$  is pointed and  $i$  is an object of  $\mathcal{P}$ . Then the *coconical weight*  $\Delta^i$  and the *conical weight*  ${}^i\Delta$  are, respectively, the conjoint and companion of the essentially unique arrow  $i \rightarrow *$ . If the colimit (resp. limit) of a diagram  $f : i \rightarrow x$  weighted by the coconical weight (resp. conical weight) exists, then these are called *conical*, and we denote them by  $\mathrm{colim} f$  (resp.  $\mathrm{lim} f$ ).

**Remark 6.31.** In the context of the definition above, note that any conjoint in  $\text{Hor}(\mathcal{P})(*, i)$  or any companion in  $\text{Hor}(\mathcal{P})(i, *)$  is a coconical or conical weight, respectively.

**Example 6.32.** The  $\infty$ -equipment  $\mathbb{C}\text{at}_\infty$  is pointed. More generally, if  $\mathcal{E}$  is an  $\infty$ -topos then  $\mathbb{C}\text{at}_\infty(\mathcal{E})$  is pointed.

**Proposition 6.33.** *Suppose that  $\mathcal{P}$  is pointed and let  $f : i \rightarrow j$  be an arrow in  $\mathcal{P}$ . The following statements are equivalent:*

- (1) *there exists an equivalence  ${}^f\Delta^i \simeq \Delta^j$  in  $\text{Hor}(\mathcal{P})(*, j)$ ,*
- (2) *the commutative square*

$$\begin{array}{ccc} i & \xrightarrow{f} & j \\ \downarrow & & \downarrow \\ * & \xlongequal{\quad} & * \end{array}$$

*is exact.*

*Dually, the following statements are equivalent:*

- (1') *there exists an equivalence  ${}^i\Delta^f \simeq j\Delta$  in  $\text{Hor}(\mathcal{P})(j, *)$ ,*
- (2') *the commutative square*

$$\begin{array}{ccc} i & \longrightarrow & * \\ f \downarrow & & \parallel \\ j & \longrightarrow & * \end{array}$$

*is exact.*

*Proof.* Clearly, (2) implies (1). Conversely, if (1) holds, then any map  ${}^f\Delta^i \rightarrow \Delta^j$  is essentially unique since  $*$  is final in  $\text{Vert}(\mathcal{P})$  and the conjoint embedding is locally fully faithful. Thus (1) implies (2). The demonstration that (1') and (2') are equivalent is similar.  $\square$

**Definition 6.34.** Suppose that  $\mathcal{P}$  is pointed. Let  $f : i \rightarrow j$  be an arrow in  $\mathcal{P}$ . Then  $f$  is called *final* if one of the equivalent conditions (1) or (2) of [Proposition 6.33](#) holds. Dually,  $f$  is called *initial* if condition (1') or (2') of [Proposition 6.33](#) holds.

Using [Proposition 6.26](#), we now immediately obtain the following result:

**Proposition 6.35.** *Suppose that  $\mathcal{P}$  is pointed. Let  $f : i \rightarrow j$  and  $g : j \rightarrow x$  be arrows in  $\mathcal{P}$ . Suppose that  $f$  is final, then the colimit of  $g$  exists if and only if the colimit of  $gf$  exists, and in this case, the canonical map*

$$\text{colim } gf \rightarrow \text{colim } g$$

*is an equivalence.*

*Dually, if  $f$  is initial, then the limit of  $g$  exists if and only if the limit of  $gf$  exists and in this case, the canonical map*

$$\text{lim } g \rightarrow \text{lim } gf$$

*is an equivalence.*

Like in the case of final and initial functors between  $\infty$ -categories ([Lur09a, Proposition 4.1.1.3]), final and initial arrows in a pointed  $\infty$ -equipment have a cancellation property:

**Proposition 6.36.** *Suppose that  $f : i \rightarrow j$  and  $g : j \rightarrow k$  are arrows in a pointed  $\infty$ -equipment  $\mathcal{P}$ . If  $f$  is final (resp. initial), then the composite  $gf$  is final (resp. initial) if and only if  $g$  is final (resp. initial).*

*Proof.* If  $f$  is final, we have

$$gf \Delta^i \simeq g(f \Delta^i) \simeq g \Delta^j.$$

From this computation, the statement for finality follows immediately. The dual statement for initiality is shown similarly.  $\square$

We can also give an analogue of *Quillen's theorem A*, for so-called *strongly pointed equipments*:

**Definition 6.37.** Suppose that  $\mathcal{P}$  is pointed. Then  $\mathcal{P}$  is called *strongly pointed* if for any proarrow  $F : x \rightarrow y$ , the following two statements hold:

- (1)  $F$  is a companion if and only if for every arrow  $p : * \rightarrow x$ , the proarrow  $F_p$  is a companion,
- (2)  $F$  is a conjoint if and only if for every arrow  $p : * \rightarrow x$ , the proarrow  ${}_p F$  is a conjoint.

**Example 6.38.** The  $\infty$ -equipment  $\mathbb{C}at_\infty$  is strongly pointed. This follows from the Yoneda lemma. Indeed, from the perspective of profunctors, a proarrow  $F : \mathcal{C} \rightarrow \mathcal{D}$  is a companion if and only if the associated functor  $\mathcal{C} \rightarrow \text{PSh}(\mathcal{D})$  factors through the Yoneda embedding. Since the Yoneda embedding is fully faithful, one just has to check that for every object  $p \in \mathcal{C}$ , the evaluation in  $p$  (which corresponds to  $F_p$ ) is representable. A similar argument shows that condition (2) also holds.

This is not true in general: if  $\mathcal{E}$  is an arbitrary  $\infty$ -topos, then  $\mathbb{C}at_\infty(\mathcal{E})$  will *virtually never* be strongly pointed.

**Proposition 6.39.** *Suppose that  $\mathcal{P}$  is strongly pointed. Then an arrow  $f : i \rightarrow j$  in  $\mathcal{P}$  is final if and only if for any  $p : * \rightarrow j$ , we have*

$${}_p(f \Delta^i) \simeq *$$

in  $\text{Hor}(\mathcal{P})(*, *)$ .

*Proof.* This follows directly from [Remark 6.31](#) and the definition of strongly pointed equipments.  $\square$

**Example 6.40.** Take  $\mathcal{P} = \mathbb{C}at_\infty$ . Suppose that  $f : I \rightarrow J$  is a functor between  $\infty$ -categories. From the perspective of two-sided discrete fibrations, the coconical weight  $W := \Delta^I$  is given by the two-sided discrete fibration

$$[0] \leftarrow I = I.$$

Hence,  ${}^g W = J_g \circ W$  corresponds to the two-sided discrete fibration

$$[0] \leftarrow I \times_{\text{Fun}(\{1\}, J)} \text{Fun}([1], J) \rightarrow J.$$

Thus in light of the proposition above,  $f$  is final if and only if

$${}_p({}^g W) \simeq (I \times_J j/J)[(I \times_J j/J)^{-1}] \simeq *$$

in  $\text{Hor}(\text{Cat}_\infty)([0], [0]) \simeq \text{DFib}([0], [0]) \simeq \mathcal{S}$  for any  $j \in J$ . This is precisely the content of Quillen’s theorem A for  $\infty$ -categories (in this full generality, originally due to Joyal).

## REFERENCES

- [AF20] D. Ayala and J. Francis, *Fibrations of  $\infty$ -categories*, Higher Structures 4 (2020), no. 1, 168–265.
- [AFR18] D. Ayala, J. Francis, and N. Rozenblyum, *Factorization homology I: Higher categories*, Advances in Mathematics 333 (2018), 1042–1177.
- [Bar05] C. Barwick,  *$(\infty, n)$ -Cat as a closed model category*, Ph.D. thesis, University of Pennsylvania, 2005.
- [BDG<sup>+</sup>16] C. Barwick, E. Dotto, S. Glasman, D. Nardin, and J. Shah, *Parametrized higher category theory and higher algebra: Exposé I*, arXiv:1608.03657, 2016.
- [BK75] F. Borceux and G. M. Kelly, *A notion of limit for enriched categories*, Bull. Aust. Math. Soc. 12 (1975), no. 1, 49–72.
- [BR13] J. Bergner and C. Rezk, *Comparison of models for  $(\infty, n)$ -categories I*, Geometry and Topology 17 (2013), no. 4, 2163–2202.
- [BR20] ———, *Comparison of models for  $(\infty, n)$ -categories II*, Journal of Topology 13 (2020), no. 4, 1554–1581.
- [BSP21] C. Barwick and C. Schommer-Pries, *On the unicity of the homotopy theory of higher categories*, J. Amer. Math. Soc. 34 (2021), no. 4, 1011–1058.
- [Cam15] T. Champion, *What is the point of pointwise Kan extensions?*, MathOverflow question, 2015, <https://mathoverflow.net/questions/220246/what-is-the-point-of-pointwise-kan-extensions>.
- [Cis19] D. Cisinski, *Higher categories and homotopical algebra*, Cambridge Studies in Advanced Mathematics, vol. 180, Cambridge University Press, 2019.
- [CM23] T. Champion and Y. Maehara, *A model-independent Gray tensor product for  $(\infty, 2)$ -categories*, arXiv:2304.05965, 2023.
- [Ehr63] C. Ehresmann, *Catégories structurées III*, Cah. Topol. Géom. Différ. Catég. 5 (1963), 1–21.
- [GHN17] D. Gepner, R. Haugseng, and T. Nikolaus, *Lax colimits and free fibrations in  $\infty$ -categories*, Documenta Mathematica 22 (2017), 1225–1266.
- [GP04] M. Grandis and R. Paré, *Adjoint for double categories*, Cah. Topol. Géom. Différ. Catég. 45 (2004), no. 3, 193–240.
- [Gra74] J. Gray, *Formal category theory: Adjointness for 2-categories*, Lecture Notes in Mathematics, vol. 391, Springer-Verlag, 1974.
- [Gui80] R. Guitart, *Relations et carrés exacts*, Ann. Sci. Math. Québec 4 (1980), no. 2, 103–125.
- [Hau13] R. Haugseng, *Weakly enriched higher categories*, Ph.D. thesis, Massachusetts Institute of Technology, 2013.
- [Hau16] ———, *Bimodules and natural transformations for enriched  $\infty$ -categories*, Homology, Homotopy and Applications 18 (2016), no. 1, 71–98.
- [Hau17] ———, *The higher Morita category of  $E_n$ -algebras*, Geometry & Topology 21 (2017), no. 3, 1631–1730.
- [Hau18] ———, *Iterated spans and classical topological field theories*, Mathematische Zeitschrift 289 (2018), no. 3, 1427–1488.
- [Hau22] ———, *On (co)ends in  $\infty$ -categories*, J. Pure Appl. Algebra 226 (2022), no. 2, 106819.
- [HHLN21] R. Haugseng, F. Hebestreit, S. Linskens, and J. Nuiten, *Lax monoidal adjunctions, two-variable fibrations and the calculus of mates*, arXiv:2011.08808, 2021.
- [Hil66] P. Hilton, *Correspondences and exact squares*, Proc. Conf. Categorical Algebra, La Jolla, Springer-Verlag, 1966, pp. 254–271.
- [Hin21] V. Hinich, *Colimits in enriched  $\infty$ -categories and Day convolution*, arXiv:2101.09538, 2021.
- [Joh02] P. T. Johnstone, *Sketches of an elephant: A topos theory compendium, vol. 1*, Oxford Logic Guides, Clarendon Press, 2002.

- [JT07] A. Joyal and M. Tierney, *Quasi-categories vs Segal spaces*, Categories in algebra, geometry and mathematical physics, Contemp. Math., vol. 431, Amer. Math. Soc., 2007, pp. 277–326.
- [Lac02] S. Lack, *A Quillen model structure for 2-categories*, K-Theory (2002), no. 2, 171–205.
- [Lor18] F. Loregian, *A Fubini rule for  $\infty$ -coends*, MPIM Preprints, preprint no. 68, 2018.
- [Lur09a] J. Lurie, *Higher topos theory*, Annals of Mathematics Studies, vol. 170, Princeton University Press, 2009.
- [Lur09b] ———,  *$(\infty, 2)$ -Categories and the Goodwillie calculus I*, arXiv:0905.0462, 2009.
- [Lur17] ———, *Higher algebra*, 2017, <https://www.math.ias.edu/~lurie/papers/HA.pdf>.
- [Mar21] L. Martini, *Yoneda’s lemma for internal higher categories*, arXiv:2103.17141, 2021.
- [Mos20] L. Moser, *A double  $(\infty, 1)$ -categorical nerve for double categories*, arXiv:2007.01848, 2020.
- [MT08] P-A. Melliès and N. Tabareau, *Free models of T-algebraic theories computed as Kan extensions*, hal-00339331, 2008.
- [MW21] L. Martini and S. Wolf, *Limits and colimits in internal higher category theory*, arXiv:2111.14495, 2021.
- [ORR] V. Ozornova, M. Rovelli, and J. Ruit, *The square-companion adjunction*, tentative title, in preparation.
- [Ras18] N. Rasekh, *Complete Segal objects*, arXiv:1805.03561, 2018.
- [Rez01] C. Rezk, *A model for the homotopy theory of homotopy theory*, Trans. Amer. Math. Soc. **353** (2001), no. 3, 973–1007.
- [Rov21] M. Rovelli, *Weighted limits in an  $(\infty, 1)$ -category*, Appl. Categ. Structures **29** (2021), no. 6, 1019–1062.
- [Rui] J. Ruit, *Formal category theory in  $\infty$ -equipments II*, in preparation.
- [Rui22] ———, *A pasting theorem for iterated Segal spaces*, arXiv:2210.04549, 2022.
- [RV16] E. Riehl and D. Verity, *Homotopy coherent adjunctions and the formal theory of monads*, Advances in Mathematics **286** (2016), 802–888.
- [RV22] ———, *Elements of  $\infty$ -category theory*, Cambridge Studies in Advanced Mathematics, vol. 194, Cambridge University Press, 2022.
- [Sha18] J. Shah, *Parametrized higher category theory*, arXiv:1809.05892, 2018.
- [Shu08] M. Shulman, *Framed bicategories and monoidal fibrations*, Theory Appl. Categ. **20** (2008), no. 18, 650–738.
- [Shu13] ———, *Enriched indexed categories*, Theory Appl. Categ. **28** (2013), 616–696.
- [Ste18] D. Stevenson, *Model structures for correspondences and bifibrations*, arXiv:1807.08226, 2018.
- [SW78] R. Street and R. Walters, *Yoneda structures on 2-categories*, Journal of Algebra **50** (1978), no. 2, 350–379.
- [Ver92] D. Verity, *Enriched categories, internal categories and change of base*, Ph.D. thesis, University of Cambridge, 1992.
- [Woo82] R.J. Wood, *Abstract proarrows I*, Cah. Topol. Géom. Différ. Catég. **23** (1982), no. 3, 279–290.

MATHEMATISCH INSTITUUT, UNIVERSITEIT UTRECHT, THE NETHERLANDS

Email address: j.c.ruit@uu.nl