Spectrally-Corrected and Regularized Global Minimum Variance Portfolio for Spiked Model

Hua Li^{a,*}, Jiafu Huang^a

^aSchool of Science, Chang Chun University, China

Abstract

Considering the shortcomings of the traditional sample covariance matrix estimation, this paper proposes an improved global minimum variance portfolio model and named spectral corrected and regularized global minimum variance portfolio (SCRGMVP), which is better than the traditional risk model. The key of this method is that under the assumption that the population covariance matrix follows the spiked model and the method combines the design idea of the sample spectrally-corrected covariance matrix and regularized. The simulation of real and synthetic data shows that our method is not only better than the performance of traditional sample covariance matrix estimation (SCME), Ledio-Wolf's shrinkage estimation (SHRE), weighteid shrinkage estimation (WSHRE) and simple spectral correction estimation (SCE), but also has lower computational complexity.

Keywords: Global minimum variance, Spiked model, High-dimensional covariance matrix, Random matrix theory, Spectrally-corrected method, Regularized technology

1. Introduction

Due to the development of the financial market, investment funds have increased significantly. Therefore, investors must know the proportion of the portfolio to maximize benefits and reduce the risks associated with investment decisions. The mean-variance portfolio proposed by Markowitz [1, 2] focuses primarily on determining the

^{*}Corresponding author

Email address: lihua@ccu.edu.cn (Hua Li)

optimal portfolio weight, i.e. the proportion of wealth invested in financially risky assets. It is one of the key components of capital market theory in dealing with financial investment decisions. By this theory, investors can construct an optimal portfolio by minimizing the portfolio variance at a given expected return or maximizing the portfolio return at a given portfolio risk. One of Markowitz's ideas was the minimization of the portfolio variance subject to the budget constraint, which is called the global minimum variance portfolio(GMVP); see arguments in [3, 4]. The GMVP is the most commonly used and efficient investment decision by researchers in the financial field. This portfolio has the smallest variance of all optimal portfolios as the solution to the Markowitz mean-variance optimization problem.

First, we would like to recall the GMVP. We have a financial market consisting of M risky assets, and let \mathbf{y}_t be the M-dimensional time series of returns at time $t = 1, \dots, n$. Thus, we have the following random vector:

$$\mathbf{y}_t = (y_{1t}, y_{2t}, \cdots, y_{Mt})^T.$$

where $\mathbf{y}_t \sim N(\boldsymbol{\mu}_t, \boldsymbol{\Sigma}_t)$. Then, we typically model the following vector stochastic processes based on expected and unexpected reward contributions:

$$\mathbf{y}_t = \boldsymbol{\mu}_t + \boldsymbol{\Sigma}_t^{\frac{1}{2}} \mathbf{x}_t. \tag{1.1}$$

where $\boldsymbol{\mu}_t$ and $\boldsymbol{\Sigma}_t = (\sigma_{ij})$ are the expected value and covariance matrix of the asset returns over an investment period, and \mathbf{x}_t is a random vector with independent and identically distributed entries having mean zero and variance one, we care about the weight corresponding to the minimum variance portfolio. Let $\mathbf{w}_t = (w_{1t}, w_{2t}, ..., w_{Mt})^T$ denotes the vector of portfolio weights, the expected value and covariance matrix of the portfolio are $E(r_t) = \mathbf{w}_t^T \boldsymbol{\mu}_t$, $Var(r_t) = \mathbf{w}_t^T \boldsymbol{\Sigma}_t \mathbf{w}_t$ respectively, where the $\mathbf{r}_t = (r_{1t}, r_{2t}, ..., r_{Mt})^T$ is random returns. During a single investment period in GMVP, under the constraint of $\mathbf{w}^T \mathbf{1}_M = 1$, the risk of the minimum variance portfolio is expressed as $\sigma_M^2(\mathbf{w}) = \mathbf{w}^T \boldsymbol{\Sigma} \mathbf{w}$. Accordingly, the global minimum variance portfolio selection problem can be formulated as the quadratic optimization problem with liner constraints, which is given by :

min:
$$\sigma_M^2(\mathbf{w}) = \mathbf{w}^T \Sigma \mathbf{w},$$

s.t.
$$\mathbf{w}^T \mathbf{1}_M = 1$$

where $\mathbf{1}_M = (1, ..., 1)^T$ stands for M dimensional vector of ones, and **w** denotes the vector of portfolio weights. The solution to the former optimization problem is given by:

$$\mathbf{w}_{GMVP} = \frac{\boldsymbol{\Sigma}^{-1} \mathbf{1}_M}{\mathbf{1}_M^T \boldsymbol{\Sigma}^{-1} \mathbf{1}_M}.$$
 (1.2)

The global minimum variance portfolio 1.2 has the smallest variance of all portfolios. Then, the global minimum variance can be given by:

$$\sigma^2(\mathbf{w}_{GMVP}) = \frac{1}{\mathbf{1}_M^T \boldsymbol{\Sigma}^{-1} \mathbf{1}_M}.$$
 (1.3)

The weights of GMVP have several desirable properties and have good applicability in practice. However, the population covariance matrix Σ is unknown and so it must be estimated using historical data of the asset returns. The quality of the estimator Σ has a strong influence on GMVP. Therefore, the estimation of GMVP is closely related to the estimation of the population covariance matrix for asset returns.

Traditional estimators are a commonly used method for the estimation of the GMVP (1.2). The traditional method is to replace the population covariance matrix Σ with a sample covariance **S**.

$$\mathbf{S} = \frac{1}{n-1} \sum_{i=1}^{n} (\mathbf{y}_i - \overline{\mathbf{y}}) (\mathbf{y}_i - \overline{\mathbf{y}})^T, \qquad (1.4)$$

where $\overline{\mathbf{y}} = \frac{1}{n} \sum_{i=1}^{n} y_i$. Then, the classic estimator of the \mathbf{w}_{GMVP} can be expressed as:

$$\hat{\mathbf{w}}_S = \frac{\mathbf{S}^{-1}\mathbf{1}_M}{\mathbf{1}_M^T\mathbf{S}^{-1}\mathbf{1}_M}.$$

In the classical asymptotic analysis, it is generally assumed that the number M of the investment asset is fixed and the sample size n is increased. In this case, the sample estimator is a consistent estimator and is often called standard asymptotics [5, 6, 7]. However, when both the asset dimension M and the sample size n tend to infinity, or more specifically, M and n have a considerable size (M/n = J > 0) classical estimator effect is not ideal, and even the consistency of the estimator is not guaranteed. Thus, it is inappropriate to use the sample covariance matrix to build a high-dimensional portfolio. To improve the performance of investment portfolios, two different methods are very popular. The first method is to directly constrain [4, 8, 9, 10] or shrink [11, 12] the portfolio weights. The second method is to use the improved covariance matrix estimator to constrain or shrink the portfolio weights, such as [13, 14]. Both dimension M and sample size n tend to infinity, which is known as high-dimensional asymptotics in [15, 16]. In this case, new asymptotic techniques must be applied for derivations. Therefore, many articles have studied the characteristics of high-dimensional global minimum variance investment portfolios, such as [17, 18, 8].

There are many articles provide different methods to improve the estimation of high-dimensional covariance matrix, which can be divided into two categories. The first category is based on additional knowledge in the estimation process, such as sparseness, graph models, and factor models [19, 20, 21, 22]. The second category is the spectrum of corrected sample covariance matrices, such as Ledoit and Wolf's (linear and nonlinear) shrinkage estimates [23, 24, 25] and other shrinkage estimators for covariance matrices [26]. In this paper, we proposed method belongs to the second category, and it integrates the design ideas of the sample spectrally-corrected covariance matrix [27] and the regularized method [28] to improve the GMVP estimation in high-dimensional settings. The details are given in the following sections.

In this paper, an extremely important and widely used model is involved, which is known as the spiked model in [29]. The model assumes that the population covariance matrix has only a fixed number of eigenvalues that are not equal to one, and all other eigenvalues are equal to one, and names those eigenvalues that are not one as "spikes". This model has been used in many real applications, we learned about mathematical finance [30], EEG signals [31, 32], and data analysis [33]. The most important result of the spiked model is the study of eigenvalues of the sample covariance matrix, such as [34, 35, 36, 37]. According to the related theory of the spiked covariance model, the population covariance matrix can be estimated by a class of sample covariance matrix is written as a finite rank perturbation of a scaled identity matrix. The principal eigenvectors of the sample covariance matrix give the direction of the low-rank perturbation,

correct the corresponding eigenvalues to one, and regularized by adding appropriate parameters.

According to the random matrix theory, derive the approximation of the portfolio variance, the regularization parameters are some of the parameters we choose to minimize the approximation of the portfolio variance. In this way, we not only preserve the structure of the spiked model, but also reduces the number of design parameters.

The rest of this paper is organized as follows: Section 2 shows the spectrallycorrected and regularized estimates of the covariance matrix. In section 3, provide a consistent estimate of our proposed method (SCRGMVP) and describe the optimal parameters. The performance of our method is studied in section 4 using synthetic and real simulations before concluding in section 5.

2. SCRGMVP under Spiked Model

In this paper, we propose a particular form of the covariance matrix, wherein Σ takes the following form:

$$\boldsymbol{\Sigma} = \sigma^2 \left(\mathbf{I}_M + \sum_{j \in \mathbb{I}_1} \lambda_j \mathbf{v}_j \mathbf{v}_j^T + \sum_{j \in \mathbb{I}_2} \lambda_j \mathbf{v}_j \mathbf{v}_j^T \right).$$
(2.1)

where $\sigma^2 > 0$, $\mathbb{I}_1 = \{1, \dots, r_1\}$, $\mathbb{I}_2 = \{-r_2, \dots, -1\}$, $r = r_1 + r_2$, $\lambda_1 \ge \dots \ge \lambda_{r_1} > 0 > \lambda_{-r_2} \ge \dots \ge \lambda_{-1} > -1$ and $\mathbf{v}_1, \dots, \mathbf{v}_{r_1}, \mathbf{v}_{-r_2}, \dots, \mathbf{v}_{-1}$ are orthonormal. This is the spiked model, which is used in many real applications, such as mathematical finance, EEG signals, and mathematical analysis. In practice, we can estimate parameters through algorithms in many relevant literature, such as [38, 39, 34, 40]. Therefore, we can assume that the parameters σ^2 , r_1 , r_2 , and $\lambda_i (i \in \mathbb{I})$ are completely known.

Based on 2.1, we consider a class of sample covariance matrix estimators that follow the same spike model, that is, the sample covariance matrix is written in the finite rank perturbation form of the identity matrix. First, we start from the eigendecomposition of the sample covariance matrix:

$$\mathbf{S} = \sum_{i=1}^{M} s_{j} \mathbf{u}_{j} \mathbf{u}_{j}^{T}$$
(2.2)

with s_i being the *j*-th largest eigenvalue and \mathbf{u}_i its corresponding eigenvector, and correcting s_i to the corresponding one of Σ as follows:

1

$$\hat{\boldsymbol{\Sigma}} = \sigma^2 \left(\mathbf{I}_M + \sum_{j \in \mathbb{I}_1} \lambda_j \mathbf{u}_j \mathbf{u}_j^T + \sum_{j \in \mathbb{I}_2} \lambda_j \mathbf{u}_j \mathbf{u}_j^T \right).$$
(2.3)

Comparing 2.1 and 2.3, we know that $\hat{\Sigma}$ is the same as Σ for the spiked model except for the spiked eigenvectors. In this way, the structure of Σ is preserved as much as possible. However, it is well known that $\hat{\Sigma}$ is a biased estimate of the population covariance matrix Σ . Accordingly, it is necessary to add regularization parameters to the sample spiked covariance matrix. So we can get the estimation of the global minimum variance portfolio weight and it is expressed by:

$$\sqrt{M}\hat{\mathbf{w}}_{SCRGMVP} = \frac{\widetilde{\mathbf{C}}\frac{\mathbf{1}_{M}}{\sqrt{M}}}{\frac{\mathbf{1}_{M}^{T}}{\sqrt{M}}\widetilde{\mathbf{C}}\frac{\mathbf{1}_{M}}{\sqrt{M}}},$$
(2.4)

where

$$\widetilde{\mathbf{C}} = \left(\mathbf{I}_M + \gamma_1 \sum_{j \in \mathbb{I}_1} \lambda_j \mathbf{u}_j \mathbf{u}_j^T + \gamma_2 \sum_{j \in \mathbb{I}_2} \lambda_j \mathbf{u}_j \mathbf{u}_j^T\right)^{-1}.$$
(2.5)

where γ_1 and γ_2 are the regularization parameters that need to be optimized. Therefore, the spectrally-corrected and regularized global minimum variance portfolio risk $(\sigma_M^2(\hat{\mathbf{w}}_{SCRGMVP}))$ is expressed as:

$$M\sigma_{M}^{2}(\hat{\mathbf{w}}_{SCRGMVP}) = \frac{\frac{\mathbf{1}_{M}^{T}}{\sqrt{M}}\widetilde{\mathbf{C}}\Sigma\widetilde{\mathbf{C}}\frac{\mathbf{1}_{M}}{\sqrt{M}}}{\left(\frac{\mathbf{1}_{M}^{T}}{\sqrt{M}}\widetilde{\mathbf{C}}\frac{\mathbf{1}_{M}}{\sqrt{M}}\right)^{2}} = \frac{\mathbf{Y}(\widetilde{\mathbf{C}},\Sigma)}{\left(\mathbf{X}(\widetilde{\mathbf{C}})\right)^{2}},$$
(2.6)

in which

$$\mathbf{Y}\left(\widetilde{\mathbf{C}}, \boldsymbol{\Sigma}\right) = \frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \widetilde{\mathbf{C}} \boldsymbol{\Sigma} \widetilde{\mathbf{C}} \frac{\mathbf{1}_{M}}{\sqrt{M}}, \qquad (2.7)$$

$$\mathbf{X}\left(\widetilde{\mathbf{C}}\right) = \frac{\mathbf{1}_{M}^{T}}{\sqrt{M}}\widetilde{\mathbf{C}}\frac{\mathbf{1}_{M}}{\sqrt{M}}.$$
(2.8)

3. Consistent Estimator of the Portfolio Risk and Parameter Optimization

In this section, the main task is to select an optimal set of parameters $\gamma = \{\gamma_1, \gamma_2\}$ to minimize the risk of SCRGMVP:

$$\gamma^* = \underset{\gamma>0}{\arg\min} \left(M \sigma_{SCRGMVP}^2(\gamma) \right)$$
(3.1)

where $\gamma^* = \{\gamma_1^*, \gamma_2^*\}$. It is usually not possible to obtain an exact optimal solution γ for 3.1. To solve this problem, we need to approximate the risk of SCRGMVP in the asymptotic growth state defined in the following assumption based on the results of random matrix theory. The parameters corresponding to the minimum approximation are the theoretical optimal parameters γ_1^*, γ_2^* .

Assumption 3.1. $M, n \to \infty$ and the following limits exist: $\frac{M}{n} \to J > 0$, and $\frac{M}{n} \to J < 1$.

Assumption 3.1 is usually a key assumption under high-dimensional random theory.

Assumption 3.2. r_1 and r_2 are fixed and $\lambda_1 > \cdots > \lambda_{r_1} > \sqrt{J} > 0 > -\sqrt{J} > \lambda_{-r_2} > \cdots > \lambda_{-1} > -1$, independently of M and $r = r_1 + r_2$.

Assumption 3.2 is the basis for our analysis results, ensuring a one-to-one mapping between sample eigenvalues s_j and unknown population eigenvalues λ_j . In fact, when $\lambda_j > \sqrt{J}(j \in \mathbb{I}_1)$ or $-1 < \lambda_j < -\sqrt{J}(j \in \mathbb{I}_2)$, λ_j can be consistently estimated using its corresponding s_j for $j \in \mathbb{I}$. Otherwise, the relation between s_j and λ_j no longer holds. *Assumption* 3.3. $\|\mu\|$ has a bounded Euclidean norm, that is $\|\mu\| = O(1)$. *Assumption* 3.4. The spectral norm of Σ is bounded, that is $\|\Sigma\| = O(1)$.

Theorem 1. Under Assumptions 3.1 to 3.4, we have

$$\mathbf{X}\left(\widetilde{\mathbf{C}}\right) - \overline{X}(\gamma_1, \gamma_2) \xrightarrow{a.s.} 0,$$
$$\mathbf{Y}\left(\widetilde{\mathbf{C}}, \mathbf{\Sigma}\right) - \overline{Y}(\gamma_1, \gamma_2) \xrightarrow{a.s.} 0$$

where

$$\overline{X}(\gamma_1, \gamma_2) = 1 - \sum_{i=1,2} \sum_{j \in \mathbb{I}} a_j b_j \gamma_{i,j} \delta_{i,j}$$
(3.2)

$$\overline{Y}(\gamma_1, \gamma_2) = 1 + \sum_{i=1,2} \sum_{j \in \mathbb{I}} \lambda_j b_j \delta_{i,j} - 2 \sum_{i=1,2} \sum_{j \in \mathbb{I}} \left(\lambda_j + 1\right) a_j b_j \gamma_{i,j} \delta_{i,j} + \sum_{i=1,2} \sum_{j \in \mathbb{I}} a_j b_j \left(\lambda_j a_j + 1\right) \gamma_{i,j}^2 \delta_{i,j}$$
(3.3)

in which

$$\gamma_{i,j} = \frac{\gamma_i \lambda_j}{1 + \gamma_i \lambda_j}, \ a_j = \frac{\lambda_j^2 - J}{\lambda_j (\lambda_j + J)}, \ b_j = \left(\frac{\mathbf{1}_M^T}{\sqrt{M}} \mathbf{v}_j\right)^2,$$

for i = 1, 2 and $j \in \mathbb{I}$ and

Theorem 2. According to theorem 1, the deterministic equivalent of the global minimum variance portfolio can be expressed as :

$$M\sigma_M^2(\hat{\mathbf{w}}_{SCRGMVP}) - M\overline{\sigma}_M^2(\hat{\mathbf{w}}_{SCRGMVP}) \xrightarrow{a.s.} 0$$

where

$$M\overline{\sigma}_{M}^{2}(\hat{\mathbf{w}}_{SCRGMVP}) = \frac{\overline{Y}(\gamma_{1},\gamma_{2})}{\left(\overline{X}(\gamma_{1},\gamma_{2})\right)^{2}}$$
(3.4)

Theorem 3. Under the setting of Assumption 1, the optimal parameters γ_1^*, γ_2^* that minimize $\overline{\sigma}_{SCRGMVP}^2(\gamma_1, \gamma_2)$ are given by:

$$\gamma_1^* = \frac{\varphi_1^*}{(1-\varphi_1^*)\lambda_1}, \text{ and } \gamma_2^* = \frac{-\varphi_2^*}{(1-\varphi_2^*)\lambda_1}.$$

where $(\varphi_1^*, \varphi_2^*)$ is the minimizer of the function $g(\varphi_1, \varphi_2), \varphi_i \in (0, 1), i = 1, 2$ given by

$$g(\varphi_1,\varphi_2) = \frac{\overline{Y}(\varphi_1,\varphi_2)}{\left(\overline{X}(\varphi_1,\varphi_2)\right)^2}$$

where

$$\begin{split} \overline{X}(\varphi_1,\varphi_2) &= 1 - \sum_{i=1,2} \sum_{j \in \mathbb{I}} a_j b_j \gamma_{i,j} \delta_{i,j} \\ \overline{Y}(\varphi_1,\varphi_2) &= 1 + \sum_{j \in \mathbb{I}} \lambda_j b_j - 2 \sum_{i=1,2} \sum_{j \in \mathbb{I}} \left(\lambda_j + 1\right) a_j b_j \gamma_{i,j} \delta_{i,j} \\ &+ \sum_{i=1,2} \sum_{j \in \mathbb{I}} a_j b_j \left(\lambda_j a_j + 1\right) \gamma_{i,j}^2 \delta_{i,j} \\ \gamma_{1,j} &= \frac{\varphi_1 \lambda_j}{(1 - \varphi_1) \lambda_1 + \varphi_1 \lambda_j}, \quad j \in \mathbb{I}_1, \\ \gamma_{2,j} &= \frac{-\varphi_2 \lambda_j}{(1 - \varphi_2) \lambda_{-1} - \varphi_2 \lambda_j}, \quad j \in \mathbb{I}_2. \end{split}$$

Theorem 4. Under Assumptions 3.1 to 3.4 and fix an $i \in \mathbb{I} = \{-r_2, ..., -1, 1, 2, ..., r_1\}$, for any deterministic unit vector $\mathbf{c} \in S_{\mathbb{R}}^{M-1}$, we have

$$\langle \mathbf{c}, \mathbf{u}_i \mathbf{u}_i^T \mathbf{c} \rangle \rightarrow \frac{\lambda_i^2 - J}{\lambda_i (\lambda_i + J)} \langle \mathbf{c}, \mathbf{v}_i \mathbf{v}_i^T \mathbf{c} \rangle,$$
 (3.5)

almost surely. This theorem has been proven in [41], so it is omitted here.

Theorem 5. Under Assumption 1, we have

$$\left|\lambda_j - \hat{\lambda}_j\right| \xrightarrow{a.s.} 0, \quad \left|b_j - \hat{b}_j\right| \xrightarrow{a.s.} 0,$$

where

$$\hat{\lambda}_j = \frac{s_j/\sigma^2 + 1 - J + \sqrt{\left(s_j/\sigma^2 + 1 - J\right)^2 - 4s_j/\sigma^2}}{2} - 1,$$
(3.6)

$$\hat{b}_{j} = \frac{1 + J/\hat{\lambda}_{j}}{1 - J/\hat{\lambda}_{j}} \frac{\frac{I_{M}}{\sqrt{M}} \mathbf{u}_{j} \mathbf{u}_{j}^{T} \frac{I_{M}}{\sqrt{M}}}{1 - J\sigma^{2}},$$
(3.7)

where, $J = \frac{M}{n}$, s_j is the jth maximum eigenvalue of the sample covariance matrix, and $j \in \mathbb{I}$.

4. Simulation

This section is mainly divided into three parts. The first part mainly compares whether the changing trends of $\sigma_M^2(\hat{\mathbf{w}}_{SCRGMVP})$ and $\overline{\sigma}_M^2(\hat{\mathbf{w}}_{SCRGMVP})$ are consistent; The second and third parts respectively compare our proposed method with other classic methods using synthetic data and multiple real stock data.

4.1. Consistent Estimation Comparison

In this paper, due to the optimization of parameter $\sigma_p^2(\hat{\mathbf{w}}_{SCRGMVP})$, we need to use the relevant results of random matrix theory to derive the consistent estimation $\overline{\sigma}_p^2(\hat{\mathbf{w}}_{SCRGMVP})$ of risk $\sigma_p^2(\hat{\mathbf{w}}_{SCRGMVP})$ in the large dimension environment of our proposed method. To verify the correctness of our derived consistent estimation, we need to compare whether the changing trends of $\sigma_p^2(\hat{\mathbf{w}}_{SCRGMVP})$ and $\overline{\sigma}_p^2(\hat{\mathbf{w}}_{SCRGMVP})$ are consistent. Next, we describe the commonly used simulation settings for experiments using synthetic data.Specifically, we set $\sigma^2 = 1$ and choose $r_1 = 3$, $r_2 = 1$. Provide the eigenvalues $\lambda_1 = 20$, $\lambda_2 = 10$, $\lambda_3 = 5$, $\lambda_M = 0.01$, and set the corresponding eigenvectors as follows: $v_1 = (1, 0, \dots, 0)'$, $v_2 = (0, 1, 0, \dots, 0)'$, $v_3 = (0, 0, 1, \dots, 0)'$, $v_M = (0, \dots, 0, 1)'$. Construct the covariance matrix of the spiked model structure according to the above settings. Using the training set, design the improved GMVP method using grid search. The performance of this risk and its estimation is illustrated in Figure 1, we can see how our asymptotic deterministic equivalence accurately approximates the true variance, gradually converging to the true variance as the sample size increases. Therefore, in high-dimensional situations, we can use grid search to find the optimal parameter γ^* .



Figure 1: Portfolio variance vs. sample size n for M = 50. Comparison of SCRGMVP(Sigma) and its consistent estimation(HatSigma) with synthetic data.

4.2. Synthetic Data Simulation

In this section, we evaluated the approximate accuracy of the following five different estimates: 1) the classical estimate, namely the sample covariance matrix estimator (SCME); 2) we propose the spectrally-corrected and regularized covariance matrix estimator (SCRE); 3) replace the unknown population covariance matrix with the shrinkage covariance matrix estimator(SHRE) proposed by Ledio-Wolf in (2004); 4) The two other methods are ordinary spectral correction estimator(SCE) and the weighted shrinkage estimator(WSHRE). We use Monte Carlo simulation methods to estimate portfolio risk and the specific steps are as follows:

- Step 1: Set σ² = 1 and choose r₁ = 3, r₂ = 1, orthogonal vectors as follows: v₁ = (1,0,...,0)', v₂ = (0,1,0,...,0)', v₃ = (0,0,1,0,...,0)', v_M = (0,...,0,1)' and their corresponding weights λ₁ = 20, λ₂ = 10, λ₃ = 5, λ_M = 0.01. Randomly generate p-dimensional mean vectors in intervals (-1, 1).
- Step 2: Generate *n* training samples.
- Step 3: Derive the spectral correction and the optimal parameters γ₁^{*}, γ₂^{*} of the sample covariance of the SCRGMVP method using grid search over {φ₁, φ₂} ∈ {[0, 1) × [0, 1)}.
- Step 4: Calculate the portfolio risks corresponding to these five population covariance estimates.
- Step 5: Repeat steps 2-4 10000 times and calculate the average portfolio risk corresponding to these five methods.



Figure 2: Portfolio variance vs. sample size n for M. Comparison for SCME, SCRE, SCE, WSHRE, MV bound and SHRE with synthetic data.

4.3. Real Data

For empirical analysis, we compare the performance of different optimal portfolio estimates by using real data from the S&P 500 index. We download stock data for the S&P 500 index from the website https://host.uniroma3.it/docenti/cesarone/DataSets.htm from October 6, 2006 to December 31, 2020 and computer their weekly logarithmic returns. There are 336 stocks and 753 weekly observations for each stock. We randomly selected 100 stocks from 336 stocks in the S&P 500 index. In the first comparison, select the top 200 as the sample set and 201 to 230 as the test set for prediction. For the second comparison, select 2 to 201 as the training set, 202 to 231 as the test set for prediction, and repeat 500 times in sequence.

To compare the performance of our proposed method SCRE with SCME, SHRE, WSHRE, and SCE. We use the following protocol for real datasets:

• Step 1: Select n weekly logarithmic returns as the training dataset and calculate the sample covariance. Spectral decomposition and correct the sample covariance under the spiked model, and estimate of spiked eigenvalues. Select the data from the next period of the training sample as the test dataset for estimating portfolio risk.

- Step 2: Using the training dataset, derive the optimal parameters γ₁^{*}, γ₂^{*} of the sample covariance of the SCRGMVP method using grid search over {φ₁, φ₂} ∈ {[0, 1) × [0, 1)}. Design the SCRGMVP estimation and the other four estimators.
- Step 3: Using the test dataset, estimate the portfolio risk for five methods.
- Step 4: Repeat steps 1-3 500 times and calculate the average portfolio risk for all estimators.

The estimated portfolio risks using different methods are shown in the following figure:



Figure 3: Portfolio variance vs. sample size n for M. Comparison for SCME, SCRE, SCE, WSHRE, and SHRE with real data.

5. Conclusion

We propose a risk model that is superior to the classical model, namely the sample covariance matrix. Due to the shortcomings of the sample covariance matrix, we propose an enhanced alternative. Thus, the purpose of this paper is to solve the Markowitz optimization problem by developing a new covariance estimation to address the essence of portfolios. The key to this method is to assume that the population covariance matrix follows a spiked model. Using this special structure, the proposed estimator uses an estimation of the covariance matrix and follows the spiked model with regularized parameters. Using the results of the random theory, the asymptotic characterization of the proposed estimator is given, and the parameter that minimizes the consistent estimation of the estimator is selected. In this way, we can use the estimated eigenvalues and corresponding eigenvectors to preserve the form of the spiked covariance model as much as possible. Meanwhile, regularization methods can further reduce investment risks. Through the simulation of synthetic and real data, we found that our proposed method not only performs well compared to other methods but also greatly reduces the number of design parameters, which greatly reduces computational complexity. In further research, the same idea can also be applied to other backgrounds of Markowitz portfolio or other fields to improve the estimation of the population covariance matrix, such as linear discriminant analysis (LDA), Markowitz maximum return, etc.

Appendix A. Proof of Theorem 1

Firstly, according to (3.5), we have

$$\begin{split} \mathbf{X}\left(\widetilde{\mathbf{C}}\right) &= \frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \widetilde{\mathbf{C}} \frac{\mathbf{1}_{M}}{\sqrt{M}} \\ &= \frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \left(\mathbf{I}_{M} + \gamma_{1} \sum_{j \in \mathbb{I}_{1}} \lambda_{j} \mathbf{u}_{j} \mathbf{u}_{j}^{T} + \gamma_{2} \sum_{j \in \mathbb{I}_{2}} \lambda_{j} \mathbf{u}_{j} \mathbf{u}_{j}^{T} \right)^{-1} \frac{\mathbf{1}_{M}}{\sqrt{M}} \\ &= \frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \left(\mathbf{I}_{M} - \sum_{j \in \mathbb{I}_{1}} \gamma_{1,j} \mathbf{u}_{j} \mathbf{u}_{j}^{T} - \sum_{j \in \mathbb{I}_{2}} \gamma_{2,j} \mathbf{u}_{j} \mathbf{u}_{j}^{T} \right) \frac{\mathbf{1}_{M}}{\sqrt{M}} \\ &= 1 - \sum_{j \in \mathbb{I}_{1}} \gamma_{1,j} \frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \mathbf{u}_{j} \mathbf{u}_{j}^{T} \frac{\mathbf{1}_{M}}{\sqrt{M}} - \sum_{j \in \mathbb{I}_{2}} \gamma_{2,j} \frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \mathbf{u}_{j} \mathbf{u}_{j}^{T} \frac{\mathbf{1}_{M}}{\sqrt{M}} \\ &\stackrel{a.s.}{\longrightarrow} 1 - \sum_{j \in \mathbb{I}_{1}} \gamma_{1,j} a_{j} b_{j} - \sum_{j \in \mathbb{I}_{2}} \gamma_{2,j} a_{j} j b_{j} \\ &\triangleq \overline{X}(\gamma_{1}, \gamma_{2}), \end{split}$$
(5.1)

in which

$$a_{j} = \frac{\lambda_{j}^{2} - J}{\lambda_{j}(\lambda_{j} + J)}, \quad b_{j} = \left(\frac{\mathbf{1}_{M}^{T}}{\sqrt{M}}\mathbf{v}_{j}\right)^{2},$$
$$\gamma_{1,j} = \frac{\varphi_{1}\lambda_{j}}{(1 - \varphi_{1})\lambda_{1} + \varphi_{1}\lambda_{j}}, \quad j \in \mathbb{I}_{1},$$
$$\gamma_{2,j} = \frac{-\varphi_{2}\lambda_{j}}{(1 - \varphi_{2})\lambda_{j}}, \quad j \in \mathbb{I}_{2}.$$
(5.2)

$$\gamma_{2,j} = \frac{-\varphi_2 \lambda_j}{(1-\varphi_2)\lambda_{-1} - \varphi_2 \lambda_j}, \quad j \in \mathbb{I}_2.$$
(5.3)

Secondly, rewrite 2.7, we have

$$\begin{aligned} \mathbf{Y}(\widetilde{\mathbf{C}}, \mathbf{\Sigma}) &= \frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \widetilde{\mathbf{C}} \mathbf{\Sigma} \widetilde{\mathbf{C}} \frac{\mathbf{1}_{M}}{\sqrt{M}} \\ &= \sigma_{M}^{2} \frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \widetilde{\mathbf{C}} \left(\mathbf{I}_{M} + \sum_{j \in \mathbb{I}_{1}} \lambda_{j} \mathbf{v}_{j} \mathbf{v}_{j}^{T} + \sum_{j \in \mathbb{I}_{2}} \lambda_{j} \mathbf{v}_{j} \mathbf{v}_{j}^{T} \right) \widetilde{\mathbf{C}} \frac{\mathbf{1}_{M}}{\sqrt{M}} \\ &= \sigma_{M}^{2} \left(\frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \widetilde{\mathbf{C}}^{2} \frac{\mathbf{1}_{M}}{\sqrt{M}} + \sum_{j \in \mathbb{I}_{1}} \lambda_{j} \left(\frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \widetilde{\mathbf{C}} \mathbf{v}_{j} \right) \right)^{2} + \sum_{j \in \mathbb{I}_{2}} \lambda_{j} \left(\frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \widetilde{\mathbf{C}} \mathbf{v}_{j} \right)^{2} \right). \end{aligned}$$

For the three elements of 5, according to 3.5, we have

$$\frac{\mathbf{1}_{M}^{T}}{\sqrt{M}}\widetilde{\mathbf{C}}^{2}\frac{\mathbf{1}_{M}}{\sqrt{M}} \xrightarrow{a.s.} 1 + \sum_{j\in\mathbb{I}_{1}} \left(\gamma_{1,j}^{2} - 2\gamma_{1,j}\right)a_{j}b_{j} + \sum_{j\in\mathbb{I}_{2}} \left(\gamma_{2,j}^{2} - 2\gamma_{2,j}\right)a_{j}b_{j},$$
$$\frac{\mathbf{1}_{M}^{T}}{\sqrt{M}}\widetilde{\mathbf{C}}\mathbf{v}_{j}\mathbf{v}_{j}^{T}\widetilde{\mathbf{C}}\frac{\mathbf{1}_{M}}{\sqrt{M}} \xrightarrow{a.s.} b_{j}\left(1 - 2\gamma_{i,j}a_{j} + \gamma_{i,j}^{2}a_{j}^{2}\right).$$

Similarly, for 5, we have

$$\begin{split} \mathbf{Y}(\widetilde{\mathbf{C}}, \mathbf{\Sigma}) &= \frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \widetilde{\mathbf{C}} \mathbf{\Sigma} \widetilde{\mathbf{C}} \frac{\mathbf{1}_{M}}{\sqrt{M}} \\ &= \sigma_{M}^{2} \left(\frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \widetilde{\mathbf{C}}^{2} \frac{\mathbf{1}_{M}}{\sqrt{M}} + \sum_{j \in \mathbb{I}_{1}} \lambda_{j} \left(\frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \widetilde{\mathbf{C}} \mathbf{v}_{j} \right)^{2} + \sum_{j \in \mathbb{I}_{2}} \lambda_{j} \left(\frac{\mathbf{1}_{M}^{T}}{\sqrt{M}} \widetilde{\mathbf{C}} \mathbf{v}_{j} \right)^{2} \right) \\ & \xrightarrow{a.s.} \sigma_{M}^{2} \left(\begin{array}{c} 1 + \sum_{j \in \mathbb{I}_{1}} \left(\gamma_{1,j}^{2} - 2\gamma_{1,j} \right) a_{j} b_{j} \\ + \sum_{j \in \mathbb{I}_{2}} \left(\gamma_{2,j}^{2} - 2\gamma_{2,j} \right) a_{j} b_{j} \\ + \sum_{j \in \mathbb{I}_{1}} \lambda_{j} b_{j} \left(1 - 2\gamma_{1,j} a_{j} + \gamma_{1,j}^{2} a_{j}^{2} \right) \\ + \sum_{j \in \mathbb{I}_{2}} \lambda_{j} b_{j} \left(1 - 2\gamma_{2,j} a_{j} + \gamma_{2,j}^{2} a_{j}^{2} \right) \end{split} \right) \end{split}$$

$$= \sigma_{M}^{2} \begin{pmatrix} 1 + \sum_{j \in \mathbb{I}_{1}} \lambda_{j}b_{j} + \sum_{j \in \mathbb{I}_{2}} \lambda_{j}b_{j} \\ -2 \sum_{j \in \mathbb{I}_{1}} \gamma_{1,j}(1 + \lambda_{j})a_{j}b_{j} \\ -2 \sum_{j \in \mathbb{I}_{2}} \gamma_{2,j}(1 + \lambda_{j})a_{j}b_{j} \\ + \sum_{j \in \mathbb{I}_{1}} \gamma_{1,j}^{2}(1 + \lambda_{j}a_{j})a_{j}b_{j} \\ + \sum_{j \in \mathbb{I}_{2}} \gamma_{2,j}^{2}(1 + \lambda_{j}a_{j})a_{j}b_{j} \end{pmatrix}$$

$$\triangleq \overline{Y}(\gamma_{1}, \gamma_{2})$$

Appendix B. Proof of Theorem 3

Let $\varphi_i = \gamma_{i,1} \in (0, 1)$ for i = 1, 2. Then, γ_1 and γ_2 can be rewritten as

$$\gamma_1 = \frac{\varphi_1}{(1-\varphi_1)\lambda_1}, \text{ and } \gamma_2 = \frac{-\varphi_2}{(1-\varphi_2)\lambda_{-1}}.$$

Further,

$$\gamma_{1,j} = \frac{\varphi_1 \lambda_j}{(1 - \varphi_1)\lambda_1 + \varphi_1 \lambda_j}, \quad j = 1, 2, ..., r_1,$$
(5.4)

$$\gamma_{2,j} = \frac{-\varphi_2 \lambda_j}{(1-\varphi_2)\lambda_{-1} - \varphi_2 \lambda_{-j}}, \quad j = -1, -2, \dots, -r_2.$$
(5.5)

Plugging (5.4) and (5.5) into (3.2) and (3.3), we have

$$\widetilde{X}(\lambda_1, \lambda_2) = \overline{G}(\gamma_1, \gamma_2) \text{ and } \widetilde{Y}(\lambda_1, \lambda_2) = \overline{D}(\gamma_1, \gamma_2).$$

Thus, the objective function can be written as

$$g(\varphi_1,\varphi_2) = \frac{\overline{Y}(\varphi_1,\varphi_2)}{\left(\overline{X}(\varphi_1,\varphi_2)\right)^2}$$

Thus, (3.1) is equivalent to

$$\varphi_1^*, \varphi_2^* = \arg\min_{\varphi_1, \varphi_2 \in (0,1)} g(\varphi_1, \varphi_2)$$

by which

$$\gamma_1^* = \frac{\varphi_1^*}{(1-\varphi_1^*)\lambda_1}, \text{ and } \gamma_2^* = \frac{-\varphi_2^*}{(1-\varphi_2^*)\lambda_{-1}}.$$

REFERENCES

- [1] Harry M Markowitz. Portfolio selection. The Journal of Finance, 7(1):77, 1952.
- [2] Review By: Alan Stuart. Portfolio selection: Efficient diversification of investmentsby harry m. markowitz. OR, 10(4):253–254, 1959.
- [3] Vijay Kumar Chopra and William T Ziemba. The effect of errors in means, variances, and covariances on optimal portfolio choice. *Journal of Portfolio Management*, 19(2):6–11, 1993.
- [4] Ravi Jagannathan, M. A. Tongshu, and Jiaqi Zhang. A note on "risk reduction in large portfolios: Why imposing the wrong constraints helps". *The Journal of Finance*, 74(5):2689–2696, 2019.
- [5] Chris Kirby and Barbara Ostdiek. Optimizing the performance of sample meanvariance efficient portfolios. *Social Science Electronic Publishing*, 2011.
- [6] Yarema, Okhrin, Wolfgang, and Schmid. Distributional properties of portfolio weights. *Journal of Econometrics*, 2006.
- [7] Taras Bodnar and Wolfgang Schmid. A test for the weights of the global minimum variance portfolio in an elliptical model. *Metrika*, 67(2), 2007.
- [8] Jianqing Fan, Jingjin Zhang, and Ke Yu. Vast portfolio selection with grossexposure constraints.
- [9] Patrick, Behr, Andre, Guettler, Felix, and Miebs. On portfolio optimization: Imposing the right constraints. *Journal of banking and finance*, 2013.
- [10] Jiahan Li. Sparse and stable portfolio selection with parameter uncertainty. *Journal of Business and Economic Statistics*, 33(3):381–392, 2015.
- [11] Vasyl Golosnoy and Yarema Okhrin. Multivariate shrinkage for optimal portfolio weights. *European Journal of Finance*, 13(5):441–458, 2007.
- [12] Taras Bodnar, Nestor Parolya, and Wolfgang Schmid. Estimation of the global minimum variance portfolio in high dimensions, 2014.

- [13] Olivier Ledoit and Michael Wolf. Improved estimation of the covariance matrix of stock returns with an application to portfolio selection. *Journal of empirical finance*, 10(5):603–621, 2003.
- [14] Olivier Ledoit, Michael Wolf, and Zhao Zhao. Efficient weighting: a more powerful test for cross-sectional anomalies. *SSRN Electronic Journal*, 2016.
- [15] Zhidong Bai and Jack W Silverstein. Spectral analysis of large dimensional random matrices, volume 20. Springer, 2010.
- [16] Zhigang Bao, Jiang Hu, Xiaocong Xu, and Xiaozhuo Zhang. Spectral statistics of sample block correlation matrices. arXiv preprint arXiv:2207.06107, 2022.
- [17] T. Tony Cai A, Jianchang Hu B, Yingying Li C, and Xinghua Zheng D. Highdimensional minimum variance portfolio estimation based on high-frequency data. *Journal of Econometrics*, 214(2):482–494, 2020.
- [18] Nikolaus Hautsch, Lada M. Kyj, and Peter Malec. Do high-frequency data improve high-dimensional portfolio allocations? *Journal of Applied Econometrics*, (2), 2015.
- [19] Bala Rajaratnam, Hélène Massam, and Carlos M Carvalho. Flexible covariance estimation in graphical gaussian models. *Ann. Statist.*, 36(6):2818–2849, 2008.
- [20] Rohde Alexandre B. Tsybakov. Estimation of high-dimensional low-rank matrices. Annals of Statistics, 39(2):887–930, 2011.
- [21] Pradeep Ravikumar, Martin J Wainwright, Garvesh Raskutti, and Bin Yu. Highdimensional covariance estimation by minimizing *ell*₁-penalized log-determinant divergence. 2008.
- [22] T Tony Cai and Harrison H Zhou. Minimax estimation of large covariance matrices under 1 -norm. *Statistica Sinica*, 22(4), 2012.
- [23] Olivier Ledoit and Michael Wolf. A well-conditioned estimator for largedimensional covariance matrices. *Academic Press*, (2), 2004.

- [24] Olivier Ledoit and Michael Wolf. Nonlinear shrinkage estimation of largedimensional covariance matrices. *The Annals of Statistics*, 40(2):1024–1060, 2011.
- [25] Olivier Ledoit and Michael Wolf. Nonlinear shrinkage estimation of the covariance matrix of asset returns. 2010.
- [26] Michael J. Daniels and Robert E. Kass. Shrinkage estimators for covariance matrices. 2001.
- [27] Zhidong Bai, Hua Li, and Wing Keung Wong. The best estimation for highdimensional markowitz mean-variance optimization. *MPRA Paper*, 2013.
- [28] Peter J. Bickel and Elizaveta Levina. Regularized estimation of large covariance matrices. arXiv e-prints, 2008.
- [29] ONATSKI and ALEXEI. Detection of weak signals in high-dimensional complex-valued data. *Random Matrices: Theory and Applications*, 03(01):1450001, 2014.
- [30] Shira Kritchman and Boaz Nadler. Determining the number of components in a factor model from limited noisy data. *Chemometrics and Intelligent Laboratory Systems*, 94(1):19–32, 2008.
- [31] D. J. Davidson. Functional mixed-effect models for electrophysiological responses. *Neurophysiology*, 41(1):71–79, 2009.
- [32] Siamac Fazli, Márton Danóczy, Jürg Schelldorfer, and Klaus Robert Müller. 1penalized linear mixed-effects models for high dimensional data with application to bci. *NeuroImage*, (4), 2011.
- [33] David C Hoyle and Magnus Rattray. Limiting form of the sample covariance eigenspectrum in pca and kernel pca. In *CiteSeer*, pages 16–23, 2004.
- [34] Zheng Tracy Ke, Yucong Ma, and Xihong Lin. Estimation of the number of spiked eigenvalues in a covariance matrix by bulk eigenvalue matching analysis. *Journal of the American Statistical Association*, pages 1–45, 2021.

- [35] Jinho Baik and Jack W Silverstein. Eigenvalues of large sample covariance matrices of spiked population models. *Journal of Multivariate Analysis*, 97, 2006.
- [36] Z. D. Bai and Jack W. Silverstein. No eigenvalues outside the support of the limiting spectral distribution of large-dimensional sample covariance matrices. *The Annals of Probability*, 1998.
- [37] Z. D. Bai and Jack W. Silverstein. Clt for linear spectral statistics of largedimensional sample covariance matrices. *The Annals of Probability*, 2004.
- [38] Zhidong Bai and Xue Ding. Estimation of spiked eigenvalues in spiked models. *Random Matrices Theory and Applications*, 1(02):1150011, 2012.
- [39] Zhidong Jiang, Dandan Bai. Generalized four moment theorem and an application to clt for spiked eigenvalues of high-dimensional covariance matrices. *Bernoulli: official journal of the Bernoulli Society for Mathematical Statistics and Probability*, 27(1), 2021.
- [40] Jianqing Fan, Jianhua Guo, and Shurong Zheng. Estimating number of factors by adjusted eigenvalues thresholding. *Journal of the American Statistical Association*, pages 1–33, 2020.
- [41] Hua Li, Wenya Luo, Zhidong Bai, Huanchao Zhou, and Zhangni Pu. Spectrallycorrected and regularized linear discriminant analysis for spiked covariance model. arXiv preprint arXiv:2210.03859, 2022.