

An explicit construction of Kaleidocycles

By Shizuo Kaji at Fukuoka, Kenji Kajiwara at Fukuoka and Shota Shigetomi at Fukuoka

Abstract. We model a family of closed kinematic chains, known as Kaleidocycles, with the theory of discrete spatial curves. By leveraging the connection between the deformation of discrete curves and the semi-discrete integrable systems, we describe the motion of a Kaleidocycle by elliptic theta functions. This study showcases an interesting example in which an integrable system generates an orbit in the space of the real solutions of polynomial equations defined by geometric constraints.

1. Introduction

The primary motivation of this work comes from the analysis of the motion of a particular class of linkage mechanisms called *Kaleidocycles* (Fig. 1 (Left)). It consists of (typically six) identical copies of equifacial tetrahedra connected by their opposing edges as hinge joints to form a ring. A Kaleidocycle can be folded from a sheet of paper and makes an example of rigid origami; it turns like a bubble ring while keeping all faces rigid. The origin of Kaleidocycle is obscure but one of the first appearances in the literature is in Bricard's paper [3], where he considers an essentially equivalent closed kinematic chain. This kinematic chain, now known as the Bricard 6R, is an *over-constrained* linkage, which violates Maxwell's degree-of-freedom counting law [8]. For nearly a hundred years, the geometry and kinematics of this object have attracted the attention of researchers (see [25] and the references therein).

For a fixed number k of tetrahedra and their shape, the state of a Kaleidocycle is identified by the sequence of the turning angles at the hinges, which we define in the following manner. Notice that the line segment, which we call the axis, connecting the centres of hinge edges for each equifacial tetrahedron is orthogonal to both hinge edges. By connecting the axes, we obtain a spatial closed polygonal curve. The angles at the vertices of this polygon determine the state of a Kaleidocycle. The angles between the hinge edges are reminiscent of the torsion angle of a polygonal curve, and the turning angles are reminiscent of the curvature angle. We will make this connection more rigorous and formulate the state of a Kaleidocycle in the language of a discrete spatial curve (§2.1).

In the geometric formulation as discrete spatial curves, the motion of a Kaleidocycle is modelled by a segment-length and torsion-angle preserving deformation of the corresponding closed polygonal curve of a constant torsional angle. Here, we ignore the self-collision of the tetrahedra. There is another algebraic perspective of Kaleidocycles. The state of a Kaleidocycle is determined by the coordinates of the vertices of the tetrahedra satisfying the distance condition specified by the edge lengths. Thus, the state space, or the configuration space, is identified with the space of real solutions to a system of quadratic equations. The motion corresponds to a one-dimensional orbit in the configuration space. The analysis of the real algebraic set obtained as the configuration space of a Kaleidocycle is mathematically challenging as well as of practical importance. For example, a simple counting of the number of variables and the constraints tells that the state space of a Kaleidocycle with k tetrahedra is generally $(k - 6)$ -dimensional (up to rigid motion). The first-named author and collaborators [19] have numerically discovered that the state space degenerates to a circle when the hinge angle is set to be maximal or minimal in some sense.¹⁾ This class of Kaleidocycles, named the *Möbius Kaleidocycles*, provides a rare example of *underconstrained* linkage mechanisms whose state space has less dimension than the expected one. Numerical experiments suggest that the Kaleidocycles constructed in this paper are indeed Möbius Kaleidocycles (Conjecture 6.1). The first-named and second-named authors have investigated the motion of Möbius Kaleidocycles and identified a particular motion of Kaleidocycles (not restricted to Möbius Kaleidocycles) that is governed by the semi-discrete modified KdV (mKdV) and the sine-Gordon equations [18](see §3.2). Given an initial state, a one-dimensional orbit in the generally $(k - 6)$ -dimensional state space is obtained by the time evolution defined by a system of ordinary differential equations. The main result of this paper can be seen as giving an explicit solution to the system by using the explicit construction in terms of the τ functions of the two-component KP hierarchy [10]. However, our approach is different from that in [18]; we start with the functional identities of elliptic theta functions in connection to the compatibility conditions of the framing of a discrete spatial curve. It is thus not trivial that these two approaches yield the same motions. We show that our construction of the motion of the Kaleidocycle (§5) indeed satisfies the system of the ordinary differential equations described in [18]. The sweeping surface of

¹⁾ The results of the patent are also published in [26].

the motion corresponds to a semi-discrete analogue of the discrete constant negative curvature surface (*K-surface*) studied in [2, 15].

Our work connects seemingly distant fields of integrable systems and kinematics of linkage via an interesting object, Kaleidocycle. It provides a remarkable example, where a flow generated by an integrable system on a real algebraic set is explicitly integrated using elliptic theta functions.

The structure of the paper is as follows. §2 introduces Kaleidocycles and their representation as discrete space curves. As essential numerical invariants to describe the state of a Kaleidocycle, the signed curvature angle and torsion angle are defined using the Frenet frame. A concise parametrisation of a Kaleidocycle as a solution to a system of quadratic equations corresponding to the geometric constraints on the binormal vectors is given. §3 reviews the connection between the theory of curves and integrable systems. The construction of discrete space curves and their Frenet frames in terms of the τ functions is presented. A previous result that describes the motion of a Kaleidocycle using semi-discrete integrable equations is recalled. §4 constructs a set of τ functions with three parameters v, r, y using elliptic theta functions. The resulting curve is shown to have a constant torsion angle. A condition for the resulting curve to be closed is also discussed; for a given k , the appropriate choice of parameters m, r, v, y yields a Kaleidocycle. By introducing another parameter, a family of time-parametrised Kaleidocycles is constructed in §5. Then, we investigate the time evolution equations satisfied by the vertex positions and curvature angle of the Kaleidocycles and show connections to previous research. §6 provides examples of Kaleidocycles and corresponding semi-discrete surfaces with figures. We present two conjectures that are suggested by numerical experiments. Most of the proofs of our theorems are deferred to Appendix since they require technically-involved computations regarding elliptic theta functions.

Acknowledgement. The second and third-named authors were partially supported by KAKENHI Grant-in-Aid for Scientific Research (C) 21K3229. All authors are partially supported by JST CREST Grant No. JP-MJCR1911.

2. Geometric and algebraic models of Kaleidocycle

2.1. Definition of Kaleidocycle. We first define a generalisation of Kaleidocycle following [17]. We consider k -copies of an equifacial tetrahedron in \mathbb{R}^3 connected by opposing edges to form a ring. Denote the vertex set by $\{0, 1, \dots, 2k-1\}$ such that the vertices $2n$ and $2n+1$ form a hinge edge and $2n, 2n+1, 2n+2$ form one of the triangle faces. Let l_{01}, l_{12}, l_{20} be the lengths of the sides of the triangle face. A state of a k -Kaleidocycles of type (l_{01}, l_{12}, l_{20}) is identified by the real solution $(x_0, x_1, \dots, x_{2k-1}) \in (\mathbb{R}^3)^{2k}$ of the quadratic equations

$$(2.1) \quad \begin{aligned} |x_{2n} - x_{2n+1}|^2 &= l_{01}^2, \\ |x_{2n} - x_{2n+2}|^2 &= |x_{2n+1} - x_{2n+3}|^2 = l_{02}^2, \\ |x_{2n+1} - x_{2n+2}|^2 &= |x_{2n} - x_{2n+3}|^2 = l_{12}^2. \end{aligned}$$

We note that we ignore collision between tetrahedra. The solution space of (2.1) admits an action of the Euclidean transformation group $E(3)$ of \mathbb{R}^3 , and the expected dimension of the quotient is $3 \cdot 2k - 5k - \dim(E(3)) = k - 6$.

The state of a Kaleidocycle can be equivalently described by a spatial polygon or a *closed discrete spatial curve*. Let $\gamma_n = (x_{2n} + x_{2n+1})/2$ ($0 \leq n < k-1$) be the mid-point of the hinge edge. We set $\gamma_k = \gamma_0$. Since the tetrahedron is equifacial, $\gamma_{n+1} - \gamma_n$ is orthogonal to both \tilde{B}_n and \tilde{B}_{n+1} , where $\tilde{B}_n = \frac{x_{2n+1} - x_{2n}}{|x_{2n+1} - x_{2n}|}$ is the hinge direction. By reordering the vertices, we can assume the inner product $\langle \tilde{B}_{n-1}, \tilde{B}_n \rangle$ is constant for $1 \leq n < k$. Denote this constant by $\cos \lambda$ with $0 < \lambda < \pi$, which we call the *torsion angle*. We have two cases $\langle \tilde{B}_{k-1}, \tilde{B}_0 \rangle = \pm \cos \lambda$. We say the Kaleidocycle is oriented (resp. non-oriented) when the value is $+\cos \lambda$ (resp. $-\cos \lambda$). We set $\tilde{B}_k = \pm \tilde{B}_0$ so that $\langle \tilde{B}_{k-1}, \tilde{B}_k \rangle = \cos \lambda$. Furthermore, since $\tilde{B}_{n+1} \times \tilde{B}_n$ is parallel to $\gamma_{n+1} - \gamma_n$, we have $\sum_{n=0}^{k-1} (\tilde{B}_{n+1} \times \tilde{B}_n) = \sum_{n=0}^{k-1} \frac{\gamma_{n+1} - \gamma_n}{|\gamma_{n+1} - \gamma_n|} = 0$. Conversely, the vertex coordinates x_n can be reconstructed from $\{\gamma_n\}$, $\{\tilde{B}_n\}$, and (l_{01}, l_{12}, l_{20}) up to translation. Therefore, $\{\tilde{B}_n \in S^2 \mid 0 \leq n \leq k\}$ satisfying the conditions

$$(2.2) \quad \begin{aligned} \langle \tilde{B}_{n-1}, \tilde{B}_n \rangle &= \cos \lambda, \\ \sum_{n=0}^{k-1} (\tilde{B}_{n+1} \times \tilde{B}_n) &= 0, \\ \tilde{B}_k &= \pm \tilde{B}_0 \end{aligned}$$




Figure 1. (Left) The classical 6-Kaleidocycle. (Right) Kaleidocycle as a discrete curve with its framing.

define a Kaleidocycle. We call the space of solutions to (2.2) the state space of the k -Kaleidocycles with the torsion angle λ .

From this perspective, a motion of a Kaleidocycle is identified with a one-dimensional (time-parametrised) orbit of the space of the quadratic system defined by (2.1) or (2.2).

2.2. Discrete spatial curve of a constant torsion angle. A discrete spatial curve with non-vanishing torsion angles can be described in terms of binormals. Let $\tilde{B} : \mathbb{Z} \rightarrow S^2$ satisfy $\langle \tilde{B}_n, \tilde{B}_{n-1} \rangle \neq 0$ for all $n \in \mathbb{Z}$. A unit segment-length discrete spatial curve having \tilde{B} as the binormals can be constructed as follows. Define the tangent $T : \mathbb{Z} \rightarrow S^2$ and the normal $\tilde{N} : \mathbb{Z} \rightarrow S^2$ by

$$(2.3) \quad T_n = \frac{\tilde{B}_{n+1} \times \tilde{B}_n}{|\tilde{B}_{n+1} \times \tilde{B}_n|}, \quad \tilde{N}_n = \tilde{B}_n \times T_n.$$

Furthermore, given $\gamma_0 \in \mathbb{R}^3$, define $\gamma : \mathbb{Z} \rightarrow \mathbb{R}^3$ by

$$(2.4) \quad \gamma_{n+1} = \gamma_n + T_n = \gamma_n + \frac{\tilde{B}_{n+1} \times \tilde{B}_n}{|\tilde{B}_{n+1} \times \tilde{B}_n|}.$$

The *signed curvature angle* κ and the *torsion angle* are defined by

$$(2.5) \quad \begin{aligned} \langle T_n, T_{n-1} \rangle &= \cos \kappa_n, & -\pi < \kappa_n < \pi, \\ \langle \tilde{B}_n, \tilde{B}_{n-1} \rangle &= \cos \lambda_n, & 0 < \lambda_n < \pi. \end{aligned}$$

The *discrete Frenet frame* $\Phi_n = [T_n, \tilde{N}_n, \tilde{B}_n] \in \text{SO}(3)$ satisfies the *discrete Frenet-Serret formula*

$$(2.6) \quad \Phi_{n+1} = \Phi_n L_n, \quad L_n = M_1(-\lambda_{n+1}) M_3(\kappa_{n+1}),$$

where

$$(2.7) \quad M_1(\theta) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{bmatrix}, \quad M_3(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Remark 2.1. Note that our Frenet frame coincides with the usual one only up to sign. We adopt this convention to avoid discontinuous change in the framing during the motion of the Kaleidocycle; its motion passes the state when the curvature vanishes.

As in the case of a smooth curve, a discrete spatial curve of a unit segment length is determined by its torsion and curvature angles. Therefore, the motion of a discrete curve of a constant torsion angle is determined by the set of functions $\kappa_n(t)$, which can be freely chosen. The fact that a Kaleidocycle is *closed* puts interesting topological constraints. For example, the self-linking number must be a half-integer, and hence, stay constant during the motion. By Călugăreanu-Fuller-White's theorem [1], the writhing number of the curve should stay constant as well. The self-linking number also tells that the state space of a Kaleidocycle of a fixed type is not connected in general.

3. Curves and integrable systems

There are deep connections between differential geometry and the theory of integrable systems, where the integrable differential or difference equations arise as compatibility conditions of the geometric objects (see, for example, [24]). In the context of the deformation of spatial and plane curves, the Frenet frame and its deformation are described by the system of linear partial differential equations. The compatibility condition yields so-called the Ablowitz-Kaup-Newell-Segur (AKNS) hierarchy for the curvature and the torsion, including the nonlinear Schrödinger (NLS) equation as a typical example. In particular, if the torsion is a constant, the integrable deformation is governed by the mKdV equation and its hierarchy [9, 21, 24]. The connections are also investigated in a discrete setting, where curves are discretised as piece-wise linear (polygonal) curves. For example, continuous deformations of discrete plane and spatial curves have been considered in [5], where the deformations are governed by the semi-discrete analogue of the mKdV and the NLS equations. Discrete deformations of the discrete plane and spatial curves have been studied in [15]. In principle, for a given solution to the nonlinear differential/difference equation for the curvature and torsion, one can reconstruct the curve by solving the associated system of linear differential/difference equations for the Frenet frame and integrating the tangent vector.

The theta functions naturally appear in so-called the quasi-periodic solutions to the nonlinear integrable equations, and the corresponding solutions to the associated linear problem are nothing but the Baker-Akhiezer functions. Reconstruction of the deformation of the spatial curve with constant torsion described by the theta function solutions to the mKdV equation has been studied in this framework in [4, 22]. Also, an explicit formula of the spatial curves in terms of the τ functions of the two-component KP hierarchy based on Hirota's bilinear formalism has been presented in [10].

Since the dynamics of the Kaleidocycles are modelled by the integrable deformation of the closed discrete spatial curves with a constant torsion angle which preserves the segment-length and torsion angle [18], we study the theta function solutions to the corresponding integrable semi-discrete equations, where the theta functions are nothing but the τ functions. Therefore we construct the explicit formula of the Kaleidocycles based on the method of direct reconstruction of the discrete spatial curves in terms of the τ functions.

Hereafter, a (discrete) curve always indicates a polygonal spatial curve.

3.1. Expression of a curve by the τ functions. To describe Kaleidocycles, we introduce a special class of curves represented by the τ functions, which is a slight modification of the one presented in [10]. We construct a family of discrete Frenet frames $\Phi_n = [T_n, \tilde{N}_n, \tilde{B}_n]$ indexed by $n \in \mathbb{Z}$ and the corresponding curve γ_n satisfying (2.6). Let $F_n, \tilde{G}_n : \mathbb{C} \rightarrow \mathbb{R}$ and $f_n, g_n, G_n, H_n : \mathbb{C} \rightarrow \mathbb{C}$ be functions indexed by $n \in \mathbb{Z}$ that satisfy the following equations

$$(3.1) \quad f_n f_n^* + g_n g_n^* = F_{n+1} F_n$$

$$(3.2) \quad \beta F_n f_n + G_n g_n^* = F_{n+1} f_{n-1}$$

$$(3.3) \quad \beta F_n g_n - G_n f_n^* = F_{n+1} g_{n-1}$$

$$(3.4) \quad H_{n+1} F_n - H_n F_{n+1} = f_n^* g_n$$

$$(3.5) \quad D_z F_{n+1} \cdot F_n = g_n g_n^*$$

$$(3.6) \quad |G_n|^2 = \tilde{G}_n^2,$$

where $\beta \in \mathbb{R}$ is some constant and D_z is Hirota's bilinear differential operator [11]. These functions are called the τ functions. Let $\varrho_n = \frac{1}{2} \frac{G_n}{\tilde{G}_n}$ and

$$(3.7) \quad \begin{aligned} T_n &= \frac{1}{f_n f_n^* + g_n g_n^*} \left(\begin{array}{c} f_n^* g_n + f_n g_n^* \\ \frac{1}{i} (f_n^* g_n - f_n g_n^*) \\ f_n f_n^* - g_n g_n^* \end{array} \right) \Bigg|_{z=0} \\ \tilde{N}_n &= \frac{1}{f_n f_n^* + g_n g_n^*} \left(\begin{array}{c} \{(f_n^*)^2 - (g_n^*)^2\} \varrho_n + \{(f_n)^2 - (g_n)^2\} \varrho_n^* \\ \frac{1}{i} [\{(f_n^*)^2 + (g_n^*)^2\} \varrho_n - \{(f_n)^2 + (g_n)^2\} \varrho_n^*] \\ -2 (f_n^* g_n \varrho_n + f_n g_n \varrho_n^*) \end{array} \right) \Bigg|_{z=0} \\ \tilde{B}_n &= \frac{1}{f_n f_n^* + g_n g_n^*} \left(\begin{array}{c} \frac{1}{i} [-\{(f_n^*)^2 - (g_n^*)^2\} \varrho_n + \{(f_n)^2 - (g_n)^2\} \varrho_n^*] \\ \{(f_n^*)^2 + (g_n^*)^2\} \varrho_n + \{(f_n)^2 + (g_n)^2\} \varrho_n^* \\ \frac{2}{i} (f_n^* g_n \varrho_n - f_n g_n \varrho_n^*) \end{array} \right) \Bigg|_{z=0} \\ \gamma_n &= \left(\begin{array}{c} \frac{H_n + H_n^*}{F_n} \\ \frac{1}{i} \frac{H_n - H_n^*}{F_n} \\ n - 2 \frac{\partial \log F_n}{\partial z} \end{array} \right) \Bigg|_{z=0}, \end{aligned}$$

where we denote by z the variable of the τ functions so that, for example, $F_n = F_n(z)$.

Similarly to [10, §3], we obtain:

Proposition 3.1. $\Phi_n = [T_n, \tilde{N}_n, \tilde{B}_n]$ defines the discrete Frenet frame for the curve γ . Its curvature and torsion angles are given by

$$(3.8) \quad \begin{aligned} \lambda_n &= \frac{1}{i} \log \left(\frac{G_n^* G_{n-1}}{\tilde{G}_n \tilde{G}_{n-1}} \right) \Bigg|_{z=0} = \frac{1}{2i} \log \left(\frac{G_n^* G_{n-1}}{G_n G_{n-1}^*} \right) \Bigg|_{z=0}, \\ \kappa_n &= 2 \arctan \left(\frac{\tilde{G}_n}{\beta F_n} \right) \Bigg|_{z=0}. \end{aligned}$$

3.2. Motion of Kaleidocycle on the osculating plane. Let γ be a Kaleidocycle, that is, a closed curve with a constant torsion angle and a unit segment length. A motion of a Kaleidocycle is identified with a time-parametrised family $\gamma : \mathbb{Z} \times \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}^3$ such that $\gamma(t)$ is a Kaleidocycle for all $t \geq 0$. In [18], a particular motion of a Kaleidocycle satisfying the following conditions is considered:

$$(3.9) \quad \langle \dot{\gamma}_n, \tilde{B}_n \rangle = 0, \quad \langle \dot{\gamma}_n, \dot{\gamma}_n \rangle = \tilde{\rho}^2, \quad |\gamma_{n+1} - \gamma_n| = 1 \quad (0 \leq \forall n \leq k),$$

where $\tilde{\rho} > 0$ is a constant and $\dot{\gamma}_n = \frac{d\gamma_n}{dt}$. These conditions mean that the velocity of each vertex is restricted within the osculating plane and has a constant magnitude. It is shown in [18] that the Frenet-Serret formula (2.6) together with (3.9) demand that there exists a function Θ_n , called the *potential function*, with $\kappa_n = \frac{\Theta_{n+1} - \Theta_{n-1}}{2}$ that satisfies either the *semi-discrete potential mKdV equation*:

$$(3.10) \quad \frac{d}{dt} \left(\frac{\Theta_{n+1} + \Theta_n}{2} \right) = (1 + \cos \lambda) \tilde{\rho} \sin \left(\frac{\Theta_{n+1} - \Theta_n}{2} \right),$$

or the *semi-discrete sine-Gordon equation*

$$(3.11) \quad \frac{d}{dt} \left(\frac{\Theta_{n+1} - \Theta_n}{2} \right) = (1 - \cos \lambda) \tilde{\rho} \sin \left(\frac{\Theta_{n+1} + \Theta_n}{2} \right).$$

The motions corresponding to these equations differ in general. However, the explicit construction discussed in §5 satisfies both equations simultaneously.

As noted in Introduction, for a fixed k , the dimension of the configurations of a Kaleidocycle defined by (2.2) is generally $k - 6$ and (3.9) may look too restrictive. However, for the non-oriented Kaleidocycle with the minimum torsional angle (the Möbius Kaleidocycle), the dimension of the configurations is conjectured to be one regardless of k . An extensive numerical simulation suggests that the motion of the Möbius Kaleidocycle satisfies (2.2). Therefore, (2.2) specifies a special yet essential motion of Kaleidocycles.

4. An explicit formula of closed discrete curve with a constant torsion angle in terms of elliptic theta functions

Based on the representation of curves discussed in §3.1, we define certain τ functions satisfying (3.1)-(3.6) using elliptic theta functions to construct a family of discrete curves with a constant torsion angle. Then, we give sufficient conditions for the curve to be closed and derive an explicit expression for the motion of Kaleidocycles.

4.1. A family of open curves with a constant torsion angle. Throughout this section, j runs through $\{1, 2, 3, 4\}$. Fix $w \in \mathbb{C}$ with $\text{Im}(w) > 0$. The elliptic theta functions $\vartheta_j : \mathbb{C} \rightarrow \mathbb{C}$, are defined as follows [20, 23]:

$$(4.1) \quad \begin{aligned} \vartheta_1(v; w) &= - \sum_{n=-\infty}^{\infty} e^{\pi i w (n + \frac{1}{2})^2 + 2\pi i (n + \frac{1}{2})(v + \frac{1}{2})}, \\ \vartheta_2(v; w) &= \sum_{n=-\infty}^{\infty} e^{\pi i w (n + \frac{1}{2})^2 + 2\pi i (n + \frac{1}{2})v}, \\ \vartheta_3(v; w) &= \sum_{n=-\infty}^{\infty} e^{\pi i w n^2 + 2\pi i n v}, \\ \vartheta_4(v; w) &= \sum_{n=-\infty}^{\infty} e^{\pi i w n^2 + 2\pi i n (v + \frac{1}{2})}. \end{aligned}$$

We often denote $\vartheta_j(v; w)$ by $\vartheta_j(v)$ when w is fixed.

We construct the τ functions satisfying (3.1)-(3.6) using these elliptic theta functions so that the torsion angle given by Proposition 3.1 is a constant. For $x \in \mathbb{C}$ and $y > 0$, we define $d_j(x; y) = \frac{\vartheta_j'(x; iy)}{\vartheta_j(x; iy)}$, where $'$ denotes the derivative with respect to the first variable of elliptic theta functions. For $r, v \in \mathbb{R} \setminus (1/2\mathbb{Z})$ and $y > 0$, we define

$$(4.2) \quad \begin{aligned} \alpha_2 &= \sqrt{\vartheta_3(2r; iy)\vartheta_3(0; iy)}, \quad \beta = \frac{\vartheta_3(iv; iy)}{\vartheta_3(0; iy)}, \quad s_2 = \vartheta_3(2r; iy)\vartheta_1(iv; iy), \\ u_2 &= \vartheta_1\left(-\frac{1}{2}iv + r; iy\right) \vartheta_3\left(-\frac{1}{2}iv + r; iy\right) \vartheta_1\left(\frac{1}{2}iv + r; iy\right) \vartheta_3\left(\frac{1}{2}iv + r; iy\right), \\ \Delta_j &= d_j\left(-\frac{1}{2}iv + r; y\right) - d_j\left(\frac{1}{2}iv + r; y\right), \\ R_j &= \frac{\vartheta_j\left(-\frac{1}{2}iv + r; iy\right)}{\vartheta_j\left(\frac{1}{2}iv + r; iy\right)}. \end{aligned}$$

Noting $\vartheta_j(z; iy)^* = \vartheta_j(z^*; iy)$ when $y > 0$, we have

$$(4.3) \quad \begin{aligned} \alpha_2^* &= \alpha_2, \quad \beta^* = \beta, \quad s_2^* = -s_2, \quad u_2^* = u_2, \\ \Delta_j^* &= -\Delta_j, \quad R_j R_j^* = 1. \end{aligned}$$

Theorem 4.1. Put $\mu_n = ivn + \frac{z}{\Delta_3 - \Delta_1}$ and define

$$(4.4) \quad \begin{aligned} F_n(z) &= \alpha_2 \exp\left(\frac{n\Delta_3}{\Delta_3 - \Delta_1} z\right) \vartheta_2\left(\mu_n - \frac{1}{2}iv\right), \\ f_n(z) &= \vartheta_3\left(-\frac{1}{2}iv + r\right) R_3^n \exp\left(\frac{(n + \frac{1}{2})\Delta_3}{\Delta_3 - \Delta_1} z\right) \vartheta_2(\mu_n + r), \\ g_n(z) &= \vartheta_1\left(\frac{1}{2}iv + r\right) R_1^{-n} \exp\left(\frac{(n + \frac{1}{2})\Delta_3}{\Delta_3 - \Delta_1} z\right) \vartheta_4(\mu_n - r), \\ G_n(z) &= \alpha_2 \frac{\vartheta_1(iv)}{\vartheta_3(0)} R_1^{-n} R_3^n \exp\left(\frac{n\Delta_3}{\Delta_3 - \Delta_1} z\right) \vartheta_4\left(\mu_n - \frac{1}{2}iv\right), \\ H_n(z) &= \frac{u_2}{\alpha_2 s_2} R_1^{-n} R_3^{-n} \exp\left(\frac{n\Delta_3}{\Delta_3 - \Delta_1} z\right) \vartheta_4\left(\mu_n - \frac{1}{2}iv - 2r\right), \\ \tilde{G}_n(z) &= -i\alpha_2 \frac{\vartheta_1(iv)}{\vartheta_3(0)} \exp\left(\frac{n\Delta_3}{\Delta_3 - \Delta_1} z\right) \vartheta_4\left(\mu_n - \frac{1}{2}iv\right). \end{aligned}$$

Then, they satisfy (3.1)-(3.6) to form τ functions. The corresponding discrete curve γ defined by (3.7) has a unit segment length $|\gamma_{n+1} - \gamma_n| = 1$. The torsion angle is given by

$$(4.5) \quad \lambda_n = \frac{1}{i} \log \left(\frac{G_n^*(0)G_{n-1}(0)}{\tilde{G}_n(0)\tilde{G}_{n-1}(0)} \right) = \frac{1}{i} \log (R_1 R_3^{-1}).$$

In particular, this does not depend on n , and γ has a constant torsion angle $\lambda_n = \lambda_0$. Moreover, the signed curvature angle is given by

$$(4.6) \quad \begin{aligned} \kappa_n &= 2 \arctan \left(\frac{\tilde{G}_n(0)}{\beta F_n(0)} \right) \\ &= 2 \arctan \left(-i \frac{\vartheta_1(iv)\vartheta_4(ivn - \frac{1}{2}iv)}{\vartheta_3(iv)\vartheta_2(ivn - \frac{1}{2}iv)} \right), \end{aligned}$$

where we take the primary value for \arctan in $(-\pi/2, \pi/2)$.

4.2. Sufficient conditions for closedness. The constant torsion-angle and unit segment-length curves given in Theorem 4.1 are not necessarily closed. We derive a sufficient condition on the parameters v, r, y for the curve γ to be closed to define a Kaleidocycle. We say a curve γ is closed with a period $k \in \mathbb{N}$ when $\gamma_{n+k} = \gamma_n$ for all n . Rewriting this condition using the τ functions, we obtain

$$(4.7) \quad \begin{aligned} \frac{H_{n+k}}{F_{n+k}} &= \frac{H_n}{F_n}, \\ n+k-2 \frac{\partial \log F_{n+k}}{\partial z} &= n-2 \frac{\partial \log F_n}{\partial z}. \end{aligned}$$

To give general solutions to (4.7) is difficult, but we can utilise the pseudo-period of elliptic theta functions (A.12) to obtain particular solutions. For a fixed k , choose $m \in \mathbb{Z} \setminus \{0\}$ and set $v = my/k$. Then, from (A.12), we see

$$(4.8) \quad \begin{aligned} \frac{H_{n+k}}{F_{n+k}} &= \frac{H_n}{F_n} (R_1 R_3)^{-k} \exp(4mr\pi i) (-1)^m, \\ F_{n+k} &= F_n \exp\left(k \frac{\Delta_3}{\Delta_3 - \Delta_1} z\right) \exp\left[-\pi i \left\{ 2m \left(\mu_n - \frac{1}{2}iv\right) + m^2 w \right\}\right], \end{aligned}$$

where $\mu_n = ivn + \frac{z}{\Delta_3 - \Delta_1}$. Using (4.8), (4.7) can be rewritten as

$$(4.9) \quad \begin{aligned} \Lambda &:= \exp\{\pi i m(4r+1)\} (R_1 R_3)^{-k} - 1 = 0, \\ \Xi &:= i(\Delta_3 + \Delta_1) + 4\pi \frac{v}{y} = 0. \end{aligned}$$

Theorem 4.2. For a fixed k , let $m \in \mathbb{Z} \setminus \{0\}$, $r, v \in \mathbb{R} \setminus (1/2\mathbb{Z})$, and $y > 0$ satisfy $kv = my$ and (4.9). Then, the curve defined by Theorem 4.1 is closed and forms a k -Kaleidocycle with a torsion angle given by (4.5).

Hereafter, we assume that the parameters v, r, y are taken to satisfy the closure condition.

Example 4.3. The classical 6-Kaleidocycle (Figure 1 Left) with $\lambda = \pi/2$ (e.g., see [8]) is obtained by $(v, r, y, m) \approx (0.640, 0.250, 1.279, 3)$.

Remark 4.4. By using (4.9), we see that $\tilde{B}_0 = (-1)^m \tilde{B}_k$. From this, we also see that m and Lk have the same parity. Furthermore, under the transformation $\tilde{B}_n \rightarrow (-1)^{n+1} \tilde{B}_n$, $\tilde{N}_n \rightarrow (-1)^{n+1} \tilde{N}_n$, the curvature becomes $(-1)^{n+1} \kappa_n$ and the torsion angle becomes $\pi + \lambda$. This indicates that this transformation corresponds to the one described in Proposition 5.16.

5. Construction of the motion of a Kaleidocycle

By introducing the deformation parameter to the curve constructed in §4.1, we build an explicit formula for the motion of a Kaleidocycle in Theorem 5.1. We then derive the evolution equations for the vertex positions of the curve and the (potential of the) curvature. Connections to the previously-known isoperimetric deformation of a curve by the semi-discrete mKdV equation [15] and by the semi-discrete potential mKdV equation and the semi-discrete sine-Gordon equation [18] are discussed and summarised in Table 2.

5.1. Explicit formula of the discrete curve with deformation parameter. The following theorem describes a Kaleidocycle together with its motion explicitly in terms of elliptic theta functions.

Theorem 5.1. Denote $\mu_n = ivn + \frac{z}{\Delta_3 - \Delta_1}$. Consider the following τ functions parameterised by $t \in \mathbb{R}$.

$$\begin{aligned}
(5.1) \quad F_n(t, z) &= \alpha_2 \exp\left(\frac{n\Delta_3}{\Delta_3 - \Delta_1}z + \frac{C}{2}tz\right) \vartheta_2\left(\mu_n - \frac{1}{2}iv + it\right), \\
f_n(t, z) &= \vartheta_3\left(-\frac{1}{2}iv + r\right) R_3^n \exp\left(\frac{(n + \frac{1}{2})\Delta_3}{\Delta_3 - \Delta_1}z + \frac{C}{2}tz - \frac{\Gamma}{2}it\right) \vartheta_2(\mu_n + r + it), \\
g_n(t, z) &= \vartheta_1\left(\frac{1}{2}iv + r\right) R_1^{-n} \exp\left(\frac{(n + \frac{1}{2})\Delta_3}{\Delta_3 - \Delta_1}z + \frac{C}{2}tz + \frac{\Gamma}{2}it\right) \vartheta_4(\mu_n - r + it), \\
G_n(t, z) &= \alpha_2 \frac{\vartheta_1(iv)}{\vartheta_3(0)} R_1^{-n} R_3^n \exp\left(\frac{n\Delta_3}{\Delta_3 - \Delta_1}z + \frac{C}{2}tz\right) \vartheta_4\left(\mu_n - \frac{1}{2}iv + it\right), \\
H_n(t, z) &= \frac{u_2}{\alpha_2 s_2} R_1^{-n} R_3^{-n} \exp\left(\frac{n\Delta_3}{\Delta_3 - \Delta_1}z + \frac{C}{2}tz + \Gamma it\right) \vartheta_4\left(\mu_n - \frac{1}{2}iv - 2r + it\right), \\
\tilde{G}_n(t, z) &= -i\alpha_2 \frac{\vartheta_1(iv)}{\vartheta_3(0)} \exp\left(\frac{n\Delta_3}{\Delta_3 - \Delta_1}z + \frac{C}{2}tz\right) \vartheta_4\left(\mu_n - \frac{1}{2}iv + it\right),
\end{aligned}$$

where $C, \Gamma \in \mathbb{R}$ are constants. Then, they satisfy (3.1)-(3.6) to form τ functions. For each t , the corresponding discrete curve γ (3.7) has a unit segment length $|\gamma_{n+1} - \gamma_n| = 1$. The torsion angle is given by

$$(5.2) \quad \lambda_n = \frac{1}{i} \log \left(\frac{G_n^*(t, 0) G_{n-1}(t, 0)}{\tilde{G}_n(t, 0) \tilde{G}_{n-1}(t, 0)} \right) = \frac{1}{i} \log (R_1 R_3^{-1}),$$

In particular, this depends neither on n nor on t , and γ has a constant torsion angle $\lambda_n(t) = \lambda_0(0)$. Moreover, the signed curvature angle is given by

$$\begin{aligned}
(5.3) \quad \kappa_n(t) &= 2 \arctan \left(\frac{\tilde{G}_n(t, 0)}{\beta F_n(t, 0)} \right) \\
&= 2 \arctan \left(-i \frac{\vartheta_1(iv) \vartheta_4(ivn - \frac{1}{2}iv + it)}{\vartheta_3(iv) \vartheta_2(ivn - \frac{1}{2}iv + it)} \right),
\end{aligned}$$

where we take the primary value for \arctan in $(-\pi/2, \pi/2)$. If r, v, y are chosen as in Theorem 4.2, γ is closed for every t .

Proposition 5.2. *The parameter C contributes only to the parallel translation of the curve along the Z -axis, whereas Γ contributes only to the rotation of the curve around the Z -axis.*

Proof. For $A \in \mathbb{C}$ and $B : \mathbb{C} \rightarrow \mathbb{C}$, we have

$$\begin{aligned}
 (5.4) \quad & \begin{pmatrix} \cos \Gamma t & -\sin \Gamma t \\ \sin \Gamma t & \cos \Gamma t \end{pmatrix} \begin{pmatrix} A + A^* \\ \frac{1}{i}(A - A^*) \end{pmatrix} \\
 &= \frac{1}{2} \begin{pmatrix} e^{i\Gamma t} + e^{-i\Gamma t} & -\frac{1}{i}(e^{i\Gamma t} - e^{-i\Gamma t}) \\ \frac{1}{i}(e^{i\Gamma t} - e^{-i\Gamma t}) & e^{i\Gamma t} + e^{-i\Gamma t} \end{pmatrix} \begin{pmatrix} A + A^* \\ \frac{1}{i}(A - A^*) \end{pmatrix} \\
 &= \begin{pmatrix} e^{i\Gamma t} A + e^{-i\Gamma t} A^* \\ \frac{1}{i}(e^{i\Gamma t} A - e^{-i\Gamma t} A^*) \end{pmatrix}
 \end{aligned}$$

and

$$(5.5) \quad 2 \frac{\partial \log \left(e^{\frac{C}{2} t z} B \right)}{\partial z} = Ct + 2 \frac{\partial \log B}{\partial z}.$$

The assertion follows from the definition (3.7). □

We will make convenient choices of these parameters to derive deformation equations.

Table 1 summarises the relationship between various variables. The first column lists the parameters that determine the curve γ through the variables listed in the first row. A circle in the table indicates that the variable shown in the first row is dependent on the parameter shown in the first column.

Table 1. Dependency of variables on the parameters $v, r, y, k, m, t, C, \Gamma$.

	γ	κ	λ	Λ	Ξ	u_2	β	s_2	R_j	Δ_j	α_2	$\beta_{1,2}$	$\epsilon_{1,2}$	ω	$\delta_{1,2,3}$
v	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○
r	○		○	○	○	○		○	○	○	○	○	○	○	○
y	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○
k	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○
m	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○
t	○	○													
Γ	○											○			
C	○														
	(3.7)	(5.3)	(5.2)	(4.9)	(4.9)	(4.2)	(4.2)	(4.2)	(4.2)	(4.2)	(4.2)	(5.6)	(5.6)	(5.6)	(5.6)

The constructed motion has a periodic curvature. The next proposition follows from (A.12).

Proposition 5.3. *We have $\kappa_n(t + y) = -\kappa_n(t)$. In other words, $2y$ is a period of the motion.*

Remark 5.4. There are certain relations between the curves constructed in Theorem 5.1 with different sets of parameters. The transformation of parameters $(v, m) \rightarrow (-v, -m)$ preserves the closure condition. We see that $\kappa_n(t; -v) = -\kappa_n(-t; v)$, $\lambda(-v, r) = -\lambda(v, r)$. In what follows, We always choose (v, m) or $(-v, -m)$ so that $\lambda \geq 0$.

5.2. Motion of vertex positions. The construction in Theorem 5.1 involves parameters C, Γ that contribute only to the global rigid transformation of the curve. In general, the velocity of the vertex position $\dot{\gamma}_n$ has a binormal component, and its magnitude is not constant as described in Theorem 5.6. However, with the right choice of C, Γ , curves that satisfy (3.9) are obtained as described in Theorem 5.9.

We derive bilinear equations satisfied by the τ functions given by Theorem 5.1 to calculate $\dot{\gamma}_n$.

For $\Gamma \in \mathbb{R}$, $r, v \in \mathbb{R} \setminus (1/2\mathbb{Z})$ and $y > 0$, we define

$$\begin{aligned}
(5.6) \quad \omega &= \frac{\vartheta_1(2r; iy) \vartheta_3(0; iy)}{\vartheta_3(2r; iy) \vartheta_1(iv; iy)}, \\
\delta_1 &= \frac{\vartheta_3(\frac{1}{2}iv + r; iy)^2 \vartheta_3(-\frac{1}{2}iv + r; iy)^2}{\vartheta_3(2r; iy) \vartheta_3(0; iy)^3}, \\
\delta_2 &= \frac{\vartheta_1(\frac{1}{2}iv + r; iy)^2 \vartheta_3(-\frac{1}{2}iv + r; iy)^2}{\vartheta_1(iv; iy)^2 \vartheta_3(2r; iy) \vartheta_3(0; iy)}, \\
\delta_3 &= \frac{-iu_2}{s_2 \vartheta_3(0; iy) \vartheta_1(iv; iy)} \left[\frac{\vartheta_1'(\frac{1}{2}iv + r; iy) \vartheta_3(-\frac{1}{2}iv + r; iy)}{\vartheta_1(-\frac{1}{2}iv + r; iy) \vartheta_3(\frac{1}{2}iv + r; iy)} \right. \\
&\quad \left. + \frac{\vartheta_1(\frac{1}{2}iv + r; iy) \vartheta_3'(-\frac{1}{2}iv + r; iy)}{\vartheta_1(-\frac{1}{2}iv + r; iy) \vartheta_3(\frac{1}{2}iv + r; iy)} \right], \\
\epsilon_1 &= \frac{2i}{\Delta_3 - \Delta_1} \frac{\vartheta_3''(0; iy)}{\vartheta_3(0; iy)}, \\
\epsilon_2 &= \frac{2i}{\Delta_3 - \Delta_1} \frac{\vartheta_1'(0; iy)^2}{\vartheta_1(iv; iy)^2}, \\
\beta_1 &= \frac{1}{2}(\delta_3 + \delta_3^*) + \frac{1}{2}i\Gamma(\delta_2 - \delta_2^*), \\
\beta_2 &= \frac{1}{2}i(\delta_3 - \delta_3^*) - \frac{1}{2}\Gamma(\delta_2 + \delta_2^*).
\end{aligned}$$

Then, it follows that

$$(5.7) \quad \omega^* = -\omega, \quad \delta_1^* = \delta_1, \quad \epsilon_1^* = \epsilon_1, \quad \epsilon_2^* = \epsilon_2, \quad \beta_1^* = \beta_1, \quad \beta_2^* = \beta_2.$$

A proof of the following proposition is given in Appendix A.

Proposition 5.5. *The τ functions (5.1) satisfy the following bilinear equations.*

$$\begin{aligned}
(5.8) \quad & f_n f_{n-1}^* + g_n^* g_{n-1} = \beta F_n F_n + \omega G_n G_n^*, \\
(5.9) \quad & g_n^* g_{n-1} - g_n g_{n-1}^* = f_n f_{n-1}^* - f_n^* f_{n-1} = \omega G_n G_n^*, \\
(5.10) \quad & \delta_2 f_n^* g_{n-1} - \delta_2^* f_{n-1}^* g_n = H_n F_n, \\
(5.11) \quad & \delta_1 F_n F_n + \delta_2 G_n G_n^* = f_n f_{n-1}^*, \\
(5.12) \quad & (\beta - \delta_1) F_n F_n + (\omega - \delta_2) G_n G_n^* = g_n^* g_{n-1}, \\
(5.13) \quad & D_t H_n \cdot F_n = (\delta_3 + i\Gamma \delta_2) f_n^* g_{n-1} + (\delta_3^* - i\Gamma \delta_2^*) f_{n-1}^* g_n, \\
(5.14) \quad & D_t D_z F_n \cdot F_n = (\epsilon_1 + C) F_n F_n + \epsilon_2 G_n G_n^*.
\end{aligned}$$

By using these equations, we obtain the evolution equation of the vertex positions of the curve:

Theorem 5.6. *The curve γ constructed in Theorem 5.1 satisfies the following deformation equation.*

$$\begin{aligned}
(5.15) \quad \dot{\gamma}_n(t) &= 2\beta \left[\beta_1 T_n - \beta_1 \tan \frac{\kappa_n}{2} \tilde{N}_n + \beta_2 \tan \frac{\kappa_n}{2} \tilde{B}_n \right] \\
&\quad + \begin{pmatrix} 0 \\ 0 \\ 2\beta_1(\beta - 2\delta_1) - \epsilon_1 - C \end{pmatrix}.
\end{aligned}$$

This is a deformation equation similar to the one presented in [15]. Starting from (5.15), the compatibility condition of the Frenet frame yields the *semi-discrete mKdV equation*.

Proposition 5.7. *When the curve with a unit segment length and a constant torsion angle deforms according to the deformation equation (5.15), the following holds.*

- (i) *The segment length is preserved.*

(ii) κ satisfies the following semi-discrete mKdV equation:

$$(5.16) \quad \dot{\kappa}_n = (2\beta\beta_1 \cos \lambda + 2\beta\beta_2 \sin \lambda) \left(\tan \frac{\kappa_{n+1}}{2} - \tan \frac{\kappa_{n-1}}{2} \right).$$

Furthermore, if the left-hand-side of the equation (5.27) does not depend on n , the following holds:

(iii) The torsion angle is preserved.

Proof. The proofs of (i) and (ii) are given by the same calculation as in Section 5 of [15]. We prove (iii). From the deformation equation (5.15) and the discrete Frenet-Serret formula (2.6), we see that

$$(5.17) \quad \begin{aligned} -\dot{\lambda} \sin \lambda &= \langle \dot{\tilde{B}}_n, \tilde{B}_{n+1} \rangle + \langle \tilde{B}_n, \dot{\tilde{B}}_{n+1} \rangle \\ &= -\sin \lambda (2\beta\beta_1 \sin \lambda - 2\beta\beta_2 \cos \lambda) \times \left(\frac{1}{\sin \kappa_{n+1}} \tan \frac{\kappa_{n+2}}{2} \right. \\ &\quad \left. + \frac{1}{\tan \kappa_{n+1}} \tan \frac{\kappa_n}{2} - \frac{1}{\tan \kappa_n} \tan \frac{\kappa_{n+1}}{2} - \frac{1}{\sin \kappa_n} \tan \frac{\kappa_{n-1}}{2} \right), \end{aligned}$$

holds, which vanishes because of (5.27). Therefore $\dot{\lambda} = 0$. \square

Remark 5.8. Proposition 5.7 (iii) does not hold in general without the assumption on (5.27). We can construct a motion of a Kaleidocycle satisfying (5.15) such that the left-hand side of (5.27) depends both on n and t . This suggests that the conservation of the left-hand side of the recurrence equation (5.27) plays a crucial role in the preservation of the torsion angle.

By adjusting the global motion of γ by C, Γ , the deformation equation of γ can be made to satisfy the equation considered in the previous studies presented in §3.2.

Theorem 5.9. Set C, Γ to be

$$(5.18) \quad \begin{aligned} \Gamma_+ &= \frac{-(\delta_3 + \delta_3^*) + (\delta_3 e^{i\lambda} + \delta_3^* e^{-i\lambda})}{i\omega}, \\ C_+ &= -\epsilon_1 - i\Gamma_+ \left(\omega(2\delta_1 - \beta) - \frac{\delta_2 + \delta_2^*}{\omega} \right) \\ &\quad - (\delta_3 + \delta_3^*)(2\delta_1 - \beta) + \frac{\delta_3 - \delta_3^*}{\omega}, \\ \text{or} \\ \Gamma_- &= \frac{-(\delta_3 + \delta_3^*) - (\delta_3 e^{i\lambda} + \delta_3^* e^{-i\lambda})}{i\omega}, \\ C_- &= -\epsilon_1 - i\Gamma_- \left(\omega(2\delta_1 - \beta) - \frac{\delta_2 + \delta_2^*}{\omega} \right) \\ &\quad - (\delta_3 + \delta_3^*)(2\delta_1 - \beta) + \frac{\delta_3 - \delta_3^*}{\omega}. \end{aligned}$$

Then, the curve constructed in Theorem 5.1 satisfies the following deformation equations.

$$(5.19) \quad \langle \dot{\gamma}_n, \tilde{B}_n \rangle = 0, \quad \langle \dot{\gamma}_n, \dot{\gamma}_n \rangle = \rho^2,$$

where $\rho^2 > 0$ is given by

$$(5.20) \quad \rho^2 = \rho_+^2 := \left(i \frac{\vartheta_1'(0)}{\vartheta_1(iv)} \frac{u_2}{\vartheta_3(r)^2 \vartheta_1(r)^2} \right)^2$$

when $(C, \Gamma) = (C_+, \Gamma_+)$ and

$$(5.21) \quad \rho^2 = \rho_-^2 := \left(-i \frac{\vartheta_1'(0)}{\vartheta_1(iv)} \frac{u_2}{\vartheta_4(r)^2 \vartheta_2(r)^2} \right)^2$$

when $(C, \Gamma) = (C_-, \Gamma_-)$.

Remark 5.10. A discrete analogue of the constant negative curvature surface, called the discrete K-surface, is introduced in [2]. It is discussed that certain discrete-time deformation of discrete curves with a constant torsion angle generates discrete K-surfaces in [15]. Deformation equation (3.9) can be seen as a continuous-time analogue of the discrete-time deformation. Theorem 5.9 shows that specialising (C, Γ) to (C_-, Γ_-) or (C_+, Γ_+) , the curve constructed in Theorem 5.1 deforms according to (3.9). Therefore, its sweeping surface forms a semi-discrete analogue of the discrete K-surface (see figures in §6).

Since the curvature and torsion angles uniquely determine a curve up to a rigid transformation, Proposition 5.3 shows that the motion constructed in Theorem 5.1 is periodic up to a global rigid transform. Here, we control the global orientation by choosing the parameters C, Γ appropriately so that the motion becomes periodic in terms of vertex positions.

Proposition 5.11. *Let $l \in \mathbb{Z}$. If we set*

$$(5.22) \quad C = \frac{4\pi i}{y} \frac{1}{\Delta_3 - \Delta_1},$$

$$(5.23) \quad \Gamma = -\frac{\pi}{y} (4r + 2l + 1),$$

we have

$$(5.24) \quad \gamma_n(t + y) = \gamma_n(t),$$

for any $n \in \mathbb{Z}$ and $t \in \mathbb{R}$.

Proof. First, we prove the periodicity for the Z -component. Denote the Z -component of γ by γ_{nZ} . From the pseudo-period of elliptic theta functions (A.12), we see

$$(5.25) \quad \begin{aligned} & \gamma_{nZ}(t + y) - \gamma_{nZ}(t) \\ &= -2 \frac{\partial}{\partial z} \log \left\{ \exp\left(\frac{C}{2}yz\right) \frac{\vartheta_2\left(iv\left(n - \frac{1}{2}\right) + \frac{z}{\Delta_3 - \Delta_1} + it + iy\right)}{\vartheta_2\left(iv\left(n - \frac{1}{2}\right) + \frac{z}{\Delta_3 - \Delta_1} + it\right)} \right\} \\ &= -2 \frac{\partial}{\partial z} \log \exp \left\{ \frac{C}{2}yz - \pi i \left(2iv\left(n - \frac{1}{2}\right) + \frac{2z}{\Delta_3 - \Delta_1} + 2it \right) + \pi y \right\} \\ &= -2 \left(\frac{C}{2}y - \frac{2\pi i}{\Delta_3 - \Delta_1} \right) = 0. \end{aligned}$$

For the X and Y components, from (3.7) we have to show $\frac{H_n(t+y)}{F_n(t+y)} \frac{F_n(t)}{H_n(t)} = 1$.

If we write $P_n = iv\left(n - \frac{1}{2}\right) + \frac{z}{\Delta_3 - \Delta_1} + it$, the left-hand side can be computed using (A.12) as

$$(5.26) \quad \begin{aligned} & \exp(i\Gamma y) \frac{\vartheta_4(P_n - 2r + iy) \vartheta_2(P_n)}{\vartheta_2(P_n + iy) \vartheta_4(P_n - 2r)} \\ &= -\exp(i\Gamma y) \frac{\exp\{-\pi i(2X_n - 4r + iy)\}}{\exp\{-\pi i(2X_n + iy)\}} \\ &= -\exp\{i(\Gamma y + 4\pi r)\} = 1. \end{aligned}$$

□

5.3. Evolution of curvature. The curve constructed in Theorem 5.1 reveals an interesting recurrence equation for κ .

Proposition 5.12. *The signed curvature angle of the curve obtained in Theorem 5.1 satisfies the following recurrence equation for any t and n :*

$$(5.27) \quad \tan \frac{\kappa_{n+1}}{2} \tan \frac{\kappa_{n-1}}{2} - \tan^2 \frac{\kappa_n}{2} + \tan \frac{\kappa_{n+1}}{2} \tan^2 \frac{\kappa_n}{2} \tan \frac{\kappa_{n-1}}{2} = -\frac{\vartheta_1(iv)^4}{\vartheta_3(iv)^4}.$$

The right-hand side of this equation depends neither on n nor on t . Therefore, the quantity on the left-hand side can be considered a conserved quantity of the signed curvature angle.

The evolution of the signed curvature angle satisfies the semi-discrete mKdV equation in Proposition 5.7.

Theorem 5.13. *The signed curvature angle (5.3) satisfies*

$$(5.28) \quad \dot{\kappa}_n = 2\beta^2 \eta_3 \left(\tan \frac{\kappa_{n+1}}{2} - \tan \frac{\kappa_{n-1}}{2} \right),$$

where

$$(5.29) \quad \eta_3 = i \frac{\vartheta_1(iv) \vartheta_3(iv)}{\vartheta_1(2iv) \vartheta_3(0)} (d_1(iv) - d_3(iv)).$$

Combined with Proposition 5.7, (5.6), and the fact that $e^{i\lambda} \delta_2 \in \mathbb{R}$, we see

$$(5.30) \quad 2\beta^2 \eta_3 = (2\beta\beta_1 \cos \lambda + 2\beta\beta_2 \sin \lambda).$$

5.4. Potential function of curvature. A family of functions Θ_n parameterised by $n \in \mathbb{Z}$ is said to be a potential of the curvature κ if the equation $2\kappa_n = \Theta_{n+1} - \Theta_{n-1}$ holds. When γ deforms by (5.19), the potential function Θ is known to satisfy the semi-discrete potential mKdV equation or the semi-discrete sine-Gordon equation [18]. Here, we derive the evolution equations satisfied by the potential function for the curve constructed by Theorem 5.1.

We construct a potential function which simultaneously satisfy (3.10) and (3.11) in terms of elliptic theta functions.

Theorem 5.14. *For fixed v, y , define the function Θ as follows.*

$$(5.31) \quad \Theta_n(t) = \Theta_n(t; v, y) = 4 \arctan \frac{\vartheta_1\left(\frac{v}{y}\left(n - \frac{1}{2}\right) + \frac{t}{y}; \frac{2i}{y}\right)}{\vartheta_4\left(\frac{v}{y}\left(n - \frac{1}{2}\right) + \frac{t}{y}; \frac{2i}{y}\right)}.$$

Then we see that κ defined in (5.3) satisfies

$$(5.32) \quad \kappa_n = \frac{\Theta_{n+1} - \Theta_{n-1}}{2}.$$

The potential function simultaneously solves the semi-discrete potential mKdV equation and the semi-discrete sine-Gordon equation.

Theorem 5.15. *Let $v \in \mathbb{R} \setminus (1/2\mathbb{Z})$ and $y > 0$. The potential function Θ defined by (5.31) simultaneously satisfies the semi-discrete potential mKdV and sine-Gordon equations*

$$(5.33) \quad \frac{d}{dt} \left(\frac{\Theta_{n+1} + \Theta_n}{2} \right) = \frac{L_1(v, y)}{y} \sin \left(\frac{\Theta_{n+1} - \Theta_n}{2} \right),$$

$$(5.34) \quad \frac{d}{dt} \left(\frac{\Theta_{n+1} - \Theta_n}{2} \right) = \frac{L_2(v, y)}{y} \sin \left(\frac{\Theta_{n+1} + \Theta_n}{2} \right),$$

where $Q_1, Q_2, L_1, L_2 \in \mathbb{R}$ are defined by

$$(5.35) \quad Q_1(v, y) = \frac{\vartheta_1'\left(0; \frac{2i}{y}\right) \vartheta_1\left(\frac{v}{y}; \frac{2i}{y}\right)}{\vartheta_4\left(0; \frac{2i}{y}\right) \vartheta_4\left(\frac{v}{y}; \frac{2i}{y}\right)}, \quad Q_2(v, y) = \frac{\vartheta_1'\left(0; \frac{2i}{y}\right) \vartheta_4\left(\frac{v}{y}; \frac{2i}{y}\right)}{\vartheta_1\left(\frac{v}{y}; \frac{2i}{y}\right) \vartheta_4\left(0; \frac{2i}{y}\right)},$$

$$(5.36) \quad L_1(v, y) = d_1\left(\frac{v}{y}; \frac{2}{y}\right) - d_4\left(\frac{v}{y}; \frac{2}{y}\right) + Q_1(v, y) + Q_2(v, y),$$

$$(5.37) \quad L_2(v, y) = d_1\left(\frac{v}{y}; \frac{2}{y}\right) - d_4\left(\frac{v}{y}; \frac{2}{y}\right) - Q_1(v, y) - Q_2(v, y).$$

The following shows two equations are related by a parameter shift.

Proposition 5.16. *By the transformation of parameters $v \rightarrow v + y$ and $r \rightarrow \frac{1}{2} - r$, the signed curvature angle κ , the torsion angles λ , the potential functions Θ , the coefficients of the semi-discrete mKdV equation L_1/y and of the semi-discrete sine-Gordon equation L_2/y , the normal \tilde{N} and the binormal \tilde{B} are transformed as*

$$(5.38) \quad \kappa_n(t) \rightarrow (-1)^{n+1} \kappa_n\left(t - \frac{y}{2}\right), \quad \lambda \rightarrow \pi + \lambda,$$

$$(5.39) \quad \Theta_n(t) \rightarrow (-1)^n \Theta_n\left(t - \frac{y}{2}\right),$$

$$(5.40) \quad \frac{L_1}{y} \rightarrow \frac{L_2}{y}, \quad \frac{L_2}{y} \rightarrow \frac{L_1}{y},$$

$$(5.41) \quad \tilde{N}_n(t) \rightarrow (-1)^{n+1} \tilde{N}_n\left(t - \frac{y}{2}\right), \quad \tilde{B}_n(t) \rightarrow (-1)^{n+1} \tilde{B}_n\left(t - \frac{y}{2}\right).$$

The transformation $\Theta_n \rightarrow \tilde{\Theta}_n := (-1)^n \Theta_n$ exchanges a solution of the semi-discrete potential mKdV equation (5.33) with one of the semi-discrete sine-Gordon equation (5.34). Proposition 5.16 shows that this equation-level transformation is also achieved at the solution level by transforming the parameters.

Remark 5.17. Rewriting (5.28) in terms of the potential function Θ , we obtain

$$(5.42) \quad \frac{d}{dt} \Theta_n = 4\beta^2 \eta_3 \tan\left(\frac{\Theta_{n+1} - \Theta_{n-1}}{4}\right),$$

which is also called the *semi-discrete potential mKdV equation* [6, 12, 13]. In other words, the potential function Θ defined by (5.31) is a simultaneous solution of three differential-difference equations, (5.33), (5.34) and (5.42). It does not usually happen that these equations are simultaneously solved. For example, it can be checked that the 2 and 3-soliton solutions of the semi-discrete sine-Gordon equation constructed in [6, 7] are not solutions of the semi-discrete potential mKdV equation.

The following describes an explicit relation between our construction and the previous work [18] reviewed in §3.2.

Proposition 5.18. *When C_+ and Γ_+ are chosen as in (5.18), we have*

$$(5.43) \quad \frac{L_1(v, y)}{y} \frac{1}{1 + \cos \lambda} = i \frac{\vartheta_1'(0)}{\vartheta_1(iv)} \frac{u_2}{\vartheta_3(r)^2 \vartheta_1(r)^2} = \rho_+.$$

Similarly, when C_- and Γ_- are chosen as in (5.18), we have

$$(5.44) \quad \frac{L_2(v, y)}{y} \frac{1}{1 - \cos \lambda} = -i \frac{\vartheta_1'(0)}{\vartheta_1(iv)} \frac{u_2}{\vartheta_4(r)^2 \vartheta_2(r)^2} = \rho_-.$$

Note the difference in approaches of [18] and this work; [18] starts with a geometric property observed in the motion of Kaleidocycles, and derive evolution equations, whereas we aim to construct a particular motion of Kaleidocycles and investigate the equations satisfied by them. They turn out to arrive at the same place; the latter gives an explicit solution to the equations derived in the former.

Table 2 summarises the relationships among the three deformation equations.

6. Numerical Experiments

In this section, we present examples of Kaleidocycles constructed by Theorem 4.2 and the semi-discrete K-surfaces obtained as their motion trajectory as explained in Remark 5.10.

Numerical experiments using our computer codes [16] suggest that the 8-, 9- and 15-Kaleidocycles in Figs. 2, 3, and 4 have the minimum torsion angles among non-orientable 8-, 9-, and 15-Kaleidocycles, respectively [17]. In other words, they are the Möbius Kaleidocycles [19].

For Fig. 5, the torsion angle is larger than the above minimum one. However, it has a self-linking number 7.5. As the self-linking number of a Kaleidocycle is a half-integer, these 15-Kaleidocycles depicted in Figs. 4 and 5 belong to different connected components of the configuration space of 15-Kaleidocycles. We numerically find that the latter attains the local minima of the torsion angle. We present the following two conjectures based on our extensive numerical experiments:

Table 2. Relation between the three deformation equations

Values of C, Γ	Deformation equation of the curve	Equation for potential function
$C, \Gamma \in \mathbb{R}$	$\dot{\gamma}_n = \begin{pmatrix} 0 \\ 0 \\ 2\beta_1(\beta - 2\delta_1) - \epsilon_1 - C \end{pmatrix} + 2\beta \left(\beta_1 T_n - \beta_1 \tan \frac{\kappa_n}{2} \tilde{N}_n + \beta_2 \tan \frac{\kappa_n}{2} \tilde{B}_n \right)$	$\dot{\Theta}_n = (4\beta\beta_1 \cos \lambda + 4\beta\beta_2 \sin \lambda) \times \tan \left(\frac{\Theta_{n+1} - \Theta_{n-1}}{4} \right)$
\downarrow Specialise the deformation equation by choosing rigid transformation parameters C, Γ .		
C_+, Γ_+	$\langle \dot{\gamma}_n, \tilde{B}_n \rangle = 0, \quad \langle \dot{\gamma}_n, \dot{\gamma}_n \rangle = \rho_+^2$	$\frac{\dot{\Theta}_{n+1} + \dot{\Theta}_n}{2} = (1 + \cos \lambda) \rho_+ \times \sin \left(\frac{\Theta_{n+1} - \Theta_n}{2} \right)$
\updownarrow By replacing the parameters of the solution with $v \rightarrow v + y, r \rightarrow 1/2 - r$, these two deformation equations transform to each other.		
C_-, Γ_-	$\langle \dot{\gamma}_n, \tilde{B}_n \rangle = 0, \quad \langle \dot{\gamma}_n, \dot{\gamma}_n \rangle = \rho_-^2$	$\frac{\dot{\Theta}_{n+1} - \dot{\Theta}_n}{2} = (1 - \cos \lambda) \rho_- \times \sin \left(\frac{\Theta_{n+1} + \Theta_n}{2} \right)$

Conjecture 6.1. Any Kaleidocycle obtained by Theorem 4.2 has a minimal or maximal torsion angle. In other words, the projection from the space defined by (2.2) onto the λ -coordinate attains a critical value.

Conjecture 6.2. The self-linking number of the closed curve γ constructed in Theorem 4.2 is $m/2$ when the curve is unknotted.

As briefly discussed in §1, these Kaleidocycles with critical torsion angles numerically exhibit special properties, including the single degree of freedom [17, 19, 26]. Our construction produces this interesting family of Kaleidocycles and may shed light on the rigorous proof for some of their unique properties.




Figure 2. (Left) An example of 8-Kaleidocycle with $(v, r, y, m) \approx (0.400, 0.717, 1.067, 3)$ and $\cos \lambda \approx 0.4700$. (Middle) 8-Kaleidocycle as a discrete curve. (Right) the semi-discrete K-surface obtained as its trajectory.

A. Appendix

In the Appendix, we prove the theorems and propositions in the paper by using the properties of elliptic theta functions listed below.




Figure 3. (Left) An example of 9-Kaleidocycle with $(v, r, y, m) \approx (0.342, 0.289, 1.027, 3)$ and $\cos \lambda \approx 0.5852$. (Middle) 9-Kaleidocycle as a discrete curve. (Right) the semi-discrete K-surface obtained as its trajectory.




Figure 4. (Left) An example of 15-Kaleidocycle with $(v, r, y, m) \approx (0.189, 0.300, 0.947, 3)$ and $\cos \lambda \approx 0.8533$. (Middle) 15-Kaleidocycle as a discrete curve. (Right) the semi-discrete K-surface obtained as its trajectory.




Figure 5. (Left) An example of 15-Kaleidocycle with $(v, r, y, m) \approx (0.401, 0.689, 1.204, 5)$ and $\cos \lambda \approx 0.6497$. (Middle) 15-Kaleidocycle as a discrete curve. (Right) the semi-discrete K-surface obtained as its trajectory.

A.1. Functional identities of elliptic theta functions. We collect some properties of elliptic theta functions which we rely heavily on our calculations.

Four term identities. Let

$$(A.1) \quad \begin{aligned} X_1 &= \frac{1}{2}(X + Y + U + V), & Y_1 &= \frac{1}{2}(X + Y - U - V), \\ U_1 &= \frac{1}{2}(X - Y + U - V), & V_1 &= \frac{1}{2}(X - Y - U + V), \end{aligned}$$

then for any $X, Y, U, V \in \mathbb{C}$, elliptic theta functions satisfy the following identities [20, 23].

$$\begin{aligned}
\text{(A.2)} \quad & \vartheta_2(X_1)\vartheta_2(Y_1)\vartheta_3(U_1)\vartheta_3(V_1) - \vartheta_4(X_1)\vartheta_4(Y_1)\vartheta_1(U_1)\vartheta_1(V_1) \\
& = \vartheta_2(X)\vartheta_2(Y)\vartheta_3(U)\vartheta_3(V) - \vartheta_4(X)\vartheta_4(Y)\vartheta_1(U)\vartheta_1(V), \\
\text{(A.3)} \quad & \vartheta_4(X_1)\vartheta_3(Y_1)\vartheta_1(U_1)\vartheta_2(V_1) - \vartheta_2(X_1)\vartheta_1(Y_1)\vartheta_3(U_1)\vartheta_4(V_1) \\
& = \vartheta_4(X)\vartheta_3(Y)\vartheta_1(U)\vartheta_2(V) - \vartheta_2(X)\vartheta_1(Y)\vartheta_3(U)\vartheta_4(V), \\
\text{(A.4)} \quad & \vartheta_2(X_1)\vartheta_2(Y_1)\vartheta_4(U_1)\vartheta_4(V_1) + \vartheta_1(X_1)\vartheta_1(Y_1)\vartheta_3(U_1)\vartheta_3(V_1) \\
& = \vartheta_4(X)\vartheta_4(Y)\vartheta_2(U)\vartheta_2(V) - \vartheta_3(X)\vartheta_3(Y)\vartheta_1(U)\vartheta_1(V), \\
\text{(A.5)} \quad & \vartheta_3(X_1)\vartheta_3(Y_1)\vartheta_2(U_1)\vartheta_2(V_1) + \vartheta_1(X_1)\vartheta_1(Y_1)\vartheta_4(U_1)\vartheta_4(V_1) \\
& = \vartheta_3(X)\vartheta_3(Y)\vartheta_2(U)\vartheta_2(V) + \vartheta_1(X)\vartheta_1(Y)\vartheta_4(U)\vartheta_4(V), \\
\text{(A.6)} \quad & \vartheta_4(X_1)\vartheta_4(Y_1)\vartheta_4(U_1)\vartheta_4(V_1) - \vartheta_1(X_1)\vartheta_1(Y_1)\vartheta_1(U_1)\vartheta_1(V_1) \\
& = \vartheta_4(X)\vartheta_4(Y)\vartheta_4(U)\vartheta_4(V) - \vartheta_1(X)\vartheta_1(Y)\vartheta_1(U)\vartheta_1(V), \\
\text{(A.7)} \quad & \vartheta_4(X_1)\vartheta_4(Y_1)\vartheta_1(U_1)\vartheta_1(V_1) - \vartheta_1(X_1)\vartheta_1(Y_1)\vartheta_4(U_1)\vartheta_4(V_1) \\
& = \vartheta_4(X)\vartheta_4(Y)\vartheta_1(U)\vartheta_1(V) - \vartheta_1(X)\vartheta_1(Y)\vartheta_4(U)\vartheta_4(V), \\
\text{(A.8)} \quad & \vartheta_4(X_1)\vartheta_4(Y_1)\vartheta_1(U_1)\vartheta_1(V_1) + \vartheta_2(X_1)\vartheta_2(Y_1)\vartheta_3(U_1)\vartheta_3(V_1) \\
& = \vartheta_3(X)\vartheta_3(Y)\vartheta_2(U)\vartheta_2(V) - \vartheta_1(X)\vartheta_1(Y)\vartheta_4(U)\vartheta_4(V), \\
\text{(A.9)} \quad & \vartheta_3(X_1)\vartheta_3(Y_1)\vartheta_1(U_1)\vartheta_1(V_1) - \vartheta_1(X_1)\vartheta_1(Y_1)\vartheta_3(U_1)\vartheta_3(V_1) \\
& = \vartheta_3(X)\vartheta_3(Y)\vartheta_1(U)\vartheta_1(V) - \vartheta_1(X)\vartheta_1(Y)\vartheta_3(U)\vartheta_3(V), \\
\text{(A.10)} \quad & \vartheta_3(X_1)\vartheta_3(Y_1)\vartheta_4(U_1)\vartheta_4(V_1) - \vartheta_1(X_1)\vartheta_1(Y_1)\vartheta_2(U_1)\vartheta_2(V_1) \\
& = \vartheta_3(X)\vartheta_3(Y)\vartheta_4(U)\vartheta_4(V) - \vartheta_1(X)\vartheta_1(Y)\vartheta_2(U)\vartheta_2(V).
\end{aligned}$$

Shifts by periods and half-periods. For any $X \in \mathbb{C}$, $n \in \mathbb{Z}$ and $y > 0$, the following holds [20, 23].

$$\begin{aligned}
\text{(A.11)} \quad & \vartheta_1(X+n) = (-1)^n \vartheta_1(X), \quad \vartheta_4(X+n) = \vartheta_4(X). \\
& \vartheta_1(X+niy; iy) = (-1)^n \exp\{-\pi i(2nX+n^2iy)\} \vartheta_1(X; iy), \\
\text{(A.12)} \quad & \vartheta_2(X+niy; iy) = \exp\{-\pi i(2nX+n^2iy)\} \vartheta_2(X; iy), \\
& \vartheta_3(X+niy; iy) = \exp\{-\pi i(2nX+n^2iy)\} \vartheta_3(X; iy), \\
& \vartheta_4(X+niy; iy) = (-1)^n \exp\{-\pi i(2nX+n^2iy)\} \vartheta_4(X; iy). \\
\text{(A.13)} \quad & \vartheta_1\left(X+\frac{1}{2}\right) = \vartheta_2(X), \quad \vartheta_2\left(X+\frac{1}{2}\right) = -\vartheta_1(X), \\
& \vartheta_3\left(X+\frac{1}{2}\right) = \vartheta_4(X), \quad \vartheta_4\left(X+\frac{1}{2}\right) = \vartheta_3(X). \\
& \vartheta_1\left(X+\frac{iy}{2}; iy\right) = i \exp\left\{-\pi i\left(X+\frac{iy}{4}\right)\right\} \vartheta_4(X; iy), \\
\text{(A.14)} \quad & \vartheta_2\left(X+\frac{iy}{2}; iy\right) = \exp\left\{-\pi i\left(X+\frac{iy}{4}\right)\right\} \vartheta_3(X; iy), \\
& \vartheta_3\left(X+\frac{iy}{2}; iy\right) = \exp\left\{-\pi i\left(X+\frac{iy}{4}\right)\right\} \vartheta_2(X; iy), \\
& \vartheta_4\left(X+\frac{iy}{2}; iy\right) = i \exp\left\{-\pi i\left(X+\frac{iy}{4}\right)\right\} \vartheta_1(X; iy).
\end{aligned}$$

Identity for the theta constants. For any $y > 0$, elliptic theta functions satisfy the following identity [20, 23].

$$\text{(A.15)} \quad \vartheta_1'(0; iy) = \pi \vartheta_2(0; iy) \vartheta_3(0; iy) \vartheta_4(0; iy).$$

Modular transformations. For any $X \in \mathbb{C}$ and $y > 0$, elliptic theta functions satisfy the following identities [20, 23].

$$(A.16) \quad \begin{aligned} \vartheta_1\left(\frac{X}{y}; i\frac{2}{y}\right) &= -i\sqrt{\frac{y}{2}} \exp\left(-\pi\frac{X^2}{2y}\right) \vartheta_1\left(\frac{iX}{2}; i\frac{y}{2}\right), \\ \vartheta_2\left(\frac{X}{y}; i\frac{2}{y}\right) &= \sqrt{\frac{y}{2}} \exp\left(-\pi\frac{X^2}{2y}\right) \vartheta_4\left(\frac{iX}{2}; i\frac{y}{2}\right), \\ \vartheta_3\left(\frac{X}{y}; i\frac{2}{y}\right) &= \sqrt{\frac{y}{2}} \exp\left(-\pi\frac{X^2}{2y}\right) \vartheta_3\left(\frac{iX}{2}; i\frac{y}{2}\right), \\ \vartheta_4\left(\frac{X}{y}; i\frac{2}{y}\right) &= \sqrt{\frac{y}{2}} \exp\left(-\pi\frac{X^2}{2y}\right) \vartheta_2\left(\frac{iX}{2}; i\frac{y}{2}\right). \end{aligned}$$

$$(A.17) \quad \frac{\vartheta_1(X; iy)}{\vartheta_2(X; iy)} = -i \frac{\vartheta_1\left(\frac{iX}{y}; \frac{i}{y}\right)}{\vartheta_4\left(\frac{iX}{y}; \frac{i}{y}\right)}.$$

Bilinear identities connecting theta functions with y and $y/2$. For any $X, Y \in \mathbb{C}$, and $y > 0$, elliptic theta functions satisfy the following identities [20, 23].

$$(A.18) \quad \begin{aligned} &2\vartheta_1(X+Y; iy) \vartheta_4(X-Y; iy) \\ &= \vartheta_1\left(X; \frac{iy}{2}\right) \vartheta_2\left(Y; \frac{iy}{2}\right) + \vartheta_2\left(X; \frac{iy}{2}\right) \vartheta_1\left(Y; \frac{iy}{2}\right), \end{aligned}$$

$$(A.19) \quad \begin{aligned} &2\vartheta_2(X+Y; iy) \vartheta_3(X-Y; iy) \\ &= \vartheta_2\left(X; \frac{iy}{2}\right) \vartheta_2\left(Y; \frac{iy}{2}\right) - \vartheta_1\left(X; \frac{iy}{2}\right) \vartheta_1\left(Y; \frac{iy}{2}\right). \end{aligned}$$

$$(A.20) \quad \begin{aligned} \vartheta_3\left(\frac{X}{2}; \frac{iy}{2}\right) \vartheta_4\left(\frac{X}{2}; \frac{iy}{2}\right) &= \vartheta_4(X; iy) \vartheta_4(0; iy), \\ \vartheta_1\left(\frac{X}{2}; \frac{iy}{2}\right) \vartheta_2\left(\frac{X}{2}; \frac{iy}{2}\right) &= \vartheta_1(X; iy) \vartheta_4(0; iy), \\ -\vartheta_1\left(\frac{X}{2}; \frac{iy}{2}\right)^2 + \vartheta_2\left(\frac{X}{2}; \frac{iy}{2}\right)^2 &= 2\vartheta_2(X; iy) \vartheta_3(0; iy). \end{aligned}$$

A.2. Proofs of Theorem 4.1 and Theorem 5.1. Here we prove that the τ functions given in Theorem 5.1 satisfy (3.1)-(3.6). Theorem 4.1 follows from Theorem 5.1 by setting $t = 0$. We assume $z = 0$ without loss of generality except for the proof of (3.5). Putting

$$(A.21) \quad \begin{cases} X = ivn + r + it, \\ Y = ivn - r + it, \\ U = \frac{1}{2}iv + r, \\ V = \frac{1}{2}iv - r, \end{cases}$$

in (A.2) yields (3.1). Putting

$$(A.22) \quad \begin{cases} X = iv\left(n - \frac{1}{2}\right) + it, \\ Y = ivn + r + it, \\ U = iv, \\ V = \frac{1}{2}iv - r, \end{cases}$$

in (A.2) yields (3.2). Putting

$$(A.23) \quad \begin{cases} X = iv \left(n - \frac{1}{2} \right) + it, \\ Y = \frac{1}{2}iv + r, \\ U = iv, \\ V = ivn - r + it, \end{cases}$$

in (A.3) yields (3.3). Putting

$$(A.24) \quad \begin{cases} X = ivn - r + it, \\ Y = -\frac{1}{2}iv + r, \\ U = ivn - r + it, \\ V = -\frac{1}{2}iv + r, \end{cases}$$

in (A.4) yields

$$(A.25) \quad \begin{aligned} & \frac{\vartheta_2(0) \vartheta_4(0)}{\vartheta_1(-\frac{1}{2}iv + r) \vartheta_3(-\frac{1}{2}iv + r)} \vartheta_2 \left(iv \left(n - \frac{1}{2} \right) + it \right) \vartheta_4 \left(iv \left(n + \frac{1}{2} \right) - 2r + it \right) \\ &= \frac{\vartheta_2(-\frac{1}{2}iv + r) \vartheta_4(-\frac{1}{2}iv + r)}{\vartheta_1(-\frac{1}{2}iv + r) \vartheta_3(-\frac{1}{2}iv + r)} \\ & \times \vartheta_2(ivn - r + it) \vartheta_4(ivn - r + it) - \vartheta_3(ivn - r + it) \vartheta_1(ivn - r + it). \end{aligned}$$

Also, putting

$$(A.26) \quad \begin{cases} X = ivn - r + it, \\ Y = \frac{1}{2}iv + r, \\ U = ivn - r + it, \\ V = \frac{1}{2}iv + r, \end{cases}$$

in (A.4) yields

$$(A.27) \quad \begin{aligned} & \frac{\vartheta_2(0) \vartheta_4(0)}{\vartheta_1(\frac{1}{2}iv + r) \vartheta_3(\frac{1}{2}iv + r)} \vartheta_2 \left(iv \left(n + \frac{1}{2} \right) + it \right) \vartheta_4 \left(iv \left(n - \frac{1}{2} \right) - 2r + it \right) \\ &= \frac{\vartheta_2(\frac{1}{2}iv + r) \vartheta_4(\frac{1}{2}iv + r)}{\vartheta_1(\frac{1}{2}iv + r) \vartheta_3(\frac{1}{2}iv + r)} \vartheta_2(ivn - r + it) \vartheta_4(ivn - r + it) \\ & - \vartheta_3(ivn - r + it) \vartheta_1(ivn - r + it). \end{aligned}$$

Subtracting (A.27) from (A.25) and substituting

$$(A.28) \quad \begin{aligned} & \vartheta_3(2r) \vartheta_4(0) \vartheta_2(0) \vartheta_1(iv) \\ &= \vartheta_1 \left(\frac{1}{2}iv + r \right) \vartheta_3 \left(\frac{1}{2}iv + r \right) \vartheta_2 \left(-\frac{1}{2}iv + r \right) \vartheta_4 \left(-\frac{1}{2}iv + r \right) \\ & - \vartheta_1 \left(-\frac{1}{2}iv + r \right) \vartheta_3 \left(-\frac{1}{2}iv + r \right) \vartheta_2 \left(\frac{1}{2}iv + r \right) \vartheta_4 \left(\frac{1}{2}iv + r \right), \end{aligned}$$

in the equation, we obtain

$$(A.29) \quad \begin{aligned} & \frac{\vartheta_1(-\frac{1}{2}iv + r) \vartheta_3(-\frac{1}{2}iv + r) \vartheta_1(\frac{1}{2}iv + r) \vartheta_3(\frac{1}{2}iv + r)}{\vartheta_3(2r) \vartheta_1(iv)} \\ & \times \left\{ \frac{\vartheta_1(\frac{1}{2}iv + r) \vartheta_3(\frac{1}{2}iv + r)}{\vartheta_1(-\frac{1}{2}iv + r) \vartheta_3(-\frac{1}{2}iv + r)} \vartheta_2 \left(iv \left(n - \frac{1}{2} \right) + it \right) \vartheta_4 \left(iv \left(n + \frac{1}{2} \right) - 2r + it \right) \right. \\ & \left. - \vartheta_2 \left(iv \left(n + \frac{1}{2} \right) + it \right) \vartheta_4 \left(iv \left(n - \frac{1}{2} \right) - 2r + it \right) \right\} \\ &= \vartheta_3 \left(\frac{1}{2}iv + r \right) \vartheta_2(ivn - r + it) \vartheta_1 \left(\frac{1}{2}iv + r \right) \vartheta_4(ivn - r + it), \end{aligned}$$

from which we see (3.4) holds. Putting

$$(A.30) \quad \begin{cases} X = ivn + z + r + it, \\ Y = ivn + z - r + it, \\ U = \frac{1}{2}iv + r + h, \\ V = \frac{1}{2}iv - r + h, \end{cases}$$

in (A.2) yields

$$(A.31) \quad \begin{aligned} & \vartheta_2 \left(iv \left(n + \frac{1}{2} \right) + z + it + h \right) \vartheta_2 \left(iv \left(n - \frac{1}{2} \right) + z + it - h \right) \vartheta_3(2r) \vartheta_3(0) \\ &= \vartheta_2 (ivn + z + r + it) \vartheta_2 (ivn + z - r + it) \vartheta_3 \left(\frac{1}{2}iv + r + h \right) \vartheta_3 \left(\frac{1}{2}iv - r + h \right) \\ & \quad - \vartheta_4 (ivn + z + r + it) \vartheta_4 (ivn + z - r + it) \vartheta_1 \left(\frac{1}{2}iv + r + h \right) \vartheta_1 \left(\frac{1}{2}iv - r + h \right). \end{aligned}$$

Differentiating (A.31) with h and then putting $h = 0$, we obtain

$$(A.32) \quad \begin{aligned} & \vartheta_3(2r) \vartheta_3(0) D_z \vartheta_2 \left(iv \left(n + \frac{1}{2} \right) + z + it \right) \cdot \vartheta_2 \left(iv \left(n - \frac{1}{2} \right) + z + it \right) \\ &= \left(\frac{\vartheta_3' \left(\frac{1}{2}iv + r \right)}{\vartheta_3 \left(\frac{1}{2}iv + r \right)} - \frac{\vartheta_3' \left(-\frac{1}{2}iv + r \right)}{\vartheta_3 \left(-\frac{1}{2}iv + r \right)} \right) \\ & \quad \times \vartheta_3 \left(-\frac{1}{2}iv + r \right) \vartheta_3 \left(\frac{1}{2}iv + r \right) \vartheta_2 (ivn + z + r + it) \vartheta_2 (ivn + z - r + it) \\ & \quad + \left(\frac{\vartheta_1' \left(\frac{1}{2}iv + r \right)}{\vartheta_1 \left(\frac{1}{2}iv + r \right)} - \frac{\vartheta_1' \left(-\frac{1}{2}iv + r \right)}{\vartheta_1 \left(-\frac{1}{2}iv + r \right)} \right) \\ & \quad \times \vartheta_1 \left(-\frac{1}{2}iv + r \right) \vartheta_1 \left(\frac{1}{2}iv + r \right) \vartheta_4 (ivn + z + r + it) \vartheta_4 (ivn + z - r + it). \end{aligned}$$

Using the equation in (A.31) with $h = 0$ and substituting (A.32), we obtain

$$(A.33) \quad \begin{aligned} & \vartheta_3(2r) \vartheta_3(0) \left\{ D_z \vartheta_2 \left(iv \left(n + \frac{1}{2} \right) + z + it \right) \cdot \vartheta_2 \left(iv \left(n - \frac{1}{2} \right) + z + it \right) \right. \\ & \quad \left. + \left(\frac{\vartheta_3' \left(-\frac{1}{2}iv + r \right)}{\vartheta_3 \left(-\frac{1}{2}iv + r \right)} - \frac{\vartheta_3' \left(\frac{1}{2}iv + r \right)}{\vartheta_3 \left(\frac{1}{2}iv + r \right)} \right) \right. \\ & \quad \left. \times \vartheta_2 \left(iv \left(n + \frac{1}{2} \right) + z + it \right) \vartheta_2 \left(iv \left(n - \frac{1}{2} \right) + z + it \right) \right\} \\ &= \left(\frac{\vartheta_1' \left(\frac{1}{2}iv + r \right)}{\vartheta_1 \left(\frac{1}{2}iv + r \right)} - \frac{\vartheta_1' \left(-\frac{1}{2}iv + r \right)}{\vartheta_1 \left(-\frac{1}{2}iv + r \right)} + \frac{\vartheta_3' \left(-\frac{1}{2}iv + r \right)}{\vartheta_3 \left(-\frac{1}{2}iv + r \right)} - \frac{\vartheta_3' \left(\frac{1}{2}iv + r \right)}{\vartheta_3 \left(\frac{1}{2}iv + r \right)} \right) \\ & \quad \times \vartheta_1 \left(-\frac{1}{2}iv + r \right) \vartheta_4 (ivn + z + r + it) \vartheta_1 \left(\frac{1}{2}iv + r \right) \vartheta_4 (ivn + z - r + it), \end{aligned}$$

from which we see (3.5) holds. By the definition of G and \tilde{G} , (3.6) follows. \square

A.3. Proof of Proposition 5.5. For the proof of Proposition 5.5, we use the following lemma:

Lemma A.1 ([10, §3]). *The following equation holds.*

$$(A.34) \quad f_n f_{n-1}^* + g_n g_{n-1}^* = f_n^* f_{n-1} + g_n^* g_{n-1} = \beta F_n F_n,$$

Proof of Proposition 5.5. Here we prove that the τ functions given in Theorem 5.1 satisfy (5.8)-(5.14). We assume $z = 0$ without loss of generality except for the proof of (5.14). Putting

$$(A.35) \quad \begin{cases} X = -\frac{1}{2}iv + r, \\ Y = \frac{1}{2}iv - r, \\ U = ivn + r + it, \\ V = iv(n-1) - r + it, \end{cases}$$

in (A.8) yields (5.8). Also, (5.9) follows from (A.34) and (5.8). Putting

$$(A.36) \quad \begin{cases} X = ivn - r + it, \\ Y = -\frac{1}{2}iv - r + h, \\ U = \frac{1}{2}iv - r + h, \\ V = iv(n-1) - r + it, \end{cases}$$

in (A.3) yields

$$(A.37) \quad \begin{aligned} & \vartheta_4 \left(iv \left(n - \frac{1}{2} \right) - 2r + it + h \right) \vartheta_3(0) \vartheta_1(iv) \vartheta_2 \left(iv \left(n - \frac{1}{2} \right) + it - h \right) \\ &= \vartheta_4(ivn - r + it) \vartheta_3 \left(-\frac{1}{2}iv - r + h \right) \\ & \times \vartheta_1 \left(\frac{1}{2}iv - r + h \right) \vartheta_2(iv(n-1) - r + it) \\ & - \vartheta_2(ivn - r + it) \vartheta_1 \left(-\frac{1}{2}iv - r + h \right) \\ & \times \vartheta_3 \left(\frac{1}{2}iv - r + h \right) \vartheta_4(iv(n-1) - r + it), \end{aligned}$$

After differentiating (A.37) by h and putting $h = 0$, we obtain (5.13) by rearrangement. Also, rearranging the equation with $h = 0$ in (A.37) yields (5.10). Putting

$$(A.38) \quad \begin{cases} X = \frac{1}{2}iv + r, \\ Y = -\frac{1}{2}iv - r, \\ U = iv \left(n - \frac{1}{2} \right) + it, \\ V = -iv \left(n - \frac{1}{2} \right) - it, \end{cases}$$

in (A.5) yields (5.11). Also, (5.12) follows from (5.8) and (5.11). Putting

$$(A.39) \quad \begin{cases} X = iv \left(n - \frac{1}{2} \right) + it + \frac{z}{\Delta_3 - \Delta_1}, \\ Y = iv \left(n - \frac{1}{2} \right) + it + \frac{z}{\Delta_3 - \Delta_1}, \\ U = h, \\ V = h, \end{cases}$$

in (A.2) yields

$$(A.40) \quad \begin{aligned} & \vartheta_2 \left(iv \left(n - \frac{1}{2} \right) + it + \frac{z}{\Delta_3 - \Delta_1} + h \right) \vartheta_2 \left(iv \left(n - \frac{1}{2} \right) + it + \frac{z}{\Delta_3 - \Delta_1} - h \right) \\ &= -\vartheta_4 \left(iv \left(n - \frac{1}{2} \right) + it + \frac{z}{\Delta_3 - \Delta_1} \right)^2 \frac{\vartheta_1(h)^2}{\vartheta_3(0)^2} \\ & + \vartheta_2 \left(iv \left(n - \frac{1}{2} \right) + it + \frac{z}{\Delta_3 - \Delta_1} \right)^2 \frac{\vartheta_3(h)^2}{\vartheta_3(0)^2}. \end{aligned}$$

After differentiating (A.40) twice with h and putting $h = 0$, we obtain (5.14). \square

A.4. Proofs of Theorem 5.6 and Theorem 5.9. For the proofs of Theorem 5.6 and Theorem 5.9, we need the following lemmas.

Lemma A.2 ([10, §3]). *Using (3.2) and (3.3), we see that the following equation holds.*

$$(A.41) \quad g_n f_{n-1} - f_n g_{n-1} = G_n F_n.$$

Lemma A.3. *The following relation holds.*

$$(A.42) \quad \omega = \delta_2 - \delta_2^*.$$

Proof. Putting

$$(A.43) \quad \begin{cases} X = -\frac{1}{2}iv + r, \\ Y = -\frac{1}{2}iv + r, \\ U = \frac{1}{2}iv + r, \\ V = \frac{1}{2}iv + r, \end{cases}$$

in (A.9) yields (A.42). □

Lemma A.4. *The following identity holds.*

$$(A.44) \quad 2\beta_1 (\delta_2 + \delta_2^*) + 2i\omega\beta_2 + \epsilon_2 = 2\delta_2\delta_3^* + 2\delta_2^*\delta_3 + \epsilon_2 = 0.$$

Proof. Putting

$$(A.45) \quad \begin{cases} X = \frac{1}{2}iv + r + h, \\ Y = -\frac{1}{2}iv + r - h, \\ U = \frac{1}{2}iv + r - h, \\ V = -\frac{1}{2}iv + r + h, \end{cases}$$

in (A.9) yields

$$(A.46) \quad \begin{aligned} & \vartheta_3 \left(\frac{1}{2}iv + r + h \right) \vartheta_3 \left(-\frac{1}{2}iv + r - h \right) \vartheta_1 \left(\frac{1}{2}iv + r - h \right) \vartheta_1 \left(-\frac{1}{2}iv + r + h \right) \\ & - \vartheta_1 \left(\frac{1}{2}iv + r + h \right) \vartheta_1 \left(-\frac{1}{2}iv + r - h \right) \vartheta_3 \left(\frac{1}{2}iv + r - h \right) \vartheta_3 \left(-\frac{1}{2}iv + r + h \right) \\ & = \vartheta_3(2r) \vartheta_3(0) \vartheta_1(iv) \vartheta_1(2h), \end{aligned}$$

After differentiating (A.46) by h and setting $h = 0$, we obtain (A.44). □

For simplicity, we define p and q as follows

$$(A.47) \quad p_n = g_n^* f_{n-1} - f_n g_{n-1}^*,$$

$$(A.48) \quad q_n = g_n^* f_{n-1} + f_n g_{n-1}^*.$$

Then we rewrite $D_t H_n \cdot F_n$ by using (5.6), (A.47) and (A.48) as follows

$$(A.49) \quad D_t H_n \cdot F_n = \beta_1 q_n^* + i\beta_2 p_n^*.$$

Also, we see the following hold.

$$(A.50) \quad \tan \frac{\kappa_n}{2} \tilde{N}_n = T_n - \frac{1}{2\beta F_n^2} \begin{pmatrix} q_n^* + q_n \\ \frac{1}{i} (q_n^* - q_n) \\ -2(\beta - 2\delta_1) F_n^2 + 2(\delta_2 + \delta_2^*) |G_n|^2 \end{pmatrix},$$

$$(A.51) \quad \tan \frac{\kappa_n}{2} \tilde{B}_n = \frac{1}{2\beta F_n^2} \begin{pmatrix} \frac{1}{i} (p_n - p_n^*) \\ p_n + p_n^* \\ -\frac{2}{i} \omega |G_n|^2 \end{pmatrix}.$$

We have the following Lemma.

Lemma A.5. *The following relations hold.*

$$(A.52) \quad p_n p_n^* = F_n^2 |G_n|^2 + \omega^2 |G_n|^4,$$

$$(A.53) \quad p_n q_n^* - p_n^* q_n = 2(2\delta_1 - \beta) \omega F_n^2 |G_n|^2 + 2\omega (\delta_2 + \delta_2^*) |G_n|^4,$$

$$(A.54) \quad q_n q_n^* = 4\delta_1 (\beta - \delta_1) F_n^4 + \{1 + 2(\beta - 2\delta_1) (\delta_2 + \delta_2^*)\} F_n^2 |G_n|^2 - (\delta_2 + \delta_2^*)^2 |G_n|^4.$$

Proof. By definition of p , we see that

$$(A.55) \quad \begin{aligned} p_n p_n^* &= (g_n^* f_{n-1} - f_n g_{n-1}^*) (g_n f_{n-1}^* - f_n^* g_{n-1}) \\ &= (f_n^* g_{n-1}^* - g_n^* f_{n-1}^*) (f_n g_{n-1} - g_n f_{n-1}) \\ &\quad + (f_n f_{n-1}^* - f_n^* f_{n-1}) (g_n^* g_{n-1} - g_n g_{n-1}^*), \end{aligned}$$

holds. Substituting (A.41) and (5.9) into (A.55) yields (A.52). By definition of p and q , we see that

$$(A.56) \quad \begin{aligned} p_n q_n^* - p_n^* q_n &= (g_n^* f_{n-1} - f_n g_{n-1}^*) (g_n f_{n-1}^* + f_n^* g_{n-1}) \\ &\quad - (g_n f_{n-1}^* - f_n^* g_{n-1}) (g_n^* f_{n-1} + f_n g_{n-1}^*) \\ &= 2(g_n^* f_{n-1} f_n^* g_{n-1} - g_n f_{n-1}^* f_n g_{n-1}^*), \end{aligned}$$

holds. On the other hand, from (A.34) and (5.9), the following holds.

$$(A.57) \quad \begin{aligned} \beta \omega F_n^2 |G_n|^2 &= (f_n f_{n-1}^* + g_n g_{n-1}^*) (g_n^* g_{n-1} - g_n g_{n-1}^*) \\ &= (f_n^* f_{n-1} + g_n^* g_{n-1}) (g_n g_{n-1} - g_n g_{n-1}^*) \\ &= (f_n f_{n-1}^* + g_n g_{n-1}^*) (f_n f_{n-1} - f_n^* f_{n-1}) \\ &= (f_n^* f_{n-1} + g_n^* g_{n-1}) (f_n f_{n-1} - f_n^* f_{n-1}). \end{aligned}$$

Using (A.57), we obtain

$$(A.58) \quad \begin{aligned} p_n q_n^* - p_n^* q_n &= f_n f_{n-1}^* f_n f_{n-1}^* - f_n^* f_{n-1} f_n^* f_{n-1} + g_n g_{n-1}^* g_n g_{n-1}^* - g_n^* g_{n-1} g_n^* g_{n-1} \\ &= (f_n f_{n-1}^*)^2 - (f_n^* f_{n-1})^2 + (g_n g_{n-1}^*)^2 - (g_n^* g_{n-1})^2 \\ &= (f_n f_{n-1}^* - f_n^* f_{n-1}) (f_n f_{n-1}^* + f_n^* f_{n-1}) \\ &\quad + (g_n g_{n-1}^* - g_n^* g_{n-1}) (g_n g_{n-1}^* + g_n^* g_{n-1}). \end{aligned}$$

Substituting (5.9), (5.11) and (5.12) into (A.58) gives

$$(A.59) \quad \begin{aligned} p_n q_n^* - p_n^* q_n &= \omega |G_n|^2 (f_n f_{n-1}^* + f_n^* f_{n-1} - g_n g_{n-1}^* - g_n^* g_{n-1}) \\ &= 2(2\delta_1 - \beta) \omega F_n^2 |G_n|^2 + 2\omega (\delta_2 + \delta_2^*) |G_n|^4, \end{aligned}$$

thus (A.53). Also from (A.41), (5.11) and (5.12), we see that

$$(A.60) \quad \begin{aligned} q_n q_n^* - F_n^2 |G_n|^2 &= (g_n^* f_{n-1} + f_n g_{n-1}^*) (g_n f_{n-1}^* + f_n^* g_{n-1}) \\ &\quad - (g_n f_{n-1}^* - f_n g_{n-1}^*) (g_n^* f_{n-1} - f_n^* g_{n-1}^*) \\ &= (f_n f_{n-1}^* + f_n^* f_{n-1}) (g_n g_{n-1}^* + g_n^* g_{n-1}) \\ &= (2\delta_1 F_n^2 + (\delta_2 + \delta_2^*) |G_n|^2) (2(\beta - \delta_1) F_n^2 - (\delta_2 + \delta_2^*) |G_n|^2) \\ &= 4\delta_1 (\beta - \delta_1) F_n^4 + 2(\beta - 2\delta_1) (\delta_2 + \delta_2^*) F_n^2 |G_n|^2 - (\delta_2 + \delta_2^*)^2 |G_n|^4. \end{aligned}$$

holds. From (A.60), we get (A.54). \square

Using Proposition 5.5, Lemma A.3, Lemma A.4 and Lemma A.5, Theorem 5.6 and Theorem 5.9 can be proved as follows.

Proof of Theorem 5.6. By differentiating γ with respect to t , we get

$$(A.61) \quad \begin{aligned} \dot{\gamma}_n(t) &= \frac{1}{F_n^2} \begin{pmatrix} D_t H_n \cdot F_n + D_t H_n^* \cdot F_n \\ \frac{1}{i} (D_t H_n \cdot F_n - D_t H_n^* \cdot F_n) \\ -D_t D_z F_n \cdot F_n \end{pmatrix} \\ &= \frac{1}{F_n^2} \begin{pmatrix} \beta_1 q_n^* + i\beta_2 p_n^* + \beta_1 q_n - i\beta_2 p_n \\ \frac{1}{i} (\beta_1 q_n^* + i\beta_2 p_n^* - \beta_1 q_n + i\beta_2 p_n) \\ -(\epsilon_1 + C) F_n^2 - \epsilon_2 |G_n|^2 \end{pmatrix}. \end{aligned}$$

On the other hand since

$$(A.62) \quad \begin{aligned} &\beta_1 \left(T_n - \tan \frac{\kappa_n}{2} \tilde{N}_n \right) + \beta_2 \tan \frac{\kappa_n}{2} \tilde{B}_n \\ &= \frac{1}{2\beta F_n^2} \begin{pmatrix} \beta_1 q_n^* + i\beta_2 p_n^* + \beta_1 q_n - i\beta_2 p_n \\ \frac{1}{i} (\beta_1 q_n^* + i\beta_2 p_n^* - \beta_1 q_n + i\beta_2 p_n) \\ -2\beta_1 (\beta - 2\delta_1) F_n^2 + 2\beta_1 (\delta_2 + \delta_2^*) |G_n|^2 + 2i\beta_2 \omega |G_n|^2 \end{pmatrix}. \end{aligned}$$

holds. From (A.44), this equation can be rewritten as follows

$$(A.63) \quad \begin{aligned} &\beta_1 \left(T_n - \tan \frac{\kappa_n}{2} \tilde{N}_n \right) + \beta_2 \tan \frac{\kappa_n}{2} \tilde{B}_n \\ &= \frac{1}{2\beta F_n^2} \begin{pmatrix} \beta_1 q_n^* + i\beta_2 p_n^* + \beta_1 q_n - i\beta_2 p_n \\ \frac{1}{i} (\beta_1 q_n^* + i\beta_2 p_n^* - \beta_1 q_n + i\beta_2 p_n) \\ -2\beta_1 (\beta - 2\delta_1) F_n^2 - \epsilon_2 |G_n|^2 \end{pmatrix}. \end{aligned}$$

Thus (5.15) is obtained. \square

Proof of Theorem 5.9. We compute the condition satisfied by C, Γ when

$$(A.64) \quad \langle \dot{\gamma}_n, \tilde{B}_n \rangle = 0,$$

$$(A.65) \quad \langle \dot{\gamma}_n, \dot{\gamma}_n \rangle = \rho^2,$$

where $\rho \in \mathbb{R}$ is a constant. From (A.51) and (A.61) we see that

$$(A.66) \quad \begin{aligned} 2iF_n^3 \tilde{G}_n \langle \dot{\gamma}_n, \tilde{B}_n \rangle &= (p_n - p_n^*) (\beta_1 q_n^* + i\beta_2 p_n^* + \beta_1 q_n - i\beta_2 p_n) \\ &\quad + (p_n + p_n^*) (\beta_1 q_n^* + i\beta_2 p_n^* - \beta_1 q_n + i\beta_2 p_n) \\ &\quad + 2\omega |G_n|^2 \{(\epsilon_1 + C) F_n^2 + \epsilon_2 |G_n|^2\} \\ &= 2F_n^2 |G_n|^2 \{2\beta_1 (2\delta_1 - \beta) \omega + 2i\beta_2 + \omega (\epsilon_1 + C)\} \\ &\quad + 2\omega |G_n|^4 \{2\beta_1 (\delta_2 + \delta_2^*) + 2i\omega \beta_2 + \epsilon_2\}, \end{aligned}$$

holds. For (A.64) to be satisfied, it is sufficient that the following equations hold.

$$(A.67) \quad 2\beta_1 (2\delta_1 - \beta) \omega + 2i\beta_2 + \omega (\epsilon_1 + C) = 0,$$

$$(A.68) \quad 2\beta_1 (\delta_2 + \delta_2^*) + 2i\omega \beta_2 + \epsilon_2 = 0.$$

By comparing (A.68) and (A.44) we see that (A.67) is a necessary and sufficient condition for the realization of (A.64). By substituting (5.6) into (A.67), it can be rewritten in terms of C, Γ as follows.

$$(A.69) \quad i\Gamma \left(\omega (2\delta_1 - \beta) - \frac{\delta_2 + \delta_2^*}{\omega} \right) + \epsilon_1 + C + (\delta_3 + \delta_3^*) (2\delta_1 - \beta) - \frac{\delta_3 - \delta_3^*}{\omega} = 0.$$

Also, since

$$(A.70) \quad \begin{aligned} & F_n^4 \langle \dot{\gamma}_n, \dot{\gamma}_n \rangle \\ &= (\beta_1 q_n^* + i\beta_2 p_n^* + \beta_1 q_n - i\beta_2 p_n)^2 \\ &\quad - (\beta_1 q_n^* + i\beta_2 p_n^* - \beta_1 q_n + i\beta_2 p_n)^2 \\ &\quad + \{(\epsilon_1 + C) F_n^2 + \epsilon_2 |G_n|^2\}^2 \\ &= F_n^4 \{16\beta_1^2 \delta_1 (\beta - \delta_1) + (\epsilon_1 + C)^2\} \\ &\quad + F_n^2 |G_n|^2 \{4\beta_1^2 (1 + 2(\beta - 2\delta_1)(\delta_2 + \delta_2^*)) \\ &\quad - 8i\beta_1 \beta_2 \omega (2\delta_1 - \beta) + 4\beta_2^2 + 2\epsilon_2 (\epsilon_1 + C)\} \\ &\quad + |G_n|^4 \left(-4\beta_1^2 (\delta_2 + \delta_2^*)^2 - 8i\beta_1 \beta_2 \omega (\delta_2 + \delta_2^*) + 4\beta_2^2 \omega^2 + \epsilon_2^2 \right), \end{aligned}$$

holds, we see that for (A.65) holds, it is sufficient that

$$(A.71) \quad 4\beta_1^2 (1 + 2(\beta - 2\delta_1)(\delta_2 + \delta_2^*)) - 8i\beta_1 \beta_2 \omega (2\delta_1 - \beta) + 4\beta_2^2 + 2\epsilon_2 (\epsilon_1 + C) = 0,$$

$$(A.72) \quad -4\beta_1^2 (\delta_2 + \delta_2^*)^2 - 8i\beta_1 \beta_2 \omega (\delta_2 + \delta_2^*) + 4\beta_2^2 \omega^2 + \epsilon_2^2 = 0,$$

hold. Since

$$(A.73) \quad \begin{aligned} & 4\beta_1^2 (\delta_2 + \delta_2^*)^2 + 8i\beta_1 \beta_2 \omega (\delta_2 + \delta_2^*) - 4\beta_2^2 \omega^2 - \epsilon_2^2 \\ &= \{2\beta_1 (\delta_2 + \delta_2^*) + 2i\omega \beta_2 + \epsilon_2\} \{2\beta_1 (\delta_2 + \delta_2^*) + 2i\omega \beta_2 - \epsilon_2\}, \end{aligned}$$

holds, (A.72) is identically satisfied. Therefore, (A.71) is a necessary and sufficient condition for the realization of (A.65). By using (A.44), (5.6) and (A.69), (A.71) can be written in terms of Γ as follows.

$$(A.74) \quad i\delta_2 \delta_2^* (\delta_2 - \delta_2^*) \Gamma^2 + 2\delta_2 \delta_2^* (\delta_3 + \delta_3^*) \Gamma + i (\delta_2 \delta_3^{*2} - \delta_2^* \delta_3^2) = 0.$$

(A.74) has the following solutions.

$$(A.75) \quad \Gamma = \frac{-\delta_2 \delta_2^* (\delta_3 + \delta_3^*) \pm \sqrt{\delta_2 \delta_2^* (\delta_2 \delta_3^* + \delta_2^* \delta_3)^2}}{i\delta_2 \delta_2^* (\delta_2 - \delta_2^*)}.$$

Since

$$(A.76) \quad \delta_2 \delta_2^* (\delta_2 \delta_3^* + \delta_2^* \delta_3)^2 = (\delta_2 \delta_2^*)^2 (\delta_3 e^{i\lambda} + \delta_3^* e^{-i\lambda})^2 = (\delta_2 \delta_2^*)^2 \left(i \frac{\vartheta_1'(0)}{\vartheta_1(iv)} \right)^2,$$

holds, (A.75) can be rewritten as

$$(A.77) \quad \Gamma = \frac{-(\delta_3 + \delta_3^*) \pm (\delta_3 e^{i\lambda} + \delta_3^* e^{-i\lambda})}{i\omega}.$$

Now, we prove equations (5.20) and (5.21). From (4.2) and (5.6), we see that when C_+, Γ_+ are chosen as in (5.18), the following relation holds

$$(A.78) \quad 16\beta_1^2 (\beta - \delta_1) \delta_1 = 4 \left(i \frac{\vartheta_1'(0)}{\vartheta_1(iv)} \right)^2 \frac{u_2^2}{\vartheta_3(2r)^2 \vartheta_3(0)^6}.$$

Also, from (4.2), (5.6), (A.67) and (A.68), we see that when C_+, Γ_+ are chosen as in (5.18), the following relation holds

$$(A.79) \quad (\epsilon_1 + C)^2 = \left(i \frac{\vartheta_1'(0)}{\vartheta_1(iv)} \right)^2 \frac{u_2^2}{\vartheta_3(2r)^2 \vartheta_3(0)^6} \frac{(\vartheta_3(r)^4 - \vartheta_1(r)^4)^2}{\vartheta_3(r)^4 \vartheta_1(r)^4}.$$

From (A.70), (A.78) and (A.79), we see that

$$(A.80) \quad \rho^2 = 16\beta_1^2 (\beta - \delta_1) \delta_1 + (\epsilon_1 + C)^2 = \left(i \frac{\vartheta_1'(0)}{\vartheta_1(iv)} \right)^2 \frac{u_2^2}{\vartheta_3(r)^4 \vartheta_1(r)^4} = \rho_+^2,$$

holds. (A.80) is equivalent to (5.20). From (4.2) and (5.6), we see that when C_- , Γ_- are chosen as in (5.18), the following relation holds

$$(A.81) \quad 16\beta_1^2 (\beta - \delta_1) \delta_1 = 4 \left(i \frac{\vartheta_1'(0)}{\vartheta_1(iv)} \right)^2 \frac{u_2^2}{\vartheta_3(2r)^2 \vartheta_3(0)^6}.$$

Also, from (4.2), (5.6), (A.67) and (A.68), we see that when C_- , Γ_- are chosen as in (5.18), the following relation holds

$$(A.82) \quad (\epsilon_1 + C)^2 = \left(i \frac{\vartheta_1'(0)}{\vartheta_1(iv)} \right)^2 \frac{u_2^2}{\vartheta_3(2r)^2 \vartheta_3(0)^6} \frac{(\vartheta_4(r)^4 - \vartheta_2(r)^4)^2}{\vartheta_4(r)^4 \vartheta_2(r)^4}.$$

From (A.70), (A.81) and (A.82), we see that

$$(A.83) \quad \rho^2 = 16\beta_1^2 (\beta - \delta_1) \delta_1 + (\epsilon_1 + C)^2 = \left(i \frac{\vartheta_1'(0)}{\vartheta_1(iv)} \right)^2 \frac{u_2^2}{\vartheta_4(r)^4 \vartheta_2(r)^4} = \rho_-^2,$$

holds. (A.83) is equivalent to (5.21). □

A.5. Proof of Proposition 5.12. Here we prove that the signed curvature angle κ satisfies (5.27). First, we state Lemma A.6.

Lemma A.6. *The τ functions (5.1) satisfy the following bilinear equations.*

$$(A.84) \quad \tilde{G}_n^2 + \beta^2 F_n^2 = F_{n+1} F_{n-1},$$

$$(A.85) \quad \beta^2 \tilde{G}_n^2 - \frac{\vartheta_1(iv)^4}{\vartheta_3(iv)^4} \beta^4 F_n^2 = \tilde{G}_{n+1} \tilde{G}_{n-1}.$$

Proof. Using (3.1)-(3.3), we see that (A.84) hold. Putting

$$(A.86) \quad \begin{cases} X = 0, \\ Y = 0, \\ U = iv \left(n + \frac{1}{2} \right) + it + \frac{z}{\Delta_3 - \Delta_1}, \\ V = -iv \left(n - \frac{3}{2} \right) - it - \frac{z}{\Delta_3 - \Delta_1}, \end{cases}$$

in (A.10), we obtain (A.85). □

Proof of Proposition 5.12. Using (A.84) and (A.85), we see that (5.27) holds. □

A.6. Proof of Theorem 5.13. Here, we show that the signed curvature angle satisfies the semi-discrete mKdV equation (5.28). For the proof of Theorem 5.13, we first state Lemma A.7.

Lemma A.7. *The τ functions (5.1) satisfy the following bilinear equation.*

$$(A.87) \quad D_t \tilde{G}_n \cdot F_n = \eta_3 \left(\tilde{G}_{n+1} F_{n-1} - \tilde{G}_{n-1} F_{n+1} \right).$$

Proof. We assume $z = 0$ without loss of generality. After differentiating the equation in (A.3) with

$$(A.88) \quad \begin{cases} X = iv \left(n + \frac{1}{2} \right) + it, \\ Y = iv + h, \\ U = -iv + h, \\ V = iv \left(n - \frac{3}{2} \right) + it, \end{cases}$$

by h and putting $h = 0$, we obtain

$$(A.89) \quad \begin{aligned} & D_t \left\{ -i\vartheta_1(iv)\vartheta_4 \left(iv \left(n - \frac{1}{2} \right) + it \right) \right\} \cdot \left\{ \vartheta_3(iv)\vartheta_2 \left(iv \left(n - \frac{1}{2} \right) + it \right) \right\} \\ &= \frac{\vartheta_1(iv)\vartheta_3(iv)}{\vartheta_1(2iv)\vartheta_3(0)} \frac{\vartheta_1'(iv)\vartheta_3(iv) - \vartheta_1(iv)\vartheta_3'(iv)}{i\vartheta_1(iv)\vartheta_3(iv)} \\ &\times \left\{ -i\vartheta_1(iv)\vartheta_4 \left(iv \left(n - \frac{3}{2} \right) + it \right) \right\} \left\{ \vartheta_3(iv)\vartheta_2 \left(iv \left(n + \frac{1}{2} \right) + it \right) \right\} \\ &- \frac{\vartheta_1(iv)\vartheta_3(iv)}{\vartheta_1(2iv)\vartheta_3(0)} \frac{\vartheta_1'(iv)\vartheta_3(iv) - \vartheta_1(iv)\vartheta_3'(iv)}{i\vartheta_1(iv)\vartheta_3(iv)} \\ &\times \left\{ -i\vartheta_1(iv)\vartheta_4 \left(iv \left(n + \frac{1}{2} \right) + it \right) \right\} \left\{ \vartheta_3(iv)\vartheta_2 \left(iv \left(n - \frac{3}{2} \right) + it \right) \right\}. \end{aligned}$$

After rearranging (A.89), we obtain (A.87). \square

Proof of Theorem 5.13. Using (A.84) and (A.87), we see that the signed curvature angle satisfies the semi-discrete mKdV equation (5.28). \square

A.7. Proof of Theorem 5.14. From (A.17), (A.18) and (A.19), we see that

$$(A.90) \quad \begin{aligned} \kappa_n &= 2 \arctan \left(-i \frac{\vartheta_1(iv; iy)\vartheta_4 \left(iv \left(n - \frac{1}{2} \right) + it; iy \right)}{\vartheta_3(iv; iy)\vartheta_2 \left(iv \left(n - \frac{1}{2} \right) + it; iy \right)} \right) \\ &= 2 \arctan \left(-i \frac{\frac{\vartheta_1 \left(i \frac{v}{2} \left(n + \frac{1}{2} \right) + \frac{it}{2}; \frac{iy}{2} \right)}{\vartheta_2 \left(i \frac{v}{2} \left(n + \frac{1}{2} \right) + \frac{it}{2}; \frac{iy}{2} \right)} - \frac{\vartheta_1 \left(i \frac{v}{2} \left(n - \frac{3}{2} \right) + \frac{it}{2}; \frac{iy}{2} \right)}{\vartheta_2 \left(i \frac{v}{2} \left(n - \frac{3}{2} \right) + \frac{it}{2}; \frac{iy}{2} \right)}}{1 - \frac{\vartheta_1 \left(i \frac{v}{2} \left(n + \frac{1}{2} \right) + \frac{it}{2}; \frac{iy}{2} \right)\vartheta_1 \left(i \frac{v}{2} \left(n - \frac{3}{2} \right) + \frac{it}{2}; \frac{iy}{2} \right)}{\vartheta_2 \left(i \frac{v}{2} \left(n + \frac{1}{2} \right) + \frac{it}{2}; \frac{iy}{2} \right)\vartheta_2 \left(i \frac{v}{2} \left(n - \frac{3}{2} \right) + \frac{it}{2}; \frac{iy}{2} \right)}} \right) \\ &= 2 \arctan \left(\frac{\frac{\vartheta_1 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right)}{\vartheta_4 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right)} - \frac{\vartheta_1 \left(\frac{v}{y} \left(n - \frac{3}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right)}{\vartheta_4 \left(\frac{v}{y} \left(n - \frac{3}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right)}}{1 + \frac{\vartheta_1 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right)\vartheta_1 \left(\frac{v}{y} \left(n - \frac{3}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right)}{\vartheta_4 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right)\vartheta_4 \left(\frac{v}{y} \left(n - \frac{3}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right)}} \right) \\ &= 2 \arctan \left(\frac{\vartheta_1 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right)}{\vartheta_4 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right)} \right) \\ &- 2 \arctan \left(\frac{\vartheta_1 \left(\frac{v}{y} \left(n - \frac{3}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right)}{\vartheta_4 \left(\frac{v}{y} \left(n - \frac{3}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right)} \right) \\ &= \frac{\Theta_{n+1} - \Theta_{n-1}}{2}, \end{aligned}$$

holds. \square

A.8. Proof of Theorem 5.15. We first construct the solution to the semi-discrete potential mKdV equation (5.33) in two stages by proving Lemma A.8 and Lemma A.9. Then we construct the solution to the semi-discrete sine-Gordon equation (5.34) in two stages: Lemma A.10 and Lemma A.11. This is based on the construction of the solution to the mKdV equation using the theory of τ functions [14].

For fixed v, y , define the following real-valued functions

$$(A.91) \quad \begin{aligned} A_n(t) &= \vartheta_4 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right), \\ B_n(t) &= \vartheta_1 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y}; \frac{2i}{y} \right). \end{aligned}$$

Then the potential function can be written as follows

$$(A.92) \quad \Theta_n(t) = 4 \arctan \frac{B_n}{A_n} = \frac{2}{i} \log \frac{A_n + iB_n}{A_n - iB_n} = \frac{2}{i} \log \frac{\tilde{\tau}_n}{\tilde{\tau}_n^*},$$

where $\tilde{\tau}_n = A_n + iB_n$. $\tilde{\tau}$ corresponds to the τ functions of the one-component KP hierarchy, while A and B correspond to the τ functions of the two-component KP hierarchy.

Lemma A.8. *Assuming that the solution of (5.33) is expressed as (A.92), we obtain the following bilinear equations as a sufficient condition for satisfying (5.33).*

$$(A.93) \quad D_t A_{n+1} \cdot A_n + D_t B_{n+1} \cdot B_n = \frac{L_1}{2y} (\eta_1 - 1) (A_{n+1} A_n + B_{n+1} B_n),$$

$$(A.94) \quad D_t A_{n+1} \cdot B_n - D_t B_{n+1} \cdot A_n = \frac{L_1}{2y} (\eta_1 + 1) (A_{n+1} B_n - B_{n+1} A_n),$$

where η_1 is a real constant.

Proof. Since

$$(A.95) \quad \begin{aligned} \frac{d}{dt} \left(\frac{\Theta_{n+1} + \Theta_n}{2} \right) &= \frac{1}{i} \frac{d}{dt} \left(\log \frac{\tilde{\tau}_{n+1}}{\tilde{\tau}_{n+1}^*} + \log \frac{\tilde{\tau}_n}{\tilde{\tau}_n^*} \right) \\ &= \frac{1}{i} \frac{d}{dt} \left(\log \frac{\tilde{\tau}_{n+1}}{\tilde{\tau}_n^*} - \log \frac{\tilde{\tau}_{n+1}^*}{\tilde{\tau}_n} \right) \\ &= \frac{1}{i} \left(\frac{D_t \tilde{\tau}_{n+1} \cdot \tilde{\tau}_n^*}{\tilde{\tau}_{n+1} \tilde{\tau}_n^*} - \frac{D_t \tilde{\tau}_{n+1}^* \cdot \tilde{\tau}_n}{\tilde{\tau}_{n+1}^* \tilde{\tau}_n} \right). \end{aligned}$$

$$(A.96) \quad \sin \left(\frac{\Theta_{n+1} - \Theta_n}{2} \right) = \frac{1}{2i} \left(\frac{\tilde{\tau}_{n+1} \tilde{\tau}_n^*}{\tilde{\tau}_{n+1}^* \tilde{\tau}_n} - \frac{\tilde{\tau}_{n+1}^* \tilde{\tau}_n}{\tilde{\tau}_{n+1} \tilde{\tau}_n^*} \right),$$

hold, it can be seen that a sufficient condition for (A.92) to be a solution of (5.33) is that

$$(A.97) \quad D_t \tilde{\tau}_{n+1} \cdot \tilde{\tau}_n^* = -\frac{L_1}{2y} \tilde{\tau}_{n+1}^* \tilde{\tau}_n + \frac{L_1}{2y} \eta_1 \tilde{\tau}_{n+1} \tilde{\tau}_n^*,$$

(η_1 is a real constant) is satisfied. Substituting $\tilde{\tau}_n = A_n + iB_n$ for this, we get

$$(A.98) \quad \begin{aligned} &D_t (A_{n+1} \cdot A_n + B_{n+1} \cdot B_n) - i D_t (A_{n+1} \cdot B_n - B_{n+1} \cdot A_n) \\ &= \frac{L_1}{2y} (\eta_1 - 1) (A_{n+1} A_n + B_{n+1} B_n) - i \frac{L_1}{2y} (\eta_1 + 1) (A_{n+1} B_n - B_{n+1} A_n). \end{aligned}$$

The real part of this equation yields (A.93), and the imaginary part yields (A.94). \square

In Lemma A.9, we construct solutions of (A.93) and (A.94) in terms of elliptic theta functions.

Lemma A.9. *The real functions A and B defined by (A.91) are the solutions of (A.93) and (A.94) with*

$$(A.99) \quad \eta_1 = \frac{d_1 \left(\frac{v}{y}; \frac{2}{y} \right) + d_4 \left(\frac{v}{y}; \frac{2}{y} \right) - Q_1(v, y) + Q_4(v, y)}{L_1(v, y)}.$$

Proof. In this proof, all elliptic theta functions have $w = 2i/y$ and we denote $\vartheta_j(v; 2i/y)$ by $\vartheta_j(v)$. After differentiating the equation in (A.6) with

$$(A.100) \quad \begin{cases} X = \frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y}, \\ Y = \frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y}, \\ U = \frac{v}{y} + h, \\ V = h, \end{cases}$$

by h and putting $h = 0$, we obtain

$$(A.101) \quad \begin{aligned} & D_t \vartheta_4 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \cdot \vartheta_4 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\ &= \frac{1}{y} \frac{\vartheta_4' \left(\frac{v}{y} \right)}{\vartheta_4 \left(\frac{v}{y} \right)} \vartheta_4 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_4 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\ & - \frac{1}{y} \frac{\vartheta_1 \left(\frac{v}{y} \right) \vartheta_1' (0)}{\vartheta_4 \left(\frac{v}{y} \right) \vartheta_4 (0)} \vartheta_1 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_1 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\ &= \frac{1}{y} d_4 \left(\frac{v}{y}; \frac{2}{y} \right) \vartheta_4 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_4 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\ & - \frac{1}{y} Q_1(v, y) \vartheta_1 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_1 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right). \end{aligned}$$

Also, after differentiating the equation in (A.7) with

$$(A.102) \quad \begin{cases} X = \frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y}, \\ Y = \frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y}, \\ U = \frac{v}{y} + h, \\ V = h, \end{cases}$$

by h and putting $h = 0$, we obtain

$$(A.103) \quad \begin{aligned} & D_t \vartheta_1 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \cdot \vartheta_1 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\ &= \frac{1}{y} \frac{\vartheta_4' \left(\frac{v}{y} \right)}{\vartheta_4 \left(\frac{v}{y} \right)} \vartheta_1 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_1 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\ & - \frac{1}{y} \frac{\vartheta_1 \left(\frac{v}{y} \right) \vartheta_1' (0)}{\vartheta_4 \left(\frac{v}{y} \right) \vartheta_4 (0)} \vartheta_4 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_4 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\ &= \frac{1}{y} d_4 \left(\frac{v}{y}; \frac{2}{y} \right) \vartheta_1 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_1 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\ & - \frac{1}{y} Q_1(v, y) \vartheta_4 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_4 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right). \end{aligned}$$

Also, after differentiating the equation in (A.7) with

$$(A.104) \quad \begin{cases} X = \frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y}, \\ Y = \frac{v}{y} + h, \\ U = \frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y}, \\ V = h, \end{cases}$$

by h and putting $h = 0$, we obtain

$$\begin{aligned}
& D_t \vartheta_1 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \cdot \vartheta_4 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\
&= \frac{1}{y} \frac{\vartheta_1' \left(\frac{v}{y} \right)}{\vartheta_1 \left(\frac{v}{y} \right)} \vartheta_1 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_4 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\
\text{(A.105)} \quad & - \frac{1}{y} \frac{\vartheta_4 \left(\frac{v}{y} \right) \vartheta_1' (0)}{\vartheta_1 \left(\frac{v}{y} \right) \vartheta_4 (0)} \vartheta_4 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_1 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\
&= \frac{1}{y} d_1 \left(\frac{v}{y}; \frac{2}{y} \right) \vartheta_1 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_4 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\
& - \frac{1}{y} Q_2(v, y) \vartheta_4 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_1 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right).
\end{aligned}$$

Also, after differentiating the equation in (A.7) with

$$\text{(A.106)} \quad \begin{cases} X = \frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y}, \\ Y = \frac{v}{y} + h, \\ U = \frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y}, \\ V = h, \end{cases}$$

by h and putting $h = 0$, we obtain

$$\begin{aligned}
& D_t \vartheta_4 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \cdot \vartheta_1 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\
&= \frac{1}{y} \frac{\vartheta_1' \left(\frac{v}{y} \right)}{\vartheta_1 \left(\frac{v}{y} \right)} \vartheta_4 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_1 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\
\text{(A.107)} \quad & - \frac{1}{y} \frac{\vartheta_4 \left(\frac{v}{y} \right) \vartheta_1' (0)}{\vartheta_1 \left(\frac{v}{y} \right) \vartheta_4 (0)} \vartheta_1 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_4 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\
&= \frac{1}{y} d_1 \left(\frac{v}{y}; \frac{2}{y} \right) \vartheta_4 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_1 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right) \\
& - \frac{1}{y} Q_2(v, y) \vartheta_1 \left(\frac{v}{y} \left(n + \frac{1}{2} \right) + \frac{t}{y} \right) \vartheta_4 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right).
\end{aligned}$$

Adding (A.101) and (A.103) yields (A.93), and subtracting (A.105) from (A.107) yields (A.94). \square

Next, we construct the solution to the semi-discrete sine-Gordon equation (5.34).

Lemma A.10. *Assuming that the solution of (5.34) is expressed as (A.92), we obtain the following bilinear equations as a sufficient condition for satisfying (5.34).*

$$\text{(A.108)} \quad D_t A_{n+1} \cdot A_n - D_t B_{n+1} \cdot B_n = \frac{L_2}{2y} (\eta_2 - 1) (A_{n+1} A_n - B_{n+1} B_n),$$

$$\text{(A.109)} \quad D_t A_{n+1} \cdot B_n + D_t B_{n+1} \cdot A_n = \frac{L_2}{2y} (\eta_2 + 1) (A_{n+1} B_n + B_{n+1} A_n),$$

where η_2 is a real constant.

Proof. Since

$$\begin{aligned}
 \frac{d}{dt} \left(\frac{\Theta_{n+1} - \Theta_n}{2} \right) &= \frac{1}{i} \frac{d}{dt} \left(\log \frac{\tilde{\tau}_{n+1}}{\tilde{\tau}_{n+1}^*} - \log \frac{\tilde{\tau}_n}{\tilde{\tau}_n^*} \right) \\
 (A.110) \quad &= \frac{1}{i} \frac{d}{dt} \left(\log \frac{\tilde{\tau}_{n+1}}{\tilde{\tau}_n} - \log \frac{\tilde{\tau}_{n+1}^*}{\tilde{\tau}_n^*} \right) \\
 &= \frac{1}{i} \left(\frac{D_t \tilde{\tau}_{n+1} \cdot \tilde{\tau}_n}{\tilde{\tau}_{n+1} \tilde{\tau}_n} - \frac{D_t \tilde{\tau}_{n+1}^* \cdot \tilde{\tau}_n^*}{\tilde{\tau}_{n+1}^* \tilde{\tau}_n^*} \right).
 \end{aligned}$$

$$(A.111) \quad \sin \left(\frac{\Theta_{n+1} + \Theta_n}{2} \right) = \frac{1}{2i} \left(\frac{\tilde{\tau}_{n+1} \tilde{\tau}_n}{\tilde{\tau}_{n+1}^* \tilde{\tau}_n^*} - \frac{\tilde{\tau}_{n+1}^* \tilde{\tau}_n^*}{\tilde{\tau}_{n+1} \tilde{\tau}_n} \right),$$

hold, it can be seen that a sufficient condition for (A.92) to be a solution of (5.34) is that

$$(A.112) \quad D_t \tilde{\tau}_{n+1} \cdot \tilde{\tau}_n = -\frac{L_2}{2y} \tilde{\tau}_{n+1}^* \tilde{\tau}_n^* + \frac{L_2}{2y} \eta_2 \tilde{\tau}_{n+1} \tilde{\tau}_n,$$

(η_2 is a real constant) is satisfied. Substituting $\tilde{\tau}_n = A_n + iB_n$ for this, we get

$$\begin{aligned}
 &D_t (A_{n+1} \cdot A_n - B_{n+1} \cdot B_n) + iD_t (A_{n+1} \cdot B_n + B_{n+1} \cdot A_n) \\
 (A.113) \quad &= \frac{L_2}{2y} (\eta_2 - 1) (A_{n+1} A_n - B_{n+1} B_n) \\
 &+ i \frac{L_2}{2y} (\eta_2 + 1) (A_{n+1} B_n + B_{n+1} A_n).
 \end{aligned}$$

The real part of (A.113) yields (A.108), and the imaginary part yields (A.109). \square

In Lemma A.11, we construct solutions of (A.108) and (A.109) in terms of elliptic theta functions.

Lemma A.11. *The real functions A and B defined by (A.91) are the solutions of (A.108) and (A.109) with*

$$(A.114) \quad \eta_2 = \frac{d_1 \left(\frac{v}{y}; \frac{2}{y} \right) + d_4 \left(\frac{v}{y}; \frac{2}{y} \right) + Q_1(v, y) - Q_4(v, y)}{L_2(v, y)}.$$

Proof. In this proof, we set $w = 2i/y$ similarly to Lemma A.9. Subtracting (A.103) from (A.101) yields (A.108), and adding (A.107) and (A.105) yields (A.109). \square

A.9. Proof of Proposition 5.16. We first show (5.40) and (5.39) by using the periodicity of elliptic theta functions (A.11). Consider replacing the parameter with $\tilde{v} = v + y$. Then we see

$$\begin{aligned}
 (A.115) \quad &Q_1(\tilde{v}, y) = -Q_1(v, y), \quad Q_2(\tilde{v}, y) = -Q_2(v, y), \\
 &d_1(\tilde{v}, y) = d_1(v, y), \quad d_4(\tilde{v}, y) = d_4(v, y), \\
 &L_1(\tilde{v}, y) = L_2(v, y).
 \end{aligned}$$

and

$$\begin{aligned}
 (A.116) \quad \Theta_n(t; \tilde{v}, y) &= 4 \arctan \frac{\vartheta_1 \left(\frac{\tilde{v}}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right)}{\vartheta_4 \left(\frac{\tilde{v}}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} \right)} \\
 &= 4 \arctan \frac{\vartheta_1 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} + n - \frac{1}{2} \right)}{\vartheta_4 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} + n - \frac{1}{2} \right)} \\
 &= (-1)^n 4 \arctan \frac{\vartheta_1 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} - \frac{1}{2} \right)}{\vartheta_4 \left(\frac{v}{y} \left(n - \frac{1}{2} \right) + \frac{t}{y} - \frac{1}{2} \right)} \\
 &= (-1)^n \Theta_n \left(t - \frac{y}{2}; v, y \right).
 \end{aligned}$$

From the relation between the signed curvature angle κ and the potential function Θ (5.32), we see

$$(A.117) \quad \kappa_n(t; \tilde{v}) = (-1)^{n+1} \kappa_n\left(t - \frac{y}{2}; v\right).$$

Next, we check the transformation of λ (5.2). From (A.13) and (A.14), we have

$$(A.118) \quad \begin{aligned} & \lambda\left(\tilde{v}, \frac{1}{2} - r\right) \\ &= \frac{1}{i} \log \left(\frac{\vartheta_1\left(-\frac{1}{2}iv + \frac{1}{2} - r - \frac{1}{2}iy\right)}{\vartheta_1\left(\frac{1}{2}iv + \frac{1}{2} - r + \frac{1}{2}iy\right)} \frac{\vartheta_3\left(\frac{1}{2}iv + \frac{1}{2} - r + \frac{1}{2}iy\right)}{\vartheta_3\left(-\frac{1}{2}iv + \frac{1}{2} - r - \frac{1}{2}iy\right)} \right) \\ &= \frac{1}{i} \log \left(\frac{\vartheta_2\left(-\frac{1}{2}iv - r - \frac{1}{2}iy\right)}{\vartheta_2\left(\frac{1}{2}iv - r + \frac{1}{2}iy\right)} \frac{\vartheta_4\left(\frac{1}{2}iv - r + \frac{1}{2}iy\right)}{\vartheta_4\left(-\frac{1}{2}iv - r - \frac{1}{2}iy\right)} \right) \\ &= \frac{1}{i} \log \left(-\frac{\vartheta_3\left(-\frac{1}{2}iv - r\right)}{\vartheta_3\left(\frac{1}{2}iv - r\right)} \frac{\vartheta_1\left(\frac{1}{2}iv - r\right)}{\vartheta_1\left(-\frac{1}{2}iv - r\right)} \right) \\ &= \pi + \frac{1}{i} \log \left(\frac{\vartheta_1\left(-\frac{1}{2}iv + r\right)}{\vartheta_1\left(\frac{1}{2}iv + r\right)} \frac{\vartheta_3\left(\frac{1}{2}iv + r\right)}{\vartheta_3\left(-\frac{1}{2}iv + r\right)} \right) \\ &= \pi + \lambda(v, r). \end{aligned}$$

Finally, we see that the transformation of the normal \tilde{N} and the binormal \tilde{B} follows from the Frenet-Serret formula (2.6). \square

A.10. Proof of Proposition 5.18. First, we calculate $L_1(v, y)/y$ and $L_2(v, y)/y$. From (5.36), (A.15), (A.16) and (A.20), we see

$$(A.119) \quad \begin{aligned} \frac{L_1(v, y)}{y} &= \frac{1}{y} \frac{\vartheta_1'\left(\frac{v}{y}, i\frac{2}{y}\right) \vartheta_4\left(\frac{v}{y}, i\frac{2}{y}\right) - \vartheta_1\left(\frac{v}{y}, i\frac{2}{y}\right) \vartheta_4'\left(\frac{v}{y}, i\frac{2}{y}\right)}{\vartheta_1\left(\frac{v}{y}, i\frac{2}{y}\right) \vartheta_4\left(\frac{v}{y}, i\frac{2}{y}\right)} \\ &+ \frac{1}{y} \frac{\vartheta_1'\left(0, i\frac{2}{y}\right) \vartheta_1\left(\frac{v}{y}, i\frac{2}{y}\right)}{\vartheta_4\left(0, i\frac{2}{y}\right) \vartheta_4\left(\frac{v}{y}, i\frac{2}{y}\right)} + \frac{1}{y} \frac{\vartheta_1'\left(0, i\frac{2}{y}\right) \vartheta_4\left(\frac{v}{y}, i\frac{2}{y}\right)}{\vartheta_4\left(0, i\frac{2}{y}\right) \vartheta_1\left(\frac{v}{y}, i\frac{2}{y}\right)} \\ &= i \frac{\vartheta_1'\left(0, iy\right) \vartheta_4(iv, iy)}{\vartheta_4\left(0, iy\right) \vartheta_1(iv, iy)} + i \frac{\vartheta_1'\left(0, iy\right) \vartheta_2(iv, iy)}{\vartheta_2\left(0, iy\right) \vartheta_1(iv, iy)} \\ &= i \frac{\vartheta_1'\left(0, iy\right) \vartheta_4(iv, iy) \vartheta_2\left(0, iy\right) + \vartheta_2(iv, iy) \vartheta_4\left(0, iy\right)}{\vartheta_1(iv, iy) \vartheta_4\left(0, iy\right) \vartheta_2\left(0, iy\right)}. \end{aligned}$$

Similarly, from (5.37), (A.15), (A.16) and (A.20), we see

$$(A.120) \quad \begin{aligned} \frac{L_2(v, y)}{y} &= \frac{1}{y} \frac{\vartheta_1'\left(\frac{v}{y}, i\frac{2}{y}\right) \vartheta_4\left(\frac{v}{y}, i\frac{2}{y}\right) - \vartheta_1\left(\frac{v}{y}, i\frac{2}{y}\right) \vartheta_4'\left(\frac{v}{y}, i\frac{2}{y}\right)}{\vartheta_1\left(\frac{v}{y}, i\frac{2}{y}\right) \vartheta_4\left(\frac{v}{y}, i\frac{2}{y}\right)} \\ &- \frac{1}{y} \frac{\vartheta_1'\left(0, i\frac{2}{y}\right) \vartheta_1\left(\frac{v}{y}, i\frac{2}{y}\right)}{\vartheta_4\left(0, i\frac{2}{y}\right) \vartheta_4\left(\frac{v}{y}, i\frac{2}{y}\right)} - \frac{1}{y} \frac{\vartheta_1'\left(0, i\frac{2}{y}\right) \vartheta_4\left(\frac{v}{y}, i\frac{2}{y}\right)}{\vartheta_4\left(0, i\frac{2}{y}\right) \vartheta_1\left(\frac{v}{y}, i\frac{2}{y}\right)} \\ &= i \frac{\vartheta_1'\left(0, iy\right) \vartheta_4(iv, iy)}{\vartheta_4\left(0, iy\right) \vartheta_1(iv, iy)} - i \frac{\vartheta_1'\left(0, iy\right) \vartheta_2(iv, iy)}{\vartheta_2\left(0, iy\right) \vartheta_1(iv, iy)} \\ &= i \frac{\vartheta_1'\left(0, iy\right) \vartheta_4(iv, iy) \vartheta_2\left(0, iy\right) - \vartheta_2(iv, iy) \vartheta_4\left(0, iy\right)}{\vartheta_1(iv, iy) \vartheta_4\left(0, iy\right) \vartheta_2\left(0, iy\right)}. \end{aligned}$$

On the other hand, from (4.2) and (5.2), we see

$$(A.121) \quad \begin{aligned} & 1 + \cos \lambda \\ &= \frac{\left\{ \vartheta_1\left(\frac{iv}{2} + r, iy\right) \vartheta_3\left(-\frac{iv}{2} + r, iy\right) + \vartheta_1\left(-\frac{iv}{2} + r, iy\right) \vartheta_3\left(\frac{iv}{2} + r, iy\right) \right\}^2}{2u_2}, \end{aligned}$$

$$(A.122) \quad \begin{aligned} & 1 - \cos \lambda \\ &= - \frac{\{\vartheta_1\left(\frac{iv}{2} + r, iy\right) \vartheta_3\left(-\frac{iv}{2} + r, iy\right) - \vartheta_1\left(-\frac{iv}{2} + r, iy\right) \vartheta_3\left(\frac{iv}{2} + r, iy\right)\}^2}{2u_2}. \end{aligned}$$

From (A.119) and (A.121), we see that (5.43) holds. Similarly, from (A.120) and (A.122), we see that (5.44) holds. \square

A.11. Special property of elliptic theta functions when w is a pure imaginary number.

Lemma A.12. *The following holds for any $z \in \mathbb{C}$ when $w = iy$.*

$$(A.123) \quad \vartheta_j(z)^* = \vartheta_j(z^*), \quad j = 1, 2, 3, 4.$$

Proof. Consider the following theta function with characteristics.

$$(A.124) \quad \vartheta_{a,b}(z) := \sum_{n \in \mathbb{Z}} \exp(2\pi i(n+a)(z+b) + \pi i(n+a)^2 \omega), \quad a, b \in \left\{0, \frac{1}{2}\right\},$$

then,

$$(A.125) \quad \vartheta_1(z) = -\vartheta_{\frac{1}{2}, \frac{1}{2}}(z), \quad \vartheta_2(z) = \vartheta_{\frac{1}{2}, 0}(z), \quad \vartheta_3(z) = \vartheta_{0,0}(z), \quad \vartheta_4(z) = \vartheta_{0, \frac{1}{2}}(z),$$

hold. Since $w^* = -w$, we have

$$(A.126) \quad \begin{aligned} \vartheta_{a,b}(z)^* &= \sum_{n \in \mathbb{Z}} \exp(-2\pi i(n+a)(z^*+b) - \pi i(n+a)^2 w^*) \\ &= \sum_{n \in \mathbb{Z}} \exp(2\pi i(-n-a)(z^*+b) + \pi i(n+a)^2 w) \\ &= \sum_{n \in \mathbb{Z}} \exp(2\pi i(n+a)(z^*+b) + \pi i(n+a)^2 w) \\ &= \vartheta_{a,b}(z^*). \end{aligned}$$

\square

This lemma guarantees that some τ functions, such as F_n , are real-valued functions.

References

- [1] *T. F. Banchoff and J. H. White*, The behavior of the total twist and self-linking number of a closed space curve under inversions, *Mathematica Scandinavica* **36** (1975), no. 2, 254–262.
- [2] *A. Bobenko and U. Pinkall*, Discrete surfaces with constant negative Gaussian curvature and the Hirota equation, *Journal of Differential Geometry* **43** (1996), no. 3, 527–611.
- [3] *R. Bricard*, *Leçons de cinématique*, volume 1, Gauthier-Villars, 1926.
- [4] *A. Calini and T. Ivey*, Bäcklund transformations and knots of constant torsion, *Journal of Knot Theory and Its Ramifications* **7** (1998), no. 06, 719–746.
- [5] *A. Doliwa and P. M. Santini*, Integrable dynamics of a discrete curve and the Ablowitz–Ladik hierarchy, *Journal of Mathematical Physics* **36** (1995), no. 3, 1259–1273.
- [6] *B.-F. Feng, J. Inoguchi, K. Kajiwara, K. Maruno and Y. Ohta*, Discrete integrable systems and hodograph transformations arising from motions of discrete plane curves, *Journal of Physics A: Mathematical and Theoretical* **44** (2011), no. 39, 395201.
- [7] *B.-F. Feng, K. Maruno and Y. Ohta*, Integrable discretizations of the short pulse equation, *Journal of Physics A: Mathematical and Theoretical* **43** (2010), no. 8, 085203.
- [8] *Fowler PW and Guest SD*, A symmetry analysis of mechanisms in rotating rings of tetrahedra, *P ROY SOC A-MATH PHY* **461** (2005), 1829–1846, , URL . Accessed on 2023/01/05.
- [9] *H. Hasimoto*, A soliton on a vortex filament, *Journal of Fluid Mechanics* **51** (1972), no. 3, 477–485.
- [10] *S. Hirose, J. Inoguchi, K. Kajiwara, N. Matsuura and Y. Ohta*, Discrete local induction equation, *Journal of Integrable Systems* **4** (2019), no. 1, .

- [11] *R. Hirota*, The Direct Method in Soliton Theory, Cambridge Tracts in Mathematics, Cambridge University Press, 2004, .
- [12] *T. Hoffmann* and *N. Kutz*, Discrete curves in and the Toda lattice, Studies in Applied Mathematics **113** (2004), no. 1, 31–55.
- [13] *J. Inoguchi*, *K. Kajiwara*, *N. Matsuura* and *Y. Ohta*, Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves, Journal of Physics A: Mathematical and Theoretical **45** (2012), no. 4, 045206.
- [14] *J. Inoguchi*, *K. Kajiwara*, *N. Matsuura* and *Y. Ohta*, Motion and Bäcklund transformations of discrete plane curves, Kyushu Journal of Mathematics **66** (2012), no. 2, 303–324.
- [15] *J. Inoguchi*, *K. Kajiwara*, *N. Matsuura* and *Y. Ohta*, Discrete mKdV and discrete sine-Gordon flows on discrete space curves, Journal of Physics A: Mathematical and Theoretical **47** (2014), no. 23, 235202.
- [16] *S. Kaji*, Computer program: Geometry of the moduli space of a closed linkage. Available online at <https://github.com/shizuo-kaji/Kaleidocycle>.
- [17] *S. Kaji*, A closed linkage mechanism having the shape of a discrete Möbius strip, Symposium Proceedings of JSPE Semestrial Meeting 2018 (2018), 62–65. English translation at arXiv:1909.02885.
- [18] *S. Kaji*, *K. Kajiwara* and *H. Park*, Linkage Mechanisms Governed by Integrable Deformations of Discrete Space Curves, CRC Press, 2019, volume 2, 356–381, .
- [19] *S. Kaji*, *J. Schönke*, *E. Fried* and *M. Grunwald*, Moebius kaleidocycle, 2023. Patent (filed on Feb. 2018), JP7261490, WO2019167941.
- [20] *S. Kharchev* and *A. Zabrodin*, Theta vocabulary I, Journal of Geometry and Physics **94** (2015), 19–31.
- [21] *G. L. Lamb*, Solitons and the motion of helical curves, Phys. Rev. Lett. **37** (1976), 235–237, , URL <https://link.aps.org/doi/10.1103/PhysRevLett.37.235>.
- [22] *J. Langer* and *D. A. Singer*, Knotted elastic curves in \mathbb{R}^3 , Journal of the London Mathematical Society **2** (1984), no. 3, 512–520.
- [23] *D. Mumford* and *C. Musili*, Tata lectures on theta. i (modern birkhäuser classics), Birkhäuser Boston Incorporated, 2007.
- [24] *C. Rogers* and *W. K. Schief*, Bäcklund and Darboux Transformations: Geometry and Modern Applications in Soliton Theory, Cambridge Texts in Applied Mathematics, Cambridge University Press, 2002, .
- [25] *C. Safsten*, *T. Fillmore*, *A. Logan*, *D. Halverson* and *L. Howell*, Analyzing the stability properties of kaleidocycles, Journal of Applied Mechanics **83** (2016), no. 5, .
- [26] *J. Schönke* and *E. Fried*, Single degree of freedom everting ring linkages with nonorientable topology, Proceedings of the National Academy of Sciences **116** (2019), no. 1, 90–95, , URL <https://www.pnas.org/doi/abs/10.1073/pnas.1809796115>.

Shizuo Kaji, Institute of Mathematics for Industry, Kyushu University, 744 Motoooka, Nishi-ku, Fukuoka
819-0395, Japan
e-mail: skaji@imi.kyushu-u.ac.jp

Kenji Kajiwara, Institute of Mathematics for Industry, Kyushu University, 744 Motoooka, Nishi-ku, Fukuoka
819-0395, Japan
e-mail: kaji@imi.kyushu-u.ac.jp

Shota Shigetomi, Institute of Mathematics for Industry, Kyushu University, 744 Motoooka, Nishi-ku, Fukuoka 819-
0395, Japan
e-mail: s-shigetomi@imi.kyushu-u.ac.jp

Eingegangen XXX, in revidierter Fassung XXX