# APPROXIMATION PROPERTIES FOR DYNAMICAL W<sup>∗</sup> -CORRESPONDENCES

K. DE COMMER AND J. DE RO

ABSTRACT. Let  $\mathbb G$  be a locally compact quantum group, and  $A, B$  von Neumann algebras on which  $\mathbb G$  acts. We refer to these as G*-dynamical W*∗*-algebras*. We make a study of G-equivariant A-B-correspondences, that is, Hilbert spaces  $H$  with an A-B-bimodule structure by \*-preserving normal maps, and equipped with a unitary representation of G which is equivariant with respect to the above bimodule structure. Such structures are a Hilbert space version of the theory of G-equivariant Hilbert C∗-bimodules. We show that there is a well-defined Fell topology on equivariant correspondences, and use this to formulate approximation properties for them. Within this formalism, we then characterize amenability of the action of a locally compact group on a von Neumann algebra, using recent results due to Bearden and Crann. We further consider natural operations on equivariant correspondences such as taking opposites, composites and crossed products, and examine the continuity of these operations with respect to the Fell topology.

## **INTRODUCTION**

In the setting of representation theory and abstract harmonic analysis, amenability properties can often be characterized by the possibility to approximate a pre-determined restricted class of 'trivial representations' within a pre-determined restricted class of 'regular representations'. We then say that the former are 'weakly contained' in the latter. These classes consist in many cases of just one object. Two examples stand out:

- If G is a locally compact group, then G is amenable if and only if the trivial representation is weakly contained in the regular representation on  $L^2(G)$ .
- If A is a von Neumann algebra, then A is amenable if and only if the trivial A-bimodule  $L^2(A)$  is weakly contained in the regular (or *coarse*) A-bimodule  $L^2(A) \otimes L^2(A)$ .

In this last example, we mean bimodules in the sense of *correspondences* [\[Con80+\]](#page-46-0) (see also [\[Pop86\]](#page-47-0), [\[Con94,](#page-46-1) Appendix B. we are given two von Neumann algebras A, B and a Hilbert space  $\mathcal H$  equipped with a normal unital ∗-representation  $\pi$  of A and a normal unital anti-∗-representation  $\rho$  of B for which H forms an A-B-bimodule.

Characterizing amenability in this setting has led to some early breakthrough results in the modern theory of von Neumann algebras by its relation to injectivity [\[Con76,](#page-46-2)[CE77,](#page-46-3)[EL77\]](#page-47-1), and continues to play a fundamental role in many aspects of operator theory. For example, the notion of *relative amenability*, introduced in [\[Pop86\]](#page-47-0) and further explored in the general context of correspondences in [\[A-D90,](#page-46-4) [A-D95\]](#page-46-5), was only recently shown to be equivalent with the corresponding notion of relative injectivity [\[BMO20\]](#page-46-6).

By means of the crossed product construction, close ties can be made between amenability of actions of locally compact groups on von Neumann algebras [\[A-D79\]](#page-46-7) and approximation properties related to correspondences constructed from the von Neumann algebra and its crossed product. In the setting of dynamical systems, such approximation properties are connected to amenability properties for certain associated groupoids [\[A-DR00\]](#page-46-8). We also mention the recent key results relating amenability of C<sup>\*</sup>-dynamical systems to amenability of W<sup>\*</sup>dynamical systems [\[BEW21,](#page-46-9) [OS21\]](#page-47-2).

In [\[KV00,](#page-47-3)[KV03\]](#page-47-4), a generalization was made of the theory of locally compact groups to that of locally compact quantum groups (we refer to these articles for more on the history of the subject). It is based on the notion of a Hopf-von Neumann algebra [\[Ern67\]](#page-47-5), which we now recall.

Definition 0.1. A Hopf-von Neumann algebra consists of a von Neumann algebra M with a unital normal faithful ∗-homomorphism  $\Delta: M \to M \bar{\otimes} M$  satisfying the coassociativity condition

$$
(\Delta \otimes id) \circ \Delta = (id \otimes \Delta) \circ \Delta.
$$

It is called involutive if it is equipped with an involutive anti-∗-automorphism  $R : M \to M$  satisfying

$$
\Delta \circ R = (R \otimes R) \circ \Delta^{\mathrm{op}},
$$

where  $\Delta^{op}$  is the composition of  $\Delta$  with the flip map  $\varsigma : x \otimes y \mapsto y \otimes x$ .

If  $(M, \Delta)$  is a Hopf-von Neumann algebra, we sometimes write  $\mathbb{G} = (M, \Delta)$  and  $M = L^{\infty}(\mathbb{G})$ . Then we say that G is a *locally compact quantum group* if moreover  $(M, \Delta)$  possesses invariant weights (Definition [1.1\)](#page-5-0). Such Hopf-von Neumann algebras are automatically involutive with respect to a canonical involution R.

The von Neumann algebraic framework for locally compact quantum groups is very flexible and high-powered. For example, consider the following notions for a Hopf-von Neumann algebra  $\mathbb{G} = (M, \Delta)$ .

<span id="page-1-0"></span>**Definition 0.2.** A unitary representation of  $\mathbb{G}$  on a Hilbert space H is a unitary  $U \in B(H) \bar{\otimes} M$  such that  $(id \otimes \Delta)U = U_{12}U_{13}.$  (0.1)

Here  $U_{12} = U \otimes 1$ , while  $U_{13}$  is  $(id \otimes \varsigma)(U \otimes 1)$ .

**Definition 0.3.** If A is a von Neumann algebra, an action of  $\mathbb{G}$  on A is a unital normal faithful  $*$ homomorphism

<span id="page-1-2"></span> $\alpha: A \to A \bar{\otimes} M$ ,

called the associated coaction, and satisfying the coaction property

$$
(\alpha \otimes id)\alpha = (id \otimes \Delta)\alpha.
$$

We also say that  $(A, \alpha)$  is a G-dynamical von Neumann algebra, or simply a G-W<sup>\*</sup>-algebra.

In the general context of Hopf-von Neumann algebras, these two notions have no apparent relation, but when  $\mathbb{G} = (M, \Delta)$  is a locally compact quantum group, there is a beautiful generalisation, due to S. Vaes [\[Vae01\]](#page-47-6), of Haagerup's canonical implementation of a group action on a von Neumann algebra [\[Haa78\]](#page-47-7): there exists a canonical unitary representation  $U_{\alpha}$  of  $\mathbb{G}$  on the standard Hilbert space  $L^2(A)$  such that

<span id="page-1-1"></span>
$$
\alpha(a) = U_{\alpha}(a \otimes 1)U_{\alpha}^*, \qquad \forall a \in A,
$$
\n(0.2)

where we view  $A \subseteq B(L^2(A))$  in standard position.

From an abstract and even an aesthetic point of view, one of the main salient features of locally compact quantum groups is that they allow an extension of the Pontryagin duality for arbitrary abelian locally compact groups to a Pontryagin duality for arbitrary locally compact quantum groups. When restricting to locally compact groups G, this is the correspondence sending the function algebra  $L^{\infty}(G)$  (with its natural coproduct encoding the group structure) into the group von Neumann algebra  $\mathcal{R}(G)$  generated by right translation operators (with its natural diagonal coproduct), and thus not an operation which gives a duality on a single category in this restricted setting. Various approximation properties such as amenability, Haagerup property, property (T),... can also be formulated naturally within the much more general setting of locally compact quantum groups, see e.g. [\[Voi77,](#page-47-8) [Rua96,](#page-47-9) [DQV02,](#page-46-10) [Tom06,](#page-47-10) [Fim10,](#page-47-11) [KyS12,](#page-47-12) [SV14,](#page-47-13) [CN15,](#page-46-11) [DFSW16\]](#page-46-12).

In this paper, we combine the above two settings by studying *equivariant correspondences*. This notion is a verbatim generalisation of the corresponding notion for  $\mathbb{G}\text{-}C^*$ -algebras with  $\mathbb{G}$  a compact quantum group, introduced in [\[AS21\]](#page-46-13).

<span id="page-1-3"></span>**Definition 0.4.** Let  $\mathbb{G} = (M, \Delta, R)$  be an involutive Hopf-von Neumann algebra, and let  $(A, \alpha), (B, \beta)$  be  $\mathbb{G}$ -W<sup>∗</sup> -algebras. A G-equivariant A-B-correspondence, or simply a G-A-B-correspondence, consists of a Hilbert space H with

• a normal unital  $*$ -representation  $\pi$  of A on H,

- a normal unital anti-∗-representation ρ of B on H, and
- a unitary representation  $U$  of  $\mathbb G$  on  $\mathcal H$ ,

satisfying for all  $a \in A$  and  $b \in B$  the compatibility relations  $\pi(a)\rho(b) = \rho(b)\pi(a)$  and

<span id="page-2-0"></span>
$$
U(\pi(a)\otimes 1)U^* = (\pi\otimes id)\alpha(a), \qquad U^*(\rho(b)\otimes 1)U = (\rho\otimes R)(\beta(b)).\tag{0.3}
$$

C<sup>\*</sup>-algebraic analogues of such equivariant correspondences, in the context of equivariant Hilbert C<sup>\*</sup>-modules, have already appeared in the original works [\[BS89,](#page-46-14) [BS93\]](#page-46-15) in relation to equivariant KK-theory, and have remained important in follow-up works (see e.g. [\[BC17\]](#page-46-16)). In [\[AS21\]](#page-46-13), concrete use was made rather of the above Hilbert space picture of equivariant correspondences, however within the context of C<sup>∗</sup> -algebras and *compact* quantum groups. On the other hand, in [\[Vae05\]](#page-47-14) there already appear specific equivariant correspondences, albeit in the particular case arising from inclusions of quantum groups, and using instead the (equivalent) notion of equivariant von Neumann bimodules.

More particular cases of interest have turned up recently within the setting of quantum automorphism *groups*  $\text{[RV22]}$  $\text{[RV22]}$  $\text{[RV22]}$  (for related results in a more classical setting, see  $\text{[AV16]}$ ), and *quantizations of semisimple* Lie groups [\[DCDT21\]](#page-46-18). It is our hope that the framework of this paper will be a convenient one to study analytic approximation properties for these important classes of examples.

Let us now come to the main results of this paper.

**Theorem** (Theorem [2.16\)](#page-13-0). Let  $\mathbb{G}$  be a locally compact quantum group, and A, B two  $\mathbb{G}$ -W<sup>\*</sup>-algebras. Then there exists a universal  $C^*$ -algebra  $C^{\mathbb{G}}(A, B)$  such that any G-equivariant A-B-correspondence H uniquely lifts to a non-degenerate  $*$ -representation  $\theta_{\mathcal{H}}$  of  $C^{\mathbb{G}}(A, B)$  on  $\mathcal{H}$ .

The C<sup>\*</sup>-algebra  $C^{G}(A, B)$  is a C<sup>\*</sup>/W<sup>\*</sup>-intermediate of the double cross product C<sup>\*</sup>-algebra appearing in [\[AS21\]](#page-46-13) (see also [\[NV02,](#page-47-16) Section 2.6] for a description in a purely algebraic context).

Due to the above theorem, we can introduce a Fell topology on the class of G-equivariant A-B-correspondences, as well as a notion of weak containment: we say G is weakly contained in H if the ∗-representation  $\theta_G$  of  $C^{\mathbb{G}}(A, B)$  factors through  $\theta_{\mathcal{H}}(C^{\mathbb{G}}(A, B))$ . When we want to emphasize that we are working in the equivariant setting, we will sometimes use the term equivariantly weakly contained.

Our next main result offers another characterisation of equivariant weak containment. If  $(\mathcal{H}, U_{\mathcal{H}}),(\mathcal{G}, U_{\mathcal{G}})$ are G-A-B-correspondences, we use the notation

<span id="page-2-1"></span>
$$
X_B(\mathcal{G}, \mathcal{H}) = \{x : \mathcal{G} \to \mathcal{H} \mid x\rho_{\mathcal{G}}(b) = \rho_{\mathcal{H}}(b)x, \forall b \in B\},\tag{0.4}
$$

and

$$
\alpha_{X_B(\mathcal{G},\mathcal{H})}: X_B(\mathcal{G},\mathcal{H}) \to X_B(\mathcal{G},\mathcal{H}) \bar{\otimes} M, \qquad x \mapsto U_{\mathcal{H}}(x \otimes 1)U_{\mathcal{G}}^*.
$$

**Theorem** (Theorem [3.7\)](#page-21-0). Let  $\mathbb{G}$  be a locally compact quantum group, and let  $A, B$  be  $\mathbb{G}$ -W<sup>\*</sup>-algebras. Let  $\mathcal{H}, \mathcal{G}$  be G-A-B-correspondences, and put  $\mathcal{H}^{\infty} = \bigoplus_{1}^{\infty} \mathcal{H}$ .

Then G is equivariantly weakly contained in H if and only if there exists a 1-bounded net  $y_i \in X_B(\mathcal{G}, H^\infty)$ such that for all  $a \in A$ 

$$
(y_i^* \pi_{\mathcal{H}^{\infty}}(a) \otimes 1)\alpha_{X_B(\mathcal{G}, \mathcal{H})}(y_i) \to \pi_{\mathcal{G}}(a) \otimes 1 \quad \sigma\text{-weakly in } B(\mathcal{G})\bar{\otimes}M. \tag{0.5}
$$

The above results naturally lead to a notion of amenability for inclusions of  $\mathbb{G}\text{-}W^*$ -algebras: if  $A \subseteq B$  is  $\mathbb{G}\text{-}$ equivariant, then we can say that this inclusion is *strongly equivariantly amenable* if  $_A L^2(A)_A$  is equivariantly weakly contained in  $_A L^2(B)_A$  (these correspondences can easily be made equivariant using the results of [\[Vae01\]](#page-47-6), see Example [2.2\)](#page-8-0). The following theorem then shows that equivariant strong amenability implies equivariant amenability (=equivariant relative injectivity).

**Theorem** (Theorem [4.3\)](#page-23-0). Let  $\mathbb{G}$  be a locally compact quantum group, and let  $(A, \alpha) \subset (B, \beta)$  be an equivariant inclusion of  $\mathbb{G}\text{-}W^*$ -algebras. If  $_A L^2(A)_A$  is equivariantly weakly contained in  $_A L^2(B)_A$ , then there exists a G-equivariant conditional expectation  $E : B \to A$ , i.e.

<span id="page-3-0"></span>
$$
(E \otimes id) \circ \beta = \alpha \circ E. \tag{0.6}
$$

As a particular case in point, one can consider for a  $\mathbb{G}\text{-}W^*$ -algebra  $(A, \alpha)$  the equivariant inclusion

$$
A \cong \alpha(A) \subseteq A \bar{\otimes} M,
$$

where  $\widehat{A} \otimes M$  is endowed with the G-action id<sub>A</sub>  $\otimes \Delta$ . Our next result is then a converse of the last result above, in the case of a usual locally compact group  $G$ . It heavily relies on [\[BC22\]](#page-46-19).

**Theorem** (Theorem [4.10\)](#page-24-0). Let G be a locally compact group, and let A be a  $G-W^*$ -algebra via

$$
\alpha: A \to A \bar{\otimes} L^{\infty}(G).
$$

Then  $_A L^2(A)_A$  is weakly contained in  $_{\alpha(A)} (L^2(A) \otimes L^2(G))_{\alpha(A)}$  if and only if there exists a G-equivariant conditional expectation  $E: A\overline{\otimes} L^{\infty}(G) \rightarrow A$ .

As mentioned, an important aspect of locally compact quantum groups concerns their Pontryagin duality theory. We recall (see Section [1](#page-5-1) for details) that if A is a G-dynamical von Neumann algebra, and  $\ddot{G}$  the Pontryagin dual, one can construct the *crossed product von Neumann algebra*  $A \rtimes \mathbb{G}$ , which becomes a  $\mathbb{G}\text{-}d$  -dynamical von Neumann algebra. We will show that this duality extends to the level of  $\mathbb{G}\text{-}$ -equivariant correspondences: to any G-equivariant A-B-correspondence H is associated a G-equivariant  $A \rtimes \mathbb{G}$ -B  $\rtimes$  Gcorrespondence  $\mathcal{H} \rtimes \mathbb{G}$ .

Our final main result is then the following topological refinement of the well-known Takesaki-Takai duality.

Theorem (Theorem [5.26](#page-33-0) and Theorem [6.4\)](#page-38-0). The assignment

$$
\mathcal{H} \mapsto \mathcal{H} \rtimes \mathbb{G}
$$

is an equivalence between the W<sup>\*</sup>-category of  $\mathbb{G}$ -equivariant A-B-correspondences and the W<sup>\*</sup>-category of  $\mathbb{G}\text{-}equivariant } A \rtimes \mathbb{G}\text{-}B \rtimes \mathbb{G}\text{-}correspondences. Moreover, this correspondence preserves the Fell topology.}$ 

The precise contents of this paper are as follows.

In the *first section*, we provide some preliminaries on locally compact quantum groups and their actions. In the second section, we introduce equivariant correspondences, provide some elementary examples, and show that the category of equivariant correspondences admits a well-defined and well-behaved Fell topology. In the third section, we introduce the notion of weak containment between equivariant correspondences, and provide an equivalent formulation for weak containment in terms of one-sided intertwiners. In the fourth section, we introduce the notion of amenability and strong amenability for equivariant correspondences, and show that these notions can be used to introduce corresponding notions for actions of locally compact quantum groups on von Neumann algebras. We show that in the setting of locally compact groups, an amenable action is automatically strongly amenable, using the results of [\[BC22\]](#page-46-19). In the fifth section, we consider various operations on equivariant correspondences such as opposites, composites and crossed products. The main result here is a generalized Takesaki-Takai-duality, providing an equivalence of W<sup>∗</sup> -bicategories between G-equivariant correspondences and  $\ddot{G}$ -equivariant correspondences. In the *sixth section*, we consider the interplay between the above constructions and the Fell topology, and consider some direct corollaries for the associated approximation properties. We end in the seventh section with a brief outlook.

### Notations and conventions

All our vector spaces are complex. If V is a vector space and  $S \subseteq V$ , we denote Span(S) for the linear span of  $S$ . If  $V, W$  are subspaces of an algebra  $A$ , we denote

$$
VW = \text{Span}\{vw \mid v \in V, w \in W\} \subseteq A.
$$

If V is a normed vector space and  $S \subseteq V$ , we denote [S] for the norm-closure of the linear span of S. We denote ⊙ for algebraic tensor products (over C). We use ⊗ for the tensor product of Hilbert spaces, or the

minimal tensor product of C<sup>∗</sup> -algebras or Hilbert C<sup>∗</sup> -modules. We use ⊗¯ for the spatial tensor product of von Neumann algebras. The inner product of Hilbert spaces is assumed anti-linear in the first variable.

When referring to linear maps between operator spaces, we use the abbrevation cb for *completely bounded*, and, when considering linear maps between operator systems, we use the abbreviation  $(u/c)$ cp for *(uni*tal/contractive) completely positive.

We will use the terms W<sup>\*</sup>-algebra and von Neumann algebra interchangeably. When  $M$  is a von Neumann algebra, we denote  $M_*$  for the predual and  $M_+$  for the positive cone. If  $\Phi$  is an nsf (= normal, semifinite, faithful) weight on  $M$ , we use the standard notations.

$$
\mathscr{M}_\Phi^+ = \{ x \in M^+ : \Phi(x) < \infty \}, \quad \mathscr{N}_\Phi = \{ x \in M \mid \Phi(x^*x) < \infty \}.
$$

We let  $L^2(M)$  be the standard Hilbert space of M, with modular conjugation  $J = J_M$  and  $\Lambda_{\Phi}: \mathcal{N}_{\Phi} \to L^2(M)$ the canonical GNS-map of  $\Phi$ . We denote  $\pi_M$  for the associated left representation of M on  $L^2(M)$ , and

<span id="page-4-0"></span>
$$
\rho_M(x) = J\pi_M(x)^* J \tag{0.7}
$$

for the associated anti- $*$ -representation of M on  $L^2(M)$ . When there is no ambiguity, we drop the symbol  $\pi_M$ , and identify  $M \subseteq B(L^2(M))$ . For the basics of weight theory on von Neumann algebras, we refer to [\[Tak03\]](#page-47-17).

If H is a Hilbert space, we denote  $\mathcal{K}(\mathcal{H})$  for the C<sup>\*</sup>-algebra of compact operators on H. If  $u \in B(\mathcal{H})$  is a unitary, we write  $\text{Ad}(u)(x) = \text{Ad}_u(x) = uxu^*$  for  $x \in B(H)$ . More generally, when  $H, G$  are Hilbert spaces, and  $x \in B(\mathcal{G}, \mathcal{H})$ , we will denote by  $\text{Ad}_x$  the completely positive map

$$
Ad_x: B(\mathcal{G}) \to B(\mathcal{H}), \qquad T \mapsto xTx^*.
$$

If  $\xi, \eta \in \mathcal{H}$ , we write

$$
\omega_{\xi,\eta} \in B(\mathcal{H})_*, \qquad B(\mathcal{H}) \ni x \mapsto \langle \xi, x\eta \rangle \in \mathbb{C}.
$$

If  $H, \mathcal{G}$  are Hilbert spaces, we denote the *flip map* by

$$
\Sigma: \mathcal{H} \otimes \mathcal{G} \to \mathcal{G} \otimes \mathcal{H}, \qquad \xi \otimes \eta \mapsto \eta \otimes \xi.
$$

We use the *leq numbering notation*: if x is an operator on a tensor product of Hilbert spaces, we write

$$
x_{12}=x\otimes 1, \qquad x_{23}=1\otimes x, \qquad x_{13}=(\mathop{\mathrm{id}}\nolimits\otimes\mathop{\mathrm{Ad}}\nolimits\Sigma)x_{12}=(\mathop{\mathrm{Ad}}\nolimits\Sigma\otimes\mathop{\mathrm{id}}\nolimits)x_{23}.
$$

In practice, there should be no ambiguity on where the unit acts.

For C a C<sup>\*</sup>-algebra, we write  $C^{**}$  for the universal W<sup>\*</sup>-envelope of C. Throughout the theory of operator algebras and quantum groups, the multiplier algebra  $M(C)$  of C plays a prominent role. However, in this paper, we will encounter a situation where the use of the multiplier  $C^*$ -algebra is no longer sufficient, and where it is necessary (cfr. Remark [2.15\)](#page-12-0) to work with an even larger object, less familiar than the multiplier algebra.

The set of quasi-multipliers of C, originally introduced in [\[AP73\]](#page-46-20), is defined as the norm-closed subspace

$$
QM(C) = \{ x \in C^{**} \mid CxC \subseteq C \} \subseteq C^{**}.
$$

In general, the space  $QM(C)$  is very complicated, partly due to its lack of multiplicative structure. To give some intuition to what kind of elements this space contains, we provide another description for it as follows: Write  $\text{Bil}_Q(C)$  for the vector space of bilinear maps  $\phi: C \times C \rightarrow C$  satisfying

$$
\phi(pq, rs) = p\phi(q, r)s, \quad p, q, r, s \in C.
$$

It is clear that there is a canonical injective linear map, given by

$$
QM(C) \to \text{Bil}_Q(C): x \mapsto (\phi_x : (p, q) \mapsto pxq).
$$

It follows from [\[AP73,](#page-46-20) Proposition 4.2] that this map is bijective. Therefore, we can identify  $QM(C) \cong$  $Bil<sub>Q</sub>(C)$  as linear spaces. In very specific examples, an explicit description of the space  $QM(C)$  can be given, see e.g. [\[Li89,](#page-47-18) Section 8].

We refer to [\[Ped79,](#page-47-19) Section 3.12] for further basic facts on quasi-multipliers, but we recall below the main properties that we will use. First of all, we emphasize that  $QM(C)$ , although not necessarily multiplicatively closed, always naturally has the structure of an operator system. Next, we recall from [\[Ped79\]](#page-47-19) that whenever  $C \subseteq B(H)$  is a faithful, non-degenerate ∗-representation, the corresponding normal unital \*-homomorphism  $C^{**} \to B(\mathcal{H})$  is faithful on  $QM(C)$ , and we may isometrically identify through this map

$$
QM(C) = \{ x \in B(\mathcal{H}) \mid CxC \subseteq C \}.
$$

For a general non-degenerate ∗-representation  $\pi : C \to B(H)$ , we will always extend it to a map  $\pi$ :  $QM(C) \to B(H)$  by restricting the natural normal extension of  $\pi$  to  $C^{**}$ . In particular, we can use the formula

$$
\pi(c)\pi(q)\pi(d) = \pi(cqd), \qquad c, d \in C, q \in QM(C).
$$

Given  $C^*$ -algebras  $C, D$  and a ccp map  $\phi : C \to D$ , the multiplicative domain of  $\phi$  [\[BO08,](#page-46-21) Proposition 1.5.7] is the C ∗ -subalgebra

$$
C_{\phi} := \{ x \in C : \phi(x^*x) = \phi(x)^* \phi(x) \text{ and } \phi(xx^*) = \phi(x)\phi(x)^* \}.
$$

<span id="page-5-1"></span>We will frequently use that if  $x \in C_{\phi}$  and  $y \in C$ , then  $\phi(xy) = \phi(x)\phi(y)$  and  $\phi(yx) = \phi(y)\phi(x)$ .

1. Locally compact quantum groups and their actions

We recall some basic results from the theory of locally compact quantum groups [\[KV00,](#page-47-3) [KV03,](#page-47-4) [VV03\]](#page-47-20), and their representations and actions [\[Kus01,](#page-47-21) [Vae01\]](#page-47-6). The following is [\[KV03,](#page-47-4) Definition 1.1].

<span id="page-5-0"></span>**Definition 1.1.** A locally compact quantum group  $\mathbb{G}$  consists of a Hopf-von Neumann algebra  $\mathbb{G} = (M, \Delta)$ possessing nsf weights  $\Phi, \Psi : M_+ \to [0, +\infty]$  satisfying the left and the right invariance conditions

$$
\Phi((\omega \otimes id)\Delta(x)) = \Phi(x)\omega(1), \quad \Psi((id \otimes \omega)\Delta(y)) = \Psi(y)\omega(1), \quad \omega \in M_*, \quad x \in \mathscr{M}_{\Phi}^+, \quad y \in \mathscr{M}_{\Psi}^+.
$$
 (1.1)

These weights are then necessarily unique up to multiplication by a positive scalar.

Whenever convenient, we write  $M = L^{\infty}(\mathbb{G}), M_*= L^1(\mathbb{G})$  and  $L^2(M) = L^2(\mathbb{G})$ . We identify  $M \subseteq B(L^2(M))$ by the standard representation, which we write explicitly as  $\pi_M$  if we want to emphasize it. We note that a locally compact quantum group automatically satisfies the density conditions

<span id="page-5-2"></span>
$$
[(M \otimes 1)\Delta(M)]^{\sigma\text{-weak}} = [\Delta(M)(1 \otimes M)]^{\sigma-\text{weak}} = M\bar{\otimes}M. \tag{1.2}
$$

We write the induced *convolution product* on  $M_*$  as

$$
\omega * \omega' := (\omega \otimes \omega')\Delta, \qquad \omega, \omega' \in M_*.
$$

Fundamental to the theory of locally compact quantum groups are the unitaries

$$
V \in B(L^2(M)) \bar{\otimes} M, \qquad W \in M \bar{\otimes} B(L^2(M)),
$$

called respectively right and left regular unitary representation. They are uniquely characterized by the identities

$$
(\mathrm{id}\otimes\omega)(V)\Lambda_{\Psi}(x) = \Lambda_{\Psi}((\mathrm{id}\otimes\omega)\Delta(x)), \qquad \omega \in M_*, x \in \mathcal{N}_{\Psi},
$$

$$
(\omega \otimes id)(W^*) \Lambda_{\Phi}(x) = \Lambda_{\Phi}((\omega \otimes id)\Delta(x)), \qquad \omega \in M_*, x \in \mathcal{N}_{\Phi}.
$$

They are *multiplicative unitaries* [\[BS93\]](#page-46-15) in the sense that as operators on  $L^2(M) \otimes L^2(M)$ , they satisfy

$$
V_{12}V_{13}V_{23} = V_{23}V_{12}, \qquad W_{12}W_{13}W_{23} = W_{23}W_{12}.
$$

These multiplicative unitaries implement the coproduct of M:

<span id="page-5-3"></span>
$$
W^*(1 \otimes x)W = \Delta(x) = V(x \otimes 1)V^*, \qquad x \in M. \tag{1.3}
$$

Moreover, we have the equality

$$
[(\omega \otimes id)V \mid \omega \in B(L^{2}(M))_{*}] = [(\mathrm{id} \otimes \omega)W \mid \omega \in B(L^{2}(M))_{*}],
$$

and this defines a  $\sigma$ -weakly dense C<sup>∗</sup>-subalgebra of M that we will write as

$$
C_0^{\text{red}}(\mathbb{G}) \subseteq M.
$$

We recall some aspects of the duality theory of locally compact quantum groups [\[KV03,](#page-47-4) Section 2]. We write

$$
\hat{M} = [(\omega \otimes id)W \mid \omega \in M_*]^{\sigma\text{-weak}},
$$

which can be shown to be a von Neumann algebra. It becomes a locally compact quantum group for the coproduct

$$
\hat{\Delta}(x) = \Sigma W(x \otimes 1)W^* \Sigma, \qquad x \in \hat{M}.
$$

When useful for the sake of intuition or terminology, we write  $\hat{M} = L^{\infty}(\hat{\mathbb{G}})$ : it is the left group von Neumann algebra of  $\mathbb{G}$ , viewed as a function algebra on what is to be considered the Pontryagin dual  $\tilde{\mathbb{G}}$  of  $\mathbb{G}$ .

As we will need to compare left and right coactions later on, it will also be convenient to introduce independently the dual  $\check{M} = M^{\text{cop}\wedge}$  of  $M^{\text{cop}} = (M, \Delta^{\text{op}})$ , where  $\Delta^{\text{op}} = \text{Ad}\Sigma \circ \Delta$ . Noting that  $(M, \Delta^{\text{op}})$  has the left and right regular unitary representation equal to respectively  $V_{21}^*$  and  $W_{21}^*$ , we have

$$
\check{M} = [(\mathrm{id}\mathop{\otimes} \omega)V \mid \omega \in M_*]^{\sigma\text{-weak}}
$$

with coproduct

$$
\check{\Delta}(x) = V^*(1 \otimes x)V, \qquad x \in \check{M}.
$$

It can be shown that  $\check{M} = \hat{M}'$ . We then also occasionally write  $\check{M} = \mathscr{R}(\mathbb{G}) = L^{\infty}(\check{\mathbb{G}})$ , and view it as either the right group von Neumann algebra of  $\mathbb{G}$ , or a function algebra on an 'opposite' Pontryagin dual  $\check{\mathbb{G}}$  of  $\mathbb{G}$ . We further note that

<span id="page-6-2"></span>
$$
C_0^{\text{red}}(\hat{\mathbb{G}}) = [(\omega \otimes \text{id})W \mid \omega \in M_*], \qquad C_0^{\text{red}}(\check{\mathbb{G}}) = [(\text{id} \otimes \omega)V \mid \omega \in M_*]. \tag{1.4}
$$

There is a canonical identification of  $L^2(M)$  and  $L^2(\check{M})$ , in such a way that

$$
\check{W}=V
$$

is the left regular representation of  $\check{G}$ . With  $\check{J} = J_{\check{M}}$  the resulting associated modular conjugation for  $\check{M}$  on  $L^2(M)$ , we have

 $\check{J}M\check{J}=M.$ 

One calls the induced anti-multiplicative involution

$$
R: M \to M, \quad x \mapsto \check{J}x^*\check{J}
$$

the unitary antipode. It turns  $(M, \Delta)$  into an involutive Hopf-von Neumann algebra. The following identities involving the modular conjugations will be useful:

<span id="page-6-0"></span>
$$
(\check{J}\otimes J)W(\check{J}\otimes J)=W^*,\qquad (J\otimes \check{J})V(J\otimes \check{J})=V^*.
$$
\n
$$
(1.5)
$$

We also recall from [\[KV03,](#page-47-4) Corollary 2.12] that one can pick a canonical unimodular number  $c \in \mathbb{C}$  such that

$$
c\check{J}J = \overline{c}J\check{J}.
$$

The resulting self-adjoint unitary  $u_{\mathbb{G}} := cJJ$  is called the *fundamental symmetry*. This unitary allows us to relate  $W$  and  $V$  directly, for we have

<span id="page-6-1"></span>
$$
(u_{\mathbb{G}} \otimes 1)V(u_{\mathbb{G}} \otimes 1) = W_{21}.
$$
\n
$$
(1.6)
$$

It also allows us to express the right regular unitary of  $\tilde{M}$  as

$$
\check{V} = \widetilde{W} := (u_{\mathbb{G}} \otimes u_{\mathbb{G}})W(u_{\mathbb{G}} \otimes u_{\mathbb{G}}).
$$

We similarly write  $\widetilde{V} = (u_{\mathbb{G}} \otimes u_{\mathbb{G}}) V (u_{\mathbb{G}} \otimes u_{\mathbb{G}})$ . The operation  $M \mapsto \widetilde{M}$  is not on the nose involutive, but we have that

$$
M^{\vee\vee} = u_{\mathbb{G}} M u_{\mathbb{G}}, \qquad \Delta^{\vee\vee} = \mathrm{Ad}(u_{\mathbb{G}} \otimes u_{\mathbb{G}}) \circ \Delta \circ \mathrm{Ad}(u_{\mathbb{G}}).
$$

Recall from Definition [0.2](#page-1-0) the notion of a unitary representation of  $\mathbb{G}$  on a Hilbert space H. If  $\omega \in M_*$ , we write

$$
U(\omega) = (\mathrm{id} \otimes \omega)U \in B(\mathcal{H}).
$$

Clearly

$$
U(\omega * \omega') = U(\omega)U(\omega'), \qquad \omega, \omega' \in M_*.
$$

<span id="page-7-3"></span>For later use, we recall the following facts, see [\[Wor96\]](#page-47-22), [\[Kus01,](#page-47-21) Theorem 1.6] and [\[BDS13,](#page-46-22) Theorem 4.12].

**Proposition 1.2.** If U is a unitary G-representation on a Hilbert space  $H$ , then the norm-closure

$$
C_U = [U(\omega) \mid \omega \in M_*]
$$

is a C<sup>∗</sup> -algebra acting non-degenerately on H, and

$$
U \in M(C_U \otimes C_0^{\text{red}}(\mathbb{G})) \subseteq M(\mathcal{K}(\mathcal{H}) \otimes C_0^{\text{red}}(\mathbb{G})) \cap M(C_U \otimes \mathcal{K}(L^2(\mathbb{G}))).
$$

The following results from [\[Vae01\]](#page-47-6) will be crucial. Assume that A is a G-W<sup>\*</sup>-algebra. The crossed product von Neumann algebra is defined as

$$
A \rtimes \mathbb{G} = \{ \alpha(a)(1 \otimes x) \mid a \in A, x \in \check{M} \}'' \subseteq A \bar{\otimes} B(L^2(\mathbb{G})),
$$

and in fact already

$$
A \rtimes \mathbb{G} = [\alpha(a)(1 \otimes x) \mid a \in A, x \in \check{M}]^{\sigma\text{-weak}}.
$$

In the following, we write

$$
A^\rtimes:=A\rtimes\mathbb{G}
$$

as a shorthand. We note that  $A^{\rtimes}$  carries the dual right  $\check{\mathbb{G}}$ -action

$$
\alpha^{\rtimes}: A^{\rtimes} \to A^{\rtimes} \bar{\otimes} \check{M}, \quad \alpha^{\rtimes}(z) = \widetilde{W}_{23} z_{12} \widetilde{W}_{23}^*, \qquad z \in A^{\rtimes}.
$$

In particular, one has

$$
\alpha^{\rtimes}(\alpha(a)) = \alpha(a) \otimes 1, \quad \alpha^{\rtimes}(1 \otimes x) = 1 \otimes \check{\Delta}(x), \qquad a \in A, x \in \check{M}.
$$

<span id="page-7-1"></span>**Theorem 1.3.** Let  $A$  be a  $\mathbb{G}-W^*$ -algebra. Then there is a canonical identification

$$
L^2(A^{\rtimes}) \cong L^2(A) \otimes L^2(\mathbb{G}),
$$

in such a way that

<span id="page-7-0"></span>
$$
U_{\alpha} = J_{A^{\rtimes}}(J_A \otimes \check{J}) \in B(L^2(A) \otimes L^2(\mathbb{G})) \tag{1.7}
$$

is a unitary G-representation implementing  $\alpha$  (in the sense of  $(0.2)$ ). Moreover, the associated GNSrepresentation  $\pi_{A^{\lambda}}$  of  $A^{\lambda}$  becomes the identity representation of  $A^{\lambda}$  on  $L^2(A) \otimes L^2(\mathbb{G})$ , while

$$
\rho_{A^{\rtimes}}(\alpha(a)) = J_{A^{\rtimes}} \pi_{A^{\rtimes}}(a)^* J_{A^{\rtimes}} = \rho_A(a) \otimes 1, \qquad a \in A,
$$
  

$$
\rho_{A^{\rtimes}}(1 \otimes x) = J_{A^{\rtimes}}(1 \otimes \pi_{\check{M}}(x)^*) J_{A^{\rtimes}} = U_{\alpha}(1 \otimes \rho_{\check{M}}(x)) U_{\alpha}^*, \qquad x \in \check{M}.
$$

The dual  $\mathfrak{S}\text{-action}$  has  $\widetilde{W}_{23}$  as its canonical unitary implementation.

Note that we make here the switch from the left coactions of [\[Vae01\]](#page-47-6) to right coactions by replacing  $(M, \Delta)$ with  $(M, \Delta^{\rm op})$ .

Proof. See [\[Vae01,](#page-47-6) Lemma 3.3, Definition 3.4, Definition 3.6 and Proposition 3.7]. The last claim follows from [\[Vae01,](#page-47-6) Proposition 4.3].

Remark 1.4. From  $(1.7)$ , we obtain the identity

<span id="page-7-2"></span>
$$
(J_A \otimes \check{J})U_{\alpha}(J_A \otimes \check{J}) = U_{\alpha}^*.
$$
\n(1.8)

**Remark 1.5.** When  $A = M$  with  $\alpha = \Delta$ , we have  $U_{\Delta} = V$ .

We end this section with a concrete computation of a canonical implementing unitary.

<span id="page-8-2"></span>**Example 1.6.** Let A be a  $\mathbb{G}\text{-}W^*$ -algebra. We can endow  $A^{\rtimes} = A \rtimes \mathbb{G}$  with the adjoint  $\mathbb{G}\text{-action}$ 

<span id="page-8-4"></span>
$$
\alpha_{\text{ad}}: A^{\rtimes} \to A^{\rtimes} \bar{\otimes} M, \quad z \mapsto V_{23} z_{12} V_{23}^*, \qquad z \in A^{\rtimes}, \tag{1.9}
$$

so

$$
\alpha_{\text{ad}}(\alpha(a)) = (\text{id} \otimes \Delta)\alpha(a) = (\alpha \otimes \text{id})\alpha(a), \qquad \alpha_{\text{ad}}(1 \otimes x) = 1 \otimes V(x \otimes 1)V^*, \qquad a \in A, x \in \check{M}.
$$

We claim that the canonical unitary implementation  $U_{ad}$  of  $\alpha_{ad}$  is given by

<span id="page-8-1"></span>
$$
U_{\rm ad} = V_{23} W_{32} U_{\alpha,13}.
$$
\n(1.10)

Indeed, note first that the crossed product von Neumann algebra  $A^{\rtimes} \rtimes \mathbb{G}$  is given by

$$
A^{\rtimes} \rtimes \mathbb{G} = [\alpha_{ad}(\alpha(a)(1 \otimes x))(1 \otimes 1 \otimes y) \mid a \in A, x, y \in \check{M}]^{\sigma-\text{weak}}
$$
  
\n
$$
= [V_{23}\alpha(a)_{12}x_2V_{23}^*y_3 \mid a \in A, x, y \in \check{M}]^{\sigma-\text{weak}}
$$
  
\n
$$
= [V_{23}\alpha(a)_{12}x_2\check{\Delta}(y)_{23}V_{23}^* \mid a \in A, x, y \in \check{M}]^{\sigma-\text{weak}}
$$
  
\n
$$
= [V_{23}\alpha(a)_{12}(1 \otimes x \otimes y)V_{23}^* \mid a \in A, x, y \in \check{M}]^{\sigma-\text{weak}}
$$
  
\n
$$
= V_{23}(A^{\rtimes} \tilde{\otimes} \check{M})V_{23}^*,
$$

Since the dual coaction of M on  $A^{\rtimes} \rtimes \mathbb{G}$  is given by conjugation with  $\widetilde{W}_{34}$ , we see that  $A^{\rtimes} \rtimes \mathbb{G}$  is just a conjugated copy of  $A^{\Join} \bar{\otimes} \check{M}$  with  $id_{A^{\Join}} \otimes \check{\Delta}$  as the coaction. In particular, under the canonical identification

$$
L^{2}(A^{\rtimes} \rtimes \mathbb{G}) = L^{2}(A^{\rtimes}) \otimes L^{2}(\mathbb{G}) = L^{2}(A) \otimes L^{2}(\mathbb{G}) \otimes L^{2}(\mathbb{G}),
$$

the associated modular conjugation of  $A^{\rtimes} \rtimes \mathbb{G}$  is given by

$$
J_{A^\rtimes \rtimes \mathbb G} = V_{23}(J_{A^\rtimes} \otimes \check J)V_{23}^*,
$$

and so the canonical unitary implementation of  $\alpha_{ad}$  is by Theorem [1.3](#page-7-1) given by

$$
U_{\alpha_{\rm ad}} = J_{A^{\times} \rtimes \mathbb{G}}(J_{A^{\times}} \otimes \check{J})
$$
  
\n
$$
= V_{23}(J_{A^{\times}} \otimes \check{J})V_{23}^{*}(J_{A^{\times}} \otimes \check{J})
$$
  
\n
$$
= V_{23}U_{\alpha,12}(J_{A} \otimes \check{J} \otimes \check{J})V_{23}^{*}(J_{A} \otimes \check{J} \otimes \check{J})U_{\alpha,12}^{*}
$$
  
\n
$$
= V_{23}U_{\alpha,12}W_{32}U_{\alpha,12}^{*}
$$
  
\n
$$
= V_{23}W_{32}U_{\alpha,13},
$$
  
\n(0.1)

<span id="page-8-3"></span>proving  $(1.10)$ .

# 2. Dynamical W<sup>∗</sup> -correspondences

2.1. Definition and Examples. Let G be a locally compact quantum group, and let  $(A, \alpha), (B, \beta)$  be G-dynamical von Neumann algebras. In Definition [0.4,](#page-1-3) we introduced the notion of a G-equivariant A-B-correspondence  $\mathcal{H} = (\mathcal{H}, U, \pi, \rho)$ . Note that if G is the trivial group, we find back the usual notion of an A-B-correspondence. On the other hand, when  $A = B = \mathbb{C}$ , we find back the notion of a unitary G-representation.

**Definition 2.1.** We denote by Corr<sup>G</sup> $(A, B)$  the category of G-A-B-correspondences, with morphisms from H to H' consisting of bounded operators  $T: \mathcal{H} \to \mathcal{H}'$  intertwining the (anti-)representations of A, B and  $M_*$ .

Clearly,  $Corr^{\mathbb{G}}(A, B)$  is a W<sup>\*</sup>-category [\[GLR85\]](#page-47-23), and we will always understand it as such.

<span id="page-8-0"></span>Here are some examples of G-equivariant correspondences.

Example 2.2. Let A be a G-dynamical von Neumann algebra. The trivial or identity G-A-A-correspondence  $E_A^{\mathbb{G}}$  is  $L^2(A)$  with its canonical representation  $\pi_A$  and anti-representation  $\rho_A$ , and the canonical unitary Grepresentation  $U_{\alpha}$  implementing  $\alpha$ .

The compatibility relations [\(0.3\)](#page-2-0) are satisfied by Theorem [1.3](#page-7-1) and the identity [\(1.8\)](#page-7-2), together with the fact that  $J$  implements the unitary antipode.

Let  $(A, \alpha)$  and  $(B, \beta)$  be G-dynamical von Neumann algebras. We say that  $A \subseteq B$  is an *equivariant inclusion* if  $\beta_{|A} = \alpha$ .

**Example 2.3.** Let  $A \subseteq C$  and  $B \subseteq D$  be G-equivariant inclusions. Then for any G-C-D correspondence H we obtain the restriction

$$
{}_A\text{Res}_B(\mathcal{H})={}_A\mathcal{H}_B.
$$

In particular, if  $A \subseteq B$  is an equivariant inclusion, we have equivariant correspondences  $_A L^2(B)_B$ ,  $_B L^2(B)_A$ and  $_A L^2(B)_A$ , each with  $U_\beta$  as unitary representation for  $\beta$  the given coaction on B.

<span id="page-9-0"></span>**Example 2.4.** Let A, B be G-dynamical von Neumann algebras, and let  $(\mathcal{H}, U, \pi, \rho)$  be a G-equivariant correspondence. Consider  $C := \rho(B)'$ . Then  $U(C \otimes 1)U^* \subseteq C \bar{\otimes} M$  by the bicommutant theorem, so that

$$
\gamma: C \to C \bar{\otimes} M, \qquad c \mapsto U(c \otimes 1)U^*
$$

is a coaction, and  $(c\mathcal{H}_B, U)$  is a G-equivariant C-B-correspondence. Moreover,  $\pi(A) \subseteq C$  is a G-equivariant inclusion.

As a particular case, let  $\pi : A \to B$  be a normal unital G-equivariant \*-homomorphism. Consider C :=  $\rho_B(\pi(A))'$ , so that  $\pi_B(\pi(A)) \subseteq \pi_B(B) \subseteq C$  is (by definition) the Jones basic construction. Then

$$
\gamma: C \to C \bar{\otimes} M, \qquad c \mapsto U_{\beta}(c \otimes 1)U_{\beta}^*
$$

is a well-defined coaction, and  $B \cong \pi_B(B) \subseteq C$  becomes a G-equivariant inclusion of von Neumann algebras.

**Example 2.5.** The coarse (or regular)  $\mathbb{G}$ -A-B-correspondence  $C_{A,B}^{\mathbb{G}}$  is  $L^2(A) \otimes L^2(\mathbb{G}) \otimes L^2(B)$  endowed with

$$
\pi(a) = (\pi_A \otimes \pi_M)(\alpha(a))_{12}, \qquad \rho(b) = (\rho_M \otimes \rho_B)(\beta^{\text{op}}(b))_{23},
$$

$$
U = V_{24} \in B(L^2(A) \otimes L^2(\mathbb{G}) \otimes L^2(B) \otimes L^2(\mathbb{G})).
$$

Here, we wrote  $\beta^{\text{op}}(b) = \varsigma \beta(b) \in M \bar{\otimes} B$  where  $\varsigma : B \bar{\otimes} M \cong M \bar{\otimes} B$  is the flip. The necessary compatibilities are satisfied by using the coaction property together with the identities [\(1.3\)](#page-5-3) and [\(1.5\)](#page-6-0). The same reasoning works for the next examples.

<span id="page-9-1"></span>**Example 2.6.** Let  $(\mathcal{H}, U, \pi, \rho)$  be a G-A-B-correspondence. We define the regular amplification to be the  $\mathbb{G}\text{-}A\text{-}B\text{-}correspondence S^{\mathbb{G}}(\mathcal{H}) = \mathcal{H}\otimes L^2(\mathbb{G})$  endowed with

$$
\pi_S(a) = (\pi \otimes \pi_M)(\alpha(a)), \qquad \rho_S(b) = (\rho \otimes \rho_M)(\beta(b)),
$$
  

$$
U_S = V_{23} \in B(\mathcal{H} \otimes L^2(\mathbb{G}) \otimes L^2(\mathbb{G})).
$$

It is a restriction of  $_{A\bar{\otimes}M}(\mathcal{H}\otimes L^2(\mathbb{G}))_{B\bar{\otimes}M}$  where  $A\bar{\otimes}M, B\bar{\otimes}M$  are endowed with the coaction id  $\otimes\Delta$ , and where we view  $L^2(\mathbb{G}) = E_M$ .

In particular, we call  $S_A^{\mathbb{G}} := S^{\mathbb{G}}(E_A)$  the G-semi-coarse (or G-semi-regular) G-A-A-correspondence.

Note that  $S_A^{\mathbb{G}}$  becomes the trivial A-A-correspondence if  $\mathbb{G}$  is trivial, and the regular  $\mathbb{G}$ -representation if A is trivial. The following example reverses this situation.

**Example 2.7.** The W<sup>\*</sup>-semi-coarse (or W<sup>\*</sup>-semi-regular) G-A-B-correspondence  $S_{A,B}^{\mathbb{G}}$  is  $L^2(A) \otimes L^2(B)$ endowed with

$$
\pi(a) = \pi_A(a) \otimes 1, \qquad \rho(b) = 1 \otimes \rho_B(b),
$$
  
 
$$
U = U_{\alpha,13}U_{\beta,23} \in B(L^2(A) \otimes L^2(B) \otimes L^2(\mathbb{G})).
$$

**Remark 2.8.** We can make the coarse G-A-B-correspondence  $C_{A,B}^G$  resemble more the W<sup>\*</sup>-semi-coarse  $\mathbb{G}\text{-}A\text{-}B\text{-}correspondence S^\mathbb{G}_{A,B}$  as follows: write

$$
\widetilde{U}_B = (u_{\mathbb{G}} \otimes 1)U^*_{\beta,21}(u_{\mathbb{G}} \otimes 1) \in \rho_M(M)\bar{\otimes}B(L^2(B)),\tag{2.1}
$$

then from the identities  $(1.5)$  and  $(1.6)$  we find that

$$
\widetilde{U}_{B,12}^* V_{13} \widetilde{U}_{B,12} = V_{13} U_{\beta,23}.
$$

So, conjugating the coarse G-A-B-correspondence with the unitary  $\widetilde{U}^*_{B,23}U^*_{\alpha,12}$ , we find that  $C^{\mathbb{G}}_{A,B}$  is equivalent to the G-A-B-correspondence on  $L^2(A) \otimes L^2(\mathbb{G}) \otimes L^2(B)$  given by

$$
\pi(a) = \pi_A(a) \otimes 1 \otimes 1, \qquad \rho(b) = 1 \otimes 1 \otimes \rho_B(b),
$$

$$
U = U_{\alpha,14} V_{24} U_{\beta,34}.
$$

<span id="page-10-2"></span>As a final example, we consider the following.

Example 2.9. Let  $(\mathcal{H}, U, \pi, \rho)$  be a G-A-B-correspondence. The adjoint amplification is the G-A-Bcorrespondence  $D^{\mathbb{G}}(\mathcal{H}) = \mathcal{H} \otimes L^2(\mathbb{G})$  endowed with

$$
\pi_D(a) = (\pi \otimes \pi_M)(\alpha(a)), \qquad \rho_D(b) = \rho(b) \otimes 1,
$$

$$
U_D = V_{23} W_{32} U_{13}.
$$

Note that  $U_D$  is indeed a unitary G-representation: it is the tensor product of  $U_{ad}$  with U, where  $U_{ad} = W_{21}V$ is the adjoint representation of  $\mathbb G$  on  $L^2(\mathbb G)$ . The compatibility of  $U_D$  and  $\pi_D$  follows from the fact that U implements  $\alpha$ , together with the easily computed commutation relation

<span id="page-10-1"></span>
$$
V_{23}W_{32}U_{13}U_{12} = U_{12}U_{13}V_{23}W_{32}.
$$
\n(2.2)

The compatibility of  $U_D$  and  $\rho_D$  is immediate.

In particular, we call  $D_A^{\mathbb{G}} := D^{\mathbb{G}}(E_A)$  the adjoint  $\mathbb{G}\text{-}A$ -A-correspondence. Using Example [1.6,](#page-8-2) we see that  $D_A^G$  is simply the two-sided restriction to A of the identity correspondence  $E_{A\rtimes G}$  for  $A\rtimes G$  with the adjoint G-action.

There are some obvious constructions one can perform on G-equivariant correspondences, such as

- taking (infinite) direct sums, and
- taking G-equivariant subcorrespondences.

Let us make the latter more precise: if  $(H, U, \pi, \rho)$  is a G-A-B-correspondence, we define  $\mathcal{G} \subseteq \mathcal{H}$  to be a G-equivariant subcorrespondence if G is stable under  $\pi(A)$ ,  $\rho(B)$  and all  $U(\omega)$  for  $\omega \in M_*$ . It follows from Proposition [1.2](#page-7-3) that G becomes a G-A-B-correspondence  $(G, U')$  in a unique way such that  $U'(\omega)$  is the restriction of  $U(\omega)$  to G. Moreover, this description also makes it clear that  $\mathcal{G}^{\perp}$  will be a G-equivariant subcorrespondence, with  $\mathcal{H} = \mathcal{G} \oplus \mathcal{G}^{\perp}$  as  $\mathbb{G}$ -A-B-correspondences.

**Definition 2.10.** We call a G-A-B-correspondence  $(\mathcal{H}, U, \pi, \rho)$  cyclic if there exists  $\xi \in \mathcal{H}$  such that

$$
\mathcal{H} = [\pi(a)U(\omega)\rho(b)\xi \mid a \in A, \omega \in M_*, b \in B].
$$

<span id="page-10-0"></span>**Proposition 2.11.** Any G-A-B-correspondence  $(\mathcal{H}, U, \pi, \rho)$  is a direct sum of cyclic G-A-B-correspondences.

*Proof.* Take  $\xi \in \mathcal{H}$ . It is sufficient to show that

$$
\mathcal{G} := [\pi(a)U(\omega)\rho(b)\xi \mid a \in A, \omega \in M_*, b \in B]
$$

is a G-equivariant subcorrespondence: since the C<sup>∗</sup>-algebra  $C_U = [\{U(\omega) | \omega \in M_*\}]$  acts non-degenerately on H, we have  $\xi \in \mathcal{G}$ , so G will indeed be cyclic, and an appeal to Zorn's lemma then lets us conclude.

Clearly G is invariant under  $\pi(A)$ . If  $e_i$  is an orthonormal basis of  $L^2(\mathbb{G})$ , then for  $\xi, \eta \in L^2(\mathbb{G})$  and  $\omega = \omega_{\xi,\eta} \in B(L^2(\mathbb{G}))_*$ , we have

$$
U(\omega)\pi(a) = (\mathrm{id}\otimes\omega)((\pi\otimes\mathrm{id})\alpha(a)U) = \sum_i \pi((\mathrm{id}\otimes\omega_{\xi,e_i})(\alpha(a)))U(\omega_{e_i,\eta}),
$$

the latter sum converging e.g.  $\sigma$ -strongly. Hence  $\mathcal G$  is invariant under the  $U(\omega)$ .

Similarly,

$$
\rho(b)U(\omega)=(\mathrm{id}\otimes\omega)(U(\rho\otimes R)(\beta(b)))=\sum_i U(\omega_{\xi,e_i})\rho((\mathrm{id}\otimes\omega_{e_i,\eta}R)(\beta(b))),
$$

making it clear that G is invariant under  $\rho(B)$  as well.

It follows easily from the above that we can make a *universal* G-A-B-correspondence  $(\mathcal{H}_u, U_u, \pi_u, \rho_u)$ , in the sense that any cyclic G-A-B-correspondence is unitarily equivalent to a direct summand of  $\mathcal{H}_u$ . In the following, we fix such a chosen universal model.

# <span id="page-11-4"></span>2.2. Associated C<sup>\*</sup>-algebras.

Definition 2.12. Let  $A$  be a  $\mathbb{G}\text{-}W^*$ -algebra. We define

$$
A_{s} = [(\mathrm{id} \otimes \omega)\alpha(a) \mid \omega \in M_{*}, a \in A] \subseteq A.
$$

It is not clear if  $A_s$  will always be a C\*-subalgebra of  $A$ , although it is easily seen to be an operator subsystem.<sup>[1](#page-11-0)</sup> In any case, we will not need any multiplicative structure of  $A_s$ . It will however be useful to observe that  $A_s$  is  $\sigma$ -weakly dense in A (see e.g. [\[KaS15,](#page-47-24) Corollary 2.7] for an explicit proof).

<span id="page-11-3"></span>The following observation will be fundamental.

**Theorem 2.13.** If  $(H, U, \pi, \rho)$  is a G-A-B-correspondence, then

<span id="page-11-2"></span>
$$
C_{\mathcal{H}}^{\mathbb{G}}(A) := [C_U \pi(A) C_U], \qquad C_{\mathcal{H}}^{\mathbb{G}}(B) := [C_U \rho(B) C_U], \qquad C_{\mathcal{H}}^{\mathbb{G}}(A, B) := [C_{\mathcal{H}}^{\mathbb{G}}(A) C_{\mathcal{H}}^{\mathbb{G}}(B)] \tag{2.3}
$$

are C<sup>∗</sup> -algebras, and

<span id="page-11-1"></span>
$$
C_{\mathcal{H}}^{\mathbb{G}}(A) = [\pi(A_{s})C_{U}] = [C_{U}\pi(A_{s})], \quad C_{\mathcal{H}}^{\mathbb{G}}(B) = [\rho(B_{s})C_{U}] = [C_{U}\rho(B_{s})].
$$
\n(2.4)

*Proof.* Clearly  $C_{\mathcal{H}}^{\mathbb{G}}(A)$  and  $C_{\mathcal{H}}^{\mathbb{G}}(B)$  are closed under the  $*$ -operation.

We now first prove  $(2.4)$ . Observe that by  $(1.4)$ , we have

$$
[(\mathrm{id} \otimes \omega)(V) \mid \omega \in B(L^2(\mathbb{G}))_*] = [(\mathrm{id} \otimes \omega)(V^*) \mid \omega \in B(L^2(\mathbb{G}))_*].
$$

We then compute (dropping the notation  $\pi$ )

$$
C_{\mathcal{H}}^{\mathbb{G}}(A) = [(\mathrm{id}\otimes\omega)(U_{12}a_1U_{13}) \mid a \in A, \omega \in B(L^2(\mathbb{G})\otimes L^2(\mathbb{G}))_*]
$$
  
\n
$$
= [(\mathrm{id}\otimes\omega)(\alpha(a)_{12}U_{12}U_{13}) \mid a \in A, \omega \in B(L^2(\mathbb{G})\otimes L^2(\mathbb{G}))_*]
$$
  
\n
$$
= [(\mathrm{id}\otimes\omega)(\alpha(a)_{12}(\mathrm{id}\otimes\Delta)(U)) \mid a \in A, \omega \in B(L^2(\mathbb{G})\otimes L^2(\mathbb{G}))_*]
$$
  
\n
$$
= [(\mathrm{id}\otimes\omega)(\alpha(a)_{12}V_{23}U_{12}V_{23}^*) \mid a \in A, \omega \in B(L^2(\mathbb{G})\otimes L^2(\mathbb{G}))_*]
$$
  
\n
$$
= [(\mathrm{id}\otimes\omega)(\alpha(a)_{12}V_{23}U_{12}) \mid a \in A, \omega \in B(L^2(\mathbb{G})\otimes L^2(\mathbb{G}))_*]
$$
  
\n
$$
= [(\mathrm{id}\otimes\omega)(\alpha(a)_{12}V_{23}U_{12}) \mid a \in A, \omega \in B(L^2(\mathbb{G})\otimes L^2(\mathbb{G}))_*]
$$
  
\n
$$
= [(\mathrm{id}\otimes\omega)(V_{23}^*(\mathrm{id}\otimes\Delta)\alpha(a)U_{12}) \mid a \in A, \omega \in B(L^2(\mathbb{G})\otimes L^2(\mathbb{G}))_*]
$$
  
\n
$$
= [(\mathrm{id}\otimes\omega)((\alpha\otimes\mathrm{id})\alpha(a)U_{12}) \mid a \in A, \omega \in B(L^2(\mathbb{G})\otimes L^2(\mathbb{G}))_*]
$$
  
\n
$$
= [(\mathrm{id}\otimes\omega)(U_{12}\alpha(a)_{13}) \mid a \in A, \omega \in B(L^2(\mathbb{G})\otimes L^2(\mathbb{G}))_*]
$$
  
\n
$$
= [C_U A_s].
$$

<span id="page-11-0"></span><sup>&</sup>lt;sup>1</sup>In case G is semi-regular, the proof of [\[BSV03,](#page-46-23) Proposition 5.7] shows that  $A_s$  will be a C<sup>∗</sup>-algebra, but it is not clear if the converse in that Proposition holds in the von Neumann algebraic setting.

Since  $C_{\mathcal{H}}^{\mathbb{G}}(A)$  is closed under the \*-operation, we obtain [\(2.4\)](#page-11-1) for A. It is now clear that  $C_{\mathcal{H}}^{\mathbb{G}}(A)$  is a C<sup>\*</sup>-algebra:  $[C_{\mathcal{H}}^{\mathbb{G}}(A)C_{\mathcal{H}}^{\mathbb{G}}(A)]=[C_UA_{\mathrm{s}}^2C_U]\subseteq C_{\mathcal{H}}^{\mathbb{G}}(A).$ 

For B, we can simply observe that  $B^{\rm op}$  has a right coaction by  $(M, \Delta^{\rm op})$  via

$$
b^{\rm op} \mapsto \beta(b)^{\rm op \otimes R},
$$

and that  $U^*$  is a unitary representation for  $(M, \Delta^{\text{op}})$  with

$$
U^*(\rho(b)\otimes 1)U=(\rho\otimes R)\beta(b),\qquad b\in B.
$$

Hence the result for B follows from that for A (since  $C_U = C_{U^*}$ ).

Finally,  $C_{\mathcal{H}}^{\mathbb{G}}(A, B)$  is closed under the \*-structure since

$$
C_{\mathcal{H}}^{\mathbb{G}}(A,B)^{*} = [C_{U}\rho(B_{s})\pi(A_{s})C_{U}] = [C_{U}\pi(A_{s})\rho(B_{s})C_{U}] = [C_{U}\pi(A)C_{U}C_{U}\rho(B)C_{U}] = C_{\mathcal{H}}^{\mathbb{G}}(A,B).
$$

A similar computation shows that  $C_{\mathcal{H}}^{\mathbb{G}}(A, B)$  is closed under multiplication.

<span id="page-12-1"></span>Remark 2.14. It also follows immediately from [\(2.3\)](#page-11-2) and [\(2.4\)](#page-11-1) that for example

$$
C^{\mathbb{G}}_{\mathcal{H}}(A) = [C_U \pi(A_{\mathrm{s}}) C_U], \qquad C^{\mathbb{G}}_{\mathcal{H}}(A, B) = [\pi(A_{\mathrm{s}}) C_U \rho(B_{\mathrm{s}})],
$$

and that

$$
\pi(A)\subseteq QM(C^{\mathbb{G}}_{\mathcal{H}}(A,B)),\qquad \rho(B)\subseteq QM(C^{\mathbb{G}}_{\mathcal{H}}(A,B)).
$$

This appearance of the space of quasi-multipliers is somewhat surprising and mysterious, but seems unavoidable as the following remark shows.

<span id="page-12-0"></span>**Remark 2.15.** In general, we will not have that  $\pi(A) \subseteq M(C_{\mathcal{H}}^{\mathbb{G}}(A,B))$ . A concrete example is obtained as follows: let  $A = B(\ell^2(\mathbb{Z} \times \mathbb{Z}))$ , let  $B = \mathbb{C}$  and let  $\mathbb{G} = \mathbb{T} = \{z \in \mathbb{C} \mid |z| = 1\}$ . Let  $e_{m,n}$  be the standard basis of  $\mathcal{H} = \ell^2(\mathbb{Z} \times \mathbb{Z})$ , and let  $\mathbb{T}$  act on  $\mathcal{H}$  via

$$
U(z)e_{n,m} = z^n e_{n,m}, \qquad z \in \mathbb{T}, m, n \in \mathbb{Z}.
$$

We also view U as a unitary in  $M(\mathcal{K}(\mathcal{H})\otimes C(\mathbb{T}))\subseteq B(\mathcal{H})\bar{\otimes}L^{\infty}(\mathbb{T})$  through

$$
(\mathrm{id}\otimes \mathrm{ev}_z)(U) = U(z), \quad z \in \mathbb{T}.
$$

 $Endow B$  with the trivial action, and endow A with the inner  $\mathbb{T}\text{-action}$ 

$$
\alpha: A \to A \bar{\otimes} L^{\infty}(\mathbb{T}): x \mapsto U(x \otimes 1)U^*.
$$

This equips H with the structure of a  $\mathbb{T}$ -equivariant A-B-correspondence, and  $C_{\mathcal{H}}^{\mathbb{T}}(A, B) = C_{\mathcal{H}}^{\mathbb{T}}(A)$ .

We write the irreducible unitary representations of  $\mathbb T$  as

$$
\chi_n: \mathbb{T} \to \mathbb{C}: z \mapsto z^n, \quad n \in \mathbb{Z}.
$$

Consider now the operator  $x \in A$  determined by

$$
xe_{n,m} = \delta_{n,0}e_{m,0}.
$$

We claim that  $x \notin M(C^{\mathbb{T}}_{\mathcal{H}}(A))$ . Indeed, let  $p_n$  be the projection onto  $\ell^2(\{n\} \times \mathbb{Z})$  for  $n \in \mathbb{Z}$ . Viewing  $\chi_n \in L^1(\mathbb{T}) = L^{\infty}(\mathbb{T})_*$ , we see that  $(id \otimes \chi_n)(U) = p_{-n}$  so that  $C_U = [p_n : n \in \mathbb{Z}]$  by the Stone-Weierstrass theorem. It is then sufficient to prove that  $x = xp_0 \notin C^{\mathbb{T}}_{\mathcal{H}}(A) = [A_s C_U]$ , or equivalently that  $x \notin [A_s p_0]$ . However, one easily verifies that the spectral subspace of  $B(\mathcal{H})$  associated to the irreducible representation  $\chi_m$  is given by

$$
A_m = \{ y \in B(\mathcal{H}) \mid \forall n \in \mathbb{Z} : y p_n = p_{n+m} y p_n \}.
$$

Consequently

$$
A_s = [A_m \mid m \in \mathbb{Z}]
$$

It is elementary to see from this description that  $\lim_{n\to\infty} ||p_ny|| = 0$  for all  $y \in [A_s p_0]$ . Since  $||p_nx|| = 1$  for all  $n \in \mathbb{Z}$ , we find that indeed  $x \notin [A_s p_0]$ .

<span id="page-13-0"></span>**Theorem 2.16.** There is a well-defined semi-norm on  $A_s \odot M_s \odot B_s$  through

<span id="page-13-1"></span>
$$
\|\sum_{i=1}^n a_i \otimes \omega_i \otimes b_i\|_u = \sup\{\|\sum_{i=1}^n \pi(a_i)U(\omega_i)\rho(b_i)\| \mid (\mathcal{H}, U) \text{ a } \mathbb{G}\text{-}A\text{-}B\text{-}correspondence\}.
$$
 (2.5)

Moreover, the associated separation-completion  $C^{C}(A, B)$  has the unique structure of a  $C^{*}$ -algebra such that, for each  $\mathbb{G}-A-B$ -correspondence  $(\mathcal{H}, U, \pi, \rho)$ , the map

<span id="page-13-4"></span>
$$
\theta = \theta_{\mathcal{H}} : C^{\mathbb{G}}(A, B) \to B(\mathcal{H}), \quad a \otimes \omega \otimes b \mapsto \pi(a)U(\omega)\rho(b) \tag{2.6}
$$

is a well-defined non-degenerate ∗-representation.

*Proof.* Clearly  $\|\text{---}\|_u$  is well-defined as a semi-norm, with

$$
\|\sum_{i=1}^n a_i \otimes \omega_i \otimes b_i\|_u \leq \sum_{i=1}^n \|a_i\| \|\omega_i\| \|b_i\|.
$$

Now by Proposition [2.11,](#page-10-0) the supremum in  $(2.5)$  does not change if we consider only cyclic G-equivariant A-B-correspondences. Hence we see that  $\| - \|_u$  is in fact realized by the universal G-equivariant A-Bcorrespondence  $(\mathcal{H}_u, U_u, \pi_u, \rho_u)$ , i.e.  $C^{\mathbb{G}}(A, B) := C^{\mathbb{G}}_{\mathcal{H}_u}(A, B)$  satisfies the requirements.

In the following, we identify  $C^{\mathbb{G}}(A, B) = C^{\mathbb{G}}_{\mathcal{H}_u}(A, B)$  whenever convenient. We do have to be a bit careful when using this identification, as we want to make sure that constructions do not actually depend on the precise model chosen. Here is a case in point: By means of Remark [2.14,](#page-12-1) we have a natural map  $\pi_{\mathcal{H}_u}: A \to QM(C_{\mathcal{H}_u}^{\mathbb{G}}(A,B)),$  which through the isomorphism

$$
\theta_{\mathcal{H}_u}: C^{\mathbb{G}}(A, B) \cong C^{\mathbb{G}}_{\mathcal{H}_u}(A, B)
$$

we can pull back to a map

<span id="page-13-2"></span>
$$
\pi_u: A \to QM(C^{\mathbb{G}}(A,B)). \tag{2.7}
$$

<span id="page-13-3"></span>We want to make sure for example that this map does not depend on the chosen model.

**Lemma 2.17.** The map  $(2.7)$  is independent of the chosen model.

*Proof.* It is enough to show the following: if  $(H, U, \pi, \rho)$  is any G-equivariant A-B-correspondence, then (using the same symbol to denote extensions of non-degenerate ∗-homomorphisms to their quasi-multipliers)

$$
\theta_{\mathcal{H}}(\theta_{\mathcal{H}_u}^{-1}(\pi_u(a))) = \pi_{\mathcal{H}}(a), \qquad \forall a \in A.
$$

Since any  $H$  is a direct sum of cyclic ones, it is enough to assume that  $H$  is cyclic as a  $\mathbb{G}\text{-}A\text{-}B\text{-}correspondence.$ But then since  $\mathcal{H}_u$  contains any cyclic G-A-B-correspondence up to unitary equivalence, we may as well assume that  $\mathcal{H} \subseteq \mathcal{H}_u$ . It is now easily seen that if  $x \in QM(C^{\mathbb{G}}(A, B))$ , then

$$
\theta_{\mathcal{H}}(x) = \theta_{\mathcal{H}_u}(x)_{|\mathcal{H}}.
$$

In particular, taking  $x = \pi_u(a)$  for  $a \in A$ , we see that

$$
\theta_{\mathcal{H}}(\theta_{\mathcal{H}_u}^{-1}(\pi_u(a))) = \pi_{\mathcal{H}_u}(a)_{|\mathcal{H}} = \pi_{\mathcal{H}}(a).
$$

 $\Box$ 

Similarly, one shows that there is a canonical map

$$
\rho_u: B \to QM(C^{\mathbb{G}}(A,B)).
$$

**Remark 2.18.** • We do not know if the map  $\pi_u$  in [\(2.7\)](#page-13-2) is multiplicative from A to  $C^{\mathbb{G}}(A, B)^{**}$ . Indeed, it is not clear if elements of  $C^{G}(A, B)^{**}$  can be separated by (normal extensions of) \*representations  $\theta$  coming from G-A-B-correspondences, while it is only in such representations that multiplicativity statements for A can be deduced.

• On the other hand, it follows from Theorem [2.13](#page-11-3) that  $\pi_u(A_s) \subseteq M(C^{\mathbb{G}}(A,B))$ , and hence the same is true for the C<sup>\*</sup>-algebra generated by A<sub>s</sub>. In particular, it is hence true that  $\pi_u(ab) = \pi_u(a)\pi_u(b)$ for  $a, b \in A_s$ .

We can now deduce a universal property characterizing \*-representations of  $C^{\mathbb{G}}(A, B)$  coming from equivariant correspondences. Let us first consider two extreme cases.

**Lemma 2.19.** Assume that  $G$  is the trivial group. Then

$$
C^{\mathbb{G}}(A, B) \cong A \otimes_{\text{bin}} B^{\text{op}}
$$

where  $\otimes_{\text{bin}}$  denotes the binormal tensor product of von Neumann algebras [\[EL77\]](#page-47-1).

Proof. This follows immediately from the discussion in [\[EL77,](#page-47-1) Section 2], together with the fact that an A-B-correspondence is a direct sum of cyclic ones.

**Lemma 2.20.** Assume that  $A = B = \mathbb{C}$ . Then

$$
C^{\mathbb{G}}(\mathbb{C},\mathbb{C})\cong C^*(\mathbb{G}),
$$

the usual universal group C<sup>∗</sup> -algebra of a locally compact quantum group [\[Kus01\]](#page-47-21).

Note: since our conventions on corepresentations are the opposite of the one in [\[Kus01\]](#page-47-21), we tacitly modify the definition of  $C^*(\mathbb{G})$  below correspondingly.

*Proof.* Recall from Proposition [1.2](#page-7-3) that any unitary G-representation  $(\mathcal{H}, U)$  is continuous, i.e.

$$
U \in M(\mathcal{K}(\mathcal{H}) \otimes C_0^{\text{red}}(\mathbb{G})).
$$

We now recall briefly the construction of  $C^*(\mathbb{G})$ . Let S be the (unbounded) antipode for  $C_0^{\text{red}}(\mathbb{G})$ , with domain  $\mathcal{D}(S)$ . For  $\omega \in L^1(\mathbb{G})$ , denote

$$
\omega^* : \mathcal{D}(S) \to \mathbb{C}, \quad x \mapsto \overline{\omega(S(x)^*)}.
$$

Then following [\[Kus01,](#page-47-21) [KV03\]](#page-47-4), we denote

$$
L^1_*(\mathbb{G}) := \{ \omega \in L^1(\mathbb{G}) \mid \exists \omega' \in L^1(\mathbb{G}) \text{ with } \omega'_{|\mathcal{D}(S)} = \omega^* \}.
$$

It then follows from [\[Kus01,](#page-47-21) Proposition 5.2] that  $C^*(\mathbb{G})$  may be identified with the universal completion of  $L^1_*(\mathbb{G})$  for the norm

 $\|\omega\|_u = \sup\{\|(\mathrm{id} \otimes \omega)U\| \mid U \text{ unitary } \mathbb{G}\text{-representation}\}.$ 

It now follows by the universal properties of the objects involved, that we obtain an isometric inclusion of C ∗ -algebras

<span id="page-14-0"></span> $C^*(\mathbb{G}) \to C^{\mathbb{G}}(\mathbb{C}, \mathbb{C}), \qquad L^1_*(\mathbb{G}) \ni \omega \mapsto \omega \in L^1$  $(2.8)$ 

It only remains to observe that, through a smoothing argument with respect to the scaling group, the space  $L^1_*(\mathbb{G})$  is norm-dense in  $L^1(\mathbb{G})$  (for the usual pre-dual norm). Hence [\(2.8\)](#page-14-0) must be an isomorphism of C ∗ -algebras.  $\Box$ 

In the following, we will simply identify  $C^*(\mathbb{G}) = C^{\mathbb{G}}(\mathbb{C}, \mathbb{C})$ . Then taking a universal model for  $C^*(\mathbb{G})$ , we obtain the associated universal multiplicative unitary

$$
V_u \in M(C^*(\mathbb{G}) \otimes C_0^{\text{red}}(\mathbb{G}))
$$

as in [\[Kus01,](#page-47-21) Proposition 5.2]. Then for  $A, B$  any pair of G-dynamical von Neumann algebras, we have a unique non-degenerate C<sup>∗</sup> -homomorphism

$$
\theta_{\mathbb{G}}: C^*(\mathbb{G}) \to C^{\mathbb{G}}(A, B), \qquad (\mathrm{id} \otimes \omega)V_u \mapsto (\mathrm{id} \otimes \omega)U_u, \qquad \omega \in M_*.
$$

<span id="page-14-1"></span>**Proposition 2.21.** Let  $\theta$  be a non-degenerate \*-representation of  $C^{\mathbb{G}}(A, B)$ . Then  $\theta = \theta_{\mathcal{H}}$  for a  $\mathbb{G}$ -A-Bcorrespondence  $(\mathcal{H}, U, \pi, \rho)$  if and only if

(1)  $\theta \circ \theta_{\mathbb{G}}$  is a non-degenerate \*-representation of  $C^*(\mathbb{G})$ ,

- <span id="page-15-1"></span><span id="page-15-0"></span>(2)  $\theta \circ \pi_u$  is a normal unital  $\ast$ -representation of A, and
- (3)  $\theta \circ \rho_u$  is a normal unital  $\ast$ -anti-representation of B.

*Proof.* If  $\theta = \theta_{\mathcal{H}}$ , a similar argument as in Lemma [2.17](#page-13-3) shows that  $\theta$  satisfies

$$
\theta(\pi_u(a)) = \pi(a), \quad \theta(\rho_u(b)) = \rho(b), \quad \theta((\mathrm{id} \otimes \omega)U_u) = (\mathrm{id} \otimes \omega)U, \qquad a \in A, b \in B, \omega \in M_*,
$$

so  $\theta$  satisfies the conditions [\(1\)](#page-14-1), [\(2\)](#page-15-0) and [\(3\)](#page-15-1).

Assume now conversely that  $(\mathcal{H}, \theta)$  satisfies the conditions [\(1\)](#page-14-1), [\(2\)](#page-15-0) and [\(3\)](#page-15-1). Write

$$
\pi = \theta \circ \pi_u, \quad \rho = \theta \circ \rho_u,
$$

<span id="page-15-2"></span>and let U be the unitary G-representation associated to the non-degenerate \*-representation  $\theta \circ \theta_{\mathbb{G}}$  of  $C^*(\mathbb{G})$ . Then we claim that

- (a)  $\pi$  and  $\rho$  commute elementwise.
- (b) U implements  $(\pi \otimes id)\alpha$  on  $\pi(A)$ .
- (c)  $U^*$  implements  $(\rho \otimes R)\beta$  on  $\rho(B)$ .

This will make  $(\mathcal{H}, U, \pi, \rho)$  into a G-equivariant A-B-correspondence, and the equality  $\theta = \theta_{\mathcal{H}}$  follows from

$$
\theta(a\otimes\omega\otimes b)=\theta(\pi_u(a)\pi_{\mathbb{G}}((\mathrm{id}\otimes\omega)(V_u))\rho_u(b))=\pi(a)(\mathrm{id}\otimes\omega)(U)\rho(b),\qquad a\in A_\mathrm{s}, b\in B_\mathrm{s}, \omega\in M_*.
$$

Let us tackle first [\(a\).](#page-15-2)

Since  $A_s$  is  $\sigma$ -weakly dense in A, and  $B_s$  is  $\sigma$ -weakly dense in B, it is by the assumption of normality of π and  $\rho$  sufficient to prove that  $\pi(A_s) = \theta(\pi_u(A_s))$  and  $\rho(B_s) = \theta(\rho_u(B_s))$  commute elementwise. But  $\pi_u(A_s)$ ,  $\rho_u(B_s)$  lie in the multiplier algebra of  $C^{\mathbb{G}}(A, B)$ , so we can check the elementwise commutation of  $\pi_u(A_s)$  and  $\rho_u(B_s)$  in a universal model, where this is obvious.

Let us now tackle the second item (the third item is similar).

Again by normality of  $\pi$ , it is sufficient to show that

<span id="page-15-3"></span>
$$
U(\pi(a)\otimes 1)U^* = (\pi\otimes \mathrm{id})(\alpha(a)), \qquad a \in A_s.
$$
 (2.9)

Then since  $\pi_u(A_s) \subseteq M(C^{\mathbb{G}}(A,B))$  and

$$
U_u = (\theta_{\mathbb{G}} \otimes \mathrm{id})V_u \in M(C^{\mathbb{G}}(A,B) \otimes C_0^{\mathrm{red}}(\mathbb{G})),
$$

we have that

$$
(\pi_u \otimes id)(\alpha(a)) = U_u(\pi_u(a) \otimes 1)U_u^* \in M(C^{\mathbb{G}}(A,B) \otimes C_0^{\text{red}}(\mathbb{G})), \quad a \in A_s,
$$

using a universal model  $(\mathcal{H}_u, U_u, \pi_u, \rho_u)$  to make sense of the left hand side as an operator on  $\mathcal{H}_u \otimes L^2(\mathbb{G})$ a priori. Applying id ⊗ $\omega$  for  $\omega \in M_*$ , we then find for  $a \in A_s$  and  $\omega \in M_*$  that

$$
\pi((\mathrm{id}\otimes\omega)\alpha(a))=\theta(\pi_u((\mathrm{id}\otimes\omega)\alpha(a)))=\theta((\mathrm{id}\otimes\omega)(U_u(\pi_u(a)\otimes 1)U_u^*))=(\mathrm{id}\otimes\omega)(U(\pi(a)\otimes 1)U^*).
$$

The identity  $(2.9)$  follows immediately from this.

2.3. Fell topology. For C a C<sup>\*</sup>-algebra, we denote by  $\text{Rep}_*(C)$  the W<sup>\*</sup>-category of its non-degenerate representations.

The following lemma states that the functor  $Corr^{\mathbb{G}}(A, B) \to \text{Rep}_*(C^{\mathbb{G}}(A, B))$  (determined by [\(2.6\)](#page-13-4)) is full.

**Lemma 2.22.** If  $(H, U, \pi, \rho), (\widetilde{H}, \widetilde{U}, \widetilde{\pi}, \widetilde{\rho})$  are G-A-B-correspondences, then an element  $x \in B(H, \widetilde{H})$  is an intertwiner of  $\mathbb{G}-A-B$ -correspondences if and only if it is an intertwiner of  $C^{\mathbb{G}}(A, B)$ -correspondences.

*Proof.* Clearly an intertwiner of G-A-B-correspondences is also an intertwiner of  $C^{\mathbb{G}}(A, B)$ -representations, by construction of  $\theta_{\mathcal{H}}$ . Conversely, if x intertwines  $\theta_{\mathcal{H}}$  and  $\theta_{\widetilde{\mathcal{H}}}$ , we have (again by construction) that

$$
x\pi(a)U(\omega)\rho(b) = \widetilde{\pi}(a)\widetilde{U}(\omega)\widetilde{\rho}(b)x, \qquad \forall a \in A_s, \omega \in M_*, b \in B_s.
$$

Taking  $a = 1$  and  $b = 1$ , we find  $xU(\omega) = \widetilde{U}(\omega)x$  for all  $\omega \in M_*$ . Also, (taking  $b = 1$ )

$$
x\pi(a)U(\omega) = \widetilde{\pi}(a)\widetilde{U}(\omega)x = \widetilde{\pi}(a)xU(\omega), \qquad \forall a \in A_s, \omega \in M_*
$$

so that by non-degeneracy of  $C_U$  we find  $x\pi(a) = \tilde{\pi}(a)x$  for  $a \in A_s$ . Since  $A_s$  is  $\sigma$ -weakly dense in A, we find  $x\pi(a) = \tilde{\pi}(a)x$  for  $a \in A$ . Similarly  $x\rho(b) = \tilde{\rho}(b)x$  for  $b \in B$ . Thus, x is an intertwiner of G-A-B-correspondences. correspondences.

We now define a topology on the objects of  $Corr^{\mathbb{G}}(A, B)$ . Since objects in a category C such as this do not form a set, we need to be careful in specifying what we mean by this. We use the formalism that a topology for objects in a category  $\mathcal C$  is determined by a set-indexed local open neighborhood system: to each object  $X \in \mathcal{C}$  is assigned a directed set  $\mathcal{F}_X$  and, to each  $F \in \mathcal{F}_X$ , a collection of objects  $\mathcal{C}_F(X)$  in  $\mathcal{C}$ , such that

- (1) X is an object of  $C_F(X)$  for each  $F \in \mathscr{F}_X$ ,
- (2)  $C_{F'}(X)$  is contained in  $C_F(X)$  if  $F \leq F'$ , and
- (3) if Y is an object of some  $C_F(X)$ , there exists  $F' \in \mathscr{F}_Y$  such that  $C_{F'}(Y)$  is contained in  $C_F(X)$ .

We then say that two such set-indexed local open neighborhood systems determine the same topology  $\tau$ , if at each object X the respective open neighborhood systems can be nested in each other.

<span id="page-16-0"></span>**Definition 2.23.** Let  $(\mathcal{H}, U, \pi, \rho)$  be a G-A-B-correspondence. Let F be a collection of a scalar  $\varepsilon > 0$  and finite subsets  $F_A, F_B, F_{\mathbb{G}}$  and  $F_{\mathcal{H}} = \{\xi_1, \ldots, \xi_n\}$  in resp.  $A, B, M_*$  and  $\mathcal{H}$ .

We say that a G-A-B-correspondence  $(\mathcal{H}', U', \pi', \rho')$  lies in the F-neighborhood  $U(\mathcal{H}; F)$  of  $(\mathcal{H}, U, \pi, \rho)$  if there exist  $\xi'_1,\ldots,\xi'_n$  in  $\mathcal{H}'$  with  $\|\xi'_i\|\leq \|\xi_i\|$  for all i, such that, for all  $a \in F_A, b \in F_B, \omega \in F_{\mathbb{G}}$  and  $1 \leq i, j \leq n,$ 

<span id="page-16-1"></span>
$$
|\langle \xi_i, \pi(a)U(\omega)\rho(b)\xi_j \rangle - \langle \xi'_i, \pi'(a)U'(\omega)\rho'(b)\xi'_j \rangle| < \varepsilon.
$$
\n(2.10)

Such  $F$  as above form a directed set in a natural way, by putting

$$
F \leq F' \quad \iff \quad \varepsilon \geq \varepsilon', F_A \subseteq F'_A, F_B \subseteq F'_B, F_{\mathbb{G}} \subseteq F'_{\mathbb{G}}, F_{\mathcal{H}} \subseteq F'_{\mathcal{H}}.
$$

It is then easily verified that the neighborhoods defined in Definition [2.23](#page-16-0) define a set-indexed local open neighborhood system, and hence a topology  $\tau$  on Corr<sup>G</sup> $(A, B)$ . Moreover, for any finite collection of Gequivariant correspondences  $\mathcal{H}_1, \ldots, \mathcal{H}_n$  and neighborhoods  $U(\mathcal{H}_i; F_i)$ , there exists F such that

$$
U(\bigoplus_{i=1}^{n} \mathcal{H}_i; F) \subseteq \bigcap_{i=1}^{n} U(\mathcal{H}_i; F_i). \tag{2.11}
$$

Remark 2.24. The ordering of the factors in  $(2.10)$  is important: in general, one seems to get a different (and less natural) notion if for example the positions of  $U(\omega)$  and  $\pi(a)$  are interchanged. However, one may interchange the positions of  $\pi(a)$  and  $\rho(b)$  at no cost, as will follow easily from Theorem [2.28.](#page-17-0)

Recall now from Theorem [2.16](#page-13-0) that any  $\mathbb{G}-A-B$ -correspondence H leads to a (unique) non-degenerate  $*$ representation  $(\mathcal{H}, \theta_{\mathcal{H}})$  of  $C^{G}(A, B)$ . We will show that the  $\tau$ -topology on  $Corr^{G}(A, B)$  is simply the restriction of the corresponding 'Zimmer topology' on the W<sup>\*</sup>-category  $\text{Rep}_*(C^{\mathbb{G}}(A, B))$  [\[Zim84\]](#page-47-25) (see also [\[BdlHV08\]](#page-46-24)Appendix F), passing from the group case to the C<sup>∗</sup>-algebra setting.

<span id="page-16-2"></span>**Definition 2.25.** Let C be a  $C^*$ -algebra, and let  $(\mathcal{H}, \theta)$  be a non-degenerate  $*$ -representation of C. Let F consist of  $\varepsilon > 0$  and finite sets  $F_C \subseteq C$  and  $F_H = \{\xi_1, \ldots, \xi_n\} \subseteq \mathcal{H}$ .

If  $(H', \theta')$  is a non-degenerate \*-representation of C, we say that  $(H', \theta')$  lies in the F-neighborhood  $V(H; F)$ of  $(\mathcal{H}, \theta)$  if there exist  $\xi'_1, \ldots, \xi'_n \in \mathcal{H}'$  such that  $\|\xi'_i\| \leq \|\xi_i\|$  for all i, and, for all  $c \in F_C$  and  $1 \leq i, j \leq n$ ,

$$
|\langle \xi_i, \theta(c)\xi_j \rangle - \langle \xi'_i, \theta'(c)\xi'_j \rangle| < \varepsilon.
$$

We denote the resulting topology on  $\text{Rep}_*(C)$  by  $\sigma$ .

**Remark 2.26.** By means of a contractive approximate unit in  $C$ , it is easily verified that if in Definition [2.25](#page-16-2) we drop the requirement that  $\|\xi_i'\| \leq \|\xi_i\|$  for all i, the resulting topology coincides with  $\sigma$ . We do not know if this still holds for Definition [2.23.](#page-16-0)

Note now that the definition of  $V(\mathcal{H}; F)$  makes sense if  $F_C \subseteq C^{**}$ , using the unique normal extension of  $\theta$ to  $C^{**}$ . We will then use the following general fact.

<span id="page-17-1"></span>**Lemma 2.27.** Let  $S \subseteq QM(C)$  be a subset such that  $C \subseteq [S]$ . Let  $\sigma_S$  be the topology on  $\text{Rep}_*(C)$  obtained by using as local neighborhoods the  $V(H; F)$  for finite  $F_C \subseteq S$ . Then

 $\sigma_S = \sigma$ .

*Proof.* Let  $(\mathcal{H}, \theta) \in \operatorname{Rep}_*(C)$  and  $F = (\varepsilon, F_C, F_H = \{\xi_1, \ldots, \xi_n\})$  with  $F_C \subseteq S$ . Choose  $d \in C$  with  $||d|| \leq 1$ and

 $\max_{i} {\{\|\xi_i - \theta(d)\xi_i\|\}} \max_{c \in F_C} {\{\|c\|\}} \max_{j} {\{\|\xi_j\|\}} < \varepsilon/4.$ 

Then for all  $1 \leq i, j \leq n$  and all  $c \in F_C$ , we have

 $|\langle \xi_i, \theta(c) \xi_j \rangle - \langle \theta(d) \xi_i, \theta(c) \theta(d) \xi_j \rangle| < \varepsilon/2.$ 

If now  $\overline{F} = (\varepsilon/2, \overline{F}_C, F_H)$  with  $\overline{F}_C = \{d^*cd \mid c \in F_C\} \subseteq C$ , then  $V(\mathcal{H}; \overline{F}) \subseteq V(\mathcal{H}; F)$ . Indeed, if  $(\mathcal{H}', \theta') \in$  $V(\mathcal{H};\tilde{F})$ , we can choose  $\xi'_1,\ldots,\xi'_n \in \mathcal{H}'$  with  $\|\xi'_i\| \le \|\xi_i\|$  such that for all  $1 \le i,j \le n$  and  $c \in F_C$ ,

$$
|\langle \theta(d)\xi_i, \theta(c)\theta(d)\xi_j \rangle - \langle \theta'(d)\xi'_i, \theta'(c)\theta'(d)\xi'_j \rangle| < \varepsilon/2,
$$

and then clearly

$$
|\langle \xi_i, \theta(c)\xi_j \rangle - \langle \theta'(d)\xi'_i, \theta'(c)\theta'(d)\xi'_j \rangle| < \varepsilon.
$$

In particular, this shows that

 $\sigma_S \subseteq \sigma$ .

We now claim conversely that  $\sigma_S$  is finer than  $\sigma$ . In fact, we have the following easy observations: the topology  $\sigma_S$  does not change if we add to S

- scalar multiples of elements of  $S$ ,
- sums of elements of  $S$ , or
- normlimits of elements of  $S$ .

In particular, we may replace S by [S], and so the assumption  $[S] \supseteq C$  immediately gives the claim.

<span id="page-17-0"></span>**Theorem 2.28.** Let  $\mathbb{G}$  be a locally compact quantum group. The topology  $\tau$  on  $\text{Corr}^{\mathbb{G}}(A, B)$  is induced by the corresponding topology  $\sigma$  on  $\text{Rep}_*(C^{\tilde{G}}(A, B))$  under the correspondence  $(\mathcal{H}, U, \pi, \rho) \mapsto (\mathcal{H}, \theta_{\mathcal{H}})$ .

Proof. Observing that

$$
C^{\mathbb{G}}(A,B) = [\pi_u(A_s)C_{U_u}\rho_u(B_s)] \subseteq [\pi_u(A)\{(\text{id}\otimes\omega)U_u \mid \omega \in M_*\}\rho_u(B)] \subseteq QM(C^{\mathbb{G}}(A,B)),
$$
  
this follows immediately from Lemma 2.27.

The topology  $\tau$  itself is not the most useful: rather, we want to consider the Fell topology<sup>[2](#page-17-2)</sup>  $\tau^{\infty}$  which has local neighorhoods

$$
U^{\infty}(\mathcal{H};F) = \{ \mathcal{G} \mid \mathcal{G}^{\infty} \in U(\mathcal{H};F) \}
$$

where here and in the following we use the shorthand notation

$$
\mathcal{G}^{\infty}:=\oplus_{1}^{\infty}\mathcal{G}.
$$

<span id="page-17-2"></span> ${}^{2}$ It is an instructive exercise to check that this indeed determines a well-defined topology, in the sense described earlier.

The corresponding topology  $\sigma^{\infty}$  on  $\text{Rep}_*(C^{\mathbb{G}}(A,B))$  indeed coincides with the usual Fell (=inner hull-kernel) topology [\[Fel62\]](#page-47-26).

### 3. Weak containment

In this section, we fix again a locally compact quantum group  $\mathbb G$  and  $\mathbb G$ -W<sup>\*</sup>-algebras  $A, B$ .

**Definition 3.1.** Let  $H, H'$  be G-A-B-correspondences. We say that  $H'$  is weakly contained in  $H$  iff  $H'$  lies in the  $\tau^{\infty}$ -closure of  $\{\mathcal{H}\}.$ 

We then write

 $\mathcal{H}' \preccurlyeq \mathcal{H}$ .

In other words,  $\mathcal{H}' \preccurlyeq \mathcal{H}$  if and only if for any F as in Definition [2.23,](#page-16-0) we have  $\bigoplus_{1}^{\infty} \mathcal{H} \in U(\mathcal{H}';F)$ .

Clearly, weak containment is a transitive and reflexive relation. We also note that in terms of the C<sup>∗</sup> -algebra  $C^{\mathbb{G}}(A, B)$ , we have  $\mathcal{G} \preccurlyeq \mathcal{H}$  if and only if the representation  $\theta_{\mathcal{G}}$  of  $C^{\mathbb{G}}(A, B)$  factors through  $\theta_{\mathcal{H}}(C^{\mathbb{G}}(A, B))$ (see e.g. [\[Dix77,](#page-46-25) Section 3.4]).

<span id="page-18-1"></span>**Remark 3.2.** Let I, J be arbitrary sets, and let  $H, H'$  be  $\mathbb{G}\text{-}A\text{-}B\text{-}correspondences$ . Then it follows immediately from the above description that

$$
\mathcal{H}' \preccurlyeq \mathcal{H} \qquad \Leftrightarrow \qquad \oplus_{j \in J} \mathcal{H}' \preccurlyeq \oplus_{i \in I} \mathcal{H},
$$

i.e. weak containment does not witness multiplicities. More intrinsically, we can write this as

$$
\mathcal{H}' \preccurlyeq \mathcal{H} \qquad \Leftrightarrow \qquad l^2(J) \otimes \mathcal{H}' \preccurlyeq l^2(I) \otimes \mathcal{H},
$$

and we may then as well replace  $l^2(I), l^2(J)$  with arbitrary multiplicity Hilbert spaces  $\mathcal{G}, \mathcal{G}'$ .

In this section, we will provide a different characterization of weak containment. This will generalize to the equivariant setting some of the techniques and ideas of [\[A-D90,](#page-46-4) [A-DH90,](#page-46-26) [A-D95\]](#page-46-5).

Before we continue, we prove a small preparatory lemma on von Neumann algebras and nsf weights. Although it must be well-known, we do not know a convenient reference, so we briefly provide a proof.

Recall that if B is a von Neumann algebra and  $\varphi$  an nsf weight on B, we can endow the opposite von Neumann algebra  $B^{\rm op}$  with the nsf weight

$$
\varphi^{\text{op}}: B^{\text{op}}_+ \to [0, +\infty], \quad b^{\text{op}} \mapsto \varphi(b).
$$
  
Then  $\mathcal{N}_{\varphi^{\text{op}}} = (\mathcal{N}_{\varphi}^*)^{\text{op}}$ , and we can identify  $L^2(B)$  with  $L^2(B^{\text{op}})$  via  

$$
\Lambda_{\varphi^{\text{op}}}(b^{\text{op}}) = J\Lambda_{\varphi}(b^*), \qquad b \in \mathcal{N}_{\varphi}^*.
$$

<span id="page-18-0"></span>**Lemma 3.3.** Let B be a von Neumann algebra. If  $\xi_1, \ldots, \xi_n \in L^2(B)$ , there exists an nsf weight  $\varphi$  on B, a vector  $\xi \in L^2(B)$  and  $b_1, \ldots, b_n \in B$  such that  $b_i^{op} \in \mathcal{N}_{\varphi^{op}}$  for all i and

$$
\xi_i = \rho_B(b_i)\xi = \Lambda_{\varphi^{\rm op}}(b_i^{\rm op}), \qquad \forall i. \tag{3.1}
$$

Proof. Let  $\mathcal{H} = [\rho_B(B)\xi_1 + \ldots + \rho_B(B)\xi_n]$ . Then  $\mathcal{H} = \pi_B(p)L^2(B)$  for some selfadjoint projection  $p \in B$ . Put  $q = 1 - p$ , and let  $\psi$  be an nsf weight on qBq. Let

$$
\omega: B \to \mathbb{C}, \qquad b \mapsto \sum_{i=1}^n \langle \xi_i, \pi_B(b)\xi_i \rangle,
$$

and put

$$
\varphi(b) = \omega(pbp) + \psi(qbq), \qquad b \in B_+.
$$

Then clearly  $\varphi$  is an nsf weight on B, with p integrable. Moreover, p is invariant under the modular automorphism group of  $\varphi$ , so we can define

$$
\xi:=\Lambda_\varphi(p)=\Lambda_{\varphi^{\rm op}}(p^{\rm op}).
$$

Now an easy calculation gives that

$$
\langle \xi, \pi_B(b)\xi \rangle = \omega(b), \qquad b \in B.
$$

Recalling the notation [\(0.7\)](#page-4-0), it follows that there exists  $b_i \in B$  such that  $\rho_B(b_i)\pi_B(b)\xi = \pi_B(b)\xi_i$ . In particular,  $\rho_B(p b_i) \xi = \xi_i$ , and clearly  $\rho_B(p b_i) \xi = \Lambda_{\varphi} \circ \mathfrak{p}((p b_i))$  $\overline{\mathcal{O}}$ .

Assume now that  $H, G$  are Hilbert spaces with a normal anti-∗-representation  $\rho_H, \rho_G$  by the von Neumann algebra  $B$ . Recall the notation  $(0.4)$ . In particular, we write

<span id="page-19-4"></span>
$$
X_B(\mathcal{H}) := X_B(L^2(B), \mathcal{H}).\tag{3.2}
$$

We can view  $X_B(\mathcal{H})$  as a self-dual (right) Hilbert B-module [\[Pas73,](#page-47-27) [Rie74\]](#page-47-28), and more generally  $X_B(\mathcal{G}, \mathcal{H})$  as a W<sup>∗</sup> -TRO [\[Zet93\]](#page-47-29), but we will not really need this more abstract viewpoint for now.

<span id="page-19-3"></span>Also the following lemma is elementary.

**Lemma 3.4.** Let G be a Hilbert space with a normal anti- $*$ -representation  $\rho_G$  by B. Then there exists a net of contractions  $x_i \in \bigcup_{n=1}^{\infty} X_B(L^2(B)^n, \mathcal{G})$  such that the maps

$$
B(\mathcal{G}) \to B(\mathcal{G}), \qquad y \mapsto x_i x_i^* y x_i x_i^*
$$

converge pointwise  $\sigma$ -weakly to the identity map.

Proof. It follows immediately from the fact that we can identify

$$
(\mathcal{G}, \rho_{\mathcal{G}}) \cong ((\pi_B \otimes id)(p)(L^2(B) \otimes l^2(I)), \rho_B \otimes id)
$$

for some set I and some  $p \in B\overline{\otimes}B(l^2)$  $(I)$ ).

We will also need a slight extension/variation of [\[A-DH90,](#page-46-26) Lemma 2.2]. Although the proof almost applies ad verbatim, we prefer to include it for the convenience of the reader.

<span id="page-19-2"></span>**Lemma 3.5.** Let B be a von Neumann algebra, and let  $H, G$  be Hilbert spaces with a normal anti- $*$ representation  $\rho_{\mathcal{H}}, \rho_{\mathcal{G}}$  of B. Let  $F \subseteq X_B(\mathcal{G}, \mathcal{H})$  be a subset which is closed under right multiplication with  $\rho_G(B)'$ . Let  $S \subseteq B(H)$  be a subset with  $1 \in S$ , and assume that  $\{ \mathrm{Ad}_{y^*} : S \to B(G) \mid y \in F \}$  is convex. Let  $x \in X_B(\mathcal{G}, \mathcal{H})$  with  $||x|| \leq 1$ , and assume that there exists a net of  $x_i \in F$  such that for each  $s \in S$ 

<span id="page-19-0"></span>
$$
x_i^* s x_i \to x^* s x \quad weakly \ in \ B(\mathcal{G}). \tag{3.3}
$$

Then there exists a net of contractive  $x_i \in F$  such that [\(3.3\)](#page-19-0) holds for the  $\sigma$ -weak topology.

*Proof.* As in [\[A-DH90,](#page-46-26) Lemma 2.2], we begin with proving the lemma under the extra condition that  $x^*x = 1$ .

Assume that a net of  $x_i \in F$  is given such that [\(3.3\)](#page-19-0) holds for each  $s \in S$ . By the assumption that  ${((\mathrm{Ad}_{y^*})_{|S} \mid y \in F)}$  is convex, we may choose this net so that  $(3.3)$  in fact holds with respect to the *strong* topology.

Put

$$
y_i = 2x_i(1 + x_i^* x_i)^{-1} \in F.
$$

Following further the computation as in [\[A-DH90,](#page-46-26) Lemma 2.2], we find

$$
(x_i - y_i)^*(x_i - y_i) \le (x_i^* x_i - 1)^2,
$$

and for each vector state  $\omega = \omega_{\xi}$  on  $B(\mathcal{G})$  and  $s \in S$  we have

<span id="page-19-1"></span>
$$
|\omega(x_i^*sx_i - y_i^*sy_i)| \le ||s||\omega((x_i^*x_i - 1)^2)^{1/2}(\omega(x_i^*x_i)^{1/2} + \omega(y_i^*y_i)^{1/2}).
$$
\n(3.4)

Since  $||y_i|| \leq 1$  and since  $\text{Ad}_{x_i^*}(1) = x_i^* x_i$  converges strongly to  $x^* x = 1$  by assumption, we find that  $(3.4)$ goes to zero in the limit. It follows that  $y_i$  is a contractive net in F such that for each  $s \in S$  the net  $Ad_{y_i^*}(s)$ converges weakly (and hence  $\sigma$ -weakly) to  $x^*sx$ , proving the lemma in this special case.

Let us now consider the case where  $x \in X_B(G, \mathcal{H})$  is an arbitrary contraction. Let  $e \in \rho_{\mathcal{G}}(B)' \subseteq B(\mathcal{G})$  be the support projection for  $x^*x$ . Then we may in fact view  $\text{Ad}_{x^*}$  as a map  $B(\mathcal{H}) \to B(e\mathcal{G})$ . By multiplying a net  $x_i$  as in [\(3.3\)](#page-19-0) with e on the right, we see that without loss of generality we may in fact assume that  $e = 1$ .

Let then  $x = u|x|$  be the polar decomposition of x, where  $u \in X_B(G, \mathcal{H})$  is isometric by the above assumption. Then u lies in the strong closure of the unit ball of  $x \rho_{\mathcal{G}}(B)'$ . It follows from this last property, together with the fact that F is closed under right multiplication with  $\rho_g(B)'$ , that  $Ad_{u^*}$  can be approximated pointwise weakly on S with maps  $\mathrm{Ad}_{w_i^*}$  for  $w_i \in F$ . By the first part of the proof, we may assume the  $w_i$  to be contractive. But then the  $y_i = w_i |x|$  will also be contractive, with  $\text{Ad}_{y_i^*}$  converging to  $\text{Ad}_{x^*}$  pointwise weakly and hence  $\sigma$ -weakly on S.

<span id="page-20-3"></span>Assume now again that  $(A, \alpha_A)$  and  $(B, \alpha_B)$  are G-W<sup>\*</sup>-algebras.

**Lemma 3.6.** Let  $H, G$  be two G-equivariant A-B-correspondences. Assume that G is weakly contained in H. Then there exists a contractive net  $y_i \in X_B(\mathcal{G}, \mathcal{H}^\infty)$  such that for all  $c \in C^{\mathbb{G}}(A, B)$ 

<span id="page-20-2"></span>
$$
y_i^* \theta_{\mathcal{H}^\infty}(c) y_i \to \theta_{\mathcal{G}}(c) \quad \sigma\text{-weakly in } B(\mathcal{G}).
$$
\n(3.5)

*Proof.* We first prove the following: if, for some n, we are given a contractive map  $T = (T_1, \ldots, T_n) \in$  $X_B(L^2(B)^n, \mathcal{G})$ , there exists a 1-bounded net  $y_i \in X_B(L^2(B)^n, \mathcal{H}^\infty)$  such that for all  $c \in C^{\mathbb{G}}(A, B)$ 

<span id="page-20-1"></span>
$$
y_i^* \theta_{\mathcal{H}^\infty}(c) y_i \to T^* \theta_{\mathcal{G}}(c) T \quad \sigma\text{-weakly in } B(L^2(B)^n). \tag{3.6}
$$

Indeed, pick a finite set  $F_{L^2(B)} \subseteq L^2(B)$  and a finite subset  $F_C \subseteq C^{\mathbb{G}}(A, B) \cup \{1\} \subseteq QM(C^{\mathbb{G}}(A, B))$ . Take  $\varepsilon > 0$ . We first look for elements  $x_1, \ldots, x_n \in X_B(\mathcal{H}^\infty)$  such that

<span id="page-20-0"></span>
$$
|\langle T_i\xi, \theta_{\mathcal{G}}(z)T_j\eta\rangle - \langle x_i\xi, \theta_{\mathcal{H}^\infty}(z)x_j\eta\rangle| < \varepsilon, \qquad \forall z \in F_C, \forall \xi, \eta \in F_{L^2(B)}, \forall i, j: 1 \to n. \tag{3.7}
$$

By Lemma [3.3,](#page-18-0) we can choose  $\zeta \in L^2(B)$  and an nsf weight  $\varphi$  on B together with a finite set  $F_B \subseteq B$  such that each  $b^{\text{op}} \in \mathcal{N}_{\varphi^{\text{op}}}$  for  $b \in F_B$  and for each  $\xi \in F_{L^2(B)}$  there exists  $b \in F_B$  with

$$
\xi = \rho_B(b)\zeta = \Lambda_{\varphi^{\rm op}}(b^{\rm op}).
$$

Since  $(\mathcal{H}, \theta_{\mathcal{H}})$  lies in an arbitrary  $\tau^{\infty}$ -neighborhood of  $(\mathcal{G}, \theta_{\mathcal{G}})$ , and since  $\rho_u(B) \subseteq QM(C^{\mathbb{G}}(A, B))$  implies that also  $\rho_u(B)(C^{\mathbb{G}}(A,B) \cup \{1\})\rho_u(B) \subseteq QM(C^{\mathbb{G}}(A,B))$ , Lemma [2.27](#page-17-1) allows us to pick  $\zeta_i \in \mathcal{H}^{\infty}$  with  $\|\zeta_i'\| \leq \|T_i\zeta\| \leq \|\zeta\|$  and, for all  $b, c \in F_B, z \in F_C$ ,

$$
\begin{aligned} |\langle T_i \rho_B(b)\zeta, \theta_{\mathcal{G}}(z)T_j \rho_B(c)\zeta\rangle - \langle \rho_{\mathcal{H}^{\infty}}(b)\zeta_i', \theta_{\mathcal{H}^{\infty}}(z)\rho_{\mathcal{H}^{\infty}}(c)\zeta_j'\rangle| \\ = |\langle T_i\zeta, \theta_{\mathcal{G}}(\rho_u(b)^*z\rho_u(c))T_j\zeta\rangle - \langle \zeta_i', \theta_{\mathcal{H}^{\infty}}(\rho_u(b)^*z\rho_u(c))\zeta_j'\rangle| < \varepsilon. \end{aligned}
$$

By norm-perturbing the  $\zeta_i'$  if necessary, we may assume that they are left bounded for  $\varphi$  (in the sense that  $\zeta_i' \in \mathfrak{D}(\mathcal{H}^{\infty}, \varphi)$ , see [\[Tak03,](#page-47-17) Section IX.3]). But then with  $x_i = L_{\zeta_i'} \in X_B(\mathcal{H}^{\infty})$ , we find that  $\rho_{\mathcal{H}^{\infty}}(b)\zeta_i' =$  $x_i \Lambda_{\varphi^{\text{op}}}(b^{\text{op}}) = x_i \rho_B(b) \zeta$ , so that [\(3.7\)](#page-20-0) is satisfied.

It follows that we can find a net  $x_i \in X_B(L^2(B)^n, \mathcal{H}^\infty)$  such that  $\text{Ad}_{x_i^*} \circ \theta_{\mathcal{H}^\infty}$  converges weakly to  $\text{Ad}_{T^*} \circ \theta_{\mathcal{G}}$ pointwise on  $C^{\mathbb{G}}(A, B) \cup \{1\}$ . Now the set

$$
F = X_B(L^2(B)^n, \mathcal{H}^\infty) \subseteq X_B(L^2(B)^n, \mathcal{G} \oplus \mathcal{H}^\infty)
$$

is stable under right multiplication with  $\rho_{L^2(B)^n}(B)$ , and with  $\text{diag}_{\infty}: B(\mathcal{H}) \to B(\mathcal{H}^{\infty})$  the diagonal inclusion, we have that

$$
\{ \mathrm{Ad}_{x^*} : B(\mathcal{G}) \oplus \mathrm{diag}_{\infty}(B(\mathcal{H})) \to B(L^2(B)^n) \mid x \in F \}
$$

is convex. By viewing also  $T \in X_B(L^2(B)^n, \mathcal{G}) \subseteq X_B(L^2(B)^n, \mathcal{G} \oplus \mathcal{H}^\infty)$  and applying Lemma [3.5](#page-19-2) to  $F \subseteq X_B(L^2(B)^n, \mathcal{G} \oplus \mathcal{H}^\infty)$  and  $S = (\theta_{\mathcal{G}} \oplus \theta_{\mathcal{H}^\infty})(C^{\mathbb{G}}(A,B) \cup \{1\}),$  we can conclude that we can also find a net  $y_i \in X_B(L^2(B)^n, \mathcal{H}^\infty)$  with  $||y_i|| \leq 1$  such that  $\text{Ad}_{y_i^*} \circ \theta_{\mathcal{H}^\infty}$  converges  $\sigma$ -weakly to  $\text{Ad}_{T^*} \circ \theta_{\mathcal{G}}$  pointwise on  $C^{\mathbb{G}}(A, B) \cup \{1\}$ . This proves in particular [\(3.6\)](#page-20-1).

We can now prove the lemma. Pick  $T \in X_B(L^2(B)^n, \mathcal{G})$  contractive for some n, and choose a contractive net  $y_i \in X_B(L^2(B)^n, \mathcal{H}^\infty)$  such that  $\text{Ad}_{y_i^*} \circ \theta_{\mathcal{H}^\infty}$  converges weakly to  $\text{Ad}_{T^*} \circ \theta_{\mathcal{G}}$  pointwise on  $C^{\mathbb{G}}(A, B)$ . Then  $y_iT^* \in X_B(G, \mathcal{H}^\infty)$  is still contractive, and for all  $c \in C^{\mathbb{G}}(A, B)$  we have

$$
(y_iT^*)^*\theta_{\mathcal{H}^\infty}(c)(y_iT^*) \to TT^*\theta_{\mathcal{G}}(c)TT^*
$$
  $\sigma$ -weakly in  $B(\mathcal{G})$ .

By Lemma [3.4,](#page-19-3) the pointwise  $\sigma$ -weak closure of the maps  $B(G) \ni x \mapsto TT^* x TT^* \in B(G)$  contains the identity map, proving  $(3.5)$ .

Given G-equivariant A-B-correspondences  $H, G$ , we obtain a coaction

$$
\alpha_{X_B(\mathcal{G},\mathcal{H})}: X_B(\mathcal{G},\mathcal{H}) \to X_B(\mathcal{G},\mathcal{H}) \bar{\otimes} M, \quad x \mapsto U_{\mathcal{H}}(x \otimes 1) U_{\mathcal{G}}^*.
$$
\n(3.8)

Here, the tensor product  $X_B(\mathcal{G}, \mathcal{H})\bar{\otimes}M$  can be interpreted in multiple ways. We can view it as the Fubini tensor product of operator spaces, or alternatively, we can view  $X_B(\mathcal{G}, \mathcal{H}) \odot M$  as a corner of

$$
X_{B}({\mathcal G}\oplus{\mathcal H},{\mathcal G}\oplus{\mathcal H})\bar{\otimes}M
$$

and then define  $X_B(\mathcal{G}, \mathcal{H})\bar{\otimes}M$  to be the  $\sigma$ -weak closure of  $X_B(\mathcal{G}, \mathcal{H})\odot M$  inside  $X_B(\mathcal{G} \oplus \mathcal{H}, \mathcal{G} \oplus \mathcal{H})\bar{\otimes}M$ . Both interpretations agree and we can make sense of the notion of a G-coaction on  $X_B(\mathcal{G}, \mathcal{H})$ , see e.g. [\[PS17,](#page-47-30) Section 1 & 3] for more details and information.

<span id="page-21-0"></span>**Theorem 3.7.** Let  $A, B$  be  $\mathbb{G}\text{-}W^*$ -algebras, and let  $\mathcal{H}, \mathcal{G}$  be  $\mathbb{G}\text{-}A\text{-}B\text{-}correspondences$ . Then  $\mathcal{G}$  is equivariantly weakly contained in H if and only if there exists a 1-bounded net  $y_i \in X_B(G, H^\infty)$  such that for all  $a \in A$ 

<span id="page-21-1"></span>
$$
(y_i^* \pi_{\mathcal{H}^{\infty}}(a) \otimes 1)\alpha_{X_B(\mathcal{G}, \mathcal{H}^{\infty})}(y_i) \to \pi_{\mathcal{G}}(a) \otimes 1 \quad \sigma\text{-weakly in } B(\mathcal{G} \otimes L^2(\mathbb{G})).
$$
\n(3.9)

*Proof.* Assume first that G is weakly contained in H. By Lemma [3.6,](#page-20-3) we can find a contractive net  $y_i \in$  $X_B(\mathcal{G}, \mathcal{H}^\infty)$  such that for all  $c \in C^{\mathbb{G}}(A, B)$  we have

$$
y_i^* \theta_{\mathcal{H}^\infty}(c) y_i \to \theta_{\mathcal{G}}(c) \quad \sigma\text{-weakly in } B(\mathcal{G}).\tag{3.10}
$$

By passing to a subnet, we may assume that

$$
\mathrm{Ad}_{y_i^*}: B(\mathcal{H}^\infty) \to B(\mathcal{G}), \qquad x \mapsto y_i^* xy_i. \tag{3.11}
$$

converges pointwise  $\sigma$ -weakly to a ccp map

$$
\widetilde{E}:B(\mathcal{H}^{\infty})\to B(\mathcal{G}).
$$

Put

<span id="page-21-2"></span>
$$
E = \widetilde{E} \circ \text{diag}_{\infty} : B(\mathcal{H}) \to B(\mathcal{G}), \tag{3.12}
$$

where  $\text{diag}_{\infty}: B(\mathcal{H}) \to B(\mathcal{H}^{\infty})$  is the diagonal inclusion. Then clearly E is a ccp map with

$$
E(\theta_{\mathcal{H}}(z)) = \theta_{\mathcal{G}}(z), \qquad \forall z \in C^{\mathbb{G}}(A, B).
$$

In particular,  $\theta_{\mathcal{H}}(C^{\mathbb{G}}(A,B))$  belongs to the multiplicative domain of E. But then also

$$
\theta_{\mathcal{G}}(x)E(\theta_{\mathcal{H}}(z))\theta_{\mathcal{G}}(y) = E(\theta_{\mathcal{H}}(xzy)) = \theta_{\mathcal{G}}(x)\theta_{\mathcal{G}}(z)\theta_{\mathcal{G}}(y), \qquad x, y \in C^{\mathbb{G}}(A, B), z \in QM(C^{\mathbb{G}}(A, B)).
$$

It follows that

$$
E(\theta_{\mathcal{H}}(z)) = \theta_{\mathcal{G}}(z), \qquad \forall z \in QM(C^{\mathbb{G}}(A,B)).
$$
\n
$$
\text{Since } \pi_u(A)C_{U_u} \subseteq QM(C^{\mathbb{G}}(A,B)), \text{ we find}
$$
\n
$$
(3.13)
$$

<span id="page-21-3"></span>
$$
y_i^* \pi_{\mathcal{H}^{\infty}}(a) U_{\mathcal{H}^{\infty}}(\omega) y_i \stackrel{\sigma\text{-weakly}}{\underset{i\to\infty}{\longrightarrow}} \pi_{\mathcal{G}}(a) U_{\mathcal{G}}(\omega), \qquad \forall a \in A, \omega \in M_*. \tag{3.14}
$$

Hence for all  $\xi, \eta \in \mathcal{G}$  and all  $f, g \in L^2(\mathbb{G})$  and  $a \in A$ , we have

$$
\langle y_i \xi \otimes f, (\pi_{\mathcal{H}^{\infty}}(a) \otimes 1) U_{\mathcal{H}^{\infty}}(y_i \eta \otimes g) \rangle \rightarrow \langle \xi \otimes f, (\pi_{\mathcal{G}}(a) \otimes 1) U_{\mathcal{G}}(\eta \otimes g) \rangle.
$$

By boundedness of the  $y_i$ , this means that for all  $a \in A$  we have

$$
(y_i^* \pi_{\mathcal{H}^{\infty}}(a) \otimes 1)U_{\mathcal{H}^{\infty}}(y_i \otimes 1) \xrightarrow{i \to \infty} \langle \pi_{\mathcal{G}}(a) \otimes 1)U_{\mathcal{G}}.
$$
 (3.15)

Multiplying to the right with  $U_{\mathcal{G}}^*$ , we find [\(3.9\)](#page-21-1).

Let us now go back. Assume that we have chosen contractive  $y_k \in X_B(\mathcal{G}, \mathcal{H}^\infty)$  satisfying [\(3.9\)](#page-21-1). Choose finitely many  $\xi_i \in \mathcal{G}$  as well as a finite set  $F_C \subseteq C^{\mathbb{G}}(A, B)$  and  $\varepsilon > 0$ . By Theorem [2.28,](#page-17-0) it is enough find  $\xi_i' \in \mathcal{H}^{\infty}$  such that  $\|\xi_i'\| \le \|\xi_i\|$  and

$$
|\langle \xi_i, \theta_{\mathcal{G}}(z)\xi_j \rangle - \langle \xi'_i, \theta_{\mathcal{H}^{\infty}}(z)\xi'_j \rangle| < \varepsilon, \qquad \forall z \in F_C, i, j \in \{1, \ldots, n\}.
$$

Now by assumption, we have for all  $\omega \in M_*$  and  $a \in A$  that

$$
y_k^*\pi_{\mathcal{H}^\infty}(a)U_{\mathcal{H}^\infty}(\omega)y_k\overset{\sigma\mathrm{-weakly}}{\underset{k\to\infty}\longrightarrow} \pi_{\mathcal{G}}(a)U_{\mathcal{G}}(\omega).
$$

Then clearly also for all  $\omega \in M_*, a \in A$  and  $b \in B$  we have

$$
y_k^* \pi_{\mathcal{H}^{\infty}}(a) U_{\mathcal{H}^{\infty}}(\omega) \rho_{\mathcal{H}^{\infty}}(b) y_k = y_k^* \pi_{\mathcal{H}^{\infty}}(a) U_{\mathcal{H}^{\infty}}(\omega) y_k \rho_{\mathcal{G}}(b) \stackrel{\sigma-\text{weakly}}{\longrightarrow} \pi_{\mathcal{G}}(a) U_{\mathcal{G}}(\omega) \rho_{\mathcal{G}}(b).
$$

Since  $[\pi_u(A_s)C_{U_u}\rho_u(B_s)]=C^{\mathbb{G}}(A,B)$ , the uniform boundedness of the  $y_k$  ensures that for all  $z \in C^{\mathbb{G}}(A,B)$ we have

$$
y_k^* \theta_{\mathcal{H}^\infty}(z) y_k \stackrel{\sigma-\text{weakly}}{\longrightarrow} \theta_{\mathcal{G}}(z).
$$

Hence for our finite collection of  $\xi_i$  and  $z \in F_C$ , we have

$$
|\langle \xi_i, \theta_{\mathcal{G}}(z)\xi_j\rangle - \langle y_k\xi_i, \theta_{\mathcal{H}^\infty}(z)y_k\xi_j\rangle| \underset{k\to\infty}{\longrightarrow} 0,
$$

so we can take  $\xi'_i = y_k \xi_i$  for k large enough.

<span id="page-22-1"></span>Let us recuperate part of the proof of Theorem [3.7](#page-21-0) as a separate corollary.

**Corollary 3.8.** If G is equivariantly weakly contained in H, there exists a ucp map  $E : B(H) \to B(G)$  such that

<span id="page-22-0"></span>
$$
E(\pi_{\mathcal{H}}(a)) = \pi_{\mathcal{G}}(a), \qquad E(\rho_{\mathcal{H}}(b)) = \rho_{\mathcal{G}}(b), \qquad E(U_{\mathcal{H}}(\omega)) = U_{\mathcal{G}}(\omega) \qquad \forall a \in A, b \in B, \omega \in M_*. \tag{3.16}
$$

*Proof.* Since the map E of [\(3.12\)](#page-21-2) satisfies [\(3.13\)](#page-21-3), and since  $\pi_u(A)$ ,  $\rho_u(A) \subseteq QM(C^{\mathbb{G}}(A,B))$ , we immediately find that  $(3.16)$  is satisfied.

## 4. Amenability for equivariant correspondences and G-actions

Let A, B, C be von Neumann algebras. Given a ucp map  $E: A \rightarrow B$  (not necessarily normal!), we use the notation

$$
E \otimes \text{id} : A \bar{\otimes} C \to B \bar{\otimes} C \tag{4.1}
$$

to denote the unique ucp map such that

$$
E((\mathrm{id}\otimes\omega)z)=(\mathrm{id}\otimes\omega)((E\otimes\mathrm{id})z),\qquad\forall z\in A\bar{\otimes}C,\omega\in C_*,
$$

using for example the identification

 $B\bar{\otimes}C \cong CB(C_*, B),$ 

where the right hand side means the set of all cb maps [\[ER00,](#page-47-31) Proposition 8.1.2]. This allows us to give meaning to [\(0.6\)](#page-3-0), which we then refer to as G-equivariance of the corresponding ucp map.

**Definition 4.1.** Let  $\pi : A \rightarrow B$  be a G-equivariant normal unital  $*$ -homomorphism.

We say that  $\pi : A \to B$  is strongly equivariantly amenable if  $E_A = A L^2(A)$  is equivariantly weakly contained in  $_{\pi(A)}L^2(B)_{\pi(A)}$ , which we view as G-A-A-correspondence. More precisely,  $_{\pi(A)}L^2(B)_{\pi(A)}$  is the  $\mathbb{G}\text{-}A\text{-}A\text{-}correspondence (L^2(B), \pi_B \circ \pi : A \to B(L^2(B)), \rho_B \circ \pi : A \to B(L^2(B)), U_\beta).$ 

We say that  $\pi: A \to B$  is equivariantly amenable if there exists a G-equivariant ucp map

 $P : B \to A$ 

which is left inverse to  $\pi$ .

We may view equivariant amenability as a relative version of G-equivariant injectivity. Note also that if  $\pi$  is (strongly) equivariantly amenable, then  $\pi$  is automatically faithful, so we are actually dealing with a G-equivariant inclusion  $A \subseteq B$ .

Remark 4.2. The notion of equivariant amenability was already introduced in [\[Moa18,](#page-47-32) Definition 4.4] in the setting of discrete quantum group actions. Note that we follow here the terminology of [\[BMO20\]](#page-46-6), which is dual to the one followed for example in [\[A-D90\]](#page-46-4).

<span id="page-23-0"></span>**Theorem 4.3.** Let  $\pi : A \to B$  be a G-equivariant unital normal \*-homomorphism between G-W<sup>\*</sup>-algebras. If  $\pi$  is strongly G-amenable, then  $\pi : A \rightarrow B$  is G-amenable.

*Proof.* By assumption, the trivial equivariant A-A-correspondence  $L^2(A)$  is equivariantly weakly contained in the equivariant A-A-correspondence  $\pi(A) L^2(B) \pi(A)$ . Consider the ucp map

$$
E: B(L^2(B)) \to B(L^2(A))
$$

as in Corollary [3.8.](#page-22-1) Then  $E(\pi_B(B))$  will lie in  $\pi_A(A)$ , as these elements commute pointwise with  $\rho_A(A)$  by a multiplicative domain argument. We can hence uniquely define a ccp map  $P : B \to A$  by

$$
\pi_A(P(z)) := E(\pi_B(z)), \qquad z \in B.
$$

It is then clear that  $P(\pi(a)) = a$  for all  $a \in A$ . Moreover, we have by [\(3.16\)](#page-22-0) that  $U_{\beta}$ , the unitary Grepresentation for the A-A-correspondence  $_A L^2(B)_A$ , lies in the multiplicative domain of  $E \otimes id_M$ , hence for  $z \in B$  and  $\omega \in M_*$  we get

$$
\pi_A(P((id \otimes \omega)\beta(z))) = (id \otimes \omega)(E \otimes id)(U_{\beta}(\pi_B(z) \otimes 1)U_{\beta}^*)
$$
  
= 
$$
(id \otimes \omega)(U_{\alpha}(E(\pi_B(z)) \otimes 1)U_{\alpha}^*)
$$
  
= 
$$
\pi_A((id \otimes \omega)\alpha(P(z))),
$$

so that P is G-equivariant.

We can use the definition of (strong) amenability for equivariant normal unital ∗-homomorphisms to define a notion of left (strong) amenability for equivariant correspondences. Recall the constructions made in Example [2.4.](#page-9-0)

<span id="page-23-2"></span>**Definition 4.4.** We say that a G-equivariant A-B-correspondence  $(H, U, \pi, \rho)$  is (strongly) left G-amenable if  $\pi : A \to C = \rho(B)'$  is (strongly)  $\mathbb{G}\text{-}amenable$ .

Remark 4.5. In [\[A-D95,](#page-46-5) Definition 2.1], strong left amenability (called left amenability loc. cit.) is rather introduced in terms of the Connes fusion product. We use for now the shortcut provided by the remarks following [\[A-D95,](#page-46-5) Definition 2.1], and will return to this point in Remark [5.13.](#page-29-0)

**Remark 4.6.** If G is a multiplicity Hilbert space, then it is easily seen that  $(\mathcal{H}, U, \pi, \rho)$  is (strongly) left Gamenable if and only if  $(G \otimes H, id_G \otimes U, id_G \otimes \pi, id_G \otimes \rho)$  is (strongly) left G-amenable. Indeed, if  $\rho(B)' = C$ , then  $(id_{\mathcal{G}} \otimes \rho(B))' = B(\mathcal{G}) \bar{\otimes} C$ , and by Remark [3.2](#page-18-1) we know that

$$
E_A \preccurlyeq_{\pi(A)} L^2(C)_{\pi(A)} \quad \iff \quad E_A \preccurlyeq L^2(B(\mathcal{G})) \otimes_{\pi(A)} L^2(C)_{\pi(A)} =_{(\text{id}_\mathbb{G} \otimes \pi)(A)} L^2(B(\mathcal{G}) \bar{\otimes} C)_{(\text{id}_\mathbb{G} \otimes \pi)(A)},
$$

showing the claim for strong left G-amenability. For left G-amenability, we notice that if  $P: C \to A$  is a G-equivariant ucp map which splits  $\pi$ , and  $\omega$  an arbitrary normal state on  $B(\mathcal{G})$ , then

<span id="page-23-1"></span>
$$
\widetilde{P}: B(\mathcal{G}) \bar{\otimes} C \to A, \quad z \mapsto P((\omega \otimes id)z) \tag{4.2}
$$

is a G-equivariant ucp map which splits  $id_G \otimes \pi$ . Conversely, given [\(4.2\)](#page-23-1), the map

$$
P: C \to A, \quad c \mapsto P(\mathrm{id}_{\mathcal{G}} \otimes c)
$$

will be a G-equivariant ucp map which splits  $\pi$ . This shows that left G-equivariant (non-)amenability is unperturbed by amplification.

<span id="page-24-2"></span>**Theorem 4.7.** If H is strongly left equivariantly amenable, then it is left equivariantly amenable.

*Proof.* This is a direct consequence of Theorem [4.3.](#page-23-0)

In [\[BMO20,](#page-46-6) Corollary A.2], it was proven that in the non-equivariant setting, the above can be strengthened: strong left amenability of  $H$  is in fact *equivalent* with left amenability. In the equivariant setting, this is currently yet unknown, as the following special situation demonstrates.

**Definition 4.8.** Let  $(A, \alpha)$  be a G-dynamical von Neumann algebra. We say that the action of G on A is (strongly) G-amenable if the inclusion  $\alpha : A \to A \bar{\otimes} M$  is (strongly) G-amenable.

Concretely, recalling the notation introduced in Example [2.6,](#page-9-1) strong G-amenability of  $\alpha$  means that  $E_A^{\mathbb{G}}$  is G-equivariantly weakly contained in  $S_A^{\mathbb{G}}$  and G-amenability of  $\alpha$  means that there exists a ucp map

$$
P: A \bar{\otimes} M \to A \tag{4.3}
$$

such that  $P(\alpha(a)) = a$  for all  $a \in A$  and

$$
P((\mathrm{id}\otimes(\mathrm{id}\otimes\omega)\Delta)z)=(\mathrm{id}\otimes\omega)\alpha(P(z)),\qquad\forall z\in A\bar{\otimes}M,\omega\in M_*.\tag{4.4}
$$

It is not clear to us if the definition of G-amenability has explicitly appeared in print before in the general setting of locally compact quantum groups, but it is the obvious one, see e.g. [\[Moa18,](#page-47-32) Definition 4.1] for the setting with G discrete, or [\[Voi77,](#page-47-8) [Rua96,](#page-47-9) [BT03,](#page-46-27) [DQV02\]](#page-46-10) when  $A = \mathbb{C}$ . On the other hand, for  $\mathbb{G} = G$  and ordinary locally compact group, the above is equivalent with the existence of a ucp map

$$
P: A\bar{\otimes}L^{\infty}(G) \to A \tag{4.5}
$$

such that  $P(a \otimes 1) = a$  for all  $a \in A$  and which is equivariant with respect to the diagonal action of G on  $A\bar{\otimes}L^{\infty}(G)$ , using the action  $\rho$  of G on  $L^{\infty}(G)$  by right shifts. This is the notion of amenability of the action  $G \curvearrowright A$  as it appears in [\[A-D79,](#page-46-7) Definition [3](#page-24-1).4].<sup>3</sup>

<span id="page-24-3"></span>Specializing Theorem [4.3,](#page-23-0) we obtain:

Theorem 4.9. Assume A is a von Neumann algebra with a strongly G-amenable G-action. Then the action of  $G$  on  $A$  is  $G$ -amenable.

*Proof.* This is just a special case of Theorem [4.3.](#page-23-0)

If we take  $A = \mathbb{C}$ , then the above simply states that if the trivial G-representation is weakly contained in the regular G-representation (= co-amenability of  $\hat{\mathbb{G}}$  [\[BT03\]](#page-46-27)), there exists a G-invariant state  $M \to \mathbb{C}$  (= amenability of G). The converse is a well-known open problem, making it hence difficult to ascertain if the converse of Theorem [4.7](#page-24-2) or even Theorem [4.9](#page-24-3) will hold. Let us now show that the converse of the result in Theorem [4.9](#page-24-3) holds for locally compact groups.

<span id="page-24-0"></span>**Theorem 4.10.** If G is a locally compact group, and A a G-W<sup>\*</sup>-algebra, then the action of G on A is strongly G-amenable if and only if it is G-amenable.

*Proof.* Note first that an action of  $G$  on  $A$  is the same thing as a coaction

$$
\alpha: A \to A \bar{\otimes} L^{\infty}(G), \qquad a \mapsto \alpha(a),
$$

where we view  $\alpha(a) \in B(L^2(A) \otimes L^2(G)) = B(L^2(G, L^2(A)))$  as multiplication by the function  $g \mapsto$  $\pi_A(\alpha_g(a))$ . Note that  $M = L^{\infty}(G)$  is endowed with the structure of a locally compact quantum group by putting  $\Delta(f)(q, h) = f(qh)$  for  $f \in L^{\infty}(G)$  (this definition makes sense locally almost everywhere).

By Theorem [4.9,](#page-24-3) it is enough to prove that G-amenability of the action  $G \cap A$  implies strong amenability, i.e. the weak containment of  $E_A$  in the G-semi-coarse correspondence  $S_A^G$ .

<span id="page-24-1"></span> ${}^{3}\text{In}$  [\[A-D79\]](#page-46-7) one uses left shifts, but this is of course inconsequential.

Let  $V_g$  for  $g \in G$  denote the right shift of g on  $L^2(G)$  (so we work with respect to dg the right invariant Haar measure). Note that this is simply the  $G$ -representation corresponding to the right regular representation  $V$ of  $L^{\infty}(G)$  as a locally compact quantum group. Similarly, we write  $g \mapsto U_g$  for the unitary G-representation associated to  $U_{\alpha}$ . Let  $Z(A)$  be the center of A, and let  $Z(A)_{\alpha}$  be as in Definition [2.12.](#page-11-4) In this case,  $Z(A)_{\alpha}$ is actually the C<sup>∗</sup> -algebra of elements which are norm-continuous for the G-action [\[Ped79,](#page-47-19) Lemma 7.5.1]. Let then  $C_c(G, Z(A_s))$  be the ∗-algebra of norm-continuous maps  $G \to Z(A_s)$  with compact support. By [\[BC22,](#page-46-19) Theorem 1.1.(1)  $\Rightarrow$  (2)], we can choose a net

$$
x_i \in C_c(G, Z(A)_s) \subseteq Z(A) \bar{\otimes} L^2(G) \subseteq B(L^2(A), L^2(A) \otimes L^2(G))
$$

which is contractive for the  $B(L^2(A), L^2(A) \otimes L^2(G))$ -norm and such that

$$
x_i^*(\alpha_g \otimes V_g)(x_i) \xrightarrow[i \to \infty]{\sigma-\text{weakly}} 1
$$
\n(4.6)

uniformly on compacts. If we define  $z_i(g) = x_i^*(\alpha_g \otimes V_g)(x_i)$ , it is elementary to verify that  $z_i \in C_c(G, Z(A)_s)$ , and that hence

$$
\int_G f(g) z_i(g) \, dg \xrightarrow[i \to \infty]{\sigma-\text{weakly}} \left( \int_G f(g) \, dg \right) 1
$$

for each  $f \in C_c(G)$ . A density argument now allows to conclude that

<span id="page-25-0"></span>
$$
(x_i^* \otimes 1)V_{23}(\alpha \otimes id)(x_i)_{132} \xrightarrow[i \to \infty]{\sigma-\text{weakly}} 1.
$$
 (4.7)

Put  $y_i = U_\alpha x_i$ . Then  $y_i \in X := X_A(L^2(A) \otimes L^2(G))$  (using the right A-representation as in Example [2.6\)](#page-9-1). Moreover, the identity  $U_{\alpha,12}^* V_{23} U_{\alpha,12} = U_{\alpha,12} V_{23}$  transforms [\(4.7\)](#page-25-0) into

<span id="page-25-1"></span>
$$
(y_i^* \otimes 1)\alpha_X(y_i) \stackrel{\sigma\text{-weakly}}{\longrightarrow} 1. \tag{4.8}
$$

But since the first leg of  $x_i$  lies in  $Z(A)$ , multiplying [\(4.8\)](#page-25-1) to the left with  $\pi_A(a) \otimes 1$  for  $a \in A$  entails that

$$
(y_i^*(\pi_A \otimes \pi_M)\alpha(a) \otimes 1)\alpha_X(y_i) \underset{i \to \infty}{\overset{\sigma-\text{weakly}}{\longrightarrow}} (\pi_A \otimes \pi_M)\alpha(a) \otimes 1, \qquad \forall a \in A. \tag{4.9}
$$

It now follows from Theorem [3.7](#page-21-0) that  $E_A$  is weakly contained in  $S_A^G$  $A$ .

To end this section, we note that Theorem [3.7](#page-21-0) seems to allow for a stronger and more symmetric condition of weak containment. Namely, let  $A, B$  be  $\mathbb{G}\text{-}W^*$ -algebras, and let  $\mathcal{H}, \mathcal{G}$  be  $\mathbb{G}\text{-}A\text{-}B\text{-}correspondences$ . Denote

$$
{}_AX_B(\mathcal{G}, \mathcal{H}) = \{ x \in X_B(\mathcal{G}, \mathcal{H}) \mid x \pi_{\mathcal{G}}(a) = \pi_{\mathcal{H}}(a)x \text{ for all } a \in A \}.
$$

Let us then say that H symmetrically weakly contains G if there exists a contractive net  $y_i \in A X_B(G, H^{\infty})$ such that

<span id="page-25-2"></span>
$$
(y_i^* \otimes 1)\alpha_X(y_i) \to 1 \quad \sigma\text{-weakly in } B(\mathcal{G} \otimes L^2(\mathbb{G})).
$$
\n(4.10)

Clearly, if H symmetrically weakly contains  $\mathcal{G}$ , then it weakly contains  $\mathcal{G}$ , but in the characterisation of weak containment in Theorem [3.7,](#page-21-0) the  $x_i$  only satisfy  $x_i\pi_G(a) = \pi_{\mathcal{H}}(a)x_i$  approximately.

If we can find  $y_i \in A X_B(G, \mathcal{H}^\infty)$  satisfying [\(4.10\)](#page-25-2), then it is easily seen that  $A \mathcal{H}_B^\infty$  must contain a (not necessarily equivariant) isometric copy of  $_{A}G_{B}$  as an A-B-correspondence. It is hence clear that symmetric weak containment is in general much stronger than ordinary weak containment, and already puts a very strong condition on the underlying correspondences. However, Theorem [4.10](#page-24-0) shows that these notions do coincide when considering the weak inclusion of  $E_A$  inside the G-semi-coarse correspondence  $S_A^G$  for G a locally compact group. We do not know if, for a general locally compact quantum group G, the equivariant weak containment  $E_A \preccurlyeq S_A^{\mathbb{G}}$  will imply the symmetric equivariant weak containment. Note that the latter, when framed appropriately in the C<sup>\*</sup>-context, coincides with how amenability is defined for actions of discrete quantum groups on unital C<sup>∗</sup>-algebras in [\[VV07,](#page-47-33) Definition 4.1].

## <span id="page-25-3"></span>5. Operations on equivariant correspondences and the Takesaki-Takai duality

In this section, we consider particular constructions which can be performed on equivariant correspondences.

5.1. Opposites of G-equivariant correspondences. Fix a locally compact quantum group  $\mathbb{G} = (M, \Delta)$ and  $\mathbb{G}\text{-}W^*$ -algebras  $A, B$ .

<span id="page-26-0"></span>**Definition 5.1.** Let H be a G-equivariant A-B-correspondence. Let  $\overline{\mathcal{H}}$  be the complex conjugate Hilbert space, and let  $C_H : \mathcal{H} \to \overline{\mathcal{H}}$  be the canonical anti-unitary conjugation map  $\xi \mapsto \overline{\xi}$ . The opposite B-Acorrespondence is obtained by endowing  $\overline{\mathcal{H}}$  with the bimodule structure

$$
\overline{\pi}(b)=C_{\mathcal{H}}\rho(b^*)C_{\overline{\mathcal{H}}},\qquad \overline{\rho}(a)=C_{\mathcal{H}}\pi(a^*)C_{\overline{\mathcal{H}}}
$$

and the unitary G-representation

$$
\overline{U} = (C_{\mathcal{H}} \otimes \check{J})U^*(C_{\overline{\mathcal{H}}} \otimes \check{J}).
$$

We can easily see that  $\overline{\mathcal{H}}$  is again a G-equivariant correspondence, using the defining relations of the Gequivariant correspondence  $\mathcal{H}$  and the fact that  $\tilde{J}$  implements the unitary antipode.

Let us consider this operation for the examples appearing in Section [2.](#page-8-3) Of course, we may in Definition [5.1](#page-26-0) use any anti-unitary operator  $C_H$  from H to (another or the same) Hilbert space, as this will clearly give an equivariant copy.

<span id="page-26-2"></span>**Example 5.2.** We have  $\overline{E_A} \cong E_A$  by taking  $C_{\mathcal{H}} = J_A$  and using [\(1.8\)](#page-7-2).

<span id="page-26-3"></span>**Example 5.3.** Let  $A \subseteq C$  and  $B \subseteq D$  be G-equivariant inclusions of G-W<sup>\*</sup>-algebras, and  $_C \mathcal{H}_D$  a Gequivariant C-D-correspondence. Then

$$
{}_B \text{Res}_A(\overline{\mathcal{H}}) = \overline{{}_A \text{Res}_B(\mathcal{H})}.
$$

**Example 5.4.** With  $D^{\mathbb{G}}(-)$  the adjoint amplification, we see that  $D^{\mathbb{G}}(\overline{\mathcal{H}}) \cong \overline{D^{\mathbb{G}}(\mathcal{H})}$ . Indeed, implementing the latter by means of the anti-unitary map

$$
C_{D^{\mathbb{G}}(\mathcal{H})}=C_{\mathcal{H}}\otimes \check{J}:\mathcal{H}\otimes L^{2}(\mathbb{G})\to \overline{\mathcal{H}}\otimes L^{2}(\mathbb{G}),
$$

we find the unitary intertwiner

$$
(C_{\mathcal{H}} \otimes \check{J})U(C_{\overline{\mathcal{H}}} \otimes \check{J}): D^{\mathbb{G}}(\overline{\mathcal{H}}) \to \overline{D^{\mathbb{G}}(\mathcal{H})},
$$

using relation [\(2.2\)](#page-10-1) for the intertwining properties of the unitary G-representations.

It is easy to see that the statement in the previous example also holds for the regular amplification.

**Example 5.5.** We have  $\overline{C_{A,B}^{\mathbb{G}}}\cong C_{B,A}^{\mathbb{G}}$ . Indeed, implementing  $C_{A,B}^{\mathbb{G}}$  using the anti-unitary  $J_A\otimes J\otimes J_B$ , one verifies that the flip

$$
L^2(A) \otimes L^2(\mathbb{G}) \otimes L^2(B) \cong L^2(B) \otimes L^2(\mathbb{G}) \otimes L^2(A)
$$

defines the desired unitary intertwiner.

5.2. Composition of  $\mathbb{G}$ -equivariant correspondences. Let  $A, B$  be von Neumann algebras. Recall that  $X_B(\mathcal{H})$  was introduced in [\(3.2\)](#page-19-4) as the self-dual Hilbert A-B-module of right B-intertwiners from  $L^2(B)$  to H. If then G is a B-C-correspondence, we can compose H and G into an A-C-correspondence by the Connes fusion product

$$
_A(\mathcal{H} \boxtimes \mathcal{G})_C := {}_A(X_B(\mathcal{H}) \underset{B}{\otimes} \mathcal{G})_C,
$$

which is the separation-completion of  $X_B(\mathcal{H}) \odot_B \mathcal{G}$  with respect to the semi-inner product

$$
\langle x \otimes_B \xi, y \otimes_B \eta \rangle = \langle \xi, x^* y \eta \rangle, \qquad x, y \in X_B(\mathcal{H}), \xi, \eta \in \mathcal{G},
$$

(where we identified  $B \cong \pi_B(B)$ ). The A-C-bimodule structure is determined by

$$
\pi_{\boxtimes}(a)(x\otimes_B \xi) = \pi_{\mathcal{H}}(a)x\otimes_B \xi, \qquad \rho_{\boxtimes}(c)(x\otimes_B \xi) = x\otimes_B \rho_{\mathcal{G}}(c)\xi, \qquad a\in A, c\in C, x\in X_B(\mathcal{H}), \xi\in\mathcal{G}.
$$

We write  $\mathcal{H} \boxtimes_B \mathcal{G}$  if we want to emphasize that we take the fusion product over B.

If  $K$  is a C-D-correspondence, we have a natural intertwining associativity unitary

<span id="page-26-1"></span>
$$
(\mathcal{H} \boxtimes \mathcal{G}) \boxtimes \mathcal{K} \cong \mathcal{H} \boxtimes (\mathcal{G} \boxtimes \mathcal{K}), \tag{5.1}
$$

using e.g. that  $X_C(\mathcal{H} \boxtimes \mathcal{G})$  is the  $\sigma$ -weak closure of the image of  $X_B(\mathcal{H}) \odot_B X_C(\mathcal{G})$  in  $B(L^2(C), \mathcal{H} \boxtimes \mathcal{G})$  by

$$
(x \otimes_B y)\eta = x \otimes_B y\eta
$$
,  $x \in X_B(\mathcal{H}), y \in X_C(\mathcal{G}), \eta \in L^2(C)$ .

The identity correspondence  $E_A = {}_A L^2(A)_A$  acts as the identity for this composition. See e.g. [\[Lan01,](#page-47-34)[Bro03\]](#page-46-28) for a detailed treatment.

We can upgrade the above construction to the G-equivariant setting. Recall that if  $(A, \alpha)$  and  $(B, \beta)$  are  $\mathbb{G}\text{-}W^*$ -algebras and  $\mathcal{H}$  a  $\mathbb{G}\text{-}$  equivariant A-B-correspondence, then  $X = X_B(\mathcal{H})$  comes equipped with the coaction

$$
\alpha_X = \alpha_{X_B(\mathcal{H})} : X \to X \bar{\otimes} M, \qquad x \mapsto U(x \otimes 1)U_{\beta}^*.
$$

<span id="page-27-2"></span>**Proposition 5.6.** Assume  $(A, \alpha), (B, \beta), (C, \gamma)$  are G-dynamical von Neumann algebras. If H is a Gequivariant A-B-correspondence and G a G-equivariant B-C-correspondence, we can endow  $\mathcal{H} \boxtimes \mathcal{G}$  with a unique unitary  $\mathbb{G}$ -representation  $U_{\boxtimes}$  such that

<span id="page-27-0"></span>
$$
U_{\boxtimes}((x \otimes_B \xi) \otimes \eta) := \alpha_X(x) \otimes_{B \bar{\otimes} M} U_{\mathcal{G}}(\xi \otimes \eta), \qquad x \in X_B(\mathcal{H}), \xi \in \mathcal{H}, \eta \in \mathcal{G}.
$$
 (5.2)

Then  $H \boxtimes G$  becomes a G-equivariant A-C-correspondence.

*Proof.* First of all, note that in [\(5.2\)](#page-27-0) one rather defines  $U_{\boxtimes}$  as a map

$$
(X_B(\mathcal{H}) \odot_B \mathcal{G}) \odot L^2(M) \rightarrow (X_B(\mathcal{H}) \bar{\otimes} M) \underset{B \bar{\otimes} M}{\otimes} (\mathcal{G} \otimes L^2(M)) \cong (X_B(\mathcal{H}) \underset{B}{\otimes} \mathcal{G}) \otimes L^2(M) = (\mathcal{H} \boxtimes \mathcal{G}) \otimes L^2(M).
$$

A direct computation shows that this indeed descends to an isometry

 $U_{\boxtimes} : (\mathcal{H} \boxtimes \mathcal{G}) \otimes L^2(M) \rightarrow (\mathcal{H} \boxtimes \mathcal{G}) \otimes L^2(M).$ 

By the bicommutant theorem, it follows straightforwardly that  $U_{\boxtimes} \in B(H \boxtimes \mathcal{G}) \bar{\otimes} M$ .

To check the corepresentation property, various identifications of Hilbert spaces have to be made, which we will not comment on explicitly. The main goal is to prove the identity

<span id="page-27-1"></span>
$$
U_{\boxtimes,13}U_{\boxtimes,12} = W_{32}^*U_{\boxtimes,12}W_{32},\tag{5.3}
$$

which we will show by applying both sides to a vector  $(x \otimes_B \xi) \otimes \eta \otimes \zeta$  with  $x \in X_B(\mathcal{H}), \xi \in \mathcal{G}$  and  $\eta, \eta \in L^2(\mathbb{G})$ . Then on the one hand, the left hand side gives

$$
U_{\boxtimes,13}U_{\boxtimes,12}((x \otimes_B \xi) \otimes \eta \otimes \zeta) \cong U_{\boxtimes,13}((\alpha_X(x) \otimes_{B \bar{\otimes} M} U_{\mathcal{G}}(\xi \otimes \eta)) \otimes \zeta)
$$
  

$$
\cong (\alpha_X \otimes \mathrm{id}) \alpha_X(x)_{132} \otimes_{B \bar{\otimes} M \bar{\otimes} M} (U_{\mathcal{G},13} U_{\mathcal{G},12}(\xi \otimes \eta \otimes \zeta)),
$$

while for the right hand side we get

$$
W_{32}^* U_{\boxtimes,12} W_{32}((x \otimes_B \xi) \otimes \eta \otimes \zeta) = W_{32}^* U_{\boxtimes,12}((x \otimes_B \xi) \otimes W_{21}(\eta \otimes \zeta))
$$
  
\n
$$
\cong (\mathrm{id} \otimes \Delta^{\mathrm{op}}) \alpha_X(x) \otimes_{B \bar{\otimes} M \bar{\otimes} M} W_{32}^* U_{\mathcal{G},12} W_{32}(\xi \otimes \eta \otimes \zeta))
$$
  
\n
$$
= (\mathrm{id} \otimes \Delta^{\mathrm{op}}) \alpha_X(x) \otimes_{B \bar{\otimes} M \bar{\otimes} M} (\mathrm{id} \otimes \Delta^{\mathrm{op}})(U_{\mathcal{G}})(\xi \otimes \eta \otimes \zeta)),
$$

so that [\(5.3\)](#page-27-1) now follows from the coaction property for  $\alpha_X$  and the corepresentation property of  $U_g$ .

One can now see that  $U_{\boxtimes}$  must be unitary since any isometric corepresentation of a locally compact quantum group is unitary, by [\[BDS13,](#page-46-22) Corollary 4.16]. Alternatively, one could more directly observe that the range of  $U_{\boxtimes}$  will contain all vectors of the form  $\alpha_X(x)(1\otimes m)\otimes_{B\bar{\otimes}M} (\xi\otimes \eta)$  for  $x\in X_B(H), m\in M$  and  $\xi\in H, \eta\in \mathcal{G},$ and then conclude from the fact that the linear span of the  $\alpha_X(x)(1 \otimes m)$  is  $\sigma$ -strongly dense in  $X_B(\mathcal{H})\bar{\otimes}M$ .

The equivariance conditions

$$
U_{\boxtimes}(\pi_{\boxtimes}(a)\otimes 1)=(\pi_{\boxtimes}\otimes \pi_M)\alpha(a)U_{\boxtimes}, \qquad (\rho_{\boxtimes}(c)\otimes 1)U_{\boxtimes}=U_{\boxtimes}(\rho_{\boxtimes}\otimes \pi_M R)\gamma(c), \qquad a\in A, c\in C
$$
  
are easily checked.

By using similar identifications as in the proof of Proposition [5.6,](#page-27-2) one can check that the associativity intertwiner [\(5.1\)](#page-26-1) is an intertwiner of G-equivariant correspondences. We refrain from spelling out the details here.

The following lemma is convenient for computations.

<span id="page-28-4"></span>**Lemma 5.7.** Assume  $(\mathcal{H} \otimes L^2(B), U, \pi, \rho)$  is endowed with the structure of a G-A-B-correspondence such that

$$
\rho(b)(\xi \otimes \eta) = \xi \otimes \rho_B(b)\eta, \qquad U = U_{\mathcal{H},13}U_{\beta,23}
$$

for some unitary G-representation  $U_{\mathcal{H}}$  on  $\mathcal{H}$ . If  $(\mathcal{G}, U_{\mathcal{G}}, \pi_{\mathcal{G}}, \rho_{\mathcal{G}})$  is a G-equivariant B-C-correspondence, we have a unitary intertwiner

$$
(\mathcal{H}\otimes L^2(B))\boxtimes\mathcal{G}\cong\mathcal{H}\otimes\mathcal{G},
$$

where the right hand side is endowed with the G-A-C-correspondence

<span id="page-28-0"></span>
$$
\pi_{\boxtimes}(a) = (\mathrm{id} \otimes \pi_{\mathcal{G}})\pi(a), \qquad \rho_{\boxtimes}(c) = 1 \otimes \rho_{\mathcal{G}}(c), \qquad a \in A, c \in C,
$$
\n
$$
U_{\boxtimes} = U_{\mathcal{H},13}U_{\mathcal{G},23}.
$$
\n
$$
(5.4)
$$

*Proof.* Note that the assumption on  $\rho$  implies that  $\pi(a) \in B(H) \bar{\otimes} \pi_B(B)$  for  $a \in A$ , so under the identification  $\pi_B(B) \cong B$  we see that  $\pi_{\boxtimes}$  in [\(5.4\)](#page-28-0) is at least a well-defined unital normal \*-homomorphism  $A \to B(\mathcal{H} \otimes \mathcal{G})$ .

Now it is immediate that  $X_B(\mathcal{H}\otimes L^2(B)) = \mathcal{H}\overline{\otimes}B$  with its obvious self-dual Hilbert A-B-bimodule structure, and with the coaction given by

$$
\alpha_X(\xi \otimes b) = U_{\mathcal{H},13}(\xi \otimes \beta(b)).
$$

We hence obtain an isometric map with dense image

$$
(\mathcal{H} \odot B) \odot_B \mathcal{G} \rightarrow \mathcal{H} \otimes \mathcal{G}, \qquad (\xi \otimes b) \otimes \zeta \mapsto \xi \otimes b\zeta.
$$

It is clear that this induces [\(5.4\)](#page-28-0) on  $\mathcal{H} \otimes \mathcal{G}$ . Also the form for  $U_{\boxtimes}$  is immediate.

**Definition 5.8.** Let  $A, B$  be  $\mathbb{G}\text{-}W^*$ -algebras. We say that  $A$  is  $\mathbb{G}\text{-}$  equivariantly W<sup>\*</sup>-Morita equivalent with B if there exists a G-equivariant A-B-correspondence  $(\mathcal{H}, U, \pi, \rho)$  which is full (or equivalently, faithful) as a right  $B$ -module<sup>[4](#page-28-1)</sup> and faithful as a left  $A$ -module, and such that

$$
\pi(A) = \rho(B)'
$$

In this case, we call  $H$  a  $\mathbb{G}-W^*$ -Morita correspondence (between A and B).

If H is a G-W<sup>\*</sup>-Morita correspondence between A and B, we put (recalling the notation  $(0.4)$ )

$$
Q_{\mathcal{H}}:=X_{B}(L^{2}(B)\oplus \mathcal{H}, L^{2}(B)\oplus \mathcal{H}).
$$

It follows from basic theory of correspondences [\[Tak03\]](#page-47-17) that we have a decomposition

$$
Q_{\mathcal{H}} \cong \begin{pmatrix} A & X_B(\mathcal{H}) \\ X_B(\mathcal{H})^* & B \end{pmatrix},
$$

and that we can then decompose

<span id="page-28-2"></span>
$$
L^2(Q_{\mathcal{H}}) = \begin{pmatrix} L^2(A) & L^2(X_B(\mathcal{H})) \\ L^2(X_B(\mathcal{H})^*) & L^2(B) \end{pmatrix} \cong \begin{pmatrix} L^2(A) & \mathcal{H} \\ \overline{\mathcal{H}} & L^2(B) \end{pmatrix}
$$
(5.5)

as an L<sup>2</sup>-direct sum. The isomorphism  $L^2(X_B(\mathcal{H})) \cong \mathcal{H}$  is uniquely determined by  $\pi_Q(x)\xi \cong x\xi$  if  $x \in X_B(\mathcal{H})$ and  $\xi \in L^2(B)$ . The isomorphism  $L^2(X_B(\mathcal{H})^*) \cong \overline{\mathcal{H}}$  is uniquely determined by  $J_Q\xi \cong \overline{\xi}$  for  $\xi \in \mathcal{H}$ .

In the sequel, we simply write  $Q = Q_H$ . From the fact that we are in the equivariant setting, Q obtains a natural G-action  $\alpha_Q$  on Q by conjugating with the unitary representation  $\hat{U}_{\beta} \oplus U$  on  $L^2(B) \oplus \mathcal{H}$ . Let  $U_Q$ be the associated canonical unitary implementing  $\alpha_Q$ .

Lemma 5.9. Under the identification [\(5.5\)](#page-28-2), we have

<span id="page-28-3"></span>
$$
U_Q \cong \begin{pmatrix} U_\alpha & U \\ \overline{U} & U_\beta \end{pmatrix} . \tag{5.6}
$$

<span id="page-28-1"></span><sup>4</sup>Concretely, this means that  $\pi_B(B) = [X_B(\mathcal{H})^* X_B(\mathcal{H})]^{\sigma\text{-weak}}$ .

Proof. It is clear that there will be a decomposition

$$
U_Q\cong \begin{pmatrix}U_{11}&U_{12}\\U_{21}&U_{22}\end{pmatrix},
$$

as  $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$  and  $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$  are invariant under  $\alpha_Q$ . The identifications on the diagonal of [\(5.6\)](#page-28-3) then follow by noticing that the construction of the unitary G-representation in [\[Vae05\]](#page-47-14) is compatible with the (G-invariant) projection maps  $L^2(Q) \to L^2(B)$  and  $L^2(Q) \to L^2(A)$ .

As  $U_Q$  implements  $\alpha_Q$ , we must have for  $x \in X_B(\mathcal{H})$  and  $\xi \in L^2(B)$ ,  $\eta \in L^2(\mathbb{G})$  that

$$
U_Q(x\xi\otimes\eta)=\alpha_X(x)U_\beta(\xi\otimes\eta)\cong U(x\xi\otimes\eta),
$$

showing the isomorphism in the top right corner of [\(5.6\)](#page-28-3). The isomorphism on the lower left corner is deduced from the observation that

$$
U_Q(J_Q\xi\otimes J\eta)=(J_Q\otimes J)U^*(\xi\otimes \eta)\cong (C_{\mathcal{H}}\otimes J)U^*(\xi\otimes \eta)=\overline{U}(C_{\mathcal{H}}\xi\otimes J\eta),\qquad \xi\in \mathcal{H},\eta\in L^2(\mathbb{G}).
$$

<span id="page-29-2"></span>**Proposition 5.10.** If  $H$  is a  $\mathbb{G}-W^*$ -Morita correspondence between A and B, then

<span id="page-29-1"></span> $\mathcal{H}\boxtimes \overline{\mathcal{H}}\cong L^2(A),\qquad \overline{\mathcal{H}}\boxtimes \mathcal{H}\cong L^2$  $(5.7)$ 

Proof. We can construct the maps in  $(5.7)$  as equivariant isometries simply by restriction of the natural unitary G-equivariant intertwiners

$$
L^2(Q) \boxtimes_Q L^2(Q) \cong L^2(Q) \cong L^2(Q) \boxtimes_Q L^2(Q).
$$

It is then well-known from the theory of (non-equivariant) correspondences that these restrictions are still unitary (cf. [\[Sau83,](#page-47-35) Proposition 3.1]).

<span id="page-29-3"></span>Corollary 5.11. If  $A, B, C$  are  $\mathbb{G}-W^*$ -algebras,  $\mathcal{H}$  is a  $\mathbb{G}$ -equivariant A-B-correspondence and  $\mathcal{G}$  is a  $\mathbb{G}$ equivariant B-C-correspondence, then

$$
\overline{\mathcal{H}\boxtimes\mathcal{G}}\cong\overline{\mathcal{G}}\boxtimes\overline{\mathcal{H}},\tag{5.8}
$$

*Proof.* We may replace G by  $\overline{G}$ , for G a C-B-correspondence. By considering  $A \oplus C$  and  $H \oplus G$ , we may moreover assume that  $A = C$  and  $\mathcal{H} = \mathcal{G}$ . Furthermore, we can then as well assume that  $\rho_{\mathcal{H}}$  is faithful, and  $A = \rho_{\mathcal{H}}(B)'$ . In other words, H is now a G-W<sup>\*</sup>-Morita correspondence between A and B, and the result follows from Proposition [5.10.](#page-29-2)

**Corollary 5.12.** Let  $A \subseteq B$  be an equivariant inclusion, and put  $C = \rho_B(A)'$ . Then C is G-equivariant by  $\gamma(c) := U_{\beta}(c \otimes 1)U_{\beta}^*, \qquad c \in C,$ 

and we obtain a  $\mathbb{G}\text{-}W^*$ -Morita C-A-correspondence  ${}_{C}L^2(B)_A$ . We then have

$$
{}_BL^2(B)_A \boxtimes {} _AL^2(B)_B \cong {} _BL^2(C)_B.
$$

*Proof.* The result is just a particular case of Proposition [5.10.](#page-29-2)

<span id="page-29-0"></span>**Remark 5.13.** More generally, we have that if H is a  $\mathbb{G}$ -equivariant A-B-correspondence and  $C = \rho(B)$ ', we can identify  $_A\mathcal{H} \boxtimes \overline{\mathcal{H}}_A \cong {}_A L^2(C)_A$ . The proof easily reduces to the above case.

Let us look at some more particular examples. We again refer back to Section [2](#page-8-3) for the notations we use.

<span id="page-29-4"></span>**Example 5.14.** Using Lemma [5.7](#page-28-4) and Corollary [5.11,](#page-29-3) it is easily seen that  $E_A^{\mathbb{G}}$  acts as an identity under  $\boxtimes$ : for any  $\mathcal{H} \in \text{Corr}^{\mathbb{G}}(A, B)$  we obtain canonical isomorphisms

$$
E_A^{\mathbb{G}} \boxtimes \mathcal{H} \cong \mathcal{H} \cong \mathcal{H} \boxtimes E_B^{\mathbb{G}}.
$$

<span id="page-30-0"></span>**Example 5.15.** If  $A \subseteq B$  is an equivariant inclusion and H is a G-equivariant B-C-correspondence, we have

$$
{}_{A}L^{2}(B)_{B}\boxtimes {}_{B}\mathcal{H}_{C}\cong {}_{A}\mathcal{H}_{C}.
$$

Similarly, if  $H$  is a  $\mathbb{G}$ -equivariant C-B-correspondence, then

$$
{}_C\mathcal{H}_B \boxtimes {}_B L^2(B)_A \cong {}_C\mathcal{H}_A.
$$

<span id="page-30-1"></span>Example 5.16. We can identify

$$
S_{A,B}^{\mathbb{G}} \cong {}_{A}L^{2}(A)_{\mathbb{C}} \boxtimes {}_{\mathbb{C}}L^{2}(B)_{B},
$$

<span id="page-30-3"></span>as follows directly by definition of the Connes fusion product.

**Example 5.17.** Combining Example [5.14,](#page-29-4) Example [5.15](#page-30-0) and Example [5.16](#page-30-1) with the associativity of  $\boxtimes$ , we see that we can identify

$$
S_{A,B}^{\mathbb{G}} \boxtimes S_B^{\mathbb{G}} \cong L^2(A) \otimes L^2(B) \otimes L^2(\mathbb{G}),
$$

where the latter is endowed with the normal (anti-)representations

$$
\pi(a) = \pi_A(a) \otimes 1 \otimes 1, \qquad \rho(b) = 1 \otimes (\rho_B \otimes \rho_M)\beta(b), \qquad a \in A, b \in B,
$$

and the unitary G-representation  $U_{\alpha,14}V_{34}$ . Conjugating with  $\Sigma_{23}U_{\alpha,13}$ , we find that

$$
S_{A,B}^{\mathbb{G}} \boxtimes S_B^{\mathbb{G}} \cong C_{A,B}^{\mathbb{G}}.
$$

It follows from the computations in Example [5.2](#page-26-2) and Corollary [5.11](#page-29-3) that also

$$
S_A^{\mathbb{G}} \boxtimes S_{A,B}^{\mathbb{G}} \cong C_{A,B}^{\mathbb{G}}.
$$

Example 5.18. Similar as in the previous Example, we obtain that

$$
S_{A,B}^{\mathbb{G}} \boxtimes S_{B,C}^{\mathbb{G}} \cong L^{2}(A) \otimes L^{2}(B) \otimes L^{2}(C)
$$

endowed with the A-C-correspondence by

$$
\pi(a) = \pi_A(a) \otimes 1 \otimes 1, \qquad \rho(c) = 1 \otimes 1 \otimes \rho_C(c), \qquad a \in A, c \in C,
$$

and the unitary G-representation  $U_{\alpha,14}U_{\beta,24}U_{\gamma,34}$ . In particular, if we take  $B = M$  with  $\beta = \Delta$ , we see that

$$
S_{A,M}^{\mathbb{G}} \boxtimes S_{M,C}^{\mathbb{G}} \cong C_{A,C}^{\mathbb{G}}.
$$

Combining this with Example [5.3,](#page-26-3) we can write this as

$$
S_{A,M}^{\mathbb G}\boxtimes \overline{S_{C,M}^{\mathbb G}}\cong C_{A,C}^{\mathbb G}.
$$

**Example 5.19.** Consider the G-regular G-A-A-correspondence  $S^A_{\mathbb{G}}$ . Since  $S^{\mathbb{G}}_A$  is the restriction of the identity correspondence  $E_{A\bar{\otimes}M}$  to  $A\cong \alpha(A)$ , we see from Example [5.15](#page-30-0) and Example [5.3](#page-26-3) that

$$
S_A^{\mathbb{G}} \cong \mathcal{H} \boxtimes \overline{\mathcal{H}}, \qquad \mathcal{H} = {}_{\alpha(A)}(L^2(A) \otimes L^2(\mathbb{G}))_{A \bar{\otimes} M}.
$$

A similar reasoning applies to the adjoint  $\mathbb{G}-A-A$ -correspondence  $D_{\mathbb{G}}^{A}$  as introduced in Example [2.9:](#page-10-2) since it is the restriction of the identity correspondence  $E_{A\rtimes\mathbb{G}}$  to  $A \cong \alpha(A)$ , where  $A\rtimes\mathbb{G}$  is endowed with the adjoint G-action, we find that

$$
D_A^{\mathbb G}\cong {\mathcal H}\boxtimes \overline{{\mathcal H}}, \qquad {\mathcal H}={_{\alpha(A)}}(L^2(A)\otimes L^2(\mathbb G))_{A\rtimes \mathbb G}.
$$

<span id="page-30-2"></span>Example 5.20. It is easily computed that the regular amplification in Example [2.6](#page-9-1) is obtained by tensoring with the  $\mathbb{G}\text{-}\mathrm{semi-converse\,\,correspondence:}\,\,\text{if}\,\mathcal{H}\,\,\text{is}\,\,\text{a}\,\,\mathbb{G}\text{-}\mathrm{equivariant}\,\,A\text{-}\mathrm{B}\text{-}\mathrm{correspondence,}\,\,\text{then}$ 

$$
S^{\mathbb{G}}(\mathcal{H}) \cong \mathcal{H} \boxtimes S_B^{\mathbb{G}} \cong S_A^{\mathbb{G}} \boxtimes \mathcal{H}.
$$

Similarly, the adjoint amplification of  $H$  is gotten by tensoring with the adjoint  $\mathbb{G}$ -correspondence,

$$
D^{\mathbb{G}}(\mathcal{H}) \cong \mathcal{H} \boxtimes D_B^{\mathbb{G}} \cong D_A^{\mathbb{G}} \boxtimes \mathcal{H}.
$$

5.3. Crossed products. We lift the crossed product construction from actions to the level of equivariant correspondences. This is an operation which is non-existent (or more precisely trivial) in the case of nonequivariant correspondences.

**Proposition 5.21.** Let  $(\mathcal{H}, U, \pi, \rho)$  be a G-A-B-correspondence. Then

$$
\mathcal{H}^{\rtimes}=\mathcal{H}\rtimes\mathbb{G}:=\mathcal{H}\otimes L^2(\mathbb{G})
$$

becomes a  $\check{G}$ - $A^{\rtimes}$ - $B^{\rtimes}$ -correspondence for the following structures:

• the normal  $A^{\rtimes}$ -representation  $\pi^{\rtimes}$  determined by

$$
\pi^{\rtimes}(z) = (\pi \otimes id)(z), \qquad z \in A^{\rtimes},
$$

• the normal anti- $*$ -representation  $\rho^{\rtimes}$  of  $B^{\rtimes}$  given by

$$
\rho^{\rtimes}(z) = U(\rho \otimes \check{J}(-)^*\check{J})(z)U^*, \qquad z \in B^{\rtimes},
$$

• the unitary  $\check{\mathfrak{G}}$ -representation  $U_{\rtimes} := \widetilde{W}_{23}$ .

Proof. It is easily seen that  $\pi^{\rtimes}$  and  $\rho^{\rtimes}$  indeed define respectively a normal \*-representation and a normal anti-∗-representation of  $A^{\rtimes}$  and  $B^{\rtimes}$ . Note that the ranges of  $\pi^{\rtimes}$  and  $\rho^{\rtimes}$  commute: indeed, we have

$$
\pi^{\rtimes}(\alpha(a)) = (\pi \otimes \pi_M)\alpha(a), \qquad \pi^{\rtimes}(1 \otimes x) = 1 \otimes \pi_{\check{M}}(x), \qquad a \in A, x \in \check{M}, \tag{5.9}
$$

$$
\rho^{\rtimes}(\beta(b)) = \rho(b) \otimes 1, \qquad \rho^{\rtimes}(1 \otimes x) = U(1 \otimes \rho_{\check{M}}(x))U^*, \qquad b \in B, x \in \check{M}, \tag{5.10}
$$

so the only non-trivial commutation to check is the one between  $U(1 \otimes \rho_{\check{M}}(\check{M}))U^*$  and  $1 \otimes \pi_{\check{M}}(\check{M})$ . But this follows easily from the identity  $U_{12}^* V_{23} U_{12} = U_{13} V_{23}$ .

It is clear that  $U_{\lambda}$  interacts in the correct way with  $\pi^{\lambda}$ . To see its  $\rho^{\lambda}$ -compatibility, simply note that

$$
\widetilde{W}_{23}^* U_{12} = U_{12} \widetilde{W}_{23}^*.
$$

<span id="page-31-2"></span>We now determine how the above crossed product constructions interact with opposites and composition.

**Proposition 5.22.** Let  $H$  be a G-A-B-correspondence. We have

$$
\overline{\mathcal{H}}^{\rtimes}\cong \overline{\mathcal{H}^{\rtimes}}.
$$

Proof. The unitary

$$
\overline{\mathcal{H}} \otimes L^{2}(\mathbb{G}) \to \overline{\mathcal{H} \otimes L^{2}(\mathbb{G})}, \qquad \overline{\xi} \otimes \eta \mapsto \overline{U(\xi \otimes J\eta)}
$$
  
does the job.

In the proof of the following theorem, it will be convenient to have a different model for the crossed product correspondence. If H is a G-A-B-correspondence, we will denote by  $\mathcal{H}^{\blacktriangleleft}$  the  $\ddot{\mathbb{G}}$ -A<sup>\n</sup>-B<sup>\n</sup>-correspondence  $\mathcal{H} \otimes L^2(\mathbb{G})$  obtained by conjugating  $\mathcal{H}^{\rtimes}$  with  $U^*$ , so in particular

$$
\pi^{\blacktriangleleft}(\alpha(a)) = \pi(a) \otimes 1, \quad \pi^{\blacktriangleleft}(1 \otimes x) = U^*(1 \otimes \pi_{\check{M}}(x))U, \qquad a \in A, x \in \check{M},
$$

$$
\rho^{\blacktriangleleft}(\beta(b)) = (\rho \otimes \pi_M R)\beta(b), \quad \rho^{\blacktriangleleft}(1 \otimes x) = 1 \otimes \rho_{\check{M}}(x), \qquad b \in B, x \in \check{M},
$$

while  $U_{\blacktriangleleft} = \widetilde{W}_{23}$  is still the associated unitary  $\check{\mathbb{G}}$ -representation. Note that under the above representation, we have for example that

<span id="page-31-0"></span>
$$
(\pi^{\blacktriangleleft} \otimes id)(V_{23}) = U_{12}^* V_{23} U_{12} = U_{13} V_{23}.
$$
\n(5.11)

<span id="page-31-1"></span>**Theorem 5.23.** Let  $(\mathcal{H}, U_{\mathcal{H}}, \pi_{\mathcal{H}}, \rho_{\mathcal{H}})$  be a G-A-B-correspondence, and let  $(\mathcal{G}, U_{\mathcal{G}}, \pi_{\mathcal{G}}, \rho_{\mathcal{G}})$  be a G-B-Ccorrespondence. Then we have

$$
(\mathcal{H}\boxtimes\mathcal{G})^{\rtimes}\cong\mathcal{H}^{\rtimes}\boxtimes\mathcal{G}^{\rtimes}.
$$

Proof. We will show rather that

$$
(\mathcal{H}\boxtimes\mathcal{G})^\blacktriangleleft\cong\mathcal{H}^\rtimes\boxtimes\mathcal{G}^\rtimes
$$

Indeed, write  $X = X_B(\mathcal{H})$  and consider the map

$$
(X \odot_B \mathcal{G}) \odot L^2(\mathbb{G}) \to \mathcal{H}^{\times} \boxtimes \mathcal{G}^{\times}, \quad (x \otimes_B \xi) \otimes \eta \mapsto \alpha_X(x) \otimes_{B^{\times}} U_{\mathcal{G}}(\xi \otimes \eta), \tag{5.12}
$$

where we have used that  $\alpha_X(X) \subseteq X_{B^{\rtimes}}(\mathcal{H}^{\rtimes})$ . A direct calculation shows that this is a well-defined isometric map, which then extends to an isometry

<span id="page-32-0"></span>
$$
T: (\mathcal{H} \boxtimes \mathcal{G}) \otimes L^2(\mathbb{G}) \to \mathcal{H}^{\times} \boxtimes \mathcal{G}^{\times}.
$$
\n
$$
(5.13)
$$

In fact, it is an easy observation that  $X_{B<sup>\times</sup>}(\mathcal{H}^{\times})$  equals the  $\sigma$ -strong closure of  $\alpha_X(X)(1 \otimes M)$ , hence [\(5.13\)](#page-32-0) will be unitary.

It is immediate that  $T$  is an intertwiner of A-representations and of right  $\dot{M}$ -representations.

To see that T is an intertwiner of  $\mathbb{G}$ -representations, we compute

$$
U_{\rtimes\boxtimes}((\alpha_X(x)\otimes_{B^{\rtimes}} U_{\mathcal{G}}(\xi\otimes\eta))\otimes\zeta) = (\alpha_X(x)\otimes 1)\otimes_{B^{\rtimes}\bar{\otimes}\tilde{M}}\overline{W}_{23}U_{\mathcal{G},12}(\xi\otimes\eta\otimes\zeta)
$$
  

$$
= (\alpha_X(x)\otimes 1)\otimes_{B^{\rtimes}\bar{\otimes}\tilde{M}} U_{\mathcal{G},12}(\xi\otimes\widetilde{W}(\eta\otimes\zeta)),
$$

from which we see that

$$
U_{\rtimes\boxtimes}(\omega)(\alpha_X(x)\otimes_{B^{\rtimes}} U_{\mathcal{G}}(\xi\otimes\eta))=\alpha_X(x)\otimes_{B^{\rtimes}} U_{\mathcal{G}}(\xi\otimes(\text{id}\otimes\omega)(\widetilde{W})\eta),\qquad\omega\in B(L^2(\mathbb{G}))_*.
$$

Hence  $TU_{\mathbb{R}}(\omega) = U_{\mathbb{R}}(\omega)T$ , and so T intertwines the two  $\check{\mathbb{G}}$ -representations.

The intertwining property for right C-representations can be obtained by observing first the identification

$$
(\mathcal{H}\boxtimes\mathcal{G})\otimes L^{2}(\mathbb{G})\cong(\mathcal{H}\otimes L^{2}(\mathbb{G}))\underset{B\bar{\otimes}M}{\boxtimes}(\mathcal{G}\otimes L^{2}(\mathbb{G})),\qquad(x\otimes_{B}\xi)\otimes\eta\mapsto(x\otimes 1)\otimes_{B\bar{\otimes}M}(\xi\otimes\eta).
$$

Under this identification, the map  $T$  can be written as

$$
T: (\mathcal{H}\otimes L^2(\mathbb{G}))\boxtimes (\mathcal{G}\otimes L^2(\mathbb{G}))\to \mathcal{H}^{\rtimes}\boxtimes \mathcal{G}^{\rtimes}, \qquad (x\otimes 1_M)\otimes_{B\bar\otimes M} \xi \mapsto \alpha_X(x)\otimes_{B^{\rtimes}}U_{\mathcal{G}}\xi,
$$

where  $x \in X_B(\mathcal{H})$  and  $\xi \in \mathcal{G} \otimes L^2(\mathbb{G})$ . With this description, it is immediate that T intertwines the right C-representations as well.

Finally, we need to check that  $T$  is an intertwiner of left  $\dot{M}$ -representations. For this, we note that we can identify

$$
(\mathcal{H} \boxtimes \mathcal{G}) \otimes L^{2}(\mathbb{G}) \otimes L^{2}(\mathbb{G}) \cong (\mathcal{H} \otimes L^{2}(\mathbb{G}) \otimes L^{2}(\mathbb{G})) \underset{B \bar{\otimes} \tilde{M} \bar{\otimes} M}{\boxtimes} (\mathcal{G} \otimes L^{2}(\mathbb{G}) \otimes L^{2}(\mathbb{G})),
$$

$$
(x \otimes \xi) \otimes \eta \otimes \zeta \mapsto (x \otimes 1 \otimes 1) \otimes_{B \bar{\otimes} \tilde{M} \bar{\otimes} M} (\xi \otimes \eta \otimes \zeta),
$$

and similarly we can identify

$$
(\mathcal{H}^{\rtimes} \underset{B^{\rtimes}}{\boxtimes} \mathcal{G}^{\rtimes}) \otimes L^{2}(\mathbb{G}) \cong (\mathcal{H}^{\rtimes} \otimes L^{2}(\mathbb{G})) \underset{B^{\rtimes} \bar{\otimes} M}{\boxtimes} (\mathcal{G}^{\rtimes} \otimes L^{2}(\mathbb{G})),
$$

$$
(x \otimes_{B^{\rtimes}} \xi) \otimes \eta \mapsto (x \otimes 1) \otimes_{B^{\rtimes} \bar{\otimes} M} (\xi \otimes \eta).
$$

Then  $T \otimes id_{L^2(\mathbb{G})}$  can be viewed as the unitary

$$
(\mathcal{H}\otimes L^{2}(\mathbb{G})\otimes L^{2}(\mathbb{G}))\underset{B\bar{\otimes}M\bar{\otimes}M}{\boxtimes}(\mathcal{G}\otimes L^{2}(\mathbb{G})\otimes L^{2}(\mathbb{G}))\to (\mathcal{H}^{\times}\otimes L^{2}(\mathbb{G}))\underset{B^{\times}\bar{\otimes}M}{\boxtimes}(\mathcal{G}^{\times}\otimes L^{2}(\mathbb{G})),
$$
  

$$
z_{13}\otimes_{B\bar{\otimes}M\bar{\otimes}M}\xi\mapsto(\alpha\otimes\mathrm{id})(z)\bar{\otimes}_{B^{\times}\bar{\otimes}M}U_{\mathcal{G},12}\xi,
$$

where  $z \in X_B(\mathcal{H}) \bar{\otimes} M$  and  $\xi \in \mathcal{G} \otimes L^2(\mathbb{G}) \otimes L^2(\mathbb{G})$ .

It is now sufficient to prove that, under these models,

<span id="page-32-1"></span>
$$
(T \otimes 1)(\pi^{\boxtimes \blacktriangleleft} \otimes \mathrm{id})(V) = (\pi^{\rtimes \boxtimes} \otimes \mathrm{id})(V)(T \otimes 1). \tag{5.14}
$$

But since by  $(5.11)$  we have

$$
(\pi^{\boxtimes \blacktriangleleft} \otimes \mathrm{id})(V) = U_{\boxtimes,13} V_{23},
$$
  

$$
33
$$

we see that [\(5.14\)](#page-32-1) follows from the following computation for  $x \in X_B(\mathcal{H})$  and  $\xi \in \mathcal{G} \otimes L^2(\mathbb{G}) \otimes L^2(\mathbb{G})$ :

$$
(T \otimes 1)(\pi^{\boxtimes} \blacktriangleleft \otimes id)(V)((x \otimes 1 \otimes 1) \otimes_{B \bar{\otimes} M \bar{\otimes} M} \xi)
$$
  
\n
$$
= (T \otimes 1)U_{\boxtimes,13}((x \otimes 1 \otimes 1) \otimes_{B \bar{\otimes} M \bar{\otimes} M} V_{23}\xi)
$$
  
\n
$$
= (T \otimes 1)(\alpha_X(x)_{13} \otimes_{B \bar{\otimes} M \bar{\otimes} M} U_{\mathcal{G},13} V_{23}\xi)
$$
  
\n
$$
= (\alpha \otimes id)\alpha(x) \otimes_{B^{\times} \bar{\otimes} M} U_{\mathcal{G},12} U_{\mathcal{G},13} V_{23}\xi
$$
  
\n
$$
= V_{23}\alpha(x)_{12} \otimes_{B^{\times} \bar{\otimes} M} V_{23}^* U_{\mathcal{G},12} U_{\mathcal{G},13} V_{23}\xi
$$
  
\n
$$
= V_{23}\alpha(x)_{12} \otimes_{B^{\times} \bar{\otimes} M} U_{\mathcal{G},12}\xi
$$
  
\n
$$
= (\pi^{\rtimes \boxtimes} \otimes id)(V)(\alpha(x)_{12} \otimes_{B^{\times} \bar{\otimes} M} U_{\mathcal{G},12}\xi)
$$
  
\n
$$
= (\pi^{\rtimes \boxtimes} \otimes id)(V)(T \otimes 1)((x \otimes 1 \otimes 1) \otimes_{B \bar{\otimes} M \bar{\otimes} M} \xi).
$$

5.4. **Takesaki-Takai duality.** Recall that  $M^{\vee\vee} \cong M$  via  $\text{Ad}(u_{\mathbb{G}})$ . Hence any  $\mathbb{G}\text{-}W^*$ -algebra A can be seen as a G∨∨-W<sup>∗</sup> -algebra by a coaction that we write

$$
\widetilde{\alpha}: A \to A \bar{\otimes} M^{\vee \vee}, \qquad a \mapsto (1 \otimes u_{\mathbb{G}}) \alpha(a) (1 \otimes u_{\mathbb{G}}).
$$

The Takesaki-Takai duality for actions of locally compact quantum groups on von Neumann algebras [\[ES80,](#page-47-36) [Vae01\]](#page-47-6) can now be stated as follows.

<span id="page-33-2"></span>**Theorem 5.24** (Takesaki-Takai duality). *Endow*  $\overline{A \otimes B}(L^2(\mathbb{G}))$  with the G-action

 $\alpha_+ = \text{Ad}(\Sigma_{23}W_{23}) \circ (\alpha \otimes \text{id}).$ 

Then there is a  $\mathbb{G}^{\vee\vee}$ -equivariant  $\ast$ -isomorphism

<span id="page-33-1"></span>
$$
\Phi: (A^{\rtimes}{}^{\rtimes}, \alpha^{\rtimes}{}^{\rtimes}) \cong (A \bar{\otimes} B(L^2(\mathbb{G})), \widetilde{\alpha_+}), \tag{5.15}
$$

 $\Box$ 

whose inverse is concretely implemented by

$$
\Phi^{-1}: A\bar{\otimes}B(L^2(\mathbb{G}))\to A^{\rtimes\rtimes}\subseteq B(L^2(A)\otimes L^2(\mathbb{G})\otimes L^2(\mathbb{G})),\quad z\mapsto V_{23}^*(\alpha\otimes\mathrm{id})(z)V_{23}.
$$

On generating elements,  $\Phi$  is determined by

$$
\alpha(a)\otimes 1\mapsto \alpha(a), \quad 1\otimes \check{\Delta}(x)\mapsto 1\otimes x, \quad 1\otimes 1\otimes y\mapsto 1\otimes y, \qquad a\in A, x\in \check{M}, y\in M',
$$

We reformulate this result as follows.

**Theorem 5.25.** Endow the Hilbert space  $_{A^{\times}} \mathcal{M}_A := L^2(A) \otimes L^2(\mathbb{G})$  with

- the normal unital  $*$ -representation  $(\pi_A \otimes id) \circ \Phi$  of  $A^{\rtimes} \times$ ,
- the normal unital anti- $*$ -representation  $\rho_A \otimes 1$  of A, and
- the  $\mathbb{G}^{\vee\vee}$ -representation  $u\in A}W_{32}U_{\alpha,13}u\in A$ .

Then  $_{A^{\rtimes}\times}\mathscr{M}_A$  becomes a  $\mathbb{G}^{\vee\vee}\text{-}W^*$ -Morita correspondence between  $(A^{\rtimes}\alpha, \alpha^{\rtimes}\alpha)$  and  $(A, \widetilde{\alpha})$ .

Note now that if  $(A, \alpha), (B, \beta)$  are G-W<sup>\*</sup>-algebras, any G-A-B-correspondence  $(\mathcal{H}, \pi, \rho, U)$  becomes a G<sup>∨∨</sup>equivariant  $(A, \tilde{\alpha})$ - $(B, \tilde{\beta})$ -correspondence by means of

$$
\mathcal{H}=(\mathcal{H},\pi,\rho,(1\otimes u_{\mathbb{G}})U(1\otimes u_{\mathbb{G}})).
$$

We can then extend the Takesaki-Takai duality to general G-equivariant correspondences.

<span id="page-33-0"></span>**Theorem 5.26.** Let  $(\mathcal{H}, U_{\mathcal{H}}, \pi_{\mathcal{H}}, \rho_{\mathcal{H}})$  be a G-equivariant A-B-correspondence. Then we have a unitary intertwiner of G∨∨-A-B-correspondences

$$
\mathcal{H}^{\rtimes \rtimes} \cong {}_{A^{\rtimes} \rtimes} \mathscr{M}_A \boxtimes \widetilde{\mathcal{H}} \boxtimes {}_B{\overline{\mathscr{M}}}_{B^{\rtimes \rtimes}}.
$$

*Proof.* Since  $\mathcal M$  is a  $\mathbb{G}^{\vee\vee}$ -W<sup>\*</sup>-Morita correspondence, it is (by associativity of ⊠ and Proposition [5.10\)](#page-29-2) enough to show that

<span id="page-34-1"></span>
$$
A^{\rtimes\rtimes} \mathcal{H}_{B^{\rtimes\rtimes}}^{\rtimes\rtimes} \boxtimes_{B^{\rtimes\rtimes}} \mathcal{M}_B \cong A^{\rtimes\rtimes} \mathcal{M}_A \boxtimes_A \widetilde{\mathcal{H}}_B. \tag{5.16}
$$

Now the representation of  $A \rtimes \mathbb{G} \rtimes \check{\mathbb{G}}$  and anti-representation of  $B \rtimes \mathbb{G} \rtimes \check{\mathbb{G}}$  on  $\mathcal{H}^{\rtimes} \rtimes = \mathcal{H} \otimes L^2(\mathbb{G}) \otimes L^2(\mathbb{G})$ are given by

$$
\pi^{\otimes \otimes}(\alpha(a) \otimes 1) = (\pi_{\mathcal{H}} \otimes \pi_M)(\alpha(a))_{12}, \quad \rho^{\otimes \otimes}(\beta(b) \otimes 1) = \rho_{\mathcal{H}}(b)_{1}, \qquad a \in A, b \in B,
$$
  

$$
\pi^{\otimes \otimes}(1 \otimes \check{\Delta}(y)) = \check{\Delta}(y)_{23}, \quad \rho^{\otimes \otimes}(1 \otimes \check{\Delta}(y)) = U_{\mathcal{H}, 12}\rho_{\check{M}}(y)_{2}U_{\mathcal{H}, 12}^{*}, \qquad y \in \check{M},
$$
  

$$
\pi^{\otimes \otimes}(1 \otimes 1 \otimes z) = \pi_{M^{\vee \vee}}(z)_{3}, \quad \rho^{\otimes \otimes}(1 \otimes 1 \otimes z) = \widetilde{W}_{23}\rho_{M^{\vee \vee}}(z)_{3}\widetilde{W}_{23}^{*}, \qquad z \in M^{\vee \vee},
$$

with associated unitary  $\mathbb{G}^{\vee\vee}$ -representation given by  $\widetilde{V}_{34}$ , where we recall that  $\widetilde{V} = \text{Ad}(u_{\mathbb{G}} \otimes u_{\mathbb{G}})(V)$ .

It is then easily verified that

$$
X_{B^{\times \times}}(\mathcal{H}^{\times \times}) = \sigma\text{-closure of the span of }\{\Delta_X(x)_{12}\check{\Delta}(y)_{23}\pi_{M^{\vee \vee}}(z)_3 \mid x \in X_B(\mathcal{H}), y \in \check{M}, z \in M^{\vee \vee}\}.
$$

From this, we find a unitary map

<span id="page-34-0"></span>
$$
\mathcal{H}^{\rtimes} \underset{B^{\rtimes}{}^{\rtimes}}{\boxtimes} (L^2(B) \otimes L^2(\mathbb{G})) \to \mathcal{H} \otimes L^2(\mathbb{G}), \quad \left(\Delta_X(x)_{12}\Delta(y)_{23}\pi_{M^{\vee}}(z)_3\right) \otimes_{B^{\rtimes}{}^{\rtimes}} \xi \mapsto \Delta_X(x)(1 \otimes yz)\xi, \tag{5.17}
$$

which hence by transport of structure endows  $\mathcal{H} \otimes L^2(\mathbb{G})$  with the structure of a  $\mathbb{G}^{\vee \vee}$ -equivariant  $A^{\rtimes}{}^{\mathcal{A}}$ -Bcorrespondence  $(\mathcal{H} \otimes L^2(\mathbb{G}), U, \pi, \rho)$ .

Clearly  $\pi$ ,  $\rho$  are given by

$$
\pi(z) = (\pi_{\mathcal{H}} \otimes \mathrm{id}_{B(L^2(\mathbb{G}))})\Phi^{-1}(z), \qquad \rho(b) = \rho_{\mathcal{H}}(b) \otimes 1, \qquad z \in A^{\times \times}, b \in B.
$$

The resulting unitary  $\mathbb{G}^{\vee\vee}$ -representation U on  $\mathcal{H}\otimes L^2(\mathbb{G})$  is given by the following transformation (dropping the notations  $\pi_{\tilde{M}}$  and  $\pi_{M} \vee \vee$ ): for  $x \in X_B(\mathcal{H}), y \in \tilde{M}, z \in M^{\vee \vee}$  and  $\xi, \eta \in L^2(\mathbb{G}),$  we have

$$
(\Delta_X(x)_{12}(1 \otimes yz)\xi) \otimes \eta \quad \mapsto \quad \Delta_X(x)_{12}y_2\Delta^{\vee\vee}(z)_{23}u_{\mathbb{G},3}W_{32}U_{\beta,13}u_{\mathbb{G},3}(\xi \otimes \eta)
$$
\n
$$
= \quad \Delta_X(x)_{12}y_2\Delta^{\vee\vee}(z)_{23}\tilde{V}_{23}u_{\mathbb{G},3}U_{\beta,13}u_{\mathbb{G},3}(\xi \otimes \eta)
$$
\n
$$
= \quad \Delta_X(x)_{12}y_2\tilde{V}_{23}z_2u_{\mathbb{G},3}U_{\beta,13}u_{\mathbb{G},3}(\xi \otimes \eta)
$$
\n
$$
= \quad \Delta_X(x)_{12}\tilde{V}_{23}u_{\mathbb{G},3}U_{\beta,13}u_{\mathbb{G},3}y_2z_2(\xi \otimes \eta)
$$
\n
$$
= \quad u_{\mathbb{G},3}\Delta_X(x)_{12}W_{32}U_{\beta,13}u_{\mathbb{G},3}y_2z_2(\xi \otimes \eta)
$$
\n
$$
= \quad u_{\mathbb{G},3}W_{32}(\mathrm{id} \otimes \Delta^{\mathrm{op}})\Delta_X(x)U_{\beta,13}u_{\mathbb{G},3}y_2z_2(\xi \otimes \eta)
$$
\n
$$
= \quad u_{\mathbb{G},3}W_{32}(\Delta_X \otimes \mathrm{id})\Delta_X(x)_{132}U_{\beta,13}u_{\mathbb{G},3}y_2z_2(\xi \otimes \eta)
$$
\n
$$
= \quad u_{\mathbb{G},3}W_{32}U_{\mathcal{H},13}u_{\mathbb{G},3}((\Delta_X(x)_{12}y_2z_2\xi) \otimes \eta).
$$

We hence find that [\(5.17\)](#page-34-0) endows  $\mathcal{H} \otimes L^2(\mathbb{G})$  with the unitary  $\mathbb{G}^{\vee \vee}$ -representation  $U = u_{\mathbb{G},3}W_{32}U_{\mathcal{H},13}u_{\mathbb{G},3}$ . But this gives precisely the right hand side of [\(5.16\)](#page-34-1) under the direct identification

$$
_{A^{\rtimes_{\rtimes}}} (L^2(A) \otimes L^2(\mathbb{G}))_A \boxtimes {}_A \widetilde{\mathcal{H}}_B \cong {}_{A^{\rtimes_{\rtimes}}} (\widetilde{\mathcal{H}} \otimes L^2(\mathbb{G}))_{B \otimes 1}.
$$

**Remark 5.27.** The above results can be upgraded slightly by noting that we obtain a W<sup>\*</sup>-bicategory Corr<sup>G</sup> whose objects are G-equivariant W<sup>\*</sup>-algebras, whose 1-cells are G-equivariant correspondences, and whose 2-cells are unitary intertwiners. Composition is determined by the Connes fusion product. Takesaki-Takai duality provides then an equivalence of W<sup>\*</sup>-bicategories between Corr<sup>G</sup> and Corr<sup>G</sup>. We do not enter into the details here.

5.5. Examples. In our next few results, we look at how the above duality interacts with our canonical G-equivariant correspondences, see Example [2.2](#page-8-0) and further.

<span id="page-35-1"></span>Proposition 5.28. We have

$$
(E_A^{\mathbb{G}})^{\rtimes} \cong E_{A^{\rtimes}}^{\check{\mathbb{G}}} \tag{5.18}
$$

<span id="page-35-2"></span>*Proof.* This follows immediately from Theorem [1.3.](#page-7-1)

**Proposition 5.29.** Let  $A \subseteq C$  and  $B \subseteq D$  be G-equivariant inclusions. Then for any G-C-D-correspondence H we have

<span id="page-35-0"></span>
$$
({}_{A}\text{Res}_{B}(\mathcal{H}))^{\rtimes} = {}_{A^{\rtimes}}\text{Res}_{B^{\rtimes}}(\mathcal{H}^{\rtimes}).\tag{5.19}
$$

Note that e.g. the inclusion  $\overline{A\otimes B}(L^2(\mathbb{G})) \subseteq C\overline{\otimes}B(L^2(\mathbb{G}))$  descends to an inclusion  $A^{\rtimes} \subseteq C^{\rtimes}$ , so the restrictions in the right hand side of [\(5.19\)](#page-35-0) are meaningful.

*Proof.* Immediate.  $\square$ 

**Proposition 5.30.** Duality interchanges the regular amplification and the adjoint amplification: if  $H$  is a G-equivariant A-B-correspondence, then

$$
(D^{\mathbb{G}}(\mathcal{H}))^{\rtimes} \cong S^{\check{\mathbb{G}}}(\mathcal{H}^{\rtimes}), \qquad S^{\mathbb{G}}(\mathcal{H})^{\rtimes} \cong D^{\check{\mathbb{G}}}(\mathcal{H}^{\rtimes}). \tag{5.20}
$$

*Proof.* By Example [5.20](#page-30-2) and Theorem [5.23,](#page-31-1) it is sufficient to show that  $(D_A^{\mathbb{G}})^{\rtimes} \cong S_{A^{\rtimes} \atop A^{\rtimes}}^{\mathbb{G}}$  and  $(S_A^{\mathbb{G}})^{\rtimes} \cong D_{A^{\rtimes}}^{\mathbb{G}}$ . Furthermore, by the observations in Example [2.6](#page-9-1) and Example [2.9](#page-10-2) that  $S_A^{\mathbb{G}}$  and  $D_A^{\mathbb{G}}$  are restrictions of identity correspondences, it is by Proposition [5.28](#page-35-1) and Proposition [5.29](#page-35-2) sufficient to show that there are G-equivariant ∗-isomorphisms

<span id="page-35-3"></span>
$$
A^{\rtimes} \rtimes_{\alpha_{ad}} \mathbb{G} \cong A^{\rtimes} \bar{\otimes} \check{M}, \qquad (A \bar{\otimes} M) \rtimes_{\mathrm{id} \otimes \Delta} \mathbb{G} \cong A^{\rtimes} \rtimes_{\alpha^{\rtimes}} \check{\mathbb{G}}, \tag{5.21}
$$

with in each case  $\alpha(A) \rtimes \mathbb{G}$  carried onto  $\alpha^*(A^*)$ . Note that all of the above von Neumann algebras are (by their canonical constructions) implemented on  $L^2(A) \otimes L^2(\mathbb{G}) \otimes L^2(\mathbb{G})$ .

Now  $A^{\rtimes} \rtimes_{\alpha_{ad}} \mathbb{G}$  is generated by  $V_{23}(\alpha(A) \otimes 1)V_{23}^*$ ,  $V_{23}(1 \otimes \check{M} \otimes 1)V_{23}^*$  and  $1 \otimes 1 \otimes \check{M}$ , with coaction implemented by the unitary representation  $\overline{W}_{34}$ . Upon conjugating with  $V_{23}^*$ , the unitary representation remains unchanged, while we now get the von Neumann algebra generated by  $\alpha(A) \otimes 1$ ,  $1 \otimes \tilde{M} \otimes 1$  and  $1\otimes\Delta(M)$ . The second and third term generate  $M\bar{\otimes}M$ , so all the terms together generate  $A^{\rtimes}\bar{\otimes}M$ . Meanwhile, the first and third term together generate  $\alpha^*(A^*)$ . Since  $\widetilde{W}_{34}$  is still an implementing unitary for the  $\check{G}$ action on  $A^{\rtimes} \bar{\otimes} \check{M}$ , this proves the existence of the first isomorphism in [\(5.21\)](#page-35-3).

Turning now to the second isomorphism in [\(5.21\)](#page-35-3), note that the right hand side can be identified with  $A\bar{\otimes}B(L^2(\mathbb{G}))$  under the Takesaki-Takai-duality isomorphism [\(5.15\)](#page-33-1). Recalling that  $\check{V}=\widetilde{W}$ , it is immediately verified that the  $\check{\mathbb{G}}$ -adjoint action on  $A^{\rtimes} \rtimes \check{\mathbb{G}}$  gets turned into the coaction

<span id="page-35-4"></span>
$$
A\overline{\otimes}B(L^2(\mathbb{G})) \to A\overline{\otimes}B(L^2(\mathbb{G}))\overline{\otimes}\check{M}, \quad z \mapsto \widetilde{W}_{23}z_{12}\widetilde{W}_{23}^*.
$$
 (5.22)

On the other hand, the left hand side of the second isomorphism in [\(5.21\)](#page-35-3) is generated by  $A \otimes 1 \otimes 1$ ,  $1 \otimes \Delta(M)$ and  $1 \otimes 1 \otimes \check{M}$ , which under conjugation with  $W_{23}$ , and the observation that M and  $\check{M}$  generate  $B(L^2(\mathbb{G}))$ , becomes  $A\bar{\otimes}1\bar{\otimes}B(L^2(\mathbb{G}))\cong A\bar{\otimes}B(L^2(\mathbb{G}))$ . Under this identification, the dual coaction gets exactly identified with [\(5.22\)](#page-35-4). It is also immediate that under these identifications, the von Neumann algebras generated by the copies of A and M<sup> $\dot{M}$ </sup> in respectively  $(A\bar{\otimes}M) \rtimes_{\mathrm{id}\otimes\Delta} \mathbb{G}$  and  $A^{\rtimes} \rtimes_{\alpha^{\rtimes}} \tilde{\mathbb{G}}$  get sent to each other.

<span id="page-35-5"></span>Proposition 5.31. We have

$$
(S_{A,M}^{\mathbb{G}})^{\rtimes} \cong S_{A^{\rtimes},\mathbb{C}}^{\check{\mathbb{G}}}\boxtimes_{\mathbb{C}} L^2(\mathbb{G})_{M^{\rtimes}},
$$

where  ${}_{\mathbb{C}} L^2(\mathbb{G})_{M^{\times}}$  is a  $\check{\mathbb{G}}$ -W<sup>\*</sup>-Morita correspondence between  $\mathbb{C}$  and  $M^{\times}$ .

Proof. By Theorem [5.23,](#page-31-1) Example [5.16,](#page-30-1) Proposition [5.28](#page-35-1) and Proposition [5.29,](#page-35-2) we have that

$$
S_{A,M}^{\mathbb{G}})^{\rtimes} \cong ({}_{A}L^{2}(A)_{\mathbb{C}} \boxtimes {}_{\mathbb{C}}L^{2}(\mathbb{G})_{M})^{\rtimes} \cong {}_{A^{\rtimes}}L^{2}(A^{\rtimes})_{\check{M}} \boxtimes {}_{\check{M}}L^{2}(M^{\rtimes})_{M^{\rtimes}}.
$$

So by Example [5.15,](#page-30-0) it is sufficient to show that

 $\left(\right)$ 

$$
{}_{\check{M}}L^2(M^\rtimes)_{M^\rtimes}\cong {}_{\check{M}}L^2(\mathbb{G})_\mathbb{C}\boxtimes {}_\mathbb{C}L^2(\mathbb{G})_{M^\rtimes}.
$$

But we have in fact that

$$
{}_{M^{\rtimes}}L^2(M^{\rtimes}){}_{M^{\rtimes}} \cong {}_{\check{M}}L^2(\mathbb{G}){}_\mathbb{C} \boxtimes {}_\mathbb{C} L^2(\mathbb{G}){}_{M^{\rtimes}}.
$$

Indeed, we may as well prove this with M replaced by  $\tilde{M}$ , in which case we can write the isomorphism as

$$
{}_{\mathbb{C}^{\rtimes M}}L^2(\mathbb{C}^{\rtimes \rtimes})_{\mathbb{C}^{\rtimes M}} \cong {}_{\mathbb{C}^{\rtimes M}}L^2(\check{\mathbb{G}})_{\mathbb{C}} \boxtimes {}_{\mathbb{C}}L^2(\check{\mathbb{G}})_{\mathbb{C}^{\rtimes M}},
$$

which is just a special case of the Takesaki-Takai duality of Theorem [5.26.](#page-33-0)

Corollary 5.32. We have

$$
(C_{A,B}^{\mathbb{G}})^{\rtimes} \cong S_{A^{\rtimes},B^{\rtimes}}^{\check{\mathbb{G}}} \tag{5.23}
$$

Proof. Using the Proposition [5.31](#page-35-5) and Theorem [5.23,](#page-31-1) we find

$$
(C_{A,B}^{\mathbb{G}})^{\rtimes} \cong (S_{A,M}^{\mathbb{G}} \boxtimes \overline{S_{B,M}^{\mathbb{G}}})^{\rtimes} \cong S_{A^{\rtimes} , \mathbb{C}}^{\check{\mathbb{G}}} \boxtimes_{\mathbb{C}} L^{2}(\mathbb{G})_{M^{\rtimes}} \boxtimes_{M^{\rtimes}} L^{2}(\mathbb{G})_{\mathbb{C}} \boxtimes S_{\mathbb{C},B^{\rtimes}}^{\check{\mathbb{G}}} \cong S_{A^{\rtimes} , \mathbb{C}}^{\check{\mathbb{G}}} \boxtimes S_{\mathbb{C},B^{\rtimes}}^{\check{\mathbb{G}}} \cong S_{A^{\rtimes},B^{\rtimes}}^{\check{\mathbb{G}}},
$$

where in the penultimate step we used that  ${}_{\mathbb{C}}L^2(\mathbb{G})_{M^{\times}}$  is a  $\check{\mathbb{G}}$ -W<sup>\*</sup>-Morita correspondence between  $\mathbb{C}$  and  $M^{\rtimes}$  and Proposition [5.10,](#page-29-2) and where in the last step we used Example [5.16.](#page-30-1)

<span id="page-36-0"></span>We also have the following observation.

Proposition 5.33. We have

$$
({}_{A}L^{2}(A)_{\mathbb{C}}\boxtimes {}_{\mathbb{C}}L^{2}(\mathbb{G})_{\mathbb{C}})^{\rtimes}\cong S_{A^{\rtimes}, \check{M}}^{\check{\mathbb{G}}},
$$

where  $_A L^2(A)_{\mathbb{C}} \boxtimes_{\mathbb{C}} L^2(\mathbb{G})_{\mathbb{C}} = {}_A \text{Res}_{\mathbb{C}}(S^{\mathbb{G}}_{A,M}).$ 

*Proof.* This follows directly from Proposition [5.31](#page-35-5) upon using Proposition [5.29.](#page-35-2)

This leads to the following, maybe more surprising corollary.

Corollary 5.34. We have

$$
(C_{A,B}^{\mathbb{G}} \otimes L^{2}(\mathbb{G}))^{\rtimes} \cong C_{A^{\rtimes},B^{\rtimes}}^{\check{\mathbb{G}}}
$$
\n
$$
(5.24)
$$

where the G-A-B-correspondence on the left is  $C_{A,B}^{\mathbb{G}}$ , amplified with a multiplicity Hilbert space  $L^2(\mathbb{G})$  (with which the structures have no interaction, indicated by the fact that we are using  $\otimes$  and not  $\boxtimes$ ).

Proof. By Fell absorption, we have

$$
{}_{\mathbb{C}} L^2(\mathbb{G})_{\mathbb{C}} \otimes L^2(\mathbb{G}) \cong {}_{\mathbb{C}} L^2(\mathbb{G})_{\mathbb{C}} \boxtimes {}_{\mathbb{C}} L^2(\mathbb{G})_{\mathbb{C}},
$$

where on the left  $L^2(\mathbb{G})$  is just a multiplicity Hilbert space, and where on the right we take the Connes fusion product of equivariant correspondences (which in this case is just the usual tensor product of the right regular G-representation with itself). We thus obtain

$$
C_{A,B}^{\mathbb{G}} \otimes L^{2}(\mathbb{G}) = (A L^{2}(A)_{\mathbb{C}} \boxtimes_{\mathbb{C}} L^{2}(\mathbb{G})_{\mathbb{C}}) \boxtimes (\mathbb{C} L^{2}(\mathbb{G})_{\mathbb{C}} \boxtimes_{\mathbb{C}} L^{2}(B)_{B}),
$$

and the corollary now follows directly from Proposition [5.33.](#page-36-0)

As a guide to the reader, we offer the following table which summarizes some of the results of this section.

$\mathcal H$	$H \boxtimes \overline{H}$	$\mathcal{H}^{\rtimes}$	$(\mathcal{H}\boxtimes\overline{\mathcal{H}})^{\rtimes}\cong\mathcal{H}^{\rtimes}\boxtimes\overline{\mathcal{H}^{\rtimes}}$
$_{\alpha(A)}L^2(A\bar{\otimes}M)_{A\bar{\otimes}M}$	$S_A^{\mathbb{G}}$	$\alpha^{\rtimes}(A^{\rtimes})L^2(A^{\rtimes}\rtimes_{\alpha^{\rtimes}}\check{\mathbb{G}})_{A^{\rtimes}\rtimes_{\alpha^{\rtimes}}\check{\mathbb{G}}}$	$D^{\tilde{\mathbb{G}}}_{A\rtimes}$
$_{\alpha(A)}L^2(A\rtimes\mathbb{G})_{A\rtimes M}$	$D_A^{\mathbb{G}}$	$_{\alpha^{\rtimes}(A^{\rtimes})}L^2(A^{\rtimes}\overline{\otimes} \check{M})_{A^{\rtimes}\overline{\otimes} \check{M}}$	$S_{A\times}^{\mathbb{G}}$
$S_{A,\mathbb{C}}^{\mathbb{G}} = {}_A L^2(A)_{\mathbb{C}}$	$S_{A,A}^{\mathbb{G}}$	$A^{\rtimes}L^2(A^{\rtimes})_{\check{M}}$	$_{A^{\rtimes}}L^2(A^{\rtimes})_{\check{M}} \boxtimes_{\check{M}}L^2(A^{\rtimes})_{A^{\rtimes}}$
$S_{A,M}^{\mathbb{G}}$	$C_{A,A}^{\mathbb{G}}$	$S_{A^{\rtimes}.\mathbb{C}}^{\mathbb{G}} \boxtimes_{\mathbb{C}} L^2(\mathbb{G})_{M^{\rtimes}}$	$S_{A^{\rtimes},A^{\rtimes}}^{\mathbb{G}}$
$_{A}L^{2}(A)_{\mathbb{C}}\boxtimes {}_{\mathbb{C}}L^{2}(\mathbb{G})_{\mathbb{C}}$	$C_{A,A}^{\mathbb{G}} \otimes L^{2}(\mathbb{G})$	$S_{A^\rtimes,\check{M}}^{\check{\mathbb{G}}}$	$C^{\tilde{\mathbb{G}}}_{A^{\rtimes},A^{\rtimes}}$

<span id="page-37-1"></span>Table 1. Operations for basic correspondences

#### 6. Preservation of weak containment and left amenability

6.1. Fell continuity for operations on equivariant correspondences. We assume by default that W<sup>\*</sup>-categories admit subobjects and arbitrary (small) direct sums. Then a functor  $F: \mathcal{C} \to \mathcal{D}$  between W<sup>\*</sup>-categories is called *normal* if all the induced maps  $F_X : End_{\mathcal{C}}(X) \to End_{\mathcal{D}}(F(X))$  are normal for any object  $X \in \mathcal{C}$ . This is the same [\[GLR85\]](#page-47-23) as each  $F_{X,Y}$ : Hom $c(X,Y) \to \text{Hom}_{\mathcal{D}}(F(X), F(Y))$  being  $\sigma$ -weakly continuous for each pair of objects  $X, Y \in \mathcal{C}$ . Moreover, this is also the same as F preserving arbitrary (small) direct sums, for a unital ∗-homomorphism between von Neumann algebras is normal if and only if it is completely additive (i.e. preserves arbitrary sums of orthogonal projections).

Normality takes care of continuity on the level of morphisms. Assume now in particular that  $C \subseteq \text{Rep}_*(C)$ and  $\mathcal{D} \subseteq \text{Rep}_*(D)$  for  $C, D$  C<sup>\*</sup>-algebras. Then on the class of objects we have the Fell topology, and we can ask when a normal functor  $F: \mathcal{C} \to \mathcal{D}$  is continuous with respect to the Fell topology. This will be the case if and only if  $F$  preserves weak containment:

$$
\pi \preccurlyeq \theta \quad \Rightarrow \quad F(\pi) \preccurlyeq F(\theta).
$$

Indeed, it is immediate that Fell continuity of  $F$  implies that  $F$  preserves weak containment. For the converse direction, one uses the well-known fact that if  $\pi \in \text{Rep}_*(C)$  and  $(\pi_\alpha)_{\alpha \in I}$  a net in  $\text{Rep}_*(C)$ , then  $\pi_\alpha \to \pi$ in the Fell topology if and only if  $\pi \preccurlyeq \oplus_{\beta \in J} \pi_{\beta}$  for each subnet  $(\pi_{\beta})_{\beta \in J}$  of  $(\pi_{\alpha})_{\alpha \in I}$  (see the introduction of [\[Fel63\]](#page-47-37) for the explicit statement, and for example [\[BdlHV08b,](#page-46-29) Proposition F.2.2] for the elements of the proof).

In light of the above, we introduce the following terminology.

**Definition 6.1.** We call a  $*$ -functor  $F : \text{Rep}_*(C) \to \text{Rep}_*(D)$  W<sup>\*</sup>-continuous if it preserves arbitrary direct sums and weak containment.

We can extend this terminology to functors  $F: \mathcal{C} \to \mathcal{D}$  of W<sup>\*</sup>-categories  $\mathcal{C} \subseteq \text{Rep}_*(C)$  and  $\mathcal{D} \subseteq \text{Rep}_*(D)$ which are closed under direct sums and subobjects. In particular, we can use this terminology for functors between categories of equivariant correspondences.

We now show the W<sup>∗</sup> -continuity of the operations introduced in Section [5.](#page-25-3) It is clear that the operations introduced there are compatible with arbitrary direct sums. We hence focus below on the preservation of weak containment.

<span id="page-37-0"></span>**Proposition 6.2.** Let  $\mathcal{G}, \mathcal{H}$  be G-A-B-correspondences. If  $\mathcal{G} \preccurlyeq \mathcal{H}$ , then  $\overline{\mathcal{G}} \preccurlyeq \overline{\mathcal{H}}$ .

*Proof.* This follows immediately from the original definition, since for  $a \in A, b \in B, \omega \in M_*$  and  $\xi, \eta \in \mathcal{G}$ , we have

$$
\langle \overline{\xi}, \overline{\pi}(b) \overline{U}(\omega) \overline{\rho}(a) \overline{\eta} \rangle = \langle \overline{\xi}, \overline{\rho(b^*)U(\omega \circ R)^* \pi(a^*)\eta} \rangle = \langle \eta, \pi(a)U(\omega \circ R) \rho(b) \xi \rangle.
$$

<span id="page-38-2"></span>**Proposition 6.3.** Let G, H be G-A-B-correspondences, and K a G-B-C-correspondence. Then

<span id="page-38-1"></span>
$$
\mathcal{G} \preccurlyeq \mathcal{H} \quad \Rightarrow \quad \mathcal{G} \boxtimes \mathcal{K} \preccurlyeq \mathcal{H} \boxtimes \mathcal{K},\tag{6.1}
$$

and this becomes  $\Leftrightarrow$  if K is a G-W<sup>\*</sup>-Morita correspondence.

Similarly for taking the fusion product on the left.

*Proof.* By Theorem [3.7,](#page-21-0) we can choose a contractive net  $z_{\alpha} \in X = X_B(\mathcal{G}, \mathcal{H}^{\infty})$  such that for all  $a \in A$ 

$$
(z_{\alpha}^*\pi_{\mathcal{H}^{\infty}}(a)\otimes 1)\alpha_X(z_{\alpha})\to \pi_{\mathcal{G}}(a)\otimes 1, \qquad \sigma\text{-weakly}.
$$

If then  $x, y \in X_B(G)$  and  $\xi, \eta \in \mathcal{K}$ , we have  $z_{\alpha}x, z_{\alpha}y \in X_B(\mathcal{H}^{\infty})$  and, for  $a \in A, c \in C$  and  $\omega = \omega_{\zeta, \zeta'} \in M_*$ , we compute that

$$
\langle (z_{\alpha}x \otimes_{B} \xi), \pi_{\boxtimes}(a)U_{\boxtimes}(\omega)\rho_{\boxtimes}(c)(z_{\alpha}y \otimes_{B} \eta) \rangle
$$
  
=  $\langle (z_{\alpha}x \otimes 1) \otimes_{B\bar{\otimes}M} (\xi \otimes \zeta), (\pi_{\mathcal{H}^{\infty}}(a) \otimes 1) \alpha_{X_{B}(\mathcal{H}^{\infty})}(z_{\alpha}y) \otimes_{B\bar{\otimes}M} U_{\mathcal{K}}(\rho_{\mathcal{K}}(c)\eta \otimes \zeta') \rangle$   
=  $\langle \xi \otimes \zeta, (x^*z^*_{\alpha}\pi_{\mathcal{H}^{\infty}}(a) \otimes 1) \alpha_X(z_{\alpha}) \alpha_{X_{B}(\mathcal{G})}(y)U_{\mathcal{K}}(\rho_{\mathcal{K}}(c)\eta \otimes \zeta') \rangle,$ 

which converges to

$$
\langle \xi \otimes \zeta, (x^*\pi_{\mathcal{G}}(a) \otimes 1)\alpha_X(y)U_{\mathcal{K}}(\rho_{\mathcal{K}}(c)\eta \otimes \zeta')\rangle = \langle (x \otimes_B \xi), \pi_{\boxtimes}(a)U_{\boxtimes}(\omega)\rho_{\boxtimes}(c)(y \otimes_B \eta)\rangle
$$

by assumption.

Together with  $\mathcal{H}^{\infty} \boxtimes \mathcal{K} \cong (\mathcal{H} \boxtimes \mathcal{K})^{\infty}$ , this immediately implies [\(6.1\)](#page-38-1). If moreover K is a G-W<sup>\*</sup>-Morita equivalence, we use the above result in combination with

$$
(\mathcal{G} \boxtimes \mathcal{K}) \boxtimes \overline{\mathcal{K}} \cong \mathcal{G} \boxtimes (\mathcal{K} \boxtimes \overline{\mathcal{K}}) \cong \mathcal{G} \boxtimes E_C^{\mathbb{G}} \cong \mathcal{G},
$$

using associativity of  $\boxtimes$  and Proposition [5.10,](#page-29-2) and similarly for  $\mathcal{H}$ .

For composition on the left, we can use Proposition [6.2](#page-37-0) and Corollary [5.11.](#page-29-3)

<span id="page-38-0"></span>We now consider the crossed product construction.

**Theorem 6.4.** If  $H, G$  are  $\mathbb{G}\text{-}A$ -B-correspondences, then

$$
\mathcal{G} \preccurlyeq \mathcal{H} \quad \Leftrightarrow \quad \mathcal{G}^{\rtimes} \preccurlyeq \mathcal{H}^{\rtimes}.
$$

*Proof.* By Theorem [5.24](#page-33-2) and Proposition [6.3,](#page-38-2) we only need to prove that  $\mathcal{G} \preccurlyeq \mathcal{H}$  implies  $\mathcal{G}^{\rtimes} \preccurlyeq \mathcal{H}^{\rtimes}$ . By replacing H with  $\mathcal{H}^{\infty}$ , we may assume that  $\mathcal{H}^{\infty} \cong \mathcal{H}$ .

By Theorem [3.7,](#page-21-0) we may then pick a contractive net  $y_i \in X = X_B(\mathcal{G}, \mathcal{H})$  such that

$$
(y_i^* \pi_{\mathcal{H}}(a) \otimes 1) \alpha_X(y_i) \stackrel{\sigma\text{-weakly}}{\underset{i \to \infty}{\longrightarrow}} \pi_{\mathcal{G}}(a) \otimes 1, \qquad a \in A.
$$

Put  $z_i = \alpha_X(y_i)$ . Then  $z_i \in X_{B^{\rtimes}}(\mathcal{G}^{\rtimes}, \mathcal{H}^{\rtimes})$  is a contractive net, and for all  $a \in A$  one has

$$
(z_i^*(\pi_{\mathcal{H}} \otimes \pi_M)\alpha(a))_{12}V_{23}z_{i,12} \stackrel{\sigma\text{-weakly}}{\longrightarrow} (\pi_{\mathcal{G}} \otimes \pi_M)(\alpha(a))_{12}V_{23}.
$$
 (6.2)

In particular, we see that for all  $a \in A$  and all  $x \in C_0^{\text{red}}(\check{\mathbb{G}})$  one has

<span id="page-38-3"></span>
$$
z_i^*(\pi_{\mathcal{H}} \otimes \pi_M)(\alpha(a))(1 \otimes \pi_{\check{M}}(x))z_i \stackrel{\sigma\text{-weakly}}{\underset{i\to\infty}{\longrightarrow}} (\pi_{\mathcal{G}} \otimes \pi_M)(\alpha(a))(1 \otimes \pi_{\check{M}}(x)).
$$
\n(6.3)

By passing to a subnet, we may assume that  $Ad_{z_i^*}$  pointwise  $\sigma$ -weakly converges to a ccp map

$$
E: B(\mathcal{H}\otimes L^2(\mathbb{G}))\to B(\mathcal{G}\otimes L^2(\mathbb{G})).
$$

Then  $1 \otimes \pi_{\tilde{M}}(C_0^{\text{red}}(\tilde{\mathbb{G}}))$  lies in the multiplicative domain of E. In particular, since  $C_0^{\text{red}}(\mathbb{G})$  acts nondegenerately on  $L^2(\mathbb{G})$ , it follows that E is ucp. It is also immediate from their definition that the  $z_i$  degenerately on  $L^2(\mathbb{G})$ , it follows that E is ucp. It is also immediate from their definition that the  $z$ intertwine  $U_{\mathcal{G}}(1\otimes x)U_{\mathcal{G}}^*$  and  $U_{\mathcal{H}}(1\otimes x)U_{\mathcal{H}}^*$  for  $x\in B(L^2(\mathbb{G}))$ . In particular,  $U_{\mathcal{H}}(1\otimes \mathcal{K}(L^2(\mathbb{G})))U_{\mathcal{H}}^*$  lies in the multiplicative domain of E.

 $\Box$ 

Now

$$
[U_{\mathcal{H}}(1 \otimes \mathcal{K}(L^{2}(\mathbb{G})))U_{\mathcal{H}}^{*}(1 \otimes C_{0}^{\text{red}}(\check{\mathbb{G}}))U_{\mathcal{H}}(1 \otimes \mathcal{K}(L^{2}(\mathbb{G})))U_{\mathcal{H}}^{*}]
$$
\n
$$
= [U_{\mathcal{H}}(1 \otimes x)(\mathrm{id} \otimes \mathrm{id} \otimes \omega)(U_{\mathcal{H},12}^{*}V_{23}U_{\mathcal{H},12})(1 \otimes y)U_{\mathcal{H}}^{*} \mid x,y \in \mathcal{K}(L^{2}(\mathbb{G})), \omega \in B(L^{2}(\mathbb{G}))_{*}]
$$
\n
$$
= [U_{\mathcal{H}}(1 \otimes x)(\mathrm{id} \otimes \mathrm{id} \otimes \omega)(U_{\mathcal{H},13}V_{23})(1 \otimes y)U_{\mathcal{H}}^{*} \mid x,y \in \mathcal{K}(L^{2}(\mathbb{G})), \omega \in B(L^{2}(\mathbb{G}))_{*}]
$$
\n
$$
= [U_{\mathcal{H}}((\mathrm{id} \otimes \omega)(U_{\mathcal{H}}(1 \otimes (\chi \otimes \mathrm{id})(V))) \otimes x)U_{\mathcal{H}}^{*} \mid x \in \mathcal{K}(L^{2}(\mathbb{G})), \omega, \chi \in B(L^{2}(\mathbb{G}))_{*}]
$$
\n
$$
= [U_{\mathcal{H}}(U_{\mathcal{H}}(\omega) \otimes x)U_{\mathcal{H}}^{*} \mid x \in \mathcal{K}(L^{2}(\mathbb{G})), \omega \in B(L^{2}(\mathbb{G}))_{*}]
$$
\n
$$
= U_{\mathcal{H}}(C_{U_{\mathcal{H}}} \otimes \mathcal{K}(L^{2}(\mathbb{G})))U_{\mathcal{H}}^{*}
$$
\n
$$
= C_{U_{\mathcal{H}}} \otimes \mathcal{K}(L^{2}(\mathbb{G})),
$$

where in the last step we used Proposition [1.2.](#page-7-3)

Letting  $C^{G}(A, B) = C_{\mathcal{H}_{u}}^{G}(A, B)$  with respect to a universal model  $(\mathcal{H}_{u}, \pi_{u}, \rho_{u}, U_{u})$ , we find from the above computation that for all  $x \in C_{U_u} \otimes \mathcal{K}(L^2(\mathbb{G}))$  we have

<span id="page-39-0"></span>
$$
E((\theta_{\mathcal{H}} \otimes \text{id})(x)) = (\theta_{\mathcal{G}} \otimes \text{id})(x). \tag{6.4}
$$

Since also  $(\pi_{\mathcal{H}} \otimes \pi_M)(\alpha(A))$  lies in the multiplicative domain of E by [\(6.3\)](#page-38-3), and

$$
[(w\otimes x)(\pi_{\mathcal{H}_u}\otimes \pi_M)(\alpha(a))(z\otimes y)\mid a\in A, w, z\in C_{U_u}, x, y\in \mathcal{K}(L^2(\mathbb{G}))]=\pi_{\mathcal{H}_u}(C_{\mathcal{H}_u}^{\mathbb{G}}(A))\otimes \mathcal{K}(L^2(\mathbb{G})),
$$

we find that [\(6.4\)](#page-39-0) in fact holds for all  $x \in C_{\mathcal{H}_u}^{\mathbb{G}}(A) \otimes \mathcal{K}(L^2(\mathbb{G}))$ .

But  $(\pi_{\mathcal{H}} \otimes id)(A \rtimes \mathbb{G}) \subseteq QM(\theta_{\mathcal{H}}(C^{\mathbb{G}}_{\mathcal{H}_{u}}(A)) \otimes \mathcal{K}(L^{2}(\mathbb{G}))),$  so we can finally conclude that for each  $x \in A \rtimes \mathbb{G}$ we have

$$
z_i^*(\pi_{\mathcal{H}} \otimes \mathrm{id})(x)z_i \xrightarrow[i \to \infty]{\sigma-\text{weakly}} E((\pi_{\mathcal{H}} \otimes \mathrm{id})(x)) = (\pi_{\mathcal{G}} \otimes \mathrm{id})(x). \tag{6.5}
$$

Since  $\alpha_{X^{\mathcal{A}}}(z_i) = z_i \otimes 1$ , we can conclude from Theorem [3.7](#page-21-0) that  $\mathcal{G}^{\mathcal{A}} \preccurlyeq \mathcal{H}^{\mathcal{A}}$ .

6.2. Left amenability revisited. Recall the definition of strong left equivariant amenability for a Gequivariant correspondence from Definition [4.4.](#page-23-2) By Remark [5.13,](#page-29-0) we obtain the following equivalent characterisation.

**Proposition 6.5.** Let  $A, B$  be  $\mathbb{G}\text{-}W^*$ -algebras. A  $\mathbb{G}\text{-}equivariant A-B-correspondence H$  is strongly left equivariantly amenable if and only if  $\mathcal{H} \boxtimes \overline{\mathcal{H}}$  equivariantly weakly contains  $E_A^{\mathbb{G}}$ .

For example, we can characterize strong equivariant amenability of an action of G on a von Neumann algebra A by the property that  $_{\alpha(A)}(L^2(A) \otimes L^2(\mathbb{G}))_{A\bar{\otimes}M}$  is strongly left G-amenable, or by the fact that the G-semi-coarse correspondence equivariantly weakly contains the trivial correspondence.

In the remainder of this section, we want to explore some other notions of 'equivariant amenability' and their interconnections. For completeness and ease of reference, we repeat the definition of (strong) G-amenability within the next definition.

<span id="page-39-1"></span>**Definition 6.6.** Let  $\mathbb{G}$  be a locally compact quantum group, and let  $(A, \alpha)$  be a  $\mathbb{G}$ -dynamical von Neumann algebra.

We say that the action of  $\mathbb G$  on A is

- (strongly) G-amenable if  $_{\alpha(A)}(L^2(A) \otimes L^2(\mathbb{G}))_{A \bar{\otimes} M}$  is (strongly) left equivariantly amenable, where  $A\bar{\otimes}M$  has the G-action determined by id  $\otimes\Delta$ ,
- (strongly) W<sup>\*</sup>-amenable if  $S_{A,\mathbb{C}}^{\mathbb{G}}$  is (strongly) left equivariantly amenable,
- (strongly)  $\mathbb{G}\text{-}\mathbf{W}^*$ -amenable if  $S^{\mathbb{G}}_{A,M}$  is (strongly) left equivariantly amenable,
- (strongly) inner G-amenable if  $_{\alpha(A)}(L^2(A) \otimes L^2(\mathbb{G}))_{A \rtimes \mathbb{G}}$  is (strongly) left equivariantly amenable, where  $A \rtimes \mathbb{G}$  has the adjoint  $\mathbb{G}\text{-action}$  [\(1.9\)](#page-8-4).

So, making use of the second column of Table [1,](#page-37-1) we see that  $(A, \alpha)$  is

- strongly G-amenable iff  $E_A^{\mathbb{G}}$  is weakly contained in the G-semi-coarse G-A-A-correspondence  $S_A^{\mathbb{G}}$ ,
- strongly W<sup>∗</sup>-amenable iff  $E_A^{\mathbb{G}}$  is weakly contained in the W<sup>∗</sup>-semi-coarse G-A-A-correspondence  $S_{A,A}^{\mathbb{G}}$ ,
- strongly  $\mathbb{G}\text{-}\mathbf{W}^*$ -amenable iff  $E_A^{\mathbb{G}}$  is weakly contained in the coarse  $\mathbb{G}\text{-}A$ -A-correspondence  $C_{A,A}^{\mathbb{G}}$ , and
- strongly inner G-amenable iff  $E_A^{\mathbb{G}}$  is weakly contained in the adjoint G-A-A-correspondence  $D_A^{\mathbb{G}}$ .

On the other hand, referring back to Definition [4.4,](#page-23-2) we see immediately from Definition [6.6](#page-39-1) that  $(A, \alpha)$  is

- G-amenable iff there exists a G-equivariant conditional expectation  $A\overline{\otimes}M \to \alpha(A)$ .
- W<sup>\*</sup>-amenable iff there exists a G-equivariant conditional expectation  $B(L^2(A)) \to A$ . Here  $B(L^2(A))$ is endowed with the G-action  $x \mapsto U_{\alpha}(x \otimes 1)U_{\alpha}^*$ .
- G-W<sup>∗</sup> -amenable iff there exists a G-equivariant conditional expectation

$$
B(L^2(A))\bar{\otimes}M\to A\otimes\mathbb{C},
$$

where the left hand side is endowed with the G-action

$$
B(L^2(A))\bar{\otimes}M \to (B(L^2(A))\bar{\otimes}M)\bar{\otimes}M, \qquad x \mapsto U_{\alpha,13}V_{23}x_{12}V_{23}^*U_{\alpha,13}^*.
$$

• inner G-amenable iff there exists a G-equivariant conditional expectation  $E: A \rtimes \mathbb{G} \to \alpha(A)$ , where  $A \rtimes \mathbb{G}$  is endowed with the adjoint G-action.

Recall that from Theorem [4.7,](#page-24-2) it follows that the latter amenability properties follow from their respective strong version.

For completeness, we include also the following obvious definition.

**Definition 6.7.** Let  $\mathbb{G}$  be a locally compact quantum group, and let  $(A, \alpha)$  be a  $\mathbb{G}$ -W<sup>\*</sup>-algebra.

We say that  $(A, \alpha)$  has G-equivariant property (T) if whenever a G-A-A-correspondence H contains  $E_A$ weakly, then  $E_A \subseteq \mathcal{H}$  as a G-equivariant subcorrespondence.

Note the following:

- If  $\mathbb{G} = \{e\}$  is trivial, then strong  $\mathbb{G}\text{-}W^*$ -amenability of A is the same as strong  $W^*$ -amenability of A, and simply means that the trivial A-A-correspondence  $L^2(A)$  is weakly contained in the coarse correspondence  $L^2(A) \otimes L^2(A)$ , i.e. A is amenable (=semi-discrete) as a von Neumann algebra [\[EL77\]](#page-47-1).
- If  $A = \mathbb{C}$ , we are considering the action of  $\mathbb{G}$  on a point •, and strong  $\mathbb{G}$ -amenability of the action is the same as strong  $\mathbb{G}\text{-}W^*$ -amenability. Moreover, the action  $\bullet \curvearrowleft \mathbb{G}$  is strongly  $\mathbb{G}\text{-}$ amenable if and only if G is strongly amenable as a locally compact quantum group, i.e. the trivial representation is weakly contained in the regular representation (this is also known as *co-amenability of*  $\hat{G}$ ), see [\[Rua96,](#page-47-9) [DQV02,](#page-46-10) [Tom06\]](#page-47-10).
- The trivial action  $\curvearrowleft \mathbb{G}$  is strongly inner G-amenable if and only if  $\mathbb{G}$  is strongly inner amenable (in the sense of locally compact quantum groups [\[OOT17\]](#page-47-38)). Similarly, the trivial action  $\bullet \curvearrowleft \mathbb{G}$  is inner G-amenable if G is inner amenable (in the sense of  $[Cr19]$ ).
- If  $A = \mathbb{C}$ , then G-equivariant property  $(T)$  is the same as the usual property  $(T)$  for locally compact quantum groups [\[CN15\]](#page-46-11) (see also [\[Fim10,](#page-47-11) [KyS12\]](#page-47-12)).

**Remark 6.8.** It is less clear to us how to introduce the Haagerup property in the setting of equivariant correspondences. It should combine the notions in [\[DFSW16\]](#page-46-12) and [\[OOT17\]](#page-47-38).

We collect now some theorems illustrating the connections between these various approximation properties.

**Theorem 6.9.** Let  $(A, \alpha)$  be a  $\mathbb{G}\text{-}W^*$ -algebra.

(1) If  $(A, \alpha)$  is strongly W<sup>\*</sup>-amenable and strongly G-amenable, then  $(A, \alpha)$  is strongly G-W<sup>\*</sup>-amenable.

(2)  $(A, \alpha)$  is W<sup>\*</sup>-amenable and G-amenable if and only if  $(A, \alpha)$  is G-W<sup>\*</sup>-amenable.

*Proof.* Under the assumptions in the first item, we have that  $E_A^{\mathbb{G}}$  is weakly contained in  $S_A^{\mathbb{G}}$  and in  $S_{A,A}^{\mathbb{G}}$ . By Example [5.14](#page-29-4) and Example [5.17,](#page-30-3) together with Proposition [6.3,](#page-38-2) we see that  $E_A^{\mathbb{G}}$  is weakly contained in  $C_{A,A}^{\mathbb{G}}$ , i.e.  $(A, \alpha)$  is strongly  $\mathbb{G}\text{-}\mathbf{W}^*$ -amenable.

Let us now prove the second item. Assume first that  $(A, \alpha)$  is both W<sup>\*</sup>-amenable and G-amenable. This means that there exists a G-equivariant ucp map  $E : (A \bar{\otimes} L^{\infty}(\mathbb{G}), id \otimes \Delta) \to (A, \alpha)$  such that  $E \circ \alpha = id_A$ and a G-equivariant ucp map  $\varphi: B(L^2(A)) \to A$  such that  $\varphi \circ \pi_A = id_A$ . Define then the ucp map

$$
\psi: B(L^2(A))\bar{\otimes}L^{\infty}(\mathbb{G})\to A: z\mapsto E((\varphi\otimes id)(U_{\alpha}zU_{\alpha}^*)).
$$

If  $a \in A$ , we have

$$
\psi(\pi_A(a)\otimes 1) = E((\varphi\otimes \mathrm{id})(\pi_A\otimes \mathrm{id})(\alpha(a))) = E(\alpha(a)) = a
$$

and if  $z \in B(L^2(A)) \bar{\otimes} L^{\infty}(\mathbb{G})$ , we have

$$
(\psi \otimes id)(U_{\alpha,13}V_{23}z_{12}V_{23}^*U_{\alpha,13}^*) = (E \otimes id)(\varphi \otimes id \otimes id)(U_{\alpha,12}U_{\alpha,13}V_{23}z_{12}V_{23}^*U_{\alpha,13}^*U_{\alpha,12}^*)
$$
  
\n
$$
= (E \otimes id)(\varphi \otimes id \otimes id)(id \otimes \Delta)(U_{\alpha}zU_{\alpha}^*)
$$
  
\n
$$
= (E \otimes id)(id \otimes \Delta)(\varphi \otimes id)(U_{\alpha}zU_{\alpha}^*)
$$
  
\n
$$
= \alpha(E(\varphi \otimes id)(U_{\alpha}zU_{\alpha}^*)) = \alpha(\psi(z))
$$

so  $\psi$  is a G-equivariant ucp map that witnesses the G-W<sup>\*</sup>-amenability of  $(A, \alpha)$ .

Conversely, assume that  $(A, \alpha)$  is  $\mathbb{G}-W^*$ -amenable, i.e. there exists a ucp  $\mathbb{G}$ -equivariant map

 $\psi: B(L^2(A))\overline{\otimes}L^{\infty}(\mathbb{G})\to A$ 

that satisfies  $\psi(\pi_A(a) \otimes 1) = a$  for all  $a \in A$ . Define

$$
E: A \bar{\otimes} L^{\infty}(\mathbb{G}) \to A: z \mapsto \psi(U_{\alpha}^{*}(\pi_{A} \otimes id)(z)U_{\alpha}).
$$

Then if  $a \in A$ , we have

$$
E(\alpha(a)) = \psi(\pi_A(a) \otimes 1) = a
$$

and if  $z \in A \bar{\otimes} L^{\infty}(\mathbb{G})$ , we have

$$
(E \otimes id)(id \otimes \Delta)(z) = (E \otimes id)(U_{\alpha,12}^*(\pi_A \otimes id \otimes id)(id \otimes \Delta)(z)U_{\alpha,12})
$$
  
\n
$$
= (\psi \otimes id)((U_{\alpha,12}^*(id \otimes \Delta)(\pi_A \otimes id)(z)U_{\alpha,12})
$$
  
\n
$$
= (\psi \otimes id)(U_{\alpha,13}(id \otimes \Delta)(U_{\alpha}^*(\pi_A \otimes id)(z)U_{\alpha})U_{\alpha,13}^*)
$$
  
\n
$$
= \alpha(\psi(U_{\alpha}^*(\pi_A \otimes id)(z)U_{\alpha})) = \alpha(E(z)).
$$

Therefore, the map E witnesses the G-amenability of  $(A, \alpha)$ . On the other hand, define the ucp map

$$
\varphi: B(L^2(A)) \to A: x \mapsto \psi(x \otimes 1).
$$

Then clearly  $\varphi \circ \pi_A = id_A$  and if  $x \in B(L^2(A))$ , we have

$$
(\varphi \otimes id)(U_{\alpha}(x \otimes 1)U_{\alpha}^{*}) = (\psi \otimes id)(U_{\alpha,13}(x \otimes 1 \otimes 1)U_{\alpha,13}^{*})
$$
  

$$
= (\psi \otimes id)(U_{\alpha,13}V_{23}(x \otimes 1 \otimes 1)V_{23}^{*}U_{\alpha,13}^{*})
$$
  

$$
= \alpha(\psi(x \otimes 1)) = \alpha(\varphi(x))
$$

so the map  $\varphi$  witnesses the W<sup>\*</sup>-amenability of  $(A, \alpha)$ .

**Remark 6.10.** We do not know if, in general, strong  $\mathbb{G}-W^*$ -amenability implies strong  $\mathbb{G}-\text{amenability}$  and strong W<sup>\*</sup>-amenability. See however statements (3) and (4) in Theorem [6.12.](#page-43-0)

In what follows, we fix a locally compact quantum group G and we will use the notations

$$
\Delta_r : B(L^2(\mathbb{G})) \to B(L^2(\mathbb{G})) \bar{\otimes} M : x \mapsto V(x \otimes 1)V^*,
$$
  

$$
\Delta_r : B(L^2(\mathbb{G})) \to B(L^2(\mathbb{G})) \bar{\otimes} \check{M} : x \mapsto \check{V}(x \otimes 1)\check{V}^*,
$$
  

$$
\Delta_l : B(L^2(\mathbb{G})) \to M \bar{\otimes} B(L^2(\mathbb{G})) : x \mapsto W^*(1 \otimes x)W.
$$

We also recall that if  $(A, \alpha)$  is a G-von Neumann algebra, then the G-fixed points of  $A \rtimes_{\alpha} \mathbb{G}$  with respect to the  $\check{\mathbb{G}}$ -coaction  $\alpha^{\mathcal{A}} = id \otimes \check{\Delta}_r$  is exactly  $\alpha(A)$  and we have

$$
A^{\rtimes} = A \rtimes_{\alpha} \mathbb{G} = \{ z \in A \bar{\otimes} B(L^{2}(\mathbb{G})) : (\alpha \otimes id)(z) = (id \otimes \Delta_{l})(z) \}.
$$

<span id="page-42-2"></span>**Theorem 6.11.** Let  $\mathbb{G}$  be a locally compact quantum group, and let  $A, B$  be  $\mathbb{G}$ -dynamical von Neumann algebras. Let  $\mathcal{H} = (\mathcal{H}, U, \pi, \rho)$  be a G-equivariant A-B-correspondence.

- (1) H is strongly left  $\mathbb{G}$ -amenable if and only if  $\mathcal{H}^{\rtimes}$  is strongly left  $\check{\mathbb{G}}$ -amenable.
- (2) H is left G-amenable if and only if  $\mathcal{H}^{\rtimes}$  is left G-amenable.

Proof. We first prove the first item. By Proposition [5.22](#page-31-2) and Theorem [5.23,](#page-31-1) we have that

$$
(\mathcal{H}\boxtimes\overline{\mathcal{H}})^{\rtimes}\cong\mathcal{H}^{\rtimes}\boxtimes\overline{\mathcal{H}^{\rtimes}}.
$$

So by Proposition [5.28](#page-35-1) and Theorem [6.4,](#page-38-0) we see that  $E_A^{\mathbb{G}}$  is equivariantly weakly contained in  $\mathcal{H} \boxtimes \overline{\mathcal{H}}$  if and only if  $E_{A^{\rtimes}}^{\mathbb{G}}$  is equivariantly weakly contained in  $\mathcal{H}^{\rtimes} \boxtimes \overline{\mathcal{H}^{\rtimes}}$ , which is exactly the statement in the first item.

Next, we prove the second item. Note that the equivalence in (2) means the equivalence of the following two items:

- (a) There exists a G-equivariant ucp conditional expectation  $\rho(B)' \to \pi(A)$ .
- (b) There exists a  $\tilde{\mathbb{G}}$ -equivariant ucp conditional expectation  $\rho^{\mathfrak{A}}(B^{\mathfrak{A}})' \to \pi^{\mathfrak{A}}(A^{\mathfrak{A}})$ .

Before proving the equivalence of these statements, we need to do some preparatory work. Let us write  $\tilde{R}$ for the anti-unitary on  $B(L^2(\mathbb{G}))$  given by  $\widetilde{R}(x) = Jx^*J$ . Given a subset S of a von Neumann algebra, we write  $[S]^\sigma$  for the  $\sigma$ -weak closure of the linear span of S. Then note that

$$
\rho^{\mathcal{A}}(B^{\mathcal{A}}) = U(\rho \otimes \widetilde{R})([\beta(B)(1 \otimes \widetilde{M})]^{\sigma})U^*
$$
  
= 
$$
[U(1 \otimes \widehat{M})U^*U(\rho \otimes R)(\beta(B))U^*]^{\sigma} = [U(1 \otimes \widehat{M})U^*(\rho(B) \otimes 1)]^{\sigma}
$$

whence  $\rho^{\aleph}(B^{\aleph})' = [U(B(\mathcal{H}) \bar{\otimes} \check{M})U^*] \cap [\rho(B)' \bar{\otimes} B(L^2(\mathbb{G}))]$ . Next, note that if  $z \in \rho^{\aleph}(B^{\aleph})'$ , then

<span id="page-42-0"></span>
$$
W_{23}^* z_{13} W_{23} = U_{12} z_{13} U_{12}^*.
$$
\n(6.6)

Indeed, if we put  $X := U^* z U$ , then [\(6.6\)](#page-42-0) is equivalent with

<span id="page-42-1"></span>
$$
W_{23}^* U_{13} X_{13} U_{13}^* W_{23} = U_{12} U_{13} X_{13} U_{13}^* U_{12}^*.
$$
\n
$$
(6.7)
$$

But since  $X \in B(H) \bar{\otimes} \check{M}$  and  $W \in M \bar{\otimes} \hat{M}$ , we find

ρ

$$
U_{12}U_{13}X_{13}U_{13}^*U_{12}^* = (\mathrm{id} \otimes \Delta)(U)X_{13}(\mathrm{id} \otimes \Delta)(U^*) = W_{23}^*U_{13}W_{23}X_{13}W_{23}^*U_{13}^*W_{23} = W_{23}^*U_{13}X_{13}U_{13}^*W_{23}
$$
 and (6.7) follows

and [\(6.7\)](#page-42-1) follows.

Assume now that (a) holds, and choose a G-equivariant ucp conditional expectation  $\phi : \rho(B)' \to \pi(A)$ . Define the map

$$
\psi := \phi \otimes id : \rho(B)' \bar{\otimes} B(L^2(\mathbb{G})) \to \pi(A) \bar{\otimes} B(L^2(\mathbb{G})).
$$

Let us write  $\gamma: B(\mathcal{H}) \to B(\mathcal{H}) \bar{\otimes} M : x \mapsto U(x \otimes 1)U^*$ . If  $z \in \rho^{\rtimes} (B^{\rtimes})' \subseteq \rho(B)' \bar{\otimes} B(L^2(\mathbb{G}))$ , then using [\(6.6\)](#page-42-0) and G-equivariance of  $\phi$ , we get

$$
(\mathrm{id}_{\pi(A)} \otimes \Delta_l)\psi(z) = (\phi \otimes \mathrm{id} \otimes \mathrm{id})(\mathrm{id} \otimes \Delta_l)(z) = (\phi \otimes \mathrm{id} \otimes \mathrm{id})(U_{12}z_{13}U_{12}^*) = (\gamma \otimes \mathrm{id})\psi(z)
$$

so  $\psi(z) \in \pi(A) \rtimes_{\gamma} \mathbb{G} = \pi^{\rtimes}(A^{\rtimes}).$  Thus,  $\psi = \phi \otimes$  id restricts to a map

$$
\psi : \rho^{\rtimes} (B^{\rtimes})' \to \pi^{\rtimes} (A^{\rtimes})
$$

which trivially is a ucp  $\check{\mathbb{G}}$ -equivariant conditional expectation. Thus, (b) holds.

Conversely, assume that (b) holds, and choose a ucp  $\check{G}$ -equivariant ucp conditional expectation  $\psi : \rho^{\rtimes} (B^{\rtimes})' \to$  $\pi^{\mathcal{A}}(A^{\mathcal{A}})$ . Since  $U(\rho(B)^{\prime} \otimes 1)U^{\ast} \subseteq \rho^{\mathcal{A}}(B^{\mathcal{A}})^{\prime}$ , it makes sense to define

$$
\widetilde{\psi}: \rho(B)' \to \pi^{\rtimes}(A^{\rtimes}) : x \mapsto \psi(U(x \otimes 1)U^*).
$$

Note that since  $\check{V}_{23}$  and  $U_{12}$  commute, we have for  $x \in \rho(B)'$  that

$$
(\mathrm{id}\otimes \Delta_r)\widetilde{\psi}(x) = (\mathrm{id}\otimes \Delta_r)\psi(U(x\otimes 1)U^*)
$$
  
\n
$$
= (\psi\otimes \mathrm{id})(\mathrm{id}\otimes \Delta_r)(U(x\otimes 1)U^*)
$$
  
\n
$$
= (\psi\otimes \mathrm{id})(\check{V}_{23}U_{12}(x\otimes 1\otimes 1)U_{12}^*\check{V}_{23}^*)
$$
  
\n
$$
= \psi(U(x\otimes 1)U^*)\otimes 1 = \widetilde{\psi}(x)\otimes 1
$$

so that  $\widetilde{\psi}(x) \in \gamma(\pi(A))$ . Next, note that since  $\psi(1 \otimes y) = 1 \otimes y$  for all  $y \in \check{M}$ , we have that  $V_{23} \in$  $M(1 \otimes C_0^{\text{red}}(\tilde{\mathbb{G}}) \otimes C_0^{\text{red}}(\mathbb{G}))$  is in the multiplicative domain of  $\psi \otimes id$ , so that for  $x \in \rho(B)'$ ,

$$
(\tilde{\psi} \otimes id)(U(x \otimes 1)U^*) = (\psi \otimes id)(U_{12}U_{13}(x \otimes 1 \otimes 1)U_{13}^*U_{12}^*)
$$
  
=  $(\psi \otimes id)(id \otimes \Delta)(U(x \otimes 1)U^*)$   
=  $V_{23}((\psi \otimes id)(U(x \otimes 1)U^*) \otimes 1)V_{23}^* = (id \otimes \Delta)\tilde{\psi}(x).$ 

Thus, the map  $\tilde{\psi}: \rho(B)' \to \gamma(\pi(A))$  is G-equivariant. Composing with the G-isomorphism

$$
\gamma^{-1} : (\gamma(\pi(A)), \mathrm{id} \otimes \Delta) \cong (\pi(A), \gamma),
$$

we find that  $\phi := \gamma^{-1} \circ \tilde{\psi} : \rho(B)' \to \pi(A)$  is G-equivariant, and it satisfies

$$
\phi(\pi(a)) = \gamma^{-1}(\widetilde{\psi}(\pi(a))) = \gamma^{-1}(\psi((\pi \otimes \text{id})(\alpha(a)))) = \gamma^{-1}((\pi \otimes \text{id})(\alpha(a))) = \gamma^{-1}(\gamma(\pi(a))) = \pi(a)
$$
  
for every  $a \in A$ . Thus, (a) is satisfied.

<span id="page-43-0"></span>**Theorem 6.12.** Let  $\mathbb{G}$  be a locally compact quantum group, and A a  $\mathbb{G}$ -dynamical von Neumann algebra. Then the following equivalences hold:

- (1) A is (strongly) G-amenable if and only if  $A \rtimes \mathbb{G}$  is (strongly) inner  $\check{G}$ -amenable.
- (2) A is (strongly) inner  $\mathbb{G}$ -amenable if and only if  $A \rtimes \mathbb{G}$  is (strongly)  $\mathbb{G}$ -amenable.
- (3) A is (strongly)  $\mathbb{G}-W^*$ -amenable if and only if  $A\rtimes\mathbb{G}$  is (strongly)  $W^*$ -amenable (with respect to the dual  $\mathbb{G}\text{-action}$ ).
- (4) A is strongly  $\mathbb{G}\text{-}W^*$ -amenable if and only if  $A \rtimes \mathbb{G}$  is strongly  $\check{\mathbb{G}}\text{-}W^*$ -amenable.

Proof. This follows directly from Table [1](#page-37-1) upon using Theorem [6.11.](#page-42-2) The only things that need a small comment, are

- that, for the last equivalence, the multiplicity factor  $L^2(\mathbb{G})$  does not matter, see Remark [3.2,](#page-18-1) and
- that, for the third equivalence, left  $\check{\mathbb{G}}$ -amenability of  $S^{\check{\mathbb{G}}}_{A^{\rtimes},\mathbb{C}} \boxtimes_{\mathbb{C}} L^2(\mathbb{G})_{M^{\rtimes}},$  whose underlying Hilbert space is  $L^2(A^{\rtimes}) \otimes L^2(\check{G})$ , is the same as left  $\check{G}$ -amenability of  $S^{\check{G}}_{A^{\rtimes},\mathbb{C}}$ , since

$$
(1 \otimes M^{\rtimes})' = B(L^2(A)) \otimes 1.
$$

 $\Box$ 

We complement the non-strong version of the above theorem with the following remark:

Remark 6.13. • The last equivalence in Theorem [6.12](#page-43-0) is also true in the non-strong setting, although we do not know a quick proof using the machinery of this paper. The proof will be given in follow-up work by one of the authors.

- Theorem [6.12](#page-43-0) unifies and generalizes several results that are already spread throughout the literature. For instance, the equivalence (1) in the above theorem was shown in [\[Moa18,](#page-47-32) Theorem 7.5] for  $\mathbb{G}$  a discrete quantum group, and was also proven in the setting of G-operator systems in [\[DH24,](#page-46-31) Theorem 4.2] where G is a discrete quantum group. A similar result also appeared in [\[BC21,](#page-46-32) Theorem 5.2, Corollary 5.3] in the setting of locally compact groups (and where the authors note that the techniques extend to co-amenable locally compact quantum groups).
- In [\[MP22\]](#page-47-39), a notion of inner amenability for actions of locally compact groups on von Neumann algebras is introduced. However, the definition given in this paper is not the same one as in Definition [6.6.](#page-39-1) That both definitions nevertheless agree (in the setting of locally compact groups) follows by combining Theorem [6.12](#page-43-0) (2) with [\[MP22,](#page-47-39) Proposition 4.3].

We end this paper by noting that it is also possible to prove the first three equivalences in the non-strong setting in Theorem [6.12](#page-43-0) directly. This is interesting, as it gives us an explicit bridge to move between certain conditional expectations. For completeness, we briefly describe for each of the equivalences how to do this. We refrain from spelling out all details.

(1)  $(A, \alpha)$  being G-amenable is equivalent to the existence of a G-equivariant ucp conditional expectation

$$
\varphi: (A \bar{\otimes} L^{\infty}(\mathbb{G}), \mathrm{id} \otimes \Delta) \to (\alpha(A), \mathrm{id} \otimes \Delta).
$$

Then the map

$$
\psi := \varphi \rtimes \mathbb{G} : A \bar{\otimes} B(L^2(\mathbb{G})) \cong (A \bar{\otimes} L^{\infty}(\mathbb{G})) \rtimes \mathbb{G} \to \alpha(A) \rtimes \mathbb{G} \cong A \rtimes \mathbb{G}
$$

is a  $\check{G}$ -equivariant ucp conditional expectation. Conversely, if  $\psi : A \bar{\otimes} B(L^2(\mathbb{G})) \to A \rtimes \mathbb{G}$  is a  $\check{G}$ equivariant ucp conditional expectation, then it restricts to a  $\mathbb{G}$ -equivariant<sup>[5](#page-44-0)</sup> conditional expectation

$$
\varphi: A\bar{\otimes} L^{\infty}(\mathbb{G}) \to \alpha(A)
$$

between the Gˇ -fixed points. On the other hand, the Takesaki-Takai isomorphism

$$
\Phi: (A \rtimes \mathbb{G}) \rtimes \check{\mathbb{G}} \to A \bar{\otimes} B(L^2(\mathbb{G}))
$$

is  $\mathbb{G}\text{-equivariant}$  (where the iterated crossed product has the adjoint  $\mathbb{G}\text{-action}$  - this follows by an explicit calculation on generators) with  $\Phi^{-1}|_{A\rtimes\mathbb{G}} = id \otimes \check{\Delta}_r$ , so the existence of a  $\check{\mathbb{G}}$ -equivariant ucp conditional expectation  $\psi : A \bar{\otimes} B(L^2(\mathbb{G})) \to A \rtimes \mathbb{G}$  is equivalent with the existence of a  $\check{\mathbb{G}}$ -equivariant ucp conditional expectation

$$
(A\rtimes\mathbb G)\rtimes\check{\mathbb G}\to\alpha^\rtimes(A\rtimes\mathbb G)
$$

which is to say that  $(A \rtimes \mathbb{G}, \alpha^{\rtimes})$  is inner  $\check{\mathbb{G}}$ -amenable.

(2)  $(A, \alpha)$  being inner G-amenable is equivalent to the existence of a G-equivariant ucp conditional expectation

$$
\varphi: A \rtimes \mathbb{G} \to \alpha(A).
$$

This is equivalent with the existence of a  $\mathbb{G}$ -equivariant ucp conditional expectation

<span id="page-44-1"></span>
$$
(A \rtimes \mathbb{G}) \rtimes \mathbb{G} \to \alpha(A) \rtimes \mathbb{G} \cong A \rtimes \mathbb{G}.\tag{6.8}
$$

Note however that the map

$$
\Psi: (A \rtimes \mathbb{G}) \bar{\otimes} L^{\infty}(\check{\mathbb{G}}) \to (A \rtimes \mathbb{G}) \rtimes \mathbb{G} : z \mapsto V_{23}zV_{23}^*
$$

is a  $\check{G}$ -equivariant \*-isomorphism such that  $\Psi \circ \alpha^{\mu} = \alpha \otimes id$  on  $A \rtimes \mathbb{G}$ . Therefore, the existence of a  $\mathbb{G}\text{-}\mathrm{equivariant}$  ucp map as in [\(6.8\)](#page-44-1) is equivalent with the existence of a  $\mathbb{G}\text{-}\mathrm{equivariant}$  ucp map

 $\psi: (A\rtimes \mathbb G)\bar{\otimes} L^\infty(\check{\mathbb G}) \to A\rtimes \mathbb G$ 

such that  $\psi \circ \alpha^{\mathcal{A}} = id$ , which is to say that  $(A \rtimes \mathbb{G}, \alpha^{\mathcal{A}})$  is  $\check{\mathbb{G}}$ -amenable.

<span id="page-44-0"></span><sup>&</sup>lt;sup>5</sup>Making use of the fact that  $V \in M(C_0^{\text{red}}(\mathbb{G})) \otimes C_0^{\text{red}}(\mathbb{G}))$ , we see that  $V_{23}$  is contained in the multiplicative domain of  $\psi \otimes \text{id}$ with  $(\psi \otimes id)(V_{23}) = V_{23}$ . Hence,  $\psi$  preserves the G-action id  $\otimes \Delta_r$  and thus its restriction  $\varphi$  preserves the G-action id  $\otimes \Delta$ .

(3) Consider the coaction

$$
\beta: B(L^2(A))\bar{\otimes}L^{\infty}(\mathbb{G})\to (B(L^2(A))\bar{\otimes}L^{\infty}(\mathbb{G}))\bar{\otimes}L^{\infty}(\mathbb{G}): z\mapsto U_{\alpha,13}V_{23}z_{12}V_{23}^*U_{\alpha,13}^*.
$$

If  $A$  is  $\mathbb{G}\text{-}W^*$ -amenable, there is a  $\mathbb{G}\text{-}\text{equivariant ucp map}$ 

$$
\varphi: (B(L^2(A))\bar{\otimes}L^{\infty}(\mathbb{G}), \beta) \to (\alpha(A), \mathrm{id}\otimes\Delta)
$$

such that  $\varphi(\pi_A(a) \otimes 1) = \alpha(a)$  for all  $a \in A$ . Define then

$$
\widetilde{\varphi}: B(L^2(A))\bar{\otimes}L^{\infty}(\mathbb{G})\to \alpha(A): z\mapsto \varphi(U_{\alpha}^*zU_{\alpha}).
$$

Then it can be checked that  $\tilde{\varphi}: (B(L^2(A))\bar{\otimes}L^{\infty}(\mathbb{G}), \mathrm{id}\otimes\Delta) \to (\alpha(A), \mathrm{id}\otimes\Delta)$  is a G-equivariant ucp map satisfying  $\tilde{\varphi}((\pi_A \otimes id)\alpha(a)) = \alpha(a)$  for  $a \in A$ . Then the map

$$
\widetilde{\varphi} \rtimes \mathbb{G}: B(L^2(A)) \bar{\otimes} B(L^2(\mathbb{G})) \cong (B(L^2(A)) \bar{\otimes} L^{\infty}(\mathbb{G})) \rtimes \mathbb{G} \to \alpha(A) \rtimes \mathbb{G} \cong A \rtimes \mathbb{G}
$$

is a  $\mathbb{G}$ -equivariant ucp map such that  $(\widetilde{\varphi} \rtimes \mathbb{G})((\pi_A \otimes id)(z)) = z$  for  $z \in A \rtimes \mathbb{G}$ . This means that  $A \rtimes \mathbb{G}$  is  $W^*$ -amenable. Conversely, if  $A \rtimes \mathbb{G}$  is  $W^*$ -amenable, then there exists a  $\check{G}$ -equivariant ucp map

$$
\psi : (B(L^2(A)) \bar{\otimes} B(L^2(\mathbb{G})), \mathrm{id} \otimes \check{\Delta}_r) \to (A \rtimes \mathbb{G}, \alpha^{\rtimes})
$$

such that  $\psi((\pi_A \otimes id)(z)) = z$  for  $z \in A \rtimes \mathbb{G}$ . Taking  $\check{G}$ -fixed points, we obtain a G-equivariant ucp map

$$
\widetilde{\psi}: B(L^2(A)) \bar{\otimes} L^{\infty}(\mathbb{G}) \to \alpha(A)
$$

and then the map

$$
\varphi: (B(L^2(A))\bar{\otimes}L^{\infty}(\mathbb{G}), \beta) \to (\alpha(A), \mathrm{id}\otimes\Delta): z \mapsto \widetilde{\psi}(U_{\alpha}zU_{\alpha}^*)
$$

is G-equivariant and satisfies  $\varphi(\pi_A(a) \otimes 1) = \alpha(a)$  for all  $a \in A$ , which is to say that A is G-W<sup>\*</sup>amenable.

## 7. OUTLOOK

In this paper, the general theory of equivariant  $W^*$ -correspondences and their weak containment was developed. Many interesting approximation properties of  $W^*$ -dynamical systems  $(A, \alpha)$  can then be formulated by considering which precise equivariant A-A-correspondences can weakly contain the trivial equivariant A-A-correspondence. In follow-up work of the authors, concrete applications of the theory developed in this paper will be investigated in the framework of compact and discrete quantum groups. We announce two such applications:

- If  $\mathbb G$  is a compact or a discrete quantum group and  $A \subseteq B$  is a  $\mathbb G$ -equivariant inclusion (of  $\sigma$ -finite von Neumann algebras), then strong equivariant amenability and equivariant amenability of the inclusion  $A \subseteq B$  are equivalent.
- There exists an amenable discrete quantum group that acts non-amenably on a von Neumann algebra.

A further venue for research is the considerations of the topics of this paper in the setting of C<sup>∗</sup> -dynamical systems. For the moment, this remains completely unexplored in the setting of locally compact quantum groups.

Acknowledgements: The work of K. De Commer was supported by Fonds voor Wetenschappelijk Onderzoek (FWO), grant G032919N. The work of J. De Ro is supported by the FWO Aspirant-fellowship, grant 1162522N. We thank J. Crann for an email exchange concerning the topics of this paper. We also thank the referees for valuable comments improving the exposition of the paper.

#### **REFERENCES**

- <span id="page-46-7"></span>[A-D79] C. Anantharaman-Delaroche, Action moyennable d'un groupe localement compact sur une algèbre de von Neumann, *Math. Scand.* 45 (1979), 289–304.
- <span id="page-46-4"></span>[A-D90] C. Anantharaman-Delaroche, On relative amenability for von Neumann algebras, *Compositio Math.* 74 (1990), 333-352.
- <span id="page-46-26"></span>[A-DH90] C. Anantharaman-Delaroche and J. F. Havet, On Approximate Factorizations of Completely Positive Maps, *J. Funct. Anal.* 90 (1990), 411–428
- <span id="page-46-5"></span>[A-D95] C. Anantharaman-Delaroche, Amenable correspondences and approximation properties for von Neumann algebras, *Pacific J. Math.* 171 (2) (1995), 309–341.
- <span id="page-46-8"></span>[A-DR00] C. Anantharaman-Delaroche, J. Renault, Amenable Groupoids, *Monogr. Enseign. Math.* 36, Enseignement Math., Geneva (2000).
- <span id="page-46-20"></span>[AP73] C.A. Akemann and G.K. Pedersen, Complications of semicontinuity in C\*-algebra theory, *Duke Math. J.* 40 (1973), 785–795.
- <span id="page-46-13"></span>[AS21] Y. Arano and A. Skalski, On the Baum–Connes conjecture for discrete quantum groups with torsion and the quantum Rosenberg conjecture, *Proc. Amer. Math. Soc.* 149 (2021), 5237–5254.
- <span id="page-46-17"></span>[AV16] Y. Arano and S. Vaes, C<sup>\*</sup>-tensor categories and subfactors for totally disconnected groups, In: Operator algebras and applications – the Abel Symposium 2015, *Abel Symp.* 12, Springer (2017), 1–43.
- <span id="page-46-14"></span>[BS89] S. Baaj and G. Skandalis, C∗-alg`ebres de Hopf et th´eorie de Kasparov ´equivariante, *K-Theory* 2 (1989), 683–721.
- <span id="page-46-15"></span>[BS93] S. Baaj and G. Skandalis, Unitaires multiplicatifs et dualité pour les produits croisés de C<sup>∗</sup>-algèbres, *Ann. scient. Ec. Norm. Sup.*, 4e série **26** (1993), 425-488.
- <span id="page-46-16"></span>[BC17] S. Baaj and J. Crespo, Equivalence monoidale de groupes quantiques et K-th´eorie ´ bivariante, *Bull. Soc. Math. France* 4 (145) (2017), 711-–802.
- <span id="page-46-23"></span>[BSV03] S. Baaj, G. Skandalis and S. Vaes, Non-Semi-Regular Quantum Groups Coming from Number Theory, *Comm. Math. Phys.* 235 (2003), 139–167.
- <span id="page-46-6"></span>[BMO20] J. Bannon, A. Marrakchi and N. Ozawa, *Full Factors and Co-amenable Inclusions*, *Comm. Math. Phys.* 378 (2020), 1107–1121.
- <span id="page-46-32"></span>[BC21] A. Bearden and J. Crann, A weak expectation property for operator modules, injectivity and amenable actions, *Internat. J. Math.* 32 (2021).
- <span id="page-46-19"></span>[BC22] A. Bearden and J. Crann, Amenable dynamical systems over locally compact groups, *Ergod. Th. & Dynam. Sys.* 42 (2022), 2468–2508.
- [BT03] E. B´edos and L. Tuset, Amenability and co-amenability for locally compact quantum groups, *Internat. J. Math.* 14 (8) (2003), 865–884.
- <span id="page-46-24"></span>[BdlHV08] B. Bekka, P. de le Harpe and A. Valette, Kazhdan's property (T), *New Mathematical Monographs* 11, Cambridge University Press (2008).
- <span id="page-46-29"></span>[BdlHV08b] B. Bekka, P. de le Harpe and A. Valette, Kazhdan's property (T), erratum, available at [https://perso.univ-rennes1.fr/bachir.bekka/Erratum-Kazhdan.pdf.](https://perso.univ-rennes1.fr/bachir.bekka/Erratum-Kazhdan.pdf)
- <span id="page-46-22"></span>[BDS13] M. Brannan, M. Daws and E. Samei, Completely bounded representations of convolution algebras of locally compact quantum groups, *M¨unster J. Math.* 6 (2013), 445–482.
- <span id="page-46-21"></span>[BO08] N.P. Brown and N. Ozawa, C\*-algebras and finite-dimensional approximations, *Graduate Studies in Mathematics* 88, American Mathematical Society, Providence, RI, (2008).
- <span id="page-46-28"></span>[Bro03] R.M. Brouwer, A bicategorical approach to Morita equivalence for von Neumann algebras, *J. Math. Phys.* 44 (2003), 2206–2214.
- <span id="page-46-9"></span>[BEW21] A. Buss, S. Echterhoff and R. Willett, Amenability and weak containment for actions of locally compact groups on C∗ -algebras, *Preprint* (2021), to appear in *Mem. Amer. Math. Soc.*
- <span id="page-46-27"></span>[BT03] E. Bédos and L. Tuset, Amenability and co-amenability for locally compact quantum groups, Internat. J. Math. 14 (2003), 865–884.
- <span id="page-46-11"></span>[CN15] X. Chen and C.K. Ng, Property T for locally compact quantum groups, *Inter. J. Math.* 26 (2015), 1550024.
- <span id="page-46-3"></span>[CE77] M.D. Choi and E.G. Effros, Nuclear C<sup>∗</sup> -algebras and injectivity: the general case, *Indiana Univ. Math. J.* 26 (1977), 443–446.
- <span id="page-46-0"></span>[Con80+] A. Connes, Correspondences, *unpublished*.
- <span id="page-46-1"></span>[Con94] A. Connes, Noncommutative Geometry, *Academic Press* (1994), 661 pages.
- <span id="page-46-2"></span>[Con76] A. Connes, Classification of injective factors, *Ann. of Math.* 104 (1976), 73–115.
- <span id="page-46-30"></span>[Cr19] J. Crann, Inner amenability and approximation properties of locally compact quantum groups, *Indiana Univ. Math. J.* 68 (2019), 1721–1766.
- <span id="page-46-12"></span>[DFSW16] M. Daws, P. Fima, A. Skalski, S. White, The Haagerup property for locally compact quantum groups, *J. reine angew. Math.* 711 (2016), 189–229.
- <span id="page-46-18"></span>[DCDT21] K. De Commer and J. R. Dzokou Talla, Invariant integrals on coideals and their Drinfeld doubles, *preprint*, arXiv:2112.07476.
- <span id="page-46-31"></span>[DH24] J. De Ro and L. Hataishi, Actions of compact and discrete quantum groups on operator systems, *Int. Math. Res. Not.* (2024), 1–30.
- <span id="page-46-10"></span>[DQV02] P. Desmedt, J. Quaegebeur and S. Vaes, Amenability and the bicrossed product construction, Illinois J. Math. 46 (4) (2002), 1259–1277.
- <span id="page-46-25"></span>[Dix77] J. Dixmier, C∗-algebras, North-Holland Publishing Co., Amsterdam (1977), Translated from the French by Francis Jellett, North-Holland Mathematical Library 15.
- <span id="page-47-1"></span>[EL77] E. Effros and C. Lance, Tensor products of operator algebras, *Adv. Math.* 25 (1977), 1–34.
- <span id="page-47-31"></span>[ER00] E. Effros and Z.-J. Ruan, Operator Spaces, *London Math. Soc. Monographs, New Series* 23, Oxford University Press, New York (2000).
- <span id="page-47-5"></span>[Ern67] J. Ernest, Hopf-von Neumann algebras, Proc. Conf. Functional analysis (Irvine, Calif.), *Académie Press, New-York* (1967), 195–215.
- <span id="page-47-26"></span>[Fel62] J.M.G. Fell, Weak containment and induced representations of groups, *Can. J. Math.* 14 (1962), 237–268.
- <span id="page-47-37"></span>[Fel63] J.M.G. Fell, Weak containment and Kronecker products of group representations, *Pac. J. Math.* 13 (1963), 503–510.
- <span id="page-47-36"></span>[ES80] M. Enock and J.-M. Schwartz, Produit croisé d'une algèbre de von Neumann par une algèbre de Kac II, *Publ. RIMS* 16 (1980), 189–232.
- <span id="page-47-11"></span>[Fim10] P. Fima, Kazhdan's property T for discrete quantum groups, *Internat. J. Math.* 21 (1) (2010), 47–65.
- <span id="page-47-23"></span>[GLR85] P. Ghez, R. Lima and J.E. Roberts, W∗-categories, *Pacific J. Math.* 120 (1) (1985), 79–109.
- <span id="page-47-7"></span>[Haa78] U. Haagerup, On the dual weights for crossed products of von Neumann algebras I., II, *Math. Scand.* 43 (1978), 99–118; 119–140.
- <span id="page-47-24"></span>[KaS15] P. Kasprzak, P.M. Soltan, *Quantum groups with projection on von Neumann algebra level*, *J. Math. Anal. Appl.* 427 (1) (2015), 289–306.
- <span id="page-47-21"></span>[Kus01] J. Kustermans, Locally compact quantum groups in the universal setting, *Int. J. Math.* 12 (3) (2001), 289–338.
- <span id="page-47-3"></span>[KV00] J. Kustermans and S. Vaes, Locally compact quantum groups, Ann. Sci. Ec. Norm. Supér. 33 (6) (2000), 837–934.
- <span id="page-47-4"></span>[KV03] J. Kustermans and S. Vaes, Locally compact quantum groups in the von Neumann algebraic setting, *Math. Scand.* 92 (1) (2003), 68–92.
- <span id="page-47-12"></span>[KyS12] D. Kyed and P. Soltan, Property (T) and exotic quantum group norms, *J. Noncomm. Geometry* 6 (4) (2012), 773–800.
- <span id="page-47-34"></span>[Lan01] N.P. Landsman, Bicategories of operator algebras and Poisson manifolds, in: *Mathematical Physics in Mathematics and Physics: Quantum and Operator Algebraic Aspects*, edited by R. Longo, *Fields Inst. Comm* 30 (2001), 271–286.
- <span id="page-47-18"></span>[Li89] H.X. Lin, The structure of quasimultipliers of C\*-algebras, *Trans. Amer. Math. Soc.* 315 (1989), 147–172.
- <span id="page-47-39"></span>[MP22] A. McKee, R. Pourshahami, Amenable and inner amenable actions and approximation properties for crossed products by locally compact groups, *Canad. Math. Bull.* 65 (2) R. (2022), 381–399.
- <span id="page-47-32"></span>[Moa18] M. S. M. Moakhar, Amenable actions of discrete quantum groups on von Neumann algebras, *preprint*, arXiv:1803.04828.
- <span id="page-47-16"></span>[NV02] D. Nikshych and L. Vainerman, Finite quantum groupoids and their applications, In: New directions in Hopf algebras, *Math. Sci. Res. Inst. Publ. Cambridge Univ. Press* 43, Cambridge (2002), 211–262.
- <span id="page-47-38"></span>[OOT17] R. Okayasu, N. Ozawa and R. Tomatsu, Haagerup approximation property via bimodules, *Math. Scand.* 121 (1) (2017), 75–91.
- <span id="page-47-2"></span>[OS21] N. Ozawa and Y. Suzuki, On characterizations of amenable C∗-dynamical systems and new examples, *Selecta Math. (N.S.)* 72 (92) (2021), 29 pages.
- <span id="page-47-27"></span>[Pas73] W.L. Paschke, Inner product modules over B<sup>∗</sup> -algebras, *Trans. Amer. Math. Soc.* 182 (1973) 443–468.
- <span id="page-47-19"></span>[Ped79] G. K. Pedersen, C<sup>∗</sup> -algebras and their automorphism groups, *LMS Monographs* 14, Academic Press (1979), ix + 416pp. [Pop86] S. Popa, Correspondences, *Preprint*.
- <span id="page-47-30"></span><span id="page-47-0"></span>[PS17] S. Pekka and A. Skalski, Actions of locally compact (quantum) groups on ternary rings of operators, their crossed products, and generalized Poisson boundaries, *Kyoto J. Math.* 57 (2017), 667–691.
- <span id="page-47-28"></span>[Rie74] M.A. Rieffel, Morita equivalence for C<sup>∗</sup> -algebras and W<sup>∗</sup> -algebras, *J. Pure Appl. Alg.* 5 (1974), 51–96.
- <span id="page-47-15"></span>[RV22] L. Rollier and S. Vaes, Quantum automorphism groups of connected locally finite graphs and quantizations of discrete groups, *preprint*, arXiv:2209.03770v2.
- <span id="page-47-9"></span>[Rua96] Z.-J. Ruan, Amenability of Hopf von Neumann algebras and Kac algebras, *J. Funct. Anal.* 139 (1996), 466–499.
- <span id="page-47-35"></span>[Sau83] J.-L. Sauvageot, Sur le produit tensoriel relatif d'espaces de Hilbert, *J. Operator Theory* 9 (2) (1983), 237–252.
- <span id="page-47-13"></span>[SV14] P.M. Soltan and A. Viselter, A Note on Amenability of Locally Compact Quantum Groups, *Canad. Math. Bull.* 57 (2) (2014), 424–430.
- <span id="page-47-17"></span>[Tak03] M. Takesaki, Theory of Operator Algebras II, *Springer, Berlin* (2003).
- <span id="page-47-10"></span>[Tom06] R. Tomatsu, Amenable discrete quantum groups, *J. Math. Soc. Japan* 58 (4) (2006), 949–964.
- <span id="page-47-6"></span>[Vae01] S. Vaes, The unitary implementation of a locally compact quantum group action, *J. Funct. Anal.* 180 (2001), 426–480.
- <span id="page-47-14"></span>[Vae05] S. Vaes, A new approach to induction and imprimitivity results, *J. Funct. Anal.* 229 (2005), 317–374.
- <span id="page-47-20"></span>[VV03] S. Vaes and A. Van Daele, The Heisenberg commutation relations, commuting squares and the Haar measure on locally compact quantum groups, In: *Operator algebras and mathematical physics: conference proceedings, Constanta (Romania), July 2-7, 2001*, Editors J.-M. Combes, J. Cuntz, G.A. Elliott, G. Nenciu, H. Siedentop and S. Stratila, *Theta Foundation*, Bucarest (2003), 379–400.
- <span id="page-47-33"></span>[VV07] S. Vaes and R. Vergnioux, The boundary of universal discrete quantum groups, exactness and factoriality, *Duke Math. J.* 140 (2007), 35–84.
- <span id="page-47-8"></span>[Voi77] D. Voiculescu, Amenability and Kac algebras, *Communication of the Colloque International du CNRS "Alg`ebres d'op´erateurs et leurs applications en physique math´ematique, Marseille, June 1977."*
- <span id="page-47-22"></span>[Wor96] S. L. Woronowicz, From multiplicative unitaries to quantum groups, *Internat. J. Math.* 7 (1996), 127–149.
- <span id="page-47-29"></span>[Zet93] H. Zettl, A characterization of ternary rings of operators, *Adv. in Math.* 48 (1983), 117–143.
- <span id="page-47-25"></span>[Zim84] R. Zimmer, Ergodic theory and semisimple groups, *Birkh¨auser* 81 (1984).

Vrije Universiteit Brussel *Email address*: kenny.de.commer@vub.be *Email address*: joeri.ludo.de.ro@vub.be