

CYCLES IN GRAPHS AND IN HYPERGRAPHS

E. ALKIN, S. DZHENZHER, O. NIKITENKO, A. SKOPENKOV, A. VOROPAEV

CONTENTS

| | |
|---|----|
| 1. One-dimensional cycles in graphs | 2 |
| 2. One-dimensional cycles in the square of a graph | 4 |
| 3. Cycles in hypergraphs | 6 |
| 4. Geometric digression: Cartesian products of graphs | 7 |
| 5. Two-dimensional cycles in the square of a graph | 8 |
| 6. Answers, hints and solutions | 10 |
| References | 17 |

Introduction

A (simplicial) **1-cycle** (modulo 2) in a graph is a set C of edges such that every vertex is contained in an even number of edges from C . E.g., a cycle in the sense of graph theory is a 1-cycle, but not vice versa. It is easy to check that the sum (modulo 2) of 1-cycles is a 1-cycle.

In this text we study the following problems: to find

- the number of all 1-cycles in a given graph;
- a small number of 1-cycles in a given graph such that any 1-cycle is the sum of some of them.

We also consider generalizations (of these problems) to graphs with symmetry, and to *2-cycles* in *2-dimensional hypergraphs* (all the italicized notions are defined later). Main problems are 1.1.f, 1.3.e, 1.4.b, 1.5.ab, and 3.1.cf, 3.2.cf. The peaks of this text are results on 1- and 2-cycles in *the square of a graph* (Problems 2.5.c, 5.5.cd, 5.6.ab and 5.10.d).

The notion of a 1-cycle and its generalizations have many applications in topology, for simpler ones see e.g. [Sk20, §§4.11, 6, 9], [Sk, §§1, 8, 9]. However elementary, this text is motivated by frontline of research, see [FH10, MS17, SS23] and the references therein. Open problems are 2.9.cdfg and 5.10.f.

Learning by doing problems

In this text we expose a theory as a sequence of problems, see e.g. [Sk21m, Introduction, Learning by doing problems] and the references therein. Most problems are useful theoretical facts. So this text could in principle be read even without solving problems or looking to §6 ‘Answers, hints and solutions’.

Problems are numbered, the words ‘problem’ are omitted. If a mathematical statement is formulated as a problem, then the objective is to prove this statement. Open-ended questions are called **riddles**; here one must come up with a clear wording, and a proof. If a problem is named ‘theorem’ (‘lemma’, ‘corollary’, etc.), then this statement is considered to be more important. Usually we *formulate* beautiful or important statement *before* giving a sequence of results (lemmas, assertions,

E. Alkin, S. Dzhenzher, A. Skopenkov: Moscow Institute of Physics and Technology.

O. Nikitenko: Altay Technical University.

A. Skopenkov: Independent University of Moscow, <https://users.mccme.ru/skopenko/>.

This text (except for problems marked by star) was presented in 2023 at the International Summer Conferences of the Tournament of Towns, see https://en.wikipedia.org/wiki/Tournament_of_the_Towns. We are grateful to I. Bogdanov, A. Ryabichev and O. Styrtyr for useful discussions, and to A. Ryabichev for preparing computer versions of some figures.

etc.) which constitute its *proof*. We give hints on that after the statements but we do not want to deprive you of the pleasure of finding the right moment when you finally are ready to prove the statement. In general, if you are stuck on a certain problem, try looking at the next ones; they may turn out to be helpful.

Important definitions are highlighted in **bold** for easy navigation.

On presentation at the Summer conference

Participants are welcomed to *consult* the jury on any questions on the project. If they successfully work on the project, they can get interesting *extra problems*.

A *team* working on this project may consist of any number of participants. For every solution *written for a user* marked with either ‘+’ or ‘+.’ a team gets five ‘beans’ (see recommendations in p. 3, ‘How to write a proof for a user’ of <https://www.mccme.ru/circles/oim/multicomb.pdf>). The jury may also award extra beans for beautiful solutions, solutions of hard problems, or solutions typeset in $\text{T}_\text{E}\text{X}$. The jury has infinitely many beans. Every team initially has five beans. A team may submit a solution *in oral form* or as *written for a developer* if the team has some beans. A team loses a bean with every attempt (successful or not).

1. ONE-DIMENSIONAL CYCLES IN GRAPHS

The rigorous definition of a graph is given at the beginning of §2. Before that you can work with graphs at an intuitive level.

Denote by

- $[n] := \{1, 2, \dots, n\}$;
- K_n the complete graph on the set $[n]$ of vertices;
- $K_{m,n}$ the complete bipartite graph with parts $[m]$ and $[n]'$ (we denote by A' a copy of A).

A **simple cycle** $v_1v_2 \dots v_kv_1$ of length k in a graph is a set $\{v_1v_2, v_2v_3, \dots, v_kv_1\}$ of edges such that the vertices v_1, \dots, v_k are pairwise distinct. A *simple cycle* is a simple cycle of some length. We omit ‘simple’ if this word is clear from the context. Clearly, any simple cycle is a 1-cycle. The definition of a 1-cycle is given in the introduction.

The **sum** (the sum modulo 2, or the symmetric difference) of sets A, B is

$$A + B := (A \cup B) \setminus (A \cap B).$$

- 1.1.** (a) The sum of 1-cycles is a 1-cycle.
 (b) Any 1-cycle in K_n is a sum of some cycles of length 3.
 (c) If every edge of a 1-cycle in K_n contains the vertex n , then the 1-cycle is empty.
 (d) How many 1-cycles in K_n are there?
 (e) For any vertices a, b, c, d in K_n we have $abc + abd + acd + bcd = 0$ (we denote $0 := \emptyset$).
 (f) Any linear relation on cycles of length 3 in K_n is a sum of some relations from (e).

E.g.

- the relation $123 + 124 + 134 + 235 + 245 + 345 = 0$ is the sum of relations $123 + 124 + 134 + 234 = 0$ and $234 + 235 + 245 + 345 = 0$;
- the sum of boundaries of triangles from any triangulation of the sphere or of the torus (Figure 1) is zero.

Rigorously, a *linear relation* is a set of cycles of length 3 in K_n such that every edge of K_n is contained in an even number of cycles from this set. Thus a rigorous formulation of (f) is as follows (we identify cycles of length 3 in K_n with 3-element subsets of $[n]$). Suppose that R is a set of 3-element subsets of $[n]$ such that any 2-element subset of $[n]$ is contained in an even number of those 3-element subsets that are elements of R . Then there is a family R' of 4-element subsets of $[n]$ such that a 3-element subset B of $[n]$ is contained in R if and only if B is contained in an odd number of those 4-element subsets that are elements of R' .

- 1.2.** (a) Any 1-cycle in a graph is a sum of some simple cycles.




FIGURE 1. A 7-vertices triangulation of the torus

- (b) Any 1-cycle in a graph is a sum of some simple cycles for which there are no chords. A *chord* in a simple cycle is an edge of the graph which does not belong to the cycle (i.e., an edge between two non-consecutive vertices).
- (c) Any 1-cycle in K_n is a sum of some of the following cycles: 123 and cycles of length 4.
- (d) The cycle 123 is not a sum of some cycles of length 4 in K_n .

- 1.3.** (a) Any 1-cycle in $K_{n,n}$ is a sum of some cycles of length 4.
- (b) If every edge of a 1-cycle in $K_{n,n}$ contains at least one of the vertices n or n' , then the 1-cycle is empty.
- (c) How many 1-cycles in $K_{n,n}$ are there?
- (d) For any pairwise distinct $a, b, c \in [n]$ and distinct $u', v' \in [n]'$ we have $au'bv' + bu'cv' + cu'av' = 0$.
- (e) Any linear relation on cycles of length 4 in $K_{n,n}$ is a sum of some relations from (d), and analogous relations $a'ub'v + b'uc'v + c'ua'v = 0$. (Rigorously, a *linear relation* is a set of cycles of length 4 in $K_{n,n}$ such that every edge of $K_{n,n}$ is contained in an even number of cycles from this set.)

1.4. How many 1-cycles are there

- (a) in a tree?
- (b) in a connected graph with V vertices and E edges?

Denote by \widetilde{K}_n the graph obtained from $K_{n,n}$ by deleting all ‘diagonal’ edges jj' , $j \in [n]$. E.g., \widetilde{K}_3 is the cycle 12'31'23' of length 6. This graph naturally appears in Problem 5.8.

- 1.5.** (a) If $n \geq 4$, then any 1-cycle in \widetilde{K}_n is a sum of some cycles of length 4.
- (b) Any 1-cycle in \widetilde{K}_n is a sum of some of the cycles $C_{ij'} := 12'31'ij'$ for edges ij' of \widetilde{K}_n such that $i, j > 1$ and $(i, j) \neq (3, 2)$.
- (c) The representation of (b) is unique.

A set B of 1-cycles is called a *base* of a set A of 1-cycles, if every 1-cycle from A has a unique representation as a sum of some 1-cycles from B .

1.6 (symmetric 1-cycles). * Denote by $t: \widetilde{K}_n \rightarrow \widetilde{K}_n$ the symmetry (involution) switching the parts, i.e., switching j and j' for every $j \in [n]$. Denote by tQ the 1-cycle symmetric to a 1-cycle Q . E.g. $t\widetilde{K}_3 = \widetilde{K}_3$ and $t(12'31'ij') = 1'23'1i'j$. A 1-cycle Q is called *t-symmetric* if $tQ = Q$. E.g. \widetilde{K}_3 is *t-symmetric*.

- (a) How many *t-symmetric* 1-cycles are there in \widetilde{K}_n ?
- (b) Any *t-symmetric* 1-cycle in \widetilde{K}_n is the sum of some of the following 1-cycles: \widetilde{K}_3 and $Q + tQ$ for cycles Q of length 4.
- (c) The set of all 1-cycles in \widetilde{K}_n has a base consisting of (the *t-symmetric* 1-cycle) \widetilde{K}_3 and pairs of 1-cycles going to each other under t .

1.7 (integer 1-cycles). * Let K be a graph with oriented edges. An assignment of integers to oriented edges of K is a (simplicial) *integer 1-cycle* if for every vertex the sum of integers assigned to incoming edges equals the sum of integers assigned to outgoing edges (Kirchhoff rule).

(a,b) State and prove the analogues of Problems 1.4.a,b for integer 1-cycles.

(c) For given set ω of orientations on edges of K denote by $H_1^\omega(K; \mathbb{Z})$ the set of the integer 1-cycles, with the componentwise sum operation. For different sets ω, ω' of orientations there is a 1–1 correspondence $\varphi: H_1^\omega(K; \mathbb{Z}) \rightarrow H_1^{\omega'}(K; \mathbb{Z})$ such that $\varphi(x + y) = \varphi(x) + \varphi(y)$ for every $x, y \in H_1^\omega(K; \mathbb{Z})$ (i.e., the groups $H_1(K; \mathbb{Z})$ for different sets of orientations are *isomorphic*).

1.8 (cohomology). * Fix an assignment of $+$ or $-$ to edges of a graph. For any vertex one can inverse the sign of every edge containing this vertex.

(a) For a tree, by the above operations for different vertices one can obtain any assignment from any other assignment.

(b) Consider a connected graph with V vertices and E edges. Find a maximal number of assignments of $+$ or $-$ to edges such that none of these assignments can be obtained from any other by the above operations.

2. ONE-DIMENSIONAL CYCLES IN THE SQUARE OF A GRAPH

A **graph** is a pair (V, E) of a finite set V and a set E of 2-element subsets (called ‘edges’) of V .

In this text K is a graph. We sometimes denote edge $\{a, b\}$ shortly by ab .

Intuitively, a *configuration* is a pair of ants (one red and one blue) sitting at two vertices of a graph (ants may occupy the same vertex). Configurations are called *adjacent* if one can be obtained from the other by one of the ants moving along one edge. This is rigorized as follows. Vertices of the graph $K^{\square 2}$ are ordered pairs (a, b) of vertices of the graph K . If vertices b and c of K are joined by an edge, then the vertices (a, b) and (a, c) of $K^{\square 2}$ are joined by an edge denoted by (a, bc) , and vertices (b, a) and (c, a) are joined by an edge denoted by (bc, a) . There are no other edges in $K^{\square 2}$.

E.g., the following are simple cycles in $K^{\square 2}$ (recall from §1 that a simple cycle is denoted by vertices it passes through):

- the *boundary*

$$ab \square uv := (a, u)(b, u)(b, v)(a, v),$$

for edges ab and uv in K (the product $ab \times uv$ is a rectangle, see §4, and $ab \square uv = \partial(ab \times uv)$ is its boundary);

- the *diagonal, off-diagonal and antidiagonal cycles*

$$\text{diag}(123) := (1, 1)(1, 2)(2, 2)(2, 3)(3, 3)(3, 1), \quad (1, 2)(1, 3)(2, 3)(2, 1)(3, 1)(3, 2),$$

$$\text{and } (1, 1)(2, 1)(2, 3)(3, 3)(3, 2)(1, 2) \text{ in } K_3^{\square 2};$$

- the *left, right, diagonal, off-diagonal, and antidiagonal cycles*

$$a \times C := (a, v_1) \dots (a, v_k), \quad C \times a := (v_1, a) \dots (v_k, a),$$

$$\text{diag } C := (v_1, v_1)(v_1, v_2)(v_2, v_2) \dots (v_k, v_k)(v_k, v_1), \quad (v_1, v_2)(v_1, v_3)(v_2, v_3) \dots (v_k, v_1)(v_k, v_2),$$

$$\text{and } (v_1, v_1)(v_2, v_1)(v_2, v_k) \dots (v_k, v_2)(v_1, v_2)$$

for a vertex a and a simple cycle $C = v_1 \dots v_k$ in K ;

• the *triodic cycle* $(1, 3)(1, 1')(1, 2)(1', 2)(3, 2)(3, 1') \dots$ in $K_{3,1}^{\square 2}$, where dots denote the part symmetric to the written part; a *triodic cycle* is also the analogous cycle corresponding to a $K_{3,1}$ -subgraph of K .

We consider the symmetry (involution) of $K^{\square 2}$ switching the factors (i.e., switching points (x, y) and (y, x)), and the corresponding map on 1-cycles. (For an application of symmetric 1-cycles see e.g. [Sk, §1.6].)

2.1 (riddle). Find graphs $K^{\square 2}$ for

- (a) $K = K_{2,1}$ a path on three vertices; (b) $K = K_3$ a cycle on three vertices;
- (c) $K = K_{3,1}$ a triod; (d) $K = K_4$.

- 2.2.** (a) Any 1-cycle in $K_{2,1}^{\square 2}$ is a sum of some boundaries.
 (b) The triodic cycle in $K_{3,1}^{\square 2}$ is a sum of some boundaries.
 (c) Any 1-cycle in $K_{3,1}^{\square 2}$ is a sum of some boundaries.
 (d) If K is a tree, then any 1-cycle in $K^{\square 2}$ is a sum of some boundaries.

Hint. Prove that if K and L are trees, then any 1-cycle in the graph $K \square L$ (defined below) is a sum of some boundaries. Prove this by induction, using deletion of a leaf vertex.

- (e) Is some left cycle a sum of some boundaries?
 (f) Is some diagonal cycle a sum of some boundaries?

The following notions are useful, in particular, for Problems 2.2.def.

Let K and L be graphs. Vertices of the *product graph* $K \square L$ are ordered pairs (a, b) of vertices a of K and b of L . If vertices b and c of L are joined by an edge, then the vertices (a, b) and (a, c) of $K \square L$ are joined by an edge denoted by (a, bc) . If vertices b and c of K are joined by an edge, then the vertices (b, a) and (c, a) of $K \square L$ are joined by an edge denoted by (bc, a) . There are no other edges in $K \square L$.

The *left projection* C_y of a 1-cycle C in $K \square L$ is the set of all edges σ in L such that there is an odd number of vertices a in K such that $a \times \sigma \in C$. (If C is a cycle, then C_y is the set of all edges passed by the blue ant an odd number of times.) The *right projection* C_x is defined analogously.

- 2.3.** (a) The diagonal cycle in $K_3^{\square 2}$ is a sum of a left cycle, a right cycle, and some boundaries.
 (b) The off-diagonal cycle in $K_3^{\square 2}$ is a sum of the diagonal cycle and some boundaries.
 (c) The antidiagonal cycle in $K_3^{\square 2}$ is a sum of the diagonal cycle and some boundaries.
 (d) The *symmetrized cycle* $a \times C + C \times a$ is the sum of $\text{diag } C$ and some boundaries.
 (e) For $K = K_4$ is the left cycle 1×234 a sum of some diagonal cycles and some boundaries?

2.4. Find the number of 1-cycles up to boundaries in

- (a) $K_3^{\square 2}$; (b) $K_{2,2}^{\square 2}$; (c) $K_{2,3}^{\square 2}$; (d) $K_4^{\square 2}$.

Hint. The number of 1-cycles up to boundaries is the maximal number of 1-cycles such that none of them is the sum of any other and some boundaries. For some parts of this problem the Künneth theorem 2.5.c will be useful.

Two 1-cycles C, C' in $K^{\square 2}$ are said to be *homologous* (or congruent modulo boundaries) if $C + C'$ is a sum of some boundaries. Notation: $C \sim C'$.

- 2.5.** (a) If K is connected and T is a tree, then for any 1-cycle C in $K \square T$ and vertex a in T there is a unique 1-cycle C_1 in K such that $C \sim C_1 \times a$. Moreover, $C \sim C_x \times a$.
 (b) If Z is a 1-cycle in $K^{\square 2}$ such that $Z_x = 0$, then $Z \sim a \times Z_y$ for any vertex a .
 (c) (Künneth theorem) If K is connected, then for any 1-cycle C in $K^{\square 2}$ and vertex a in K there are unique 1-cycles C_1, C_2 in K such that $C \sim C_1 \times a + a \times C_2$. Moreover, $C \sim C_x \times a + a \times C_y$.

The following (and $K^{\square 2}$ defined in §5) is a graph-theoretical analogue of the set of arrangements. Consider two ants on K as above, but now forbidden to sit in the same vertex. Rigorously speaking, vertices of the graph $K^{\square 2}$ are ordered pairs of distinct vertices of K . Vertices of $K^{\square 2}$ are joined by an edge in $K^{\square 2}$ if they are joined by an edge in $K^{\square 2}$.

2.6. (riddle) Find graphs $K^{\square 2}$ in cases (a,b,c,d) of Problem 2.1.

2.7. Find the number of 1-cycles in

- (a) $K_3^{\square 2}$; (b) $K_{2,2}^{\square 2}$; (c) $K_{2,3}^{\square 2}$; (d) $K_4^{\square 2}$; (e) $K_{3,3}^{\square 2}$; (f) $K_5^{\square 2}$.

(a'-f') Find the number of 1-cycles in $K^{\square 2}$ up to boundaries (contained) in $K^{\square 2}$ (i.e., up to boundaries corresponding to pairs of non-adjacent edges).

Hints. (a'-d') If a set of boundaries in $K^{\square 2}$ has zero sum, then the set is empty.

(e'), (f') The sum of all boundaries in $K^{\square 2}$ is zero. This is the only non-empty set of boundaries in $K^{\square 2}$ that has zero sum.

2.8. * (a) If a sum of boundaries is symmetric, it is a *symmetrized boundary*, i.e., a sum of $B + B'$ for some pairs of boundaries B, B' symmetric to each other.

(b) Any symmetric 1-cycle in $K^{\square 2}$ is a sum of some symmetrized cycles and some symmetrized boundaries. (Hence it is a sum of some diagonal cycles and some boundaries; note that a diagonal cycle is symmetric up to adding boundaries.)

Hint. Follows by (a) and the Künneth Theorem 2.5.c.

(c) For K connected, the map $C \mapsto C \times a + a \times C$ defines a 1–1 correspondence between 1-cycles in K and symmetric 1-cycles in $K^{\square 2}$ up to symmetrized boundaries. The inverse correspondence is given by $C \mapsto C_x = C_y$.

(d) For K connected, is the number of symmetric 1-cycles in $K^{\square 2}$ uniquely defined by the number V of vertices and the number E of edges? (We do not know solution of this problem.)

2.9. * Let K be a connected graph.

(a) A triodic cycle is not a sum of some boundaries in $K_{3,1}^{\square 2}$.

(b) Is an antidiagonal cycle a sum of a 1-cycle in $K^{\square 2}$ and some boundaries (in $K^{\square 2}$)?

(c) (conjecture) In $K^{\square 2}$ any off-diagonal cycle is not a sum of symmetrized cycles, and boundaries.

(d) In $K_4^{\square 2}$ is any off-diagonal cycle a sum of symmetrized cycles, triodic cycles, and boundaries?

(e) In $K^{\square 2}$ any symmetrized cycle is a sum of off-diagonal cycles, triodic cycles, and boundaries.

(f) (conjecture) In $K^{\square 2}$ any 1-cycle is a sum of left cycles, right cycles, off-diagonal cycles, triodic cycles, and boundaries.

(g) (conjecture) Any symmetric 1-cycle in $K^{\square 2}$ is a sum of off-diagonal cycles, triodic cycles, and boundaries (in $K^{\square 2}$)? The same with ‘... boundaries in $K^{\square 2}$ ’. The same with ‘boundaries’ replaced by ‘symmetrized boundaries’.

See more information in [CPG].

3. CYCLES IN HYPERGRAPHS

Higher-dimensional cycles in hypergraphs appear, in particular, as relations on 1-cycles in a graph, see Assertions 1.1.ef.

3.1 (cf. Problem 1.1). A *2-cycle* is a set C of 3-element subsets (called ‘faces’) of $[n]$ such that every 2-element subsets of $[n]$ is contained in an even number of subsets from C . E.g. the empty set is a 2-cycle. For a 4-element subset $A \subset [n]$ let the *tetrahedron* T_A be the set of all 3-element subsets of A . In other words, for pairwise distinct $a, b, c, d \in [n]$ define the *tetrahedron*

$$T_{\{a,b,c,d\}} := \{\{a, b, c\}, \{a, b, d\}, \{a, c, d\}, \{b, c, d\}\}.$$

Clearly, any tetrahedron is a 2-cycle.

(a) The sum of 2-cycles is a 2-cycle.

(b) If every face of a 2-cycle contains the number n , then the 2-cycle is empty.

(c) Any 2-cycle is a sum of some tetrahedra.

(d) How many 2-cycles for $[n]$ are there?

(e) For any 5-element subset $A \subset [n]$ we have $\sum_{j \in A} T_{A-\{j\}} = 0$.

(f) Any linear relation on tetrahedra is a sum of some relations from (e). (Give a rigorous formulation analogous to Assertion 1.1.f.)

3.2. Recall that $[n]^\ell$ is the set of all vectors of length ℓ whose entries are numbers from $[n]$. A *line* is a subset of $[n]^\ell$ given by fixing all coordinates except one. A *rook cycle* is a subset of $[n]^\ell$ containing an even number of vertices of every line. E.g. the empty set is a rook cycle. A *parallelepiped* is a subset $P_1 \times \dots \times P_\ell \subset [n]^\ell$, where P_i are 2-element subsets of $[n]$. Clearly, any parallelepiped is a rook cycle.

(a) The sum of rook cycles is a rook cycle.

(b) If a rook cycle contains no elements from $[n-1]^\ell$, then the rook cycle is empty.

- (c) Any rook cycle is a sum of some parallelepipeds.
 (d) How many rook cycles for $[n]^\ell$ are there?
 (e) For pairwise distinct $a, b, c \in [n]$ and parallelepiped $P \subset [n]^{\ell-1}$ we have $P \times \{a, b\} + P \times \{b, c\} + P \times \{c, a\} = 0$.
 (f) Any linear relation on parallelepipeds in $[n]^\ell$ is a sum of some relations from (e), and relations obtained from that by permutations of coordinates. (A rigorous formulation is analogous to Assertions 1.3.e and 3.1.f.)

Hint. First try the case $\ell = 2$, cf. Problem 1.3.

3.3 (riddle). Invent and prove a higher-dimensional analogue of Problem 3.1.

3.4. * A *2-hypergraph* (a two-dimensional hypergraph, or a 3-uniform hypergraph) is a pair (V, F) of a finite set V and a set F of 3-element subsets ('faces') of V . A (simplicial) *2-cycle* (modulo 2) in a 2-hypergraph (V, F) is a set C of faces such that every 2-element subset of V is contained in an even number of subsets from C .

(a) There are two *connected* hypergraphs having the same numbers of vertices, edges (i.e., 2-element subsets of faces), and faces, but different numbers of 2-cycles. (Cf. Problem 1.4.b.)

(b) Let $L = (V, F)$ be a hypergraph. Denote by E the set of edges, by b_0 the number of connected components, by 2^{b_1} the number of 1-cycles up to sums of some boundaries of faces, by 2^{b_2} the number of 2-cycles. Then $b_0 - b_1 + b_2 = |V| - |E| + |F|$.

4. GEOMETRIC DIGRESSION: CARTESIAN PRODUCTS OF GRAPHS

First we give an intuitive definition of a cylinder, and then a rigorous one.

Take rectangle bands $aa'bb'$ corresponding to edges ab of a graph K . Glue the end segments of the bands corresponding to the same vertex, so that the hatched letters would stick together with the hatched ones. The obtained two-dimensional figure is called a *cylinder* over graph K .

Recall that \mathbb{R}^d is the d -dimensional Euclidean space (for $d = 2$ and $d = 3$ this is the usual plane and space which one studies in elementary geometry class). The *cylinder* over a subset $U \subset \mathbb{R}^d$ is

$$U \times K_2 := \{(x, t) \in \mathbb{R}^{d+1} : x \in U, t \in [-1, 1]\}.$$

For instance, the cylinders over K_2 , K_3 , $K_{3,1}$, and K_5 are shown in Figure 2; the cylinder over $K_{k,1}$ 'looks like' the book with k sheets.




FIGURE 2. The cylinders over K_2 , K_3 , $K_{3,1}$, K_5

4.1. Any graph can be drawn without edges crossing on a book with a certain number of sheets (Figure 2, $K_{3,1}$) depending on the graph. More precisely, for any n there exists an integer k , as well as n points and $n(n-1)/2$ non-self-intersecting polygonal lines on a book with k sheets such that

- every pair of points is joined by a polygonal line, and
- no polygonal line intersects the interior of another polygonal line.

4.2 (riddle). The cylinder over any graph is realizable without self-intersections in \mathbb{R}^3 .

The (*geometric*) *product* of subsets $U, V \subset \mathbb{R}^d$ is

$$U \times V := \{(x, y) \in \mathbb{R}^{2d} : x \in U, y \in V\}.$$

In particular, the *square* of a subset $U \subset \mathbb{R}^d$ is $U^2 = U \times U$. For instance, the square K_2^2 is the ordinary square in the plane, and the square K_3^2 (called the *torus*) is shown in Figure 4, middle.

4.3 (riddle). Draw in \mathbb{R}^3

- (a) the product $K_3 \times K_{3,1}$ without self-intersections.
- (b) the square $K_{3,1}^2$; self-intersections are allowed.

Remark. Sometimes the word ‘graph’ is used for ‘the body of a graph’ defined as follows. Suppose that a subset of \mathbb{R}^3 is in 1–1 correspondence with the set of vertices of K , and no segment corresponding to an edge of K intersects the interior of any other such segment. The *body* of K is the union of the subset, and of all such segments.

In this language $K^{\square 2}$ is the set (the configuration space) of ordered pairs (x, y) of points of a graph K (or rather of its body) such that either x or y is a vertex of K . The set $K^{\square 2}$ is a union of a finite number of segments, i.e., is a graph.

Analogously, $K^{\square 2}$ is the set (the configuration space) of ordered pairs (x, y) of points of a graph K (or rather of its body) such that either x or y is a vertex of K , and x, y do not belong to the same edge. The set $K^{\square 2}$ is a union of a finite number of segments, i.e., is a graph.

5. TWO-DIMENSIONAL CYCLES IN THE SQUARE OF A GRAPH

Let the (*combinatorial*) *product* of graphs K and L be

$$K \times L := \{(\sigma, \tau) : \sigma \text{ is an edge of } K, \tau \text{ is an edge of } L\}.$$

Cells (σ, τ) and (γ, δ) in $K \times L$ are said to be *adjacent* if either

- $\sigma = \gamma$, and τ, δ have a common vertex, or
- $\tau = \delta$, and σ, γ have a common vertex.

(i.e., if the cells ‘share a common edge’ in $K \square L$). If we glue adjacent cells along their ‘common edges’ in $K \square L$, we obtain the geometric product discussed in §4.

Let the *square* of K be $K^2 := K \times K$.

A **cellular 2-cycle** (in K^2) is a subset $C \subset K^2$ such that for every vertex a and edge β of K

- there is an even number of edges α of K such that $\alpha \ni a$ and $(\alpha, \beta) \in C$, and
- there is an even number of edges α of K such that $\alpha \ni a$ and $(\beta, \alpha) \in C$.

In other words, $C \subset K^2$ is a cellular 2-cycle if every edge in $K^{\square 2}$ is ‘contained’ in an even number of cells $(\alpha, \beta) \in C$.

5.1. (a) For subset C of K^2 , the sum of boundaries $\sigma \square \tau$ over $(\sigma, \tau) \in C$ is zero if and only if C is a cellular 2-cycle.

(b) A subset $C \subset K^2$ is a cellular 2-cycle if and only if for every edge σ of K both sets (‘column section’ and ‘row section’)

$$C_{\sigma, \cdot} := \{\tau : (\sigma, \tau) \in C\} \quad \text{and} \quad C_{\cdot, \sigma} := \{\tau : (\tau, \sigma) \in C\}$$

are 1-cycles in K .

5.2. (a) The product of any two simple cycles in K (the *torus*) is a cellular 2-cycle.

(b) The sum modulo 2 of cellular 2-cycles is a cellular 2-cycle.

(c) Does there exist a tree T , and a non-empty cellular 2-cycle in T^2 ?

(d) Does there exist a graph K , a tree $T \subset K$, and a non-empty cellular 2-cycle contained in $\overline{T} := T \times K \cup K \times T$?

5.3. Find the number of cellular 2-cycles in

- (a) K_3^2 ; (b) $K_{2,2}^2$; (c) $K_{2,3}^2$; (d) K_4^2 .

Hint. For some parts of this problem the following problems will be useful.

Intuitively (and not precisely) speaking, the *combinatorial deleted product* $K^{\square 2}$ of K is a set of ordered pairs (x, y) of points in the graph K such that x, y do not belong to adjacent edges. The set $K^{\square 2}$ could be represented as a union of rectangles. Rigorously, let the *combinatorial deleted product* of K be

$$K^{\square 2} := \{(\sigma, \tau) : \sigma, \tau \text{ are non-adjacent edges of } K\}.$$

Remark. This set and its generalizations have many applications, notably to algorithm recognizing realizability of hypergraphs in higher-dimensional Euclidean spaces, see surveys [Sk06, Sk18]. The subset $K^{\square 2} \subset K^2$ is smaller than K^2 , and so is sometimes easier-to-draw. E.g. $K_2^{\square 2} = K_3^{\square 2} = K_{n,1}^{\square 2} = \emptyset$.

Glue adjacent cells along their ‘common edges’ in $K^{\square 2}$. Then the combinatorial deleted product of the path on 5 vertices, of the cycle on 5 vertices, of K_4 , of $K_{3,3}$, and of K_5 look like the disjoint union of two disks, the annulus, the cuboctahedron without triangular faces (Figure 5), the sphere with 4 handles, and the sphere with 6 handles, respectively.

(For $K = K_{3,3}$ and $K = K_5$ one proves that $K^{\square 2}$ looks like a connected orientable two-dimensional surface, and uses the classification theorem for two-dimensional surfaces; see details in [Sa91, the text after 3.4.1].)

5.4. (a) Are there any non-empty cellular 2-cycles in $K^{\square 2}$ for a *cycle* K ?

(b) Analogous question for a *wheel* K , i.e., for the graph with the set $\{0\} \cup [n]$ of vertices, and edges $\{n, 1\}$, $\{0, j\}$ and $\{j, j+1\}$ for $j \in [n-1]$.

(c) The subset $K_{3,3}^{\square 2}$ of $K_{3,3}^2$ is a cellular 2-cycle.

(d) Analogous statement for K_5 .

(e) There is a graph K such that $K^{\square 2}$ has a non-empty proper subset that is a cellular 2-cycle.

5.5. (a) Express $K_{3,3}^{\square 2}$ as a sum of some tori.

(b) Express $K_5^{\square 2}$ as a sum of some tori.

(c) Any cellular 2-cycle (in K^2) is a sum of some tori.

(d) How many cellular 2-cycles in K^2 are there, for a connected graph K with V vertices and E edges?

Hint to (c, d). The 1-cycle $\widehat{\varphi}\sigma$ is defined in the [solution](#) of Problem 1.4.b. By answer to Problem 5.2.d any cellular 2-cycle C in K^2 equals $\sum_{(\sigma, \tau) \in C - \overline{T}} \widehat{\varphi}\sigma \times \widehat{\varphi}\tau$, and is uniquely defined by its cells

outside \overline{T} .

Denote by $H_1(K)$ the set of all 1-cycles in K , with the sum operation.

Denote by $H_2(K^2)$ the set of all cellular 2-cycles in K^2 , with the sum operation.

5.6 (Künneth theorem). (a) There is a base C_1, \dots, C_q in $H_1(K)$ such that $C_i \times C_j$, $i, j \in [q]$, is a base in $H_2(K^2)$.

(b) If C_1, \dots, C_q is a base in $H_1(K)$, then $C_i \times C_j$, $i, j \in [q]$, is a base in $H_2(K^2)$.

(I.e., $H_2(K^2) \cong H_1(K) \otimes H_1(K)$. Base is defined analogously to the definition before Problem 1.6.)

5.7. (a) The product of any two vertex-disjoint cycles in K (i.e., two cycles which do not have common vertices) is a cellular 2-cycle in K^2 .

(b) Is any cellular 2-cycle in K^2 a sum of some products of vertex-disjoint cycles?

5.8. (a) How many cellular 2-cycles in $K_{n,n}^{\square 2}$ are there?

(b) There is a 1–1 correspondence $H_2(K_{n,n}^{\square 2}) \rightarrow H_2(\widetilde{K}_n^2)$.

(c) There is a 1–1 correspondence between $K_{n,n}^{\square 2}$ and \widetilde{K}_n^2 that respects the adjacency.

Hint: solve parts (a-c) in the reverse order.

Consider the symmetry (involution) of K^2 switching the factors (i.e., switching points (x, y) and (y, x)), and the map induced by this symmetry on cellular 2-cycles.

5.9. * (a) For any simple cycles Q, R in K the *symmetrized torus* $Q \times R + R \times Q$ is a symmetric cellular 2-cycle.

(b) The sum modulo 2 of symmetric cellular 2-cycles is a symmetric cellular 2-cycle.

(c) Is any symmetric cellular 2-cycle a sum of some symmetrized tori?

(e) For K connected, is the number of symmetric cellular 2-cycles in K^2 uniquely defined by the number V of vertices and the number E of edges?

5.10. * (a) Any symmetric cellular 2-cycle contained in K^2 is a sum of some symmetrized tori.

(b) How many symmetric cellular 2-cycles in $K_{n,n}^2$ are there?

(c) The 1–1 correspondence of Assertion 5.8.b moves cellular 2-cycles in $K_{n,n}^2$ transposed by the symmetry to cellular 2-cycles in \widetilde{K}_n^2 transposed by the symmetry t^2 . Here the symmetry $t : \widetilde{K}_n \rightarrow \widetilde{K}_n$ is defined in Problem 1.6, and $t^2(x, y) := (tx, ty)$.

(d) Any symmetric cellular 2-cycle in $K_{n,n}^2$ is a sum of some of the following ones:

- symmetrized tori $Q \times R + R \times Q$ for vertex-disjoint cycles Q, R of length 4 in $K_{n,n}$;
- the combinatorial deleted product $K_{3,3}^2$ of the subgraph $K_{3,3}$ of $K_{n,n}$.

(e) Any t^2 -symmetric cellular 2-cycle in \widetilde{K}_n^2 is a sum of some of the following ones: t^2 -symmetrized tori $Q \times R + tQ \times tR$, and \widetilde{K}_3^2 .

(f) Is the analogue of (d) for K_n and the subgraph (homeomorphic to) K_5 of K_n correct?

Hints. (d) One needs Künneth Theorem 5.6, Assertion 5.8.b and (e).

(e) One needs Assertion 1.6.c.

5.11. * (a) The following assertion [Sa91, 3.4.2] is wrong, even for connected graphs: there is a cellular 2-cycle C in K^2 such that any cellular 2-cycle in K^2 is a sum of some of the following cellular 2-cycles: C and products of vertex-disjoint cycles.

(b) (open problem) Is the assertion of (a) correct for 3-connected graphs?

(c) Any cellular 2-cycle in K^2 is a sum of some products of vertex-disjoint cycles, and some combinatorial deleted products of subgraphs homeomorphic to K_5 or to $K_{3,3}$.

(d) (riddle) How many cellular 2-cycles in K^2 are there, for a connected graph K with V vertices and E edges (use additional data of K , if required)?

Hints. (a) Take disjoint union of two copies of K_5 . Make the disjoint union connected by joining the copies by an edge.

6. ANSWERS, HINTS AND SOLUTIONS

1.1. (b) The assertion follows from Assertion 1.2.b.

Remark. A specific representation of a 1-cycle as a sum of some cycles of length 3 is given in (d).

(d) *Answer:* $2^{\frac{(n-1)(n-2)}{2}}$.

Hint. An arbitrary 1-cycle C in K_n equals the sum $\widehat{C} := \sum_{ij \in C, i, j < n} ijn$ of cycles ijn over all edges $ij \in C$ not containing the vertex n . This follows because every edge in $C + \widehat{C}$ contains the vertex n , so by (c) we have $C + \widehat{C} = 0$, i.e., $C = \widehat{C}$.

Moreover, such a representation is unique, see details in the proof of Assertion 1.4.b.

(f) In the rigorous formulation this is equivalent to Assertion 3.1.c.

1.2. (b) Any 1-cycle is a sum of some simple cycles. Any chord for a simple cycle of length l gives a representation of the cycle as the sum of two simple cycles of lengths less than l .

(c) The sum of two cycles of length 3 having a common edge is a cycle of length 4. So if a cycle α of length 3 is a sum of 123 and cycles of length 4, then any cycle of length 3 having a common edge with α is such. Therefore, all cycles of length 3 are such. Now (c) follows by Assertion 1.1.b.

(d) For a finite set X denote by $|X|_2$ the size of X modulo 2. Then for any 1-cycles C, C' we have $|C + C'|_2 = |C|_2 + |C'|_2$.

We often shorten $\varphi(x)$ to φx for the image of x under φ , and $\{\sigma\}$ to σ for 1-element subset.

1.3. (a) The assertion follows from Assertion 1.2.b.

Remark. Another way of representing 1-cycle C as a sum of some cycles of length 4 is $\widehat{\varphi}\varphi C$ defined in our solution of (c).

(c) *Answer:* $2^{(n-1)^2}$.

Remark. Compare to Assertion 1.4.b.

Proof. It suffices to prove that there exists a 1–1 correspondence between 1-cycles in $K_{n,n}$ and subsets¹ of $K_{n-1,n-1}$. Define a map

$$\varphi: H_1(K_{n,n}) \rightarrow 2^{K_{n-1,n-1}} \quad \text{by} \quad \varphi C := C \cap K_{n-1,n-1}.$$

For any edge $\sigma = ab'$ in $K_{n-1,n-1}$ denote by $\widehat{\varphi}\sigma$ the cycle $ab'nn'$ of length 4. Define a map

$$\widehat{\varphi}: 2^{K_{n-1,n-1}} \rightarrow H_1(K_{n,n}) \quad \text{by} \quad \widehat{\varphi}D := \sum_{\sigma \in D} \widehat{\varphi}\sigma.$$

Since $\varphi\widehat{\varphi}\sigma = \sigma$, we have $\varphi\widehat{\varphi}D = D$ for any $D \subset K_{n-1,n-1}$.

Take any $C \in H_1(K_{n,n})$. Since both $\widehat{\varphi}\varphi C$ and C are 1-cycles, $C' := \widehat{\varphi}\varphi C + C$ is a 1-cycle. Applying $\varphi\widehat{\varphi}D = D$ for $D = \varphi C$, we obtain $\varphi C' = 0$. So $C' \subset K_{n,n} \setminus K_{n-1,n-1}$. Hence by (b) we have $C' = 0$, so $\widehat{\varphi}\varphi C = C$.

Thus, φ is a 1–1 correspondence.

(e) See Assertion 3.2.f.

Remark. Problem 1.3 is equivalent to a particular case of Problem 3.2 for $\ell = 2$. An edge $ab' \in K_{n,n}$ corresponds to $(a, b) \in [n]^2$. A vertex $a \in [n]$ corresponds to the line $x = a$, a vertex $b' \in [n]'$ corresponds to the line $y = b$. An edge containing a vertex corresponds to an element of $[n]^2$ belonging to a line. Then 1-cycles in $K_{n,n}$ correspond to rook cycles in $[n]^2$. In particular, cycles of length 4 correspond to parallelepipeds (which for $\ell = 2$ may be called parallelograms).

1.4. *Answers:* (a) 1; (b) 2^{E-V+1} .

(b) *Proof.* Let T be a maximal tree of a connected graph K . It suffices to prove that there exists a 1–1 correspondence between $H_1(K)$ and subsets of $K \setminus T$.

Define a map

$$\varphi: H_1(K) \rightarrow 2^{K \setminus T} \quad \text{by} \quad \varphi C := C \cap (K \setminus T).$$

For any edge $\sigma \in K \setminus T$ denote by $\widehat{\varphi}\sigma$ the simple cycle in K formed by σ and the simple path in T joining ends of σ .

Define a map

$$\widehat{\varphi}: 2^{K \setminus T} \rightarrow H_1(K) \quad \text{by} \quad \widehat{\varphi}D := \sum_{\sigma \in D} \widehat{\varphi}\sigma.$$

Since $\varphi\widehat{\varphi}\sigma = \sigma$, we have $\varphi\widehat{\varphi}D = D$ for any $D \in 2^{K \setminus T}$.

Take any $C \in H_1(K)$. Since both $\widehat{\varphi}\varphi C$ and C are 1-cycles, $C' := \widehat{\varphi}\varphi C + C$ is a 1-cycle. Applying $\varphi\widehat{\varphi}D = D$ for $D = \varphi C$, we obtain $\varphi C' = 0$. So $C' \subset T$. Hence by (a) we have $C' = 0$, so $\widehat{\varphi}\varphi C = C$.

Thus, φ is a 1–1 correspondence.

1.5. (a) A simple cycle in \widetilde{K}_n is formed by an l -element cyclic sequence of vertices such that

- the number l is even and greater than 2;
- the parts of vertices alternate in the sequence;
- within each separate part, the vertices are not repeated;
- there are no two consecutive vertices m and m' .

¹Sometimes the notation for a graph K is used for the set of edges of K (or, alternatively, the notation ‘ $\cap K$ ’ is used for the ‘restriction to K ’ operation on the sets of edges).

Assume that such a cycle has no chords. Then any two non-consecutive vertices are not adjacent, i.e., they either are in the same part or are m and m' . Any vertex has $\frac{l}{2} - 2$ non-consecutive vertices from the other part. Then $\frac{l}{2} - 2 \leq 1$, so $l \in \{4, 6\}$.




FIGURE 3. A cycle of length 6 is the sum of three cycle of length 4

In a simple cycle of length 6 having no chords, any two opposite vertices are m and m' . Hence the cycle is $(m_1, m'_2, m_3, m'_1, m_2, m'_3)$ for some pairwise distinct m_1, m_2, m_3 . Since $n \geq 4$, there is $a \in [n] - \{m_1, m_2, m_3\}$. Then the cycle equals

$$(m_1, m'_2, m_3, a') + (m_2, m'_3, m_1, a') + (m_3, m'_1, m_2, a'),$$

see Figure 3. Now the required result holds by Assertion 1.2.b.

(b,c) Let S be the set of all edges ij' of \widetilde{K}_n such that $i, j > 1$ and $(i, j) \neq (3, 2)$. The assertions follow (analogously to Problem 1.4.b) because

- an edge $e' \in S$ is contained in the cycle C_e if and only if $e = e'$;
- the graph $\widetilde{K}_n - S$ obtained from \widetilde{K}_n by deleting all edges of S is a tree.

The second of these assertions follows because the edges of $\widetilde{K}_n - S$ are exactly $32'$ and $1i', i1'$, where $i > 1$, so $\widetilde{K}_n - S$ is obtained by connecting with edge $32'$ the ‘star’ trees consisting of edges $1i'$ with $i > 1$, and of edges $1'i$ with $i > 1$.

1.6. (a,b) The natural projection $\widetilde{K}_n \rightarrow K_n$ gives a 1–1 correspondence between t -symmetric 1-cycles in \widetilde{K}_n and 1-cycles in K_n . Use Assertions 1.1.d and 1.2.c.

(c) Take the base given by Assertions 1.5.bc. Then the 1-cycles

$$\widetilde{K}_3 = C_{23'}, \quad C_{ij'}, \quad C_{23'} - C_{ji'}, \quad \text{where } i > j > 1 \quad \text{and} \quad (i, j) \neq (3, 2),$$

also form a base. This base is as required because $C_{23'} - C_{ji'} = 1i'j1'32'12'31'23' = 1i'j1'23' = tC_{ij'}$.

Alternatively, one can take the cycles $\widetilde{K}_3, 1j'i2', 1'j'i'2$, where $j > i > 2$, and $1i'23', 1'i2'3$, where $i > 3$.

1.7. (a) *Statement.* There is only one integer 1-cycle in a tree. It assigns 0 to all edges.

Hint. Start with proving that any integer 1-cycle on a tree assigns 0 to all terminal edges.

(b) *Statement.* Let T be a maximal tree of a connected graph K . Every assignment of integers to edges of $K \setminus T$ has a unique extension to integer 1-cycle in K .

Proof. Note that the sum of integer 1-cycles is an integer 1-cycle.

Denote by $\widehat{\varphi}_{\mathbb{Z}}(\sigma)$ the integer 1-cycle in K

- assigning 1 to all edges of $\widehat{\varphi}\sigma$ oriented in $\widehat{\varphi}\sigma$ in the same way as σ ,
- assigning -1 to all edges of $\widehat{\varphi}\sigma$ oriented in $\widehat{\varphi}\sigma$ the opposite way, and
- assigning 0 to all edges in $K \setminus \widehat{\varphi}\sigma$.

For every assignment D of integers to edges of $K \setminus T$ denote $\widehat{\varphi}D := \sum_{\sigma \in K \setminus T} D(\sigma) \widehat{\varphi}_{\mathbb{Z}}(\sigma)$. Then $\widehat{\varphi}D$

is an integer 1-cycle extending D .

Take any integer 1-cycle C in K extending D . Then $C - \widehat{\varphi}D$ is an integer 1-cycle that assigns 0's to all edges in $K \setminus T$. So by (a) integer 1-cycle $C - \widehat{\varphi}D$ also assigns 0's to all edges in T . So $C = \widehat{\varphi}D$.

1.8. (b) *Answer:* 2^{E-V+1} .

2.1. (a) The 3×3 grid graph, see Figure 4, left.

(b) See Figure 4, middle (where boundary cycles are drawn in grey). Alternatively, $K_3^{\square 2}$ is obtained from the graph in (a) by adding edges between corresponding vertices:




FIGURE 4. Left: $K_{2,1}^{\square 2}$ is the 3×3 grid graph. Middle: $K_3^{\square 2}$ on the torus K_3^2 . Right: $K_{3,1}^{\square 2}$.

- of the 1st and the 3rd rows;
- of the 1st and the 3rd column.

(Compare to obtaining K_3 from the path 123 by adding an edge between 1 and 3.)

(c) See Figure 4, right.

(d) Vertices are nodes of a 4×4 grid. Two vertices are joined by an edge if and only if they are in the same row or in the same column.

2.2. (b) The triodic cycle in $K_{3,1}^{\square 2}$ equals $\sum_{i,j \in [3], i \neq j} i'j1'$.

(e,f) *Answers:* no.

(e) *Hint.* The left projection of any boundary (and hence of any sum of boundaries) is empty.

2.3. (a) We have $\text{diag}(123) = 1 \times K_3 + K_3 \times 1 + 12 \square 23 + 12 \square 31 + 23 \square 31$.

(b) The off-diagonal cycle in $K_3^{\square 2}$ equals $1 \times K_3 + K_3 \times 1 + 12 \square 31 + 31 \square 12$.

(c) The antidiagonal cycle in $K_3^{\square 2}$ equals $1 \times K_3 + K_3 \times 1 + 23 \square 31 + 31 \square 23 + 31 \square 31$.

(d) We have $1 \times C + C \times 1 = \text{diag } C + \sum_{i,j \in [n], i < j} i(i+1) \square j(j+1)$, where $n+1 := 1$.

(e) *Answer:* no.

2.4. *Answers:* (a) 2^2 ; (b) 2^2 ; (c) 2^4 ; (d) 2^6 .

2.5. (a) Prove this by induction on the number of vertices of T , using deletion of a leaf vertex.

(b) This can be proved by induction on the number of pairs of edges of Z left-projected to the same edge of K . The inductive base when the number is zero can be proved by induction on the number of vertices b of K such that $Z \cap (b \times K) \neq \emptyset$. For the inductive step, take such a pair $\{(\sigma, v_1), (\sigma, v_k)\}$, take a path $v_1 \dots v_k$ in K , and replace Z by $Z + \sigma \square v_1 v_2 + \dots + \sigma \square v_{k-1} v_k$.

(c) By (b), if $Z_x = Z_y = 0$, then $Z \sim 0$. Applying this to $Z = C + C_x \times a + a \times C_y$, we obtain the representation from (c). The uniqueness follows because the left and the right projections of sum of boundaries are equal to 0.

2.6. *Answers.* (a) The disjoint union of two copies of $K_{2,1}$.

(b) The cycle on 6 vertices.

(c) The cycle on 12 vertices.

(d) The union of edges of a cuboctahedron, see Figure 5.

2.7. *Answers.*

(a) 2^1 ; (b) 2^5 ; (c) 2^{17} ; (d) 2^{13} ; (e) 2^{43} ; (f) 2^{41} .

(a') 2^1 ; (b') 2^1 ; (c') 2^5 ; (d') 2^7 ; (e') 2^8 ; (f') 2^{12} .

Hint. Apply 1.4.b to obtain (a)-(f).

Remark. Cf. examples before Problem 5.4.

2.8. (a) It suffices to prove the result for K connected. It suffices to prove that if a sum of pairwise distinct boundaries is a symmetric 1-cycle, then either the sum is zero, or there are two summands symmetric to each other. The analogous result for $K^{\square 2}$ replaced by $K \square T \bigcup_{T \square T} T \square K$, where T is




FIGURE 5. Left: cuboctahedron; the union of edges is $K_4^{\square 2}$ (three of the six boundaries are highlighted); the complement to triangular faces is K_4^2 . Right: the same with some explanations; the figure does not show the invisible part whose projection is obtained from the pictured projection by rotation through $\pi/3$

a tree, is proved by induction on the number of vertices in T , using deletion of a leaf vertex. For reduction of (a) to this analogous result, use the equality $\sum_{i,j} \sigma_i \square \tau_j = 0$ for two simple cycles in K having consecutive edges $\sigma_1 \dots \sigma_k$ and $\tau_1 \dots \tau_\ell$. Using this equality we replace the given sum of boundaries by an equal sum not containing summands $\sigma \square \tau$ corresponding to edges σ, τ outside a maximal tree of K .

3.1. (c) It suffices to prove that an arbitrary 2-cycle C equals the sum $\widehat{C} := \sum_{\{i,j,k\} \in C, i,j,k < n} T_{i,j,k,n}$ of tetrahedra $T_{i,j,k,n}$ over 3-element subsets $\{i, j, k\} \in C$ not containing n . This follows because every face in $C + \widehat{C}$ contains n , so by (b) we have $C + \widehat{C} = 0$, i.e., $C = \widehat{C}$.

(f) A rigorous formulation is as follows (we identify tetrahedra in $[n]$ with 4-element subsets of $[n]$). Suppose that R is a set of 4-element subsets of $[n]$ such that any 3-element subset of $[n]$ is contained in an even number of those 4-element subsets that are elements of R . Then there is a family R' of 5-element subsets of $[n]$ such that a 4-element subset B of $[n]$ is contained in R if and only if B is contained in an odd number of those 5-element subsets that are elements of R' .

This statement is a higher dimensional analogue of Assertion 3.1.c; the proof is analogous.

3.2. (c) For $a \in [n-1]^\ell$ denote by $P(a) := \{n, a_1\} \times \dots \times \{n, a_\ell\}$ the parallelepiped with opposite vertices a and (n, \dots, n) . It suffices to prove that any rook cycle $C \subset [n]^\ell$ equals the sum \widehat{C} of parallelepipeds $P(a)$ over $a \in C \cap [n-1]^\ell$. The sum $C + \widehat{C}$ is a rook cycle. Since $P(a) \cap [n-1]^\ell = \{a\}$, we have $(C + \widehat{C}) \cap [n-1]^\ell = \emptyset$. By (b), we have $C + \widehat{C} = \emptyset$, i.e., $C = \widehat{C}$.


(f) *Hint.* Any parallelepiped P with some $P_i \subset [n-1]$ is a sum of some given relations and two parallelepipeds obtained from P by replacing with n one of the two elements in P_i . So in each of the two parallelepipeds the number of pairs P_j not containing n is smaller than in P . Hence any relation between parallelepipeds is a sum of some given relations and a relation $P(a_1) + \dots + P(a_s) = 0$ for some pairwise distinct $a_1, \dots, a_s \in [n-1]^\ell$. The latter relation is trivial because

$$\emptyset = (P(a_1) + \dots + P(a_s)) \cap [n-1]^\ell = \{a_1, \dots, a_s\}.$$

Here the second equality holds because $P(a) \cap [n-1]^\ell = \{a\}$.

4.2. Let $O, V, A_{11}, \dots, A_{1n}$ be points in \mathbb{R}^3 of which no 4 lie in the same plane. For every $p \in [n]$ take a point A_{2p} such that $\overrightarrow{OA_{2p}} = \overrightarrow{OV} + \overrightarrow{OA_{1p}}$. If V is close enough to O , then the points A_{jp} , $j \in \{1, 2\}$, $p \in [n]$, are as required, i.e., the union of parallelograms $A_{1p}A_{2p}A_{2q}A_{1q}$ forms the drawing of the cylinder over K_n in \mathbb{R}^3 .

4.3. (b) See Figure 6.

FIGURE 6. To drawing of $K_{3,1}^2$ in 3-space

5.1. (b) In this proof by σ, τ we denote edges of K , and by v we denote a vertex of K :

$$\begin{aligned}
 C \text{ is a 2-cycle} &\Leftrightarrow \begin{cases} |\{\sigma : \sigma \ni v, (\sigma, \tau) \in C\}| \text{ is even for any } v \text{ and } \tau \\ |\{\tau : \tau \ni v, (\sigma, \tau) \in C\}| \text{ is even for any } v \text{ and } \sigma \end{cases} \Leftrightarrow \\
 &\Leftrightarrow \begin{cases} |\{\sigma \in C_{\cdot, \tau} : \sigma \ni v\}| \text{ is even for any } v \text{ and } \tau \\ |\{\tau \in C_{\sigma, \cdot} : \sigma \ni v\}| \text{ is even for any } v \text{ and } \sigma \end{cases} \Leftrightarrow \begin{cases} C_{\cdot, \tau} \text{ is a 1-cycle for any } \tau \\ C_{\sigma, \cdot} \text{ is a 1-cycle for any } \sigma. \end{cases}
 \end{aligned}$$

5.2. (a) Let $C := \tilde{C}_1 \times \tilde{C}_2 \subset K^2$ be the product of simple cycles \tilde{C}_1 and \tilde{C}_2 in K . For every edge σ of K the set $C_{\sigma, \cdot}$ is either \tilde{C}_2 or the empty set. For every edge τ of K the set $C_{\cdot, \tau}$ is either \tilde{C}_1 or the empty set. Hence by Assertion 5.1.b, C is a cellular 2-cycle.

(c) *Answer:* no.

Solution. Let C be a cellular 2-cycle in T^2 . Since T is a tree, by Assertion 5.1.b $C_{\sigma, \cdot} = 0$ for any edge σ of T . Hence $C = 0$.

(d) *Answer:* no.

Solution. Let C be a cellular 2-cycle in \overline{T} . Since T is a tree, by Assertion 5.1.b $C_{\sigma, \cdot} = 0$ for any edge σ of $K \setminus T$. Hence $C \subset T \times K$. Since T is a tree, by Assertion 5.1.b $C_{\cdot, \sigma} = 0$ for any edge σ of K . Hence $C = 0$.

5.3. *Answers:* (a) 2; (b) 2; (c) 2^4 ; (d) 2^9 .

Solutions.

(a, b) Let C be a cellular 2-cycle in K^2 such that $C \neq K$, i.e., $(\sigma, \tau) \notin C$ for some edges σ, τ in K . By Assertion 5.1.b, $C_{\sigma, \cdot}$ is a 1-cycle. Since K is a cycle graph, $C_{\sigma, \cdot} = 0$. Then $(\sigma, \sigma) \notin C$. Hence $C \subset \overline{K - \sigma}$. By answer to Problem 5.2.d, $C = 0$.

(c) By answer to Problem 5.5.d, the number of 2-cycles in $K_{2,3}^2$ equals $2^{(E-V+1)^2} = 2^{2^2} = 16$.

(d) By answer to Problem 5.5.d, the number of 2-cycles in K_4^2 equals $2^{(E-V+1)^2} = 2^{3^2} = 2^9 = 512$.

5.4. (a) *Answer:* no.

Solution. Let C be a cellular 2-cycle in K^2 . Take an edge σ of K . By Assertion 5.1.b, $C_{\sigma, \cdot}$ is a 1-cycle. Since $C_{\sigma, \cdot}$ is a subset of the path graph $K - \sigma$, then $C_{\sigma, \cdot} = 0$. Hence $C = 0$.

(b) *Answer:* no.

Solution. Let C be a cellular 2-cycle in K^2 . By Assertion 5.1.b, $C_{\sigma, \cdot}$, $C_{\cdot, \sigma}$ are 1-cycles for any edge σ of K . Since both $C_{0j, \cdot}$ and $C_{\cdot, 0j}$ are subsets of the path graph $K - 0 - j$ for any $j \in [n]$, then $C_{0j, \cdot} = C_{\cdot, 0j} = 0$ for any $j \in [n]$. Then $C \subset (K - 0)^2$ for the cycle graph $K - 0$. Hence by (a), $C = 0$.

(c) Denote $C := K_{3,3}^2$. For any edge σ in $K_{3,3}$ both $C_{\sigma, \cdot}$ and $C_{\cdot, \sigma}$ are cycles of length 4. Then the required result follows by Assertion 5.1.b.

(d) Denote $C := K_5^2$. For any edge σ in K_5 both $C_{\sigma, \cdot}$ and $C_{\cdot, \sigma}$ are cycles of length 3. Then the required result follows by Assertion 5.1.b.

(e) Take $K := K_6$ and $C := 123 \times 456$. Since $(12, 34) \notin C$, the product C is a non-empty proper subset of K^2 . By Assertion 5.2.a, the product C of cycles is a cellular 2-cycle.

5.5. (a, b) *Hint:* the required representations can be obtained from the proof of (c).

(c) Part (c) easily follows from the particular case of connected K , which we further assume.

Let T be a maximal tree of K . Define a map

$$\varphi: H_2(K^2) \rightarrow 2^{K^2 - \bar{T}} \quad \text{by} \quad \varphi C := C \cap (K^2 - \bar{T}).$$

For any edge $\sigma \in K \setminus T$ the 1-cycle $\widehat{\varphi}\sigma$ is defined in the proof of Assertion 1.4.b. Define a map

$$\widehat{\varphi}: 2^{K^2 - \bar{T}} \rightarrow H_2(K^2) \quad \text{by} \quad \widehat{\varphi}D := \sum_{(\sigma, \tau) \in D} \widehat{\varphi}\sigma \times \widehat{\varphi}\tau.$$

By Assertion 5.2.d, $\widehat{\varphi}\varphi C = C$ for any $C \in H_2(K^2)$, i.e., an arbitrary cellular 2-cycle C in K^2 is the sum of tori $\widehat{\varphi}\sigma \times \widehat{\varphi}\tau$ over $(\sigma, \tau) \in \varphi C$.

(d) *Answer:* $2^{(E-V+1)^2}$.

Solution. In (c), we defined maps $\varphi: H_2(K^2) \rightarrow 2^{K^2 - \bar{T}}$ and $\widehat{\varphi}: 2^{K^2 - \bar{T}} \rightarrow H_2(K^2)$ for a maximal tree T of connected graph K and proved that $\widehat{\varphi}\varphi C = C$ for any $C \in H_2(K^2)$. In addition, $\varphi\widehat{\varphi}D = D$ for any $D \in 2^{K^2 - \bar{T}}$. Thus, φ is a 1–1 correspondence. Hence

$$|H_2(K^2)| = |2^{K^2 - \bar{T}}| = |2^{(K \setminus T)^2}| = 2^{(E-V+1)^2}.$$

5.6. (b) Denote $B := \{C_i \times C_j : i, j \in [q]\}$. In the following two paragraphs we prove that every 2-cycle in K^2 is a sum of some 2-cycles from B , and that such a representation is unique, respectively.

Apply Assertion 5.5.c. Every torus $\widetilde{C}_1 \times \widetilde{C}_2 \subset K^2$ is in turn a sum of some 2-cycles from B , because both \widetilde{C}_1 and \widetilde{C}_2 are sums of some 1-cycles from C_1, \dots, C_q .

Denote by N the number of connected components of K . Analogously to Assertion 1.4.b, $|H_1(K)| = 2^{E-V+N}$. Since C_1, \dots, C_q is a base in $H_1(K)$, it follows that $2^q = |H_1(K)|$, so $q = E - V + N$. Therefore $|B| = q^2 = (E - V + N)^2$. By generalization of the answer to Problem 5.5.d to an arbitrary graph K , we have $|H_2(K^2)| = 2^{(E-V+N)^2}$, which is the number of linear combinations of 2-cycles from B . This implies the uniqueness.

5.7. (a) *Hint:* this follows by Assertion 5.1.b.

(b) *Answer:* no.

Solution. Take $K := K_5$ (or $K := K_{3,3}$) and $C := K^2$. By Assertion 5.4.c (or 5.4.d), C is a cellular 2-cycle. Since there are no two non-empty vertex-disjoint cycles in K , the 2-cycle C is not a sum of some products of vertex-disjoint cycles.

5.8. (a) *Answer:* $2^{(n^2-3n+1)^2}$.

(c) *Hint.* The 1–1 correspondence is defined by the formula $f(\sigma_1\sigma'_2, \tau_1\tau'_2) := (\sigma_1\tau'_1, \sigma_2\tau'_2)$ for disjoint edges $\sigma, \tau \in [n]^2$. (Recall that for $a, b \in [n]$ we denote by ab' the edge joining a and b' .) Cells $\alpha, \beta \in [n] \times [n]' \times [n] \times [n]'$ in $K_{n,n}^2$ are adjacent if and only if they differ only at one of their four coordinates. Since $K_{n,n}^2, \widetilde{K}_n^2 \subset K_{n,n}^2$, it follows that α and β in $K_{n,n}^2$ are adjacent if and only if $f\alpha$ and $f\beta$ in \widetilde{K}_n^2 are adjacent, i.e., f respects the adjacency.

Remark. Define a 1–1 correspondence \widehat{f} between edges of $K_{n,n}^{\square 2}$ and edges of $\widetilde{K}_n^{\square 2}$ by skipping one of $\sigma_1, \sigma'_2, \tau_1, \tau'_2$ in the formula for f (e.g. $\widehat{f}(\sigma_1, \tau_1\tau'_2) = (\sigma_1\tau'_1, \tau'_2)$). One can check that the correspondences f, \widehat{f} respect incidence. (Recall that an edge η and a 2-cell $\alpha = (\xi, \zeta)$ are *incident* if $\eta = \xi$ or $\eta = \zeta$.) Note that there is no 1–1 correspondence between vertices that saves incidence. Indeed, there is no 1–1 correspondence between 1-cycles in $K_{n,n}^{\square 2}$ and in $\widetilde{K}_n^{\square 2}$. E.g. for $n = 3$ we have $|H_1(\widetilde{K}_3^{\square 2})| = 2^{37}$, as opposed to the answer to Problem 2.7.e; besides, in $\widetilde{K}_3^{\square 2}$ there are four 1-cycles up to boundaries, as opposed to the answer to Problem 2.7.e'.

5.9. (c) *Answer:* no.

Solution. Take $K = K_3$ and $C = K^2$. Then C is a non-empty symmetric cellular 2-cycle in K^2 . Since $K \times K + K \times K = 0$, there are no non-empty symmetrized tori in K^2 . Hence C is not a sum of some symmetrized tori.

5.10. (a) For any symmetric cellular 2-cycle C in K^2 we have

$$C \stackrel{(1)}{=} \sum_{(\sigma, \tau) \in C - \bar{T}} \widehat{\varphi}_\sigma \times \widehat{\varphi}_\tau \stackrel{(2)}{=} \sum_{\{\sigma, \tau\}: (\sigma, \tau) \in C - \bar{T}} (\widehat{\varphi}_\sigma \times \widehat{\varphi}_\tau + \widehat{\varphi}_\tau \times \widehat{\varphi}_\sigma).$$

Here

- (1) holds by the formula in the hint to Assertion 5.5.c, and
 - (2) holds because $C - \bar{T}$ is symmetric, and $C - \bar{T} \subset C \subset K^2$, so $\sigma \neq \tau$ for any pair $(\sigma, \tau) \in C - \bar{T}$.
- (c,d,e) See the details in [SS23].

REFERENCES

- [CPG] * https://en.wikipedia.org/wiki/Cartesian_product_of_graphs
- [FH10] *M. Farber, E. Hanbury.* Topology of Configuration Space of Two Particles on a Graph, II. *Algebr. Geom. Topol.* 10 (2010) 2203–2227. arXiv:1005.2300.
- [MS17] *T. Maciazek, A. Sawicki.* Homology groups for particles on one-connected graphs *J. Math. Phys.* 58, 062103 (2017). arXiv:1606.03414.
- [Sa91] *K. S. Sarkaria.* A one-dimensional Whitney trick and Kuratowski’s graph planarity criterion, *Israel J. Math.* 73 (1991), 79–89. <http://kssarkaria.org/docs/One-dimensional.pdf>.
- [Sk06] * *A. Skopenkov,* Embedding and knotting of manifolds in Euclidean spaces, *London Math. Soc. Lect. Notes*, 347 (2008) 248–342. arXiv:math/0604045.
- [Sk14] * *A. Skopenkov,* Realizability of hypergraphs and intrinsic linking theory, *Mat. Prosveschenie*, to appear, arXiv:1402.0658.
- [Sk18] * *A. Skopenkov.* Invariants of graph drawings in the plane. *Arnold Math. J.*, 6 (2020) 21–55; full version: arXiv:1805.10237.
- [Sk20] * *A. Skopenkov,* Algebraic Topology From Geometric Standpoint (in Russian), MCCME, Moscow, 2020 (2nd edition). Part of the book: <http://www.mccme.ru/circles/oim/obstruct.pdf>. Part of the English translation: <https://www.mccme.ru/circles/oim/obstructeng.pdf>.
- [Sk21m] * *A. Skopenkov.* Mathematics via Problems. Part 1: Algebra. Amer. Math. Soc., Providence, 2021. Preliminary version: https://www.mccme.ru/circles/oim/algebra_eng.pdf
- [Sk] * *A. Skopenkov.* Algebraic Topology From Algorithmic Standpoint, draft of a book, mostly in Russian, <http://www.mccme.ru/circles/oim/algor.pdf>.
- [SS23] *A. Skopenkov and O. Styrť,* Embeddability of join powers and minimal rank of partial matrices, arXiv:2305.06339.

Books, surveys and expository papers in this list are marked by the stars.

arXiv is an open-access archive of papers on mathematics, computer science, and other topics. To access a paper enter the identifier (e.g. arXiv:1005.2300) in the search bar of arxiv.org. Googling the identifier usually works as well.

