

FIXED POINT SETS OF INVOLUTIONS ON THE PRODUCT OF THREE SPHERES

DIMPI AND HEMANT KUMAR SINGH

ABSTRACT. Let $G = \mathbb{Z}_2$ act on a finitistic space X having mod 2 cohomology of the product of three spheres $\mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, $1 \leq n \leq m \leq l$. In this paper, we have determined the fixed point sets of involutions on X . This generalizes J. C. Su [12] results for involutions on the product of two sphere $\mathbb{S}^n \times \mathbb{S}^m$, $n \leq m$.

1. INTRODUCTION

Let (G, X) be a transformation group, where G is compact Lie group and X is compact Hausdroff space, with the fixed point set F . The study of the cohomological structure of the fixed point set has been an interesting problem in transformation groups. J. C. Su [12] gave a conjecture which states that if $G = \mathbb{Z}_p$, p a prime, act on a finitistic space X which satisfies poincaré duality with respect to Čech cohomology then each component of the fixed point set also satisfies poincaré duality. Bredon [4] proved Su's conjecture, in the case when X is totally nonhomologous to zero in X_G (Borel space). In a particular case, Puppe [11] has proved the Bredon's conjecture: if $H^*(X, \mathbb{Z}_p)$ generated by k elements as an algebra, then any component F_0 of F , $H^*(F_0, \mathbb{Z}_p)$ generated by at most k elements. Puppe proved that if X is totally nonhomologous to zero in X_G , then the number of generators of the cohomology ring of each component of the fixed point set with \mathbb{Z}_p -coefficient is at most the number of generators of $H^*(X, \mathbb{Z}_p)$.

For the fixed point sets of actions of a group G on the spheres \mathbb{S}^n and on the product of two spheres $\mathbb{S}^n \times \mathbb{S}^m$, a lot of work has been done in the literature. Here, we recall some of the results. The fixed point sets of $G = \mathbb{Z}_p$, p a prime, or $G = \mathbb{T}^k$, $k \geq 1$ (\mathbb{T}^k denotes k -dimensional torus) on a finitistic space X having mod p or integral cohomology n -sphere are mod p or integral cohomology r -sphere, where $-1 \leq r \leq n$ and $n - r$ is even for $G = \mathbb{Z}_p$, $p > 2$ and $G = \mathbb{T}^k$ (see Chapter III, [3]). According

2020 *Mathematics Subject Classification*. Primary 57S17; Secondary 55M35.

Key words and phrases. Finitistic space; Fixed point set; Totally nonhomologous to zero; Leray-Serre spectral sequence.

The first author of the paper is supported by SRF of UGC, New Delhi, with reference no.: 201610039267.

to this well known result, the fixed point sets of transformation group $(G, X_1 \times X_2)$ (with diagonal action), where X_1 and X_2 has mod p or integral cohomology of spheres, has the same mod p or integral cohomology algebra of the product of two spheres. Swan [8] gave sufficient conditions which insure that the fixed point set of $G = \mathbb{Z}_p$, p a prime, actions on mod p cohomology product of two even dimensional spheres have mod p cohomology of product of two even dimensional spheres of lower dimensions. Su [12] determined the fixed point sets of $G = \mathbb{Z}_p$ actions on a space X with mod p cohomology algebra product of spheres $\mathbb{S}^m \times \mathbb{S}^n$. Further, Chang et al. [6] and David [7] discussed the fixed point sets of circle actions and torus actions on the product of even dimensional spheres and the product of odd dimensional spheres, respectively. Allday [1] discussed the fixed point sets of torus actions on cohomology product of three odd dimensional spheres. In continuation, it seems to be an interesting problem to determine the fixed point sets of involutions on a finitistic space X having mod 2 cohomology of the arbitrary product of three spheres $\mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, $1 \leq n \leq m \leq l$.

In this paper, we have determined the possibilities of the fixed point sets of involutions on a finitistic space X having mod 2 cohomology of the arbitrary product of three spheres $\mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, $1 \leq n \leq m \leq l$.

2. PRELIMINARIES

We recall some known facts that will be used in this paper. A paracompact Hausdorff space is called finitistic if every open covering has a finite dimension refinement. For example: (1) Compact Hausdorff spaces are finitistic spaces. (2) $\prod_{n \geq 1} (\mathbb{S}^n \times \mathbb{R}^k)$ is noncompact finitistic space. (3) Paracompact spaces with finite covering dimension are finitistic spaces. Let G be a finite cyclic group acting on a finitistic space X with the fixed point set F . Note that G act freely on contractible space E_G (an infinite join of G with itself) and the orbit map $E_G \rightarrow B_G$ is the universal G -bundle. The associated fibration $X \xrightarrow{i} X_G \xrightarrow{\pi} B_G$ is called the Borel fibration, where $X_G = (X \times E_G)/G$ (Borel space) and $B_G = E_G/G$ (classifying space). If F is nonempty and $x \in F$, then $\eta_x : B_G \hookrightarrow X_G$ is cross section of π , where $B_G \approx \{x\} \times_G E_G$, and we get $H^*(X_G) = \ker \eta_x^* \oplus im \pi^*$. The induced homomorphism η_x^* depends on the component F_0 of F in which x lies. If $\alpha \in H^n(X_G)$ such that $\alpha \in \ker \eta_x^*$, then the image of α under the restriction of $j : F_G \hookrightarrow X_G$ on $(F_0)_G$ does not involve the term of $H^n(B_G) \otimes H^0(F_0)$. For details, we refer [3, 4].

A space X is said to be totally nonhomologous to zero (TNHZ) in X_G with respect to the ring R if the inclusion map $i : X \hookrightarrow X_G$ induces a surjection in the cohomology $i^* : H^*(X_G; R) \rightarrow H^*(X; R)$. So, if X is TNHZ in X_G , then for $a \in H^n(X)$ there exist

$\alpha' \in H^n(X_G)$ such that $i^*(\alpha') = a$. Define $\alpha = \alpha' - \pi^*\eta_x^*(\alpha')$. Then it is easy to see that $i^*(\alpha) = a$ and $\eta_x^*(\alpha) = 0$. We have used the following propositions to determine the fixed point sets of product three of spheres.

Proposition 2.1. ([3]) Let $G = \mathbb{Z}_2$ act on finitistic space X and $\sum \text{rk } H^i(X, \mathbb{Z}_2) < \infty$, then the following statements are equivalent:

- (a) X is TNHZ in X_G .
- (b) $\sum_{i \geq 0} \text{rk } H^i(F, \mathbb{Z}_2) = \sum_{i \geq 0} \text{rk } H^i(X, \mathbb{Z}_2)$.
- (c) $\pi_1(B_G)$ acts trivially on $H^*(X; \mathbb{Z}_2)$ and spectral sequence $E_2^{r,q}$ of fibration $X \hookrightarrow X_G \rightarrow B_G$ degenerates.

Proposition 2.2. ([3]) Let $G = \mathbb{Z}_2$ act on the finitistic space X . Suppose that A be a closed and invariant subset of X , and $H^i(X, A; \mathbb{Z}_2) = 0$ for $i > n$. Then $j^* : H^k(X_G, A_G; \mathbb{Z}_2) \rightarrow H^k(F_G, A_G \cap F_G; \mathbb{Z}_2)$ is an isomorphism for $k > n$. If (X, A) is TNHZ in (X_G, A_G) , then j^* is a monomorphism for all k .

Proposition 2.3. ([5]) Let X be TNHZ in X_G and $\{\gamma_j\}$ be a set of homogeneous elements in $H^*(X_G; \mathbb{Z}_2)$ such that $\{i^*(\gamma_j)\}$ forms \mathbb{Z}_2 -basis of $H^*(X; \mathbb{Z}_2)$. Then, $H^*(X_G; \mathbb{Z}_2)$ is the free $H^*(B_G)$ -module generated by $\{\gamma_j\}$.

Proposition 2.4. ([3]) For a finitistic G -space X , where G is cyclic group of prime order p , we have $\sum_{i \geq j} \text{rk } H^i(F; \mathbb{Z}_2) \leq \sum_{i \geq j} \text{rk } H^i(X; \mathbb{Z}_2)$, for each j .

Definition 2.5. [3] A space X is said to be poincaré duality space over a field k of formal dimension r if the following conditions are satisfied:

- (a) $H^*(X; k)$ is finitely generated.
- (b) $H^i(X; k) = 0$ for $i > r$ and $H^r(X; k) \approx k$.
- (c) An element $u \in H^i(X; k)$ is nonzero if and only if there exists an element $u' \in H^{r-i}(X; k)$ called poincaré dual of u , such that the cup product $0 \neq uu' \in H^r(X; k)$.

Recall that the poincaré duals of basis elements are unique [10].

Proposition 2.6. ([3]) Let X be a finitistic poincaré duality space over \mathbb{Z}_2 of formal dimension n , where p is prime. Let $G = \mathbb{Z}_2$ act on X , where X is TNHZ in X_G . Then, for each component F_0 of F is a poincaré duality space of formal dimension $r \leq n$. If $r = n$, then $F = F_0$ is connected and the restriction $H^*(X; \mathbb{Z}_2) \rightarrow H^*(F; \mathbb{Z}_2)$ are isomorphisms.

Proposition 2.7. [11] Let $G = \mathbb{Z}_2$ act on a finitistic space X , where X is TNHZ in X_G . If $H^*(X; \mathbb{Z}_2)$ is generated by k elements as an algebra, then each component of the fixed point set is generated by at most k elements.

Proposition 2.8. ([3]) Let $G = \mathbb{Z}_2$ act on a finitistic space X and $\pi_1(B_G)$ acts nontrivially on $H^*(X)$. Then, the E_2 term of the Leray-Serre spectral sequence of the fibration $X \xrightarrow{i} X_G \xrightarrow{\pi} B_G$ is given by

$$E_2^{k,i} = \begin{cases} \ker \tau & \text{for } k = 0, \\ \ker \tau / \text{im } \sigma & \text{for } k > 0, \end{cases}$$

where $\tau = \sigma = 1 + g^*$, g^* is induced by a generator g of G .

Proposition 2.9. ([3]) Let $G = \mathbb{Z}_2$ act on a finitistic space X . Then, for any $a \in H^n(X)$, the element $ag^*(a)$ is permanent cocycle in spectral sequence of $X_G \rightarrow B_G$.

Recall that $H^*(\mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l; \mathbb{Z}_2) = \mathbb{Z}_2[a, b, c] / \langle a^2, b^2, c^2 \rangle$, where $\deg a = n$, $\deg b = m$ and $\deg c = l$, $1 \leq n \leq m \leq l$.

Throughout the paper, $H^*(X)$ will denote the Čech cohomology of a space X with coefficient group $G = \mathbb{Z}_2$, and $X \sim_2 Y$, means $H^*(X; \mathbb{Z}_2) \cong H^*(Y; \mathbb{Z}_2)$.

3. FIXED POINT SETS OF INVOLUTIONS ON THE PRODUCT OF THREE SPHERES

Let $G = \mathbb{Z}_2$ act on a finitistic space $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, $1 \leq n \leq m \leq l$. If the fixed point set is empty, then the cohomology ring of the orbit space X/G has been discussed in [9]. In this section, we determine the possibilities of nonempty fixed points sets of involutions on X . First, we determine the possibilities of nonempty connected fixed point sets of involutions on X .

3.1. Connected fixed point sets of involutions on X . Suppose that X is TNHZ in $X_{\mathbb{Z}_2}$, then we have $\sum \text{rk } H^i(F) = \sum \text{rk } H^i(X) = 8$. By Proposition 2.6, we get F is Poincaré duality space of formal dimension $r \leq n + m + l$. Let $0 < q_1 \leq q_2 \leq q_3 \leq q_4 \leq q_5 \leq q_6 < r$ be the non vanishing dimensions of $H^*(F)$, and u_i is generator of q_i^{th} cohomology group of F , $1 \leq i \leq 6$ and v is generator of $H^r(F)$. Assume that $\alpha \in H^n(X_G)$, $\beta \in H^m(X_G)$ and $\gamma \in H^l(X_G)$ represents generators a, b and c of $H^*(X, x)$, respectively, such that $\eta_x(\alpha) = \eta_x(\beta) = \eta_x(\gamma) = 0$. By Proposition 2.3, we get $\{1, \alpha, \beta, \gamma, \alpha\beta, \alpha\gamma, \beta\gamma, \alpha\beta\gamma\}$ forms a basis for $H^*(X_G)$ over $H^*(B_G)$ -module. Note that the images of generators of $H^*(X_G)$ under the homomorphism $j^* : H^*(X_G) \rightarrow H^*(F_G)$ involves generators of $H^*(F)$. Since $a^2 = b^2 = c^2 = 0$, we get α^2, β^2 and γ^2 may be zero or nonzero, and hence $u_i u_j$, $1 \leq i, j \leq 6$, may be zero or nonzero. Further, if each generator u_i is Poincaré dual of itself or any two generators are Poincaré dual of itself or any four generators are Poincaré dual of itself, then we get $H^*(F)$ must have more than three generators, which contradicts Proposition 2.7. As $q_i \leq q_{i+1}$, $1 \leq i \leq 5$, we get u_1, u_2 and u_3 are Poincaré duals of u_6, u_5 and u_4 , respectively. So, we have $u_1 u_6 = u_2 u_5 =$

$u_3u_4 = v \in H^r(F)$. Clearly, $q_4 = r - q_3$, $q_5 = r - q_2$ and $q_6 = r - q_1$, and $u_2u_6 = u_3u_5 = u_3u_6 = u_4u_5 = u_4u_6 = u_5u_6 = v^2 = 0$ & $u_i v = 0, 1 \leq i \leq 6$. Thus, the possibilities of the fixed point set will depend on the cup products $u_1u_2, u_1u_3, u_1u_4, u_1u_5, u_2u_3$ and u_2u_4 which may be zero or nonzero. So, we have considered the cases depending upon how many cup products out of these six are simultaneously nonzero.

Theorem 3.1. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, where $1 \leq n \leq m \leq l$ and X is TNHZ in X_G . If F is connected and none of the cup products $u_1u_2, u_1u_3, u_1u_4, u_1u_5, u_2u_3$ and u_2u_4 of generators of $H^*(F)$ is nonzero, then $F \sim_2 \#_3 \mathbb{P}^3(q), q \leq n$ and $q = 1, 2, 4, 8$.

Proof. As discussed above, we have $H^*(X_G)$ is the free $H^*(B_G)$ -module with basis $\{1, \alpha, \beta, \gamma, \alpha\beta, \alpha\gamma, \beta\gamma, \alpha\beta\gamma\}$. Then we write

$$\begin{aligned} j^*(\alpha) &= A_1 t^{n-q_1} \otimes u_1 + A_2 t^{n-q_2} \otimes u_2 + A_3 t^{n-q_3} \otimes u_3 + A_4 t^{n-q_4} \otimes u_4 + \\ &\quad A_5 t^{n-q_5} \otimes u_5 + A_6 t^{n-q_6} \otimes u_6 + A_7 t^{n-r} \otimes v, \\ j^*(\beta) &= A'_1 t^{m-q_1} \otimes u_1 + A'_2 t^{m-q_2} \otimes u_2 + A'_3 t^{m-q_3} \otimes u_3 + A'_4 t^{m-q_4} \otimes u_4 + \\ &\quad A'_5 t^{m-q_5} \otimes u_5 + A'_6 t^{m-q_6} \otimes u_6 + A'_7 t^{m-r} \otimes v, \\ j^*(\gamma) &= A''_1 t^{l-q_1} \otimes u_1 + A''_2 t^{l-q_2} \otimes u_2 + A''_3 t^{l-q_3} \otimes u_3 + A''_4 t^{l-q_4} \otimes u_4 + \\ &\quad A''_5 t^{l-q_5} \otimes u_5 + A''_6 t^{l-q_6} \otimes u_6 + A''_7 t^{l-r} \otimes v, \end{aligned}$$

where all $A_i, A'_i, A''_i \in \mathbb{Z}_2 \forall 1 \leq i \leq 7$. As all cup products are zero, we get $u_4^2 = u_5^2 = u_6^2 = 0$. So, we have

$$j^*(\alpha\beta\gamma) = A_1 A'_1 A''_1 t^{l-q_1} \otimes u_1^3 + A_2 A'_2 A''_2 t^{l-q_2} \otimes u_2^3 + A_3 A'_3 A''_3 t^{l-q_3} \otimes u_3^3.$$

As j^* is injective and $\alpha\beta\gamma \neq 0$, we get $j^*(\alpha\beta\gamma) \neq 0$. Thus, at least one of u_1^3, u_2^3 and u_3^3 must be nonzero. Now, we consider following cases:

Case (i) Any one of u_1^3, u_2^3 and u_3^3 is nonzero.

Assume $u_1^3 \neq 0$ and $u_2^3 = u_3^3 = 0$. We must have $u_2^2 = u_3^2 = 0$ and $u_6 = u_1^2$. Thus, we get $u_1^3 = u_3u_4 = u_2u_5 = v$ is generator of $H^r(F)$. Clearly, the cohomology algebra of the fixed point set F has five generators which contradicts Proposition 2.7. So, this case not possible.

Case (ii) Any two of u_1^3, u_2^3 and u_3^3 are nonzero.

Assume $u_1^3 \neq 0, u_2^3 \neq 0$ and $u_3^3 = 0$. Then, we must have $u_6 = u_1^2, u_5 = u_2^2, u_3^2 = 0$ and $u_1^3 = u_2^3 = u_3u_4 = v$ generator of $H^r(F)$. Clearly, $q_1 = q_2$ and $q_5 = q_6$. It is easy to observe that the cohomological algebra of the fixed point set F has four generators, a contradiction. So, this case is also not possible.

Case (iii) u_1^3, u_2^3 and u_3^3 are nonzero.

Clearly, u_1^2, u_2^2 and u_3^2 are all nonzero, and we must have $u_4 = u_3^2, u_5 = u_2^2$ and $u_6 = u_1^2$.

Therefore, $u_1^3 = u_2^3 = u_3^3$ is generator of $H^r(F)$. In this case, we must have $q_1 = q_2 = q_3$. So, we get u_1, u_2, u_3 are generator of degree q , and u_1^2, u_2^2, u_3^2 are generators of degree $2q$. Thus, the cohomological algebra of fixed point set F is given by

$$\mathbb{Z}_2[u_1, u_2, u_3]/\langle u_1^4, u_2^4, u_3^4, u_1^3 + u_2^3, u_2^3 + u_3^3, u_1^3 + u_3^3, u_1u_2, u_2u_3, u_1u_3 \rangle,$$

where $\deg u_1 = \deg u_2 = \deg u_3 = q$. Thus, $F \sim_2 \#_3 \mathbb{P}^3(q)$. By Proposition 2.4, we get $q \leq n$, where $q = 1, 2, 3, 4$ [2]. \square

Theorem 3.2. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, where $1 \leq n \leq m \leq l$ and X is TNHZ in X_G . If F is connected and one cup product from $u_1u_2, u_1u_3, u_1u_4, u_1u_5, u_2u_3$ and u_2u_4 of generators of $H^*(F)$ is nonzero, then $F \sim_2 \mathbb{P}^3(r_1) \# (\mathbb{P}^2(r_2) \times \mathbb{S}^{r_3})$, where $\min \{r_1, r_2\} \leq n$ and $r_1, r_2 = 1, 2, 4, 8$.

Proof. First, assume that $u_1u_2 \neq 0$ and $u_1u_3 = u_1u_4 = u_1u_5 = u_2u_3 = u_2u_4 = 0$.

It is easy to see that either $u_1u_2 = u_5$ or $u_1u_2 = u_6$. Suppose that $u_1u_2 = u_5$. Then, we have $u_1u_6 = u_2^2u_1 = u_3u_4$ is generator of $H^r(F)$. As all the cup products u_1u_4, u_1u_5, u_2u_3 and u_2u_4 are zero, we must have $u_2^2 = u_6$ and $u_4^2 = 0$. Now, we consider four cases: (i) $u_1^2 = u_3^2 = 0$ (ii) $u_1^2 \neq 0, u_3^2 = 0$ (iii) $u_1^2 = 0, u_3^2 \neq 0$, and (iv) $u_1^2 \neq 0, u_3^2 \neq 0$.

Case (i): $u_1^2 = u_3^2 = 0$.

In this case, the cohomology algebra of the fixed point set has four generators, namely, u_1, u_2, u_3 and u_4 , which contradicts Proposition 2.7.

Case (ii): $u_1^2 \neq 0$ and $u_3^2 = 0$.

Since, $u_1u_3 = u_1u_4 = u_1u_5 = 0$, we must have $u_6 = u_1^2$. So, $u_2^2u_1 = u_1^3 = u_3u_4$ is generator of $H^r(F)$. Thus, the cohomological algebra of the fixed point set is generated by four generators u_1, u_2, u_3 and u_4 , a contradiction.

Case (iii): $u_1^2 = 0$ and $u_3^2 \neq 0$.

In this case, we get $u_4 = u_3^2$ and $q_1 = q_2 = q_3$. Thus, $H^{2q_1}(F)$ is generated by u_3^2, u_1u_2, u_2^2 , and $u_2^2u_1 = u_3^3$ generates $H^{3q_1}(F)$. Hence, the cohomological algebra of the fixed point set is given by

$$\mathbb{Z}_2[u_1, u_2, u_3]/\langle u_1^2, u_2^3, u_3^4, u_3^3 + u_2^2u_1, u_1u_3, u_2u_3 \rangle,$$

where $\deg u_1 = \deg u_2 = \deg u_3 = q_1$. Thus, $F \sim_2 \mathbb{P}^3(q_1) \# (\mathbb{P}^2(q_1) \times \mathbb{S}^{q_1})$. By Proposition 2.4, we get $q_1 \leq n$ and $q_1 = 1, 2, 4, 8$ [2].

Case (iv): $u_1^2 \neq 0$ and $u_3^2 \neq 0$.

In this case, we get $u_4 = u_3^2$ and $u_6 = u_1^2 = u_2^2$. So, $u_2^2u_1 = u_1^3 = u_3^3$ is generator of $H^r(F)$, and $q_1 = q_2 = q_3$. By the change of basis, $d = u_1 + u_2$, we get $d^2 = 0$, $u_2d = u_2u_1 + u_2^2 \neq 0$, $u_6d = u_1u_6 \neq 0$. Clearly, $u_3d = u_4d = u_5d = 0$ and $u_2^2d = u_3^2$ is generator of $H^{3q_1}(F)$. Hence, $F \sim_2 \mathbb{P}^3(q_1) \# (\mathbb{P}^2(q_1) \times \mathbb{S}^{q_1})$.

Now, we suppose $u_1u_2 = u_6$. Then, $u_1^2u_2 = u_2u_5 = u_3u_4$ generates $H^r(F)$. Clearly, $u_5 = u_1^2$ and $u_4^2 = 0$. Now, u_2^2 and u_3^2 are either zero or nonzero. Similarly, as discussed above, if $\{u_2^2 = 0 \text{ and } u_3^2 = 0\}$ or $\{u_2^2 \neq 0 \text{ and } u_3^2 = 0\}$, then the cohomology ring $H^*(F)$ has four generators, which is not possible. Now, if $u_2^2 = 0$ and $u_3^2 \neq 0$, then we must have $u_4 = u_3^2$. Thus, $q_1 = q_2 = q_3$, and $u_1^2u_2 = u_3^3$ generates $H^{3q_1}(F)$. Hence, the cohomological algebra of the fixed point set is given by

$$\mathbb{Z}_2[u_1, u_2, u_3]/\langle u_1^3, u_2^2, u_3^4, u_3^3 + u_1^2u_2, u_1u_3, u_2u_3 \rangle,$$

where $\deg u_1 = \deg u_2 = \deg u_3 = q_1$. Thus, $F \sim_2 \mathbb{P}^3(q_1)\#(\mathbb{P}^2(q_1) \times \mathbb{S}^{q_1})$, $q_1 \leq n$. Finally, if u_2^2 and u_3^2 both are nonzero, then we get $u_4 = u_3^2$ and $u_5 = u_1^2 = u_2^2$. By the change of basis, $d = u_1 + u_2$, we get $F \sim_2 \mathbb{P}^3(q_1)\#(\mathbb{P}^2(q_1) \times \mathbb{S}^{q_1})$, $q_1 \leq n$.

Similarly, if $u_1u_3 \neq 0$ or $u_2u_3 \neq 0$, then $F \sim_2 \mathbb{P}^3(q_1)\#(\mathbb{P}^2(q_1) \times \mathbb{S}^{q_1})$.

Next, assume that $u_1u_4 \neq 0$. Here, we must have $u_1u_4 = u_6$, and $u_1^2u_4 = u_2u_5 = u_3u_4$ is generator of $H^r(F)$. Clearly, $u_4^2 = u_5^2 = 0$ and $u_3 = u_1^2$. If $u_2^2 = 0$, then cohomology algebra of the fixed point set has four generators, which is not possible. If $u_2^2 \neq 0$, then $u_5 = u_2^2$ and we get $F \sim_2 \mathbb{P}^3(q_2)\#(\mathbb{P}^2(q_1) \times \mathbb{S}^{q_4})$, $q_1 \leq n$ and $q_1, q_2 = 1, 2, 4, 8$.

Similarly, if $u_1u_5 \neq 0$, then $F \sim_2 \mathbb{P}^3(q_2)\#(\mathbb{P}^2(q_1) \times \mathbb{S}^{q_5})$, and if $u_2u_4 \neq 0$, then $F \sim_2 \mathbb{P}^3(q_1)\#(\mathbb{P}^2(q_2) \times \mathbb{S}^{q_4})$. This completes the proof. \square

Theorem 3.3. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, where $1 \leq n \leq m \leq l$ and X is TNHZ in X_G . If F is connected and two cup products from $u_1u_2, u_1u_3, u_1u_4, u_1u_5, u_2u_3$ and u_2u_4 of generators of $H^*(F)$ are nonzero, then F must be one of the following:

- (1) $F \sim_2 (\mathbb{P}^5(r_1)\#(\mathbb{S}^{r_2} \times \mathbb{S}^{r_3}))$, where $r_1 = 1, 2, 4, 8$ and $\min\{r_1, r_2, r_3\} \leq n$.
- (2) $F \sim_2 (\mathbb{P}^2(r_1)\#\mathbb{P}^2(r_1)) \times \mathbb{S}^{r_2}$, where $r_1 = 1, 2, 4, 8$ and $\min\{r_1, r_2\} \leq n$.
- (3) $F \sim_2 \mathbb{Z}_2[x, y, z]/\langle x^3, u^3, v^2, xv + u^2, yz \rangle$, where $(u, v) \in \{(y, z), (z, y)\}$ and $(\deg x, \deg y, \deg z) \in \{(q_1, q_2, q_3), (q_1, q_2, q_4), (q_2, q_1, q_3), (q_1, q_3, q_5), (q_3, q_1, q_2)\}$.
- (4) $F \sim_2 \mathbb{Z}_2[x, y, z]/\langle x^2, y^3, z^3, y^2 + z^2 + ux, yz \rangle$, where $u \in \{y, z\}$ and $(\deg x, \deg y, \deg z) \in \{(q_1, q_2, q_3), (q_2, q_1, q_3), (q_3, q_1, q_2)\}$.

Proof. Clearly, we have $\binom{6}{2} = 15$ cases depending on pairs of nonzero cup products. It is easy to see that out of these fifteen cases, the following six nonzero pairs are not possible: (1) u_1u_2, u_2u_4 (2) u_1u_3, u_2u_4 (3) u_1u_4, u_1u_5 (4) u_1u_4, u_2u_3 (5) u_1u_5, u_2u_3 , and (6) u_1u_5, u_2u_4 . A contradiction of these cases occurs from the generators whose cup products are zero. Now, we consider remaining nonzero pairs of cup products:

Case (7): $u_1u_2 \neq 0$ & $u_1u_5 \neq 0$.

As $u_1u_3 = u_1u_4 = u_2u_3 = u_2u_4 = 0$, we must have either $u_1u_5 = u_1u_5 = u_6$ or $\{u_1u_2 = u_5 \text{ and } u_1u_5 = u_6\}$. First, suppose that $u_1u_2 = u_1u_5 = u_6$. Then, $u_1^2u_2 = u_1^2u_5 = u_2u_5 = u_3u_4$ is generator of $H^r(F)$. So, $u_1^2 \neq 0$. As $u_1u_3 = u_1u_4 = 0$, we have

either $u_1^2 = u_2$ or $u_1^2 = u_5$. Further, in either cases, u_2 or u_5 have two poincaré duals, a contradiction.

Next, suppose that $u_1u_2 = u_5$ and $u_1u_5 = u_6$. Then, we have $u_1u_5 = u_1^2u_2$ and $u_1^3u_2 = u_2^2u_1 = u_3u_4$ is generator of $H^r(F)$. Since, $u_1u_3 = u_1u_4 = 0$, we have $u_2 = u_1^2$. Thus, $u_1u_2 = u_1^3$, $u_1^2u_2 = u_1^4$ and $u_1^5 = u_3u_4$ are generators of $H^{q_5}(F)$, $H^{q_6}(F)$ and $H^r(F)$, respectively. Clearly, $u_4^2 = 0$. If $u_3^2 \neq 0$, then we must have $u_4 = u_3^2$, and hence $F \sim_2 \mathbb{P}^5(q_1) \# \mathbb{P}^3(q_3)$, where $5q_1 = 3q_3$. By Adam [2], this is not possible. If $u_3^2 = 0$, then we get $F \sim_2 \mathbb{P}^5(q_1) \# (\mathbb{S}^{q_3} \times \mathbb{S}^{q_4})$. By Proposition 2.4, $q_1 \leq n$. This realizes possibility (1).

Similarly, for Case (8) $u_1u_3 \neq 0$ & $u_1u_4 \neq 0$, we get $F \sim_2 \mathbb{P}^5(q_1) \# (\mathbb{S}^{q_2} \times \mathbb{S}^{q_5})$, and for Case (9) $u_2u_3 \neq 0$ & $u_2u_4 \neq 0$, we get $F \sim_2 \mathbb{P}^5(q_6) \# (\mathbb{S}^{q_1} \times \mathbb{S}^{q_6})$. So, these cases also realizes possibility (1).

Case (10): $u_1u_4 \neq 0$ & $u_2u_4 \neq 0$.

In this case, we must have $u_2u_4 = u_5$ and $u_1u_4 = u_6$. Thus, we have $u_1^2u_4 = u_2^2u_4 = u_3u_4$ is a generator of $H^r(F)$. As $u_1u_2 = u_1u_3 = u_1u_5 = u_2u_3 = 0$, we get $u_1^2 = u_2^2 = u_3$. Clearly, $q_1 = q_2$ and $u_4^2 = 0$. Hence, $F \sim_2 (\mathbb{P}^2(q_1) \# \mathbb{P}^2(q_1)) \times \mathbb{S}^{q_4}$, $q_1 \leq n$. This realizes possibility (2).

Case (11): $u_1u_2 \neq 0$ and $u_1u_4 \neq 0$.

In this case, we get either $u_1u_4 = u_1u_5 = u_6$ or $\{u_1u_2 = u_5$ and $u_1u_4 = u_6\}$. If $u_1u_2 = u_1u_4 = u_6$, then by uniqueness of poincaré dual, this is not possible.

Next, Suppose that $u_1u_2 = u_5$ and $u_1u_4 = u_6$. Then, we have $u_1^2u_4 = u_2^2u_1 = u_3u_4$ generates of $H^r(F)$. Clearly, $u_1^2 = u_3$, $u_2^2 = u_6$ and $u_4^2 = 0$. Consequently, $u_1^3 = u_2^3 = u_4^2 = 0$. Hence, the cohomology ring of the fixed point set is isomorphic to

$$\mathbb{Z}_2[u_1, u_2, u_4] / \langle u_1^3, u_2^3, u_4^2, u_1u_4 + u_2^2, u_2u_4 \rangle$$

where $\deg u_1 = q_1$, $\deg u_2 = q_2$ and $\deg u_4 = q_4$. This realizes possibility (3) for $(u, v) = (y, z)$, $x = u_1$, $y = u_2$ and $z = u_4$.

Similarly, for Case (12) $u_1u_3 \neq 0$ & $u_1u_5 \neq 0$, the cohomology ring of the fixed point set is isomorphic to

$$\mathbb{Z}_2[u_1, u_3, u_5] / \langle u_1^3, u_3^3, u_5^2, u_1u_5 + u_3^2, u_3u_5 \rangle$$

where $\deg u_1 = q_1$, $\deg u_3 = q_3$ and $\deg u_5 = q_5$. This also realizes possibility (3) for $(u, v) = (y, z)$, $x = u_1$, $y = u_3$ and $z = u_5$.

Case (13): $u_1u_2 \neq 0$ and $u_1u_3 \neq 0$.

In this case, we have following subcases (i) $u_1u_2 = u_1u_3 = u_6$ (ii) $u_1u_2 = u_5$ & $u_1u_3 = u_4$ (iii) $u_1u_2 = u_5$ & $u_1u_3 = u_6$, and (iv) $u_1u_2 = u_6$ & $u_1u_3 = u_4$. Note that subcase (i) is not possible.

Subcase (ii): $u_1u_2 = u_5$ and $u_1u_3 = u_4$. Then, we have $u_2^2u_1 = u_3^2u_1 = u_1u_6$ generates $H^r(F)$. It is easy to see that $u_6 = u_3^2 = u_2^2$ is generator of $H^{q_6}(F)$. Then, $q_2 = q_3$. Now, if $u_1^2 = 0$, then $F \sim_2 (\mathbb{P}^3(q_2) \# \mathbb{P}^3(q_2)) \times \mathbb{S}^{q_1}$. If $u_1^2 \neq 0$, then we must have $u_1^2 = u_6$. So, we get $q_1 = q_2 = q_3$. By the change of basis $d = u_1 + u_2$, we get $F \sim_2 (\mathbb{P}^3(q_1) \# \mathbb{P}^3(q_1)) \times \mathbb{S}^{q_1}$. These realizes possibility (2).

Subcase (iii): $u_1u_2 = u_5$ and $u_1u_3 = u_6$. Then, $u_1^2u_3 = u_2^2u_1 = u_3u_4$ generates $H^r(F)$. Clearly, $u_4 = u_1^2$ and $u_1u_3 = u_2^2$. Consequently, $u_1^3 = u_3^3 = 0$. If $u_2^2 = 0$, then the cohomology ring of the fixed point set is isomorphic to

$$\mathbb{Z}_2[u_1, u_2, u_3]/\langle u_1^3, u_2^3, u_3^2, u_1u_3 + u_2^2, u_2u_3 \rangle,$$

where $\deg u_1 = q_1$, $\deg u_2 = q_2$ and $\deg u_3 = q_3$. This realizes possibility (3) for $(u, v) = (y, z)$, $x = u_1, y = u_2$ and $z = u_3$. If $u_2^2 \neq 0$, then we get $u_4 = u_3^2 = u_1^2$. Thus, $q_1 = q_2 = q_3$, and by change of basis $d = u_1 + u_3$, we get $\{d, u_2, u_3\}$ generates $H^{q_1}(F)$, $\{u_3d, u_2d, u_2^2 = u_3^2 + u_3d\}$ generates $H^{2q_1}(F)$ and $u_2^2d = u_3^2d$ generates $H^{3q_1}(F)$. Thus, the cohomology ring $H^*(F)$ is isomorphic to

$$\mathbb{Z}_2[d, u_2, u_3]/\langle d^2, u_2^3, u_3^3, u_3d + u_2^2 + u_3^2, u_2u_3 \rangle,$$

where $\deg d = \deg u_2 = \deg u_3 = q_1$. This realizes possibility (4) for $u = z, x = d, y = u_2$ and $z = u_3$.

Subcase (iv): $u_1u_2 = u_6$ and $u_1u_3 = u_4$. Then, $u_1^2u_2 = u_3^2u_1$ generates $H^r(F)$. It is easy to see that $u_5 = u_1^2$ and $u_6 = u_3^2$. If $u_2^2 = 0$, then the cohomology ring of the fixed point set is isomorphic to

$$\mathbb{Z}_2[u_1, u_2, u_3]/\langle u_1^3, u_2^2, u_3^3, u_1u_2 + u_3^2, u_2u_3 \rangle,$$

where $\deg u_1 = q_1$, $\deg u_2 = q_2$ and $\deg u_3 = q_3$. This realizes possibility (3) for $(u, v) = (z, y)$ and $x = u_1, y = u_2$ & $z = u_3$. Now, if $u_2^2 \neq 0$, then $u_5 = u_1^2 = u_2^2$, and again by the change of basis $d = u_1 + u_2$, we get $H^*(F)$ is isomorphic to

$$\mathbb{Z}_2[d, u_2, u_3]/\langle d^2, u_2^3, u_3^3, u_2d + u_2^2 + u_3^2, u_2u_3 \rangle,$$

where $\deg d = \deg u_2 = \deg u_3 = q_1$. This realizes possibility (4) for $u = y, x = d, y = u_2$ and $z = u_3$.

Similarly, for Case (14) $u_1u_3 \neq 0$ & $u_2u_3 \neq 0$, we realize possibility (2), possibility (3) for $(u, v) = (y, z)$ & (z, y) ; $x = u_2, y = u_1$ & $z = u_3$, and possibility (3) for $u = y$ & z ; $x = u_2, y = d$ & $z = u_3$, and for Case (15) $u_1u_2 \neq 0$ & $u_2u_3 \neq 0$, we realize possibility (2), possibility (3) for $(u, v) = (y, z)$ & (z, y) ; $x = u_3, y = u_2$ & $z = u_1$, and possibility (3) for $u = y$ & z ; $x = u_1, y = u_2$ & $z = d$. \square

Theorem 3.4. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, where $1 \leq n \leq m \leq l$ and X is TNHZ in X_G . If F is connected and three cup products from $u_1u_2, u_1u_3, u_1u_4, u_1u_5, u_2u_3$ and u_2u_4 of generators of $H^*(F)$ are nonzero. Then, F must be one of the following:

- (1) $F \sim_2 \mathbb{Z}_2[x, y]/\langle x^6, y^3, x^4 + y^2 \rangle$, where $\deg x = q_1$ and $\deg y \in \{q_2, q_3\}$.
- (2) $F \sim_2 \mathbb{Z}_2[x, y, z]/\langle x^{2+i}, y^{2+j}, z^{2+k}, a_i(yz + x^i), a_j(xz + y^j), a_k(xy + z^k) \rangle$, where $i, j, k \in \{0, 2\}$ or $i = j = k = 1, a_0 = 0, a_1 = a_2 = 1$ and $\deg x = q_1, \deg y = q_2$ & $\deg z \in \{q_3, q_4\}$. In particular, for $i = j = k = 0, F \sim_2 \mathbb{S}^{r_1} \times \mathbb{S}^{r_2} \times \mathbb{S}^{r_3}$, where $r_1 \leq n, r_2 \leq m$ and $r_3 \leq l$.

Proof. Here, we have $\binom{6}{3} = 20$ cases depending on which three nonzero cup products are taken from $u_1u_2, u_1u_3, u_1u_4, u_1u_5, u_2u_3$ and u_2u_4 . Out of these twenty cases, the following four cases are possible: (1) $u_1u_2 \neq 0, u_1u_3 \neq 0$ & $u_1u_4 \neq 0$, (2) $u_1u_2 \neq 0, u_1u_3 \neq 0$ & $u_1u_5 \neq 0$, (3) $u_1u_2 \neq 0, u_1u_3 \neq 0$ & $u_2u_3 \neq 0$, and (4) $u_1u_2 \neq 0, u_1u_4 \neq 0$ & $u_2u_4 \neq 0$.

Case (1) $u_1u_2 \neq 0, u_1u_3 \neq 0$ and $u_1u_4 \neq 0$.

In this case, we must have $u_1u_3 = u_4, u_1u_2 = u_5$, and $u_1u_4 = u_6$. Thus, $u_1u_4 = u_1^2u_3$ and $u_1u_2^2 = u_1^3u_3 = u_1u_3^2$ is the generator of $H^r(F)$. So, u_3^2, u_2^2 and u_1^3 are nonzero. Since, $u_1u_5 = 0$, we must get $u_3 = u_1^2$. Thus, $u_4 = u_1^3, u_6 = u_1^4$, and $u_5^5 = u_2^2u_1$. Hence, the cohomology ring $H^*(F)$ is isomorphic to

$$\mathbb{Z}_2[u_1, u_2]/\langle u_1^6, u_2^3, u_2^2 + u_1^4 \rangle,$$

where $\deg u_1 = q_1$ and $\deg u_2 = q_2$. This realize possibility (1) for $x = u_1, y = u_2$ and $\deg y = q_2$.

Similarly, for Case (2) $u_1u_2 \neq 0, u_1u_3 \neq 0, u_1u_5 \neq 0$, we realizes possibility (1) for $x = u_1, y = u_3$ and $\deg y = q_3$.

Case(3) $u_1u_2 \neq 0, u_1u_3 \neq 0$ and $u_2u_3 \neq 0$.

As the cup products u_1u_4, u_1u_5 and u_2u_3 are zero, only the following subcases are possible: (i) $u_1u_2 = u_4, u_1u_3 = u_5$ & $u_2u_3 = u_6$, (ii) $u_1u_2 = u_5, u_1u_3 = u_6$ & $u_2u_3 = u_4$, and (iii) $u_1u_2 = u_6, u_1u_3 = u_4$ & $u_2u_3 = u_5$.

Subcase (i) $u_1u_2 = u_4, u_1u_3 = u_5$ and $u_2u_3 = u_6$. Then, $u_1u_2u_3$ is the generator of $H^r(F)$. Now, if u_1^2, u_2^2 and u_3^2 are all zero, then $F \sim_2 \mathbb{S}^{q_1} \times \mathbb{S}^{q_2} \times \mathbb{S}^{q_3}$. By Proposition 2.4, we get $r_1 \leq n, r_2 \leq m$ and $r_3 \leq l$. This realizes possibility (2) for $i = j = k = 0$.

Suppose that u_1^2, u_2^2 and u_3^2 are all nonzero. Then, we must get $u_4 = u_3^2, u_5 = u_2^2$ and $u_6 = u_3^2$. Clearly, $u_1^3 = u_2^3 = u_3^3 = u_1u_2u_3$ and $u_1^4 = u_2^4 = u_3^4 = 0$. Thus, the cohomology ring $H^*(F)$ is isomorphic to

$$\mathbb{Z}_2[u_1, u_2, u_3]/\langle u_1^4, u_2^4, u_3^4, u_1u_2 + u_3^2, u_1u_3 + u_2^2, u_2u_3 + u_1^2 \rangle,$$

where $\deg u_1 = q_1$, $\deg u_2 = q_2$ and $\deg u_3 = q_3$. This realizes possibility (2) for $i = j = k = 2$.

Suppose that any two of u_1^2, u_2^2 and u_3^2 are nonzero. Assume u_1^2 and u_2^2 are nonzero and $u_3^2 = 0$. Then, $u_5 = u_2^2$ and $u_6 = u_1^2$. So, the cohomology ring $H^*(F)$ is isomorphic to

$$\mathbb{Z}_2[u_1, u_2, u_3]/\langle u_1^4, u_2^4, u_3^2, u_1u_3 + u_2^2, u_2u_3 + u_1^2 \rangle,$$

where $\deg u_1 = q_1$, $\deg u_2 = q_2$ and $\deg u_3 = q_3$. This realizes possibility (2) for $i = j = 2$ & $k = 0$.

Finally, suppose that one of u_1^2, u_2^2 and u_3^2 is nonzero. Assume $u_1^2 \neq 0$ and $u_2^2 = u_3^2 = 0$. Then, $u_6 = u_1^2$. So, the cohomology ring $H^*(F)$ is isomorphic to

$$\mathbb{Z}_2[u_1, u_2, u_3]/\langle u_1^4, u_2^2, u_3^2, u_2u_3 + u_1^2 \rangle,$$

where $\deg u_1 = q_1$, $\deg u_2 = q_2$ and $\deg u_3 = q_3$. This realizes possibility (2) for $i = 2$ & $j = k = 0$.

Subcase (ii) $u_2u_3 = u_4, u_1u_2 = u_5$ and $u_1u_3 = u_6$. Then, $u_1^2u_3 = u_2^2u_1 = u_3^2u_2$ generates $H^r(F)$. Clearly, $u_1^2 = u_4, u_3^2 = u_5$ and $u_2^2 = u_6$. Thus, the cohomology ring $H^*(F)$ is isomorphic to

$$\mathbb{Z}_2[u_1, u_2, u_3]/\langle u_1^3, u_2^3, u_3^3, u_2u_3 + u_1^2, u_1u_2 + u_3^2, u_1u_3 + u_2^2 \rangle,$$

where $\deg u_1 = q_1$, $\deg u_2 = q_2$ and $\deg u_3 = q_3$. This realizes possibility (2) for $i = j = k = 1$.

Subcase (iii) $u_2u_3 = u_5, u_1u_2 = u_6$ and $u_1u_3 = u_4$. In this subcase, we get same cohomology ring $H^*(F)$ as in subcase (ii).

Case (4) $u_1u_2 \neq 0, u_1u_4 \neq 0, u_2u_4 \neq 0$.

In this case, we must have $u_1u_2 = u_3, u_1u_4 = u_5$ and $u_2u_4 = u_6$. So, $u_1u_2u_4$ is generator of $H^r(F)$. If $u_4^2 \neq 0$, then either $u_4^2 = u_5$ or u_6 , it fails the uniqueness of Poincaré dual of a generating element. So, $u_4^2 = 0$. Now, if $u_1^2 = u_2^2 = 0$, then $F \sim_2 \mathbb{S}^{q_1} \times \mathbb{S}^{q_2} \times \mathbb{S}^{q_3}$. This realizes possibility (2) for $i = j = k = 0$. If both u_1^2 and u_2^2 are nonzero, then we get $u_6 = u_1^2$ and $u_5 = u_2^2$. Thus, the cohomology ring $H^*(F)$ is isomorphic to

$$\mathbb{Z}_2[u_1, u_2, u_4]/\langle u_1^4, u_2^4, u_4^2, u_1u_4 + u_2^2, u_2u_4 + u_1^2 \rangle,$$

where $\deg u_1 = q_1$, $\deg u_2 = q_2$ and $\deg u_4 = q_4$. This realizes possibility (2) for $i = j = 2$ & $k = 0$.

Finally, if any one of u_1^2 and u_2^2 is nonzero, say $u_1^2 \neq 0$, then $u_6 = u_1^2$. Thus, the cohomology ring $H^*(F)$ is isomorphic to

$$\mathbb{Z}_2[u_1, u_2, u_4]/\langle u_1^4, u_2^2, u_4^2, u_2u_4 + u_1^2 \rangle,$$

where $\deg u_1 = q_1$, $\deg u_2 = q_2$ and $\deg u_4 = q_4$. This realizes possibility (2) for $i = 2$ & $j = k = 0$.

It is easy to observe that the remaining sixteen cases can be discarded by the zero cup products or by the uniqueness of poincaré duals. \square

Theorem 3.5. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, where $1 \leq n \leq m \leq l$ and X is TNHZ in X_G . If F is connected and any four cup products from $u_1u_2, u_1u_3, u_1u_4, u_1u_5, u_2u_3$ and u_2u_4 of generators of $H^*(F)$ are nonzero. Then, F must be one of the following:

- (1) $F \sim_2 \mathbb{P}^3(r_1) \times \mathbb{S}^{r_2}$, where $r_1 = 1, 2, 4, 8$.
- (2) $F \sim_2 \mathbb{Z}_2[x, y]/\langle x^4, y^4, x^3 + y^2, y^2x \rangle$, where $\deg x = q_1$ and $\deg y = q_2$.

Proof. We consider $\binom{6}{4} = 15$ cases, depending on which pair of cup products $u_1u_2, u_1u_3, u_1u_4, u_1u_5, u_2u_3, u_2u_4$ are zero and rest four cup products are nonzero. Out of these fifteen cases, the following six cases are possible: (i) $u_1u_3 = u_2u_3 = 0$, (ii) $u_1u_3 = u_1u_5 = 0$, (iii) $u_1u_4 = u_1u_5 = 0$, (iv) $u_1u_4 = u_2u_4 = 0$, (v) $u_1u_5 = u_2u_3 = 0$, and (vi) $u_1u_5 = u_2u_4 = 0$.

Case (i) $u_1u_3 = u_2u_3 = 0$.

We have u_1u_2, u_1u_4, u_1u_5 and u_2u_4 are nonzero cup products. Clearly, $u_1u_5 = u_6$. By the uniqueness of poincaré dual of u_1^2 , we must have $u_1u_4 = u_5$. Similarly, we get $u_1u_2 = u_3$ and $u_1u_5 = u_2u_4 = u_6$. Then, $u_1^3u_4 = u_1u_2u_4$ is the generator of $H^r(F)$. It is easy to see that, $u_2 = u_1^2$ and $u_4^2 = 0$. Thus, $0 = u_1u_3 = u_1^4$, and hence $F \sim_2 \mathbb{P}^3(q_1) \times \mathbb{S}^{q_4}$. Note that $q_1 = 1, 2, 4, 8$. This realize possibility (1)

Similarly, for Case (ii) & (iii), we get $F \sim_2 \mathbb{P}^3(q_2) \times \mathbb{S}^{q_1}$, for Case (iv), we get $F \sim_2 \mathbb{P}^3(q_1) \times \mathbb{S}^{q_3}$, and for Case (v), we get $F \sim_2 \mathbb{P}^3(q_1) \times \mathbb{S}^{q_2}$. This also realizes possibility (1).

Case (vi) $u_1u_5 = u_2u_4 = 0$.

As $u_1u_5 = u_2u_4 = 0$, we get $u_1u_4 = u_6$, $u_1u_2 = u_4$, $u_1u_3 = u_5$ and $u_1u_4 = u_2u_3 = u_6$. Thus, $u_1^3u_2 = u_1u_2u_3$ generates $H^r(F)$. Clearly, $u_1^2 = u_3$. So, if $u_2^2 = 0$, then $F \sim_2 \mathbb{P}^3(q_1) \times \mathbb{S}^{q_2}$. This realizes possibility (1). If $u_2^2 \neq 0$, then we must have $u_5 = u_1^3 = u_2^2$ and $u_1^3u_2 = u_2^3$ generates $H^r(F)$. We have $0 = u_1u_5 = u_1^4$ and $0 = u_2u_4 = u_2^2u_1$. Hence, the cohomology ring $H^*(F)$ is isomorphic to

$$\mathbb{Z}_2[u_1, u_2]/\langle u_1^4, u_2^4, u_1^3 + u_2^2, u_2^2u_1 \rangle,$$

where $\deg u_1 = q_1$ and $\deg u_2 = q_2$. This realizes possibility (2).

It is easy to observe that the remaining nine cases are not possible. \square

Theorem 3.6. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, where $1 \leq n \leq m \leq l$ and X is TNHZ in X_G . If F is connected and any five cup products from $u_1u_2, u_1u_3, u_1u_4, u_1u_5, u_2u_3$

and u_2u_4 of generators of $H^*(F)$ are nonzero. Then, $F \sim_2 \mathbb{P}^3(r_1) \times \mathbb{S}^{r_1}$, where $r_1 = 1, 2, 4, 8$.

Proof. Here, we have six cases depending on which cup product out of $u_1u_2, u_1u_3, u_1u_4, u_1u_5, u_2u_3$ and u_2u_4 is zero, and rest five cup products are nonzero. Clearly, the only possible case is $u_1u_5 = 0$. We have either $u_2u_4 = u_6$ or $u_2u_4 = u_5$.

If $u_2u_4 = u_6$, then we must have $u_1u_3 = u_6, u_1u_4 = u_2u_3 = u_5$ and $u_1u_2 = u_3$. Thus, $u_1^3u_2 = u_2^3u_1$ generates $H^r(F)$. Clearly, $u_1^2 = u_2^2 = u_4$. So, we get $u_1^4 = u_2^4 = 0$. By the change of basis $d = u_1 + u_2$, we get $\{d, u_2\}$ generates $H^{q_1}(F)$, $\{u_2d, u_2^2\}$ generates $H^{q_3}(F)$, $\{u_2^2d, u_2^3\}$ generates $H^{q_5}(F)$, and u_2^3d generates $H^r(F)$. Hence, $F \sim_2 \mathbb{P}^3(q_1) \times \mathbb{S}^{q_1}$.

Now, if $u_2u_4 = u_5$, then we must have $u_1u_3 = u_5, u_1u_4 = u_2u_3 = u_6$, and $u_1u_2 = u_4$. So, $u_1^3u_2 = u_2^3u_1$ generates $H^r(F)$. Similarly, by the change of basis, we get $F \sim_2 \mathbb{P}^3(q_1) \times \mathbb{S}^{q_1}$. Hence, our claim. \square

If all cup products $u_1u_2, u_1u_3, u_1u_4, u_1u_5, u_2u_3$ and u_2u_4 are nonzero, then we get $u_1u_2 = u_3, u_1u_3 = u_4, u_1u_4 = u_2u_3 = u_5, u_1u_5 = u_2u_4 = u_6$. So, we must have $u_1^2 = u_2$, and hence $u_1^7 = 0$. Thus, we have:

Theorem 3.7. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, where $1 \leq n \leq m \leq l$ and X is TNHZ in X_G . If F is connected and all cup products $u_1u_2, u_1u_3, u_1u_4, u_1u_5, u_2u_3$ and u_2u_4 of generators of $H^*(F)$ are nonzero, then $F \sim_2 \mathbb{P}^7(q)$, $q = 1, 2, 4, 8$.

Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^{n_1} \times \mathbb{S}^{n_2} \times \cdots \times \mathbb{S}^{n_k}$, where $1 \leq n_1 \leq n_2 \leq \cdots \leq n_k$ and X is TNHZ in X_G . Then, $\text{rk}H^*(F) = 2^k$. Suppose that F is connected. Let $0 < q_1 \leq q_2 \leq \cdots \leq q_{2^k-2} < r$ be non vanishing dimensions of $H^*(F)$ and u_i is generator of q_i^{th} cohomology group, and v is generator of $H^r(F)$. WLOG, we may assume that $u_1, \cdots, u_{2^{k-1}-1}$ has poincaré duals $u_{2^k-2}, \cdots, u_{2^{k-1}}$, respectively. Thus, $v = u_1u_{2^k-2} = \cdots = u_{2^{k-1}-1}u_{2^{k-1}}$. Clearly, $u_iu_j = 0$ if $i + j \geq 2^k$. It is easy to derive the following results:

Theorem 3.8. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^{n_1} \times \mathbb{S}^{n_2} \times \cdots \times \mathbb{S}^{n_k}$, where $1 \leq n_1 \leq n_2 \leq \cdots \leq n_k$ and X is TNHZ in X_G . If F is connected and all the cup products of generators of $H^*(F)$ other than generator of r^{th} cohomology group are zero, then $F \sim_2 \#_k \mathbb{P}^k(q)$, $q \leq n_1$ and $q = 1, 2, 4, 8$.

Theorem 3.9. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^{n_1} \times \mathbb{S}^{n_2} \times \cdots \times \mathbb{S}^{n_k}$, where $1 \leq n_1 \leq n_2 \leq \cdots \leq n_k$ and X is TNHZ in X_G . If F is connected and all the cup products u_iu_j , where $i + j \leq 2^k - 2, i \neq j$, of generators of $H^*(F)$ other than r^{th} cohomology group are nonzero, then $F \sim_2 \mathbb{P}^{2^k-1}(q)$, $q = 1, 2, 4, 8$.

Next, suppose that X is not TNHZ in X_G and F is nonempty connected fixed point set of involutions on X . So, we have $0 < \text{rk } H^*(F) < 8$. By the Floyd's formula [3], we get $\text{rk } H^*(F) = 2, 4$ or 6 . If $\pi_1(B_G)$ acts nontrivially on $H^*(X)$, then we have proved the following theorem:

Theorem 3.10. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l, 1 \leq n \leq m \leq l$ and X is not TNHZ in X_G . If F is nonempty connected fixed point set and $\pi_1(B_G)$ acts nontrivially on $H^*(X)$, then F must be the one of the following:

- (1) $F \sim_2 \mathbb{S}^{r_1}, 0 < r_1 \leq n + m + l$.
- (2) $F \sim_2 \mathbb{S}^{r_1} \times \mathbb{S}^{r_2}, 0 < r_1 \leq \min\{2n, l\}$ or $n, 0 < r_2 \leq \max\{2n, l\}$ or $2m$.
- (3) $F \sim_2 \mathbb{P}^3(r_1), r_1 \leq \min\{2n, l\}$ or $n, r_1 = 1, 2, 4, 8$.
- (4) $F \sim_2 \mathbb{P}^2(r_1) \# \mathbb{P}^2(r_1), r_1 \leq \min\{2n, l\}$ or $n, r_1 = 1, 2, 4, 8$.
- (5) $F \sim_2 \mathbb{P}^2(r_1) \vee \mathbb{S}^{r_2}, r_1 \leq \min\{2n, l\}$ or $n, 0 < r_2 \leq 2n + l$ or $n + 2m, r_1 = 1, 2, 4, 8$.

Proof. As $\pi_1(B_G) = G$ acts nontrivially on $H^*(X)$, we must have one of the following conditions: (i) $n = m < l$ (ii) $n < m = l$ (iii) $n = m = l$.

If $n = m < l$, then G acts nontrivially on $H^n(X)$. Let g be generator of G . We have exactly three possibilities of nontrivial actions: (1) $g^*(a) = b$ & $g^*(b) = a$, (2) $g^*(a) = a$ & $g^*(b) = a + b$ and (3) $g^*(a) = a + b$ & $g^*(b) = b$. First, we consider nontrivial action defined by $g^*(a) = b$ & $g^*(b) = a$. By the naturality of cup product, we get G also acts nontrivially on $H^{n+l}(X)$. Note that $\sigma = \tau = 1 + g^*$. By Proposition 2.8, we get $E_2^{0,i} \cong \mathbb{Z}_2$ and $E_2^{k,i} = 0 \forall k > 0, i = n, n+l$. For $i \neq n, n+l, \pi_1(B_G)$ acts trivially on $H^i(X)$, and hence, $E_2^{k,i} \cong H^k(B_G) \otimes H^i(X) \cong \mathbb{Z}_2 \forall k \geq 0, i = 0, l, 2n, 2n + l$. By Proposition 2.9, $1 \otimes ab$ is a permanent cocycle.

First, suppose that $E_2^{*,*} \neq E_\infty^{*,*}$. As $F \neq \emptyset$, we must have $d_{l+1}(1 \otimes c) = 0$. If d_{l-2n+1} is nontrivial, then d_{2n+l+1} must be trivial. Thus $\text{rk } H^k(X_G) = 2$ for large value of k , and by Proposition 2.2, we get $\text{rk } H^*(F) = 2$. So, $F \sim_2 \mathbb{S}^q$, and by Proposition 2.4, we get $0 < q \leq n + m + l$. Similarly, if d_{l-2n+1} is trivial, then d_{2n+l+1} must be nontrivial. Clearly, $F \sim_2 \mathbb{S}^q, 0 < q \leq n + m + l$. This realizes possibility (1).

Next, suppose that $E_2^{*,*} = E_\infty^{*,*}$. Then $\text{rk } H^*(F) = 4$. Assume that $x \in F$, and $\alpha \in H^*(X_G, x_G)$ and $\beta \in H^*(X_G, x_G)$ represent $1 \otimes ab \in E_2^{0,2n}$ and $1 \otimes c \in E_2^{0,l}$, respectively. Clearly, $\{t^i \alpha, t^i \beta, t^i \alpha \beta\}$ forms a basis for $H^k(X_G, x_G)$ over \mathbb{Z}_2 -module. Let $0 < q_1 \leq q_2 \leq q_3$ be the non vanishing dimensions of $H^*(F)$ and u_i generates q_i^{th} cohomology group of $F, 1 \leq i \leq 3$. Then we write

$$j^*(\alpha) = A_1 t^{2n-q_1} \otimes u_1 + A_2 t^{2n-q_2} \otimes u_2 + A_3 t^{2n-q_3} \otimes u_3,$$

$$j^*(\beta) = B_1 t^{l-q_1} \otimes u_1 + B_2 t^{l-q_2} \otimes u_2 + B_3 t^{l-q_3} \otimes u_3,$$

So, we get

$$j^*(\alpha\beta) = A_1B_1t^{2n-2q_1} \otimes u_1^2 + (A_1B_2 + A_2B_1)t^{2n+l-q_1-q_2} \otimes u_1u_2 + A_2B_2t^{2n+l-2q_2} \otimes u_2^2,$$

where A_i & B_i are in \mathbb{Z}_2 . Clearly, $u_1u_3 = u_2u_3 = u_3^2 = 0$. As $\alpha\beta \neq 0$, $j^*(\alpha\beta)$ is one of the following: (1) $t^{2n+l-2q_1} \otimes u_1^2$, (2) $t^{2n+l-2q_2} \otimes u_2^2$, (3) $t^{2n+l-q_1-q_2} \otimes u_1u_2$, (4) $t^{2n+l-2q_1} \otimes u_1^2 + t^{2n+l-q_1-q_2} \otimes u_1u_2$, (5) $t^{2n+l-2q_2} \otimes u_2^2 + t^{2n+l-q_1-q_2} \otimes u_1u_2$, and (6) $t^{2n+l-2q_1} \otimes u_1^2 + t^{2n+l-2q_2} \otimes u_2^2$.

It is easy to observe that the cases (5) and (6) are not possible. If $j^*(\alpha\beta) = t^{2n+l-2q_1} \otimes u_1^2 + t^{2n+l-2q_2} \otimes u_2^2$, then either $\text{rk } H^*(F) > 4$ or $j^*(\alpha\beta) = 0$, a contradiction. And, if $j^*(\alpha\beta) = t^{2n+l-2q_2} \otimes u_2^2 + t^{2n+l-q_1-q_2} \otimes u_1u_2$, then $j^*(\alpha\beta) = 0$, again a contradiction.

Now, suppose that $j^*(\alpha\beta) = t^{2n+l-2q_1} \otimes u_1^2 + t^{2n+l-q_1-q_2} \otimes u_1u_2$. Then, if $u_1u_2 = u_1^2$, then $j^*(\alpha\beta) = 0$, a contradiction. So, $u_2 = u_2^2$, and hence $F \sim_2 \mathbb{P}^3(q_1)$, $q_1 < 2n$ or $q_1 < l$, according as $2n < l$ or $l < 2n$, respectively. Note that, if $q_1 = 2n$ or $q_1 = l$, then $u_1 = a|F$, and we get $u_1^2 = 0$, a contradiction. This realizes possibility (3).

Next, suppose that $j^*(\alpha\beta) = t^{2n+l-q_1-q_2} \otimes u_1u_2$. In this case, we have four conditions (i) $u_1^2 = 0$ & $u_2^2 = 0$, (ii) $u_1^2 \neq 0$ & $u_2^2 = 0$, (iii) $u_1^2 = 0$ & $u_2^2 \neq 0$, and (iv) $u_1^2 \neq 0$ & $u_2^2 \neq 0$. Clearly, if $u_1^2 = 0$ & $u_2^2 = 0$, then $F \sim_2 \mathbb{S}^{q_1} \times \mathbb{S}^{q_2}$. Now, if $u_1^2 \neq 0$ & $u_2^2 = 0$, then for $u_2 = u_2^2$, we have $F \sim_2 \mathbb{P}^3(q)$, and for $u_1u_2 = u_1^2$, we get $q_1 = q_2$ and by the change of basis $d = u_1 + u_2$, we have $F \sim_2 \mathbb{P}^2(q) \# \mathbb{P}^2(q)$. Next, if $u_1^2 = 0$ & $u_2^2 \neq 0$, then $u_1u_2 = u_2^2$, and we get $F \sim_2 \mathbb{P}^2(q) \# \mathbb{P}^2(q)$. Finally, if $u_1^2 \neq 0$ & $u_2^2 \neq 0$, then we must have $u_1u_2 = u_2^2 = u_1^2$, and we get $F \sim_2 \mathbb{P}^2(q) \vee \mathbb{S}^q$.

Now, suppose that $j^*(\alpha\beta) = t^{2n+l-2q_1} \otimes u_1^2$, then as above we have four conditions: (i) $u_2^2 = 0$ & $u_1u_2 = 0$, (ii) $u_2^2 \neq 0$ & $u_1u_2 = 0$, (iii) $u_2^2 = 0$ & $u_1u_2 \neq 0$, and (iv) $u_2^2 \neq 0$ & $u_1u_2 \neq 0$. If $u_2^2 = 0$ & $u_1u_2 = 0$, then $F \sim_2 \mathbb{P}^2(q_1) \vee \mathbb{S}^{q_2}$ or $F \sim_2 \mathbb{P}^2(q_1) \vee \mathbb{S}^{q_3}$, accordingly $u_1^2 = u_3$ or $u_1^2 = u_2$, respectively. If $u_2^2 \neq 0$ & $u_1u_2 = 0$, then we must have $u_1^2 = u_2^2$, and hence $F \sim_2 \mathbb{P}^2(q) \# \mathbb{P}^2(q)$. If $u_2^2 = 0$ & $u_1u_2 \neq 0$, then for $u_2 = u_1^2$ we get $F \sim_2 \mathbb{P}^3(q)$, and for $u_1u_2 = u_1^2$, we get $q_1 = q_2$ and by the change of basis $d = u_1 + u_2$, we have $F \sim_2 \mathbb{P}^2(q) \# \mathbb{P}^2(q)$. If $u_2^2 \neq 0$ & $u_1u_2 \neq 0$, then we must have $u_1u_2 = u_2^2 = u_1^2$, and we get $F \sim_2 \mathbb{P}^2(q) \vee \mathbb{S}^q$. Clearly, all the inequalities hold by Propostion 2.4.

Finally, suppose that $j^*(\alpha\beta) = t^{2n+l-2q_1} \otimes u_1^2$. This case is similar to the case when $j^*(\alpha\beta) = t^{2n+l-2q_1} \otimes u_1^2$.

For the other two nontrivial actions of G on $H^n(X)$, we get the same result.

Next, if $n < m = l$ or $n = m = l$, then we get the same possibilities as in the case when $n = m < l$. \square

Remark 3.11. If $\pi_1(B_G)$ acts trivially on $H^*(X)$ and $\text{rk } H^*(F) = 2$ or 4 , then we get same possibilities of nonempty connected fixed point sets as in Theorem 3.10. Note that for the smooth manifolds in Theorem 3.10, we get possibilities (1)-(4) only.

3.2. Possibilities of disconnected fixed point sets. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l, 1 \leq n \leq m \leq l$. In this section, we determine the possibilities of a disconnected fixed point set F when X is TNHZ in X_G . We have $\sum \text{rk } H^i(F) = 8$. Clearly, F has at most eight components. By Proposition 2.6, each component F_0 of F is a poincaré duality space of formal dimension, say $r \leq n + m + l$. If $\text{rk } H^*(F_0) = 1$ or 2 or 3 , then F_0 have the mod 2 cohomology of a point or a sphere or a projective space of height two, respectively. Now, if $\text{rk } H^*(F_0) = 4$, then F_0 have the mod 2 cohomology of either a product of spheres $\mathbb{S}^{q_1} \times \mathbb{S}^{q_2}$, or a projective space of height three $\mathbb{P}^3(q)$ or a connected sum of two projective spaces of height two $\mathbb{P}^2(q) \# \mathbb{P}^2(q)$ [4]. For any component F_0 of F with $\text{rk } H^*(F_0) \leq 4$, we have following theorem:

Theorem 3.12. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, where $1 \leq n \leq m \leq l$ and X is TNHZ in X_G . If F is disconnected and the rank of cohomology ring of a component of F is at most four, then F must be one of the following:

- (1) $F \sim_2 \bigsqcup_{i=0}^3 F_i$, where $F_i \sim_2 \mathbb{S}^{r_i}, 0 \leq i \leq 3$.
- (2) $F \sim_2 \bigsqcup_{i=0}^2 F_i$, where $F_0 \sim_2 \mathbb{P}^2(r_1)$ or $\mathbb{S}^{r_2} + pt$, $F_1 \sim_2 \mathbb{P}^2(r_3)$ and $F_2 \sim_2 \mathbb{S}^{r_4}$.
- (3) $F \sim_2 \bigsqcup_{i=0}^2 F_i$, where $F_0 \sim_2 \mathbb{S}^{r_1} \times \mathbb{S}^{r_2}$ or $\mathbb{P}^3(r_1)$ or $\mathbb{P}^2(r_1) \# \mathbb{P}^2(r_2)$ and $F_i \sim_2 \mathbb{S}^{r_i}, i = 1, 2$.
- (4) $F \sim_2 \bigsqcup_{i=0}^2 F_i$, where $F_0 \sim_2 \mathbb{S}^{r_1} \times \mathbb{S}^{r_2}$ or $\mathbb{P}^3(r_1)$ or $\mathbb{P}^2(r_1) \# \mathbb{P}^2(r_1)$, $F_1 \sim_2 \mathbb{P}^2(r_2)$ and $F_2 \sim_2 pt$.
- (5) $F \sim_2 \bigsqcup_{i=0}^1 F_i$, where $F_i \sim_2 \mathbb{S}^{r_1} \times \mathbb{S}^{r_2}$ or $\mathbb{P}^3(r_1)$ or $\mathbb{P}^2(r_1) \# \mathbb{P}^2(r_1), i = 0, 1$.

Further, if $\text{rk } H^*(F_0) > 4$, where F_0 is a component of F , then we have following theorems:

Theorem 3.13. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, where $1 \leq n \leq m \leq l$ and X is TNHZ in X_G . If F is disconnected and the rank of cohomology ring of a component of F is five, then F must be one of the following:

- (1) $F \sim_2 \bigsqcup_{i=0}^1 F_i$, where $F_0 \sim_2 \#_3 \mathbb{P}^2(r_1)$ or $\mathbb{P}^2(r_1) \# (\mathbb{S}^{r_2} \times \mathbb{S}^{r_3})$ or $\mathbb{P}^4(r_1)$ and $F_1 \sim_2 \mathbb{P}^2(r_1)$.
- (2) $F \sim_2 \bigsqcup_{i=0}^2 F_i$, where $F_0 \sim_2 \#_3 \mathbb{P}^2(r_1)$ or $\mathbb{P}^2(r_1) \# (\mathbb{S}^{r_2} \times \mathbb{S}^{r_3})$ or $\mathbb{P}^4(r_1)$ and $F_1 \sim_2 \mathbb{S}^{r_4}$ or $F_2 \sim_2 pt$.

Proof. Let F_0 be the component of F such that $\text{rk } H^*(F_0) = 5$. Suppose that $0 < q_1 \leq q_2 \leq q_3 < r$ are non vanishing dimensions of $H^*(F_0)$, and u_i generates $H^{q_i}(F_0)$, $1 \leq i \leq 3$ and v generates $H^r(F_0)$. Clearly, $u_i v = 0$, $1 \leq i \leq 3$.

First, we consider that each generator u_i , $1 \leq i \leq 3$, is poincaré dual of itself. In this case, we get $u_1^2 = u_2^2 = u_3^2$ generates $H^r(F)$, and hence $F_0 \sim_2 \# \mathbb{P}^2(q_1)$.

Now, we consider that exactly one generator is poincaré dual of itself. Let u_2 be poincaré dual of itself. So, we have u_3 is the poincaré dual of u_1 . Then, $u_1 u_3 = u_2^2$ generates $H^r(F_0)$. Clearly, $u_2 u_3 = u_3^2 = 0$. Suppose that $u_1 u_2 = 0$. Further, if $u_1^2 \neq 0$, then we must have $u_3 = u_1^2$, and hence $F_0 \sim_2 \mathbb{P}^3(q_1) \# \mathbb{P}^2(q_2)$, which is not possible. If $u_1^2 = 0$, then $F_0 \sim_2 \mathbb{P}^2(q_2) \# (\mathbb{S}^{q_1} \times \mathbb{S}^{q_3})$. Now, suppose $u_1 u_2 \neq 0$. Then, we get $u_3 = u_1 u_2$. So, $u_1^2 u_2 = u_2^2$ generates $H^r(F_0)$ and we must have $u_2 = u_1^2$. Hence, $F_0 \sim_2 \mathbb{P}^4(q_1)$, where $q_1 = 1, 2, 4, 8$. For the remaining cases, when u_1 or u_3 is poincaré dual of itself, we only get $F_0 \sim_2 \mathbb{P}^2(q) \# (\mathbb{S}^q \times \mathbb{S}^q)$.

Since $\text{rk } H^*(F) = 8$ and $\text{rk } H^*(F_0) = 5$, then F have at most four components. Clearly, the components other than F_0 have the mod 2 cohomology of a point or a sphere or a projective space of height two. Hence, our claim. \square

Theorem 3.14. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, where $1 \leq n \leq m \leq l$ and X is TNHZ in X_G . If F is disconnected and one of the rank of cohomology ring of a component of F is six, then $F \sim_2 \bigsqcup_{i=1}^2 F_i$, where $F_1 \sim_2 \# \mathbb{P}^3(r_1)$ or $\mathbb{P}^3(r_1) \# (\mathbb{S}^{r_2} \times \mathbb{S}^{r_3})$ or $\mathbb{P}^2(r_1) \times \mathbb{S}^{r_2}$ or $\mathbb{P}^5(r_1)$ and $F_2 \sim_2 \mathbb{S}^{r_2}$.

Proof. Let F_0 be the component of F with $\text{rk } H^*(F_0) = 6$. Assume that $0 < q_1 \leq q_2 \leq q_3 \leq q_4 < r$ are non vanishing dimensions of $H^*(F_0)$, and u_i generates $H^{q_i}(F_0)$, $1 \leq i \leq 4$ and v generates $H^r(F_0)$. If each generator u_i is poincaré dual of itself or any two generators are poincaré dual of itself, then this contradicts Proposition 2.7. As $q_i \leq q_{i+1}$, $1 \leq i \leq 3$, we must have u_4 and u_3 are poincaré duals of u_1 and u_2 , respectively. So, $u_1 u_4 = u_2 u_3$ generates $H^r(F_0)$. Clearly, $u_2 u_4 = u_3 u_4 = 0$. So, the cup products $u_1 u_2$ and $u_1 u_3$ are either zero or nonzero.

First, Suppose that $u_1 u_2 = u_1 u_3 = 0$. Clearly, $u_3^2 = u_4^2 = 0$. Now, we have u_1^2 and u_2^2 are either zero or nonzero. If $u_1^2 = u_2^2 = 0$, then $F \sim_2 (\mathbb{S}^{q_2} \times \mathbb{S}^{q_3}) \# (\mathbb{S}^{q_1} \times \mathbb{S}^{q_4})$, which contradicts Proposition 2.7. If $u_1^2 \neq 0$ and $u_2^2 \neq 0$, then we get $u_4 = u_1^2$ and $u_3 = u_2^2$. Thus, $F \sim_2 \mathbb{P}^3(q_1) \# \mathbb{P}^3(q_2)$. If $u_1^2 \neq 0$ and $u_2^2 = 0$, then we get $F \sim_2 \mathbb{P}^3(q_1) \# (\mathbb{S}^{q_2} \times \mathbb{S}^{q_3})$. Similarly, if $u_1^2 = 0$ and $u_2^2 \neq 0$, then $F \sim_2 \mathbb{P}^3(q_1) \# (\mathbb{S}^{q_1} \times \mathbb{S}^{q_4})$.

Now, suppose that $u_1 u_2 \neq 0$ and $u_1 u_3 = 0$. Then, we have either $u_1 u_2 = u_3$ or $u_1 u_2 = u_4$. If $u_1 u_2 = u_3$, then $u_1 u_4 = u_1 u_2^2$ generates $H^r(F_0)$. So, $u_2^2 \neq 0$, and we get $u_4 = u_2^2$. Further, if $u_1^2 = 0$, then we get $F \sim_2 \mathbb{P}^2(q_1) \times \mathbb{S}^{q_2}$. For $u_1^2 \neq 0$, we get $u_1^2 = u_2^2$.

By the change of basis $d = u_1 + u_2$, we get $F \sim_2 \mathbb{P}^2(q_1) \times \mathbb{S}^{q_1}$. Similarly, if $u_1u_2 = u_4$, then $F \sim_2 \mathbb{P}^2(q_1) \times \mathbb{S}^{q_2}$.

Now, suppose that $u_1u_2 = 0$ and $u_1u_3 \neq 0$. Then, $u_1u_3 = u_4$ and $u_1^2u_3$ generates $H^r(F_0)$. Clearly, $u_3^2 = 0$ and $u_2 = u_1^2$. Thus, $F \sim_2 \mathbb{P}^2(q_1) \times \mathbb{S}^{q_3}$.

Finally, suppose that both u_1u_2 and u_1u_3 are nonzero. In this case, we get $u_3 = u_1u_2$ and $u_4 = u_1u_3$. Clearly, $u_2 = u_1^2$, and $F \sim_2 \mathbb{P}^5(q_1)$, where $q_1 = 1, 2, 4, 8$.

Note that, the component other than F_0 have sphere cohomology. Hence, our claim. \square

Theorem 3.15. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, where $1 \leq n \leq m \leq l$ and X is TNHZ in X_G . If F is disconnected and one of the rank of cohomology ring of component of F is seven, then F must be one of the following:

- (1) $F \sim_2 \bigsqcup_{i=1}^2 F_i$, where $F_1 \sim_2 (\mathbb{P}^2(r_1) \times \mathbb{S}^{r_2}) \# \mathbb{P}^2(r_3)$ or $\mathbb{P}^4(r_1) \# (\mathbb{S}^{r_2} \times \mathbb{S}^{r_3})$ or $\mathbb{P}^6(r_1)$ and $F_2 \sim_2 pt$.
- (2) $F \sim_2 \mathbb{Z}_2[u_1, u_2] / \langle u_1^5, u_2^3, u_1^3 + u_2^2, u_1^2u_2 \rangle + pt$, where $\deg u_1 = r_1$ and $\deg u_2 = r_2$.

Proof. Let F_0 be the component of F such that $\text{rk } H^*(F_0) = 7$. Clearly, the component of F other than F_0 have point cohomology. Assume that $0 < q_1 \leq q_2 \leq q_3 \leq q_4 \leq q_5 < r$ are the non vanishing dimensions of $H^*(F_0)$ with $u_i, 1 \leq i \leq 5$ is generator of $H^{q_i}(F_0)$ and v generates $H^r(F_0)$. If each generator u_i is poincaré dual of itself or any three genertors are poincaré duals of itself, then we get a contradiction of Proposition 2.7. Clearly, u_3 is poincaré dual of itself and u_5 and u_4 are poincaré duals of u_1 and u_2 , respectively. Therefore, $u_1u_5 = u_2u_4 = u_3^2$ generates $H^r(F_0)$. Note that the cup products $v^2 = u_2u_5 = u_3u_4 = u_3u_5 = u_4^2 = u_5^2 = u_iv = 0, 1 \leq i \leq 5$. The remaining cup products u_1u_2, u_1u_3, u_1u_4 , and u_2u_3 may be zero or nonzero. Now, we considered cases accordingly to how many cup products are nonzero.

It is easy to observe that, if any three cup products are nonzero or none of the cup products are nonzero then these cases are not possible. So, we consider the following cases:

Case(1): If any one cup product from $u_1u_2, u_1u_3, u_1u_4, u_2u_3$ is nonzero.

First, Suppose that $u_1u_2 \neq 0$, then either $u_1u_2 = u_4$ or $u_1u_2 = u_5$. Let $u_1u_2 = u_4$. Then, $v = u_1u_5 = u_2^2u_1 = u_3^2$. If $u_2^2 = u_3$, then $u_3^2 = u_2^4 \neq 0$ but $u_2u_3 = u_3^3 = 0$, a contradiction. By the uniqueness of poincaré dual of u_2 , we get $u_2^2 = u_5$. If $u_1^2 = 0$, then we get $F \sim_2 (\mathbb{P}^2(q_1) \times \mathbb{S}^{q_2}) \# \mathbb{P}^2(q_3)$, and $q_1, q_3 \in \{1, 2, 4, 8\}$. If $u_1^2 \neq 0$, then we must have $u_2^2 = u_1^2$, and by the change of basis $d = u_1 + u_2$, we get $F \sim_2 (\mathbb{P}^2(q_1) \times \mathbb{S}^{q_1}) \# \mathbb{P}^2(q_3)$. Similarly, if $u_1u_2 = u_5$, then $F \sim_2 (\mathbb{P}^2(q_1) \times \mathbb{S}^{q_2}) \# \mathbb{P}^2(q_3)$.

Suppose that $u_1u_3 \neq 0$, then $u_1u_3 = u_5$. So, $u_1^2u_3 = u_2u_4 = u_3^2 = v$. Clearly, $u_1^2 = u_3$ and $u_2^2 = 0$. Thus, $F \sim_2 \mathbb{P}^4(q_1) \# (\mathbb{S}^{q_2} \times \mathbb{S}^{q_2})$, $q_1 = 1, 2, 4, 8$.

Similarly, if $u_1u_4 \neq 0$, then we get $F \sim_2 (\mathbb{P}^2(q_1) \times \mathbb{S}^{q_4}) \# \mathbb{P}^2(q_3)$, and if $u_2u_3 \neq 0$, then we get $F \sim_2 \mathbb{P}^4(q_2) \# (\mathbb{S}^{q_1} \times \mathbb{S}^{q_5})$. These realizes possibility (1).

Case(2): If any two cup products from $u_1u_2, u_1u_3, u_1u_4, u_2u_3$ are nonzero.

It is easy observe that the only possible nonzero pair is u_1u_2 & u_1u_3 . We must have $u_1u_3 = u_6$ and $u_1u_2 = u_4$. Thus, $u_1^2u_3 = u_2^2u_1 = u_3^2 = v$. Clearly, $u_3 = u_1^2$ and $u_1u_3 = u_2^2$. Thus, the cohomology ring of $H^*(F_0)$ isomorphic to

$$\mathbb{Z}_2[u_1, u_2] / \langle u_1^5, u_2^3, u_1^3 + u_2^2, u_1^2u_2 \rangle,$$

where $\deg u_1 = q_1$ and $\deg u_2 = q_2$. This realizes possibility (2).

Case(3): If all cup products $u_1u_2, u_1u_3, u_1u_4, u_2u_3$ are nonzero.

In this case, we must have $u_1u_2 = u_3$, $u_1u_3 = u_4$, $u_1u_4 = u_2u_3 = u_5$ and $u_1^2 = u_2$. Hence, $F_0 \sim_2 \mathbb{P}^6(q)$, where $q = 1, 2, 4, 8$. This realizes possibility (1). \square

Next, suppose that X is not TNHZ in X_G and F is disconnected fixed point set of involutions on X . A proof of the following theorem is similar to proof of Theorem 3.10.

Theorem 3.16. Let $G = \mathbb{Z}_2$ act on $X \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, $1 \leq n \leq m \leq l$ and X is not TNHZ in X_G . If F is disconnected fixed point set and $\pi_1(B_G)$ acts nontrivially on $H^*(X)$, then F must be the one of the following:

- (1) $F \sim_2 \mathbb{S}^0$.
- (2) $F \sim_2 \mathbb{S}^{r_1} + \mathbb{S}^{r_2}$, $0 \leq r_1, r_2 \leq n + m + l$.
- (3) $F \sim_2 \mathbb{P}^2(r_1) + pt$, $r_1 = 1, 2, 4, 8$, $r_1 \leq n + m + l$.
- (4) $F \sim_2 \mathbb{S}^{r_1} \vee \mathbb{S}^{r_2} + pt$, $0 \leq r_1, r_2 \leq n + m + l$.

Note that if $\pi_1(B_G)$ acts trivially on $H^*(X)$ and $\text{rk } H^*(F) = 2$ or 4 , then we get same possibilities of disconnected fixed point sets as in Theorem 3.16. For the smooth manifolds possibilities (3) and (4) will not occur.

Next, we give some examples to realizes above Theorems:

Example 3.17. Let $G = \mathbb{Z}_2$ act on $Y \sim_2 \mathbb{S}^n \times \mathbb{S}^m$, $n \leq m$. If Y is TNHZ in Y_G , then the possibilities of mod 2 cohomology of fixed point sets are $\mathbb{S}^{r_1} \times \mathbb{S}^{r_2}$, $\mathbb{P}^2(r_1) \# \mathbb{P}^2(r_1)$, $\mathbb{P}^3(r_1)$, $\mathbb{P}^2(r_1) + pt$ and $\mathbb{S}^{r_1} + \mathbb{S}^{r_2}$, and if Y is not TNHZ in Y_G , then $F \sim_2 \mathbb{S}^{r_1}$ [4]. Recall that if G acts on $Z \sim_2 \mathbb{S}^l$, $l > 0$, then mod 2 cohomology of the fixed point set is \mathbb{S}^{r_3} , $r_3 \geq 0$. After taking the diagonal action of G on $X = Y \times Z \sim_2 \mathbb{S}^n \times \mathbb{S}^m \times \mathbb{S}^l$, we get the possibilities of mod 2 cohomology of fixed point sets are $\mathbb{S}^{r_1} \times \mathbb{S}^{r_2} \times \mathbb{S}^{r_3}$, $(\mathbb{P}^2(r_1) \# \mathbb{P}^2(r_1)) \times \mathbb{S}^{r_3}$, $\mathbb{P}^3(r_1) \times \mathbb{S}^{r_3}$, $(\mathbb{P}^2(r_1) + pt) \times \mathbb{S}^{r_3}$ and $(\mathbb{S}^{r_1} + \mathbb{S}^{r_2}) \times \mathbb{S}^{r_3}$, in the case when X is TNHZ in X_G and $F \sim_2 \mathbb{S}^{r_1} \times \mathbb{S}^{r_3}$, in the case when X is not TNHZ in X_G . This realizes possibility (2)

of Theorems 3.3, 3.4 and 3.10, possibility (1) of Theorems 3.5 and 3.6, possibilities (1) and (5) of Theorem 3.12, and Theorem 3.14.

REFERENCES

- [1] C. Allday, Torus actions on a cohomology product of three odd spheres, *Trans. Amer. Math. Soc.* **203**(1975), 343-358.
- [2] J. F. Adam, On the non-existence of elements of Hopf invariants one, *Ann. of Math.* **72**(1960), 20-104.
- [3] G. E. Bredon, Introduction to Compact Transformation Groups (New York, USA: Academic Press) (1972).
- [4] G. E. Bredon, Cohomological aspects of transformation groups, Proceedings of the Conference on Transformation Groups (New Orleans, 1967), Springer-Verlag, New York, 1968, 245-280.
- [5] G. E. Bredon, The cohomology ring structure of a fixed point set. *Ann. of Math.* **80**(1964), 524-537.
- [6] T. Chang and M. Comenetz, Group actions on cohomology projective planes and products of spheres. *Q. J. Math.* **28** (1977).
- [7] G. David, Torus actions on a product of two odd spheres. *Topology* **10** (1971), 313-326.
- [8] R. G. Swan, A new method in fixed point theory, *Comment. Math. Helv.* **34** (1960), 1-16.
- [9] Dimpi and H. K. Singh, Orbit spaces of free involutions on the product of three spheres, arXiv:2303.16474v1,(2023).
- [10] J. W. Milnor and J. D. Stasheff, Characteristic classes, Princeton University Press and University of Tokyo Press, Princeton, New Jersey, (1974).
- [11] P. Volker, On a conjecture of Bredon, *Manuscripta mathematica*, **12** (1974), 11-16.
- [12] J. C. Su, Periodic transformations on the product of two spheres, *Trans. Amer. Math. Soc.* **112** (1964), 369-380.

DIMPI
 DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF DELHI
 DELHI – 110007, INDIA.
Email address: dimpipaul2@gmail.com

HEMANT KUMAR SINGH
 DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF DELHI
 DELHI – 110007, INDIA.
Email address: hemantksingh@maths.du.ac.in