

COMMUTING KRAUS OPERATORS ARE NORMAL

MARTIN FRAAS

Department of Mathematics, University of California, Davis, CA 95616, USA

ABSTRACT. Let $\{V_1, \dots, V_n\}$ be a set of mutually commuting matrices. We show that if $V_1^*V_1 + \dots + V_n^*V_n = \text{Id}$ then the matrices are normal and, in particular, simultaneously diagonalizable.

1. RESULT

Let \mathcal{H} be a finite dimensional complex Hilbert space. A matrix A is called normal if it commutes with its Hermitian conjugate matrix A^* . A set of mutually commuting normal matrices is simultaneously diagonalizable, and there exists an orthonormal basis whose elements are joint eigenvectors of all the matrices. The condition that the matrices are normal is necessary, if two matrices A, B commute, $[A, B] = 0$, then without further assumptions they only need to have one common eigenvector [10]. We provide a criteria that implies this condition.

Theorem 1. *Let $\{V_1, \dots, V_n\}$ be a set of commuting matrices such that $\sum_{\alpha=1}^n V_\alpha^*V_\alpha = \text{Id}$. Then the matrices are normal and, in particular, simultaneously diagonalizable.*

The motivation for the theorem comes from the theory of quantum trajectories. In this context, a set of matrices satisfying the normalization condition

$$(1) \quad \sum_{\alpha=1}^n V_\alpha^*V_\alpha = \text{Id}$$

are called Kraus operators or jump operators. The operators describe evolution of an initial state $\psi \in \mathcal{H}$ conditioned on outcomes $\alpha_1, \dots, \alpha_n$ of a repeated measurement, and the probability of obtaining these outcomes. The probability is given by

$$(2) \quad \mathbb{P}(\alpha_1, \dots, \alpha_n) := \|V_{\alpha_n} \dots V_{\alpha_1} \psi\|^2,$$

which is a probability measure provided that $\|\psi\| = 1$. Following the experiments of the quantum optics group of S. Haroche [9], the case of non-demolition measurements attracted a lot of attention in physics and math literature [5, 6, 4, 3]. A standard definition of a non-demolition measurement postulates which observable N is not being demolished.

Condition 1. *A set of jump operators V_α is called non-demolition if there exists a Hermitian operator N and complex functions f_α such that $V_\alpha = f_\alpha(N)$.*

E-mail address: `mfraas@ucdavis.edu`.

Date: August 11, 2023.

In particular, in the non-demolition case the jump operators commute, and the probability measure (2) is exchangeable. This suggests a natural intrinsic definition of the non-demolition case.

Condition 2. *A set of jump operators V_α is called non-demolition if the operators V_α are mutually commuting.*

An immediate corollary of Theorem 1 is that these two definitions give the same class of Kraus operators.

Corollary 2. *Conditions 1 and 2 are equivalent.*

Proof. As discussed above Condition 1 implies Condition 2. In the opposite direction, if Kraus operators V_α are mutually commuting then by Theorem 1 they are normal and hence there exists a Hermitian matrix N such that all matrices V_α are functions of N . \square

2. COMPLETELY POSITIVE MAPS

The proof of the theorem that we give below involves some results about completely positive maps. We recall these results. A completely positive map $\Phi : B(\mathcal{H}) \rightarrow B(\mathcal{H})$ has a form

$$\Phi(X) = \sum_{\alpha} V_{\alpha}^{*} X V_{\alpha},$$

for some finite set of operators V_{α} (that do not necessarily commute). We will also use the dual completely positive map

$$\Phi^{*}(A) = \sum_{\alpha} V_{\alpha} A V_{\alpha}^{*}.$$

The first fact, see [11], about completely positive maps that we will need is that

$$(3) \quad \|\Phi\| = \left\| \sum_{\alpha} V_{\alpha}^{*} V_{\alpha} \right\|,$$

where both norms are the operator norms. If Φ satisfies the normalization condition Eq. (1) then $\|\Phi\| = 1$ belongs to the spectrum of Φ . Furthermore, there exists an positive eigenmatrix ρ such that $\Phi^{*}(\rho) = \rho$. We will call such eigenmatrix a stationary state.

We now assume the normalization condition Eq. (1). The second fact, see [8] for items (i), (ii) and [1, Proposition 3] for (iii), that we will use is that there exists a decomposition of the Hilbert space $\mathcal{H} = \mathcal{H}_F \oplus \mathcal{H}_D$ such that with respect to this decomposition the matrices have a form

$$(4) \quad V_{\alpha} = \begin{pmatrix} A_{\alpha} & B_{\alpha} \\ 0 & C_{\alpha} \end{pmatrix},$$

and

(i) Φ^{*} restricted to $B(\mathcal{H}_F)$,

$$\Phi_F^{*}(X) = \sum_{\alpha} A_{\alpha} X A_{\alpha}^{*},$$

possesses a faithful (full rank) stationary state,

(ii) Φ restricted to $B(\mathcal{H}_D)$,

$$\Phi_D(X) = \sum_{\alpha} C_{\alpha}^{*} X C_{\alpha},$$

has a spectral radius strictly less than 1,

(iii) Any solution of $\Phi(X) = X$ is diagonal with respect to the decomposition, i.e. it has a form

$$X = \begin{pmatrix} X_F & 0 \\ 0 & X_D \end{pmatrix}.$$

If the decomposition is trivial, $\mathcal{H}_D = 0$, we define $B_\alpha = C_\alpha = 0$. The decomposition is trivial if and only if $B_\alpha \equiv 0$.

We will not need the full claim (iii). We will use a claim that any solution of $\Phi(X) = X$ that acts as zero on \mathcal{H}_F is a zero matrix. This also follows from (ii). By (ii), subspace \mathcal{H}_D is transient, i.e. for any matrix A ,

$$\lim_{n \rightarrow \infty} P_D \Phi^{*n}(A) P_D = 0,$$

where P_D is the projection on \mathcal{H}_D . By positivity of Φ , also $P_D \Phi^{*n}(A)$ and $\Phi^{*n}(A) P_D$ go to zero. Since $\text{tr}(XA) = \text{tr}(X\Phi^{*n}(A))$ this implies that $\text{tr}(XA) = 0$ so X is indeed zero.

3. PROOF OF THE THEOREM

We use the decomposition Eq. (4). The normalization Eq. (1) is equivalent to

$$(5) \quad \sum_{\alpha} A_{\alpha}^* A_{\alpha} = \text{Id},$$

$$(6) \quad \sum_{\alpha} A_{\alpha}^* B_{\alpha} = 0,$$

$$(7) \quad \sum_{\alpha} B_{\alpha}^* B_{\alpha} + C_{\alpha}^* C_{\alpha} = \text{Id}.$$

The commutation assumption is equivalent to equations

$$(8) \quad A_{\alpha} A_{\beta} = A_{\beta} A_{\alpha},$$

$$(9) \quad A_{\alpha} B_{\beta} + B_{\alpha} C_{\beta} = A_{\beta} B_{\alpha} + B_{\beta} C_{\alpha},$$

$$(10) \quad C_{\alpha} C_{\beta} = C_{\beta} C_{\alpha},$$

holding for all α, β . Eqs. (5), (8) imply that the matrices A_1, \dots, A_n are mutually commuting and normalized as in Eq. (1). By the property of the decomposition, the corresponding map Φ_F^* has a faithful (full rank) stationary solution ρ , i.e. a matrix such that $\Phi_F^*(\rho) = \rho$. Let Φ_F be the dual map,

$$\Phi_F(X) = \sum_{\alpha} A_{\alpha}^* X A_{\alpha}.$$

For any β , we then have

$$\Phi_F(A_{\beta} A_{\beta}^*) - A_{\beta} \Phi_F(A_{\beta}^*) - \Phi_F(A_{\beta}) A_{\beta}^* + A_{\beta} A_{\beta}^* = \sum_{\alpha} |[A_{\beta}^*, A_{\alpha}]|^2.$$

Since the matrices commute we have that $\Phi_F(A_{\beta}) = A_{\beta}$ and $\Phi_F(A_{\beta}^*) = A_{\beta}^*$. Hence we obtain

$$\sum_{\alpha} \text{tr}(\rho |[A_{\beta}^*, A_{\alpha}]|^2) = 0,$$

and it follows that the matrices A_{α} are normal.

We now show that $B_\alpha = 0$ which implies that $\mathcal{H}_D = 0$ and finishes the proof. Multiplying Eq. (9) from the left by A_α^* and summing over α we get

$$B_\beta = \sum_{\alpha} A_\alpha^* B_\beta C_\alpha,$$

where we used that $[A_\beta^*, A_\alpha] = 0$ and Eqs. (6), (5). The equation implies that

$$\Phi \begin{pmatrix} 0 & B_\beta \\ 0 & Y \end{pmatrix} = \begin{pmatrix} 0 & B_\beta \\ 0 & \Phi_D(Y) + \sum_{\alpha} B_\alpha^* B_\beta B_\alpha \end{pmatrix}.$$

Since Φ_D has spectral radius less than 1, the sum

$$X_D := \sum_{n=0}^{\infty} \Phi_D^n \left(\sum_{\alpha} B_\alpha^* B_\beta B_\alpha \right)$$

is convergent. Hence the matrix

$$X = \begin{pmatrix} 0 & B_\beta \\ 0 & X_D \end{pmatrix}$$

satisfies the equation $\Phi X = X$. Any solution of the equation has to be diagonal so we got $B_\beta = 0$ as announced. This finishes the proof.

Remark 3. *A related result is given in [3]. Assuming that V_α 's are commuting and that the decomposition Eq. (4) is trivial, [3] shows that the quantum trajectory purifies on the joint spectral decomposition of $A_\alpha^* A_\alpha$. For the notion of purification on the spectrum we refer the reader to the aforementioned article, and only note that this, in particular, means that the spectral measure of ψ associated to the joint spectral decomposition gives the de Finetti decomposition of the measure Eq. (2). For an example, see Example 4.*

4. DISCUSSION

The assumption that the Hilbert space is finite dimensional cannot be relaxed. Any partial isometry gives a counterexample. For example, the right shift R on $L^2(\mathbb{N})$,

$$R(x_1, x_2, x_3, \dots) = (0, x_1, x_2, x_3, \dots),$$

satisfies the normalization condition $R^* R = \text{Id}$ (and obviously $[R, R] = 0$) but R is not normal. The reason for the failure of the theorem is that the decomposition Eq. (4) is more complicated in infinite dimensions, see [7], there are invariant subspaces that do not possess any stationary state.

The theory of non-demolition quantum trajectories in infinite dimensional Hilbert spaces assuming the definition given by Condition 1 was developed in [2]. It would be interesting to understand the behavior of measure Eq. (2) under Condition 2. We illustrate the difference on two examples.

Example 4. *Consider the space $L^2(\mathbb{Z})$ and jump operators*

$$V_1 = \frac{1}{2}(1 + R), \quad V_2 = \frac{1}{2}(1 - R),$$

where R is the right shift operator on \mathbb{Z} . Then V_1, V_2 are mutually commuting and normal. Let $\psi \in L^2(\mathbb{Z})$ and let $\hat{\psi} \in L^2[0, 2\pi]$ be its Fourier transform, i.e.

$$\psi_n = \frac{1}{\sqrt{2\pi}} \int_0^{2\pi} e^{ink} \hat{\psi}(k) dk.$$

Then for the measure Eq. (2) we get

$$\begin{aligned} \mathbb{P}(\alpha_1, \dots, \alpha_n) &= \int_0^{2\pi} \left| \frac{1 + e^{-ik}}{2} \right|^{2n_1} \left| \frac{1 - e^{-ik}}{2} \right|^{2n_2} |\hat{\psi}(k)|^2 dk \\ &= 2^{-n} \int_0^{2\pi} (1 + \cos k)^{n_1} (1 - \cos k)^{n_2} |\hat{\psi}(k)|^2 dk \end{aligned}$$

where n_1, n_2 are the number of ones resp. twos in the sequence $\alpha_1, \dots, \alpha_n$. The i.i.d. measures under the integral are the same for k and $\pi - k$ so that de Finetti decomposition of the measure is

$$\mathbb{P}(\alpha_1, \dots, \alpha_n) = 2^{-n} \int_{\pi/2}^{3\pi/2} (1 + \cos k)^{n_1} (1 - \cos k)^{n_2} (|\hat{\psi}(k)|^2 + |\hat{\psi}(\pi - k)|^2) dk.$$

The measure $(|\hat{\psi}(k)|^2 + |\hat{\psi}(\pi - k)|^2) dk$ in the decomposition is the spectral measure of ψ with respect to the discrete Laplacian $R + R^*$. This demonstrates Remark 3, it is a general feature that assuming Condition 1, the measure in the de Finetti decomposition is the spectral measure associated with N , see [2].

This example should be contrasted with

Example 5. Consider the space $L^2(\mathbb{N})$ and jump operators

$$V_1 = \frac{1}{2}(1 + R), \quad V_2 = \frac{1}{2}(1 - R),$$

where R is the right shift operator on \mathbb{N} . Then V_1, V_2 are mutually commuting but not normal. The action of V_α on any ψ is the same as if V_α was defined on the whole \mathbb{Z} and ψ was extended by zero on the negative numbers. Hence de Finetti decomposition given in the previous example holds true also here. However, the measure is not anymore associated to the spectral measure of $R + R^*$ on $L^2(\mathbb{N})$.

ACKNOWLEDGEMENTS

The author thanks Tristan Benoist for comments on the draft version of the article. The research has been supported by ANR project QTraj (ANR-20-CE40-0024-01) of the French National Research Agency (ANR).

REFERENCES

- [1] V. V. Albert. Asymptotics of quantum channels: conserved quantities, an adiabatic limit, and matrix product states. *Quantum*, 3:151, June 2019. 2
- [2] M. Ballesteros, N. Crawford, M. Fraas, J. Fröhlich, and B. Schubnel. Non-demolition measurements of observables with general spectra. *Mathematical Problems in Quantum Physics, Contemp. Math*, 717:241–256, 2018. 4, 5
- [3] M. Ballesteros, M. Fraas, J. Fröhlich, and B. Schubnel. Indirect acquisition of information in quantum mechanics. *Journal of Statistical Physics*, pages 1–35, 2015. 1, 4
- [4] M. Bauer, T. Benoist, and D. Bernard. Repeated quantum non-demolition measurements: convergence and continuous time limit. In *Ann. H. Poincaré*, volume 14, pages 639–679. Springer, 2013. 1
- [5] M. Bauer and D. Bernard. Convergence of repeated quantum nondemolition measurements and wavefunction collapse. *Phys. Rev. A*, 84(4):044103, 2011. 1
- [6] M. Bauer, D. Bernard, and T. Benoist. Iterated stochastic measurements. *J. Phys. A Math Th*, 45(49):494020, 2012. 1
- [7] R. Carbone and Y. Pautrat. Irreducible decompositions and stationary states of quantum channels. *Rep. Math. Phys.*, 77(3):293–313, 2016. 4

- [8] G. I. Cirillo and F. Ticozzi. Decompositions of hilbert spaces, stability analysis and convergence probabilities for discrete-time quantum dynamical semigroups. *Journal of Physics A: Mathematical and Theoretical*, 48(8):085302, 2015. 2
- [9] C. Guerlin, J. Bernu, S. Deleglise, C. Sayrin, S. Gleyzes, S. Kuhr, M. Brune, J.-M. Raimond, and S. Haroche. Progressive field-state collapse and quantum non-demolition photon counting. *Nature*, 448(7156):889–893, 2007. 1
- [10] Wikipedia contributors. Commuting matrices — Wikipedia, the free encyclopedia, 2023. [Online; accessed 1-August-2023]. 1
- [11] M. M. Wolf. Quantum channels & operations: Guided tour. <http://www-m5.ma.tum.de/foswiki/pub/M5/Allgemeines/MichaelWolf/QChannelLecture.pdf>, 2012. Lecture notes based on a course given at the Niels-Bohr Institute. 2