

# Lossy Kernelization for (Implicit) Hitting Set Problems

Fedor V. Fomin ✉

University of Bergen, Bergen, Norway.

Tien-Nam Le ✉

École Normale Supérieure de Lyon, Lyon, France.

Daniel Lokshtanov ✉

University of California Santa Barbara, USA.

Saket Saurabh ✉

The Institute of Mathematical Sciences, HBNI, Chennai, India, and University of Bergen, Bergen, Norway.

Stéphan Thomassé ✉

École Normale Supérieure de Lyon, Lyon, France.

Meirav Zehavi ✉

Ben-Gurion University of the Negev, Beersheba, Israel.

---

## Abstract

We re-visit the complexity of polynomial time pre-processing (kernelization) for the  $d$ -HITTING SET problem. This is one of the most classic problems in Parameterized Complexity by itself, and, furthermore, it encompasses several other of the most well-studied problems in this field, such as VERTEX COVER, FEEDBACK VERTEX SET IN TOURNAMENTS (FVST) and CLUSTER VERTEX DELETION (CVD). In fact,  $d$ -HITTING SET encompasses any deletion problem to a hereditary property that can be characterized by a finite set of forbidden induced subgraphs. With respect to bit size, the kernelization complexity of  $d$ -HITTING SET is essentially settled: there exists a kernel with  $\mathcal{O}(k^d)$  bits ( $\mathcal{O}(k^d)$  sets and  $\mathcal{O}(k^{d-1})$  elements) and this is tight by the result of Dell and van Melkebeek [STOC 2010, JACM 2014]. Still, the question of whether there exists a kernel for  $d$ -HITTING SET with *fewer elements* has remained one of the most major open problems in Kernelization.

In this paper, we first show that if we allow the kernelization to be *lossy* with a qualitatively better loss than the best possible approximation ratio of polynomial time approximation algorithms, then one can obtain kernels where the number of elements is linear for every fixed  $d$ . Further, based on this, we present our main result: we show that there exist approximate Turing kernelizations for  $d$ -HITTING SET that even beat the established bit-size lower bounds for exact kernelizations—in fact, we use a *constant* number of oracle calls, each with “near linear” ( $\mathcal{O}(k^{1+\epsilon})$ ) bit size, that is, almost the best one could hope for. Lastly, for two special cases of implicit 3-HITTING SET, namely, FVST and CVD, we obtain the “best of both worlds” type of results— $(1 + \epsilon)$ -approximate kernelizations with a linear number of vertices. In terms of size, this substantially improves the exact kernels of Fomin et al. [SODA 2018, TALG 2019], with simpler arguments.

**2012 ACM Subject Classification** Theory of computation → Parameterized complexity and exact algorithms

**Keywords and phrases** Hitting Set, Lossy Kernelization

**Digital Object Identifier** 10.4230/LIPIcs.CVIT.2016.23

**Funding** *Fedor V. Fomin*: Research Council of Norway via the project BWCA (grant no. 314528).

*Daniel Lokshtanov*: Supported by NSF award CCF-2008838.

*Saket Saurabh*: European Research Council (ERC) grant agreement no. 819416, and Swarnajayanti Fellowship no. DST/SJF/MSA01/2017-18.

*Stéphan Thomassé*: ANR projects TWIN-WIDTH (CE48-0014-01) and DIGRAPHS (CE48-0013-01).

*Meirav Zehavi*: European Research Council (ERC) grant titled PARAPATH.



© Fedor V. Fomin, Tien-Nam Le, Daniel Lokshtanov, Saket Saurabh, Stéphan Thomassé and Meirav Zehavi;

licensed under Creative Commons License CC-BY 4.0

42nd Conference on Very Important Topics (CVIT 2016).

Editors: John Q. Open and Joan R. Access; Article No. 23; pp. 23:1–23:42



Leibniz International Proceedings in Informatics

LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

## 1 Introduction

In  $d$ -HITTING SET, the input consists of a universe  $U$ , a family  $\mathcal{F}$  of sets over  $U$ , where each set in  $\mathcal{F}$  is of size at most  $d$ , and an integer  $k$ . The task is to determine whether there exists a set  $S \subseteq U$ , called a *hitting set*, of size at most  $k$  that has a nonempty intersection with every set of  $\mathcal{F}$ . The  $d$ -HITTING SET problem is a classical optimization problem whose computational complexity has been studied for decades from the perspectives of different algorithmic paradigms. Notably,  $d$ -HITTING SET is a generic problem, and hence, in particular, various computational problems can be re-cast in terms of it. Of course, VERTEX COVER, the most well-studied problem in Parameterized Complexity, is the special case of  $d$ -HITTING SET with  $d = 2$ . More generally,  $d$ -HITTING SET encompasses a variety of (di)graph modification problems, where the task is to delete at most  $k$  vertices (or edges) from a graph such that the resulting graph does not contain an induced subgraph (or a subgraph) from a family of forbidden graphs  $\mathcal{F}$ . Examples of some such well-studied problems include CLUSTER VERTEX DELETION,  $d$ -PATH VERTEX COVER,  $d$ -COMPONENT ORDER CONNECTIVITY,  $d$ -BOUNDED-DEGREE VERTEX DELETION, SPLIT VERTEX DELETION and FEEDBACK VERTEX SET IN TOURNAMENTS.

Kernelization, a subfield of Parameterized Complexity, provides a mathematical framework to capture the performance of polynomial time preprocessing. It makes it possible to quantify the degree to which polynomial time algorithms succeed at reducing input instances of NP-hard problems. More formally, every instance of a parameterized problem  $\Pi$  is associated with an integer  $k$ , which is called the *parameter*, and  $\Pi$  is said to admit a *kernel* if there is a polynomial-time algorithm, called a *kernelization algorithm*, that reduces the input instance of  $\Pi$  down to an equivalent instance of  $\Pi$  whose size is bounded by a function  $f(k)$  of  $k$ . (Here, two instances are equivalent if both of them are either Yes-instances or No-instances.) Such an algorithm is called an  $f(k)$ -*kernel* for  $\Pi$ . If  $f(k)$  is a polynomial function of  $k$ , then we say that the kernel is a *polynomial kernel*. Over the last decade, Kernelization has become a central and active field of study, which stands at the forefront of Parameterized Complexity, especially with the development of complexity-theoretic lower bound tools for kernelization. These tools can be used to show that a polynomial kernel [3, 12, 18, 23], or a kernel of a specific size [9, 10, 21] for concrete problems would imply an unlikely complexity-theoretic collapse. We refer to the recent book on kernelization [17] for a detailed treatment of the area of kernelization. In this paper, we provide a number of positive results on the kernelization complexity of  $d$ -HITTING SET, as well as on several special cases of 3-HITTING SET.

The most well-known example of a polynomial kernel, which, to the best of our knowledge, is taught in the first class/chapter on kernelization of any course/book that considers this subject, is the classic kernel for VERTEX COVER (2-HITTING SET) that is based on Buss rule. More generally, one of the most well-known examples of a polynomial kernel is a kernel with  $\mathcal{O}(k^d)$  sets and elements for  $d$ -HITTING SET (when  $d$  is a fixed constant) using the Erdős-Rado Sunflower lemma.<sup>1</sup> Complementing this positive result, originally in 2010, a celebrated result by Dell and van Melkebeek [10] showed that unless  $\text{co-NP} \subseteq \text{NP/poly}$ , for any  $d \geq 2$  and any  $\epsilon > 0$ ,  $d$ -HITTING SET does not admit a kernel with  $\mathcal{O}(k^{d-\epsilon})$  sets. Hence, the kernel with  $\mathcal{O}(k^d)$  sets is essentially tight with respect to size. However, when it comes to the bound on the number of elements in a kernel, the situation is unclear. Abu-Khazam [1]

<sup>1</sup> The origins of this result are unclear. The first kernel with  $\mathcal{O}(k^d)$  sets appeared in 2004 [13], but the authors do not make use of the Sunflower Lemma. To the best of our knowledge, the first exposition of the kernel based on the Sunflower Lemma appears in the book of Flum and Grohe [15].

showed that  $d$ -HITTING SET admits a kernel with at most  $(2d-1)k^{d-1}+k$  elements. However, we do not know whether this bound is tight or even close to that. As it was written in [17, page 470]:

*Could it be that  $d$ -HITTING SET admits a kernel with a polynomial in  $k$  number of elements, where the degree of the polynomial does not depend on  $d$ ? This does not look like a plausible conjecture, but we do not know how to refute it either.*

The origins of this question can be traced back to the open problems from WorKer 2010 [4, page 4]. Moreover, in the list of open problems from WorKer 2013 and FPT School 2014 [7, page 4], the authors asked whether  $d$ -HITTING SET admits a kernel with  $f(d) \cdot k$  elements for some function  $f$  of  $d$  only. After being explicitly stated at these venues, this question and its variants have been re-stated in a considerable number of papers (see, e.g., [11, 17, 30, 2]), and is being repeatedly asked in annual meetings centered around parameterized complexity. Arguably, this question has become the most major and longstanding open problem in kernelization for a specific problem. In spite of many attempts, even for  $d = 3$ , the question whether  $d$ -HITTING SET admits a kernel with  $\mathcal{O}(k^{2-\epsilon})$  elements, for some  $\epsilon > 0$ , has still remained open.

From an approximation perspective, the optimization version of  $d$ -HITTING SET admits a trivial  $d$ -approximation. Up to the Unique Game Conjecture, this bound is tight—for any  $\epsilon > 0$ ,  $d$ -HITTING SET does not admit a polynomial time  $(d - \epsilon)$ -approximation [22]. So, at this front, the problem is essentially resolved.

With respect to kernelization, firstly, the barrier in terms of number of sets, and secondly, the lack of progress in terms of the number of elements, coupled with the likely impossibility of  $(d - \epsilon)$ -approximation of  $d$ -HITTING SET, bring lossy kernelization as a natural tool for further exploring of the complexity of this fundamental problem. We postpone the formal definition of lossy kernelization to Section 2. Informally, a polynomial size  $\alpha$ -approximate kernel consists of two polynomial-time procedures. The first is a pre-processing algorithm that takes as input an instance  $(I, k)$  to a parameterized problem, and outputs another instance  $(I', k')$  to the same problem, such that  $|I'| + k' \leq k^{\mathcal{O}(1)}$ . The second transforms, for every  $c \geq 1$ , a  $c$ -approximate solution  $S'$  to the pre-processed instance  $(I', k')$  into a  $(c \cdot \alpha)$ -approximate solution  $S$  to the original instance  $(I, k)$ . Then, the main question(s) that we address in this paper is:

Is it possible to obtain a lossy kernel for  $d$ -HITTING SET with a qualitatively better loss than  $d$  and with  $\mathcal{O}(k^{d-1-\epsilon})$  bit-size, or at least with  $\mathcal{O}(k^{d-1-\epsilon})$  elements?

In this paper, we present a surprising answer: *not only the number of elements can be bounded by  $\mathcal{O}(k)$  (rather than just  $\mathcal{O}(k^{d-1-\epsilon})$ ), but even the bit-size can “almost” be bounded by  $\mathcal{O}(k)$ !* From the perspective of the size of the kernel, this is essentially the best that one could have hoped for. Still, we only slightly (though non-negligibly) improve on the approximation ratio  $d$ . For example, for  $d = 2$  (VERTEX COVER), we attain an approximation ratio of 1.721. So, while we make a critical step that is also the first—in particular, we show that, conceptually, the combination of kernelization and approximation breaks their independent barriers—we also open up the door for further research of this kind, on this problem as well as other problems.

More precisely, we present the following results and concept. We remark that for all of our results, we use an interesting fact about the natural Linear Programming (LP) relaxation

of  $d$ -HITTING SET: the support of any optimal LP solution to the LP-relaxation of  $d$ -HITTING SET is of size at most  $d \cdot \text{frac}$  where  $\text{frac}$  is the optimum (minimum value) of the LP [20]. Furthermore, to reduce bit-size rather than only element number, we introduce an “adaptive sampling strategy” that is, to the best of our knowledge, also novel in parameterized complexity. We believe that these ideas will find further applications in kernelization in the future. More information on our methods can be found in the next section.

- **Starting Point: Linear-Element Lossy Kernel for  $d$ -HITTING SET.** First, we show that  $d$ -HITTING SET admits a  $(d - \frac{d-1}{d})$ -approximate  $d \cdot \text{opt}$ -element kernel, where  $\text{opt} \leq k$  is the (unknown) optimum (that is, size of smallest solution).<sup>2</sup> For example, when  $d = 3$ , the approximation ratio is  $d - \frac{d-1}{d} = 2\frac{1}{3}$ , which is a notable improvement over 3. When  $d = 2$ , this result encompasses the classic (exact)  $2 \cdot \text{opt}$ -vertex kernel for VERTEX COVER [6, 28]. We also remark that our linear-element lossy kernel for  $d$ -HITTING SET is a critical component (used as a black box) in all of our other results.
- **Conceptual Contribution: Lossy Kernelization Protocols.** We extend the notions of lossy kernelization and kernelization protocols<sup>3</sup> to *lossy kernelization protocols*. Roughly speaking, an  $\alpha$ -approximate kernelization protocol can perform a bounded in  $k$  number of calls (called *rounds*) to an oracle that solves the problem on instances of size (called *call size*) bounded in  $k$ , and besides that it runs in polynomial time. Ideally, the number of calls is bounded by a fixed constant, in which case the protocol is called *pure*. Then, if the oracle outputs  $c$ -approximate solutions to the instances it is given, the protocol should output a  $(c \cdot \alpha)$ -approximate solution to the input instance. In particular, a lossy kernel is the special case of a lossy protocol with one oracle call. The *volume* of a lossy kernelization protocol is the sum of the sizes of the calls it performs.
- **Main Contribution: Near-Linear Volume and Pure Lossy Kernelization Protocol for  $d$ -HITTING SET.** We remark that the work of Dell and van Melkebeek [10] further asserts that also the existence of an exact (i.e., 1 approximate in our terms) kernelization protocol for  $d$ -HITTING SET of volume  $\mathcal{O}(k^{d-\epsilon})$  is impossible unless  $\text{co-NP} \subseteq \text{NP/poly}$ . First, we show that VERTEX COVER admits a (randomized) 1.721-approximate kernelization protocol of 2 rounds and call size  $\mathcal{O}(k^{1.5})$ . This special case is of major interest in itself: VERTEX COVER is the most well-studied problem in Parameterized Complexity, and, until now, no result that breaks both bit-size and approximation ratio barriers simultaneously has been known.

Then, we build upon the ideas exemplified for the case of VERTEX COVER to significantly generalize the result: while VERTEX COVER corresponds to  $d = 2$ , we are able to capture *all* choices of  $d$ . Thereby, we prove our main result: for any  $\epsilon > 0$ ,  $d$ -HITTING SET admits a (randomized) pure  $(d - \delta)$ -approximate kernelization protocol of call size  $\mathcal{O}(k^{1+\epsilon})$ . Here, the number of rounds and  $\delta$  are fixed constants that depend only on  $d$  and  $\epsilon$ . While the improvement over the barrier of  $d$  in terms of approximation is minor (though still notable when  $d = 2$ ), it is a *proof of concept*—that is, it asserts that  $d$  is not an impassable barrier.<sup>4</sup> Moreover, it does so with almost the best possible (being almost linear) output size.

<sup>2</sup> In fact, when the parameter is  $k$ , we show that the bound is better.

<sup>3</sup> We remark that kernelization protocols are a highly restricted special case of Turing kernels, that yet generalizes kernels.

<sup>4</sup> Possibly, building upon our work, further improvements on the approximation factor (though perhaps at the cost of an increase in the output size) may follow.

- **Outlook: Relation to Ruzsa-Szemerédi Graphs.** Lastly, we present a connection between the possible existence of a  $(1 + \epsilon)$ -approximate kernelization protocol for VERTEX COVER of call size  $\mathcal{O}(k^{1.5})$  and volume  $\mathcal{O}(k^{1.5+o(1)})$  and a known open problem about Ruzsa-Szemerédi graphs (defined in Section 4). We discuss this result in more detail in Section 3.

**Kernels for Implicit 3-Hitting Set Problems.** Lastly, we provide better lossy kernels for two well-studied graph problems, namely, CLUSTER VERTEX DELETION and FEEDBACK VERTEX SET IN TOURNAMENTS, which are known to be implicit 3-HITTING SET problems [8]. Notably, both our algorithms are based on some of the ideas and concepts that are part of our previous results, and, furthermore, we believe that the approach underlying the parts common to both these algorithms may be useful when dealing also with other hitting and packing problems of constant-sized objects. In the CLUSTER VERTEX DELETION problem, we are given a graph  $G$  and an integer  $k$ . The task is to decide whether there exists a set  $S$  of at most  $k$  vertices of  $G$  such that  $G - S$  is a cluster graph. Here, a cluster graph is a graph where every connected component is a clique. It is known that this problem can be formulated as a 3-HITTING SET problem where the family  $\mathcal{F}$  contains the vertex sets of all induced  $P_3$ 's of  $G$ . (An induced  $P_3$  is a path on three vertices where the first and last vertices are non-adjacent in  $G$ .) In the FEEDBACK VERTEX SET IN TOURNAMENTS problem, we are given a tournament  $G$  and an integer  $k$ . The task is to decide whether there is a set  $S$  of  $k$  vertices such that each directed cycle of  $G$  contains a member of  $S$  (i.e.,  $G - S$  is acyclic). It is known that FEEDBACK VERTEX SET IN TOURNAMENTS can be formulated as a 3-HITTING SET problem as well, where the family  $\mathcal{F}$  contains the vertex sets of all directed cycles on three vertices (triangles) of  $G$ .

In [16], it was shown that FEEDBACK VERTEX SET IN TOURNAMENTS and CLUSTER VERTEX DELETION admit kernels with  $\mathcal{O}(k^{\frac{3}{2}})$  vertices and  $\mathcal{O}(k^{\frac{5}{3}})$  vertices, respectively. This answered an open question from WorKer 2010 [4, page 4], regarding the existence of kernels with  $\mathcal{O}(k^{2-\epsilon})$  vertices for these problems. The question of the existence of linear-vertex kernels for these problems is open. In the realm of approximation algorithms, for FEEDBACK VERTEX SET IN TOURNAMENTS, Cai, Deng and Zang [5] gave a factor 2.5 approximation algorithm, which was later improved to  $7/3$  by Mnich, Williams and Végé [27]. Recently, Lokshtanov, Misra, Mukherjee, Panolan, Philip and Saurabh [24] gave a 2-approximation algorithm for FEEDBACK VERTEX SET IN TOURNAMENTS. For CLUSTER VERTEX DELETION, You, Wang and Cao [30] gave a factor 2.5 approximation algorithm, which later was improved to  $7/3$  by Fiorini, Joret and Schaudt [14]. It is open whether CLUSTER VERTEX DELETION admits a 2-approximation algorithm. We remark that both problems admit approximation-preserving reductions from VERTEX COVER, and hence they too do not admit  $(2 - \epsilon)$ -approximation algorithms up to the Unique Games Conjecture.

We provide the following results for FEEDBACK VERTEX SET IN TOURNAMENTS and CLUSTER VERTEX DELETION.

- **Cluster Vertex Deletion.** For any  $0 < \epsilon < 1$ , the CLUSTER VERTEX DELETION problem admits a  $(1 + \epsilon)$ -approximate  $\mathcal{O}(\frac{1}{\epsilon} \cdot \text{opt})$ -vertex kernel.
- **Feedback Vertex Set in Tournaments.** For any  $0 < \epsilon < 1$ , the FEEDBACK VERTEX SET IN TOURNAMENTS problem admits a  $(1 + \epsilon)$ -approximate  $\mathcal{O}(\frac{1}{\epsilon} \cdot \text{opt})$ -vertex kernel.

**Reading Guide.** First, in Section 2, we present the concept lossy kernelization. Then, in Section 3, we present an overview of our proofs. In Section 4, we present some basic terminology used throughout the paper. In Section 5, we present a known result regarding

the support of optimum LP solutions to the LP-relaxation of  $d$ -HITTING SET. In Section 6, we present our lossy linear-element kernel for  $d$ -HITTING SET. In Section 7, we present our three lossy kernelization protocols (for VERTEX COVER, its generalization to  $d$ -HITTING SET with near-linear call size, and a protocol relating the problem to Ruzsa-Szemerédi graphs). In Section 8, we present our  $(1 + \epsilon)$ -approximate linear-vertex kernels for CLUSTER VERTEX DELETION and FEEDBACK VERTEX SET IN TOURNAMENTS. Lastly, in Section 9, we conclude with some open problems. For easy reference, problem definitions can be found in Appendix A.

## 2 Lossy Kernelization: Algorithms and Protocols

**Lossy Kernelization Algorithms.** We follow the framework of lossy kernelization presented in [25]. Here, we deal only with minimization problems where the value of a solution is its size, and where the computation of an arbitrary solution (where no optimization is enforced) is trivial. Thus, for the sake of clarity of presentation, we only formulate the definitions for this context, and remark that the definitions can be extended to the more general setting in the straightforward way (for more information, see [25]). To present the definitions, consider a parameterized problem  $\Pi$ . Given an instance  $I$  of  $\Pi$  with parameter  $k = \kappa(I)$ , denote: if  $k$  is a structural parameter, then  $\pi_I(\text{opt}) = \text{opt}$ , and otherwise (if  $k$  is a bound on the solution size given as part of the input)  $\pi_I(\text{opt}) = \min\{\text{opt}, k + 1\}$ . Moreover, for any solution  $S$  to  $I$ , denote: if  $k$  is a structural parameter, then  $\pi_I(S) = |S|$ , and otherwise  $\pi_I(S) = \min\{|S|, k + 1\}$ . We remark that when  $\pi$  is irrelevant (e.g., when the parameter is structural), we will drop it. A discussion of the motivation behind this definition of  $\pi_I$  can be found in [25]; here, we only briefly note that it signifies that we “care” only for solutions of size at most  $k$ —all other solutions are considered equally bad, treated as having size  $k + 1$ .

► **Definition 1.** *Let  $\Pi$  be a parameterized minimization problem. Let  $\alpha \geq 1$ . An  $\alpha$ -approximate kernelization algorithm for  $\Pi$  consists of two polynomial-time procedures: **reduce** and **lift**. Given an instance  $I$  of  $\Pi$  with parameter  $k = \kappa(I)$ , **reduce** outputs another instance  $I'$  of  $\Pi$  with parameter  $k' = \kappa(I')$  such that  $|I'| \leq f(k, \alpha)$  and  $k' \leq k$ . Given  $I, I'$  and a solution  $S'$  to  $I'$ , **lift** outputs a solution  $S$  to  $I$  such that  $\frac{\pi_I(S)}{\pi_I(\text{opt}(I))} \leq \alpha \frac{\pi_{I'}(S')}{\pi_{I'}(\text{opt}(I'))}$ . If  $\frac{\pi_I(S)}{\pi_I(\text{opt}(I))} \leq \max\{\alpha, \frac{\pi_{I'}(S')}{\pi_{I'}(\text{opt}(I'))}\}$  holds, then the algorithm is termed *strict*.*

In case  $\Pi$  admits an  $\alpha$ -approximate kernelization algorithm where the output has size  $f(k, \alpha)$ , or where the output has  $g(k, \alpha)$  “elements” (e.g., vertices), we say that  $\Pi$  admits an  $\alpha$ -approximate kernel of size  $f(k, \alpha)$ , or an  $\alpha$ -approximate  $g(k, \alpha)$ -element kernel, respectively. When it is clear from context, we simply write  $f(k)$  and  $g(k)$ . When it is guaranteed that  $|I'| \leq f(k', \alpha)$  rather than only  $|I'| \leq f(k, \alpha)$ , then we say that the lossy kernel is *output-parameter sensitive*.

We only deal with problems that have constant-factor polynomial-time approximation algorithms, and where we may directly work with (the unknown)  $\text{opt}$  as the parameter (then,  $\pi$  can be dropped). However, working with  $k$  (and hence  $\pi$ ) has the effect of artificially altering kernel sizes, but not so if one remembers that  $k$  and  $\text{opt}$  are different parameterizations. The following lemma clarifies a relation between these two parameterizations.

► **Lemma 2.** *Let  $\Pi$  be a minimization problem that, when parameterized by the optimum, admits an  $\alpha$ -approximate kernelization algorithm  $\mathfrak{A}$  of size  $f(\text{opt})$  (resp., an  $\alpha$ -approximate  $g(\text{opt})$ -element kernel). Then, when parameterized by  $k$ , a bound on the solution size that*

is part of the input, it admits an  $\alpha$ -approximate kernelization algorithm  $\mathfrak{B}$  of size  $f(\frac{k+1}{\alpha})$  (resp., an  $\alpha$ -approximate  $g(\frac{k+1}{\alpha})$ -element kernel).

**Proof.** We design  $\mathfrak{B}$  as follows. Given an instance  $(I, k)$  of  $\Pi$ , **reduce** of  $\mathfrak{B}$  calls **reduce** of  $\mathfrak{A}$  on  $I$ . If the output instance size is at most  $f(\frac{k+1}{\alpha})$  (resp., the output has at most  $g(\frac{k+1}{\alpha})$  elements), then it outputs this instance with parameter  $k' = k$ . Otherwise, it outputs a trivial constant-sized instance. Given  $(I, k), (I', k')$  and a solution  $S'$  to  $(I', k')$ , if  $I'$  is the output of the **reduce** procedure of  $\mathfrak{A}$  on  $I$ , then **lift** of  $\mathfrak{B}$  calls **lift** of  $\mathfrak{A}$  on  $I, I', S'$  and outputs the result. Otherwise, it outputs a trivial solution to  $I$ .

The **reduce** and **lift** procedures of  $\mathfrak{B}$  clearly have polynomial time complexities, and the definition of  $\mathfrak{B}$  implies the required size (or element) bound on the output of **reduce**. It remains to prove that the approximation ratio is  $\alpha$ . To this end, consider an input  $(I, k), (I', k'), S'$  to **lift** of  $\mathfrak{B}$ . Let  $S$  be its output. We differentiate between two cases.

- First, suppose that  $\text{opt}(I) \geq \frac{k+1}{\alpha}$ . Then,  $\frac{\pi_I(S)}{\pi_I(\text{opt}(I))} \leq \frac{k+1}{\frac{k+1}{\alpha}} = \alpha \leq \alpha \frac{\pi_{I'}(S')}{\pi_{I'}(\text{opt}(I'))}$  (where the last inequality follows because  $|S'| \geq \text{opt}(I')$  and hence  $\pi_{I'}(S') \geq \pi_{I'}(\text{opt}(I'))$ ).
- Second, suppose that  $\text{opt}(I) < \frac{k+1}{\alpha}$ . Then, it necessarily holds that  $I'$  is the output of the **reduce** procedure of  $\mathfrak{A}$  on  $I$ . Moreover, note that  $\text{opt}(I') \leq \text{opt}(I)$  and  $k' = k$ . So, if  $|S'| \geq k' + 2$ , then  $\frac{\pi_I(S)}{\pi_I(\text{opt}(I))} \leq \frac{k+1}{\pi_I(\text{opt}(I))} = \frac{k'+1}{\text{opt}(I)} \leq \frac{k'+1}{\text{opt}(I')} = \frac{\pi_{I'}(S')}{\pi_{I'}(\text{opt}(I'))}$ . Else, we suppose that  $|S'| \leq k' + 1$  and hence  $\pi_{I'}(S') = |S'|$ . Then,

$$\frac{\pi_I(S)}{\pi_I(\text{opt}(I))} \leq \frac{|S|}{\pi_I(\text{opt}(I))} = \frac{|S|}{\text{opt}(I)} \leq \alpha \frac{|S'|}{\text{opt}(I')} = \alpha \frac{\pi_{I'}(S')}{\pi_{I'}(\text{opt}(I'))}.$$

Here, the second inequality follows because the approximation ratio of  $\mathfrak{A}$  is  $\alpha$ .

This completes the proof.  $\blacktriangleleft$

Approximate kernelization algorithm often use strict reduction rules, defined as follows.

► **Definition 3.** Let  $\Pi$  be a parameterized minimization problem. Let  $\alpha \geq 1$ . An  $\alpha$ -strict reduction rule for  $\Pi$  consists of two polynomial-time procedures: **reduce** and **lift**. Given an instance  $I$  of  $\Pi$  with parameter  $k = \kappa(I)$ , **reduce** outputs another instance  $I'$  of  $\Pi$  with parameter  $k' = \kappa(I') \leq k$ . Given  $I, I'$  and a solution  $S'$  to  $I'$ , **lift** outputs a solution  $S$  to  $I$  such that  $\frac{\pi_I(S)}{\pi_I(\text{opt}(I))} \leq \max\{\alpha, \frac{\pi_{I'}(S')}{\pi_{I'}(\text{opt}(I'))}\}$ .

► **Proposition 4** ([25]). Let  $\Pi$  be a parameterized problem. For any  $\alpha \geq 1$ , an approximate kernelization algorithm for  $\Pi$  that consists only of  $\alpha$ -strict reduction rules has approximation ratio  $\alpha$ . Furthermore, it is strict.

**Lossy Kernelization Protocols.** We extend the notion of lossy kernelization algorithms to lossy kernelization protocols as follows.

► **Definition 5 (Lossy Kernelization Protocol).** Let  $\Pi$  be a parameterized minimization problem with parameter  $k$ . Let  $\alpha \geq 1$ . An  $\alpha$ -approximate kernelization protocol of call size  $f(k, \alpha)$  and  $g(k, \alpha)$  rounds for  $\Pi$  is defined as follows. First, the protocol assumes to have access to an oracle  $\mathfrak{D}$  that, given an instance  $I'$  of  $\Pi$  of size at most  $f(k, \alpha)$ , returns a solution  $S'$  to  $I'$  such that  $\pi_{I'}(S') \leq \beta \pi_{I'}(\text{opt}(I'))$  for minimization and  $\pi_{I'}(S') \geq \frac{1}{\beta} \pi_{I'}(\text{opt}(I'))$  for maximization, for some fixed  $\beta > 0$ . Second, for the same fixed  $\beta > 0$ , given an instance  $I$  of  $\Pi$ , the protocol may perform  $g(k, \alpha)$  calls to  $\mathfrak{D}$  and other operations in polynomial time, and then output a solution  $S$  to  $I$  such that  $\frac{\pi_I(S)}{\pi_I(\text{opt}(I))} \leq \alpha\beta$ .

The volume (or size) of the protocol is  $f(k, \alpha)g(k, \alpha)$ . In case  $g(k, \alpha) = g(\alpha)$  (i.e.,  $g$  depends only on  $\alpha$ ), the protocol is called pure.

Notice that an  $\alpha$ -approximate kernelization algorithm is the special case of an  $\alpha$ -approximate kernelization protocol when the number of rounds is 1.

Practically, we think that (lossy) kernelization protocols can often be as useful as standard (lossy) kernels, and, in some cases, more useful. Like standard (lossy) kernels, they reduce the total size of what we need to solve, only that now what we need to solve is split into several instances, to be solved one after another. On the one hand, this relaxation seems to, in most cases, not be restrictive (as what we really care about is the total size of what we need to solve). On the other hand, it might be helpful if by using this relaxation one can achieve better bounds than what is known (or, even, what is possible) on the sizes of the reduced instances, or to simplify the algorithm. For example, for the case of  $d$ -HITTING SET, we do not know how to beat  $\mathcal{O}(k^d)$  using a lossy kernel rather than a protocol.

### 3 Overview of Our Proof Ideas

In this section, we present a high-level overview of our proof ideas. For standard terminology not defined here or earlier, we refer the reader to Section 4.

#### 3.1 Linear-Element Lossy Kernel for $d$ -HITTING SET


We make use of a known result about the natural LP relaxation of  $d$ -HITTING SET: the support of any optimal LP solution to the LP-relaxation of  $d$ -HITTING SET is of size at most  $d \cdot \text{frac}$  where  $\text{frac}$  is the optimum (minimum value) of the LP [20]. For the sake of completeness, we provide a proof. We then provide a lossy reduction rule that computes an optimal LP solution, and deletes all vertices assigned values at least  $\frac{1}{d-1}$ . Having applied this rule exhaustively, we arrive at an instance having an optimal LP solution that assigns only values strictly smaller than  $\frac{1}{d-1}$ . Then, it can be shown that all hitting sets are contained within the support of this LP solution. In turn, in light of the aforementioned known result, this yields an approximate  $d \cdot \text{frac}$ -element and  $(d\text{frac})^d$ -set kernel that is output-parameter sensitive.

The analysis that the approximation factor is  $d - \frac{d-1}{d}$  is slightly more involved, and is based on case distinction. In case the number of vertices deleted is “small enough”, the cost of adding them is “small enough” as well. In the more difficult case where the number of vertices deleted is “large”, by making use of the already established bound on the output size as well as the drop in the fractional optimum, we are able to show that, in fact, we return a solution of approximation factor  $d - \frac{d-1}{d}$  irrespective of the approximation ratio of the solution we are given. More generally, the definition of “small enough” and “large” gives rise to a trade-off that is critical for our kernelization protocol for  $d$ -HITTING SET, which in particular yields that we can either obtain a *negligible additive error* or directly a solution of the desired (which is some fixed constant better than  $d$  but worse than  $d - \frac{d-1}{d}$ ) approximation ratio. Specifically, this means that it is “safe” to compose our element kernel as part of other kernelization algorithms or protocols.

#### 3.2 2-Round $\mathcal{O}(\text{frac}^{1.5})$ -Call Size Lossy Kernelization Protocol for VERTEX COVER

Towards the presentation of our near-linear call size lossy kernelization protocol for  $d$ -HITTING SET, we abstract some of the ideas using a simpler 2-round  $\mathcal{O}(\text{frac}^{1.5})$ -call size






■ **Figure 1** The three cases encountered by our 2-call lossy kernelization protocol for VERTEX COVER: (I)  $|S_1|$  is large, and we return  $V(G)$ ; (II)  $|S_1|$  is small and  $|S_2|$  is small, and we return  $S_1 \cup S_2$ ; (III)  $|S_1|$  is small and  $|S_2|$  is large, and we return  $(V(G) \setminus S_1) \cup A$ .

1.721-approximate kernelization protocol for VERTEX COVER (where  $\text{frac} \leq \text{opt} \leq k$  is the optimum of the natural LP relaxation of VERTEX COVER). First, we apply an (exact) kernelization algorithm to have a graph  $G$  on at most  $2\text{frac}$  vertices. The purpose of having only  $2\text{frac}$  vertices is twofold. First, it means that to obtain a “good enough” approximate solution, it suffices that we do not pick a “large enough” (linear fraction) of vertices of  $G$  to our solution. Second, it is required for a probability bound derived using union bound over vertex subsets to hold. Then, roughly speaking, the utility of the first oracle call is mainly, indirectly, to uncover a “large” (linear in  $n = |V(G)|$ ) induced subgraph of  $G$  that is “sparse”, and hence can be sent to the second oracle call to be solved optimally.

More precisely, after applying the initial kernelization, we begin by sampling roughly  $\text{frac}^{1.5}$  edges from  $G$ . Then, we call the oracle on the sampled graph to obtain a solution  $S_1$  to it (but not to  $G$ ). In case that solution  $S_1$  is “large” compared to the size of the vertex set of  $G$  (that is, sufficiently larger than  $n/2 \leq \text{frac}$ ), we can just return the entire vertex set of  $G$  (see Fig. 1). Else, we know that the subgraph of the sampled graph that is induced by  $V(G) \setminus S_1$  is edgeless. In addition, we can show (due to the initial kernelization) that with high probability, every set of edges of size (roughly) at least  $\text{frac}^{1.5}$  that is the edge set of some induced subgraph of  $G$  has been hit by our edge sample. Together, this implies that the subgraph of  $G$  induced by  $V(G) \setminus S_1$  has at most  $\text{frac}^{1.5}$  edges, and hence can be solved optimally by a second oracle call. Then, because we know that this subgraph is large compared to  $G$  (else  $S_1$  is large), if the oracle returned a “small” solution  $S_2$  to it, we may just take this solution together with  $S_1$  (which will form a vertex cover), and yet not choose sufficiently many vertices so that this will be good enough in terms of the approximation ratio achieved. Else, also because we know that this subgraph is large compared to  $G$ , if the second oracle returned a “large” solution  $S_2$ , then we know that every optimal solution must take many vertices from this subgraph, and hence, to compensate for this, the optimum of  $G[S_1]$  must be “very small”. So, we compute a 2-approximate solution  $A$  to  $G[S_1]$ , which we know should not be “too large”, and output the union of  $A$  and  $V(G) \setminus S_1$  (which yields a vertex cover).

### 3.3 Near-Linear Volume and Pure Lossy Kernelization Protocol for $d$ -HITTING SET

For any fixed  $\epsilon > 0$ , we present a pure  $d(1 - h(d, \epsilon))$ -approximate (randomized) kernelization protocol for  $d$ -HITTING SET with call size  $\mathcal{O}((\text{frac})^{1+\epsilon})$  where  $h(d, \epsilon)$  is a fixed positive constant that depends only on  $d, \epsilon$ . On a high-level, the idea of our more general lossy



■ **Figure 2** The nested solutions computed by oracle calls in our lossy kernelization protocol for  $d$ -HITTING SET. Each  $S_i$  is a solution to a subinstance  $(U_{i-1}, \mathcal{F}_{i-1})$  sampled from  $(U_{i-1}, \mathcal{T}_{i-1})$ .

kernelization protocol is to compute a nested family of solutions based on the approach described above for VERTEX COVER (see Fig. 2). Intuitively, as we now can sample only few sets (that is,  $\text{frac}^{1+\epsilon}$ ), when we compute a solution that hits them using an oracle call, the number of sets it misses can still be huge, and hence we will need to iteratively use the oracle (a constant number of times) until we reach a subuniverse such that we can optimally solve the subinstance induced by it by a single oracle call. Below, we give a more elaborate overview.

First, we apply our linear-element lossy kernel to have an instance  $I_0 = (U_0, \mathcal{T}_0)$  where the universe  $U_0$  consists of at most  $d \cdot \text{frac}$  elements. Here, the error of this application is not multiplied by the error attained next, but will only yield (as mentioned earlier) a negligible additive error (or directly a solution of the desired approximation ratio). The purpose of having only  $d \cdot \text{frac}$  elements is twofold, similarly as it is in the protocol described earlier for VERTEX COVER. Afterwards, we begin by sampling a family  $\mathcal{F}_1$  of roughly  $\text{frac}^{1+\epsilon}$  sets from  $\mathcal{T}_0$ . Then, we call the oracle on the sampled family  $\mathcal{F}_1$  to obtain a solution  $S_1$  to it. In case that solution  $S_1$  is “large” (sufficiently larger than  $|U_0|/d \leq \text{frac}$ ), we can just return  $U_0$ . Else, we know that the family of sets corresponding to the subinstance  $I_1$  induced by  $U_1 = U_0 \setminus S_1$ —that is, the family of all sets in  $\mathcal{T}_0$  contained in  $U_1$ , which we denote by  $\mathcal{T}_1$ —was missed by our set sample. In addition, we can show (due to the initial kernelization) that with high probability, every family of sets of size (roughly) at least  $\text{frac}^{d-\epsilon}$  that corresponds to a subinstance induced by a subset of  $U_0$  has been hit by our set sample. Together, this implies that  $\mathcal{T}_1$  has at most  $\text{frac}^{d-\epsilon}$  (rather than  $\text{frac}^d$ ) sets. Hence, in some sense, we have made progress towards the discovery of a sparse subinstance that we can optimally solve.

Due to important differences, let us describe also the second iteration—among at most  $\frac{1}{\epsilon}(d-1)$  iterations performed in total—before skipping to the (last) one where we have a subinstance that we can optimally solve by an oracle call. The last iteration may not even be reached, if we find a “good enough” solution earlier. We remark that it is critical to stop and return a solution as soon as we find a “large enough” one by an oracle call<sup>5</sup> as for our arguments to work, we need to always deal with subinstances whose universe is large (a linear fraction of  $|U_0|$ ), and these are attained by removing oracle solutions we got along the way. We begin the second iteration by sampling a family  $\mathcal{F}_2$  of roughly  $\text{frac}^{1+\epsilon}$  sets from  $\mathcal{T}_1$ . Then, we call the oracle on the sampled family  $\mathcal{F}_2$  to obtain a solution  $S_2$  to it. On the

<sup>5</sup> The solution we return is not the one given by the oracle call, but its union with another solution, as will be clarified immediately, or just  $U_0$  in case of the first iteration describe above.

one hand, in case that solution  $S_2$  is “large” (sufficiently larger than  $|U_1|/d$ ), we cannot just return  $U_0$  as in the first iteration, as now it may not be true that the optimum of  $I_0$  is large compared to  $|U_0|$ . Still, it is true that the optimum of  $I_1$  is large compared to  $|U_1|$ . So, every optimal solution (to  $I_0$ ) must take many elements from  $U_1 \setminus S_2$ , and hence, to compensate for this, the optimum of the subinstance induced by  $S_1$  must be “very small”. So, we compute a  $d$ -approximate solution to this subinstance, which we know should not be “too large”, and output the union of it and  $U_1$  (which yields a hitting set). On the other hand, in case  $S_2$  is “small”, we proceed as follows. We observe that the family of sets corresponding to the subinstance  $I_2$  induced by  $U_2 = U_1 \setminus S_2$ , whose family of sets we denote by  $\mathcal{T}_2$ , was missed by our set sample. In addition, we can show (due to the initial kernelization) that with high probability, every family of sets of size (roughly) at least  $\text{frac}^{d-2\epsilon}$  that corresponds to a subinstance induced by a subset of  $U_1$  has been hit by our set sample. Together, this implies that  $\mathcal{T}_2$  has at most  $\text{frac}^{d-2\epsilon}$  (rather than just  $\text{frac}^{d-\epsilon}$  as in the first iteration) sets. Hence, in some sense, we have made further progress towards the discovery of a sparse subinstance that we can optimally solve.

Finally, we arrive at a subinstance  $I'$  induced by a subuniverse  $U' \subseteq U_0$  that is of size linear in  $U_0$  (else we should have returned a solution earlier) and where the family of sets,  $\mathcal{F}'$ , is of size at most  $\text{frac}^{1+\epsilon}$ . Then, we call the oracle on  $I'$  to obtain a solution  $S'$  to it. On the one hand, in case that solution  $S'$  is “large” (sufficiently larger than  $|U'|/d$ ), we compute a  $d$ -approximate solution to the subinstance induced by  $U_0 \setminus U'$  (which is the union of all solutions returned by oracle calls except the last one), and output the union of it and  $U'$ . Otherwise, we output  $(U_0 \setminus U') \cup S'$ , which is “good enough” because  $U'$  is sufficiently large while  $S'$  is sufficiently small compared to it, it does not contain a “large enough” number of elements from  $U_0$ .

### 3.4 Outlook: Relation to Ruzsa-Szemerédi Graphs

A graph  $G$  is an  $(r, t)$ -Ruzsa-Szemerédi graph if its edge set can be partitioned into  $t$  edge-disjoint induced matchings, each of size  $r$ . These graphs were introduced in 1978 [29], and have been extensively studied since then. When  $r$  is a function of  $n$ , let  $\gamma(r)$  denote the maximum  $t$  (which is a function of  $n$ ) such that there exists an  $(r, t)$ -Ruzsa-Szemerédi graph. In [19], the authors considered the case where  $r = cn$ . They showed that when  $c = \frac{1}{4}$ ,  $\gamma(r) \in \Theta(\log n)$ , and when  $\frac{1}{5} \leq c \leq \frac{1}{4}$ ,  $t \in \mathcal{O}(\frac{n}{\log n})$ . It is an open problem whether whenever  $c$  is a fixed constant,  $t \in \mathcal{O}(n^{1-\epsilon})$ . For any fixed constant  $0 < c < \frac{1}{4}$ , we present a  $(1 + 4c)$ -approximate (randomized) kernelization protocol for VERTEX COVER with  $t + 1$  rounds and call size  $\mathcal{O}(t(\text{frac})^{1.5})$ . Clearly, this result makes sense only when  $t \in o(\sqrt{n})$ , preferably  $t \in \mathcal{O}(n^{\frac{1}{2}-\lambda})$  for  $\lambda$  as close to  $1/2$  as possible, because the volume is  $\mathcal{O}(\text{opt}^{2-\lambda})$ . If  $t$  is “sufficiently small” (depending on the desired number of rounds) whenever  $c$  is a fixed constant (specifically, substitute  $c = \frac{\epsilon}{4}$ ), this yields a  $(1 + \epsilon)$ -approximate kernelization protocol.

We observe that, for a graph  $G$ ,  $r = r(n), t = t(n) \in \mathbb{N}$  and  $U_1, U_2, \dots, U_t \subseteq V(G)$  such that for all  $i \in \{1, 2, \dots, t\}$ ,  $G[U_i]$  has a matching  $M_i$  of size at least  $r$ , and for all distinct  $i, j \in \{1, 2, \dots, t\}$ ,  $E(G[U_i]) \cap E(G[U_j]) = \emptyset$ , we have that  $G$  is a supergraph of an  $(r, t)$ -Ruzsa-Szemerédi graph. Having this observation in mind, we devise our protocol as follows. After applying an exact  $2\text{frac}$ -vertex kernel, we initialize  $E' = \emptyset$ , and we perform  $t + 1$  iterations of the following procedure. We sample a set of roughly  $\text{frac}^{1.5}$  edges from  $G$ , and call the oracle on the subgraph of  $G$  whose edge set is the set of sampled edges union  $E'$  to obtain a solution  $S$  to it (but not to  $G$ ), and compute a maximal matching  $M$  in  $G - S$ . If  $|M|$  is smaller than  $cn \leq 2\text{frac}$ , then we return the union of the set of vertices incident

to edges in  $M$  (which is a solution to  $G - S$ ) and  $S$ . Else, similarly to the first protocol we described for VERTEX COVER, we can show that with high probability,  $G - S$  has (roughly) at most  $k^{1.5}$  edges, and we add this set of edges to  $E'$ . The crux of the proof is in the argument that, at the latest, at the  $(t + 1)$ -st iteration the computed matching will be of size smaller than  $cn \leq 2c\text{frac}$ , as otherwise we can use the matchings we found, together with the vertex sets (of the form  $G - S$ ) we found them in, to construct an  $(r, t + 1)$ -Ruzsa-Szemerédi graph based on the aforementioned observation, which contradicts the choice of  $t$ .

### 3.5 $(1 + \epsilon)$ -Approximate $\mathcal{O}(\frac{1}{\epsilon} \cdot \text{opt})$ -Vertex Kernel for Implicit 3-HITTING SET Problems

Both of our lossy kernels share a common scheme, which might be useful to derive  $(1 + \epsilon)$ -approximate linear-vertex kernels for other implicit hitting and packing problems as well. Essentially, they both consist of two rules (although in the presentation, they are merged for simplicity). To present them, we remind that a module (in a graph) is a set of vertices having the same neighborhood relations with all vertices outside the set. Now, our first rule reveals some modules in the graph, and our second rule shrinks their size. The first rule in both of our lossy kernels is essentially the same.

Now, we elaborate on the first rule. We start by computing an optimal solution  $\alpha$  to the LP-relaxation of the corresponding 3-HITTING SET problem. Notice that  $\text{support}(\alpha)$  is a solution, and its size is at most  $3\text{frac}$  (in fact, we show that it is at most  $3\text{frac} - 2|\alpha^{-1}(1)|$ ). Then, the first rule is as follows. At the beginning, no vertex is marked. Afterwards, one-by-one, for each vertex  $v$  assigned 1 by  $\alpha$  (i.e., which belongs to  $\alpha^{-1}(1)$ ), we construct a graph whose vertex set is the set of yet unmarked vertices in  $V(G) \setminus \text{support}(\alpha)$  and where there is an edge between every two vertices that create an obstruction together with  $v$  (that is, an induced  $P_3$  in CLUSTER VERTEX DELETION and a triangle in FEEDBACK VERTEX SET IN TOURNAMENTS). We compute a maximal matching in this graph, and decrease its size to  $\frac{1}{\epsilon}$  if it is larger (in which case, it is no longer maximal). The vertices incident to the edges in the matching are then considered marked. We prove that among the vertices in  $\alpha^{-1}(1)$  whose matching size was decreased, whose set is denoted by  $D$ , any solution can only exclude an  $\epsilon$  fraction of its size among the vertices in  $D$ , and hence it is “safe” (in a lossy sense) to delete  $D$ . Let  $M$  be the set of all marked vertices. Then, we show that  $(\text{support}(\alpha) \cup M) \setminus \{v\}$ , for any  $v \in \text{support}(\alpha)$  (including those not in  $\alpha^{-1}(1)$ ), is also a solution.

For CLUSTER VERTEX DELETION, we prove that the outcome of the first rule means that the vertex set of every clique in  $G - (\text{support}(\alpha) \cup M)$  is a module in  $G - D$ , and that for every vertex in  $\text{support}(\alpha)$ , the set of its neighbors in  $V(G - (\text{support}(\alpha) \cup M))$  is the vertex set of exactly one of these cliques. So, for CLUSTER VERTEX DELETION, this gives rise to the following second reduction rule (which is, in fact, exact) to decrease the size of module. For every clique among the aforementioned cliques whose size is larger than that of its neighborhood, we arbitrarily remove some of its vertices so that its size will be equal to the size of its neighborhood. This rule is safe since if at least one of the vertices in such a clique is deleted by a solution, then because it is a module, either that deletion is irrelevant or the entire clique is deleted, and in the second case we might just as well delete its neighborhood instead. Because the neighborhoods of the cliques are pairwise-disjoint (since for every vertex in  $\text{support}(\alpha)$ , the set of its neighbors in  $V(G - (\text{support}(\alpha) \cup M))$  is the vertex set of exactly one of the cliques), this means that now their total size is at most  $(\text{support}(\alpha) \setminus D) \cup M$ , and hence we arrive at the desired kernel.

For FEEDBACK VERTEX SET IN TOURNAMENTS, we consider the unique (because  $G$  is a tournament) topological ordering of the vertices in  $G - \text{support}(\alpha)$ , so that all arcs are “forward”

arcs. We prove that the outcome of the first rule means that each vertex  $v \in \text{support}(\alpha)$  has a unique position within this ordering when restricted to  $G - (\text{support}(\alpha) \cup M)$ , so that still all arcs (that is, including those incident to  $v$ ) are forward arcs in  $G - (\text{support}(\alpha) \cup M) \cup \{v\}$ . (Further, the vertex set of each subtournament induced by the vertices “between” any two marked vertices in  $G - \text{support}(\alpha)$  is a module in  $G - D$ .) We are thus able to characterize all triangles in  $G - D$  as follows: each either consists of three vertices in  $(\text{support}(\alpha) \setminus D) \cup M$ , or it consists of a vertex  $v \in \text{support}(\alpha) \setminus D$ , a vertex  $u \in (\text{support}(\alpha) \setminus D) \cup M$  and a vertex  $w \in V(G) \setminus (\text{support}(\alpha) \cup M)$  with a backward arc between  $v$  and  $u$  and where  $w$  is “in-between” the positions of  $v$  and  $u$ . This gives rise to a reduction rule for module shrinkage whose presentation and analysis are more technical than that of CLUSTER VERTEX DELETION (in particular, unlike the second rule of CLUSTER VERTEX DELETION, the second rule of FEEDBACK VERTEX SET IN TOURNAMENTS is lossy) and of the first rule, and hence we defer them to the appropriate Section 8.2.

## 4 Preliminaries

### 4.1 General Notation

The *support* of a function  $f : A \rightarrow \mathbb{R}$  is  $\{a \in A : f(a) \neq 0\}$ , denoted by  $\text{support}(f)$ .

Given an instance  $I$  of some optimization problem  $\Pi$ , we denote by  $\text{opt}(I)$  the optimum (value of an optimal solution, if one exists) of  $I$ . When  $I$  is clear from context, we simply write  $\text{opt}$ .

To bound the approximation ratios of our algorithms, we will use the following fact.

► **Proposition 6** (Folklore, see, e.g., [25]). *For any positive reals  $x, y, p$  and  $q$ ,  $\min\left(\frac{x}{p}, \frac{y}{q}\right) \leq \frac{x+y}{p+q} \leq \max\left(\frac{x}{p}, \frac{y}{q}\right)$ .*

We now present a well-known Chernoff bound, to be used in the analysis of our (randomized) lossy kernelization protocols.

► **Proposition 7.** *Let  $X_1, \dots, X_n$  be independent random variables over  $\{0, 1\}$ . Let  $X$  denote their sum and let  $\mu = \mathbb{E}[X]$  denote the expected value of  $X$ . Then, for any  $0 \leq \delta \leq 1$ ,*

$$\text{Prob}[X \geq (1 + \delta)\mu] \leq e^{-\frac{\delta^2 \mu}{3}}.$$

### 4.2 Graph Notation

Given a graph  $G$ , let  $V(G)$  and  $E(G)$  denote its vertex set and edge (or arc) set, respectively. When clear from context,  $n = |V(G)|$  and  $m = |E(G)|$ . Given a vertex  $v \in V(G)$ , let  $N_G(v)$  denote the set of neighbors of  $v$  in  $G$ , and given a subset  $U \subseteq V(G)$ , let  $N_G(U)$  denote the open neighborhood of  $U$  in  $G$ . Given a subset  $U \subseteq V(G)$ , let  $G[U]$  denote the subgraph of  $G$  induced by  $U$ , that is, the graph on vertex set  $U$  and edge set  $\{\{u, v\} \in E(G) : u, v \in U\}$ . Moreover, given a subgraph  $G'$  of  $G$  (possibly  $G' = G$ ) and a subset  $U \subseteq V(G)$  (possibly  $U \setminus V(G') \neq \emptyset$ ), let  $G' - U$  denote the graph on vertex set  $V(G') \setminus U$  and edge set  $\{\{u, v\} \in E(G') : u, v \notin U\}$ . A *module* in  $G$  is a subset  $U \subseteq V(G)$  such that for every vertex  $v \in V(G) \setminus U$  either  $U \subseteq N_G(v)$  or  $U \cap N_G(v) = \emptyset$ . Given a subset  $W \subseteq E(G)$ , let  $G - W$  denote the graph on vertex set  $V(G)$  and edge set  $E(G) \setminus W$ . An *induced  $P_3$*  in  $G$  is a path on three vertices in  $G$  whose endpoints are not adjacent in  $G$ . A *cluster graph* is a graph in which every connected component is a clique. An *acyclic digraph* is a digraph that contains no directed cycles. A *tournament* is a digraph where for every two vertices  $u, v$ , exactly one among the arcs  $(u, v)$  and  $(v, u)$  belongs to the digraph.

► **Definition 8.** A graph  $G$  is an  $(r, t)$ -Ruzsa-Szemerédi graph if its edge set can be partitioned into  $t$  edge-disjoint induced matchings, each of size  $r$ .

These graphs were introduced in 1978 [29], and have been extensively studied since then. In [19], the authors considered the case where  $r = cn$ . They showed that when  $c = \frac{1}{4}$ , the maximum  $t$ , which we denote by  $\gamma(r)$ , is  $\Theta(\log n)$ , and when  $\frac{1}{5} \leq c \leq \frac{1}{4}$ ,  $t = \mathcal{O}(\frac{n}{\log n})$ . It is an open problem whether when  $c$  is a fixed constant,  $t = \mathcal{O}(n^{1-\epsilon})$ .

### 4.3 Linear Programming

A canonical form of a linear program (LP) is  $[\max \sum_{i=1}^n c_i x_i \text{ s.t. } \forall j = 1, \dots, m : \sum_{i=1}^n a_{ji} x_i \leq b_j; \forall i = 1, \dots, n : x_i \geq 0]$ , or  $[\min \sum_{j=1}^m b_j y_j \text{ s.t. } \forall i = 1, \dots, n : \sum_{j=1}^m a_{ij} y_j \geq c_i; \forall j = 1, \dots, m : y_j \geq 0]$ . Here, the  $x_i$ 's ( $y_j$ 's) are variables. Moreover, two programs of the aforementioned forms that refer to the same set of coefficients  $\{c_i\}_{i=1}^n, \{a_{ij} : i \in \{1, \dots, n\}, j \in \{1, \dots, m\}\}, \{b_j\}_{j=1}^m$  are *dual* of each other. A *solution* to an LP is an assignment of real values to its variables so that all constraints are satisfied. Further, a solution is *optimal* if it also optimizes (maximizes or minimizes) the value of the objective function. The optimum (value of an optimal solution, if one exists) of an LP  $I$  (or which is associated with some entity  $I$ , where no confusion can arise) is denoted by  $\text{frac}(I)$ . When  $I$  is clear from context, we simply write  $\text{frac}$ .

► **Proposition 9** ([26]). Any LP (with rational coefficients) that admits a solution, admits an optimal solution that assigns only rational values. Furthermore, such an optimal solution can be computed in polynomial time.

We will need a well-known proposition relating optimal solutions to LPs and their duals, known as strong duality and complementary slackness:

► **Proposition 10** ([26]). Let **(P)**  $[\max \sum_{i=1}^n c_i x_i \text{ s.t. } \forall j = 1, \dots, m : \sum_{i=1}^n a_{ji} x_i \leq b_j; \forall i = 1, \dots, n : x_i \geq 0]$  be a primal LP; **(D)**  $[\min \sum_{j=1}^m b_j y_j \text{ s.t. } \forall i = 1, \dots, n : \sum_{j=1}^m a_{ij} y_j \geq c_i; \forall j = 1, \dots, m : y_j \geq 0]$  be the dual LP. Let  $\alpha$  and  $\beta$  be solutions to **(P)** and **(D)**, respectively. Then,  $\alpha$  and  $\beta$  are both optimal if and only if  $\sum_{i=1}^n c_i \alpha(x_i) = \sum_{j=1}^m b_j \beta(y_j)$  [strong duality]. Moreover,  $\alpha$  and  $\beta$  are both optimal if and only if [complementary slackness]:

- For  $i = 1, \dots, n$ :  $\alpha(x_i) > 0$  if and only if  $\sum_{j=1}^m a_{ij} \beta(y_j) = c_i$ .
- For  $j = 1, \dots, m$ :  $\beta(y_j) > 0$  if and only if  $\sum_{i=1}^n a_{ji} \alpha(x_i) = b_j$ .

## 5 The Support Size of Any Optimal Solution to the LP of $d$ -HITTING SET

In this section, we present a tight bound on the support size of any optimal solution to the classic LP of the  $d$ -HITTING SET problem, defined as follows.

► **Definition 11.** Let  $(U, \mathcal{F})$  be an instance of  $d$ -HITTING SET. Then, the classic LP that corresponds to  $(U, \mathcal{F})$  is defined as follows:  $[\min \sum_{u \in U} y_u \text{ s.t. } \forall S \in \mathcal{F} : \sum_{u \in S} y_u \geq 1; \forall u \in U : y_u \geq 0]$ .

We will re-name  $y$  by  $x$  when it is more convenient (in Section 6) and no confusion arises.

We present the following theorem, which has been originally proved in [20]. For the sake of completeness, we present a short proof here.

► **Theorem 12** ([20]). Let  $I = (U, \mathcal{F})$  be an instance of  $d$ -HITTING SET. Let  $\beta$  be an optimal solution to its classic LP. Then,  $|\text{support}(\beta)| \leq d \cdot \text{frac}(I)$ . In particular,  $|\text{support}(\beta)| \leq d \cdot \text{opt}(I)$ .

**Proof.** Let us denote the classic LP that corresponds to  $(U, \mathcal{F})$  by  $(\mathbf{D})$ . We note that the dual LP of  $(\mathbf{D})$ , which we denote by  $(\mathbf{P})$ , is defined as follows:  $[\max \sum_{S \in \mathcal{F}} x_S \text{ s.t. } \forall u \in U : \sum_{S \in \mathcal{F}: u \in S} x_S \leq 1; \forall S \in \mathcal{F} : x_S \geq 0]$ . Let  $\alpha$  be an optimal solution to  $(\mathbf{P})$ . Then,

$$\begin{aligned}
\text{frac}(I) &= \sum_{u \in U} \beta(y_u) && [\beta \text{ is optimal}] \\
&= \sum_{S \in \mathcal{F}} \alpha(x_S) && [\text{strong duality}] \\
&= \frac{1}{d} \cdot \sum_{S \in \mathcal{F}} (d \cdot \alpha(x_S)) \\
&\geq \frac{1}{d} \cdot \sum_{S \in \mathcal{F}} \sum_{u \in S} \alpha(x_S) && [\forall S \in \mathcal{F} : |S| \leq d] \\
&= \frac{1}{d} \cdot \sum_{u \in U} \sum_{S \in \mathcal{F}: u \in S} \alpha(x_S) \\
&\geq \frac{1}{d} \cdot \sum_{u \in \text{support}(\beta)} \sum_{S \in \mathcal{F}: u \in S} \alpha(x_S) \\
&= \frac{1}{d} \cdot \sum_{u \in \text{support}(\beta)} 1 && [\text{complementary slackness}] \\
&= \frac{1}{d} \cdot |\text{support}(\beta)|
\end{aligned}$$

We conclude that  $|\text{support}(\beta)| \leq d \cdot \text{opt}(I)$ . Because  $\text{frac}(I) \leq \text{opt}(I)$ , the proof is complete.  $\blacktriangleleft$

Observe that the bound in Theorem 12 is tight, that is, it is satisfied with equality for infinitely many instances of  $d$ -HITTING SET. To see this, for any  $n \in \mathbb{N}$  that is a multiple of  $d$ , consider an instance  $I = (U, \mathcal{F})$  where  $|U| = n$  and  $\mathcal{F}$  is a partition of  $U$  into parts of equal size  $d$  (so,  $|\mathcal{F}| = n/d$ ). Then, the optimum of the corresponding classic LP is easily seen to be  $n/d$ , and it can be attained by an assignment that assigns  $1/d$  to each variable, and thus has support size  $n = d \cdot \text{opt}(I)$ .

## 6 A $(d - \frac{d-1}{d})$ -Approximate Linear-Element Kernel for $d$ -HITTING SET

We first present the following reduction rule that is the basis of our kernelization algorithm.

► **Definition 13.** *The  $d$ -HITTING SET element reduction rule is defined as follows:*

- **reduce:** *Let  $I = (U, \mathcal{F})$  be an instance of  $d$ -HITTING SET. Use the algorithm in Proposition 9 to compute an optimal solution  $\alpha$  to the classic LP corresponding to it (Definition 11). Let  $H = \{u \in U : \alpha(u) \geq \frac{1}{d-1}\}$ . Output  $I' = (U', \mathcal{F}')$  where  $\mathcal{F}' = \{S \in \mathcal{F} : S \cap H = \emptyset\}$  and  $U' = \bigcup \mathcal{F}'$ .*
- **lift:** *Given  $I, I'$  and a solution  $S'$  to  $I'$ , output  $S = S' \cup H$ .*

Essentially, our approximate kernelization algorithm will consist of exhaustive (i.e., as long as  $|H| \geq 1$ ) application of the  $d$ -HITTING SET element reduction rule. Unfortunately, the  $d$ -Hitting Set rule is *not*  $(1 - \frac{d-1}{d})$ -strict, and hence, unlike other lossy kernelization algorithms that consist of repetitive applications of one or more reduction rules, we cannot make direct use of Proposition 4. So, we present the algorithm explicitly in order to ease its analysis.

► **Definition 14.** *The  $d$ -HITTING SET element kernelization algorithm is defined as follows:*

## 23:16 Lossy Kernelization for (Implicit) Hitting Set Problems

- **reduce:** Let  $I = (U, \mathcal{F})$  be an instance of  $d$ -HITTING SET. Let  $i = 1, \mathcal{F}_1 = \mathcal{F}$  and  $U_1 = \bigcup \mathcal{F}_1$ . As long as a break command is not reached:
  1. Use the algorithm in Proposition 9 to compute an optimal solution  $\alpha_i$  to the classic LP corresponding to  $I_i = (U_i, \mathcal{F}_i)$  (Definition 11).
  2. Let  $H_i = \{u \in U : \alpha(u) \geq \frac{1}{d-1}\}$ . If  $H_i = \emptyset$ , then break the loop.
  3. Increase  $i$  by 1, and let  $\mathcal{F}_i = \{S \in \mathcal{F}_{i-1} : S \cap H_{i-1} = \emptyset\}$  and  $U_i = \bigcup \mathcal{F}_i$ . Let  $H^* = \bigcup_{j=1}^{i-1} H_j$ . Output  $I' = (U', \mathcal{F}')$  where  $\mathcal{F}' = \{S \in \mathcal{F} : S \cap H^* = \emptyset\}$  (which equals  $\mathcal{F}_i$ ) and  $U' = \bigcup \mathcal{F}'$  (which equals  $U_i$ ).
- **lift:** Given  $I, I'$  and a solution  $S'$  to  $I'$ , output  $S = S' \cup H^*$ .

In order to bound the output size of our kernelization algorithm, we will make use of the following lemma, whose proof is based on Theorem 12.

► **Lemma 15.** Let  $I = (U, \mathcal{F})$  be an instance of  $d$ -HITTING SET where  $U = \bigcup_{S \in \mathcal{F}} S$ , and let  $\alpha$  be an optimal solution to its classic LP that assigns only values strictly smaller than  $\frac{1}{d-1}$ . Then,  $|U| \leq d \cdot \text{frac}(I)$ .

**Proof.** We first claim that every  $S \in \mathcal{F}$  is a subset of  $\text{support}(\alpha)$ . To this end, consider some set  $S \in \mathcal{F}$ . Then, because  $\alpha$  is a solution, it satisfies  $\sum_{u \in S} \alpha(x_u) \geq 1$ . Targeting a contradiction, suppose that there exists  $u' \in S \setminus \text{support}(\alpha)$ . Then, because  $|S| \leq d$  and  $\alpha$  assigns only values strictly smaller than  $\frac{1}{d-1}$ , we have that

$$\sum_{u \in S} \alpha(x_u) = \sum_{u \in S \setminus \{u'\}} \alpha(x_u) < \sum_{u \in S \setminus \{u'\}} \frac{1}{d-1} \leq 1,$$

which yields a contradiction.

We conclude that  $\bigcup \mathcal{F} \subseteq \text{support}(\alpha)$ . By Theorem 12,  $|\text{support}(\alpha)| \leq d \cdot \text{frac}(I)$ , and hence  $|\bigcup \mathcal{F}| \leq d \cdot \text{frac}(I)$ . Because  $U = \bigcup_{S \in \mathcal{F}} S$ , the proof is complete. ◀

In particular, we now show this lemma yields the desired bound on the number of elements in the output instance of our kernelization algorithm:

► **Lemma 16.** Let  $I = (U, \mathcal{F})$  be an instance of  $d$ -HITTING SET. Consider a call to **reduce** of the  $d$ -HITTING SET element kernelization algorithm on input  $I = (U, \mathcal{F})$  and whose output is  $I' = (U', \mathcal{F}')$ . Then,  $|U'| \leq d \cdot \text{frac}(I)$  and  $|\mathcal{F}'| \leq (d \cdot \text{frac}(I))^d$ .

**Proof.** Due to the condition to break the loop in **reduce**, we have an instance  $I'$  whose classic LP admits an optimal solution  $\alpha'$  that assigns only values strictly smaller than  $\frac{1}{d-1}$ . Moreover, recall that  $U' = \bigcup_{S \in \mathcal{F}'} S$ . So, by Lemma 15,  $|U'| \leq d \cdot \text{frac}(I')$ . Clearly, this also implies that  $|\mathcal{F}'| \leq \binom{|U'|}{d} \leq \binom{d \cdot \text{frac}(I)}{d} \leq (d \cdot \text{frac}(I))^d$ . ◀

We now justify the approximation ratio of our kernelization algorithm. We remark that the particular way in which we phrase it, in particular distinguishing between the two items in its statement rather than only in its proof, is required for later purposes, as we explain before stating Theorem 18.

► **Lemma 17.** Let  $I = (U, \mathcal{F})$  be an instance of  $d$ -HITTING SET. Consider a call to **lift** of the  $d$ -HITTING SET element kernelization algorithm on input  $I = (U, \mathcal{F}), I' = (U', \mathcal{F}'), S'$  and whose output is  $S$ . For any  $0 < \rho$ , at least one of the following conditions holds:

1.  $|S| - |S'| \leq \rho \cdot \text{opt}(I)$ .
2.  $\frac{|S|}{\text{opt}(I)} \leq d - \frac{\rho}{d-1}$ .



Furthermore,  $\frac{|S|}{\text{opt}(I)} \leq (d - \frac{d-1}{d}) \frac{|S'|}{\text{opt}(I')}$ .

**Proof.** We consider two cases, depending on  $|H^*|$ .

1. First, suppose that  $|H^*| \leq \rho \cdot \text{opt}(I)$ . Then, because  $|S| - |S'| = |H^*|$ , we directly have that  $|S| - |S'| \leq \rho \cdot \text{opt}(I)$ .
2. Second, suppose that  $|H^*| \geq \rho \cdot \text{opt}(I)$ . Let  $t$  denote the number of iterations performed by the  $d$ -HITTING SET element kernelization algorithm. For every  $i \in \{1, 2, \dots, t-1\}$ , observe that  $\alpha_i|_{\{x_u: u \in U_{i+1}\}}$  is a solution to the classic LP corresponding to  $I_{i+1}$ , therefore
 
$$\sum_{u \in U_{i+1}} \alpha_{i+1}(x_u) \leq \sum_{u \in U_{i+1}} \alpha_i(x_u) = \sum_{u \in U_i} \alpha_i(x_u) - \sum_{u \in H_i} \alpha_i(x_u) \leq \sum_{u \in U_i} \alpha_i(x_u) - \frac{1}{d-1} |H_i|.$$
 (Here, the last inequality follows since  $\alpha_i(x_u) \geq \frac{1}{d-1}$  for every  $u \in H_i$ .) So,

$$\begin{aligned} \sum_{u \in U_t} \alpha_t(x_u) &\leq \sum_{u \in U_{t-1}} \alpha_{t-1}(x_u) - \frac{1}{d-1} |H_{t-1}| \\ &\leq \sum_{u \in U_{t-2}} \alpha_{t-2}(x_u) - \frac{1}{d-1} |H_{t-1}| - \frac{1}{d-1} |H_{t-2}| \\ &\quad \dots \\ &\leq \sum_{u \in U_1} \alpha_1(x_u) - \frac{1}{d-1} |H_{t-1}| - \frac{1}{d-1} |H_{t-2}| - \dots - \frac{1}{d-1} |H_1| \\ &= \sum_{u \in U_1} \alpha_1(x_u) - \frac{1}{d-1} |H^*|. \end{aligned}$$

In particular,  $\text{frac}(I') \leq \text{frac}(I) - \frac{1}{d-1} |H^*|$ . Moreover, by Lemma 16 and because  $S' \subseteq U'$ , we know that  $|S'| \leq d \cdot \text{frac}(I')$ . So,

$$\begin{aligned} |S| = |S'| + |H^*| &\leq d \cdot \text{frac}(I') + |H^*| \\ &\leq d \cdot (\text{frac}(I) - |H^*|/(d-1)) + |H^*| \\ &\leq d \cdot \text{opt}(I) - |H^*|/(d-1) \\ &\leq (d - \rho/(d-1)) \cdot \text{opt}(I). \end{aligned}$$

This directly implies that  $\frac{|S|}{\text{opt}(I)} \leq d - \frac{\rho}{d-1}$ .

This proves the first part of the lemma. For the second part, we choose  $\rho = \frac{(d-1)^2}{d}$ . Now, we show that in each of the aforementioned two cases,  $\frac{|S|}{\text{opt}(I)} \leq (d - \frac{d-1}{d}) \frac{|S'|}{\text{opt}(I')}$ . For the second case, this directly follows by substituting  $\rho$  by  $\frac{(d-1)^2}{d}$ . So, in what follows, we only consider the first case, where  $|S| - |S'| \leq \rho \cdot \text{opt}(I) = \frac{(d-1)^2}{d} \cdot \text{opt}(I)$ , and hence  $|S| \leq |S'| + \frac{(d-1)^2}{d} \cdot \text{opt}(I)$ . Then,

$$\begin{aligned} \frac{|S|}{\text{opt}(I)} &\leq \frac{|S'| + \frac{(d-1)^2}{d} \cdot \text{opt}(I)}{\text{opt}(I)} \\ &\leq \frac{|S'|}{\text{opt}(I')} + \frac{(d-1)^2}{d} \\ &\leq (1 + \frac{(d-1)^2}{d}) \frac{|S'|}{\text{opt}(I')} \end{aligned}$$

Here, the second inequality follows since  $\text{opt}(I') \leq \text{opt}(I)$ , and the third inequality follows since  $|S'| \geq \text{opt}(I')$ . Now, observe that  $1 + \frac{(d-1)^2}{d} = \frac{d}{d} + \frac{d^2-2d+1}{d} = \frac{d^2-d+1}{d} = d - \frac{d-1}{d}$ . So, indeed  $\frac{|S|}{\text{opt}(I)} \leq (d - \frac{d-1}{d}) \frac{|S'|}{\text{opt}(I')}$ . ◀

## 23:18 Lossy Kernelization for (Implicit) Hitting Set Problems

We are now ready to prove the main theorem of this subsection. In particular, while we prove that our kernelization algorithm is a  $(d - \frac{d-1}{d})$ -approximate  $d \cdot \text{frac}$ -element and  $(d \cdot \text{frac})^d$ -set kernel, we also state that it is output-parameter sensitive, and we should keep in mind that it also satisfies the two conditions in Lemma 17. In particular, we will need the two conditions in this lemma for the purpose of being able to compose it later: rather than incurring a  $(d - \frac{d-1}{d})$  multiplicative error, it can be used so that it either incurs an (*essentially*) *negligible additive* error, or returns a solution  $S$  of approximation ratio better than  $d$  (though not  $(d - \frac{d-1}{d})$ , but depending on how “negligible” the additive error in the first case should be) irrespective of the approximation ratio of the solution  $S'$  given to it. These conditions will be necessary for the correctness of our approximate kernelization protocol for  $d$ -HITTING SET that is given in the next section.

► **Theorem 18.** *The  $d$ -HITTING SET problem, parameterized by the fractional optimum of the classic LP, admits a  $(d - \frac{d-1}{d})$ -approximate  $d \cdot \text{frac}$ -element and  $(d \cdot \text{frac})^d$ -set kernel. Furthermore, it is output-parameter sensitive.*

**Proof.** Clearly, the **lift** procedure of the kernelization algorithm is performed in polynomial time. Further, the loop of the **reduce** procedure can perform at most  $|U|$  iterations before the one where it breaks (since each of them removes at least one element from the universe), and each is performed in polynomial time, so overall this procedure is performed in polynomial time. The bounds on the number of elements in the output as well as its size, along with the property of being output-parameter sensitive, follow from Lemma 16. Lastly, the approximation ratio follows from Lemma 17. This completes the proof. ◀

Because parameterization by the fractional optimum of the classic LP is lower bounded by parameterization by the optimum, and due to Lemma 2, we have the following corollaries of Theorem 18.

► **Corollary 19.** *The  $d$ -HITTING SET problem, parameterized by the optimum, admits a  $(d - \frac{d-1}{d})$ -approximate  $d \cdot \text{opt}$ -element  $(d \cdot \text{opt})^d$ -set kernel.*

► **Corollary 20.** *The  $d$ -HITTING SET problem, parameterized by a bound  $k$  on the solution size, admits a  $(d - \frac{d-1}{d})$ -approximate  $\frac{d}{d-1} \cdot (k+1)$ -element  $(\frac{d}{d-1} \cdot (k+1))^d$ -set kernel.*

It is noteworthy that when  $d = 2$ , in which  $d$ -HITTING SET equals VERTEX COVER, we retrieve the classic result that VERTEX COVER admits a 1-approximate (i.e., exact)  $2k$ -vertex kernelization algorithm [17]. This does not follow directly from the stated approximation ratio of  $d - \frac{d-1}{d}$  (which equals  $1\frac{1}{2}$  rather than 1 when  $d = 2$ ). However, the argument used to prove the correctness of the classic result, that is, that there exists a solution that contains all vertices whose variables are assigned 1, also implies for our kernel that it is exact (see, e.g., [17]). Thus, our theorem regarding  $d$ -HITTING SET can be viewed as a generalization of this classic result.

### 7 A Pure $d'$ -Approximate Kernelization Protocol for $d$ -HITTING SET of Almost Linear Call Size where $d' < d$

For the sake of clarity, we first give a warm-up example. Afterwards, we present our general result that is based on the approach presented by that warm-up example, non-trivial insights regarding how to apply that approach in a recursive manner, and critically also on Theorem 12 (via Theorem 18). Lastly, we present some further outlook by relating a method to prove the existence of a  $(1 + \epsilon)$ -approximate kernelization protocol for VERTEX COVER to the

non-existence of  $(r, t)$ -Ruzsa-Szemerédi graphs where  $r$  is linear in  $n$  (the number of vertices) and  $t$  is “large”, which is an open problem.

We will make use of a polynomial-time  $d$ -approximation algorithm for  $d$ -HITTING SET:

► **Proposition 21** (Folklore). *The  $d$ -HITTING SET problem admits a polynomial-time  $d$ -approximation algorithm.*

## 7.1 Warm-Up Example: A 1.721-Approximate Kernelization Protocol for VERTEX COVER of 2 Rounds and Call Size $(2k)^{1.5}$

We start with a warm-up and, in a sense, toy example which exemplifies a main insight behind our more general result, that is, that essentially we may use the oracle to find a “large subinstance” that is “sparse”, and hence which (with another oracle call), we can solve optimally. We will make use of Theorem 12 (as to stay as close as possible to the proof of the more general result, where it is necessary), though here, as  $d = 2$ , one can equally use the classic 1-approximate  $2k$ -vertex kernel for VERTEX COVER [17].

► **Theorem 22.** *The VERTEX COVER problem, parameterized by the fractional optimum of the classic LP, admits a pure, having 2 rounds,  $\frac{2}{\sqrt{10}-2}$ -approximate<sup>6</sup> (randomized)<sup>7</sup> kernelization protocol with call size  $2\text{frac} + 2(2\text{frac})^{1.5}$  (where the number of edges is at most  $2(2\text{frac})^{1.5}$ ).*

**Proof.** We first describe the algorithm. To this end, consider some input  $\hat{I} = (\hat{U}, \hat{\mathcal{F}})$  (in terms of graphs,  $\hat{U}$  is the vertex set and  $\hat{\mathcal{F}}$  is the edge set of the input graph).<sup>8</sup> Then:

1. Call the **reduce** procedure of the algorithm in Theorem 18 on  $\hat{I}$  to obtain a new instance  $I = (U, \mathcal{F})$  where  $|U| \leq 2\text{frac}(I)$ . (Recall that when  $d = 2$ , this algorithm is exact.)<sup>9</sup>
2. Let  $0 < \nu < 1$  (analogous to  $\frac{\mu}{2}$  in the general result) be a fixed constant that will be determined later.
3. Sample  $\mathcal{F}_1$  from  $\mathcal{F}$  as follows: Insert each set  $S \in \mathcal{F}$  to  $\mathcal{F}_1$  independently at random with probability  $p_1 = \frac{1}{(2\text{frac}(I))^{0.5}}$ .
4. If  $|\mathcal{F}_1| > 2p_1|\mathcal{F}|$ , then let  $S$  be an arbitrary solution to  $I$ , and proceed directly to Step 11. [#Failure]
5. Call the oracle on  $(U, \mathcal{F}_1)$ , and let  $S_1$  denote its output.
6. If  $|S_1| \geq \nu|U|$ , then let  $S = U$ , and proceed directly to Step 11. [#Success]
7. Let  $U_1 = U \setminus S_1$  and  $\mathcal{T}_1 = \{S \in \mathcal{F} : S \subseteq U_1\}$ .
8. If  $|\mathcal{T}_1| > 2(2\text{frac}(I))^{1.5}$ , then let  $S$  be an arbitrary solution to  $I$ , and proceed directly to Step 11. [#Failure]
9. Call the oracle on  $I' = (U_1, \mathcal{T}_1)$ , and let  $S_2$  denote its output.
10. Let  $S' = S_2 \cup S_1$  and  $S'' = U_1 \cup T$  where  $T$  is a 2-approximate solution to  $\tilde{I} = (S_1, \{S \in \mathcal{F} : S \subseteq S_1\})$  (computed using Proposition 21). Let  $S$  be a minimum-sized set among  $S'$  and  $S''$ . [#Success]
11. Call the **lift** procedure of the algorithm in Theorem 18 on  $\hat{I}, I, S$  to obtain a solution  $\hat{S}$  to  $\hat{I}$ . Output  $\hat{S}$ .

<sup>6</sup> Note that  $\frac{2}{\sqrt{10}-2} \leq 1.721$ .

<sup>7</sup> Here, randomization means that we may fail to return a  $(\frac{2}{\sqrt{10}-2})$ -approximate solution (i.e., we may return a “worse” solution), but we must succeed with probability, say, at least  $9/10$ . It should be clear that the success probability can be boosted to any constant arbitrarily close to 1.

<sup>8</sup> We represent the input using a universe and sets so that it will resemble our more general protocol more.

<sup>9</sup> See the discussion at the end of Section 6.

Clearly, the algorithm runs in polynomial time, and only two oracle calls are performed. Further, when we call the oracle on  $(U, \mathcal{F}_1)$ ,  $|\mathcal{F}_1| \leq 2p_1|\mathcal{F}| \leq 2 \cdot \frac{1}{(2\text{frac}(I))^{0.5}} \cdot (2\text{frac}(I))^2 = 2(2\text{frac}(I))^{1.5}$  (due to **reduce**). Thus, each oracle call is performed on an instance with at most  $2\text{frac}(I)$  vertices (as  $|U| \leq 2\text{frac}(I)$  due to **reduce**) and  $2(2\text{frac}(I))^{1.5}$  edges, and since  $\text{frac}(I) \leq \text{frac}(\widehat{I})$ , the statement in the lemma regarding the call size is satisfied.

We now consider the probability of failure. By Chernoff bound (Proposition 7), the probability that  $|\mathcal{F}_1| > 2p_1|\mathcal{F}|$  is at most  $e^{-\frac{p_1|\mathcal{F}|}{3}}$ . Further, by union bound, the probability that there exists a subset  $U' \subseteq U$  such that  $\mathcal{F}_1 \cap \mathcal{F}_{U'} = \emptyset$  (where  $\mathcal{F}_{U'} = \{S \in \mathcal{F} : S \subseteq U'\}$ ) under the assumption that  $|\mathcal{F}_{U'}| > 2(2\text{frac}(I))^{1.5}$  is at most  $2^{2\text{frac}(I)} \cdot (1 - p_1)^{2(2\text{frac}(I))^{1.5}} = 2^{2\text{frac}(I)} \cdot (1 - \frac{1}{(2\text{frac}(I))^{0.5}})^{2(2\text{frac}(I))^{1.5}} \leq 2^{2\text{frac}(I)} \cdot e^{-4\text{frac}(I)}$ . Thus, by union bound, under the implicit supposition that  $\text{frac}$  (and  $|\mathcal{F}|$ ) is a large enough constant (e.g., 10),<sup>10</sup> the probability that at least one of the events in the steps marked by “failure” occurs is at most  $1/10$ . Notice that if these events occur,  $S$  is a solution. Further, we now claim that if these events do not occur, then we compute a set  $S$  that is a solution to  $I$  and, furthermore, it is  $\frac{2}{\sqrt{10-2}}$ -approximate. Then, by the correctness of **lift** (in particular, since the kernelization algorithm in Theorem 18 is 1-approximate, that is, exact, for  $d = 2$ ), this will conclude the proof. For this purpose, we have the following case distinction, where  $\beta$  is the approximation ratio of the oracle.

First, suppose that  $S$  is computed in Step 6. Then,  $|S_1| \geq \nu|U|$  and  $S = U$ . Clearly,  $S$  is a solution to  $I$ . Because  $S_1$  is a  $\beta$ -approximate solution to  $(U, \mathcal{F}_1)$ , which is a subinstance of  $(U, \mathcal{F})$ , this means that  $\text{opt}(I) \geq \frac{\nu}{\beta}|U|$ . So, in this case, the approximation ratio is  $\frac{|S|}{\text{opt}(I)} \leq \frac{|U|}{\frac{\nu}{\beta}|U|} = \beta \frac{1}{\nu}$ .

Second, suppose that  $S$  is computed in Step 10. Then,  $|S_1| < \nu|U|$ . On the one hand, because  $S_2$  is a solution to  $I' = (U_1, \mathcal{T}_1)$ , and, as  $\mathcal{T}_1 = \{S \in \mathcal{F} : S \subseteq U_1\}$ , every set in  $\mathcal{F} \setminus \mathcal{T}_1$  contains at least one vertex from  $U \setminus U_1 = S_1$ , we have that  $S' = S_2 \cup S_1$  is a solution to  $I$ . Further, since  $S_2$  is a  $\beta$ -approximate solution to  $I'$ ,  $|S'| \leq \beta \text{opt}(I') + |S_1|$ . On the other hand, because  $T$  is a solution to  $\widetilde{I} = (S_1, \{S \in \mathcal{F} : S \subseteq S_1\})$ , and every set in  $\mathcal{F} \setminus \{S \in \mathcal{F} : S \subseteq S_1\}$  contains at least one vertex from  $U_1$ , we have that  $S'' = U_1 \cup T$  is also a solution to  $I$ . Further, because  $T$  is a 2-approximate solution to  $\widetilde{I}$ ,  $|S''| \leq 2\text{opt}(\widetilde{I}) + |U_1| = 2\text{opt}(\widetilde{I}) + |U| - |S_1|$ .

Consider some optimal solution  $S^*$  to  $I$ . Then,  $S^* \setminus S_1$  is a solution to  $I'$ , and  $S^* \cap S_1$  is a solution to  $\widetilde{I}$ , which means that  $\text{opt}(I') \leq |S^* \setminus S_1|$  and  $\text{opt}(\widetilde{I}) \leq |S^* \cap S_1|$ . So, denoting  $\lambda = \frac{|S^* \cap S_1|}{|S_1|}$  ( $0 \leq \lambda \leq 1$ ) and  $\rho = \frac{|S_1|}{|U|}$  ( $0 \leq \rho < \nu$ ), we know that

$$\blacksquare |S'| \leq \beta|S^* \setminus S_1| + |S_1| = \beta \text{opt}(I) - \beta|S^* \cap S_1| + |S_1| = \beta \text{opt}(I) + (1 - \beta\lambda)\rho|U| \leq (\beta + 2\rho - 2\beta\lambda\rho)\text{opt}(I).$$

$$\blacksquare |S''| \leq 2|S^* \cap S_1| + |U| - |S_1| = (1 + 2\lambda\rho - \rho)|U| \leq (2 + 4\lambda\rho - 2\rho)\text{opt}(I).$$

As  $\lambda$  grows larger, the first term becomes better, and as it grows smaller, the second term is better. So, the worst case is such that equality is attained when  $\lambda = \frac{\beta + 4\rho - 2}{2(2 + \beta)\rho}$ . Then, the approximation ratio is  $2 + 4\frac{\beta + 4\rho - 2}{2(2 + \beta)\rho}\rho - 2\rho = 2 + \frac{2(\beta + 4\rho - 2)}{2 + \beta} - 2\rho = 2 - \frac{4 - 2\beta}{2 + \beta} + (\frac{4 - 2\beta}{2 + \beta})\rho$ . When  $\beta \geq 2$ , the correctness of the approximation ratio is trivial, since then even returning all of  $U$  is a  $\beta$ -approximation. So, suppose that  $\beta < 2$ . Then, the aforementioned function grows larger as  $\rho$  grows larger (since when  $\beta < 2$ , its coefficient is positive), and as  $\rho < \nu$ , an upper bound on the maximum is  $2 + (\frac{4 - 2\beta}{2 + \beta})\nu - \frac{4 - 2\beta}{2 + \beta}$ . Now, we fix  $\nu$  such that  $2 + (\frac{4 - 2\beta}{2 + \beta})\nu - \frac{4 - 2\beta}{2 + \beta} = \frac{\beta}{\nu}$  when  $\beta = 1$ . So, we require  $\frac{4}{3} + \frac{2}{3}\nu = \frac{1}{\nu}$ , that is,  $2\nu^2 + 4\nu - 3 = 0$ , which is satisfied when  $\nu = \frac{\sqrt{10}}{2} - 1$ . Then, in the first case, the approximation ratio is at most  $\beta \frac{1}{\nu} = \beta \frac{2}{\sqrt{10-2}}$  as required. In the

<sup>10</sup> Otherwise, the instance can be solved optimally in polynomial time using brute-force.

second case, the approximation ratio is also  $2 + \left(\frac{4-2\beta}{2+\beta}\right)\nu - \frac{4-2\beta}{2+\beta} \leq \beta\left(\frac{4}{3} + \frac{2}{3}\nu\right) = \beta\frac{2}{\sqrt{10-2}}$  as required. This completes the proof.  $\blacktriangleleft$

► **Corollary 23.** *The VERTEX COVER problem, parameterized by the optimum, admits a pure, having 2 rounds,  $\frac{2}{\sqrt{10-2}}$ -approximate (randomized) kernelization protocol with call size  $2\text{opt} + 2(2\text{opt})^{1.5}$  (where the number of edges is at most  $2(2\text{opt})^{1.5}$ ).*

## 7.2 Generalization to Almost Linear Call Size and $d \geq 2$

A critical part of our algorithm is Theorem 18. First, after calling its algorithm to reduce the number of elements, there will only be  $2^{\text{frac}}$  many subsets of  $U$  such that, if the instance induced by them is not “sparse enough” (where the definition of sparse enough becomes stricter and stricter as the execution of our algorithm proceeds), then with high probability we will “hit” at least one of their sets when using an oracle call. Further, Theorem 18 will be used to prove that, after calling its algorithm to reduce the number of elements, once we find a “sufficiently” large (linear in  $k$ ) subset of  $U$  along with a solution to the instance induced by that subset that is large compared to its size (in particular, consisting of more than a fraction of  $1/d$  of its elements), we are essentially done. Our algorithm will repeatedly try to find subsets as mentioned above, while, if it fails at every step, it eventually arrives at a “sufficiently” large (linear in  $k$ ) subset of  $U$  such that it can optimally solve the instance induced by that subset.

► **Theorem 24.** *For any fixed  $\epsilon > 0$ , the  $d$ -HITTING SET problem, parameterized by the fractional optimum of the classic LP, admits a pure  $d(1 - h(d, \epsilon))$ -approximate (randomized)<sup>11</sup> kernelization protocol with call size  $d \cdot \text{frac} + 2^{\frac{d}{\epsilon}}(d \cdot \text{frac})^{1+\epsilon}$  (where the number of sets is at most  $(d \cdot \text{frac})^{1+\epsilon}$ ) where  $h(d, \epsilon) = \frac{1}{10d} \left(\frac{1}{4}\right)^{\frac{d}{\epsilon}}$  is a fixed positive constant that depends only on  $d, \epsilon$ .<sup>12</sup>*

**Proof.** We first describe the algorithm. To this end, consider some input  $\widehat{I} = (\widehat{U}, \widehat{\mathcal{F}})$ . Then:

1. Call the **reduce** procedure of the algorithm in Theorem 18 on  $I$  to obtain a new instance  $I = (U, \mathcal{F})$  where  $|U| \leq d \cdot \text{frac}(I)$ .
2. Denote  $\mu = \mu(d) = \frac{d+1}{2}$ , and  $\tau = \tau(d, \epsilon) = \frac{1}{\epsilon}(d-1)$ .
3. Initialize  $U_0 = U$  and  $\mathcal{T}_0 = \mathcal{F}$ .
4. For  $i = 1, 2, \dots, \tau$ :
  - a. Sample  $\mathcal{F}_i$  from  $\mathcal{T}_{i-1}$  as follows: Insert each set  $S \in \mathcal{T}_{i-1}$  to  $\mathcal{F}_i$  independently at random with probability  $p_i = \frac{1}{(d \cdot \text{frac}(I))^{d-1-i\epsilon}}$ .
  - b. If  $|\mathcal{F}_i| > 2^i(d \cdot \text{frac}(I))^{1+\epsilon}$ , then let  $S$  be an arbitrary solution to  $I$ , and proceed directly to Step 6. [#Failure]
  - c. Call the oracle on  $(U_{i-1}, \mathcal{F}_i)$ , and let  $S_i$  denote its output. [#We will verify in the proof that all calls are done with at most  $(d \cdot \text{frac}(\widehat{I}))^{1+\epsilon}$  sets.]
  - d. If  $|S_i| \geq \frac{\mu}{d}|U_{i-1}|$ , then:

<sup>11</sup> Here, randomization means that we may fail to return a  $(d - h(d, \epsilon))$ -approximate solution (i.e., we may return a “worse” solution), but we must succeed with probability, say, at least  $9/10$ . It should be clear that the success probability can be boosted to any constant arbitrarily close to 1.

<sup>12</sup> We remark that we preferred to simplify the algorithm and its analysis rather than to optimize  $h(d, \epsilon)$  (in fact, the same algorithms with slightly more careful analysis already yields a much better yet “uglier” constant). In particular, our approximation ratio is a *fixed constant* (under the assumption that  $d, \epsilon$  are fixed) strictly smaller than  $d$ .

## 23:22 Lossy Kernelization for (Implicit) Hitting Set Problems

- i. Call the algorithm in Proposition 21 on  $(U \setminus U_{i-1}, \{S \in \mathcal{F} : S \subseteq U \setminus U_{i-1}\})$ , and let  $T$  denote its output.
- ii. Let  $S = T \cup U_{i-1}$  and proceed directly to Step 6. [#Success]
- e. Let  $U_i = U_{i-1} \setminus S_i$  and  $\mathcal{T}_i = \{S \in \mathcal{T}_{i-1} : S \subseteq U_i\}$ .
5. Let  $S = S_\tau \cup (U \setminus U_{\tau-1})$ . [#Success]
6. Call the **lift** procedure of the algorithm in Theorem 18 on  $\widehat{I}, I, S$  to obtain a solution  $\widehat{S}$  to  $\widehat{I}$ . Output  $\widehat{S}$ .

Clearly, the algorithm runs in polynomial time. Further, each oracle call has at most  $d \cdot \text{frac}(I)$  many elements. We first verify that it also has at most  $(d \cdot \text{frac}(I))^{1+\epsilon}$  sets. To this end, we have two preliminary claims.

▷ **Claim 25.** For all  $i = 1, 2, \dots, \tau$ , if the algorithm reaches iteration  $i$  and  $|\mathcal{T}_{i-1}| \leq 2^{i-1}(d \cdot \text{frac}(I))^{d-(i-1)\epsilon}$ , then  $|\mathcal{F}_i| \leq 2^i(d \cdot \text{frac}(I))^{1+\epsilon}$  with probability at least  $1 - e^{-\frac{2^{i-1}(d \cdot \text{frac}(I))^{1+\epsilon}}{3}}$ .

**Proof.** Observe that the expected size of  $\mathcal{F}_i$  is:

$$\begin{aligned} \mathbb{E}[|\mathcal{F}_i|] &= |\mathcal{T}_{i-1}| \cdot p_i \\ &\leq 2^{i-1}(d \cdot \text{frac}(I))^{d-(i-1)\epsilon} \cdot \frac{1}{(d \cdot \text{frac}(I))^{d-1-i\epsilon}} \\ &= 2^{i-1}(d \cdot \text{frac}(I))^{1+\epsilon}. \end{aligned}$$

Thus, Chernoff bound (Proposition 7) implies that

$$\text{Prob}[|\mathcal{F}_i| > 2^i(d \cdot \text{frac}(I))^{1+\epsilon}] \leq e^{-\frac{2^{i-1}(d \cdot \text{frac}(I))^{1+\epsilon}}{3}}.$$

This completes the proof of the claim. ◀

▷ **Claim 26.** For all  $i = 1, 2, \dots, \tau$ , if the algorithm reaches Step 4e in iteration  $i$  and  $|\mathcal{T}_{i-1}| \leq 2^{i-1}(d \cdot \text{frac}(I))^{d-(i-1)\epsilon}$ , then  $|\mathcal{T}_i| \leq 2^i(d \cdot \text{frac}(I))^{d-i\epsilon}$  with probability at least  $1 - (\frac{2}{e})^{2^i \cdot d \cdot \text{frac}(I)}$ .

**Proof.** Consider some iteration  $i \in \{1, 2, \dots, \tau\}$ , and suppose that the algorithm reaches Step 4e iteration  $i$ . Hence,  $|\mathcal{F}_i| \leq 2^i(d \cdot \text{frac}(I))^{1+\epsilon}$ . Consider some subfamily  $\mathcal{T}' \subseteq \mathcal{T}_{i-1}$  such that  $|\mathcal{T}'| > 2^i(d \cdot \text{frac}(I))^{d-i\epsilon}$ . Then,

$$\begin{aligned} \text{Prob}(\mathcal{T} \cap \mathcal{F}_i = \emptyset) &= (1 - p_i)^{|\mathcal{T}'|} \\ &\leq \left(1 - \frac{1}{(d \cdot \text{frac}(I))^{d-1-i\epsilon}}\right)^{2^i(d \cdot \text{frac}(I))^{d-i\epsilon}} \\ &\leq e^{-2^i \cdot d \cdot \text{frac}(I)}. \end{aligned}$$

Because there exist at most  $2^{d \cdot \text{frac}(I)}$  subsets of  $U_{i-1}$ , union bound implies that the probability that there exists  $U' \subseteq U_{i-1}$  such that the subfamily  $\{S \in \mathcal{T}_{i-1} : S \subseteq U'\}$  is of size larger than  $2^i(d \cdot \text{frac}(I))^{d-i\epsilon}$  and has empty intersection with  $\mathcal{F}_i$  is at most  $(\frac{2}{e})^{2^i \cdot d \cdot \text{frac}(I)}$ . Recall that  $\mathcal{T}_i = \{S \in \mathcal{T}_{i-1} : S \subseteq U_i\}$  and note that  $\mathcal{T}_i$  has empty intersection with  $\mathcal{F}_i$  because  $S_i = U_{i-1} \setminus U_i$  is a solution to  $(U_i, \mathcal{F}_i)$  (by the correctness of the oracle). This completes the proof of the claim. ◀

We now prove the desired bound on each call size, based on Claims 25 and 26.

▷ **Claim 27.** The following statement holds with probability at least 9/10: For all  $i = 1, 2, \dots, \tau$ , if the algorithm reaches iteration  $i$  and calls the oracle, then  $|\mathcal{F}_i| \leq 2^i(d \cdot \text{frac}(I))^{1+\epsilon}$  and the algorithm does not exit the loop in Step 4b.

**Proof.** We claim that for every  $j \in \{0, 1, \dots, \tau\}$ , the following holds with probability at least  $1 - \sum_{i=1}^j \left(\frac{2}{e}\right)^{2^i \cdot d \cdot \text{frac}(I)}$ : for every  $i \in \{0, 1, \dots, j\}$  such that the algorithm reaches Step 4e in iteration  $i$  (when  $i = 0$ , we mean the initialization),  $|\mathcal{T}_i| \leq 2^i (d \cdot \text{frac}(I))^{d-i\epsilon}$ . The proof is by induction on  $j$ . At the basis, where  $j = 0$ ,  $\mathcal{T}_0 = \mathcal{F}$ , and hence due to **reduce**, with probability 1,  $|\mathcal{T}| \leq (d \cdot \text{frac}(I))^d$ . Now, suppose that the claim is true for  $j - 1$ , and let us prove it for  $j$ . By the inductive hypothesis, with probability at least  $1 - \sum_{i=1}^{j-1} \left(\frac{2}{e}\right)^{2^i \cdot d \cdot \text{frac}(I)}$ , the following holds: for every  $i \in \{0, 1, \dots, j-1\}$  such that the algorithm reaches Step 4e in iteration  $i$ ,  $|\mathcal{T}_i| \leq 2^i (d \cdot \text{frac}(I))^{d-i\epsilon}$ . Now, if the algorithm further reaches Step 4e in iteration  $j$ , Claim 26 implies that  $|\mathcal{T}_j| \leq 2^j (d \cdot \text{frac}(I))^{d-j\epsilon}$  with probability at least  $1 - \left(\frac{2}{e}\right)^{2^j \cdot d \cdot \text{frac}(I)}$ . So, by union bound, the claim for  $j$  is true.

In particular, by setting  $j = \tau$ , we have that with probability at least  $1 - \sum_{i=1}^{\tau} \left(\frac{2}{e}\right)^{2^i \cdot d \cdot \text{frac}(I)}$ , the following holds: for every  $i \in \{0, 1, \dots, \tau\}$  such that the algorithm reaches Step 4e in iteration  $i$  (when  $i = 0$ , we mean the initialization),  $|\mathcal{T}_i| \leq 2^i (d \cdot \text{frac}(I))^{d-i\epsilon}$ . However, by Claim 25 and union bound, this directly extends to the following statement: with probability at least  $1 - \sum_{i=1}^{\tau} \left(\frac{2}{e}\right)^{2^i \cdot d \cdot \text{frac}(I)} - \sum_{i=1}^{\tau} e^{-\frac{2^{i-1} (d \cdot \text{frac}(I))^{1+\epsilon}}{3}}$ , the following holds: for every  $i \in \{0, 1, \dots, \tau\}$  such that the algorithm reaches iteration  $i$  and calls the oracle, then  $|\mathcal{F}_i| \leq 2^i (d \cdot \text{frac}(I))^{1+\epsilon}$  and the algorithm does not exit the loop in Step 4b. Now, observe that

$$\begin{aligned} \sum_{i=1}^{\tau} \left(\frac{e}{2}\right)^{-2^i \cdot d \cdot \text{frac}(I)} + \sum_{i=1}^{\tau} e^{-\frac{2^{i-1} (d \cdot \text{frac}(I))^{1+\epsilon}}{3}} &\leq \tau \cdot \left( \left(\frac{e}{2}\right)^{-d \cdot \text{frac}(I)} + e^{-\frac{(d \cdot \text{frac}(I))^{1+\epsilon}}{3}} \right) \\ &\leq 2\tau \cdot \left(\frac{e}{2}\right)^{-\frac{(d \cdot \text{frac}(I))^{1+\epsilon}}{3}} \leq \frac{1}{10}. \end{aligned}$$

Here, the last inequality follows by assuming that  $\text{frac}(I)$  is large enough (to ensure that the inequality is satisfied) compared to  $d, \epsilon$ . Indeed, if this is not the case, then  $\text{frac}(I)$  (and hence also  $\text{opt}(I)$ , because it is bounded by  $d \cdot \text{frac}(I)$ ) is a fixed constant (that depends only on  $d, \epsilon$ ), and hence the problem can just be a-priori solved in polynomial time by, e.g., brute force search. We thus conclude that the failure probability is at most  $1/10$ , which completes the proof of the claim.  $\blacktriangleleft$

Let  $\beta \geq 1$  denote the approximation ratio of the oracle. We now turn to analyze the approximation ratio. Towards that, we present a lower bound on the size of each universe  $U_i$ .

$\triangleright$  **Claim 28.** For all  $i = 1, 2, \dots, \tau$ , if the algorithm reaches iteration  $i$  and computes  $U_i$ , then  $|U_i| \geq (1 - \frac{\mu}{d})^i |U| \geq \left(\frac{1}{4}\right)^{\frac{1}{\epsilon} d} |U|$ .

**Proof.** We first claim that for all  $i \in \{1, 2, \dots, \tau\}$ , if the algorithm reaches iteration  $i$  and computes  $U_i$ , then  $|U_i| \geq (1 - \frac{\mu}{d})^i |U|$ . The proof is by induction on  $i$ , where we let  $i = 0$  be the basis. Then, in the basis,  $U_0 = U$  and the claim trivially holds. Now, suppose that the claim holds for  $i - 1$ , and let us prove it for  $i$ . By the inductive hypothesis,  $|U_{i-1}| \geq (1 - \frac{\mu}{d})^{i-1} |U|$ . Further, by the definition of  $U_i$ ,  $U_i = U_{i-1} \setminus S_i$ , and as the algorithm reaches the computation of  $U_i$ ,  $|S_i| < \frac{\mu}{d} |U_{i-1}|$ . Thus, we have that

$$|U_i| \geq |U_{i-1}| - |S_i| > \left(1 - \frac{\mu}{d}\right) |U_{i-1}| \geq \left(1 - \frac{\mu}{d}\right)^i |U|.$$

Hence, the claim holds for  $i$ , and therefore our (sub)claim holds.

## 23:24 Lossy Kernelization for (Implicit) Hitting Set Problems

Lastly, observe that for all  $i \in \{1, 2, \dots, \tau\}$ ,  $|U_i| \geq |U_\tau|$ . Moreover, due to our (sub)claim and substitution of  $\tau$  and  $\mu$ , and because  $\frac{x+1}{2x} \leq \frac{3}{4}$  for all  $x \geq 2$  (the maximum is achieved when  $x = 2$ ), we have that

$$|U_\tau| \geq \left(1 - \frac{\mu}{d}\right)^\tau |U| = \left(1 - \frac{d+1}{2d}\right)^{\frac{1}{\epsilon}(d-1)} |U| \geq \left(\frac{1}{4}\right)^{\frac{1}{\epsilon}d} |U|.$$

This completes the proof of the claim.  $\blacktriangleleft$

Now, having the property that each universe  $U_i$  is “large enough”, we argue that if  $S$  is computed in Step 4(d)ii, then it is a solution of the approximation ratio  $d(1 - d \cdot h(d, \epsilon))$ .

$\triangleright$  **Claim 29.** For all  $i = 1, 2, \dots, \tau$ , if the algorithm reaches iteration  $i$  and Step 4(d)ii of that iteration, then  $S$  is a solution to  $I$  such that  $\frac{|S|}{\text{opt}(I)} \leq \beta d(1 - d \cdot h(d, \epsilon))$ .

**Proof.** Let  $i \in \{1, 2, \dots, \tau\}$  such that the algorithm reaches iteration  $i$  and Step 4(d)ii of that iteration. Then,  $|S_i| \geq \frac{\mu}{d}|U_{i-1}|$  and  $S = T \cup U_{i-1}$  (I). Let  $S^*$  be an optimal solution to  $I$ , so  $|S^*| = \text{opt}(I)$  (II). Consider the following subinstances of  $I$ :

- $I' = (U_{i-1}, \mathcal{F}_i)$ . Because  $S_i$  is a  $\beta$ -approximate solution to  $I'$ , we have that  $\text{opt}(I') \geq \frac{|S_i|}{\beta} \geq \frac{\mu}{\beta d}|U_{i-1}|$ .
- $I'' = (U_{i-1}, \{S \in \mathcal{F} : S \subseteq U_{i-1}\})$ . Because  $I'$  is a subinstance of  $I''$ , we have that  $\text{opt}(I'') \geq \text{opt}(I')$ , and hence  $\text{opt}(I'') \geq \frac{\mu}{\beta d}|U_{i-1}|$ . In particular, since  $S^* \cap U_{i-1}$  is a solution to  $I''$ , we have that  $|S^* \cap U_{i-1}| \geq \frac{\mu}{\beta d}|U_{i-1}|$ . This has two consequences: first,  $|U_{i-1}| \leq \frac{\beta d}{\mu}|S^* \cap U_{i-1}|$  (III); second, due to Claim 28,  $|S^* \cap U_{i-1}| \geq \frac{\mu}{\beta d}|U_{i-1}| \geq \frac{\mu}{\beta d}\left(\frac{1}{4}\right)^{\frac{1}{\epsilon}d}|U|$  (IV).
- $I''' = (U \setminus U_{i-1}, \{S \in \mathcal{F} : S \subseteq U \setminus U_{i-1}\})$ . Due to Proposition 21,  $T$  is a solution to  $I'''$  such that  $|T| \leq d \cdot \text{opt}(I''')$ . Note that all sets in  $\mathcal{F}$  that do not occur in this instance have non-empty intersection with  $U_{i-1}$ , and hence  $S$  is a solution to  $I$ . Further,  $S^* \setminus U_{i-1}$  is a solution to  $I'''$ , and hence  $|S^* \setminus U_{i-1}| \geq \text{opt}(I''')$ . Thus,  $|T| \leq d|S^* \setminus U_{i-1}|$  (V).

So, we have proved that  $S$  is a solution to  $I$ , and we have that

$$\begin{aligned} |S| &= |T| + |U_{i-1}| && \text{[(I)]} \\ &\leq d|S^* \setminus U_{i-1}| + \frac{\beta d}{\mu}|S^* \cap U_{i-1}| && \text{[(III)+(V)]} \\ &= d(|S^*| - (1 - \frac{\beta}{\mu})|S^* \cap U_{i-1}|) \\ &\leq d(|S^*| - (1 - \frac{\beta}{\mu})\frac{\mu}{\beta d}\left(\frac{1}{4}\right)^{\frac{1}{\epsilon}d}|U|) && \text{[(IV)]} \\ &= d(\text{opt}(I) - (1 - \frac{\beta}{\mu})\frac{\mu}{\beta d}\left(\frac{1}{4}\right)^{\frac{1}{\epsilon}d}|U|) && \text{[(II)]} \\ &\leq d(\text{opt}(I) - (1 - \frac{\beta}{\mu})\frac{\mu}{\beta d}\left(\frac{1}{4}\right)^{\frac{1}{\epsilon}d}d \cdot \text{frac}(I)) && \text{[Application of reduce]} \\ &\leq \left(1 - (1 - \frac{\beta}{\mu})\frac{\mu}{\beta d}\left(\frac{1}{4}\right)^{\frac{1}{\epsilon}d}\right) d \cdot \text{opt}(I) \\ &= \left(\frac{1}{\beta} - \left(\frac{d+1}{2\beta^2} - \frac{1}{\beta}\right)\left(\frac{1}{4}\right)^{\frac{d}{\epsilon}}\right) \beta d \cdot \text{opt}(I). \end{aligned}$$

Hence,  $\frac{|S|}{\text{opt}(I)} \leq \left(\frac{1}{\beta} - \left(\frac{d+1}{2\beta^2} - \frac{1}{\beta}\right)\left(\frac{1}{4}\right)^{\frac{d}{\epsilon}}\right) \beta d$ . So, because  $h(d, \epsilon) = \frac{1}{10d}\left(\frac{1}{4}\right)^{\frac{d}{\epsilon}}$ , to conclude that  $\frac{|S|}{\text{opt}(I)} \leq \beta d(1 - d \cdot h(d, \epsilon))$ , it suffices to prove that  $\frac{1}{\beta} - \left(\frac{d+1}{2\beta^2} - \frac{1}{\beta}\right)\left(\frac{1}{4}\right)^{\frac{d}{\epsilon}} \leq 1 - \frac{1}{10}\left(\frac{1}{4}\right)^{\frac{d}{\epsilon}}$ . For this, we have the following case distinction.

- Suppose that  $\beta \geq \frac{10}{9}$ . Then,  $\frac{1}{\beta} - \left(\frac{d+1}{2\beta^2} - \frac{1}{\beta}\right)\left(\frac{1}{4}\right)^{\frac{d}{\epsilon}} \leq \frac{1}{\beta} \leq \frac{9}{10} \leq 1 - \frac{1}{10}\left(\frac{1}{4}\right)^{\frac{d}{\epsilon}}$ .
- Suppose that  $\beta \leq \frac{10}{9}$ . As  $\frac{1}{\beta} \leq 1$ , it suffices to prove that  $\frac{d+1}{2\beta^2} - \frac{1}{\beta} \geq \frac{1}{10}$ , and as  $d \geq 2$ , it further suffices to prove that  $\frac{3}{2\beta^2} - \frac{1}{\beta} \geq \frac{1}{10}$ . Because  $\beta \leq \frac{10}{9}$ , we have that  $\frac{3}{2\beta^2} - \frac{1}{\beta} \geq \frac{3}{2\left(\frac{10}{9}\right)^2} - \frac{9}{10} \geq \frac{1}{10}$ .

This completes the proof.  $\blacktriangleleft$



Further, we argue that if  $S$  is computed in Step 5, then also it is a solution of this approximation ratio. Towards that, we have the following trivial claim.

▷ **Claim 30.** For all  $i = 1, 2, \dots, \tau$ , if the algorithm reaches iteration  $i$  and computes  $\mathcal{T}_i$ , then  $\mathcal{T}_i = \{S \in \mathcal{F} : S \subseteq U_i\}$ .

**Proof.** The proof is by induction on  $i$  (where we use  $i = 0$  as basis). When  $i = 0$ ,  $U_0 = U$  and  $\mathcal{T}_0 = \mathcal{F}$ , thus the claim trivially holds. Now, suppose that it holds for  $i - 1$ , and let us prove it for  $i$ . By the inductive hypothesis and the definition of  $\mathcal{T}_i$ , we have that

$$\mathcal{T}_i = \{S \in \mathcal{T}_{i-1} : S \subseteq U_i\} = \{S \in \mathcal{F} : S \subseteq U_i\}.$$

This completes the proof of the claim. ◀

We now present the promised claim.

▷ **Claim 31.** For all  $i = 1, 2, \dots, \tau$ , if the algorithm reaches Step 5, then  $S$  is a solution to  $I$  such that  $\frac{|S|}{\text{opt}(I)} \leq d(1 - d \cdot h(d, \epsilon))$ .

**Proof.** In this case,  $S = S_\tau \cup (U \setminus U_{\tau-1})$ . We first argue that  $S$  is a solution to  $I$ . To this end, notice that  $p_\tau = 1$ , so  $\mathcal{F}_\tau = \mathcal{T}_{\tau-1}$ . This means, by the correctness of the oracle, that  $S_\tau$  is a solution to  $(U_{\tau-1}, \mathcal{T}_{\tau-1})$ . That is, it has non-empty intersection with every set in  $\mathcal{T}_{\tau-1}$ . By Claim 30,  $\mathcal{T}_{\tau-1} = \{S \in \mathcal{F} : S \subseteq U_{\tau-1}\}$ , so  $U \setminus U_{\tau-1}$  has non-empty intersection with every set in  $\mathcal{F} \setminus \mathcal{T}_{\tau-1}$ . Thus,  $S$  has non-empty intersection with every set in  $\mathcal{F}$ , and is therefore a solution to  $I$ .

For the approximation ratio, note that the condition in Step 4d is false when  $i = \tau$ , else the algorithm would not have reached Step 5. Thus,

$$\begin{aligned} |S| &= |S_\tau| + |U| - |U_{\tau-1}| && [S_\tau \subseteq U_{\tau-1} \subseteq U] \\ &< |U| - (1 - \frac{\mu}{d})|U_{\tau-1}| && [\text{The condition in Step 4d is false}] \\ &\leq |U| - (1 - \frac{\mu}{d})(\frac{1}{4})^{\frac{1}{\epsilon}d}|U| && [\text{Claim 28}] \\ &= \left(1 - (1 - \frac{\mu}{d})(\frac{1}{4})^{\frac{1}{\epsilon}d}\right)|U| \\ &\leq \left(1 - (1 - \frac{\mu}{d})(\frac{1}{4})^{\frac{1}{\epsilon}d}\right)d \cdot \text{frac}(I) && [\text{Application of reduce}] \\ &\leq \left(1 - (1 - \frac{\mu}{d})(\frac{1}{4})^{\frac{1}{\epsilon}d}\right)d \cdot \text{opt}(I). \end{aligned}$$

Hence,  $\frac{|S|}{\text{opt}(I)} \leq \left(1 - (1 - \frac{\mu}{d})(\frac{1}{4})^{\frac{1}{\epsilon}d}\right)d$ . So, because  $h(d, \epsilon) = \frac{1}{10d}(\frac{1}{4})^{\frac{d}{\epsilon}}$ , to conclude that  $\frac{|S|}{\text{opt}(I)} \leq d(1 - d \cdot h(d, \epsilon))$ , it suffices to prove that  $(1 - \frac{\mu}{d})(\frac{1}{4})^{\frac{d}{\epsilon}} \geq \frac{1}{10}(\frac{1}{4})^{\frac{d}{\epsilon}}$ , which follows by substitution of  $\mu = \frac{d+1}{2}$ . This completes the proof of the claim. ◀

Lastly, we turn to conclude the proof of the theorem. First, because  $\text{frac}(I) \leq \text{frac}(\widehat{I})$  (by the correctness of **reduce**), Claim 27 implies that each call is of size as stated in the theorem. Further, this claim implies that with probability at least 9/10, the algorithm does not exit in Step 4b. Under the assumption that the algorithm does not exit in Step 4b, notice that Claims 29 and 31 ensure that  $S$  is a solution and that  $\frac{|S|}{\text{opt}(I)} \leq \beta d(1 - d \cdot h(d, \epsilon))$ . So, by Lemma 17 with  $\rho = \frac{d-1}{10}(\frac{1}{4})^{\frac{d}{\epsilon}}$ , at least one of the following conditions holds:

## 23:26 Lossy Kernelization for (Implicit) Hitting Set Problems

1.  $|\widehat{S}| - |S| \leq \rho \cdot \text{opt}(\widehat{I})$ , and hence  $|\widehat{S}| \leq |S| + \frac{d-1}{10} \left(\frac{1}{4}\right)^{\frac{d}{\epsilon}} \cdot \text{opt}(\widehat{I})$ . Then,

$$\begin{aligned} \frac{|\widehat{S}|}{\text{opt}(\widehat{I})} &\leq \frac{|S| + \frac{d-1}{10} \left(\frac{1}{4}\right)^{\frac{d}{\epsilon}} \cdot \text{opt}(\widehat{I})}{\text{opt}(\widehat{I})} \\ &\leq \frac{|S|}{\text{opt}(I)} + \frac{d-1}{10} \left(\frac{1}{4}\right)^{\frac{d}{\epsilon}} \\ &\leq \beta d(1 - d \cdot h(d, \epsilon)) + \frac{d-1}{10} \left(\frac{1}{4}\right)^{\frac{d}{\epsilon}} \\ &= \beta d(1 - d \cdot h(d, \epsilon)) + d(d-1) \cdot h(d, \epsilon) \\ &\leq \beta d(1 - h(d, \epsilon)). \end{aligned}$$

2.  $\frac{|\widehat{S}|}{\text{opt}(\widehat{I})} \leq d - \frac{\rho}{d-1} = d - \frac{1}{10} \left(\frac{1}{4}\right)^{\frac{d}{\epsilon}} = d(1 - h(d, \epsilon)) \leq \beta d(1 - h(d, \epsilon))$ .

So, in both cases we got that  $\frac{|\widehat{S}|}{\text{opt}(\widehat{I})} \leq \beta d(1 - h(d, \epsilon))$ . This completes the proof.  $\blacktriangleleft$

► **Corollary 32.** *For any fixed  $\epsilon > 0$ , the  $d$ -HITTING SET problem, parameterized by the optimum, admits a pure  $d(1 - h(d, \epsilon))$ -approximate (randomized) kernelization protocol with call size  $d \cdot \text{opt} + 2^{\frac{d}{\epsilon}} (d \cdot \text{opt})^{1+\epsilon}$  (where the number of sets is at most  $2^{\frac{d}{\epsilon}} (d \cdot \text{opt})^{1+\epsilon}$ ) where  $h(d, \epsilon) = \frac{1}{10d} \left(\frac{1}{4}\right)^{\frac{d}{\epsilon}}$  is a fixed positive constant that depends only on  $d, \epsilon$ .*

### 7.3 Relation Between a $(1 + \epsilon)$ -Approximate Kernelization Protocol for VERTEX COVER and the Ruzsa-Szemerédi Problem

We first present the following simple lemma.

► **Lemma 33.** *Let  $G$  be an  $n$ -vertex graph. Let  $r = r(n), t = t(n) \in \mathbb{N}$ . Let  $U_1, U_2, \dots, U_t \subseteq V(G)$  such that*

- *for all  $i \in \{1, 2, \dots, t\}$ ,  $G[U_i]$  has a matching  $M_i$  of size at least  $r$ , and*
- *for all distinct  $i, j \in \{1, 2, \dots, t\}$ ,  $E(G[U_i]) \cap E(G[U_j]) = \emptyset$ .*

*Then,  $G$  is a supergraph of an  $(r, t)$ -Ruzsa-Szemerédi graph.*

**Proof.** For all  $i \in \{1, 2, \dots, t\}$ , let  $M'_i$  be a matching in  $G[U_i]$  of size exactly  $r$ , and let  $U'_i \subseteq U_i$  be the vertices incident to at least one edge in  $M'_i$ . Let  $G'$  be the graph on vertex set  $\bigcup_{i=1}^t U'_i$  and edge set  $\bigcup_{i=1}^t M'_i$ . Notice that  $M'_1, M'_2, \dots, M'_t$  are matchings in  $G'$ . Because for all distinct  $i, j \in \{1, 2, \dots, t\}$ ,  $E(G[U_i]) \cap E(G[U_j]) = \emptyset$ , we have that  $M'_1, M'_2, \dots, M'_t$  are pairwise disjoint, and hence form a partition of  $E(G')$ . Lastly, we claim that for all  $i \in \{1, 2, \dots, t\}$ ,  $M'_i$  is an induced matching in  $G'$ . Targeting a contradiction, suppose that this is false for some  $i \in \{1, 2, \dots, t\}$ . So, there exist  $u, v \in U'_i$  such that  $\{u, v\} \notin M'_i$  but  $\{u, v\} \in M'_j$  for some  $j \in \{1, 2, \dots, t\} \setminus \{i\}$ . However, this means that  $\{u, v\} \in E(G[U_i]) \cap E(G[U_j])$ , which is a contradiction. This completes the proof.  $\blacktriangleleft$

We are now ready to present our main theorem, which follows the lines of the kernelization protocol presented in Section 7.1. Clearly, this result makes sense only for choices of  $c < \frac{1}{4}$  (so that the approximation ratio will be below 2) and when  $t = o(\sqrt{n})$ , preferably  $t = \mathcal{O}(n^{\frac{1}{2}-\lambda})$  for  $\lambda$  as close to  $1/2$  as possible, so that the volume will be  $\mathcal{O}(\text{opt}^{2-\lambda})$ . Further, if  $t$  is “sufficiently small” (depending on the desired number of rounds) whenever  $c$  is a fixed constant, this yields a  $(1 + \epsilon)$ -approximate kernelization protocol.

► **Theorem 34.** *Let  $0 < c < \frac{1}{4}$  be a fixed constant. For  $r = r(n) = cn$ , let  $t = t(n) = \gamma(r)$ .<sup>13</sup> Then, the VERTEX COVER problem, parameterized by the optimum, admits a  $(1 + 4c)$ -approximate (randomized) kernelization protocol with  $t + 1$  rounds and call size  $2\text{frac} + 2(t + 1)(2\text{frac})^{1.5}$  (where the number of edges is at most  $2(t + 1)(2\text{frac})^{1.5}$ ).*

**Proof.** We first describe the algorithm. To this end, consider some input  $\widehat{I} = \widehat{G}$ . Then:

1. Call the **reduce** procedure of the algorithm in Theorem 18 on  $I$  to obtain a new instance  $I = G$  where  $|V(G)| \leq 2\text{frac}(I)$ .
2. Initialize  $E_0 = \emptyset$ .
3. For  $i = 1, 2, \dots, t + 1$ :
  - a. Sample  $W_i$  from  $E(G)$  as follows: Insert each edge  $e \in E(G)$  to  $W_i$  independently at random with probability  $p = \frac{1}{(2\text{frac}(I))^{0.5}}$ .
  - b. If  $|W_i| > 2p|E(G)|$ , then let  $S$  be an arbitrary solution to  $I$ , and proceed directly to Step 5. [#Failure]
  - c. Call the oracle on  $G_i = G - E(G) \setminus (E_{i-1} \cup W_i)$ , and let  $S_i$  denote its output.
  - d. Let  $M_i$  be some maximal matching in  $G - S_i$ , and let  $T_i = E(G - S_i)$ .
  - e. If  $|M_i| < c|V(G)|$ , then let  $S = S_i \cup (\bigcup M_i)$ ,<sup>14</sup> and proceed directly to Step 5. [#Success]
  - f. If  $|T_i| > 2(2\text{frac}(I))^{1.5}$ , then let  $S$  be an arbitrary solution to  $I$ , and proceed directly to Step 5. [#Failure]
  - g. Let  $E_i = E_{i-1} \cup T_i$ .
4. Let  $S$  be an arbitrary solution to  $I$ , and proceed directly to Step 6. [#Never Reach]
5. Call the **lift** procedure of the algorithm in Theorem 18 on  $\widehat{I}, I, S$  to obtain a solution  $\widehat{S}$  to  $\widehat{I}$ . Output  $\widehat{S}$ .

Clearly, the algorithm runs in polynomial time, and only  $t + 1$  oracle calls are performed. Further, when we call the oracle on  $G_i$ , then  $|E(G_i)| \leq i \cdot 2p|E(G)| \leq 2(t + 1)(2\text{frac}(I))^{1.5}$  (due to **reduce**). Thus, each oracle call is performed on an instance with at most  $2\text{frac}(I)$  vertices (as  $|V(G)| \leq 2\text{frac}(I)$  due to **reduce**) and  $2(t + 1)(2\text{frac}(I))^{1.5}$  edges, and since  $\text{frac}(I) \leq \text{frac}(\widehat{I})$ , the statement in the lemma regarding the call size is satisfied.

Now, due to the correctness of **lift**, it remains to show that we compute a solution  $S$  to  $I$  that, with probability at least  $9/10$ , is a  $\beta(1 + 4c)$ -approximate solution to  $I$ , where  $\beta$  is the approximation ratio of the solutions returned by the oracle. Notice that if  $S$  is computed in the step marked “success”, say, at some iteration  $i$ , then clearly  $|S| = |S_i| + 2|M_i| < \beta\text{opt}(I) + 2c|V(G)| \leq \beta\text{opt}(I) + 4c\text{opt}(I) \leq \beta(1 + 4c)\text{opt}(I)$ . Moreover, since  $M_i$  is a maximal matching, every edge in  $G$  that is not incident to  $S_i$  must share an endpoint with at least one edge in  $M_i$ . So,  $S$  is then a solution to  $I$ . Thus, it suffices to show that with probability at least  $9/10$ ,  $S$  is computed in the step marked by “success”.

Just like in the proof of Theorem 22, we can show that the probability that the conditions in the steps marked by “failure” are not satisfied with probability at least  $9/10$ . So, it remains to show that we never reach Step 4. Targeting a contradiction, suppose that we reach this step. For all  $i \in \{1, 2, \dots, t + 1\}$ , let  $U_i = V(G) \setminus S_i$ . Then,  $G[U_i]$  has a matching of size at least  $r = c|V(G)|$  (that is  $M_i$ ). Further, for all  $1 \leq i < j \leq t + 1$ , because  $S_j$  is a vertex cover of  $G_j$  and  $E(G[U_i]) = T_i \subseteq E(G_j)$ ,  $E(G[U_i]) \cap E(G[U_j]) = \emptyset$ . By Lemma 33, this means

<sup>13</sup>That is,  $t$  is the maximum value (as a function of  $n$ ) such that there exists a  $(r, t)$ -Ruzsa-Szemerédi graph where  $r = cn$  (see Definition 8 and the discussion below it).

<sup>14</sup>That is,  $S$  is the set that contains every vertex in  $S_i$  as well every vertex incident to an edge in  $M_i$ .

that  $G$  is a supergraph of an  $(r, t + 1)$ -Ruzsa-Szemerédi graph. However, this contradicts the definition of  $t$ . Thus, the proof is complete. ◀

► **Corollary 35.** *Let  $0 < c < \frac{1}{4}$  be a fixed constant. For  $r = r(n) = cn$ , let  $t = t(n) = \gamma(r)$ . Then, the VERTEX COVER problem, parameterized by the optimum, admits a  $(1 + 4c)$ -approximate (randomized) kernelization protocol with  $t + 1$  rounds and call size  $2\text{opt} + 2(t + 1)(2\text{opt})^{1.5}$  (where the number of edges is at most  $2(t + 1)(2\text{opt})^{1.5}$ ).*

## 8 $(1 + \epsilon)$ -Approximate Linear-Vertex Kernels for Implicit 3-HITTING SET Problems

In this section, we present lossy kernels for two well-known implicit 3-HS problems, called CLUSTER VERTEX DELETION, and FEEDBACK VERTEX SET IN TOURNAMENTS. In CLUSTER VERTEX DELETION, given a graph  $G$ , the task is to compute a minimum-sized subset  $S \subseteq V(G)$  such that  $G - S$  is a cluster graph. In FEEDBACK VERTEX SET IN TOURNAMENTS, given a tournament  $G$ , the task is to compute a minimum-sized subset  $S \subseteq V(G)$  such that  $G - S$  is acyclic. We attain a linear number of vertices at an approximation cost of only  $(1 + \epsilon)$  rather than 2 as is given for 3-HS in Section 6. Notably, both our algorithms follow similar lines, and we believe that the approach underlying their common parts may be useful when dealing also with other hitting and packing problems of constant-sized objects. In particular, in both algorithms we first “reveal modules” using essentially the same type of marking scheme, which yields a lossy rule, and afterwards we shrink the size of these modules using yet another rule that, unlike the first one, is problem-specific.

### 8.1 CLUSTER VERTEX DELETION

Our lossy kernel will use Theorem 12 and consist of two rules, one lossy rule and one exact rule, each to be applied only once. The first rule (to which we will refer as the “module revealing operation”) will ensure that all unmarked vertices in a clique (in some subgraph of the original graph, obtained by the removal of an approximate solution) form a module and furthermore that certain vertices among those removed have neighbors in only one of them, and the second one (“module shrinkage operation”) will reduce the size of each such module. For simplicity, we will actually merge them together to a single rule. We begin by reminding that CLUSTER VERTEX DELETION can be interpreted as a special case of 3-HITTING SET:

► **Proposition 36** ([8]). *A graph  $G$  is a cluster graph if and only if it does not have any induced  $P_3$ .*

► **Definition 37.** *Given a graph  $G$ , define the 3-HITTING SET instance corresponding to  $G$  by  $\text{HS}(G) = (V(G), \{\{u, v, w\} \subseteq V(G) : G[\{u, v, w\}] \text{ is an induced } P_3\})$ .*

► **Corollary 38.** *Let  $G$  be a graph. Then, a subset  $S \subseteq V(G)$  is a solution to the 3-HITTING SET instance corresponding to  $G$  if and only if  $G - S$  is a cluster graph.*

To perform the module revealing operation, given a graph  $G$ , we will be working with an optimal solution  $\alpha$  to the classic LP of the 3-HITTING SET instance corresponding to  $G$ . The approximate solution we will be working with will be the support of  $\alpha$ . For the sake of clarity, we slightly abuse notation and use vertices to refer both to vertices and to the variables corresponding to them, as well as use an instance of CLUSTER VERTEX DELETION to refer also to the 3-HITTING SET instance corresponding to it when no confusion arises. We first show that the cliques in  $G - \text{support}(\alpha)$  are already modules in  $G - \alpha^{-1}(1)$  (i.e., in

the graph obtained by removing all vertices to which  $\alpha$  assigns 1). Thus, to reveal modules, we will only deal with vertices in  $\alpha^{-1}(1)$ .

► **Lemma 39.** *Let  $G$  be a graph, and let  $\alpha$  be a solution to the 3-HITTING SET instance corresponding to  $G$ . Let  $C$  be a clique in  $G - \text{support}(\alpha)$ . Then,  $V(C)$  is a module in  $G - \alpha^{-1}(1)$ .*

**Proof.** First, notice that as  $\alpha$  is optimal, it does not assign values greater than 1. Targeting a contradiction, suppose that  $V(C)$  is not a module in  $G - \alpha^{-1}(1)$ . So, there exist vertices  $v \in \text{support}(\alpha) \setminus \alpha^{-1}(1)$  and  $u, w \in V(C)$  such that  $\{u, v\} \in E(G)$  and  $\{w, v\} \notin E(G)$ . Then, as  $\{u, w\} \in E(G)$  (since  $C$  is a clique),  $G[\{v, u, w\}]$  is an induced  $P_3$ . However,  $\alpha$  assigns 0 to the variables of  $u$  and  $w$ , and a value smaller than 1 to the variable of  $v$ , while the sum of these variables should be at least 1 (because  $\alpha$  is a solution). Thus, we have reached a contradiction. ◀

To deal with the vertices in  $\alpha^{-1}(1)$ , we now define a marking procedure that will be used by the first (implicit) rule.

► **Definition 40.** *Given  $0 < \epsilon < 1$ , a graph  $G$  and an optimal solution  $\alpha$  to the 3-HITTING SET instance corresponding to  $G$ ,  $\text{Marking}(\epsilon, G, \alpha)$  is defined as follows.*

1. For every vertex  $v \in \alpha^{-1}(1)$ , initialize  $\text{mark}(v) = \emptyset$ .
2. For every vertex  $v \in \alpha^{-1}(1)$ :
  - a. Let  $H_v$  be the graph defined as follows:  $V(H_v) = V(G) \setminus (\text{support}(\alpha) \cup (\bigcup_{u \in \alpha^{-1}(1)} \text{mark}(u)))$ , and  $E(H_v) = \{\{w, r\} \subseteq V(H_v) : G[\{v, w, r\}] \text{ is an induced } P_3\}$ .
  - b. Compute a maximal matching  $\mu_v$  in  $H_v$ .<sup>15</sup>
  - c. If  $|\mu_v| > \frac{1}{\epsilon}$ , then let  $\nu_v$  be some (arbitrary) subset of  $\mu_v$  of size exactly  $\frac{1}{\epsilon}$ , and otherwise let  $\nu_v = \mu_v$ . Let  $\text{mark}(v) = \bigcup \nu_v$  (i.e.,  $\text{mark}(v)$  is the set of vertices incident to edges in  $\nu_v$ ).
3. For every vertex  $v \in \alpha^{-1}(1)$ , output  $\text{mark}(v)$ . Moreover, output  $D = \{v \in \alpha^{-1}(1) : |\text{mark}(v)| = \frac{1}{\epsilon}\}$ .

We now prove that when all marked vertices are removed, the remainders of the cliques form modules in  $G - D$ .

► **Lemma 41.** *Given  $0 < \epsilon < 1$ , a graph  $G$  and an optimal solution  $\alpha$  to the 3-HITTING SET instance corresponding to  $G$ , let  $\{\text{mark}(v)\}_{v \in \alpha^{-1}(1)}$ ,  $D$  be the output of  $\text{Marking}(\epsilon, G, \alpha)$ . Then, the vertex set of every clique  $C$  in  $G - (\text{support}(\alpha) \cup (\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v)))$  is a module in  $G - D$ .*

**Proof.** Consider some clique  $C$  in  $G - (\text{support}(\alpha) \cup (\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v)))$ . Clearly, every vertex in  $G - \text{support}(\alpha)$  is adjacent to either all vertices in  $C$  (when they belong to the same clique in  $G - \text{support}(\alpha)$ ) or to none (when they belong to different cliques). Further, due to Lemma 39, every vertex in  $\text{support}(\alpha) \setminus \alpha^{-1}(1)$  also has this property. So, it remains to prove that every vertex in  $\alpha^{-1}(1) \setminus D$  also has this property. To this end, consider some vertex  $v \in \alpha^{-1}(1) \setminus D$ . Targeting a contradiction, suppose that there exist vertices  $u, w \in V(C)$  such that  $\{u, v\} \in E(G)$  but  $\{w, v\} \notin E(G)$ . As  $V(C) \cap (\bigcup_{v' \in \alpha^{-1}(1)} \text{mark}(v')) = \emptyset$ ,  $H_v$  (in Definition 40) contained the edge  $\{u, w\}$ . Moreover, neither  $u$  nor  $w$  was inserted into  $\text{mark}(v)$  and hence, as  $\nu_v = \mu_v$  (because  $v \notin D$ ), none of them is incident to an edge in  $\mu_v$ .

<sup>15</sup>For example, by greedily picking edges so that the collection of edges remains a matching as long as it possible.

## 23:30 Lossy Kernelization for (Implicit) Hitting Set Problems

However, this contradicts that  $\mu_v$  is a maximal matching, as we can insert  $\{u, w\}$  to it and it would remain a matching.  $\blacktriangleleft$

We now argue that every optimal solution contains all of the vertices of  $D$  except of an  $\epsilon$ -fraction of the optimum, and hence it is not “costly” to seek only solutions that contain  $D$ .

► **Lemma 42.** *Let  $I = G$  be an instance of CLUSTER VERTEX DELETION. Given  $0 < \epsilon < 1$ ,  $G$  and an optimal solution  $\alpha$  to the 3-HITTING SET instance corresponding to  $G$ , let  $\{\text{mark}(v)\}_{v \in \alpha^{-1}(1), D}$  be the output of  $\text{Marking}(\epsilon, G, \alpha)$ . Let  $S^*$  be an optimal solution to  $I$ . Then,  $|D \setminus S^*| \leq \epsilon \text{opt}(I)$ .*

**Proof.** Consider some vertex  $v \in D$ . Notice that  $v$  together with any edge in  $\nu_v$  form an induced  $P_3$  in  $G$ . Thus, if  $v \notin S^*$ , then from every edge in  $\nu_v$ , at least one vertex must belong to  $S^*$ . As  $\nu_v$  is a matching, and its size is  $\frac{1}{\epsilon}$ , this means that  $S^*$  has to contain at least  $\frac{1}{\epsilon}$  vertices from  $\text{mark}(v)$ . As the sets assigned by  $\text{mark}$  are pairwise disjoint, we have that  $|D \setminus S^*|$  can be of size at most  $\epsilon|S^*| = \epsilon \text{opt}(I)$ .  $\blacktriangleleft$

Intuitively, the arguments above naturally give rise to a reduction rule that deletes  $D$ . However, a minor technicality arises—that is, we will need to transmit  $\alpha$  and the marked sets to the reduced instance in order for our next arguments to work, which, when complying with necessary formalities, requires to define an annotated version of the problem. We avoid this by merging the rule implicitly in our main rule later, which simplifies the presentation.

Before we proceed to shrink the size of the modules, we argue that every vertex outside them (except for those in  $D$ ) has neighbors in at most one of them.

► **Lemma 43.** *Given  $0 < \epsilon < 1$ , a graph  $G$  and an optimal solution  $\alpha$  to the 3-HITTING SET instance corresponding to  $G$ , let  $\{\text{mark}(v)\}_{v \in \alpha^{-1}(1), D}$  be the output of  $\text{Marking}(\epsilon, G, \alpha)$ . Then, for every vertex  $v \in \text{support}(\alpha) \setminus D$ ,  $N_G(v) \setminus (\text{support}(\alpha) \cup (\bigcup_{v' \in \alpha^{-1}(1)} \text{mark}(v')))$  is either empty or equals the vertex set of exactly one clique  $C$  in  $G - (\text{support}(\alpha) \cup (\bigcup_{v' \in \alpha^{-1}(1)} \text{mark}(v')))$ .*

**Proof.** Consider some vertex  $v \in \text{support}(\alpha) \setminus D$ . Targeting a contradiction, suppose that the lemma is false with respect to  $v$ . Due to Lemma 41, this necessarily means that there exist two distinct cliques  $C, C'$  in  $G - (\text{support}(\alpha) \cup (\bigcup_{v' \in \alpha^{-1}(1)} \text{mark}(v')))$  such that  $v$  has neighbors in both. So, let  $u \in V(C)$  and  $w \in V(C')$  be such that  $\{u, v\}, \{w, v\} \in E(G)$ . Observe that as  $u, w$  belong to different cliques,  $\{u, w\} \notin E(G)$ . As  $(V(C) \cup V(C')) \cap (\bigcup_{v' \in \alpha^{-1}(1)} \text{mark}(v')) = \emptyset$ ,  $H_v$  (in Definition 40) contained the edge  $\{u, w\}$ . Moreover, neither  $u$  nor  $w$  was inserted into  $\text{mark}(v)$  and hence, as  $\nu_v = \mu_v$  (because  $v \notin D$ ), none of them is incident to an edge in  $\mu_v$ . However, this contradicts that  $\mu_v$  is a maximal matching, as we can insert  $\{u, w\}$  to it and it would remain a matching.  $\blacktriangleleft$

We now proceed to shrink the size of the modules we have just revealed. We note that this part is, in fact, exact (i.e. 1-approximate). For this purpose, we start with the following observation.

► **Lemma 44.** *Let  $I = G$  be an instance of CLUSTER VERTEX DELETION. Let  $T \subseteq V(G)$  be a module in  $G$  such that  $G[T]$  is a clique. Then, any induced  $P_3$  in  $G$  that contains at least one vertex from  $T$ , contains exactly one vertex from  $T$  and at least one vertex from  $N_G(T)$ .*

**Proof.** Let  $P$  be an induced  $P_3$  in  $G$  that contains at least one vertex from  $T$ . Clearly,  $P$  cannot contain three from  $T$  as  $T$  induces a clique. Further, if it contains exactly two vertices from  $T$ , then they must be adjacent in  $P$  as  $T$  induces a clique, but then the third vertex in  $P$  will be a neighbor of one but non-neighbor of the other, which is a contradiction because

$T$  is a module. So,  $P$  contains exactly one vertex from  $T$ , which also implies that it must contain at least one vertex from  $N_G(T)$ . ◀

We use the above lemma to give a simple lemma that says that if at least one vertex of a module that induces a clique is deleted, then unless that deletion is unnecessary, all of its vertices are deleted.

► **Lemma 45.** *Let  $I = G$  be an instance of CLUSTER VERTEX DELETION. Let  $T \subseteq V(G)$  be a module in  $G$  such that  $G[T]$  is a clique. Then, for every solution  $S$  to  $I$ , either  $T \subseteq S$  or  $S \setminus T$  is also a solution to  $I$ .*

**Proof.** Consider a solution  $S$  to  $I$  such that  $T$  is not contained in  $S$ . So, there exists some vertex  $v \in T \setminus S$ . We claim that  $S \setminus T$  is also a solution to  $I$ , which will complete the proof. Targeting a contradiction, suppose that  $S \setminus T$  is not a solution to  $I$ , thus there exists an induced  $P_3$ , say,  $P$ , in  $G - (S \setminus T)$ . Observe that  $P$  contains at least one vertex from  $T$ , as  $S$  is a solution. By Lemma 44, this means that it must contain exactly one vertex from  $T$ . So, let  $u$  denote the only vertex in  $T \cap V(P)$ . As  $T$  is a module, by replacing  $u$  by  $v$ , we obtain yet another induced  $P_3$ , and this one belongs to  $G - S$ . This is a contradiction as  $S$  is a solution to  $I$ . ◀

Further, given a solution that contains  $D$ , we may exchange a clique by its neighborhood and still have a solution, as stated below.

► **Lemma 46.** *Let  $I = G$  be an instance of CLUSTER VERTEX DELETION. Given  $0 < \epsilon < 1$ ,  $G$  and an optimal solution  $\alpha$  to the 3-HITTING SET instance corresponding to  $G$ , let  $\{\text{mark}(v)\}_{v \in \alpha^{-1}(1)}, D$  be the output of  $\text{Marking}(\epsilon, G, \alpha)$ . Let  $S^*$  be a solution to  $I$ . Then, for every clique  $C$  in  $G - (\text{support}(\alpha) \cup (\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v)))$ , we have that  $(S^* \setminus V(C)) \cup N_G(V(C))$  is a solution to  $I$ .*

**Proof.** Consider some clique  $C$  in  $G - (\text{support}(\alpha) \cup (\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v)))$ . As  $S^*$  is a solution to  $I$ , every induced  $P_3$  in  $G - (S^* \setminus V(C))$  must contain at least one vertex from  $C$ . However, in  $G - N_G(V(C))$ ,  $C$  is an isolated clique and hence there exists no induced  $P_3$  that contains at least one of its vertices. So,  $G - (S^* \setminus V(C)) \cup N_G(V(C))$  does not have any induced  $P_3$ , and hence  $(S^* \setminus V(C)) \cup N_G(V(C))$  is a solution to  $I$ . ◀

So, Lemmata 45 and 46 naturally give rise to a reduction operation where each clique whose size is larger than the size of its neighborhood is shrunk to the size of its neighborhood.

Finally, we devise our merged reduction rule:

► **Definition 47.** *The CLUSTER VERTEX DELETION reduction rule is defined as follows:*

- **reduce:** *Let  $I = G$  be an instance of CLUSTER VERTEX DELETION. Use the algorithm in Proposition 9 to compute an optimal solution  $\alpha$  to the classic LP corresponding to it (Definitions 11 and 37). Let  $\{\text{mark}(v)\}_{v \in \alpha^{-1}(1)}, D$  be the output of  $\text{Marking}(\epsilon, G, \alpha)$ . Output  $I' = G'$  where  $G'$  is obtained from  $G - D$  as follows: for every clique  $C$  in  $G - (\text{support}(\alpha) \cup (\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v)))$ , delete (arbitrarily chosen)  $\max\{|V(C)| - |N_{G-D}(V(C))|\}$  vertices from  $C$ .*
- **lift:** *Given  $I, I'$  and a solution  $S'$  to  $I'$ , output  $S = D \cup S''$  where  $S''$  is obtained from  $S'$  as follows: for every clique  $C$  in  $G - (\text{support}(\alpha) \cup (\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v)))$  such that  $V(C)$  is not contained in  $G'$  and  $V(C) \subseteq S'$ , remove  $V(C) \cap S'$  and add  $N_G(V(C))$  instead.*

Before we prove our main theorem, we present a simple lemma that will help us derive a tighter bound on the number of vertices in the output graph.

## 23:32 Lossy Kernelization for (Implicit) Hitting Set Problems

► **Lemma 48.** *Let  $I = G$  be an instance of CLUSTER VERTEX DELETION, and let  $\alpha$  be a solution to the 3-HITTING SET instance corresponding to  $G$ . Then,  $|\text{support}(\alpha)| \leq 3\text{frac}(I) - 2|\alpha^{-1}(1)|$ .*

**Proof.** Let  $\beta$  denote the restriction of  $\alpha$  to  $G - \alpha^{-1}(1)$ . Observe that  $\beta$  is a solution to the classic LP of the 3-HITTING SET instance corresponding to  $G$  and its value is  $\text{frac}(I) - |\alpha^{-1}(1)|$ . Further, consider some solution  $\beta'$  to the classic LP of the 3-HITTING SET instance corresponding to  $G' = G - \alpha^{-1}(1)$ , thought of as an instance  $I'$  of CLUSTER VERTEX DELETION. Then, by extending  $\beta'$  to assign 1 to each vertex in  $\alpha^{-1}(1)$ , we obtain a solution  $\alpha'$  to the classic LP of the 3-HITTING SET instance corresponding to  $G$ . As  $\alpha$  is optimal, its value is at most that of  $\alpha'$ . Thus, it must hold that the value of  $\beta'$  is at least that of  $\beta$ . Since the choice of  $\beta'$  was arbitrary, this implies that  $\beta$  is optimal. Hence, by Theorem 12,

$$|\text{support}(\beta)| \leq 3 \cdot \text{frac}(I') = 3 \cdot (\text{frac}(I) - |\alpha^{-1}(1)|).$$

Thus, we have that

$$|\text{support}(\alpha)| = |\text{support}(\beta)| + |\alpha^{-1}(1)| \leq 3\text{frac}(I) - 2|\alpha^{-1}(1)|.$$

This completes the proof. ◀

Based on Lemmata 41, 42, 43, 45, 46 and 48, we are now ready to prove the main theorem of this subsection.

► **Theorem 49.** *Let  $0 < \epsilon < 1$ . The CLUSTER VERTEX DELETION problem, parameterized by the fractional optimum of the classic LP, admits a  $(1 + \epsilon)$ -approximate  $\max(6, \frac{4}{\epsilon}) \cdot \text{frac}$ -vertex kernel.*

**Proof.** Our lossy kernelization algorithm consists only of the CLUSTER VERTEX DELETION reduction rule. Clearly, it runs in polynomial time.

First, we consider the number of vertices in the output graph  $G'$  of **reduce**. By Lemma 48,  $|\text{support}(\alpha)| \leq 3 \cdot \text{frac}(I) - 2|\alpha^{-1}(1)|$  (I). Moreover,  $|\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v)| = 2|\bigcup_{v \in \alpha^{-1}(1)} \nu_v| \leq \frac{2}{\epsilon}|\alpha^{-1}(1)|$  (II). By the definition of the reduction rule, for every clique  $C$  in  $G' - (\text{support}(\alpha) \cup (\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v)))$ ,  $|V(C)| \leq |N_{G'}(V(C))|$ . Additionally, by Lemma 43, the neighborhood sets of these cliques are pairwise vertex disjoint. This implies that, altogether, these cliques contain at most  $|\text{support}(\alpha) \setminus \alpha^{-1}(1) \cup (\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v))|$  vertices. Thus, because the vertex set of  $G'$  consists only of these cliques and of  $\text{support}(\alpha) \setminus \alpha^{-1}(1) \cup (\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v))$ , we conclude that

$$\begin{aligned} |V(G')| &\leq 2|\text{support}(\alpha)| - 2|\alpha^{-1}(1)| + 2|\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v)| && \text{[Last two sentences]} \\ &\leq 2|\text{support}(\alpha)| - 2|\alpha^{-1}(1)| + \frac{4}{\epsilon}|\alpha^{-1}(1)| && \text{[(II)]} \\ &\leq 6 \cdot \text{frac}(I) - 4|\alpha^{-1}(1)| - 2|\alpha^{-1}(1)| + \frac{4}{\epsilon}|\alpha^{-1}(1)| && \text{[(I)]} \\ &= 6 \cdot \text{frac}(I) + (\frac{4}{\epsilon} - 6)|\alpha^{-1}(1)| \\ &\leq \max(6, \frac{4}{\epsilon}) \cdot \text{frac}(I) && [|\alpha^{-1}(1)| \leq \text{frac}(I)]. \end{aligned}$$

We turn to prove that **lift** returns a solution having the desired approximation ratio. To this end, suppose that it is given  $I, I', S'$  where  $S'$  is a solution to  $I'$ . First, notice that  $S' \cup (V(G) \setminus V(G'))$  is a solution to  $I$ . Thus, because  $S$  can be obtained from  $S' \cup (V(G) \setminus V(G'))$  by doing deletion and exchange operations as described in Lemmata 45 and 46, these lemmata imply that  $S$  is a solution to  $I$ .

Now, we consider the approximation ratio of  $S$ . For this, on the one hand, let  $\widehat{S}$  be a minimal solution contained in  $S'$ . By Lemmata 41 and 45, for every clique  $C'$  in



$G' - (\text{support}(\alpha) \cup (\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v)))$ , either  $V(C') \subseteq \widehat{S}$  or  $V(C') \cap \widehat{S} = \emptyset$ . So, because every clique  $C$  in  $G - (\text{support}(\alpha) \cup (\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v)))$  such that  $V(C)$  is not contained in  $G'$  satisfies that  $|V(C) \cap V(G')| = |N_{G-D}(V(C))|$ , we know that  $|S \setminus D| = |\widehat{S}| \leq |S'|$ . Hence, (i)  $|S| \leq |S'| + |D|$ . On the other hand, let  $S^*$  be an optimal solution to  $I$ . By Lemma 42,  $|D \setminus S^*| \leq \epsilon \text{opt}(I)$ . Thus,  $|S^* \setminus D| = \text{opt}(I) - |S^* \cap D| \geq \text{opt}(I) - (|D| - \epsilon \text{opt}(I)) = (1 + \epsilon)\text{opt}(I) - |D|$ . Further, as  $S^* \cap V(G')$ , which is a subset of  $S^* \setminus D$ , is a solution to  $I'$ , we have that  $\text{opt}(I') \leq |S^* \setminus D|$ , and hence (ii)  $\text{opt}(I') \leq (1 + \epsilon)\text{opt}(I) - |D|$ . From (i) and (ii), we conclude that

$$\frac{|S|}{\text{opt}(I)} \leq (1 + \epsilon) \frac{|S'| + |D|}{\text{opt}(I') + |D|} \leq (1 + \epsilon) \max\left\{\frac{|S'|}{\text{opt}(I')}, \frac{|D|}{|D|}\right\} = (1 + \epsilon) \frac{|S'|}{\text{opt}(I')}.$$

Here, the last inequality follows from Proposition 6. This completes the proof.  $\blacktriangleleft$

► **Corollary 50.** *Let  $0 < \epsilon < 1$ . The CLUSTER VERTEX DELETION problem, parameterized by the optimum, admits a  $(1 + \epsilon)$ -approximate  $\max(6, \frac{4}{\epsilon}) \cdot \text{opt}$ -vertex kernel.*

Due to Lemma 2, we also have the following corollary of Theorem 49.

► **Corollary 51.** *Let  $0 < \epsilon < 1$ . The CLUSTER VERTEX DELETION problem, parameterized by a bound  $k$  on the solution size, admits a  $(1 + \epsilon)$ -approximate  $\max(\frac{6}{1+\epsilon}, \frac{4}{(1+\epsilon)\epsilon}) \cdot (k + 1)$ -vertex kernel.*

## 8.2 FEEDBACK VERTEX SET IN TOURNAMENTS

Our lossy kernel will use Theorem 12 and consist of two lossy rules, each to be applied only once. The first rule (to which we will refer as the “module revealing operation”) will ensure that, with respect to some linear order on the vertices not in some approximate solution, all consecutive unmarked vertices between two marked vertices form a module and furthermore that there is an essentially unique position to place each vertex (including those in the approximate solution) between them, and the second one (“module shrinkage operation”) will reduce the size of each such module. For simplicity, we will actually merge them together to a single rule. We begin by reminding that FEEDBACK VERTEX SET IN TOURNAMENTS can be interpreted as a special case of 3-HITTING SET:

► **Proposition 52** ([8]). *A tournament  $G$  is acyclic if and only if it does not have any triangle (i.e., a directed cycle on three vertices).*

► **Definition 53.** *Given a tournament  $G$ , define the 3-HITTING SET instance corresponding to  $G$  by  $\text{HS}(G) = (V(G), \{\{u, v, w\} \subseteq V(G) : G[\{u, v, w\}] \text{ is triangle}\})$ .*

► **Corollary 54.** *Let  $G$  be a tournament. Then, a subset  $S \subseteq V(G)$  is a solution to the 3-HITTING SET instance corresponding to  $G$  if and only if  $G - S$  is acyclic.*

To perform the module revealing operation, given a graph  $G$ , we will be working with an optimal solution  $\alpha$  to the classic LP of the 3-HITTING SET instance corresponding to  $G$ . The approximate solution we will be working with will be the support of  $\alpha$ . For the sake of clarity, we slightly abuse notation and use vertices to refer both to vertices and to the variables corresponding to them, as well as use an instance of FEEDBACK VERTEX SET IN TOURNAMENTS to refer also to the 3-HITTING SET instance corresponding to it when no confusion arises. We will use the following well-known characterization of acyclic digraphs.

► **Proposition 55** (Folklore). *A digraph  $G$  is acyclic if and only if there exists a linear order  $<$  on  $V(G)$  such that for every arc  $(u, v) \in E(G)$ ,  $u < v$ . Moreover, given an acyclic digraph  $G$ , such an order is computable in linear time, and if  $G$  is a tournament, then this order is unique.*

This gives rise to the following definition.

► **Definition 56.** *Let  $G$  be a tournament, and let  $\alpha$  be a solution to the 3-HITTING SET instance corresponding to  $G$ . Then, the linear order induced by  $\alpha$ , denoted  $<_\alpha$ , is the unique linear ordering of  $V(G) \setminus \text{support}(\alpha)$  such that for every arc  $(u, v) \in E(G - \text{support}(\alpha))$ ,  $u <_\alpha v$ . We say that two vertices  $u, v \in V(G - \text{support}(\alpha))$  are consecutive in  $<_\alpha$  if  $u <_\alpha v$  and there is no vertex  $w \in V(G - \text{support}(\alpha))$  such that  $u <_\alpha w <_\alpha v$ ; then,  $u$  is called the successor of  $v$ , and  $v$  is called the predecessor of  $u$ .*

We further define the notion of a *position* based on this order.

► **Definition 57.** *Let  $G$  be a tournament, and let  $\alpha$  be a solution to the 3-HITTING SET instance corresponding to  $G$ . Let  $M \subseteq V(G) \setminus \text{support}(\alpha)$ . Then, a vertex  $v \in \text{support}(\alpha) \cup M$   $M$ -fits  $<_\alpha$  if one of the following conditions holds.*

- *For all  $u \in V(G) \setminus (\text{support}(\alpha) \cup M)$ ,  $(v, u) \in E(G)$ . In this case, we say that  $v$  has 0-position (with respect to  $M$ ).*
- *There exists  $u \in V(G) \setminus (\text{support}(\alpha) \cup M)$  such that for every  $r \in V(G) \setminus (\text{support}(\alpha) \cup M)$  where  $r \leq_\alpha u$ ,  $(r, v) \in E(G)$ , and for every  $r \in V(G) \setminus (\text{support}(\alpha) \cup M)$  where  $r >_\alpha u$ ,  $(v, r) \in E(G)$ . In this case, we say that  $v$  has  $u$ -position (with respect to  $M$ ).*

We suppose that a 0-position is the lowest possible, that is,  $0 <_\alpha u$  for all  $u \in V(G) \setminus (\text{support}(\alpha) \cup M)$ . The following observations are immediate.

► **Observation 58.** *Let  $G$  be a tournament, and let  $\alpha$  be a solution to the 3-HITTING SET instance corresponding to  $G$ . Let  $M \subseteq V(G) \setminus \text{support}(\alpha)$ . Let  $v \in \text{support}(\alpha) \cup M$  be a vertex that  $M$ -fits  $<_\alpha$ . Then, there exists exactly one element  $u \in \{0\} \cup (V(G) \setminus (\text{support}(\alpha) \cup M))$  such that  $v$  has  $u$ -position.*

► **Observation 59.** *Let  $G$  be a tournament, and let  $\alpha$  be a solution to the 3-HITTING SET instance corresponding to  $G$ . Let  $M \subseteq V(G) \setminus \text{support}(\alpha)$ . Then, every vertex in  $M$   $M$ -fits  $<_\alpha$ .*

We first show that the vertices in  $\text{support}(\alpha) \setminus \alpha^{-1}(1)$  already  $\emptyset$ -fit  $<_\alpha$  (so, they also  $M$ -fit  $<_\alpha$  with respect to any  $M \subseteq V(G) \setminus \text{support}(\alpha)$ ). Thus, to reveal modules that give rise to unique positions, we will only deal with vertices in  $\alpha^{-1}(1)$ .

► **Lemma 60.** *Let  $G$  be a tournament, and let  $\alpha$  be a solution to the 3-HITTING SET instance corresponding to  $G$ . Let  $v \in \text{support}(\alpha) \setminus \alpha^{-1}(1)$ . Then,  $v$   $\emptyset$ -fits  $<_\alpha$ .*

**Proof.** First, notice that as  $\alpha$  is optimal, it does not assign values greater than 1. Thus,  $\alpha(x_v) < 1$ . So,  $G - (\text{support}(\alpha) \setminus \{v\})$  does not have a triangle, else the sum of the variables of its vertices will be less than 1, contradicting that  $\alpha$  is a solution. By Corollary 54 and Proposition 55, this means that  $G - (\text{support}(\alpha) \setminus \{v\})$  admits a unique linear order  $<$  such that for every arc  $(x, y) \in E(G - (\text{support}(\alpha) \setminus \{v\}))$ ,  $x < y$ , and its restriction to  $G - \text{support}(\alpha)$  must equal  $<_\alpha$ . This directly implies the lemma, where if  $v$  is first in  $<$  then it has 0-position, and otherwise it has  $u$ -position where  $u$  is its predecessor in  $<$ . ◀

To deal with the vertices in  $\alpha^{-1}(1)$ , we define the following marking procedure.

► **Definition 61.** Given  $0 < \delta < 1$ , a tournament  $G$  and an optimal solution  $\alpha$  to the 3-HITTING SET instance corresponding to  $G$ ,  $\text{Marking}(\delta, G, \alpha)$  is defined as follows.

1. For every vertex  $v \in \alpha^{-1}(1)$ , initialize  $\text{mark}(v) = \emptyset$ .
2. For every vertex  $v \in \alpha^{-1}(1)$ :
  - a. Define the graph  $H_v$  as follows:  $V(H_v) = V(G) \setminus (\text{support}(\alpha) \cup (\bigcup_{u \in \alpha^{-1}(1)} \text{mark}(u)))$ , and  $E(H_v) = \{\{w, r\} \subseteq V(H_v) : G[\{v, w, r\}] \text{ is a triangle}\}$ .
  - b. Compute a maximal matching  $\mu_v$  in  $H_v$ .
  - c. If  $|\mu_v| > \frac{1}{\delta}$ , then let  $\nu_v$  be some (arbitrary) subset of  $\mu_v$  of size exactly  $\frac{1}{\delta}$ , and otherwise let  $\nu_v = \mu_v$ . Let  $\text{mark}(v) = \bigcup \nu_v$  (i.e.,  $\text{mark}(v)$  is the set of vertices incident to edges in  $\nu_v$ ).
3. For every vertex  $v \in \alpha^{-1}(1)$ , output  $\text{mark}(v)$ . Moreover, output  $M = \bigcup \{\text{mark}(v)\}_{v \in \alpha^{-1}(1)}$ ,  $D = \{v \in \alpha^{-1}(1) : |\text{mark}(v)| = \frac{1}{\epsilon}\}$ .

We define *regions* based on marked vertices as follows. We will not need this definition for our proof, but we still give it since it provides some intuition regarding which modules are created. We remark that this is the only notion/argument in this subsection that is not necessary.

► **Definition 62.** Given  $0 < \delta < 1$ , a tournament  $G$  and an optimal solution  $\alpha$  to the 3-HITTING SET instance corresponding to  $G$ , let  $\{\text{mark}(v)\}_{v \in \alpha^{-1}(1)}, M, D$  be the output of  $\text{Marking}(\delta, G, \alpha)$ . Then, an  $(M, D)$ -region (region for short) is a maximal subset  $U \subseteq V(G) \setminus (\text{support}(\alpha) \cup M)$  such that there do not exist vertices  $v \in M$ ,  $u, w \in V(G) \setminus (\text{support}(\alpha) \cup M)$  such that  $u <_\alpha v <_\alpha w$ . The collection of regions is denoted by  $\mathcal{R}$ .

We prove that all vertices except for those in  $D$ , and not just those in  $\text{support}(\alpha) \setminus \alpha^{-1}(1)$ , now have unique positions when marked vertices are removed.

► **Lemma 63.** Given  $0 < \delta < 1$ , a tournament  $G$  and an optimal solution  $\alpha$  to the 3-HITTING SET instance corresponding to  $G$ , let  $\{\text{mark}(v)\}_{v \in \alpha^{-1}(1)}, M, D$  be the output of  $\text{Marking}(\delta, G, \alpha)$ . Then, every vertex  $v \in (\text{support}(\alpha) \setminus D) \cup M$   $M$ -fits  $<_\alpha$ .

**Proof.** By Observation 59, the lemma is true for vertices in  $M$ . So, let  $v \in \text{support}(\alpha) \setminus D$ . Due to Lemma 60, the lemma is correct if  $v \notin \alpha^{-1}(1)$ , so we next suppose that  $v \in \alpha^{-1}(1)$ . We claim that  $G - ((\text{support}(\alpha) \cup M) \setminus \{v\})$  does not have a triangle. Targeting a contradiction, suppose that it has a triangle  $T$ . Then, as  $\text{support}(\alpha)$  is a solution, necessarily  $v$  belongs to the triangle. So, denote  $V(T) = \{v, u, w\}$ . However, we have that  $(u, w) \in \mu_v$  but  $(u, w) \notin \nu_v$ . This is a contradiction since  $v \notin D$ . So far, we conclude  $G - ((\text{support}(\alpha) \cup M) \setminus \{v\})$  does not have a triangle. Thus, by Corollary 54 and Proposition 55, this means that  $G - ((\text{support}(\alpha) \cup M) \setminus \{v\})$  admits a unique linear order  $<$  such that for every arc  $(x, y) \in E(G - ((\text{support}(\alpha) \cup M) \setminus \{v\}))$ ,  $x < y$ , and its restriction to  $G - \text{support}(\alpha)$  must equal  $<_\alpha$ . This directly implies the lemma, where if  $v$  is first in  $<$  then it has 0-position, and otherwise it has  $u$ -position where  $u$  is its predecessor in  $<$ . ◀

We remark that Lemma 63 will be implicitly used throughout, specifically when we consider vertices  $v \in \text{support}(\alpha) \setminus D$  and implicitly suppose that the definition of their position is valid. An easy consequence of Lemma 63 is that all regions are modules. However, we will not need to directly use this, but rather use Lemma 63. Moreover, as a consequence of Lemma 63, we can characterize the triangles in  $G$  as follows.

► **Lemma 64.** Given  $0 < \delta < 1$ , a tournament  $G$  and an optimal solution  $\alpha$  to the 3-HITTING SET instance corresponding to  $G$ , let  $\{\text{mark}(v)\}_{v \in \alpha^{-1}(1)}, M, D$  be the output of  $\text{Marking}(\delta, G, \alpha)$ . Then, every triangle in  $G - D$  consists of either

1. three vertices of  $(\text{support}(\alpha) \setminus D) \cup M$ , or
2. a vertex  $v \in \text{support}(\alpha) \setminus D$ , a vertex  $u \in (\text{support}(\alpha) \setminus D) \cup M$  and a vertex  $w \in V(G) \setminus (\text{support}(\alpha) \cup M)$  such that either (i)  $(u, v) \in E(G)$ ,  $v$  is of position 0 or  $r <_\alpha w$ , and  $u$  is of position  $r' \geq_\alpha w$ , or (ii)  $(v, u) \in E(G)$ ,  $u$  is of position 0 or  $r <_\alpha w$ , and  $v$  is of position  $r' \geq_\alpha w$ .

**Proof.** Let  $T$  be a triangle in  $G - D$ . Because  $\text{support}(\alpha)$  is a solution to  $G$ ,  $T$  must contain at least one vertex from  $\text{support}(\alpha) \setminus D$ , which we will denote by  $v$ . In case the other two vertices of  $T$  belong to  $(\text{support}(\alpha) \setminus D) \cup M$ , then the proof is complete. Thus, suppose that  $T$  contains at least one vertex  $w \in V(G) \setminus (\text{support}(\alpha) \cup M)$ . Because  $v$   $M$ -fits  $<_\alpha$  (by Lemma 63), the third vertex of  $T$  cannot also belong to  $V(G) \setminus (\text{support}(\alpha) \cup M)$ , as otherwise  $G - ((\text{support}(\alpha) \cup M) \setminus \{v\})$  contains a triangle (which contradicts that  $v$   $M$ -fits  $<_\alpha$  due to Proposition 55). So, the third vertex, which we denote by  $u$ , belongs to  $(\text{support}(\alpha) \setminus D) \cup M$ . We suppose that  $v$  is of position 0 or  $r <_\alpha w$ , as the proof for the other case, where  $v$  is of position  $r' \geq_\alpha w$ , is symmetric. Then, by the definition of position,  $(v, w) \in E(G)$ . So, because  $T$  is a triangle, this implies that  $(w, u), (u, v) \in E(G)$ . Now, because  $u$   $M$ -fits  $<_\alpha$  (by Lemma 63), having the arc  $(w, u) \in E(G)$  implies that  $u$  is of position  $r' \geq_\alpha w$ . This completes the proof.  $\blacktriangleleft$

We now argue that  $|D|$  is only a  $\delta$ -fraction of the optimum, and hence it is not “costly” to seek only solutions that contain  $D$ . We remark that as we will apply another (non-strict) lossy rule later, we will need to call Marking with  $\delta < \epsilon$ .

► **Lemma 65.** *Given  $0 < \delta < 1$ , a tournament  $G$  and an optimal solution  $\alpha$  to the 3-HITTING SET instance corresponding to  $G$ , let  $\{\text{mark}(v)\}_{v \in \alpha^{-1}(1)}, M, D$  be the output of Marking( $\delta, G, \alpha$ ). Let  $S^*$  be a solution to  $I$ . Then,  $|D \setminus S^*| \leq \delta |S^*|$ .*

**Proof.** Consider some vertex  $v \in D$ . Notice that  $v$  together with any edge in  $\nu_v$  form a triangle in  $G$ . Thus, if  $v \notin S^*$ , then from every edge in  $\nu_v$ , at least one vertex must belong to  $S^*$ . As  $\nu_v$  is a matching, and its size is  $\frac{1}{\delta}$ , this means that  $S^*$  had to contain at least  $\frac{1}{\delta}$  vertices from  $\text{mark}(v)$ . As the sets assigned by  $\text{mark}$  are pairwise disjoint, we have that  $|D \setminus S^*|$  can be of size at most  $\delta \text{opt}(I)$ .  $\blacktriangleleft$

Intuitively, the arguments above naturally give rise to a reduction rule that deletes  $D$ . This will be part of our merged rule given later on.

In order to shrink the size of modules, we will need another marking procedure.

► **Definition 66.** *Given  $0 < \delta, \delta' < 1$ , a tournament  $G$  and an optimal solution  $\alpha$  to the 3-HITTING SET instance corresponding to  $G$ , let  $\{\text{mark}(v)\}_{v \in \alpha^{-1}(1)}, M, D$  be the output of Marking( $\delta, G, \alpha$ ). Then, ExtraMarking( $\delta', G, \alpha, M$ ) is defined as follows.*

1. For every vertex  $v \in \text{support}(\alpha)$ , initialize  $\text{backw}(v) = \emptyset$  and  $\text{forw}(v) = \emptyset$ .
2. For every vertex  $v \in \text{support}(\alpha) \setminus D$ :
  - a. Let  $p$  be the position of  $v$ .
  - b. Let  $\rho_v^{\text{backw}} = \{u \in V(G) \setminus (\text{support}(\alpha) \cup M \cup \bigcup_{r \in \text{support}(\alpha)} (\text{forw}(r) \cup \text{backw}(r))) : u \leq_\alpha p\}$ . If  $|\rho_v^{\text{backw}}| > \frac{1}{\delta'}$ , then let  $\text{backw}(v)$  be the subset of the  $\frac{1}{\delta'}$  largest (according to  $<_\alpha$ ) vertices in  $\rho_v^{\text{backw}}$ , and otherwise let  $\text{backw}(v) = \rho_v^{\text{backw}}$ .
  - c. Let  $\rho_v^{\text{forw}} = \{u \in V(G) \setminus (\text{support}(\alpha) \cup M \cup \bigcup_{r \in \text{support}(\alpha)} (\text{forw}(r) \cup \text{backw}(r))) : p <_\alpha u\}$ . If  $|\rho_v^{\text{forw}}| > \frac{1}{\delta'}$ , then let  $\text{forw}(v)$  be the subset of the  $\frac{1}{\delta'}$  smallest (according to  $<_\alpha$ ) vertices in  $\rho_v^{\text{forw}}$ , and otherwise let  $\text{forw}(v) = \rho_v^{\text{forw}}$ .
3. For every  $v \in \text{support}(\alpha)$ , output  $\text{backw}(v), \text{forw}(v)$ , and  $\widehat{M} = \bigcup_{v \in \text{support}(\alpha) \setminus D} (\text{backw}(v) \cup \text{forw}(v))$ .

The main utility of this marking scheme is given by the following lemma.

► **Lemma 67.** *For  $0 < \delta, \delta' < 1$ , a tournament  $G$  and an optimal solution  $\alpha$  to the 3-HITTING SET instance corresponding to  $G$ , let  $\{\text{mark}(v)\}_{v \in \alpha^{-1}(1)}, M, D$  be the output of  $\text{Marking}(\delta, G, \alpha)$ , and  $\{b_v, f_v, \text{backw}(v), \text{forw}(v)\}_{v \in \text{support}(\alpha)}, \widehat{M}$  be the output of  $\text{ExtraMarking}(\delta', G, \alpha, M)$ . Let  $v \in \text{support}(\alpha) \setminus D, u \in (\text{support}(\alpha) \setminus D) \cup M, w \in V(G) \setminus (\text{support}(\alpha) \cup M \cup \widehat{M})$  such that  $G[\{v, u, w\}]$  is a triangle. Then, the following conditions hold.*

- *If  $(v, u) \in E(G)$ , then  $|\text{backw}(v)| = \frac{1}{\delta'}$  and for every  $r \in \text{backw}(v)$ ,  $G[\{v, u, r\}]$  is a triangle.*
- *Otherwise (when  $(u, v) \in E(G)$ ), then  $|\text{forw}(v)| = \frac{1}{\delta'}$  and for every  $r \in \text{forw}(v)$ ,  $G[\{v, u, r\}]$  is a triangle.*

**Proof.** We only give a proof for the case where  $(v, u) \in E(G)$ , as the proof for the case where  $(u, v) \in E(G)$  is symmetric. Then,  $(w, v) \in E(G)$ . So, Lemma 64 implies that  $u$  is of position 0 or  $p' <_\alpha w$ , and  $v$  is of position  $p \geq_\alpha w$ . Thus,  $w \in \rho^{\text{backw}(v)}$ . Having  $w \in V(G) \setminus (\text{support}(\alpha) \cup M \cup \widehat{M})$  also means that  $w \notin \text{backw}(v)$ , and therefore necessarily  $|\text{backw}(v)| = \frac{1}{\delta'}$ . Now, consider some  $r \in \text{backw}(v)$ . Because  $w \in \rho^{\text{backw}(v)}$  but  $w \notin \text{backw}(v)$ , this means that  $w <_\alpha r$  (because we insert the largest vertices from  $\rho^{\text{backw}(v)}$  into  $\text{backw}(v)$ ). Hence, since  $u$  is of position 0 or  $p' <_\alpha w$ , we have that  $(u, r) \in E(G)$ . Further, by the definition of  $\rho^{\text{backw}(v)}$ , we know that  $r \leq p$ , and therefore  $(r, v) \in E(G)$ . Thus, indeed  $G[\{v, u, r\}]$  is a triangle. This completes the proof. ◀

We now argue that if either all vertices in  $\text{backw}(v)$  are deleted or all vertices in  $\text{forw}(v)$  are deleted (or both), then it is not “costly” to seek only solutions that delete  $v$  as well.

► **Lemma 68.** *Given  $0 < \delta, \delta' < 1$ , a tournament  $G$  and an optimal solution  $\alpha$  to the 3-HITTING SET instance corresponding to  $G$ , let  $\{\text{mark}(v)\}_{v \in \alpha^{-1}(1)}, M, D$  be the output of  $\text{Marking}(\delta, G, \alpha)$ , and  $\{\text{backw}(v), \text{forw}(v)\}_{v \in \text{support}(\alpha)}, \widehat{M}$  be the output of  $\text{ExtraMarking}(\delta', G, \alpha, M)$ . Let  $S'$  be a solution to  $G' = G - (D \cup X)$  for  $X = V(G) \setminus (\text{support}(\alpha) \cup M \cup \widehat{M})$ . Let  $Y = \{v \in \text{support}(\alpha) \setminus D : |\text{backw}(v)| = \frac{1}{\delta'}, \text{backw}(v) \subseteq S'\} \cup \{v \in \text{support}(\alpha) \setminus D : |\text{forw}(v)| = \frac{1}{\delta'}, \text{forw}(v) \subseteq S'\}$ . Then,  $|Y| \leq \delta' |S'|$ .*

**Proof.** Because the collection of the sets  $\text{backw}(v)$  and  $\text{forw}(v)$  taken over all vertices  $v \in \text{support}(\alpha) \setminus D$  are pairwise disjoint,  $S'$  can contain at most  $\delta' |S'|$  such sets of size  $\frac{1}{\delta'}$ . As  $|Y|$  is precisely the number of such sets of size  $\frac{1}{\delta'}$  that  $S'$  contains, the lemma follows. ◀

► **Definition 69.** *Given  $0 < \delta, \delta' < 1$ , the FVST( $\delta, \delta'$ ) reduction rule is defined as follows:*

- **reduce:** *Let  $I = G$  be an instance of FEEDBACK VERTEX SET IN TOURNAMENTS. Use the algorithm in Proposition 9 to compute an optimal solution  $\alpha$  to the classic LP corresponding to it (Definitions 11 and 53). Let  $\{\text{mark}(v)\}_{v \in \alpha^{-1}(1)}, M, D$  be the output of  $\text{Marking}(\epsilon, G, \alpha)$ . Let  $\{\text{backw}(v), \text{forw}(v)\}_{v \in \text{support}(\alpha)}, \widehat{M}$  be the output of  $\text{ExtraMarking}(\delta', G, \alpha, M)$ . Output  $I' = G'$  where  $G' = G - (D \cup X)$  for  $X = V(G) \setminus (\text{support}(\alpha) \cup M \cup \widehat{M})$ .*
- **lift:** *Given  $I, I'$  and a solution  $S'$  to  $I'$ , output  $S = S' \cup D \cup Y$  where  $Y = \{v \in \text{support}(\alpha) \setminus D : |\text{backw}(v)| = \frac{1}{\delta'}, \text{backw}(v) \subseteq S'\} \cup \{v \in \text{support}(\alpha) \setminus D : |\text{forw}(v)| = \frac{1}{\delta'}, \text{forw}(v) \subseteq S'\}$ .*

Just like Lemma 48 in Section 8.1, here also we present a simple lemma that will help us derive a tighter bound on the number of vertices in the output graph. Since the proof follows the exact same arguments as the proof of Lemma 48, it is omitted.

► **Lemma 70.** *Let  $I = G$  be an instance of FEEDBACK VERTEX SET IN TOURNAMENTS, and let  $\alpha$  be a solution to the 3-HITTING SET instance corresponding to  $G$ . Then,  $|\text{support}(\alpha)| \leq 3\text{frac}(I) - 2|\alpha^{-1}(1)|$ .*

Based on Lemmata 64, 65, 67, 68 and 70, we are now ready to prove the main theorem of this subsection.

► **Theorem 71.** *Let  $0 < \epsilon < 1$ . The FEEDBACK VERTEX SET IN TOURNAMENTS problem, parameterized by the fractional optimum of the classic LP, admits a  $(1 + \epsilon)$ -approximate  $(13 + \frac{9}{\epsilon})\text{frac}(I)$ -vertex kernel.*

**Proof.** Our lossy kernelization algorithm consists only of the FVST( $\delta, \delta'$ ) reduction rule where  $\delta = \frac{\epsilon}{3} - \frac{2\epsilon^2}{9}, \delta' = \frac{2\epsilon}{3}$ . Clearly, it runs in polynomial time.

First, we consider the number of vertices in the output graph  $G'$  of **reduce**. By Lemma 70,  $|\text{support}(\alpha)| \leq 3\text{frac}(I) - 2|\alpha^{-1}(1)|$  (I). Moreover,  $|M| = |\bigcup_{v \in \alpha^{-1}(1)} \text{mark}(v)| = 2|\bigcup_{v \in \alpha^{-1}(1)} \nu_v| \leq \frac{2}{\delta}|\alpha^{-1}(1)|$  (II). Additionally,  $|\widehat{M}| = |\bigcup_{v \in \text{support}(\alpha) \setminus D} (\text{backw}(v) \cup \text{forw}(v))| \leq \frac{2}{\delta'}|\text{support}(\alpha)|$  (III). As  $V(G') \subseteq \text{support}(\alpha) \cup M \cup \widehat{M}$  (more precisely,  $V(G') = (\text{support}(\alpha) \setminus D) \cup M \cup \widehat{M}$ ), we have that

$$\begin{aligned}
 |V(G')| &\leq |\text{support}(\alpha)| + |M| + |\widehat{M}| && \text{[Last sentence]} \\
 &\leq |\text{support}(\alpha)| + \frac{2}{\delta}|\alpha^{-1}(1)| + \frac{2}{\delta'}|\text{support}(\alpha)| \\
 &= (1 + \frac{2}{\delta'})|\text{support}(\alpha)| + \frac{2}{\delta}|\alpha^{-1}(1)| && \text{[(I)]} \\
 &\leq (1 + \frac{2}{\delta'}) (3\text{frac}(I) - 2|\alpha^{-1}(1)|) + \frac{2}{\delta}|\alpha^{-1}(1)| && \text{[(II)+(III)]} \\
 &\leq 3(1 + \frac{2}{\delta'})\text{frac}(I) + 2(\frac{1}{\delta} - \frac{2}{\delta'} - 1)|\alpha^{-1}(1)| \\
 &= 3(1 + \frac{3}{\epsilon})\text{frac}(I) + 2(\frac{9}{\epsilon(3-2\epsilon)} - \frac{3}{\epsilon} - 1)|\alpha^{-1}(1)| && \text{[Substitute } \delta \text{ and } \delta'] \\
 &= 3(1 + \frac{3}{\epsilon})\text{frac}(I) + 2(\frac{6}{3-2\epsilon} - 1)|\alpha^{-1}(1)| \\
 &\leq 3(1 + \frac{3}{\epsilon})\text{frac}(I) + 10|\alpha^{-1}(1)| && [\epsilon < 1] \\
 &\leq (13 + \frac{9}{\epsilon})\text{frac}(I) && [|\alpha^{-1}(1)| \leq \text{frac}(I)].
 \end{aligned}$$

We turn to prove that **lift** returns a solution having the desired approximation ratio. To this end, suppose that it is given  $I, I', S'$  where  $S'$  is a solution to  $I'$ . We first show that  $S = S' \cup D \cup Y$  is a solution to  $I$ . Targeting a contradiction, suppose that this is false, and hence there exists a triangle  $T$  in  $G - S$ . As  $D \subseteq S$ , this triangle also exists in  $G - D$ , and hence by Lemma 64,  $T$  consists of either

1. three vertices of  $(\text{support}(\alpha) \setminus D) \cup M$ , or
2. a vertex  $v \in \text{support}(\alpha) \setminus D$ , a vertex  $u \in (\text{support}(\alpha) \setminus D) \cup M$  and a vertex  $w \in V(G) \setminus (\text{support}(\alpha) \cup M)$  such that either (i)  $(u, v) \in E(G)$ ,  $v$  is of position 0 or  $r <_{\alpha} w$ , and  $u$  is of position  $r' \geq_{\alpha} w$ , or (ii)  $(v, u) \in E(G)$ ,  $u$  is of position 0 or  $r <_{\alpha} w$ , and  $v$  is of position  $r' \geq_{\alpha} w$ .

Since  $S'$  is a solution to  $I'$ ,  $T$  must consist of at least one vertex from  $V(G) \setminus V(G') = X = V(G) \setminus (\text{support}(\alpha) \cup M \cup \widehat{M})$ , and therefore the first case is impossible. Moreover, this implies that in the second case,  $w \in X$ . We only consider the case where  $(v, u) \in E(G)$ , as the proof for the other case (when  $(u, v) \in E(G)$ ) follows symmetric arguments. So,  $T = G[\{v, u, w\}]$  where  $v \in \text{support}(\alpha) \setminus D$ ,  $u \in (\text{support}(\alpha) \setminus D) \cup M$ ,  $w \in V(G) \setminus (\text{support}(\alpha) \cup M \cup \widehat{M})$ ,  $(v, u) \in E(G)$ ,  $u$  is of position 0 or  $r <_{\alpha} w$ , and  $v$  is of position  $r' \geq_{\alpha} w$ . By Lemma 67, this means that  $|\text{backw}(v)| = \frac{1}{\delta'}$  and for every  $r \in \text{backw}(v)$ ,  $G[\{v, u, r\}]$  is a triangle. As  $S'$  is a solution to  $I'$  that excludes  $u$  and  $v$ , and as for every  $r \in \text{backw}(v)$ ,  $G[\{v, u, r\}]$  exists in  $G'$ , we have that  $\text{backw}(v) \subseteq S'$ . However, this implies that  $v \in Y$ , and hence  $v \in S$ , so  $T$  cannot exist in  $G - S$ . As we have reached a contradiction,  $S$  is indeed a solution to  $I$ .

It remains to consider the approximation ratio of  $S$ . To this end, first note that  $|S| \leq |S'| + |D| + |Y|$ . So, by Lemma 68, (i)  $|S| \leq (1 + \delta')|S'| + |D|$ . On the other hand, let  $S^*$  be an optimal solution to  $I$ . Observe that, as  $G'$  is a subgraph of  $G$ ,  $S^* \cap V(G')$  is a solution to  $I'$ . So,  $\text{opt}(I') \leq |S^* \cap V(G')|$ . Further,  $S^* \cap V(G) \subseteq S^* \setminus D$ , and by Lemma 65,  $|D \setminus S^*| \leq \delta |S^*|$ . Thus,  $|S^* \cap V(G')| \leq |S^* \setminus D| = |S^*| - |S^* \cap D| = |S^*| - (|D| - |D \setminus S^*|) \leq (1 + \delta)|S^*| - |D|$ , which means that (ii)  $\text{opt}(I') \leq (1 + \delta)\text{opt}(I) - |D|$ . Notice that  $(1 + \delta)(1 + \delta') = (1 + \frac{\epsilon}{3} - \frac{2\epsilon^2}{9})(1 + \frac{2\epsilon}{3}) \leq (1 + \epsilon)$ . Then, from (i) and (ii), we conclude that

$$\begin{aligned} \frac{|S|}{\text{opt}(I)} &\leq (1 + \delta) \frac{(1 + \delta')|S'| + |D|}{\text{opt}(I') + |D|} \\ &\leq (1 + \delta)(1 + \delta') \frac{|S'| + |D|}{\text{opt}(I') + |D|} \\ &\leq (1 + \epsilon) \frac{|S'| + |D|}{\text{opt}(I') + |D|} \\ &\leq (1 + \epsilon) \max\left\{\frac{|S'|}{\text{opt}(I')}, \frac{|D|}{|D|}\right\} \\ &= (1 + \epsilon) \frac{|S'|}{\text{opt}(I')}. \end{aligned}$$

Here, the last inequality follows from Proposition 6. This completes the proof.  $\blacktriangleleft$

► **Corollary 72.** *Let  $0 < \epsilon < 1$ . The FEEDBACK VERTEX SET IN TOURNAMENTS problem, parameterized by the optimum, admits a  $(1 + \epsilon)$ -approximate  $(13 + \frac{9}{\epsilon})\text{opt}$ -vertex kernel.*

Due to Lemma 2, we also have the following corollary of Theorem 71.

► **Corollary 73.** *Let  $0 < \epsilon < 1$ . The FEEDBACK VERTEX SET IN TOURNAMENTS problem, parameterized by a bound  $k$  on the solution size, admits a  $(1 + \epsilon)$ -approximate  $\frac{13 + \frac{9}{\epsilon}}{1 + \epsilon}k$ -vertex kernel.*

## 9 Conclusion

In this paper, we presented positive results on the kernelization complexity of  $d$ -HITTING SET, as well as for its special cases CLUSTER VERTEX DELETION and FEEDBACK VERTEX SET IN TOURNAMENTS. First, we proved that if we allow the kernelization to be *lossy* with a qualitatively better loss than the best possible approximation ratio of polynomial time approximation algorithms, then one can obtain kernels where the number of elements is linear for every fixed  $d$ . Further, we extended the notion of lossy kernelization algorithms to *lossy kernelization protocols* and, then, presented our main result: For any  $\epsilon > 0$ ,  $d$ -HITTING SET admits a (randomized) pure  $(d - \delta)$ -approximate kernelization protocol of call size  $\mathcal{O}(k^{1+\epsilon})$ . Here, the number of rounds and  $\delta$  are fixed constants (that depend only on  $d$  and  $\epsilon$ ). Finally, we complemented the aforementioned results as follows: for the special cases of 3-HITTING SET, namely, CLUSTER VERTEX DELETION and FEEDBACK VERTEX SET IN TOURNAMENTS, we showed that for any  $0 < \epsilon < 1$ , they admit a  $(1 + \epsilon)$ -approximate  $\mathcal{O}(\frac{1}{\epsilon} \cdot \text{opt})$ -vertex kernel.

We conclude the paper with a few interesting open problems.

1. Does  $d$ -HITTING SET admit a kernel with  $f(d) \cdot k^{d-1-\epsilon}$  elements for some fixed  $\epsilon > 0$ , or, even, with just  $f(d) \cdot k$  elements?
2. Does  $d$ -HITTING SET admit a  $(1 + \epsilon)$ -approximate  $\mathcal{O}(f(\epsilon) \cdot k)$ -element kernel (or protocol)?
3. Does  $d$ -HITTING SET admit a  $(1 + \epsilon)$ -approximate  $\mathcal{O}(f(\epsilon) \cdot k)$ -bits kernel (or protocol)?
4. Do FEEDBACK VERTEX SET IN TOURNAMENTS and CLUSTER VERTEX DELETION admit linear vertex kernels?
5. Are lossy kernelization protocols “more powerful” than lossy kernelization algorithms?

## References

- 1 Faisal N. Abu-Khzam. A kernelization algorithm for  $d$ -Hitting Set. *J. Computer and System Sciences*, 76(7):524–531, 2010.
- 2 Stéphane Bessy, Fedor V. Fomin, Serge Gaspers, Christophe Paul, Anthony Perez, Saket Saurabh, and Stéphan Thomassé. Kernels for feedback arc set in tournaments. *J. Comput. Syst. Sci.*, 77(6):1071–1078, 2011.
- 3 Hans L. Bodlaender, Rodney G. Downey, Michael R. Fellows, and Danny Hermelin. On problems without polynomial kernels. *Journal of Computer and System Sciences*, 75(8):423–434, 2009.
- 4 Hans L. Bodlaender, Fedor V. Fomin, and Saket Saurabh. Open problems, worker 2010. Available at <http://fpt.wikidot.com/open-problems>, 2010.
- 5 Mao-cheng Cai, Xiaotie Deng, and Wenan Zang. An approximation algorithm for feedback vertex sets in tournaments. *SIAM J. Comput.*, 30(6):1993–2007, 2000.
- 6 Jianer Chen, Iyad A Kanj, and Weijia Jia. Vertex cover: further observations and further improvements. *Journal of Algorithms*, 41(2):280–301, 2001.
- 7 Marek Cygan, Fedor V. Fomin, Bart MP Jansen, Lukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michał Pilipczuk, and Saket Saurabh. Open problems for fpt school 2014. URL: <http://fptschool.mimuw.edu.pl/opl.pdf>, 2014.
- 8 Marek Cygan, Fedor V. Fomin, Lukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michał Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015.
- 9 Holger Dell and Dániel Marx. Kernelization of packing problems. In *Proceedings of the 23rd Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pages 68–81, 2012.
- 10 Holger Dell and Dieter van Melkebeek. Satisfiability allows no nontrivial sparsification unless the polynomial-time hierarchy collapses. *Journal of the ACM*, 61(4):23:1–23:27, 2014.
- 11 Michael Dom, Jiong Guo, Falk Hüffner, Rolf Niedermeier, and Anke Truß. Fixed-parameter tractability results for feedback set problems in tournaments. *J. Discrete Algorithms*, 8(1):76–86, 2010.
- 12 Andrew Drucker. New limits to classical and quantum instance compression. *SIAM Journal on Computing*, 44(5):1443–1479, 2015.
- 13 Michael R. Fellows, Christian Knauer, Naomi Nishimura, Prabhakar Ragde, Frances A. Rosamond, Ulrike Stege, Dimitrios M. Thilikos, and Sue Whitesides. Faster fixed-parameter tractable algorithms for matching and packing problems. *Algorithmica*, 52(2):167–176, 2008.
- 14 Samuel Fiorini, Gwenaél Joret, and Oliver Schaudt. Improved approximation algorithms for hitting 3-vertex paths. In *Integer Programming and Combinatorial Optimization - 18th International Conference, IPCO 2016, Liège, Belgium, June 1-3, 2016, Proceedings*, volume 9682 of *Lecture Notes in Computer Science*, pages 238–249, 2016.
- 15 Jörg Flum and Martin Grohe. *Parameterized Complexity Theory*. Texts in Theoretical Computer Science. An EATCS Series. Springer-Verlag, Berlin, 2006.
- 16 Fedor V. Fomin, Tien-Nam Le, Daniel Lokshtanov, Saket Saurabh, Stéphan Thomassé, and Meirav Zehavi. Subquadratic kernels for implicit 3-Hitting Set and 3-Set Packing problems. *ACM Trans. Algorithms*, 15(1):13:1–13:44, 2019. doi:10.1145/3293466.
- 17 Fedor V. Fomin, Daniel Lokshtanov, Saket Saurabh, and Meirav Zehavi. *Kernelization. Theory of parameterized preprocessing*. Cambridge University Press, Cambridge, 2019.
- 18 Lance Fortnow and Rahul Santhanam. Infeasibility of instance compression and succinct PCPs for NP. *Journal of Computer and System Sciences*, 77(1):91–106, 2011.
- 19 Jacob Fox, Hao Huang, and Benny Sudakov. On graphs decomposable into induced matchings of linear sizes. *Bulletin of the London Mathematical Society*, 49(1):45–57, 2017.
- 20 Zoltán Füredi. Matchings and covers in hypergraphs. *Graphs and Combinatorics*, 4(1):115–206, 1988.
- 21 Danny Hermelin and Xi Wu. Weak compositions and their applications to polynomial lower bounds for kernelization. In *Proceedings of the 23rd Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pages 104–113, 2012.



- 22 Subhash Khot and Oded Regev. Vertex cover might be hard to approximate to within 2-epsilon. *J. Comput. Syst. Sci.*, 74(3):335–349, 2008. doi:10.1016/j.jcss.2007.06.019.
- 23 Stefan Kratsch and Magnus Wahlström. Representative sets and irrelevant vertices: new tools for kernelization. In *Proceedings of the 53rd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 450–459. IEEE, 2012.
- 24 Daniel Lokshtanov, Pranabendu Misra, Joydeep Mukherjee, Fahad Panolan, Geevarghese Philip, and Saket Saurabh. 2-approximating feedback vertex set in tournaments. In *Proceedings of the 2020 ACM-SIAM Symposium on Discrete Algorithms, SODA 2020, Salt Lake City, UT, USA, January 5-8, 2020*, pages 1010–1018, 2020.
- 25 Daniel Lokshtanov, Fahad Panolan, M. S. Ramanujan, and Saket Saurabh. Lossy kernelization. *CoRR*, abs/1604.04111, 2016.
- 26 Jiri Matousek and Bernd Gärtner. *Understanding and using linear programming*. Springer Science & Business Media, 2007.
- 27 Matthias Mnich, Virginia Vassilevska Williams, and László A. Végh. A 7/3-approximation for feedback vertex sets in tournaments. In *24th Annual European Symposium on Algorithms, ESA 2016, August 22-24, 2016, Aarhus, Denmark*, volume 57 of *LIPICs*, pages 67:1–67:14, 2016.
- 28 George L Nemhauser and Leslie Earl Trotter. Properties of vertex packing and independence system polyhedra. *Mathematical programming*, 6(1):48–61, 1974.
- 29 Imre Z Ruzsa and Endre Szemerédi. Triple systems with no six points carrying three triangles. *Combinatorics (Keszthely, 1976), Coll. Math. Soc. J. Bolyai*, 18:939–945, 1978.
- 30 Jie You, Jianxin Wang, and Yixin Cao. Approximate association via dissociation. *Discrete Applied Mathematics*, 219:202–209, 2017.

**A** Problem Definitions

**Vertex Cover (VC).** Given a graph  $G$ , compute a minimum-sized vertex cover  $S$  of  $G$ , that is, a subset  $S \subseteq V(G)$  such that every edge in  $G$  is incident to at least one vertex in  $S$ .

**$d$ -Hitting Set ( $d$ -HS).** Given a universe  $U$  and a family of sets  $\mathcal{F} \subseteq 2^U$  where each set in  $\mathcal{F}$  has size  $d$ , compute a minimum-sized hitting set  $S$  of  $\mathcal{F}$ , that is, a subset  $S \subseteq U$  such that every set in  $\mathcal{F}$  has non-empty intersection with  $S$ .

Note that VERTEX COVER is equivalent to 2-HITTING SET.

**Cluster Vertex Deletion (CVD).** Given a graph  $G$ , compute a minimum-sized subset  $S \subseteq V(G)$  such that  $G - S$  is a cluster graph.

**Feedback Vertex Set in Tournaments (FVST).** Given a tournament  $G$ , compute a minimum-sized subset  $S \subseteq V(G)$  such that  $G - S$  is acyclic.