Symplectic self-orthogonal and LCD codes from the Plotkin sum construction

Shixin Zhu, Yang Li[∗] , Shitao Li †‡

Abstract

In this work, we propose two criteria for linear codes obtained from the Plotkin sum construction being symplectic self-orthogonal (SO) and linear complementary dual (LCD). As specific constructions, several classes of symplectic SO codes with good parameters including symplectic maximum distance separable codes are derived via ℓ-intersection pairs of linear codes and generalized Reed-Muller codes. Also symplectic LCD codes are constructed from general linear codes. Furthermore, we obtain some binary symplectic LCD codes, which are equivalent to quaternary trace Hermitian additive complementary dual codes that outperform best-known quaternary Hermitian LCD codes reported in the literature. In addition, we prove that symplectic SO and LCD codes obtained in these ways are asymptotically good.

Keywords: Symplectic inner product, Self-orthogonal code, LCD code, Symplectic maximum distance separable code, Plotkin sum construction

Mathematics Subject Classification 94B05 15B05 12E10

1 Introduction

Throughout this paper, let $q = p^m$ be a prime power and \mathbb{F}_q be the finite field with size q. An $[n,k]_q$ *linear code* C is a k-dimensional linear subspace of \mathbb{F}_q^n . Let C^{\perp} be the dual code of ${\mathcal C}$ with respect to a certain inner product (such as the Euclidean, Hermitian or symplectic inner product). A linear code C is said to be *self-orthogonal (SO)* if $C \subset C^{\perp}$ and *linear complementary dual (LCD)* if $C \cap C^{\perp} = \{0\}$. Both SO and LCD codes have attracted significant attention in recent years due to their theoretical and practical importance.

On one hand, constructing, enumerating, characterizing and classifying SO codes have remained as four essential and dynamic research problems since the beginning of coding

[∗]Corresponding author

[†]This research is supported by the National Natural Science Foundation of China (U21A20428, 12171134 and 12001002).

[‡]Shixin Zhu and Yang Li are with the School of Mathematics, Hefei University of Technology, Hefei, 230601, China (email: zhushixinmath@hfut.edu.cn, yanglimath@163.com). Shitao Li is with the School of Mathematical Sciences, Anhui University, Hefei, 230601, China, (email: lishitao0216@163.com).

theory (see [\[6,](#page-14-0) [13,](#page-15-0) [17,](#page-15-1) [31,](#page-16-0) [33,](#page-16-1) [34,](#page-16-2) [37,](#page-16-3) [39,](#page-16-4) [45\]](#page-17-0) and the references therein). Two main factors contribute to the intriguing and broad appeal of SO codes. First, Ding [\[16\]](#page-15-2), Zhang et al. [\[46\]](#page-17-1) and Jin et al. [\[30\]](#page-16-5) respectively proved that binary Euclidean SO codes, q-ary Euclidean SO codes $(q \text{ is odd})$ and $q\text{-ary symplectic SO}$ codes are asymptotically good. Second, extensive researches have established strong correlations between SO codes and various mathematical fields such as combinatorial t-design theory [\[9\]](#page-14-1), group theory [\[15\]](#page-15-3), lattice theory [\[8,](#page-14-2)[15,](#page-15-3)[24\]](#page-15-4), modular forms [\[41\]](#page-16-6), and quantum error-correcting codes (QECCs) [\[14,](#page-15-5) [32\]](#page-16-7). Specifically, finite groups like the Mathieu groups were found to be associated with some SO codes and the extended binary SO Golay code was linked to the Conway group. Many 5-designs were also obtained from SO codes [\[5\]](#page-14-3).

On the other hand, LCD codes were first introduced by Massey [\[38\]](#page-16-8) in 1992, which can provide an optimum linear coding solution for two-user binary adder channel. Subsequently, Sendrier $[40]$ and Güneri *et al.* $[23]$ showed that Euclidean and Hermitian LCD codes are asymptotically good. Carlet et al. [\[10\]](#page-14-4) also further developed LCD codes to combat side channel attacks (SCAs) and fault injection attacks (FIAs). In 2018, Carlet et al. [\[12\]](#page-15-7) proved that any linear code over \mathbb{F}_q is equivalent to some Euclidean LCD code for $q > 3$ and any linear code over \mathbb{F}_{q^2} is equivalent to some Hermitian LCD code for $q > 2$. Since then, the focus has been on binary, ternary Euclidean LCD codes, quaternary Hermitian LCD codes and q-ary symplectic LCD codes (see for example $[2-4, 21, 25, 26, 28, 36, 37, 44]$ $[2-4, 21, 25, 26, 28, 36, 37, 44]$ $[2-4, 21, 25, 26, 28, 36, 37, 44]$ $[2-4, 21, 25, 26, 28, 36, 37, 44]$ $[2-4, 21, 25, 26, 28, 36, 37, 44]$ $[2-4, 21, 25, 26, 28, 36, 37, 44]$ $[2-4, 21, 25, 26, 28, 36, 37, 44]$ $[2-4, 21, 25, 26, 28, 36, 37, 44]$. Note that Xu *et al.* [\[44\]](#page-16-12) and Huang *et al.* [\[2](#page-14-5)8] also employed symplectic LCD symplectic maximum distance separable (MDS) codes to construct maximal entanglement MDS entanglement-assisted QECCs.

Hence, it is always interesting to construct new SO and LCD codes with good parameters and it should also be emphasized that a variety of effective techniques have been developed in the literature. In particular, one of such excellent methods is the so-called Plotkin sum construction [\[29\]](#page-16-13), also referred to as the $(\mathbf{u}, \mathbf{u} + \mathbf{v})$ construction, which can generate new linear codes from old ones. Very recently, by employing the Plotkin sum construction, Li et al. [\[37\]](#page-16-3) constructed many good Euclidean and Hermitian SO and LCD codes. Motivated by this work and the growing interest in SO and LCD codes, a natural problem arises: Can symplectic SO and LCD codes with good parameters be constructed from the Plotkin sum construction? In this paper, we provide an affirmative answer. Main contributions of ours are summarized as follows:

- (1) Criteria for linear codes obtained from the Plotkin sum construction being symplectic SO and symplectic LCD codes are characterized in Theorems [3.2](#page-4-0) and [4.1.](#page-11-0) Also symplectic SO and LCD codes obtained from these ways are proved to be asymptotically good with respect to the symplectic distance in Theorems [3.3](#page-6-0) and [4.2.](#page-12-0)
- (2) Using these two criteria, we further obtain many symplectic SO and LCD codes with good parameters.
	- By utilizing ℓ -intersection pairs of linear codes and generalized Reed-Muller codes, we construct several families of symplectic SO codes with explicit parameters in Theorems [3.6,](#page-7-0) [3.7,](#page-8-0) [3.8](#page-8-1) and [3.13](#page-10-0) as well as Corollary [3.9.](#page-8-2) Many

symplectic dual-containing (DC) codes are also produced by considering symplectic dual codes of these symplectic SO codes. As results, lots of symplectic SO and DC codes with good parameters including symplectic MDS codes are deduced.

• By employing general linear codes, we present a method to construct symplectic LCD codes in Theorem [4.3.](#page-12-1) Based on this method, we further obtain some binary symplectic LCD codes, which are equivalent to the so-called quaternary trace Hermitian additive complementary dual codes that outperform best-known quaternary Hermitian LCD codes in Table 2.

The paper is organized as follows. In Section [2,](#page-2-0) we review some necessary knowledge. In Sections [3](#page-4-1) and [4,](#page-11-1) we apply the Plotkin sum construction to obtain symplectic SO, DC and LCD codes with explicit and good parameters. We also study the asymptotic results of symplectic SO and LCD codes from the Plotkin sum construction. In Section [5,](#page-13-0) we conclude this paper.

2 Preliminaries

2.1 Linear codes

From now on, we always denote 0 as an appropriate zero vector and O as a proper zero matrix. Let $\mathbf{x}_1 = (x_1, x_2, \ldots, x_n), \mathbf{x}_2 = (x_{n+1}, x_{n+2}, \ldots, x_{2n}), \mathbf{y}_1 = (y_1, y_2, \ldots, y_n)$ and $\mathbf{y}_2 = (y_{n+1}, y_{n+2}, \ldots, y_{2n})$ be any four vectors in \mathbb{F}_q^n . The *Hamming weight* of \mathbf{x}_1 is $\mathbf{wt}_{\mathcal{H}}(\mathbf{x}_1) = |\{i \mid x_i \neq 0, 1 \leq i \leq n\}|$ and the *minimum Hamming distance* of an $[n, k]_q$ linear code C is $d_H(\mathcal{C}) = \min\{\textbf{wt}_H(\textbf{x}_1) \mid \textbf{x}_1 \in \mathcal{C} \text{ and } \textbf{x}_1 \neq \textbf{0}\}\.$ Denote $\textbf{x} = (\textbf{x}_1 \mid \textbf{x}_2) =$ $(x_1, \ldots, x_n, x_{n+1}, \ldots, x_{2n})$ and $y = (y_1 | y_2) = (y_1, \ldots, y_n, y_{n+1}, \ldots, y_{2n})$. The *symplectic weight* of **x** is $\mathbf{wt}_s(\mathbf{x}) = |\{i \mid (x_i, x_{n+i}) \neq (0, 0), 1 \leq i \leq n\}|$ and the *minimum symplectic distance* of a $[2n, k]_q$ linear code C is $d_s(\mathcal{C}) = \min\{\mathbf{wt}_s(\mathbf{x}) \mid \mathbf{x} \in \mathcal{C} \text{ and } \mathbf{x} \neq \mathbf{0}\}\)$. In this paper, we denote C by $[n, k, d_H]_q^H$ (resp. $[2n, k, d_s]_q^s$) if C is an $[n, k]_q$ (resp. a $[2n, k]_q$) linear code with minimum Hamming (resp. symplectic) distance d_H (resp. d_s). It is well-known that for an $[n, k, d_H]^H_q$ (resp. a $[2n, k, d_s]^s_q$) linear code C, the *Hamming (resp. symplectic) Singleton bound* says that $d_H \leq n - k + 1$ (resp. $d_s \leq \lfloor \frac{2n-k+2}{2} \rfloor$). Then such a linear code C is called a *Hamming (resp. symplectic)* MDS code if $d_H = n - k + 1$ (resp. $d_s = \lfloor \frac{2n-k+2}{2} \rfloor$ $\frac{-k+2}{2}$]).

Let $\Omega = \begin{bmatrix} 0 & I_n \\ I_n & Q \end{bmatrix}$ $-I_n$ O 1 , where I_n is the identity matrix of size $n \times n$. The *Euclidean inner product* of x_1 and y_1 is defined by

$$
\langle \mathbf{x}_1, \mathbf{y}_1 \rangle_E = \sum_{i=1}^n x_i y_i \tag{1}
$$

and the *symplectic inner product* of x and y is defined by

$$
\langle \mathbf{x}, \mathbf{y} \rangle_s = \mathbf{x} \Omega \mathbf{y}^T = \sum_{i=1}^n (x_i y_{n+i} - x_{n+i} y_i).
$$
 (2)

If C is an $[n, k, d_H]^H$ linear code, its *Euclidean dual code* is given by

$$
\mathcal{C}^{\perp_{\mathcal{E}}} = \{ \mathbf{y}_1 \mid \langle \mathbf{x}_1, \mathbf{y}_1 \rangle_{\mathcal{E}} = 0, \ \forall \ \mathbf{x}_1 \in \mathcal{C} \}
$$
(3)

and C^{\perp_E} has parameters $[n, n-k, d_H^{\perp_E}]_q^H$, where $d_H^{\perp_E}$ denotes the minimum Hamming distance of $C^{\perp_{\mathbb{E}}}$. If C is a $[2n, k, d_s]_q^s$ linear code, its *symplectic dual code* is given by

$$
\mathcal{C}^{\perp_{\rm s}} = \{ \mathbf{y} \mid \langle \mathbf{x}, \mathbf{y} \rangle_{\rm s} = 0, \ \forall \ \mathbf{x} \in \mathcal{C} \}
$$
 (4)

and $C^{\perp_{\rm s}}$ has parameters $[n, n-k, d_{\rm s}^{\perp_{\rm s}}]_q^s$, where $d_{\rm s}^{\perp_{\rm s}}$ denotes the minimum symplectic distance of $C^{\perp_{\rm s}}$. As defined previously, we call C *Euclidean* (resp. *symplectic*) *SO* if $C \subseteq C^{\perp_{\rm E}}$ (resp. $C \subseteq C^{\perp_s}$), call C Euclidean (resp. *symplectic*) DC if $C^{\perp_E} \subseteq C$ (resp. $C^{\perp_s} \subseteq C$), and call C *Euclidean* (resp. *symplectic*) LCD if $C \cap C^{\perp_{E}} = \{0\}$ (resp. $C \cap C^{\perp_{S}} = \{0\}$). In addition, it is easy to check that $(\mathcal{C}^{\perp_{\mathcal{E}}})^{\perp_{\mathcal{E}}} = \mathcal{C}$ and $(\mathcal{C}^{\perp_{\mathcal{S}}})^{\perp_{\mathcal{S}}} = \mathcal{C}$. Note that symplectic SO and LCD codes are characterized by the following two lemmas.

Lemma 2.1 (Theorem 1 in [\[45\]](#page-17-0)). Let C be a $[2n, k]_q$ linear code with a generator matrix G. Then $\mathcal C$ is a symplectic SO code if and only if $G\Omega G^T$ is a zero matrix.

Lemma 2.2 (Theorem 1 in [\[44\]](#page-16-12)). Let C be a $[2n, k]_q$ linear code with a generator matrix G. Then $\mathcal C$ is a symplectic LCD code if and only if $G\Omega G^T$ is nonsingular.

2.2 Additive codes

An $(n, q^k)_{q^2}$ *additive code* is an \mathbb{F}_q -linear subgroup of \mathbb{F}_q^n q^2 , which has size q^k for some integer k satisfying $0 \leq k \leq 2n$. An $(n, q^k, d_H)_{q^2}^H$ *additive code* is an $(n, q^k)_{q^2}$ additive code with minimum Hamming distance d_{H} . Let \mathbb{F}_q^* $\psi_{q^2}^* = \langle \omega \rangle$. For $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v}=(v_1,v_2,\ldots,v_n)\in\mathbb{F}_{q^2}^n$, the *alternating form* of \mathbf{u} and \mathbf{v} is defined by

$$
\langle \mathbf{u}, \mathbf{v} \rangle_{\text{a}} = \sum_{i=1}^{n} \frac{u_i v_i^q - u_i^q v_i}{\omega - \omega^q}.
$$
\n
$$
(5)
$$

We use C^{\perp_a} to denote the dual code of an additive code C under the alternating form and call C *additive SO* if $C \subseteq C^{\perp_{\text{a}}}$ and *additive complementary dual (ACD)* if $C \cap C^{\perp_{\text{a}}} = \{0\}.$ Consider the following map

$$
\phi: \mathbb{F}_q^{2n} \to \mathbb{F}_{q^2}^n,
$$

$$
(a_1, \ldots, a_n, a_{n+1}, \ldots, a_{2n}) \mapsto (a_1 + \omega a_{n+1}, \ldots, a_n + \omega a_{2n}).
$$

It can be checked that $\langle \mathbf{u}, \mathbf{v} \rangle_s = \langle \phi(\mathbf{u}), \phi(\mathbf{v}) \rangle_a$ for any $\mathbf{u}, \mathbf{v} \in \mathbb{F}_q^{2n}$ (see [\[32\]](#page-16-7)). It is also easy to verify that ϕ is an isomorphic map from $(\mathbb{F}_q^{2n}, d_s, \langle \cdot, \cdot \rangle_s)$ to (\mathbb{F}_q^{n}) $\frac{n}{q^2}$, $d_{\rm H}$, $\langle \cdot, \cdot \rangle_{\rm a}$). Therefore, a $[2n, k, d]_q^s$ linear code is equivalent to an $(n, q^k, d)_{q^2}^H$ additive code.

2.3 Asymptotic results

Finally, we end this section with the concept of asymptotic results. For an infinite subset of $[n_i, k_i, d_i]_q^s$ codes from a family of linear codes with $\lim_{i\to\infty} n_i = \infty$, if both of its *asymptotic rate*

$$
r = \liminf_{i \to \infty} \frac{k_i}{n_i}
$$

and *asymptotic relative Hamming (resp. symplectic) distance*

$$
\delta = \liminf_{i \to \infty} \frac{d_i}{n_i}
$$

are greater than 0, we call this family of linear code *asymptotically good with respect to the Hamming (resp. symplectic) distance*. As pointed out in [\[29\]](#page-16-13), asymptotically good linear codes are generally considered interesting and one prefers to delve into a family of asymptotically good codes in coding theory.

3 Symplectic SO codes

3.1 The characterization and the asymptotic result of symplectic SO codes from the Plotkin sum construction

Definition 3.1 ([29]). Let \mathcal{C}_i be an $[n, k_i]_q$ linear code for $i = 1, 2$. Let G_i and H_i be respectively a generator matrix and a parity check matrix of C_i for $i = 1, 2$. The *Plotkin sum* of C_1 and C_2 is a $[2n, k_1 + k_2]_q$ linear code $\mathcal{P}(C_1, C_2)$ defined by

$$
\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2) = \{ (\mathbf{u}, \mathbf{u} + \mathbf{v}) \mid \mathbf{u} \in \mathcal{C}_1, \ \mathbf{v} \in \mathcal{C}_2 \},\tag{6}
$$

whose generator matrix and parity check matrix are respectively

$$
G = \begin{pmatrix} G_1 & G_1 \\ O & G_2 \end{pmatrix} \text{ and } H = \begin{pmatrix} H_1 & O \\ -H_2 & H_2 \end{pmatrix} . \tag{7}
$$

Theorem 3.2. Let C_i be an $[n, k_i, d_i]_q^H$ linear code for $i = 1, 2$. Then the following three *statements are equivalent.*

- (1) $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)$ *is a symplectic SO* $[2n, k_1 + k_2, \min\{d_1, d_2\}]_q^s$ *code.*
- (2) $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_{\rm s}}$ is a symplectic DC [2n, 2n k₁ k₂, min{d_H($\mathcal{C}_1^{\perp_{\rm E}}$) $(\mathcal{C}_1^{\perp_{\mathbb{E}}}),d_{\mathrm{H}}(\mathcal{C}_2^{\perp_{\mathbb{E}}})$ $\binom{\perp_{\rm E}}{2}$ } $\}$ ⁸_{*q} code.*</sub>

$$
(3) \ \mathcal{C}_1 \subseteq \mathcal{C}_2^{\perp_E}.
$$

Proof. First, we prove that $\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)$ and $\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)^{\perp_s}$ have parameters $[2n, k_1 +$ $\{k_2, \min\{d_1, d_2\}\}_{q}^s$ and $[2n, 2n - k_1 - k_2, \min\{d_{\rm H}(C_1^{\perp_{\rm E}})\}$ $(\frac{1}{1}\mathbf{E}), d_{\mathrm{H}}(\mathcal{C}_{2}^{\perp_{\mathrm{E}}})$ $\left\{\frac{1}{2}E\right\}\right\}^s_{q}$, respectively. Then we reduce the three statements to three simpler statements. With Definition [3.1,](#page-4-2) $\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)$ is a $[2n, k_1 + k_2]_q$ linear code, and then $\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)^{\perp_s}$ is a $[2n, 2n - k_1 - k_2]_q$ linear code. Hence, it suffices to prove that $d_s(\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)) = \min\{d_1, d_2\}$ and $d_s(\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2))^{\perp_s} =$ $\min\{d_{\rm H}(\mathcal{C}_1^{\perp_{\rm E}}% ,d_{\rm H}(\mathcal{C}_2^{\perp_{\rm E}}\,,d_{\rm H}(\mathcal{C}_1^{\perp_{\rm E}})\})\}\label{eq:dimless}$ $\mathcal{d}_1^{\perp_{\operatorname{E}}}),d_{\operatorname{H}}(\mathcal{C}_2^{\perp_{\operatorname{E}}})$ $\binom{\perp E}{2}$.

Case 1: the proof of $d_s(\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)) = \min\{d_1, d_2\}$. Let $(u, u + v) \in \mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)$ be any nonzero codeword with $u \in C_1$ and $v \in C_2$. Then $u \neq 0$ or $v \neq 0$. We have two subcases.

• If $\mathbf{v} = \mathbf{0}$, since **u** is a nonzero codeword in \mathcal{C}_1 , we have that

$$
\mathbf{wt}_s((\mathbf{u},\mathbf{u}+\mathbf{v})) = \mathbf{wt}_s((\mathbf{u},\mathbf{u})) = \mathbf{wt}_H(\mathbf{u}) \geq d_1.
$$

• If $\mathbf{v} \neq \mathbf{0}$, let $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v} = (v_1, v_2, \dots, v_n)$. Note that for each integer $1 \leq i \leq n$, if $v_i \neq 0$, then $(u_i, u_i + v_i) \neq (0, 0)$ whether $u_i = 0$ or not; if $v_i = 0$, then $(u_i, u_i + v_i) = (0, 0)$ if and only if $u_i = 0$. Hence, we have that

 $\mathbf{wt}_{s}((\mathbf{u},\mathbf{u}+\mathbf{v})) > \mathbf{wt}_{\mathbf{H}}(\mathbf{v}) > d_2.$

Then it follows that $d_s(\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)) > \min\{d_1, d_2\}.$

Conversely, for $i = 1, 2$, since $d_H(\mathcal{C}_i) = d_i$, there is a codeword $c_i \in \mathcal{C}_i$ such that $\textbf{wt}_{\text{H}}(\textbf{c}_i) = d_i$. Note that both $(\textbf{c}_1, \textbf{c}_1)$ and $(\textbf{0}, \textbf{c}_2)$ are codewords in $\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)$. Then

$$
d_{s}(\mathcal{P}(\mathcal{C}_{1}, \mathcal{C}_{2})) \leq \min \{ \mathbf{wt}_{s}((\mathbf{c}_{1}, \mathbf{c}_{1})), \mathbf{wt}_{s}((\mathbf{0}, \mathbf{c}_{2})) \}
$$

=
$$
\min \{ \mathbf{wt}_{H}(\mathbf{c}_{1}), \mathbf{wt}_{H}(\mathbf{c}_{2}) \}
$$

=
$$
\min \{ d_{1}, d_{2} \}.
$$

In summary, $d_s(\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)) = \min\{d_1,d_2\}$. This completes the proof of **Case 1.**

Case 2: the proof of $d_s(P(C_1, C_2))^{\perp_s} = \min\{d_H(C_1^{\perp_E}), d_H(C_2^{\perp_E})\}$. From Defi-nition [3.1,](#page-4-2) $\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)^{\perp_{\text{E}}}$ has a generator matrix $H = \begin{pmatrix} H_1 & O \\ H & H \end{pmatrix}$ $-H_2$ H_2 . On the other hand, from the definitions of Euclidean dual codes and symplectic dual codes, we can deduce that $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_{\mathcal{S}}} = \mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_{\mathcal{E}}}\Omega_n = \{\mathbf{c}\Omega_n \mid \mathbf{c} \in \mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_{\mathcal{E}}}\}\.$ This turns out that

$$
H\Omega_n = \begin{pmatrix} H_1 & O \\ -H_2 & H_2 \end{pmatrix} \begin{pmatrix} O & I_n \\ -I_n & O \end{pmatrix} = \begin{pmatrix} O & H_1 \\ -H_2 & -H_2 \end{pmatrix}
$$

is a generator matrix of $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_s}$. Since $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_s}$ is linear and H_i is a generator matrix of $C_i^{\perp_E}$ ^{\perp}E for $i = 1, 2$, it is again inferred from Definition [3.1](#page-4-2) that $\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)^{\perp_s}$ can be viewed as the Plotkin sum of $\mathcal{C}_2^{\perp_{\mathcal{E}}}$ and $\mathcal{C}_1^{\perp_{\mathcal{E}}}$ $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_{\mathrm{s}}}= \mathcal{P}(\mathcal{C}_2^{\perp_{\mathrm{E}}})$ $\frac{\perp_{\rm E}}{2}, \widetilde{\mathcal{C}}^{\perp_{\rm E}}_1$ $_1^{\perp_{\rm E}}$). Hence, $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_{\rm s}}$ has parameters $[2n,2n-k_1-k_2,\min\{d_{\rm H}(\mathcal{C}_1^{\perp_{\rm E}})$ $\mathcal{d}_1^{\perp_{\rm E}}), \mathcal{d}_{\rm H}(\mathcal{C}_2^{\perp_{\rm E}})$ $\binom{\perp_{\rm E}}{2}$ } $]_q^{\rm s}$ according to **Case** 1. This completes the proof of Case 2.

By the results above, we only need to prove that the following three simpler statements are equivalent: (1') $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)$ is a symplectic SO code; (2') $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_s}$ is a symplectic DC code; and (3') $\mathcal{C}_1 \subseteq \mathcal{C}_2^{\perp_{\mathbf{E}}}$.

 $(1') \Leftrightarrow (2')$ Since $(\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)^{\perp_s})^{\perp_s} = \mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)$, we have $\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2) \subseteq \mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)^{\perp_s}$ if and only if $(\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_s})^{\perp_s} \subseteq \mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_s}$. Hence $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)$ is symplectic SO if and only if $\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)^{\perp_{\mathbf{s}}}$ is symplectic DC.

 $(1') \Leftrightarrow (3')$ By Definition [3.1,](#page-4-2) $\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)$ has a generator matrix $G =$ $\int G_1$ G_1 O G_2 \setminus . Then we have

$$
G\Omega_n G^T = \begin{pmatrix} G_1 & G_1 \\ O & G_2 \end{pmatrix} \begin{pmatrix} O & I_n \\ -I_n & O \end{pmatrix} \begin{pmatrix} G_1^T & O \\ G_1^T & G_2^T \end{pmatrix} = \begin{pmatrix} O & G_1 G_2^T \\ -G_2 G_1^T & O \end{pmatrix} . \tag{8}
$$

It follows from Lemma [2.1](#page-3-0) and Equation [\(8\)](#page-6-1) that $\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)$ is symplectic SO if and only if $G\Omega_n G^T = O$, if and only if $-G_2 G_1^T = -(G_1 G_2^T)^T = O$, if and only if $C_1 \subseteq C_2^{\perp_E}$. Therefore, the statements $(1')$ and $(3')$ are equivalent.

In summary, the statements (1), (2) and (3) are equivalent. This completes the proof. \Box

We next show that symplectic SO codes obtained by Theorem [3.2](#page-4-0) are asymptotically good. It is therefore an interesting class of linear codes.

Theorem 3.3. *Let* q = 2 *or* q *be an odd prime power. Then* q*-ary symplectic SO codes derived from Theorem [3.2](#page-4-0) are asymptotically good with respect to the symplectic distance.*

Proof. It follows from [\[16,](#page-15-2) Remark 3] and [\[46,](#page-17-1) Corollary 4.2] that there exist asymptotically good binary and q -ary $(q$ is odd) Euclidean SO codes with respect to the Hamming distance. Hence, we assume that $\{\mathcal{C}_i\}_{i=0}^{\infty}$ is an infinite subset of $[n_i, k_i, d_i]_q^{\text{H}}$ codes from asymptotically good q -ary Euclidean SO codes, where $q = 2$ or q is odd. Then it implies that $\lim_{i\to\infty} n_i = \infty$ with $\liminf_{i\to\infty} \frac{k_i}{n_i}$ $\frac{k_i}{n_i} > 0$ and $\liminf_{i \to \infty} \frac{d_i}{n_i}$ $\frac{d_i}{n_i} > 0.$

Since \mathcal{C}_i is Euclidean SO, *i.e.*, $\mathcal{C}_i \subseteq \mathcal{C}_i^{\perp_{\mathrm{E}}},$ it follows form Theorem [3.2](#page-4-0) that $\mathcal{P}(\mathcal{C}_i, \mathcal{C}_i)$ is a $[2n_i, 2k_i, d_i]_q^s$ symplectic SO code, denoted by \mathcal{P}_i . Note that $\lim_{i\to\infty} 2n_i = \infty$. Also for the infinite sequence of symplectic SO codes $\{P_i\}_{i=0}^{\infty}$, the asymptotic rate of $\{P_i\}_{i=0}^{\infty}$ is

$$
r = \liminf_{i \to \infty} \frac{2k_i}{2n_i} = \liminf_{i \to \infty} \frac{k_i}{n_i} > 0
$$
\n
$$
(9)
$$

and the asymptotic relative symplectic distance of $\{\mathcal{P}_i\}_{i=0}^{\infty}$ is

$$
\delta = \liminf_{i \to \infty} \frac{d_i}{2n_i} = \frac{1}{2} \liminf_{i \to \infty} \frac{d_i}{n_i} > 0.
$$
\n(10)

This proves the expected result.

3.2 The first construction related to ℓ -intersection pairs of linear codes

Let \mathcal{C}_i be an $[n, k_i]_q$ linear code for $i = 1, 2$. Then they are said to be an ℓ -intersection *pair* if $\dim(\mathcal{C}_1 \cap \mathcal{C}_2) = \ell$. Here we recall some important results on ℓ -intersection pairs of linear codes.

Lemma 3.4 (Lemma 2.2 in [\[20\]](#page-15-11)). Let \mathcal{C}_i be an $[n, k_i]_q$ linear code for $i = 1, 2$. Suppose *that* dim($C_1 \cap C_2$) = ℓ . Then max{ $k_1 + k_2 - n$, 0 } $\leq \ell \leq \min\{k_1, k_2\}$.

 \Box

Lemma 3.5 (Theorem 7 in [\[27\]](#page-15-12)). Let $q \geq 3$ be a prime power. Then there exist two *Hamming MDS codes* C_1 *and* C_2 *with parameters* $[n, k_1, n-k_1+1]_q^H$ *and* $[n, k_2, n-k_2+1]_q^H$ such that $\dim(\mathcal{C}_1 \cap \mathcal{C}_2) = \ell$ *if any one of the following conditions holds:*

- (1) $2 \leq n \leq q+1, \ 1 \leq k_1, k_2 \leq n-1, \ \max\{k_1+k_2-n, 0\} \leq \ell \leq \min\{k_1, k_2\}$ and $\overline{(n, k_1, k_2, \ell)} \neq (q + 1, 2, 1, 1)$ *or* $(q + 1, 1, 2, 1)$;
- (2) $q = 2^m > 4$, $0 \le \ell \le 3$, $n = q + 2$ and $(k_1, k_2) = (3, q 1)$, $(q 1, 3)$ or $(3, 3)$;
- (3) $q = 2^m \ge 4, q 4 \le \ell \le q 1, n = q + 2 \text{ and } (k_1, k_2) = (q 1, q 1).$

Next, we apply the Plotkin sum construction to ℓ -intersection pairs of linear codes. By Theorem [3.2,](#page-4-0) we can transform ℓ -intersection pairs of linear codes to symplectic SO codes, which is not effectively applicable to the Euclidean and Hermitian inner products under the Hamming distance. In addition, these constructions have explicit parameters.

Theorem 3.6. Let $q \geq 3$ be a prime power. Let n, k_1 and k_2 be any three positive integers *satisfying* $2 \leq n \leq q+1$, $1 \leq k_1 \leq k_2 \leq n-1$ *and* $k_1 + k_2 \geq n$ *. Then the following statements hold.*

- (1) *There exists a symplectic SO* $[2n, n + k_1 k_2, n k_1 + 1]_q^s$ *code.*
- (2) *There exists a symplectic DC* $[2n, n + k_2 k_1, n k_2 + 1]_q^s$ *code.*

Proof. For each integer $2 \leq n \leq q+1$ and $1 \leq k_1 \leq k_2 \leq n-1$ satisfying $(n, k_1, k_2) \neq$ $(q + 1, 2, 1)$ or $(q + 1, 1, 2)$, it follows from Lemma [3.5](#page-7-1) (1) that there exist two Hamming MDS codes, denoted by C_1 and C_2 , with respective parameters $[n, k_1, n - k_1 + 1]_q^H$ and $[n, k_2, n-k_2+1]_q^{\text{H}}$ such that $\dim(\mathcal{C}_1 \cap \mathcal{C}_2) = k_1$. Then $\mathcal{C}_2^{\perp_{\text{E}}}$ $i_2^{\perp_{\rm E}}$ is a Hamming MDS $[n, n$ $k_2, k_2 + 1$ ^H_q code and $\mathcal{C}_1 \subseteq \mathcal{C}_2 = (\mathcal{C}_2^{\perp_{\mathrm{E}}}$ $\binom{\perp_E}{2}^{\perp_E}$.

Hence, $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2^{\perp_{\text{E}}}$ $\binom{1}{2}$ is a symplectic SO $[2n, n + k_1 - k_2, \min\{n - k_1 + 1, k_2 + 1\}]^s_q$ code and $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2^{\perp_{\text{E}}}$ $\frac{1}{2}$ ^L₂</sub> is a symplectic DC $[2n, n+k_2-k_1, \min\{k_1+1, n-k_2+1\}]_q^s$ code from Theorem [3.2.](#page-4-0) Since $d_s(\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2^{\perp_{\mathrm{E}}})$ $(\frac{1}{2}E)$) = min{n - k₁ + 1, k₂ + 1} and d_s ($\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2^{\perp_E})$ $(\frac{\perp_E}{2})^{\perp_s}$ = $\min\{k_1 + 1, n - k_2 + 1\}$, we have the following two cases.

- Case 1: If $k_1 + k_2 \ge n$, we have $d_s(\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2^{\perp_{\text{E}}})$ $(\mathcal{P}_2^{\perp_{\mathrm{E}}})$) = $n - k_1 + 1$ and $d_{\mathrm{s}}(\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2^{\perp_{\mathrm{E}}})$ $\frac{\perp_E}{2}$ $)$ $\frac{\perp_S}{2}$ $) =$ $n - k_2 + 1$, *i.e.*, a symplectic SO $[2n, n + k_1 - k_2, n - k_1 + 1]_q^s$ code and a symplectic DC $[2n, n + k_2 - k_1, n - k_2 + 1]_q^s$ code exist.
- Case 2: If $k_1 + k_2 \leq n$, we have $d_s(\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2^{\perp_{\text{E}}})$ $(\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2^{\perp_{\mathbf{E}}})) = k_2 + 1$ and $d_{\mathbf{s}}(\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2^{\perp_{\mathbf{E}}}))$ $(\frac{\perp_E}{2})^{\perp_s}$ = $k_1 + 1$, *i.e.*, a symplectic SO $[2n, n + k_1 - k_2, k_2 + 1]_q^s$ code and a symplectic DC $[2n, n + k_2 - k_1, k_1 + 1]_q^s$ code exist.

Due to the flexibility of the ranges of k_1 and k_2 , it is not difficult to check that **Case 1** and Case 2 will yield the same families of symplectic SO (resp. DC) codes. Hence, we only consider Case 1 for our construction. Moreover, since $1 \leq k_1 \leq k_2 \leq n-1$ and $k_1 + k_2 \ge n$ are limited in **Case 1**, then (n, k_1, k_2) are always not equal to $(q + 1, 2, 1)$ or $(q+1,1,2)$. Therefore, the desired results (1) and (2) follow. \Box **Theorem 3.7.** Let $q \geq 3$ be a prime power. Let $2 \leq n \leq q+1$ be a positive integer. Then *the following statements hold.*

- (1) *There exists a symplectic MDS symplectic SO* $[2n, 2k, n-k+1]_q^s$ *code and a symplectic MDS symplectic DC* $[2n, 2n-2k, k+1]_q^s$ *code for each integer* $1 \leq k \leq \lfloor \frac{n}{2} \rfloor$ $\frac{n}{2}$.
- (2) *There exists a symplectic MDS symplectic SO* $[2n, 2k+1, n-k]_q^s$ *code and a symplectic MDS* symplectic $DC [2n, 2n - 2k - 1, k + 1]_q^s$ code for each integer $0 \le k \le \lfloor \frac{n-1}{2} \rfloor$ $\frac{-1}{2}$.
- (3) *There exists a symplectic MDS symplectic SO* $[2^{m+1} + 4, 6, 2^m]_{2^m}^s$ *code and a symplectic MDS symplectic DC* $[2^{m+1} + 4, 2^{m+1} - 2, 4]_{2^m}^s$ *code for each* $m \ge 2$ *.*

Proof. (1) Let $k_1 + k_2 = n$ and $1 \leq k_1 \leq \lfloor \frac{n}{2} \rfloor$ $\frac{n}{2}$ in Theorem [3.6.](#page-7-0) Then we can obtain a symplectic SO $[2n, 2k_1, n-k_1+1]_q^s$ code and a symplectic DC $[2n, 2n-2k_1, k_1+1]_q^s$ code. It follows from the facts $n - k_1 + 1 = \left\lfloor \frac{2n - 2k_1 + 2}{2} \right\rfloor$ $\left\lfloor \frac{2k_1+2}{2} \right\rfloor$ and $k_1 + 1 = \left\lfloor \frac{2n-(2n-2k_1)+2}{2} \right\rfloor$ $\frac{-2k_1+2}{2}$ that these two codes are symplectic MDS. This completes the proof of (1) by taking $k = k_1$.

(2) Let $k_1 + k_2 = n + 1 > n$ and $1 \leq k_1 \leq \left\lfloor \frac{n+1}{2} \right\rfloor$ $\frac{+1}{2}$ in Theorem [3.6.](#page-7-0) Then we can obtain a symplectic SO $[2n, 2k_1 - 1, n - k_1 + 1]_q^s$ code and a symplectic DC $[2n, 2n - 2k_1 + 1, k_1]_q^s$ code. It follows from the facts $n-k_1+1=\left\lfloor\frac{2n-(2k_1-1)+2}{2}\right\rfloor$ $\left\lfloor \frac{n(n-1)+2}{2} \right\rfloor$ and $k_1 = \left\lfloor \frac{2n-(2n-2k_1+1)+2}{2} \right\rfloor$ $\frac{2k_1+1+2}{2}$ that these two codes are symplectic MDS. This completes the proof of (2) by taking $k = \bar{k}_1 - 1$.

(3) Similar to the proof of Theorem [3.6,](#page-7-0) for each $m \geq 2$, it follows from taking $(k_1, k_2, \ell) = (3, 2^m - 1, 3)$ in Lemma [3.5](#page-7-1) (2) and Theorem [3.2](#page-4-0) that a symplectic SO $[2^{m+1}+4, 6, 2^m]_{2^m}$ code and a symplectic DC $[2^{m+1}+4, 2^{m+1}-2, 4]_{2^m}$ code exist. Similar to (1) above, it is easy to check that these two codes are symplectic MDS. This completes the desired result (3). \Box

Theorem 3.8. There exists a symplectic self-dual $[2^{m+1} + 4, 2^m + 2, 4]_{2^m}^s$ code for each *integer* $m > 2$ *.*

Proof. Similar to the proof of Theorem [3.6](#page-7-0) again, the desired result follows by taking $(k_1, k_2, \ell) = (3, 3, 3)$ in Lemma [3.5](#page-7-1) (2) or by taking $(k_1, k_2, \ell) = (2^m - 1, 2^m - 1, 2^m - 1)$ in Lemma [3.5](#page-7-1) (3). \Box

In particular, symplectic MDS symplectic self-dual codes can be further deduced from Theorems [3.7](#page-8-0) and [3.8](#page-8-1) as follows.

Corollary 3.9. Let $q \geq 3$ be a prime power. Then the following statements hold.

- (1) *There exists a symplectic MDS symplectic self-dual* $[2n, n, \frac{n}{2} + 1]_q^s$ *code for each even* $2 \leq n \leq q+1$.
- (2) *There exists a symplectic MDS symplectic self-dual* $[2n, n, \frac{n+1}{2}]_q^s$ *code for each odd* $2 \leq n \leq q+1$.
- (3) *There exists a symplectic MDS symplectic self-dual* $[12, 6, 4]_4^s$ *code.*

Proof. Taking $k = \frac{n}{2}$ with even n and $\frac{n-1}{2}$ with odd n in Theorems [3.7](#page-8-0) (1) and (2), respectively, the desired results (1) and (2) follow immediately. Also the desired result (3) holds by taking $m = 2$ in Theorem [3.7](#page-8-0) (3) or Theorem [3.8.](#page-8-1) \Box

Remark 3.10*.* As mentioned earlier, this construction is not effectively applicable to the Euclidean and Hermitian inner products under the Hamming distance based on ℓ intersection pairs of linear codes. Currently, almost all known Hamming MDS Euclidean and Hermitian SO codes are constructed from (extended) generalized Reed-Solomon codes. These known Hamming MDS SO codes can yield additive (in fact, linear) SO codes of generalized Reed-Solomon type by [\[32,](#page-16-7) Lemma 18]. It is not difficult to check that symplectic SO codes $\mathcal C$ we obtain from ℓ -intersection pairs of linear codes are not equivalent to known Hamming MDS Hermitian SO codes, even if they have the same parameters because our method is more universal and $\phi(\mathcal{C})$ is nonlinear additive. In fact, Adriaensen and Ball [\[1\]](#page-14-7) also motivated us to construct additive MDS codes with length less than or equal to $q + 1$.

Example 3.11. Let $q = 8$. By Part (1) of Theorem [3.7,](#page-8-0) we can obtain symplectic MDS symplectic SO $[6, 2, 3]_8^s$, $[8, 2, 4]_8^s$, $[10, 2, 5]_8^s$, $[10, 4, 4]_8^s$, $[12, 2, 6]_8^s$, $[12, 4, 5]_8^s$, $[14, 2, 7]_8^s$ $[14, 4, 6]_{8}^{s}$, $[14, 6, 5]_{8}^{s}$, $[16, 2, 8]_{8}^{s}$, $[16, 4, 7]_{8}^{s}$, $[16, 6, 6]_{8}^{s}$, $[18, 2, 9]_{8}^{s}$, $[18, 4, 8]_{8}^{s}$, $[18, 6, 7]_{8}^{s}$ $[18, 8, 6]_8^s$ and $[20, 6, 8]_8^s$ codes. By Part (2) of Theorem [3.7,](#page-8-0) we can obtain symplectic MDS symplectic SO $[8, 3, 3]_{8}^{s}$, $[10, 3, 4]_{8}^{s}$, $[12, 3, 5]_{8}^{s}$, $[12, 5, 4]_{8}^{s}$, $[14, 3, 6]_{8}^{s}$, $[14, 5, 5]_{8}^{s}$, $[16, 3, 7]_{8}^{s}$, $[16, 5, 6]_{8}^{s}$, $[16, 7, 5]_{8}^{s}$, $[18, 3, 8]_{8}^{s}$, $[18, 5, 7]_{8}^{s}$ and $[18, 7, 6]_{8}^{s}$ codes. By Part (1) of Corollary [3.9,](#page-8-2) we can obtain symplectic MDS symplectic self-dual $[4, 2, 2]_8^s$, $[8, 4, 3]_8^s$, $[12, 6, 4]_8^s$ and [16, 8, 5]^s codes. By Part (2) of Corollary [3.9,](#page-8-2) we can obtain symplectic MDS symplectic self-dual $[6, 3, 2]_{8}^{s}$, $[10, 5, 3]_{8}^{s}$, $[14, 7, 4]_{8}^{s}$ and $[18, 9, 5]_{8}^{s}$ codes. In addition, Theorem [3.8](#page-8-1) also yields a symplectic self-dual $[20, 10, 4]$ ^s code.

3.3 The second construction related to generalized Reed-Muller codes

We recall some notions on *generalized Reed-Muller* (GRM) codes and refer to [\[29\]](#page-16-13) for more details. Let $R = \mathbb{F}_q[x_1, x_2, \ldots, x_m]$ be the ring of polynomials over \mathbb{F}_q and $I =$ $\langle x_1^q-x_1, x_2^q-x_2, \ldots, x_m^q-x_m \rangle$ be the ideal of R. Denote the corresponding \mathbb{F}_q -algebra by $\overline{AL} = \mathbb{F}_q[x_1, x_2, \dots, x_m]/I$ and the set of zeros in \mathbb{F}_q of I by $O(I) = \mathbb{F}_q^m = \{P_1, P_2, \dots, P_n\}.$ Then for any integer $r \geq 0$, the r-th order GRM code GRM (r, m) is defined as

$$
GRM(r, m) = \{ (f(P_1), f(P_2), \dots, f(P_n)) \mid f \in AL, \deg(f) \le r \}
$$
(11)

and we always select a canonical representative of f excluding power x_i^j i^j for $j \geq q$. Then some known results of GRM codes from [\[29\]](#page-16-13) are shown as follows.

Lemma 3.12 ([\[29\]](#page-16-13)). Let notations be the same as above. Suppose $0 \le r < m(q-1)$ and *write* $m(q-1) - r = a(q-1) + b$ *with integers* $a, b \ge 0$ *and* $b < q-1$ *. Then the following statements hold.*

(1) GRM(r, m) *has parameters*

$$
\left[q^m, \sum_{j=0}^m (-1)^j \binom{m}{j} \binom{m+r-jq}{r-jq}, (b+1)q^a\right]_q.
$$
\n(12)

(2) *The Euclidean dual code of a GRM code is still a GRM code. Specifically, we have*

$$
GRM(r, m)^{\perp_E} = GRM(m(q-1) - r - 1, m). \tag{13}
$$

(3) For any integers $0 \leq i \leq r < m(q-1)$, we have

$$
GRM(i, m) \subseteq GRM(r, m). \tag{14}
$$

.

Note that Theorem [3.2](#page-4-0) generally gives q -ary symplectic SO (resp. DC) codes of length up to $2q + 2$ (some can take up to $2q + 4$ when $q = 2^m \ge 4$). Next, we apply the Plotkin sum construction to GRM codes. This permits us to construct q -ary symplectic SO (resp. DC) codes of length exceeding $2q + 2$.

Theorem 3.13. Let r, i and m be three positive integers satisfying $0 \le r \le m(q-1)$ *and* $0 \le i \le m(q-1) - r - 1$ *. Write* $m(q-1) - r = a_r(q-1) + b_r$, $m(q-1) - i =$ $a_i(q-1) + b_i, r+1 = a'_i$ $f'_r(q-1) + b'_r$ and $i+1 = a'_r$ $i(q-1)+b'_i$ a'_{i} , where $a_{r}, a_{i}, a'_{r}, a'_{i} \geq 0$ and $0 \leq b_r, b_i, b'_r, b'_i < q-1$. Then the following statements hold.

(1) *There exists a symplectic SO code with parameters*

$$
\left[2q^m,\sum_{j=0}^m(-1)^j\binom{m}{j}\left[\binom{m+i-jq}{i-jq}+\binom{m+r-jq}{r-jq}\right],\min\left\{(b_i+1)q^{a_i},(b_r+1)q^{a_r}\right\}\right]_q^s.
$$

(2) *There exists a symplectic DC code with parameters*

$$
\left[2q^m, 2q^m - \sum_{j=0}^m (-1)^j \binom{m}{j} \left[\binom{m+i-jq}{i-jq} + \binom{m+r-jq}{r-jq}\right], \min\left\{(b_i'+1)q^{a_i'}, (b_r'+1)q^{a_r'}\right\}\right]_q^s.
$$

Proof. (1) Let notations be the same as before. Let $C_2 = \text{GRM}(r, m)$. Since r is a positive integer satisfying $0 \le r < m(q-1)$, then it follows from Lemma [3.12](#page-9-0) (1) and the representation of r that C_2 has parameters $[q^m, \sum_{j=0}^m (-1)^j \binom{m}{j} \binom{m+r-jq}{r-jq}, (b_r+1)q^{a_r}]^H_q$. From Lemma [3.12](#page-9-0) (2), we have $C_2^{\perp_{\rm E}} = \text{GRM}(m(q-1) - r - 1, m)$.

Note that i is a positive integer satisfying $0 \le i \le m(q-1)-r-1$, then $0 \le i \le m(q-1)$. On one hand, from Lemma [3.12](#page-9-0) (1) and the representation of i, $\text{GRM}(i, m)$ has parameters

$$
\left[q^m, \sum_{j=0}^m (-1)^j \binom{m}{j} \binom{m+i-jq}{i-jq}, (b_i+1)q^{a_i}\right]_q^H
$$

On the other hand, from Lemma [3.12](#page-9-0) (3), we have $\text{GRM}(i, m) \subseteq \text{GRM}(m(q - 1) - r - 1)$ $1, m$) = $\mathcal{C}_2^{\perp_{\mathrm{E}}}$ ^{\perp}E. Let $\mathcal{C}_1 = \text{GRM}(i, m)$. Then $\mathcal{C}_1 \subseteq \mathcal{C}_2^{\perp_E}$. Therefore, from Theorem [3.2](#page-4-0) (1), $\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)$ is a symplectic SO code with parameters

$$
\left[2q^m, \sum_{j=0}^m (-1)^j \binom{m}{j} \left[\binom{m+i-jq}{i-jq} + \binom{m+r-jq}{r-jq}\right], \min\left\{(b_i+1)q^{a_i}, (b_r+1)q^{a_r}\right\}\right]_q^s
$$

This completes the proof of (1).

(2) Let notations be the same as (1) above. From Lemma [3.12](#page-9-0) (2), we have $\mathcal{C}_1^{\perp_{\rm E}}$ = GRM $(m(q-1) - i - 1, m)$. Since $0 \le r < m(q-1)$ and $0 \le i \le m(q-1) - r - 1$, we have $0 \le m(q-1) - r - 1 < m(q-1)$ and $0 \le r \le m(q-1) - i - 1 < m(q-1)$. It follows from Lemma [3.12](#page-9-0) (1) and the representations of $r + 1$ and $i + 1$ that the respective minimum Hamming distances of $C_1^{\perp_E}$ and $C_2^{\perp_E}$ are $(b'_i + 1)q^{a'_i}$ and $(b'_r + 1)q^{a'_r}$. Therefore, from Theorem [3.2](#page-4-0) (2), $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_{\mathbf{s}}}$ is a symplectic DC code with parameters

$$
\left[2q^m, 2q^m - \sum_{j=0}^m (-1)^j \binom{m}{j} \left[\binom{m+i-jq}{i-jq} + \binom{m+r-jq}{r-jq} \right], \min\left\{ (b_i'+1)q^{a_i'}, (b_r'+1)q^{a_r'} \right\} \right]_q^s
$$

 \Box

This completes the proof of (2).

Example 3.14. We list some symplectic SO and DC codes in Table 1 obtained by Theorem [3.13.](#page-10-0) Note that the differences between the symplectic Singleton bound and their minimum symplectic distances are at most 2 and some of them are 0. Furthermore, it is well-known that the Singleton bound is a rough bound when the length is relatively large compared to q. Hence, these codes listed indeed have good parameters.

q	$\,m$	\boldsymbol{r}	ι	л. Symplectic SO	Bound	q	m	$\,r$	ı	Symplectic DC	Bound
$\overline{2}$	2	\cup		$[8, 4, 2]_2^s$	3	3	$\overline{2}$			$[18, 12, 3]_3^s$	4
$\overline{2}$	3			$[16, 8, 4]_2^s$	5	3	3	\mathbf{I}		$[54, 46, 3]_3^s$	5
3			θ	$[6, 3, 2]_3^s$	$\mathcal{D}_{\mathcal{L}}$	4	$\overline{2}$	1		$[32, 26, 3]_4^s$	
3	$\overline{2}$			$[18, 6, 6]_3^{\rm s}$		4	3	\mathbf{I}		$[128, 120, 3]_4^s$	5
3	$\overline{2}$	$\overline{2}$		$[18, 9, 3]_3^s$	5	5	1	1		$[10, 6, 3]_5^s$	3
4				$[8, 4, 3]_4^s$	3	5	$\overline{2}$	\mathbf{I}		$[50, 44, 3]_5^s$	
4	$\overline{2}$			$[32, 6, 12]_4^s$	14	5	3			$[250, 242, 3]_5^s$	5
5				$[10, 4, 4]_5^s$	4	⇁	1	л.		$[14, 10, 3]_{7}^{s}$	3
5		$\overline{2}$		$[10, 5, 3]_5^s$	3	⇁		$\overline{2}$	$\overline{2}$	$[14, 8, 4]_{7}^{s}$	
				$[14, 4, 6]_7^8$	6	⇁	$\overline{2}$	1 \mathbf{I}		$[98, 92, 3]_{7}^{s}$	
		റ	$\overline{2}$	$[14, 6, 5]_{7}^{s}$	5		3			$[686, 678, 3]_{7}^{s}$	5

Table 1: Some good symplectic SO and DC codes from Theorem [3.13](#page-10-0)

The "Bound" in the sixth and twelfth columns denote the symplectic Singleton bound.

4 Symplectic LCD codes

In this section, we construct symplectic LCD codes from the Plotkin sum construction. The following criterion is important for us.

Theorem 4.1. Let \mathcal{C}_i be an $[n, k_i, d_i]_q^{\text{H}}$ linear code for $i = 1, 2$. Then the following three *statements are equivalent.*

- (1) $\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)$ *is a symplectic LCD* $[2n, k_1 + k_2, \min\{d_1, d_2\}]_q^s$ *code.*
- (2) $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_{\rm s}}$ *is a symplectic LCD* [2n, 2n k₁ k₂, min{d_H($\mathcal{C}_1^{\perp_{\rm E}}$) $\frac{1}{1}$ E), $d_{\rm H}$ ($\mathcal{C}_2^{\perp_{\rm E}}$ $\binom{\perp_{\rm E}}{2}$ } $\}$ ⁸_{*q} code.*</sub>

(3) $C_1 \cap C_2^{\perp_{\mathcal{E}}} = \{0\}$ *and* $k_1 = k_2$.

Proof. The parameters of $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)$ and $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)^{\perp_s}$ are straightforward from the proof of Theorem [3.2.](#page-4-0)

 $(1) \Leftrightarrow (2)$ The result is obvious since $(\mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)^{\perp_s})^{\perp_s} = \mathcal{P}(\mathcal{C}_1, \mathcal{C}_2)$.

(1) \Leftrightarrow (3) By Equation [\(8\)](#page-6-1), $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)$ is symplectic LCD if and only if rank $(G\Omega_nG^T)$ = $2\text{rank}(G_1G_2^T) = k_1 + k_2$. Since $\text{rank}(G_1G_2^T) = k_1 - \text{dim}(\mathcal{C}_1 \cap \mathcal{C}_2^{\perp_{\text{E}}})$, then $\mathcal{P}(\mathcal{C}_1,\mathcal{C}_2)$ is symplectic LCD is further equivalent to $2 \dim(C_1 \cap C_2^{\perp_{\mathcal{E}}}) = k_1 - k_2$. Note that C_1 and $C_2^{\perp_{\mathcal{E}}}$ 2 have respective parameters $[n, k_1]_q$ and $[n, n - k_2]_q$. Then according to Lemma [3.4,](#page-6-2) we have $2\dim(\mathcal{C}_1 \cap \mathcal{C}_2^{\perp_{\mathbb{E}}}) \ge \max\{2k_1 - 2k_2, 0\}$. Hence, $2\dim(\mathcal{C}_1 \cap \mathcal{C}_2^{\perp_{\mathbb{E}}}) = k_1 - k_2$ holds if and only if $k_1 - k_2 \ge 0$ and $k_1 - k_2 \ge 2k_1 - 2k_2$, *i.e.*, dim($C_1 \cap C_2^{\perp_{\text{E}}}$) = 0 (*i.e.*, $C_1 \cap C_2^{\perp_{\text{E}}}$ = {0}) and $k_1 = k_2$, which implies that the statements (1) and (3) are equivalent.

In summary, the statements (1), (2) and (3) are equivalent. This completes the proof. \Box

Similar to Theorem [3.3,](#page-6-0) we demonstrate that symplectic LCD codes derived from Theorem [4.1](#page-11-0) also exhibit asymptotic goodness. Therefore, these symplectic LCD codes are also meaningful.

Theorem 4.2. *Let* q *be a prime power. Then* q*-ary symplectic LCD codes derived from Theorem [4.1](#page-11-0) are asymptotically good with respect to the symplectic distance.*

Proof. By [\[38\]](#page-16-8), q-ary Euclidean LCD codes are asymptotically good with respect to the Hamming distance. Let $\{\mathcal{C}_i\}_{i=0}^{\infty}$ be an infinite subset of $[n_i, k_i, d_i]_q^{\text{H}}$ codes from asymptotically good q-ary Euclidean LCD codes. Since $\dim(\mathcal{C}_i \cap \mathcal{C}_i^{\perp_{\mathrm{E}}}) = 0$, it follows from Theorem [4.1](#page-11-0) that $\mathcal{P}(\mathcal{C}_i, \mathcal{C}_i)$ gives a $[2n_i, 2k_i, d_i]_q^s$ symplectic LCD code, denoted by \mathcal{P}_i . Then the rest of the proof is similar to that of Theorem [3.3.](#page-6-0) Therefore, the excepted result follows. **D**

Furthermore, we have the following result.

Theorem 4.3. Let C_1 be an $[n, k, d]_q^H$ linear code. If there is a permutation matrix P such that $C_2 = C_1 P$ and $C_1 \cap C_2^{\perp_E} = \{0\}$, then there is a symplectic LCD $[2n, 2k, d]_q^s$, i.e., $an \, ACD \, (n, q^{2k}, d)_{q^2}^{\rm H}$ *code and a symplectic LCD* $[2n, 2n - 2k, d_{\rm H}(C^{\perp_{\rm E}})]_q^s$ *, i.e., an ACD* $(n, q^{2n-2k}, d_{\rm H}(\mathcal{C}^{\perp_{\rm E}}))_{q^2}^{\rm H}$ code.

Proof. Since $C_2 = C_1 P$ and P is a permutation matrix, then C_2 is also an $[n, k, d]_q^H$ linear code. Then it follows from Theorem [4.1](#page-11-0) that there exists a symplectic LCD $[2n, 2k, d]_q^s$ code \mathcal{C}' and a symplectic LCD $[2n, 2n-2k, d_H(\mathcal{C}^{\perp_{\mathcal{E}}})]_q^s$ code \mathcal{C}'' . Consider $\phi(\mathcal{C}')$ and $\phi(\mathcal{C}'')$. The desired result follows immediately from [\[32,](#page-16-7) Lemma 14]. \Box

For convenience, we use an array to represent a permutation matrix. For example, $P =$ $(3\ 5\ 2\ 1\ 6\ 4)$ denotes a permutation matrix P whose $(1, 3), (2, 5), (3, 2), (4, 1), (5, 6), (6, 4)$ entries are 1 and others are 0. Note also that a $[2n, 2k, d]_2^s$ binary symplectic LCD code is further equivalent to a so-called $(n, 2^{2k}, d)^{H}_{4}$ quaternary trace Hermitian (TrH) ACD code [\[14\]](#page-15-5). For more details on TrH ACD codes, one can refer to recent papers [\[22,](#page-15-13)[42,](#page-16-14)[43\]](#page-16-15). In the following, we give some specific constructions of binary symplectic LCD codes, i.e., quaternary TrH ACD codes that outperform best-known quaternary Hermitian LCD codes reported in the literature.

Example 4.4. In [\[35\]](#page-16-16), Li constructed a best-known $[63, 56, 3]_4^{\text{H}}$ quaternary Hermitian LCD code. Take C_1 as the best-known $[63, 56, 4]_2^H$ binary linear code in the current MAGMA BKLC database [\[7,](#page-14-8) [19\]](#page-15-14). Let

$$
P_{65} = (53\ 29\ 63\ 14\ 44\ 47\ 46\ 4\ 51\ 59\ 11\ 20\ 10\ 23\ 13\ 37\ 42\ 9\ 26\ 34\ 12
$$

49\ 38\ 30\ 62\ 56\ 16\ 55\ 28\ 33\ 3\ 61\ 40\ 6\ 5\ 35\ 22\ 24\ 52\ 50\ 25\ 7
18\ 39\ 36\ 31\ 8\ 21\ 27\ 57\ 17\ 60\ 41\ 58\ 19\ 43\ 54\ 48\ 1\ 32\ 15\ 45\ 2)

and $C_2 = C_1 P_{65}$. Verified by the Magma software package [\[7\]](#page-14-8), $C_1 \cap C_2^{\perp_{\rm E}} = \{0\}$. Hence, Theorem [4.3](#page-12-1) yields a $[126, 112, 4]_2^s$ binary symplectic LCD code, which is equivalent to a $(63, 2^{56}, 4)_4^{\text{H}}$ TrH ACD code. Note that the $(63, 2^{56}, 4)_4^{\text{H}}$ TrH ACD code outperforms the best-known $[63, 56, 3]_4^H$ Hermitian LCD code. In Table 2, we list more quaternary TrH ACD codes derived from Theorem [4.3](#page-12-1) outperform linear counterparts.

Table 2: Some quaternary TrH ACD codes outperform linear counterparts

	\boldsymbol{P}	Binary symplectic	Quaternary TrH	Known quaternary	
C_1		LCD codes	ACD codes	Hermitian LCD codes	
$[46, 23, 11]_2^{\rm H}$	P_{46}	$[92, 46, 11]_2^s$	$(46,2^{23},11)_{4}^{\rm H}$	$[46, 23, 7]_4^H$ [18]	
$[52, 26, 10]_2^{\rm H}$	P_{52}	$[104, 52, 10]_2^s$	$(52, 2^{26}, 10)^{\text{H}}_{4}$	$[52, 26, 2]_4^H$ [18]	
$[56, 28, 12]_2^{\rm H}$	$P_{\rm 56}$	$[112, 56, 12]_2^s$	$(56, 2^{28}, 12)^{\text{H}}_{4}$	$[56, 28, 4]_4^H$ [18]	
$[58, 29, 12]_2^{\rm H}$	P_{58}	$[116, 58, 12]_2^s$	$(58, 2^{29}, 12)^{\text{H}}_4$	$[58, 29, 4]_4^H$ [18]	
$[62, 31, 12]_2^{\rm H}$	P_{62}	$[114, 62, 12]_2^s$	$(62, 2^{31}, 12)^{\text{H}}_4$	$[62, 31, 7]_4^H$ [11]	
$[63, 56, 4]^{\text{H}}_{2}$	P_{63}	$[126, 112, 4]_2^s$	$(63, 2^{56}, 4)$ ^H	$[63, 56, 3]_4^H$ [35]	
$[64, 32, 12]_2^{\rm H}$	P_{64}	$[128, 64, 12]_2^s$	$(64, 2^{32}, 12)^{\text{H}}_{4}$	$[64, 32, 8]_4^H$ [11]	
$[70, 35, 14]_2^{\rm H}$	P_{70}	$[140, 70, 14]_2^s$	$(70, 2^{35}, 14)^{\text{H}}_4$	$[70, 35, 5]_4^H$ [18]	
$[72, 36, 15]_2^{\rm H}$	P_{72}	$[144, 72, 15]_2^s$	$(72, 2^{36}, 15)^{\text{H}}_4$	$[72, 36, 6]_4^H$ [18]	
$[74, 37, 14]_2^{\rm H}$	P_{74}	$[148, 74, 14]_2^s$	$(74, 2^{37}, 14)^{\text{H}}_{4}$	$[74, 37, 4]_4^H$ [18]	

 \cdot C_1 is the best-known binary linear code in the current MAGMA BKLC database [\[7,](#page-14-8) [19\]](#page-15-14).

 $P_{46}, P_{52}, P_{56}, P_{58}, P_{62}, P_{63}, P_{64}, P_{70}, P_{72}, P_{74}$ are permutation matrices listed in Appendix.

5 Conclusions

In this paper, we study the Plotkin sum construction with respect to the symplectic inner product and symplectic distance. Two criteria for linear codes derived from the Plotkin sum construction being symplectic SO and LCD codes are proposed. Symplectic SO and LCD codes constructed by these two ways are also proved to be asymptotically good. Based on these criteria, we further present several explicit constructions of symplectic SO and LCD codes via ℓ -intersection pairs of linear codes, GRM codes and general linear codes. As a result, many symplectic SO (resp. DC) and LCD codes with good parameters including symplectic MDS codes are obtained. In particular, we also construct some binary symplectic LCD codes, which are equivalent to quaternary TrH ACD codes that outperform best-known quaternary Hermitian LCD codes.

Generally, it is difficult to determine the minimum symplectic distance or the symplectic weights distribution of a symplectic SO code or a symplectic LCD code. Although there are explicit minimum symplectic distances of symplectic SO and LCD codes obtained in this paper, it would be interesting to discuss symplectic weights distributions of these symplectic SO codes and symplectic LCD codes or to construct more symplectic SO codes and symplectic LCD codes with explicit minimum symplectic distances. In addition, it would also be interesting to apply symplectic SO and LCD codes obtained in this paper to construct QECCs and maximal entanglement EAQECCs.

Conflict of Interest

The authors affirm that no conflicts of interest exist.

Data Availability

Data used for the research has been described in the article.

References

- [1] Adriaensen S., Ball S.: On additive MDS codes with linear projections. Finite Fields Appl. 91, 102255 (2023).
- [2] Araya M., Harada M.: On the minimum weights of binary linear complementary dual codes. Cryptogr. Commun. 12(2), 285-300 (2020).
- [3] Araya M., Harada M., Saito K.: Quaternary Hermitian linear complementary dual codes. IEEE Trans. Inf. Theory 66(5), 2751-2759 (2020).
- [4] Araya M., Harada M., Saito K.: On the minimum weights of binary LCD codes and ternary LCD codes. Finite Fields Appl. 76, 101925 (2021).
- [5] Assmus Jr E.F., Mattson Jr H.F.: New 5-designs. J. Comb. Theory 6(2), 122-151 (1969).
- [6] Bouyukliev I., Bouyuklieva S., Gulliver T.A., Ostergard P.R.J.: Classification of optimal binary self-orthogonal codes. J. Comb. Math. Comb. Comput. 59(33), (2006).
- [7] Bosma W., Cannon J., Playoust C.: The Magma algebra system I: The user language. J. Symbolic Comput. 24(3-4), 235-265 (1997).
- [8] Bannai E., Dougherty S.T., Harada M., Oura M.: Type II codes, even unimodular lattices, and invariant rings. IEEE Trans. Inf. Theory 45(4), 1194-1205 (1999).
- [9] Bachoc C., Gaborit P.: Designs and self-dual codes with long shadows. J. Comb. Theory Ser. A 105(1), 15-34 (2004).
- [10] Carlet C., Guilley S.: Complementary dual codes for counter-measures to sidechannel attacks. Adv. Math. Commun. 10(1), 131-150 (2016).
- [11] Crnković D., Grbac A., Švob A.: Formally self-dual LCD codes from two-class association schemes. Appl. Algebra Eng. Commun. Comput. 34, 183-200 (2023).
- [12] Carlet C., Mesnager S., Tang C., Qi Y., Pellikaan R.: Linear codes over \mathbb{F}_q are equivalent to LCD codes for $q > 3$. IEEE Trans. Inf. Theory $64(4)$, 3010-3017 (2018).
- [13] Conway J.H., Pless V., Sloane N.J.A.: The binary self-dual codes of length up to 32: A revised enumeration. J. Combin. Theory Ser. A 60(2), 183-195 (1992).
- [14] Calderbank A.R., Rains E.M., Shor P.M., Sloane N.J.A.: Quantum error correction via codes over GF(4). IEEE Trans. Inf. Theory 44(4), 1369-1387 (1998).
- [15] Conway J.H., Sloane N.J.A.: *Sphere Packings, Lattices and Groups*, 3rd edn. Springer, New York (1998).
- [16] Ding Y.: Asymptotic bound on binary self-orthogonal codes. Sci. China Series A: Math. **52**(4), 631-638 (2009).
- [17] Fu, Y., Liu, H.: Galois self-orthogonal constacyclic codes over finite fields. Des. Codes Cryptogr. 90(11), 2703-2733 (2022).
- [18] Grbac, A.: Self-dual and LCD codes from two class association schemes (Doctoral dissertation, University of Zagreb. Faculty of Science. Department of Mathematics). (2020).
- [19] Grassl M.: Bounds on the minimum distance of linear codes and quantum codes. Online available at http://www.codetables.de. (Accessed 20 May 2023).
- [20] Guenda K., Gulliver T.A., Jitman S., Thipworawimon S.: Linear ℓ-intersection pairs of codes and their applications. Des. Codes Cryptogr. 88, 133-152 (2020).
- [21] Galvez L., Kim J.-K., Lee N., Roe Y.G., Won B.S.: Some bounds on binary LCD codes. Cryptogr. Commun. 10(4), 719-728 (2018).
- [22] Guan C., Li R., Liu Y., Ma Z.: Some quaternary additive codes outperform linear counterparts. IEEE Trans. Inf. Theory 69(11), 7122-7131 (2023).
- [23] Güneri C., Özkaya B., Solé P.: Quasi-cyclic complementary dual codes. Finite Fields Appl. 42, 67-80 (2016).
- [24] Harada M.: On the existence of frames of the Niemeier lattices and self-dual codes over \mathbb{F}_p . J. Algebra **321**(8), 2345-2352 (2009).
- [25] Harada M.: Construction of binary LCD codes, ternary LCD codes and quaternary Hermitian LCD codes. Des. Codes Cryptogr. 89(10), 2295-2312 (2021).
- [26] Harada M., Saito K.: Binary linear complementary dual codes. Cryptogr. Commun. 11(4), 677-696 (2019).
- [27] Huang Z., Fang W., Fu F.W.: Linear ℓ -intersection pairs of MDS codes and their applications to AEAQECCs. Cryptogr. Commun. 14(5), 1189-1206 (2022).
- [28] Huang X., Li J, Huang S.: Constructions of symplectic LCD MDS codes from quasicyclic codes. Adv. Math. Commun. 16(4), 1-12 (2022).
- [29] Huffman W.C., Pless V.: *Fundamentals of error-correcting codes*. Cambridge University Press (2003).
- [30] Jin, L., Xing, C.: Quantum Gilbert-Varshamov bound through symplectic selforthogonal codes. in: IEEE Int. Symp. Inf. Theory (ISIT), pp. 455-458 (2011).
- [31] Kim J.-L., Choi W.-H.: Self-orthogonality matrix and Reed-Muller codes. IEEE Trans. Inf. Theory 68(11), 7159-7164 (2022).
- [32] Ketkar A., Klappenecker A., Kumar S., Sarvepalli P.K.: Nonbinary stabilizer codes over finite fields. IEEE Trans. Inf. Theory 52(11), 4892-4914 (2006).
- [33] Kim J.-L, Kim Y.-H., Lee N.: Embedding linear codes into self-orthogonal codes and their optimal minimum distances. IEEE Trans. Inf. Theory $67(6)$, $3701-3707$ (2021).
- [34] Kim J.-L., Lee, Y.: Euclidean and Hermitian self-dual MDS codes over large finite fields. J. Combin. Theory Ser. A 105(1), 79-95 (2004).
- [35] Li C.: Hermitian LCD codes from cyclic codes. Des. Codes Cryptogr. 86, 2261-2278 (2018).
- [36] Liu X., Liu H., Yu L. New binary and ternary LCD codes from matrix-product codes. Linear Multilinear A. 70(5), 809-823 (2022).
- [37] Li Y., Zhu S., Martínez-Moro E.: The hull of two classical propagation rules and their applications. IEEE Trans. Inf. Theory $69(10)$, $6500-6511$ (2023).
- [38] Massey J.L.: Linear codes with complementary duals. Discrete Math. 106, 337-342 (1992).
- [39] Pless V.: A classification of self-orthogonal codes over GF(2). Discrete Math. 3(1-3), 209-246 (1972).
- [40] Sendrier N.: Linear codes with complementary duals meet the Gilbert-Varshamov bound. Discrete Math. 285(1), 345-347 (2004).
- [41] Shi M., Choie Y.J., Sharma A., Solé P.: Codes and modular forms: a dictionary. World Scientific, 2020.
- [42] Shi, M., Liu, N., Kim, J.-L., Solé, P.: Additive complementary dual codes over \mathbb{F}_4 . Des. Codes Cryptogr. 91(1), 273-284 (2023).
- [43] Shi, M., Liu, N., Ozbudak, F., Solé, P.: Additive cyclic complementary dual codes over F4. Finite Fields Appl. 83, 102087 (2022).
- [44] Xu H., Du W.: Constructions of symplectic LCD MDS codes. Bull. Malaysian Math. Sci. Soc. 44, 3377-3390 (2021).
- [45] Xu H., Du W.: On some binary symplectic self-orthogonal codes. Appl. Algebra Eng. Commun. Comput. 33(3), 321-337 (2022).
- [46] Zhang G., Chen B.: Self-orthogonal quasi-abelian codes are asymptotically good. Finite Fields Appl. 78, 101958 (2022).

Appendix

Here, we give more details on permutation matrices used in Table 2 as follows:

 P_{46} = (23 28 2 20 32 18 22 3 17 6 46 15 36 27 14 43 16 39 26 38 12 42 4 10 40 31 37 24 7 29 5 19 44 1 41 13 11 45 21 35 30 9 25 33 34 8),

- $P_{52} = (22\ 29\ 50\ 46\ 35\ 37\ 42\ 45\ 18\ 30\ 36\ 2\ 25\ 43\ 10\ 5\ 26\ 11\ 24\ 27$ 20 49 17 23 41 47 7 6 51 48 12 16 34 31 44 15 3 13 52 19 14 33 38 40 9 8 32 39 4 21 28 1),
- $P_{56} = (42 \ 15 \ 52 \ 24 \ 38 \ 11 \ 12 \ 18 \ 6 \ 39 \ 51 \ 53 \ 2 \ 30 \ 46 \ 21 \ 29 \ 3 \ 13 \ 19$ 49 36 48 9 7 31 41 50 40 8 4 25 1 47 34 5 33 45 10 27 14 28 16 23 26 37 54 43 35 32 55 56 44 22 20 17),
- $P_{58} = (11\ 38\ 18\ 51\ 42\ 52\ 19\ 48\ 56\ 55\ 22\ 20\ 7\ 5\ 4\ 21\ 34\ 14\ 40\ 35$ 27 24 41 44 9 10 23 6 8 31 46 39 57 47 36 45 54 3 50 33 49 1 15 58 29 16 26 2 13 25 17 28 12 53 43 37 32 30),
- P_{62} = (1 23 42 44 62 3 25 27 36 48 47 29 11 34 20 16 30 18 51 31 4 52 33 7 57 19 49 58 2 43 8 26 5 37 59 40 6 55 61 9 46 15 21 22 38 32 13 28 53 12 17 24 56 10 60 39 50 35 14 41 45 54),
- $P_{63} = (53\ 29\ 63\ 14\ 44\ 47\ 46\ 4\ 51\ 59\ 11\ 20\ 10\ 23\ 13\ 37\ 42\ 9\ 26\ 34\ 12$ 49 38 30 62 56 16 55 28 33 3 61 40 6 5 35 22 24 52 50 25 7 18 39 36 31 8 21 27 57 17 60 41 58 19 43 54 48 1 32 15 45 2),
- $P_{64} = (48\ 32\ 16\ 55\ 9\ 43\ 23\ 46\ 49\ 10\ 60\ 62\ 40\ 22\ 3\ 38\ 8\ 34\ 59\ 35\ 36\ 30$ 1 6 63 57 5 44 15 33 53 4 25 31 29 45 58 17 64 11 14 24 52 42 18 20 51 37 56 61 41 13 19 7 50 2 12 21 26 54 39 27 47 28),
- P_{70} = (56 23 40 64 63 61 39 41 55 33 13 22 26 17 48 18 36 20 67 46 59 25 19 45 31 51 38 65 50 57 43 60 9 54 10 52 70 14 30 44 62 66 58 5 35 3 42 6 7 21 16 47 53 28 8 49 29 34 4 27 15 37 69 11 32 1 68 12 2 24),
- P_{72} = (36 25 56 57 22 35 29 23 8 31 3 38 19 70 45 68 10 72 53 2 42 58 18 51 61 63 27 54 52 46 13 43 11 62 17 28 71 41 66 47 14 6 12 48 26 65 67 30 40 24 7 33 32 34 64 55 4 50 49 69 37 60 15 21 16 1 44 20 39 9 59 5),
	- $P_{74} = (49 6 67 11 21 36 71 64 52 7 59 10 42 66 28 27 30 43 3 38$ 20 14 16 40 60 15 5 41 54 73 17 19 26 57 62 32 2 53 61 34 9 74 8 1 31 55 22 44 29 68 39 63 33 56 50 13 72 58 69 35 70 51 24 47 4 12 45 48 46 18 25 37 23 65).