

$(1 - \varepsilon)$ -Approximation of Knapsack in Nearly Quadratic Time

Xiao Mao
Stanford University
xiaomao@stanford.edu

Abstract

Knapsack is one of the most fundamental problems in theoretical computer science. In the $(1 - \varepsilon)$ -approximation setting, although there is a fine-grained lower bound of $(n + 1/\varepsilon)^{2-o(1)}$ based on the $(\min, +)$ -convolution hypothesis ([Künnemann, Paturi and Stefan Schneider, ICALP 2017] and [Cygan, Mucha, Wegrzycki and Włodarczyk, 2017]), the best algorithm is randomized and runs in $\tilde{O}\left(n + \left(\frac{1}{\varepsilon}\right)^{11/5} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})}\right)^1$ time [Deng, Jin and Mao, SODA 2023], and it remains an important open problem whether an algorithm with a running time that matches the lower bound (up to a sub-polynomial factor) exists. We answer the question positively by showing a deterministic $(1 - \varepsilon)$ -approximation scheme for knapsack that runs in $\tilde{O}(n + (1/\varepsilon)^2)$ time. We first extend a known lemma in a recursive way to reduce the problem to $n\varepsilon$ -additive approximation for n items with profits in $[1, 2)$. Then we give a simple efficient geometry-based algorithm for the reduced problem.

1 Introduction

Knapsack is one of the most fundamental problems in computer science and mathematical optimization, and is actively being studied in fields such as integer programming and fine-grained complexity. In the Knapsack problem (sometimes also called 0-1 Knapsack), we are given a set I of n items where each item $i \in I$ has weight $w_i > 0$ and profit $p_i > 0$, as well as a knapsack capacity W , and we want to choose a subset $J \subseteq I$ satisfying the weight constraint $\sum_{j \in J} w_j \leq W$ such that the total profit $\sum_{j \in J} p_j$ is maximized.

Knapsack is well-known to be hard: it appeared in Karp’s original list of 21 NP-hard problems [Kar72]. To cope with NP-hardness, a natural direction is to study its *approximation algorithms*. Given a parameter $\varepsilon > 0$, and an input instance with optimal value OPT , a $(1 - \varepsilon)$ -approximation algorithm is required to output a number SOL such that $(1 - \varepsilon)\text{OPT} \leq \text{SOL} \leq \text{OPT}$. Fortunately, Knapsack is well-known to have *fully polynomial-time approximation schemes (FPTASes)*, namely $(1 - \varepsilon)$ -approximation algorithm that runs in $\text{poly}(n, 1/\varepsilon)$ time, for any $\varepsilon > 0$.

¹Throughout this paper, we use $\tilde{O}(f)$ to denote $O(f \cdot \text{poly} \log(n/\varepsilon))$.

Table 1: FPTASes for 0-1 knapsack

$O\left(n \log n + \left(\frac{1}{\varepsilon}\right)^4 \log \frac{1}{\varepsilon}\right)$	Ibarra and Kim [IK75]	1975
$O\left(n \log \frac{1}{\varepsilon} + \left(\frac{1}{\varepsilon}\right)^4\right)$	Lawler [Law79]	1979
$O\left(n \log \frac{1}{\varepsilon} + \left(\frac{1}{\varepsilon}\right)^3 \log^2 \frac{1}{\varepsilon}\right)$	Kellerer and Pferschy [KP04]	2004
$O\left(n \log \frac{1}{\varepsilon} + \left(\frac{1}{\varepsilon}\right)^{5/2} \log^3 \frac{1}{\varepsilon}\right)$ (randomized)	Rhee [Rhe15]	2015
$O\left(n \log \frac{1}{\varepsilon} + \left(\frac{1}{\varepsilon}\right)^{12/5} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})}\right)$	Chan [Cha18]	2018
$O\left(n \log \frac{1}{\varepsilon} + \left(\frac{1}{\varepsilon}\right)^{9/4} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})}\right)$	Jin [Jin19]	2019
$O\left(n \log \frac{1}{\varepsilon} + \left(\frac{1}{\varepsilon}\right)^{11/5} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})}\right)$ (randomized)	Deng, Jin and Mao [DJM23]	2023
$\tilde{O}\left(n + \left(\frac{1}{\varepsilon}\right)^2\right)$	This work	

Reductions showed by Cygan et al. [CMWW19] and Künnemann et al. [KPS17] imply that 0-1 knapsack and unbounded knapsack have no FPTAS in $O\left((n + \frac{1}{\varepsilon})^{2-\delta}\right)$ time, unless $(\min, +)$ -convolution admits a truly subquadratic algorithm [MWW19]. Experts in this field have been in pursuit of an algorithm that matches this lower bound long before such a lower bound is known. The history of FPTASes for the Knapsack problem is summarized in Table 1. The first algorithm with only subcubic dependence on $1/\varepsilon$ was due to Rhee [Rhe15], which reduced the problem to a linear programming instance that can be solved using special randomized methods. In 2018, Chan [Cha18] gave an elegant algorithm for the 0-1 knapsack problem in deterministic $O\left(n \log \frac{1}{\varepsilon} + \left(\frac{1}{\varepsilon}\right)^{5/2} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})}\right)$ via simple combination of the SMAWK algorithm [AKM⁺87] and a standard divide-and-conquer technique. The speedup of superpolylogarithmic factor $2^{\Omega(\sqrt{\log(1/\varepsilon)})}$ was due to progress on $(\min, +)$ -convolution [BCD⁺14, Wil14, CW16]. Using an elementary number-theoretic lemma, Chan further improved the algorithm to $O\left(n \log \frac{1}{\varepsilon} + \left(\frac{1}{\varepsilon}\right)^{12/5} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})}\right)$ time. Jin extended this algorithm by using a greedy lemma and extending the number theoretical lemma to multiple layers, obtaining a running time of $O\left(n \log \frac{1}{\varepsilon} + \left(\frac{1}{\varepsilon}\right)^{9/4} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})}\right)$ [Jin19]. Very recently, Deng, Jin and Mao designed an algorithm that used an improved greedy lemma using a result in additive combinatorics, as well as a procedure based on random partitioning, obtaining a running time of $O\left(n \log \frac{1}{\varepsilon} + \left(\frac{1}{\varepsilon}\right)^{11/5} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})}\right)$, but the algorithm was randomized. In this work, we obtain a running time of $\tilde{O}\left(n + (1/\varepsilon)^2\right)$ deterministically, putting an end to this decades-long effort to match the lower bound. Although our work uses somewhat complicated results from previous work as black boxes at the beginning, our core geometry-based procedure is fairly simple and is quite different from the framework of techniques made popular by Chan.

For the related *unbounded knapsack* problem, where every item has infinitely many copies, Jansen and Kraft [JK18] obtained an $O\left(n + \left(\frac{1}{\varepsilon}\right)^2 \log^3 \frac{1}{\varepsilon}\right)$ -time algorithm; the unbounded version can be reduced to 0-1 knapsack with only a logarithmic blowup in the problem size [CMWW19].

There are also some recent advances for the related problems Subset Sum and Partition. The Subset Sum problem is a special case of Knapsack, where the weight of an item is always equal to its profit. The *partition* problem is an interesting special case of the Subset Sum problem where $W = \frac{1}{2} \sum_{i \in I} w_i$. For Subset Sum, the best known algorithm by Bringmann and Nakos [BN21]

runs in $\tilde{O}(n + \varepsilon^{-2}/2^{\Omega(\sqrt{\log(1/\varepsilon)})})$ time (improving [KMPS03] by low-order factors). Bringmann and Nakos [BN21] showed a matching lower bound based on the $(\min, +)$ -convolution hypothesis. For Partition, the best FPTAS by Deng, Jin and Mao runs in $\tilde{O}(n + (\frac{1}{\varepsilon})^{5/4})$ time [DJM23]. Abboud, Bringmann, Hermelin, and Shabtay [ABHS19] showed that Partition cannot be approximated in $\text{poly}(n)/\varepsilon^{1-\delta}$ time for any $\delta > 0$, under the Strong Exponential Time Hypothesis.

We can see that before this work, the complexity of Subset Sum is already settled, but for Knapsack and Partition there still remain gaps between the best-known algorithm and its conditional lower bound. As we have mentioned, in this paper, we make a breakthrough that closes this gap for the Knapsack problem:

Theorem 1.1. *There is a deterministic algorithm for $(1 - \varepsilon)$ -approximating Knapsack with running time $\tilde{O}(n + \varepsilon^{-2})$.*

2 Preliminaries

We write $\mathbb{N} = \{0, 1, 2, \dots\}$ and $\mathbb{N}^+ = \{1, 2, \dots\}$. For $n \in \mathbb{N}$ we write $[n] = \{1, 2, \dots, n\}$. For a finite set S we let $|S|$ be the size of S (i.e. the number of elements in S). For a sequence $A = \{a_i\}$, we let $|A|$ be the length of A and for an interval $[l, r] \in [1, |A|]$ we write $A_{l,r}$ as a shorthand for the subsequence $\{a_l, a_{l+1}, \dots, a_r\}$.

2.1 Problem Statements

In the Knapsack problem, the input is a list of n items $(p_1, w_1), \dots, (p_n, w_n) \in \mathbb{N} \times \mathbb{N}$ together with a knapsack capacity $W \in \mathbb{N}$, and the optimal value is

$$\text{OPT} := \max_{J \subseteq [n]} \left\{ \sum_{j \in J} p_j \mid \sum_{j \in J} w_j \leq W \right\}.$$

Given a Knapsack instance and a parameter $\varepsilon \in (0, 1)$, an $(1 - \varepsilon)$ -approximation algorithm is required to output a number SOL such that $(1 - \varepsilon)\text{OPT} \leq \text{SOL} \leq \text{OPT}$.

In this problem, we can assume $n = O(\varepsilon^{-4})$ and hence $\log n = O(\log \varepsilon^{-1})$. For larger n , Lawler's algorithm [Law79] for Knapsack in $O(n \log \frac{1}{\varepsilon} + (\frac{1}{\varepsilon})^4)$ time is already near-optimal.

We will sometimes describe algorithms with approximation ratio $1 - O(\varepsilon)$ (or $1 - \tilde{O}(\varepsilon)$), which can be made $1 - \varepsilon$ by scaling down ε by a constant factor (or a logarithmic factor) at the beginning.

2.2 Knapsack Problem and Profit functions

In the knapsack problem, assume $0 < w_i \leq W$ and $p_i > 0$ for every item i . Then a trivial lower bound of the maximum total profit is $\max_j p_j$. At the beginning, we can discard all items i with $p_i \leq \frac{\varepsilon}{n} \max_j p_j$, reducing the total profit by at most $\varepsilon \max_j p_j$, which is only an $O(\varepsilon)$ fraction of the optimal total profit. Therefore, we can assume $\frac{\max_j p_j}{\min_j p_j} \leq \frac{n}{\varepsilon}$.

We adopt the terminology of Chan [Cha18] with some modification. We use the term *profit function* to refer to any monotone non-decreasing step function over some domain range $x \in [l, r]$. The

complexity of a monotone step function refers to the number of its steps. The $(\max, +)$ -convolution of two functions f with domain $x \in [l_f, r_f]$ and g with domain $x \in [l_g, r_g]$ is a function with domain $x \in [l_f + l_g, r_f + r_g]$ and is defined to be:

$$(f \oplus g)(x) = \max_{x_f \in [l_f, r_f], x_g \in [l_g, r_g], x_f + x_g = x} f(x_f) + g(x_g).$$

The contour of a profit function with domain $x \in [x_1, x_k]$ can be described by a set $P(f)$ of $(2k - 1)$ 2-D points

$$\{(x_1, y_1), (x_2, y_1), (x_2, y_2), (x_3, y_2), \dots, (x_k, y_{k-1}), (x_k, y_k)\}$$

with $x_1 < x_2 < \dots < x_k$ and $y_1 < y_2 < \dots < y_k$, where $f(x) = y_i$ for $x_i \leq x < x_{i+1}$ ($1 \leq i \leq n$).

For a set I of items, we use f_I to denote the profit function defined as:

$$f_I(x) = \max \left\{ \sum_{i \in J} p_i : \sum_{i \in J} w_i \leq x, \quad J \subseteq I \right\}$$

over $x \in [0, +\infty)$.

Let I_1, I_2 be two disjoint subsets of items, and $I = I_1 \cup I_2$. It is straightforward to see that $f_I = f_{I_1} \oplus f_{I_2}$.

For each item $i \in I$, we let its *unit profit* be p_i/w_i .

2.3 $(1 - \delta, \Delta)$ approximation up to t

We sometimes uses the notion of $(1 - \delta, \Delta)$ -approximation up to t , as stated below.

Definition 2.1 (Approximation for Profit Functions). For functions \tilde{f}, f with identical domains and real numbers $t, \Delta \in \mathbb{R}_{\geq 0}, \delta \in [0, 1)$, we say that \tilde{f} is a $(1 - \delta, \Delta)$ approximation of f up to t , if for all x in their domain,

$$\tilde{f}(x) \leq f(x)$$

holds, and

$$\tilde{f}(x) \geq (1 - \delta)f(x) - \Delta$$

holds whenever $f(x) \leq t$.

For the case of $t = +\infty$, we simply omit the phrase “up to t .”

We also refer to $(1, \Delta)$ approximation as Δ -additive approximation, and refer to $(1 - \delta, 0)$ approximation as $(1 - \delta)$ -multiplicative approximation, or simply $(1 - \delta)$ approximation.

We now prove the following fact:

Proposition 2.2. *Let I_1, I_2 be two disjoint subsets of items. For $i \in \{1, 2\}$, suppose \tilde{f}_i is a $(1 - \delta, \Delta_i)$ approximation of the profit function f_{I_i} up to t . Then, $f = \tilde{f}_1 \oplus \tilde{f}_2$ is a $(1 - \delta, \Delta_1 + \Delta_2)$ approximation of $f_{I_1 \cup I_2}$ up to t .*

Proof. For any $b \in [0, t]$ where $f(b) = f_1(b_1) + f_2(b_2)$ for $b_1 + b_2 = b$ and $b_1, b_2 \in [0, t]$, we have for $i \in [1, 2]$, $(1 - \delta)f_i(b_i) - \Delta_i \leq f_{I_i}(b_i)$. Since I_1 and I_2 are disjoint, $f_{I_1 \cup I_2}(b) \geq f_{I_1}(b_1) + f_{I_2}(b_2)$. Hence, $f_{I_1}(b_1) + f_{I_2}(b_2) \leq (1 - \delta)f_1(b_1) - \Delta_1 + (1 - \delta)f_2(b_2) - \Delta_2 = (1 - \delta)(f_1(b_1) + f_2(b_2)) - (\Delta_1 + \Delta_2) = (1 - \Delta)f(b) - (\Delta_1 + \Delta_2)$.

The converse direction can be verified similarly. \square

Following Chan [Cha18] and Jin [Jin19], given a monotone step function f with range contained in $\{-\infty, 0\} \cup [A, B]$, one can round the positive values of f down to powers of $1/(1 - \varepsilon)$, and obtain another profit function \tilde{f} which has complexity only $O(\varepsilon^{-1} \log(B/A))$, and $(1 - \varepsilon)$ -approximates f . In our algorithm we will always have $B/A \leq \text{poly}(n/\varepsilon)$, so we may always assume that the intermediate profit functions computed during our algorithm are monotone step functions with complexity $\tilde{O}(\varepsilon^{-1})$ after rounding.

2.4 Technical Overview

We will first reduce our problem of approximating Knapsack to the case where we only have a few items (Section 4.1). Namely, we reduce our problem to the following problem:

Problem 1. *Given a list I of $n = O(1/\varepsilon)$ items with p_i being multiples of ε in interval $[1, 2]^2$, compute a profit function that $(n\varepsilon)$ -additively approximates f_I .*

Lemma 2.3. *Problem 1 can be solved in $\tilde{O}(\varepsilon^{-2})$ time.*

The reduction is based on an extension of the Greedy Exchange Lemma in [DJM23], which reduces the computation to three parts. We observe that one of the parts is identical to the original problem on a smaller scale, and thus we can apply the Greedy Exchange Lemma recursively.

The proof of Lemma 2.3 is the crux of our algorithm (Section 4.2). Recall that profit functions can be described by a series of points. We will show that one can remove some of these points, obtain a new profit function described by these points, and still not lose too much precision (Lemma 4.9). Moreover, our point removal scheme guarantees that the remaining points consist of few blocks of upper or lower convex hulls, and merging profit functions determined by these points can be done more efficiently (Lemma 4.11). Readers can compare these ideas with the ideas based on “sparsification” described in Bringmann and Nako’s FPTAS for Subset Sum [BN21], which are much simpler analogs of our ideas in 1D.

3 Summary of Previous Techniques

3.1 Known Lemmas

By known reductions (e.g., [Cha18, Jin19, DJM23]), we can focus on solving the following cleaner problem, which already captures the main difficulty of knapsack:

²The input profit values in the Knapsack problem are integers, but in intermediate problems, the profits can be real numbers due to rounding.

Problem 2. Assume $\varepsilon \in (0, 1/2)$ and $1/\varepsilon \in \mathbb{N}^+$. Given a list I of items $(p_1, w_1), \dots, (p_n, w_n)$ with weights $w_i \in \mathbb{N}$ and profits p_i being multiples of ε in the interval $[1, 2)$, compute a profit function that $(1 - \varepsilon)$ -approximates f_I up to $2/\varepsilon$.

Lemma 3.1. If for some $c \geq 2$, Problem 2 can be solved in $\tilde{O}(n + 1/\varepsilon^c)$ time, then $(1 - \varepsilon)$ -approximating Knapsack can also be solved in $\tilde{O}(n + 1/\varepsilon^c)$ time.

The proof of Lemma 3.1 can be seen in the appendix.

The following useful lemma allows us to merge multiple profit functions, which was proven by Chan using divide-and-conquer and improved algorithms for $(\min, +)$ -convolution [BCD⁺14, Wil14, CW16].

Lemma 3.2 ([Cha18, Lemma 2(i)]). Let f_1, \dots, f_m be monotone step functions with total complexity N and ranges contained in $\{-\infty, 0\} \cup [A, B]$. Then we can compute a monotone step function that has complexity $\tilde{O}(\frac{1}{\varepsilon} \log B/A)$ and $(1 - O(\varepsilon))$ -approximates $f_1 \oplus \dots \oplus f_m$, in $O(N) + \tilde{O}((\frac{1}{\varepsilon})^2 m / 2^{\Omega(\sqrt{\log(1/\varepsilon)})} \log B/A)$ time.

The following reduction was used in [DJM23].

Lemma 3.3. Given a list I of n items with p_i being multiples of ε in interval $[1, 2)$, and integer $1 \leq m \leq n$ with $m = O(1/\varepsilon)$, if for some $c \geq 2$ one can compute a profit function that $(m\varepsilon)$ -additively approximates f_I up to $2m$ in $\tilde{O}(n + 1/\varepsilon^c)$ time. Then one can $(1 - \varepsilon)$ -approximate Knapsack in $\tilde{O}(n + 1/\varepsilon^c)$ time.

Proof. From Lemma 3.1, it suffices to approximate Problem 2. We divide $[1, 2\varepsilon^{-1})$ into $O(\log(1/\varepsilon))$ many intervals $[m, 2m)$ where m 's are powers of 2, and for each m , compute profit functions f_m achieving $m\varepsilon$ -additive approximation up to $2m$. For each $x \geq 0$, let m_x be such that $m_x \leq f_I(x) < 2m_x$. Then $f_{m_x}(x) \in [f_I(x) - m_x\varepsilon, f_I(x)]$. Since $f_I(x) \geq m_x$, $f_{m_x}(x) \in [f_I(x)(1 - \varepsilon), f_I(x)]$. Thus taking the pointwise maxima of all f_m 's yields a $(1 - \varepsilon)$ approximation of f_I . Note that by using the rounding scheme mentioned before we can make the complexity of each profit function $\tilde{O}(1/\varepsilon)$. Therefore taking the pointwise maxima of all f_m 's takes $O(\log(1/\varepsilon)) \times \tilde{O}(1/\varepsilon) = o(n + 1/\varepsilon^c)$ time. \square

As an ingredient, we use a corollary of Lemma 7 of [Cha18]. The corollary states that profit functions can be approximately more efficiently when there are few distinct profit values or when there is a good upper bound on profit values. We first present the lemma in its original form. A step function f is p -uniform if the function values are of the form $-\infty, 0, p, 2p, \dots, lp$ for some non-negative integer l . A p -uniform function is pseudo-concave if the sequence of differences of consecutive x -breakpoints is non-decreasing.

Lemma 3.4 (Lemma 7 of [Cha18]). Let f_1, \dots, f_m be monotone step functions with ranges contained in $\{-\infty, 0\} \cup [1, B]$. If every f_i is p_i -uniform and pseudo-concave for some $p_i \in [1, 2]$, then we can compute a monotone step function that $(1 - \varepsilon)$ -approximates $\min(f_1 \oplus \dots \oplus f_m, B)$ and complexity $\tilde{O}(\frac{1}{\varepsilon})$ in $\tilde{O}(\sqrt{B}m/\varepsilon)$ deterministic³ time, assuming $B = \tilde{O}(1/\varepsilon)$.

³The original lemma used the word "expected" since it was initially randomized due to a randomized construction for a number theoretical lemma, but Chan later derandomized it in the same paper. A simpler deterministic construction can be seen in Lemma 3.11 of [DJM23].

Corollary 3.5. *Given a list I of items $(p_1, w_1), \dots, (p_n, w_n)$ with weights $w_i \in \mathbb{N}$ and profits p_i being multiples of ε in the interval $[1, 2)$, if there are only Δ distinct profit values p_i , then for any $B = \tilde{O}(1/\varepsilon)$, one can $(1 - \varepsilon)$ -approximate $\min(f_I, B)$ (i.e. f_I up to B) in $\tilde{O}(n + \Delta\sqrt{B}\varepsilon^{-1})$ time.*

Proof. Note that $I = I_1 \cup I_2 \cup \dots \cup I_\Delta$ where each I_j contains items of the same profit value. It suffices to compute $\min(f_{I_1} \oplus \dots \oplus f_{I_\Delta}, B)$. Fix $j \in [1, \Delta]$. If we let w'_k be the k -th smallest weight in I_j , then:

- For every $0 \leq k \leq |I_j| - 1$, for $x \in [\sum_{1 \leq k' \leq k} w'_{k'}, \sum_{1 \leq k' \leq k+1} w'_{k'})$ we have $f_{I_j}(x) = kp_j$.
- $f_{I_j}(\sum_{1 \leq k' \leq |I_j|} w'_{k'}) = |I_j|p_j$.

It is easy to see that f_{I_j} is p_j -uniform and pseudo-concave with $p_j \in [1, 2]$. Furthermore, for each j it suffices to compute $\min(f_{I_j}, B)$, whose range is contained in $\{-\infty, 0\} \cup [1, B]$. From Lemma 3.4 we can compute $\min(f_{I_1} \oplus \dots \oplus f_{I_\Delta}, B)$ in $\tilde{O}(\sqrt{B}m/\varepsilon)$ time. Constructing $\min(f_{I_j}, B)$ all j 's takes $\tilde{O}(n)$ time. The total time complexity is $\tilde{O}(n + \Delta\sqrt{B}/\varepsilon)$. \square

3.2 Greedy Exchange Lemma

We now introduce the Greedy Exchange Lemma previously employed in [DJM23]. We will follow the framework in [DJM23]. Given items $(p_1, w_1), \dots, (p_n, w_n)$, where $p_i \in [1, 2)$ are multiples of ε , we sort them by non-increasing order of unit profit: $p_1/w_1 \geq p_2/w_2 \geq \dots \geq p_n/w_n$.

Then, we consider prefixes of this sequence of items, and define the following measure of diversity:

Definition 3.6 ($D(i)$). For $1 \leq i \leq n$, let $D(i) = \min_J C([i] \setminus J)$, where the minimization is over all subsets $J \subseteq [i]$ with $|J| \leq 2m$, and $C([i] \setminus J)$ denote the number of distinct values in $\{p_j : j \in [i] \setminus J\}$.

We have the following immediate observations about $D(i)$:

Observation 3.7. 1. For all $2 \leq i \leq n$, $0 \leq D(i) - D(i-1) \leq 1$.

2. $D(i)$ (and the minimizer J) can be computed in $\tilde{O}(i)$ time by the following greedy algorithm: start with all values p_1, p_2, \dots, p_i . Repeat the following up to $2m$ times: remove the value p_j with the minimum multiplicity, and add j into J .

Definition 3.8 (i^Δ). For some parameter Δ , define $i^\Delta \in \{1, 2, \dots, n\}$ be the maximum such that $D(i^\Delta) \leq \Delta$, which can be found using Observation 3.7 with a binary search in $\tilde{O}(n)$ time.

In [DJM23], the following lemma was proven. Recall that Corollary 3.5 guarantees a faster run time when we have a bound on profit values. The goal of the lemma is to impose a bound on the total profit items in $[n] \setminus [i^\Delta]$ can contribute. On a high level, it argues that, given any subset S of items, we can swap out some items in $S \cap ([n] \setminus [i^\Delta])$ with items in $[i^\Delta]$, resulting in a new subset \tilde{S} that is not too much worse than S , but the contribution from $[n] \setminus [i^\Delta]$ to \tilde{S} is small.

Lemma 3.9. [Greedy Exchange Lemma, Lemma 3.9 of [DJM23]] Let $S \subseteq [n]$ be any item set with total profit $\sum_{s \in S} p_s \leq 2m$. Let $\Delta = \omega(\varepsilon^{-1/2} \log^{1/2}(1/\varepsilon))$ be some parameter. Let $B := c\varepsilon^{-1}/\Delta$ for some universal constant $c \geq 1$.

Then, there exists an item set $\tilde{S} \subseteq [n]$, such that the total profit \tilde{p} contributed by items $[n] \setminus [i^\Delta]$ in \tilde{S} satisfies

$$\tilde{p} := \sum_{s \in \tilde{S} \cap ([n] \setminus [i^\Delta])} p_s \leq B, \quad (1)$$

and

$$\sum_{s \in \tilde{S}} p_s \geq (1 - \varepsilon) \sum_{s \in S} p_s, \quad (2)$$

and

$$\sum_{s \in \tilde{S}} w_s \leq \sum_{s \in S} w_s. \quad (3)$$

The proof is included in the appendix for completeness.

In [DJM23], we have $\Delta := \lfloor \varepsilon^{-5/8} \rfloor$, which satisfies the condition in Lemma 3.9.

The reduction in [DJM23] uses the following lemma for $(1 - \varepsilon)$ -approximating knapsack up to a small B :

Lemma 3.10 (Follows from Lemma 17 of [Jin19]). Given a list I of items $(p_1, w_1), \dots, (p_n, w_n)$ with weights $w_i \in \mathbb{N}$ and profits p_i being multiples of ε in the interval $[1, 2)$, one can $(1 - \varepsilon)$ -approximate the profit function f_I up to B in $\tilde{O}(n + \varepsilon^{-2} B^{1/3} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})})$ time.

Finally, a weaker version of Lemma 2.3 was proven in [DJM23] using random partitioning.

Lemma 3.11. [Lemma 3.6 of [DJM23]]

Problem 1 can be solved in randomized $\tilde{O}(n^{4/5} \varepsilon^{-7/5} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})})$ time.

In [DJM23], a randomized running time of $\tilde{O}(n + \varepsilon^{-11/5} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})})$ was achieved for Problem 2 by combining the following lemma with Lemma 3.3. To finish our introduction for techniques in [DJM23], we will also include its proof.

Lemma 3.12 (Lemma 3.5 of [DJM23]). Given a list I of n items with p_i being multiples of ε in interval $[1, 2)$, and integer $1 \leq m \leq n$ with $m = O(1/\varepsilon)$, one can compute a profit function that $(m\varepsilon)$ -additively approximates f_I up to $2m$ in $O(n + \varepsilon^{-11/5} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})})$ time.

Proof of Lemma 3.12. Recall that we have $\Delta := \lfloor \varepsilon^{-5/8} \rfloor$, and that $i^\Delta \in \{1, 2, \dots, n\}$ is the maximum such that $D(i^\Delta) \leq \Delta$, which can be found using Observation 3.7 with a binary search in $\tilde{O}(n)$ time. Let $J \subset [i^\Delta]$ with $|J| \leq 2m$ be the minimizer for $D(i^\Delta)$.

Now, we approximately compute the profit functions $f_J, f_{[i^\Delta] \setminus J}, f_{[n] \setminus [i^\Delta]}$ for three item sets $J, [i^\Delta] \setminus J, [n] \setminus [i^\Delta]$ using different algorithms, described as follows:

1. Use Lemma 3.11 to compute f_1 that $(2m\varepsilon)$ -additively approximates f_J in randomized $O(m^{4/5} \varepsilon^{-7/5} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})}) \leq O(\varepsilon^{-11/5} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})})$ time.

2. By the definition of i^Δ , items in $[i^\Delta] \setminus J$ have no more than Δ distinct profit values. Hence we can use Corollary 3.5 to compute f_2 that $(1 - \varepsilon)$ -approximates $f_{[i^\Delta] \setminus J}$ up to $2m$, in $\tilde{O}(n + \Delta\sqrt{2m\varepsilon^{-1}}) = \tilde{O}(\varepsilon^{-17/8})$ time.
3. Use Lemma 3.10 to compute f_3 that $(1 - \varepsilon)$ -approximates the $f_{[n] \setminus [i^\Delta]}$ up to $B = c\varepsilon^{-1}/\Delta = \Theta(\varepsilon^{-1}/\Delta)$ where c is the universal constant in Lemma 3.9, in $\tilde{O}(B^{1/3}\varepsilon^{-2}) \leq \tilde{O}(\varepsilon^{-17/8})$ time.

Finally, merge the three parts f_1, f_2, f_3 using Lemma 3.2 in $\tilde{O}(\varepsilon^{-2})$ time, and return the result.

In the third part, the correctness of only computing up to B is justified by Lemma 3.9, which shows that if we only consider approximating sets with total profit up to $2m$, then we can assume the items in $[n] \setminus [i^\Delta]$ only contributes profit at most B (1), at the cost of only incurring an $(1 - \varepsilon)$ approximation factor (2).

To analyze the error, notice that in the first part we incur an additive error of $2m\varepsilon$. In the second and third part and the final merging step we incur $(1 - O(\varepsilon))$ multiplicative error, which turns into $O(m\varepsilon)$ additive error since we only care about approximating up to $2m$. Hence the overall additive error is $O(m\varepsilon)$, which can be made $m\varepsilon$ by lowering the value of ε . \square

Our new algorithm will use a new reduction that uses the Greedy Exchange Lemma recursively. Intuitively, observe that the computation of f_3 is a reduced version of the original problem with a stricter upper bound, so instead of computing it directly, we can apply the procedure recursively.

3.3 Past Algorithms for Knapsack

3.3.1 Chan's Framework

We can easily see that the extra running time incurred by the reduction in Lemma 3.12 on top of solving Problem 1 is $\tilde{O}(\varepsilon^{-17/8})$, which is dominated by the factor $\varepsilon^{-11/5}$ in Lemma 3.11. Therefore, the most important reason that we are able to obtain a speedup in our new algorithm is due to the improvement from Lemma 3.11 to Lemma 2.3, namely a better running time for Problem 1:

Problem 1. *Given a list I of $n = O(1/\varepsilon)$ items with p_i being multiples of ε in interval $[1, 2)$ ⁴, compute a profit function that $(n\varepsilon)$ -additively approximates f_I .*

Improving the running time for Problem 1 and for Knapsack have been very challenging tasks in the recent years. This is mainly due to the inability to go around the trade-off between two different approaches: the sequential approach that is efficient at the cost of precision, and the divide-and-conquer approach that is precise at the cost of efficiency. We now briefly introduce them and will limit our discussion on Problem 1 for now for simplicity.

In the sequential approach, we go through items in I one by one and update the profit function. The classical dynamic programming solution to *knapsack* can be seen as an analog in the deterministic case. Suppose the total profit value of items seen so far is $P_i := \sum_{j \leq i} p_j$. The profit function for the first i items has a complexity of $O(P_i)$ and updating the profit function with the

⁴The input profit values in the Knapsack problem are integers, but in intermediate problems, the profits can be real numbers due to rounding.

addition of a single item can easily be done in $O(P_i)$ time. In the approximate case, we perform rounding on the profit function after each addition. Since errors from sequential rounding add up, to guarantee a final additive error of $n\varepsilon$, one has to round to multiples of ε . Since $P_i = O(n)$, this unfortunately makes the complexities of profit functions $O(n\varepsilon^{-1})$ and the total running time becomes $O(n^2\varepsilon^{-1}) = O(\varepsilon^{-3})$.

An alternative method is to use divide and conquer. Without loss of generality assume that $|I|$ is a power of 2, We divide I into two equal parts $I = X \cup Y$. We approximate the profit functions f_X and f_Y recursively and approximate $f_I = f_X \oplus f_Y$. Every time we are in a sub-problem with total profit value P , we round to multiples of $P\varepsilon$, and we can show that the total additive error is $O(n\varepsilon \log n) = \tilde{O}(n\varepsilon)$. Thus we can keep the complexities of profit functions $O(\varepsilon^{-1})$. However, we have to merge two profit functions of complexity $O(\varepsilon^{-1})$ for each of the $O(n)$ sub-problems in the entire recursion tree, which is a $(\max, +)$ -convolution that generally takes quadratic time. Therefore we still only have a running time of $\tilde{O}(n\varepsilon^{-2}) = \tilde{O}(\varepsilon^{-3})$.

As shown by Chan [Cha18], assuming that $n = O(\varepsilon^{-1})$, one can achieve a running time of $\tilde{O}(\varepsilon^{-2.5})$ by combining these two ideas: divide I into $\varepsilon^{-0.5}$ chunks of size $O(\varepsilon^{-0.5})$, solve for each chunk sequentially and use divide-and-conquer to merge the answers for chunks. Chan's original version was a slightly more complicated algorithm directly for Knapsack with a running time of $\tilde{O}(n + \varepsilon^{-2.5})$ as the reduction of the form of Lemma 3.12 did not exist back then.

Obtaining further improvements for either Problem 1 or Knapsack has been an arduous endeavor, and such improvements have all been based on Chan's framework. In the same work, Chan observed that in the sequential method, instead of repeated rounding at each item, if all p_i 's are close to multiples of some large number p' , we can simply round everything to multiples of p' at the beginning. Chan employed a number theoretical lemma stating that it is possible to group items into groups where items in each group is close to multiples of some large number, and achieved a running time of $\tilde{O}(n + \varepsilon^{-12/5})$ for Knapsack and $\tilde{O}(\varepsilon^{-7/3})$ for Problem 1. Jin [Jin19] extended the technique to multiple layers and combined it with greedy, resulting in a running time of $\tilde{O}(n + \varepsilon^{-9/4})$ for Knapsack and $\tilde{O}(\varepsilon^{-9/4})$ for Problem 1. The reduction in Lemma 3.12 was proposed in the algorithm by Deng, Jin and Mao [DJM23], which used Jin's previous results as ingredients. They also used a randomized partitioning technique for limiting the scope of $(\max, +)$ -convolution to achieve a running time of $\tilde{O}(\varepsilon^{-11/5})$ for Problem 1 and a running time of $\tilde{O}(n + \varepsilon^{-11/5})$ for Knapsack. As we can see, not only have improvements been more and more marginal, but the algorithms that achieve them have also increased in complexity.

3.3.2 Sparsification for Subset Sum and Its Potential on Knapsack

Researchers have struggled much less on the related problem of approximating subset-sum. The first algorithm with a running time of $\tilde{O}(n/\varepsilon)$ was known in the 70s [GL79], and the first algorithm that achieves a running time of $\tilde{O}(n + \varepsilon^{-2})$ was known in 1997 [KPS97]. Recently, Bringmann and Nakos [BN21] improved the running time to $\tilde{O}(n + \varepsilon^{-2}/2^{\Omega(\sqrt{\log(1/\varepsilon)})})$ and showed a matching lower bound based on the $(\min, +)$ -convolution hypothesis. Bringmann and Nakos showed the sparsification technique for reducing the size of sets of subset-sums we need to store without losing any precision, which, if employed to knapsack, could give us a best-of-both-worlds solution that achieves efficiency and precision at the same time.

We will give a brief introduction of this technique. For simplicity, we only discuss additive approximation. A Δ -additive approximation of a set S is a set S' such that $\forall x \in S, \exists x' \in S', x - \Delta \leq x' \leq x$ and $\forall x' \in S', \exists x \in S, x - \Delta \leq x' \leq x$. Let S be a set of numbers. To “sparsify” S , we recursively do the following: if there exists $u, v, w \in S$ such that $u < v < w \leq u + \Delta$, remove v from S . In the resulting set, for every three consecutive numbers $u < v < w$, we have $w - u > \Delta$. Thus if $S \subset [0, B]$, then the sparsification S' has a size of $O(B/\Delta)$. Suppose we have two sets S_X and S_Y representing possible subset sums of two disjoint sets X and Y , and we want to approximate $\{x + y \mid x \in S_X, y \in S_Y\}$ which is the set of possible subset sums of $X \cup Y$. Suppose the sparsifications of S_X and S_Y are \hat{S}_X and \hat{S}_Y . It is proven in [BN21] that the set $\{x + y \mid x \in \hat{S}_X, y \in \hat{S}_Y\}$ is a Δ -approximation (rather than 2Δ -approximation) of the set $\{x + y \mid x \in S_X, y \in S_Y\}$. In other words, individual rounding errors do not accumulate.

Unfortunately, the simplest analog of sparsification for profit function fails to prevent accumulation of rounding errors. Suppose that to “sparsify” a profit function f , we recursively do the following: if there exist three consecutive x -breakpoints u, v, w such that $f(u) < f(v) < f(w) \leq f(u) + \Delta$, we remove the x -breakpoint at v so that in the sparsification f' , $f'(i) = f(u)$ for $i \in [u, w)$. In the following example, the sparsification f' for profit functions f is such that $f' \oplus f'$ is only a $(2\Delta - 2)$ -additive approximation for $f \oplus f$. Suppose $f(x) = -\infty$ for $x < 0$, $f(x) = 0$ for $0 \leq x < 1$, $f(x) = \Delta - 1$ for $1 \leq x < 2$, and $f(x) = \Delta$ for $x \geq 2$. In the sparsification f' , the x -breakpoint for at $x = 1$ is removed and we have $f'(x) = -\infty$ for $x < 0$, $f'(x) = 0$ for $0 \leq x < 2$, and $f'(x) = \Delta$ for $x \geq 2$. Then we have $(f \oplus f)(2) = 2\Delta - 2$ but $(f' \oplus f')(2) = 0$.

The core idea in this paper is a sophisticated way of sparsifying profit functions without accumulation of rounding errors. The sparsified results can be described by a small amount of convex hulls on a 2D plane. We will use techniques in computational geometry to merge these convex hulls, resulting in the final quadratic running time.

4 Main Algorithm

4.1 Reduction to the Sparse Case via Recursive Greedy Exchange

In this subsection, we will prove the following:

Lemma 4.1. *Given a list I of n items with p_i being multiples of ε in interval $[1, 2)$, and integer $1 \leq m \leq n$ with $m = O(1/\varepsilon)$, one can compute a profit function with complexity $\tilde{O}(\varepsilon^{-1})$ that $(m\varepsilon)$ -additively approximates f_I up to $2m$ in $\tilde{O}(n + \varepsilon^{-2})$ time.*

Lemma 4.1 implies Theorem 1.1 due to Lemma 3.3. Our core ingredient is Lemma 2.3 which will be proved in Section 4.2:

Problem 1. *Given a list I of $n = O(1/\varepsilon)$ items with p_i being multiples of ε in interval $[1, 2)$, compute a profit function that $(n\varepsilon)$ -additively approximates f_I .*

Lemma 2.3. *Problem 1 can be solved in $\tilde{O}(\varepsilon^{-2})$ time.*

Also recall Corollary 3.5:

Corollary 3.5. *Given a list I of items $(p_1, w_1), \dots, (p_n, w_n)$ with weights $w_i \in \mathbb{N}$ and profits p_i being multiples of ε in the interval $[1, 2)$, if there are only Δ distinct profit values p_i , then for any $B = \tilde{O}(1/\varepsilon)$, one can $(1 - \varepsilon)$ -approximate $\min(f_I, B)$ (i.e. f_I up to B) in $\tilde{O}(n + \Delta\sqrt{B}\varepsilon^{-1})$ time.*

To prove Lemma 4.1, we will use the proof in Lemma 3.12 in a recursive manner. We solve the following problem for a variety of parameters B . Assume that $C > 0$ is a universal large constant.

Problem 3. *Given a list I of n items with p_i being multiples of ε in interval $[1, 2)$, integer $1 \leq m \leq n$ with $m = O(1/\varepsilon)$, integer $k > 0$, as well as a number $1 \leq B \leq 2m$, compute a profit function that $(Ckm\varepsilon)$ -additively approximates f_I up to B .*

Lemma 4.2. *Let $T(n, m, B, k)$ be the running time for Problem 3 with parameters n, m, k and B . Then there exists $B' = o(B^{1/2})$ and $n' \leq n$ such that:*

$$T(n, m, B) = \tilde{O}(n + \varepsilon^{-2}) + T(n', m, B').$$

Proof. To solve the problem for (n, m, k, B) , we set

$$\Delta := \Theta((\varepsilon^{-1}/B^{1/2}) \log(1/\varepsilon)) = \omega(\varepsilon^{-1/2} \log^{1/2}(1/\varepsilon)).$$

Recall Definition 3.6, Observation 3.7 and Definition 3.8. $i^\Delta \in \{1, 2, \dots, n\}$ is the maximum such that $D(i^\Delta) \leq \Delta$, which can be found using with a binary search in $\tilde{O}(n)$ time. Let $J \subset [i^\Delta]$ with $|J| \leq 2m$ be the minimizer for $D(i^\Delta)$.

As in the proof of Lemma 3.12, we approximately compute the profit functions $f_J, f_{[i^\Delta] \setminus J}, f_{[n] \setminus [i^\Delta]}$ for three item sets $J, [i^\Delta] \setminus J, [n] \setminus [i^\Delta]$ using different algorithms, described as follows:

1. Use Lemma 2.3 to compute f_1 that $(2m\varepsilon)$ -additively approximates f_J , in $\tilde{O}(\varepsilon^{-2})$ time.
2. By definition of i^Δ , items in $[i^\Delta] \setminus J$ have no more than Δ distinct profit values. Since we only need to approximate up to B , we can use Corollary 3.5 to compute f_2 that $(1 - \varepsilon)$ -approximates $\min(f_{[i^\Delta] \setminus J}, B)$ in $\tilde{O}(n + \Delta\sqrt{B}\varepsilon^{-1}) = \tilde{O}(n + \varepsilon^{-2})$ time.
3. We $C(k - 1)m\varepsilon$ -additively approximate $\min(f_3, B')$ for $B' = c\varepsilon^{-1}/\Delta = o(B^{1/2})$ where c is the universal constant in Lemma 3.9. This is equivalent to Problem 3 with parameters $(n', m, k - 1, B')$ where $n' = n - i^\Delta$, and can be done in $T(n', m, k - 1, B')$ time.

Finally, merge the three parts f_1, f_2, f_3 using Lemma 3.2 in $\tilde{O}(n + \varepsilon^{-2})$ time, and return the result. The total running time is $\tilde{O}(n + \varepsilon^{-2}) + T(n', m, B')$.

In the third part, the correctness of only computing up to B' is justified by Lemma 3.9, which shows that if we only consider approximating sets with total profit up to $2m \geq B$, then we can assume the items in $[n] \setminus [i^\Delta]$ only contributes profit at most B' (Inequality 1), at the cost of only incurring an $(1 - \varepsilon)$ -approximation factor (Inequality 2).

To analyze the error, notice that in the first part, we incur an additive error of $2m\varepsilon$. In the second part and the final merging step we incur $(1 - O(\varepsilon))$ multiplicative error, which turns into $O(m\varepsilon)$ additive error since we only care about approximating up to $B \leq 2m$. Hence the additive error for these parts is $O(m\varepsilon)$. By letting the constant C be large we can assume that the overall additive error is at most $Cm\varepsilon$. Finally, the third part incurs an additive error of $C(k - 1)m\varepsilon$. Hence the overall additive error is at most $C(k - 1)m\varepsilon + Cm\varepsilon = Ckm\varepsilon$. \square

We are now ready to prove Lemma 4.1

Proof of Lemma 4.1. Note that Problem 3 is trivial when $B < 2$. Note that for any value no more than $2m$, by recursively taking its square root at most $O(\log \log m)$ times, we get a value less than 2. Note that in Lemma 4.2 we always have $B' < B^{1/2}$. Thus the running time for $T(n, m, k, 2m)$ is a summation of at most $O(\log \log m)$ repetitions of the term $\tilde{O}(n + \varepsilon^{-2})$, as long as k is larger than the number of repetitions. Hence, for $k = \omega(\log \log m)$ we have:

$$T(n, m, k, 2m) \leq O(\log \log m) \times \tilde{O}(n + \varepsilon^{-2}) = \tilde{O}(n + \varepsilon^{-2}).$$

Namely, we can solve Problem 3 in $\tilde{O}(n + \varepsilon^{-2})$ time for $(n, m, k, 2m)$ as long as $k = \omega(\log \log m)$. Thus for $k = O(\log(1/\varepsilon))$, we can get an $O(m\varepsilon \log(1/\varepsilon))$ -additive approximation for the problem in Lemma 4.1 $\tilde{O}(n + \varepsilon^{-2})$ time, which can be made into an $m\varepsilon$ -additive approximation by lowering the value of ε by a polylogarithmic factor. \square

4.2 Approximation for the Sparse Case via Geometry

4.2.1 Definitions

In this section, we prove our core lemma:

Lemma 2.3. *Problem 1 can be solved in $\tilde{O}(\varepsilon^{-2})$ time.*

First we recall the relevant definitions. The contour of a profit function with domain $x \in [x_1, x_k]$ with a complexity of k can be described by a set⁵ $P(f)$ of $(2k - 1)$ 2-D points

$$\{(x_1, y_1), (x_2, y_1), (x_2, y_2), (x_3, y_2), \dots, (x_k, y_{k-1}), (x_k, y_k)\}$$

with $x_1 < x_2 < \dots < x_k$ and $y_1 < y_2 < \dots < y_k$, where $f(x) = y_i$ for $x_i \leq x < x_{i+1}$ ($1 \leq i \leq k$).

A set of points $P = \{(x_1, y_1), (x_2, y_2), \dots, (x_k, y_k)\}$ satisfying $x_i \leq x_{i+1}$ and $y_i \leq y_{i+1}$ for $1 \leq i \leq k - 1$ and $x_{i+2} > x_i$ for $1 \leq i \leq k - 2$ is called a *monotone point set*. The profit function $f(P)$ with domain $[x_1, x_k]$ is defined as $(f(P))(x) = y_i$ for $x_i \leq x < x_{i+1}$ for $1 \leq i < k$ and $(f(P))(x_k) = y_k$.

For a set S of points in the 2D plane, let $\min_x S$, $\max_x S$, $\min_y S$ and $\max_y S$ be the minimum x-coordinate, the maximum x-coordinate, the minimum y-coordinate, and the maximum y-coordinate of points in S .

For a point p on the 2D plane, we let p_x and p_y be its x-coordinate and y-coordinate respectively. For two distinct points p and q , we say $p < q$ if either $p_x < q_x$ or $p_x = q_x$ and $p_y < q_y$. Given two points p and q , let $r := p + q$ be the point with $r_x = p_x + q_x$ and $r_y = p_y + q_y$, and let $r := p - q$ be the point with $r_x = p_x - q_x$ and $r_y = p_y - q_y$. Given a point p and a point set S , let $p + S := S + p := \{p + s \mid s \in S\}$. Given two sets of points A and B , their *Minkowski sum* is defined as $A + B = \{a + b \mid a \in A, b \in B\}$.

For a monotone point set P , the set of segments $E(P)$ are the line segments between adjacent points in P : $E(P) := \{(p_i, p_{i+1}) \mid 1 \leq i < |P|\}$. The region $R(P)$ is defined to be the set of points

⁵A set must contain distinct elements.

$(x, y)(x_1 \leq x \leq x_k, y \geq 0)$ on or below the segments $E(P)$. For $x_1 \leq x \leq x_k$, let $g[P](x)$ be equal to the maximum y -coordinate of a point on the upper boundary of $R(P)$ with x -coordinate equal to x . For any $1 \leq i \leq |P|$, if $i \in \{1, |P|\}$ or the point p_i is above the line segment connecting p_{i-1} and p_{i+1} , we call the point p_i an *apex* of P . If $i \notin \{1, |P|\}$ and the point p_i is below the line segment connecting p_{i-1} and p_{i+1} , we call the point p_i a *base* of P . If $i \notin \{1, |P|\}$ and the point p_i is on the line segment connecting p_{i-1} and p_{i+1} , we call the point p_i *degenerate*.




Figure 1: Monotone Point Set $P = \{A, B, C, D, E, F, G\}$

For example, in Fig. 1, the set $P = \{A, B, C, D, E, F, G\}$ is a monotone point set. The region $R(P)$ is the shaded region below the thick red curve, which is the plot of $g[P]$. The plot of $f(P)$ is the thin blue line. For instance, C is an apex since it is above the line connecting B and D , and B is a base since it is below the line connecting A and C . We will see an example of a degenerate point later.

Given two monotone point sets P_a and P_b , let $P_a \oplus P_b$ be the minimum set V of points such that $R(V) = R(P_a) + R(P_b)$. It is easy to verify that V is a monotone point set without degenerate points.

It is well-known [dBvKOS00] that the Minkowski sum of two polygons P and Q with $|P|$ and $|Q|$ vertices is the union of $|P|$ translations of Q and $|Q|$ translations of P consisting of:

- $p + Q$ for each $p \in P$, and
- $P + q$ for each $q \in Q$.

It is also easy to verify that $g[P_a \oplus P_b] = g[P_a] \oplus g[P_b]$.

4.2.2 Roadmap

We will show an analog of the “sparsification” technique in [BN21] in 2D. We can approximate a profit function f with a monotone point set. To start, we can see that $P(f)$ can approximate f exactly. We now show a way to reduce the monotone point set associated with a profit function so that we can still approximate the profit function with the reduced monotone point set, and the rounding errors do not accumulate when combining two profit functions and their monotone point set approximations.

Definition 4.3 (F -generatability). For a parameter $\delta := \varepsilon n$, and a set of m profit functions $F = \{f_1, f_2, \dots, f_m\}$, we define the notion of F -generatability on a pair consisting of a profit function and a monotone point set based on the three operations below:

1. For each $f \in F$, $(f, P(f))$ is F -generatable.
2. If (f_a, P_a) and (f_b, P_b) are both F -generatable, $(f_a \oplus f_b, P_a \oplus P_b)$ is also F -generatable;
3. Suppose (f, P) is F -generatable. Suppose we perform a *reduction* on P : find two apexes $x, y \in P$ on P with $0 < y_y - x_y \leq \delta$ such that all points strictly between x and y are below the line segment connecting x and y , let \bar{P} be P with all points between x and y removed. Let P' be the minimum subset of \bar{P} with $R(P') = R(\bar{P})$ (i.e. \bar{P} with degenerate points removed). Then the new pair (f, P') is F -generatable.

To explain Operation 3, suppose $P = \{A, B, C, D, E, F, G\}$ as shown in Fig. 1, and suppose δ is equal to the difference in y-coordinates between C and F . For $(x, y) = (C, F)$, since both D and E are below the line segment connecting C and F , we can remove them and get $\bar{P} = \{A, B, C, F, G\}$. Note that since F is on the segment connecting C and G , it is a degenerate point of \bar{P} and is removed from \bar{P} to obtain $P' = \{A, B, C, G\}$. We can not perform Operation 3 for $(x, y) = (A, F)$ despite all points between them being below the segment since $F_y - A_y > \delta$.

It is easy to see that if (f, P) is F -generatable for some F , then P contains no degenerate points.

Solving Problem 1 is equivalent to combining n profit functions $\bar{F} = \{\bar{f}[1], \bar{f}[2], \dots, \bar{f}[n]\}$ where $\bar{f}[i]$ has a domain of $[0, w_i]$ and $\bar{f}[i](x) := 0$ for $0 \leq x < w_i$ and $\bar{f}[i](x) := p_i$ for $x = w_i$. We can see that $f_I = \bar{f}[1] \oplus \bar{f}[2] \oplus \dots \oplus \bar{f}[n]$. In our algorithm, since we can not afford to combine the actual profit functions themselves, we will instead combine monotone point sets and reduce the combined monotone point sets using Operation 3 whenever necessary. Definition 4.3 provides us a tool to analyze our algorithm. Every time the algorithm computes a monotone point set P , the set is intended to approximate some unknown profit function f . Our algorithm will make sure that at any time, (f, P) is F -generatable for some set F that can be used to approximate f_I (e.g. \bar{F}). Our goal is to show:

- **Correctness:** an F -generatable (f, P) should be such that P can be used to approximate f .
- **Efficiency:** if (f_a, P_a) and (f_b, P_b) are both F -generatable, then there is an efficient way to obtain a P such that $(f_a \oplus f_b, P)$ is F -generatable. This allows us to efficiently approximate $f_a \oplus f_b$ if both functions are already approximated by P_a and P_b .

For two functions \tilde{f} and f and a constant c , we say that \tilde{f} c -subtractively approximates f if f c -additively approximates \tilde{f} . If \tilde{f} c -subtractively approximates f , then the function \hat{f} with identical domain as \tilde{f} defined as $\hat{f}(x) := \tilde{f}(x) - \delta$ c -additively approximates f . Thus subtractive approximation is as powerful as additive approximation. The following shows that individual rounding errors do not accumulate:

Lemma 4.9. *Given a set F of profit functions. For any set of m profit functions $\{f_1, f_2, \dots, f_m\}$ and any set of m monotone point sets $\{P_1, P_2, \dots, P_m\}$ such that $(f_1, P_1), (f_2, P_2), \dots, (f_m, P_m)$ are all F -generatable, $(g[P_1] \oplus g[P_2] \oplus \dots \oplus g[P_m])$ δ -subtractively approximates $f_1 \oplus f_2 \oplus \dots \oplus f_m$.*

For efficiency, we will first define a measure of complexity on monotone point sets. Given two monotone point sets $U = \{u_1, u_2, \dots, u_{|U|}\}$ and $V = \{v_1, v_2, \dots, v_{|V|}\}$, if $u_{|U|} = v_1$, define the concatenation $U \circ V := \{u_1, u_2, \dots, u_{|U|} = v_1, v_2, \dots, v_{|V|}\}$. A monotone point set $P = \{p_1, p_2, \dots, p_k\}$ is an *upper convex hull* if for all $1 < i < k$, the point p_i is strictly above the line segment connecting p_{i-1} and p_{i+1} . P is a *lower convex hull* if for all $1 < i < k$, the point p_i is strictly below the line segment connecting p_{i-1} and p_{i+1} . A *decomposition* of monotone point set P into b blocks is a way to write $P = B[1] \circ B[2] \circ \dots \circ B[b]$ such that each $B[i]$ is either an upper convex hull or a lower convex hull. For example, the monotone point set $P = \{A, B, C, D, E, F, G\}$ in Fig. 1 can be decomposed into 3 blocks $P = \{A, B, C\} \circ \{C, D, E, F\} \circ \{F, G\}$, where $\{A, B, C\}$, $\{C, D, E, F\}$ are lower convex hulls, and $\{F, G\}$ can be considered either an upper convex hull or a lower convex hull.

Given a monotone point set $P = \{p_1, p_2, \dots, p_k\}$ with no degenerate point, let t_i be 1 if p_i is an apex and 0 if p_i is a base. Let the *convex number* of P be the number of maximal consecutive subsequences of 1's and 0's of the sequence $T = \{t_i\}$. For example, for the monotone point set $P = \{A, B, C, D, E, F, G\}$ in Fig. 1, we have $T = \{1, 0, 1, 0, 0, 1, 1\}$, which consists of 5 maximal consecutive subsequences of 1's and 0's: $\{t_1\}$, $\{t_2\}$, $\{t_3\}$, $\{t_4, t_5\}$ and $\{t_6, t_7\}$ and thus the convex number of P is 5. For the relation between convex number and decomposition, we have the following observation:

Observation 4.4. *Let τ be the convex number of P . We have a decomposition into τ blocks: for $1 \leq i \leq k$, there is a block starting from p_i for if and only if $i = 1$ or $t_i \neq t_{i-1}$.*

For example, for the monotone point set $P = \{A, B, C, D, E, F, G\}$ in Fig. 1, this gives the valid decomposition $P = \{A, B\} \circ \{B, C\} \circ \{C, D\} \circ \{D, E, F\} \circ \{F, G\}$.

We first show that there is a way to reduce the convex number of a monotone point set while still being able to approximate the associated profit function:

Lemma 4.10. *Fix a set F of profit functions. Suppose (f, P) is F -generatable and P contains l points. We can compute another point set P' with a convex number of $O((\max_y P - \min_y P)/\delta)$ in $O(l)$ time such that (f, P') is F -generatable.*

We say a monotone point set \hat{P} σ -accurately approximates another monotone point set if the upper boundary of $R(\hat{P})$ and $R(P)$ match whenever the y-coordinate is a multiple of σ , namely if:

$$\bullet \left\lfloor \min_y \hat{P} / \sigma \right\rfloor = \left\lfloor \min_y P / \sigma \right\rfloor \text{ and } \left\lfloor \max_y \hat{P} / \sigma \right\rfloor = \left\lfloor \max_y P / \sigma \right\rfloor.$$

- For every $y \in [\min_y P, \max_y P]$ such that $\sigma \mid y$, the minimum x-coordinate such that $g[P](x) \geq y$ is the same as the minimum x-coordinate such that $g[\hat{P}](x) \geq y$.

We show that if two monotone point sets have low convex numbers, we can combine them efficiently:

Lemma 4.11. *Suppose we have two monotone point sets P_a and P_b . Suppose the y-coordinates of points in both P_a and P_b are multiples of σ in $[0, l\sigma)$, and suppose both P_a and P_b have a convex number of at most m . We can compute a set that σ -accurately approximates $P_a \oplus P_b$ in $\tilde{O}(lm)$ time.*

The problem we are solving here is different from the one in [BL23] where all blocks must be upper convex hulls⁶.

Lemma 4.10 and Lemma 4.11 give us a good way to achieve efficiency: suppose (f_a, P_a) and (f_b, P_b) are both F -generatable for some F . We first use Lemma 4.10 to obtain (f_a, P'_a) and (f_b, P'_b) that are both F -generatable where both P'_a and P'_b have low convex numbers, and then we will Lemma 4.11 to obtain $P'_a \oplus P'_b$ (σ -accurately with regards to some parameter σ). By definition, $(f_a \oplus f_b, P'_a \oplus P'_b)$ is F -generatable.

With these lemmas, we can prove Lemma 2.3.

Proof of Lemma 2.3. First, we can reduce ε so that $1/\varepsilon$ becomes an integer.

We can assume $n = 2^K$ since otherwise we can pad the set with items of very large weights. For $i = K, K-1, \dots, 0$ in order, we will compute a set of profit functions $F_i = \{f_i[1], f_i[2], \dots, f_i[2^i]\}$ such that each function has values being multiples of $\varepsilon 2^{K-i}$ in $[0, 2^{K-i+1}]$ and that for each i , $f_i[1] \oplus f_i[2] \oplus \dots \oplus f_i[2^i]$ ($2\varepsilon n(K-i+1)$)-subtractively approximates f_I for $x \in [0, \sum_{j \in I} w_j]$. By extending the domain of $f_0[1]$ to all $x \geq 0$ trivially we can get a $(2\varepsilon n \log_2 n)$ -subtractive approximation of f_I , which can be made εn by lowering the value of ε by a logarithmic factor.

For $i = K$, it suffices to let $f_i[j]$ be $\bar{f}[j]$ with values rounded up to multiples of ε , which introduces a subtractive error of $n\varepsilon$. The values of $\bar{f}[j]$'s are contained in $[0, 2]$ since $p_j \in [1, 2]$ for $j \in I$. Since $1/\varepsilon$ is an integer, we never round a value over 2, and thus the values of $f_i[j]$'s are contained in $[0, 2]$.

Suppose we correctly computed F_{i+1} . For each $J \in [1, 2^{i+1}]$. Let $\hat{P}_{i+1}[J] = P(f_{i+1}[J])$. Note that since values of $f_{i+1}[J]$ are multiples of $\varepsilon 2^{K-i}$ no more than $2^{K-(i+1)+1}$, the y-coordinates in $\hat{P}_i[J]$ are also multiples of $\varepsilon 2^{K-i}$ no more than $2^{K-(i+1)+1}$, and thus $\hat{P}_i[J]$ has no more than $O(2^{K-(i+1)+1}/(2^{K-i}\varepsilon)) = O(1/\varepsilon)$ points. We use Lemma 4.10 on each $\hat{P}_{i+1}[J]$ to get $P'_{i+1}[J]$ with a convex number of $O(2^{K-(i+1)+1}/\delta)$, which takes total time

$$O(2^{i+1}(1/\varepsilon)) = O(\varepsilon^{-2}).$$

We then use Lemma 4.11 to $(\varepsilon 2^{K-(i+1)})$ -accurately approximate $P_i[j] = P'_{i+1}[2j-1] \oplus P'_{i+1}[2j]$ for each $j \in [1, 2^i]$ in time

$$\tilde{O}(2^i \times (1/\varepsilon) \times (2^{K-(i+1)+1}/\delta)) = \tilde{O}(2^k \varepsilon^{-2}/n) = \tilde{O}(\varepsilon^{-2}).$$

⁶See the part regarding convex numbers. The definition in the paper is that blocks are lower convex hulls, but the paper deals with $(\min, +)$ -convolution instead of $(\max, +)$ -convolution.

We set $\hat{F}_i = \{g[P_i[1]], g[P_i[2]], \dots, g[P_i[2^i]]\}$. For each j , we let $\tilde{f}_i[j] = f_{i+1}[2j-1] \oplus f_{i+1}[2j]$. Then each $(\tilde{f}_i[j], P_i[j])$ is F_{i+1} -generatable. By Lemma 4.9 we can see that $g[P_i[1]] \oplus g[P_i[2]] \oplus \dots \oplus g[P_i[2^i]]$ εn -subtractively approximates $f_i[1] \oplus f_i[2] \oplus \dots \oplus f_i[2^i]$. Thus \hat{F}_i only introduces an extra additive error of εn compared to F_{i+1} . To get the set $F_i = \{f_i[1], f_i[2], \dots, f_i[2^i]\}$, we round values of each $g[P_i[j]]$ up to multiples of $\varepsilon 2^{K-i}$ to obtain $f_i[j]$, which is possible from a $(\varepsilon 2^{K-(i+1)})$ -accurate approximation of $P_i[j]$ since $\varepsilon 2^{K-i}$ is a multiple of $(\varepsilon 2^{K-(i+1)})$. Such rounding introduces an additive error of at most $\varepsilon 2^{K-i} \times 2^i = \varepsilon n$. The total additive error for F_i is at most:

$$2\varepsilon n(K - (i + 1) + 1) + \varepsilon n + \varepsilon n = 2\varepsilon n(K - i + 1).$$

It is easy to see that

$$\max_y P_i[j] = \max_y P_{i+1}[2j-1] + \max_y P_{i+1}[2j] \leq 2^{K-(i+1)+1} \times 2 = 2^{K-i+1},$$

and

$$\min_x P_i[j] = \min_x P_{i+1}[2j-1] + \min_x P_{i+1}[2j] \geq 0.$$

Thus the values of the function $g[P_i[j]]$ are contained in $[0, 2^{K-i+1}]$. Since $1/\varepsilon$ is an integer, we never round a value over 2^{K-i+1} . Thus the values of the function $f_i[j]$ are also contained in $[0, 2^{K-i+1}]$. Since there are $K = \log_2 n = \tilde{O}(1)$ layers, the total running time is $\tilde{O}(\varepsilon^{-2})$. \square

4.2.3 Proof of Lemma 4.9, Lemma 4.10 and Lemma 4.11

For correctness, we need to prove Lemma 4.9:

Lemma 4.9. *Given a set F of profit functions. For any set of m profit functions $\{f_1, f_2, \dots, f_m\}$ and any set of m monotone point sets $\{P_1, P_2, \dots, P_m\}$ such that $(f_1, P_1), (f_2, P_2), \dots, (f_m, P_m)$ are all F -generatable, $(g[P_1] \oplus g[P_2] \oplus \dots \oplus g[P_m])$ δ -subtractively approximates $f_1 \oplus f_2 \oplus \dots \oplus f_m$.*

To derive Lemma 4.9, we need to prove some ingredients.

Lemma 4.5. *For any two monotone point sets P and Q with no degenerate points, let $V = P \oplus Q$. Then for every v on the upper boundary of $R(V)$, $v = p + q$ such that either:*

- p is on the upper boundary of P and q is an apex of Q , or
- q is on the upper boundary of Q and p is an apex of P .

Proof. We know that for every v on the upper boundary of $R(V)$, $v = p + q$ such that either:

- p is on the upper boundary of P and $q \in Q$, or
- q is on the upper boundary of Q and $p \in P$.

Without loss of generality suppose that p is on the upper boundary of P and $q \in Q$.

If p is an apex of P , then since q is on the upper boundary of Q , we are already done. Otherwise, it suffices to show that q is an apex of Q .

Since p is not an apex of P , we have an infinitesimal segment $e \subset R(P)$ such that p is strictly contained in e . Thus $p + q$ is strictly contained in $e + q$, which is contained in $R(P) + q$. Suppose that e connects p_l and p_r with $p_l < p < p_r$.




Figure 2: Proof of Lemma 4.5

Figure 2 shows the immediately neighborhood of $p + q$ after zooming into the infinitesimal segment $e + q$, the shaded region is $R(P) + q$. Suppose q is not an apex of Q , then q is not an endpoint of Q . Suppose q_l and q_r are the point before and after q in Q . The points $p + q_l$ and $p + q_r$ can not both be below $e + q$ since otherwise it would make q an apex of Q . Without loss of generality suppose $p + q_l$ is above the segment $e + q$, as shown in the figure. Then a non-zero fraction of the segment connecting $p_r + q_l$ and $p_r + q$, which is entirely in $R(V)$, is above $p + q$. Thus as shown in the figure, we can find a point p' with $p'_x < p_x$ and $p'_y > p_y$ on the segment. Since $p' \in R(V)$, $v = p + q$ must not be on the upper boundary of $R(V)$ and thus must not be in V , which is a contradiction. \square

Lemma 4.6. For any two monotone point set P and Q with no degenerate points, if we let $V = P \oplus Q$, then every apex v of V must be the sum of an apex p of P and an apex q of Q .

Proof. Since v is an apex of V , it is easy to see that there must not be a non-degenerate segment (i.e. segment with a length greater than 0) $e \subset V$ such that $v \in e$.

Suppose $v = p + q$. Suppose p is not an apex of P . Then there exists an infinitesimal segment $e \subset R(P)$ such that p is strictly contained in e . We have a non-degenerate segment $e + q \subset V$ that contains $v = p + q$, which is a contradiction. Therefore p must be an apex of P . Symmetrically, q must be an apex of Q . \square

Lemma 4.7. For any set F of profit functions, and an F -generatable (f, P) , we have the following:

1. For every x in the domain of f , $g[P](x) \geq f(x)$. Equivalently, $R(P(f)) \subset R(P)$.

2. For any apex $p \in P$, $f(p_x) = p_y$.

3. For any point p on the upper boundary of $R(P)$, there exist points a and b ($0 \leq a_y < b_y \leq a_y + \delta$) such that $f(a_x) \geq a_y$, $f(b_y) \geq b_y$, and p is on the line segment connecting a and b .

Proof. It suffices to argue that each of the three operations in Definition 4.3 does not introduce new F -generatable pairs that violate each of the invariants.

For operation (1), since $P = P(f)$, Invariants (1), (2) are obviously satisfied. For Invariant (3), since we must have $f(p_x) \geq p_y$, we can set $a = b = p$.

For operation (2), fix (f_a, P_a) and (f_b, P_b) . Since $R(P(f_a)) \subset R(P_a)$ and $R(P(f_b)) \subset R(P_b)$, it is easy to see that $R(P(f_a \oplus f_b)) \subset R(P_a) + R(P_b) = R(P_a \oplus P_b)$, and therefore invariant (1) must hold. For Invariant (2), suppose a point $p = (p_x, p_y)$ is an apex of $P_a \oplus P_b$. From Lemma 4.6, p can be expressed as the sum of two apexes u on P_a and v on P_b . Since $f_a(u_x) = u_y$ and $f_b(v_x) = v_y$, we must have $p_y = u_y + v_y = f_a(u_x) + f_b(v_x) \leq (f_a \oplus f_b)(p_x)$. Since $p_y = g[P](p_x) \geq (f_a \oplus f_b)(p_x)$ (from Invariant (1)), we must have $(f_a \oplus f_b)(p_x) = p_y$. For Invariant (3), let p be a point on the boundary of $R(P)$, then from Lemma 4.5, $p = u + v$ such that either u is on the upper boundary of $R(P_a)$ and v is an apex of P_b , or v is on the upper boundary of $R(P_b)$ and u is an apex of P_a . Without loss of generality suppose that u is on the upper boundary of $R(P_a)$ and v is an apex of P_b . Then $f_b(v_x) = v_y$ (Invariant (2)). Note that Invariant (3) is satisfied for (f_a, P_a) . Let a_u and b_u be the points in Invariant (3) for u on the upper boundary of $R(P_a)$. Then it is not hard to verify that setting $a := a_u + v$ and $b := b_u + v$ satisfies Invariant (3) for p on the upper boundary of $R(P)$.

For operation (3), since we only remove non-apexes in P to obtain P' , $R(P) \subset R(P') = R(\hat{P})$. Therefore $P(f) \subset R(P) \subset R(\hat{P})$, and thus Invariant (1) must hold. It is easy to see that no new apexes can be created, and therefore Invariant (2) does not get broken. Consider Invariant (3) for \hat{P} . If p is on the upper boundary of \hat{P} , then Invariant (3) holds since it holds for (f, P) . Otherwise, it is easy to see that p must be on a segment connecting two apexes a and b with y -coordinates at most δ apart and it is easy to verify that a and b satisfy Invariant (3) for p . \square

Recall that \tilde{f} c -subtractively approximates f if f c -additively approximates \tilde{f} .

Corollary 4.8. For any set F of profit functions, and an F -generatable (f, P) , $g[P]$ δ -subtractively approximates $f(x)$.

Proof. We show that for every x in the domain of f , we have $g[P](x) - \delta \leq f(x) \leq g[P](x)$.

$f(x) \leq g[P](x)$ follows from Invariant (1) of Lemma 4.7. To see why $g[P](x) - \delta \leq f(x)$. Apply Invariant (3) on $(x, g[P](x))$ and obtain a and b . Since $a_y \leq g[P](x) \leq b_y$ and f is non-decreasing, we have $f(x) \geq f(a_x) \geq a_y \geq b_y - \delta \geq g[P](x) - \delta$. \square

Now we are ready to prove Lemma 4.9:

Lemma 4.9. Given a set F of profit functions. For any set of m profit functions $\{f_1, f_2, \dots, f_m\}$ and any set of m monotone point sets $\{P_1, P_2, \dots, P_m\}$ such that $(f_1, P_1), (f_2, P_2), \dots, (f_m, P_m)$ are all F -generatable, $(g[P_1] \oplus g[P_2] \oplus \dots \oplus g[P_m])$ δ -subtractively approximates $f_1 \oplus f_2 \oplus \dots \oplus f_m$.

Proof. Since with $m - 1$ repetitions of Operation 2 we can F -generate $(f_1 \oplus f_2 \oplus \dots \oplus f_m, P_1 \oplus P_2 \oplus \dots \oplus P_m)$, $g[P_1 \oplus P_2 \oplus \dots \oplus P_m]$ δ -subtractively approximates $f_1 \oplus f_2 \oplus \dots \oplus f_m$. It is easy to see that $g[P_1] \oplus g[P_2] \oplus \dots \oplus g[P_m] = g[P_1 \oplus P_2 \oplus \dots \oplus P_m]$. Therefore $(g[P_1] \oplus g[P_2] \oplus \dots \oplus g[P_m])$ δ -subtractively approximates $f_1 \oplus f_2 \oplus \dots \oplus f_m$. \square

Now we move on to efficiency and prove Lemma 4.10:

Lemma 4.10. *Fix a set F of profit functions. Suppose (f, P) is F -generatable and P contains l points. We can compute another point set P' with a convex number of $O((\max_y P - \min_y P)/\delta)$ in $O(l)$ time such that (f, P') is F -generatable.*

Proof. We can obtain P' in the following way. Initially set $P' = P$. Scan the apexes in order and store them in a stack. Every time we encounter a new apex x . If the stack is not empty, the top of the stack is y , and $|x_y - y_y| \leq \delta$, we can simulate an Operation 3 on P' with x and y , which involves:

- Removing the vertices between x and y in P' , and
- If y becomes degenerate, removing y from P' .

After this, if y is no longer an apex of P' or gets removed from P' , we remove y from the stack and repeat the procedure above. Otherwise, we push x to the end of the stack.

After processing each apex x , let $P'_{1\dots k}$ be the prefix of P' containing the remaining points in P' up to x . It is easy to verify that the following two invariants are true:

- No points of $P'_{1\dots k}$ are degenerate.
- The stack contains exactly all apexes of $P'_{1\dots k}$.
- For every maximal consecutive subsequence of bases in $P'_{1\dots k}$ enclosed by two apexes $a < b$, $b_y - a_y > \delta$.

The procedure can easily be done in $O(l)$ time, and the resulting P' is clearly F -generatable. Since every maximal consecutive subsequence of bases in P' is enclosed by two apexes δ with a difference larger than δ in y -coordinates, it is easy to see that P' has a convex number of $O((\max_y P - \min_y P)/\delta)$. \square

Finally, we prove Lemma 4.11:

Lemma 4.11. *Suppose we have two monotone point sets P_a and P_b . Suppose the y -coordinates of points in both P_a and P_b are multiples of σ in $[0, l\sigma)$, and suppose both P_a and P_b have a convex number of at most m . We can compute a set that σ -accurately approximates $P_a \oplus P_b$ in $\tilde{O}(lm)$ time.*

Proof. We can suppose $\sigma = 1$ since otherwise we can re-scale all the y -coordinates. We obtain decompositions $P_a = B_a[1] \circ B_a[2] \circ \dots \circ B_a[m_a]$ with $m_a \leq m$ and $P_b = B_b[1] \circ B_b[2] \circ \dots \circ B_b[m_b]$ with $m_b \leq m$ from Observation 4.4.

Suppose that for each two blocks $B_a[i] \in P_a$ and $B_b[j] \in P_b$, we can compute $A[i, j] = B_a[i] \oplus B_b[j]$. Then by taking the union of $R(A[i, j])$'s, we get $R(P_a \oplus P_b)$ which can give us $P_a \oplus P_b$. The

total number of y-coordinates involved in $R(A[i, j])$ does not exceed $(\max_y B_a[i] - \min_y B_a[i]) + (\max_y B_b[j] - \min_y B_b[j]) + 2$. Since we only need a 1-accurate approximation, to take the union of $R(A[i, j])$'s, we only need to find the minimum x-coordinate for every y-coordinate. Therefore every time we take the union with a new $R(A[i, j])$, we only need to go through all the involved y-coordinates and update the minimum x-coordinate. For all $R(A[i, j])$'s in total this takes time

$$O\left(\sum_{i,j} (\max_y B_a[i] - \min_y B_a[i]) + (\max_y B_b[j] - \min_y B_b[j]) + 2\right) = O(lm).$$

We claim that the following claim is true: For two blocks $B_a[i]$ and $B_b[j]$ with the total span of y-coordinates being \mathcal{Y} , $B_a[i] \oplus B_b[j]$ can be computed in $\tilde{O}(\mathcal{Y})$ time.

If the claim is true, then the total running time for computing every $A[i, j]$ is

$$\tilde{O}\left(\sum_{i,j} (\max_y B_a[i] - \min_y B_a[i]) + (\max_y B_b[j] - \min_y B_b[j]) + 2\right) = \tilde{O}(lm),$$

which finishes the proof of the lemma. Now we prove that for two blocks $B_a[i]$ and $B_b[j]$ with the total span of y-coordinates being \mathcal{Y} , we can indeed compute $B_a[i] \oplus B_b[j]$ in $\tilde{O}(\mathcal{Y})$ time.

For the case where both $B_a[i]$ and $B_b[j]$ are upper convex hulls, it is well known that the Minkowski sum of two convex polygons with v vertices in total can be computed in $\tilde{O}(v)$ time [dBvKOS00]. Since the number of vertices is bounded by the number of different y-coordinates, such running time is sufficient.

It suffices to deal with computations where one of the blocks is a lower convex hull. Without loss of generality, suppose $B_a[i]$ is a lower convex hull. Let $X = \max_y B_a[i] - \min_y B_a[i] + 1$, and $Y = \max_y B_b[j] - \min_y B_b[j] + 1$. We will now introduce a procedure that computes $B_a[i] \oplus B_b[j]$ in $\tilde{O}(X + Y)$ time.

Without loss of generality, suppose the y-coordinates of $B_a[i]$ span $[0, X - 1]$ and those of $B_b[j]$ span $[0, Y - 1]$. For $y \in [0, X - 1]$, define $\phi(y)$ as the minimum x with $g[B_a[i]](x) \geq y$. For $y \in [0, Y - 1]$, define $\gamma(y)$ as the minimum x with $g[B_b[j]](x) \geq y$. It suffices to compute the convolution

$$\eta(y) = \min_{y_f \in [0, X-1], y_g \in [0, Y-1], y_f + y_g = y} \phi(y_f) + \gamma(y_g),$$

where we know that ϕ is convex (i.e. the difference $\phi(y + 1) - \phi(y)$ decreases as y increases), in time linear in X and Y .

For the computation of $\eta(y)$ for $y \in [0, X - 1]$, we use a ‘‘reversed’’ version of a well-known technique in competitive programming⁷. We treat values of $\gamma(y)$ where $y > Y - 1$ to be ∞ . We compute $\eta(y)$ in increasing order of y . We have different ‘‘candidates’’ c which we can use to update $\eta(y)$ with $\phi(y - c) + \gamma(c)$. For two candidates c_a and c_b such that $c_a < c_b$, if $\phi(y - c_a) + \gamma(c_b) < \phi(y - c_b) + \gamma(c_b)$, then since $\phi(y - c_a)$ increases more slowly than $\phi(y - c_b)$ as y increases,

⁷See e.g. <https://codeforces.com/blog/entry/8219?#comment-139241>. We believe that the reversed version has also been featured in some competitive programming task in the past, but unfortunately, we could not find a reference.

c_b will never be a better candidate than c_a ⁸. For two candidates $c_a < c_b$, we can compute the minimum \hat{y} at which $\phi(\hat{y} - c_a) + \gamma(c_a) < \phi(\hat{y} - c_b) + \gamma(c_b)$ (or lack thereof) by binary search, and such \hat{y} is the “time” when c_a becomes better than c_b . We maintain a stack that stores the candidates $c_1 < c_2 < c_3 < \dots < c_k$ from bottom to top. We also maintain the timestamps $t_1 < t_2 < \dots < t_{k-1}$ where t_i is the time at which c_i becomes better than c_{i+1} , namely the smallest t_i such that $\phi(t_i - c_i) + \gamma(c_i) < \phi(t_i - c_{i+1}) + \gamma(c_{i+1})$. After computing $\eta(y)$ for $y = c$, c becomes available as a new candidate for future y 's. we compute the time \hat{t}_k at which the top of the stack c_k becomes better than c . If $\hat{t}_k \geq t_{k-1}$, then c_k can never become the best candidate and we can remove it from the stack. We repeat until at some k' we have $t_{k'-1} < \hat{t}_{k'}$ and then add c as $c_{k'+1}$ into the stack and set $t_{k'} = \hat{t}_{k'}$. As each candidate gets removed at most once, the total running time is $\tilde{O}(X)$.

This algorithm will not work for $y \geq X$ as a candidate c becomes unavailable when $y - c > X - 1$. To deal with this issue, we split integers in $[0, X + Y - 1)$ into consecutive blocks of size $X - 1$, where block k contains integers in $\{[k(X - 1), \min(X + Y - 1, (k + 1)(X - 1))]\}$. For each y , a candidate c within the same block will always be available. Thus if we call the procedure above on each block we can cover all pairs of (y, c) in the same block. It remains to deal with the case where y and c are in adjacent blocks. To do this, for every two adjacent blocks $[(k-1)(X-1), k(X-1)), [k(X-1), (k+1)(X-1))$, in decreasing order of $i = X - 2, X - 3, \dots, 0$, we compute the contribution of candidates in $[(k-1)(X-1) + i, k(X-1))$ to $y = k(X-1) + i$ (if $y < X + Y - 1$). To do this, we first add $c = (k-1)(X-1) + i$ to the set of available candidates and then find the best candidate for y from the set. It is not hard to maintain and find the best candidate for all y 's in near-linear time using the same technique described before. The total running time is $\tilde{O}(X + Y)$. \square

5 Open Problems

Although this paper closes the gap between the best algorithm and the conditional lower bound for the Knapsack problem, there are still some open problems in this field.

- Can we make our algorithm simple? Intuitively, a “simple” algorithm should not use any results from additive combinatorics, or a technique analogous to Chan’s number theoretical lemma. Despite the fact that our algorithm for Lemma 2.3 is simple, the reduction to the special case in the proof of Lemma 4.1 requires Lemma 3.9, which uses results from additive combinatorics, as well as Corollary 3.5, which uses the number theoretical lemma.
- Can we shave an $\Omega(2\sqrt{\log^{-1}(1/\varepsilon)})$ factor from our algorithm? The previous algorithm from [DJM23] has a running time of $\tilde{O}\left(n + \left(\frac{1}{\varepsilon}\right)^{11/5} / 2^{\Omega(\sqrt{\log(1/\varepsilon)})}\right)$, which has such a factor shaved. Moreover, can we establish an equivalence between $(1 - \varepsilon)$ -approximation of Knapsack and

⁸In the more common, “non-reversed” version, the observation is true for $c_a > c_b$. Thus a double-ended queue is used, but here we will use a stack instead. The “non-reversed” version can also be directly solved using the SMAWK algorithm.

(min, +)-convolution? Alternatively, can we prove a stronger conditional lower bound that rules out such an improvement?

References

- [ABHS19] Amir Abboud, Karl Bringmann, Danny Hermelin, and Dvir Shabtay. SETH-based lower bounds for subset sum and bicriteria path. In *Proceedings of the 30th ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 41–57, 2019. doi:[10.1137/1.9781611975482.3](https://doi.org/10.1137/1.9781611975482.3). 3
- [AKM⁺87] Alok Aggarwal, Maria M. Klawe, Shlomo Moran, Peter Shor, and Robert Wilber. Geometric applications of a matrix-searching algorithm. *Algorithmica*, 2(1):195–208, November 1987. doi:[10.1007/BF01840359](https://doi.org/10.1007/BF01840359). 2
- [BCD⁺14] David Bremner, Timothy M. Chan, Erik D. Demaine, Jeff Erickson, Ferran Hurtado, John Iacono, Stefan Langerman, Mihai Pătraşcu, and Perouz Taslakian. Necklaces, convolutions, and $x+y$. *Algorithmica*, 69(2):294–314, June 2014. doi:[10.1007/s00453-012-9734-3](https://doi.org/10.1007/s00453-012-9734-3). 2, 6
- [BL23] Cornelius Brand and Alexandra Lassota. Fast Convolutions for Near-Convex Sequences. In Satoru Iwata and Naonori Kakimura, editors, *34th International Symposium on Algorithms and Computation (ISAAC 2023)*, volume 283 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 16:1–16:16, Dagstuhl, Germany, 2023. Schloss Dagstuhl – Leibniz-Zentrum für Informatik. URL: <https://drops-dev.dagstuhl.de/entities/document/10.4230/LIPIcs.ISAAC.2023.16>, doi:[10.4230/LIPIcs.ISAAC.2023.16](https://doi.org/10.4230/LIPIcs.ISAAC.2023.16). 17
- [BN21] Karl Bringmann and Vasileios Nakos. A fine-grained perspective on approximating subset sum and partition. In *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms, SODA 2021, Virtual Conference, January 10 - 13, 2021*, pages 1797–1815. SIAM, 2021. doi:[10.1137/1.9781611976465.108](https://doi.org/10.1137/1.9781611976465.108). 2, 3, 5, 10, 11, 15
- [BW21] Karl Bringmann and Philip Wellnitz. On near-linear-time algorithms for dense subset sum. In Dániel Marx, editor, *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms, SODA 2021, Virtual Conference, January 10 - 13, 2021*, pages 1777–1796. SIAM, 2021. doi:[10.1137/1.9781611976465.107](https://doi.org/10.1137/1.9781611976465.107). 26
- [Cha18] Timothy M. Chan. Approximation Schemes for 0-1 Knapsack. In *Proceedings of the 1st Symposium on Simplicity in Algorithms (SOSA)*, pages 5:1–5:12, 2018. doi:[10.4230/OASIcs.SOSA.2018.5](https://doi.org/10.4230/OASIcs.SOSA.2018.5). 2, 3, 5, 6, 10
- [CMWW19] Marek Cygan, Marcin Mucha, Karol Węgrzycki, and Michał Włodarczyk. On problems equivalent to (min,+)-convolution. *ACM Trans. Algorithms*, 15(1):14:1–14:25, January 2019. doi:[10.1145/3293465](https://doi.org/10.1145/3293465). 2

- [CW16] Timothy M. Chan and Ryan Williams. Deterministic apsp, orthogonal vectors, and more: Quickly derandomizing razborov-smolensky. In *Proceedings of the 27th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1246–1255, 2016. doi:[10.1137/1.9781611974331.ch87](https://doi.org/10.1137/1.9781611974331.ch87). 2, 6
- [dBvKOS00] Mark de Berg, Marc van Kreveld, Mark Overmars, and Otfried Schwarzkopf. *Computational Geometry: Algorithms and Applications*. Springer-Verlag, second edition, 2000. URL: <http://www.cs.uu.nl/geobook/>. 14, 22
- [DJM23] Mingyang Deng, Ce Jin, and Xiao Mao. Approximating knapsack and partition via dense subset sums. In *Proceedings of the 2023 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 2961–2979, 2023. URL: <https://epubs.siam.org/doi/abs/10.1137/1.9781611977554.ch113>, arXiv:<https://epubs.siam.org/doi/pdf/10.1137/1.9781611977554.ch113>, doi:[10.1137/1.9781611977554.ch113](https://doi.org/10.1137/1.9781611977554.ch113). 2, 3, 5, 6, 7, 8, 10, 23, 26
- [GL79] George Gens and Eugene Levner. Computational complexity of approximation algorithms for combinatorial problems. In Jirí Běcvar, editor, *Mathematical Foundations of Computer Science 1979, Proceedings, 8th Symposium, Olomouc, Czechoslovakia, September 3-7, 1979*, volume 74 of *Lecture Notes in Computer Science*, pages 292–300. Springer, 1979. doi:[10.1007/3-540-09526-8_26](https://doi.org/10.1007/3-540-09526-8_26). 10
- [IK75] Oscar H. Ibarra and Chul E. Kim. Fast approximation algorithms for the knapsack and sum of subset problems. *Journal of the ACM (JACM)*, 22(4):463–468, October 1975. doi:[10.1145/321906.321909](https://doi.org/10.1145/321906.321909). 2
- [Jin19] Ce Jin. An improved FPTAS for 0-1 knapsack. In Christel Baier, Ioannis Chatzigiannakis, Paola Flocchini, and Stefano Leonardi, editors, *46th International Colloquium on Automata, Languages, and Programming, ICALP 2019, July 9-12, 2019, Patras, Greece*, volume 132 of *LIPICs*, pages 76:1–76:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2019. doi:[10.4230/LIPICs.ICALP.2019.76](https://doi.org/10.4230/LIPICs.ICALP.2019.76). 2, 5, 8, 10
- [JK18] Klaus Jansen and Stefan E.J. Kraft. A faster fptas for the unbounded knapsack problem. *European Journal of Combinatorics*, 68:148 – 174, 2018. doi:[10.1016/j.ejc.2017.07.016](https://doi.org/10.1016/j.ejc.2017.07.016). 2
- [Kar72] Richard M Karp. Reducibility among combinatorial problems. In *Complexity of computer computations*, pages 85–103. Springer, 1972. 1
- [KMPS03] Hans Kellerer, Renata Mansini, Ulrich Pferschy, and Maria Grazia Speranza. An efficient fully polynomial approximation scheme for the subset-sum problem. *J. Comput. Syst. Sci.*, 66(2):349–370, 2003. doi:[10.1016/S0022-0000\(03\)00006-0](https://doi.org/10.1016/S0022-0000(03)00006-0). 3

- [KP04] Hans Kellerer and Ulrich Pferschy. Improved dynamic programming in connection with an fptas for the knapsack problem. *Journal of Combinatorial Optimization*, 8(1):5–11, March 2004. doi:10.1023/B:JOCO.0000021934.29833.6b. 2
- [KPS97] Hans Kellerer, Ulrich Pferschy, and M.Grazia Speranza. An efficient approximation scheme for the subset-sum problem. volume 1350, pages 394–403, 12 1997. doi:10.1007/3-540-63890-3_42. 10
- [KPS17] Marvin Künnemann, Ramamohan Paturi, and Stefan Schneider. On the fine-grained complexity of one-dimensional dynamic programming. In *Proceedings of the 44th International Colloquium on Automata, Languages, and Programming (ICALP)*, pages 21:1–21:15, 2017. doi:10.4230/LIPIcs.ICALP.2017.21. 2
- [Law79] Eugene L. Lawler. Fast approximation algorithms for knapsack problems. *Mathematics of Operations Research*, 4(4):339–356, 1979. doi:10.1287/moor.4.4.339. 2, 3
- [MWW19] Marcin Mucha, Karol Węgrzycki, and Michał Włodarczyk. Subquadratic approximation scheme for partition. In *Proceedings of the 30th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 70–88, 2019. doi:10.1137/1.9781611975482.5. 2
- [Rhe15] Donguk Rhee. Faster fully polynomial approximation schemes for knapsack problems. Master’s thesis, Massachusetts Institute of Technology, 2015. URL: <http://hdl.handle.net/1721.1/98564>. 2
- [Wil14] Ryan Williams. Faster all-pairs shortest paths via circuit complexity. In *Proceedings of the 46th Annual ACM Symposium on Theory of Computing (STOC)*, pages 664–673, 2014. doi:10.1145/2591796.2591811. 2, 6

A Proof of Lemma 3.9

The following structural lemma follows from Theorem 4.1 and Theorem 4.2 of Bringmann and Wellnitz [BW21].

Lemma A.1. *Let n distinct positive integers $X = \{x_1, \dots, x_n\} \subseteq [\ell, 2\ell]$ be given, where $\ell = o(n^2 / \log n)$.*

Then, for a universal constant $c \geq 1$, for every $c\ell^2/n \leq t \leq \Sigma(X)/2$, there exists $t' \in \mathcal{S}(X)$ such that $0 \leq t' - t \leq 8\ell/n$.

Now we are ready to prove Lemma 3.9

Lemma 3.9. *[Greedy Exchange Lemma, Lemma 3.9 of [DJM23]] Let $S \subseteq [n]$ be any item set with total profit $\sum_{s \in S} p_s \leq 2m$. Let $\Delta = \omega(\varepsilon^{-1/2} \log^{1/2}(1/\varepsilon))$ be some parameter. Let $B := c\varepsilon^{-1}/\Delta$ for some universal constant $c \geq 1$.*

Then, there exists an item set $\tilde{S} \subseteq [n]$, such that the total profit \tilde{p} contributed by items $[n] \setminus [i^\Delta]$ in \tilde{S} satisfies

$$\tilde{p} := \sum_{s \in \tilde{S} \cap ([n] \setminus [i^\Delta])} p_s \leq B, \quad (1)$$

and

$$\sum_{s \in \tilde{S}} p_s \geq (1 - \varepsilon) \sum_{s \in S} p_s, \quad (2)$$

and

$$\sum_{s \in \tilde{S}} w_s \leq \sum_{s \in S} w_s. \quad (3)$$

Proof. If $D(i) < \Delta$, then by the definition of i we have $i = n$, and we can simply let $\tilde{S} = S$, since $\tilde{p} = 0$ always holds. So in the following we assume $D(i) = \Delta$.

We define $\tilde{S} \subseteq [n]$ as the maximizer of

$$\sum_{s \in \tilde{S} \cap [i]} p_s + \sum_{s \in \tilde{S} \cap ([n] \setminus [i])} (1 - \varepsilon) p_s$$

among all \tilde{S} satisfying $\sum_{s \in \tilde{S}} w_s \leq \sum_{s \in S} w_s$ and $\sum_{s \in \tilde{S}} p_s \leq \sum_{s \in S} p_s$. We claim that \tilde{S} satisfies the properties (1), (2), (3). Observe that (2), (3) immediately follow from the definition of \tilde{S} . The main part is to verify (1).

Suppose for contradiction that (1) does not hold. Then, we can find a subset $K \subseteq \tilde{S} \cap ([n] \setminus [i])$ with total profit $p^* = \sum_{k \in K} p_k \in (B, B + 2]$, which can be obtained by removing items from $\tilde{S} \cap ([n] \setminus [i])$ (recall that each item has profit in $[1, 2)$).

Define item set $I' := [i] \setminus \tilde{S}$. Since $|\tilde{S}| < \sum_{s \in \tilde{S}} p_s / \min_{s \in \tilde{S}} p_s \leq \sum_{s \in \tilde{S}} p_s \leq \sum_{s \in S} p_s \leq 2m$, by the definition of $D(i)$, we know that $\{p_i : i \in I'\}$ contains at least $D(i) = \Delta$ distinct elements.

Suppose c is the universal constant in Lemma A.1. We apply Lemma A.1 on the set of integers $X = \{p_i/\varepsilon : i \in I'\} \subseteq [1/\varepsilon, 2/\varepsilon)$ which contains at least Δ distinct integers, where the premise $1/\varepsilon = o(\Delta^2/\log \Delta)$ in Lemma A.1 is satisfied due to the fact that $\Delta = \omega(\varepsilon^{-1/2} \log^{1/2}(1/\varepsilon))$. Lemma A.1 states that for every $t \in [c\varepsilon^{-2}/\Delta, 0.5\Delta/\varepsilon]$, there exists $t' \in \mathcal{S}(X)$ such that $0 \leq t' - t \leq 8\varepsilon^{-1}/\Delta$. Here we set

$$t := \frac{(1 - \varepsilon)p^*}{\varepsilon} + \frac{\varepsilon^{-1}}{\Delta},$$

which satisfies $t > p^*(1 - \varepsilon)/\varepsilon > (1 - \varepsilon)B/\varepsilon = (1 - \varepsilon)(9c\varepsilon^{-1}/\Delta)/\varepsilon > c\varepsilon^{-2}/\Delta$, and $t < (B + 2)/\varepsilon + \varepsilon^{-1}/\Delta = 9c\varepsilon^{-2}/\Delta + 2/\varepsilon + \varepsilon^{-1}/\Delta \leq o(\varepsilon^{-3/2} \log^{-1/2}(1/\varepsilon)) \leq 0.5\Delta/\varepsilon$. Then the conclusion of Lemma A.1 says that there is a subset $R \subseteq I'$ of items with total profit $\tilde{p} := \varepsilon \cdot t'$, satisfying

$$1/\Delta \leq \tilde{p} - p^*(1 - \varepsilon) \leq 9/\Delta. \quad (4)$$

Note that (4) implies

$$\begin{aligned}
p^* - \tilde{p} &\geq \varepsilon \cdot p^* - 9/\Delta \\
&> \varepsilon \cdot B - 9/\Delta \\
&= \varepsilon \cdot 9c\varepsilon^{-1}/\Delta - 9/\Delta \\
&\geq 0.
\end{aligned}$$

Recall that $R \subseteq I' = [i] \setminus \tilde{S}$ and $K \subseteq \tilde{S} \cap ([n] \setminus [i])$, which must both be non-empty. Since the efficiency of items are sorted in non-increasing order, we have $\min_{r \in R} p_r/w_r \geq \max_{k \in K} p_k/w_k$. Now we define the set of items

$$\tilde{S}' := (\tilde{S} \setminus K) \cup R.$$

Then, we have

$$\begin{aligned}
\sum_{s \in \tilde{S}} p_s - \sum_{s \in \tilde{S}'} p_s &= \sum_{k \in K} p_k - \sum_{r \in R} p_r \\
&= p^* - \tilde{p} \\
&\geq 0,
\end{aligned}$$

and

$$\begin{aligned}
\sum_{s \in \tilde{S}} w_s - \sum_{s \in \tilde{S}'} w_s &= \sum_{k \in K} w_k - \sum_{r \in R} w_r \\
&\geq \frac{\sum_{k \in K} p_k}{\max_{k \in K} (p_k/w_k)} - \frac{\sum_{r \in R} p_r}{\min_{r \in R} (p_r/w_r)} \\
&\geq \frac{1}{\min_{r \in R} (p_r/w_r)} \cdot \left(\sum_{k \in K} p_k - \sum_{r \in R} p_r \right) \\
&= \frac{1}{\min_{r \in R} (p_r/w_r)} \cdot (p^* - \tilde{p}) \\
&\geq 0.
\end{aligned}$$

Hence, $\sum_{s \in \tilde{S}'} p_s \leq \sum_{s \in \tilde{S}} p_s$ and $\sum_{s \in \tilde{S}'} w_s \leq \sum_{s \in \tilde{S}} w_s$. On the other hand, by (4), we know that

$$\begin{aligned}
&\left(\sum_{s \in \tilde{S}' \cap [i]} p_s + \sum_{s \in \tilde{S}' \cap ([n] \setminus [i])} (1 - \varepsilon) p_s \right) - \left(\sum_{s \in \tilde{S} \cap [i]} p_s + \sum_{s \in \tilde{S} \cap ([n] \setminus [i])} (1 - \varepsilon) p_s \right) \\
&= \sum_{r \in R} p_r - \sum_{k \in K} (1 - \varepsilon) p_k \\
&= \tilde{p} - (1 - \varepsilon) p^* \\
&\geq 1/\Delta > 0,
\end{aligned}$$

contradicting the definition of \tilde{S} being a maximizer.

Hence, we have established that \tilde{S} satisfies (1). □

B Proof of Lemma 3.1

Recall that we defined the following simpler problem.

Problem 2. Assume $\varepsilon \in (0, 1/2)$ and $1/\varepsilon \in \mathbb{N}^+$. Given a list I of items $(p_1, w_1), \dots, (p_n, w_n)$ with weights $w_i \in \mathbb{N}$ and profits p_i being multiples of ε in the interval $[1, 2)$, compute a profit function that $(1 - \varepsilon)$ -approximates f_I up to $2/\varepsilon$.

Lemma 3.1. If for some $c \geq 2$, Problem 2 can be solved in $\tilde{O}(n + 1/\varepsilon^c)$ time, then $(1 - \varepsilon)$ -approximating Knapsack can also be solved in $\tilde{O}(n + 1/\varepsilon^c)$ time.

Proof. First, we can reduce ε so that $1/\varepsilon$ becomes an integer.

We will restrict the profit values into small intervals, as follows: divide the items into $O(\log \frac{\max_j p_j}{\min_j p_j}) = O(\log \varepsilon^{-1})$ groups (see Section 2.2), each containing items with $p_i \in [2^j, 2^{j+1}]$ for some j (which can be rescaled to $[1, 2]$). Finally, use the merging lemma Lemma 3.2 to merge the profit functions of all groups, in $\tilde{O}(n + \varepsilon^{-2})$ overall time.

Now, having restricted the profit values into $[1, 2)$, we can round every profit value to a multiple of ε , which incurs only $(1 - O(\varepsilon))$ approximation factor in total.

Finally, the following greedy lemma takes care of the case with total profit above $\Omega(\varepsilon^{-1})$.

Lemma B.1. Suppose $p_i \in [1, 2]$ for all $i \in I$. For $B = \Omega(\varepsilon^{-1})$, the profit function f_I can be approximated with additive error $O(\varepsilon B)$ in $O(n \log n)$ time.

Proof. Simply sort the items in nonincreasing order of efficiency p_i/w_i , and define the profit function \tilde{f} resulting from greedy, with function values $0, p_1, p_1 + p_2, \dots, p_1 + \dots + p_n$ and x -breakpoints $0, w_1, w_1 + w_2, \dots, w_1 + \dots + w_n$. It clearly approximates f_I with an additive error of $\max_i p_i \leq 2 \leq O(\varepsilon B)$ for $B = \Omega(\varepsilon^{-1})$. \square

This greedy approach achieves $(1 - O(\varepsilon))$ -approximation for large profit values. Hence, it is sufficient to approximate f_I up to $2/\varepsilon$. \square