BOX DIMENSION OF GENERALIZED AFFINE FRACTAL INTERPOLATION FUNCTIONS (II)

LAI JIANG AND HUO-JUN RUAN

ABSTRACT. Let f be a generalized affine fractal interpolation function with vertical scaling functions. In this paper, we prove the monotonicity of spectral radii of vertical scaling matrices without additional assumptions. We also obtain the irreducibility of these matrices under certain conditions. By these results, we estimate $\dim_B \Gamma f$, the box dimension of the graph of f, by the limits of spectral radii of vertical scaling matrices. We also estimate $\dim_B \Gamma f$ directly by the sum function of vertical scaling functions. As an application, we study the box dimension of the graph of a generalized Weierstrass-type function.

1. INTRODUCTION

Fractal interpolation functions (FIFs) were introduced by Barnsley [4] in 1986. Basically, an FIF f is a function which interpolates given data and its graph is the invariant set of an iterated function system (IFS).

The generalized affine FIFs are an important class of FIFs. There are many works on these FIFs, including theoretical analysis [2, 3, 6, 10, 11, 26] and applications [21, 25]. In particular, there are some works on the box dimension of the graphs of generalized affine FIFs [7, 8, 9, 13, 17, 23]. In [18], the authors introduced vertical scaling matrices. We also obtained the monotonicity of spectral radii and irreducibility of these matrices under certain conditions. Then we estimated the box dimension of generalized affine FIFs by the limits of the spectral radii of these matrices.

In the present paper, we continue the study of vertical scaling matrices and the box dimension of generalized affine FIFs. Mainly, we prove the monotonicity of spectral radii of vertical scaling matrices without additional assumptions, and obtain the irreducibility of vertical scaling matrices under weaker conditions than that in [18]. By using these results, we estimate the box dimension by the limits of vertical scaling matrices under weaker conditions. We also estimate the box dimension directly by the sum function of vertical scaling functions. We remark that the class of generalized affine FIFs in the present paper is more general than the setting in [18], so that our results are applicable to some classical fractal functions, including the classical Weierstrass functions.

The paper is organized as follows. In section 2, we recall some basic definitions and present main results. In section 3, we study the monotonicity of spectral radii

²⁰²⁰ Mathematics Subject Classification. Primary 28A80; Secondary 41A30.

Key words and phrases. Fractal interpolation functions, box dimension, iterated function systems, vertical scaling functions, spectral radius, vertical scaling matrices.

The research was supported in part by NSFC grant 12371089, ZJNSF grant LY22A010023, and the Fundamental Research Funds for the Central Universities of China grant 226-2024-00136.

Corresponding author: Huo-Jun Ruan.

and irreducibility of vertical scaling matrices. By using these results, in section 4, we estimate the box dimension of generalized affine FIFs by the limits of radii of vertical scaling matrices under certain conditions. In section 5, we estimate the box dimension of generalized affine FIFs by the sum function of the vertical scaling functions. In section 6, we apply our results to study the box dimension of a generalized Weierstrass-type function, and make some further remarks.

2. Preliminaries and main results

2.1. The definition of generalized affine FIFs. Let $N \ge 2$ be a positive integer. Given a data set $\{(x_n, y_n)\}_{n=0}^N \subset \mathbb{R}^2$ with $x_0 < x_1 < \ldots < x_N$, we define a family of functions $\{W_n\}_{n=1}^N$ from $[x_0, x_N] \times \mathbb{R}$ to $[x_0, x_N] \times \mathbb{R}$ by

$$W_n(x,y) = (a_n x + b_n, S_n(x)y + q_n(x)), \quad 1 \le n \le N,$$

such that for each n, a_n and b_n are real numbers, S_n and q_n are continuous functions on $[x_0, x_N]$ with $|S_n(x)| < 1$ for all $x \in [x_0, x_N]$ and

$$W_n(x_0, y_0) = (x_{n-1}, y_{n-1}), \quad W_n(x_N, y_N) = (x_n, y_n)$$

According to Barnsley's classical result [4], there exists a unique continuous function f on $[x_0, x_N]$ such that its graph $\Gamma f := \{(x, f(x)) : x \in [x_0, x_N]\}$ is the invariant set of the iterated function system (IFS for short) $\{W_n : 1 \le n \le N\}$, i.e.,

(2.1)
$$\Gamma f = \bigcup_{n=1}^{N} W_n(\Gamma f).$$

Furthermore, the function f always interpolates the data set, i.e., $f(x_n) = y_n$ for all $0 \le n \le N$. The function f is called the *generalized affine fractal interpolation* function (generalized affine FIF for short) determined by the IFS $\{W_n\}_{n=1}^N$.

In the present paper, we will study $\dim_B \Gamma f$, the box dimension of the graph of f, where the following conditions are satisfied for each n:

- (A1) $x_n x_{n-1} = (x_N x_0)/N$,
- (A2) S_n is of bounded variation on $[x_0, x_N]$ and $|S_n(x)| < 1$ for all $x \in [x_0, x_N]$,
- (A3) q_n is of bounded variation on $[x_0, x_N]$.

2.2. Main results. In the rest of the paper, we write $I = [x_0, x_N]$ for simplicity. We define a function γ on I by

$$\gamma(x) = \sum_{n=1}^{N} |S_n(x)|.$$

We call γ the sum function of the family of vertical scaling functions $\mathbf{S} = \{S_n\}_{n=1}^N$. Write $\gamma^* = \max_{x \in I} \gamma(x)$ and $\gamma_* = \min_{x \in I} \gamma(x)$.

Given a closed interval J = [a, b], for each $k \in \mathbb{Z}^+$ and $1 \leq j \leq N^k$, we write

(2.2)
$$J_j^k = \left[a + \frac{j-1}{N^k}(b-a), a + \frac{j}{N^k}(b-a)\right]$$

Given $k \in \mathbb{Z}^+$, $1 \le n \le N$ and $1 \le j \le N^k$, we set

$$\overline{s}_{n,j}^k = \max_{x \in I_j^k} |S_n(x)|, \qquad \underline{s}_{n,j}^k = \min_{x \in I_j^k} |S_n(x)|.$$

Now, for every $k \in \mathbb{Z}^+$, we define a matrix \overline{M}_k by setting for $1 \le n \le N$, $1 \le \ell \le N^{k-1}$ and $1 \le j \le N^k$,

$$(\overline{M}_k)_{(n-1)N^{k-1}+\ell,j} = \begin{cases} \overline{s}_{n,j}^k, & \text{if } (\ell-1)N < j \le \ell N, \\ 0, & \text{otherwise.} \end{cases}$$

Similarly, we define another $N^k \times N^k$ matrix \underline{M}_k by replacing $\overline{s}_{n,j}^k$ with $\underline{s}_{n,j}^k$. We call \overline{M}_k (resp. \underline{M}_k) the upper (resp. the lower) vertical scaling matrix with level-k.

In this paper, we prove the monotonicity of spectral radii of vertical scaling matrices without additional assumptions. We also obtain the irreducibility of lower vertical scaling matrices under a weaker condition than that in [18].

Theorem 2.1. With previous notations, we have

- (1) $\rho(\overline{M}_k)$, the spectral radius of \overline{M}_k , is decreasing with respect to k. As a result, $\rho^* = \lim_{k \to \infty} \rho(\overline{M}_k)$ exists.
- (2) $\rho(\underline{M}_k)$ is increasing with respect to k. As a result, $\rho_* = \lim_{k \to \infty} \rho(\underline{M}_k)$ exists.
- (3) If $|S_n|$ is positive on I for all $1 \le n \le N$, then $\rho_* = \rho^*$.
- (4) If $\gamma_* \geq 1$ and S_n has only finitely many zero points on I for all $1 \leq n \leq N$, then \underline{M}_k is primitive for sufficiently large enough $k \in \mathbb{Z}^+$.

By using same arguments in [18], we can show that \overline{M}_k is primitive for all $k \in \mathbb{Z}^+$ if the function S_n is not identically zero on every subinterval of I for all $1 \leq n \leq N$.

Let $\operatorname{Var}(f, I)$ be the classical total variation of f on I. By using Theorem 2.1 and the irreduciblity of \overline{M}_k , we can obtain the estimate of the box dimension of Γf .

Theorem 2.2. Let f be a generalized affine FIF satisfying conditions (A1)-(A3). Then we have the following results on the box dimension of Γf .

- (1) Assume that the function \underline{S}_n is not identically zero on every subinterval of I for all $1 \le n \le N$. Then $\overline{\dim}_B \Gamma f \le \max\{1, 1 + \log_N \rho^*\}$.
- (2) Assume that $\gamma_* \geq 1$ and the function S_n has only finitely many zero points on I for all $1 \leq n \leq N$. If $\operatorname{Var}(f, I) = \infty$, then $\dim_B \Gamma f \geq 1 + \log_N \rho_*$.
- (3) Under the assumption of the previous item and the additional assumption that $\rho_* = \rho^*$, if $\operatorname{Var}(f, I) = \infty$ and $\rho_{\mathbf{S}} > 1$, then

$$\dim_B \Gamma f = 1 + \log_N \rho_{\mathbf{S}},$$

otherwise dim_B $\Gamma f = 1$. Here $\rho_{\mathbf{S}}$ is the common value of ρ_* and ρ^* .

We remark that the assumption $\gamma_* \geq 1$ in Theorems 2.1 and 2.2 can be replaced by a weaker assumption $\mathcal{ZM}(\mathbf{S}) \geq N-2$. Please see section 3 for the definition of $\mathcal{ZM}(\mathbf{S})$.

It is easy to see that

$$\gamma_* \le \rho_* \le \rho^* \le \gamma^*.$$

In section 5, we estimate the box dimension of Γf by γ_* and γ^* under weaker assumptions than these in Theorem 2.2. Akhtar, Prasad and Navascués [1] used $S_{\max} = \max\{|S_i(x)| : x \in [0, 1], 1 \le i \le N\}$ and $S_{\min} = \min\{|S_i(x)| : x \in [0, 1], 1 \le i \le N\}\}$ to estimate the box dimension of α -fractal functions, which is a special class of generalized affine FIFs. Our results are better than that in [1].

3. Analysis on vertical scaling matrices

3.1. Some well known theorems and definitions. We recall some notations and definitions in matrix analysis [15]. Given a matrix $A = (a_{ij})_{n \times n}$, we say A is non-negative (resp. positive), denoted by $A \ge 0$ (resp. A > 0), if $a_{ij} \ge 0$ (resp. $a_{ij} > 0$) for all i and j. Let $B = (b_{ij})_{n \times n}$ be another matrix. We write $A \ge B$ (resp. A > B) if $a_{ij} \ge b_{ij}$ (resp. $a_{ij} > b_{ij}$) for all i and j. Similarly, given $u = (u_1, \ldots, u_n), v = (v_1, \ldots, v_n) \in \mathbb{R}^n$, we write $u \ge v$ (resp. u > v) if $u_i \ge v_i$ (resp. $u_i > v_i$) for all i.

A nonnegative matrix $A = (a_{ij})_{n \times n}$ is called *irreducible* if for any $i, j \in \{1, \ldots, n\}$, there exists a finite sequence $i_0, \ldots, i_t \in \{1, \ldots, n\}$ such that $i_0 = i, i_t = j$ and $a_{i_{\ell-1},i_{\ell}} > 0$ for all $1 \leq \ell \leq t$. A is called *primitive* if there exists $k \in \mathbb{Z}^+$ such that $A^k > 0$. It is clear that a primitive matrix is irreducible.

Given an $n \times n$ matrix A, we write $\sigma(A)$ the set of all eigenvalues of A and define $\rho(A) = \max\{|\lambda| : \lambda \in \sigma(A)\}$. We call $\rho(A)$ the spectral radius of A.

The following two lemmas are well known. Please see [15, Chapter 8] for details.

Lemma 3.1 (Perron-Frobenius Theorem). Let $A = (a_{ij})_{n \times n}$ be an irreducible nonnegative matrix. Then

(1) $\rho(A)$ is positive,

(2) $\rho(A)$ is an eigenvalue of A and has a positive eigenvector,

(3) $\rho(A)$ increases if any element of A increases.

Lemma 3.2. Let $A = (a_{ij})_{n \times n}$ be a nonnegative matrix. Then $\rho(A)$ is an eigenvalue of A and there is a nonnegative nonzero vector x such that $Ax = \rho(A)x$.

3.2. Monotonicity of spectral radii of vertical scaling matrices.

Theorem 3.3. For all $k \in \mathbb{Z}^+$,

$$\rho(\overline{M}_{k+1}) \le \rho(\overline{M}_k).$$

As a result, $\lim_{k\to\infty} \rho(\overline{M}_k)$ exists, denoted by ρ^* .

In [18], we proved this theorem under an additional assumption. Essentially, we required that \overline{M}_k are irreducible for all k.

Proof. Similarly as in [18], we introduce another $N^{k+1} \times N^{k+1}$ matrix \overline{M}_k^* as follows:

$$(\overline{M}_k^*)_{(n-1)N^k+\ell,j} = \begin{cases} \overline{s}_{n,\ell}^k, & \text{if } (\ell-1)N < j \le \ell N \\ 0, & \text{otherwise,} \end{cases}$$

for $1 \leq n \leq N$, $1 \leq \ell \leq N^k$ and $1 \leq j \leq N^{k+1}$. Using the same arguments in the proof of [18, Theorem 3.3], we have $\overline{M}_{k+1} \leq \overline{M}_k^*$ so that $\rho(\overline{M}_{k+1}) \leq \rho(\overline{M}_k^*)$.

Now we prove that $\rho(\overline{M}_k^*) = \rho(\overline{M}_k)$. The proof is divided into two parts. Firstly we show that $\rho(\overline{M}_k) \ge \rho(\overline{M}_k^*)$. Write $\lambda = \rho(\overline{M}_k^*)$. From Lemma 3.2, λ is an eigenvalue of \overline{M}_k^* and there is a nonnegative nonzero vector $u = (u_1, \ldots, u_{N^{k+1}})^T$ such that $\overline{M}_k^* u = \lambda u$. We define a vector $u' = (u'_1, \ldots, u'_{N^k})^T$ by

$$u'_{j} = \sum_{p=(j-1)N+1}^{jN} u_{p}, \quad 1 \le j \le N^{k}.$$

It is clear that u' is also nonnegative and nonzero. By using the same arguments in the proof of [18, Theorem 3.3], we can obtain that $\overline{M}_k u' = \lambda u'$ so that λ is an eigenvalue of \overline{M}_k . Hence, $\rho(\overline{M}_k^*) = \lambda \leq \rho(\overline{M}_k)$.

Secondly we show that $\rho(\overline{M}_k) \leq \rho(\overline{M}_k^*)$. Without loss of generality, we may assume that $\mu := \rho(\overline{M}_k) > 0$. From Lemma 3.2, μ is an eigenvalue of \overline{M}_k and there is a nonnegative nonzero vector $v = (v_1, \ldots, v_{N^k})^T$ such that $\overline{M}_k v = \mu v$. We define a vector $v' = (v'_1, \ldots, v'_{N^{k+1}})^T$ by

$$v'_{(n-1)N^k+\ell} = \overline{s}^k_{n,\ell} v_\ell, \quad 1 \le n \le N, 1 \le \ell \le N^k$$

It is clear that v' is nonnegative. Furthermore, it follows from $\overline{M}_k v = \mu v$ that for all $1 \leq n \leq N$ and $1 \leq j \leq N^{k-1}$,

(3.1)
$$\mu v_{(n-1)N^{k-1}+j} = \sum_{t=(j-1)N+1}^{jN} \overline{s}_{n,t}^k v_t = \sum_{t=(j-1)N+1}^{jN} v'_{(n-1)N^k+t}.$$

Thus v' is a nonzero vector since otherwise, v is a zero vector which is a contradiction. For any $1 \le n \le N$ and $1 \le \ell \le N^k$, there exist $1 \le n' \le N$ and $1 \le j' \le N^{k-1}$ such that $\ell = (n'-1)N^{k-1} + j'$. Thus,

$$(\overline{M}_{k}^{*}v')_{(n-1)N^{k}+\ell} = \overline{s}_{n,\ell}^{k} \sum_{p=(\ell-1)N+1}^{\ell N} v'_{p} = \overline{s}_{n,\ell}^{k} \sum_{t=(j'-1)N+1}^{j'N} v'_{(n'-1)N^{k}+t}$$
$$= \overline{s}_{n,\ell}^{k} \mu v_{(n'-1)N^{k-1}+j'} \qquad (By (3.1))$$
$$= \mu \overline{s}_{n,\ell}^{k} v_{\ell} = \mu v'_{(n-1)N^{k}+\ell},$$

which implies $\overline{M}_{k}^{*}v' = \mu v'$ so that μ is an eigenvalue of \overline{M}_{k}^{*} . Hence, $\rho(\overline{M}_{k}) = \mu \leq \rho(\overline{M}_{k}^{*})$.

From the above arguments, $\rho(\overline{M}_{k+1}) \leq \rho(\overline{M}_k) = \rho(\overline{M}_k)$. Since $\rho(\overline{M}_k) \geq 0$ for all k, we know that $\lim_{k\to\infty} \rho(\overline{M}_k)$ exists.

We remark that compared to [18, Theorem 3.3], the different part in the above proof is the proof of $\rho(\overline{M}_k) \leq \rho(\overline{M}_k^*)$. Similarly, we can obtain the following result.

Theorem 3.4. For all $k \in \mathbb{Z}^+$,

 $\rho(\underline{M}_{k+1}) \ge \rho(\underline{M}_k).$

As a result, $\lim_{k\to\infty} \rho(\underline{M}_k)$ exists, denoted by ρ_* .

In the case that $\rho_* = \rho^*$, we denote the common value by $\rho_{\mathbf{S}}$. The following result has been proved in [18, Proposition 3.5] under the assumption that S_n is Lipschitz for all n. Our present proof only use the fact that S_n is continuous for all n.

Theorem 3.5. Assume that $|S_n|$ is positive on I for all $1 \le n \le N$. Then $\rho_* = \rho^*$. *Proof.* For any $1 \le n \le N$, from the fact that S_n is continuous and nonzero on I, we have

$$\underline{S}_n := \min\{|S_n(x)| : x \in I\} > 0.$$

Fix $\varepsilon > 0$. Since S_n is uniformly continuous on I for all n, we know that for sufficiently large k,

$$\overline{s}_{n,j}^k \leq \underline{s}_{n,j}^k + \varepsilon \underline{S}_n \leq (1+\varepsilon) \underline{s}_{n,j}^k, \quad 1 \leq n \leq N, \ 1 \leq j \leq N^k,$$

so that $\underline{M}_k \leq \overline{M}_k \leq (1 + \varepsilon)\underline{M}_k$. Thus,

$$\rho(\underline{M}_k) \le \rho(\overline{M}_k) \le (1+\varepsilon)\rho(\underline{M}_k)$$

for sufficiently large k. By letting k tend to infinity, $\rho_* \leq \rho^* \leq (1+\varepsilon)\rho_*$. From the arbitrariness of ε , we have $\rho_* = \rho^*$.

3.3. The irreducibility of vertical scaling matrices. Recall that

$$\gamma(x) = \sum_{n=1}^{N} |S_n(x)|, \quad x \in I,$$

and $\gamma^* = \max_{x \in I} \gamma(x), \ \gamma_* = \min_{x \in I} \gamma(x)$. For any $k \in \mathbb{Z}^+$, we define

$$\overline{\gamma}_k = \max_{1 \le j \le N^k} \sum_{n=1}^N \overline{s}_{n,j}^k, \quad \underline{\gamma}_k = \min_{1 \le j \le N^k} \sum_{n=1}^N \underline{s}_{n,j}^k.$$

Using the similar arguments in Theorem 3.5, we can obtain that

$$\gamma^* = \lim_{k \to \infty} \overline{\gamma}_k, \quad \gamma_* = \lim_{k \to \infty} \underline{\gamma}_k.$$

For every $k \in \mathbb{Z}^+$, from [15, Theorem 8.1.22], $\underline{\gamma}_k \leq \rho(\underline{M}_k) \leq \rho(\overline{M}_k) \leq \overline{\gamma}_k$. Hence,

$$\gamma_* \le \rho_* \le \rho^* \le \gamma^*.$$

Thus, if γ is a constant function on I, then $\gamma(x) = \rho_{\mathbf{S}}$ for all $x \in I$.

Using the same arguments in the proof of [18, Lemma 3.2], we have the following result.

Lemma 3.6. Assume that for each $1 \leq n \leq N$, vertical scaling function S_n is not identically zero on every subinterval of I. Then $(\overline{M}_k)^k > 0$ for all $k \in \mathbb{Z}^+$. As a result, \overline{M}_k is primitive for all $k \in \mathbb{Z}^+$.

Similarly, we can show that \underline{M}_k is primitive if $|S_n|$ is positive for each $1 \leq n \leq N$. However, it is much more involved to prove the primitivity of \underline{M}_k under general setting. In this paper, we will show that \underline{M}_k is primitive for sufficiently large k if $\gamma_* \geq 1$ and S_n has finitely many zero points for each $1 \leq n \leq N$.

Define the multiplicity of zero points of $\mathbf{S} = \{S_n : 1 \leq n \leq N\}$ at $x \in I$ by

$$\mathcal{ZM}(\mathbf{S}, x) = \operatorname{card} \{ n : S_n(x) = 0, 1 \le n \le N \},\$$

where card (A) is the cardinality of a set A. Write $\mathcal{ZM}(\mathbf{S}) = \max_{x \in I} \mathcal{ZM}(\mathbf{S}, x)$. We have the following simple fact.

Lemma 3.7. If $\gamma_* \geq 1$, then $\mathcal{ZM}(\mathbf{S}) \leq N-2$.

Proof. We prove this lemma by contradiction. Assume that there exists $\tilde{x} \in I$ such that $\mathcal{ZM}(\mathbf{S}, \tilde{x}) \geq N - 1$. Then there exists $1 \leq n_0 \leq N$, such that $S_n(\tilde{x}) = 0$ for all $n \neq n_0$. Hence,

$$\gamma_* \leq \gamma(\widetilde{x}) = \sum_{n=1}^N |S_n(\widetilde{x})| = |S_{n_0}(\widetilde{x})| < 1,$$

which contradicts the fact that $\gamma_* \geq 1$.

Lemma 3.8. Assume that $\mathcal{ZM}(\mathbf{S}) \leq N-2$ and the function S_n has finitely many zero points for all $1 \leq n \leq N$. Then there exists $k_1 \in \mathbb{Z}^+$ such that for all $k > k_1$, every row of \underline{M}_k has at least N-1 positive entries, and every column of \underline{M}_k has at least 2 positive entries.

Proof. Let Z_n be the set of zero points of S_n for $1 \le n \le N$ and write $Z = \bigcup_{n=1}^N Z_n$. Let E be the set of endpoints of all I_j^k for $1 \le j \le N^k$ and $k \ge 1$, i.e.,

$$E = \bigcup_{k \ge 1} \{ x_0 + j N^{-k} (x_N - x_0) : 0 \le j \le N^k \}.$$

Since Z is a finite set, there exists a positive integer k_1 satisfying the following two conditions:

- (1) $I_j^{k_1}$ contains at most one element of Z for all $1 \le j \le N^{k_1}$,
- (2) for every point $x \in Z \cap E$, there exists $1 \le j \le N^{k_1}$ such that x is the endpoint of $I_i^{k_1}$.

Then it is easy to see that for all $k > k_1$, every row of \underline{M}_k has at least N - 1 positive entries.

Notice that $\mathcal{ZM}(\mathbf{S}, x) \leq N - 2$ for all $x \in \mathbb{Z}$. Thus every column of \underline{M}_k has at least 2 positive entries.

Lemma 3.9. Under the assumptions of Lemma 3.8, for all $k > k_1$, every row of $(\underline{M}_k)^k$ has at least $(N-1)^k$ positive entries and every column of $(\underline{M}_k)^k$ has at least 2^k positive entries. Here k_1 is the constant in Lemma 3.8.

Proof. Fix $k > k_1$. For all $m \ge 1$ and $1 \le i \le N^k$, we define

$$row_m(i) = \{j : \left((\underline{M}_k)^m\right)_{ij} > 0\}.$$

Notice that $((\underline{M}_k)^{m+1})_{ij} = \sum_{t=1}^{N^k} (\underline{M}_k)_{it} ((\underline{M}_k)^m)_{tj}$ for all $m \ge 1$ and $1 \le i, j \le N^k$. Thus for all $m \ge 1$ and $1 \le i \le N^k$,

 $row_{m+1}(i) = \{j : \text{there exists } t \in row_1(i) \text{ such that } j \in row_m(t)\}.$

It follows from the definition of \underline{M}_k that for all $1 \leq i \leq N, 1 \leq \ell \leq N^{k-1}$,

(3.2)
$$row_1((i-1)N^{k-1}+\ell) \subset \{(\ell-1)N+1, (\ell-1)N+2, \dots, \ell N\}.$$

We claim that for each $1 \le m \le k - 1$,

$$row_m((i-1)N^{k-m}+\ell) \subset \{(\ell-1)N^m+1, (\ell-1)N^m+2, \dots, \ell N^m\}$$

for all $1 \le i \le N^m$ and $1 \le \ell \le N^{k-m}$.

It follows from (3.2) that the claim holds for m = 1. Assume that the claim holds for some $1 \le m \le k - 2$. Now given $1 \le i \le N^{m+1}$ and $1 \le \ell \le N^{k-(m+1)}$, we write $i' = (i-1)N^{k-(m+1)} + \ell$. If $j \in row_{m+1}(i')$, then there exists $t \in row_1(i')$ such that $j \in row_m(t)$. Notice that there exist unique integer pair (i_1, i_2) with $1 \le i_1 \le N$ and $1 \le i_2 \le N^m$ such that $i = (i_1 - 1)N^m + i_2$. Thus

$$i' = (i_1 - 1)N^{k-1} + (i_2 - 1)N^{k-(m+1)} + \ell.$$

Hence, from (3.2), $(i_2 - 1)N^{k-m} + (\ell - 1)N + 1 \le t \le (i_2 - 1)N^{k-m} + \ell N$. Combining this with the inductive assumption, we have $(\ell - 1)N^{m+1} + 1 \le j \le \ell N^{m+1}$ so that the claim holds for m + 1. This completes the proof of the claim.

It directly follows from the claim that for all $1 \leq m \leq k-1$ and $1 \leq i \leq N^k$, if $t_1 \neq t_2 \in row_1(i)$, then $row_m(t_1) \cap row_m(t_2) = \emptyset$, which implies that

(3.3)
$$\operatorname{card}\left(row_{m+1}(i)\right) = \sum_{t \in row_1(i)} \operatorname{card}\left(row_m(t)\right).$$

From Lemma 3.8, card $(row_1(i)) \ge N - 1$ for all $1 \le i \le N^k$. Combining this with (3.3), we can use inductive arguments to obtain that card $(row_m(i)) \ge (N-1)^m$ for all $1 \le m \le k$ and $1 \le i \le N^k$. Thus every row of $(\underline{M}_k)^k$ has at least $(N-1)^k$ positive entries.

Similarly, for all $m \ge 1$ and $1 \le j \le N^k$, we define

$$col_m(j) = \{i : \left((\underline{M}_k)^m\right)_{ij} > 0\}.$$

Then for all $m \ge 1$ and $1 \le j \le N^k$,

 $col_{m+1}(j) = \{i : \text{there exists } t \in col_1(j) \text{ such that } i \in col_m(t)\}.$

By using similar arguments as above, we can obtain that for each $1 \le m \le k-1$,

$$col_m((j-1)N^m + \ell) \subset \{j, j+N^{k-m}, \dots, j+(N^m-1)N^{k-m}\}$$

for all $1 \leq j \leq N^{k-m}$ and $1 \leq \ell \leq N^m$. Hence, for all $1 \leq m \leq k-1$ and $1 \leq j \leq N^k$, if $t_1 \neq t_2 \in col_1(j)$, then $col_m(t_1) \cap col_m(t_2) = \emptyset$. As a result, we have $card(col_m(j)) \geq 2^m$ for all $1 \leq m \leq k$ and $1 \leq j \leq N^k$, which implies that every column of $(\underline{M}_k)^k$ has at least 2^k positive entries. \Box

The following result is part of the statement in [15, 8.5.P5]. We will use it to prove that \underline{M}_k is primitive for sufficiently large k under certain conditions.

Lemma 3.10 ([15]). Let $A = (a_{ij})_{n \times n}$ be an irreducible nonnegative matrix. Assume that at least one of its main diagonal entry $a_{ii}(1 \le i \le n)$ is positive. Then A is primitive.

Theorem 3.11. Assume that $\mathcal{ZM}(\mathbf{S}) \leq N-2$ and the function S_n has finitely many zero points for each $1 \leq n \leq N$. Then there exists $k_0 \in \mathbb{Z}^+$ such that \underline{M}_k is primitive for all $k > k_0$.

Proof. For every $1 \le n \le N$ and $1 \le j \le N^k$, we call $\underline{s}_{n,j}^k$ a basic entry of the matrix \underline{M}_k . If all basic entries are positive, then using the same arguments in the proof of [18, Lemma 3.2], we can obtain that $(\underline{M}_k)^k > 0$.

In the case that N = 2, we have $\mathcal{ZM}(\mathbf{S}) = 0$ so that $\mathcal{ZM}(\mathbf{S}, x) = 0$ for all $x \in I$. Thus $S_n(x) \neq 0$ for n = 1, 2 and all $x \in I$. It follows that all basic entries are positive so that $(\underline{M}_k)^k > 0$. Hence \underline{M}_k is primitive for all $k \in \mathbb{Z}^+$.

Now we assume that $N \geq 3$. Let m_n be the number of zero points of S_n on I. Write $m = \sum_{n=1}^{N} m_n$. Then for any $k \geq 1$, there are at most 2m basic entries equal to zero. Notice that for every $k \geq 1$ and $1 \leq i, j \leq N^k$, the (i, j) entry of $(\underline{M}_k)^k$ is

(3.4)
$$\left((\underline{M}_k)^k\right)_{ij} = \sum_{\substack{1 \le t_2, \dots, t_k \le N^k \\ t_1 = i, t_{k+1} = j}} \prod_{\ell=1}^k (\underline{M}_k)_{t_\ell, t_{\ell+1}},$$

and both every row and every column of \underline{M}_k have N basic entries. Hence, a zero basic entry of \underline{M}_k can make at most kN^{k-1} entries of $(\underline{M}_k)^k$ to be zero. Thus there are at most $2mkN^{k-1}$ zero entries in $(\underline{M}_k)^k$.

Let k_1 be the constant in Lemma 3.9 and $k_0 = \max\{2, m, k_1\}$. We claim that $(\underline{M}_k)^k$ is irreducible for all $k > k_0$.

We prove the claim by contradiction. Assume that $(\underline{M}_k)^k$ is reducible. Then there are nonempty and disjoint subsets A, B of $\{1, \ldots, N^k\}$ satisfying $A \cup B = \{1, \ldots, N^k\}$, and for all $i \in A$ and $j \in B$, the (i, j) entry of $(\underline{M}_k)^k$ is zero. From Lemma 3.9 and $N-1 \geq 2$, there are at least 2^k elements in both A and B. Hence

$$\operatorname{card}(A) \cdot \operatorname{card}(B) = \operatorname{card}(A) \cdot (N^k - \operatorname{card}(B)) \ge 2^k (N^k - 2^k)$$

so that $(\underline{M}_k)^k$ has at least $2^k(N^k - 2^k)$ zero entries. From $N \ge 3$ and $k > k_0$, we have $2^k > k^2 > mk$ and $N^k - 2^k > 2N^{k-1}$ so that $2^k(N^k - 2^k) > 2mkN^{k-1}$, which is a contradiction. This completes the proof of the claim.

Now we will show that for all $k > k_0$, at least one of the main diagonal entry of $(\underline{M}_k)^k$ is positive, so that $(\underline{M}_k)^k$ is primitive by Lemma 3.10. As a result, \underline{M}_k is primitive for all $k > k_0$.

For any $j \in \{1, 2, ..., N^k\}$, there exists a unique finite sequence $j_1, ..., j_k \in \{1, ..., N\}$ such that

$$j = (j_1 - 1)N^{k-1} + (j_2 - 1)N^{k-2} + \dots + (j_{k-1} - 1)N + j_k.$$

We define $\sigma(j) = (j - (j_1 - 1)N^{k-1})N + j_1$. Then $\sigma(j) \in \{1, 2, ..., N^k\}$. Thus we can define $\sigma^p(j) = \sigma(\sigma^{p-1}(j))$ for $p \ge 2$. It is easy to see that $\sigma^k(j) = j$ and $(\underline{M}_k)_{j,\sigma(j)}$ is a basic entry of \underline{M}_k for all $j \in \{1, ..., N^k\}$.

Write $\sigma^0(j) = j$. From (3.4), $\left((\underline{M}_k)^k\right)_{jj} \geq \prod_{p=1}^k (\underline{M}_k)_{\sigma^{p-1}(j),\sigma^p(j)}$. Hence, a zero basic entry of \underline{M}_k can make at most k main digonal entries of $(\underline{M}_k)^k$ to be zero. Thus there are at most 2mk zero main digonal entries in $(\underline{M}_k)^k$.

Notice that $k_0 \ge \max\{2, m\}$ and $N \ge 3$. Hence, for $k > k_0$, we have $N^k \ge 3^k > 2k^2 > 2mk$ so that $(\underline{M}_k)^k$ contains at least one positive main diagonal entry. \Box

From Lemma 3.7 and Theorems 3.3, 3.4, 3.5 and 3.11, we know that Theorem 2.1 holds.

4. Proof of Theorem 2.2

In the rest of the paper, we always assume that f is a generalized affine FIFs satisfying conditions (A1)-(A3).

4.1. Box dimension estimate of the graph of continuous functions. Given a bounded subset E of \mathbb{R}^d , we use $\overline{\dim}_B E$ and $\underline{\dim}_B E$ to denote the upper box dimension and the lower box dimension of E, respectively. If $\overline{\dim}_B E = \underline{\dim}_B E$, then we use $\dim_B E$ to denote the common value and call it the *box dimension* of E. It is well known that $\underline{\dim}_B E \geq 1$ when E is the graph of a continuous function on a closed interval of \mathbb{R} . Please see [12] for details.

Let g be a continuous function on J. For any $U \subset J$, we use O(g, U) to denote the oscillation of g on U, that is,

$$O(g, U) = \sup_{x', x'' \in U} |g(x') - g(x'')|.$$

Write

$$O_k(g,J) = \sum_{j=1}^{N^k} O(g,J_j^k),$$

where J_i^k is defined by (2.2).

The following lemma presents a method to estimate the upper and lower box dimensions of the graph of a function by its oscillation. Similar results can be found in [12, 20, 23].

Lemma 4.1 ([18]). Let g be a continuous function on a closed interval J. Then

$$\underline{\dim}_B \Gamma g \ge 1 + \underline{\lim}_{k \to \infty} \frac{\log \left(O_k(g, J) + 1 \right)}{k \log N}, \quad and$$
$$\overline{\dim}_B \Gamma g \le 1 + \underline{\lim}_{k \to \infty} \frac{\log \left(O_k(g, J) + 1 \right)}{k \log N}.$$

We remark that J = [0, 1] in the original version of the above lemma in [18]. However, it is straightforward to see that the lemma still holds in the present version.

It is clear that $\{O_k(g, J)\}_{k=1}^{\infty}$ is increasing with respect to k. Thus $\lim_{k\to\infty} O_k(g, J)$ always exists. Write $\operatorname{Var}(g, J)$ the classical total variation of g on J. We have the following simple fact.

Lemma 4.2. Let g be a continuous function on a closed interval J = [a, b]. Then $\lim_{k\to\infty} O_k(g, J) = \operatorname{Var}(g, J)$.

Proof. Clearly, $O_k(g, J) \leq \operatorname{Var}(g, J)$ for all $k \in \mathbb{Z}^+$. Thus $\lim_{k \to \infty} O_k(g, J) \leq \operatorname{Var}(g, J)$. Now we prove the another inequality.

Arbitrarily pick a partition $T = \{a = t_0 < t_1 < \cdots < t_n = b\}$ of J. Fix $k \in \mathbb{Z}^+$ large enough such that $N^{-k} < \min\{t_i - t_{i-1} : 1 \le i \le n\}$. For every $0 \le i \le n$, there exists $\alpha_i \in \{1, \ldots, N^k\}$ such that $t_i \in J_{\alpha_i}^k$. Furthermore, it is easy to see that $1 = \alpha_0 < \alpha_1 < \cdots < \alpha_n = N^k$. Notice that for any $1 \le i \le n$,

$$|g(t_i) - g(t_{i-1})| \le \sum_{p=\alpha_{i-1}}^{\alpha_i} O(g, J_p^k).$$

Thus

$$\sum_{i=1}^{n} |g(t_i) - g(t_{i-1})| \le \sum_{i=1}^{n} \sum_{p=\alpha_{i-1}}^{\alpha_i} O(g, J_p^k)$$
$$= O_k(g, J) + \sum_{i=1}^{n-1} O(g, J_{\alpha_i}^k) \le \lim_{k \to \infty} O_k(g, J) + \sum_{i=1}^{n-1} O(g, J_{\alpha_i}^k).$$

Since g is continuous on I, we can choose k large enough such that $\sum_{i=1}^{n-1} O(g, J_{\alpha_i}^k)$ as small as possible. Hence

$$\sum_{i=1}^{n} |g(t_i) - g(t_{i-1})| \le \lim_{k \to \infty} O_k(g, J).$$

By the arbitrariness of the partition T, $\operatorname{Var}(g, J) \leq \lim_{k \to \infty} O_k(g, J)$.

10

4.2. Estimate of oscillations. By the definition of W_n , it is easy to see that $W_n(x,y) = (L_n(x), F_n(x,y))$, where

$$L_n(x) = (x - x_0)/N + x_{n-1}, \quad F_n(x, y) = S_n(x)y + q_n(x).$$

From (2.1), $W_n(x, f(x)) = (L_n(x), f(L_n(x)))$. Thus, we have the following useful equality:

(4.1)
$$f(L_n(x)) = S_n(x)f(x) + q_n(x), \quad x \in [x_0, x_N], \ n = 1, 2, \dots, N.$$

Write $M_f = \max_{x \in I} |f(x)|$. By using the similar arguments in the proof of [18, Lemma 4.2], we can obtain the following lemma.

Lemma 4.3. For any $1 \le n \le N$ and $D \subset I$,

$$O(f, L_n(D)) \le \sup_{x \in D} |S_n(x)| O(f, D) + M_f O(S_n, D) + O(q_n, D), \quad and \\ O(f, L_n(D)) \ge \inf_{x \in D} |S_n(x)| O(f, D) - M_f O(S_n, D) - O(q_n, D).$$

Proof. From (4.1),

$$O(f, L_n(D)) = \sup_{x', x'' \in D} \left| S_n(x')f(x') - S_n(x'')f(x'') + q_n(x') - q_n(x'') \right|$$

$$\leq \sup_{x', x'' \in D} \left| S_n(x') \left(f(x') - f(x'') \right) \right| + \sup_{x', x'' \in D} \left| f(x'') \left(S_n(x') - S_n(x'') \right) \right|$$

$$+ \sup_{x', x'' \in D} \left| q_n(x') - q_n(x'') \right|$$

$$\leq \sup_{x \in D} \left| S_n(x) \left| O(f, D) + M_f O(S_n, D) + O(q_n, D) \right|.$$

On the other hand, we choose $x', x'' \in D$ such that O(f, D) = |f(x') - f(x'')|. Then

$$O(f, L_n(D)) \ge |f(L_n(x')) - f(L_n(x''))| \ge |S_n(x')(f(x') - f(x''))| - |f(x'')(S_n(x') - S_n(x''))| - |q_n(x') - q_n(x'')| \ge \inf_{x \in D} |S_n(x)|O(f, D) - M_fO(S_n, D) - O(q_n, D).$$

Thus the lemma holds.

Using the argument similar to the proof of the first part of the above lemma, we have the following result.

Lemma 4.4. For any $1 \le n \le N$, $D \subset I$ and $t \in D$,

$$|O(f, L_n(D)) - |S_n(t)|O(f, D)| \le 2M_f O(S_n, D) + O(q_n, D).$$

Proof. For any $x', x'' \in D$, we have

$$\left|S_n(x')f(x') - S_n(x'')f(x'') - S_n(t)(f(x') - f(x''))\right| \le 2M_f O(S_n, D).$$

Thus

$$|S_n(t)|O(f,D) - 2M_f O(S_n,D) \le |S_n(x')f(x') - S_n(x'')f(x'')| \le |S_n(t)|O(f,D) + 2M_f O(S_n,D)|$$

so that the lemma holds.

Given $k, p \in \mathbb{Z}^+$ and $g \in C(I)$, we define

$$V(g,k,p) = \left(O_p(g,I_1^k), O_p(g,I_2^k), \dots, O_p(g,I_{N^k}^k)\right)^T \in \mathbb{R}^{N^k},$$

and call it an oscillation vector of g with respect to (k, p). It is obvious that

$$O_{k+p}(g, I) = ||V(g, k, p)||_1,$$

where $||v||_1 := \sum_{i=1}^d |v_i|$ for any $v = (v_1, \dots, v_d) \in \mathbb{R}^d$. Define a vectors ξ_k in \mathbb{R}^{N^k} by

(4.2)
$$(\xi_k)_{(n-1)N^{k-1}+\ell} = M_f \operatorname{Var}(S_n, I_\ell^{k-1}) + \operatorname{Var}(q_n, I_\ell^{k-1}),$$

where $1 \le n \le N, 1 \le \ell \le N^{k-1}$.

Lemma 4.5. For any $k \in \mathbb{Z}^+$ and any $p \in \mathbb{Z}^+$,

(4.3)
$$-\xi_k + \underline{M}_k V(f,k,p) \le V(f,k,p+1) \le \xi_k + \overline{M}_k V(f,k,p).$$

Proof. From Lemma 4.3, for any $1 \le n \le N$, $k \in \mathbb{Z}^+$, $1 \le j \le N^k$ and any $D \subset I_j^k$,

$$O(f, L_n(D)) \le \overline{s}_{n,j}^k O(f, D) + O(q_n, D) + M_f O(S_n, D).$$

Notice that $(L_n(I_j^k))_m^p = L_n((I_j^k)_m^p)$ for $1 \le m \le N^p$. Thus,

$$O_{p}(f, L_{n}(I_{j}^{k})) = \sum_{m=1}^{N^{p}} O\left(f, \left(L_{n}(I_{j}^{k})\right)_{m}^{p}\right)$$

$$\leq \sum_{m=1}^{N^{p}} \left(\overline{s}_{n,j}^{k} O(f, (I_{j}^{k})_{m}^{p})\right) + O(q_{n}, (I_{j}^{k})_{m}^{p}) + M_{f} O(S_{n}, (I_{j}^{k})_{m}^{p})\right)$$

$$= \overline{s}_{n,j}^{k} O_{p}(f, I_{j}^{k}) + O_{p}(q_{n}, I_{j}^{k}) + M_{f} O_{p}(S_{n}, I_{j}^{k}).$$

Hence, from $I_{\ell}^{k-1} = \bigcup_{j=(\ell-1)N+1}^{\ell N} I_j^k$,

$$O_{p+1}(f, L_n(I_{\ell}^{k-1})) = \sum_{j=(\ell-1)N+1}^{\ell N} O_p(f, L_n(I_j^k))$$

$$\leq \sum_{j=(\ell-1)N+1}^{\ell N} \left(\overline{s}_{n,j}^k O_p(f, I_j^k) + O_p(q_n, I_j^k) + M_f O_p(S_n, I_j^k) \right)$$

$$\leq M_f O_{p+1}(S_n, I_{\ell}^{k-1}) + O_{p+1}(q_n, I_{\ell}^{k-1}) + \sum_{j=(\ell-1)N+1}^{\ell N} \overline{s}_{n,j}^k O_p(f, I_j^k).$$

By the definitions of ξ_k and \overline{M}_k , we can rewrite this inequality as

$$O_{p+1}(f, I_{(n-1)N^{k-1}+\ell}^k) \le (\xi_k)_{(n-1)N^{k-1}+\ell} + (\overline{M}_k V(f, k, p))_{(n-1)N^{k-1}+\ell}$$

so that $V(f, k, p + 1) \leq \xi_k + \overline{M}_k V(f, k, p)$. Similarly, we can prove that another inequality in (4.3) holds.

4.3. Estimate the box dimension of Γf by ρ_* and ρ^* .

Theorem 4.6. Assume that the function S_n is not identically zero on every subinterval of I for all $1 \le n \le N$. Then

(4.4)
$$\overline{\dim}_B \Gamma f \le \max\left\{1, 1 + \log_N \rho^*\right\}.$$

Proof. Fix $k \in \mathbb{Z}^+$. Let ξ_k be the vector in \mathbb{R}^{N^k} defined by (4.2). From Lemma 3.6, \overline{M}_k is primitive so that it is irreducible. By Lemma 3.1, we can choose a positive eigenvector w_k of \overline{M}_k such that $w_k \geq \xi_k$ and $w_k \geq V(f, p, 1)$. Hence, from Theorem 4.5, we have

$$V(f, k, p+1) \le w_k + \overline{M}_k V(f, k, p)$$

for all $p \in \mathbb{Z}^+$. Thus,

$$V(f,k,p) \le w_k + \overline{M}_k w_k + \dots + (\overline{M}_k)^{p-2} w_k + (\overline{M}_k)^{p-1} V(f,k,1)$$
$$\le \sum_{\ell=0}^{p-1} \rho(\overline{M}_k)^{\ell} w_k$$

for all $p \in \mathbb{Z}^+$. It follows that

$$O_{k+p}(f,I) = ||V(f,k,p)||_1 \le ||w_k||_1 \sum_{\ell=0}^{p-1} (\rho(\overline{M}_k))^{\ell} \le ||w_k||_1 p(\rho(\overline{M}_k)^p + 1).$$

Hence,

$$\overline{\lim_{p \to \infty} \frac{\log(O_{k+p}(f, I) + 1)}{p \log N}} \le \max\left\{0, \frac{\log \rho(\overline{M}_k)}{\log N}\right\}.$$

Thus, from Lemma 4.1,

$$\overline{\dim}_B \Gamma f \le 1 + \lim_{p \to \infty} \frac{\log(O_{k+p}(f, I) + 1)}{p \log N} \le \max\left\{1, 1 + \frac{\log \rho(\overline{M}_k)}{\log N}\right\}.$$

By the arbitrariness of k, we know from Theorem 3.3 that (4.4) holds.

Theorem 4.7. Assume that $\operatorname{Var}(f, I) = \infty$, $\mathcal{ZM}(\mathbf{S}) \leq N - 2$ and the function S_n has finitely many zero points on I for all $1 \leq n \leq N$. Then

(4.5)
$$\underline{\dim}_B \Gamma f \ge 1 + \log_N \rho_*.$$

Proof. Notice that $\underline{\dim}_B \Gamma f \geq 1$ always holds. Thus, without loss of the generality, we may assume that $\rho_* > 1$. From $\rho_* = \lim_{k \to \infty} \rho(\underline{M}_k)$, there exists a positivie integer k_2 , such that $\rho(\underline{M}_k) > 1$ for all $k > k_2$. Let k_0 be the constant in Theorem 3.11. From Theorem 3.11, \underline{M}_k is primitive so that it is irreducible for all $k > k_0$.

Fix $k > \max\{k_0, k_2\}$. Given $1 < \tau < \rho(\underline{M}_k)$, from Lemma 3.1, we can find a positive eigenvector w_k of \underline{M}_k with eigenvalue $\rho(\underline{M}_k)$ such that $w_k \ge \xi_k/(\rho(\underline{M}_k) - \tau)$. Since \underline{M}_k is primitive, there exists $\ell_k \in \mathbb{Z}^+$ such that $(\underline{M}_k)^{\ell_k} > 0$. Let α_k be the minimal entry of the matrix $(\underline{M}_k)^{\ell_k}$. Then $\alpha_k > 0$. From Theorem 4.5,

(4.6)
$$V(f,k,p+1) \ge \underline{M}_k V(f,k,p) - \xi_k$$

for all $p \in \mathbb{Z}^+$. Repeatedly using this inequality, we can obtain that for all $p \in \mathbb{Z}^+$,

(4.7)
$$V(f,k,p+\ell_k) \ge (\underline{M}_k)^{\ell_k} V(f,k,p) - \sum_{q=0}^{\ell_k-1} (\underline{M}_k)^q \xi_k$$

Notice that the maximal entry of V(f, k, p) is at least $N^{-k} ||V(f, k, p)||_1$. Thus,

$$(\underline{M}_k)^{\ell_k} V(f,k,p) \ge (\alpha'_k,\ldots,\alpha'_k),$$

where $\alpha'_k = \alpha_k N^{-k} ||V(f,k,p)||_1$. Notice that

$$\lim_{p \to \infty} \|V(f, k, p)\|_1 = \lim_{p \to \infty} O_{k+p}(f, I) = \operatorname{Var}(f, I) = \infty.$$

Hence, we can choose p_* large enough such that

$$(\underline{M}_k)^{\ell_k} V(f,k,p_*) \ge w_k + \sum_{q=0}^{\ell_k - 1} (\underline{M}_k)^q \xi_k.$$

Let $p_k = p_* + \ell_k$. Then from (4.7),

$$V(f,k,p_k) \ge w_k \ge \frac{1}{\rho(\underline{M}_k) - \tau} \xi_k.$$

From (4.6),

 $V(f,k,p_k+1) \ge \rho(\underline{M}_k)w_k - \xi_k \ge \rho(\underline{M}_k)w_k - (\rho(\underline{M}_k) - \tau)w_k = \tau w_k.$ Notice that for all $\ell \in \mathbb{Z}^+$,

$$\rho(\underline{M}_k)\tau^{\ell}w_k - \xi_k = \rho(\underline{M}_k)(\tau^{\ell} - 1)w_k + \rho(\underline{M}_k)w_k - \xi_k$$

$$\geq \tau(\tau^{\ell} - 1)w_k + \tau w_k = \tau^{\ell+1}w_k.$$

Thus, by induction, $V(f, k, p_k + \ell) \ge \tau^{\ell} w_k$ for all $\ell \in \mathbb{Z}^+$. Hence

$$O_{k+p_k+\ell}(f,I) = \|V(f,k,p_k+\ell)\|_1 \ge \tau^{\ell} \|w_k\|_1$$

which implies that

$$\lim_{\ell \to \infty} \frac{\log \left(O_{\ell}(f, I) + 1 \right)}{\ell \log N} = \lim_{\ell \to \infty} \frac{\log \left(O_{k+p_k+\ell}(f, I) + 1 \right)}{\ell \log N} \ge \frac{\log \tau}{\log N}$$

It follows from the arbitrariness of τ that $\log \rho(\underline{M}_k) / \log N$ is less than the left hand side of this inequality. Combining this with Lemma 4.1, we have

$$\underline{\dim}_B \Gamma f \ge 1 + \frac{\log \rho(\underline{M}_k)}{\log N}$$

Since this result holds for all $k > \max\{k_0, k_2\}$, we know from Theorem 3.4 that (4.5) holds.

Remark 4.8. From the proof of the above theorem, it is easy to see that under the assumptions of the theorem, $\operatorname{Var}(f, I_i^k) = \infty$ for any $k \in \mathbb{Z}^+$ and $1 \leq j \leq N^k$.

Theorem 4.9. Under the assumption of Theorem 4.7 and the additional assumption that $\rho_* = \rho^*$, if $\operatorname{Var}(f, I) = \infty$ and $\rho_{\mathbf{S}} > 1$, then

(4.8)
$$\dim_B \Gamma f = 1 + \log_N \rho_{\mathbf{S}},$$

otherwise $\dim_B \Gamma f = 1$.

Proof. In the case that $\operatorname{Var}(f, I) < \infty$, we know from Lemma 4.1 that $\dim_B \Gamma f \leq 1$. In the case that $\rho_{\mathbf{S}} \leq 1$, we know from Theorem 4.6 that $\overline{\dim}_B \Gamma f \leq 1$. Since $\underline{\dim}_B \Gamma f \geq 1$ always holds, $\dim_B \Gamma f = 1$ if $\operatorname{Var}(f, I) < \infty$ or $\rho_{\mathbf{S}} \leq 1$.

In the case that $\operatorname{Var}(f, I) = \infty$ and $\rho_{\mathbf{S}} > 1$, we know from Theorems 4.6 and 4.7 that (4.8) holds.

From Lemma 3.7 and Theorems 4.6, 4.7 and 4.9, we know that Theorem 2.2 holds. Furthermore, from Theorems 3.5 and 4.9, we have the following result.

Corollary 4.10. Assume that the function $|S_n|$ is positive on I for each $1 \le n \le N$. Then in the case that $\operatorname{Var}(f, I) = \infty$ and $\rho_{\mathbf{S}} > 1$, (4.8) holds, otherwise $\dim_B \Gamma f = 1$.

5. Estimate the box dimension of FIFs by γ^* and γ_*

In this section, we will estimate the box dimension of Γf by the sum function of vertical scaling functions. By Lemma 4.4, we can obtain the following result.

Lemma 5.1. For all $k \in \mathbb{Z}^+$,

(5.1)
$$O_{k+1}(f,I) \le \gamma^* \cdot O_k(f,I) + \sum_{n=1}^N (\operatorname{Var}(q_n,I) + 2M_f \operatorname{Var}(S_n,I)), \text{ and}$$

(5.2)
$$O_{k+1}(f,I) \ge \gamma_* \cdot O_k(f,I) - \sum_{n=1}^N \left(\operatorname{Var}(q_n,I) + 2M_f \operatorname{Var}(S_n,I) \right).$$

Proof. Given $D \subset I$, we know from Lemma 4.4 that for any $t \in D$,

$$\sum_{n=1}^{N} O(f, L_n(D)) \le \gamma(t) \cdot O(f, D) + \sum_{n=1}^{N} \left(O(q_n, D) + 2M_f O(S_n, D) \right)$$
$$\le \gamma^* \cdot O(f, D) + \sum_{n=1}^{N} \left(O(q_n, D) + 2M_f O(S_n, D) \right).$$

For any $k \in \mathbb{Z}^+$ and $1 \leq j \leq N^k$, by letting $D = I_j^k$ in the above inequality, we have

$$\sum_{n=1}^{N} O(f, L_n(I_j^k)) \le \gamma^* \cdot O(f, I_j^k) + \sum_{n=1}^{N} \left(O(q_n, I_j^k) + 2M_f O(S_n, I_j^k) \right).$$

Hence

$$O_{k+1}(f,I) = \sum_{n=1}^{N} \sum_{j=1}^{N^{k}} O(f, L_{n}(I_{j}^{k}))$$

$$\leq \gamma^{*} \cdot O_{k}(f,I) + \sum_{n=1}^{N} \left(O_{k}(q_{n},I) + 2M_{f}O_{k}(S_{n},I) \right)$$

$$\leq \gamma^{*} \cdot O_{k}(f,I) + \sum_{n=1}^{N} \left(\operatorname{Var}(q_{n},I) + 2M_{f}\operatorname{Var}(S_{n},I) \right),$$

so that (5.1) holds. Similarly, we can prove that (5.2) holds.

From this lemma, we can obtain the upper box dimension estimate by γ^* and the lower box dimension estimate by γ_* .

Theorem 5.2. We have $\overline{\dim}_B \Gamma f \leq \max\{1, 1 + \log_N \gamma^*\}$. Furthermore, if $\gamma_* > 1$ and $\operatorname{Var}(f, I) = \infty$, then $\underline{\dim}_B \Gamma f \geq 1 + \log_N \gamma_*$.

Proof. Write $\eta = \sum_{n=1}^{N} (\operatorname{Var}(q_n, I) + 2M_f \operatorname{Var}(S_n, I))$. It is clear that $\eta < \infty$ since S_n and q_n are of bounded variation on I for each n. If $\gamma^* \leq 1$, from Lemma 5.1,

$$O_{k+1}(f,I) \le O_k(f,I) + \eta, \quad \forall k \ge 1,$$

so that

$$O_k(f,I) \le O_1(f,I) + (k-1)\eta, \quad \forall k \ge 1.$$

Thus from Lemma 4.1, $\overline{\dim}_B \Gamma f \leq 1 = \max\{1, 1 + \log_N \gamma^*\}.$

In the case that $\gamma^* > 1$, we know from Lemma 5.1 that

$$O_{k+1}(f,I) + \frac{\eta}{\gamma^* - 1} \le \gamma^* \Big(O_k(f,I) + \frac{\eta}{\gamma^* - 1} \Big), \quad \forall k \ge 1,$$

so that

$$O_k(f,I) + \frac{\eta}{\gamma^* - 1} \le (\gamma^*)^{k-1} \Big(O_1(f,I) + \frac{\eta}{\gamma^* - 1} \Big), \quad \forall k \ge 1.$$

Thus from Lemma 4.1, $\overline{\dim}_B \Gamma f \leq 1 + \log_N \gamma^* = \max\{1, 1 + \log_N \gamma^*\}.$

Now we assume that $\gamma_* > 1$ and $Var(f, I) = \infty$. Using Lemma 5.1 again, we have

(5.3)
$$O_{k+1}(f,I) - \frac{\eta}{\gamma_* - 1} \ge \gamma_* \Big(O_k(f,I) - \frac{\eta}{\gamma_* - 1} \Big), \quad \forall k \ge 1.$$

Since $\operatorname{Var}(f, I) = \infty$, from Lemma 4.2, there exists $k_0 \in \mathbb{Z}^+$ such that $O_{k_0}(f, I) > \eta/(\gamma_* - 1)$. From (5.3),

$$O_k(f,I) - \frac{\eta}{\gamma_* - 1} \ge (\gamma_*)^{k - k_0} \Big(O_{k_0}(f,I) - \frac{\eta}{\gamma_* - 1} \Big), \quad \forall k \ge k_0.$$

Thus from Lemma 4.1, $\underline{\dim}_B \Gamma f \ge 1 + \log_N \gamma_*$. Hence, the theorem holds.

Remark 5.3. From the proof of Theorem 5.2, it is easy to see that under the condition $\gamma_* > 1$, the following two properties are equivalent:

(1) $\operatorname{Var}(f, I) = \infty$,

(2) there exists $k_0 \in \mathbb{Z}^+$ such that

$$O_{k_0}(f,I) > (\gamma_* - 1)^{-1} \sum_{n=1}^{N} (\operatorname{Var}(q_n,I) + 2M_f \operatorname{Var}(S_n,I)).$$

Remark 5.4. Under the condition that the function S_n is nonnegative for each n, from (4.1),

$$\sum_{n=1}^{N} f(L_n(x)) = \sum_{n=1}^{N} \left(S_n(x) f(x) + q_n(x) \right) = \gamma(x) f(x) + \sum_{n=1}^{N} q_n(x).$$

Thus, by using arguments similar to the proof of [18, Theorem 4.10], we have

$$O_{k+1}(f,I) \ge \gamma_* O_k(f,I) - M_f \operatorname{Var}(\gamma,I) - \operatorname{Var}\Big(\sum_{n=1}^N q_n,I\Big).$$

Thus, if $\gamma_* > 1$ and the function S_n is nonnegative on I for each $1 \leq n \leq N$, then $\operatorname{Var}(f, I) = \infty$ if and only if there exists $k_0 \in \mathbb{Z}^+$ satisfying

$$O_{k_0}(f,I) > (\gamma_* - 1)^{-1} \Big(M_f \operatorname{Var}(\gamma,I) + \operatorname{Var}(\sum_{n=1}^N q_n,I) \Big).$$

From Theorem 5.2, we can obtain the following result.

Theorem 5.5. Assume that $\gamma(x) \equiv \gamma_0$ for all $x \in I$. Then in the case that $\gamma_0 > 1$ and $\operatorname{Var}(f, I) = \infty$,

(5.4)
$$\dim_B \Gamma f = 1 + \log_N \gamma_0,$$

otherwise $\dim_B \Gamma f = 1$.

Proof. Notice that $\underline{\dim}_B \Gamma f \geq 1$ always holds since f is a continuous function on I. In the case that $\gamma_0 \leq 1$, it follows from Theorem 5.2 that $\overline{\dim}_B \Gamma f \leq 1$. In the case that $\operatorname{Var}(f, I) < \infty$, we have $\lim_{k \to \infty} O_k(f, I) < \infty$. Thus, from Lemma 4.1, $\overline{\dim}_B \Gamma f \leq 1$. Hence $\dim_B \Gamma f = 1$ if $\gamma_0 \leq 1$ or $\operatorname{Var}(f, I) < \infty$.

Now we assume that $\gamma_0 > 1$ and $\operatorname{Var}(f, I) = \infty$. From Theorem 5.2,

$$\dim_B \Gamma f \le 1 + \log_N \gamma_0 \le \underline{\dim}_B \Gamma f.$$

so that $\dim_B \Gamma f = 1 + \log_N \gamma_0$. Thus (5.4) holds.

From Remark 5.4 and Theorem 5.5, we have the following result.

Corollary 5.6. Assume that the function S_n is nonnegative for each n, and both γ and $\sum_{n=1}^{N} q_n$ are constant functions on I. Then in the case that $\gamma(0) > 1$ and f is not a constant function, dim_B $\Gamma f = 1 + \log_N \gamma(0)$, otherwise dim_B $\Gamma f = 1$.

6. An example and further remarks

6.1. An example: generalized Weierstrass-type functions. Weierstrass functions are classical fractal functions. There are many works on fractal dimensions of their graphs, including the box and Hausdorff dimension. Please see [16, 19, 22] and the references therein. For example, Ren and Shen [22] studied the following Weierstrass-type functions

$$g^{\phi}_{\lambda,N}(x) = \sum_{k=0}^{\infty} \lambda^k \phi(N^k x), \quad x \in \mathbb{R},$$

where $N \geq 2$ is an integer, $1/N < \lambda < 1$ and $\phi : \mathbb{R} \to \mathbb{R}$ is a \mathbb{Z} -periodic real analytic function. They proved that either such a function is real analytic, or the Hausdorff dimension of its graph is equal to $2 + \log_N \lambda$.

It is well known that $f = g_{\lambda,N}^{\phi}|_{[0,1]}$ is a generalized affine FIF. In fact, for $n \in \{1, 2, \ldots, N\}$ and $x \in [0, 1]$, we have

$$f\left(\frac{x+n-1}{N}\right) = \phi\left(\frac{x+n-1}{N}\right) + \lambda \sum_{k=0}^{\infty} \lambda^k \phi(N^k x) = \phi\left(\frac{x+n-1}{N}\right) + \lambda f(x).$$

Thus, $\Gamma f = \bigcup_{n=1}^{N} W_n(\Gamma f)$, where for $n = 1, 2, \dots, N$,

$$W_n(x,y) = \left(\frac{x+n-1}{N}, \lambda y + \phi\left(\frac{x+n-1}{N}\right)\right), \quad (x,y) \in [0,1] \times \mathbb{R}.$$

Let $\phi(x) = \cos(2\pi x)$. Then $g^{\phi}_{\lambda,N}$ is the classical Weierstrass function. Shen [24] proved that the Hausdorff dimension of its graph is equal to $2 + \log_N \lambda$. Let $q_n(x) =$ $\cos(2\pi(x+n-1)/N), 1 \le n \le N$. It is easy to check that $\sum_{n=1}^{N} q_n(x) = 0$ for all $x \in [0,1]$. Thus, from Corollary 5.6, we obtain the well known result dim_B $\Gamma f =$ $2 + \log_N \lambda$, where $f = g^{\phi}_{\lambda,N}|_{[0,1]}$ and $\phi(x) = \cos(2\pi x)$.

By Theorem 2.2, we can study the box dimension of generalized Weierstrass-type functions by replacing vertical scaling factor λ with vertical scaling functions.



FIGURE 1. The FIF in Example 6.1

Example 6.1. Let I = [0, 1], N = 3, and $x_n = n/3$, n = 0, 1, 2, 3. Let vertical scaling functions S_n , $1 \le n \le 3$ on [0, 1] are defined by

$$S_1(x) = S_2(x) = \frac{1}{2} + \frac{\sin(2\pi x)}{4}, \qquad S_3(x) = \frac{1}{2} - \frac{\sin(2\pi x)}{4}$$

Then each function S_n is positive on I so that $\rho_* = \rho^*$.

Let $\phi(x) = \cos(2\pi x)$ and define maps W_n , $1 \le n \le 3$ by

$$W_n(x,y) = \left(\frac{x+n-1}{3}, S_n(x)y + \phi\left(\frac{x+n-1}{3}\right)\right), \quad (x,y) \in [0,1] \times \mathbb{R}.$$

Let $x_n = n/3$ for $0 \le n \le 3$. Let $y_0 = y_2 = 2$ and $y_1 = y_3 = 1/2$. Then it is easy to check that

$$W_n(x_0, y_0) = (x_{n-1}, y_{n-1}), \quad W_n(x_3, y_3) = (x_n, y_n)$$

for n = 1, 2, 3. Thus $\{W_n\}_{n=1}^3$ determines a generalized affine FIF f. Please see Figure 1 for the graph of f.

Notice that $\gamma(x) = \sum_{n=1}^{3} |S_n(x)| = 3/2 + \sin(2\pi x)/4$ for $x \in [0, 1]$. Hence, $\gamma_* = 5/4$, $\gamma^* = 7/4$ and $\lambda' = \pi/2$ is a Lipschitz constant of $\gamma(x)$.

Let $q_n(x) = \phi((x+n-1)/3), x \in [0,1], n = 1, 2, 3$. Then $\sum_{n=1}^{3} q_n(x) = 0$ for all

 $x \in [0,1] \text{ so that } \operatorname{Var}(\sum_{n=1}^{3} q_n, I) = 0.$ Now we calculate $M_f = \max\{|f(x)| : x \in I\}$. Notice that for any $x \in I$, there exists $n_1 n_2 \cdots \in \{1,2,3\}^{\infty}$ such that $x \in \bigcap_{k=1}^{\infty} L_{n_1} \circ L_{n_2} \circ \cdots \circ L_{n_k}(I)$. Thus from

(4.1), we have

$$f(x) = q_{n_1}(L_{n_1}^{-1}(x)) + S_{n_1}(L_{n_1}^{-1}(x))f(L_{n_1}^{-1}(x))$$

= $q_{n_1}(L_{n_1}^{-1}(x)) + \sum_{k=2}^{\infty} \left(\prod_{t=1}^{k-1} S_{n_t}(L_{n_t}^{-1} \circ \dots \circ L_{n_1}^{-1}(x))\right) q_{n_k}(L_{n_k}^{-1} \circ \dots \circ L_{n_1}^{-1}(x)).$

Hence, from $q^* := \max\{|q_n(x)| : x \in [0,1], n = 1,2,3\} = 1$ and

$$S^* := \max\{S_n(x) : x \in [0,1], n = 1,2,3\} = \frac{3}{4},$$

we have $M_f \le q^* \sum_{k=0}^{\infty} (S^*)^k = q^*/(1-S^*) = 4$. Thus,

$$\frac{\lambda' M_f |I| + \operatorname{Var}(\sum_{n=1}^3 q_n, I)}{\gamma_* - 1} \le \frac{(\pi/2) \times 4 \times 1 + 0}{5/4 - 1} = 8\pi.$$

By calculation, $O_6(f, I) > 8\pi$. Thus, from Remark 5.4, $Var(f, I) = \infty$. By definition of vertical scaling matrices, we have

$$\overline{M}_1 = \begin{pmatrix} \frac{3}{4} & \frac{1}{2} + \frac{\sqrt{3}}{8} & \frac{1}{2} \\ \frac{3}{4} & \frac{1}{2} + \frac{\sqrt{3}}{8} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} + \frac{\sqrt{3}}{8} & \frac{3}{4} \end{pmatrix}, \quad \underline{M}_1 = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} - \frac{\sqrt{3}}{8} & \frac{1}{4} \\ \frac{1}{2} & \frac{1}{2} - \frac{\sqrt{3}}{8} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{2} - \frac{\sqrt{3}}{8} & \frac{1}{2} \end{pmatrix}.$$

In general, by calculation, we can obtain the spectral radii of vertical scaling matrices $\rho(\overline{M}_k)$ and $\rho(\underline{M}_k)$, k = 1, 2, 4, 5, 7, 8 as in Tabel 1. Thus, from Theorem 2.2,

$$\dim_B \Gamma f = 1 + \log_N \rho_{\mathbf{S}} \approx 1 + \log 1.516 / \log 3 \approx 1.379.$$

k	1	2	4	5	7	8
$\rho(\overline{M}_k)$	1.95688	1.68984	1.53627	1.52277	1.51675	1.51625
$\rho(\underline{M}_k)$	1.05567	1.33590	1.49577	1.50926	1.51525	1.51575

TABLE 1. $\rho(\overline{M}_k)$ and $\rho(\underline{M}_k)$ in Example 6.1

6.2. Further remarks. From the proof of Theorem 3.3, we essentially prove that $\rho^* = \lim_{k\to\infty} \rho(\overline{M}_k)$ exists without any restrictions on vertical scaling functions. This also holds for the existence of $\rho_* = \lim_{k\to\infty} \rho(\underline{M}_k)$. Hence, from Theorem 2.2, we have the following conjecture.

Conjecture 6.1. Let f be a generalized affine FIF satisfying conditions (A1)-(A3). Then $\rho_* = \rho^*$. Furthermore, in the case that $\operatorname{Var}(f, I) = \infty$ and $\rho_{\mathbf{S}} > 1$, $\dim_B \Gamma f = 1 + \log_N \rho_{\mathbf{S}}$, otherwise $\dim_B \Gamma f = 1$.

References

- M. N. AKHTAR, M. G. P. PRASAD AND M. A. NAVASCUÉS, Box dimension of α-fractal function with variable scaling factors in subintervals, Chaos Solitons Fractals, 103 (2017), 440– 449.
- P. ALLAART The pointwise Hölder spectrum of general self-affine functions on an interval, J. Math. Anal. Appl., 488 (2020), Paper No. 124096, 35 pp.
- B. BÁRÁNY, M. RAMS AND K. SIMON, Dimension of the repeller for a piecewise expanding affine map. Ann. Acad. Sci. Fenn. Math., 45 (2020), 1135–1169.

- 4. M. F. BARNSLEY, Fractal functions and interpolation, Constr. Approx., 2 (1986), 303–329.
- M. F. BARNSLEY, J. ELTON, D. HARDIN AND P. MASSOPUST, Hidden variable fractal interpolation functions, SIAM J. Math. Anal., 20 (1989), 1218–1242.
- M. F. BARNSLEY AND A. N. HARRINGTON, The calculus of fractal interpolation functions, J. Approx. Theory, 57 (1989), 14–34.
- M. F. BARNSLEY AND P. R. MASSOPUST, Bilinear fractal interpolation and box dimension, J. Approx. Theory, 192 (2015), 362–378.
- T. BEDFORD, Hölder exponents and box dimension for self-affine fractal functions, Constr. Approx., 5 (1989), 33–48.
- 9. T. BEDFORD, The box dimension of self-affine graphs and repellers, Nonlinearity, 2 (1989), 53–71.
- A. K. B. CHAND, S. K. KATIYAR, AND P. V. VISWANATHAN, Approximation using hidden variable fractal interpolation function, J. Fractal Geom., 2 (2015), 81–114.
- S. DUBUC, Non-differentiability and Hölder properties of self-affine functions, Expo. Math., 36 (2018), 119–142.
- 12. K. J. FALCONER, Fractal geometry: Mathematical foundation and applications (Third edition), John Wiley & Sons, 2014.
- Z. FENG, Variation and Minkowski dimension of fractal interpolation surfaces, J. Math. Anal. Appl., 345 (2008), 322–334.
- D. P. HARDIN AND P. R. MASSOPUST, The capacity for a class of fractal functions, Comm. Math. Phys., 105 (1986), 455–460.
- 15. R. A. HORN AND C. R. JOHNSON, *Matrix analysis (Second edition)*, Cambridge University Press, 2013.
- T. Y. HU AND K.-S. LAU, Fractal dimensions and singularities of the Weierstrass type functions, Trans. Amer. Math. Soc., 335 (1993), 649–665.
- 17. S. JHA AND S. VERMA, Dimensional analysis of α -fractal functions, Results Math., **76** (2021), Paper No. 186, 24 pp.
- L. JIANG AND H.-J. RUAN, Box dimension of generalized affine fractal interpolation functions, J. Fractal Geom., to appear. (DOI 10.4171/JFG/136)
- 19. J. L. KAPLAN, J. MALLET-PARET AND J. A. YORKE, The Lyapunov dimension of a nowhere differentiable attracting torus, Ergodic Theory Dynam. Systems, 4 (1984), 261–281.
- Q.-G. KONG, H.-J. RUAN AND S. ZHANG, Box dimension of bilinear fractal interpolation surfaces, Bull. Aust. Math. Soc., 98 (2018), 113–121.
- 21. D. S. MAZEL AND M. H. HAYES, Using iterated function systems to model discrete sequences, IEEE Trans. Signal Process, 40 (1992), 1724–1734.
- H. REN AND W. SHEN, A dichotomy for the Weierstrass-type functions, Invent. Math., 226 (2021), 1057–1100.
- H.-J. RUAN, W.-Y. SU AND K. YAO, Box dimension and fractional integral of linear fractal interpolation functions, J. Approx. Theory, 161 (2009), 187–197.
- 24. W. SHEN, Hausdorff dimension of the graphs of the classical Weierstrass functions, Math. Z., **289** (2018), 223–266.
- 25. J. L. VÉHEL, K. DAOUDI AND E. LUTTON, Fractal modeling of speech signals, Fractals, 2 (1994), 379–382.
- H. Y. WANG AND J. S. YU, Fractal interpolation functions with variable parameters and their analytical properties, J. Approx. Theory, 175 (2013), 1–18.

SCHOOL OF MATHEMATICAL SCIENCES, ZHEJIANG UNIVERSITY, HANGZHOU 310058, CHINA *Email address*: jianglai@zju.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, ZHEJIANG UNIVERSITY, HANGZHOU 310058, CHINA *Email address:* ruanhj@zju.edu.cn