

# Infinite chains in the tree of numerical semigroups

Mariana Rosas-Ribeiro<sup>1</sup> and Maria Bras-Amorós<sup>2</sup>

<sup>1</sup>Universitat Rovira i Virgili, mariana.rosas@urv.cat

<sup>2</sup>Universitat Rovira i Virgili, maria.bras@urv.cat

August 2023

## Abstract

One major problem in the study of numerical semigroups is determining the growth of the semigroup tree. In the present work, infinite chains of numerical semigroups in the semigroup tree, firstly introduced in [4], are studied. Computational results show that these chains are rare, but without them the tree would not be infinite. It is proved that for each genus  $g \geq 5$  there are more semigroups of that genus not belonging to infinite chains than semigroups belonging. The reference [4] presented a characterization of the semigroups that belong to infinite chains in terms of the coprimality of the left elements of the semigroup as well as a result on the cardinality of the set of infinite chains to which a numerical semigroup belongs in terms of the primality of the greatest common divisor of these left elements. We revisit these results and fix an imprecision on the cardinality of the set of infinite chains to which a semigroup belongs in the case when the greatest common divisor of the left elements is a prime number. We then look at infinite chains in subtrees with fixed multiplicity. When the multiplicity is a prime number there is only one infinite chain in the tree of semigroups with such multiplicity. When the multiplicity is 4 or 6 we prove a self-replication behavior in the subtree and prove a formula for the number of semigroups in infinite chains of a given genus and multiplicity 4 and 6, respectively.

## 1 Introduction

A numerical semigroup  $\Lambda$  is a subset of the non-negative integers  $\mathbb{N}_0$ , containing 0, closed under addition and with finite complement in  $\mathbb{N}_0$ . The elements of  $\mathbb{N}_0$  that are not in  $\Lambda$  are the *gaps* and the number of gaps is the *genus*  $g(\Lambda)$  of the numerical semigroup. Intuitively, the possibilities of obtaining numerical semigroups with  $g$  gaps seem to increase as the number  $g$  increases and indeed computational results have shown this ([2], [5]). It was conjectured in [2], 2008, that the sequence  $n_g$  formed by the number of numerical semigroups of genus  $g$ , not only is increasing but also grows like the Fibonacci sequence. In 2012 it was proved in [7] one such conjecture, namely  $\frac{n_g}{\varphi^g} \xrightarrow{g \rightarrow \infty} S$ , where  $\varphi$  is the golden

ratio and  $S$  is at least 3.78. The conjecture that remains unproved is that  $n_{g-2} + n_{g-1} \leq n_g$  and the weaker one that  $n_{g-1} \leq n_g$ , already announced in 2007 [1].

One way to label all elements of the numerical semigroup  $\Lambda$  is to enumerate them. The enumeration  $\lambda$  of a numerical semigroup  $\Lambda$  is the unique increasing bijection between  $\mathbb{N}_0$  and  $\Lambda$ , and we denote by  $\lambda_i$  the image of  $i \in \mathbb{N}_0$ . The *multiplicity* of  $\Lambda$  is  $m(\Lambda) = \lambda_1$ . Obviously,  $m(\Lambda) \leq g(\Lambda) + 1$ . The *conductor* of  $\Lambda$  is  $c(\Lambda) = \lambda_k$ , where  $\lambda_k + n \in \Lambda$  for all  $n \in \mathbb{N}$  and  $\lambda_k - 1 \notin \Lambda$ . The number  $c(\Lambda) - 1$  is called the *Frobenius number* of  $\Lambda$  and denoted by  $F(\Lambda)$ . The elements of  $\Lambda$  that are smaller than its conductor are the *left elements*, denoted by  $\mathcal{L}(\Lambda)$ .

Let  $[a, \infty)$  stand for  $\{i \in \mathbb{N}_0 : i \geq a\}$ . Numerical semigroups of the form  $\{0\} \cup [\lambda_1, \infty)$  are called *ordinary* and those of the form  $\{2n : n \geq 0\} \cup [\lambda_k, \infty)$ ,  $\lambda_k$  even, are called *hyperelliptic semigroups*.

It is well known that each numerical semigroup has a unique minimal system of generators. The cardinality of the minimal system of generators is called the *embedding dimension* of  $\Lambda$  and we denote it by  $e(\Lambda)$ . Clearly,  $e(\Lambda) \leq \lambda_1$ . The minimal generators of  $\Lambda$  that are not in  $\mathcal{L}(\Lambda)$  are the *effective generators* or *right generators* of  $\Lambda$  and with them we can organize the collection of all numerical semigroups in a tree rooted in  $\mathbb{N}_0$ .

That tree is constructed by removing one by one the effective generators from each numerical semigroup. In this way, the level of the tree is equal to the genus of the numerical semigroups that are at that level (assume that the root has level zero). For more details of this tree, we suggest consulting [3], [7] and [5]. From the tree perspective, numerical semigroups can also be seen as nodes. In this way the *children* of a numerical semigroup  $\Lambda$  are those numerical semigroups that are obtained by taking away one right generator from  $\Lambda$ . Conversely, the *parent* of a numerical semigroup is obtained by adding its Frobenius number (the parent is one level lower in the tree). The number of children of a node  $\Lambda$  is called *efficacy* and is denoted by  $h(\Lambda)$ . By the previous remarks, one can deduce that  $h(\Lambda) \leq g(\Lambda) + 1$ . As can be seen in the references cited above, this tree has been used to study the growth of the  $n_g$  sequence.

Every numerical semigroup of genus larger than  $g(\Lambda)$  connected to  $\Lambda$  through edges in the tree is said to be a *descendant* of  $\Lambda$ . A node that has no descendants is said to be a *leaf* and a node that has exactly one child is said to be a *stick*. An infinite set of numerical semigroups is said to be an *infinite chain* if the semigroups in the set can be enumerated in such a way that each one is the parent of the next one. A numerical semigroup is in an infinite chain if and only if it has infinitely many descendants.

The set of ordinary semigroups and the set of hyperelliptic semigroups each form an infinite chain and, as we will see throughout this paper, these two chains are well positioned in the numerical semigroup tree.

Infinite chains of numerical semigroups were first investigated in [4]. This reference presented a characterization of the semigroups that belong to infinite chains in terms of the coprimality of the left elements of the semigroup as well as a result on the cardinality of the set of infinite chains to which a numerical semigroup belongs in terms of the primality of the greatest common divisor of these left elements.

In Section 2 we state some preliminary results and in Section 3 we revisit the main result of [4] and fix an imprecision on the cardinality of the set of infinite chains to which

a semigroup belongs in the case when the greatest common divisor of the left elements is a prime number. The resulting fixed theorem is Theorem 3.4.

Computational results show that infinite chains are rare, but without them the tree would not be infinite. For this reason, we direct the study on infinite chains in order to investigate the broader problem of the tree growth. In Section 4 we prove that for each genus  $g \geq 5$  there are more semigroups of that genus not belonging to infinite chains than semigroups belonging (Theorem 4.10).

Another aspect that is evident in the tree is that semigroups of the same multiplicity form subtrees. In Section 5 we look at infinite chains in subtrees with fixed multiplicity. When the multiplicity is a prime number, one can prove, as a consequence of Theorem 3.4, that there is only one infinite chain in the tree of semigroups with such multiplicity. For the cases of multiplicity 4 and multiplicity 6, we prove a self-replication behavior in the subtree formed by the semigroups that are in infinite chains with fixed multiplicity equal to 4 (Subsection 5.2) and a different self-replication behavior in the subtree formed by the semigroups that are in infinite chains with fixed multiplicity equal to 6 (Subsection 5.3). This enables us to prove a formula for the amount  $i_g$  of semigroups of genus  $g$  that belong to infinite chains, of any given prime multiplicity (Theorem 5.3), of multiplicity 4 (Theorem 5.4), or of multiplicity 6 (Theorem 5.10), respectively.

## 2 Preliminary results

The next lemma was proved in [3] but we enunciate it here in a different way, underlying the result used in the ordinary proof. Among other things, it allows to study the children of the children, or grandchildren, of most numerical semigroups.

**Lemma 2.1** ([4], Lemma 1). If  $\lambda_{i_1} < \lambda_{i_2} < \dots < \lambda_{i_n}$  are the effective generators of a non-ordinary numerical semigroup  $\Lambda$  then the effective generators of  $\Lambda \setminus \{\lambda_{i_j}\}$  are either  $\lambda_{i_{j+1}}, \dots, \lambda_{i_n}$  or  $\lambda_{i_{j+1}}, \dots, \lambda_{i_n}, \lambda_{i_j} + \lambda_1$ .

Next proposition gives a bound for the maximum effective generator.

**Proposition 2.2.** Every effective generator is at most the conductor plus the multiplicity minus one.

*Proof.* Indeed, for a numerical semigroup  $\Lambda$ , if  $a \geq c(\Lambda) + m(\Lambda)$ , then  $a = m(\Lambda) + (a - m(\Lambda))$ , with both  $m(\Lambda)$  and  $a - m(\Lambda)$  nongaps. □

The last preliminary result is a characterization of ordinary semigroups by the number of children in terms of the genus.

**Lemma 2.3.** If a numerical semigroup  $\Lambda$  with genus  $g$  has  $g+1$  children, then  $\Lambda$  is ordinary.

*Proof.* The following sequence of inequalities must indeed be a sequence of equalities:

$$g + 1 = h(\Lambda) \leq e(\Lambda) \leq m(\Lambda) \leq g + 1.$$

Hence,  $m(\Lambda) = g + 1$ , meaning that all gaps of  $\Lambda$  are smaller than  $m(\Lambda)$ , and, hence,  $\Lambda$  must be ordinary. □

### 3 Infinite chains

In this section we are interested in studying the nodes of the tree of numerical semigroups that have infinitely many descendants. An infinite chain is a sequence of semigroups  $\Lambda_0, \Lambda_1, \Lambda_2, \dots$  such that  $\Lambda_i$  is the parent of  $\Lambda_{i+1}$  in the semigroup tree. Notice that a numerical semigroup belongs to an infinite chain if and only if it has infinitely many descendants.

First, we analyze which nodes in the semigroup tree have an infinite number of descendants. For the nodes having a finite number of descendants we give a way to determine the descendant at largest distance; for the nodes having an infinite number of descendants we determine the number of infinite chains in which the semigroup lies. It turns out here that primality and coprimality of integers appear in the scene as discriminating factors.

Let  $\mathbb{S}$  be the set of all numerical semigroups, and let  $\mathbb{I}$  be the set of all infinite chains. One element of  $\mathbb{I}$  contains infinitely many elements in  $\mathbb{S}$ . As an example, the sequence of ordinary semigroups,  $\mathcal{O}_g = \{0, g + 1, \rightarrow\}$  for any  $g \in \mathbb{N}_0$ , is in  $\mathbb{I}$ . We denote it  $l_{\mathcal{O}}$ . That is,

$$l_{\mathcal{O}} = \{\mathcal{O}_g : g \geq 0\}.$$

Another example is the sequence of hyperelliptic semigroups,  $\mathcal{H}_g = \{0, 2, 4, \dots, 2g, \rightarrow\}$ . We denote it  $l_{\mathcal{H}}$ . That is,

$$l_{\mathcal{H}} = \{\mathcal{H}_g : g \geq 0\}.$$

Each element in  $\mathbb{S}$  may be contained in none, one, several, or infinitely many elements of  $\mathbb{I}$ . The aim of this section is analyzing what circumstances give what of these cases. Then, we analyze what kind of generators appear in infinite chains.

For the proof of the next lemma we use that the integers  $l_1, \dots, l_m$  generate a numerical semigroup if and only if they are coprime.

**Lemma 3.1.** Given an infinite chain  $l = (\Lambda_i)_{i \geq 0}$  different than  $l_{\mathcal{O}}$ , it holds that

$$\bigcap_{i \geq 0} \Lambda_i = d \cdot \Lambda$$

for some integer  $d > 1$  and some numerical semigroup  $\Lambda$ .

*Proof.* The intersection  $\bigcap_{i \geq 0} \Lambda_i$  satisfies  $0 \in \bigcap_{i \geq 0} \Lambda_i$ . Since there exists a non-ordinary semigroup in  $l$ , there must be a non-zero element  $x \in \bigcap_{i \geq 0} \Lambda_i$  and  $x + y \in \bigcap_{i \geq 0} \Lambda_i$  for all  $x, y \in \bigcap_{i \geq 0} \Lambda_i$ .

Furthermore, all elements in  $\bigcap_{i \geq 0} \Lambda_i$  must be divisible by an integer  $d > 1$ . Indeed, otherwise we could find a finite set of coprime elements which would generate a numerical semigroup, and this numerical semigroup should be a subset of  $\bigcap_{i \geq 0} \Lambda_i$ . Then the infinite chain would not contain any semigroup with genus larger than that of this semigroup, giving a contradiction. Let  $d$  be the greatest of the common divisors of  $\bigcap_{i \geq 0} \Lambda_i$ . Then  $\frac{1}{d}(\bigcap_{i \geq 0} \Lambda_i)$  must be a numerical semigroup. □

**Lemma 3.2.** Given an integer  $d > 1$  and a numerical semigroup  $\Lambda$ , the infinite chain obtained by deleting repetitions in the sequence  $\Lambda_j = d \cdot \Lambda \cup \{l \in \mathbb{N} : l \geq j\}$  has intersection  $d \cdot \Lambda$ .

Lemma 3.1 and its proof suggest the map

$$\begin{aligned} \sigma : \mathbb{I} \setminus I_{\mathcal{O}} &\longrightarrow \mathbb{N}_{\geq 2} \times \mathbb{S} \\ \{\Lambda_i\}_{i \in \mathbb{N}_0} &\mapsto (\gcd(\cap_{i \geq 0} \Lambda_i), \cap_{i \geq 0} \Lambda_i / \gcd(\cap_{i \geq 0} \Lambda_i)). \end{aligned}$$

Lemma 3.2 proves that the map

$$\begin{aligned} \omega : \mathbb{N}_{\geq 2} \times \mathbb{S} &\longrightarrow \mathbb{I} \setminus I_{\mathcal{O}} \\ (d, \Lambda) &\mapsto \{d \cdot \Lambda \cup \{l \in \mathbb{N} : l \geq i\}\}_{i \notin d \cdot \Lambda} \end{aligned}$$

is the inverse of  $\sigma$ .

Consequently,  $\mathbb{I} \setminus I_{\mathcal{O}}$  and  $\mathbb{N}_{\geq 2} \times \mathbb{S}$  are in a one-to-one correspondence. For example, in this correspondence, the image of  $I_{\mathcal{H}}$  would be  $(2, \mathbb{N}_0)$ .

**Remark 3.3.** Any ordinary numerical semigroup belongs to infinitely many infinite chains. Indeed,  $\{0, m, \rightarrow\}$  belongs, among others to  $\omega(m', \mathbb{N}_0)$  for any  $m' \geq m$ .

In the next lemma we show that the greatest common divisor of the first elements of a non-ordinary numerical semigroup determine whether the numerical semigroup has infinite number of descendants. For that we define a descendant of a numerical semigroup *beyond* a given nongap of the semigroup as a descendant that contains all nongaps up to the given nongap. For example,  $\{0, 4, 5, 8, 9, 10, 12, \rightarrow\}$  and  $\{0, 4, 5, 6, 8, \rightarrow\}$  are descendants of  $\{0, 4, \rightarrow\}$ , but only  $\{0, 4, 5, 6, 8, \rightarrow\}$  is a descendant beyond 6.

We use the fact that a finite number of coprime elements generate a numerical semigroup.

**Theorem 3.4.** Let  $\Lambda$  be a non-ordinary numerical semigroup with enumeration  $\lambda$ , genus  $g$ , and conductor  $c$ , and let  $d$  be the greatest common divisor of  $\lambda_1, \dots, \lambda_{c-g-1}$ . Then,

- $\Lambda$  lies in an infinite chain if and only if  $d \neq 1$ .
- If  $d = 1$ , then the descendant of  $\Lambda$  with largest genus is the numerical semigroup generated by  $\lambda_1, \dots, \lambda_{c-g-1}$ .
- If  $d \neq 1$  and  $d$  is not prime, then  $\Lambda$  lies in infinitely many infinite chains.
- If  $d$  is a prime then the number of infinite chains in which  $\Lambda$  lies is one plus the number of descendants of  $\{0, \frac{\lambda_1}{d}, \dots, \frac{\lambda_{c-g-1}}{d}\} \cup \{l \in \mathbb{N}_0 : l \geq \lceil \frac{c}{d} \rceil\}$  beyond  $\frac{\lambda_{c-g-1}}{d}$ .

*Proof.* a. If  $c - g - 1 = 1$ , then  $\Lambda$  belongs at least to the infinite chain  $\omega(\lambda_1, \mathbb{N}_0)$ , while  $d = \lambda_1 \neq 1$ . So, we can assume that  $c - g - 1 \geq 2$ .

If  $d = 1$  and  $c - g - 1 \geq 2$ , then  $\lambda_1, \dots, \lambda_{c-g-1}$  generate a numerical semigroup  $\Lambda'$  with no descendants in the semigroup tree. Now, each descendant of  $\Lambda$  must contain  $\Lambda'$ . Thus, the maximum of the genera of the descendants of  $\Lambda$  is the genus of  $\Lambda'$ , which is finite and, so, there is no infinite chain containing  $\Lambda$ .

On the other hand, if  $d \neq 1$  and  $c - g - 1 \geq 2$ , then

$$\lambda_0 = d\tilde{\lambda}_0, \lambda_1 = d\tilde{\lambda}_1, \dots, \lambda_{c-g-1} = d\tilde{\lambda}_{c-g-1}$$

with  $\tilde{\lambda}_1, \dots, \tilde{\lambda}_{c-g-1}$  coprime. Let  $\tilde{\Lambda}$  be the numerical semigroup generated by  $\tilde{\lambda}_1, \dots, \tilde{\lambda}_{c-g-1}$ . Then,  $\Lambda$  belongs to the infinite chain  $\omega(d, \tilde{\Lambda})$ .

- b. As we have seen in the previous statement, the semigroup  $\Lambda'$ , which is a descendant itself of  $\Lambda$ , must be contained in each descendant of  $\Lambda$ . And the result follows from the fact that  $\Lambda'$  has no descendants.
- c. If  $d$  is not prime, then  $d = d_1 d_2$  for some  $d_1, d_2 > 1$  and, as before,

$$\lambda_0 = d_1 d_2 \tilde{\lambda}_0, \lambda_1 = d_1 d_2 \tilde{\lambda}_1, \dots, \lambda_{c-g-1} = d_1 d_2 \tilde{\lambda}_{c-g-1}$$

with  $\tilde{\lambda}_1, \dots, \tilde{\lambda}_{c-g-1}$  coprime. Let  $\tilde{\Lambda}$  be  $\mathbb{N}_0$  if  $c-g-1 = 1$  or the numerical semigroup generated by  $\tilde{\lambda}_1, \dots, \tilde{\lambda}_{c-g-1}$ , otherwise. For each  $a \geq 0$  define

$$\Lambda_a = d_2 \tilde{\Lambda} \cup \{l \in \mathbb{N}_0 : l \geq a\}.$$

Then  $\Lambda$  belongs to the infinite chains  $\omega(d_1, \Lambda_a)$  for all  $a$  larger than  $\frac{c}{d_1}$  and not multiple of  $d_2$ , which are all different. So, it belongs to an infinite number of infinite chains.

- d. Suppose that  $d$  is prime and that an infinite chain  $(\Lambda_i)_{i \geq 0}$  contains  $\Lambda$ . The infinite chain must satisfy  $\cap_{i \geq 0} \Lambda_i = d \cdot \tilde{\Lambda}$  for a unique numerical semigroup  $\tilde{\Lambda}$  such that
- $d \tilde{\lambda}_0 = \lambda_0, d \tilde{\lambda}_1 = \lambda_1, \dots, d \tilde{\lambda}_{c-g-1} = \lambda_{c-g-1}$ ,
  - $d \tilde{\lambda}_{c-g} \geq c$ , since  $d \tilde{\Lambda} \subseteq \Lambda$ .

Now, the conductor  $\tilde{c}$  of  $\tilde{\Lambda}$  must be  $\tilde{c} \geq \tilde{\lambda}_{c-g} \geq \lceil \frac{c}{d} \rceil$ . Thus,  $\tilde{\Lambda}$  is either  $\Lambda/d := \{0, \frac{\lambda_1}{d}, \dots, \frac{\lambda_{c-g-1}}{d}\} \cup \{l \in \mathbb{N}_0 : l \geq \lceil \frac{c}{d} \rceil\}$  or one of its descendants beyond  $\frac{\lambda_{c-g-1}}{d}$ .

Conversely, if  $\tilde{\Lambda}$  is either  $\{0, \frac{\lambda_1}{d}, \dots, \frac{\lambda_{c-g-1}}{d}\} \cup \{l \in \mathbb{N}_0 : l \geq \lceil \frac{c}{d} \rceil\}$  or one of its descendants beyond  $\frac{\lambda_{c-g-1}}{d}$ , then  $\Lambda$  belongs to the infinite chain  $\omega(d, \tilde{\Lambda})$ .

It remains to prove that  $\{0, \frac{\lambda_1}{d}, \dots, \frac{\lambda_{c-g-1}}{d}\} \cup \{l \in \mathbb{N}_0 : l \geq \lceil \frac{c}{d} \rceil\}$  has a finite number of descendants beyond  $\frac{\lambda_{c-g-1}}{d}$ . If  $c-g-1 = 1$ , then  $d = \lambda_1 \neq 1$  and  $\frac{\lambda_{c-g-1}}{d} = 1$ , and the semigroup  $\{0, \frac{\lambda_1}{d}\} \cup \{l \in \mathbb{N}_0 : l \geq \lceil \frac{c}{d} \rceil\}$  is  $\mathbb{N}_0$  has no descendants beyond 1. On the other hand, if  $c-g-1 \geq 2$ , then any descendant beyond  $\frac{\lambda_{c-g-1}}{d}$  contains  $\frac{\lambda_1}{d}, \dots, \frac{\lambda_{c-g-1}}{d}$  which is a set of at least two elements which are coprime. Hence, any such descendant contains the numerical semigroup generated by  $\frac{\lambda_1}{d}, \dots, \frac{\lambda_{c-g-1}}{d}$ . Now, the result follows from the fact that the number of semigroups that contain a given numerical semigroup is finite. □

## 4 Minority of semigroups in infinite chains

In this section we prove that for each fixed genus  $g \geq 5$  the majority of semigroups do not belong to any infinite chain.

From the characterization given by Theorem 3.4 of the semigroups that are in infinite chains, we are able to investigate which children of such semigroups remain in infinite chains.

**Corollary 4.1.** If the numerical semigroup  $\Lambda$  lies in an infinite chain, then it has at most two children in infinite chains.

*Proof.* Let  $\Lambda$  be a numerical semigroup belonging to an infinite chain,  $c$  its conductor,  $\mathcal{L}(\Lambda)$  the set of left elements and  $d = \gcd(\mathcal{L}(\Lambda))$ . By Theorem 3.4,  $d \neq 1$  and by definition an effective generator of  $\Lambda$  is  $c + i$  for some  $i \geq 0$ . So every child of  $\Lambda$  is of the type  $\Lambda \setminus \{c + i\}$  for some  $i \geq 0$  and the left elements of this child are either  $\mathcal{L}(\Lambda \setminus \{c\}) = \mathcal{L}(\Lambda)$ , if  $i = 0$ , or  $\mathcal{L}(\Lambda \setminus \{c + i\}) = \mathcal{L}(\Lambda) \cup \{c, \dots, c + i - 1\}$ , if  $i > 0$ . Now, suppose  $i \geq 2$ . This implies  $\mathcal{L}(\Lambda \setminus \{c + i\}) \supseteq \{c, c + 1\}$  and, since  $\gcd(c, c + 1) = 1$ , we have  $\gcd(\mathcal{L}(\Lambda \setminus \{c + i\})) = 1$ . So,  $\Lambda \setminus \{c + i\}$ , for  $i \geq 2$ , does not have infinitely many descendants. □

**Remark 4.2.** The proof above makes it clear that the only children of a semigroup  $\Lambda$  that may have infinitely many descendants are  $\Lambda \setminus \{c(\Lambda)\}$  and  $\Lambda \setminus \{c(\Lambda) + 1\}$ .

**Proposition 4.3.** Except for  $\mathbb{N}_0$  and hyperelliptic semigroups, every numerical semigroup that is in an infinite chain has at least one child that is not.

*Proof.* Let  $\Lambda$  be a numerical semigroup such that  $\gcd(\mathcal{L}(\Lambda)) \neq 1$ . Let us suppose  $\Lambda$  only has children that are in infinite chains. By Corollary 4.1, there are three possible cases:  $c = c(\Lambda)$  is the unique effective generator,  $c + 1$  is the unique effective generator or  $c$  and  $c + 1$  are all the effective generators of  $\Lambda$ .

In the first case, the only child of  $\Lambda$  is  $\Lambda_1 = \Lambda \setminus \{c\}$  and, since  $\Lambda$  is in an infinite chain,  $\Lambda_1$  cannot be a leaf. This implies that  $c$  is a strong generator of  $\Lambda$  and, by Lemma 2.1,  $\Lambda_1$  has a (unique) effective generator which is  $c + m$ , where  $m = m(\Lambda)$ . So, the child of  $\Lambda_1$  is  $\Lambda_2 = \Lambda_1 \setminus \{c + m\} = \Lambda \setminus \{c, c + m\}$ . If  $m > 2$  then, as  $c$  is the conductor of  $\Lambda$ , we have  $\{c + 1, c + 2\} \subset \Lambda \setminus \{c, c + m\} = \Lambda_2$ . However, since  $\Lambda_2$  is the only child of the only child of  $\Lambda$ , it must also be in an infinite chain. That is,  $\gcd(\mathcal{L}(\Lambda_2)) \neq 1$  but  $\gcd(c + 1, c + 2) = 1$ , a contradiction.

Similarly, in the second case if the only effective generator of  $\Lambda$  is  $c + 1$  then all its descendants are sticks, the next ones being  $\Lambda_1 = \Lambda \setminus \{c + 1\}$  and  $\Lambda_2 = \Lambda_1 \setminus \{c + 1 + m\} = \Lambda \setminus \{c + 1, c + 1 + m\}$ . And if  $m > 2$ ,  $\{c + 2, c + 3\} \subset \Lambda_2$ , contradicting  $\gcd(\mathcal{L}(\Lambda_2)) \neq 1$ .

Finally, for the third case we assume that  $c$  and  $c + 1$  are the only effective generators of  $\Lambda$ . So, when we take out  $c + 1$  we are in the same situation as in the second case, where we need  $m \leq 2$ .

Therefore the only possibility for  $\Lambda$  to have all children in infinite chains is if the multiplicity of  $\Lambda$  is equal to 1 or 2, that is, if  $\Lambda$  is  $\mathbb{N}_0$  or a hyperelliptic semigroup. □

A direct conclusion from Proposition 4.3 is that, with the exception of  $\mathbb{N}_0$  and hyperelliptic semigroups, a numerical semigroup that is in an infinite chain has at least two children.

The way the tree of numerical semigroups is constructed generates subtrees of numerical semigroups with the same multiplicity. We will comment more on this in the next section. For now, consider the following result.

**Lemma 4.4.** For every pair  $(g, m)$ ,  $g \geq 1, 2 \leq m \leq g + 1$ , there exists a numerical semigroup  $\Lambda$  with  $g(\Lambda) = g$  that lies in an infinite chain formed by numerical semigroups of multiplicity  $m$ .

*Proof.* Observe that

$$\left\lfloor \frac{g}{m-1} \right\rfloor (m-1) < g+1 < \left( \left\lfloor \frac{g}{m-1} \right\rfloor + 1 \right) (m-1) + 1.$$

Hence,  $\lfloor \frac{g}{m-1} \rfloor m < g+1 + \lfloor \frac{g}{m-1} \rfloor < (\lfloor \frac{g}{m-1} \rfloor + 1)m$ . Now, one can take the numerical semigroup  $\{0, m, 2m, \dots, \lfloor \frac{g}{m-1} \rfloor m, g+1 + \lfloor \frac{g}{m-1} \rfloor, \rightarrow\}$ . It belongs to the infinite chain formed by the semigroups  $\{m\mathbb{N} \cup \{0, c, \rightarrow\} : c \in \mathbb{N}\}$  and its genus is clearly  $g$ . □

**Lemma 4.5.** Let  $\Lambda$  be a numerical semigroup with enumeration  $\lambda$ , genus  $g$ , conductor  $c$ , and  $\gcd(\mathcal{L}(\Lambda)) = d > 1$ . Then the number of children of  $\Lambda$  is at least  $d - 1$  and if  $\lambda_1 = d$  then the number of children of  $\Lambda$  is exactly  $d - 1$ .

*Proof.* Since  $d$  is a divisor of the left elements of  $\Lambda$ , we have  $d \leq \lambda_1$ , thus,  $d - 1 < \lambda_1$ . Since  $d \neq 1$ , the set  $\{c, c+1, \dots, c+d-1\}$  has more than one element and it has exactly one multiple of  $d$ . A non-multiple of  $d$  of the form  $c+i$  with  $0 \leq i \leq d-1$  can not be generated by  $\{\lambda_1, \dots, \lambda_{c-g-1}\} \cup \{c, c+1, \dots, c+i-1\}$ . Indeed, let us call  $\mathcal{R} = \{c, c+1, \dots, c+i-1\}$ . Suppose that  $c+i = \lambda_{j_1} + \dots + \lambda_{j_k}$ . But none of the  $\lambda_{j_s}$  is in  $\mathcal{R}$  since, otherwise, if  $\lambda_{j_s} \in \mathcal{R}$ , then  $\lambda_{j_s} \geq c$  and, so,  $c+i - \lambda_{j_s} \leq i \leq d-1 < \lambda_1$ , a contradiction. Therefore, every numerical semigroup that is in an infinite chain has at least  $d - 1$  children. Moreover, if  $\lambda_1 = d$ , then, by Proposition 2.2,  $\Lambda$  has exactly  $d - 1$  effective generators. □

**Definition 4.6.** A numerical semigroup is *fertile* if most of its children are in infinite chains.

**Proposition 4.7.** A numerical semigroup  $\Lambda$  is fertile if and only if one of the options below holds

- a.  $\Lambda = \mathbb{N}_0$ ;
- b.  $\Lambda$  is a hyperelliptic semigroup;
- c.  $h(\Lambda) = 3$  and two of its children are in infinite chains.

*Proof.* Let  $\Lambda$  be a fertile numerical semigroup. By Corollary 4.1, if  $h(\Lambda) > 3$  then it has at most 2 children in infinite chains, so it can not be fertile. If  $h(\Lambda) = 3$ , then it is fertile if it has 2 children in infinite chains, which corresponds to the case c. If  $h(\Lambda) = 2$ , then  $\Lambda$  is fertile if both its children are in infinite chains and, by Proposition 4.3, that occurs when  $\Lambda = \mathbb{N}_0 \setminus \{1\}$ . Finally, if  $h(\Lambda) = 1$  and its only child is in an infinite chain, then  $\Lambda$  is a hyperelliptic semigroup or  $\mathbb{N}_0$ . □



There is one hyperelliptic semigroup at each level and we would like to explore what is the frequency of numerical semigroups of type  $c$  at each level of the semigroup tree. Consider the semigroup sequence defined by  $M_n := \{4k : k \geq 0\} \cup [4n + 2, \infty)$ .

Note that, if  $\Lambda = M_n$  for some  $n \geq 1$ , then  $c(\Lambda) = 4n + 2$  is a minimal generator of  $\Lambda$ ,  $\gcd(\mathcal{L}(\Lambda)) = \gcd(0, 4, \dots, 4n) = 4$  and  $\gcd(\mathcal{L}(\Lambda \setminus \{c(\Lambda) + 1\})) = \gcd(\mathcal{L}(\Lambda), 4n + 2) = 2$ . So, by Theorem 3.4,  $\Lambda \setminus \{c\}$  and  $\Lambda \setminus \{c + 1\}$  are in infinite chains. In addition,  $c(\Lambda) + 3 = 4n + 5$  is also an effective generator and, by Proposition 2.2, it is the last of them. Therefore,  $\Lambda = M_n$  is of type  $c$ .

**Proposition 4.8.** A numerical semigroup  $\Lambda$  of genus  $g > 2$  is of type  $c$  if and only if  $\Lambda = M_n$  for some  $n \geq 1$ , where  $M_n = \{4k : k \geq 0\} \cup [4n + 2, \infty)$ .

*Proof.* It is obvious that the only ordinary numerical semigroup which is of type  $c$  has genus equal to 2 since the number of effective generators of the ordinary semigroup of genus  $g$  is  $g + 1$ , while type  $c$  semigroups have efficacy equal to 3. Let  $\Lambda$  be a numerical semigroup of type  $c$ , and let  $g = g(\Lambda) > 2$ ,  $c = c(\Lambda)$  be its genus and conductor and let  $\lambda$  be its enumeration. Thus,  $\gcd(\lambda_0, \dots, \lambda_{c-g-1}) = d > 1$  and, since  $h(\Lambda) = 3$ , by Lemma 4.5, we have  $d \leq 4$ . Moreover, by Remark 4.2,  $c$  and  $c + 1$  are generators of  $\Lambda$ , with  $\gcd(\lambda_0, \dots, \lambda_{c-g-1}, \lambda_{c-g}) = d' > 1$ .

If  $d = 2$ , then  $d' = 2$ ,  $c$  is even, and  $\lambda_1 \neq 2$  by Lemma 4.5. In this case, all nongaps before  $c + 1$  are even, so if  $c + 3$  is not a generator,  $2 \in \Lambda$ , which is a contradiction. Thus  $c + 3$  is the largest effective generator and the only possibility to write  $c + 5$  as the sum of nongaps is  $c + 5 = (c + 1) + 4$ . Then,  $4 \in \Lambda$  and  $c$  must be congruent with 2 modulo 4, in order to be a generator. Now, since  $d \neq 4$ , there must exist  $\lambda_i \in \mathcal{L}(\Lambda)$  such that  $4 \nmid \lambda_i$ . That is,  $\lambda_i$  is congruent with 2 modulo 4. But then  $c - \lambda_i$  is a multiple of 4 and so, it belongs to  $\Lambda$ , contradicting the fact that  $c$  is a generator.

If  $d = 3$ , similarly we have  $3 \mid c$  and so  $3 \nmid c + 2$ . Thus,  $c + 2$  is not equal to the sum of any elements of  $\Lambda$  smaller than  $c$ , so  $c + 2$  is the third effective generator. Consequently  $c, c + 1, c + 2$  are the three effective generators of  $\Lambda$  and, since  $c + 4$  is not multiple of 3 and it is not a generator, writing  $c + 4$  as the sum of nongaps we get  $c + 4 = (c + 1) + 3$  or  $c + 4 = (c + 2) + 2$ . So, either  $2 \in \Lambda$  or  $3 \in \Lambda$  but  $2 \in \Lambda$  contradicts  $d = 3$  and  $3 \in \Lambda$  contradicts  $c$  being a generator.

If  $d = 4$  and  $m > 4$ , then  $c$  is even because  $d' \mid 4$  and  $d' \neq 1$ . Thus,  $c + 3$  and  $c + 5$  are odd and, so, they are generators of  $\Lambda$ , contradicting  $h(\Lambda) = 3$ .

If  $d = 4$  and  $m = 4$ , then all left elements are multiples of 4, and  $c$  needs to be even and not multiple of 4 since we have  $d' \neq 1$  and  $c$  a generator. This description corresponds exactly to the semigroups of the sequence  $M_n$ . □

It is easy to see that  $g(M_n) = 3n + 1$ . Thus we obtain the following result.

**Theorem 4.9.** The unique fertile semigroups of genus  $g$  are:

- a.  $\mathbb{N}_0$  if  $g = 0$ .
- b.  $\mathbb{N}_0 \setminus \{1\}$  if  $g = 1$ .

- c.  $H_g$  if  $g > 1$  and  $g \not\equiv 1 \pmod{3}$ .
- d.  $H_g$  and  $M_{\frac{g-1}{3}}$  if  $g > 1$  and  $g \equiv 1 \pmod{3}$ .

*Proof.* Direct from Propositions 4.7 and 4.8. □

Let  $i_g$  be the number of numerical semigroups of genus  $g$  that are in an infinite chain.

**Theorem 4.10.** For  $g \geq 5$  we have  $i_g \leq \frac{1}{2}n_g$ .

*Proof.* Computations show that  $i_5 = 6$  and  $n_5 = 12$ . Fixed  $g \geq 6$ , let us prove that there are more numerical semigroups with genus  $g$  that are not in infinite chains than the other way around. To do so, we partition the set of numerical semigroups of genus  $g$  into three disjoint sets:

$$\begin{aligned} F_g &:= \{\Lambda : g(\Lambda) = g \text{ and } \Lambda \cup \{F(\Lambda)\} \text{ is fertile}\} \\ O_g &:= \{\Lambda : g(\Lambda) = g \text{ and } \Lambda \cup \{F(\Lambda)\} \text{ is ordinary}\} \\ P_g &:= \{\Lambda : g(\Lambda) = g \text{ and } \Lambda \notin F_g \sqcup O_g\} \end{aligned}$$

By Lemma 4.4,  $P_g \neq \emptyset$ . By Theorem 4.9 the difference between the number of elements of  $P_g$  which are not in an infinite chain and the number of those that are is non-negative.

If  $g - 1 \equiv 1 \pmod{3}$  then, by Theorem 4.9, there exist 3 numerical semigroups in  $F_g$  which are in an infinite chain and 1 which is not. In  $O_g$  there are 2 elements in an infinite chain and  $g - 2 \geq 4$  that are not. In this way, the difference between the number of elements of  $F_g \sqcup O_g$  which are not in an infinite chain and the number of those that are is non-negative as well. If  $g - 1 \not\equiv 1 \pmod{3}$  then  $F_g$  has only 1 element and it belongs to an infinite chain and the result holds similarly. □

## 5 Fixing the multiplicity

In the tree that organizes all semigroups by genus, the set of descendants of  $\mathbb{N}_0 \setminus \{1, 3\}$  is a single infinite chain of hyperelliptic semigroups. Indeed,  $\mathbb{N}_0 \setminus \{1, 3\}$  is the third semigroup in  $I_{\mathcal{H}}$ . On the other hand  $\mathbb{N}_0 \setminus \{1, 2\}$  is in  $I_{\mathcal{O}}$ , the infinite chain formed by the ordinary semigroups. From each ordinary semigroup in  $I_{\mathcal{O}}$  it emerges a subtree that contains all the semigroups of the same multiplicity (as one can see in [4]). From now on, we focus on these subtrees with fixed multiplicity. One way to transit in such a subtree is, from a semigroup in it, to move to another numerical semigroup of higher genus while maintaining the structure of the first semigroup, that is, maintaining the structure of its non-gaps. The construction explained in next definition does this: it pushes the non-gaps of a numerical semigroup while keeping its multiplicity.

**Definition 5.1.** The *push* of a numerical semigroup  $\Lambda$ , with enumeration  $\lambda$ , by its multiplicity is  $\lambda_1 \oplus \Lambda := \{0\} \cup \{\lambda_1 + \lambda_j; \lambda_j \in \Lambda\}$ .

The set  $\lambda_1 \oplus \Lambda$  as above is a numerical semigroup. In fact, if  $x, y \in \lambda_1 \oplus \Lambda$  then, either  $x, y$  or both are zero, in which case the sum belongs to  $\lambda_1 \oplus \Lambda$ , or there exist  $\lambda_a, \lambda_b \in \Lambda$  such that  $x = \lambda_1 + \lambda_a$  and  $y = \lambda_1 + \lambda_b$ . In this case,  $x + y = \lambda_1 + (\lambda_a + \lambda_1 + \lambda_b) \in \lambda_1 \oplus \Lambda$ . In addition, if  $x \geq \lambda_1 + c(\Lambda)$ , then  $x - \lambda_1 \in \Lambda$  and, so,  $x \in \lambda_1 \oplus \Lambda$ . That is,  $\lambda_1 + c(\Lambda) - 1$  is the maximum of  $(\lambda_1 \oplus \Lambda)^c$  and, thus,  $(\lambda_1 \oplus \Lambda)^c$  is finite. Moreover,  $m(\lambda_1 \oplus \Lambda) = \lambda_1$ ,  $g(\lambda_1 \oplus \Lambda) = g(\Lambda) + \lambda_1 - 1$ ,  $c(\lambda_1 \oplus \Lambda) = \lambda_1 + c(\Lambda)$ ,  $\mathcal{L}(\lambda_1 \oplus \Lambda) = \{0\} \cup \{\lambda_1 + \lambda_j; \lambda_j \in \mathcal{L}(\Lambda)\}$  and  $\#\mathcal{L}(\lambda_1 \oplus \Lambda) = \#\mathcal{L}(\Lambda) + 1$ .

**Lemma 5.2.** Let  $\Lambda$  be a numerical semigroup with enumeration  $\lambda$  and let  $\Pi$  be a non-ordinary numerical semigroup. Then,

- a. If  $\lambda_k$  is a (effective) generator of  $\Lambda$ , then  $\lambda_1 + \lambda_k$  is a (effective) generator of  $\lambda_1 \oplus \Lambda$ .
- b.  $\Pi$  is a child of  $\Lambda$  if and only if  $\lambda_1 \oplus \Pi$  is a child of  $\lambda_1 \oplus \Lambda$ .
- c.  $\Lambda$  lies in an infinite chain if and only if so does  $\lambda_1 \oplus \Lambda$ .

*Proof.* Direct by Definition 5.1. □

The reciprocal of implication a is not true in general. For instance,  $\langle 4, 5, 6 \rangle$  is a leaf, while  $4 \oplus \langle 4, 5, 6 \rangle$  is a stick. From now on, we use the notation  $\lambda_1 \oplus^n \Lambda$ ,  $n \geq 2$ , to represent  $\lambda_1 \oplus (\underbrace{\cdots (\lambda_1 \oplus (\lambda_1 \oplus \Lambda))}_{n \text{ times}})$ .

## 5.1 Prime multiplicity

From Theorem 3.4 one can deduce that, for each prime multiplicity, there is only one infinite chain containing semigroups of that multiplicity. Indeed, this unique chain is  $\omega(m, \mathbb{N}_0)$ . Hence, we can state the following theorem.

**Theorem 5.3.** If  $m$  is a prime, then the number  $i_g(m)$  of numerical semigroups of genus  $g$  and multiplicity  $m$  that are in an infinite chain is:

- a. 0, if  $g < m - 1$ ,
- b. 1, otherwise.

## 5.2 Multiplicity 4


The following tree represents the numerical semigroups of multiplicity 4, genus from 4 to 7 and which are in infinite chains. The edges connect semigroups as usual, being a child of a given semigroup the given semigroup minus one of its effective generators.

Specifically:

$$A = \{0, 4, 6, \rightarrow\}$$

$$B = \{0, 4, 7, \rightarrow\}$$

Figure 1: Tree structure of numerical semigroups of multiplicity 4 that are in an infinite chain, from genus 4 to 7.



$$C = \{0, 4, 6, 8, \rightarrow\}$$

$$D = \{0, 4, 8, \rightarrow\}$$

$$E = \{0, 4, 6, 8, 10, \rightarrow\}$$

$$F = \{0, 4, 8, 10, \rightarrow\}$$

$$G = \{0, 4, 6, 8, 10, 12, \rightarrow\}.$$

As we can observe,  $F = 4 \oplus A$ . So, by Lemma 5.2 and the fact that at most two of the children of a node belong to infinite chains, the children in infinite chains of  $F$  are exactly  $4 \oplus B$  and  $4 \oplus C$ . It is easy to see that, for any positive integer  $n$ ,  $c(4 \oplus^n B) + 1$ ,  $c(4 \oplus^n C)$ ,  $c(4 \oplus^n D)$  and  $c(4 \oplus^n G)$  are multiples of 4. Moreover,  $c(4 \oplus^n E) = 4(n + 2) + 2$  and  $4(n + 1) + 2 \in 4 \oplus^n E$ . With this we obtain that any push by 4 of the numerical semigroups of the tree above has the same number of children in infinite chains as the initial numerical semigroup.

On the other hand, the efficacy of  $4 \oplus^n C = \{0, 4, \dots, 4n, 4n + 2, 4n + 4, \rightarrow\}$ , with  $n \geq 1$ , is equal to 2. So the efficacy of any descendant of  $4 \oplus^n C$  is at most 2. Then by Proposition 4.3 every descendant of  $4 \oplus^n C$  has exactly 1 child in an infinite chain. So, the infinite chain emerging from a node of the form  $4 \oplus^n C$  is made of sticks in the tree of semigroups that are in infinite chains. In that way we have the following theorem.


**Theorem 5.4.** The number  $i_g(4)$  of numerical semigroups of genus  $g$  and multiplicity 4 that are in an infinite chain is:

- a. 0, if  $g \leq 2$
- b. 1 if  $g = 3, 4$
- c.  $\left\lfloor \frac{g + 1}{3} \right\rfloor$ , if  $g \geq 5$ .

*Proof.* The tree in Figure 1 is self-replicated in the tree of infinite chains of semigroups of multiplicity 4. Moreover, as observed before, the numerical semigroups of multiplicity 4 that are in an infinite chain and do not participate in self-replication are sticks in the tree of semigroups that are in infinite chains. □

**Example 5.5.** The tree in Figure 2 was generated with `github.com/mbrasamoros/drawsgtree`, by command `./drawsgtree -g40 -m4 -e infinitechains -etrim -d .2`. In it we can observe eleven complete self-replications of the tree in Figure 1.

Figure 2: Tree of numerical semigroups in infinite chains, with multiplicity 4, from genus 3 up to 40.



### 5.3 Multiplicity 6


The numerical semigroup of multiplicity 6 and with the smallest genus is the ordinary semigroup  $\{0, 6, \rightarrow\}$ . It has two children in infinite chains: the ordinary semigroup  $\{0, 7, \rightarrow\}$  and the semigroup  $\{0, 6, 8, \rightarrow\}$ . To study the numerical semigroups that are in infinite chains and have multiplicity 6, we study the tree  $\nu$  formed by  $\{0, 6, \rightarrow\}$  in its root and connected to  $\{0, 6, 8, \rightarrow\}$  and all semigroups descending from it that are in infinite chains. In other words,  $\nu$  is formed by the subtree rooted at  $\{0, 6, \rightarrow\}$  that contains all numerical semigroups in infinite chains, where the node  $\{0, 7, \rightarrow\}$  and all its descendants have been trimmed.

The semigroups of  $\nu$  are of type  $\Lambda_n = \{0, 6, \dots, 6n, 6n + \beta_1, \dots, 6n + \beta_j, \rightarrow\}$ . In this case we can use the  $j$ -tuple  $(\beta_1, \dots, \beta_j)_n$  to represent  $\Lambda_n$ . For example, we represent  $\{0, 6, 8, \rightarrow\}$  by  $(2)_1$  and  $\{0, 6, 12, 14, 16, 18, 20, 22, 24, \rightarrow\}$  by  $(2, 4, 6, 8, 10, 12)_2$ .

#### 5.3.1 The tree $\tau_n$ and its replications

In Table 5.3.1 we have 23 numerical semigroups that have multiplicity 6, are in infinite chains and have some parenting relationships that can be observed in the tree  $\tau_n$  in Figure 3.

Figure 3: Finite tree  $\tau_n$



Note that  $R_n = 6 \oplus A_n$  and  $6 \oplus \Delta_n = \Delta_{n+1}$  for all  $n \geq 1$  and  $\Delta$  a numerical semigroup from Table 5.3.1. Then, by statements b and c of Lemma 5.2 there is an injection of the finite tree  $\tau_n$  to the tree rooted in  $R_n$  and formed by the numerical semigroups that are in infinite chains. Furthermore,  $\tau_1$  is exactly the tree rooted in  $\{0, 6, 8, \rightarrow\}$  and formed by the numerical semigroups that are in infinite chains up to genus 11. Let us verify that, in fact,  $\tau_n$  is mapped onto the infinite descent of  $R_n$  up to level  $g(R_n) + 5$ . For this, it

Table 1: Some numerical semigroups that have multiplicity 6 and are in infinite chains.

Label	Numerical Semigroup	Only children
$A_n$	$(2)_n$	
$B_n$	$(3)_n$	
$C_n$	$(2, 4)_n$	
$D_n$	$(4)_n$	
$E_n$	$(3, 5)_n$	$c(E_n) + 1 = 6n + 6$ is not a minimal generator
$F_n$	$(2, 5)_n$	$c(F_n) + 1 = 6n + 6$ is not a minimal generator
$G_n$	$(2, 4, 6)_n$	$c(G_n) = 6n + 6$ is not a minimal generator
$H_n$	$(5)_n$	$c(H_n) + 1 = 6n + 6$ is not a minimal generator
$I_n$	$(4, 6)_n$	$c(I_n) = 6n + 6$ is not a minimal generator
$J_n$	$(3, 6)_n$	$c(J_n) = 6n + 6$ is not a minimal generator
$K_n$	$(2, 6)_n$	$c(K_n) = 6n + 6$ is not a minimal generator
$L_n$	$(2, 4, 6, 8)_n$	$c(L_n) = 6n + 8$ is not a minimal generator
$M_n$	$(6)_n$	$c(M_n) = 6n + 6$ is not a minimal generator
$N_n$	$(4, 6, 8)_n$	
$O_n$	$(3, 6, 8)_n$	$c(O_n) + 1 = 6n + 9$ is not a minimal generator
$P_n$	$(2, 6, 8)_n$	$c(P_n) = 6n + 8$ is not a minimal generator
$Q_n$	$(2, 4, 6, 8, 10)_n$	$c(Q_n) = 6n + 10$ is not a minimal generator
$R_n$	$(6, 8)_n$	
$S_n$	$(4, 6, 9)_n$	
$T_n$	$(4, 6, 8, 10)_n$	
$U_n$	$(3, 6, 9)_n$	
$V_n$	$(2, 6, 8, 10)_n$	
$X_n$	$(2, 4, 6, 8, 10, 12)_n$	

suffices to prove that if a numerical semigroup  $\Delta_n = (\beta_1, \dots, \beta_j)_n$  of  $\tau_n$  has only one child in an infinite chain and  $g(\Delta_n) < g(R_n)$ , then  $\Delta_{n+1} = (\beta_1, \dots, \beta_j)_{n+1}$  has only one child in an infinite chain as well. The last column in Table 5.3.1 has been added to clarify this point.

Therefore, the finite tree in Figure 3 is self-replicated in  $\mathcal{v}$ .

### 5.3.2 Outside the replications of $\tau_n$

Now we dedicate to the numerical semigroups of multiplicity 6 that are in an infinite chain and do not participate in that self-replication, i.e., the descendants of  $S_n, T_n, U_n, V_n,$  and  $X_n$ .

**Proposition 5.6.** The numerical semigroups  $T_n = \{0, 6, \dots, 6n, 6n+4, 6n+6, 6n+8, 6n+10, \rightarrow\}$ ,  $U_n = \{0, 6, \dots, 6n, 6n+3, 6n+6, 6n+9, \rightarrow\}$ , and  $X_n = \{0, 6, \dots, 6n, 6n+2, 6n+4, 6n+6, 6n+8, 6n+10, 6n+12, \rightarrow\}$  each belong to one infinite chain.

*Proof.* By Theorem 3.4 the number of infinite chains to which  $T_n$  belongs is one plus the number of descendants of  $\tilde{T}_n = \{0, 3, \dots, 3n, 3n+2, 3n+3, 3n+4, 3n+5, \rightarrow\}$  beyond

$3n + 4$ . Now,  $\tilde{T}_n$  has no generators larger than  $3n + 4$  and, so, it has no descendants beyond  $3n + 4$ . Thus,  $T_n$  only belongs to one infinite chain.

Similarly, the number of infinite chains to which  $U_n$  belongs is one plus the number of descendants of  $\tilde{U}_n = \{0, 2, \dots, 2n, 2n + 1, 2n + 2, 2n + 3, \rightarrow\}$  beyond  $2n + 2$ . But  $\tilde{U}_n$  has no generators larger than  $2n + 2$  and, so, it has no descendants beyond  $2n + 2$ . Thus,  $U_n$  only belongs to one infinite chain.

Finally, the number of infinite chains to which  $X_n$  belongs is one plus the number of descendants of  $\tilde{X}_n = \{0, 3, \dots, 3n, 3n + 1, 3n + 2, 3n + 3, 3n + 4, 3n + 5, 3n + 6, \rightarrow\}$  beyond  $3n + 5$ . But  $\tilde{X}_n$  has no generators larger than  $3n + 5$  and, so, it has no descendants beyond  $3n + 5$ . Thus,  $X_n$  only belongs to one infinite chain.  $\square$

**Proposition 5.7.** The numerical semigroups  $S_n = \{0, 6, \dots, 6n, 6n + 4, 6n + 6, 6n + 9, \rightarrow\}$ , and  $V_n = \{0, 6, \dots, 6n, 6n + 2, 6n + 6, 6n + 8, 6n + 10, \rightarrow\}$  each belong to  $n$  infinite chains.

*Proof.* By Theorem 3.4, the number of infinite chains in which  $S_n$  lies is one plus the number of descendants of  $\tilde{S}_n = \{0, 3, \dots, 3n, 3n + 2, 3n + 3, 3n + 5, \rightarrow\}$  beyond  $3n + 3$ .

If  $n = 1$ ,  $\tilde{S}_n$  has no descendants beyond 6. Hence,  $S_1$  lies in exactly one infinite chain.

Suppose  $n > 1$ . Observe that  $\tilde{S}_n$  has one unique child, which is  $\tilde{S}_n \setminus \{3n + 7\}$ . Observe also that any semigroup containing  $\{0, 3, \dots, 3n, 3n + 2\}$  must contain the semigroup

$$\{0, 3, \dots, 3n, 3n + 2, 3(n + 1), 3(n + 1) + 2, \dots, 3(2n + 1), 3(2n + 1) + 1, 3(2n + 1) + 2, \rightarrow\}.$$

One can deduce that all descendants of  $\tilde{S}_n$  beyond  $3n + 3$  are exactly the semigroups  $\tilde{S}_{n,t} = \{0, 3, \dots, 3n, 3n + 2, 3(n + 1), 3(n + 1) + 2, \dots, 3(n + t), 3(n + t) + 1, 3(n + t) + 2, \rightarrow\}$  for  $3 \leq t \leq n + 1$ . Hence,  $S_n$  lies in exactly  $n$  infinite chains.

By Theorem 3.4, the number of infinite chains in which  $V_n$  lies is one plus the number of descendants of  $\tilde{V}_n = \{0, 3, \dots, 3n, 3n + 1, 3n + 3, 3n + 4, 3n + 5, \rightarrow\}$  beyond  $3n + 4$ .

If  $n = 1$ ,  $\tilde{V}_n$  has no descendants beyond 7. Hence,  $V_1$  lies in exactly one infinite chain.

Suppose  $n > 1$ . Observe that  $\tilde{V}_n$  has one unique child, which is  $\tilde{V}_n \setminus \{3n + 5\}$ . Observe also that any semigroup containing  $\{0, 3, \dots, 3n, 3n + 1\}$  must contain the semigroup

$$\{0, 3, \dots, 3n, 3n + 1, 3(n + 1), 3(n + 1) + 1, \dots, 3(2n), 3(2n) + 1, 3(2n) + 2, \rightarrow\}.$$

One can deduce that all descendants of  $\tilde{V}_n$  beyond  $3n + 4$  are exactly the semigroups  $\tilde{V}_{n,t} = \{0, 3, \dots, 3n, 3n + 1, 3(n + 1), 3(n + 1) + 1, \dots, 3(n + t), 3(n + t) + 1, 3(n + t) + 2, \rightarrow\}$  for  $2 \leq t \leq n$ . Hence,  $V_n$  lies in exactly  $n$  infinite chains.  $\square$

In order to know at which levels of the tree of numerical semigroups the descendants of the semigroups  $S_n$  and  $V_n$  have two children in infinite chains, let us define in the next two lemmas some numerical semigroups using the notation that we explained initially.

**Lemma 5.8.** For every integer  $t$  with  $1 \leq t \leq n + 1$ , the set

$$\begin{aligned} \gamma_{n,t} &= \{0, 6, \dots, 6n, 6n + 4, 7n, 7n + 4, \dots, 6(n + t - 1), 6(n + t - 1) + 4, \\ &\quad 6(n + t), 6(n + t) + 2, \rightarrow\} \\ &= \{0, 6, \dots, 6n, 6n + a_1, 6n + a_2, \dots, 6n + a_{2t+1}, \rightarrow\}, \end{aligned}$$

with

$$a_i = \begin{cases} 3i + 1 & \text{if } i \text{ is odd and } i \neq 2t + 1 \\ 3i, & \text{if } i \text{ is even} \\ 3i - 1, & \text{if } i = 2t + 1. \end{cases}$$

satisfies:

- a.  $\gamma_{n,t}$  is a numerical semigroup.
- b.  $\gamma_{n,1}$  is  $N_n$ , which is the parent of  $S_n$  (and of  $T_n$ ) and, for  $t \geq 2$ ,  $\nu_{n,t}$  is a descendant of  $S_n$ .
- c. If  $t \leq n$  then  $\gamma_{n,t}$  has two children in infinite chains.
- d. The semigroups  $S_n, \gamma_{n,2}, \gamma_{n,3}, \dots, \gamma_{n,n}$  are in the same infinite chain.
- e. The genus of  $\gamma_{n,t}$  is  $5n + 4t + 1$ .

The proof is straightforward and is left to the reader.

Let  $\zeta_n$  be the infinite chain where all the semigroups  $\{S_n, \gamma_{n,t}\}_{2 \leq t \leq n}$  of Lemma 5.8 belong. Since each  $\gamma_{n,t}$  has two children in infinite chains, one child belongs to  $\zeta_n$  and the other one belongs to a different infinite chain. Thus, the  $n$  infinite chains from Proposition 5.7 to which  $S_n$  belongs are  $\zeta_n$  and  $n-1$  other chains, each of them containing  $\gamma_{t,n} \setminus \{c(\gamma_{t,n}) + 1\}$ , for  $t$  between 2 and  $n$ .

**Lemma 5.9.** For every integer  $t$  with  $1 \leq t \leq n$ , the set

$$\begin{aligned} \nu_{n,t} &= \{0, 6, \dots, 6n, 6n + 2, 7n, 7n + 2, \dots, 6(n + t - 1), 6(n + t - 1) + 2, \\ &\quad 6(n + t), 6(n + t) + 2, 6(n + t) + 4 \rightarrow\} \\ &= \{0, 6, \dots, 6n, 6n + b_1, 6n + b_2, \dots, 6n + b_{2t+2}, \rightarrow\}, \end{aligned}$$

with

$$b_i = \begin{cases} 3i - 1 & \text{if } i \text{ is odd} \\ 3i, & \text{if } i \text{ is even and } i \neq 2t + 2 \\ 3i - 2, & \text{if } i = 2t + 2. \end{cases}$$

satisfies:

- a.  $\nu_{n,t}$  is a numerical semigroup.
- b.  $\nu_{n,1} = V_n$  and, for  $t \geq 2$ ,  $\nu_{n,t}$  is a descendant of  $V_n$ .
- c. If  $t \leq n - 1$ , then  $\nu_{n,t}$  has two children in infinite chains.
- d. The semigroups  $V_n, \nu_{n,2}, \nu_{n,3}, \dots, \nu_{n,n-1}$  are in the same infinite chain.
- e. The genus of  $\nu_{n,t}$  is  $5n + 4t + 2$ .



Again, the proof is straightforward and is left to the reader.

Let  $\eta_n$  be the infinite chain where all the semigroups  $\{V_n, \nu_{n,t}\}_{2 \leq t \leq n-1}$  of Lemma 5.9 belong. For  $n$  fixed, since each  $\nu_{n,t}$  and  $V_n$  have two children in infinite chains, one child belongs to  $\eta_n$  and the other one belongs to a different infinite chain. Thus, the  $n$  infinite chains from Proposition 5.7 to which  $V_n$  belongs are  $\eta_n$ , another infinite chain containing  $V_n \setminus \{c(V_n) + 1\}$  and  $n - 2$  other chains, each one containing  $\nu_{t,n} \setminus \{c(\nu_{t,n}) + 1\}$ , for  $t$  between 2 and  $n - 1$ .

### 5.3.3 Growth of the tree $\nu$

In Figure 4 we draw the edges and nodes of  $\nu$  corresponding to semigroups from genus 5 up to genus 31. The tree  $\tau_3$  and the infinite chains  $\zeta_3$  and  $\eta_3$  are highlighted. Notice also that one can observe five full self-replications of the tree from Figure 3. The drawing was generated with the help of <https://github.com/mbrasamoros/drawsg>, with the command `./drawsgtree -g31 -m6 -e infinitechains -etrim -d .1`.

The number  $i_g(6)$  of numerical semigroups of genus  $g$  and multiplicity 6 that are in an infinite chain whenever  $g \geq 5$ , is equal to the number of numerical semigroups on level  $g - 5$  of the tree  $\nu$ . From what has been developed in Subsection 5.3, the growth of the tree  $\nu$  is well known and we are ready to state it in the next theorem.


**Theorem 5.10.** The number  $i_g(6)$  of numerical semigroups of genus  $g$  and multiplicity 6 that are in an infinite chain is

$$\left\{ \begin{array}{ll} 0 & \text{if } g \leq 4 \\ 1 & \text{if } g = 5 \\ g - 5 & \text{if } 5 < g \leq 15 \text{ and } (g \bmod 5) \leq 2 \\ g - 4 & \text{if } 5 < g \leq 15 \text{ and } (g \bmod 5) > 2 \\ g - 5 + \sum_{n=1}^{\lfloor \frac{g-11}{9} \rfloor} n + \sum_{n=1}^{\lfloor \frac{g-8}{9} \rfloor} n + \sum_{n=\lfloor \frac{g-2}{9} \rfloor}^{\lfloor \frac{g-15}{5} \rfloor} \lfloor \frac{g-11-5n}{4} \rfloor + \sum_{n=\lfloor \frac{g+1}{9} \rfloor}^{\lfloor \frac{g-12}{5} \rfloor} \lfloor \frac{g-8-5n}{4} \rfloor & \text{if } g \geq 16 \text{ and } (g \bmod 5) \leq 2. \\ g - 4 + \sum_{n=1}^{\lfloor \frac{g-11}{9} \rfloor} n + \sum_{n=1}^{\lfloor \frac{g-8}{9} \rfloor} n + \sum_{n=\lfloor \frac{g-2}{9} \rfloor}^{\lfloor \frac{g-15}{5} \rfloor} \lfloor \frac{g-11-5n}{4} \rfloor + \sum_{n=\lfloor \frac{g+1}{9} \rfloor}^{\lfloor \frac{g-12}{5} \rfloor} \lfloor \frac{g-8-5n}{4} \rfloor & \text{if } g \geq 16 \text{ and } (g \bmod 5) > 2 \end{array} \right.$$

*Proof.* First of all, notice that the number of self-replications of the structure of  $\tau_1$ , from Figure 3, which lie before level  $g - 5$  of  $\nu$  is  $\lfloor \frac{g-6}{5} \rfloor$ .

If  $g \geq 16$ , then  $\lfloor \frac{g-6}{5} \rfloor \geq 2$  and with this we can define four sets of numerical semi-

Figure 4: Numerical semigroups of multiplicity 6 in infinite chains, from genus 5 up to genus 31. The subtree  $\tau_3$  is highlighted in blue. The parts of the chains  $\eta_3$  and  $\zeta_3$  of genus larger than or equal to 21 are highlighted in green.



groups:

$$\begin{aligned}
A &:= \left\{ \Lambda : \Lambda \in \tau_{\lfloor \frac{g-6}{5} \rfloor + 1} \text{ and } g(\Lambda) = g \right\} \\
B &:= \left\{ \Lambda : \Lambda \text{ is an descendant of } T_n, U_n, \text{ or } X_n, 1 \leq n \leq \left\lfloor \frac{g-6}{5} \right\rfloor, \text{ and } g(\Lambda) = g \right\} \cup \\
&\quad \left\{ \Lambda : \Lambda \text{ is an descendant of } S_n \text{ or } V_n, 1 \leq n \leq \left\lfloor \frac{g-6}{5} \right\rfloor, \Lambda \in \zeta_n \cup \eta_n, \text{ and } g(\Lambda) = g \right\} \\
C &:= \left\{ \Lambda : \Lambda \text{ is an descendant of } S_n, 2 \leq n \leq \left\lfloor \frac{g-6}{5} \right\rfloor, \Lambda \notin \zeta_n, \text{ and } g(\Lambda) = g \right\} \\
D &:= \left\{ \Lambda : \Lambda \text{ is an descendant of } V_n, 2 \leq n \leq \left\lfloor \frac{g-6}{5} \right\rfloor, \Lambda \notin \eta_n, \text{ and } g(\Lambda) = g \right\}.
\end{aligned}$$

In the definitions of  $C$  and  $D$  the case  $n = 1$  has been excluded since  $S_1$  belongs only to one infinite chain which is exactly  $\nu_1$ , while  $V_1$  belongs only to one infinite chain which is exactly  $\eta_1$ .

Now, the numerical semigroups of genus  $g$ ,  $g \geq 16$ , and multiplicity 6 which are in an infinite chain are the numerical semigroups of  $A \sqcup B \sqcup C \sqcup D$ .

To set the cardinality of  $A$ , notice that the remainder  $r$  such that  $g - 6 = 5 \lfloor \frac{g-6}{5} \rfloor + r$  tells us at which level of the tree  $\tau_{\lfloor \frac{g-6}{5} \rfloor + 1}$  are the numerical semigroups of genus  $g$  that participate in self-replication. So, the cardinality of  $A$  is equal to  $r + 1$ , if  $r = 0, 1, 4$  and equal to  $r + 2$  when  $r = 2, 3$ .

By Proposition 5.6, Lemma 5.8, and Lemma 5.9, the cardinality of  $B$  is equal to  $5 \lfloor \frac{g-6}{5} \rfloor$ .

On the other hand,  $5 \lfloor \frac{g-6}{5} \rfloor = g - 6 - r$ . So,

$$\#A + \#B = \begin{cases} g - 5 & \text{if } r = 0, 1, 4 \\ g - 4 & \text{if } r = 2, 3. \end{cases}$$

Now,  $r = 0, 1, 4$  is equivalent to  $g - 6 \equiv 0, 1, 4 \pmod{5}$ , that is,  $g \equiv 1, 2, 0 \pmod{5}$ , while  $r = 2, 3$  is equivalent to  $g \equiv 3, 4 \pmod{5}$ . Hence,

$$\#A + \#B = \begin{cases} g - 5 & \text{if } (g \pmod{5}) = 0, 1, 2 \\ g - 4 & \text{if } (g \pmod{5}) = 3, 4. \end{cases} \quad (1)$$

The descendants of  $S_n$ , for  $2 \leq n \leq \lfloor \frac{g-6}{5} \rfloor$ , which belong to infinite chains but do not belong to  $\zeta_n$  and have genus  $g$ ,  $g \geq 16$ , are in bijection with the numerical semigroups  $\gamma_{n,t}$  from Lemma 5.8 such that  $g(\gamma_{n,t}) < g$ . Since  $g(\gamma_{n,t}) = 5n + 4t + 1$ , we have that the

cardinality of  $C$  is equal to

$$\begin{aligned}
& \# \left\{ (\tilde{n}, \tilde{t}) : 5\tilde{n} + 4\tilde{t} + 1 < g, \text{ where } 2 \leq \tilde{n} \leq \left\lfloor \frac{g-6}{5} \right\rfloor \text{ and } 2 \leq \tilde{t} \leq \tilde{n} \right\} \\
&= \# \left\{ (n, t) : 1 \leq t \leq \left\lfloor \frac{g-11-5n}{4} \right\rfloor, 1 \leq t \leq n \text{ for } 1 \leq n \leq \left\lfloor \frac{g-6}{5} \right\rfloor - 1 \right\} \\
&= \# \left\{ (n, t) : 1 \leq t \leq \min \left( \left\lfloor \frac{g-11-5n}{4} \right\rfloor, n \right) \text{ for } 1 \leq n \leq \left\lfloor \frac{g-6}{5} \right\rfloor - 1 \right\} \\
&= \sum_{n=1}^{\left\lfloor \frac{g-11}{9} \right\rfloor} n + \sum_{n=\left\lfloor \frac{g-2}{9} \right\rfloor}^{\left\lfloor \frac{g-11}{5} \right\rfloor} \left\lfloor \frac{g-11-5n}{4} \right\rfloor \\
&= \sum_{n=1}^{\left\lfloor \frac{g-11}{9} \right\rfloor} n + \sum_{n=\left\lfloor \frac{g-2}{9} \right\rfloor}^{\left\lfloor \frac{g-15}{5} \right\rfloor} \left\lfloor \frac{g-11-5n}{4} \right\rfloor.
\end{aligned}$$

On the other hand, by Lemma 5.9, the set  $D$  is in bijection with the set  $\{\nu_{n,t} : 2 \leq n \leq \left\lfloor \frac{g-6}{5} \right\rfloor, 2 \leq t \leq n-1 \text{ and } g(\nu_{n,t}) < g\} \cup \{V_n : 2 \leq n \leq \left\lfloor \frac{g-6}{5} \right\rfloor\}$ . Now, since  $V_n = \nu_{n,1}$ ,  $D$  is in bijection with the set  $\{\nu_{n,t} : 2 \leq n \leq \left\lfloor \frac{g-6}{5} \right\rfloor, 1 \leq t \leq n-1 \text{ and } g(\nu_{n,t}) < g\}$ . Furthermore, since  $g(\nu_{n,t}) = 5n + 4t + 2$ , we have that the cardinality of  $D$  is equal to

$$\begin{aligned}
& \# \left\{ (\tilde{n}, t) : 5\tilde{n} + 4t + 2 < g, \text{ where } 2 \leq \tilde{n} \leq \left\lfloor \frac{g-6}{5} \right\rfloor \text{ and } 1 \leq t \leq \tilde{n} - 1 \right\} \\
&= \# \left\{ (n, t) : 1 \leq t \leq \left\lfloor \frac{g-8-5n}{4} \right\rfloor, 1 \leq t \leq n \text{ for } 1 \leq n \leq \left\lfloor \frac{g-6}{5} \right\rfloor - 1 \right\} \\
&= \sum_{n=1}^{\left\lfloor \frac{g-8}{9} \right\rfloor} n + \sum_{n=\left\lfloor \frac{g+1}{9} \right\rfloor}^{\left\lfloor \frac{g-12}{5} \right\rfloor} \left\lfloor \frac{g-8-5n}{4} \right\rfloor.
\end{aligned}$$

We omitted two steps since they can be parallelized from the previous sequence of equalities.

If  $11 \leq g \leq 15$ , then  $\left\lfloor \frac{g-6}{5} \right\rfloor = 1$  and, by Proposition 5.6 and Proposition 5.7, the numerical semigroups with that genus and multiplicity 6 which are in an infinite chain are exactly one descendant of each  $S_1, T_1, U_1, V_1, X_1$  and the numerical semigroups in  $\tau_2$  of genus  $g$ . And, if  $6 \leq g \leq 10$ , then  $\left\lfloor \frac{g-6}{5} \right\rfloor = 0$  and the numerical semigroups with that genus and multiplicity 6 which are in an infinite chain are only the numerical semigroups in  $\tau_1$  with genus  $g$ . Both collections add up  $g-5$  numerical semigroups when  $(g \bmod 5) = 0, 1$  or  $2$ , and  $g-4$  otherwise. □

## 6 Acknowledgments

The first author was supported by the Catalan government under grant 2021 FIDUR 00189. Both authors were supported by the Spanish government under grant PID2021-124928NB-I00, and by the Catalan government under grant 2021 SGR 00115.

All the graphs have been drawn using the `drawsgtree` tool, which can be downloaded from <https://github.com/mbrasamoros/drawsgtree>.

## References

- [1] M. Bras-Amorós: On numerical semigroups and their applications to algebraic geometry codes, *Thematic Seminar on Algebraic Geometry, Coding and Computing*, Universidad de Valladolid, Segovia, 2007.
- [2] M. Bras-Amorós, Fibonacci-like behavior of the number of numerical semigroups of a given genus, *Semigroup Forum* **76** (2008) 379–384, <http://dx.doi.org/10.1007/s00233-007-9014-8>.
- [3] M. Bras-Amorós, Bounds on the number of numerical semigroups of a given genus, *Journal of Pure and Applied Algebra* **213** (2008) 997–1001.
- [4] M. Bras-Amorós and S. Bulygin, Towards a better understanding of the semigroup tree, *Semigroup Forum* **79** (3) (2009) 561–574.
- [5] J. Fromentin, F. Hivert, Exploring the tree of numerical semigroups, *Math. Comp.* **85** (2016) 2553–2568, <http://dx.doi.org/10.1090/mcom/3075>.
- [6] P.A. Garcia-Sanchez and J.C. Rosales, *Numerical semigroups*. New York: Springer, 2009.
- [7] A. Zhai, Fibonacci-like growth of numerical semigroups with a given genus, *Semigroup Forum* **86** (2013) 634–662, <http://dx.doi.org/10.1007/s00233-012-9456-5>.