

Incorporating Connectivity among Internet Search Data for Enhanced Influenza-like Illness Tracking

Shaoyang Ning^{1,*}, Ahmed Hussain^{1,+}, and Qing Wang²

¹Department of Mathematics and Statistics, Williams College, Williamstown, MA, 01267, USA

²Department of Mathematics, Wellesley College, Wellesley, MA, 02481, USA

*sn9@williams.edu

+This author contributed to the project as an undergraduate research assistant at Williams College

ABSTRACT

Big data collected from the Internet possess great potential to reveal the ever-changing trends in society. In particular, accurate infectious disease tracking with Internet data has grown in popularity, providing invaluable information for public health decision makers and the general public. However, much of the complex connectivity among the Internet search data is not effectively addressed among existing disease tracking frameworks. To this end, we propose ARGO-C (Augmented Regression with Clustered GOogle data), an integrative, statistically principled approach that incorporates the clustering structure of Internet search data to enhance the accuracy and interpretability of disease tracking. Focusing on multi-resolution %ILI (influenza-like illness) tracking, we demonstrate the improved performance and robustness of ARGO-C over benchmark methods at various geographical resolutions. We also highlight the adaptability of ARGO-C to track various diseases in addition to influenza, and to track other social or economic trends.

Introduction

Big data collected from the Internet, recording billions of people's digital footprints, possess great potential to reveal the ever-changing trends in society. A growing number of attempts have been made to harness the potential of Internet data to address issues in a wide range of fields, including public health [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11], economics [12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23], business [24, 25, 26], finance [17, 27, 28], social policy [29], popular culture trends [30], among others. In particular, digital disease detection, which utilizes big data from online source to provide accurate and up-to-date tracking of infectious diseases, has grown in popularity, especially since the 2009 H1N1 influenza pandemic and the 2020 global pandemic of COVID-19 [31, 32, 33, 34, 35, 36]. However, despite the existing efforts in enhancing infectious disease tracking with Internet search data, many challenges and limitations still remain. In particular, there lacks a statistically rooted and integrative approach in existing digital disease tracking frameworks that effectively accounts for the connectivity within the Internet search data. The purpose of our paper is to pioneer a statistical learning method, ARGO-C (Augmented Regression with Clustered GOogle data). ARGO-C takes advantage of the interconnections among the Internet search data and aims to improve the accuracy and interpretability of the disease tracking framework. Our method focuses on influenza (flu) tracking, but has the generality and flexibility to be adapted to tracking other infectious diseases or other social/economic trends.

Influenza (flu) epidemics occur every year with varying timing and intensity. It may claim up to 650,000 deaths per year worldwide [37], and, on average, results in 610,660 life-years loss in the US [38]. Our ability to prepare for and respond to epidemics or pandemics depends on the timely tracking and forecasting of the infectious disease activities [39, 40, 41]. Traditionally, the tracking and surveillance of flu activities in the US rely mainly on the US Outpatient Influenza-like Illness Surveillance Network (ILINet) by the Centers for Disease Control (CDC). Each week, ILINet collects outpatients information from thousands of healthcare providers across the nation and reports the percentage of Influenza-like Illness patients (%ILI). However, due to the time incurred by data collection, aggregation, and administrative processing, the CDC's weekly flu report usually lags behind real time by 1-2 weeks, which is far from optimal for tracking a fast-spreading, ever-changing disease epidemic such as flu.

In order to eliminate the time lag between CDC's flu report and the real time event, digital disease detection [42], a new disease tracking framework based on Internet data, was proposed and has since revolutionized the landscape of flu tracking. In particular, methods for digital flu detection employ statistical or mechanistic models to harness Internet-derived data from various sources [1, 2, 6, 43, 44, 45, 46, 47, 48, 49, 50, 50, 51, 52, 53, 54, 55] to provide current estimation of future prediction of flu activity (usually in terms of %ILI). This is also referred to as "nowcasting", in contrast to forecasting (i.e., predicting future). Among these approaches, Google Flu Trends (GFT) [2], which uses the volume of selected Google search terms to estimate current influenza-like illnesses (ILI) activity, has attracted the most attention. However, the significant prediction

errors by GFT in the following flu season, as well as its lack of transparency and reproducibility, has incurred many criticisms. This has also inspired a growing literature [31, 56, 57, 58, 59, 60] in digital disease detection, with the aim to identify what Google had done wrong and improve from there.

In particular, the ARGO framework (AutoRegression with GOogle search data) [61] provides robust and highly accurate ILI estimates at the national level by directly addressing the limitations of GFT. Through a linear model design that is justified by a hidden Markov model, the ARGO framework effectively integrates multi-source information from the CDC's flu reports and Google's search volume data while accounting for dynamics in flu epidemics and people's search patterns. Due to its flexibility and generality, ARGO has been well-adapted to multi-resolution, multi-disease tracking based on multi-source data [36, 52, 62, 63, 64, 65, 66, 67].

Nevertheless, among the existing methods for digital disease tracking, few have directly addressed the complex connectivity observed within the Internet data. Particularly, many of the Internet search terms included in ARGO share semantic similarities, such as phrases like "treat flu", "how to treat the flu", and "treat the flu", and consequently, their search volumes may be closely related. Such connection among Internet data has never been explicitly investigated in the existing ARGO-derived frameworks. Our goal here is to propose ARGO-C, a general framework that incorporates underlying interconnections among the Internet data and improves flu tracking's accuracy and interpretability. Our contribution is significant in that (i) ARGO-C provides an innovative statistical learning framework that explicitly models the connectivity among Internet search data and utilizes the information to improve the accuracy of disease tracking; (ii) it enhances the interpretability of the predictive model by revealing the clustering structure of search terms and each cluster's contribution to the model; (iii) it provides a general framework that is readily adaptable to tracking other diseases and/or social/economic trends with Internet search data.

Methods

The ARGO model

The model of ARGO targets the time series of logit-transformed ILI activity level Y_t (i.e., the logit-transformed %ILI) at the national-level from CDC's flu report at week t . It assumes a Markovian structure in a period M of flu activities $\mathbf{Y}_{(t-M+1):t}$ (note that $\{(t-M+1):t\} = \{t-M+1, t-M+2, \dots, t-1, t\}$, the set of all integer indices in between), which takes the form of an autoregressive model. Consequently, $\mathbf{X}_t = (X_{1t}, X_{2t}, \dots, X_{pt})^\top$, a vector of p log-transformed search volumes of flu-related queries from Google, depends solely on the the flu activities at time t . Jointly, the structure of the search volumes \mathbf{X}_t and recent flu activities $\mathbf{Y}_{(t-M+1):t}$ can be summarized by a Hidden Markov model, as presented in Equation (1).

$$\cdots \rightarrow \mathbf{Y}_{(t-M):(t-1)} \rightarrow \mathbf{Y}_{(t-M+1):t} \rightarrow \mathbf{Y}_{(t-M+2):(t+1)} \rightarrow \cdots$$

$$\downarrow \qquad \downarrow \qquad \downarrow$$

$$\mathbf{X}_{t-1} \qquad \mathbf{X}_t \qquad \mathbf{X}_{t+1} \tag{1}$$

With further assumptions on stationarity, normality of the observations, and linear dependence of search volumes \mathbf{X}_t on flu activities $\mathbf{Y}_{(t-M+1):t}$, the prediction distribution $Y_t | \mathbf{Y}_{(t-M+1):(t-1)}, \mathbf{X}_t$ is Normal with a mean linear in $\mathbf{Y}_{(t-M+1):(t-1)}$ and \mathbf{X}_t and a stationary variance. This leads to a linear predictive model for ARGO:

$$Y_t = \mu_y + \sum_{s=1}^M \alpha_s Y_{t-s} + \sum_{j=1}^p \beta_j X_{jt} + \varepsilon_t, \tag{2}$$

where the random errors ε_t are i.i.d., with mean 0 and constant variance σ_ε^2 . Given the large number of predictors in model (2), ARGO adopts an L_1 -regularized regression [68] to achieve an adaptive selection of predictors. To further account for the dynamic changes in search patterns and flu epidemics, ARGO employs a rolling window prediction scheme, with a sliding training set of $N = 104$ weeks. Therefore, at a given week T , the coefficients μ_y , $\beta = (\beta_1, \beta_2, \dots, \beta_p)^\top$, and $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_M)^\top$ are estimated as follows:

$$(\hat{\mu}_y, \hat{\beta}, \hat{\alpha}) = \arg \min_{\mu_y, \beta, \alpha} \left\{ \frac{1}{2} \sum_{t=T-N}^{T-1} (Y_t - \mu_y - \mathbf{X}_t^\top \beta)^2 + \lambda \|\beta\|_1 + \lambda \|\alpha\|_1 \right\}, \tag{3}$$

where $\|\cdot\|$ represents the L_1 norm, and λ ($\lambda \geq 0$) is the tuning parameter for penalization.

The ARGO2 and ARGOX models for localized flu tracking

ARGO2 [63] generalizes the national ARGO model to localized, regional flu tracking (the US Health and Human Services (HHS) regions). It is operated in two steps:

- Step One is to extract Internet search information. It employs the framework of ARGO model and applies to each region individually (with autoregressive terms left out) to obtain a preliminary raw estimate for each region's %ILI of the week.
- Step Two integrates multi-source, multi-resolution information to boost the regional %ILI prediction. Specifically, the best linear predictor based on a structured covariance matrix is used to provide joint %ILI prediction for all 10 regions, which incorporates Google search information from Step one's raw estimates, the national flu baseline level estimated by the ARGO model, and the recent flu time series trends from the latest CDC's flu reports.

ARGOX [65] further extends the ARGO framework to state-level, thereby establishing a coherent multi-resolution framework for digital flu tracking. Specifically, ARGOX first dichotomizes all the states into two groups: the epidemically connected and the disconnected, and then customizes different prediction models in the Step Two algorithm accordingly.

Penalization methods with group-wise sparsity

As discussed in the previous sections, due to the high dimension of the Google Search terms, penalization technique is integrated into the ARGO, ARGO2, and ARGOX algorithms. In this paper, we are interested in adapting penalization methods with a group-wise regularization to these frameworks.

In linear and generalized linear regressions, penalization methods are among popular practical tools that aim at reducing the variability of parameter estimation by conventional methods, such as the ordinary least squares (OLS) method or the maximum likelihood (ML) algorithm, or decreasing the dimensionality of a given feature space. Over the past several decades, a number of penalization techniques have been developed. For instance, Horel and Kennard[69] proposed the ridge estimator that shrinks the OLS estimator towards zero so as to alleviate its large sampling variation. Later, Lasso and elastic net algorithms were proposed [68, 70, 71, 72, 73], both of which can realize simultaneous variable selection and parameter estimation. These methods each is defined on a different penalty function. However, they all impose regularization on individual model parameters and intend to drive each estimated parameter towards zero.

Motivated by multi-factor analysis of variance (ANOVA), Yuan and Lin [74] proposed a group lasso method that is designed to select groups of indicator variables associated with the same factor. They combine the dummy indicator variables defined for a given factor together, and then select a subset of important factors through a penalty term imposed on the corresponding grouped model parameters. More specifically, suppose there are p predictor variables which can be partitioned into K groups with group size p_k ($1 \leq k \leq K; \sum_{k=1}^K p_k = p$). In the context of linear regression, the group lasso solution for the regression coefficients β can be expressed as

$$\hat{\beta}^{GL} = \arg \min_{\beta} \left\{ \frac{1}{2} \sum_{i=1}^n \left(Y_i - \sum_{k=1}^K (\mathbf{X}_i^{(k)})^\top \beta^{(k)} \right)^2 + \lambda \sum_{k=1}^K \sqrt{p_k} \|\beta^{(k)}\|_2 \right\},$$

where Y_i is the response variable for the i -th observation, $\mathbf{X}_i^{(k)}$ is the i -th observation's corresponding predictors in group k , $\beta^{(k)}$ is the vector of coefficients in group k , and λ ($\lambda \geq 0$) is the tuning parameter for penalization. Later, Meier et al.[75] applied the group lasso method to logistic regression and other generalized linear regression models.

More recently, Simon et al.[76] studied a sparse group lasso (SGL) algorithm, imposing a convex combination of the L_1 - and L_2 -norm penalties on the grouped and individual parameters respectively. In linear regression, the SGL solution for the regression coefficients β is

$$\hat{\beta}^{SGL} = \arg \min_{\beta} \left\{ \frac{1}{2} \sum_{i=1}^n \left(Y_i - \sum_{k=1}^K \mathbf{X}_i^{(k)} \beta^{(k)} \right)^2 + (1-\alpha)\lambda \sum_{i=1}^m \sqrt{p_k} \|\beta^{(k)}\|_2 + \alpha\lambda \|\beta\|_1 \right\},$$

where $\alpha \in [0, 1]$ is a weight tuning parameter, $\lambda \geq 0$, and $\|\cdot\|_1$ is the L_1 norm. When $\alpha = 0$, the solution is reduced to the group lasso solution; when $\alpha = 1$, it becomes the lasso solution. Simon et al.[76] also discussed the application of SGL in other model settings, such as generalized linear regression.

ARGO-C: the proposed method

In this subsection, we illustrate in detail the methodology of our proposed approach, ARGO-C.

We first recognize that some of the flu-related Google search terms often share a common theme. For example, search terms concerning flu treatments may include phrases such as "treat flu", "how to treat the flu", and "treat the flu"; or a specific

sub-type of flu may contain terms like “influenza a”, “influenza type a”, etc. It is then natural to consider clustering similar search terms together, and then fit a penalized linear regression model with a group-wise penalty. This direction motivated our project.

In particular, our approach enhances the infrastructure of ARGO. Here we focus on the ILI tracking at the national level as an illustration. Additionally, to showcase the generality of our approach, we provide an exemplary integration of our approach to the ARGO2 and ARGOX frameworks [63] for localized (regional and state-level) flu tracking. Our methodology can be readily integrated to existing methods to track other infectious diseases [36, 62, 66, 67], or other social and/or economic trends [23].

National level

For national level %ILI prediction, our proposed ARGO-C is realized in two steps: in Step 1, we identify the connectivity structure among the candidate flu-related Google search terms by unsupervised statistical learning; in Step 2, we integrate the identified cluster structure into the predictive model of weekly flu activities (in terms of %ILI) using a penalized regression with group-wise regularization.

We start with defining some notations that we will refer to for the rest of the paper. Let Y_t be the logit-transformed percentage of influenza-like illness (%ILI) at the national level at week t from CDC’s weekly flu reports; $\mathbf{X}_t = (X_{1t}, X_{2t}, \dots, X_{pt})^\top$ be the log-transformed Google search volumes of p flu-related terms at week t . Due to the delay of CDC’s reports, at the current week T , we are only able to observe $Y_{1:(T-1)}$, up to the previous week; the Google search data are instead up-to-date, with $\mathbf{X}_{1:T}$ all available. Our proposed method can be summarized as follows:

- **Step 1: clustering Internet search terms.**

Partition the p search terms into K groups, denoted by G_1, \dots, G_K ($K \geq 2$). This can be done using some standard clustering method, such as hierarchical clustering [77], k-means [78], or model-based clustering [79]. And, standard statistical programming languages, such as *R* [80] or *Python* [81], have existing packages that can realize these methods easily. In practice, we recommend using hierarchical clustering based on correlation as the distance metric between the search terms’ time series. The number of clusters K can be determined through investigating measures such as the within-group variance, silhouette [82], or the gap statistic [83].

- **Step 2: nowcasting using group-structured, penalized regression.**

Predict the current week’s (logit-transformed) ILI%, Y_T , by

$$\hat{Y}_T = \mu_y + \sum_{s=1}^m \gamma_s Y_{T-s} + \sum_{j=1}^p \beta_j X_{jT}, \quad (4)$$

where m is the length of the lagged time series terms, μ_y is the intercept, $\gamma = \gamma_{m:1}$ is the autoregressive coefficients, and $\beta = \beta_{1:p}$ are the exogenous coefficients of Google search terms. The coefficients β are partitioned into K groups as identified in the previous step. Then, the model parameters in Equation (4), (μ_y, β, γ) , can be estimated by minimizing the following penalized sum of squares quantity,

$$\frac{1}{2} \sum_{t=T-N}^{T-1} \left(Y_t - \mu_y - \sum_{s=1}^m \gamma_s Y_{t-s} - \sum_{j=1}^p \beta_j X_{jt} \right)^2 + \alpha \lambda (\|\gamma\|_1 + \|\beta\|_1) + (1 - \alpha) \lambda \left(\sum_{k=1}^K \sqrt{p_k} \|\beta^{(k)}\|_2 \right)$$

where N is the training window length, $\beta^{(k)} = \{\beta_j, j \in G_k\}$ is the coefficients of search terms in cluster k , and p_k is the size of cluster k . Specifically, the cluster structure of search terms is incorporated through the sparse group lasso (SGL) regularization $\|\beta^{(k)}\|_2 = \sqrt{\sum_{j \in G_k} \beta_j^2}$, which intends to impose sparsity on all the coefficients of terms in each cluster simultaneously. Note that α and λ are the tuning parameters, with α determining the weights between the individual and group-wise regularization and λ controlling the strength of regularization. In practice, we use SGL’s default setting $\alpha = 0.95$ and use cross-validation to select λ . We also follow the ARGO’s default setting and set the training windows to be two years, i.e., $N = 104$ weeks.

Regional level

For ARGO-C’s regional flu tracking, we break down ARGO2’s Internet data extraction step (i.e., Step 1) into two sub-steps.

- **Step 1.1: clustering Internet search terms.**

For each region r ($r = 1, \dots, 10$), follow the similar procedure in Step 1 of the national %ILI to partition the p search terms into K groups, denoted by $G_1^{(r)}, \dots, G_K^{(r)}$. We here keep the cluster number K the same as the national level for consistency.

- **Step 1.2: extracting regional Google information based on clustered search terms.**

Obtain preliminary estimates for regional level %ILI based completely on the region-wise Google search data and the clustering structure learned in Step 1.1. Specifically, the raw estimate for region r 's (log-transformed) %ILI at the current week T , $\hat{Y}_T^{(r)}$, is given by

$$\hat{Y}_T^{(r)} = \mu_Y^{(r)} + \sum_{j=1}^p \beta_j^{(r)} X_{jt}^{(r)}, \quad (5)$$

where the superscript (r) indicates the region-specific parameters and data. And, similar to Step 2 of ARGO-C at the national level, the parameters are estimated through a sparse group lasso regularization:

$$\frac{1}{2N} \sum_{t=T-N}^{T-1} \left(Y_t^{(r)} - \mu_y^{(r)} - \sum_{j=1}^p \beta_j^{(r)} X_{jt}^{(r)} \right)^2 + \alpha \lambda \|\beta^{(r)}\|_1 + (1-\alpha) \lambda \left(\sum_{k=1}^K \sqrt{p_k} \|\beta^{(k,r)}\|_2 \right)$$

where N is the training window length, $\beta^{(k,r)} = \{\beta_j^{(r)}, j \in G_k\}$ is the region-specific coefficients of search terms in cluster k , and p_k is the size of cluster k .

- **Step 2: cross-regional boosting.**

This follows exactly the same as the original Step 2 in ARGO2, to prediction 10 regions' %ILI jointly based on multi-source, multi-resolution information. More details can be found in [63].

State level

At the state level, ARGO-C is readily adaptable to fit into the ARGOX framework. Step 1.1 and 1.2 follow the regional level procedure above but applied to each state; Step 2 directly inherits the original Step 2 of ARGOX [65], with the same dichotomic treatment of the 51 states/district/city.

Data

CDC's %ILI Data

The CDC's weekly flu report is released every Friday, listing the percent of outpatient visits with influenza-like illness (%ILI) in the previous week [84] (<https://www.cdc.gov/flu/weekly/overview.htm>). Therefore, the CDC's %ILI always lags behind real time by at least one week. The CDC's report includes %ILI at the national level, of the 10 US Health and Human Services (HHS) regions, and the 51 states/district/city (50 states plus Washington DC, excluding Florida but including New York City). The CDC's %ILI data for this study were collected on January 29, 2023.

Google Search Data

The Internet search data from Google are publicly available through Google Trends (trends.google.com). Once a user specifies a desired query term (or a topic), a geographical indicator, and a time range on Google Trends, the website will return a time series of the term's weekly search volumes. With Google Trends API, we are able to obtain the un-normalized search frequencies for the specified term, which includes all the searches that contain the entire term.

The search query terms in this study are selected based on previous works [61, 63, 65]. Notably, we included 161 flu-related search terms/topics, with 71 terms identified by March 29, 2009 and the remaining by May 22, 2010 to account for the 2009 H1N1 outbreak. Table S4 lists these search terms. The regional level Google search volumes is aggregated based on state-level data, following the ARGO2 framework [63].

We admit that the Google search data may only be representative of the search interests among Google users rather than the entire population. The ARGO (including ARGO2 and ARGOX) framework [65] attempts to correct for such potential bias in the modeling.

As one benchmark method for comparison, we downloaded the discontinued Google Flu Trends (GFT) data (<https://www.google.org/flutrends/about/data/flu/us/data.txt>). GFT has the weekly %ILI prediction from January 1, 2004 to August 9, 2015.

Evaluation Metrics

We use three metrics to evaluate the accuracy of an estimate against the actual %ILI released by the CDC: the root mean squared error (RMSE), the mean absolute error (MAE), and the Pearson correlation (Correlation). RMSE between an estimate \hat{p}_t and the true value p_t over period $t = 1, \dots, N$ is given by

$$\text{RMSE} = \sqrt{\frac{1}{N} \sum_{t=1}^N (\hat{p}_t - p_t)^2}.$$

The MAE between an estimate \hat{p}_t and the true value p_t over period $t = 1, \dots, N$ is defined as

$$\text{MAE} = \frac{1}{N} \sum_{t=1}^N |\hat{p}_t - p_t|.$$

And, the correlation we considered is the Pearson correlation coefficient between $\hat{\mathbf{p}} = (\hat{p}_1, \dots, \hat{p}_N)$ and $\mathbf{p} = (p_1, \dots, p_N)$.

Results

We first applied the ARGO-C model to retrospectively estimate the weekly %ILI at the US national level from March 29, 2009 to January 28, 2023. Figures S1 - S2 illustrate the clustering structures identified among the flu-related Google search terms by ARGO-C's Step 1. The results were obtained via hierarchical clustering with an average linkage function using correlation as the distance metric, based on the Google data available prior to the earliest prediction date. In particular, we realized the clustering analysis twice, using two sets of search terms of 71 and 161 terms/topics, respectively (see Table S4). Then, the identified clusters of search terms were incorporated into ARGO-C's Step 2 to predict %ILI after the collection date of the corresponding set of search terms (March 29, 2009 - May 21, 2010 for the first 71 terms, and May 22, 2010 onward for all 161 terms, respectively). More specifically, 53 clusters were identified among the 71 search terms originally collected by March 29, 2009 (see Figure S1, where the number of clusters was determined by minimizing the sum of within-group variation while preserving interpretability). As can be seen, the clustering method successfully grouped search terms with close semantics into the same group. For example, one cluster consists of search terms related to flu treatments, containing phrases such as "treat flu", "how to treat the flu", and "treat the flu". In addition, another cluster includes search terms about respiratory illness related to flu, such as "sinus", "bronchitis", "pneumonia", and "walking pneumonia". In the same fashion, based on the second set of 161 flu-related search terms/topics, 45 clusters were identified, as illustrated in Figure S2.

Figure 1 and Table 1 summarize the national-level prediction performance of our proposed model, ARGO-C, in comparison with benchmark methods, including Google's original GFT (discontinued on July 11, 2015), vector autoregression model with lag 1 (VAR1), the original ARGO model [61], and the naive method which simply carries over the previous week's %ILI to predict the current week. We first focus on the period prior to the influence of COVID-19. That is, we exclude the period when the %ILI reported by CDC was highly confounded and contaminated by COVID-19 symptoms and cases, thus not accurately reflecting the flu activities anymore [67, 84]. During this whole period from 2009 to 2020 (Figure 1), our method ARGO-C shows the leading prediction accuracy compared to all other benchmarks across all three performance metrics. In particular, the improvement of ARGO-C from ARGO confirms the potential of integrating the interconnection among Internet search data into the modeling process while showcasing the effectiveness of ARGO-C in utilizing such information to enhance disease prediction. The effective use of connectivity among Internet data is further confirmed by a closer look at the evolving patterns of search terms included and excluded in the ARGO-C model over time. Among three exemplary clusters highlighted in Figure S3, we observe that ARGO-C frequently selects/filters out an entire cluster of search terms thanks to the introduced group-penalty structure, and thus fully takes advantage of the interconnection among Internet data; on the other hand, each search term can also be selected/filtered out individually within a cluster, indicating a good balance between the individual and group-wise penalization. Breaking down the results into each flu season, ARGO-C's performance is consistent, giving the most accurate predictions in majority of the flu seasons. Notably, ARGO-C is the single leading method in every flu season since 2010 by the measure of correlation (Table 1). This highlights ARGO-C's strength in predicting the flu epidemic trends. Moreover, ARGO-C's advantage over the vanilla ARGO is significantly evident in certain difficult flu seasons, yielding up to 30% of error reduction (i.e., there are error reductions of 30.7% for '10-'11, 18.4% for '14-'15, and 18.9% for '16-'17 in terms of RSME). Additionally, the 95% prediction interval given by ARGO-C (based on the stationary bootstrap [61]) has an empirical coverage of 95.08%.

We also applied the proposed ARGO-C model to localized flu tracking. At the regional level, ARGO-C again shows the strongest performance across all 10 regions in all three accuracy metrics. More specifically, compared to the naive method, ARGO-C reduces the RMSE by 12% to 29%, and improves the MAE measure by 11% to 25% (Table 2). In addition, the correlation measure based on the ARGO-C method is uniformly higher than other benchmarks in all 10 regions. Breaking

down into individual flu seasons (Table S5-S14), ARGO-C still leads in vast majority of all evaluated periods. The strengths of ARGO-C at the regional level also reaffirms our projection that interconnection among Internet search data would contribute effectively in improving flu tracking performance. Additionally, Table 3 reports the empirical interval coverage for the regional %ILI prediction (ranging from 93% to 96%, given a nominal level of 95%), further confirming the reliability of ARGO-C.

Table 4 and S2 summarize ARGO-C's state-level flu tracking performance in comparison with the benchmarks. Averaging over 51 states, ARGO-C again is the best performing method compared to all benchmarks, showcasing its adaptability and robustness in high-resolution disease tracking. The strength of ARGO-C attributes to our effective modeling of interconnection among search terms, which is even more relevant for efficient extracting of low-quality, high-noise Internet data at high resolution. More detailed reports on each individual state and each flu season are given in Tables S15 - S65. During the period from 2014 to 2020 (pre-COVID, Table 4), ARGO-C leads the chart for the majority of the states (ARGO-C outperforms other models for 42 states in terms of MSE). Notably, after including the irregular flu seasons since COVID (2014-2023, Table S2), ARGO-C shows even more remarkable strength over benchmarks (ARGO-C outperforms other models for 47 states in terms of MSE). To further confirm the reliability of ARGO-C, we report the actual coverage rate of the 95% prediction interval given by ARGO-C for each state in Table S3: Overall, the average coverage rate over the 51 states is 92.6%, close to the nominal level.

Discussion

In summary, to account for the interconnection among the Internet search data, we proposed an innovative statistical learning framework, ARGO-C. By applying ARGO-C to both national and localized flu tracking, we observe that ARGO-C enhances the original ARGO/ARGO2/ARGOX framework by effectively and efficiently extracting and utilizing the inherent grouping structure of Google search terms.

The first step model of our proposed ARGO-C identifies the interconnection structure among Google search terms through clustering. In general, any clustering technique may be employed at this stage, and for each given method, several fitting criteria may also be considered, such as different choices of the distance metric and/or other tuning parameters (e.g., linkage function for hierarchical clustering). Specifically for the task of flu prediction, we recommend using hierarchical clustering based on the correlation distance metric and the average linkage function. This is through our empirical explorations of multiple classic clustering methods, including hierarchical clustering, k-means, and PAM [85] with various tuning configurations (linkage, distance metric, etc.) In practice, one may also explore a few different clustering methods and choose the one that gives the most interpretable results in the given context. It is also possible to realize clustering in a more dynamic fashion, updating the search term clusters periodically over time, which may possibly improve the accuracy of the %ILI predictions in the following modeling step. In addition, we also explored various criteria to determine the number of clusters K . Our final choice of K was a joint decision based on the within-cluster distance with consideration of model interpretability. In practice, we recommend choosing a relatively large K for effective incorporation of the clustering information.

We acknowledge that there are limitations of utilizing Google search data into %ILI predictions during the COVID-19 pandemic or the post-COVID period in the near future. First of all, %ILI is only a proxy for the actual flu incidence in the population. Since the seasonal flu and COVID-19 share many symptoms in common, the reported %ILI may as well include visits due to COVID-19. Consequently, the %ILI predictions may be largely influenced by simultaneous COVID-19 seasonal surges or underlying COVID-19 cases. In the main focus of our data analysis, we excluded the period that was possibly contaminated by COVID-19, as we suspect that the %ILI target and the existing set of Google search terms may not well represent flu activities during that period. Nevertheless, the %ILI surveillance data can still provide valuable insights on the general trend of influenza activity [84] (so, we also presented the results during the post-COVID period in the Supplementary Information, which also showcases the robustness of ARGO-C). It could be among our future endeavors to update the Google search terms after accounting for the effect of COVID-19, and/or to target alternative flu indicators [41], such as laboratory-confirmed influenza hospital admissions [86]. In addition, it will also be an interesting future project to explore the possibility of predicting seasonal flu cases and COVID-19 cases simultaneously by accounting for their interactive effects on each other [67].

Although not presented in this paper, ARGO-C is highly robust and can be easily adapted to digital tracking of other diseases or social/economic trends (as we have done to ILI prediction at various geographical resolution). We hope that our proposed framework can improve the real-time tracking of various infectious diseases and potentially contribute to the area of public health by saving more people's lives.

Acknowledgements

The work by A.H. was supported by the Finnerty Fund for summer undergraduate research in the Department of Mathematics and Statistics, Williams College, and by the College's DFRC fund for undergraduate research assistant.

Author contributions statement

S.N. and Q.W. conceived the research; S.N. and A.H. conducted the data analysis and experiments; S.N., A.H., and Q.W. analyzed the results; S.N. and Q.W. wrote the paper. All authors reviewed the manuscript.

Data availability statement

The datasets generated and analysed during the current study, as well as the codes generating the results, are available at <https://github.com/shaoyangning/argo-c>.

Additional information

Competing Interests

The authors declare no competing interests.

References

1. Polgreen, P. M., Chen, Y., Pennock, D. M., Nelson, F. D. & Weinstein, R. A. Using internet searches for influenza surveillance. *Clin. infectious diseases* **47**, 1443–1448 (2008).
2. Ginsberg, J. *et al.* Detecting influenza epidemics using search engine query data. *Nature* **457**, 1012–1014 (2009).
3. Althouse, B. M., Ng, Y. Y. & Cummings, D. A. Prediction of dengue incidence using search query surveillance. *PLoS neglected tropical diseases* **5**, e1258 (2011).
4. Chan, E. H., Sahai, V., Conrad, C. & Brownstein, J. S. Using web search query data to monitor dengue epidemics: a new model for neglected tropical disease surveillance. *PLoS Neglected Trop. Dis.* **5**, e1206 (2011).
5. Murdoch, T. B. & Detsky, A. S. The inevitable application of big data to health care. *Jama* **309**, 1351–1352 (2013).
6. Lee, K., Agrawal, A. & Choudhary, A. Real-time disease surveillance using twitter data: demonstration on flu and cancer. In *Proceedings of the 19th ACM SIGKDD international conference on Knowledge discovery and data mining*, 1474–1477 (2013).
7. Khoury, M. J. & Ioannidis, J. P. Big data meets public health. *Science* **346**, 1054–1055 (2014).
8. Rufai, S. R. & Bunce, C. World leaders' usage of twitter in response to the covid-19 pandemic: a content analysis. *J. public health* **42**, 510–516 (2020).
9. Effenberger, M. *et al.* Association of the covid-19 pandemic with internet search volumes: a google trendstm analysis. *Int. J. Infect. Dis.* **95**, 192–197 (2020).
10. Aiello, A. E., Renson, A. & Zivich, P. Social media-and internet-based disease surveillance for public health. *Annu. review public health* **41**, 101 (2020).
11. Lampos, V. *et al.* Tracking covid-19 using online search. *NPJ digital medicine* **4**, 17 (2021).
12. Ettredge, M., Gerdes, J. & Karuga, G. Using web-based search data to predict macroeconomic statistics. *Commun. ACM* **48**, 87–92 (2005).
13. Goel, S., Hofman, J. M., Lahaie, S., Pennock, D. M. & Watts, D. J. Predicting consumer behavior with web search. *Proc. Natl. Acad. Sci.* **107**, 17486–17490 (2010).
14. McLaren, N. & Shanbhogue, R. Using internet search data as economic indicators. *Bank Engl. Q. Bull.* Q2 (2011).
15. Bollen, J., Mao, H. & Zeng, X. Twitter mood predicts the stock market. *J. computational science* **2**, 1–8 (2011).
16. Choi, H. & Varian, H. Predicting the present with google trends. *Econ. Rec.* **88**, 2–9 (2012).
17. Preis, T., Moat, H. S. & Stanley, H. E. Quantifying trading behavior in financial markets using google trends. *Sci. reports* **3**, 1–6 (2013).

18. Scott, S. L. & Varian, H. R. Predicting the present with bayesian structural time series. *Int. J. Math. Model. Numer. Optimisation* **5**, 4–23 (2014).
19. Einav, L. & Levin, J. Economics in the age of big data. *Science* **346**, 1243089 (2014).
20. Wu, L. & Brynjolfsson, E. The future of prediction: How google searches foreshadow housing prices and sales. In *Economic analysis of the digital economy*, 89–118 (University of Chicago Press, 2015).
21. Vicente, M. R., López-Menéndez, A. J. & Pérez, R. Forecasting unemployment with internet search data: Does it help to improve predictions when job destruction is skyrocketing? *Technol. Forecast. Soc. Chang.* **92**, 132–139 (2015).
22. Scott, S. L. & Varian, H. R. Bayesian variable selection for nowcasting economic time series. In *Economic analysis of the digital economy*, 119–135 (University of Chicago Press, 2015).
23. Yi, D., Ning, S., Chang, C.-J. & Kou, S. Forecasting unemployment using internet search data via prism. *J. Am. Stat. Assoc.* **116**, 1662–1673 (2021).
24. Manyika, J. *et al.* *Big data: The next frontier for innovation, competition, and productivity* (McKinsey & Company, 2011).
25. McAfee, A. & Brynjolfsson, E. Big data: The management revolution. *Harv. Bus. Rev.* **90**, 60–68 (2012).
26. Chen, H., Chiang, R. H. & Storey, V. C. Business intelligence and analytics: From big data to big impact. *MIS Q.* **36**, 1165–1188 (2012).
27. Risteski, D. & Davcev, D. Can we use daily internet search query data to improve predicting power of egarch models for financial time series volatility. In *Proceedings of the International Conference on Computer Science and Information Systems (ICSI'S'2014), October 17–18, 2014, Dubai (United Arab Emirates)* (2014).
28. Zhu, C. Big data as a governance mechanism. *The Rev. Financial Stud.* **32**, 2021–2061 (2019).
29. Kim, G.-H., Trimis, S. & Chung, J.-H. Big-data applications in the government sector. *Commun. ACM* **57**, 78–85 (2014).
30. Bennett, J. & Lanning, S. The netflix prize. In *Proceedings of KDD Cup and Workshop 2007* (2007).
31. Santillana, M., Zhang, D. W., Althouse, B. M. & Ayers, J. W. What can digital disease detection learn from (an external revision to) google flu trends? *Am. journal preventive medicine* **47**, 341–347 (2014).
32. Wójcik, O. P., Brownstein, J. S., Chunara, R. & Johansson, M. A. Public health for the people: participatory infectious disease surveillance in the digital age. *Emerg. themes epidemiology* **11**, 1–7 (2014).
33. Bates, M. Tracking disease: digital epidemiology offers new promise in predicting outbreaks. *IEEE pulse* **8**, 18–22 (2017).
34. Li, C. *et al.* Retrospective analysis of the possibility of predicting the covid-19 outbreak from internet searches and social media data, china, 2020. *Eurosurveillance* **25**, 2000199 (2020).
35. Ma, S., Sun, Y. & Yang, S. Using internet search data to forecast covid-19 trends: A systematic review. *Analytics* **1**, 210–227 (2022).
36. Ma, S. & Yang, S. Covid-19 forecasts using internet search information in the united states. *Sci. Reports* **12**, 11539 (2022).
37. Iuliano, A. D. *et al.* Estimates of global seasonal influenza-associated respiratory mortality: a modelling study. *The Lancet* **391**, 1285–1300 (2018).
38. Molinari, N.-A. M. *et al.* The annual impact of seasonal influenza in the US: measuring disease burden and costs. *Vaccine* **25**, 5086–5096 (2007).
39. Lipsitch, M., Finelli, L., Heffernan, R. T., Leung, G. M. & Redd; for the 2009 H1N1 Surveillance Group, S. C. Improving the evidence base for decision making during a pandemic: the example of 2009 influenza a/h1n1. *Biosecurity bioterrorism: biodefense strategy, practice, science* **9**, 89–115 (2011).
40. Nsoesie, E. O., Brownstein, J. S., Ramakrishnan, N. & Marathe, M. V. A systematic review of studies on forecasting the dynamics of influenza outbreaks. *Influ. other respiratory viruses* **8**, 309–316 (2014).

- 41.** Chretien, J.-P., George, D., Shaman, J., Chitale, R. A. & McKenzie, F. E. Influenza forecasting in human populations: a scoping review. *PloS One* **9**, e94130 (2014).
- 42.** Brownstein, J. S., Freifeld, C. C. & Madoff, L. C. Digital disease detection—harnessing the web for public health surveillance. *The New Engl. journal medicine* **360**, 2153 (2009).
- 43.** Dalton, C. *et al.* Flutracking: a weekly australian community online survey of influenza-like illness in 2006, 2007 and 2008. *Commun. diseases intelligence quarterly report* **33**, 316–322 (2009).
- 44.** Achrekar, H., Gandhe, A., Lazarus, R., Yu, S.-H. & Liu, B. Predicting flu trends using twitter data. In *2011 IEEE conference on computer communications workshops (INFOCOM WKSHPS)*, 702–707 (IEEE, 2011).
- 45.** Yuan, Q. *et al.* Monitoring influenza epidemics in china with search query from baidu. *PloS one* **8**, e64323 (2013).
- 46.** Paul, M. J., Dredze, M. & Broniatowski, D. Twitter improves influenza forecasting. *PLoS currents* **6** (2014).
- 47.** McIver, D. J. & Brownstein, J. S. Wikipedia usage estimates prevalence of influenza-like illness in the united states in near real-time. *PLoS computational biology* **10**, e1003581 (2014).
- 48.** Santillana, M., Nsoesie, E. O., Mekaru, S. R., Scales, D. & Brownstein, J. S. Using clinicians' search query data to monitor influenza epidemics. *Clin. Infect. Dis.* **59**, 1446–1450 (2014).
- 49.** Paolotti, D. *et al.* Web-based participatory surveillance of infectious diseases: the influenzanet participatory surveillance experience. *Clin. Microbiol. Infect.* **20**, 17–21 (2014).
- 50.** Smolinski, M. S. *et al.* Flu near you: crowdsourced symptom reporting spanning 2 influenza seasons. *Am. journal public health* **105**, 2124–2130 (2015).
- 51.** Santillana, M. *et al.* Combining search, social media, and traditional data sources to improve influenza surveillance. *PLoS Comput. Biol* **11**, e1004513 (2015).
- 52.** Yang, S. *et al.* Using electronic health records and internet search information for accurate influenza forecasting. *BMC infectious diseases* **17**, 1–9 (2017).
- 53.** Bradshaw, B. *et al.* Influenza surveillance using wearable mobile health devices. *Online J. Public Heal. Informatics* **11** (2019).
- 54.** Hassan Zadeh, A., Zolbanin, H. M., Sharda, R. & Delen, D. Social media for nowcasting flu activity: Spatio-temporal big data analysis. *Inf. Syst. Front.* **21**, 743–760 (2019).
- 55.** Viboud, C. & Santillana, M. Fitbit-informed influenza forecasts. *The Lancet Digit. Heal.* **2**, e54–e55 (2020).
- 56.** Cook, S., Conrad, C., Fowlkes, A. L. & Mohebbi, M. H. Assessing google flu trends performance in the united states during the 2009 influenza virus a (h1n1) pandemic. *PloS one* **6**, e23610 (2011).
- 57.** Pervaiz, F., Pervaiz, M., Rehman, N. A., Saif, U. *et al.* Flubreaks: early epidemic detection from google flu trends. *J. medical Internet research* **14**, e2102 (2012).
- 58.** Butler, D. When google got flu wrong: Us outbreak foxes a leading web-based method for tracking seasonal flu. *Nature* **494**, 155–157 (2013).
- 59.** Olson, D. R., Konty, K. J., Paladini, M., Viboud, C. & Simonsen, L. Reassessing google flu trends data for detection of seasonal and pandemic influenza: a comparative epidemiological study at three geographic scales. *PLoS computational biology* **9**, e1003256 (2013).
- 60.** Lazer, D., Kennedy, R., King, G. & Vespignani, A. The parable of google flu: traps in big data analysis. *science* **343**, 1203–1205 (2014).
- 61.** Yang, S., Santillana, M. & Kou, S. C. Accurate estimation of influenza epidemics using google search data via argo. *Proc. Natl. Acad. Sci.* **112**, 14473–14478 (2015).
- 62.** Yang, S. *et al.* Advances in using internet searches to track dengue. *PLoS computational biology* **13**, e1005607 (2017).

- 63.** Ning, S., Yang, S. & Kou, S. Accurate regional influenza epidemics tracking using internet search data. *Sci. reports* **9**, 5238 (2019).
- 64.** Lu, F. S., Hattab, M. W., Clemente, C. L., Biggerstaff, M. & Santillana, M. Improved state-level influenza nowcasting in the united states leveraging internet-based data and network approaches. *Nat. communications* **10**, 147 (2019).
- 65.** Yang, S., Ning, S. & Kou, S. Use internet search data to accurately track state level influenza epidemics. *Sci. reports* **11**, 1–10 (2021).
- 66.** Wang, T., Ma, S., Baek, S. & Yang, S. Covid-19 hospitalizations forecasts using internet search data. *Sci. Reports* **12**, 9661 (2022).
- 67.** Ma, S., Ning, S. & Yang, S. Joint covid-19 and influenza-like illness forecasts in the united states using internet search information. *Commun. Medicine* **3**, 39 (2023).
- 68.** Tibshirani, R. Regression shrinkage and selection via the lasso. *J. Royal Stat. Soc. B* **58**, 267–288 (1996).
- 69.** Hoerl, A. & Kennard, R. Ridge regression: biased estimation for nonorthogonal problems. *Technometrics* **12**, 55–67 (1970).
- 70.** Tibshirani, R. The lasso method for variable selection in the Cox model. *Stat. Medicine* **16**, 385–395 (1997).
- 71.** Lokhorst, J. The lasso and generalized linear models. Tech. Rep., University of Adelaide (1999).
- 72.** Roth, V. The generalized lasso. *IEEE Transactions on Neural Networks* **15**, 16–28 (2004).
- 73.** Zou, H. & Hastie, T. Regularization and variable selection via the elastic net. *J. Royal Stat. Soc. B* **67**, 301–320 (2005).
- 74.** Yuan, M. & Lin, Y. Model selection and estimation in regression with grouped variables. *J. Royal Stat. Soc. B* **68**, 49–67 (2006).
- 75.** Meier, L., van de Geer, S. & Bhlmann, P. The grouped lasso for logistic regression. *J. Royal Stat. Soc. B* **70**, 53–71 (2008).
- 76.** Simon, N., Friedman, J., Hastie, T. & Tibshirani, R. A sparse-group lasso. *J. Comput. Graph. Stat.* **22**, 231–245 (2013).
- 77.** Ward, J. H. Hierarchical grouping to optimize an objective function. *J. Am. Stat. Assoc.* **58**, 236–244 (1963).
- 78.** MacQueen, J. Classification and analysis of multivariate observations. In *5th Berkeley Symp. Math. Statist. Probability*, 281–297 (University of California Los Angeles LA USA, 1967).
- 79.** Banerjee, A. & Shan, H. *Model-Based Clustering*. In: Sammut, C., Webb, G.I. (eds) *Encyclopedia of Machine Learning*. (Springer, 2011).
- 80.** R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria (2022).
- 81.** Van Rossum, G., Drake, F. L. et al. *Python reference manual* (Centrum voor Wiskunde en Informatica Amsterdam, 1995).
- 82.** Rousseeuw, P. J. Silhouettes: a graphical aid to the interpretation and validation of cluster analysis. *J. computational applied mathematics* **20**, 53–65 (1987).
- 83.** Tibshirani, R., Walther, G. & Hastie, T. Estimating the number of clusters in a data set via the gap statistic. *J. Royal Stat. Soc. Ser. B (Statistical Methodol.)* **63**, 411–423 (2001).
- 84.** Center for Disease Control and Preventions. Flu activity & surveillance (2023). <https://www.cdc.gov/flu/weekly/overview.htm#ILINet>, Last accessed on 2023-04-13.
- 85.** Kaufman, L. Partitioning around medoids (program pam). *Find. groups data* **344**, 68–125 (1990).
- 86.** Center for Disease Control and Preventions. Flu activity & surveillance (2023). <https://www.cdc.gov/flu/weekly/flusight/index.html>, Last accessed on 2023-04-13.




Figure 1. Comparison of %ILI estimation between ARGO-C and other benchmarks. The evaluation is based on the national level %ILI in three accuracy metrics: RMSE, MAE, and correlation; the evaluation period is the overall period from March 29, 2009 to March 21, 2020 (excluding the influence of COVID-19). Detailed numbers can be found in Table 1.

	Whole period	GFT period	'09-'10	'10-'11	'11-'12	'12-'13	'13-'14	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20
RSME													
ARGO-C	0.246	0.283	0.457	0.233	0.153	0.489	0.152	0.206	0.149	0.270	0.204	0.171	0.425
GFT	–	0.770	0.437	0.376	0.493	2.221	0.346	0.316	–	–	–	–	–
VAR1	0.355	0.348	0.521	0.333	0.164	0.503	0.353	0.455	0.237	0.364	0.561	0.346	0.695
ARGO	0.261	0.294	0.454	0.336	0.132	0.455	0.184	0.244	0.159	0.333	0.179	0.204	0.436
naive	0.352	0.347	0.520	0.339	0.163	0.499	0.350	0.457	0.234	0.343	0.556	0.344	0.691
MAE													
ARGO-C	0.156	0.178	0.300	0.176	0.126	0.306	0.119	0.153	0.108	0.188	0.152	0.122	0.297
GFT	–	0.366	0.294	0.327	0.441	1.634	0.189	0.214	–	–	–	–	–
VAR1	0.210	0.199	0.322	0.256	0.136	0.314	0.211	0.283	0.191	0.267	0.377	0.280	0.538
ARGO	0.158	0.172	0.290	0.265	0.104	0.260	0.120	0.178	0.135	0.218	0.141	0.149	0.334
naive	0.210	0.201	0.323	0.259	0.135	0.325	0.212	0.290	0.187	0.256	0.384	0.282	0.533
Correlation													
ARGO-C	0.985	0.977	0.982	0.980	0.922	0.959	0.987	0.988	0.981	0.978	0.997	0.992	0.975
GFT	–	0.876	0.995	0.968	0.833	0.926	0.969	0.987	–	–	–	–	–
VAR1	0.968	0.960	0.967	0.955	0.885	0.922	0.921	0.938	0.932	0.950	0.965	0.956	0.933
ARGO	0.983	0.971	0.975	0.963	0.922	0.940	0.978	0.986	0.970	0.978	0.997	0.988	0.974
naive	0.968	0.961	0.968	0.954	0.887	0.924	0.923	0.939	0.935	0.956	0.964	0.957	0.933

Table 1. Comparison of national %ILI estimation between ARGO-C and other benchmarks. The evaluation is based on the national level %ILI in multiple periods and multiple metrics. RMSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period. Methods considered here include ARGO-C, VAR1, GFT, the original ARGO, and the naive method. All comparisons are conducted on the original scale of the CDC's %ILI. The whole period is March 29, 2009 to March 21, 2020, excluding the period with COVID influence. The second column (GFT period) is the period when the estimation by GFT is available, i.e., March 29, 2009 to July 11, 2015. The remaining columns contain yearly regular flu seasons, from week 40 to week 20 next year, as defined by CDC's Morbidity and Mortality Weekly Report. (The 19'-20' season is up to March 21, 2020).

	Region 1	Region 2	Region 3	Region 4	Region 5	Region 6	Region 7	Region 8	Region 9	Region 10
RMSE										
ARGO-C	0.301	0.399	0.338	0.334	0.281	0.584	0.423	0.310	0.364	0.372
ARGO2	0.308	0.410	0.367	0.347	0.302	0.604	0.454	0.319	0.376	0.382
VAR1	0.369	0.487	0.476	0.462	0.391	0.730	0.565	0.382	0.413	0.438
naive	0.366	0.487	0.473	0.460	0.389	0.722	0.558	0.374	0.415	0.432
MAE										
ARGO-C	0.162	0.267	0.210	0.203	0.159	0.353	0.247	0.175	0.224	0.228
ARGO2	0.164	0.275	0.226	0.212	0.167	0.371	0.266	0.180	0.229	0.230
VAR1	0.191	0.303	0.266	0.264	0.208	0.440	0.313	0.206	0.253	0.247
naive	0.192	0.304	0.267	0.266	0.211	0.440	0.316	0.205	0.255	0.256
Correlation										
ARGO-C	0.965	0.969	0.971	0.978	0.972	0.973	0.972	0.973	0.946	0.955
ARGO2	0.964	0.968	0.967	0.976	0.969	0.972	0.969	0.972	0.943	0.953
VAR1	0.946	0.954	0.941	0.957	0.945	0.957	0.948	0.958	0.931	0.938
naive	0.948	0.955	0.943	0.958	0.946	0.959	0.950	0.960	0.932	0.941

Table 2. Comparison of regional %ILI estimation between ARGO-C and other benchmarks. The evaluation is based on the %ILI at 10 HHS regional level in multiple metrics. RMSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period. Methods considered here include ARGO-C, VAR1, the original ARGO2, and the naive method. All comparisons are conducted on the original scale of the CDC's %ILI. The evaluation period is March 29, 2009 to March 21, 2020, excluding the period with COVID-19 influence.

Region 1	Region 2	Region 3	Region 4	Region 5	Region 6	Region 7	Region 8	Region 9	Region 10
0.935	0.930	0.953	0.946	0.953	0.942	0.958	0.939	0.944	0.951

Table 3. Actual coverage of prediction intervals by ARGO-C for regional %ILI prediction. The coverage is for 95% nominal level. The average coverage over the ten US HHS regions is 94.5%. The evaluation period is March 29, 2009 to March 21, 2020, excluding the period with COVID-19 influence.

	Whole period	GFT period	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20
RMSE							
ARGO-C	0.569	0.714	0.473	0.662	0.680	0.548	0.877
ARGOX	0.578	0.733	0.474	0.673	0.680	0.553	0.936
VAR1	1.855	2.300	1.350	1.992	2.703	1.822	2.785
GFT	—	1.546	—	—	—	—	—
naive	0.680	0.849	0.507	0.743	0.894	0.659	1.063
MAE							
ARGO-C	0.336	0.402	0.321	0.412	0.440	0.357	0.561
ARGOX	0.339	0.410	0.318	0.421	0.438	0.362	0.590
VAR1	1.186	1.489	1.032	1.396	1.738	1.277	2.044
GFT	—	1.001	—	—	—	—	—
naive	0.388	0.467	0.340	0.464	0.558	0.443	0.691
Correlation							
ARGO-C	0.950	0.911	0.821	0.872	0.934	0.921	0.905
ARGOX	0.948	0.907	0.824	0.869	0.935	0.920	0.896
VAR1	0.728	0.696	0.636	0.660	0.785	0.759	0.700
GFT	—	0.902	—	—	—	—	—
naive	0.930	0.877	0.803	0.842	0.899	0.890	0.873

Table 4. Comparison of state-level %ILI estimation between ARGO-C and other benchmarks. The evaluation is based on the average of 51 US state/district in multiple periods and multiple metrics. RMSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period. Methods considered here include ARGO-C, VAR1, GFT, the original ARGOX, and the naive method. All comparisons are conducted on the original scale of the CDC's %ILI. The whole period is January 10, 2014 (first available estimate by ARGOX) to March 21, 2020, excluding the period with COVID influence. The second column (GFT period) is the period when the estimation by GFT is available, i.e., January 10, 2014 to July 11, 2015. Each regular flu season ('14-'15 overlaps with GFT period) is from week 40 to week 20 next year, as defined by CDC's Morbidity and Mortality Weekly Report. (The 19'-20' season is up to March 21, 2020).

Supplementary Information

	Overall '09-'23	'20-'21	'21-'22	'22-'23
RSME				
ARGO-C	0.252	0.113	0.224	0.490
GFT	–	–	–	–
VAR1	0.359	0.081	0.357	0.769
ARGO	0.258	0.100	0.232	0.487
naive	0.355	0.082	0.358	0.734
MAE				
ARGO-C	0.161	0.089	0.181	0.429
GFT	–	–	–	–
VAR1	0.211	0.066	0.245	0.644
ARGO	0.158	0.086	0.168	0.374
naive	0.210	0.067	0.239	0.591
Correlation				
ARGO-C	0.984	0.927	0.983	0.975
GFT	–	–	–	–
VAR1	0.968	0.941	0.899	0.909
ARGO	0.983	0.940	0.962	0.969
naive	0.968	0.939	0.900	0.907

Table S1. Comparison of % ILI estimation between ARGO-C and other benchmarks at the national level, for flu seasons since COVID-19. The evaluation is based at the national level %ILI in multiple periods and multiple metrics. RMSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period. Methods considered here include ARGO-C, VAR1, GFT, the original ARGO, and the naive method. All comparisons are conducted on the original scale of the CDC's %ILI. The overall period '09-'23 is March 29, 2009 to January 28, 2023, including the period since COVID. Each regular flu season is from week 40 to week 20 next year, as defined by CDC's Morbidity and Mortality Weekly Report. (The '22-'23 season is up to January 28, 2023).

Step 1: Clustering of Google Search Terms




Figure S1. Clustering of the 71 Google search terms, collected by March 29, 2009. 53 clusters were identified. Hierarchical clustering with average linkage and correlation distance metric were used. The clustering was conducted based on the time series of Google search data from January 10, 2004 (the earliest available Google Trends data) to March 29, 2009 (the earliest prediction date by these 71 terms).




Figure S2. Clustering of the 161 Google search terms, collected by May 22, 2010. 45 clusters were identified. Hierarchical clustering with average linkage and correlation distance metric were adopted. The clustering was conducted based on the time series of Google search data from January 10, 2004 (the earliest available Google Trends data) to May 22, 2010 (the earliest prediction date by these 161 terms).




Figure S3. Traceplots of clustered predictors included in the ARGO-C model at the national level over time. The heatmaps indicate whether each predictor was included in the predictive ARGO-C model at each week. Three exemplary clusters were highlighted. Each of the top two clusters contains three search terms (identified among 71 search terms by March 29, 2009, used for predictions before 2010), while the last one includes four search terms (identified among 161 topics/terms by May 22, 2010 and used for predictions since 2010). The entire cluster was penalized and excluded from the model when an entire column in the traceplot is colored grey.

	Overall '14-'23	post-COVID	'20-'21	'21-'22	'22-'23
RMSE					
ARGO-C	0.564	0.554	0.287	0.480	1.048
ARGOX	0.576	0.574	0.283	0.488	1.101
VAR1	1.789	1.660	0.736	1.368	3.398
GFT	—	—	—	—	—
naive	0.662	0.626	0.279	0.551	1.224
MAE					
ARGO-C	0.325	0.304	0.186	0.306	0.685
ARGOX	0.329	0.309	0.182	0.305	0.735
VAR1	1.114	0.979	0.511	0.954	2.364
GFT	—	—	—	—	—
naive	0.370	0.336	0.166	0.346	0.849
Correlation					
ARGO-C	0.947	0.933	0.703	0.862	0.890
ARGOX	0.944	0.929	0.709	0.855	0.877
VAR1	0.675	0.719	0.478	0.563	0.591
GFT	—	—	—	—	—
naive	0.929	0.919	0.710	0.816	0.844

Table S2. Comparison of % ILI estimation between ARGO-C and other benchmarks at the state level, for flu seasons since COVID-19. The evaluation is based on the average of 51 US state/district in multiple periods and multiple metrics. RMSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period. Methods considered here include ARGO-C, VAR1, GFT, the original ARGOX, and the naive method. All comparisons are conducted on the original scale of the CDC's %ILI. The overall period '14-'23 is January 10, 2014 (first available estimate by ARGO framework) to January 28, 2023, including the period since COVID. The post-COVID period is the period since COVID, March 21, 2020 to January 28, 2023. Each regular flu season is from week 40 to week 20 next year, as defined by CDC's Morbidity and Mortality Weekly Report. (The '22-'23 season is up to January 28, 2023).

AL	AK	AZ	AR	CA	CO	CT	DE	DC	GA	HI	ID	IL	IN	IA
0.931	0.926	0.915	0.929	0.947	0.903	0.906	0.947	0.908	0.933	0.942	0.933	0.931	0.924	0.931
KS	KY	LA	ME	MD	MA	MI	MN	MS	MO	MT	NE	NV	NH	NJ
0.938	0.894	0.922	0.917	0.933	0.919	0.940	0.942	0.938	0.935	0.935	0.910	0.903	0.908	0.929
NM	NY	NC	ND	OH	OK	OR	PA	RI	SC	SD	TN	TX	UT	VT
0.915	0.899	0.952	0.949	0.924	0.924	0.926	0.940	0.919	0.938	0.940	0.915	0.933	0.931	0.945
VA	WA	WV	WI	WY	NYC									
0.954	0.908	0.917	0.945	0.885	0.915									

Table S3. Actual coverage of prediction intervals by ARGO-C for state-level %ILI prediction. The coverage is for 95% nominal level. The average coverage over 51 states/city/district is 92.6%. The evaluation period is January 10, 2014 to March 21, 2020, excluding the period with COVID-19 influence.

acute bronchitis	body temperature	break a fever	bronchitis
cold or flu	cold vs flu	cough fever	cure the flu
dangerous fever	fever cough	fever flu	fever reducer
flu contagious period	flu contagious	flu duration	flu fever
flu how long	flu in children	flu incubation	flu medicine
flu or cold	flu report	flu test	flu treatment
flu treatments	flu vs cold	get over the flu	high fever
how long is the flu contagious	how long is the flu	how to treat the flu	incubation period for the flu
influenza a and b	influenza a	influenza contagious	influenza incubation period
influenza incubation	influenza symptoms	influenza treatment	influenza type a
is flu contagious	low body	normal body temperature	normal body
over the counter flu	painful cough	pneumonia	reduce a fever
remedies for the flu	robitussin	signs of the flu	sinus infections
sinus	strep	symptoms of bronchitis	symptoms of flu
symptoms of influenza	symptoms of pneumonia	symptoms of the flu	treat flu
treat the flu	treating flu	treating the flu	treatment for flu
treatment for the flu	tussin	tussionex	type a flu
type a influenza	upper respiratory	walking pneumonia	
a influenza	braun thermoscan	chest cold	cold and flu
cold versus flu	contagious flu	cure flu	do i have the flu
ear thermometer	early flu symptoms	expectorant	exposed to flu
fever breaks	fight the flu	flu a symptoms	flu and cold
flu and fever	flu care	flu children	flu complications
flu cough	flu germs	flu headache	flu how long are you contagious
flu incubation period	flu lasts	flu length	flu recovery
flu relief	flu remedies	flu remedy	flu reports
flu symptoms	flu texas	flu versus cold	flu
get rid of the flu	having the flu	how long am i contagious with the flu	how long contagious
how long does flu last	how long does the flu last	how long flu	how long is flu contagious
how to break a fever	how to bring a fever down	how to get rid of the flu	how to treat flu
how to treat the flu at home	human temperature	i have the flu	incubation period for flu
medicine for flu	medicine for the flu	oscillococcinum	over the counter flu medicine
rapid flu	reduce fever	remedies for flu	respiratory flu
signs of flu	strep throat	taking temperature	tessalon
the flu virus	the flu	thermoscan	type a flu symptoms
what to do if you have the flu			
Influenza vaccine	Influenza	Fever	Influenza A virus
Influenza B virus	Common cold	Cough	Sore throat
Virus	Avian influenza	Spanish flu	Headache
Nausea	Flu season	Oseltamivir	Nasal congestion
Canine influenza	Rapid influenza diagnostic test	Theraflu	Dextromethorphan
Rhinorrhea			

Table S4. All search query terms used in this study. The first 71 terms were collected on March 29, 2009, the remaining terms/topics were identified on May 22, 2010. The last 21 terms separated by a horizontal line from the first 140 terms were “Related topics” from Google Trends.

	'09-'23	'09-'15	'09-'10	'10-'11	'11-'12	'12-'13	'13-'14	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE																
ARGO-C	0.307	0.349	0.903	0.174	0.115	0.253	0.214	0.321	0.183	0.214	0.256	0.180	0.432	0.092	0.297	0.784
ARGO2	0.316	0.353	0.904	0.172	0.104	0.284	0.216	0.330	0.190	0.225	0.273	0.187	0.450	0.096	0.306	0.829
VAR1	0.382	0.409	1.069	0.194	0.104	0.372	0.214	0.371	0.212	0.276	0.440	0.271	0.636	0.067	0.365	1.011
GFT	–	0.941	0.466	0.237	0.245	2.819	0.395	0.418	–	–	–	–	–	–	–	–
naive	0.377	0.406	1.062	0.193	0.105	0.362	0.214	0.360	0.213	0.264	0.436	0.271	0.628	0.068	0.360	0.970
MAE																
ARGO-C	0.164	0.169	0.486	0.133	0.095	0.157	0.171	0.203	0.141	0.169	0.186	0.141	0.300	0.069	0.209	0.548
ARGO2	0.168	0.171	0.487	0.127	0.087	0.163	0.178	0.204	0.147	0.173	0.188	0.145	0.306	0.076	0.221	0.597
VAR1	0.198	0.187	0.569	0.147	0.086	0.221	0.182	0.213	0.170	0.199	0.323	0.186	0.488	0.052	0.282	0.725
GFT	–	0.327	0.267	0.195	0.176	1.330	0.334	0.322	–	–	–	–	–	–	–	–
naive	0.198	0.188	0.583	0.143	0.087	0.224	0.180	0.221	0.165	0.193	0.324	0.192	0.497	0.053	0.278	0.690
Correlation																
ARGO-C	0.968	0.934	0.913	0.956	0.630	0.960	0.885	0.931	0.938	0.964	0.984	0.983	0.975	0.949	0.931	0.943
ARGO2	0.966	0.933	0.913	0.955	0.690	0.951	0.880	0.929	0.933	0.960	0.983	0.982	0.975	0.951	0.923	0.935
VAR1	0.949	0.909	0.876	0.931	0.689	0.915	0.882	0.907	0.921	0.929	0.951	0.961	0.942	0.948	0.888	0.900
GFT	–	0.768	0.988	0.911	0.699	0.831	0.786	0.917	–	–	–	–	–	–	–	–
naive	0.951	0.914	0.884	0.932	0.693	0.919	0.884	0.914	0.923	0.936	0.951	0.961	0.944	0.947	0.892	0.899

Table S5. Comparison of different methods for regional level %ILI estimation in Region 1. The RMSE, MAE, and correlation measures are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	'09-'23	'09-'15	'09-'10	'10-'11	'11-'12	'12-'13	'13-'14	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE																
ARGO-C	0.446	0.418	0.646	0.385	0.134	0.340	0.337	0.374	0.354	0.540	0.599	0.228	0.953	0.229	0.552	0.885
ARGO2	0.458	0.430	0.647	0.394	0.125	0.395	0.330	0.415	0.352	0.577	0.588	0.241	0.929	0.230	0.579	0.990
VAR1	0.544	0.480	0.793	0.415	0.120	0.498	0.366	0.496	0.372	0.639	0.880	0.333	1.250	0.222	0.665	1.078
GFT	–	1.037	0.463	1.093	0.510	2.370	0.665	0.662	–	–	–	–	–	–	–	–
naive	0.541	0.482	0.799	0.421	0.121	0.498	0.367	0.489	0.381	0.627	0.868	0.333	1.229	0.225	0.667	1.047
MAE																
ARGO-C	0.280	0.268	0.430	0.272	0.110	0.242	0.285	0.314	0.292	0.442	0.429	0.186	0.567	0.174	0.412	0.626
ARGO2	0.288	0.275	0.434	0.282	0.101	0.260	0.283	0.345	0.296	0.460	0.439	0.193	0.575	0.177	0.417	0.684
VAR1	0.325	0.293	0.450	0.313	0.097	0.347	0.308	0.355	0.307	0.462	0.577	0.274	0.896	0.177	0.462	0.822
GFT	–	0.632	0.364	0.936	0.440	1.310	0.611	0.487	–	–	–	–	–	–	–	–
naive	0.323	0.297	0.460	0.304	0.097	0.358	0.309	0.360	0.301	0.449	0.572	0.275	0.879	0.180	0.442	0.796
Correlation																
ARGO-C	0.969	0.950	0.944	0.938	0.664	0.968	0.848	0.951	0.921	0.946	0.971	0.974	0.950	0.768	0.916	0.937
ARGO2	0.968	0.947	0.946	0.934	0.693	0.959	0.849	0.945	0.922	0.936	0.974	0.971	0.958	0.767	0.903	0.921
VAR1	0.952	0.934	0.912	0.927	0.678	0.930	0.831	0.907	0.902	0.912	0.940	0.944	0.901	0.765	0.866	0.878
GFT	–	0.783	0.981	0.870	0.398	0.915	0.871	0.932	–	–	–	–	–	–	–	–
naive	0.954	0.935	0.914	0.927	0.684	0.931	0.831	0.910	0.902	0.917	0.940	0.945	0.904	0.763	0.867	0.883

Table S6. Comparison of different methods for regional level %ILI estimation in Region 2. The RMSE, MAE, and correlation measures are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	'09-'23	'09-'15	'09-'10	'10-'11	'11-'12	'12-'13	'13-'14	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE																
ARGO-C	0.324	0.393	0.755	0.264	0.175	0.597	0.230	0.458	0.219	0.367	0.379	0.194	0.295	0.128	0.303	0.500
ARGO2	0.350	0.425	0.799	0.291	0.181	0.667	0.243	0.539	0.232	0.403	0.381	0.217	0.342	0.125	0.335	0.564
VAR1	0.460	0.522	0.998	0.327	0.193	0.786	0.311	0.745	0.320	0.530	0.652	0.334	0.708	0.104	0.450	0.698
GFT	–	0.859	0.604	0.501	0.926	2.188	0.514	0.380	–	–	–	–	–	–	–	–
naive	0.457	0.522	0.993	0.337	0.196	0.785	0.312	0.745	0.323	0.506	0.645	0.329	0.699	0.106	0.454	0.673
MAE																
ARGO-C	0.201	0.241	0.458	0.208	0.143	0.396	0.173	0.286	0.176	0.277	0.272	0.152	0.218	0.092	0.209	0.361
ARGO2	0.213	0.255	0.484	0.237	0.143	0.421	0.188	0.339	0.178	0.306	0.293	0.173	0.249	0.097	0.214	0.396
VAR1	0.258	0.282	0.555	0.245	0.153	0.477	0.204	0.410	0.254	0.336	0.407	0.269	0.512	0.080	0.284	0.594
GFT	–	0.567	0.474	0.428	0.886	1.760	0.480	0.312	–	–	–	–	–	–	–	–
naive	0.258	0.286	0.555	0.256	0.153	0.482	0.210	0.430	0.260	0.319	0.411	0.262	0.513	0.081	0.281	0.548
Correlation																
ARGO-C	0.973	0.957	0.960	0.975	0.725	0.916	0.975	0.964	0.939	0.961	0.984	0.991	0.988	0.922	0.954	0.952
ARGO2	0.969	0.951	0.955	0.969	0.712	0.894	0.974	0.954	0.936	0.953	0.986	0.989	0.986	0.932	0.935	0.938
VAR1	0.944	0.923	0.924	0.950	0.671	0.852	0.924	0.886	0.868	0.905	0.944	0.960	0.924	0.866	0.891	0.892
GFT	–	0.865	0.991	0.953	0.732	0.891	0.975	0.985	–	–	–	–	–	–	–	–
naive	0.946	0.925	0.927	0.947	0.676	0.859	0.926	0.891	0.870	0.916	0.945	0.962	0.927	0.866	0.892	0.892

Table S7. Comparison of different methods for regional level %ILI estimation in Region 3. The RMSE, MAE, and correlation measures are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	'09-'23	'09-'15	'09-'10	'10-'11	'11-'12	'12-'13	'13-'14	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE																
ARGO-C	0.335	0.356	0.313	0.377	0.180	0.524	0.243	0.476	0.179	0.390	0.421	0.239	0.531	0.139	0.286	0.726
ARGO2	0.353	0.357	0.276	0.394	0.192	0.524	0.267	0.517	0.210	0.415	0.418	0.300	0.662	0.141	0.321	0.776
VAR1	0.463	0.422	0.286	0.477	0.203	0.568	0.386	0.728	0.281	0.505	0.817	0.483	0.923	0.098	0.466	0.994
GFT	–	0.821	0.522	0.908	0.270	2.023	0.407	0.358	–	–	–	–	–	–	–	–
naive	0.460	0.423	0.312	0.484	0.205	0.569	0.387	0.734	0.275	0.482	0.809	0.492	0.916	0.100	0.465	0.958
MAE																
ARGO-C	0.204	0.205	0.225	0.291	0.139	0.325	0.166	0.278	0.153	0.297	0.284	0.168	0.398	0.119	0.206	0.456
ARGO2	0.214	0.206	0.213	0.309	0.138	0.329	0.172	0.298	0.171	0.327	0.276	0.211	0.468	0.126	0.219	0.505
VAR1	0.267	0.234	0.219	0.357	0.158	0.351	0.241	0.416	0.208	0.386	0.529	0.367	0.699	0.080	0.308	0.790
GFT	–	0.474	0.462	0.740	0.233	1.468	0.319	0.255	–	–	–	–	–	–	–	–
naive	0.268	0.238	0.235	0.365	0.157	0.363	0.244	0.428	0.206	0.368	0.520	0.383	0.695	0.082	0.308	0.751
Correlation																
ARGO-C	0.977	0.963	0.986	0.966	0.830	0.931	0.973	0.950	0.978	0.963	0.987	0.985	0.974	0.880	0.973	0.919
ARGO2	0.975	0.962	0.987	0.962	0.806	0.923	0.967	0.945	0.974	0.959	0.988	0.975	0.958	0.882	0.960	0.907
VAR1	0.957	0.947	0.986	0.945	0.792	0.907	0.920	0.880	0.926	0.931	0.952	0.931	0.908	0.904	0.844	0.844
GFT	–	0.835	0.984	0.975	0.771	0.892	0.977	0.979	–	–	–	–	–	–	–	–
naive	0.958	0.948	0.986	0.944	0.793	0.909	0.922	0.881	0.929	0.937	0.951	0.930	0.911	0.916	0.906	0.847

Table S8. Comparison of different methods for regional level %ILI estimation in Region 4. The RMSE, MAE, and correlation measures are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	'09-'23	'09-'15	'09-'10	'

	'09-'23	'09-'15	'09-'10	'10-'11	'11-'12	'12-'13	'13-'14	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE																
ARGO-C	0.557	0.615	0.893	0.466	0.239	0.541	0.548	1.117	0.362	0.593	0.803	0.517	1.017	0.298	0.311	0.625
ARGO2	0.578	0.631	0.884	0.507	0.273	0.612	0.572	1.108	0.362	0.647	0.763	0.553	1.144	0.314	0.303	0.663
VAR1	0.685	0.713	0.977	0.664	0.309	0.759	0.759	1.160	0.441	0.762	1.143	0.795	1.315	0.249	0.451	0.785
GFT	–	1.356	0.899	0.639	0.908	3.607	0.770	1.146	–	–	–	–	–	–	–	–
naive	0.678	0.708	1.012	0.665	0.306	0.770	0.747	1.164	0.444	0.727	1.131	0.791	1.300	0.251	0.456	0.753
MAE																
ARGO-C	0.338	0.370	0.593	0.366	0.175	0.407	0.361	0.627	0.300	0.399	0.424	0.353	0.707	0.240	0.222	0.499
ARGO2	0.352	0.385	0.579	0.402	0.188	0.442	0.368	0.678	0.293	0.417	0.441	0.373	0.788	0.250	0.220	0.542
VAR1	0.410	0.421	0.580	0.477	0.212	0.506	0.493	0.732	0.364	0.555	0.759	0.608	0.978	0.164	0.304	0.648
GFT	–	0.714	0.618	0.439	0.848	2.567	0.544	0.661	–	–	–	–	–	–	–	–
naive	0.411	0.424	0.625	0.461	0.207	0.534	0.498	0.752	0.359	0.544	0.750	0.618	0.983	0.168	0.323	0.624
Correlation																
ARGO-C	0.973	0.959	0.961	0.974	0.909	0.967	0.963	0.896	0.890	0.972	0.981	0.984	0.954	0.691	0.949	0.952
ARGO2	0.971	0.958	0.962	0.969	0.879	0.958	0.959	0.899	0.886	0.968	0.982	0.980	0.942	0.675	0.946	0.947
VAR1	0.958	0.945	0.952	0.943	0.850	0.934	0.924	0.890	0.833	0.944	0.957	0.947	0.916	0.684	0.878	0.911
GFT	–	0.878	0.980	0.978	0.914	0.943	0.956	0.928	–	–	–	–	–	–	–	–
naive	0.960	0.947	0.954	0.944	0.855	0.933	0.927	0.893	0.837	0.950	0.957	0.949	0.920	0.681	0.877	0.916

Table S10. Comparison of different methods for regional level %ILI estimation in Region 6. The RMSE, MAE, and correlation measures are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	'09-'23	'09-'15	'09-'10	'10-'11	'11-'12	'12-'13	'13-'14	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE																
ARGO-C	0.423	0.463	0.921	0.387	0.298	0.470	0.263	0.377	0.211	0.448	0.461	0.487	0.584	0.260	0.268	0.781
ARGO2	0.453	0.493	0.969	0.402	0.324	0.535	0.309	0.459	0.212	0.489	0.510	0.515	0.709	0.266	0.255	0.840
VAR1	0.552	0.572	1.122	0.491	0.348	0.631	0.459	0.621	0.245	0.583	0.908	0.656	0.917	0.231	0.305	1.006
GFT	–	0.917	1.968	1.077	0.603	0.718	0.322	0.856	–	–	–	–	–	–	–	–
naive	0.546	0.569	1.111	0.495	0.343	0.637	0.468	0.623	0.246	0.555	0.903	0.651	0.903	0.236	0.312	0.964
MAE																
ARGO-C	0.244	0.261	0.600	0.302	0.229	0.290	0.179	0.242	0.166	0.300	0.314	0.357	0.405	0.140	0.205	0.481
ARGO2	0.260	0.280	0.641	0.312	0.242	0.313	0.203	0.302	0.164	0.319	0.342	0.375	0.488	0.157	0.186	0.516
VAR1	0.300	0.304	0.698	0.346	0.262	0.365	0.278	0.376	0.200	0.404	0.595	0.481	0.630	0.116	0.214	0.705
GFT	–	0.506	1.385	0.791	0.436	0.539	0.200	0.493	–	–	–	–	–	–	–	–
naive	0.303	0.313	0.716	0.354	0.259	0.390	0.298	0.412	0.195	0.397	0.600	0.467	0.630	0.119	0.224	0.658
Correlation																
ARGO-C	0.969	0.959	0.943	0.953	0.941	0.961	0.970	0.975	0.901	0.961	0.988	0.958	0.970	0.798	0.963	0.931
ARGO2	0.965	0.955	0.938	0.949	0.931	0.952	0.960	0.967	0.903	0.954	0.986	0.953	0.955	0.792	0.956	0.918
VAR1	0.946	0.936	0.913	0.916	0.917	0.929	0.899	0.921	0.855	0.930	0.951	0.913	0.924	0.817	0.928	0.876
GFT	–	0.846	0.985	0.960	0.953	0.934	0.964	0.963	–	–	–	–	–	–	–	–
naive	0.948	0.938	0.918	0.917	0.919	0.929	0.898	0.919	0.858	0.936	0.952	0.917	0.928	0.813	0.924	0.876

Table S11. Comparison of different methods for regional level %ILI estimation in Region 7. The RMSE, MAE, and correlation measures are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	'09-'23	'09-'15	'09-'10	'10-'11	'11-'12	'12-'13	'13-'14	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE																
ARGO-C	0.307	0.336	0.812	0.237	0.152	0.213	0.283	0.294	0.132	0.351	0.169	0.470	0.361	0.134	0.244	0.710
ARGO2	0.319	0.347	0.831	0.219	0.169	0.252	0.300	0.323	0.151	0.337	0.172	0.476	0.415	0.139	0.259	0.758
VAR1	0.392	0.410	0.985	0.226	0.180	0.341	0.369	0.389	0.189	0.363	0.260	0.562	0.727	0.088	0.317	0.919
GFT	–	0.622	1.502	0.382	0.387	0.912	0.282	0.455	–	–	–	–	–	–	–	–
naive	0.385	0.402	0.969	0.229	0.177	0.341	0.374	0.388	0.186	0.369	0.250	0.532	0.721	0.088	0.315	0.894
MAE																
ARGO-C	0.177	0.172	0.403	0.189	0.105	0.161	0.169	0.199	0.109	0.262	0.130	0.372	0.262	0.105	0.187	0.554
ARGO2	0.182	0.177	0.411	0.181	0.117	0.170	0.177	0.225	0.121	0.250	0.132	0.376	0.290	0.114	0.195	0.595
VAR1	0.212	0.196	0.496	0.156	0.130	0.234	0.217	0.227	0.150	0.269	0.193	0.427	0.516	0.072	0.242	0.734
GFT	–	0.315	0.863	0.273	0.339	0.506	0.234	0.289	–	–	–	–	–	–	–	–
naive	0.211	0.199	0.512	0.162	0.124	0.241	0.226	0.239	0.147	0.273	0.179	0.397	0.513	0.072	0.236	0.720
Correlation																
ARGO-C	0.976	0.959	0.947	0.952	0.905	0.983	0.921	0.968	0.969	0.824	0.986	0.952	0.985	0.871	0.939	0.957
ARGO2	0.974	0.957	0.945	0.961	0.885	0.978	0.910	0.964	0.960	0.833	0.986	0.950	0.979	0.870	0.921	0.950
VAR1	0.960	0.938	0.923	0.941	0.874	0.955	0.866	0.930	0.931	0.797	0.963	0.924	0.933	0.891	0.848	0.917
GFT	–	0.859	0.988	0.929	0.872	0.926	0.955	0.930	–	–	–	–	–	–	–	–
naive	0.961	0.941	0.927	0.941	0.877	0.955	0.867	0.931	0.935	0.803	0.963	0.930	0.935	0.889	0.848	0.915

Table S12. Comparison of different methods for regional level %ILI estimation in Region 8. The RMSE, MAE, and correlation measures are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	'09-'23	'09-'15	'09-'10	'10-'11	'11-'12	'12-'13	'13-'14	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE																
ARGO-C	0.347	0.437	0.665	0.358	0.382	0.474	0.297	0.284	0.224	0.199	0.404	0.181	0.306	0.135	0.229	0.639
ARGO2	0.359	0.453	0.654	0.353	0.394	0.525	0.319	0.330	0.244	0.186	0.413	0.178	0.288	0.134	0.250	0.700
VAR1	0.406	0.470	0.664	0.395	0.419	0.538	0.382	0.380	0.315	0.208	0.622	0.224	0.462	0.137	0.353	0.877
GFT	–	1.091	1.101	1.168	0.567	2.536	0.593	0.540	–	–	–	–	–	–	–	–
naive	0.405	0.475	0.676	0.382	0.438	0.544	0.385	0.382	0.312	0.210	0.606	0.222	0.461	0.139	0.358	0.828
MAE																
ARGO-C	0.213	0.275	0.502	0.261	0.308	0.325	0.204	0.201	0.172	0.142	0.265	0.132	0.230	0.109	0.183	0.453
ARGO2	0.217	0.282	0.488	0.270	0.321	0.339	0.214	0.219	0.186	0.149	0.266	0.127	0.209	0.109	0.184	0.480
VAR1	0.247	0.297	0.510	0.320	0.349	0.327	0.247	0.258	0.243	0.161	0.357	0.181	0.338	0.106	0.214	0.726
GFT	–	0.690	0.995	1.090	0.500	1.296	0.393	0.410	–	–	–	–	–	–	–	–
naive	0.247	0.299	0.514	0.298	0.368	0.354	0.253	0.260	0.244	0.167	0.365	0.182	0.333	0.107	0.221	0.656
Correlation																
ARGO-C	0.956	0.921	0.891	0.944	0.768	0.922	0.963	0.967	0.965	0.945	0.959	0.972	0.971	0.948	0.957	0.959
ARGO2	0.953	0.915	0.893	0.944	0.751	0.902	0.958	0.960	0.960	0.946	0.955	0.974	0.949	0.946	0.947	0.947
VAR1	0.940	0.909	0.887	0.922	0.728	0.894	0.925	0.930	0.930	0.933	0.897	0.957	0.929	0.943	0.887	0.895
GFT	–	0.806	0.945	0.905	0.844	0.900	0.937	0.917	–	–	–	–	–	–	–	–
naive	0.941	0.910	0.890	0.924	0.728	0.894	0.926	0.931	0.933	0.933	0.901	0.958	0.930	0.942	0.886	0.896

Table S13. Comparison of different methods for regional level %ILI estimation in Region 9. The RMSE, MAE, and correlation measures are reported.. The method with the best performance is highlighted in boldface for each metric in each period.

	'09-'23	'09-'15	'09-'10	'10-'11	'11-'12	'12-'13	'13-'14	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE																
ARGO-C	0.383	0.419	0.697	0.466	0.225	0.379	0.376	0.259	0.254	0.438	0.345	0.282	0.453	0.122	0.205	1.144
ARGO2	0.396	0.428	0.687	0.469	0.221	0.406	0.399	0.267	0.259	0.452	0.350	0.308	0.530	0.120	0.179	1.194
VAR1	0.471	0.469	0.733	0.477	0.235	0.438	0.463	0.330	0.277	0.493	0.504	0.410	0.774	0.085	0.267	1.526
GFT	–	1.083	0.800	0.423	0.629	2.880	0.869	0.339	–	–	–	–	–	–	–	–
naive	0.457	0.457	0.705	0.493	0.237	0.451	0.467	0.325	0.271	0.506	0.505	0.400	0.783	0.085	0.259	1.393
MAE																
ARGO-C	0.223	0.253	0.513	0.340	0.174	0.254	0.221	0.177	0.201	0.282	0.241	0.208	0.340	0.094	0.170	0.751
ARGO2	0.225	0.256	0.494	0.325	0.179	0.260	0.226	0.187	0.206	0.282	0.220	0.232	0.380	0.091	0.144	0.759
VAR1	0.253	0.260	0.454	0.306	0.202	0.263	0.270	0.245	0.213	0.282	0.292	0.282	0.569	0.063	0.185	1.184
GFT	–	0.644	0.610	0.355	0.597	2.175	0.764	0.254	–	–	–	–	–	–	–	–
naive	0.255	0.270	0.474	0.335	0.208	0.286	0.280	0.248	0.208	0.330	0.321	0.277	0.564	0.064	0.176	0.997
Correlation																
ARGO-C	0.958	0.944	0.964	0.869	0.852	0.920	0.926	0.959	0.903	0.880	0.962	0.968	0.969	0.900	0.932	0.937
ARGO2	0.956	0.942	0.964	0.866	0.860	0.905	0.916	0.960	0.899	0.872	0.962	0.962	0.958	0.906	0.946	0.930
VAR1	0.936	0.931	0.956	0.859	0.847	0.890	0.882	0.933	0.885	0.849	0.919	0.927	0.899	0.927	0.877	0.891
GFT	–	0.804	0.985	0.919	0.887	0.911	0.974	0.933	–	–	–	–	–	–	–	–
naive	0.942	0.935	0.962	0.859	0.850	0.887	0.883	0.933	0.892	0.850	0.919	0.932	0.898	0.925	0.882	0.891

Table S14. Comparison of different methods for regional level %ILI estimation in Region 10. The RMSE, MAE, and correlation measures are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.831	0.752	0.577	1.756	1.922	0.625	0.544	0.846	0.504	0.698	0.198	0.274	1.507
ARGOX	0.873	0.797	0.632	1.865	2.045	0.607	0.596	0.864	0.530	0.779	0.192	0.289	1.653
VARI	2.407	2.383	2.336	4.624	5.083	1.324	2.231	3.394	1.693	3.559	0.351	0.687	5.138
GFT	—	—	—	2.891	3.183	—	—	—	—	—	—	—	—
naive	1.115	0.977	0.646	2.271	2.495	0.591	0.831	1.315	0.785	1.043	0.142	0.453	1.645
MAE													
ARGO-C	0.462	0.400	0.284	1.107	1.262	0.480	0.407	0.459	0.342	0.492	0.155	0.213	1.067
ARGOX	0.477	0.413	0.294	1.119	1.292	0.463	0.467	0.482	0.369	0.530	0.154	0.196	1.210
VARI	1.349	1.271	1.125	2.847	3.324	1.156	1.665	2.169	1.072	2.374	0.293	0.599	4.203
GFT	—	—	—	1.934	2.321	—	—	—	—	—	—	—	—
naive	0.595	0.501	0.325	1.316	1.552	0.444	0.639	0.757	0.573	0.782	0.121	0.307	1.189
Correlation													
ARGO-C	0.956	0.956	0.945	0.911	0.894	0.815	0.958	0.976	0.970	0.962	0.709	0.980	0.832
ARGOX	0.952	0.951	0.936	0.898	0.878	0.831	0.950	0.976	0.966	0.951	0.721	0.970	0.799
VARI	0.675	0.517	0.632	0.715	0.655	0.820	0.639	0.615	0.859	0.278	-0.105	0.777	-0.040
GFT	—	—	—	0.960	0.952	—	—	—	—	—	—	—	—
naive	0.923	0.928	0.933	0.855	0.826	0.851	0.906	0.941	0.925	0.908	0.795	0.906	0.808

Table S15. Comparison of different methods for state-level %ILI estimation in Alabama. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.961	0.849	0.586	0.663	0.630	0.828	0.705	0.634	1.097	1.962	0.301	0.631	0.625
ARGOX	0.966	0.855	0.596	0.654	0.622	0.832	0.707	0.635	1.108	1.979	0.298	0.655	0.630
VARI	1.058	0.945	0.690	0.689	0.656	1.159	0.929	1.095	1.215	1.847	0.305	0.789	1.071
GFT	—	—	—	1.015	1.042	—	—	—	—	—	—	—	—
naive	1.002	0.896	0.654	0.646	0.605	0.843	0.783	0.645	1.126	2.047	0.287	0.788	0.660
MAE													
ARGO-C	0.594	0.519	0.381	0.493	0.460	0.565	0.519	0.508	0.765	1.267	0.208	0.363	0.437
ARGOX	0.595	0.521	0.382	0.488	0.454	0.569	0.524	0.515	0.774	1.265	0.200	0.361	0.436
VARI	0.684	0.601	0.449	0.538	0.502	0.749	0.686	0.938	0.836	1.148	0.214	0.447	0.860
GFT	—	—	—	0.867	0.877	—	—	—	—	—	—	—	—
naive	0.619	0.552	0.429	0.479	0.438	0.596	0.604	0.519	0.821	1.259	0.210	0.479	0.485
Correlation													
ARGO-C	0.871	0.873	0.821	0.731	0.777	0.818	0.796	0.915	0.881	0.629	0.604	0.795	0.838
ARGOX	0.870	0.871	0.808	0.737	0.782	0.816	0.796	0.914	0.879	0.620	0.605	0.782	0.828
VARI	0.856	0.851	0.701	0.756	0.807	0.802	0.743	0.893	0.870	0.614	0.552	0.741	0.678
GFT	—	—	—	0.623	0.776	—	—	—	—	—	—	—	—
naive	0.864	0.864	0.774	0.768	0.812	0.820	0.769	0.907	0.880	0.603	0.533	0.712	0.759

Table S16. Comparison of different methods for state-level %ILI estimation in Alaska. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.396	0.378	0.342	0.246	0.256	0.391	0.319	0.587	0.315	0.662	0.272	0.336	0.483
ARGOX	0.404	0.382	0.338	0.253	0.266	0.413	0.311	0.645	0.309	0.629	0.259	0.335	0.513
VARI	2.372	2.080	1.384	4.487	4.938	1.218	2.423	2.891	1.615	2.398	0.365	0.539	3.543
GFT	—	—	—	1.160	1.253	—	—	—	—	—	—	—	—
naive	0.454	0.432	0.387	0.280	0.298	0.401	0.304	0.787	0.363	0.707	0.279	0.424	0.679
MAE													
ARGO-C	0.290	0.277	0.254	0.187	0.193	0.286	0.267	0.448	0.267	0.533	0.227	0.243	0.370
ARGOX	0.292	0.278	0.251	0.182	0.192	0.304	0.259	0.484	0.266	0.505	0.212	0.246	0.401
VARI	1.409	1.164	0.708	2.910	3.479	0.976	1.960	1.892	0.976	1.839	0.316	0.343	2.430
GFT	—	—	—	1.010	1.131	—	—	—	—	—	—	—	—
naive	0.309	0.294	0.266	0.216	0.233	0.293	0.254	0.539	0.269	0.542	0.217	0.292	0.525
Correlation													
ARGO-C	0.953	0.952	0.938	0.959	0.955	0.945	0.862	0.942	0.921	0.846	0.770	0.888	0.954
ARGOX	0.951	0.951	0.940	0.958	0.953	0.938	0.864	0.928	0.925	0.855	0.780	0.882	0.945
VARI	0.613	0.660	0.754	0.725	0.666	0.468	0.680	0.719	0.707	0.799	0.516	0.861	0.484
GFT	—	—	—	0.923	0.912	—	—	—	—	—	—	—	—
naive	0.939	0.939	0.924	0.941	0.932	0.939	0.865	0.884	0.901	0.816	0.749	0.816	0.904

Table S17. Comparison of different methods for state-level %ILI estimation in Arizona. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.821	0.741	0.564	1.050	1.144	0.622	0.842	0.905	0.830	1.576	0.195	0.405	0.718
ARGOX	0.853	0.780	0.622	1.102	1.201	0.612	0.897	0.973	0.805	1.716	0.199	0.382	0.798
VARI	1.103	0.950	0.565	1.382	1.514	0.656	1.154	1.597	1.041	1.812	0.167	0.417	0.885
GFT	—	—	—	1.391	1.466	—	—	—	—	—	—	—	—
naive	1.049	0.919	0.609	1.328	1.453	0.647	1.080	1.444	0.987	1.871	0.168	0.421	0.846
MAE													
ARGO-C	0.518	0.450	0.324	0.601	0.657	0.456	0.596	0.698	0.569	1.159	0.155	0.292	0.579
ARGOX	0.530	0.463	0.339	0.649	0.716	0.446	0.605	0.732	0.584	1.228	0.159	0.274	0.672
VARI	0.636	0.521	0.306	0.809	0.923	0.464	0.713	1.012	0.771	1.202	0.142	0.309	0.729
GFT	—	—	—	1.179	1.237	—	—	—	—	—	—	—	—
naive	0.627	0.518	0.316	0.747	0.847	0.472	0.708	1.042	0.729	1.253	0.136	0.326	0.704
Correlation													
ARGO-C	0.946	0.943	0.923	0.922	0.907	0.865	0.937	0.975	0.938	0.826	0.722	0.859	0.927
ARGOX	0.942	0.937	0.910	0.913	0.896	0.872	0.928	0.969	0.940	0.795	0.723	0.856	0.910
VARI	0.911	0.908	0.914	0.877	0.853	0.860	0.885	0.914	0.899	0.753	0.712	0.814	0.892
GFT	—	—	—	0.971	0.967	—	—	—	—	—	—	—	—
naive	0.913	0.914	0.911	0.878	0.854	0.866	0.893	0.918	0.903	0.755	0.711	0.812	0.900

Table S18. Comparison of different methods for state-level %ILI estimation in Arkansas. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.252	0.267	0.292	0.255	0.275	0.278	0.247	0.389	0.235	0.342	0.142	0.248	0.649
ARGOX	0.260	0.279	0.312	0.276	0.298	0.303	0.243	0.416	0.217	0.326	0.133	0.261	0.738
VARI	0.793	0.872	1.003	1.287	1.347	0.648	0.640	0.987	0.595	0.810	0.264	0.671	2.477
GFT	—	—	—	0.535	0.555	—	—	—	—	—	—	—	—
naive	0.344	0.372	0.419	0.370	0.402	0.354	0.252	0.640	0.262	0.492	0.150	0.410	0.937
MAE													
ARGO-C	0.182	0.185	0.192	0.197	0.216	0.219	0.179	0.263	0.188	0.251	0.115	0.206	0.513
ARGOX	0.181	0.186	0.194	0.205	0.225	0.241	0.183	0.274	0.165	0.239	0.109	0.208	0.559
VARI	0.636	0.646	0.665	1.206	1.264	0.472	0.570	0.836	0.475	0.620	0.206	0.513	2.311
GFT	—	—	—	0.398	0.400	—	—	—	—	—	—	—	—
naive	0.220	0.228	0.244	0.247	0.275	0.277	0.194	0.390	0.220	0.347	0.119	0.243	0.739
Correlation													
ARGO-C	0.974	0.977	0.981	0.967	0.961	0.942	0.934	0.959	0.950	0.970	0.947	0.955	0.953
ARGOX	0.972	0.975	0.978	0.963	0.956	0.930	0.930	0.953	0.959	0.973	0.952	0.948	0.933
VARI	0.774	0.750	0.812	0.902	0.883	0.879	0.482	0.902	0.853	0.836	0.807	0.597	0.900
GFT	—	—	—	0.941	0.936	—	—	—	—	—	—	—	—
naive	0.952	0.956	0.960	0.928	0.914	0.907	0.917	0.889	0.940	0.934	0.940	0.873	0.890

Table S19. Comparison of different methods for state-level %ILI estimation in California. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.422	0.410	0.386	0.410	0.450	0.151	0.608	0.225	0.790	0.422	0.150	0.370	0.862
ARGOX	0.417	0.411	0.401	0.402	0.441	0.155	0.589	0.207	0.783	0.443	0.146	0.384	0.906
VARI	1.414	1.289	1.019	1.671	1.669	1.512	0.880	1.602	1.921	1.829	0.621	1.340	1.582
GFT	—	—	—	0.809	0.886	—	—	—	—	—	—	—	—
naive	0.461	0.489	0.536	0.478	0.525	0.208	0.685	0.270	0.725	0.829	0.130	0.376	1.143
MAE													
ARGO-C	0.272	0.262	0.244	0.242	0.279	0.117	0.445	0.180	0.629	0.305	0.117	0.281	0.673
ARGOX	0.270	0.262	0.248	0.245	0.282	0.113	0.436	0.156	0.621	0.332	0.114	0.276	0.706
VARI	1.202	1.039	0.735	1.611	1.599	1.438	0.771	1.368	1.624	1.488	0.429	1.162	1.242
GFT	—	—	—	0.458	0.514	—	—	—	—	—	—	—	—
naive	0.303	0.305	0.309	0.304	0.352	0.166	0.506	0.216	0.562	0.596	0.100	0.303	0.915
Correlation													
ARGO-C	0.970	0.975	0.981	0.930	0.921	0.939	0.450	0.977	0.913	0.983	0.866	0.918	0.955
ARGOX	0.971	0.974	0.979	0.933	0.924	0.935	0.481	0.980	0.913	0.982	0.875	0.913	0.949
VARI	0.630	0.725	0.864	0.854	0.832	0.865	0.626	0.790	0.846	0.872	-0.210	0.189	0.849
GFT	—	—	—	0.841	0.816	—	—	—	—	—	—	—	—
naive	0.964	0.964	0.964	0.902	0.886	0.885	0.374	0.959	0.904	0.930	0.854	0.903	0.917

Table S20. Comparison of different methods for state-level %ILI estimation in Colorado. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.534	0.489	0.393	0.677	0.743	0.471	0.409	0.462	0.445	0.724	0.162	0.337	0.734
ARGOX	0.536	0.493	0.400	0.680	0.746	0.471	0.395	0.447	0.412	0.769	0.158	0.371	0.762
VARI	1.425	1.274	0.933	1.684	1.840	1.548	1.589	2.358	0.828	1.163	0.267	1.012	2.018
GFT	—	—	—	1.475	1.614	—	—	—	—	—	—	—	—
naive	0.576	0.537	0.456	0.625	0.683	0.462	0.492	0.575	0.493	0.865	0.149	0.414	0.935
MAE													
ARGO-C	0.365	0.316	0.226	0.440	0.514	0.376	0.314	0.351	0.352	0.543	0.127	0.211	0.520
ARGOX	0.358	0.313	0.231	0.433	0.504	0.382	0.300	0.336	0.311	0.540	0.120	0.221	0.575
VARI	1.078	0.907	0.589	1.311	1.496	1.363	1.253	2.055	0.614	1.020	0.220	0.865	1.595
GFT	—	—	—	1.092	1.254	—	—	—	—	—	—	—	—
naive	0.376	0.336	0.261	0.348	0.388	0.360	0.379	0.423	0.367	0.640	0.110	0.281	0.741
Correlation													
ARGO-C	0.959	0.963	0.972	0.901	0.874	0.884	0.948	0.966	0.946	0.967	0.904	0.937	0.935
ARGOX	0.959	0.963	0.970	0.901	0.875	0.881	0.952	0.968	0.954	0.965	0.903	0.913	0.928
VARI	0.729	0.755	0.907	0.642	0.564	0.697	0.679	0.950	0.840	0.902	0.598	0.697	0.904
GFT	—	—	—	0.748	0.686	—	—	—	—	—	—	—	—
naive	0.952	0.956	0.963	0.914	0.894	0.887	0.925	0.948	0.934	0.945	0.898	0.850	0.889

Table S21. Comparison of different methods for state-level %ILI estimation in Connecticut. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.421	0.404	0.370	0.549	0.604	0.252	0.203	0.689	0.256	0.808	0.190	0.293	0.315
ARGOX	0.421	0.405	0.373	0.553	0.608	0.243	0.186	0.684	0.254	0.836	0.186	0.295	0.324
VARI	1.898	1.755	1.453	1.747	1.882	1.808	2.227	2.454	2.420	3.279	0.426	0.696	2.519
GFT	—	—	—	2.085	2.232	—	—	—	—	—	—	—	—
naive	0.453	0.453	0.452	0.509	0.560	0.256	0.230	0.785	0.293	0.954	0.202	0.376	0.502
MAE													
ARGO-C	0.217	0.218	0.221	0.272	0.320	0.189	0.159	0.372	0.196	0.547	0.153	0.229	0.254
ARGOX	0.217	0.218	0.222	0.280	0.330	0.184	0.147	0.354	0.201	0.561	0.151	0.235	0.254
VARI	1.529	1.302	0.880	1.504	1.644	1.659	2.022	2.240	2.300	2.877	0.374	0.586	2.096
GFT	—	—	—	1.880	2.032	—	—	—	—	—	—	—	—
naive	0.226	0.233	0.247	0.245	0.288	0.173	0.173	0.468	0.208	0.618	0.163	0.277	0.396
Correlation													
ARGO-C	0.873	0.887	0.896	0.825	0.811	0.829	0.767	0.883	0.907	0.809	0.661	0.897	0.935
ARGOX	0.873	0.886	0.894	0.823	0.809	0.840	0.810	0.885	0.912	0.800	0.671	0.883	0.932
VARI	0.708	0.671	0.709	0.449	0.380	0.494	0.835	0.781	0.775	0.609	0.735	0.845	0.875
GFT	—	—	—	0.737	0.714	—	—	—	—	—	—	—	—
naive	0.859	0.864	0.846	0.859	0.847	0.832	0.739	0.848	0.882	0.735	0.638	0.809	0.844

Table S22. Comparison of different methods for state-level %ILI estimation in Delaware. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	1.421	1.264	0.904	1.515	1.596	1.595	2.054	1.190	0.494	0.802	0.508	0.728	1.660
ARGOX	1.418	1.262	0.902	1.509	1.587	1.615	2.062	1.226	0.459	0.798	0.512	0.714	1.673
VARI	1.523	1.346	0.934	1.560	1.651	1.632	2.157	1.604	0.542	0.861	0.459	0.777	1.601
GFT	—	—	—	5.236	5.015	—	—	—	—	—	—	—	—
naive	1.447	1.276	0.872	1.588	1.676	1.669	2.193	0.846	0.529	0.770	0.496	0.796	1.465
MAE													
ARGO-C	0.977	0.860	0.645	1.153	1.200	1.329	1.422	0.956	0.348	0.647	0.352	0.561	1.409
ARGOX	0.978	0.863	0.648	1.148	1.198	1.295	1.447	1.018	0.315	0.642	0.380	0.546	1.432
VARI	1.069	0.925	0.657	1.220	1.289	1.265	1.505	1.483	0.438	0.638	0.338	0.588	1.313
GFT	—	—	—	4.935	4.686	—	—	—	—	—	—	—	—
naive	0.890	0.788	0.601	1.274	1.342	1.277	1.509	0.453	0.342	0.607	0.335	0.606	1.224
Correlation													
ARGO-C	0.874	0.886	0.923	0.831	0.848	0.797	0.601	0.649	0.642	0.731	0.432	0.745	0.847
ARGOX	0.874	0.886	0.924	0.832	0.851	0.791	0.592	0.664	0.688	0.738	0.403	0.742	0.846
VARI	0.850	0.867	0.924	0.807	0.819	0.816	0.552	0.557	0.486	0.774	0.398	0.700	0.864
GFT	—	—	—	0.766	0.833	—	—	—	—	—	—	—	—
naive	0.873	0.888	0.928	0.816	0.828	0.810	0.562	0.721	0.605	0.787	0.390	0.700	0.884

Table S23. Comparison of different methods for state-level %ILI estimation in District of Columbia. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.577	0.543	0.473	0.860	0.945	0.318	0.421	1.031	0.401	0.700	0.247	0.446	0.856
ARGOX	0.593	0.560	0.492	0.871	0.957	0.330	0.437	1.017	0.459	0.753	0.249	0.482	0.951
VARI	3.205	2.992	2.550	2.220	2.421	1.629	3.280	5.718	3.521	4.928	1.583	2.289	4.099
GFT	—	—	—	0.819	0.896	—	—	—	—	—	—	—	—
naive	0.816	0.758	0.638	0.999	1.098	0.358	0.564	1.501	0.672	1.360	0.206	0.740	0.935
MAE													
ARGO-C	0.330	0.326	0.318	0.418	0.480	0.239	0.343	0.622	0.300	0.544	0.204	0.313	0.655
ARGOX	0.339	0.332	0.319	0.430	0.497	0.243	0.368	0.600	0.314	0.574	0.201	0.292	0.739
VARI	2.431	2.339	2.170	1.560	1.741	1.554	3.056	4.344	3.335	4.445	1.539	1.950	3.503
GFT	—	—	—	0.505	0.574	—	—	—	—	—	—	—	—
naive	0.430	0.419	0.397	0.486	0.564	0.275	0.418	0.856	0.488	1.041	0.164	0.461	0.700
Correlation													
ARGO-C	0.970	0.968	0.961	0.864	0.845	0.870	0.950	0.969	0.961	0.965	0.841	0.971	0.897
ARGOX	0.969	0.966	0.958	0.854	0.833	0.866	0.947	0.970	0.950	0.960	0.828	0.960	0.874
VARI	0.543	0.551	0.662	0.185	0.093	0.622	0.665	0.526	0.647	0.587	0.423	0.770	-0.188
GFT	—	—	—	0.892	0.877	—	—	—	—	—	—	—	—
naive	0.941	0.939	0.929	0.812	0.787	0.837	0.909	0.933	0.878	0.858	0.872	0.885	0.869

Table S24. Comparison of different methods for state-level %ILI estimation in Georgia. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.903	0.775	0.450	1.659	1.702	0.769	0.688	1.155	0.611	0.648	0.354	0.349	0.471
ARGOX	0.906	0.778	0.453	1.660	1.704	0.773	0.684	1.164	0.625	0.643	0.350	0.356	0.485
VARI	2.945	2.647	1.978	5.999	6.525	1.870	0.973	3.918	1.713	3.541	1.481	1.985	2.306
GFT	—	—	—	4.131	4.457	—	—	—	—	—	—	—	—
naive	0.930	0.796	0.449	1.597	1.627	0.787	0.740	1.269	0.707	0.618	0.374	0.347	0.504
MAE													
ARGO-C	0.618	0.517	0.330	1.132	1.185	0.611	0.568	0.865	0.468	0.487	0.297	0.241	0.369
ARGOX	0.620	0.519	0.332	1.137	1.192	0.621	0.564	0.861	0.474	0.492	0.295	0.246	0.379
VARI	1.636	1.632	1.623	4.806	5.521	1.371	0.747	2.152	1.262	2.811	1.397	1.668	1.804
GFT	—	—	—	3.590	3.997	—	—	—	—	—	—	—	—
naive	0.654	0.540	0.328	1.100	1.153	0.642	0.609	0.977	0.510	0.475	0.311	0.236	0.398
Correlation													
ARGO-C	0.935	0.942	0.925	0.915	0.900	0.907	0.745	0.770	0.870	0.905	0.108	0.862	0.558
ARGOX	0.934	0.942	0.924	0.915	0.901	0.905	0.743	0.762	0.864	0.901	0.111	0.860	0.532
VARI	0.299	0.310	0.700	0.490	0.399	0.796	0.708	0.528	0.457	0.491	-0.050	0.614	0.445
GFT	—	—	—	0.906	0.942	—	—	—	—	—	—	—	—
naive	0.932	0.940	0.927	0.923	0.908	0.905	0.716	0.721	0.833	0.890	0.113	0.865	0.470

Table S25. Comparison of different methods for state-level %ILI estimation in Hawaii. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.508	0.622	0.791	0.651	0.670	0.530	0.411	0.568	0.730	0.906	0.233	0.578	1.475
ARGOX	0.495	0.623	0.809	0.632	0.651	0.499	0.420	0.519	0.739	0.938	0.231	0.569	1.515
VARI	2.281	2.027	1.443	4.065	4.298	2.339	1.798	2.449	1.559	1.851	0.500	1.146	3.160
GFT	—	—	—	1.243	1.355	—	—	—	—	—	—	—	—
naive	0.566	0.669	0.826	0.827	0.850	0.470	0.478	0.639	0.820	0.808	0.234	0.595	1.672
MAE													
ARGO-C	0.361	0.394	0.456	0.475	0.477	0.435	0.330	0.429	0.522	0.622	0.153	0.405	0.976
ARGOX	0.347	0.389	0.465	0.451	0.452	0.401	0.329	0.388	0.527	0.643	0.146	0.398	1.018
VARI	1.860	1.545	0.962	3.381	3.654	1.923	1.676	2.294	1.379	1.624	0.403	0.938	2.588
GFT	—	—	—	1.035	1.168	—	—	—	—	—	—	—	—
naive	0.380	0.402	0.442	0.623	0.632	0.375	0.362	0.474	0.583	0.528	0.162	0.376	1.128
Correlation													
ARGO-C	0.916	0.900	0.871	0.939	0.931	0.563	0.801	0.846	0.786	0.685	0.808	0.645	0.902
ARGOX	0.920	0.899	0.864	0.941	0.933	0.611	0.786	0.870	0.778	0.669	0.814	0.653	0.891
VARI	0.794	0.508	0.528	0.719	0.652	0.587	0.637	0.791	0.679	-0.270	0.428	0.283	0.430
GFT	—	—	—	0.953	0.955	—	—	—	—	—	—	—	—
naive	0.899	0.890	0.868	0.901	0.888	0.692	0.750	0.814	0.754	0.781	0.822	0.674	0.867

Table S26. Comparison of different methods for state-level %ILI estimation in Idaho. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.284	0.284	0.284	0.460	0.503	0.233	0.230	0.328	0.269	0.430	0.156	0.293	0.555
ARGOX	0.291	0.292	0.292	0.443	0.485	0.233	0.241	0.354	0.272	0.471	0.146	0.273	0.615
VARI	1.628	1.552	1.399	2.173	2.332	1.926	1.832	1.962	0.955	2.258	0.715	0.730	2.283
GFT	—	—	—	0.672	0.704	—	—	—	—	—	—	—	—
naive	0.401	0.396	0.387	0.459	0.501	0.271	0.360	0.626	0.393	0.765	0.114	0.319	0.756
MAE													
ARGO-C	0.182	0.182	0.180	0.280	0.314	0.189	0.186	0.218	0.184	0.308	0.127	0.217	0.413
ARGOX	0.188	0.186	0.182	0.275	0.311	0.183	0.197	0.231	0.200	0.338	0.122	0.212	0.459
VARI	1.330	1.175	0.888	1.792	1.926	1.814	1.755	1.609	0.829	1.959	0.382	0.576	2.074
GFT	—	—	—	0.604	0.633	—	—	—	—	—	—	—	—
naive	0.251	0.241	0.223	0.287	0.318	0.221	0.268	0.406	0.285	0.579	0.091	0.223	0.613
Correlation													
ARGO-C	0.979	0.979	0.978	0.938	0.925	0.954	0.976	0.983	0.953	0.977	0.910	0.922	0.943
ARGOX	0.978	0.978	0.977	0.939	0.924	0.956	0.972	0.980	0.950	0.972	0.920	0.928	0.929
VARI	0.750	0.541	0.501	0.195	-0.003	0.588	0.913	0.860	0.818	0.844	0.726	0.485	0.349
GFT	—	—	—	0.945	0.937	—	—	—	—	—	—	—	—
naive	0.959	0.960	0.962	0.931	0.914	0.939	0.937	0.889	0.922	0.920	0.895	0.892	—

Table S27. Comparison of different methods for state-level %ILI estimation in Illinois. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.544	0.512	0.446	0.656	0.714	0.416	0.718	0.571	0.596	0.810	0.163	0.230	0.690
ARGOX	0.566	0.537	0.478	0.714	0.778	0.431	0.713	0.640	0.613	0.884	0.165	0.207	0.728
VARI	1.770	1.584	1.162	1.788	1.936	0.968	1.866	2.985	1.888	2.675	0.172	0.633	2.147
GFT	—	—	—	0.478	0.521	—	—	—	—	—	—	—	—
naive	0.728	0.654	0.486	0.916	1.004	0.562	0.830	0.978	0.783	1.018	0.103	0.248	0.863
MAE													
ARGO-C	0.370	0.333	0.264	0.422	0.464	0.299	0.508	0.412	0.430	0.540	0.126	0.157	0.523
ARGOX	0.377	0.339	0.268	0.465	0.518	0.303	0.499	0.459	0.428	0.516	0.133	0.144	0.564
VARI	1.240	1.057	0.717	1.341	1.494	0.727	1.294	2.375	1.547	2.281	0.131	0.481	1.727
GFT	—	—	—	0.379	0.429	—	—	—	—	—	—	—	—
naive	0.457	0.397	0.285	0.525	0.607	0.433	0.596	0.624	0.540	0.664	0.080	0.193	0.675
Correlation													
ARGO-C	0.951	0.948	0.936	0.937	0.931	0.870	0.896	0.974	0.915	0.927	0.911	0.963	0.939
ARGOX	0.947	0.943	0.928	0.923	0.915	0.860	0.898	0.967	0.909	0.913	0.922	0.957	0.929
VARI	0.515	0.505	0.663	0.511	0.444	0.447	0.604	0.878	0.667	-0.235	0.878	0.189	0.771
GFT	—	—	—	0.970	0.968	—	—	—	—	—	—	—	—
naive	0.914	0.917	0.926	0.877	0.864	0.776	0.865	0.923	0.857	0.873	0.934	0.913	0.901

Table S28. Comparison of different methods for state-level %ILI estimation in Indiana. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.399	0.686	1.023	0.668	0.735	0.280	0.362	0.394	0.409	0.537	0.930	0.534	1.034
ARGOX	0.397	0.699	1.050	0.652	0.718	0.267	0.388	0.386	0.408	0.632	0.923	0.597	1.119
VARI	1.442	1.466	1.510	1.982	2.101	1.811	1.730	1.630	0.869	2.133	0.915	0.985	1.368
GFT	—	—	—	2.296	2.519	—	—	—	—	—	—	—	—
naive	0.463	0.786	1.169	0.646	0.711	0.294	0.425	0.554	0.478	0.749	0.995	0.608	1.386
MAE													
ARGO-C	0.244	0.319	0.457	0.355	0.420	0.236	0.260	0.304	0.288	0.362	0.537	0.383	0.756
ARGOX	0.240	0.324	0.479	0.347	0.414	0.223	0.273	0.285	0.298	0.396	0.546	0.426	0.802
VARI	1.206	1.109	0.928	1.643	1.721	1.740	1.577	1.348	0.755	1.834	0.463	0.795	1.110
GFT	—	—	—	1.346	1.534	—	—	—	—	—	—	—	—
naive	0.261	0.350	0.514	0.314	0.377	0.244	0.302	0.366	0.347	0.466	0.419	0.429	1.041
Correlation													
ARGO-C	0.917	0.845	0.759	0.677	0.600	0.706	0.907	0.939	0.925	0.926	0.495	0.833	0.880
ARGOX	0.918	0.838	0.745	0.683	0.606	0.731	0.892	0.940	0.924	0.902	0.508	0.791	0.854
VARI	0.790	0.592	0.502	0.396	0.261	0.802	0.710	0.886	0.821	0.781	0.488	0.379	0.835
GFT	—	—	—	0.575	0.507	—	—	—	—	—	—	—	—
naive	0.891	0.811	0.725	0.723	0.668	0.702	0.875	0.878	0.895	0.852	0.516	0.796	0.784

Table S29. Comparison of different methods for state-level %ILI estimation in Iowa. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.523	0.508	0.478	0.702	0.770	0.283	0.864	0.533	0.507	0.635	0.212	0.482	0.931
ARGOX	0.526	0.515	0.494	0.729	0.799	0.277	0.902	0.481	0.456	0.668	0.213	0.472	1.012
VARI	1.444	1.251	0.774	1.783	1.958	0.470	2.102	2.230	0.877	1.951	0.249	0.614	1.280
GFT	—	—	—	1.472	1.620	—	—	—	—	—	—	—	—
naive	0.678	0.633	0.542	0.898	0.986	0.298	0.971	1.021	0.554	0.874	0.196	0.441	1.151
MAE													
ARGO-C	0.332	0.322	0.303	0.420	0.487	0.230	0.569	0.402	0.320	0.503	0.157	0.352	0.623
ARGOX	0.328	0.320	0.305	0.412	0.475	0.229	0.590	0.361	0.307	0.516	0.157	0.349	0.676
VARI	0.879	0.751	0.513	1.168	1.366	0.350	1.201	1.638	0.643	1.542	0.220	0.459	0.925
GFT	—	—	—	1.090	1.302	—	—	—	—	—	—	—	—
naive	0.404	0.376	0.326	0.528	0.614	0.222	0.674	0.723	0.381	0.649	0.142	0.323	0.860
Correlation													
ARGO-C	0.979	0.974	0.942	0.964	0.956	0.884	0.944	0.990	0.949	0.970	0.892	0.788	0.903
ARGOX	0.979	0.973	0.939	0.962	0.953	0.890	0.939	0.992	0.960	0.968	0.891	0.778	0.882
VARI	0.883	0.863	0.845	0.860	0.845	0.764	0.758	0.957	0.836	0.909	0.923	0.688	0.814
GFT	—	—	—	0.960	0.957	—	—	—	—	—	—	—	—
naive	0.964	0.960	0.931	0.939	0.924	0.874	0.929	0.964	0.937	0.943	0.899	0.785	0.850

Table S30. Comparison of different methods for state-level %ILI estimation in Kansas. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.670	0.631	0.550	0.279	0.306	0.254	1.071	1.018	0.837	0.997	0.215	0.391	1.210
ARGOX	0.683	0.656	0.603	0.293	0.322	0.244	1.075	0.996	0.875	1.163	0.217	0.347	1.280
VARI	1.698	1.507	1.067	2.646	2.906	0.741	1.284	1.900	2.522	2.259	0.275	0.636	2.434
GFT	—	—	—	2.912	3.180	—	—	—	—	—	—	—	—
naive	0.850	0.774	0.607	0.393	0.432	0.281	1.222	1.300	1.038	1.384	0.174	0.395	1.438
MAE													
ARGO-C	0.379	0.353	0.306	0.178	0.206	0.164	0.798	0.692	0.504	0.658	0.168	0.306	0.860
ARGOX	0.391	0.361	0.306	0.188	0.216	0.158	0.830	0.668	0.533	0.746	0.171	0.261	0.907
VARI	1.019	0.869	0.590	2.027	2.365	0.697	0.834	1.229	1.544	1.768	0.235	0.464	1.738
GFT	—	—	—	2.101	2.363	—	—	—	—	—	—	—	—
naive	0.465	0.420	0.337	0.220	0.265	0.190	0.913	0.871	0.680	0.975	0.136	0.324	1.099
Correlation													
ARGO-C	0.971	0.968	0.953	0.952	0.947	0.925	0.925	0.965	0.957	0.955	0.848	0.921	0.903
ARGOX	0.970	0.965	0.945	0.946	0.941	0.931	0.924	0.968	0.953	0.940	0.860	0.923	0.889
VARI	0.797	0.797	0.800	0.777	0.749	0.916	0.894	0.872	0.723	0.772	0.808	0.728	0.657
GFT	—	—	—	0.962	0.959	—	—	—	—	—	—	—	—
naive	0.954	0.952	0.943	0.906	0.895	0.908	0.898	0.941	0.934	0.909	0.862	0.859	0.867

Table S31. Comparison of different methods for state-level %ILI estimation in Kentucky. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.511	0.487	0.438	0.488	0.534	0.276	0.274	0.617	0.870	0.896	0.507	0.350	0.569
ARGOX	0.524	0.496	0.438	0.455	0.498	0.264	0.277	0.679	0.866	0.937	0.491	0.317	0.655
VARI	2.297	2.099	1.671	2.946	3.199	1.567	2.045	2.380	1.876	4.363	0.345	0.769	2.875
GFT	—	—	—	1.015	1.106	—	—	—	—	—	—	—	—
naive	0.641	0.612	0.553	0.567	0.621	0.249	0.355	0.917	0.981	1.373	0.323	0.385	0.680
MAE													
ARGO-C	0.306	0.299	0.287	0.325	0.371	0.230	0.200	0.429	0.491	0.682	0.318	0.225	0.471
ARGOX	0.318	0.308	0.290	0.310	0.355	0.220	0.215	0.476	0.515	0.741	0.303	0.198	0.562
VARI	1.888	1.599	1.065	2.661	2.979	1.501	1.598	2.032	1.666	3.708	0.289	0.686	2.146
GFT	—	—	—	0.748	0.838	—	—	—	—	—	—	—	—
naive	0.386	0.361	0.315	0.384	0.441	0.195	0.272	0.608	0.705	1.121	0.144	0.267	0.592
Correlation													
ARGO-C	0.979	0.980	0.976	0.964	0.952	0.871	0.968	0.980	0.951	0.952	0.252	0.955	0.898
ARGOX	0.978	0.979	0.976	0.968	0.955	0.878	0.968	0.976	0.950	0.948	0.244	0.951	0.866
VARI	0.769	0.711	0.621	0.735	0.708	0.725	0.714	0.774	0.906	0.659	0.442	0.775	0.408
GFT	—	—	—	0.981	0.978	—	—	—	—	—	—	—	—
naive	0.968	0.968	0.965	0.946	0.924	0.876	0.948	0.952	0.927	0.884	0.422	0.878	0.849

Table S32. Comparison of different methods for state-level %ILI estimation in Louisiana. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.276	0.391	0.542	0.327	0.352	0.335	0.274	0.248	0.293	0.828	0.127	0.287	1.136
ARGOX	0.278	0.398	0.556	0.328	0.353	0.332	0.280	0.251	0.296	0.857	0.117	0.317	1.142
VARI	3.012	2.731	2.112	2.174	2.376	1.319	2.017	5.069	3.685	5.899	0.617	0.840	2.727
GFT	—	—	—	0.744	0.813	—	—	—	—	—	—	—	—
naive	0.296	0.428	0.602	0.354	0.380	0.375	0.297	0.253	0.297	0.865	0.114	0.255	1.354
MAE													
ARGO-C	0.203	0.222	0.257	0.253	0.273	0.269	0.216	0.197	0.236	0.440	0.099	0.218	0.805
ARGOX	0.204	0.224	0.262	0.252	0.272	0.266	0.220	0.197	0.238	0.453	0.093	0.230	0.811
VARI	2.244	1.931	1.350	1.792	2.047	1.229	1.786	4.150	3.133	5.433	0.514	0.680	2.165
GFT	—	—	—	0.482	0.547	—	—	—	—	—	—	—	—
naive	0.216	0.238	0.278	0.273	0.292	0.290	0.229	0.208	0.232	0.451	0.091	0.193	1.034
Correlation													
ARGO-C	0.952	0.944	0.940	0.886	0.859	0.295	0.804	0.792	0.945	0.894	0.693	0.878	0.921
ARGOX	0.951	0.943	0.939	0.885	0.858	0.306	0.793	0.787	0.943	0.887	0.702	0.856	0.921
VARI	0.677	0.534	0.506	0.696	0.600	0.328	0.497	0.625	0.845	0.571	0.281	0.211	0.272
GFT	—	—	—	0.917	0.916	—	—	—	—	—	—	—	—
naive	0.945	0.933	0.926	0.871	0.844	0.257	0.789	0.760	0.940	0.885	0.645	0.897	0.882

Table S33. Comparison of different methods for state-level %ILI estimation in Maine. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.479	0.462	0.431	0.540	0.514	0.568	0.593	0.446	0.511	0.646	0.135	0.428	0.791
ARGOX	0.494	0.476	0.440	0.558	0.522	0.580	0.621	0.438	0.510	0.639	0.126	0.459	0.893
VARI	1.055	1.168	1.352	1.089	1.098	1.237	1.329	0.993	0.775	1.705	0.417	0.806	3.443
GFT	—	—	—	1.282	1.387	—	—	—	—	—	—	—	—
naive	0.601	0.565	0.491	0.714	0.743	0.657	0.721	0.619	0.624	0.810	0.134	0.564	0.893
MAE													
ARGO-C	0.353	0.318	0.253	0.364	0.326	0.440	0.463	0.336	0.403	0.451	0.108	0.278	0.638
ARGOX	0.362	0.327	0.262	0.359	0.311	0.450	0.492	0.348	0.402	0.498	0.103	0.280	0.745
VARI	0.768	0.791	0.832	0.730	0.674	0.990	0.939	0.672	0.640	1.279	0.377	0.642	3.007
GFT	—	—	—	1.023	1.137	—	—	—	—	—	—	—	—
naive	0.416	0.371	0.289	0.447	0.434	0.494	0.510	0.444	0.496	0.620	0.106	0.335	0.743
Correlation													
ARGO-C	0.942	0.950	0.962	0.859	0.894	0.745	0.924	0.963	0.914	0.956	0.916	0.924	0.900
ARGOX	0.939	0.947	0.960	0.841	0.884	0.730	0.913	0.964	0.915	0.957	0.920	0.905	0.875
VARI	0.765	0.714	0.687	0.358	0.396	0.223	0.849	0.794	0.814	0.789	0.824	0.694	-0.081
GFT	—	—	—	0.853	0.926	—	—	—	—	—	—	—	—
naive	0.911	0.928	0.952	0.753	0.775	0.695	0.872	0.924	0.876	0.918	0.905	0.838	0.869

Table S34. Comparison of different methods for state-level %ILI estimation in Maryland. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.242	0.326	0.442	0.275	0.296	0.230	0.190	0.369	0.188	0.621	0.115	0.426	0.897
ARGOX	0.249	0.331	0.446	0.275	0.297	0.223	0.204	0.381	0.213	0.619	0.101	0.445	0.911
VARI	0.720	0.864	1.080	0.648	0.621	0.816	0.892	0.419	0.652	1.012	0.415	0.920	2.514
GFT	—	—	—	0.344	0.353	—	—	—	—	—	—	—	—
naive	0.329	0.415	0.540	0.335	0.363	0.258	0.271	0.541	0.324	0.711	0.085	0.500	1.229
MAE													
ARGO-C	0.161	0.188	0.239	0.186	0.196	0.171	0.154	0.263	0.146	0.393	0.092	0.311	0.604
ARGOX	0.163	0.191	0.242	0.177	0.189	0.163	0.159	0.276	0.162	0.404	0.079	0.326	0.635
VARI	0.577	0.638	0.752	0.506	0.461	0.681	0.653	0.316	0.540	0.835	0.386	0.756	1.997
GFT	—	—	—	0.266	0.260	—	—	—	—	—	—	—	—
naive	0.208	0.234	0.282	0.237	0.258	0.181	0.203	0.368	0.231	0.550	0.061	0.376	0.861
Correlation													
ARGO-C	0.975	0.970	0.966	0.916	0.908	0.888	0.954	0.965	0.973	0.936	0.918	0.922	0.952
ARGOX	0.973	0.969	0.965	0.920	0.913	0.894	0.947	0.963	0.964	0.938	0.923	0.913	0.948
VARI	0.846	0.824	0.806	0.534	0.571	0.533	0.819	0.950	0.801	0.943	0.879	0.590	0.426
GFT	—	—	—	0.940	0.931	—	—	—	—	—	—	—	—
naive	0.952	0.949	0.946	0.877	0.861	0.870	0.908	0.919	0.919	0.912	0.927	0.890	0.892

Table S35. Comparison of different methods for state-level %ILI estimation in Massachusetts. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.276	0.250	0.194	0.366	0.398	0.207	0.332	0.338	0.298	0.316	0.132	0.223	0.329
ARGOX	0.275	0.252	0.201	0.384	0.419	0.199	0.328	0.319	0.285	0.344	0.130	0.238	0.326
VARI	1.020	1.029	1.046	0.980	1.048	0.646	1.471	2.000	0.605	1.188	0.173	0.408	2.644
GFT	—	—	—	1.203	1.302	—	—	—	—	—	—	—	—
naive	0.359	0.329	0.266	0.538	0.591	0.257	0.411	0.489	0.329	0.437	0.132	0.245	0.520
MAE													
ARGO-C	0.193	0.173	0.137	0.247	0.278	0.165	0.226	0.273	0.226	0.230	0.106	0.163	0.255
ARGOX	0.194	0.175	0.140	0.264	0.302	0.158	0.238	0.248	0.221	0.259	0.108	0.171	0.253
VARI	0.673	0.610	0.493	0.741	0.783	0.447	1.138	1.582	0.501	0.760	0.146	0.331	1.845
GFT	—	—	—	1.017	1.126	—	—	—	—	—	—	—	—
naive	0.234	0.212	0.171	0.304	0.353	0.184	0.299	0.403	0.255	0.321	0.103	0.176	0.412
Correlation													
ARGO-C	0.967	0.969	0.972	0.939	0.931	0.950	0.947	0.973	0.861	0.962	0.238	0.855	0.939
ARGOX	0.967	0.969	0.971	0.932	0.925	0.957	0.949	0.977	0.875	0.954	0.254	0.835	0.940
VARI	0.479	0.542	0.848	0.382	0.250	0.124	0.333	0.627	0.632	0.812	0.389	0.620	0.816
GFT	—	—	—	0.978	0.976	—	—	—	—	—	—	—	—
naive	0.944	0.947	0.949	0.860	0.839	0.917	0.936	0.837	0.924	0.174	0.804	0.849	—

Table S36. Comparison of different methods for state-level %ILI estimation in Michigan. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.571	0.568	0.561	0.611	0.658	0.560	0.415	0.407	0.503	1.187	0.239	0.963	0.666
ARGOX	0.578	0.593	0.619	0.653	0.706	0.565	0.420	0.375	0.505	1.248	0.253	1.016	0.768
VARI	1.153	1.164	1.185	1.557	1.693	0.758	0.812	0.934	1.185	2.129	0.198	2.103	1.641
GFT	—	—	—	0.780	0.850	—	—	—	—	—	—	—	—
naive	0.721	0.671	0.567	0.898	0.970	0.714	0.526	0.502	0.538	1.437	0.092	0.984	0.770
MAE													
ARGO-C	0.362	0.342	0.306	0.417	0.454	0.476	0.318	0.330	0.363	0.654	0.182	0.575	0.438
ARGOX	0.365	0.356	0.339	0.455	0.498	0.475	0.331	0.310	0.364	0.678	0.202	0.626	0.548
VARI	0.816	0.764	0.666	1.156	1.283	0.565	0.681	0.757	1.002	1.679	0.149	1.509	1.275
GFT	—	—	—	0.573	0.653	—	—	—	—	—	—	—	—
naive	0.431	0.375	0.271	0.584	0.641	0.544	0.378	0.392	0.402	0.775	0.076	0.475	0.563
Correlation													
ARGO-C	0.924	0.926	0.924	0.928	0.919	0.741	0.936	0.976	0.835	0.786	0.665	0.827	0.901
ARGOX	0.922	0.919	0.908	0.917	0.905	0.734	0.936	0.979	0.835	0.760	0.688	0.806	0.869
VARI	0.814	0.786	0.704	0.791	0.755	0.689	0.768	0.957	0.500	0.684	0.055	0.532	0.241
GFT	—	—	—	0.927	0.916	—	—	—	—	—	—	—	—
naive	0.882	0.900	0.926	0.844	0.825	0.629	0.888	0.959	0.807	0.670	0.864	0.829	0.861

Table S37. Comparison of different methods for state-level %ILI estimation in Minnesota. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.568	0.561	0.547	0.652	0.713	0.353	0.486	0.939	0.632	0.691	0.400	0.428	1.131
ARGOX	0.610	0.604	0.591	0.755	0.828	0.354	0.488	1.027	0.636	0.763	0.400	0.438	1.270
VARI	2.273	2.162	1.938	2.985	3.274	1.781	1.877	4.023	2.191	2.034	1.610	1.553	3.783
GFT	—	—	—	1.454	1.561	—	—	—	—	—	—	—	—
naive	0.790	0.751	0.673	1.022	1.123	0.448	0.650	1.320	0.840	0.946	0.446	0.634	1.384
MAE													
ARGO-C	0.381	0.387	0.398	0.467	0.531	0.285	0.388	0.666	0.411	0.555	0.313	0.333	0.983
ARGOX	0.403	0.406	0.412	0.519	0.598	0.289	0.393	0.730	0.429	0.595	0.314	0.316	1.115
VARI	1.605	1.600	1.591	2.362	2.740	1.676	1.593	2.812	1.533	1.643	1.499	1.343	3.178
GFT	—	—	—	1.284	1.394	—	—	—	—	—	—	—	—
naive	0.507	0.495	0.473	0.665	0.781	0.344	0.519	0.906	0.601	0.769	0.354	0.442	1.178
Correlation													
ARGO-C	0.969	0.964	0.944	0.964	0.952	0.891	0.933	0.966	0.958	0.939	0.663	0.932	0.854
ARGOX	0.964	0.958	0.935	0.951	0.935	0.891	0.933	0.960	0.957	0.926	0.663	0.931	0.820
VARI	0.681	0.666	0.661	0.706	0.650	0.641	0.638	0.432	0.673	0.659	-0.259	0.550	0.365
GFT	—	—	—	0.956	0.949	—	—	—	—	—	—	—	—
naive	0.940	0.936	0.918	0.910	0.881	0.835	0.883	0.928	0.912	0.874	0.617	0.852	0.784

Table S38. Comparison of different methods for state-level %ILI estimation in Mississippi. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.658	0.584	0.410	0.531	0.564	0.370	0.624	0.949	0.807	1.223	0.164	0.372	0.553
ARGOX	0.670	0.608	0.472	0.561	0.600	0.360	0.676	0.946	0.840	1.335	0.154	0.331	0.580
VARI	0.876	0.759	0.470	0.895	0.971	0.394	0.761	1.378	1.012	1.542	0.107	0.396	0.741
GFT	—	—	—	0.728	0.786	—	—	—	—	—	—	—	—
naive	0.843	0.744	0.511	0.911	0.989	0.393	0.692	1.368	1.008	1.474	0.112	0.406	0.726
MAE													
ARGO-C	0.393	0.348	0.264	0.355	0.370	0.302	0.410	0.614	0.561	0.841	0.130	0.260	0.407
ARGOX	0.396	0.352	0.271	0.375	0.396	0.284	0.455	0.610	0.571	0.891	0.126	0.241	0.423
VARI	0.488	0.405	0.249	0.491	0.542	0.311	0.489	0.884	0.719	1.045	0.084	0.259	0.553
GFT	—	—	—	0.604	0.670	—	—	—	—	—	—	—	—
naive	0.484	0.407	0.265	0.528	0.586	0.311	0.466	0.907	0.718	1.035	0.085	0.271	0.533
Correlation													
ARGO-C	0.960	0.960	0.966	0.961	0.953	0.771	0.932	0.970	0.940	0.929	0.845	0.961	0.943
ARGOX	0.959	0.957	0.958	0.957	0.947	0.784	0.920	0.970	0.929	0.917	0.846	0.961	0.934
VARI	0.930	0.933	0.947	0.886	0.857	0.756	0.912	0.934	0.886	0.894	0.908	0.926	0.894
GFT	—	—	—	0.959	0.964	—	—	—	—	—	—	—	—
naive	0.934	0.936	0.946	0.886	0.858	0.761	0.919	0.936	0.890	0.900	0.904	0.919	0.896

Table S39. Comparison of different methods for state-level %ILI estimation in Missouri. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.227	0.284	0.368	0.249	0.271	0.233	0.141	0.157	0.439	0.390	0.166	0.349	0.811
ARGOX	0.229	0.291	0.381	0.254	0.277	0.233	0.143	0.157	0.440	0.398	0.167	0.362	0.850
VARI	3.395	2.943	1.829	4.147	4.503	2.903	3.537	6.030	2.562	2.865	1.351	1.787	3.343
GFT	—	—	—	1.100	1.192	—	—	—	—	—	—	—	—
naive	0.241	0.317	0.424	0.296	0.323	0.259	0.154	0.174	0.429	0.457	0.166	0.394	0.953
MAE													
ARGO-C	0.136	0.168	0.225	0.181	0.204	0.160	0.097	0.117	0.320	0.276	0.123	0.262	0.557
ARGOX	0.137	0.169	0.229	0.184	0.208	0.161	0.099	0.117	0.321	0.283	0.124	0.265	0.586
VARI	2.805	2.369	1.559	3.685	4.120	2.838	3.325	5.030	2.358	2.514	1.237	1.619	2.868
GFT	—	—	—	0.930	1.029	—	—	—	—	—	—	—	—
naive	0.140	0.179	0.253	0.188	0.213	0.169	0.112	0.123	0.320	0.325	0.125	0.287	0.682
Correlation													
ARGO-C	0.961	0.958	0.942	0.898	0.887	0.421	0.575	0.751	0.919	0.958	0.816	0.818	0.885
ARGOX	0.960	0.956	0.939	0.893	0.881	0.412	0.562	0.753	0.919	0.956	0.816	0.808	0.875
VARI	0.497	0.460	0.735	0.703	0.655	0.689	0.276	0.357	0.945	0.639	0.087	0.659	0.332
GFT	—	—	—	0.938	0.930	—	—	—	—	—	—	—	—
naive	0.956	0.948	0.925	0.861	0.845	0.341	0.499	0.725	0.916	0.942	0.823	0.766	0.852

Table S40. Comparison of different methods for state-level %ILI estimation in Montana. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.605	0.600	0.591	0.698	0.766	0.326	0.466	0.823	0.828	0.789	0.239	0.362	1.489
ARGOX	0.617	0.617	0.617	0.707	0.777	0.303	0.492	0.817	0.894	0.850	0.230	0.389	1.536
VARI	2.160	1.991	1.632	1.568	1.711	0.491	1.314	3.915	3.141	2.887	0.415	1.269	4.152
GFT	—	—	—	0.866	0.951	—	—	—	—	—	—	—	—
naive	0.704	0.702	0.697	0.835	0.918	0.286	0.521	0.913	0.972	1.100	0.193	0.479	1.744
MAE													
ARGO-C	0.388	0.377	0.357	0.401	0.464	0.259	0.330	0.606	0.538	0.575	0.203	0.285	1.091
ARGOX	0.388	0.382	0.371	0.397	0.461	0.242	0.343	0.595	0.583	0.623	0.195	0.295	1.162
VARI	1.435	1.287	1.013	1.206	1.356	0.403	0.951	3.412	2.361	2.144	0.353	1.121	3.299
GFT	—	—	—	0.530	0.618	—	—	—	—	—	—	—	—
naive	0.433	0.417	0.387	0.479	0.564	0.211	0.353	0.693	0.684	0.692	0.160	0.338	1.336
Correlation													
ARGO-C	0.950	0.951	0.952	0.840	0.812	0.536	0.883	0.927	0.927	0.942	0.607	0.878	0.892
ARGOX	0.948	0.948	0.947	0.838	0.810	0.591	0.870	0.929	0.913	0.936	0.628	0.850	0.881
VARI	0.589	0.623	0.751	0.807	0.776	0.380	0.660	0.662	0.256	0.381	0.462	0.434	0.577
GFT	—	—	—	0.842	0.816	—	—	—	—	—	—	—	—
naive	0.934	0.935	0.934	0.785	0.752	0.608	0.859	0.900	0.901	0.872	0.674	0.765	0.846

Table S41. Comparison of different methods for state-level %ILI estimation in Nebraska. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.308	0.333	0.374	0.365	0.400	0.332	0.245	0.215	0.368	0.518	0.403	0.319	0.370
ARGOX	0.310	0.338	0.384	0.371	0.406	0.332	0.254	0.221	0.369	0.527	0.400	0.321	0.430
VARI	1.351	1.313	1.240	0.557	0.580	1.095	2.185	2.248	1.059	1.985	0.479	0.887	2.850
GFT	—	—	—	1.658	1.705	—	—	—	—	—	—	—	—
naive	0.330	0.359	0.408	0.447	0.489	0.363	0.245	0.354	0.429	0.394	0.418	0.354	0.550
MAE													
ARGO-C	0.209	0.236	0.285	0.272	0.315	0.235	0.187	0.174	0.231	0.418	0.303	0.238	0.294
ARGOX	0.210	0.237	0.288	0.278	0.321	0.240	0.195	0.175	0.230	0.424	0.298	0.230	0.325
VARI	0.830	0.820	0.800	0.447	0.454	0.921	1.507	1.457	0.759	1.626	0.389	0.777	2.361
GFT	—	—	—	1.569	1.602	—	—	—	—	—	—	—	—
naive	0.220	0.248	0.300	0.310	0.356	0.251	0.185	0.270	0.283	0.320	0.316	0.265	0.423
Correlation													
ARGO-C	0.940	0.931	0.902	0.950	0.941	0.886	0.885	0.976	0.923	0.809	0.700	0.860	0.939
ARGOX	0.939	0.929	0.897	0.950	0.942	0.886	0.872	0.975	0.922	0.803	0.700	0.860	0.919
VARI	0.618	0.549	0.492	0.909	0.891	0.655	0.281	0.618	0.769	0.057	0.679	0.787	0.635
GFT	—	—	—	0.930	0.919	—	—	—	—	—	—	—	—
naive	0.932	0.922	0.888	0.922	0.909	0.859	0.865	0.930	0.898	0.866	0.696	0.827	0.862

Table S42. Comparison of different methods for state-level %ILI estimation in Nevada. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.407	0.407	0.406	0.285	0.312	0.152	0.340	0.719	0.605	0.662	0.230	0.584	0.269
ARGOX	0.406	0.411	0.421	0.272	0.298	0.148	0.327	0.727	0.595	0.709	0.236	0.581	0.284
VARI	0.872	1.390	2.027	1.228	1.350	0.375	0.604	1.318	0.810	1.884	1.259	2.589	3.477
GFT	—	—	—	1.106	1.196	—	—	—	—	—	—	—	—
naive	0.436	0.419	0.385	0.241	0.264	0.165	0.346	0.797	0.612	0.620	0.251	0.598	0.366
MAE													
ARGO-C	0.243	0.240	0.235	0.183	0.211	0.109	0.227	0.476	0.363	0.423	0.161	0.374	0.215
ARGOX	0.243	0.243	0.243	0.176	0.204	0.106	0.223	0.479	0.365	0.453	0.164	0.374	0.229
VARI	0.541	0.888	1.533	0.802	0.946	0.303	0.439	0.731	0.583	1.497	1.171	1.990	2.725
GFT	—	—	—	0.889	0.972	—	—	—	—	—	—	—	—
naive	0.258	0.248	0.228	0.156	0.181	0.116	0.242	0.476	0.384	0.452	0.177	0.373	0.274
Correlation													
ARGO-C	0.918	0.895	0.807	0.840	0.813	0.897	0.663	0.909	0.732	0.919	0.222	0.639	0.914
ARGOX	0.919	0.893	0.793	0.858	0.835	0.904	0.688	0.907	0.743	0.906	0.232	0.636	0.906
VARI	0.660	0.454	0.530	0.888	0.870	0.636	0.669	0.793	0.473	0.600	0.497	-0.197	0.839
GFT	—	—	—	0.909	0.897	—	—	—	—	—	—	—	—
naive	0.907	0.894	0.823	0.894	0.879	0.883	0.692	0.886	0.743	0.928	0.192	0.661	0.844

Table S43. Comparison of different methods for state-level %ILI estimation in New Hampshire. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.446	0.623	0.859	0.470	0.511	0.485	0.620	0.716	0.196	1.575	0.266	0.810	0.773
ARGOX	0.461	0.638	0.876	0.498	0.541	0.485	0.676	0.718	0.238	1.590	0.252	0.848	0.812
VARI	1.548	1.842	2.290	1.157	1.221	1.580	2.953	1.932	1.062	4.069	1.344	1.738	2.497
GFT	—	—	—	0.828	0.860	—	—	—	—	—	—	—	—
naive	0.566	0.704	0.905	0.593	0.641	0.503	0.605	1.004	0.431	1.625	0.234	0.973	0.947
MAE													
ARGO-C	0.319	0.352	0.413	0.367	0.410	0.381	0.510	0.526	0.165	0.892	0.206	0.466	0.590
ARGOX	0.322	0.356	0.419	0.386	0.435	0.381	0.552	0.506	0.195	0.876	0.193	0.485	0.636
VARI	1.145	1.248	1.439	0.987	1.035	1.231	2.780	1.316	0.941	2.654	1.268	1.166	2.084
GFT	—	—	—	0.688	0.699	—	—	—	—	—	—	—	—
naive	0.361	0.402	0.477	0.438	0.483	0.376	0.456	0.609	0.303	1.113	0.182	0.556	0.783
Correlation													
ARGO-C	0.972	0.959	0.944	0.895	0.879	0.922	0.927	0.969	0.988	0.916	0.889	0.895	0.934
ARGOX	0.970	0.957	0.942	0.884	0.866	0.923	0.910	0.970	0.982	0.915	0.882	0.883	0.928
VARI	0.790	0.728	0.648	0.802	0.771	0.580	0.777	0.827	0.898	0.409	0.370	0.525	0.617
GFT	—	—	—	0.888	0.872	—	—	—	—	—	—	—	—
naive	0.955	0.947	0.937	0.837	0.810	0.909	0.917	0.929	0.940	0.901	0.881	0.831	0.892

Table S44. Comparison of different methods for state-level %ILI estimation in New Jersey. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.502	0.573	0.685	0.430	0.466	0.449	0.362	0.731	0.586	0.856	0.248	0.535	1.687
ARGOX	0.498	0.574	0.694	0.449	0.488	0.465	0.347	0.689	0.567	0.928	0.249	0.498	1.691
VARI	2.238	2.942	3.927	2.220	2.402	2.092	1.849	2.396	2.605	4.151	1.208	3.901	9.287
GFT	—	—	—	1.456	1.510	—	—	—	—	—	—	—	—
naive	0.647	0.726	0.852	0.665	0.729	0.484	0.409	0.911	0.749	1.136	0.256	0.706	2.174
MAE													
ARGO-C	0.329	0.355	0.402	0.304	0.335	0.345	0.267	0.441	0.390	0.637	0.195	0.412	1.270
ARGOX	0.322	0.351	0.405	0.299	0.331	0.360	0.244	0.415	0.381	0.678	0.197	0.393	1.271
VARI	1.610	2.016	2.769	1.886	2.080	1.831	1.560	1.692	2.070	3.290	1.137	3.667	7.937
GFT	—	—	—	1.285	1.322	—	—	—	—	—	—	—	—
naive	0.389	0.418	0.471	0.392	0.443	0.368	0.310	0.592	0.501	0.759	0.201	0.533	1.618
Correlation													
ARGO-C	0.967	0.972	0.975	0.948	0.938	0.929	0.949	0.952	0.955	0.962	0.837	0.935	0.952
ARGOX	0.967	0.971	0.973	0.943	0.933	0.923	0.953	0.958	0.957	0.956	0.834	0.946	0.947
VARI	0.675	0.500	0.493	0.480	0.376	0.661	0.530	0.710	0.696	0.636	0.272	0.455	0.451
GFT	—	—	—	0.947	0.943	—	—	—	—	—	—	—	—
naive	0.945	0.954	0.959	0.868	0.841	0.916	0.935	0.924	0.924	0.932	0.813	0.887	0.907

Table S45. Comparison of different methods for state-level %ILI estimation in New Mexico. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.585	0.575	0.555	0.665	0.702	0.523	0.990	0.801	0.321	0.871	0.179	0.444	1.103
ARGOX	0.589	0.576	0.551	0.690	0.734	0.520	1.064	0.719	0.294	0.839	0.167	0.447	1.131
VARI	1.260	1.593	2.074	1.217	1.310	0.886	1.402	1.417	1.764	3.741	1.135	1.597	1.939
GFT	—	—	—	0.860	0.916	—	—	—	—	—	—	—	—
naive	0.688	0.669	0.632	0.838	0.899	0.580	1.060	0.992	0.379	1.044	0.150	0.430	1.384
MAE													
ARGO-C	0.392	0.362	0.306	0.502	0.517	0.413	0.750	0.582	0.233	0.565	0.143	0.317	0.866
ARGOX	0.390	0.360	0.304	0.513	0.534	0.401	0.781	0.546	0.232	0.552	0.137	0.315	0.902
VARI	1.010	1.145	1.394	0.858	0.942	0.694	1.161	1.002	1.690	2.529	1.079	1.114	1.763
GFT	—	—	—	0.651	0.696	—	—	—	—	—	—	—	—
naive	0.415	0.391	0.347	0.553	0.588	0.432	0.732	0.662	0.291	0.775	0.119	0.339	1.146
Correlation													
ARGO-C	0.960	0.954	0.937	0.926	0.926	0.825	0.928	0.963	0.956	0.936	0.758	0.909	0.802
ARGOX	0.959	0.954	0.938	0.922	0.920	0.822	0.916	0.972	0.962	0.943	0.769	0.898	0.791
VARI	0.841	0.784	0.830	0.870	0.870	0.618	0.926	0.880	0.905	0.684	0.705	0.681	0.867
GFT	—	—	—	0.900	0.894	—	—	—	—	—	—	—	—
naive	0.945	0.939	0.921	0.886	0.878	0.794	0.915	0.942	0.940	0.901	0.814	0.867	0.702

Table S46. Comparison of different methods for state-level %ILI estimation in New York. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.460	0.428	0.363	0.537	0.589	0.344	0.711	0.716	0.335	0.372	0.121	0.202	0.941
ARGOX	0.492	0.458	0.387	0.696	0.765	0.359	0.726	0.739	0.328	0.352	0.112	0.205	1.012
VARI	1.197	1.910	2.785	1.171	1.250	0.474	1.204	0.990	1.307	2.437	0.686	2.770	6.903
GFT	—	—	—	0.644	0.704	—	—	—	—	—	—	—	—
naive	0.693	0.617	0.442	1.022	1.125	0.456	0.917	1.065	0.541	0.634	0.093	0.324	1.078
MAE													
ARGO-C	0.292	0.256	0.189	0.372	0.434	0.268	0.478	0.466	0.210	0.277	0.091	0.166	0.635
ARGOX	0.297	0.262	0.197	0.421	0.497	0.269	0.495	0.500	0.190	0.236	0.085	0.163	0.725
VARI	0.772	1.079	1.647	0.849	0.887	0.392	0.892	0.719	0.987	1.639	0.617	2.539	5.125
GFT	—	—	—	0.452	0.510	—	—	—	—	—	—	—	—
naive	0.409	0.343	0.221	0.509	0.606	0.348	0.680	0.725	0.374	0.454	0.079	0.204	0.781
Correlation													
ARGO-C	0.967	0.966	0.964	0.964	0.960	0.937	0.929	0.962	0.983	0.977	0.942	0.971	0.870
ARGOX	0.962	0.961	0.960	0.936	0.928	0.930	0.927	0.959	0.980	0.979	0.950	0.967	0.852
VARI	0.795	0.670	0.690	0.823	0.796	0.897	0.936	0.919	0.759	0.848	0.674	0.646	0.163
GFT	—	—	—	0.976	0.972	—	—	—	—	—	—	—	—
naive	0.923	0.930	0.948	0.852	0.831	0.881	0.881	0.908	0.940	0.925	0.935	0.896	0.833

Table S47. Comparison of different methods for state-level %ILI estimation in North Carolina. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.767	0.889	1.079	0.899	0.990	0.432	1.099	0.776	0.610	1.284	0.394	0.999	2.578
ARGOX	0.786	0.911	1.106	0.902	0.992	0.435	1.114	0.796	0.648	1.335	0.396	1.061	2.631
VARI	1.158	1.427	1.824	1.061	1.161	0.618	1.377	0.729	0.885	2.886	0.855	1.935	3.853
GFT	—	—	—	1.020	1.112	—	—	—	—	—	—	—	—
naive	0.891	1.011	1.202	0.960	1.056	0.495	1.342	0.988	0.775	1.480	0.412	1.077	2.861
MAE													
ARGO-C	0.478	0.513	0.576	0.510	0.606	0.353	0.761	0.596	0.459	0.915	0.301	0.645	1.610
ARGOX	0.491	0.526	0.591	0.512	0.608	0.357	0.776	0.622	0.489	0.966	0.301	0.686	1.644
VARI	0.696	0.815	1.035	0.632	0.709	0.479	1.033	0.585	0.605	2.112	0.641	1.161	2.691
GFT	—	—	—	0.709	0.788	—	—	—	—	—	—	—	—
naive	0.556	0.603	0.690	0.557	0.663	0.405	0.899	0.720	0.604	1.128	0.340	0.799	1.949
Correlation													
ARGO-C	0.875	0.868	0.847	0.870	0.850	0.739	0.670	0.867	0.881	0.835	0.744	0.877	0.700
ARGOX	0.869	0.862	0.840	0.869	0.849	0.733	0.661	0.857	0.862	0.822	0.746	0.857	0.695
VARI	0.726	0.672	0.570	0.887	0.871	0.580	0.567	0.901	0.753	0.331	-0.046	0.704	0.804
GFT	—	—	—	0.834	0.808	—	—	—	—	—	—	—	—
naive	0.839	0.840	0.826	0.859	0.839	0.687	0.575	0.748	0.814	0.793	0.768	0.859	0.675

Table S48. Comparison of different methods for state-level %ILI estimation in North Dakota. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.301	0.343	0.410	0.517	0.568	0.222	0.235	0.347	0.317	0.416	0.187	0.407	0.888
ARGOX	0.321	0.374	0.455	0.580	0.636	0.214	0.239	0.352	0.315	0.478	0.187	0.436	1.015
VARI	1.528	1.458	1.320	1.907	2.055	0.733	2.348	2.549	0.952	2.274	0.248	0.911	2.513
GFT	—	—	—	0.954	1.039	—	—	—	—	—	—	—	—
naive	0.426	0.473	0.549	0.783	0.861	0.255	0.322	0.526	0.379	0.515	0.189	0.515	1.334
MAE													
ARGO-C	0.185	0.210	0.256	0.243	0.278	0.159	0.175	0.251	0.236	0.307	0.138	0.335	0.631
ARGOX	0.190	0.219	0.273	0.284	0.320	0.148	0.178	0.244	0.227	0.316	0.141	0.351	0.742
VARI	0.994	0.892	0.702	1.424	1.548	0.570	1.763	1.807	0.758	1.571	0.200	0.752	1.910
GFT	—	—	—	0.774	0.870	—	—	—	—	—	—	—	—
naive	0.233	0.261	0.314	0.388	0.453	0.192	0.225	0.339	0.268	0.391	0.142	0.408	0.951
Correlation													
ARGO-C	0.951	0.972	0.976	0.925	0.920	0.866	0.961	0.962	0.899	0.939	0.750	0.930	0.952
ARGOX	0.944	0.966	0.970	0.903	0.896	0.880	0.959	0.961	0.899	0.918	0.751	0.916	0.934
VARI	0.777	0.648	0.721	0.431	0.376	0.838	0.882	0.867	0.789	0.656	0.573	0.810	0.727
GFT	—	—	—	0.971	0.969	—	—	—	—	—	—	—	—
naive	0.903	0.946	0.957	0.826	0.811	0.835	0.922	0.910	0.860	0.899	0.744	0.878	0.887

Table S49. Comparison of different methods for state-level %ILI estimation in Ohio. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.955	0.901	0.790	1.444	1.576	0.890	1.552	0.853	0.725	1.018	0.579	1.014	1.099
ARGOX	0.933	0.889	0.801	1.435	1.564	0.887	1.500	0.733	0.747	1.068	0.577	0.996	1.121
VARI	3.239	2.751	1.465	4.118	4.533	2.011	5.221	3.008	3.988	4.228	0.960	1.142	1.758
GFT	—	—	—	2.056	2.208	—	—	—	—	—	—	—	—
naive	0.987	0.925	0.796	1.375	1.508	0.887	1.457	1.015	0.994	1.176	0.618	1.002	1.130
MAE													
ARGO-C	0.601	0.573	0.523	0.982	1.119	0.726	1.087	0.598	0.495	0.685	0.451	0.656	0.761
ARGOX	0.589	0.569	0.532	0.996	1.126	0.724	1.050	0.536	0.508	0.707	0.456	0.640	0.816
VARI	2.027	1.678	1.032	2.770	3.328	1.801	3.975	2.103	3.007	3.647	0.784	0.973	1.506
GFT	—	—	—	1.424	1.504	—	—	—	—	—	—	—	—
naive	0.650	0.609	0.533	0.970	1.123	0.716	1.091	0.763	0.706	0.812	0.500	0.692	0.916
Correlation													
ARGO-C	0.952	0.947	0.915	0.914	0.892	0.506	0.921	0.972	0.973	0.941	0.774	0.533	0.915
ARGOX	0.955	0.949	0.913	0.917	0.897	0.498	0.925	0.978	0.971	0.935	0.777	0.529	0.909
VARI	0.709	0.491	0.709	0.541	0.479	0.257	0.743	0.765	0.513	0.864	0.571	0.431	0.869
GFT	—	—	—	0.842	0.809	—	—	—	—	—	—	—	—
naive	0.950	0.946	0.919	0.923	0.904	0.535	0.931	0.941	0.940	0.921	0.751	0.534	0.898

Table S50. Comparison of different methods for state-level %ILI estimation in Oklahoma. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.428	0.397	0.332	0.371	0.402	0.569	0.817	0.339	0.358	0.413	0.158	0.222	0.775
ARGOX	0.425	0.393	0.324	0.368	0.399	0.579	0.810	0.290	0.367	0.387	0.144	0.197	0.804
VARI	2.425	2.061	1.102	3.854	4.239	1.258	3.884	2.004	2.512	2.624	0.986	1.147	1.567
GFT	—	—	—	0.570	0.487	—	—	—	—	—	—	—	—
naive	0.500	0.478	0.433	0.323	0.348	0.632	0.946	0.442	0.450	0.757	0.127	0.283	0.891
MAE													
ARGO-C	0.260	0.237	0.195	0.275	0.306	0.419	0.465	0.231	0.253	0.296	0.120	0.177	0.580
ARGOX	0.258	0.234	0.188	0.273	0.303	0.429	0.464	0.205	0.265	0.297	0.114	0.161	0.589
VARI	1.527	1.253	0.745	2.792	3.300	1.129	2.993	1.324	2.038	2.044	0.781	0.927	1.275
GFT	—	—	—	0.473	0.396	—	—	—	—	—	—	—	—
naive	0.292	0.273	0.238	0.254	0.280	0.428	0.560	0.319	0.275	0.580	0.097	0.189	0.672
Correlation													
ARGO-C	0.940	0.945	0.959	0.794	0.726	0.741	0.781	0.983	0.953	0.969	0.938	0.919	0.900
ARGOX	0.940	0.946	0.960	0.795	0.724	0.731	0.781	0.986	0.952	0.973	0.945	0.936	0.889
VARI	0.727	0.720	0.823	0.593	0.469	0.658	0.579	0.917	0.856	0.713	0.533	0.405	0.763
GFT	—	—	—	0.755	0.822	—	—	—	—	—	—	—	—
naive	0.918	0.922	0.931	0.799	0.737	0.713	0.728	0.950	0.919	0.892	0.939	0.874	0.867

Table S51. Comparison of different methods for state-level %ILI estimation in Oregon. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.328	0.306	0.262	0.492	0.529	0.341	0.431	0.413	0.190	0.442	0.143	0.261	0.322
ARGOX	0.336	0.309	0.250	0.535	0.575	0.332	0.456	0.388	0.190	0.404	0.131	0.264	0.352
VARI	1.117	0.962	0.574	2.077	2.247	1.147	1.112	1.065	0.706	0.878	0.209	0.670	1.090
GFT	—	—	—	0.589	0.640	—	—	—	—	—	—	—	—
naive	0.465	0.429	0.354	0.764	0.832	0.454	0.564	0.573	0.286	0.673	0.138	0.340	0.498
MAE													
ARGO-C	0.216	0.193	0.151	0.340	0.365	0.250	0.314	0.267	0.146	0.241	0.105	0.183	0.241
ARGOX	0.216	0.191	0.145	0.366	0.393	0.241	0.329	0.235	0.153	0.235	0.097	0.169	0.259
VARI	0.751	0.626	0.392	1.605	1.757	1.019	0.859	0.587	0.546	0.660	0.168	0.491	0.981
GFT	—	—	—	0.480	0.535	—	—	—	—	—	—	—	—
naive	0.288	0.256	0.197	0.504	0.565	0.336	0.396	0.368	0.212	0.500	0.101	0.211	0.376
Correlation													
ARGO-C	0.969	0.969	0.967	0.951	0.949	0.871	0.940	0.977	0.977	0.961	0.917	0.965	0.958
ARGOX	0.968	0.969	0.971	0.943	0.940	0.877	0.932	0.982	0.978	0.969	0.926	0.956	0.948
VARI	0.707	0.719	0.871	0.714	0.678	0.669	0.705	0.808	0.746	0.845	0.824	0.412	0.805
GFT	—	—	—	0.946	0.942	—	—	—	—	—	—	—	—
naive	0.937	0.940	0.943	0.880	0.867	0.793	0.900	0.944	0.943	0.904	0.915	0.869	0.892

Table S52. Comparison of different methods for state-level %ILI estimation in Pennsylvania. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.445	0.473	0.522	0.475	0.522	0.225	0.471	0.639	0.509	0.911	0.149	0.595	0.899
ARGOX	0.457	0.488	0.539	0.483	0.531	0.232	0.486	0.638	0.520	1.003	0.144	0.579	0.897
VARI	1.227	1.181	1.090	1.614	1.707	1.538	0.914	0.907	1.226	2.481	0.523	0.790	1.319
GFT	—	—	—	0.674	0.709	—	—	—	—	—	—	—	—
naive	0.542	0.528	0.501	0.475	0.522	0.239	0.542	0.825	0.654	0.959	0.135	0.577	1.051
MAE													
ARGO-C	0.260	0.265	0.276	0.261	0.305	0.164	0.315	0.468	0.362	0.556	0.121	0.409	0.595
ARGOX	0.266	0.270	0.279	0.270	0.313	0.169	0.325	0.465	0.372	0.607	0.118	0.396	0.625
VARI	0.976	0.877	0.693	1.376	1.445	1.480	0.785	0.696	0.913	1.910	0.466	0.596	1.021
GFT	—	—	—	0.593	0.619	—	—	—	—	—	—	—	—
naive	0.304	0.289	0.261	0.285	0.335	0.153	0.365	0.578	0.490	0.618	0.105	0.376	0.725
Correlation													
ARGO-C	0.969	0.954	0.906	0.916	0.906	0.920	0.935	0.952	0.963	0.950	0.776	0.626	0.928
ARGOX	0.967	0.951	0.900	0.914	0.905	0.913	0.928	0.953	0.960	0.940	0.802	0.626	0.928
VARI	0.787	0.746	0.630	0.763	0.735	0.767	0.890	0.900	0.867	0.549	0.702	0.333	0.933
GFT	—	—	—	0.943	0.937	—	—	—	—	—	—	—	—
naive	0.951	0.944	0.919	0.911	0.901	0.907	0.904	0.915	0.936	0.946	0.746	0.589	0.890

Table S53. Comparison of different methods for state-level %ILI estimation in Rhode Island. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.567	0.583	0.612	0.512	0.560	0.332	0.913	0.679	0.689	0.790	0.338	0.316	1.498
ARGOX	0.594	0.620	0.665	0.593	0.648	0.330	0.942	0.709	0.682	0.858	0.320	0.319	1.683
VARI	2.333	2.324	2.308	1.559	1.715	0.799	2.941	4.113	2.165	4.152	1.123	1.449	5.240
GFT	—	—	—	1.584	1.725	—	—	—	—	—	—	—	—
naive	0.875	0.841	0.775	0.796	0.873	0.376	1.006	1.358	0.933	1.589	0.246	0.503	1.788
MAE													
ARGO-C	0.347	0.352	0.362	0.265	0.301	0.258	0.733	0.447	0.433	0.578	0.270	0.238	1.081
ARGOX	0.363	0.364	0.366	0.284	0.317	0.261	0.763	0.457	0.450	0.622	0.243	0.222	1.258
VARI	1.471	1.486	1.513	0.834	0.989	0.633	2.505	3.045	1.807	3.584	0.987	1.136	3.904
GFT	—	—	—	1.319	1.483	—	—	—	—	—	—	—	—
naive	0.514	0.481	0.420	0.421	0.481	0.292	0.795	0.903	0.683	1.174	0.195	0.342	1.365
Correlation													
ARGO-C	0.980	0.975	0.961	0.944	0.939	0.855	0.913	0.985	0.961	0.970	0.920	0.979	0.849
ARGOX	0.978	0.972	0.955	0.919	0.910	0.856	0.906	0.984	0.958	0.965	0.931	0.971	0.813
VARI	0.782	0.737	0.596	0.295	0.208	0.600	0.711	0.751	0.840	0.753	0.758	0.563	-0.169
GFT	—	—	—	0.977	0.975	—	—	—	—	—	—	—	—
naive	0.952	0.949	0.939	0.851	0.834	0.823	0.888	0.939	0.915	0.878	0.930	0.905	0.791

Table S54. Comparison of different methods for state-level %ILI estimation in South Carolina. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.326	0.306	0.265	0.315	0.329	0.292	0.418	0.492	0.358	0.355	0.182	0.283	0.430
ARGOX	0.329	0.310	0.270	0.318	0.332	0.295	0.425	0.495	0.362	0.368	0.179	0.288	0.443
VARI	2.064	2.162	2.333	0.832	0.888	0.722	1.882	3.319	2.914	4.225	1.169	1.176	5.356
GFT	—	—	—	1.033	1.132	—	—	—	—	—	—	—	—
naive	0.354	0.330	0.282	0.362	0.380	0.309	0.427	0.542	0.376	0.403	0.184	0.302	0.468
MAE													
ARGO-C	0.233	0.215	0.183	0.235	0.243	0.218	0.297	0.369	0.279	0.233	0.148	0.204	0.336
ARGOX	0.235	0.217	0.184	0.238	0.245	0.217	0.299	0.371	0.283	0.247	0.147	0.209	0.350
VARI	1.264	1.352	1.514	0.588	0.615	0.559	1.481	2.117	2.444	3.700	0.978	0.965	4.631
GFT	—	—	—	0.512	0.584	—	—	—	—	—	—	—	—
naive	0.257	0.233	0.188	0.263	0.270	0.244	0.318	0.428	0.284	0.257	0.154	0.226	0.367
Correlation													
ARGO-C	0.950	0.950	0.937	0.921	0.912	0.851	0.925	0.923	0.876	0.960	0.381	0.907	0.905
ARGOX	0.949	0.949	0.933	0.919	0.910	0.847	0.923	0.922	0.873	0.957	0.380	0.902	0.898
VARI	0.812	0.750	0.702	0.806	0.780	0.434	0.759	0.906	0.851	0.849	-0.145	0.803	0.359
GFT	—	—	—	0.898	0.886	—	—	—	—	—	—	—	—
naive	0.941	0.941	0.925	0.898	0.885	0.840	0.924	0.903	0.844	0.948	0.378	0.893	0.880

Table S55. Comparison of different methods for state-level %ILI estimation in South Dakota. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.703	0.683	0.644	0.572	0.620	0.478	0.883	1.061	0.467	1.195	0.232	0.400	1.620
ARGOX	0.728	0.718	0.698	0.591	0.641	0.482	0.870	1.026	0.505	1.369	0.222	0.368	1.775
VARI	1.943	2.166	2.529	1.593	1.736	0.667	1.983	1.858	1.331	4.675	0.810	1.494	6.530
GFT	—	—	—	1.239	1.341	—	—	—	—	—	—	—	—
naive	0.866	0.820	0.727	0.890	0.974	0.507	0.991	1.212	0.707	1.546	0.126	0.512	1.837
MAE													
ARGO-C	0.472	0.429	0.350	0.410	0.453	0.391	0.684	0.670	0.360	0.896	0.176	0.249	1.273
ARGOX	0.477	0.438	0.365	0.407	0.451	0.394	0.663	0.651	0.366	0.981	0.171	0.249	1.429
VARI	1.145	1.311	1.618	0.970	1.078	0.528	1.746	1.462	0.940	3.731	0.743	1.289	5.572
GFT	—	—	—	0.990	1.089	—	—	—	—	—	—	—	—
naive	0.507	0.464	0.385	0.534	0.612	0.358	0.711	0.726	0.522	1.055	0.103	0.362	1.494
Correlation													
ARGO-C	0.951	0.954	0.961	0.954	0.950	0.844	0.857	0.907	0.950	0.944	0.914	0.972	0.891
ARGOX	0.947	0.950	0.955	0.952	0.948	0.844	0.862	0.915	0.936	0.926	0.918	0.968	0.870
VARI	0.658	0.604	0.672	0.695	0.656	0.770	0.887	0.631	0.732	0.725	0.350	0.823	0.225
GFT	—	—	—	0.966	0.964	—	—	—	—	—	—	—	—
naive	0.927	0.936	0.951	0.891	0.879	0.836	0.822	0.866	0.869	0.898	0.932	0.902	0.861

Table S56. Comparison of different methods for state-level %ILI estimation in Tennessee. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.851	0.751	0.520	1.473	1.609	0.564	0.761	1.087	0.647	1.147	0.342	0.386	0.816
ARGOX	0.852	0.763	0.565	1.466	1.603	0.552	0.792	1.049	0.623	1.301	0.347	0.360	0.840
VARI	2.766	2.323	1.106	3.051	3.298	2.776	1.873	4.760	3.334	2.524	0.391	0.889	2.359
GFT	—	—	—	1.253	1.368	—	—	—	—	—	—	—	—
naive	0.996	0.871	0.570	1.544	1.690	0.577	0.796	1.398	0.822	1.536	0.338	0.555	0.841
MAE													
ARGO-C	0.509	0.447	0.331	0.848	0.953	0.451	0.574	0.664	0.443	0.833	0.264	0.270	0.660
ARGOX	0.509	0.448	0.336	0.870	0.989	0.448	0.590	0.671	0.415	0.868	0.256	0.271	0.715
VARI	2.134	1.637	0.717	2.567	2.819	2.411	1.319	3.934	2.974	2.063	0.276	0.736	1.752
GFT	—	—	—	0.835	0.951	—	—	—	—	—	—	—	—
naive	0.595	0.513	0.359	0.893	1.016	0.454	0.577	0.957	0.634	1.100	0.218	0.378	0.690
Correlation													
ARGO-C	0.961	0.963	0.967	0.860	0.837	0.824	0.953	0.970	0.973	0.955	0.568	0.928	0.923
ARGOX	0.961	0.962	0.962	0.861	0.837	0.830	0.949	0.972	0.975	0.942	0.571	0.937	0.918
VARI	0.820	0.763	0.902	0.848	0.831	-0.195	0.799	0.886	0.893	0.836	0.528	0.887	0.876
GFT	—	—	—	0.928	0.916	—	—	—	—	—	—	—	—
naive	0.947	0.951	0.958	0.854	0.832	0.816	0.945	0.949	0.946	0.914	0.513	0.854	0.909

Table S57. Comparison of different methods for state-level %ILI estimation in Texas. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.430	0.410	0.372	0.296	0.312	0.330	0.413	0.397	0.619	0.803	0.181	0.355	0.398
ARGOX	0.445	0.425	0.386	0.300	0.317	0.375	0.449	0.392	0.627	0.857	0.170	0.338	0.458
VARI	3.032	2.584	1.417	2.404	2.607	1.991	3.754	6.236	1.856	3.097	0.660	0.830	3.268
GFT	—	—	—	1.083	1.163	—	—	—	—	—	—	—	—
naive	0.507	0.489	0.454	0.512	0.550	0.429	0.439	0.451	0.658	0.997	0.154	0.371	0.695
MAE													
ARGO-C	0.296	0.280	0.251	0.228	0.240	0.263	0.293	0.301	0.474	0.583	0.148	0.277	0.315
ARGOX	0.302	0.282	0.246	0.233	0.244	0.289	0.307	0.296	0.477	0.593	0.135	0.260	0.348
VARI	2.105	1.703	0.957	1.795	1.977	1.850	2.877	5.066	1.532	2.543	0.583	0.503	2.991
GFT	—	—	—	1.002	1.106	—	—	—	—	—	—	—	—
naive	0.334	0.313	0.273	0.355	0.381	0.329	0.289	0.304	0.515	0.724	0.124	0.291	0.517
Correlation													
ARGO-C	0.962	0.960	0.957	0.970	0.965	0.930	0.909	0.891	0.938	0.938	0.601	0.884	0.971
ARGOX	0.959	0.957	0.955	0.970	0.965	0.909	0.889	0.893	0.935	0.930	0.617	0.876	0.957
VARI	0.693	0.702	0.842	0.920	0.914	0.651	0.048	0.853	0.934	0.906	0.335	0.531	0.734
GFT	—	—	—	0.951	0.957	—	—	—	—	—	—	—	—
naive	0.947	0.945	0.937	0.911	0.894	0.885	0.889	0.868	0.921	0.899	0.541	0.809	0.894

Table S58. Comparison of different methods for state-level %ILI estimation in Utah. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.493	0.514	0.550	0.753	0.810	0.430	0.325	0.537	0.540	0.545	0.230	0.246	1.456
ARGOX	0.493	0.515	0.554	0.754	0.809	0.429	0.320	0.533	0.539	0.555	0.232	0.248	1.468
VARI	1.101	1.103	1.106	1.289	1.355	0.591	0.959	1.417	1.214	1.463	0.335	1.746	1.350
GFT	—	—	—	1.220	1.330	—	—	—	—	—	—	—	—
naive	0.536	0.531	0.520	0.820	0.896	0.469	0.357	0.579	0.590	0.619	0.246	0.224	1.325
MAE													
ARGO-C	0.348	0.320	0.269	0.517	0.557	0.324	0.257	0.405	0.406	0.387	0.174	0.170	0.910
ARGOX	0.348	0.320	0.268	0.518	0.553	0.324	0.254	0.406	0.409	0.389	0.175	0.168	0.915
VARI	0.854	0.867	0.891	0.908	0.907	0.488	0.781	1.166	1.008	1.167	0.292	1.685	1.210
GFT	—	—	—	0.833	0.938	—	—	—	—	—	—	—	—
naive	0.381	0.342	0.271	0.560	0.639	0.360	0.264	0.450	0.447	0.445	0.187	0.165	0.845
Correlation													
ARGO-C	0.914	0.902	0.863	0.904	0.872	0.790	0.797	0.920	0.878	0.934	0.716	0.773	0.691
ARGOX	0.914	0.901	0.863	0.904	0.872	0.788	0.798	0.921	0.878	0.932	0.712	0.763	0.691
VARI	0.618	0.616	0.730	0.763	0.693	0.811	0.075	0.834	0.745	0.867	0.594	0.560	0.801
GFT	—	—	—	0.836	0.814	—	—	—	—	—	—	—	—
naive	0.901	0.897	0.873	0.884	0.849	0.753	0.754	0.910	0.859	0.918	0.691	0.743	0.717

Table S59. Comparison of different methods for state-level %ILI estimation in Vermont. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.418	0.439	0.475	0.623	0.684	0.251	0.476	0.658	0.431	0.364	0.149	0.483	0.885
ARGOX	0.431	0.467	0.528	0.682	0.748	0.233	0.495	0.636	0.418	0.469	0.136	0.596	0.984
VARI	0.638	0.761	0.947	0.933	1.026	0.289	0.688	1.023	0.565	0.926	0.118	0.938	2.222
GFT	—	—	—	1.045	1.147	—	—	—	—	—	—	—	—
naive	0.628	0.636	0.650	0.947	1.042	0.290	0.646	0.990	0.561	0.914	0.120	0.849	1.050
MAE													
ARGO-C	0.238	0.240	0.243	0.300	0.349	0.189	0.319	0.443	0.292	0.266	0.124	0.269	0.682
ARGOX	0.243	0.254	0.273	0.328	0.377	0.172	0.331	0.416	0.286	0.337	0.113	0.345	0.776
VARI	0.330	0.377	0.464	0.437	0.510	0.232	0.396	0.624	0.416	0.667	0.097	0.542	1.722
GFT	—	—	—	0.519	0.593	—	—	—	—	—	—	—	—
naive	0.332	0.333	0.335	0.466	0.547	0.240	0.381	0.613	0.411	0.674	0.098	0.497	0.837
Correlation													
ARGO-C	0.980	0.976	0.971	0.957	0.953	0.918	0.942	0.980	0.978	0.991	0.888	0.945	0.927
ARGOX	0.979	0.973	0.965	0.945	0.939	0.936	0.936	0.981	0.977	0.985	0.908	0.915	0.912
VARI	0.951	0.928	0.895	0.885	0.872	0.887	0.871	0.939	0.954	0.930	0.921	0.816	0.776
GFT	—	—	—	0.976	0.974	—	—	—	—	—	—	—	—
naive	0.953	0.950	0.946	0.886	0.873	0.889	0.885	0.939	0.956	0.933	0.920	0.836	0.884

Table S60. Comparison of different methods for state-level %ILI estimation in Virginia. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.420	0.487	0.590	0.347	0.380	0.323	0.527	0.658	0.428	0.602	0.149	0.298	1.650
ARGOX	0.433	0.516	0.643	0.347	0.379	0.315	0.568	0.663	0.443	0.733	0.139	0.252	1.762
VARI	1.241	1.477	1.837	1.384	1.517	0.701	1.061	1.381	0.992	2.428	0.267	1.054	4.934
GFT	—	—	—	0.784	0.840	—	—	—	—	—	—	—	—
naive	0.494	0.583	0.721	0.393	0.430	0.338	0.568	0.753	0.549	0.976	0.122	0.294	1.896
MAE													
ARGO-C	0.258	0.256	0.254	0.233	0.263	0.267	0.347	0.370	0.277	0.454	0.112	0.237	1.025
ARGOX	0.258	0.260	0.263	0.236	0.267	0.262	0.365	0.352	0.286	0.510	0.103	0.204	1.097
VARI	0.907	0.901	0.890	1.043	1.207	0.637	0.827	1.039	0.896	1.799	0.203	0.915	3.355
GFT	—	—	—	0.695	0.759	—	—	—	—	—	—	—	—
naive	0.295	0.302	0.315	0.284	0.325	0.283	0.352	0.422	0.358	0.697	0.099	0.211	1.384
Correlation													
ARGO-C	0.950	0.955	0.956	0.939	0.933	0.824	0.845	0.877	0.925	0.965	0.642	0.874	0.913
ARGOX	0.947	0.949	0.948	0.940	0.934	0.831	0.817	0.876	0.920	0.951	0.661	0.910	0.897
VARI	0.521	0.454	0.518	0.617	0.549	0.604	0.012	0.373	0.731	0.362	0.557	0.508	0.225
GFT	—	—	—	0.961	0.959	—	—	—	—	—	—	—	—
naive	0.932	0.936	0.936	0.924	0.915	0.809	0.834	0.847	0.876	0.903	0.621	0.881	0.884

Table S61. Comparison of different methods for state-level %ILI estimation in Washington. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.499	0.504	0.512	0.621	0.678	0.367	0.399	0.818	0.627	0.678	0.393	0.461	0.744
ARGOX	0.519	0.523	0.530	0.633	0.690	0.357	0.389	0.870	0.614	0.815	0.384	0.447	0.760
VARI	1.344	1.442	1.607	1.742	1.916	0.963	1.270	2.004	1.517	1.859	0.387	0.815	4.273
GFT	—	—	—	1.002	1.004	—	—	—	—	—	—	—	—
naive	0.645	0.611	0.543	0.765	0.838	0.391	0.426	1.117	0.692	0.968	0.435	0.431	0.836
MAE													
ARGO-C	0.312	0.319	0.331	0.412	0.459	0.309	0.331	0.509	0.406	0.396	0.232	0.360	0.609
ARGOX	0.320	0.325	0.333	0.405	0.450	0.295	0.316	0.548	0.419	0.487	0.223	0.332	0.625
VARI	0.843	0.832	0.811	1.151	1.360	0.868	0.902	1.318	1.118	1.453	0.210	0.647	2.912
GFT	—	—	—	0.915	0.900	—	—	—	—	—	—	—	—
naive	0.356	0.352	0.346	0.491	0.563	0.281	0.298	0.665	0.437	0.659	0.261	0.285	0.703
Correlation													
ARGO-C	0.958	0.946	0.892	0.958	0.953	0.798	0.931	0.931	0.923	0.955	0.368	0.333	0.779
ARGOX	0.954	0.941	0.887	0.956	0.951	0.811	0.935	0.922	0.927	0.934	0.403	0.297	0.771
VARI	0.737	0.595	0.723	0.900	0.886	0.728	0.455	0.748	0.695	0.630	0.245	-0.214	0.627
GFT	—	—	—	0.966	0.961	—	—	—	—	—	—	—	—
naive	0.930	0.922	0.884	0.930	0.919	0.798	0.921	0.872	0.907	0.910	0.317	0.396	0.712

Table S62. Comparison of different methods for state-level %ILI estimation in West Virginia. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.371	0.357	0.329	0.529	0.562	0.333	0.396	0.385	0.270	0.410	0.126	0.209	0.394
ARGOX	0.372	0.356	0.324	0.555	0.596	0.330	0.380	0.354	0.267	0.405	0.131	0.203	0.404
VARI	0.474	0.445	0.386	0.759	0.827	0.351	0.447	0.552	0.297	0.616	0.089	0.232	0.505
GFT	—	—	—	0.784	0.842	—	—	—	—	—	—	—	—
naive	0.446	0.428	0.392	0.717	0.781	0.347	0.424	0.459	0.294	0.575	0.085	0.236	0.486
MAE													
ARGO-C	0.273	0.243	0.186	0.407	0.430	0.278	0.322	0.295	0.212	0.346	0.097	0.169	0.312
ARGOX	0.271	0.239	0.181	0.415	0.445	0.268	0.319	0.269	0.211	0.334	0.103	0.160	0.314
VARI	0.322	0.278	0.196	0.467	0.519	0.299	0.360	0.415	0.237	0.461	0.073	0.171	0.409
GFT	—	—	—	0.635	0.686	—	—	—	—	—	—	—	—
naive	0.305	0.266	0.192	0.461	0.517	0.298	0.344	0.353	0.224	0.414	0.067	0.177	0.383
Correlation													
ARGO-C	0.953	0.954	0.954	0.935	0.909	0.737	0.887	0.947	0.894	0.965	0.919	0.896	0.939
ARGOX	0.953	0.954	0.955	0.929	0.896	0.745	0.895	0.956	0.897	0.969	0.913	0.897	0.934
VARI	0.931	0.931	0.937	0.881	0.816	0.738	0.866	0.915	0.880	0.917	0.931	0.798	0.903
GFT	—	—	—	0.896	0.914	—	—	—	—	—	—	—	—
naive	0.933	0.936	0.938	0.883	0.822	0.743	0.869	0.921	0.885	0.923	0.932	0.799	0.905

Table S63. Comparison of different methods for state-level %ILI estimation in Wisconsin. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.487	0.613	0.795	0.357	0.392	0.348	0.533	0.728	0.735	0.773	0.254	0.795	1.812
ARGOX	0.493	0.607	0.776	0.342	0.374	0.348	0.558	0.695	0.745	0.793	0.240	0.763	1.792
VARI	0.909	0.994	1.134	1.372	1.500	0.779	0.860	1.139	0.815	1.230	0.376	1.416	2.262
GFT	—	—	—	0.588	0.596	—	—	—	—	—	—	—	—
naive	0.551	0.685	0.881	0.357	0.391	0.373	0.582	0.810	0.825	0.929	0.256	0.802	2.118
MAE													
ARGO-C	0.324	0.374	0.468	0.280	0.327	0.233	0.390	0.509	0.559	0.575	0.184	0.582	1.307
ARGOX	0.326	0.373	0.460	0.275	0.318	0.237	0.416	0.490	0.558	0.602	0.175	0.565	1.307
VARI	0.644	0.666	0.707	0.917	1.031	0.652	0.668	0.912	0.623	0.883	0.235	1.042	1.739
GFT	—	—	—	0.503	0.496	—	—	—	—	—	—	—	—
naive	0.353	0.411	0.520	0.266	0.310	0.265	0.448	0.601	0.593	0.696	0.199	0.566	1.653
Correlation													
ARGO-C	0.956	0.926	0.866	0.939	0.918	0.891	0.862	0.954	0.911	0.913	0.281	0.696	0.724
ARGOX	0.955	0.928	0.873	0.945	0.927	0.891	0.850	0.959	0.908	0.908	0.297	0.722	0.727
VARI	0.847	0.796	0.744	0.900	0.886	0.710	0.632	0.912	0.888	0.810	0.241	0.517	0.882
GFT	—	—	—	0.877	0.856	—	—	—	—	—	—	—	—
naive	0.943	0.911	0.847	0.940	0.921	0.881	0.842	0.938	0.890	0.871	0.159	0.731	0.656

Table S64. Comparison of different methods for state-level %ILI estimation in Wyoming. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.

	Whole period	'14-'23	post-COVID	GFT period	'14-'15	'15-'16	'16-'17	'17-'18	'18-'19	'19-'20	'20-'21	'21-'22	'22-'23
RMSE													
ARGO-C	0.296	0.520	0.780	0.205	0.217	0.196	0.357	0.516	0.196	1.406	0.192	0.571	1.128
ARGOX	0.308	0.527	0.786	0.202	0.215	0.204	0.392	0.528	0.214	1.397	0.165	0.596	1.186
VARI	1.284	1.751	2.386	0.941	0.910	1.384	1.391	1.102	1.524	3.637	0.386	1.543	4.954
GFT	—	—	—	0.531	0.554	—	—	—	—	—	—	—	—
naive	0.406	0.609	0.868	0.231	0.243	0.262	0.393	0.773	0.324	1.585	0.085	0.641	1.338
MAE													
ARGO-C	0.197	0.255	0.361	0.163	0.173	0.163	0.268	0.359	0.165	0.751	0.150	0.338	0.905
ARGOX	0.201	0.257	0.360	0.160	0.170	0.173	0.283	0.355	0.177	0.756	0.133	0.345	0.978
VARI	1.092	1.202	1.406	0.841	0.793	1.250	1.161	0.905	1.391	2.641	0.339	1.197	4.388
GFT	—	—	—	0.465	0.487	—	—	—	—	—	—	—	—
naive	0.240	0.303	0.421	0.180	0.188	0.207	0.273	0.466	0.246	1.105	0.069	0.365	1.100
Correlation													
ARGO-C	0.978	0.968	0.964	0.958	0.950	0.959	0.922	0.970	0.975	0.912	0.363	0.907	0.947
ARGOX	0.976	0.967	0.963	0.960	0.952	0.956	0.904	0.969	0.970	0.916	0.375	0.893	0.939
VARI	0.804	0.715	0.734	0.824	0.771	0.584	0.571	0.854	0.498	0.655	-0.005	-0.215	0.636
GFT	—	—	—	0.953	0.955	—	—	—	—	—	—	—	—
naive	0.959	0.955	0.953	0.939	0.929	0.929	0.902	0.927	0.929	0.879	0.696	0.815	0.909

Table S65. Comparison of different methods for state-level %ILI estimation in New York NY. The MSE, MAE, and correlation are reported. The method with the best performance is highlighted in boldface for each metric in each period.