

New Solutions of Coupled Nonlocal NLS and Coupled Nonlocal mKdV Equations

Avinash Khare

Physics Department, Savitribai Phule Pune University
Pune 411007, India

Avadh Saxena

Theoretical Division and Center for Nonlinear Studies, Los Alamos
National Laboratory, Los Alamos, New Mexico 87545, USA

Abstract

We provide several novel solutions of the coupled Ablowitz-Musslimani (AM) version of the nonlocal nonlinear Schrödinger (NLS) equation and the coupled nonlocal modified Korteweg-de Vries (mKdV) equations. In each case we compare and contrast the corresponding solutions of the relevant coupled local equations. Further, we provide new solutions of the coupled local NLS and the coupled local mKdV equations which are not the solutions of the corresponding nonlocal equations.

1 Introduction

In recent years the nonlocal nonlinear equations have received considerable attention in the literature. It primarily started with the papers of Ablowitz and Musslimani (AM) [1, 2] about the nonlocal NLS equation and its integrability [3]. Subsequently, several other nonlocal equations have been proposed [4, 5, 6, 7, 8]. It is hoped that some of these nonlocal equations may have applications in areas of physics such as optics, photonics, etc. [9]. Besides one hopes to understand the nature of the relevant nonlocality. It is clearly possible that in some cases coupled nonlocal equations could be relevant in a number of physical settings.

It is with this motivation that recently we have studied coupled AM type nonlocal NLS as well as nonlocal coupled mKdV equations [10, 11, 12] and obtained a large number of solutions of these coupled equations. We thought that perhaps one has essentially covered most of the allowed solutions. Of course, we were aware that nonlinear equations being rich one can never be sure about it. Recently, we were able to obtain new solutions of the uncoupled symmetric ϕ^4 equation [13] as well as the coupled ϕ^4 equations [14] and that has inspired us to look for similar solutions of the

coupled nonlocal equations. The purpose of this paper is to report on several new solutions of the coupled AM variant of the nonlocal NLS as well as the coupled nonlocal mKdV equations [4, 5]. In each case we also enquire if the corresponding local equations also admit such a solution, and if yes, under what conditions.

The plan of the paper is the following. In Sec. II we present new solutions of the coupled nonlocal AM variant of the NLS equation. In each case we compare and contrast with the solutions of the (local) coupled NLS equations (in case they are admitted). Particular mention may be made of a truncated coupled nonlocal NLS equation which admits novel nonreciprocal solutions where one field is in terms of the Lamé polynomial of order two while the other field is in terms of the Lamé polynomial of order one. It may be noted that the full coupled nonlocal or local NLS equations do not admit similar solutions. In Appendix A we present those solutions of the local coupled NLS equations which are not the solutions of the nonlocal coupled NLS equations. In Sec. III we present new solutions of the nonlocal coupled mKdV equations and compare and contrast them with the corresponding solutions of the local coupled mKdV equations. Particular mention may be made of the plane wave solutions as well as the soliton solutions multiplied by the plane wave factor of the coupled nonlocal mKdV equations. In contrast, the corresponding local coupled mKdV equations do not admit either the plane wave or the soliton solutions multiplied by the plane wave factor. In Appendix B we present those solutions of the local coupled mKdV equations which are not the solutions of the nonlocal coupled mKdV.

2 Novel Solutions of a Coupled nonlocal Ablowitz-Musslimani NLS Model

Let us consider the following coupled nonlocal Ablowitz-Musslimani NLS model [11, 12]

$$iu_t(x, t) + u_{xx}(x, t) + [g_{11}u(x, t)u^*(-x, t) + g_{12}v(x, t)v^*(-x, t)]u(x, t) = 0, \quad (1)$$

$$iv_t(x, t) + v_{xx}(x, t) + [g_{21}u(x, t)u^*(-x, t) + g_{22}v(x, t)v^*(-x, t)]v(x, t) = 0. \quad (2)$$

In order to find exact solutions to the coupled Eqs. (1) and (2), we start with the ansatz

$$u(x, t) = e^{i\omega_1(t+t_0)}u(x), \quad v(x, t) = e^{i\omega_2(t+t_1)}, \quad (3)$$

where t_0, t_1 are arbitrary real constants. In that case Eqs. (1) and (2) take the form

$$u_{xx}(x) = \omega_1 u(x) - [g_{11}u(x)u^*(-x) + g_{12}v(x)v^*(-x)]u(x), \quad (4)$$

$$v_{xx}(x) = \omega_2 v(x) - [g_{21}u(x)u^*(-x) + g_{22}v(x)v^*(-x)]v(x). \quad (5)$$

For simplicity, now onwards we will put $t_0, t_1 = 0$. However, it is understood that such a shift in time t is always allowed. However, in view of the nonlocality, similar shift in x is not allowed. This is unlike the local NLS case where such a shift is allowed.

We now discuss new solutions of these coupled equations and point out whether the corresponding coupled local NLS equations

$$iu_t(x, t) + u_{xx}(x, t) + [g_{11}u^2(x, t) + g_{12}v^2(x, t)]u(x, t) = 0, \quad (6)$$

$$iv_t(x, t) + v_{xx}(x, t) + [g_{21}u^2(x, t) + g_{22}v^2(x, t)]v(x, t) = 0, \quad (7)$$

also admit these solutions and if yes under what conditions. In Appendix A, we discuss those solutions which are admitted by the local coupled Eqs. (6) and (7) but not by the coupled nonlocal Eqs. (1) and (2).

Solution I

It is not difficult to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A\sqrt{m}\operatorname{sn}(\beta x, m)}{B + \operatorname{dn}(\beta x, m)}, \quad v(x, t) = e^{i\omega_2 t} \frac{D\sqrt{m}\operatorname{cn}(\beta x, m)}{B + \operatorname{dn}(\beta x, m)}, \quad (8)$$

with $B > 0$, is an exact solution of the coupled Eqs. (1) and (2) provided

$$\begin{aligned} \omega_1 = \omega_2 = -\frac{(2-m)\beta^2}{2}, \quad g_{12}D^2 = 3g_{22}D^2 = \frac{3(B^2-1)\beta^2}{2}, \\ 3g_{11}A^2 = g_{21}A^2 = \frac{3(B^2-1+m)\beta^2}{2}. \end{aligned} \quad (9)$$

Thus $g_{12}, g_{22} > (<) 0$ provided $B^2 > (<) 1$ while $g_{11}, g_{21} > (<) 0$ provided $B^2 < (>) 1 - m$. On the other hand $\omega_1, \omega_2 < 0$. Here m is the modulus of the Jacobi elliptic functions [15]. It may be noted that $B > 0$ is true for all the thirteen solutions given below.

It is worth pointing out that the local coupled NLS Eqs. (6) and (7) also admit the solution (8) provided relations as given by Eq. (9) are satisfied except that g_{11} and g_{21} have opposite values compared to those given in (9).

Solution II

In the limit $m = 1$, the solution I goes over to the hyperbolic solution

$$u(x, t) = e^{i\omega_1 t} \frac{A \tanh(\beta x)}{B + \operatorname{sech}(\beta x)}, \quad v(x, t) = e^{i\omega_2 t} \frac{D \operatorname{sech}(\beta x)}{B + \operatorname{sech}(\beta x)}, \quad (10)$$

provided

$$\begin{aligned} \omega_1 = \omega_2 = -\frac{\beta^2}{2}, \quad g_{12}D^2 = 3g_{22}D^2 = \frac{3(B^2 - 1)\beta^2}{2}, \\ B > 0, \quad 3g_{11}A^2 = g_{21}A^2 = \frac{3B^2\beta^2}{2}. \end{aligned} \quad (11)$$

Thus $g_{12}, g_{22} > (<) 0$ provided $B^2 > (<) 1$ while $g_{11}, g_{21} > 0$. On the other hand $\omega_1, \omega_2 < 0$.

It is worth pointing out that the local coupled NLS Eqs. (6) and (7) also admit the hyperbolic solution (10) provided relations as given by Eq. (11) are satisfied except that g_{11} and g_{21} have opposite values compared to those given in (11).

Solution III

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A \operatorname{sn}(\beta x, m)}{B + \operatorname{cn}(\beta x, m)}, \quad v(x, t) = e^{i\omega_2 t} \frac{D \operatorname{dn}(\beta x, m)}{B + \operatorname{cn}(\beta x, m)}, \quad (12)$$

is an exact solution of the coupled Eqs. (1) and (2) provided

$$\begin{aligned} \omega_1 = \omega_2 = -\frac{(2m - 1)\beta^2}{2}, \quad g_{12}D^2 = 3g_{22}D^2 = \frac{3(B^2 - 1)\beta^2}{2}, \\ B > 1, \quad 3g_{11}A^2 = g_{21}A^2 = \frac{3(mB^2 + 1 - m)\beta^2}{2}. \end{aligned} \quad (13)$$

Thus $g_{12}, g_{22}, g_{11}, g_{21}$ are all > 0 . On the other hand $\omega_1, \omega_2 < 0 (>) 0$ depending on if $m > (<) 1/2$.

It is worth pointing out that the local coupled NLS Eqs. (6) and (7) also admit the solution (12) provided relations as given by Eq. (13) are satisfied except that g_{11} and g_{21} have opposite values compared to those given in (9).

Solution IV

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A}{B + \cos(\beta x)}, \quad v(x, t) = e^{i\omega_2 t} \frac{D \sin(\beta x)}{B + \cos(\beta x)}, \quad (14)$$

with $B > 1$, is an exact solution of the coupled Eqs. (1) and (2) provided

$$\begin{aligned}\omega_1 = \omega_2 &= \frac{\beta^2}{2}, \quad 3g_{11}A^2 = g_{21}A^2 = 3(B^2 - 1)\frac{\beta^2}{2}, \\ g_{12}D^2 &= 3g_{22}D^2 = \frac{3\beta^2}{2}.\end{aligned}\quad (15)$$

It is worth pointing out that the local coupled NLS Eqs. (6) and (7) also admit the solution (21) provided relations as given by Eq. (15) are satisfied except that g_{12} and g_{22} have opposite values compared to those given in (15).

Solution V

It is straightforward to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A \cos(\beta x)}{1 + B \cos^2(\beta x)}, \quad v(x, t) = e^{i\omega_2 t} \frac{D \sin(\beta x)}{1 + B \cos^2(\beta x)}, \quad B > 0, \quad (16)$$

is an exact solution of the nonlocal coupled NLS Eqs. (1) and (2) provided

$$\omega_1 = \omega_2 = -\beta^2, \quad g_{12}D^2 = 3g_{22}D^2 = -6B\beta^2, \quad g_{21}A^2 = 3g_{11}A^2 = -6B(B+1)\beta^2. \quad (17)$$

Note that the condition $B > 0$ is also valid for the solutions VI to XIII presented below and we will not mention it again.

It is worth noting that the solution (16) is also a solution of the coupled local NLS Eqs. (6) and (7) except that the signs of g_{12} and g_{22} are opposite to those given in Eq. (17).

Solution VI

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A \cos(\beta x)}{1 + B \cos^2(\beta x)}, \quad v(x, t) = e^{i\omega_2 t} \frac{D}{1 + B \cos^2(\beta x)}, \quad (18)$$

is an exact solution of the nonlocal coupled Eqs. (1) and (2) provided

$$\begin{aligned}\omega_1 = -\beta^2, \quad \omega_2 &= -4\beta^2, \quad g_{12}D^2 = -6B\beta^2, \quad g_{11}A^2 = -2B(B+4)\beta^2, \\ g_{22}D^2 &= -2(2-B)\beta^2, \quad g_{21}A^2 = -6B(B+2)\beta^2.\end{aligned}\quad (19)$$

It is worth noting that the solution (18) is also a solution of the coupled local NLS Eqs. (6) and (7).

Solution VII

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A}{1 + B \cos^2(\beta x)}, \quad v(x, t) = e^{i\omega_2 t} \frac{D \sin(\beta x)}{1 + B \cos^2(\beta x)}, \quad (20)$$

is an exact solution of the nonlocal coupled Eqs. (1) and (2) provided

$$\begin{aligned}\omega_1 &= -4\beta^2, \quad g_{12}D^2 = -6B(B+2)\beta^2, \quad g_{11}A^2 = -2(B+1)(3B+2)\beta^2, \\ \omega_2 &= -\beta^2, \quad g_{22}D^2 = -2B(3B+4)\beta^2, \quad g_{21}A^2 = -6B(B+1)\beta^2.\end{aligned}\quad (21)$$

It is worth noting that the solution (20) is also a solution of the coupled local NLS Eqs. (6) and (7) except that the signs of g_{12} and g_{22} are opposite to those given in Eq. (21).

Solution VIII

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A}{1 + B \cos^2(\beta x)}, \quad v(x, t) = e^{i\omega_2 t} \frac{D \sin(\beta x) \cos(\beta x)}{1 + B \cos^2(\beta x)}, \quad (22)$$

is an exact solution of the nonlocal coupled Eqs. (1) and (2) provided

$$\begin{aligned}\omega_1 &= \omega_2 = 2\beta^2, \quad g_{12}D^2 = 3g_{22}D^2 = 6B^2\beta^2, \\ g_{21}A^2 &= 3g_{11}A^2 = 6(B+1)\beta^2.\end{aligned}\quad (23)$$

It is worth noting that the solution (22) is also a solution of the coupled local NLS Eqs. (6) and (7) except that the signs of g_{12} and g_{22} are opposite to those given in Eq. (23).

Solution IX

It is not difficult to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A \cos(\beta x)}{1 + B \cos^2(\beta x)}, \quad v(x, t) = e^{i\omega_2 t} \frac{D \sin(\beta x) \cos(\beta x)}{1 + B \cos^2(\beta x)}, \quad (24)$$

is an exact solution of the nonlocal coupled Eqs. (1) and (2) provided

$$\begin{aligned}\omega_1 &= -6(B+1)\beta^2, \quad g_{12}D^2 = -6B^3\beta^2, \\ g_{11}A^2 &= -2B(B+1)(3B+4)\beta^2, \quad \omega_2 = -6(B+4)\beta^2, \\ g_{21}A^2 &= -6B(B+1)(B+2)\beta^2, \quad g_{22}D^2 = -2(3B+2)\beta^2.\end{aligned}\quad (25)$$

It is worth noting that the solution (24) is also a solution of the coupled local NLS Eqs. (6) and (7) except the signs of g_{12} and g_{22} are opposite to those given in Eq. (25).

Solution X

It is easy to check that

$$u(x, t) = \frac{A \sin(\beta x)}{1 + B \cos^2(\beta x)}, \quad v(x, t) = \frac{D \sin(\beta x) \cos(\beta x)}{1 + B \cos^2(\beta x)}, \quad (26)$$

is an exact solution of coupled Eqs. (1) and (2) provided

$$\begin{aligned}(1+B)a_1 &= (5B-1)\beta^2, & (1+B)d_1D^2 &= 6B^3\beta^2, \\ (1+B)b_1A^2 &= -2B(B+4)\beta^2, & (1+B)a_2 &= 2(B-2)\beta^2, \\ (1+B)d_2D^2 &= 2B^2(B-2)\beta^2, & (1+B)b_2A^2 &= -6B(B+2)\beta^2.\end{aligned}\quad (27)$$

It is worth noting that the solution (26) is also a solution of the coupled local NLS Eqs. (6) and (7) except the signs of g_{11}, g_{12}, g_{21} and g_{22} are all opposite to those given in Eq. (27).

Solution XI

We now present three soliton solutions with a power law tail. It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A(F+x^2)}{B+x^2}, \quad v(x, t) = e^{i\omega_2 t} \frac{Dx}{B+x^2}, \quad (28)$$

is an exact solution of the coupled Eqs. (1) and (2) provided

$$\begin{aligned}\omega_1 = \omega_2 < 0, & \quad 3g_{11} = 4g_{21} < 0, \quad 5g_{12} = 6g_{22} > 0, \quad \omega_1 = -\frac{9}{B}, \\ A^2 = \frac{|\omega_1|}{|g_{11}|}, & \quad 6d_2 = 5d_1, \quad D^2 = \frac{24}{g_{12}}, \quad F = -B/3.\end{aligned}\quad (29)$$

It is worth pointing out that the local coupled NLS Eqs. (6) and (7) also admit the solution (28) provided the relations as given by Eq. (29) are satisfied except that g_{12} and g_{22} have opposite values compared to those given in (29).

Solution XII

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A(F+x^2)}{B+x^2}, \quad v(x, t) = e^{i\omega_2 t} \frac{Dx}{\sqrt{B+x^2}}, \quad (30)$$

is an exact solution of the coupled Eqs. (1) and (2) provided

$$\begin{aligned}\omega_1 = 8\omega_2 < 0, & \quad 3g_{11} = 8g_{21} > 0, \quad 7g_{12} = 32g_{22} > 0, \quad \omega_1 = -\frac{15}{2B}, \\ 5A^2 = \frac{3|\omega_1|}{|g_{11}|}, & \quad D^2 = \frac{12}{Bg_{12}}, \quad F = -B/3.\end{aligned}\quad (31)$$

It is worth pointing out that the local coupled NLS Eqs. (6) and (7) also admit the solution (30) provided the relations as given by Eq. (31) are

satisfied except that g_{12} and g_{22} have opposite values compared to those given in (31).

Solution XIII

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{Ax}{B + x^2}, \quad v(x, t) = e^{i\omega_2 t} \frac{Dx}{\sqrt{B + x^2}}, \quad (32)$$

is an exact solution of the coupled Eqs. (1) and (2) provided

$$\begin{aligned} \omega_1 = 2\omega_2 < 0, \quad 3g_{11} = 8g_{21} > 0, \quad 7g_{12} = 32g_{22} > 0, \quad \omega_1 = -\frac{6}{B}, \\ g_{11}A^2 = 8, \quad D^2 = \frac{6}{Bg_{12}}. \end{aligned} \quad (33)$$

It is worth pointing out that the local coupled NLS Eqs. (6) and (7) also admit the solution (32) provided the relations as given by Eq. (33) are satisfied except that g_{11}, g_{12}, g_{21} and g_{22} have opposite values compared to those given in (33).

2.1 Nonreciprocal Solutions of The Model

We now show that the coupled nonlocal NLS Eqs. (1) and (2) admit rather novel *nonreciprocal* solutions. Without any loss of generality, if we always choose $u(x, t)$ to be a Lamé (inverse Lamé) polynomial of order two and $v(x, t)$ to be a Lamé (inverse Lamé) polynomial of order one, then nonreciprocal solutions exist provided $g_{11} = g_{21} = 0$. Thus in this case one is effectively solving simpler coupled equations

$$iu_t(x, t) + u_{xx}(x, t) + g_{12}v(x, t)v^*(-x, t)u(x, t) = 0, \quad (34)$$

$$iv_t(x, t) + v_{xx}(x, t) + g_{22}v^2(x, t)v^*(-x, t) = 0. \quad (35)$$

Notice that while $v(x, t)$ satisfies an uncoupled Eq. (35), on the other hand $u(x, t)$ satisfies the coupled Eq. (34). It is worth remembering that if instead we consider the full coupled Eqs. (1) and (2), then $u(x, t)$ being Lamé (inverse Lamé) polynomial of order two and $v(x, t)$ being Lamé (inverse Lamé) polynomial of order one is *not* a solution of these coupled equations. On the other hand, if both $u(x, t)$ and $v(x, t)$ are Lamé (inverse Lamé) polynomials of order one and are solutions of the full coupled Eqs. (1) and (2), then they are also the solutions of the truncated coupled Eqs. (34) and (35). Hence we will only consider those reciprocal solutions where u is in terms of Lamé (inverse Lamé) polynomial of order two while v is in

terms of Lamé (inverse Lamé) polynomial of order one. We first present twelve nonreciprocal periodic solutions and the corresponding six hyperbolic nonreciprocal solutions in terms of Lamé polynomials of order two in u and order one in v . Later we will present 12 periodic solutions in terms of inverse Lamé polynomials of order two in u and inverse Lamé polynomials of order one in v .

Solution XIV

It is not difficult to check that

$$u(x, t) = Ae^{i\omega_1 t} \sqrt{m} \operatorname{sn}(\beta x, m) \operatorname{dn}(\beta x, m), \quad v(x, t) = Be^{i\omega_2 t} \sqrt{m} \operatorname{sn}(\beta x, m), \quad (36)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$g_{12}B^2 = \beta^2, \quad 3g_{22}B^2 = \beta^2, \quad (37)$$

$$\omega_1 = -(4m + 1)\beta^2, \quad \omega_2 = -(1 + m)\beta^2. \quad (38)$$

What is remarkable about this solution is that it is independent of the value of A . It is also independent of whether the u field is odd or even under parity-time reversal (PT) symmetry. In fact this feature is common to all the reciprocal solutions presented below and hence will not be repeated.

It is worth pointing out that the solution (36) is also a solution of the local NLS Eqs.

$$iu_t(x, t) + u_{xx}(x, t) + 6g_{12}|v(x, t)|^2 u(x, t) = 0, \quad (39)$$

$$iv_t(x, t) + v_{xx}(x, t) + 6g_{22}|v(x, t)|^2 v(x, t) = 0, \quad (40)$$

except that the signs of g_{12} and g_{22} are opposite compared to those given in Eq. (37).

Solution XV

It is easy to check that

$$u(x, t) = Ae^{i\omega_1 t} m \operatorname{sn}(\beta x, m) \operatorname{cn}(\beta x, m), \quad v(x, t) = Be^{i\omega_2 t} \sqrt{m} \operatorname{sn}(\beta x, m), \quad (41)$$

is an exact solution of coupled Eqs. (34) and (35) provided relations (37) are satisfied and further

$$\omega_1 = -(4 + m)\beta^2, \quad \omega_2 = -(1 + m)\beta^2. \quad (42)$$

It is worth pointing out that the solution (41) is also a solution of the local NLS Eqs. (39) and (40) except that the signs of g_{12} and g_{22} are opposite compared to those given in Eq. (37).

Solution XVI

In the limit $m = 1$, both the solutions XIV and XV go over to the hyperbolic nonreciprocal solution

$$u(x, t) = Ae^{i\omega_1 t} \operatorname{sech}(\beta x) \tanh(\beta x), \quad v(x, t) = Be^{i\omega_2 t} \tanh(\beta x), \quad (43)$$

provided relations (37) are satisfied and further

$$\omega_1 = -5\beta^2, \quad \omega_2 = -2\beta^2. \quad (44)$$

It is worth pointing out that the solution (43) is also a solution of the local NLS Eqs. (39) and (40) except that the signs of g_{12} and g_{22} are opposite compared to those given in Eq. (37).

Solution XVII

It is straightforward to check that

$$u(x, t) = Ae^{i\omega_1 t} \sqrt{m} \operatorname{sn}(\beta x, m) \operatorname{dn}(\beta x, m), \quad v(x, t) = Be^{i\omega_2 t} \sqrt{m} \operatorname{cn}(\beta x, m), \quad (45)$$

is an exact solution of coupled Eqs. (34) and (35) provided relations (37) are satisfied and further

$$\omega_1 = \omega_2 = (2m - 1)\beta^2. \quad (46)$$

It is worth pointing out that the solution (45) is also a solution of the local NLS Eqs. (39) and (40).

Solution XVIII

It is not difficult to check that

$$u(x, t) = Ae^{i\omega_1 t} \sqrt{m} \operatorname{sn}(\beta x, m) \operatorname{dn}(\beta x, m), \quad v(x, t) = Be^{i\omega_2 t} \operatorname{dn}(\beta x, m), \quad (47)$$

is an exact solution of coupled Eqs. (34) and (35) provided the relations (37) are satisfied and further

$$\omega_1 = (5 - 4m)\beta^2, \quad \omega_2 = (2 - m)\beta^2. \quad (48)$$

It is worth pointing out that the solution (47) is also a solution of the local NLS Eqs. (39) and (40).

Solution XIX

It is easy to check that

$$u(x, t) = Ae^{i\omega_1 t} m \operatorname{sn}(\beta x, m) \operatorname{cn}(\beta x, m), \quad v(x, t) = Be^{i\omega_2 t} \sqrt{m} \operatorname{cn}(\beta x, m), \quad (49)$$

is an exact solution of coupled Eqs. (34) and (35) provided the relations (37) are satisfied and further

$$\omega_1 = (5m - 4)\beta^2, \quad \omega_2 = (2m - 1)\beta^2. \quad (50)$$

It is worth pointing out that the solution (49) is also a solution of the local NLS Eqs. (39) and (40).

Solution XX

It is easy to check that

$$u(x, t) = Ae^{i\omega_1 t} m \operatorname{sn}(\beta x, m) \operatorname{cn}(\beta x, m), \quad v(x, t) = Be^{i\omega_2 t} \operatorname{dn}(\beta x, m), \quad (51)$$

is an exact solution of coupled Eqs. (34) and (35) provided the relations (37) are satisfied and further

$$\omega_1 = \omega_2 = (2 - m)\beta^2. \quad (52)$$

It is worth pointing out that the solution (51) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXI

In the limit $m = 1$, all the four solutions XVII to XX go over to the hyperbolic solution

$$u(x, t) = Ae^{i\omega_1 t} \operatorname{sech}(\beta x) \tanh(\beta x), \quad v(x, t) = Be^{i\omega_2 t} \operatorname{sech}(\beta x), \quad (53)$$

provided the relations (37) are satisfied and further

$$\omega_1 = \omega_2 = \beta^2. \quad (54)$$

It is worth pointing out that the solution (53) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXII

It is easy to check that

$$u(x, t) = Ae^{i\omega_1 t} \sqrt{m} \operatorname{cn}(\beta x, m) \operatorname{dn}(\beta x, m), \quad v(x, t) = Be^{i\omega_2 t} \sqrt{m} \operatorname{sn}(\beta x, m), \quad (55)$$

is an exact solution of coupled Eqs. (34) and (35) provided the relations (37) are satisfied and further

$$\omega_1 = \omega_2 = -(1 + m)\beta^2. \quad (56)$$

It is worth pointing out that the solution (55) is also a solution of the local NLS Eqs. (39) and (40) except that the signs of g_{12} and g_{22} are opposite compared to those given in Eq. (37).

Solution XXIII

It is easy to check that

$$u(x, t) = Ae^{i\omega_1 t}[\text{dn}^2(\beta x, m) + y], \quad v(x, t) = Be^{i\omega_2 t}\sqrt{m}\text{sn}(\beta x, m), \quad (57)$$

is an exact solution of coupled Eqs. (34) and (35) provided relations (37) are satisfied and further

$$\omega_1 = 2(3y + 1 - 2m)\beta^2, \quad \omega_2 = -(1 + m)\beta^2, \quad (58)$$

where

$$y = \frac{-(2 - m) \pm \sqrt{1 - m + m^2}}{3}. \quad (59)$$

It is worth pointing out that the solution (57) is also a solution of the local NLS Eqs. (39) and (40) except that the signs of g_{12} and g_{22} are opposite compared to those given in Eq. (37).

Solution XXIV

In the limit $m = 1$, $y = 0$ or $y = -2/3$. In case $y = 0$ and $m = 1$, both the solutions XXII and XXIII go over to the hyperbolic solution

$$u(x, t) = Ae^{i\omega_1 t}\text{sech}^2(\beta x), \quad v(x, t) = B\sqrt{m}\tanh(\beta x), \quad (60)$$

provided relations (37) are satisfied and further

$$\omega_1 = \omega_2 = -2\beta^2. \quad (61)$$

It is worth pointing out that the solution (60) is also a solution of the local NLS Eqs. (39) and (40) except that the signs of g_{12} and g_{22} are opposite compared to those given in Eq. (37).

Solution XXV

On the other hand, in case $y = -2/3$ and $m = 1$, the solution XXIII goes over to the hyperbolic solution

$$u(x, t) = Ae^{i\omega_1 t}[\text{sech}^2(\beta x) - 2/3], \quad v(x, t) = B\sqrt{m}\tanh(\beta x), \quad (62)$$

provided relations (37) are satisfied and further

$$\omega_1 = -6\beta^2, \quad \omega_2 = -2\beta^2. \quad (63)$$

It is worth pointing out that the solution (62) is also a solution of the local NLS Eqs. (39) and (40) except that the signs of g_{12} and g_{22} are opposite compared to those given in Eq. (37).

Solution XXVI

It is easy to check that

$$u(x, t) = Ae^{i\omega_1 t} \sqrt{m} \operatorname{cn}(\beta x, m) \operatorname{dn}(\beta x, m), \quad v(x, t) = Be^{i\omega_2 t} \sqrt{m} \operatorname{cn}(\beta x, m), \quad (64)$$

is an exact solution of coupled Eqs. (34) and (35) provided the relations (37) are satisfied and further

$$\omega_1 = (5m - 1)\beta^2, \quad \omega_2 = (2m - 1)\beta^2. \quad (65)$$

It is worth pointing out that the solution (64) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXVII

It is straightforward to check that

$$u(x, t) = Ae^{i\omega_1 t} \sqrt{m} \operatorname{cn}(\beta x, m) \operatorname{dn}(\beta x, m), \quad v(x, t) = Be^{i\omega_2 t} \operatorname{dn}(\beta x, m), \quad (66)$$

is an exact solution of coupled Eqs. (34) and (35) provided the relations (37) are satisfied and further

$$\omega_1 = (5 - m)\beta^2, \quad \omega_2 = (2 - m)\beta^2. \quad (67)$$

It is worth pointing out that the solution (66) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXVIII

It is easy to check that

$$u(x, t) = Ae^{i\omega_1 t} [\operatorname{dn}^2(\beta x, m) + y], \quad v(x, t) = Be^{i\omega_2 t} \sqrt{m} \operatorname{cn}(\beta x, m), \quad (68)$$

is an exact solution of coupled Eqs. (34) and (35) provided the relations (37) are satisfied and further

$$\omega_1 = 2[(1 + m) + 3y]\beta^2, \quad \omega_2 = (2m - 1)\beta^2, \quad (69)$$

where y is again as given by Eq. (59).

It is worth pointing out that the solution (68) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXIX

It is easy to check that

$$u(x, t) = Ae^{i\omega_1 t} [\operatorname{dn}^2(\beta x, m) + y], \quad v(x, t) = Be^{i\omega_2 t} \operatorname{dn}(\beta x, m), \quad (70)$$

is an exact solution of coupled Eqs. (34) and (35) provided the relations (37) are satisfied and further

$$\omega_1 = 2[2(2 - m) + 3y]\beta^2, \quad \omega_2 = (2 - m)\beta^2, \quad (71)$$

where y is again as given by Eq. (59).

It is worth pointing out that the solution (70) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXX

In the limit $m = 1$, $y = 0$ or $y = -2/3$. In case $y = 0$ and $m = 1$, all the four solutions XXVI to XXIX go over to the hyperbolic solution

$$u(x, t) = Ae^{i\omega_1 t} \operatorname{sech}^2(\beta x), \quad v(x, t) = Be^{i\omega_2 t} \operatorname{sech}(\beta x), \quad (72)$$

provided relations (37) are satisfied and further

$$\omega_1 = 4\beta^2, \quad \omega_2 = \beta^2. \quad (73)$$

It is worth pointing out that the solution (72) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXXI

On the other hand, in case $y = -2/3$ and $m = 1$, the solutions XXVIII and XXIX go over to the solution

$$u(x, t) = Ae^{i\omega_1 t} [\operatorname{sech}^2(\beta x) - 2/3], \quad v(x, t) = Be^{i\omega_2 t} \operatorname{sech}(\beta x), \quad (74)$$

provided relations (37) are satisfied and further

$$\omega_1 = 0, \quad \omega_2 = \beta^2. \quad (75)$$

It is worth pointing out that the solution (74) is also a solution of the local NLS Eqs. (39) and (40).

I now present two novel superposed solutions.

Solution XXXII

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} [A \operatorname{dn}^2(\beta x, m) + Ay + B\sqrt{m} \operatorname{cn}(\beta x, m) \operatorname{dn}(\beta x, m)], \quad (76)$$

$$v(x, t) = De^{i\omega_2 t} [\operatorname{dn}(\beta x, m) + E\sqrt{m} \operatorname{cn}(\beta x, m)], \quad (77)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$\begin{aligned} 4g_{12}D^2 &= \beta^2, \quad \omega_1 = \left[\frac{(7+m)}{2} + 3y \right] \beta^2, \quad E = \pm D, \quad B = \pm A, \\ \omega_2 &= \frac{(1+m)\beta^2}{2}, \quad 12g_{22}D^2 = \beta^2, \\ y &= \frac{-(5-m) \pm \sqrt{1+14m+m^2}}{6}. \end{aligned} \quad (78)$$

Note that the \pm signs in D, E are correlated with the signs in B and A . Further, the solution is independent of the values of B and A except that they are constrained by $B = A$. In the limit $m = 1$, this solution goes over to the solution XXX.

It is worth pointing out that the solution (76) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXXIII

It is easy to check that

$$u(x, t) = Ae^{i\omega_1 t}[\sqrt{m}\operatorname{dn}(\beta x, m)\operatorname{sn}(\beta x, m) + Bm\operatorname{cn}(\beta x, m)\operatorname{sn}(\beta x, m)], \quad (79)$$

$$v(x, t) = e^{i\omega_2 t}[D\operatorname{dn}(\beta x, m) + E\sqrt{m}\operatorname{cn}(\beta x, m)], \quad (80)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$\begin{aligned} 4g_{12}D^2 &= \beta^2, \quad \omega_1 = \omega_2 = \frac{(1+m)\beta^2}{2}, \quad E = \pm D, \quad B = \pm A, \\ 12g_{22}D^2 &= \beta^2. \end{aligned} \quad (81)$$

Note that the \pm signs in D, E are correlated with the signs in B and A . Further, the solution is independent of the values of A and B except that they must satisfy the constraint $B = \pm A$. In the limit $m = 1$, this solution goes over to the hyperbolic solution XXI.

It is worth pointing out that the solution (79) is also a solution of the local NLS Eqs. (39) and (40) except that the signs of g_{12} and g_{22} are opposite to those given in Eq. (81).

Remarkably, these coupled equations also admit several solutions in terms of inverse Lamé polynomial solutions of order two in $u(x, t)$ and of inverse Lamé polynomial solutions of order one in $v(x, t)$. We list these solutions one by one. It may be noted that all these solutions are only valid for $0 < m < 1$.

Solution XXXIV

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A\sqrt{m}\operatorname{cn}(\beta x, m)}{\operatorname{dn}^2(\beta x, m)}, \quad v(x, t) = e^{i\omega_2 t} \frac{B\sqrt{m}\operatorname{cn}(\beta x, m)}{\operatorname{dn}(\beta x, m)}, \quad (82)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$g_{12}B^2 = -6\beta^2, \quad g_{22}B^2 = -2\beta^2, \quad (83)$$

$$\omega_1 = -(4m + 1)\beta^2, \quad \omega_2 = -(1 + m)\beta^2. \quad (84)$$

It is worth pointing out that the solution (82) is also a solution of the local NLS Eqs. (39) and (40). Note that this solution is independent of the magnitude of A . In fact this feature is common to all the solutions given below.

Solution XXXV

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A\sqrt{m}\text{cn}(\beta x, m)}{\text{dn}^2(\beta x, m)}, \quad v(x, t) = e^{i\omega_2 t} \frac{B\sqrt{m}\text{sn}(\beta x, m)}{\text{dn}(\beta x, m)}, \quad (85)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$g_{12}B^2 = -6(1 - m)\beta^2, \quad g_{22}B^2 = -2(1 - m)\beta^2, \quad (86)$$

$$\omega_1 = \omega_2 = (2m - 1)\beta^2. \quad (87)$$

It is worth pointing out that the solution (85) is also a solution of the local NLS Eqs. (39) and (40) except that the signs of g_{12} and g_{22} are opposite to those given in Eq. (86).

Solution XXXVI

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A\sqrt{m}\text{cn}(\beta x, m)}{\text{dn}^2(\beta x, m)}, \quad v(x, t) = e^{i\omega_2 t} \frac{B}{\text{dn}(\beta x, m)}, \quad (88)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$g_{12}B^2 = 6(1 - m)\beta^2, \quad g_{22}B^2 = 2(1 - m)\beta^2, \quad (89)$$

$$\omega_1 = (5 - 4m)\beta^2, \quad \omega_2 = (2 - m)\beta^2. \quad (90)$$

It is worth pointing out that the solution (88) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXXVII

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A\sqrt{m}\text{sn}(\beta x, m)}{\text{dn}^2(\beta x, m)}, \quad v(x, t) = e^{i\omega_2 t} \frac{B\text{cn}(\beta x, m)}{\text{dn}(\beta x, m)}, \quad (91)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$g_{12}B^2 = -6\beta^2, \quad g_{22}B^2 = -2\beta^2, \quad (92)$$

$$\omega_1 = \omega_2 = -(1 + m)\beta^2. \quad (93)$$

It is worth pointing out that the solution (91) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXXVIII

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A\sqrt{m}\operatorname{sn}(\beta x, m)}{\operatorname{dn}^2(\beta x, m)}, \quad v(x, t) = e^{i\omega_2 t} \frac{B\operatorname{sn}(\beta x, m)}{\operatorname{dn}(\beta x, m)}, \quad (94)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$g_{12}B^2 = -6(1-m)\beta^2, \quad g_{22}B^2 = -2(1-m)\beta^2, \quad (95)$$

$$\omega_1 = (5m-1)\beta^2, \quad \omega_2 = (2m-1)\beta^2. \quad (96)$$

It is worth pointing out that the solution (94) is also a solution of the local NLS Eqs. (39) and (40) except that the signs of g_{12} and g_{22} are opposite to those given in Eq. (95).

Solution XXXIX

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A\sqrt{m}\operatorname{sn}(\beta x, m)}{\operatorname{dn}^2(\beta x, m)}, \quad v(x, t) = e^{i\omega_2 t} \frac{B}{\operatorname{dn}(\beta x, m)}, \quad (97)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$g_{12}B^2 = 6(1-m)\beta^2, \quad g_{22}B^2 = 2(1-m)\beta^2, \quad (98)$$

$$\omega_1 = -(5-m)\beta^2, \quad \omega_2 = (2-m)\beta^2. \quad (99)$$

It is worth pointing out that the solution (97) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXXX

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A\sqrt{m}\operatorname{sn}(\beta x, m)\operatorname{cn}(\beta x, m)}{\operatorname{dn}^2(\beta x, m)}, \quad (100)$$

$$v(x, t) = e^{i\omega_2 t} \frac{B\sqrt{m}\operatorname{cn}(\beta x, m)}{\operatorname{dn}(\beta x, m)}, \quad (101)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$g_{12}B^2 = -6\beta^2, \quad g_{22}B^2 = -2\beta^2, \quad (102)$$

$$\omega_1 = -(4+m)\beta^2, \quad \omega_2 = -(1+m)\beta^2. \quad (103)$$

It is worth pointing out that the solution (100) and (101) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXXXI

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A\sqrt{m}\operatorname{sn}(\beta x, m)\operatorname{cn}(\beta x, m)}{\operatorname{dn}^2(\beta x, m)}, \quad (104)$$

$$v(x, t) = e^{i\omega_2 t} \frac{B\sqrt{m}\operatorname{sn}(\beta x, m)}{\operatorname{dn}(\beta x, m)}, \quad (105)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$g_{12}B^2 = -6(1-m)\beta^2, \quad g_{22}B^2 = -2(1-m)\beta^2, \quad (106)$$

$$\omega_1 = -(5m-4)\beta^2, \quad \omega_2 = (2m-1)\beta^2. \quad (107)$$

It is worth pointing out that the solution (104) and (105) is also a solution of the local NLS Eqs. (39) and (40) except that the signs of g_{12} and g_{22} are opposite to those given in Eq. (106).

Solution XXXXII

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A\sqrt{m}\operatorname{sn}(\beta x, m)\operatorname{cn}(\beta x, m)}{\operatorname{dn}^2(\beta x, m)}, \quad (108)$$

$$v(x, t) = e^{i\omega_2 t} \frac{B}{\operatorname{dn}(\beta x, m)}, \quad (109)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$g_{12}B^2 = 6(1-m)\beta^2, \quad g_{22}B^2 = 2(1-m)\beta^2, \quad (110)$$

$$\omega_1 = \omega_2 = (2-m)\beta^2. \quad (111)$$

It is worth pointing out that the solution (108) and (109) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXXXIII

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} A \left[\frac{1}{\operatorname{dn}^2(\beta x, m)} + y \right], \quad (112)$$

$$v(x, t) = e^{i\omega_2 t} \frac{B\sqrt{m}\operatorname{cn}(\beta x, m)}{\operatorname{dn}(\beta x, m)}, \quad (113)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$g_{12}B^2 = -6\beta^2, \quad g_{22}B^2 = -2\beta^2, \quad (114)$$

$$\omega_1 = -2\left(3 + \frac{1}{y}\right)\beta^2, \quad \omega_2 = (2 - m)\beta^2, \quad (115)$$

where

$$y = \frac{-(2 - m) \pm \sqrt{1 - m + m^2}}{3(1 - m)}. \quad (116)$$

It is worth pointing out that the solution (112) and (113) is also a solution of the local NLS Eqs. (39) and (40).

Solution XXXXIV

It is not difficult to check that

$$u(x, t) = e^{i\omega_1 t} A \left[\frac{1}{\operatorname{dn}^2(\beta x, m)} + y \right], \quad (117)$$

$$v(x, t) = e^{i\omega_2 t} \frac{B\sqrt{m}\operatorname{sn}(\beta x, m)}{\operatorname{dn}(\beta x, m)}, \quad (118)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$g_{12}B^2 = -6(1 - m)\beta^2, \quad g_{22}B^2 = -2(1 - m)\beta^2, \quad (119)$$

$$\omega_1 = -2\left[3(1 - m) + \frac{1}{y}\right]\beta^2, \quad \omega_2 = (2m - 1)\beta^2, \quad (120)$$

while y is again as given by Eq. (116).

It is worth pointing out that the solution (117) and (118) is also a solution of the local NLS Eqs. (39) and (40) except that the signs of g_{12} and g_{22} are opposite to those given in Eq. (119).

Solution XXXXV

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} A \left[\frac{1}{\operatorname{dn}^2(\beta x, m)} + y \right], \quad (121)$$

$$v(x, t) = e^{i\omega_2 t} \frac{B}{\operatorname{dn}(\beta x, m)}, \quad (122)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$g_{12}B^2 = 6(1 - m)\beta^2, \quad g_{22}B^2 = (2 - m)\beta^2, \quad (123)$$

$$\omega_1 = -\frac{2}{y}\beta^2, \quad \omega_2 = (2-m)\beta^2, \quad (124)$$

while y is again as given by Eq. (116).

It is worth pointing out that the solution (121) and (122) is also a solution of the local NLS Eqs. (39) and (40).

Finally, we present a novel solution where while ϕ_1 is a combination of Lamé polynomials of order two and inverse Lamé polynomials of order two, ϕ_2 is a combination of Lamé polynomials of order one and inverse Lamé polynomials of order one.

Solution XXXXVI

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \left(A[\text{dn}^2(\beta x, m) + \sqrt{1-m}y] + \frac{B(1-m)}{\text{dn}^2(\beta x, m)} \right), \quad (125)$$

$$v(x, t) = e^{i\omega_2 t} \left[D\text{dn}(\beta x, m) + \frac{E\sqrt{1-m}}{\text{dn}(\beta x, m)} \right], \quad (126)$$

is an exact solution of coupled Eqs. (34) and (35) provided

$$\begin{aligned} g_{12}D^2 &= 6\beta^2, \quad \omega_2 = (2-m \pm 6\sqrt{1-m})\beta^2, \quad E = \pm D, \quad B = \pm A, \\ g_{22}D^2 &= 2\beta^2, \quad \omega_1 = 2[2(2-m) + 3y \pm 6\sqrt{1-m}]\beta^2, \\ y &= \frac{-(2-m) \pm 6\sqrt{1-m}}{3}, \quad 0 < m < 1. \end{aligned} \quad (127)$$

Note that the \pm signs in the various relations are correlated. Further, the solution is independent of the values of A and B except that they must satisfy the constraint $B = \pm A$.

It is worth pointing out that the solution (125) and (126) is also a solution of the local NLS Eqs. (39) and (40).

3 New Solutions of a Coupled Nonlocal mKdV Model

Let us consider the following nonlocal coupled mKdV model [12]

$$u_t(x, t) + u_{xxx}(x, t) + 6[g_{11}u(x, t)u(-x, -t) + g_{12}v(x, t)v(-x, -t)]u_x(x, t) = 0, \quad (128)$$

$$v_t(x, t) + v_{xxx}(x, t) + 6[g_{21}u(x, t)u(-x, -t) + g_{22}v(x, t)v(-x, -t)]v_x(x, t) = 0. \quad (129)$$

First of all, notice that unlike the coupled local mKdV, the nonlocal coupled mKdV Eqs. (128) and (129) admit a plane wave solution

$$u(x, t) = Ae^{i(k_1x - \omega_1t)}, \quad v(x, t) = Be^{i(k_2x - \omega_2t)}, \quad (130)$$

provided

$$\omega_1 + k_1^3 = 6k_1[g_{11}A^2 + g_{12}B^2], \quad \omega_2 + k_2^3 = 6k_2[g_{21}A^2 + g_{22}B^2]. \quad (131)$$

It is obvious from here that even the uncoupled nonlocal mKdV equation

$$u_t(x, t) + u_{xxx}(x, t) + 6gu(x, t)u(-x, -t)u_x(x, t) = 0, \quad (132)$$

admits the plane wave solution

$$u(x, t) = Ae^{i(kx - \omega t)}, \quad (133)$$

provided

$$\omega_1 + k_1^3 = 6k_1A^2. \quad (134)$$

We now show that unlike the coupled local mKdV equations

$$u_t(x, t) + u_{xxx}(x, t) + 6[g_{11}u^2(x, t) + g_{12}v^2(x, t)]u_x(x, t) = 0, \quad (135)$$

$$v_t(x, t) + v_{xxx}(x, t) + 6[g_{21}u^2(x, t) + g_{22}v^2(x, t)]v_x(x, t) = 0, \quad (136)$$

the nonlocal mKdV Eqs. (128) and (129) admit solutions similar to those obtained by Zakharov and Shabat [16] for the local uncoupled NLS case.

Solution I

It is straightforward to check that

$$\begin{aligned} u(x, t) &= \sqrt{n_1}[B \tanh(\xi) + iA]e^{i(k_1x - \omega_1t)}, \\ v(x, t) &= \sqrt{n_2}[D \tanh(\xi) + iE]e^{i(k_2x - \omega_2t)}, \quad \xi = \beta(x - ct), \end{aligned} \quad (137)$$

is an exact solution of the nonlocal coupled Eqs. (128) and (129) provided

$$\beta^2 = g_{11}n_1B^2 + g_{12}n_2D^2 = g_{21}n_1B^2 + g_{22}n_2D^2, \quad (138)$$

$$\omega_1 + k_1^3 = -6\beta^2k_1 - 6k_1[g_{11}n_1A^2 + g_{12}n_2E^2], \quad (139)$$

$$\omega_2 + k_2^3 = -6\beta^2k_2 - 6k_2[g_{21}n_1A^2 + g_{22}n_2E^2], \quad (140)$$

$$Bck_1 = -2Bk_1^3 + 4k_1B\beta^2 - 6k_1^2A\beta + B\omega_1, \quad (141)$$

$$Dck_2 = -2Dk_2^3 + 4k_2D\beta^2 - 6k_2^2E\beta + D\omega_2. \quad (142)$$

Solution II

It is easy to check that

$$\begin{aligned} u(x, t) &= \sqrt{n_1}[B \tanh(\xi) + iA]e^{i(k_1x - \omega_1t)}, \\ v(x, t) &= \sqrt{n_2}[E + iD \tanh(\xi)]e^{i(k_2x - \omega_2t)}, \end{aligned} \quad (143)$$

is an exact solution of the nonlocal coupled Eqs. (128) and (129) provided

$$\beta^2 = g_{11}n_1B^2 - g_{12}n_2D^2 = g_{21}n_1B^2 - g_{22}n_2D^2, \quad (144)$$

$$\omega_1 + k_1^3 = -6\beta^2k_1 - 6k_1[g_{11}n_1A^2 - g_{12}n_2E^2], \quad (145)$$

$$\omega_2 + k_2^3 = -6\beta^2k_2 - 6k_2[g_{21}n_1A^2 - g_{22}n_2E^2], \quad (146)$$

$$Bck_1 = -2Bk_1^3 + 4k_1B\beta^2 - 6k_1^2A\beta + B\omega_1, \quad (147)$$

$$Dck_2 = -2Dk_2^3 + 4k_2D\beta^2 + 6k_2^2E\beta + D\omega_2. \quad (148)$$

Solution III

It is not difficult to check that

$$\begin{aligned} u(x, t) &= \sqrt{n_1}[A + iB \tanh(\xi)]e^{i(k_1x - \omega_1t)}, \\ v(x, t) &= \sqrt{n_2}[E + iD \tanh(\xi)]e^{i(k_2x - \omega_2t)}, \end{aligned} \quad (149)$$

is an exact solution of the nonlocal coupled Eqs. (128) and (129) provided

$$\beta^2 = -[g_{11}n_1B^2 + g_{12}n_2D^2] = -[g_{21}n_1B^2 + g_{22}n_2D^2], \quad (150)$$

$$\omega_1 + k_1^3 = -6\beta^2k_1 + 6k_1[g_{11}n_1A^2 + g_{12}n_2E^2], \quad (151)$$

$$\omega_2 + k_2^3 = -6\beta^2k_2 + 6k_2[g_{21}n_1A^2 + g_{22}n_2E^2], \quad (152)$$

$$Bck_1 = -2Bk_1^3 + 4k_1B\beta^2 + 6k_1^2A\beta + B\omega_1, \quad (153)$$

$$Dck_2 = -2Dk_2^3 + 4k_2D\beta^2 + 6k_2^2E\beta + D\omega_2. \quad (154)$$

Note that the coupled local mKdV Eqs. (135) and (136) do not admit the solutions I, II and III.

We now show that corresponding to any solution of the coupled nonlocal Eqs. (128) and (129) of the form

$$u(x, t) = u(\xi), \quad v(x, t) = v(\xi), \quad \xi = \beta(x - ct), \quad (155)$$

where $u(\xi)$ and $v(\xi)$ are real, there are always solutions of the form

$$u(x, t) = u(\xi)e^{i(k_1x - \omega_1t)}, \quad v(x, t) = v(\xi)e^{i(k_2x - \omega_2t)}, \quad \xi = \beta(x - vt). \quad (156)$$

The proof is straightforward. On substituting the ansatz as given by Eq. (156) in the coupled Eqs. (128) and (129) and equating the real and imaginary parts we obtain the following four relations

$$(\omega_1 + k_1^3)u(\xi) = 3\beta^2 k_1 u''(\xi) + 6k_1 [g_{11}u(\xi)u(-\xi) + g_{12}v(\xi)v(-\xi)]u(\xi), \quad (157)$$

$$(c + 3k_1^2)u'(\xi) = \beta^2 u'''(\xi) + 6[g_{11}u(\xi)u(-\xi) + g_{12}v(\xi)v(-\xi)]u'(\xi), \quad (158)$$

$$(\omega_2 + k_2^3)v(\xi) = 3\beta^2 k_2 v''(\xi) + 6k_2 [g_{21}u(\xi)u(-\xi) + g_{12}v(\xi)v(-\xi)]v(\xi), \quad (159)$$

$$(c + 3k_2^2)v'(\xi) = \beta^2 v'''(\xi) + 6[g_{21}u(\xi)u(-\xi) + g_{22}v(\xi)v(-\xi)]v'(\xi). \quad (160)$$

In the limit $k_1 = k_2 = \omega_1 = \omega_2 = 0$, the two Eqs. (157) as well as (159) disappear while the other two equations lead to the two coupled equations for $u(\xi)$ and $v(\xi)$

$$cu'(\xi) = \beta^2 u'''(\xi) + 6[g_{11}u(\xi)u(-\xi) + g_{12}v(\xi)v(-\xi)]u'(\xi), \quad (161)$$

$$cv'(\xi) = \beta^2 v'''(\xi) + 6[g_{11}u(\xi)u(-\xi) + g_{12}v(\xi)v(-\xi)]v'(\xi). \quad (162)$$

We reemphasize that the corresponding local mKdV coupled Eqs. (135) and (136) do not admit either the plane wave type solutions of the form (130) nor the solutions of the form (156). As an illustration, we now present six well known solutions of the nonlocal coupled mKdV Eqs. (128) and (129) without the plane wave factor but reconsider them multiplied by the extra plane wave factor.

Solution IV

It is not difficult to check that

$$u(x, t) = A\sqrt{m}\operatorname{sn}(\xi, m)e^{i(k_1x - \omega_1t)}, \quad v(x, t) = B\sqrt{m}\operatorname{cn}(\xi, m)e^{i(k_2x - \omega_2t)}, \quad (163)$$

is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} \beta^2 &= g_{11}A^2 + g_{12}B^2 = g_{21}A^2 + g_{22}B^2, \\ c + 3k_1^2 &= (5m - 1)\beta^2 - 6mg_{11}A^2, \quad \omega_2 + k_2^3 = 3(2m - 1)k_2\beta^2 - 6mk_2g_{21}A^2, \\ c + 3k_2^2 &= (2m - 1)\beta^2 - 6mg_{21}A^2, \quad \omega_1 + k_1^3 = -3(1 - m)k_1\beta^2 - 6mk_1g_{11}A^2. \end{aligned} \quad (164)$$

Solution V

It is easy to check that

$$u(x, t) = A\sqrt{m}\operatorname{sn}(\xi, m)e^{i(k_1x - \omega_1t)}, \quad v(x, t) = B\operatorname{dn}(\xi, m)e^{i(k_2x - \omega_2t)}, \quad (165)$$

is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned}
\beta^2 &= g_{11}A^2 + g_{12}B^2 = g_{21}A^2 + g_{22}B^2, \\
c + 3k_1^2 &= (5 - m)\beta^2 - 6g_{11}A^2, \quad \omega_2 + k_2^3 = 3(2 - m)k_2\beta^2 - 6k_2g_{21}A^2, \\
c + 3k_2^2 &= (2 - m)\beta^2 - 6g_{21}A^2, \quad \omega_1 + k_1^3 = 3(1 - m)k_1\beta^2 - 6k_1g_{11}A^2.
\end{aligned} \tag{166}$$

Solution VI

In the limit $m = 1$, both the solutions I and II go over to the hyperbolic solution

$$u(x, t) = A \tanh(\xi)e^{i(k_1x - \omega_1t)}, \quad v(x, t) = B \operatorname{sech}(\xi)e^{i(k_2x - \omega_2t)}, \tag{167}$$

provided

$$\begin{aligned}
\beta^2 &= g_{11}A^2 + g_{12}B^2 = g_{21}A^2 + g_{22}B^2, \quad \omega_1 + k_1^3 = -6k_1g_{11}A^2, \\
c + 3k_1^2 &= 4\beta^2 - 6g_{11}A^2, \quad \omega_2 + k_2^3 = 3k_2\beta^2 - 6k_2g_{21}A^2, \\
c + 3k_2^2 &= \beta^2 - 6g_{21}A^2.
\end{aligned} \tag{168}$$

Solution VII

It is easy to check that

$$u(x, t) = A \operatorname{dn}(\xi, m)e^{i(k_1x - \omega_1t)}, \quad v(x, t) = B\sqrt{m} \operatorname{cn}(\xi, m)e^{i(k_2x - \omega_2t)}, \tag{169}$$

is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned}
\beta^2 &= g_{11}A^2 + g_{12}B^2 = g_{21}A^2 + g_{22}B^2, \\
c + 3k_1^2 &= (2 - m)\beta^2 - 6(1 - m)g_{12}B^2, \\
\omega_2 + k_2^3 &= 3k_2\beta^2 - 6(1 - m)k_2g_{22}B^2, \\
c + 3k_2^2 &= (5 - 4m)\beta^2 - 6(1 - m)g_{22}B^2, \\
\omega_1 + k_1^3 &= 3(2 - m)k_1\beta^2 - 6(1 - m)k_1g_{12}A^2.
\end{aligned} \tag{170}$$

Solution VIII

In the limit $m = 1$, the solution IV goes over to the hyperbolic solution

$$u(x, t) = A \operatorname{sech}(\xi)e^{i(k_1x - \omega_1t)}, \quad v(x, t) = B \operatorname{sech}(\xi)e^{i(k_2x - \omega_2t)}, \tag{171}$$

provided

$$\begin{aligned}
\beta^2 &= g_{11}A^2 + g_{12}B^2 = g_{21}A^2 + g_{22}B^2, \quad \omega_1 + k_1^3 = 3k_1\beta^2, \\
c + 3k_1^2 &= \beta^2, \quad k_2 = \pm k_1, \quad \omega_2 = \pm \omega_1.
\end{aligned} \tag{172}$$

Note that the \pm signs in Eq. (172) are correlated.

Solution IX

It is straightforward to check that

$$u(x, t) = A \tanh(\xi) e^{i(k_1 x - \omega_1 t)}, \quad v(x, t) = B \tanh(\xi) e^{i(k_2 x - \omega_2 t)}, \quad (173)$$

is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} \beta^2 &= g_{11}A^2 + g_{12}B^2 = g_{21}A^2 + g_{22}B^2, \quad \omega_1 + k_1^3 = -6k_1\beta^2, \\ c + 3k_1^2 &= -2\beta^2, \quad k_2 = \pm k_1, \quad \omega_2 = \pm\omega_1. \end{aligned} \quad (174)$$

Note that the \pm signs in Eq. (174) are correlated. As mentioned above, the corresponding local coupled mKdV Eqs. (135) and (136) do not admit any of the above six solutions unless, of course, if $k_1 = k_2 = \omega_1 = \omega_2 = 0$.

It easily follows from here that corresponding to any solution of the uncoupled nonlocal Eq. (132) of the form

$$u(x, t) = u(\xi), \quad \xi = \beta(x - ct), \quad (175)$$

where $u(\xi)$ is real, there is always a solution of the form

$$u(x, t) = u(\xi) e^{i(k_1 x - \omega_1 t)}, \quad (176)$$

We now show that the coupled Eqs. (128) and (129) admit several new solutions which we present one by one. For simplicity, we present these solutions without the plane wave factor. However, we emphasize that all the coupled solutions discussed below are also valid with the plane wave factor as discussed in the above six solutions.

Solution X

It is easy to check that

$$u = \frac{A[F + \operatorname{sech}(\xi)]}{[D + \operatorname{sech}(\xi)]}, \quad v = \frac{B \operatorname{sech}(\xi)}{[D + \operatorname{sech}(\xi)]}, \quad D > 0, \quad (177)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} c(D - F) &= (D + 2F)\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2F(D - F) &= D^2\beta^2, \quad 2g_{12}B^2F = (2D^2F - D - F)\beta^2. \end{aligned} \quad (178)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (177) provided the relations (178) are satisfied.

Solution XI

It is easy to check that

$$u = \frac{A[F + \operatorname{sech}(\xi)]}{[D + \operatorname{sech}(\xi)]}, \quad v = \frac{B}{[D + \operatorname{sech}(\xi)]}, \quad D > 0, \quad (179)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} c(D - F) &= (6FD^2 - 2D - F)\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2(D - F) &= 3D(1 - 2D^2)\beta^2, \quad 2g_{12}B^2 = D(2D^2F - D - F)\beta^2. \end{aligned} \quad (180)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (179) provided the relations (180) hold good.

Solution XII

It is not difficult to check that

$$u = \frac{A \operatorname{sech}(\xi)}{[D + \operatorname{sech}(\xi)]}, \quad v = \frac{B}{[D + \operatorname{sech}(\xi)]}, \quad D > 0, \quad (181)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} c &= -2\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2 &= 3(2D^2 - 1)\beta^2, \quad 2g_{12}B^2 = -D^2\beta^2. \end{aligned} \quad (182)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (181) provided the relations (182) hold good.

Solution XIII

It is easy to check that

$$u = \frac{A[F + \cos(\xi)]}{[D + \cos(\xi)]}, \quad v = \frac{B \cos(\xi)}{[D + \cos(\xi)]}, \quad D > 1, \quad (183)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} D(F - D)c &= (D^2 - 6 + 2DF)\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2F(F - D) &= (D^2 - 2)\beta^2, \quad 2g_{12}B^2DF = (D^2 + DF - 2)\beta^2. \end{aligned} \quad (184)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (183) provided the relations (184) hold good.

Solution XIV

It is straightforward to check that

$$u = \frac{A[F + \cos(\xi)]}{[D + \cos(\xi)]}, \quad v = \frac{B}{[D + \cos(\xi)]}, \quad D > 1, \quad (185)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} (D - F)c &= (2D + F)\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2(D - F) &= D\beta^2, \quad 2g_{12}B^2 = (D^2 + DF - 2)\beta^2. \end{aligned} \quad (186)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (185) provided the relations (186) hold good.

Solution XV

It is easy to check that

$$u = \frac{A \cos(\xi)}{[D + \cos(\xi)]}, \quad v = \frac{B}{[D + \cos(\xi)]}, \quad D > 1, \quad (187)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} c &= 2\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2 &= \beta^2, \quad 2g_{12}B^2 = (D^2 - 2)\beta^2. \end{aligned} \quad (188)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (187) provided the relations (188) hold good.

Solution XVI

It is not difficult to check that

$$u = \frac{A[F + \tanh(\xi)]}{[D + \tanh(\xi)]}, \quad v = \frac{B \tanh(\xi)}{[D + \tanh(\xi)]}, \quad D > 1, \quad (189)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} cD(F - D) &= 2(2DF + D^2 - 3)\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ g_{11}A^2F(F - D) &= (D^2 - 1)\beta^2, \quad g_{12}B^2DF = (DF + D^2 - 1 - D^3F)\beta^2. \end{aligned} \quad (190)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (189) provided the relations (190) hold good.

Solution XVII

It is easy to check that

$$u = \frac{A[F + \tanh(\xi)]}{[D + \tanh(\xi)]}, \quad v = \frac{B}{[D + \tanh(\xi)]}, \quad D > 1, \quad (191)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} c(D - F) &= 2(2D + F - 3FD^2)\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ g_{11}A^2(D - F) &= D(1 - D^2)\beta^2, \quad g_{12}B^2 = (D^2 - 1)(1 - DF)\beta^2. \end{aligned} \quad (192)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (191) provided the relations (192) hold good.

Solution XVIII

It is straightforward to check that

$$u = \frac{A[F + \operatorname{dn}(\xi, m)]}{[D + \operatorname{dn}(\xi, m)]}, \quad v = \frac{B\operatorname{dn}(\xi, m)}{[D + \operatorname{dn}(\xi, m)]}, \quad D > 0, \quad (193)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} cD(D - F) &= [(2 - m)D(D + 2F) - 6(1 - m)]\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2F(D - F) &= [(2 - m)D^2 - 2(1 - m)]\beta^2, \\ 2g_{12}B^2DF &= [2(1 - m) - (2 - m)D(D + F) + 2D^3F]\beta^2. \end{aligned} \quad (194)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (193) provided the relations (194) hold good.

Solution XIX

It is easy to check that

$$u = \frac{A[F + \operatorname{dn}(\xi, m)]}{[D + \operatorname{dn}(\xi, m)]}, \quad v = \frac{B}{[D + \operatorname{dn}(\xi, m)]}, \quad D > 0, \quad (195)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} c(D - F) &= [6FD^2 - (2 - m)(2D + F)]\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2(D - F) &= D[2D^2 - (2 - m)]\beta^2, \\ 2g_{12}B^2 &= [2(1 - m) - (2 - m)D(D + F) - 2D^3F]\beta^2. \end{aligned} \quad (196)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (195) provided the relations (196) hold good.

Solution XX

It is not difficult to check that

$$u = \frac{A \operatorname{dn}(\xi, m)}{[D + \operatorname{dn}(\xi, m)]}, \quad v = \frac{B}{[D + \operatorname{dn}(\xi, m)]}, \quad D > 0, \quad (197)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} c &= -2(2 - m)\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2 &= [2D^2 - (2 - m)]\beta^2, \quad 2g_{12}B^2 = [2(1 - m) - (2 - m)D^2]\beta^2. \end{aligned} \quad (198)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (197) provided the relations (198) hold good.

Solution XXI

It is easy to check that

$$u = \frac{A[F + \operatorname{cn}(\xi, m)]}{[D + \operatorname{cn}(\xi, m)]}, \quad v = \frac{B \operatorname{cn}(\xi, m)}{[D + \operatorname{cn}(\xi, m)]}, \quad D > 1, \quad (199)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} D(D - F)c &= [6(1 - m) + (2m - 1)D(D + 2F)]\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2F(D - F) &= [2(1 - m) + (2m - 1)D^2]\beta^2, \\ 2g_{12}B^2DF &= [2mD^3F - (2m - 1)D(D + F) - 2(1 - m)]\beta^2. \end{aligned} \quad (200)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (199) provided the relations (200) hold good.

Solution XXII

It is easy to check that

$$u = \frac{A[F + \operatorname{cn}(\xi, m)]}{[D + \operatorname{cn}(\xi, m)]}, \quad v = \frac{B}{[D + \operatorname{cn}(\xi, m)]}, \quad D > 1, \quad (201)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} (D - F)c &= [6mFD^2 - (2m - 1)(2D + F)]\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2(D - F) &= D[2m - 1 - 2mD^2]\beta^2, \\ 2g_{12}B^2 &= -[2(1 - m) + D(F + D) + 2mD^3F]\beta^2. \end{aligned} \quad (202)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (201) provided the relations (202) hold good.

Solution XXIII

It is easy to check that

$$u = \frac{A \cos(\xi)}{[D + \cos(\xi)]}, \quad v = \frac{B}{[D + \cos(\xi)]}, \quad D > 1, \quad (203)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} c &= -2(2m - 1)\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2 &= [2mD^2 - (2m - 1)]\beta^2, \quad 2g_{12}B^2 = -[(2m - 1)D^2 + 2(1 - m)]\beta^2. \end{aligned} \quad (204)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (203) provided the relations (204) hold good.

We now present six soliton solutions with a power law tail.

Solution XXIV

It is not difficult to check that

$$u = \frac{A(F + x^2)}{D + x^2}, \quad v = \frac{Bx^2}{D + x^2}, \quad D > 0, \quad (205)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} cD(F - D) &= 6(F + 2D), \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ F(F - D)g_{11}A^2 &= 3D, \quad FDg_{12}B^2 = F + 3D. \end{aligned} \quad (206)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (205) provided the relations (206) hold good.

Solution XXV

It is straightforward to check that

$$u = \frac{A(F + x^2)}{D + x^2}, \quad v = \frac{B}{D + x^2}, \quad D > 0, \quad (207)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} c(D - F) &= 6, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ (F - D)g_{11}A^2 &= 1, \quad (D - F)g_{12}B^2 = F + 3D. \end{aligned} \quad (208)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (207) provided the relations (208) hold good.

Solution XXVI

It is easy to check that

$$u = \frac{Ax^2}{D + x^2}, \quad v = \frac{B}{D + x^2}, \quad D > 0, \quad (209)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} cD &= 6, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ Dg_{11}A^2 &= 1, \quad g_{12}B^2 = 3D. \end{aligned} \quad (210)$$

It is worth pointing out that the corresponding local coupled mKdV Eqs. (135) and (136) also admit the solution (209) provided the relations (210) hold good.

Solution XXVII

It is not difficult to check that

$$u = \frac{A(F + x^2)}{D + x^2}, \quad v = \frac{Bx}{\sqrt{D + x^2}}, \quad D > 0, \quad (211)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} cD(F - D)^2 &= 12(F^2 + 2DF - D^2) = 3D(4F^2 - D^2 + 2FD), \\ (F - D)^2g_{11}A^2 &= 4D, \quad D(F - D)g_{12}B^2 = 2(F + 3D), \\ (F - D)^2g_{21}A^2 &= (5/2)D^2, \quad (F - D)g_{22}B^2 = 2F + 3D. \end{aligned} \quad (212)$$

On equating the two expressions for c we find that $D \neq 4$ and further if $D = 1$ then $F = 1/2$. Moreover, if $D \neq 1, 4$, then F is given in terms of D by

$$F = \frac{D(4 - D) \pm D\sqrt{32 - 28D + 5D^2}}{4(D - 1)}. \quad (213)$$

Notice that F is real provided $D < 8/5$ or $D > 4$.

Solution XXVIII

It is easy to check that

$$u = \frac{Ax^2}{D + x^2}, \quad v = \frac{Bx}{\sqrt{D + x^2}}, \quad D > 0, \quad (214)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} c = -3, \quad D = 4, \quad g_{11}A^2 = 1, \quad g_{12}B^2 = -3/2, \\ g_{21}A^2 = 5/2, \quad g_{22}B^2 = -3. \end{aligned} \quad (215)$$

Solution XXIX

It is straightforward to check that

$$u = \frac{A}{D + x^2}, \quad v = \frac{B}{\sqrt{D + x^2}}, \quad D > 0, \quad (216)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (128) and (129) provided

$$\begin{aligned} c = 12, \quad D = 1, \quad g_{11}A^2 = 4, \quad g_{12}B^2 = 2, \\ g_{21}A^2 = 5/2, \quad g_{22}B^2 = 2. \end{aligned} \quad (217)$$

4 Conclusion And Open Problems

In this paper we have obtained several new solutions of the coupled AM variant of the nonlocal NLS equation as well as the coupled mKdV equations. Moreover, we compared and contrasted these solutions with the solutions (in case they are admitted) of the corresponding coupled NLS or mKdV equations, respectively. In the nonlocal NLS case, particular mention may be made of the host of the periodic as well as the hyperbolic nonreciprocal solutions in a special subclass of the coupled nonlocal NLS solutions. In the nonlocal mKdV case, we have shown that in contrast to the local mKdV case, the nonlocal coupled mKdV equations admit plane wave solutions as well as moving soliton solutions. This paper raises several open questions, some of which are as follows.

1. While we have obtained a large number of nonreciprocal solutions for the coupled nonlocal (as well as local) NLS equations, we have so far not been able to obtain similar solutions for either the coupled nonlocal or the local mKdV equations. It is worthwhile asking if nonreciprocal solutions can also be obtained for the mKdV or other coupled nonlocal or even local nonlinear equations.
2. Unlike the local mKdV equations we have shown that the nonlocal mKdV equations admit plane wave solutions as well as soliton solutions multiplied by the plane wave factor. It is worth asking if similar solutions can also be obtained for the other nonlocal equations.
3. For the full coupled nonlocal NLS equations as well as coupled nonlocal mKdV equations we have obtained novel periodic as well as hyperbolic solutions. One obvious question is, can one similarly obtain exact solutions of the other coupled nonlocal equations, e.g., coupled Hirota or coupled KdV equations or coupled Yang-type nonlocal NLS? Further, in the coupled nonlocal KdV case do we again get plane wave solutions as well as soliton solutions multiplied by the plane wave factor?

5 Acknowledgment

One of us (AK) is grateful to Indian National Science Academy (INSA) for the award of INSA Honorary Scientist position at Savitribai Phule Pune University. The work at Los Alamos National Laboratory was carried out under the auspices of the US DOE and NNSA under contract No. DEAC52-06NA25396.

6 Appendix A: Novel Periodic Superposed Solutions of a Coupled Local NLS Model

We now present those solutions of the coupled local NLS Eqs. (6) and (7) which are, however, not the solutions of the nonlocal coupled Eqs. (1) and (2).

In this context, it is interesting to observe that unlike the coupled nonlocal AM Eqs. (1) and (2), the (local) coupled NLS Eqs. (6) and (7) admit the plane wave solution

$$u(x, t) = Ae^{i(\omega_1 t - k_1 x)}, \quad v(x, t) = Be^{i(\omega_2 t - k_2 x)}, \quad (218)$$

provided

$$\begin{aligned}\omega_1 + k_1^2 &= [g_{11}A^2 + g_{12}B^2], \\ \omega_2 + k_2^2 &= [g_{21}A^2 + g_{22}B^2].\end{aligned}\quad (219)$$

We now show that corresponding to any stationary soliton solution of the coupled local NLS Eqs. (6) and (7), one always has a corresponding moving soliton solution. The proof is straightforward. Let us assume that there is a stationary soliton solution of the form

$$u(x, t) = u(\beta x)e^{i(\omega_1 t - k_1 x)}, \quad v(x, t) = v(\beta x)e^{i(\omega_2 t - k_2 x)}, \quad \xi = \beta(x - ct), \quad (220)$$

which is an exact solution of the coupled Eqs. (6) and (7). Using the ansatz (220) in Eqs. (6) and (7), we find that it is an exact solution provided

$$\begin{aligned}\beta^2 u''(\eta) &= \omega_1 u + [g_{11}|u|^2 + g_{12}|v|^2]u, \\ \beta^2 v''(\eta) &= \omega_2 v + [g_{21}|u|^2 + g_{22}|v|^2]v.\end{aligned}\quad (221)$$

Let us now consider the moving soliton solutions. Let

$$u(x, t) = u(\xi)e^{i(\omega_1 t - k_1 x)}, \quad v(x, t) = v(\xi)e^{i(\omega_2 t - k_2 x)}, \quad \xi = \beta(x - ct), \quad (222)$$

be an exact solution of the coupled local NLS Eqs. (6) and (7). On substituting this ansatz in Eqs. (6) and (7), we find that it is an exact solution provided

$$\begin{aligned}k_1 &= k_2, \quad c = -2k_1, \quad \beta^2 u''(\xi) = (\omega_1 + k_1^2)u + [g_{11}|u|^2 + g_{12}|v|^2]u, \\ \beta^2 v''(\xi) &= (\omega_2 + k_1^2)v + [g_{21}|u|^2 + g_{22}|v|^2]v.\end{aligned}\quad (223)$$

Now notice that in the limit $k_1 = k_2 = c = 0$, the coupled Eqs. (223) indeed go over to Eqs. (221).

For simplicity, we now only present the new stationary soliton solutions of the local coupled NLS Eqs. (6) and (7). But as proved above, it is understood that there are corresponding moving soliton solutions.

Solution IA

It is not difficult to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A \operatorname{dn}(\beta x, m)}{D + \operatorname{sn}(\beta x, m)}, \quad v(x, t) = e^{i\omega_2 t} \frac{B \operatorname{cn}(\beta x, m)}{D + \operatorname{sn}(\beta x, m)}, \quad (224)$$

with $A, B > 0, D > 1$, is an exact solution of the coupled Eqs. (6) and (7) provided

$$\begin{aligned}\omega_1 = \omega_2 &= \frac{(1+m)\beta^2}{2}, \quad g_{12}B^2 = \frac{3(mD^2-1)\beta^2}{2}, \quad g_{11}A^2 = \frac{(D^2-1)\beta^2}{2}, \\ g_{22}B^2 &= \frac{(mD^2-1)\beta^2}{2}, \quad g_{21}A^2 = \frac{3(D^2-1)\beta^2}{2}.\end{aligned}\quad (225)$$

Thus $g_{12}, g_{22} > 0$ if $1 < D^2 < 1/m$ while $g_{11}, g_{21} > 0$. On the other hand $\omega_1, \omega_2 > 0$.

Solution IIA

It is easy to check that

$$u(x, t) = e^{i\omega_1 t} \frac{A[F + \tanh(\beta x)]}{D + \tanh(\beta x)}, \quad v(x, t) = e^{i\omega_2 t} \frac{B \operatorname{sech}(\beta x)}{D + \tanh(\beta x)}, \quad (226)$$

with $A, B > 0, D > 1$, is an exact solution of the coupled local NLS Eqs. (6) and (7) (but not nonlocal coupled Eqs. (1) and (2)) provided

$$\begin{aligned} \omega_1 &= 4\beta^2, \quad \omega_2 = \beta^2, \quad F = \pm 1, \quad g_{11}A^2 = (D \pm 1)^2\beta^2, \\ g_{12}B^2 &= 2(D^2 - 1)\beta^2, \quad g_{22} = g_{12}, \quad g_{21} = 0. \end{aligned} \quad (227)$$

Note that the \pm signs here are correlated.

Solution IIIA

It is straightforward to check that

$$\phi_1 = A \left[\frac{1 - m}{\operatorname{dn}^2(\beta x, m)} + y \right] + \frac{B\sqrt{m}(1 - m)\operatorname{sn}(\beta x, m)}{\operatorname{dn}^2(\beta x, m)}, \quad (228)$$

$$\phi_2 = \frac{D\sqrt{1 - m}}{\operatorname{dn}(\beta x, m)} + \frac{E\sqrt{m(1 - m)}\operatorname{sn}(\beta x, m)}{\operatorname{dn}(\beta x, m)}, \quad (229)$$

is an exact nonreciprocal solution of the unusual coupled local NLS Eqs. (39) and (40) (but not the nonlocal Eqs. (34) and (35)) provided $0 < m < 1$ and further

$$\begin{aligned} d_1 D^2 &= -\frac{3\beta^2}{2}, \quad a_1 = \left[\frac{(7 + m)}{2} + 3(1 - m)y \right] \beta^2, \quad E = \pm D, \quad B = \pm A, \\ a_2 &= \frac{(1 + m)\beta^2}{2}, \quad d_2 D^2 = -\frac{\beta^2}{2}, \quad y = \frac{-(5 - m) \pm \sqrt{1 + 14m + m^2}}{6} \end{aligned} \quad (230)$$

Note that the \pm signs in D, E are correlated with the signs in B and A . Further, the solution is valid for arbitrary values of A .

Solution IVA

It is easy to check that

$$\phi_1 = \frac{A\sqrt{m}\operatorname{cn}(\beta x, m)}{\operatorname{dn}^2(\beta x, m)} + \frac{Bm\operatorname{cn}(\beta x, m)\operatorname{sn}(\beta x, m)}{\operatorname{dn}^2(\beta x, m)}, \quad (231)$$

$$\phi_2 = \frac{D\sqrt{1 - m}}{\operatorname{dn}(\beta x, m)} + \frac{E\sqrt{m(1 - m)}\operatorname{sn}(\beta x, m)}{\operatorname{dn}(\beta x, m)}, \quad (232)$$

is an exact nonreciprocal solution of the unusual coupled local NLS Eqs. (39) and (40) (but not the solutions of the coupled nonlocal Eqs. (34) and (35)) provided

$$\begin{aligned} d_1 D^2 &= -\frac{3\beta^2}{2}, \quad a_1 = a_2 = \frac{(1+m)\beta^2}{2}, \quad E = \pm D, \quad B = \pm A, \\ d_2 D^2 &= -\frac{\beta^2}{2}, \quad 0 < m < 1. \end{aligned} \quad (233)$$

Note that the \pm signs in D, E are correlated with the signs in B and A . Further, the solution is valid for arbitrary values of A .

We now present three solutions of the coupled local NLS where in general $k_1 \neq k_2$. Further, in these cases even solutions with $c = 0$ are allowed even though k_1, k_2 are still nonzero and in general unequal.

Solution VA

It is not difficult to check that

$$\begin{aligned} u(x, t) &= \sqrt{n_1} [\cos(\theta_1) \tanh(\xi) + i \sin(\theta_1)] e^{i(k_1 x - \omega_1 t)}, \\ v(x, t) &= \sqrt{n_2} [\cos(\theta_2) \tanh(\xi) + i \sin(\theta_2)] e^{i(k_2 x - \omega_2 t)}, \quad \xi = \beta(x - ct), \end{aligned} \quad (234)$$

is an exact solution of the local NLS coupled Eqs. (6) and (7) provided

$$n_1 g_{11} \cos^2(\theta_1) + n_2 g_{12} \cos^2(\theta_2) = n_1 g_{21} \cos^2(\theta_1) + n_2 g_{22} \cos^2(\theta_2) = -2\beta^2, \quad (235)$$

$$\omega_1 - k_1^2 + n_1 g_{11} + n_2 g_{12} = 0, \quad (236)$$

$$\omega_2 - k_2^2 + n_1 g_{21} + n_2 g_{22} = 0, \quad (237)$$

$$c = 2k_1 + 2\beta \tan(\theta_1) = 2k_2 + 2\beta \tan(\theta_2). \quad (238)$$

Solution VIA

It is easy to check that

$$\begin{aligned} u(x, t) &= \sqrt{n_1} [\cos(\theta_1) \tanh(\xi) + i \sin(\theta_1)] e^{i(k_1 x - \omega_1 t)}, \\ v(x, t) &= \sqrt{n_2} [\sin(\theta_2) + i \cos(\theta_2) \tanh(\xi)] e^{i(k_2 x - \omega_2 t)}, \quad \xi = \beta(x - ct), \end{aligned} \quad (239)$$

is an exact solution of the local NLS coupled Eqs. (6) and (7) provided Eqs. (235) to Eqs. (237) are satisfied and further

$$c = 2k_1 + 2\beta \tan(\theta_1) = 2k_2 - 2\beta \tan(\theta_2). \quad (240)$$

Solution VIIA

It is straightforward to check that

$$\begin{aligned} u(x, t) &= \sqrt{n_1} [\sin(\theta_1) + i \cos(\theta_1) \tanh(\xi)] e^{i(k_1 x - \omega_1 t)}, \\ v(x, t) &= \sqrt{n_2} [\sin(\theta_2) + i \cos(\theta_2) \tanh(\xi)] e^{i(k_2 x - \omega_2 t)}, \quad \xi = \beta(x - ct), \end{aligned} \quad (241)$$

is an exact solution of the local NLS coupled Eqs. (6) and (7) provided Eqs. (235) to Eqs. (237) are satisfied and further

$$c = 2k_1 - 2\beta \tan(\theta_1) = 2k_2 - 2\beta \tan(\theta_2). \quad (242)$$

7 Appendix B: Novel Periodic Superposed Solutions of a Coupled Local mKdV Model

We now present those solutions of the coupled local mKdV Eqs. (135) and (136) which are however not the solutions of the coupled nonlocal Eqs. (128) and (129).

Solution IB

It is easy to check that

$$u(x, t) = \frac{A \tanh(\xi)}{[D + \tanh(\xi)]}, \quad v(x, t) = \frac{B}{[D + \tanh(\xi)]}, \quad D > 1, \quad (243)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (135) and (136) provided

$$\begin{aligned} c &= 4\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ g_{11}A^2 &= (1 - D^2)\beta^2, \quad g_{12}B^2 = (D^2 - 1)\beta^2. \end{aligned} \quad (244)$$

Solution IIB

It is not difficult to check that

$$u(x, t) = \frac{A[F + \operatorname{sn}(\xi, m)]}{[D + \operatorname{sn}(\xi, m)]}, \quad v(x, t) = \frac{B \operatorname{sn}(\xi, m)}{[D + \operatorname{sn}(\xi, m)]}, \quad D > 1, \quad (245)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (135) and (136) provided

$$\begin{aligned} cD(F - D) &= 2[(1 + m)D(D + 2F) - 3]\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2F(F - D) &= [(1 + m)D^2 - 2]\beta^2, \\ 2g_{12}B^2DF &= [(1 + m)D(D + F) - 2 - 2mD^3F]\beta^2. \end{aligned} \quad (246)$$

Solution IIIB

It is easy to check that

$$u(x, t) = \frac{A[F + \operatorname{sn}(\xi, m)]}{[D + \operatorname{sn}(\xi, m)]}, \quad v(x, t) = \frac{B}{[D + \operatorname{sn}(\xi, m)]}, \quad A, B > 0, \quad D > 1, \quad (247)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (135) and (136) provided

$$\begin{aligned} c(D - F) &= [(2D + F)(1 + m) - 6mFD^2]\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2(D - F) &= D[1 + m - 2mD^2]\beta^2, \\ 2g_{12}B^2 &= [(1 + m)(D^2 + DF) - 2 - 2mFD^3]\beta^2. \end{aligned} \quad (248)$$

Solution IVB

It is straightforward to check that

$$u(x, t) = \frac{A\operatorname{sn}(\xi, m)}{[D + \operatorname{sn}(\xi, m)]}, \quad v(x, t) = \frac{B}{[D + \operatorname{sn}(\xi, m)]}, \quad D > 1, \quad (249)$$

where $\xi = \beta(x - ct)$ is an exact solution of the coupled Eqs. (135) and (136) provided

$$\begin{aligned} c &= 2(1 + m)\beta^2, \quad g_{11} = g_{21}, \quad g_{12} = g_{22}, \\ 2g_{11}A^2 &= (1 + m - 2mD^2)\beta^2, \quad 2g_{12}B^2 = [(1 + m)D^2 - 2]\beta^2. \end{aligned} \quad (250)$$

References

- [1] M. J. Ablowitz and Z. H. Musslimani, Phys. Rev. Lett. **110** (2013) 064105.
- [2] M. J. Ablowitz and Z. H. Musslimani, Nonlinearity **29** (2016) 915.
- [3] V. S. Gerdjikov and A. Saxena, J. Math. Phys. **58** (2017) 013502.
- [4] F. He, E. Fan and J. Xu, Commun. Theor. Phys. **71** (2019) 475; arXiv:1804.10863.
- [5] M. Gürses and A. Peçkan, Comm. Nonlin. Sc. Num. Simul. **67** (2019) 427; arXiv:1711.01588.
- [6] J. Cen, F. Correa and A. Fring, J. Math. Phys. **60** (2019) 081508.
- [7] Y. Zia, R. Yao and X. Xin, Chinese Physics **B31** (2021) 020401

- [8] J. Yang, Phys. Rev. **E98** (2018) 042202.
- [9] Z. H. Musslimani et al., Phys. Rev. Lett. **100** (2008) 030402; K. G. Makris et al., ibid. **100** (2008) 103904; A. Ruschaupt, F. Delgado and J. G. Muga, J. Phys. **A38** (2005), L171; A. Guo et al., Phys. Rev. Lett. **103** (2009) 093902; C. E. Rutter, K. G. Makris, R. El-Ganainy, R. N. Christodulides, M. Segev and D. Kip, Nature Phys. **6** (2010) 192.
- [10] A. Khare and A. Saxena, J. Math. Phys. **55** (2014) 032701.
- [11] A. Khare and A. Saxena, J. Math. Phys. **56** (2015) 032104.
- [12] A. Khare and A. Saxena, J. Math. Phys. **63** (2022) 122903.
- [13] A. Khare, S. Banerjee and A. Saxena, Ann. Phys. **455** (2023) 169362.
- [14] A. Khare, S. Banerjee and A. Saxena, Ann. Phys. **457** (2023) 169433.
- [15] See, for example, M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs and Mathematical Tables*, Dover, NY (1964).
- [16] V. S. Zakharov and A. B. Shabat, JETP **37** (1973) 823.