On Positivity Preservers and their Generators

Philipp J. di Dio

Department of Mathematics and Statistics, University of Konstanz, Universitätsstraße 10, D-78464 Konstanz, Germany

Zukunftskolleg, Universitity of Konstanz, Universitätsstraße 10, D-78464 Konstanz, Germany

Abstract

We study K-positivity preserves $T : \mathbb{R}[x_1, \ldots, x_n] \to \mathbb{R}[x_1, \ldots, x_n]$ with constant coefficients and $K \subseteq \mathbb{R}^n$; especially their generators A, i.e., $T = e^A$. We completely describe the set of generators A such that e^{tA} is a positivity preserver for all $t \ge 0$ (Theorem 4.8). We solve the long standing open problem of describing the K-positivity preservers with constant coefficients when $K \subsetneq \mathbb{R}^n$ (Theorem 3.4). The complete description of generators of $[0, \infty)$ -positivity preservers is given in Theorem 6.4. We give an example of a strange positivity action which on $[0, \infty)$ maps odd non-negative polynomials to odd non-negative polynomials but even non-negative polynomials are mapped to non-negative functions which are not polynomials. We introduce the concept of a convolution of (moment) sequences. The techniques we use come from (infinite dimensional matrix) Lie group theory, measure theory (infinitely divisible measures and the Lévy–Khinchin formula), and of course the theory of moments.

Keywords: positivity preserver, generator, constant coefficient, moments 2020 MSC: Primary 44A60, 47A57, 15A04; Secondary 12D15, 45P05, 47B38.

1. Introduction

Non-negative polynomials are widely applied and studied. A first step to investigate non-negative polynomials it to study the cone

$$\operatorname{Pos}(K) := \{ p \in \mathbb{R}[x_1, \dots, x_n] \mid p \ge 0 \text{ on } K \}$$

of non-negative polynomials on some $K \subseteq \mathbb{R}^n$.

The second logical step is to study (linear) maps

$$T: \mathbb{R}[x_1, \dots, x_n] \to \mathbb{R}[x_1, \dots, x_n] \tag{1}$$

between polynomials which preserve non-negativity [1-3], i.e.,

$$T\operatorname{Pos}(K) \subseteq \operatorname{Pos}(K).$$
 (2)

Preprint submitted to working project

August 22, 2023

It is folklore that every linear operator (1) is of the form

$$T = \sum_{\alpha \in \mathbb{N}_0^n} q_\alpha \cdot \partial^\alpha \tag{3}$$

with unique polynomial coefficients $q_{\alpha} \in \mathbb{R}[x_1, \ldots, x_n]$, see e.g. [3, Lem. 2.3]. An operator T of the form (3) is called a *differential operator with polynomial coefficients*. Furthermore, T is called a *positivity preserver* if it preserves nonnegative polynomials, i.e., (2) holds with $K = \mathbb{R}^n$. If $K \subsetneq \mathbb{R}^n$, then T in (2) is called a *K*-positivity preserver. T is called a positivity preserver with constant coefficients if additionally $q_{\alpha} \in \mathbb{R}$ for all $\alpha \in \mathbb{N}_0^n$. By [2, Thm. 3.1] T is a positivity preserver if and only if $(\alpha! \cdot q_{\alpha}(y))_{\alpha \in \mathbb{N}_0^n}$ is a moment sequence for all $y \in \mathbb{R}^n$.

The following is a linear operator T which is not a positivity preserver.

Example 1.1. Let $K \geq 3$ and $a \in \mathbb{R} \setminus \{0\}$. Then

$$e^{a\partial_x^K} = \sum_{k=0}^{\infty} \frac{a^{k \cdot K} \cdot \partial_x^{k \cdot K}}{k!}$$

is not a positive preserver since $q_{2K} = a^{2K} \neq 0$ but $q_{2K+2} = 0$, i.e., $(k! \cdot q_k)_{k \in \mathbb{N}_0}$ is not a moment sequence.

From elementary calculations and measure theory we get the following additional properties of positivity preservers. Let T be a positivity preserver with constant coefficients and μ be a representing measure of the corresponding moment sequence $s = (s_{\alpha})_{\alpha \in \mathbb{N}_{0}^{n}} = (\alpha ! q_{\alpha})_{\alpha \in \mathbb{N}_{0}^{n}}$, then

$$(Tf)(x) = \int_{\mathbb{R}^n} f(x+y) \, \mathrm{d}\mu(y)$$

for all $f \in \mathbb{R}[x_1, \ldots, x_n]$. For short we call μ also a representing measure of T. If follows for two such operators T and T' with representing measures μ and μ' that

$$(TT'f)(x) = \int f(x+y) \, \mathrm{d}(\mu * \mu')(y)$$
 (4)

where $\mu * \mu'$ is the convolution of the two measures μ and μ' :

$$(\mu * \mu')(A) := \int_{\mathbb{R}^n \times \mathbb{R}^n} \chi_A(x+y) \, \mathrm{d}\mu(x) \, \mathrm{d}\mu'(y)$$
$$= \int_{\mathbb{R}^n} \mu(A-y) \, \mathrm{d}\mu'(y) = \int_{\mathbb{R}^n} \mu'(A-x) \, \mathrm{d}\mu(x) \quad (5)$$

for any $A \in \mathfrak{B}(\mathbb{R}^n)$, see e.g. [4, Sect. 3.9]. Here, χ_A is the characteristic function of the Borel set A. For the supports we have

$$\operatorname{supp}\left(\mu \ast \mu'\right) = \operatorname{supp}\mu + \operatorname{supp}\mu'. \tag{6}$$

We abbreviate

$$\mu^{*k} := \underbrace{\mu * \cdots * \mu}_{k\text{-times}}$$

for all $k \in \mathbb{N}$ and set $\mu^{*0} := \delta_0$ with δ_0 the Dirac measure centered at 0.

Previously we investigated the heat semi-group and its action on (nonnegative) polynomial [5, 6]. In [6] we observed the very strange behavior that several non-negative polynomials which are not sums of squares (e.g. the Motzkin and the Choi–Lam polynomial) become a sum of squares in finite time under the heat equation. In [6, Thm. 3.20] we showed that every non-negative $p \in \mathbb{R}[x, y, z]_{\leq 4}$ becomes a sum of squares in finite time. All these observations hold for the heat equation, i.e., the family $(e^{t\Delta})_{t\geq 0}$ of positivity preservers with constant coefficients with the generator Δ . So a natural question to further study these effects is to understand any family $(e^{tA})_{t\geq 0}$ of positivity preservers and hence to determine all possible generators A. This is the main question of the current work.

The paper is structured as follows. In (4) we have seen that the composition of two positivity preservers results in the convolution of the representing measures. Hence, in Section 2 we introduce and study the convolution of (moment) sequences. From this investigation we then give the full description of K-positivity preservers with constant coefficients in Section 3. In Section 4 we give the full description of positivity generators and in Section 6 we describe all generators of $[0, \infty)$ -positivity preservers with constant coefficients. The Section 5 is devoted to a strange action on polynomials non-negative on $[0, \infty)$, i.e., we give a family of actions which map odd polynomials non-negative on $[0, \infty)$ on odd polynomials non-negative on $[0, \infty)$ but for even polynomials we leave the polynomial ring $\mathbb{R}[x]$. We end our investigation with open problems in Section 7.

2. Convolution of Sequences

Definition 2.1. Let $n \in \mathbb{N}$. We set

$$\mathfrak{S} := \left\{ s = (s_{\alpha})_{\alpha \in \mathbb{N}_{0}^{n}} \, \middle| \, s_{\alpha} \in \mathbb{R} \text{ for all } \alpha \in \mathbb{N}_{0}^{n} \text{ with } s_{0} = 1 \right\}$$

to be the set of sequences with $s_0 = 1$ and

$$\mathfrak{D} := \left\{ \sum_{\alpha \in \mathbb{N}_0^n} q_\alpha \partial^\alpha \, \middle| \, q_\alpha \in \mathbb{R} \text{ for all } \alpha \in \mathbb{N}_0^n \text{ with } q_\alpha = 1 \right\}$$

to be the set of linear operators $T : \mathbb{R}[x_1, \ldots, x_n] \to \mathbb{R}[x_1, \ldots, x_n]$ with constant coefficients and $q_0 = 1$. Additionally, we set

$$\mathfrak{d} := \left\{ \sum_{\alpha \in \mathbb{N}_0^n} a_\alpha \partial^\alpha \, \middle| \, a_\alpha \in \mathbb{R} \text{ for all } \alpha \in \mathbb{N}_0^n \text{ with } a_0 = 0 \right\}.$$

For $d \in \mathbb{N}_0$ we denote by $\mathfrak{S}|_d$, $\mathfrak{D}|_d$, and $\mathfrak{d}|_d$ the truncated cases $|\alpha| \leq d$. We set $\mathfrak{S}|_{\infty} = \mathfrak{S}$, $\mathfrak{D}|_{\infty} = \mathfrak{D}$, and $\mathfrak{d}|_{\infty} = \mathfrak{d}$.

The map

$$D: \mathfrak{S}|_d \to \mathfrak{D}|_d, \quad s = (s_\alpha)_\alpha \mapsto D(s) := \sum_\alpha \frac{s_\alpha}{\alpha!} \cdot \partial^\alpha$$
(7)

is bijective for all $d \in \mathbb{N}_0 \cup \{\infty\}$.

Definition 2.2. Let $n \in \mathbb{N} \cup \{\infty\}$. On $\mathfrak{S}|_d$ we define the *convolution* of sequences by

$$*:\mathfrak{S}|_d\times\mathfrak{S}|_d\to\mathfrak{S}|_d,\quad (s,s')\mapsto s*s':=\mathcal{D}^{-1}(\mathcal{D}(s)\mathcal{D}(s')|_d),$$

i.e., $\mathcal{D}(s * s')|_d = \mathcal{D}(s)\mathcal{D}(s')|_d$. We abbreviate

$$s^{*k} := \underbrace{s * \cdots * s}_{k-\text{times}}$$

for all $k \in \mathbb{N}$ and we set $s^{*0} = (1, 0, 0, \dots) \in \mathfrak{S}|_d$.

For two sequences $s = (s_{\alpha})$ and $t = (t_{\alpha})$ we have that $w = s * t = (w_{\alpha})$ is explicitly given by

$$w_{\alpha} = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \cdot s_{\beta} \cdot t_{\alpha-\beta}$$

where $\binom{\alpha}{\beta} = \frac{\alpha!}{\beta! \cdot (\alpha - \beta)!}$ and $\beta \le \alpha$ if and only if $\beta_i \le \alpha_i$ for all $i = 1, \ldots, n$.

Lemma 2.3. Let $d \in \mathbb{N} \cup \{\infty\}$ and $s, s' \in \mathfrak{S}|_d$ be moment sequences with representing measures μ and μ' . Then $s * s' \in \mathfrak{S}|_d$ is a moment sequence with representing measure $\mu * \mu'$.

Proof. Follows immediately from (4) and Definition 2.2.

The following is a consequence of the well-known facts that the Fourier transform $\hat{\mu}$ of a measure μ uniquely determines the measure, $\widehat{\mu * \nu} = \hat{\mu}\hat{\nu}, \hat{\mu}$ is uniformly continuous, and $\hat{\mu}$ is generated by the moments of μ , see [4, Sect. 3.8].

Corollary 2.4. Let $s, t \in S$ be moment sequences such that t is indeterminate. Then s * t is indeterminate.

Proof. Since s is a moment sequence we find a representing measure μ . Write $\mu = \mu_1 + \mu_2$ where μ_1 has compact support and let s_1 and s_2 be the corresponding moment sequences of μ_1 and μ_2 . Then $\hat{\mu}_1$ is analytic and hence $s_1 * t$ is indeterminate. Therefore, $s * t = s_1 * t + s_2 * t$ is indeterminate.

3. K-Positivity Preservers

Already the simple introduction of the concept of a convolution of two (moment) sequences and their relation to the convolution of two measures and the composition of two positivity preservers (4) enables us to solve a long standing open problem: to characterize all K-positivity presevers with constant coefficients. With that we extend [3].

We need the following definition.

Definition 3.1. Let $n \in \mathbb{N}$ and $K \subseteq \mathbb{R}^n$ be closed. We define the set

$$K^{\sharp} := \{ x \in \mathbb{R}^n \, | \, x + K \subseteq K \}.$$

We have the following properties.

Lemma 3.2. Let $n \in \mathbb{N}$ and $K \subseteq \mathbb{R}^n$ be closed. The following hold:

- (i) $0 \in K^{\sharp} \neq \emptyset$.
- (ii) $(y+K)^{\sharp} = K^{\sharp}$ for all $y \in \mathbb{R}^n$.
- (iii) $x + y \in K^{\sharp}$ for all $x, y \in K^{\sharp}$.

Proof. Follow immediately from Definition 3.1.

Depending on $K \subseteq \mathbb{R}^n$ the set K^{\sharp} can be very diverse, i.e., K^{\sharp} is in general not a cone.

Examples 3.3. (a) Let $c \in \mathbb{R}$ and $K = [c, \infty) \times [-1, 1]$. Then $K^{\sharp} = [0, \infty) \times \{0\}$.

- (b) For a closed convex cone $K \subseteq \mathbb{R}^n$ with apex 0 we have $K^{\sharp} = K$.
- (c) Let $K = \bigcup_{z \in \mathbb{Z}^n} B_{1/4}(z)$. Then $K^{\sharp} = \mathbb{Z}^n$.

We get the following complete description of all K-positivity preservers with constant coefficients.

Theorem 3.4. Let $n \in \mathbb{N}$, $K \subseteq \mathbb{R}^n$ be closed, and $s \in \mathfrak{S}$ be a sequence. The following are equivalent:

- (i) D(s) is a K-positivity preserver.
- (ii) s is a K^{\sharp} -moment sequence.

If one of the equivalent conditions (i) or (ii) hold and μ is a representing measure of s, then

$$[D(s)p](x) = \int p(x+y) \, \mathrm{d}\mu(y)$$

for all $p \in \mathbb{R}[x_1, \ldots, x_n]$.

Proof. (ii) \Rightarrow (i): Since *s* is a K^{\sharp} -moment sequence, it is also a \mathbb{R}^{n} -moment sequence and its action on $\mathbb{R}[x_{1}, \ldots, x_{n}]$ is given by $D(s)p = \int p(\cdot + y) d\mu(y)$ for some representing measure μ of *s*. If $p \in \text{Pos}(K)$, then $p(\cdot + y) \in \text{Pos}(K)$ for all $y \in K^{\sharp}$ since by Definition 3.1 we have $y + K \subseteq K$. Hence, the integral over μ is an integral over *K*-positive polynomials which proves (i).

(i) \Rightarrow (ii): D(s) is a K-positivity preserver and therefore $D(s)^k = D(s^{*k})$ is a K-positivity preserver for all $k \in \mathbb{N}$. Hence, s^{*k} is a K-moment sequence for all $k \in \mathbb{N}$. Let μ be any representing measure of s and assume that $\sup \mu \not\subseteq K^{\sharp}$, then there exists a $x_0 \in \operatorname{supp} \mu \setminus K^{\sharp}$ and a $k \in \mathbb{N}$ such that $kx_0 \notin K$. Hence, s^{*k} has a representing measure μ^{*k} such that $\sup \mu^{*k} \not\subseteq K$ which is a contradiction to the fact that s^{*k} is a K-moment sequence for any $k \in \mathbb{N}$.

Note, for the representing measure μ of s in the previous theorem we do not require $\operatorname{supp} \mu \subseteq K^{\sharp}$. This inclusion always holds since s is a K^{\sharp} -moment sequence.

With $K = \mathbb{R}^n$ we have $K^{\sharp} = \mathbb{R}^n$ and hence the constant coefficient case in [2] is recovered. For compact $K \subset \mathbb{R}^n$ the implication of Theorem 3.4 is severe.

Corollary 3.5. Let $n \in \mathbb{N}$ and $K \subset \mathbb{R}^n$ be compact. Then every K-positivity preserver T with constant coefficients is of the form T = c for some $c \ge 0$.

Proof. Since K is compact we have $K^{\sharp} = \{0\}$ and by Theorem 3.4 every K-positivity preserver D(s) comes from a K^{\sharp} -moment sequence, i.e., s has the unique representing measure $c\delta_0$ for some $c \geq 0$.

In the next section we will learn more about the K-positivity preservers by looking at their generators, if they exist.

4. Generators of Positivity Preservers

The convolution * on $\mathfrak{S}|_d$ and the composition on $\mathfrak{D}|_d$ give both sets the structure of a commutative (infinite dimensional) Lie group and $\mathfrak{d}|_d$ is the corresponding Lie algebra. With the map (7) $\mathfrak{S}|_d$ and \mathfrak{D}_d are isomorphic Lie groups.

Theorem 4.1. For all $d \in \mathbb{N}_0 \cup \{\infty\}$ the set $\mathfrak{D}|_d$ is a commutative Lie group with Lie algebra $\mathfrak{d}|_d$. The map

$$\exp: \mathfrak{d} \to \mathfrak{D}, \ A \mapsto \exp(A) := \sum_{k=0}^{\infty} \frac{A^k}{k!}$$

is bijective with inverse

$$\log: \mathfrak{D} \to \mathfrak{d}, \ T \mapsto \log(T) := \sum_{k=0}^{\infty} \frac{(1-T)^k}{k}$$

Proof. $d \in \mathbb{N}_0$: By identifying $\mathfrak{D}|_d$ as a matrix group, the statement follows immediately from matrix valued Lie groups and Lie algebras [7, p. 134, No. 15].

 $\underline{d = \infty}$: Since \mathfrak{D} acts on $\mathbb{R}[x_1, \ldots, x_n]$ and every $p \in \mathbb{R}[x_1, \ldots, x_n]$ has finite degree, the claim follows from the case $d = \deg p \in \mathbb{N}_0$ for every $p \in \mathbb{R}[x_1, \ldots, x_n]$.

Note that a Lie group as a differentialble manifold and Lie algebra as the tangent space at the unit element are usually finite dimensional. But also infinite dimensional Lie groups and algebras have been investigated. In our case we have an infinite dimensional Lie group \mathfrak{D} and an infinite dimensional Lie algebra \mathfrak{d} . However, they are commutative and we can always restrict ourselves to the finite dimensional case $\mathfrak{D}|_d$ and $\mathfrak{d}|_d$, $d \in \mathbb{N}$, by the previous theorem and then go to the limit $d \to \infty$.

We want to point out that Hirschman and Widder [8] extensively investigated the inversion theory of convolution transforms. But from Theorem 4.1 we see that the convolution of measures (and therefore functions) is from a moment theoretic point of view trivial when we allow signed moment sequences. For any $s \in \mathfrak{S}$ there is a $s^{-1} \in \mathfrak{S}$ such that $s * s^{-1} * t = t$ for all $t \in \mathfrak{S}$ since \mathfrak{S} is a (Lie) group.

Definition 4.2. By $\mathfrak{D}_+ \subsetneq \mathfrak{D}$ we denote the set of *positivity preservers with* constant coefficients with $q_0 = 1$. Additionally, we set

$$\mathfrak{S}_+ := D^{-1}(\mathfrak{D}_+)$$

and we define the set of *positive generators* \mathfrak{d}_+ by

$$\mathfrak{d}_+ := \left\{ A \in \mathfrak{d} \, \big| \, e^{tA} \in \mathfrak{D}_+ \text{ for all } t \ge 0 \right\}.$$

With the map (7) we see that \mathfrak{S}_+ is a base of the moment cone. Hence, by investigating \mathfrak{D}_+ and \mathfrak{d}_+ we also investigate the moment cone. We have the following examples for elements in \mathfrak{d}_+ .

Examples 4.3 (n = 1).

- (a) Let $a \in \mathbb{R}$. Then $(e^{a\partial_x}f)(x) = f(x+a)$ for all $f \in \mathbb{R}[x]$, i.e., $a\partial_x \in \mathfrak{d}_+$.
- (b) Let $a \in [0, \infty)$. Then $e^{a\partial_x^2} f = \Theta_a * f$ for all $f \in \mathbb{R}[x]$ where Θ_a is the heat kernel, i.e., $a\partial_x^2 \in \mathfrak{d}_+$.
- (c) $\mathfrak{D}_+ 1 \subsetneq \mathfrak{d}_+$. For $\subseteq \text{let } A \in \mathfrak{D}_+$ and $f \in \mathbb{R}[x]_{\leq 2d}$. Then $(A-1)^{2d+k}f = 0$ for all $k \in \mathbb{N}$ and hence

$$e^{t \cdot (A-1)} f = \sum_{k=0}^{2d} \frac{t^k \cdot (A-1)^k}{k!} f \in \mathbb{R}[x]_{\leq 2d}$$

for all $t \ge 0$. If additionally $f \ge 0$, then

$$e^{t \cdot (A-1)} f = e^{-t} \cdot e^{tA} f = e^{-t} \cdot \sum_{k=0}^{\infty} \frac{t^k}{k!} \underbrace{A^k f}_{\geq 0} \geq 0$$

for all $t \ge 0$ since [A, 1] = 0. $\mathfrak{D}_+ - 1 \neq \mathfrak{d}_+$ follows from (a) and (b).

Hence, we have

$$\mathbb{R}_{\geq 0} \cdot \partial_x^2 + \mathbb{R} \cdot \partial_x + \mathbb{R}_{\geq 0} \cdot (\mathfrak{D}_+ - 1) \subseteq \mathfrak{d}_+.$$
(8)

The next results show that (a) and (b) are the only finite positive generators.

Lemma 4.4. Let $A = \sum_{k=1}^{K} a_k \partial_x^k \in \mathfrak{d}_+$. Then $K \leq 2$.

Proof. Let $K \geq 3$ and $a_K = 1$. By Example 1.1 we have that $e^{\partial_x^K}$ is not a positivity preserver, i.e., there is a $f_0 \in \mathbb{R}[x]$ with $f_0 \geq 0$ and $x_0 \in \mathbb{R}$ such that $(e^{\partial_x^K} f_0)(x_0) = -1$.

Assume to the contrary that $A \in \mathfrak{d}_+$. By scaling x and A we have that also $A_{\lambda} := \sum_{k=1}^{K} \lambda^{K-k} a_k \partial_x^k \in \mathfrak{d}_+$ for all $\lambda > 0$. Since $(e^{A_{\lambda}} f_0)(x_0)$ is continuous in λ and $(e^{A_0} f_0)(x_0) = -1$ there exists a $\lambda_0 \in (0, \infty)$ such that $(e^{A_{\lambda_0}} f_0)(x_0) < 0$. Hence, $A \notin \mathfrak{d}_+$.

Corollary 4.5. Let $n \ge 2$. The following are equivalent:

(i) $A = \sum_{\alpha \in \mathbb{N}_0^n : |\alpha| = 1, \dots, K} a_\alpha \partial^\alpha \in \mathfrak{d}_+ \text{ for some } K \ge 2.$ (ii) $A = y \cdot \nabla + \sum_{i, j=1}^n a_{i,j} \cdot \partial_{x_i} \partial_{x_j} \text{ with } (a_{i,j})_{i,j=1}^n \succeq 0 \text{ and } y \in \mathbb{R}^n.$

In Theorem 4.8 we completely describe \mathfrak{d}_+ and hence showing that the inclusion (8) is strict. Theorem 4.1 and the description of \mathfrak{D}_+ through moment sequences and hence through non-negative polynomials suggested an algebraic approach. This seems not possible or is at least too complicated. We solve it with a measure theoretic approach using the Lévy–Khinchin formula on \mathbb{R}^n [9, Satz 16.17]

$$\psi(t) = -\frac{1}{2}t^T \cdot \Sigma \cdot t + i\langle b, t \rangle + \int (e^{i\langle t, x \rangle} - 1 - i\langle t, x \rangle \cdot \mathbb{1}_{\|x\|_2 < 1}) \, \mathrm{d}\nu(x). \tag{9}$$

Here, we have $\Sigma \succeq 0$ and $\|\cdot\|_2$ the l^2 -norm on \mathbb{R}^n . ν is called the *Lévy measure*. This gives us the main theorem, the complete description of \mathfrak{d}_+ . We need the following lemma.

Lemma 4.6. $A \in \mathfrak{d}_+$ iff e^A has an infinitely divisible representing measure.

Proof. \Rightarrow : Since $A \in \mathfrak{d}_+$ we have that for every $k \in \mathbb{N}$ the operator $e^{A/k}$ is a positivity preserver and has a representing measure μ_k . Then $\mathcal{F} = (\mu_k^{*k})_{k \in \mathbb{N}}$ is a family of representing measures of e^A and since $\mathbb{R}[x_1, \ldots, x_n]$ is an adapted space the family \mathcal{F} is vaguely compact [10, Thm. 1.19]. Hence, \mathcal{F} has an accumulation point μ which is an infinitely divisible measure and a representing measure of e^A by construction.

 \leq : Let μ be an infinitely divisible representing measure of e^A . Then μ^{*q} exists for all $q \in \mathbb{Q} \cap [0, 1]$ and is a representing measure of e^{qA} . Hence, e^{qA} is a positivity preserver for all $q \in \mathbb{Q} \cap [0, 1]$. By continuity of the exp-function (Theorem 4.1) and since $\mathbb{Q} \cap [0, 1]$ is dense in [0, 1] we have that e^{tA} is a positivity preserver for all $t \in [0, 1]$ and hence $A \in \mathfrak{d}_+$. □

Note that $\mathfrak{d}_+ \subsetneq \log \mathfrak{D}_+$. To see that, take a positivity preserver which is represented by a compact measure.

Corollary 4.7. Let $s \in \mathfrak{S}_+$ be an indeterminate moment sequence. Then $e^{D(s)} = D(t)$ is a positivity preserver which has infinitely many infinitely divisible representing measures, i.e., $t = e^{*s}$ is an indeterminate moment sequence.

Proof. Let μ_0 and μ_1 be two representing measures of s. Then $e^{D(s)}$ is represented by the measures

$$e^{*\mu_{\lambda}} := \sum_{k=0}^{\infty} \frac{\mu_{\lambda}^{*k}}{k!}$$

for $\mu_{\lambda} := \lambda \cdot \mu_0 + (1 - \lambda) \cdot \mu_1$ and $\lambda \in [0, 1]$.

We now know that $A \in \mathfrak{d}_+$ is related to e^A having an infinitely divisible measure. From the Lévy-Khinchin formula (9) on \mathbb{R}^n we can now determine A.

Theorem 4.8. Let $n \in \mathbb{N}$ and e_1, \ldots, e_n be the canonical basis of \mathbb{R}^n . The following are equivalent:

- (i) $A = \sum_{\alpha \in \mathbb{N}_0^n} \frac{a_\alpha}{\alpha!} \cdot \partial^\alpha \in \mathfrak{d}_+.$
- (ii) There exists a matrix $\Sigma = (\sigma_{i,j})_{i,j=1}^n \in \mathbb{R}^n$ with $\Sigma \succeq 0$, a vector $b = (b_1, \ldots, b_n) \in \mathbb{R}^n$, and a measure ν on \mathbb{R}^n such that

$$a_{0} = 0,$$

$$a_{e_{i}} = b_{i} + \int_{\|x\|_{2} \ge 1} x_{i} \, d\nu(x) \qquad \text{for all } i = 1, \dots, n,$$

$$a_{e_{i}+e_{j}} = \frac{1}{2}\sigma_{i,j} + \int_{\mathbb{R}^{n}} x^{e_{i}+e_{j}} \, d\nu(x) \qquad \text{for all } i, j = 1, \dots, n,$$

and

$$a_{\alpha} = \int_{\mathbb{R}^n} x^{\alpha} \, \mathrm{d}\nu(x) \qquad \qquad \text{for all } \alpha \in \mathbb{N}^n_0 \, \text{ with } |\alpha| \ge 3.$$

Proof. By Lemma 4.6 we know that for every $A \in \mathfrak{d}_+$ there exists an infinitely divisible measure μ and that this measure μ is uniquely determined by (9) with the canonical triple (Σ, b, ν) . We show that the entries a_{ij} of A come from the characteristic function $\psi(t)$ since $\hat{\mu}(-it)$ is the moment generating function of μ we have that $\psi = \log \hat{\mu}$ generates A. Since the characteristic function ψ comes from the Fourier transform

$$\psi(t) := \log \int_{\mathbb{R}^n} e^{i\langle x,t\rangle} \,\mathrm{d}\mu(x)$$

of the measure μ . This immediately implies (i) $\Leftrightarrow i^{|\alpha|}a_{\alpha} = \partial^{\alpha}\psi|_{t=0} \Leftrightarrow$ (ii). \Box

The following diagram organizes the relations between the representing measure μ , its Fourier transform $\hat{\mu}$, its characteristic function $\psi = \log \hat{\mu}$, the Lévy–Khinchin formula, the positivity preservers T, and the positive generators A used in the proof of the main theorem:

$$\begin{split} \mu \\ & \updownarrow \\ \hat{\mu}(t) = \int e^{i \langle t, x \rangle} \, \mathrm{d}\mu(x) \\ & \Leftrightarrow \quad T = \sum_{\alpha \in \mathbb{N}_0^n} \frac{s_\alpha}{\alpha!} \cdot \partial^\alpha \\ & \log \Uparrow \exp : \hat{\mu} = e^{\psi} \\ & \log \Uparrow \exp : T = e^A \end{split}$$

Lévy–Khinchin for inifinitely divisible measures:

$$\psi(t) = (it)^T \Sigma(it) + itb + \int e^{i\langle t, x \rangle} - 1 - i\langle t, x \rangle \mathbb{1}_{\{|x| < 1\}} \mathrm{d}\nu(x) \quad \Leftrightarrow \quad A = \sum_{\alpha \in \mathbb{N}_0^n} \frac{a_\alpha}{\alpha!} \cdot \partial^\alpha$$

The horizontal \Leftrightarrow can be seen as (formal) Fourier transformations, i.e., replacing $(it)^{\alpha}$ by ∂^{α} in the formal expansion of $e^{i\langle t,x\rangle}$.

Note, the difference between \mathfrak{d}_+ and a moment sequence is that the representing (Lévy) measure can have a singularity of order ≤ 2 at the origin. This is the reason for the strict inclusion in (8).

5. A Strange Action on $Pos([0,\infty))$

We want to discuss an example of a strange positivity action on $K = K^{\sharp} = [0, \infty)$. We take the following example. Let $\Delta = \partial_x^2$ on $L^2((0, \infty))$ with Dirichlet boundary conditions, i.e.,

$$\partial_t u = \Delta u$$

$$u(0,t) = 0 \quad \text{for all } t \ge 0$$

$$u(\cdot,0) = u_0 \in L^2((0,\infty)).$$

Then

$$u(x,t) = \int_0^\infty k_t(x,y) u_0(y) \, \mathrm{d}y$$

with

$$k_t(x,y) = \frac{1}{\sqrt{4\pi t}} \cdot e^{-(x-y)^2/4} \cdot \left[1 - e^{xy/t}\right]$$

for all $x, y \ge 0$ and t > 0, see [11, Exm. 4.1.1].

For $u_0 \in \mathcal{S}([0,\infty))$ we have that $k_t * u_0 \in \mathcal{S}([0,\infty))$ for all t > 0 and hence we look at the time-dependent moments

$$s_j(t) = \int_K x^j \cdot u(x,t) \, \mathrm{d}x = \int_{[0,\infty)^2} x^j \cdot k_t(x,y) \cdot u_0(y) \, \mathrm{d}x \, \mathrm{d}y$$

for all $j \in \mathbb{N}_0$ and t > 0.

The action on $\mathbb{R}[x]$ is then given by $(T_t p)(x) = \int_0^\infty p(y) \cdot k_t(x,y) \, \mathrm{d}y$. We find for p = 1 that

$$(T_t 1)(x) = \int_0^\infty k_t(x, y) \, \mathrm{d}y$$

is a continuous, non-decreasing function in $x \in [0, \infty)$ with $(T_t 1)(0) = 0$ and $\lim_{x\to\infty} (T_t 1)(x) = 1$, i.e., $T_t 1 \notin \mathbb{R}[x]$ for any $t \in (0, \infty)$. While for $p = x^j$ with $j \in \mathbb{N}$ we get for the time-dependent moments by partial integration

$$\begin{aligned} \partial_t s_j(t) &= \partial_t \int_0^\infty x^j \cdot u(x,t) \, \mathrm{d}x \\ &= \int_0^\infty x^j \cdot u(x,t)' \, \mathrm{d}x \\ &= \underbrace{x^j \cdot u(x,t)'}_{=0} \Big|_{x=0}^\infty - \int_0^\infty j \cdot x^{j-1} \cdot u(x,t)' \, \mathrm{d}x \\ &= \underbrace{-j \cdot x^{j-1} \cdot u(x,t)}_{=0} \Big|_{x=0}^\infty + \int_0^\infty j \cdot (j-1) \cdot x^{j-2} \cdot u(x,t) \, \mathrm{d}x \\ &= \underbrace{j \cdot (j-1) \cdot s_{j-2}(t).} \end{aligned}$$

These are the recursive relations we already encountered with the heat equation on \mathbb{R} , see [5, 6]. Therefore, for *odd* polynomials $p \in \mathbb{R}[x]$ we have

$$(T_t p)(x) = \int_0^\infty p(y) \cdot k_t(x, y) \, \mathrm{d}y = (e^{t\partial_x^2} p)(x) \in \mathbb{R}[x]$$

with $T_t x^{2d+1} = \mathfrak{p}_{2d+1}(x,t)$ for all $d \in \mathbb{N}_0$ in [6, Dfn. 3.1] and if additionally $p \geq 0$ on $[0,\infty)$, then $T_t p \geq 0$ on $[0,\infty)$. On the other hand for *even* polynomials $p \in \mathbb{R}[x]$ we find that $T_t p \notin C^{\infty}([0,\infty)) \setminus \mathbb{R}[x]$ but at least $T_t p \geq 0$ on $[0,\infty)$. The reason for this strange behavior is of course the Dirichlet boundary condition and the resulting reflection principle. This effect needs further investigations.

This example shall also serve as a warning. Just because the operator (symbol) is ∂_x^2 does not mean that the positivity preservers are $e^{t\partial_x^2}$. The domain and the corresponding boundary conditions have to be taken into account as is done and is well-known in the partial differential equation literature and semi-group theory. With that in mind we recommend that the reader revisits Section 4 to see that the results there are non-trivial since these boundary effects are already included as long as it is a positivity preserver with constant coefficients.

6. Generators of $[0, \infty)$ -Positivity Preservers

We extend the results from Section 4 to $K \subsetneq \mathbb{R}^n$. They are based on the full description of K-positivity preservers with constant coefficients in Theorem 3.4.

Definition 6.1. Let $n \in \mathbb{N}$ and $K \subseteq \mathbb{R}^n$. We define the generators $\mathfrak{d}_+(K)$ of *K*-positivity preservers by

 $\mathfrak{d}_+(K) := \left\{ A \in \mathfrak{d}_+ \mid e^{tA} \text{ is a } K \text{-positivity preserver for all } t \ge 0 \right\}.$

Similar to Lemma 4.6 we get the following result.

Lemma 6.2. $A \in \mathfrak{d}_+(K)$ if and only if e^A has an infinitely divisible K^{\sharp} -representing measure.

Proof. Combine Theorem 3.4 with Lemma 4.6.

Corollary 6.3. $\mathfrak{d}_+(K) = \mathfrak{d}_+(K^{\sharp}).$

We want to describe all $[0, \infty)$ -positivity preservers.

Theorem 6.4. Let $K = K^{\sharp} = [0, \infty)$. The following are equivalent:

- (i) $A = \sum_{k \in \mathbb{N}_0} \frac{a_k}{k!} \cdot \partial^k \in \mathfrak{d}_+(K).$
- (ii) There exist $b \ge 0$, and a measure ν on $(0, \infty)$ such that

$$a_0 = 0,$$

$$a_1 = b + \int x \, \mathrm{d}\nu(x) < \infty,$$

and

$$a_k = \int_0^\infty x^k \, \mathrm{d}\nu(x) < \infty \quad \text{for all } k \in \mathbb{N} \text{ with } k \ge 2.$$

Proof. By Lemma 6.2 we have to describe all infinitely divisible measures μ on $K^{\sharp} = [0, \infty)$. These measures are fully characterized by the log-Laplace transformation [9, Satz 16.14]:

Let μ be a finite measure on $[0,\infty)$ and

$$u: [0,\infty) \to [0,1], t \mapsto -\log \int e^{-tx} d\mu(x)$$

be the log-Laplace transform of μ . μ is infinitely divisible if and only if there exists a $\alpha \geq 0$ and a σ -finite measure ν on $(0, \infty)$ with $\int \min(1, x) d\nu(x) < \infty$ such that

$$u(t) = \alpha t + \int (1 - e^{-tx}) \, \mathrm{d}\nu(x)$$

for all $t \geq 0$.

With this representation follow the proof of Theorem 4.8 and replace the Fourier transform with the Laplace transform. $\hfill\square$

Comparing Theorem 4.8 with Theorem 6.4 we get the following differences between $K = \mathbb{R}^n$ and $K = [0, \infty)$:

- (a) On $[0,\infty)$ there is only one positivity generator of finite order: $A = b\partial_x$ with $(e^{b\partial_x}p)(x) = p(x+b)$ for all $b \ge 0$.
- (b) On $[0,\infty)$ the support of ν has to be restricted to $[0,\infty)$.
- (c) On $[0,\infty)$ by the Lévy–Khinchin description only a singularity of order at most 1 at 0 is allowed for ν while order up to 2 was allowed on \mathbb{R}^n .

Both results coincide in the fact that if $a = (a_{\alpha})$ is a \mathbb{R}^{n} - resp. $[0, \infty)$ -moment sequence then $D(a) \in \mathfrak{d}_{+}$ resp. $\mathfrak{d}_{+}([0, \infty))$.

7. Open Problems

If

$$A = \sum_{\alpha \in \mathbb{N}_0^n} q_\alpha \cdot \partial^\alpha \quad \text{with} \quad \deg q_\alpha \le |\alpha| \tag{10}$$

then

$$A\mathbb{R}[x_1, \dots, x_n]_{\leq d} \subseteq \mathbb{R}[x_1, \dots, x_n]_{\leq d} \quad \text{for all} \quad d \in \mathbb{N}_0, \tag{11}$$

i.e., A preserves the degree. In fact for any linear $A : \mathbb{R}[x_1, \ldots, x_n] \to \mathbb{R}[x_1, \ldots, x_n]$ we have (10) \Leftrightarrow (11). (10) \Rightarrow (11) is clear. For (11) \Rightarrow (10) take in (10) the smallest α with respect to the lex-order such that deg $q_{\alpha} > |\alpha|$. Then deg $Ax^{\alpha} > |\alpha|$ is a contradiction to (11).

The partial differential equation

$$\partial_t p(x,t) = Ap(x,t)$$
 with $p(x,0) = p_0(x) \in \mathbb{R}[x_1,\ldots,x_n]$

is equivalent to a linear system of ordinary differential equations in the coefficients of p and hence has a unique solution $p(\cdot, t) = e^{tA}p_0$ for all $t \in \mathbb{R}$, i.e., e^{tA} is well-defined for any A as in (10) and $t \in \mathbb{R}$. If additionally A is a positivity preserver then e^{tA} is a positivity preserver with polynomial coefficients for all $t \ge 0$. In Theorem 4.8 we have already seen for the positivity preservers with constant coefficients that additional A appear. The restriction that e^{tA} for all $t \ge 0$ is degree preserving is a natural restriction since any degree jump is a discontinuity in e^{tA} in contradiction to the fact that e^{tA} is or needs to be continuous in t. So we arrive at the following open problem

Open Problem 1. What is the description of the set

$$\{A : \mathbb{R}[x_1, \dots, x_n] \to \mathbb{R}[x_1, \dots, x_n] \mid e^{tA} \text{ is a positivity preserver for all } t \ge 0\}$$

of all positivity generators, i.e., not necessarily with constant coefficients?

The constant coefficient part is completely described by Theorem 4.8. Part of the non-constant coefficient cases are covered by the previous case that A is a degree preserving positivity preserver. For the non-constant coefficient cases note that we are then in the framework of a non-commutative infinite dimensional Lie group and Lie algebra. Here the theory is much richer and far from being trivial.

Secondly, in Theorem 3.4 we completely described all K-positivity preservers with constant coefficients. A step in [2, 3] to go to the general case was to use the relation $(Tp)(x) = (T_xp)(x)$ with $T_x = \sum_{\alpha \in \mathbb{N}_0^n} q_\alpha(x) \partial^\alpha$, i.e., with that all \mathbb{R}^n positivity preservers are described. However, in Theorem 3.4 this is not possible. Let e.g. $K = B_1(0)$. Then since K is compact only $T = c \ge 0$ are positivity preservers with constant coefficients. However, rotation or inversion $x \mapsto -x$ are positivity preservers in this case but do not come from multiplication with a constant. Hence, the following open question remains.

Open Problem 2. What is a description of all K-positivity preserves, $K \subsetneq \mathbb{R}^n$, similar to Theorem 3.4 with non-constant coefficients?

Open Problem 1 and 2 lead to the following.

Open Problem 3. For $K \subsetneq \mathbb{R}^n$ what is the description of the set

 $\{A: \mathbb{R}[x_1,\ldots,x_n] \to \mathbb{R}[x_1,\ldots,x_n] \mid e^{tA} \text{ is a } K \text{-positivity preserver for all } t \ge 0\}$

of all K-positivity generators with non-constant coefficients?

Funding

The author and this project are supported by the Deutsche Forschungsgemeinschaft DFG with the grant DI-2780/2-1 and his research fellowship at the Zukunfskolleg of the University of Konstanz, funded as part of the Excellence Strategy of the German Federal and State Government.

References

- A. Guterman, B. Shapiro, On linear operators preserving the set of positive polynomials, J. Fixed Point Theory Appl. 3 (2008) 411–429.
- [2] J. Borcea, Classification of linear operators presevering elliptic, positive and non-negative polynomials, J. reine angew. Math. 650 (2011) 67–82.
- [3] T. Netzer, Representation and Approximation of Positivity Preservers, J. Geom. Anal. 20 (2010) 751–770.
- [4] V. I. Bogachev, Measure Theory, Springer-Verlag, Berlin, 2007.
- [5] R. Curto, P. J. di Dio, Time-dependent moments from the heat equation and a transport equation, Int. Math. Res. Notices (2022) https://doi.org/10.1093/imrn/rnac244.

- [6] R. Curto, P. J. di Dio, M. Korda, V. Magron, Time-dependent moments from partial differential equations and the time-dependent set of atoms, arXiv:2211.04416.
- [7] F. W. Warner, Foundation of Differential Manifolds and Lie Groups, Springer-Verlag, New York, Berlin, Heidelberg, 1983.
- [8] I. I. Hirschman, D. V. Widder, The Convolution Transform, Princeton University Press, Princeton, N.J., 1955.
- [9] A. Klenke, Wahrscheinlichkeitstheorie, Springer, Berlin, 2006.
- [10] K. Schmüdgen, The Moment Problem, Springer, New York, 2017.
- [11] E. B. Davis, Heat kernels and spectral theory, Cambridge University Press, Cambridge, UK, 1989.