# Dimension Independent Helly Theorem for Lines and Flats

Sutanoya Chakraborty<sup>∗</sup> Arijit Ghosh<sup>∗</sup> Soumi Nandi<sup>∗</sup>

#### Abstract

We give a generalization of *dimension independent* Helly Theorem of Adiprasito, Bárány, Mustafa, and Terpai (*Discrete & Computational Geometry* 2022) to higher dimensional transversal. We also prove some impossibility results that establish the tightness of our extension.

### 1 Introduction

Helly Theorem is a fundamental result in discrete and convex geometry. The theorem states that given any finite collection  $\mathcal F$  of convex sets in  $\mathbb R^d$ , if the sets in every subfamily of  $\mathcal F$  of size  $d+1$  have a point in common, then all the sets in the entire family  $\mathcal F$  have a point in common. Note that one can relax the finiteness of F can be relaxed if we assume that the convex sets in F are compact, therefore unless otherwise stated explicitly, we will not assume anything about the cardinality of a collection. Since its discovery [\[Hel23\]](#page-9-0), Helly Theorem has found multiple applications and generalizations [\[Eck93,](#page-9-1) [ADLS17,](#page-9-2) [DLGMM19,](#page-9-3) [BK22\]](#page-9-4).

We say a set  $T \subseteq \mathbb{R}^d$  pierces a family  $\mathcal F$  of subsets of  $\mathbb{R}^d$  if all sets in  $\mathcal F$ , has a non-empty intersection with the set  $T$ . A natural generalization of Helly Theorem would be to consider the problem of piercing a family of convex sets with a  $k$ -flats<sup>[1](#page-0-0)</sup>. Results of the above form are called Helly-type results. Generally, we want to show that there exists an integer  $h(k, d)$  such that if any  $h(k, d)$  convex sets from a collection  $\mathcal F$  can be pierced by a k-flat then the whole collection  $\mathcal F$  can be pierced by a k-flat. Unfortunately, Santaló  $\text{San40}$  showed the impossibility of getting such a result for even piercing a collection of convex sets by a line. Hadwiger [\[Had56\]](#page-9-6) showed that for a countable collection  $\mathcal F$  of disjoint convex sets in  $\mathbb R^d$  with non-empty interior and all congruent to a fixed compact convex set C, if every  $d+1$  sets from F can be pierced by a line then the whole family can be pierced by a line. Later, Danzer, Grünbaum and Klee [\[DGK63\]](#page-9-7) showed that the "congruent" assumption in Hadwiger's result [\[Had56\]](#page-9-6) can be weakened if the convex sets have bounded diameters. Aronov, Goodman, Pollack, and Wenger [\[AGPW00\]](#page-9-8) proved the first Helly-type result for hyperplanes about families of well-separated compact convex sets of arbitrary shapes in higher dimensions. Later, Aronov, Goodman, and Pollack [\[AGP02\]](#page-9-9) showed a Helly-type result for k-flat for families of convex bodies that are unbounded in k-independent directions.

Adiprasito, Bárány, Mustafa, and Terpai [\[ABMT20\]](#page-9-10) in a breakthrough paper proved the first dimension independent variant of the classical Helly Theorem:

**Theorem 1** (Adiprasito, Bárány, Mustafa, and Terpai [\[ABMT20\]](#page-9-10)). Let  $C_1, \ldots, C_n$  be convex sets in  $\mathbb{R}^d$  and  $r \in \{1, \ldots, n\}$ . For  $J \subseteq \{1, \ldots, n\}$ , let  $\mathcal{C}(J) := \bigcap_{j \in J} C_j$ . If  $B(b, 1)$  intersects  $\mathcal{C}(J)$  for

<span id="page-0-0"></span><sup>∗</sup> Indian Statistical Institute, India

<sup>&</sup>lt;sup>1</sup>By k-flat we mean affine subspace of  $\mathbb{R}^d$  of dimension k. Note that by lines and hyperplanes, we will mean 1-flats and  $(d-1)$ -flats respectively.

every  $J \subseteq \{1, \ldots, n\}$  with  $|J| = r$ , then there is a point  $q \in \mathbb{R}^d$  such that for all  $i \in \{1, \ldots, n\}$  we have

$$
d(q, C_i) \le \sqrt{\frac{n-r}{r(n-1)}}.
$$
\n(1)

Observe that for a fixed k and  $n \to \infty$ , the right-hand side of the above equation approaches  $\frac{1}{\sqrt{2}}$  $\frac{1}{r}$ .

Note that for all  $q \in \mathbb{R}^d$  and  $S \subseteq \mathbb{R}^d$ ,  $d(q, S)$  denotes the distance between the point q and the set S, that is,  $d(q, S) := \inf_{x \in S} ||q - x||$ . Also, note that  $B(x, R)$  denotes a closed Euclidean ball centered at  $x \in \mathbb{R}^d$  with radius R.

They also proved the following colorful variant of the above theorem.

<span id="page-1-1"></span>**Theorem 2** (Adiprasito, Bárány, Mustafa, and Terpai [\[ABMT20\]](#page-9-10)). Let  $\mathcal{F}_1, \ldots, \mathcal{F}_r$  be families of convex sets in  $\mathbb{R}^d$  with  $r \leq d$ , and  $b \in \mathbb{R}^d$ . Assume that for any r-tuple  $(C_1, \ldots, C_r)$  with  $C_i \in \mathcal{F}_i$ for all  $i \in \{1,\ldots,r\}$ , we have  $\left(\bigcap_{1 \leq i \leq r} C_i\right) \bigcap B(b,1) \neq \emptyset$ . Then there exists a point  $q \in \mathbb{R}^d$  and  $\exists i \in \{1, \ldots, r\}$  such that  $\forall C \in \mathcal{F}_i$  we have  $d(q, C) \leq \frac{1}{\sqrt{r}}$ .

In this short note, we will be generalizing the above results for k-flats, and also prove some impossibility results which will establish the optimality of our generalization.

Notations. Throughout the rest of the paper we will be using the following notations:

- Origin of  $\mathbb{R}^d$  will be denoted by  $\mathcal{O}$ .
- For all  $n \in \mathbb{N}$ ,  $[n]$  denotes the set  $\{1, \ldots, n\}$ .
- For any two points p and q in  $\mathbb{R}^d$ , the Euclidean distance between p and q will be denoted by  $||p - q||$ .
- |X| will denote the size of a set X.
- The diameter of the set  $C \subseteq \mathbb{R}^d$  will be denoted by diam(*C*).
- For all  $b \in \mathbb{R}^d$  and  $r > 0$ , closed and open balls centered at the point b and radius r will be denoted by  $B(b,r) := \{p \in \mathbb{R}^d : ||p-b|| \leq r\}$  and  $B^o(b,r) := \{p \in \mathbb{R}^d : ||p-b|| < r\}$ respectively.
- By  $\mathbb{S}^{d-1}$  we will denote the  $(d-1)$ -dimensional unit sphere centered at the origin.
- The *distance*  $d(S_1, S_2)$  between two subsets  $S_1$  and  $S_2$  of  $\mathbb{R}^d$  is defined as  $d(S_1, S_2) :=$  $\inf_{(p_1,p_2)\in S_1\times S_2} \|p_1-p_2\|.$
- Given points p and q in  $\mathbb{R}^d$ ,  $\overline{pq}$  denotes the closed line segment connecting p and q.

### <span id="page-1-0"></span>2 Our results

Before we can give the statements of our results we need to first introduce some definitions which will be required to state our results. We define the *central projection map*  $f : \mathbb{R}^d \to \mathbb{S}^{d-1}$ , where  $\mathbb{S}^{d-1}$  is the unit sphere centered at O, in the following way: for all  $x \in \mathbb{R}^d$ 

$$
f(x) := \frac{x}{\|x\|}.
$$

We say  $y \in \mathbb{S}^{d-1}$  is a limiting direction of the collection F if there exists a infinite sequence of points  ${s_n}_{n\in\mathbb{N}}$  and infinite sequence of sets  ${S_n}_{n\in\mathbb{N}}$  satisfying the following properties:

- for all  $n \in \mathbb{N}$ ,  $S_n \in \mathcal{F}$
- for all  $n \neq m$  in N,  $S_n \neq S_m$ .
- for all  $n \in \mathbb{N}$ ,  $s_n \in S_n$
- $\lim_{n \to \infty} ||s_n|| = \infty$  and  $\lim_{n \to \infty} f(s_n) = y$

For a collection F of subsets of  $\mathbb{R}^d$ , limiting direction set  $LDS(\mathcal{F})$  of F is defined as

$$
LDS(\mathcal{F}) := \left\{ y \in \mathbb{S}^{d-1} \, : \, y \text{ is a limiting direction of } \mathcal{F} \right\}.
$$

We will call the collection  $\mathcal{F}$  k-unbounded if the vector space spanned by set  $LDS(\mathcal{F})$  has dimension at least k. To give an idea of how k-unbounded framework will be used in a proof we will first give a colorful generalization of Helly Theorem for k-flats by Aronov, Goodman and Pollack [\[AGP02\]](#page-9-9).

<span id="page-2-2"></span>**Theorem 3** (Colorful Helly Theorem for k-flats). Suppose for each  $i \in [d+1], S_i$  is a k-unbounded collection of compact convex sets in  $\mathbb{R}^d$ , and there exists  $R > 0$  such that  $\forall C \in S_1 \cup \cdots \cup S_{d+1}$  we have diam(C)  $\lt R$ . Also, assume that for every  $(d+1)$ -tuple  $(C_1, \ldots, C_{d+1}) \in \mathcal{S}_1 \times \cdots \times \mathcal{S}_{d+1}$  there exists a k-flat H such that H intersects  $C_i$  for all  $i \in [d+1]$ . Then  $\exists j \in [d+1]$  and a k-flat H that intersects every sets in  $S_i$ .

The following theorem is the main technical result in this paper, and we will show that the rest of the results will be a direct consequence of this result.

<span id="page-2-0"></span>**Theorem 4** (Dimension independent colorful Helly Theorem for k-flats). Let  $\mathcal{F}_1, \ldots, \mathcal{F}_r$  be families of convex sets in  $\mathbb{R}^d$ ,  $k < r \leq d$ , and the family  $\mathcal{F} := \mathcal{F}_1 \cup \cdots \cup \mathcal{F}_r$  of convex sets satisfy the following properties:

- (i)  $\exists R > 0$  such that for all  $C \in \mathcal{F}$  we have  $\text{diam}(C) < R$
- (ii)  $\exists J_k = \{j_1, ..., j_k\} \subset [r]$  and  $\{y_{j_1}, ..., y_{j_k}\} \subset \mathbb{S}^{d-1}$  such that
	- for all  $i \in J_k$  we have  $y_i \in LDS(\mathcal{F}_i)$ , and
	- the collection of vectors  $\{y_{j_1}, \ldots, y_{j_k}\}$  are linearly independent

If for any r-tuple  $(C_1, \ldots, C_r) \in \mathcal{F}_1 \times \cdots \times \mathcal{F}_r$  there exists a k-flat that intersects the closed unit ball  $B(0,1)$  and every convex set  $C_i$  for all  $i \in [r]$ , then there exists a k-flat K and  $j \in [r]$  such that, for all  $C \in \mathcal{F}_j$ , we have

$$
d(C, K) \le \sqrt{\frac{1}{r - k}}.\tag{2}
$$

Observe that by substituting  $\mathcal{F}_1 = \mathcal{F}_2 = \cdots = \mathcal{F}_r = \mathcal{F}$  in Theorem [4](#page-2-0) we get the following dimension independent Helly Theorem for k-flats.

<span id="page-2-1"></span>**Theorem 5** (Dimension independent Helly Theorem for k-flats). Let  $\mathcal F$  be a k-unbounded family of convex sets in  $\mathbb{R}^d$  and there exists an  $R > 0$  such that  $\forall C \in \mathcal{F}$  we have have  $\text{diam}(C) < R$ . For  $r \in \mathbb{N}$  with  $k < r \leq d$  and  $b \in \mathbb{R}^d$ , if for every  $C_1, C_2, \ldots, C_r$  in F there exists a k-flat H that intersects  $B(0,1)$  and every  $C_i$  for all  $i \in \{1,2,\ldots,r\}$ , then there exists a k-flat K such that, for all  $C \in \mathcal{F}$ , we have

$$
d(C, K) \le \sqrt{\frac{1}{r-k}}.
$$

If each  $\mathcal{F}_i$  in Theorem [4](#page-2-0) is k-unbounded then we get the following colorful generalization of the above Theorem [5.](#page-2-1)

<span id="page-3-2"></span>**Theorem 6** (Colorful generalization of Theorem [5\)](#page-2-1). Let  $\mathcal{F}_1, \ldots, \mathcal{F}_r$  be k-unbounded families of convex sets in  $\mathbb{R}^d$  where  $k < r \leq d$ , and there exists  $R > 0$  such that  $\forall C \in \mathcal{F}_1 \cup \cdots \cup \mathcal{F}_r$  we have diam(C) < R. If for any r-tuple  $(C_1, ..., C_r) \in \mathcal{F}_1 \times \cdots \times \mathcal{F}_r$  there exists a k-flat that intersects the closed unit ball  $B(0,1)$  and every convex set  $C_i$  for all  $i \in \{1,\ldots,r\}$ , then there exists a k-flat K and  $j \in \{1, \ldots, r\}$  such that, for all  $C \in \mathcal{F}_j$ , we have

$$
d(C, K) \le \sqrt{\frac{1}{r - k}} \; .
$$

In the above theorems we require the convex sets to have bounded diameter. Note that this condition cannot be relaxed. Consider hyperplanes in  $\mathbb{R}^d$  and observe that any finite collection of hyperplanes can be pierced by a line passing through the origin  $\mathcal O$  in  $\mathbb R^d$ . But, for any k-flat K, with  $k \leq d-1$ , and  $\forall \Delta > 0$  there exists a hyperplane H such that  $d(K, H) > \Delta$ . The following two results will complement our results on dimension independent Helly Theorem by showing the tightness of the bound guaranteed by our results and also show that the k-unboundedness condition is unavoidable.

<span id="page-3-0"></span>**Theorem 7** (On families not being k-unbounded). There exists a family F of convex sets in  $\mathbb{R}^3$ such that

- there exists  $R > 0$  such that  $\text{diam}(C) < R$  for all  $C \in \mathcal{F}$ ,
- $F$  is 1-unbounded,
- any three convex sets in  $\mathcal F$  can be pierced by a plane (2-dimensional affine space) passing through the origin  $\mathcal{O}$ , and
- for any plane K in  $\mathbb{R}^3$  there exists a  $C_K \in \mathcal{F}$  such that  $d(K, C_K) > 1$ .

<span id="page-3-1"></span>**Theorem 8** (Tightness of the bound in Theorem [4\)](#page-2-0). There exist families  $\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3$  of convex sets in  $\mathbb{R}^3$  satisfying the following properties:

- $\forall C \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3$ , diam $(C) = \sqrt{2}$ ,
- both  $\mathcal{F}_1$  and  $\mathcal{F}_2$  are 1-unbounded,
- $\forall (C_1, C_2, C_3) \in \mathcal{F}_1 \times \mathcal{F}_2 \times \mathcal{F}_2$  there exists a line L that pierces  $C_1, C_2, C_3$ , and  $d(L, \mathcal{O}) \leq 1$ , and
- for every line K in  $\mathbb{R}^3$  there exists  $j \in [3]$  such that

$$
\max_{C \in \mathcal{F}_j} d(C, K) \ge \frac{1}{\sqrt{2}}.
$$

#### 3 Proofs of the claimed results

In this section we will give the proofs of the results claimed in Section [2.](#page-1-0) We will first begin by proving the following colorful generalization of the Helly Theorem for k-flats proved by Aronov, Goodman and Pollack [\[AGP02\]](#page-9-9).

**Theorem 9** (Restatement of Theorem [3\)](#page-2-2). Suppose for each  $i \in [d+1]$ ,  $S_i$  is a k-unbounded collection of compact convex sets in  $\mathbb{R}^d$ , and there exists  $R > 0$  such that  $\forall C \in \mathcal{S}_i$  we have  $\text{diam}(C) < R$ . Also, assume that for every  $(d+1)$ -tuple  $(C_1, \ldots, C_{d+1}) \in S_1 \times \cdots \times S_{d+1}$  there exists a k-flat H such that H intersects  $C_i$  for all  $i \in [d+1]$ . Then  $\exists j \in [d+1]$  and a k-flat H that intersects every sets in  $S_i$ .

*Proof.* Suppose for each  $i \in [d+1]$ ,  $\mathcal{L}_i$  is a set of k linearly independent vectors in the limiting directions set  $LDS(\mathcal{S}_i)$  of  $\mathcal{S}_i$ . Then there exists a linearly independent set of k vectors, say  $\mathcal{L} =$  $\{z_1,\ldots,z_k\}$ , such that for each  $i \in [k], z_i \in \mathcal{L}_i$ . Suppose K is the k-flat generated by the linear span of L. Now for each  $i \in [k]$ , since  $z_i \in \mathcal{L}_i$ , so  $\exists$  a sequence  $\{S_{i,n}\}_{n\in\mathbb{N}}$  in  $\mathcal{S}_i$  and for each  $n \in \mathbb{N}$ ,  $\exists x_{i,n} \in S_{i,n}$  such that the sequence  $\{f(x_{i,n})\}_{n \in \mathbb{N}}$  converges to  $z_i$ , as  $n \to \infty$ .

Now if  $\exists i \in [k]$  such that  $\mathcal{S}_i$  has a k-transversal then there is nothing to prove. Otherwise for each  $i \in [d+1], i > k$ , we take any  $B_i \in \mathcal{S}_i$  and take any  $n \in \mathbb{N}$ . Then  $B_{k+1}, \ldots, B_{d+1}$  together with  $S_{1,n},\ldots,S_{k,n}$ , as a colorful tuple, is pierceable by a k-flat. This follows that  $B_{k+1},\ldots,B_{d+1}$ can be pierced by a k-flat arbitrarily close to the direction of  $K$ . So by compactness of  $B_i$ 's we can say that,  $B_{k+1}, \ldots, B_{d+1}$  can be pierced by a k-flat in the direction of K, i.e, parallel to K.

Now for each  $i \in [d+1], i > k$ , suppose  $\mathcal{S}'_i$  is the projected family of  $\mathcal{S}_i$  on the  $(d-k)$  dimensional space  $\mathcal{K}^{\perp}$ , orthogonal to  $\mathcal{K}$ . Then every colorful  $(d-k+1)$  tuple from  $\mathcal{S}'_{k+1}, \ldots, \mathcal{S}'_{d+1}$  is pierceable by a point in the space  $\mathcal{K}^{\perp}$ . So by *Colorful Helly's Theorem*,  $\exists i \in [d+1], i > k$  such that  $\mathcal{S}'_i$  is pierceable by a point in  $\mathcal{K}^{\perp}$ . Hence there exists a k-flat parallel to K that hits all the members of the family  $S_i$ .  $\Box$ 

**Theorem 10** (Restatement of [4\)](#page-2-0). Let  $\mathcal{F}_1, \ldots, \mathcal{F}_r$  families of convex sets in  $\mathbb{R}^d$ ,  $k < r \leq d$ , and the family  $\mathcal{F} := \mathcal{F}_1 \cup \cdots \cup \mathcal{F}_r$  of convex sets satisfy the following properties:

- (i)  $\exists R > 0$  such that for all  $C \in \mathcal{F}$  we have  $\text{diam}(C) < R$
- (ii)  $\exists J_k = \{j_1, \ldots, j_k\}$  ⊂ [r] and  $\{y_{j_1}, \ldots, y_{j_k}\}$  ⊂  $\mathbb{S}^{d-1}$  such that
	- for all  $i \in J_k$  we have  $y_i \in LDS(\mathcal{F}_i)$ , and
	- the collection of vectors  $\{y_{j_1}, \ldots, y_{j_k}\}$  are linearly independent

If for any r-tuple  $(C_1, \ldots, C_r) \in \mathcal{F}_1 \times \cdots \times \mathcal{F}_r$  there exists a k-flat that intersects the closed unit ball  $B(0,1)$  and every convex set  $C_i$  for all  $i \in \{1,\ldots,r\}$ , then there exists a k-flat K and  $j \in \{1,\ldots,r\}$ such that, for all  $C \in \mathcal{F}_i$ , we have

$$
d(C, K) \le \sqrt{\frac{1}{r - k}}.
$$

*Proof.* Suppose for any set  $F \subset \mathbb{R}^d$ ,  $\overline{F}$  is the closure of F. Since for any point  $p \in \mathbb{R}^d$ ,  $d(p, F) =$  $d(p, \overline{F})$ , it is enough to show that  $\exists j \in [r]$  such that there exists a k-flat K satisfying  $\forall C \in \mathcal{F}_j$ ,

$$
d(\bar{C}, K) \le \sqrt{\frac{1}{r - k}}.
$$

Now without loss of generality, we assume that  $J_k = [k]$ . Then for each  $j \in J$ ,  $y_j \in \text{LDS}(\mathcal{F}_j)$ implies that  $\exists$  a sequence  $\{F_{j,n}\}_{n\in\mathbb{N}}$  in  $\mathcal{F}_j$  and for each  $n \in \mathbb{N}$ ,  $\exists x_{j,n} \in \mathcal{F}_{j,n}$  such that the sequence  ${f(x_{j,n})}_{{n\in\mathbb{N}}}$  converges to  $y_j$ .

Now if  $\exists j \in [r]$  such that there exists a k-flat K satisfying  $\forall C \in \mathcal{F}_j$ ,  $d(C, K) \leq \sqrt{\frac{1}{r-k}}$ , then there is nothing to prove. Otherwise, suppose K is the k-flat generated by the linear span of  $\{y_1, \ldots, y_k\}$ . Now for each  $i \in [r]$ ,  $i > k$ , take any  $F_i \in \mathcal{F}_i$ . Then for each  $n \in \mathbb{N}$ , there exists a k-flat  $K_n$ 

intersecting  $B(0,1)$  and piercing  $F_{1,n}, F_{2,n}, \ldots, F_{k,n}, F_{k+1}, \ldots, F_r$ . This implies that  $F_{k+1}, \ldots, F_r$ can be pierced by a k-flat arbitrarily close to the direction of K. Now by compactness of  $\bar{F}_i$ 's, we conclude that  $\exists a \in K^{\perp}$  such that the k-flat  $a + K$  pierces  $\bar{F}_{k+1}, \ldots, \bar{F}_r$ . Since  $a + K$  intersects  $B(O, 1)$ , we must have  $||a|| \leq 1$ .

Now let for any set A in  $\mathbb{R}^d$ ,  $\pi(A)$  denote the orthogonal projection of A onto the  $(d-k)$ dimensional space  $K^{\perp}$ . Then for any  $(C_{k+1}, \ldots C_r) \in \mathcal{F}_{k+1} \times \cdots \times \mathcal{F}_r$ , we have

$$
\left(\bigcap_{i=k+1}^r \pi(\bar{C}_i)\right)\bigcap B(0,1)\neq\emptyset.
$$

Then, by Theorem [2,](#page-1-1)  $\exists q \in K^{\perp}$  and  $\exists i \in \{k+1,\ldots,r\}$  such that  $\forall C \in \mathcal{F}_i$  we have

$$
d\left(q,\pi\left(\overline{C}\right)\right)<\sqrt{\frac{1}{r-k}}.
$$

Suppose  $q' \in \pi^{-1}(q)$  and consider the k-flat  $K' = q' + K$ . Then we have  $\forall C \in \mathcal{F}_i$ ,

$$
d\left(K',\overline{C}\right)<\sqrt{\frac{1}{r-k}}.
$$

 $\Box$ 

**Theorem 11** (On families not being k-unbounded: Restatement of Theorem [7\)](#page-3-0). There exists a family F of convex sets in  $\mathbb{R}^3$  such that

- there exists  $R > 0$  such that  $\text{diam}(C) < R$  for all  $C \in \mathcal{F}$ ,
- $F$  is 1-unbounded,
- any three convex sets in  $\mathcal F$  can be pierced by a plane (2-dimensional affine space) passing through the origin  $\mathcal{O}$ , and
- for any plane K in  $\mathbb{R}^3$  there exists a  $C_K \in \mathcal{F}$  such that  $d(K, C_K) > 1$ .

*Proof.* To establish the necessity of  $k$ -unboundedness we will be using a construction that is closely related to the one given by Aronov et al. [\[AGP02\]](#page-9-9).

Consider the eight shaded convex regions in Figure [1](#page-6-0) created by four circles and four squares centered at a point  $\mathcal{O}$ . Let  $x_3 = 0$  be the plane that contains the Figure [1,](#page-6-0) and without loss of generality assume that  $\mathcal O$  be the origin in  $\mathbb R^3$ . We will call these eight shaded convex regions  $a_1$ ,  $a_2, \ldots, a_8$ , respectively. Observe that any 3 of these convex sets can be intersected by a straight line passing  $\mathcal{O}$ .

We will now create additional convex sets in the following way: we choose the eight convex sets in a fixed order, and in each step elevate the sets in increasing heights in that order along the  $x_3$ -axis such that for any  $n \in \mathbb{N}$  there are infinitely many sets of this collection that lie outside  $B(\mathcal{O}, n)$ . This gives us a countably infinite sequence F of sets, where F is 1-unbounded, but not 2-unbounded. Clearly, for any three sets in  $\mathcal{F}$ , there exists a plane that passes through  $\mathcal{O}$  and intersects these sets.

Let  $\ell$  denote the length of the side of the smallest square in Figure [1.](#page-6-0) We show that it is not possible to find a plane that is at most a distance 1 unit away from all the sets in  $\mathcal F$  when  $\ell$  is large enough. Let K be a plane for which the maximum distance from the sets in  $\mathcal F$  is minimized, and

<span id="page-6-0"></span>

Figure 1: An example demonstrating the necessity of  $k$ -unboundedness condition. This figure has been taken from [\[AGP02\]](#page-9-9).

 $l_K$  be the intersection of K with the plane  $x_3 = 0$ . Clearly, K must be perpendicular to the plane that contains the first 8 sets, because otherwise for any  $R > 0$  we would find a set C in F for which  $d(K, C) > R$ . Consider the straight line  $l_K$  that is the intersection of K and the plane given by the equation  $x_3 = 0$ . If  $l_K$  is moved on the  $x_3 = 0$  plane closer to  $\mathcal O$  along the line perpendicular to  $l_K$  from  $\mathcal{O}$ , the quantity max  $\{d(a_{2n}, l_K), d(a_{2n-1}, l_K)\}\)$  does not increase for  $n \in \{1, 2, 3, 4\}$ . Since  $d(K, a_i) = d(l_K, a_i)$   $\forall i \in [8]$ , we can take K to be passing through O. Let the side-lengths of the 4 squares in Figure [1](#page-6-0) be  $\ell_1 (= \ell), \ell_3, \ell_5, \ell_7$ , where the side of side-length  $\ell_i$  is shared by the set  $a_i$ . Let the diagonals of the largest square in Figure [1](#page-6-0) lie on the  $x_1$ -axis and the  $x_2$ -axis respectively. Then, if  $l_K$  makes an angle  $\theta \in [0, \pi)$  with the x<sub>1</sub>-axis, then we have the following: for  $i \in \{1, 3, 5, 7\}$  we have

$$
d(a_i, l_K) = d(a_{i+1}, l_k),
$$
\n(3)

and

$$
d(a_1, l_K) = \begin{cases} \frac{\ell_1}{\sqrt{2}} \sin(\pi/4 - \theta) & \text{if } 0 \le \theta \le \pi/4\\ \frac{\ell_1}{\sqrt{2}} \sin(\pi - \theta) & \text{if } 3\pi/4 \le \theta \le \pi\\ 0 & \text{otherwise} \end{cases}
$$
(4)

$$
d(a_3, l_K) = \begin{cases} \frac{\ell_3}{\sqrt{2}} \sin \theta & \text{if } 0 \le \theta \le \pi/4\\ \frac{\ell_3}{\sqrt{2}} \sin (\pi/2 - \theta) & \text{if } \pi/4 \le \theta \le \pi/2\\ 0 & \text{otherwise} \end{cases}
$$
(5)

$$
d(a_5, l_K) = \begin{cases} \frac{\ell_5}{\sqrt{2}} \sin(\theta - \pi/4) & \text{if } \pi/4 \le \theta \le \pi/2\\ \frac{\ell_5}{\sqrt{2}} \sin(3\pi/4 - \theta) & \text{if } \pi/2 \le \theta \le 3\pi/4\\ 0 & \text{otherwise} \end{cases} \tag{6}
$$

$$
d(a_7, l_K) = \begin{cases} \frac{\ell_7}{\sqrt{2}} \sin (\theta - \pi/2) & \text{if } \pi/2 \le \theta \le 3\pi/4\\ \frac{\ell_7}{\sqrt{2}} \sin (\pi - \theta) & \text{if } 3\pi/4 \le \theta \le \pi\\ 0 & \text{otherwise} \end{cases}
$$
(7)

<span id="page-7-0"></span>

Figure 2: An example demonstrating the tightness of the bound given in Theorem [8.](#page-3-1)

Now, observe that

$$
\sum_{i=1}^{8} d(a_i, l_K) \ge \frac{\ell}{\sqrt{2}} \times \min_{\theta \in [0, \pi/4]} (\sin \theta + \sin (\pi/4 - \theta))
$$

$$
\ge \ell \sqrt{2} \sin (\pi/8)
$$

Thus, for  $\ell > \sqrt{2}/\sin(\pi/8)$ , there are no planes that are at most 1 distance away from each set in  ${\mathcal F}.$  $\Box$ 

Theorem 12 (Tightness of the bound in Theorem [4:](#page-2-0) Restatement of Theorem [8\)](#page-3-1). There exist families  $\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3$  of convex sets in  $\mathbb{R}^3$  satisfying the following properties:

- $\forall C \in \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3$ , diam $(C) = \sqrt{2}$ ,
- both  $\mathcal{F}_1$  and  $\mathcal{F}_2$  are 1-unbounded,
- $\forall (C_1, C_2, C_3) \in \mathcal{F}_1 \times \mathcal{F}_2 \times \mathcal{F}_2$  there exists a line L that pierces  $C_1, C_2, C_3$ , and  $d(L, \mathcal{O}) \leq 1$ and
- for every line K in  $\mathbb{R}^3$  there exists  $j \in [3]$  such that  $\max_{C \in \mathcal{F}_j} d(C, K) \geq \frac{1}{\sqrt{2}}$  $\frac{1}{2}$ .

Proof. We provide an example where the bound given in Theorem [6](#page-3-2) is tight.

Let  $\{A, B, C, D, E, F, G, H\}$  be the 8 vertices cube in  $\mathbb{R}^3$  whose centroid is the origin  $\mathcal O$  and Let  $\{A, B, C, D, E, F, G, H\}$  be the 8 vertices cube in  $\mathbb{R}^2$  whose centroid is the origin  $C$  and side length is  $\sqrt{2}$  and  $EFGH$  is parallel to the plane  $x_3 = 0$  (see Figure [2\)](#page-7-0). Define  $\mathcal{F}_i :=$  $\{S_{i,1}, S_{i,2}, S_{i,3}, ...\}$  for  $i = 1, 2, 3$  in the following way: Set  $S_{1,1} = \overline{AB}, S_{1,2} = \overline{CD}$  and for  $n > 2$ ,  $S_{1,n}$  is the line segment  $\overline{CD}$  raised to the height  $x_3 = n$ . Similarly, set  $S_{2,1} = \overline{GH}$ ,

 $S_{1,2} = \overline{EF}$  and for  $n > 2$ ,  $S_{2,n}$  is the line segment  $\overline{EF}$  lowered to the height  $x_3 = -n$ . Now set  $S_{3,1} = \overline{BC}, S_{3,2} = \overline{DA}, S_{3,3} = \overline{FG}, S_{3,4} = \overline{HE}$ , and let  $S_{3,4n+j}$  be the set  $S_{3,j}$ , for all  $j \in \{0,1,2,3\}$ (see Figure [2\)](#page-7-0). Clearly, any colorful 3-tuple  $(C_1, C_2, C_3)$ ,  $C_i \in \mathcal{F}_i$ ,  $i \in [3]$ , can be hit by a straight line that is at most at a distance 1 away from the centroid. Let  $\mathcal L$  denote the set of all straight line transversals l of colorful 3-tuples such that l is as close to  $\mathcal O$  as possible. Let  $l_n$  denote the straight line transversal in L that passes through  $S_{1,n}$ ,  $S_{2,1}$ ,  $S_{3,1}$ . Clearly,  $d(l_n, \mathcal{O}) \to 1$  as  $n \to \infty$ . Then

$$
\sup_{l\in\mathcal{L}}d(\mathcal{O},l)=1.
$$

Now note that for the straight line L that is perpendicular to the plane on which ABCD lies and passes through  $\mathcal{O}$ , we have

$$
\inf_{i \in [3]} \sup_{C \in \mathcal{F}_i} d(l, C) \ge \inf_{i \in [3]} \sup_{C \in \mathcal{F}_i} d(L, C),
$$

for any straight line l in  $\mathbb{R}^3$ . This we can show in the following way: let  $l_1$  be a straight line such that

$$
\sup_{C \in \mathcal{F}_1} d(l_1, C) = \inf_{l \in \mathcal{L}} \sup_{C \in \mathcal{F}_1} d(C, l).
$$

Then  $l_1$  must be perpendicular to the plane on which  $ABCD$  lies, otherwise, the supremum of its distances from sets in  $\mathcal{F}_1$  would be infinity. Then,  $l_1$  must be equidistant from both  $S_{1,1}$  and  $S_{1,2}$ , and therefore, we can take  $l_1$  to be L. Similar arguments show that

$$
\sup_{C \in \mathcal{F}_2} d(L, C) = \inf_{l \in \mathcal{L}} \sup_{C \in \mathcal{F}_1} d(C, l).
$$

We have

$$
\inf_{i \in [3]} \sup_{C \in \mathcal{F}_i} d(L, C) = \frac{1}{\sqrt{2}}.
$$

To see that

$$
\inf_{l \in \mathcal{L}} \sup_{C \in \mathcal{F}_3} d(C, l) = \frac{1}{\sqrt{2}},
$$

project  $\overline{BC}$ ,  $\overline{AD}$ ,  $\overline{FG}$ ,  $\overline{EH}$  onto the plane P that contains ABEF. If there is a straight line  $l_3$ such that

$$
\sup_{C \in \mathcal{F}_3} d(l_3, C) = \inf_{l \in \mathcal{L}} \sup_{C \in \mathcal{F}_3} d(C, l),
$$

then let the projection of  $l_3$  onto P be  $l'_3$ . If L is not the straight line that minimizes

$$
\inf_{l\in\mathcal{L}}\sup_{C\in\mathcal{F}_6}d(C,l),
$$

then the perpendicular distance from  $l'_3$  to  $A, B, E$  and F must be smaller than  $\frac{1}{\sqrt{2}}$  $\frac{1}{2}$ . Let, without loss of generality,  $A$  be the point from which  $l'_3$  is the farthest. Then, we must have another point among  $B, E$ , and F from which  $l'_3$  has the same distance as A. This point then must be E, because otherwise, we could have taken  $\tilde{L}$  to be  $l_3$ . This means that  $l'_3$  passes through the centroid of the square ABEF and two points from  $A, B, E$ , and F lie on each side of  $l'_3$ . But this implies that  $l'_3$ must be parallel to  $AB$  since  $l'_3$  has the minimum distance from both  $\overline{A}$  and  $\overline{B}$  and is at least as close to  $E$  and  $F$ , which is a contradiction. We have

$$
\inf_{i \in [3]} \sup_{C \in \mathcal{F}_i} d(L, C) = \frac{1}{\sqrt{2}},
$$

which is what we get by plugging in the values of r and k in the inequality given in Theorem [6.](#page-3-2)  $\Box$ 

## References

- <span id="page-9-10"></span>[ABMT20] K. A. Adiprasito, I. Bárány, N. H. Mustafa, and T. Terpai. Theorems of Carathéodory, Helly, and Tverberg Without Dimension. Discrete  $\mathcal{B}$  Computational Geometry,  $64(2):233 - 258, 2020.$
- <span id="page-9-2"></span>[ADLS17] N. Amenta, J. A. De Loera, and P. Soberón. Helly's Theorem: New Variations and Applications. Algebraic and Geometric Methods in Discrete Mathematics, 685:55–95, 2017.
- <span id="page-9-9"></span>[AGP02] B. Aronov, J. E. Goodman, and R. Pollack. A Helly-type Theorem for Higherdimensional Transversals. Computational Geometry: Theory and Applications, 21(3):177–183, 2002.
- <span id="page-9-8"></span>[AGPW00] B. Aronov, J. E. Goodman, R. Pollack, and R. Wenger. A Helly-Type Theorem for Hyperplane Transversals to Well-Separated Convex Sets. In Proceedings of the 16th Annual Symposium on Computational Geometry, SoCG, page 57–63, 2000.
- <span id="page-9-4"></span>[BK22] I. Bárány and G. Kalai. Helly-type Problems. *Bulletin of the American Mathematical* Society, 59(4):471–502, 2022.
- <span id="page-9-7"></span>[DGK63] L. Danzer, B. Grünbaum, and V. Klee. *Helly's Theorem and Its Relatives*. Proceedings of Symposia in Pure Mathematics: Convexity. American Mathematical Society, 1963.
- <span id="page-9-3"></span>[DLGMM19] J. De Loera, X. Goaoc, F. Meunier, and N. Mustafa. The discrete yet ubiquitous theorems of Carathéodory, Helly, Sperner, Tucker, and Tverberg. Bulletin of the American Mathematical Society, 56(3):415–511, 2019.
- <span id="page-9-1"></span>[Eck93] J. Eckhoff. Helly, Radon, and Carathéodory Type Theorems. In Handbook of Convex Geometry, pages 389 – 448. Elsevier, 1993.
- <span id="page-9-6"></span>[Had56] H. Hadwiger. Über einen Satz Hellyscher Art. Arch. Math., 7:377–379, 1956.
- <span id="page-9-0"></span>[Hel23] E. Helly. Uber Mengen konvexer Körper mit gemeinschaftlichen Punkten. *Jahres*bericht der Deutschen Mathematiker-Vereinigung, 32:175 – 176, 1923.
- <span id="page-9-5"></span>[San40] L Santaló. Un teorema sobre conjuntos de paralelepipedos de aristas paralelas. Publ. Inst. Mat. Univ. Nac. Litoral, 2:49–60, 1940.