

# Quasiclassical integrability condition in AKNS scheme

A. M. Kamchatnov, D. V. Shaykin

*Institute of Spectroscopy, Russian Academy of Sciences, Troitsk, Moscow, 142190, Russia*

*Moscow Institute of Physics and Technology, Institutsky lane 9, Dolgoprudny, Moscow region, 141700, Russia*

*Skolkovo Institute of Science and Technology, Skolkovo, Moscow, 143026, Russia*

---

## Abstract

In this paper, we study the condition of quasiclassical integrability of soliton equations. This condition states that the Hamiltonian structure of equations, which govern propagation of high-frequency wave packets, is preserved by the dispersionless flow independently of initial conditions. If this condition is fulfilled, then the carrier wave number of any packet is a certain function of local values of the dispersionless variables pertained to the soliton equations under consideration. We show for several examples that this function together with the dispersion relation for linear harmonic waves determine the quasiclassical limit of the Lax pair functions in the scalar representation of the Ablowitz-Kaup-Newell-Segur scheme.

*Keywords:* integrable nonlinear wave equations, quasiclassical approximation, AKNS scheme

*PACS:* 02.30.Ik, 05.45.Yv, 43.20.Bi

---

## 1. Introduction

The general definition of the “integrability” property of nonlinear wave equations is apparently impossible (see, e.g., discussion in Ref. [1]). In a narrow sense, integrability is often related with existence of the Lax pair of linear systems whose compatibility condition leads to the nonlinear wave equations under consideration. First discovered for the Korteweg-de Vries (KdV) equation [2, 3] and nonlinear Schrödinger (NLS) equation [4], Lax pairs were found for a number of wave equations and they were widely used in various physical applications (see, e.g., Refs. [5, 6, 7] and reference therein). However, the relationship between the physical properties of wave equations and associated with them Lax pairs still remains unclear.

As was noticed in Ref. [8] in the theory of the generalized NLS equation

$$i\psi_t + \frac{1}{2}\psi_{xx} - f(|\psi|^2)\psi = 0, \quad (1)$$

the integrable case with  $f(|\psi|^2) = |\psi|^2$  is distinguished by a very special property of propagation of high-frequency wave packets along large scale background waves, and this property can be formulated in purely physical characteristics of Eq. (1). In the problem of packets propagation we have two very different characteristic length parameters: a small wavelength  $\sim k^{-1}$  of harmonics that compose the packet ( $k$  is the carrier wave number) and the size  $\sim l$  of the background wave, so  $(kl)^{-1}$  is a small parameter of the theory. Its existence allows one to separate dispersionless evolution of a background wave from propagation of a linear wave packet. The dispersionless evolution is described by the hydrodynamic equations

$$\rho_t + (\rho u)_x = 0, \quad u_t + uu_x + \frac{c^2}{\rho}\rho_x = 0, \quad (2)$$

where  $\rho$  and  $u$  are defined according to the formula

$$\psi(x, t) = \sqrt{\rho(x, t)} \exp\left(i \int^x u(x', t) dx'\right) \quad (3)$$

and

$$c^2 = \rho f'(\rho). \quad (4)$$

The small amplitude wave packet propagates along smooth distributions  $\rho = \rho(x, t)$ ,  $u = u(x, t)$ , given

---

*Email addresses:* kamch@isan.troitsk.ru  
(A. M. Kamchatnov), shaykin.dv@phystech.edu  
(D. V. Shaykin)

by some specific solution of Eqs. (2), according to the Bogoliubov dispersion relation

$$\omega = k \left( u \pm \sqrt{c^2 + \frac{k^2}{4}} \right), \quad (5)$$

where due to the condition  $kl \gg 1$  one can consider  $u$  and  $c = c(\rho)$  constant within the packet's width. This means that the packet is considered as a point-like particle with coordinate  $x = x(t)$  whose motion obeys the Hamilton equations (see, e.g., [9, 10])

$$\frac{dx}{dt} = \frac{\partial \omega}{\partial k}, \quad \frac{dk}{dt} = -\frac{\partial \omega}{\partial x}. \quad (6)$$

As we showed in Ref. [8], the system (2), (6) admits the solution in the form  $k = k(\rho, u)$ , where the wave number depends on  $(x, t)$ -coordinates via the local values  $\rho$  and  $u$  of the background wave, only in case of the integrable NLS equation with  $f(\rho) = \rho$ . In this case we obtain

$$k^2 = (q - u)^2 - 4\rho, \quad (7)$$

where  $q$  is an integration constant. As was shown in Ref. [11], the expression in the right-hand side of this formula is related with the quasiclassical limit of the Zakharov-Shabat spectral problem [4] ( $q$  plays the role of the spectral parameter), whereas the quasiclassical limit of the other equation of the corresponding Lax pair reduces to the number of waves conservation law (see [12, 13])

$$k_t + \omega_x = 0. \quad (8)$$

Thus, both equations of the Lax pair in their quasiclassical limit for the NLS equation turn out to be related with dispersion relation  $\omega = \omega(k)$  of linear waves and nonlinearity function  $f(\rho) = \rho$  which appears in the last term of the right-hand side of Eq. (7). Substitution of this formula into the first Hamilton equation (6) yields very simple equation for propagation of wave packets,

$$\frac{dx}{dt} = q - \frac{2\rho(x, t)}{q - u(x, t)}, \quad (9)$$

where  $q$  is determined by the initial condition  $k(x = x_0, t = t_0) = k_0$  for Eq. (7).

In this paper, we formulate the observation of Ref. [11] in more general terms and confirm validity of such a general formulation by some other examples of integrable equations which belong to the Ablowitz-Kaup-Newell-Segur (AKNS) scheme [14].

We derive for these examples the relations similar to Eq. (7), demonstrate their connection with the quasiclassical limit of the corresponding Lax pairs, and obtain equations of motion of high-frequency wave packets along non-uniform and time dependent background waves of the dispersionless limit.

## 2. General theory

We assume that a physical system is characterized by two variables which in many situations have physical sense of “density”  $\rho(x, t)$  and “flow velocity”  $u(x, t)$ . The two natural limits can be distinguished in dynamics of such systems. First, we can consider infinitely small amplitude waves propagating along uniform stationary background  $\rho = \rho_0$ ,  $u = u_0$ . Then we obtain the dispersion relation for harmonic waves  $\rho - \rho_0, u - u_0 \propto \exp[i(kx - \omega t)]$ ,

$$\omega = \omega(k, \rho_0, u_0), \quad (10)$$

(see, e.g., Eq. (5)). This dispersion relation often includes some parameter  $k_D$ , so that Eq. (10) becomes linear for  $k \ll k_D$ ,

$$\omega \approx (u_0 \pm c_0)k, \quad k \ll k_D, \quad (11)$$

where  $c_0$  is called “sound velocity” and Eq. (11) corresponds to long sound waves propagating upstream or downstream the flow moving with velocity  $u_0$ .

As another typical limit, we can consider long waves that change little on distances about  $\sim k_D^{-1}$  (i.e.,  $|\partial\rho/\partial x| \sim \rho/l$ ,  $|\partial u/\partial x| \sim u/l$ , and  $lk_D \gg 1$ ). In this case we arrive at the so-called “dispersionless” or “hydrodynamic” limit of our equations (see, e.g., Eqs. (2)). Generally speaking, in case of two wave variables the hydrodynamic equations can be transformed to the diagonal form

$$\begin{aligned} \frac{\partial r_+}{\partial t} + v_+(r_+, r_-) \frac{\partial r_+}{\partial x} &= 0, \\ \frac{\partial r_-}{\partial t} + v_-(r_+, r_-) \frac{\partial r_-}{\partial x} &= 0 \end{aligned} \quad (12)$$

for “Riemann invariants”  $r_{\pm} = r_{\pm}(\rho, u)$ , where the characteristic velocities are given by the formulas

$$v_{\pm} = u \pm c \quad (13)$$

for sound waves propagation in the long wavelength limit (11) and they can also be expressed as functions of the Riemann invariants,  $v_{\pm} = v_{\pm}(r_+, r_-)$ .

Now, if we consider a high frequency wave packet with carrier wave number  $k$  and width  $d$  such that

$$2\pi/k \ll d \ll l, \quad (14)$$

then we can define the packet's coordinate  $x(t)$  with good enough accuracy. As is well known, such a packet propagates with the group velocity and its local wavelength changes due to refraction (dependence of the phase velocity  $\omega/k$  on the coordinate  $x$ ), so the packet's dynamics obeys the Hamilton equations (6) with

$$\omega = \omega(k, r_+, r_-). \quad (15)$$

As was shown in Ref. [8], if we demand that the Hamiltonian form of Eqs. (6) is preserved by the hydrodynamic flow (12), then the carrier wave number  $k$  must only depend on the local values of the dispersionless variables  $r_+, r_-$ ,

$$k = k(r_+, r_-, q), \quad (16)$$

which satisfied the equations

$$\frac{\partial k}{\partial r_+} = \frac{\partial \omega / \partial r_+}{v_+ - \partial \omega / \partial k}, \quad \frac{\partial k}{\partial r_-} = \frac{\partial \omega / \partial r_-}{v_- - \partial \omega / \partial k}, \quad (17)$$

and  $q$  is an integration constant. Naturally, solution of these equations does only exist if

$$\frac{\partial}{\partial r_+} \left( \frac{\partial k}{\partial r_-} \right) = \frac{\partial}{\partial r_-} \left( \frac{\partial k}{\partial r_+} \right). \quad (18)$$

Usually this condition is only fulfilled in the limit of large  $k$ , so then one can obtain an approximate solution which is enough for discussion of propagation of high frequency wave packets. However, as was noticed in Ref. [8], Eq. (18) is satisfied identically for the integrable case of Eq. (1) with  $f(\rho) = \rho$ , so that the exact solution of Eqs. (17) can be found (see Eq. (7) with  $\rho$  and  $u$  expressed in terms of the Riemann invariants). We will show below that the same statement turns out to be true for other examples of integrable equations in AKNS scheme.

The above theory refers to any wave packet. Let us consider now a particular case of wave packets corresponding to the small amplitude edges of dispersive shock waves propagating along smooth parts of large wave pulses. As was noticed in Ref. [15], oscillations enter into the dispersive shock region with velocity equal to difference between the group and phase velocities of this small-amplitude edge wave packet. In situations when these oscillations transform eventually into separate solitons,

this remark allows one to write the formula for the number of solitons as (see Refs. [16, 17])

$$N = \frac{1}{2\pi} \int_0^\infty \left( k \frac{\partial \omega}{\partial k} - \omega \right) dt = \frac{S}{2\pi}, \quad (19)$$

where  $S$  is the action produced by a point-like particle moving according to the Hamilton equations (6) with Hamiltonian  $\omega$ . The integral here is taken over the whole packet's path  $x = x(t), k = k(t)$  from the wave breaking moment  $t = 0$  till large enough time  $t \rightarrow \infty$  of full transformation of the pulse into solitons. We can also take the path back with  $k$  replaced by  $-k$ , so the direct and inverse paths are combined to a closed contour  $C$  in the phase space  $(x, k)$  and Eq. (19) can be written in the form of the Poincaré-Cartan integral invariant (see Ref. [11] and references therein)

$$N = \frac{1}{\pi} \oint (k \delta x - \omega \delta t), \quad (20)$$

where  $\delta x = (\partial \omega / \partial k) \delta t$ . As was shown in Ref. [11], the Poincaré-Cartan integral invariant is preserved by the hydrodynamic flow (12) for  $k$  given by solution (16) of Eqs. (17). Consequently, we can transform the contour  $C$  to the initial state  $t = 0$  and then Eq. (20) gives the expression

$$N = \frac{1}{2\pi} \int_{-\infty}^\infty k(r_+^0(x), r_-^0(x)) dx, \quad (21)$$

where  $r_\pm^0(x)$  are the initial distributions of the dispersionless Riemann invariants. For a particular case of simple wave distributions, when one of the Riemann invariants is constant, such a formula was obtained by a different method in Refs. [18, 19, 20] and by direct calculation of the integral in Eq. (19) in Refs. [17, 21, 22].

Now we will relate Eq. (21) with the Lax pair in AKNS scheme. It is convenient to discuss its quasiclassical limit in scalar representation [23], when the integrable equation under consideration is expressed as a compatibility condition of linear equations

$$\phi_{xx} = \mathcal{A}\phi, \quad \phi_t = -\frac{1}{2}\mathcal{B}_x\phi + \mathcal{B}\phi_x. \quad (22)$$

We will give concrete examples of this scheme in the next Section and here we will consider some its general properties. To this end, we notice that the second order differential equation (22) has two basis solutions  $\phi_+, \phi_-$ , so we define the 'squared basis function',

$$g = \phi_+ \phi_-, \quad (23)$$

which satisfies the equations

$$g_{xxx} - 2\mathcal{A}_x g - 4\mathcal{A}g_x = 0, \quad g_t = \mathcal{B}g_x - \mathcal{B}_x g. \quad (24)$$

The first equation can be integrated to give

$$\frac{1}{2}gg_{xx} - \frac{1}{4}g_x^2 - \mathcal{A}g^2 = P, \quad (25)$$

where  $P$  is an integration constant. Here  $\mathcal{A}$  depends, besides the wave variables, on the spectral parameter  $\lambda$  and we normalize  $\mathcal{A}$  in such a way that

$$\mathcal{A} \rightarrow -\sigma\lambda^r \quad \text{as} \quad \lambda \rightarrow \infty, \quad (26)$$

and  $\sigma = \pm 1$  denotes the sign of this leading term in this series expansion of  $\mathcal{A}$ ,  $r$  being the exponent of the leading term. Then in case of polynomial dependence of  $g$  on  $\lambda$  we get

$$\frac{\sqrt{P/\lambda^r}}{g} \rightarrow 1 \quad \text{as} \quad \lambda \rightarrow \infty. \quad (27)$$

The second equation (24) yields the generating function of conservation laws

$$\left(\frac{\sqrt{P}}{g}\right)_t - \left(\frac{\sqrt{P}}{g}\mathcal{B}\right)_x = 0, \quad (28)$$

where inessential factor  $\lambda^{-r/2}$  is dropped out and the expressions  $\sqrt{P}/g$  and  $\sqrt{P}\mathcal{B}/g$  can be expanded in series in powers of  $\lambda^{-1}$  to give a series of particular conservation laws. This makes Eq. (28) a convenient tool for derivation of the Whitham modulation equations [24, 25].

Solutions of Eqs. (22) can also be expressed in terms of  $g$  [26]

$$\phi_{\pm} = \sqrt{g} \exp\left(\pm i \int^x \frac{\sqrt{P}}{g} dx\right). \quad (29)$$

The Whitham averaging method in Krichever's formulation [27] assumes that Eq. (29) remains correct during the whole process of transformation of the initially smooth pulse to a train of separate solitons. At the initial moment of time the wave variables are smooth functions of  $x$  and they can be approximated by the dispersionless Riemann invariants  $r_{\pm}^0$ . Then the function  $g$  is also smooth and the terms with its derivatives in Eq. (25) can be neglected, so we arrive at the approximate expression

$$\frac{\sqrt{P}}{g} \approx \sqrt{-\sigma\mathcal{A}} \equiv \bar{k}(r_+, r_-, \lambda), \quad (30)$$

where  $\bar{\mathcal{A}} = \bar{\mathcal{A}}(r_+, r_-, \lambda)$  is to be obtained from  $\mathcal{A}$  in the same approximation with omitted derivatives of  $r_+^0$  and  $r_-^0$  (or  $\rho$  and  $u$ ). Consequently, we obtain a quasiclassical limit of Eq. (29),

$$\phi_{\pm} \approx \sqrt{g} \exp\left(\pm i \int^x \bar{k}(r_+, r_-, \lambda) dx\right). \quad (31)$$

The condition that  $\phi_{\pm}$  are single-valued functions of  $x$  gives the Bohr-Sommerfeld quantization rule (see, e.g., [28, 29, 30, 31]) and, in particular, the formula for the number of solitons

$$\int_{-\infty}^{\infty} \bar{k}(r_+(x), r_-(x), \lambda_n) dx = \pi N, \quad (32)$$

where  $N$  is the largest quantum number equal to the number of solitons and we replaced the corresponding turning points  $x_1^N, x_2^N$  by  $\mp\infty$ , since contribution of the intervals  $-\infty < x < x_1^N, x_2^N < x < \infty$  is supposed to be negligibly small in this quasiclassical approximation. Comparison of Eqs. (21) and (32) gives the expression

$$k(r_+, r_-, q) = 2\bar{k}(r_+, r_-, \lambda), \quad (33)$$

which defines the relationship

$$\lambda = \lambda(q) \quad (34)$$

between the integration constant  $q$  in (16) and the spectral parameter  $\lambda$  in the Lax pair (22).

Substitution of  $\sqrt{P}/g = k/2$  into Eq. (28) gives

$$k_t - (k\bar{\mathcal{B}})_x = 0, \quad (35)$$

where  $\bar{\mathcal{B}}(r_+, r_-, \lambda)$  is obtained from  $\mathcal{B}$  in the same quasiclassical approximation with neglected  $x$ -derivatives of the Riemann invariants. This equation must coincide with Eq. (8), so we get

$$\frac{\omega}{k} = -\bar{\mathcal{B}}. \quad (36)$$

Thus, we arrive at the following expressions for the quasiclassical limits of  $\mathcal{A}$  and  $\mathcal{B}$ :

$$\begin{aligned} \bar{\mathcal{A}}(r_+, r_-, \lambda(q)) &= -\frac{1}{4}k^2(r_+, r_-, q), \\ \bar{\mathcal{B}}(r_+, r_-, \lambda(q)) &= -\frac{\omega(k(r_+, r_-, q), r_+, r_-)}{k(r_+, r_-, q)}, \end{aligned} \quad (37)$$

where  $k^2$  is defined by solution of Eqs. (17). The condition (18) of existence of such a solution becomes the condition of existence of the functions  $\bar{\mathcal{A}}$  and  $\bar{\mathcal{B}}$ , that is it plays the role of the 'integrability

test' for the equation under consideration in framework of AKNS scheme at least in the quasiclassical limit.

Validity of the above expressions was illustrated in Ref. [11] by an example of NLS equation. Here we will consider several other examples.

### 3. Examples

#### 3.1. Kaup-Boussinesq system

At first we will consider the Kaup-Boussinesq system

$$h_t + (hu)_x - \frac{1}{4}h_{xxx} = 0, \quad u_t + uu_x + h_x = 0, \quad (38)$$

which describes propagation of waves on shallow water [32, 33] and in two-component Bose-Einstein condensates [34]. In the shallow-water context,  $h(x, t)$  is the local depth of water and  $u(x, t)$  is its mean horizontal flow velocity. Linearized equations lead to the dispersion relation for harmonic waves,

$$\omega(k, h, u) = k \left( u \pm \sqrt{h + \frac{k^2}{4}} \right). \quad (39)$$

Equations of the dispersionless limit

$$h_t + (hu)_x = 0, \quad u_t + uu_x + h_x = 0 \quad (40)$$

are cast to the diagonal form (12) by introduction of the Riemann invariants

$$r_{\pm} = \frac{u}{2} \pm \sqrt{h}, \quad (41)$$

so that

$$v_+ = \frac{1}{2}(3r_+ + r_-), \quad v_- = \frac{1}{2}(r_+ + 3r_-) \quad (42)$$

and

$$u = r_+ + r_-, \quad h = \frac{1}{4}(r_+ - r_-)^2. \quad (43)$$

It is convenient to transform Eqs. (17) to independent variables  $h$  and  $u$  instead of  $r_+$  and  $r_-$ . Straightforward calculation gives

$$\frac{\partial k^2}{\partial h} = -4, \quad \frac{\partial k^2}{\partial u} = -2\sqrt{k^2 + 4h}, \quad (44)$$

and we get the solution in the form

$$k^2 = (q - u)^2 - 4h, \quad (45)$$

$q$  is an integration constant. Consequently, Eqs. (37) give the quasiclassical limit of the Lax pair functions (we assume  $q > u$  and the upper sign in the dispersion relation (39))

$$\bar{\mathcal{A}} = h - \left( \frac{q}{2} - \frac{u}{2} \right)^2, \quad \bar{\mathcal{B}} = - \left( \frac{q}{2} + \frac{u}{2} \right). \quad (46)$$

It is remarkable that in this case the quasiclassical limit coincides with the exact expressions [33] for the functions  $\mathcal{A}$  and  $\mathcal{B}$ , if we identify  $q/2$  with the spectral parameter  $\lambda$ .

The theory of propagation of high-frequency wave packets actually coincides with that theory for the defocusing NLS equation [8], since the dispersion relation (39) and dispersionless equations (40) are mathematically identical. Therefore substitution of Eq. (45) into the first Hamilton equation (6) gives Eq. (9) with  $\rho(x, t)$  replaced by  $h(x, t)$ .

Eqs. (39) and (40) satisfy the 'integrability test', that is the derivatives (44) commute and define the same function (45) as (7) in the NLS equation case, but we have two different dispersive generalizations with different Lax pairs. It is worth noticing that these two integrable systems, NLS and Kaup-Boussinesq equations, can be obtained as two different approximations to the integrable Landau-Lifshitz equations with an easy-plane anisotropy [34].

#### 3.2. Derivative nonlinear Schrödinger equation

Derivative nonlinear Schrödinger (DNLS) equation

$$i\psi_t + \frac{1}{2}\psi_{xx} - i(|\psi|^2\psi)_x = 0 \quad (47)$$

plays an important role in the theory of nonlinear Alfvén waves (see, e.g., Ref. [35]). It is transformed by means of the substitution (3) to a hydrodynamic-like form

$$\begin{aligned} \rho_t + \left[ \rho \left( u - \frac{3}{2}\rho \right) \right]_x &= 0, \\ u_t + uu_x - (\rho u)_x + \left( \frac{\rho_x^2}{8\rho^2} - \frac{\rho_{xx}}{4\rho} \right)_x &= 0. \end{aligned} \quad (48)$$

Linearization of these equations yields the dispersion relation for harmonic waves

$$\omega = k \left( u - 2\rho \pm \sqrt{\rho(\rho - u) + \frac{k^2}{4}} \right). \quad (49)$$

The uniform state is modulationally stable for  $\rho > u$ . Then the dispersionless limit equations

$$\begin{aligned} \rho_t + \left[ \rho \left( u - \frac{3}{2}\rho \right) \right]_x &= 0, \\ u_t + uu_x - (\rho u)_x &= 0 \end{aligned} \quad (50)$$

have real characteristic velocities

$$v_{\pm} = u - 2\rho \pm \sqrt{\rho(\rho - u)} \quad (51)$$

and can be transformed to a diagonal form (12) by introduction of the Riemann invariants

$$r_{\pm} = \frac{u}{2} - \rho \pm \sqrt{\rho(\rho - u)}. \quad (52)$$

The velocities (51) are expressed in terms of the Riemann invariants by Eqs. (42). Again we transform Eqs. (17) to independent variables  $\rho, u$  and obtain

$$\begin{aligned} \frac{\partial k^2}{\partial \rho} &= 4 \left[ \sqrt{k^2 + 4\rho(\rho - u)} - (2\rho - u) \right], \\ \frac{\partial k^2}{\partial u} &= -2 \left[ \sqrt{k^2 + 4\rho(\rho - u)} - 2\rho \right]. \end{aligned} \quad (53)$$

These derivatives commute and admit the solution

$$\begin{aligned} k^2 &= (q - u)^2 + 4q\rho \\ &\equiv (2\rho - u + q)^2 - 4\rho(\rho - u), \end{aligned} \quad (54)$$

$q$  is an integration constant. Assuming  $2\rho - u + q > 0$  and taking the upper sign in Eq. (49), we obtain from Eqs. (37)

$$\bar{\mathcal{A}} = -\frac{1}{4}(q - u)^2 - q\rho, \quad \bar{\mathcal{B}} = -\frac{q}{2} + \frac{u}{2} + \rho. \quad (55)$$

The Lax pair for the DNLS equation was found in Ref. [36] and in the scalar form it reads [23]

$$\begin{aligned} \mathcal{A} &= -\frac{1}{4} \left( \lambda^2 - \frac{i\psi_x}{\psi} \right)^2 + 4\lambda^2\rho - \left( \frac{i\psi_x}{2\psi} \right)_x, \\ \mathcal{B} &= 2\lambda^2 + \rho + \frac{i\psi_x}{2\psi}. \end{aligned} \quad (56)$$

After substitution of Eq. (3) and taking the dispersionless limit of negligibly small derivatives of  $\rho$  and  $u$ , we get

$$\bar{\mathcal{A}} = -\frac{1}{4}(\lambda^2 + u)^2 + 4\lambda^2\rho, \quad \bar{\mathcal{B}} = 2\lambda^2 - \frac{u}{2} + \rho. \quad (57)$$

These formulas coincide with Eqs. (55) for  $q = -4\lambda^2$ .

Substitution of Eq. (54) into the first Hamilton equation (6) gives a convenient form of equation of motion for high-frequency wave packets

$$\frac{dx}{dt} = q - \frac{2\rho(\rho - u)}{2\rho - u + q}, \quad (58)$$

where  $\rho = \rho(x, t), u = u(x, t)$  are given by some solution of the dispersionless Eqs. (50) and  $q$  is determined by the initial conditions.

### 3.3. Landau-Lifshitz equation

Landau-Lifshitz equation

$$\mathbf{M}_t = (\mathbf{M}_{xx} - M_3\mathbf{e}_3) \wedge \mathbf{M} \quad (59)$$

for the magnetization vector  $\mathbf{M} = (M_1, M_2, M_3)$ ,  $|\mathbf{M}| = 1$ , with an easy-plane anisotropy ( $\mathbf{e}_3$  is a unit vector of the  $z$  axis normal to the plane of anisotropy) can be transformed to more convenient variables  $w, v$  according to the formulas [34, 37, 38]

$$\begin{aligned} M_3 &= -w, \\ M_- &\equiv M_1 - iM_2 \\ &= \sqrt{1 - w^2} \exp\left(-i \int^x v dx'\right), \end{aligned} \quad (60)$$

so we get

$$\begin{aligned} w_t - [v(1 - w^2)]_x &= 0, \\ v_t - [w(1 - v^2)]_x \\ &+ \left[ \frac{1}{\sqrt{1 - w^2}} \left( \frac{w_x}{\sqrt{1 - w^2}} \right) \right]_x = 0. \end{aligned} \quad (61)$$

As usual, we obtain the dispersion relation

$$\omega = k \left( 2wv \pm \sqrt{k^2 + (1 - v^2)(1 - w^2)} \right) \quad (62)$$

for harmonic waves, that is we get the condition  $|v| \leq 1$  of modulation stability of a uniform state (the condition  $|w| \leq 1$  is satisfied by definition). The dispersionless equations

$$\begin{aligned} w_t - [v(1 - w^2)]_x &= 0, \\ v_t - [w(1 - v^2)]_x &= 0 \end{aligned} \quad (63)$$

have characteristic velocities

$$v_{\pm} = 2wv \pm \sqrt{(1 - v^2)(1 - w^2)} \quad (64)$$

and can be diagonalized by introduction of the Riemann invariants

$$r_{\pm} = wv \pm \sqrt{(1 - v^2)(1 - w^2)}. \quad (65)$$

As a result, Eqs. (63) are cast to the form (12) with  $v_{\pm}$  given by Eqs. (42). Eqs. (17) transformed to independent variables  $w, v$  have simple symmetrical form

$$\begin{aligned}\frac{\partial k^2}{\partial w} &= 2 \left[ w(1-v^2) - v\sqrt{k^2 + (1-v^2)(1-w^2)} \right], \\ \frac{\partial k^2}{\partial v} &= 2 \left[ v(1-w^2) - w\sqrt{k^2 + (1-v^2)(1-w^2)} \right].\end{aligned}\quad (66)$$

These derivatives commute and yield the solution

$$k^2 = (q - vw)^2 - (1 - v^2)(1 - w^2). \quad (67)$$

Assuming  $q - vw > 0$ , we obtain the expressions

$$\begin{aligned}\bar{\mathcal{A}} &= \frac{1}{4} \left[ (1 - v^2)(1 - w^2) - (q - vw)^2 \right], \\ \bar{\mathcal{B}} &= -q - vw.\end{aligned}\quad (68)$$

The Lax pair for the Landau-Lifshitz equation was found in Ref. [39] and in a scalar representation it takes the form [23]

$$\begin{aligned}\mathcal{A} &= \frac{1}{4} \left( i\lambda M_3 - \frac{(M_-)_x}{M_-} \right)^2 + \frac{1}{4} (1 - \lambda^2)(1 - M_3^2) \\ &\quad + \frac{1}{2} \left( i\lambda M_3 - \frac{(M_-)_x}{M_-} \right)_x, \\ \mathcal{B} &= -\lambda - i \left[ (M_3)_x - M_3 \frac{(M_-)_x}{M_-} \right].\end{aligned}\quad (69)$$

Substitution of Eqs. (60) and taking the dispersionless limit easily reproduce Eqs. (68) with  $\lambda = q$ .

At last, substitution of Eq. (67) into the expression for the group velocity yields the equation of motion for high-frequency wave packets

$$\frac{dx}{dt} = 2q - \frac{(1 - v^2)(1 - w^2)}{q - vw}, \quad (70)$$

where  $v = v(x, t)$ ,  $w = w(x, t)$  are given by some solution of Eqs. (63) and  $q$  is defined by initial conditions.

#### 4. Conclusion

We showed in this paper that the condition of quasiclassical integrability leads to important relationship between the carrier wave number  $k$  of high-frequency wave packets propagating along smooth background waves and the local values  $(r_+, r_-)$

of the background variables. This relationship does not depend on initial conditions for the background wave evolution. We showed that the function  $k = k(r_+, r_-)$  and the dispersion relation  $\omega = \omega(k, r_+, r_-)$  are related with the functions  $\bar{\mathcal{A}}, \bar{\mathcal{B}}$  which represent the Lax pairs for the corresponding equations in AKNS scheme in quasiclassical limit.

The obtained here relationship  $k = k(r_+, r_-)$  has several important applications. Its substitution into expression for the group velocity yields a convenient equation of motion for wave packets. On the other side, an analytical continuation of this formula on complex values of  $k = i\kappa$  gives the expression for the inverse half-width  $\kappa$  of solitons as a function of the local dispersionless variables, so we obtain the soliton's equation of motion along a smooth background wave. Besides that, the function  $k(r_+, r_-)$  is used in formulation of the Bohr-Sommerfeld quantization rule for parameters of solitons emerging from an intensive initial pulse. At last, if the equation under consideration is not completely integrable, but Eqs. (17) admit an approximate solution in the limit of large  $k$ , then we can apply it to non-integrable equations on the same footing as for integrable ones. Some examples of such an extended theory were considered in Refs. [8, 11, 40]. We believe that such an approach will find many other applications.

#### Acknowledgments

This research is funded by the research project FFUU-2021-0003 of the Institute of Spectroscopy of the Russian Academy of Sciences (Section II) and by the RSF grant number 19-72-30028 (Section III).

#### References

- [1] H. Flaschka, A. C. Newell and M. Tabor, *Integrability, in What Is Integrability?* Ed. V. E. Zakharov, p. 73, Springer, Berlin, 1991.
- [2] C. S. Gardner, J. M. Green, M. D. Kruskal and R. M. Miura, *Phys. Rev. Lett.*, 19 (1967) 1095.
- [3] P. D. Lax, *Commun. Pure Appl. Math.*, 21 (1968) 467.
- [4] V. E. Zakharov and A. B. Shabat, *Zh. Eksp. Teor. Fiz.*, 61 (1971) 118; *Sov. Phys. JETP*, 34 (1972) 62.
- [5] G. L. Lamb, jr., *Elements of Soliton Theory*, Wiley, N. y., 1980.
- [6] P. G. Drazin and R. S. Johnson, *Solitons: An Introduction*, CUP, Cambridge, 1989.
- [7] A. Scott, *Nonlinear Science. Emergence and Dynamics of Coherent Structures*, Oxford University Press, Oxford, 2003.
- [8] D. V. Shaykin and A. M. Kamchatnov, *Phys. Fluids*, 35 (2023) 062108.

- [9] J. L. Synge, Geometrical Optics. An Introduction into Hamilton's method, CUP, Cambridge, 1937.
- [10] Yu. A. Kravtsov, Yu. I. Orlov, Geometrical Optics of Inhomogeneous Media, Springer, Berlin, 1990.
- [11] A. M. Kamchatnov, Asymptotic theory of not completely integrable soliton equations, preprint arXiv:2305.12346 (2023).
- [12] G. B. Whitham, Proc. Roy. Soc. Lond., A 283 (1965) 238.
- [13] G. B. Whitham, Linear and Nonlinear Waves, Wiley, New York, 1974.
- [14] M. J. Ablowitz, D. J. Kaup, A. C. Newell, H. Segur, Stud. Appl. Math., 53 (1974) 249.
- [15] A. V. Gurevich and L. P. Pitaevskii, Zh. Eksp. Teor. Fiz., 93 (1987) 871 [Sov. Phys.-JETP, 93 (1987) 871].
- [16] A. M. Kamchatnov, Usp. Fiz. Nauk., 191 (2021) 52 [Phys.-Uspekhi, 64 (2021) 48].
- [17] A. M. Kamchatnov, Chaos 30 (2020) 123148.
- [18] G. A. El, A. Gammal, E. G. Khamis, R. A. Kraenkel, A. M. Kamchatnov, Phys. Rev. A 76 (2007) 053813.
- [19] G. A. El, R. H. J. Grimshaw, N. F. Smyth, Physica D, 237 (2008) 2423.
- [20] M. D. Maiden, N. A. Franco, E. G. Webb, G. A. El, and M. A. Hoefer, J. Fluid Mech. 883 (2020) A10.
- [21] A. M. Kamchatnov, Zh. Eksp. Teor. Fiz. 159 (2021) 76 [JETP, 132 (2021) 63].
- [22] L. F. Calazans de Brito, A. M. Kamchatnov, Phys. Rev. E 104 (2021) 054203.
- [23] A. M. Kamchatnov and R. A. Kraenkel, J. Phys. A: Math. Gen., 35 (2002) L13.
- [24] A. M. Kamchatnov, Phys. Lett. A, 186 (1994) 387.
- [25] A. M. Kamchatnov, Physica D, 188 (2004) 247.
- [26] A. M. Kamchatnov, J. Phys. A: Math. Gen., 34 (2001) L441.
- [27] I. M. Krichever, Funk. Analiz. Prilozh. 22 (1988) 37 [Funct. Anal. Appl., 22 (1988) 200].
- [28] V. I. Karpman, Phys. Lett. A, 25 (1967) 708.
- [29] V. I. Karpman, *Non-Linear Waves in Dispersive Media*, (Nauka, Moscow, 1973) (English translation: Pergamon Press, Oxford, 1975).
- [30] S. Jin, C. D. Levermore, D. W. McLaughlin, Comm. Pure Appl. Math., 52 (1999) 613.
- [31] A. M. Kamchatnov, R. A. Kraenkel, B. A. Umarov, Phys. Rev. E 66 (2002) 036609.
- [32] J. Boussinesq, Mém. Prés. Div. Sav. Acad. Sci. Inst. Fr. 23 (1877) 1.
- [33] D. J. Kaup, Prog. Theor. Phys. 54 (1975) 396.
- [34] S. K. Ivanov, A. M. Kamchatnov, T. Congy, and N. Pavloff, Phys. Rev. E 96 (2017) 062202.
- [35] C. F. Kennel, B. Buti, T. Hada, and R. Pellat, Phys. Fluids, 31 (1988) 1949.
- [36] D. J. Kaup and A. C. Newell, J. Math. Phys. 19 (1978) 798.
- [37] T. Congy, A. M. Kamchatnov, and N. Pavloff, SciPost Phys., 1 (2016) 006.
- [38] E. Iacocca, T. J. Silva, and M. A. Hoefer, Phys. Rev. Lett. 118 (2017) 017203.
- [39] A. E. Borovik and V. N. Robuk, Teor. Mat. Fiz. 46 (1981) 371 [Theor. Math. Phys. 46 (1981) 242].
- [40] A. M. Kamchatnov and D. V. Shaykin, Propagation of generalized Korteweg-de Vries solitons along large scale waves, preprint arXiv:2307.12321 (2023).