

The third homotopy module of a 2-complex

W.H.Mannan

ABSTRACT: Given a connected 2- complex X with fundamental group G , we show how $\pi_3(X)$ may be computed as a module over $\mathbb{Z}[G]$. Further we show that if X is a finite connected 2- complex with $G (= \pi_1(X))$ finite of odd order then the stable class of $\pi_3(X)$ is determined by G . In this case we also show that $\pi_3(X) \otimes \mathbb{Q}$ is free over $\mathbb{Q}[G]$.

MSC: 55Q15, 55Q20 55Q91 **Keywords** Homotopy groups, 2-complexes

email: wajid@mannan.info

§1 Introduction

Let X be a connected 2- complex with fundamental group G . Let $C_*(\tilde{X})$ denote the algebraic complex over $\mathbb{Z}[G]$ associated to the universal cover of X . We may write this algebraic complex:

$$F_2 \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0$$

From the Hurewicz isomorphism theorem, we know that

$$\pi_2(X) \cong H_2(C_*(\tilde{X}); \mathbb{Z}[G]) \cong \text{Ker} \partial_2$$

as (right) modules over $\mathbb{Z}[G]$. Hence $\pi_2(X)$ is easily attained from $C_*(\tilde{X})$.

Now let J be a module over $\mathbb{Z}[G]$, which is freely generated over \mathbb{Z} . There is a G - action on $J \otimes_{\mathbb{Z}} J$, given by $(a \otimes b)g = ag \otimes bg$, making it a $\mathbb{Z}[G]$ module. Let σ be the $\mathbb{Z}[G]$ - linear automorphism of $J \otimes_{\mathbb{Z}} J$ defined by $\sigma(a \otimes b) = b \otimes a$.

Definition 1.1 $S^2(J) = \{x \in J \otimes J \mid \sigma x = x\}$

We now state our main result:

Theorem A We have an isomorphism of $\mathbb{Z}[G]$ modules: $\pi_3(X) \cong S^2(\pi_2(X))$

When X is a finite connected 2- complex, Schanuel's lemma tells us that the stable class of $\pi_2(X)$ is determined by G . We prove an analogous result for $\pi_3(X)$, under the added assumption that G is finite, of odd order:

Theorem B Let X and X' be finite connected 2- complexes, with finite fundamental group G . If the order of G is odd, then $\pi_3(X)$ and $\pi_3(X')$ are stably equivalent.

Also if G is finite, of odd order we have the following result:

Theorem C Let X be a finite connected 2- complex with finite fundamental group G . If the order of G is odd, then $\pi_3(X) \otimes \mathbb{Q}$ is free over $\mathbb{Q}[G]$.

Definition 1.2 G -quadratic A map $f : A \rightarrow B$ between $\mathbb{Z}[G]$ modules A, B , is G -quadratic if given $a, b, c \in A$ and $g \in G$ we have:

- i) $f(-a) = f(a)$
- ii) $f(a + b + c) = f(a + b) + f(a + c) + f(b + c) - f(a) - f(b) - f(c)$
- iii) $f(ag) = f(a)g$

Lemma 1.3 Let $f : A \rightarrow B$ be G -quadratic;

- i) $f(0) = 0$
- ii) $f(a + b) + f(a - b) = 2f(a) + 2f(b)$

Proof: i) $f(0) = f(0 + 0 + 0) = 0$

$$\begin{aligned} \text{ii) } f(a + b) + f(a - b) &= f(a + b) + f(a - b) + f(b - b) \\ &= f(a + b - b) + f(a) + f(b) + f(-b) = 2f(a) + 2f(b) \quad \square \end{aligned}$$


Take J as before. We have a G -quadratic map $q : J \rightarrow S^2(J)$ which maps $\alpha \mapsto \alpha \otimes \alpha$, for any $\alpha \in J$. The main argument of this paper will depend on a certain universal property of q , which we mention as a preliminary.

Let $\{\alpha_i \mid i \in I\}$ be a basis for J over \mathbb{Z} for some set I . Given $\alpha \in J$ we may write $\alpha = \sum_{i \in I} \alpha_i \lambda_i$ with each $\lambda_i \in \mathbb{Z}$ and only finitely many λ_i non-zero. We define the norm of α to be $\sum_{i \in I} |\lambda_i|$.

The $\{\alpha_i \otimes \alpha_j \mid i, j \in I\}$ are a basis for $J \otimes_{\mathbb{Z}} J$ over \mathbb{Z} . So given $\omega \in S^2(J)$, we may write $\omega = \sum_{i, j \in I} \lambda_{ij} \alpha_i \otimes \alpha_j$. Clearly for each $i, j \in I$ we must have $\lambda_{ij} = \lambda_{ji}$. Hence $S^2(J)$ is generated freely over \mathbb{Z} by the $\alpha_i \otimes \alpha_i$ and the $\alpha_i \otimes \alpha_j + \alpha_j \otimes \alpha_i, i \neq j$.

For each $i, j \in I$ we have $\alpha_i \otimes \alpha_j + \alpha_j \otimes \alpha_i = q(\alpha_i + \alpha_j) - q(\alpha_i) + q(\alpha_j)$. Hence another basis for $S^2(J)$ over \mathbb{Z} is given by the $q(\alpha_i)$ and the $q(\alpha_i + \alpha_j), i \neq j$.

Lemma 1.4 The map $q : J \rightarrow S^2(J)$ is universal: Given any G -quadratic map $f : J \rightarrow M$, there exists a unique $\mathbb{Z}[G]$ -linear map $\phi : S^2(J) \rightarrow M$ making the following diagram commute:



Proof: We define ϕ linearly on a basis by:

$$\begin{aligned} \phi q(\alpha_i) &= f(\alpha_i) \\ \phi q(\alpha_i + \alpha_j) &= f(\alpha_i + \alpha_j), \quad i \neq j \end{aligned}$$

Clearly no other map can make the diagram commute. It remains to show that

$\phi q = f$ and that ϕ is $\mathbb{Z}[G]$ -linear.

By lemma 1.3 ii) we have $\phi q(\alpha_i - \alpha_j) = f(\alpha_i - \alpha_j)$. Hence, by construction we have $\phi q(\alpha) = f(\alpha)$ for any $\alpha \in J$ with norm less than or equal to 2.

We proceed by induction. Suppose we have verified that $\phi q(\alpha) = f(\alpha)$ for all $\alpha \in J$ with norm less than r , where $r \geq 3$. Then consider any $\beta \in J$ with norm equal to r . We may write $\beta = \alpha + x + y$ with $x = \pm\alpha_i$ and $y = \pm\alpha_j$ for some $i, j \in I$, such that the norms of $\alpha, \alpha + x, \alpha + y$ are all less than r .

Then property ii) of a G -quadratic map implies that

$$\phi q(\beta) = \phi q(\alpha + x + y) = f(\alpha + x + y) = f(\beta)$$

So we have verified that $\phi q(\alpha) = f(\alpha)$ for all $\alpha \in J$ with norm less than $r + 1$. Hence by induction $\phi q = f$. It remains to show that ϕ is $\mathbb{Z}[G]$ -linear.

By construction ϕ is \mathbb{Z} -linear. Given $\alpha \in J$ we have $\phi(q(\alpha)g) = \phi q(\alpha g) = f(\alpha g) = f(\alpha)g = (\phi q(\alpha))g$. Any element of $S^2(J)$ is a \mathbb{Z} -linear combination of elements of the form $q(\alpha), \alpha \in J$. Hence ϕ is $\mathbb{Z}[G]$ -linear on the whole of $S^2(J)$. \square

In §2 we consider a general topological space X with $\pi_1(X) = G$. We give a construction for a G -quadratic map $\pi_2(X) \rightarrow \pi_3(X)$. Whenever $\pi_2(X)$ is freely generated over \mathbb{Z} lemma 1.4 will therefore give us a $\mathbb{Z}[G]$ -linear map $\phi : S^2(\pi_2(X)) \rightarrow \pi_3(X)$. In §3 we show that when X is a connected 2-complex, this map is an isomorphism.

§2 The Hopf fibration

Let X be an arbitrary topological space with basepoint $*$. Let G denote $\pi_1(X)$. We will consider the map which the Hopf fibration $h : S^3 \rightarrow S^2$ induces on homotopy: $h^* : \pi_2(X) \rightarrow \pi_3(X)$. In particular we will show that it is G -quadratic.

We may regard S^3 as $\{(z, w) \in \mathbb{C}^2 \mid z\bar{z} + w\bar{w} = 1\}$. Define a relation \sim , by setting $(z, w) \sim (z\lambda, w\lambda)$ whenever $\lambda \in \mathbb{C}$ satisfies $\lambda\bar{\lambda} = 1$.

Definition 2.1 *Hopf fibration* The Hopf fibration, $h : S^3 \rightarrow S^2$ is the natural map $S^3 \rightarrow S^3/\sim$ composed with the identifications: $S^3/\sim \cong \mathbb{C}P^1 \cong S^2$.

We take $(0, 1)$ as basepoint of S^3 and let $+ = h(0, 1) \in S^2$ be the basepoint of S^2 . We define $h^* : \pi_2(X) \rightarrow \pi_3(X)$ to be the map given by precomposition with h . As composition of maps is associative we know that h^* is natural:

Lemma 2.2 *Let $f : X \rightarrow Y$ be a basepoint preserving continuous map of based topological spaces. Then $h^* f_* = f_* h^*$.*

We first show that h^* satisfies property i) of a G -quadratic map:

Lemma 2.3 *Let $\alpha \in \pi_2(X)$. Then $h^*(-\alpha) = h^*(\alpha)$.*

Proof: Let $R : S^3 \rightarrow S^3$ be the rotation given by $(z, w) \mapsto (\bar{z}, \bar{w})$. Let $\hat{R} : S^2 \rightarrow S^2$ be the reflection induced on S^3/\sim . Then R is homotopic to the identity (keeping the basepoint $(0, 1)$ fixed). So as elements of $\pi_3(X)$ we have:

$$h^*(-\alpha) = \alpha \circ \hat{R} \circ h = \alpha \circ h \circ R = \alpha \circ h = h^*(\alpha)$$

□

Given $\alpha, \beta \in \pi_2(X)$, we follow [5], Chap. X, §7 by denoting their Whitehead product $[\alpha, \beta]$. We also recall the following results from that section:

Proposition 2.4 *The Whitehead product satisfies:*

i) (Naturality, see (7.2)) Given $\alpha, \beta \in \pi_2(X)$ we have $f_*[\alpha, \beta] = [f_*\alpha, f_*\beta]$ for any continuous basepoint preserving map $f : X \rightarrow Y$.

ii) (Commutativity, see (7.5)) Given $\alpha, \beta \in \pi_2(X)$ we have $[\alpha, \beta] = [\beta, \alpha]$.

iii) (Biadditivity, see (7.12)) Given $\alpha_1, \alpha_2, \beta \in \pi_2(X)$ we have:

$[\alpha_1 + \alpha_2, \beta] = [\alpha_1, \beta] + [\alpha_2, \beta]$ and $[\beta, \alpha_1 + \alpha_2] = [\beta, \alpha_1] + [\beta, \alpha_2]$.

Whitehead products are related to h^* by the following result:

Lemma 2.5 $h^*(\alpha + \beta) = h^*(\alpha) + h^*(\beta) + [\alpha, \beta]$

Proof: See [5], Chap. XI, (1.16) and from (4.4) note that we may choose conventions so that the Hopf invariant of h is 1. □

We may now prove that h^* satisfies property ii) of a G -quadratic map:

Lemma 2.6 $h^*(\alpha + \beta + \gamma) = h^*(\alpha + \beta) + h^*(\alpha + \gamma) + h^*(\beta + \gamma) - h^*(\alpha) - h^*(\beta) - h^*(\gamma)$

Proof: $h^*(\alpha + \beta + \gamma) = h^*(\alpha + \beta) + h^*(\gamma) + [\alpha + \beta, \gamma]$

$= h^*(\alpha + \beta) + h^*(\alpha + \gamma) + h^*(\beta + \gamma) - h^*(\alpha) - h^*(\beta) - h^*(\gamma)$

□

Finally it remains to show that h^* respects the action of G . Let $\alpha : S^2 \rightarrow X$ represent an element of $\pi_2(X)$. We may identify the complement of $+$ in S^2 with the interior of the closed unit disk $D^2 \subset \mathbb{R}^2$. α extends naturally to a map $\alpha' : D^2 \rightarrow X$ which maps the boundary ∂D^2 to the basepoint $* \in X$. Let R_θ denote a positive rotation of D^2 through the angle θ .

We consider $h^*(\alpha) : S^3 \rightarrow X$. For each $(z, w) \in S^3$ we have a fibre $\{(\lambda z, \lambda w) \mid \lambda \bar{\lambda} = 1\}$. If $z \neq 0$ this fibre completes a revolution of the origin in the w -plane as it passes round the z -plane. The fibre $\{(0, \lambda) \mid \lambda \bar{\lambda} = 1\}$ gets mapped to $* \in X$ by $h^*(\alpha)$. We may deform $h^*(\alpha)$ to thicken this fibre. The remaining fibres then form a solid torus. We may therefore represent $h^*(\alpha) \in \pi_3(X)$ by the map $\alpha^T : S^3 \rightarrow X$ defined as follows:

Consider an unknotted solid torus $N \subset S^3$, not containing the basepoint. We make an untwisted topological identification of N with $S^1 \times D^2$. Then:

$$\begin{aligned} \alpha^T(\theta, d) &= \alpha'(R_\theta(d)) & (\theta, d) \in N \\ \alpha^T(p) &= * & p \notin N \end{aligned}$$

Let I denote the closed unit interval and let $g : I \rightarrow X$ represent an element of $\pi_1(X)$, so $g(0) = g(1) = *$. Then g acts on α by stretching it round the loop g . Formally we have $\alpha g : D^2 \rightarrow X$ defined by:

$$\begin{aligned} \alpha g(d) &= \alpha'(2d) & |d| &\leq \frac{1}{2} \\ \alpha g(d) &= g(2|d| - 1) & |d| &\geq \frac{1}{2} \end{aligned}$$

Let $K \subset S^3$ be a thickened torus (parameterized $T^2 \times I$), bounding N and not enclosing the basepoint. Then $h^*(\alpha g)$ is represented by the map $(\alpha g)^T : S^3 \rightarrow X$, given by:

$$\begin{aligned} (\alpha g)^T(\theta, d) &= \alpha'(R_\theta(d)) & (\theta, d) &\in N \\ (\alpha g)^T(t, i) &= g(i) & (t, i) &\in K \\ (\alpha g)^T(p) &= * & p &\notin N, K \end{aligned}$$

Let $L \subset S^3$ be a thickened sphere (parameterized $S^2 \times I$), enclosing N and not enclosing the basepoint. The group element g acts on an element of $\pi_3(X)$ by stretching it round the loop g . Formally, $h^*(\alpha)g$ is represented by the map $\alpha^T g : S^3 \rightarrow X$, given by:

$$\begin{aligned} \alpha^T g(\theta, d) &= \alpha'(R_\theta(d)) & (\theta, d) &\in N \\ \alpha^T g(s, i) &= g(i) & (s, i) &\in L \\ \alpha^T g(p) &= * & p &\notin N, L \end{aligned}$$

Lemma 2.7 *Let g be an element of $\pi_1(X)$ and let α be an element of $\pi_2(X)$. Then $h^*(\alpha)g = h^*(\alpha g)$.*

Proof: We must show that $\alpha^T g$ is homotopic to $(\alpha g)^T$. Let $C \subset S^3$ be a solid cylinder which passes through the hole in the solid torus N and passes through the thickened sphere L on either side of this hole.

We may parameterize C as $D^2 \times I$ so that for $(d, i) \in C$, we have

$$\alpha^T g(d, i) = m(i)$$

where $m = e \cdot g^{-1} \cdot e \cdot g \cdot e$. We define a homotopy $m_t, t \in I$, from m to e :

$$\begin{aligned} m_t(i) &= g(5r - .5) & r &\in [.1, .3] \\ m_t(i) &= * & r &\notin [.1, .3] \end{aligned}$$

where $r = \sqrt{t^2 + (i - .5)^2}$.

We next construct a homotopy f_t with $f_0 = \alpha^T g$, which is constant outside C . Inside C it is given by:

$$f_t(d, i) = m_{(1-|d|)t}(i)$$

Thus intuitively f pulls together two regions of the thickened sphere L through the hole in the solid torus N , so that they cancel out, leaving a thickened torus as required. □

We may conclude:

Proposition 2.8 *h^* is a G -quadratic map.*

Proof: We get property i) from lemma 2.3, property ii) from lemma 2.6 and property iii) from lemma 2.7. □

§3 π_3 of a connected 2- complex

We now restrict to the case where X is a connected 2- complex with basepoint $*$ and $\pi_1(X) = G$. The universal cover of X is a simply connected 2- complex and hence a Moore space.

As $H_2(\tilde{X}; \mathbb{Z})$ is a submodule of the free \mathbb{Z} - module generated by the 2- cells of \tilde{X} , we have $H_2(\tilde{X}; \mathbb{Z}) \cong \mathbb{Z}^I$ for some set I (see [3], Chap. IV, Theorem 6.1).

From [1], Lemma 1.3.1, we know that up to homotopy there is a unique CW complex of type $M(\mathbb{Z}^I, 2)$. Hence we have a homotopy equivalence:

$$\bigvee_{i \in I} S_i^2 \rightarrow \tilde{X}$$

preserving basepoint (which we take to be $\bigcap_{i \in I} S_i^2$).

Composing with the covering map $\tilde{X} \rightarrow X$, we obtain a map $m : \bigvee_i S_i^2 \rightarrow X$. For $n > 1$, the map m induces an isomorphism $\pi_n(\bigvee_i S_i^2) \rightarrow \pi_n(X)$.

For $j \in I$ let $w_j : S^2 \rightarrow \bigvee_i S_i^2$ be a homeomorphism onto S_j^2 , preserving basepoint.

Lemma 3.1 *i) $\pi_2(\bigvee_i S_i^2)$ is freely generated over \mathbb{Z} by the w_i .*

ii) $\pi_3(\bigvee_i S_i^2)$ is freely generated over \mathbb{Z} by the $h^(w_i)$ and the $[w_i, w_j]$, $i \neq j$.*

Proof: For i) see [2], Example 4.26 and for ii) see [2], Example 4.52. \square

Let $\alpha_i = m_*(w_i)$ for $i \in I$. As m_* is an isomorphism $\pi_2(\bigvee_i S_i^2) \rightarrow \pi_2(X)$, we know the α_i freely generate $\pi_2(X)$ over \mathbb{Z} .

Lemma 3.2 *We have a \mathbb{Z} - basis for $\pi_3(X)$ given by the $h^*(\alpha_i)$ and the $h^*(\alpha_i + \alpha_j)$, $i \neq j$.*

Proof: As m_* is an isomorphism $\pi_3(\bigvee_i S_i^2) \rightarrow \pi_3(X)$, the $m_*(h^*(w_i))$ and the $m_*[w_i, w_j]$ freely generate $\pi_3(X)$ over \mathbb{Z} . By lemma 2.2 and proposition 2.4i) as well as lemma 2.5 we have:

$$\begin{aligned} m_*(h^*(w_i)) &= h^*(\alpha_i) & \forall i \\ m_*[w_i, w_j] &= [\alpha_i, \alpha_j] = h^*(\alpha_i + \alpha_j) - h^*(\alpha_i) - h^*\alpha_j & \forall i \neq j \end{aligned} \quad \square$$

By lemma 1.4 we have a $\mathbb{Z}[G]$ - linear map $\phi : S^2(\pi_2(X)) \rightarrow \pi_3(X)$ which satisfies $\phi q = h^*$.

Lemma 3.3 *ϕ is an isomorphism.*

Proof: Recall that the $q(\alpha_i)$ and the $q(\alpha_i + \alpha_j)$, $i \neq j$ form a \mathbb{Z} - basis for $S^2(\pi_2(X))$. It is sufficient to show that ϕ maps this basis to a \mathbb{Z} - basis of $\pi_3(X)$. By construction, $\phi q(\alpha_i) = h^*(\alpha_i)$ and $\phi q(\alpha_i + \alpha_j) = h^*(\alpha_i + \alpha_j)$. By lemma 3.2 the $h^*(\alpha_i)$ and the $h^*(\alpha_i + \alpha_j)$, $i \neq j$, form a \mathbb{Z} - basis of $\pi_3(X)$ as required. \square

From this lemma we may conclude our Theorem A:

Theorem A We have an isomorphism of $\mathbb{Z}[G]$ modules: $\pi_3(X) \cong S^2(\pi_2(X))$

§4 The effect of stabilizing π_2 .

Let X be a finite connected 2- complex with finite fundamental group G . Let X' be another finite connected 2- complex with fundamental group G . By Schanuel's lemma we know that there exist integers a, b such that

$$\pi_2(X) \oplus \mathbb{Z}[G]^a \cong \pi_2(X') \oplus \mathbb{Z}[G]^b$$

In this section we investigate the corresponding relationship between $\pi_3(X)$ and $\pi_3(X')$.

If we let $J = \pi_2(X)$ and $J' = \pi_2(X')$, then from Theorem A we have $\pi_3(X) \cong S^2(J)$ and $\pi_3(X') \cong S^2(J')$. We know that

$$S^2(J \oplus \mathbb{Z}[G]^a) \cong S^2(J' \oplus \mathbb{Z}[G]^b)$$

The next few lemmas give us an expansion of this.

Lemma 4.1 *Let $A_i, i = 1, \dots, n$, be modules over $\mathbb{Z}[G]$ with finitely generated free underlying Abelian groups. Then*

$$S^2\left(\bigoplus_{i=1}^n A_i\right) = \bigoplus_{i=1}^n S^2(A_i) \oplus \bigoplus_{i < j} A_i \otimes_{\mathbb{Z}} A_j$$

Proof: For each i , let the $e_{i,r}$ be a basis over \mathbb{Z} for A_i . Then $S^2\left(\bigoplus_{i=1}^n A_i\right)$ is freely generated over \mathbb{Z} , by elements of the form:

$$\begin{aligned} & e_{i,r} \otimes e_{i,r}, \\ & e_{i,r} \otimes e_{i,s} + e_{i,s} \otimes e_{i,r} && r < s, \\ & e_{i,r} \otimes e_{j,s} + e_{j,s} \otimes e_{i,r} && i < j, \end{aligned}$$

For each i , the \mathbb{Z} - linear span of the

$$\begin{aligned} & e_{i,r} \otimes e_{i,r}, \\ & e_{i,r} \otimes e_{i,s} + e_{i,s} \otimes e_{i,r} && r < s, \end{aligned}$$

is closed under the group action and is isomorphic (over $\mathbb{Z}[G]$) to $S^2(A_i)$.

Similarly, for each pair i, j , with $i < j$, the \mathbb{Z} - linear span of the

$$e_{i,r} \otimes e_{j,s} + e_{j,s} \otimes e_{i,r} \quad \forall r, s$$

is closed under the group action. Furthermore we have an isomorphism (over $\mathbb{Z}[G]$) from it to $A_i \otimes_{\mathbb{Z}} A_j$, which maps:

$$e_{i,r} \otimes e_{j,s} + e_{j,s} \otimes e_{i,r} \quad \mapsto \quad e_{i,r} \otimes e_{j,s}$$

So over $\mathbb{Z}[G]$, the module $S^2\left(\bigoplus_{i=1}^n A_i\right)$ decomposes into the sum:

$$\bigoplus_{i=1}^n S^2(A_i) \oplus \bigoplus_{i < j} A_i \otimes_{\mathbb{Z}} A_j$$

□

Lemma 4.2 *Let A be a $\mathbb{Z}[G]$ module whose underlying Abelian group is isomorphic to \mathbb{Z}^k . Then $A \otimes_{\mathbb{Z}} \mathbb{Z}[G] \cong \mathbb{Z}[G]^k$.*

Proof: The action of an element $g \in G$ on A is a \mathbb{Z} -linear isomorphism. Hence, if e_1, \dots, e_k is a \mathbb{Z} -linear basis for A , then so is e_1g, \dots, e_kg . Hence $A \otimes_{\mathbb{Z}} \mathbb{Z}[G]$ is freely generated over \mathbb{Z} by elements of the form $e_i g \otimes g$, for $g \in G$ and $i \in \{1, \dots, k\}$.

For each i , the \mathbb{Z} -linear span of the $e_i g \otimes g$, $g \in G$, is closed under the action of G . Furthermore, we have a $\mathbb{Z}[G]$ -linear isomorphism from it to $\mathbb{Z}[G]$, which maps $e_i g \otimes g \mapsto g$.

Hence,

$$A \otimes_{\mathbb{Z}} \mathbb{Z}[G] \cong \bigoplus_{i=1}^k \mathbb{Z}[G] \cong \mathbb{Z}[G]^k$$

□

Let p denote the number of pairs, $\{g, g^{-1}\} \in G$, with $g \neq g^{-1}$. Let T denote the set of $g \in G$ with g having order 2.

Definition 4.3 For a finite group G , we set

$$V_G = \bigoplus_{t \in T} (1+t)\mathbb{Z}[G]$$

Lemma 4.4 $S^2(\mathbb{Z}[G]) \cong \mathbb{Z}[G]^{1+p} \oplus V_G$

Proof: Let the elements of G be ordered. We have a \mathbb{Z} -basis for $S^2(\mathbb{Z}[G])$ given by:

$$\begin{array}{ll} (e \otimes e)h & h \in G \\ (e \otimes g + g \otimes e)h & g < g^{-1}, \quad h \in G \\ (e \otimes t + t \otimes e)h & t \in T, \quad h < th \end{array}$$

The \mathbb{Z} -linear span of the $\{(e \otimes e)h \mid h \in G\}$ is isomorphic (over $\mathbb{Z}[G]$) to $\mathbb{Z}[G]$. Also, for $g < g^{-1}$, the \mathbb{Z} -linear span of the $\{(e \otimes g + g \otimes e)h \mid h \in G\}$ is isomorphic (over $\mathbb{Z}[G]$) to $\mathbb{Z}[G]$.

Given $t \in T$, the \mathbb{Z} -linear span of the $\{(e \otimes t + t \otimes e)h \mid h < th\}$ is isomorphic to $(e \otimes t)(1+t)\mathbb{Z}[G] \subset \mathbb{Z}[G] \otimes_{\mathbb{Z}} \mathbb{Z}[G]$.

Finally note that as $(e \otimes t)\mathbb{Z}[G] \cong \mathbb{Z}[G]$, we have $(e \otimes t)(1+t)\mathbb{Z}[G] \cong (1+t)\mathbb{Z}[G]$.

□

Note that for $\mathbb{Z}[G]$ -modules A, B , we have a $\mathbb{Z}[G]$ -linear isomorphism $A \otimes_{\mathbb{Z}} B \rightarrow B \otimes_{\mathbb{Z}} A$, given by sending $a \otimes b \mapsto b \otimes a$.

Using lemma 4.1 to expand the identity $S^2(J \oplus \mathbb{Z}[G]^a) \cong S^2(J' \oplus \mathbb{Z}[G]^b)$ we obtain:

$$\begin{aligned} & S^2(J) \oplus S^2(\mathbb{Z}[G]^a) \oplus (J \otimes_{\mathbb{Z}} \mathbb{Z}[G])^a \oplus (\mathbb{Z}[G] \otimes_{\mathbb{Z}} \mathbb{Z}[G])^{a(a-1)/2} \\ & \cong S^2(J') \oplus S^2(\mathbb{Z}[G]^b) \oplus (J' \otimes_{\mathbb{Z}} \mathbb{Z}[G])^b \oplus (\mathbb{Z}[G] \otimes_{\mathbb{Z}} \mathbb{Z}[G])^{b(b-1)/2} \end{aligned}$$

Let k denote the \mathbb{Z} -rank of J , and let k' denote the \mathbb{Z} -rank of J' . Also let n denote the order of G . By Theorem A we have $S^2(J) \cong \pi_3(X)$ and $S^2(J') \cong \pi_3(X')$. Applying lemmas 4.2 and 4.4 to the equation above gives:

$$\begin{aligned} & \pi_3(X) \oplus (\mathbb{Z}[G]^{1+p} \oplus V_G)^a \oplus \mathbb{Z}[G]^{ka} \oplus \mathbb{Z}[G]^{na(a-1)/2} \\ & \cong \pi_3(X') \oplus (\mathbb{Z}[G]^{1+p} \oplus V_G)^b \oplus \mathbb{Z}[G]^{k'b} \oplus \mathbb{Z}[G]^{nb(b-1)/2} \end{aligned}$$

So:

$$\pi_3(X) \oplus \mathbb{Z}[G]^{(1+p+k+n(a-1)/2)a} \oplus V_G^a \cong \pi_3(X') \oplus \mathbb{Z}[G]^{(1+p+k'+n(b-1)/2)b} \oplus V_G^b$$

Theorem 4.5 *Let X and X' be finite connected 2- complexes, with finite fundamental group G . Then there exist integers a, b, q, r such that:*

$$\pi_3(X) \oplus \mathbb{Z}[G]^q \oplus V_G^a \cong \pi_3(X') \oplus \mathbb{Z}[G]^r \oplus V_G^b$$

Note that if the order of G is odd then it does not contain any elements of order 2. Hence $V_G = 0$ and we have obtained Theorem B:

Theorem B *Let X and X' be finite connected 2- complexes, with finite fundamental group G . If the order of G is odd, then $\pi_3(X)$ and $\pi_3(X')$ are stably equivalent.*

§5 Calculating $S^2(IG^*)$

Again let X be a finite connected 2- complex with finite fundamental group G . Rationally we have $\pi_2(X) \otimes \mathbb{Q} \cong IG^* \otimes \mathbb{Q} \oplus \mathbb{Q}[G]^m$, for some $m \geq 0$. Consequently $\pi_3(X) \otimes \mathbb{Q} \cong S^2(IG^* \oplus \mathbb{Z}[G]^m) \otimes \mathbb{Q}$ as modules over $\mathbb{Q}[G]$. For this reason we examine $S^2(IG^*)$.

We will then conclude by considering some examples where $\pi_2(X)$ actually equals IG^* . For this to happen the group G must necessarily have a balanced presentation and free period 4.

IG^* is defined by the short exact sequence:

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}[G] \xrightarrow{\delta} IG^* \rightarrow 0$$

where $1 \in \mathbb{Z}$ is mapped to $\sum_{g \in G} g \in \mathbb{Z}[G]$, denoted Σ .

We now consider the $\mathbb{Z}[G]$ - linear surjection $\delta' : S^2(\mathbb{Z}[G]) \rightarrow S^2(IG^*)$ defined by:

$$\begin{aligned} g \otimes g & \mapsto \delta(g) \otimes \delta(g) \\ g \otimes h + h \otimes g & \mapsto \delta(g) \otimes \delta(h) + \delta(h) \otimes \delta(g) \end{aligned}$$

Lemma 5.1 *The kernel of δ' is generated over $\mathbb{Z}[G]$, by $\Sigma \otimes e + e \otimes \Sigma$ and $\Sigma \otimes \Sigma$.*

Proof: We may take a \mathbb{Z} - basis for $\mathbb{Z}[G]$ given by $\{g \in G \mid g \neq e\} \sqcup \{\Sigma\}$. We have a \mathbb{Z} - basis for IG^* given by $\{p(g) \mid g \in G, g \neq e\}$.

These basis' induce basis' on $S^2(\mathbb{Z}[G])$ and $S^2(IG^*)$ respectively. We have:

$$\begin{aligned} \delta'(\Sigma \otimes \Sigma) &= 0 \\ \delta'(g \otimes \Sigma + \Sigma \otimes g) &= 0 \quad g \neq e \end{aligned}$$

and δ' maps the other elements of the induced basis for $S^2(\mathbb{Z}[G])$ to elements of the induced basis for $S^2(IG^*)$.

Hence the kernel of δ' is generated over \mathbb{Z} by $\Sigma \otimes \Sigma$ and $g \otimes \Sigma + \Sigma \otimes g$, $g \neq e$. Clearly this is contained in the $\mathbb{Z}[G]$ linear span of $\Sigma \otimes e + e \otimes \Sigma$ and $\Sigma \otimes \Sigma$. \square

Let $u = e \otimes \Sigma + \Sigma \otimes e$. Then we have the kernel of δ' generated over $\mathbb{Z}[G]$ by $\{u, u\Sigma/2\}$. Note that $u\mathbb{Z}[G] \cong \mathbb{Z}[G]$.

As $u\Sigma/2$ is in the $\mathbb{Q}[G]$ -linear span of u , the kernel of $\delta' \otimes \mathbb{Q} : S^2(\mathbb{Q}[G]) \rightarrow S^2(IG^*) \otimes \mathbb{Q}$ is generated by u . We therefore have a short exact sequence:

$$0 \rightarrow \mathbb{Q}[G] \rightarrow S^2(\mathbb{Q}[G]) \rightarrow S^2(IG^*) \otimes \mathbb{Q} \rightarrow 0$$

As surjections over $\mathbb{Q}[G]$ split and cancellation of finitely generated modules holds over $\mathbb{Q}[G]$, we may write:

$$S^2(IG^*) \otimes \mathbb{Q} \cong S^2(\mathbb{Q}[G])/\mathbb{Q}[G]$$

Rationally, for some $a \in \mathbb{Z}$, $\pi_2(X) \otimes \mathbb{Q} \cong (IG^* \otimes \mathbb{Q}) \oplus \mathbb{Q}[G]^a$. Consequentially, we have:

$$\pi_3(X) \otimes \mathbb{Q} \cong S^2(\pi_2(X)) \otimes \mathbb{Q} \cong S^2(\pi_2(X) \otimes \mathbb{Q}) \cong S^2(IG^* \oplus \mathbb{Z}[G]^a) \otimes \mathbb{Q}$$

From lemmas 4.1, 4.2 and 4.4, for some $q \in \mathbb{Z}$ we have:

$$S^2(IG^* \oplus \mathbb{Z}[G]^a) \otimes \mathbb{Q} \cong (S^2(IG^*) \otimes \mathbb{Q}) \oplus \mathbb{Q}[G]^q \oplus (V_G \otimes \mathbb{Q})^a$$

As $S^2(IG^*) \otimes \mathbb{Q} \cong (S^2(\mathbb{Z}[G]) \otimes \mathbb{Q})/\mathbb{Q}[G]$, from lemma 4.4 we may conclude:

Theorem 5.2

$$\pi_3(X) \otimes \mathbb{Q} \cong \mathbb{Q}[G]^{p+q} \oplus (V_G \otimes \mathbb{Q})^{a+1}$$

Again if the order of G is odd then $V_G = 0$ so we obtain our Theorem C:

Theorem C *Let X be a finite connected 2-complex with finite fundamental group G . If the order of G is odd, then $\pi_3(X) \otimes \mathbb{Q}$ is free over $\mathbb{Q}[G]$.*

Let S be a subset of G , containing precisely one of g or g^{-1} , for each pair g, g^{-1} , with $g \neq g^{-1}$. Let $e_g = g \otimes e + e \otimes g$ for each $g \in S$. Let $e_t = t \otimes e + e \otimes t$ for each $t \in T$. Let $e_0 = e \otimes e$. Then from the proof of lemma 4.4 we have:

$$S^2(\mathbb{Z}[G]) = e_0\mathbb{Z}[G] \oplus \bigoplus_{g \in S} e_g\mathbb{Z}[G] \oplus \bigoplus_{t \in T} e_t\mathbb{Z}[G]$$

Also,

$$u = e \otimes \Sigma + \Sigma \otimes e = e_0 2 + \sum_{g \in S} e_g(1 + g^{-1}) + \sum_{t \in T} e_t$$

and

$$S^2(IG^*) = \frac{S^2(\mathbb{Z}[G])}{\langle u, u\frac{\Sigma}{2} \rangle}$$

The relation $u = 0$ is equivalent to $e_0 2 = -(\sum_{g \in S} e_g(1 + g^{-1}) + \sum_{t \in T} e_t)$.

For $t \in T$, we let $S_t \in \mathbb{Z}[G]$ be some element satisfying $(1 + t)S_t = \Sigma$. Hence we have $e_t S_t = (e \otimes t)(1 + t)S_t = (e \otimes t)\Sigma = (e \otimes t)(1 + t)\Sigma/2 = e_t \Sigma/2$.

The relation $u\Sigma/2 = 0$ is equivalent to $e_0 \Sigma = -(\sum_{g \in S} e_g \Sigma + \sum_{t \in T} e_t S_t)$.

Let

$$M = \bigoplus_{g \in S} e_g \mathbb{Z}[G] \oplus \bigoplus_{t \in T} e_t \mathbb{Z}[G]$$

and let $u_M = -(\sum_{g \in S} e_g(1 + g^{-1}) + \sum_{t \in T} e_t)$.

Proposition 5.3 *If $\pi_2 X = IG^*$ then $\pi_3(X) = M[\frac{u_M}{2}]$.*

Proof: We need to check that $\frac{u_M}{2}$ satisfies the relations which e_0 was subject to:

$$\frac{u_M}{2} 2 = -(\sum_{g \in S} e_g(1 + g^{-1}) + \sum_{t \in T} e_t)$$

$$\frac{u_M}{2} \Sigma = -(\sum_{g \in S} e_g \Sigma + \sum_{t \in T} e_t S_t) \quad \square$$

Example 5.4 $G = C_3$ and X is the Cayley complex associated to the balanced presentation: $\langle x \mid x^3 = e \rangle$.

From [4], 41(i) we have $\pi_2(X) = IG^*$. Hence:

$$\pi_3(X) \cong S^2(IG^*) \cong e_{x^2} \mathbb{Z}[G][[(1 + x)/2] \cong \mathbb{Z}[G][[\frac{(1 + x)}{2}]]$$

Example 5.5 $G = Q_8$ and X is the Cayley complex associated to the balanced presentation: $\langle x, y \mid x^2 = y^2, xyx = y \rangle$.

Again from [4], 41(iii) we have $\pi_2(X) = IG^*$. There is only one element of order two in Q_8 so we have:

$$\pi_3(X) \cong S^2(IG^*) \cong (e_x \mathbb{Z}[G] \oplus e_y \mathbb{Z}[G] \oplus e_{xy} \mathbb{Z}[G] \oplus e_{y^2} \mathbb{Z}[G])[\frac{u_M}{2}]$$

where $u_M = -(e_x(1 + x^3) + e_y(1 + y^3) + e_{xy}(1 + xy^3) + e_{y^2})$.

Recall from the proof of lemma 4.4 that:

$$e_x \mathbb{Z}[G] \cong e_y \mathbb{Z}[G] \cong e_{xy} \mathbb{Z}[G] \cong \mathbb{Z}[G], \quad e_{y^2} \mathbb{Z}[G] \cong (1 + y^2) \mathbb{Z}[G]$$

References

- [1] Hans-Joachim Baues, *Homotopy Type and Homology* (OUP, 1996)
- [2] Allen Hatcher, *Algebraic Topology* (CUP, 2002)
- [3] Thomas W. Hungerford, *Algebra* (GTM 73, 1974)
- [4] F.E.A. Johnson, *Stable Modules and the D(2)- Problem* (LMS 301, 2003)
- [5] George W. Whitehead, *Elements of Homotopy Theory* (GTM 61, 1978)