

Structural and microscopic relaxation processes in liquid Hydrogen Fluoride

R. Angelini¹, P. Giura¹, G. Monaco¹, G. Ruocco³, F. Sette¹, and R. Verbeni¹.

¹ *European Synchrotron Radiation Facility. B.P. 220 F-38043 Grenoble, Cedex France.*

³ *Università di Roma "La Sapienza" and Istituto Nazionale di Fisica della Materia, I-00185, Roma, Italy.*

(Dated: August 28, 2023)

The high frequency collective dynamics of liquid hydrogen fluoride (HF) is studied by inelastic x-ray scattering on the coexistence curve at $T=239$ K. The comparison with existing molecular dynamics simulations shows the existence of two active relaxation processes with characteristic time-scales in the sub-picosecond range. The observed scenario is very similar to that found in liquid water. This suggests that hydrogen bonded liquids behave similarly to other very different systems as simple and glass forming liquids, thus indicating that these two relaxation processes are universal features of the liquid state.

PACS numbers: 61.20.-p, 63.50.+x, 61.10.Eq, 78.70.Ck

A marked characteristic of the liquid state is that the dynamics of the density fluctuations is controlled by relaxation processes. For example, in liquids, the structural re-arrangement of the particles takes place with a characteristic time-scale, τ_α , dictated by the local inter-particles interaction and the actual thermodynamic state. The presence of these re-arrangements affects the dynamics of the density fluctuations as it allows energy exchanges between different density fluctuation modes. In this context, one calls *relaxation process* the mechanism governing these energy flows, and, in this specific example, we have described the well known *structural- or α -relaxation process*[1]. Other relaxation processes may be active, each one characterized by a specific underlying physical mechanism. One of the open problems in the physics of the liquid state is to understand on a general ground the common features of these relaxation processes. In simple monatomic liquids both kinetic and mode-coupling theories predict the existence of two distinct relaxation processes [2] and, this prediction has been demonstrated both by numerical simulations [3, 4] and by experiments [5]. In these theories, one of the two processes is the α -relaxation, and the other is a faster process (*microscopic or instantaneous process*) which is thought to be associated with the interactions between an atom and the "cage" of its nearest neighbors. Other relaxation processes, beyond the α and the instantaneous processes, associated with the internal molecular degrees of freedom may be observed in molecular liquids. In glass-forming systems, one also finds the same relaxation processes pattern. In this case, by driving the system to the glassy state, where the structural arrest freezes the α -process and τ_α becomes extremely large, is still possible to clearly observe the microscopic process [6].

The outlined scenario seems to point that the α - and μ - relaxation processes are universal features of the liquid state. In this context, however, no attempt has been made so far to include the important class of hydrogen

bonded liquids. This is the aim of the present work.

The operational method to identify relaxation processes is the study of the dynamic structure factor, $S(Q, \omega)$, and of its inelastic features due to collective excitations - at momentum transfer Q these are observed at energy $\hbar\Omega(Q)$. The dispersion relation of $\Omega(Q)$ allows to define an "apparent" sound velocity, $c(Q) = \Omega(Q)/Q$, which is a constant in the low Q -limit, and decreases with Q approaching the inverse of the inter-particle distance d . Whenever a relaxation process with characteristic time τ is active, $c(Q)$ has a further Q -dependence which shows up as a transition from a low frequency value, c_o , to a high frequency one, c_∞ . This transition takes place when the condition $\Omega(Q) \approx 1/\tau$ is fulfilled. The amplitude of the jump between c_∞ and c_o defines the strength, Δ , of the coupling between the density fluctuations and the degrees of freedom involved in the relaxation process. Multiple relaxation processes will give rise to multiple "jumps" in $c(Q)$. Specifically, in the case of α - and μ - relaxation processes, one expects that $c(Q)$ goes from c_o to $c_{\infty\alpha}$ because of the α -process, and then from $c_{o\mu} = c_{\infty\alpha}$ to c_∞ because of the μ -process [7]. In real systems, this picture is often an over-simplification. Indeed: *i*) The sound velocities c_o , $c_{\infty\alpha}$ and c_∞ , as well as the associated quantities τ and Δ are Q -dependent, especially when $Qd \approx 1$. *ii*) In certain cases the condition $\omega\tau_\mu = 1$ cannot be reached because $1/\tau_\mu(Q)$ is always larger than the maximum value of $\Omega(Q)$.

In the framework of hydrogen bonded liquids, water has been extensively studied and we report in Fig. 1 a summary of our current understanding of the collective dynamics in terms of apparent sound velocity. Experimental and numerical simulation data points for ambient conditions fall between the $c_o(Q)$ and $c_\infty(Q)$ curves determined by computer simulations. We observe a positive dispersion of $c(Q)$ from $c_o(Q)$ (≈ 1500 m/s at $Q = 0$) to a value of ≈ 3200 m/s in the Q region $Q=2\div 4$ nm⁻¹. Extensive temperature and density dependent studies have

shown that this positive dispersion is associated to the α -process [1]. It is then tempting to associate the large difference between $c_{\infty\alpha}$ and $c_{\infty}(Q)$, e. g. between ≈ 3200 and ≈ 5000 m/s at $Q=6$ nm $^{-1}$, to the μ -process. This observation, evident in the low Q region, is less obvious at large Q where the difference between $c_{\infty\alpha}$ and $c_{\infty}(Q)$ is much smaller.

To clarify whether a μ -process can be clearly identified in a hydrogen bonded liquid we studied hydrogen fluoride. We report the first experimental data of the $S(Q, \omega)$ of HF, and compare them to existing simulation data [8]. The picture emerging from our analysis shows that in HF both the α - and μ - process are active, and they can be clearly identified because their relative strength is substantially weaker than in liquid water. This study confirms, therefore, that these two relaxation processes are active also in hydrogen bonded liquids.

The $S(Q, \omega)$ of HF has been studied at T=239 K by Inelastic X-ray Scattering (IXS) as a function of wave vector Q in the range $2 \div 31$ nm $^{-1}$. The HF sample (≈ 1 cm 3) was confined in a stainless steel cell equipped with two sapphire windows (total thickness 500 μ m) to allow the passage of the incident and scattered x-rays. The experiment has been carried out at the very high energy resolution IXS beam-line ID16 at the European Synchrotron Radiation Facility. The instrument consists of a back-scattering monochromator and five independent analyzer systems, held one next to each other with a constant angular offset on a 6.5 m long analyzer arm. The used Si(11 11 11) configuration [9], gives an instrumental energy resolution of 1.6 meV full width half maximum (FWHM) and an offset of 3 nm $^{-1}$ between two neighbor analyzers. The momentum transfer is selected by rotating the analyzer arm. The spectra at constant Q and as a function of energy were measured with a Q resolution of 0.4 nm $^{-1}$ FWHM. The energy scans were performed varying the monochromator temperature with respect to that of the analyzer crystals. Further details on the beam-line are reported elsewhere [10]. Each scan took about 180 min and each spectrum at fixed Q was obtained by summing up to 6 scans.

In Fig. 2(a) we report selected spectra at the indicated momentum transfer and the corresponding measured resolution functions aligned and scaled to the central peak. The contribution of the empty cell was found to be negligible. The spectra consist of a band centered at zero energy transfer which becomes broader with increasing Q and with a characteristic asymmetry due to the detailed balance. A preliminary determination of $c(Q)$ was obtained fitting the spectra to the convolution of the resolution function with a model function for $S(Q, \omega)$ composed by a Lorentzian for the central peak ($S_o(Q, \omega)$) and a Damped Harmonic Oscillator (DHO) [11] for the inelastic signal ($S_i(Q, \omega)$). The deconvoluted inelastic part of the current spectra, $\omega^2/Q^2 S_i(Q, \omega)$, -whose maxima correspond to the parameter $\Omega(Q)$ - is reported in Fig.2b.

The dispersion curve $\Omega(Q)$ vs. Q is shown in Fig. 3 for the low Q region data. For Q between 4 and 7 nm $^{-1}$, it shows a linear dependence with a slope corresponding to a sound speed of 1080 m/s. This value is substantially higher than the adiabatic sound speed $c_o = 580$ m/s as obtained from Brillouin light scattering [12]. Moreover, in the 2 to 4 nm $^{-1}$, the IXS data are compatible with a transition of $c(Q)$ from its low frequency value c_o to the higher value. This increase of $c(Q)$ is then interpreted as due to the α -relaxation, and it comes out to be quite similar to that of water, where the ratio $c_{\infty\alpha}/c_o$ was also close to two. A more formal procedure to describe the effect of a relaxation process in the $S(Q, \omega)$ is based on the viscoelastic model. In this approach the $S(Q, \omega)$ is expressed as:

$$S(Q, \omega) = I(Q) \frac{\omega_0(Q)^2 M'(Q, \omega)}{[\omega^2 - \omega_0(Q)^2 - \omega M''(Q, \omega)]^2 + [\omega M'(Q, \omega)]^2} \quad (1)$$

where $\omega_0(Q)^2 = (K_B T / m S(Q)) Q^2$ is the normalized second frequency moment of $S(Q, \omega)$, K_B is the Boltzmann constant, m is the mass of the molecule and $M'(Q, \omega)$, $M''(Q, \omega)$ are respectively the real and the imaginary part of the Laplace transform of the memory function $M(Q, t)$. In the two relaxation processes scenario, we model $M(Q, t)$ by the sum of two exponential decay contributions:

$$M(Q, t) = \Delta_\alpha^2(Q) e^{-t/\tau_\alpha(Q)} + \Delta_\mu^2(Q) e^{-t/\tau_\mu(Q)} \quad (2)$$

where $\Delta_\alpha^2(Q) = [c_{\infty\alpha}(Q)^2 - c_o(Q)^2] Q^2$, and $\Delta_\mu^2(Q) = [c_\infty(Q)^2 - c_{\infty\alpha}(Q)^2] Q^2$ are the strengths of the two processes. As, similarly to water, one expects that the μ -process is very fast with respect to the investigated timescale [13], the second term is approximated by a δ -function:

$$M(Q, t) = \Delta_\alpha^2(Q) e^{-t/\tau_\alpha(Q)} + \Gamma_\mu(Q) \delta(t) \quad (3)$$

with $\Gamma_\mu(Q) = \Delta_\mu^2 \tau_\mu(Q)$. The experimental data were fitted to the convolution of the experimental resolution function with the dynamic structure factor model given by Eq. 1. The relevant independent parameters are $c_{\infty\alpha}(Q)$, $c_o(Q)$, $\tau_\alpha(Q)$ and $\Gamma_\mu(Q)$. In Fig. 4 we show the values obtained for $c_\infty(Q)$ and $c_o(Q)$, together with those for $c(Q)$ as deduced from the DHO model. We observe that the positive dispersion found for $c(Q)$ by the DHO analysis takes place between the values of $c_o(Q)$ and $c_{\infty\alpha}(Q)$, derived from the viscoelastic analysis. Therefore, this finding confirms the hypothesis that the transition of $c(Q)$ is governed by the α - process. In particular, for Q larger than 4 nm $^{-1}$, the coincidence of $c(Q)$ and $c_{\infty\alpha}(Q)$ tells us that the α - process is fully accomplished for Q larger than 4 nm $^{-1}$, and, in the explored Q region,

there is no evidence for a further dispersion of $c(Q)$ that could be associated to the μ - process. The fit provides $\tau_\alpha(Q)$ to be 0.30 ± 0.05 ps at low Q , a value consistent with the hydrodynamic limit $\tau(0) = 0.25 \pm 0.05$ ps as calculated through the relation [13]:

$$\tau(0) = \frac{\nu_L - \frac{\Gamma_\mu(0)}{2Q^2}}{c_{\infty\alpha}^2(0) - c_0^2(0)} \quad (4)$$

where ν_L and c_0 are respectively the kinematic longitudinal viscosity and the adiabatic sound velocity measured by Brillouin light scattering [12]. This numerical equivalence gives further support to the validity of the employed viscoelastic model.

Recent molecular dynamics simulation studies on HF [8, 14, 15] provide the necessary information to discuss the present experimental results in the framework of the two relaxation processes scenario. Similarly to Fig. 1 for liquid water, in Fig. 5 we report the sound velocities of Fig. 4 together with numerical simulation results [8]: *i*) $c_o(Q)$ as derived from the simulated static structure factor, *ii*) $c(Q)$ as derived from the maxima of the simulated longitudinal current spectra, and *iii*) $c_\infty(Q)$ as derived from the fourth moment of the dynamic structure factor. It is worth to note that the quantity $c_{\infty\alpha}(Q)$ does not have a simple expression in terms of microscopic variables, and cannot be directly evaluated numerically. In spite of the slightly different thermodynamic points between the experiment ($T=239$ K) and the simulation ($T=203$ K), we observe an excellent agreement between the two common sets of data, i. e. for $c_o(Q)$ and $c(Q)$. This agreement implies that the interaction potential model used in the simulation matches well the properties of the real system or that these quantities are relatively insensitive to the interaction potential. The important information emerging from the comparison of the data in Fig. 5, is the very large difference between the (measured) $c_{\infty\alpha} = c_{o\mu}$ and the (calculated) c_∞ . This, in turn, implies not only the existence of the μ -process, but also that, in HF, this process has a relative strength substantially larger than in water over the whole considered Q range, which extends beyond the first peak in the static structure factor.

In conclusion we provide a strong indication that, similarly to simple and glass-forming liquids, also hydrogen bonded liquids present *two* relaxation processes affecting their high frequency collective dynamics. These two processes produce a phenomenology consistent with the α - and μ - processes, thus suggesting their universality in the liquid state. The strengths of these two processes are,

however, dependent on the specific system. In particular - contrary to simple liquids where $c_\infty/c_o \approx 1.2$ and $c_{\infty\alpha}$ differs from c_o by few percent [5] - the strengths of these relaxation processes are much larger in hydrogen bonded liquids: at low Q in both water and HF $c_{\infty\alpha}/c_o \approx 2$, while $c_\infty/c_o \approx 3$ in water and ≈ 7 in HF. It is intriguing to understand the origin of these quantitative differences, and to see whether they are correlated the different hydrogen bond networks existing in water and HF.

We acknowledge D. Fioretto for his help during the Brillouin light scattering measurements, C. Henriquet for the design, development and assembly of the hydrogen-fluoride cell, C. Lapras for technical help and R. Vallauri for useful discussions.

-
- [1] G. Harrison, *The Dynamic Properties of Supercooled Liquid* (Academic Press - London, New York, San Francisco, 1976).
 - [2] U. Balucani and M. Zoppi, *Dynamics of the Liquid State* (Clarendon Press - Oxford, 1994).
 - [3] A. Rahman, Phys. Rev. A **9**, 1667 (1974).
 - [4] R. D. Mountain, Phys. Rev. A **26**, 2859 (1982).
 - [5] T. Scopigno, U. Balucani, G. Ruocco, and F. Sette, J. Phys. C. **12**, 8009 (2000), and references therein.
 - [6] W. Götze and M. R. Mayr, Phys. Rev. E **61**, 587 (2000).
 - [7] This ordering is based on the observation that τ_α is always larger than τ_μ .
 - [8] D. Bertolini, G. Sutmann, A. Tani, and R. Vallauri, Phys. Rev. Lett. **81**, 2080 (1998).
 - [9] C. Masciovecchio, U. Bergmann, M. Krisch, G. Ruocco, F. Sette, and R. Verbeni, Nucl. Instrum. Methods Phys. Res. B **117**, 339 (1996).
 - [10] C. Masciovecchio, U. Bergmann, M. Krisch, G. Ruocco, F. Sette, and R. Verbeni, Nucl. Instrum. Methods Phys. Res. B **111**, 181 (1996).
 - [11] B. Fak and B. Dorner (1992), institute Laue Langevin (Grenoble, France) Technical Report ILL 92FA008S1992.
 - [12] R. Angelini *et al*, to be published.
 - [13] G. Monaco, A. Cunsolo, G. Ruocco, and F. Sette, Phys. Rev. E **60**, 5505 (1999).
 - [14] U. Balucani, D. Bertolini, A. Tani, and R. Vallauri, J. Chem. Phys. **112**, 9025 (2000).
 - [15] G. Garberoglio and R. Vallauri, Phys. Rev. Lett. **84**, 4878 (2000).
 - [16] M. Sampoli, G. Ruocco, and F. Sette, Phys. Rev. Lett. **79**, 1678 (1997).
 - [17] F. Sette, G. Ruocco, M. Krisch, C. Masciovecchio, R. Verbeni, and U. Bergmann, Phys. Rev. Lett. **77**, 83 (1996).
U. Balucani, G. Ruocco, A. Torcini, and R. Vallauri, Phys. Rev. E **47**, 1677 (1993).




FIG. 1: Q dependence of experimental and theoretical velocities of sound in water. Open circles, MD simulations [16]; full squares, circles and diamonds, IXS data [17]; solid (dashed) line, zero (infinite) frequency limit [?].




FIG. 2: *a*) IXS spectra of HF at fixed temperature $T = 239K$ and at the indicated moment transfer Q plotted together with the corresponding experimental resolutions (solid lines). *b*) Resolution deconvoluted inelastic part (DHO) of the longitudinal spectral current.




FIG. 3: Dispersion curve obtained from the maxima of the current spectra shown in Fig. 2. The upper dashed line is the linear fit to the data, the lower dashed line indicates the adiabatic sound velocity as measured by Brillouin light scattering [12].




FIG. 4: Q -dependence of the sound velocities $c_0(Q)$ (open diamonds) and $c_{\infty\alpha}(Q)$ (open circles) from the viscoelastic fit, and $c(Q)$ from Fig. 3. The value of the adiabatic sound velocity c_0 is indicated by the arrow [12].




FIG. 5: Q -dependence of the sound velocities at $T=239$ K from this work: $c_0(Q)$ (open diamonds), $c_{\infty\alpha}$ (open circles), $c(Q)$ (full circles) and at $T=203$ K from the MD results of Ref. [8]: $c_0(Q)$ (solid line), $c_{\infty\mu}$ (dashed line), $c(Q)$ (stars).