

Article

Nanozyme-based biosensing for clinical diagnosis of COVID-19: A mini review

Saeed Reza Hormozi Jangi

Hormozi Laboratory of Chemistry and Biochemistry, Zabol 9861334367, Iran; E-mail: saeedrezahormozi@gmail.com (S.R. Hormozi Jangi)

Abstract

Several clinical methods had been utilized for diagnosis of COVID-19 for instance, real-time reverse transcription-polymerase chain reaction (rRT-PCR), hematology examination, polymerase chain reaction (PCR), diagnostic guidelines based on clinical features, and Chest CT scans. However, the accurate current methods are time-consuming and expensive and other methods are inaccurate. To solve these drawbacks, nanozyme-based sensors have been developed for the reliable, accurate, and rapid detection of SARS-CoV-2. The main basis of these sensors is the detection of color variation of a nanozyme-mediated oxidation reaction in the presence and the absence of antigens of COVID-19. Besides, some of methods are based on probing the fluorescence of these systems as the clinical signal toward detection of SARS-CoV-2. This mini review focused on overviewing the nanozymes-based methods toward COVID-19 diagnosis. The historical background of COVID-19 was reviewed. Thereafter, the biomedical applications of nanozymes was discussed and finally, the recent progress of early diagnosis of COVID-19 based on nanozymatic systems was briefly reviewed.

Keywords: COVID-19; Nanozymes; COVID-19diagnosis; Nanozyme-based sensors

1. Introduction

On December 31, 2019, the first case of a novel infectious disease with unknown origin (causative agent), features, duration of human transmission, and epidemiological parameters was confirmed in a designated hospital in Wuhan, a major city in China [1, 2]. The studies on this new infectious disease revealed that a new generation of coronavirus, SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2), is its causative agent [3-5]. Coronaviruses are a group of Coronaviridae families with a broad distribution in mammals which are known as the non-segmented positive-sense RNA viruses [6]. This novel disease caused by SARS-CoV-2 was called Coronavirus disease 2019 and termed COVID-19 by WHO on 11 Feb 2020 [7]. Although the human infections resulting from coronavirus are mild in most cases [6], shortly after the first report of COVID-19, the novel COVID-19 exhibited a high potential for outbreaks and becoming an epidemic disease and even a pandemic, as now we see in the world [8-10]. Currently, there are several methods for diagnosis of COVID-19 including real-time reverse transcription-polymerase chain reaction (rRT-PCR), hematology examination, polymerase chain reaction (PCR), diagnostic guidelines based on clinical features, Chest CT scans [1]. Here, it is challenging to develop effective diagnostics and therapeutics against SARS-CoV-2 [11]. Since the first report of COVID-19 in 2019, nanozymes-based systems have been applied for reliable fast diagnosis of COVID-19. Hence, this mini

review focused on overviewing the the nanozyme-based methods toward COVID-19 diagnosis.

2. Nanozymes and their biomedical applications

The fast development of nanoscience and material chemistry has increased interest in researching new and innovative synthesis methods to produce new nanomaterials with unique catalytic activity [12, 13], unique optical properties [14-16], high active area [17], antibacterial properties [18], and high biocompatibility [19]. The new field of nanozyme-based catalysis, which has been introduced as an alternative to enzyme-based catalysis, is called nanozyme chemistry. On the other hand, nanozymes are known as nanomaterials with high enzyme-like activity and can be used to simulate enzymatic reactions in harsh environmental conditions (for example, higher temperature or wider pH range) [20-27]. As previously reported in the literature [27], native enzymes, for instance, native peroxidases or ureases suffer from several disadvantages and drawbacks such as low pH stability, low thermal stability, low recoverability, and no reusability. Commonly, to solve these difficulties and drawbacks of native enzymes, the development of enzyme immobilization protocols has been widely considered in the literature [27-30]. Hence to solve these difficulties, the design and development of low-cost nanozymes were considered as an interesting way for performing enzyme-catalyzed reactions in harsh conditions [21, 31]. Nanozymes have been used for several applications in catalysis [32], biomedical imaging [33], tumor therapy [34, 35], and sensing and

detection [36-38]. For instance, up to date, different types of nanozyme-based sensors such as single nanozymatic sensors, enzyme-nanozyme hybrid sensors, etc. have been developed [39]. Recently a new generation of nanozyme-based systems called “multinanozyme system” was introduced by Hormozi Jangi et al. (2020) [40, 41]. During the last years, a wide variety of nanozyme-based colorimetric sensors have been developed for the detection and quantification of a variety of analytes for instance, tryptophan [42], glutathione (GSH) [43], dopamine [44], tetracycline [45], metal cations [46], glucose [47], H₂O₂ [48], explosives [49], and cysteine [50]. Besides, some of the nanozyme-based sensors with fluorescence-based response had been developed and utilized for detecting several analytes [51, 52].

3. Nanozymes application for diagnosis of COVID-19

There is several reports in the literature regarding application of nanozymes for diagnosis of COVID-19. For example, Liang et al. (2021) [53] developed a nanozyme-linked nanosensor for the rapid and quantitative diagnosis of COVID-19 by detecting the SARS-CoV-2 nucleocapsid protein in human blood. Besides, Fu et al. (2021) [54] used porous metallic gold@platinum nanozymes for the diagnosis of COVID-19 via colorimetric detection of spike (S1) protein of SARS-CoV-2, obtaining a wide linear working range over 10–100 ng mL⁻¹ along with a low limit of detection (LOD) of 11 ng mL⁻¹.

In 2021, Liu et al. [55] developed a paper-based chemiluminescence nanozymatic strip test for sensitive detection of SARS-CoV-2 antigen which the core of their paper test was a Co-Fe@hemin-peroxidase nanozyme that can catalyze chemiluminescence and amplify immune reaction signal. In addition, Liu et al. (2021) [56] introduced a smartphone-based nanozyme-linked immunosorbent assay for quantitative sensing of SARS-CoV-2 nucleocapsid phosphoprotein in 37 serum samples from 20 patients infected with COVID-19. In 2022, Ali and Omer [57] introduced an ultrasensitive aptamer-functionalized Cu-MOF fluorescent nanozyme and utilized it for optical detection of C-reactive protein toward the diagnosis of COVID-19 via colorimetric and fluorometric dual mode responses of a nanozymatic process based on TMB oxidation for colorimetric and variation of fluorescence intensity of Cu-MOF for fluorometric mode detection of COVID-19. Moreover, Zhao et al. (2022) [58] employed MIL-101(CuFe) nanozymes for accurate visual naked-eye diagnosis of COVID-19 via detecting the universal receptor of CD147, providing a very low detection limit of 3 PFU/mL and a detection time as short as 30 min. Besides, Wu et al. (2022) [59] developed a MnO₂ nanozyme-mediated CRISPR-Cas12a system for naked-eye diagnosis of COVID-19. In this system, the MnO₂ nanorods were initially linked to magnetic beads using a single-stranded DNA (ssDNA). Also, in 2023, He et al. [60] performed a nanozyme-based colorimetric method for naked-eye diagnosis of COVID-19 by iron manganese silicate nanozymes as peroxidase-like nanozymes.

5. Conclusions

Several clinical methods had been utilized for diagnosis of COVID-19 for instance, real-time reverse transcription-polymerase chain reaction (rRT-PCR), hematology examination, polymerase chain reaction (PCR), diagnostic guidelines based on clinical features, and Chest CT scans. However, the accurate current methods are time-consuming and expensive and other methods are inaccurate. To solve these drawbacks, nanozyme-based sensors have been developed for the reliable, accurate, and rapid detection of SARS-CoV-2. The main basis of these sensors is the detection of color variation of a nanozyme-mediated oxidation reaction in the presence and the absence of antigens of COVID-19. Besides, some of methods are based on probing the fluorescence of these systems as the clinical signal toward detection of SARS-CoV-2. This mini review focused on overviewing the nanozymes-based methods toward COVID-19 diagnosis. The historical background of COVID-19 was reviewed. Thereafter, the biomedical applications of nanozymes was discussed and finally, the recent progress of early diagnosis of COVID-19 based on nanozymatic systems was briefly reviewed.

Acknowledgments

The authors gratefully thank the Hormozi Laboratory of Chemistry and Biochemistry (Zabol, Iran) for the support of this work.

Conflict of interest

None.

6. References

1. Hormozi Jangi, S. R. (2023). A Brief Overview on Clinical and Epidemiological Features, Mechanism of Action, and Diagnosis of Novel Global Pandemic Infectious Disease, Covid-19, And its Comparison with Sars, Mers, And H1n1. *World J Clin Med Img*, 2(1), 45-52.
2. Jangi, S. R. H. (2023). Natural Polyphenols of Pomegranate and Black Tea Juices can Combat COVID-19 through their SARS-CoV-2 3C-like Protease-inhibitory Activity. *Qeios*.
3. Hao, W., & Li, M. (2020). Clinical diagnostic value of CT imaging in COVID-19 with multiple negative RT-PCR testing. *Travel medicine and infectious disease*, 34, 101627.
4. Li, R., Tian, J., Yang, F., Lv, L., Yu, J., Sun, G., ... & Ding, J. (2020). Clinical characteristics of 225 patients with COVID-19 in a tertiary Hospital near Wuhan, China. *Journal of Clinical Virology*, 127, 104363.
5. Tang, X., Du, R. H., Wang, R., Cao, T. Z., Guan, L. L., Yang, C. Q., ... & Shi, H. Z. (2020). Comparison of hospitalized patients with ARDS caused by COVID-19 and H1N1. *Chest*, 158(1), 195-205.
6. Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., ... & Cao, B. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The lancet*, 395(10223), 497-506.

7. <https://www.who.int/emergencies/diseases/novelcoronavirus-2019>.
8. Ye, G., Li, Y., Lu, M., Chen, S., Luo, Y., Wang, S., ... & Wang, X. (2020). Experience of different upper respiratory tract sampling strategies for detection of COVID-19. *Journal of Hospital Infection*, 105(1), 1-2.
9. Wu, Z., & McGoogan, J. M. (2020). Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *Jama*, 323(13), 1239-1242.
10. Hill, M. A., Mantzoros, C., & Sowers, J. R. (2020). Commentary: COVID-19 in patients with diabetes. *Metabolism-Clinical and Experimental*, 107.
11. Verma, M. K., Sharma, P. K., Verma, H. K., Singh, A. N., Singh, D. D., Verma, P., & Siddiqui, A. H. (2021). Rapid diagnostic methods for SARS-CoV-2 (COVID-19) detection: an evidence-based report. *Journal of Medicine and Life*, 14(4), 431.
12. Hormozi Jangi, S. R. (2023). Low-temperature destructive hydrodechlorination of long-chain chlorinated paraffins to diesel and gasoline range hydrocarbons over a novel low-cost reusable ZSM-5@ Al-MCM nanocatalyst: a new approach toward reuse instead of common mineralization. *Chemical Papers*, 1-15.
13. HORMOZI JANGI, S. R., & Akhond, M. (2020). High throughput green reduction of tris (p-nitrophenyl) amine at ambient temperature over homogenous AgNPs as H-transfer catalyst. *Journal of Chemical Sciences*, 132, 1-8.

14. Dehghani, Z., Akhond, M., Jangi, S. R. H., & Absalan, G. (2024). Highly sensitive enantioselective spectrofluorimetric determination of R-/S-mandelic acid using l-tryptophan-modified amino-functional silica-coated N-doped carbon dots as novel high-throughput chiral nanoprobe. *Talanta*, 266, 124977.
15. Hormozi Jangi, S. R., & Gholamhosseinzadeh, E. (2023). Developing an ultra-reproducible and ultrasensitive label-free nanoassay for L-methionine quantification in biological samples toward application in homocystinuria diagnosis. *Chemical Papers*, 1-13.
16. Jangi, S. R. H., & Akhond, M. (2021). Ultrasensitive label-free enantioselective quantification of d-/l-leucine enantiomers with a novel detection mechanism using an ultra-small high-quantum yield N-doped CDs prepared by a novel highly fast solvent-free method. *Sensors and Actuators B: Chemical*, 339, 129901.
17. Thakkar, K. N., Mhatre, S. S., & Parikh, R. Y. (2010). Biological synthesis of metallic nanoparticles. *Nanomedicine: nanotechnology, biology and medicine*, 6(2), 257-262.
18. Hajipour, M. J., Fromm, K. M., Ashkarran, A. A., de Aberasturi, D. J., de Larramendi, I. R., Rojo, T., ... & Mahmoudi, M. (2012). Antibacterial properties of nanoparticles. *Trends in biotechnology*, 30(10), 499-511.

19. Hormozi Jangi, S. R. (2023). Synthesis and characterization of magnesium-based metal-organic frameworks and investigating the effect of coordination solvent on their biocompatibility. *Chemical Research and Nanomaterials*, 1(4), 1-9.
20. Hormozi Jangi, S. R. (2023). Biochemical characterization of enzyme-like silver nanoparticles toward nanozyme-catalysed oxidation reactions, *Micromaterials and Interfaces* 1 (1), 2170.
21. Hormozi Jangi, S. R. (2023). Evaluation of Biochemical Behavior and Stability of Gold Nanoparticles with High Intrinsic Peroxidase-Like Activity. *Petro Chem Indus Intern*, 6(4), 234-239.
22. Jangi, S. R. H. (2023). Introducing a High Throughput Nanozymatic Method for Eco-Friendly Nanozyme-Mediated Degradation of Methylene Blue in Real Water Media. *Sustainable Chemical Engineering*, 90-99.
23. Hormozi Jangi, S. R., & Dehghani, Z. (2023). Kinetics and biochemical characterization of silver nanozymes and investigating impact of storage conditions on their activity and shelf-life. *Chemical Research and Nanomaterials*, 1(4), 25-33.
24. Jangi, S. R. H. (2023). Determining kinetics parameters of bovine serum albumin-protected gold nanozymes toward different substrates. *Qeios*.

25. Hormozi Jangi, S. R. (2023). Effect of daylight and air oxygen on nanozymatic activity of unmodified silver nanoparticles: Shelf-stability. *Qeios*. doi: 10.32388/9RWVYI.3
26. Ahmadi-Leilakouhi, B., Hormozi Jangi, S. R., & Khorshidi, A. (2023). Introducing a novel photo-induced nanozymatic method for high throughput reusable biodegradation of organic dyes. *Chemical Papers*, 77(2), 1033-1046.
27. Hormozi Jangi, S. R. (2023). A Comparative Study on Kinetics Performances of BSA-gold Nanozymes for Nanozyme-mediated Oxidation of 3,3',5,5'-Tetramethylbenzidine and 3,3'-Diaminobenzidine. *Biochemistry & Molecular Biology Journal* 9(3), 21.
28. Jangi, S. R. H., & Akhond, M. (2021). High throughput urease immobilization onto a new metal-organic framework called nanosized electroactive quasi-coral-340 (NEQC-340) for water treatment and safe blood cleaning. *Process Biochemistry*, 105, 79-90.
29. Jangi, S. R. H., & Akhond, M. (2022). Introducing a covalent thiol-based protected immobilized acetylcholinesterase with enhanced enzymatic performances for biosynthesis of esters. *Process Biochemistry*, 120, 138-155.
30. Jangi, S. R. H., Akhond, M., & Dehghani, Z. (2020). High throughput covalent immobilization process for improvement of shelf-life, operational cycles, relative

activity in organic media and enzymatic kinetics of urease and its application for urea removal from water samples. *Process Biochemistry*, 90, 102-112.

31. Wang, Q., Wei, H., Zhang, Z., Wang, E., & Dong, S. (2018). Nanozyme: An emerging alternative to natural enzyme for biosensing and immunoassay. *TrAC Trends in Analytical Chemistry*, 105, 218-224.
32. Lu, W., Guo, Y., Zhang, J., Yue, Y., Fan, L., Li, F., ... & Shuang, S. (2022). A High Catalytic Activity Nanozyme Based on Cobalt-Doped Carbon Dots for Biosensor and Anticancer Cell Effect. *ACS Applied Materials & Interfaces*, 14(51), 57206-57214.
33. Ren, X., Chen, D., Wang, Y., Li, H., Zhang, Y., Chen, H., ... & Huo, M. (2022). Nanozymes-recent development and biomedical applications. *Journal of Nanobiotechnology*, 20(1), 92.
34. Tang, G., He, J., Liu, J., Yan, X., & Fan, K. (2021, August). Nanozyme for tumor therapy: Surface modification matters. In *Exploration* (Vol. 1, No. 1, pp. 75-89).
35. Li, Menghuan, Hui Zhang, Yanhua Hou, Xuan Wang, Chencheng Xue, Wei Li, Kaiyong Cai, Yanli Zhao, and Zhong Luo. "State-of-the-art iron-based nanozymes for biocatalytic tumor therapy." *Nanoscale Horizons* 5, no. 2 (2020): 202-217.
36. Yu, R., Wang, R., Wang, Z., Zhu, Q., & Dai, Z. (2021). Applications of DNA-nanozyme-based sensors. *Analyst*, 146(4), 1127-1141.

37. Chang, Y., Gao, S., Liu, M., & Liu, J. (2020). Designing signal-on sensors by regulating nanozyme activity. *Analytical Methods*, 12(39), 4708-4723.
38. Arshad, F., Mohd-Naim, N. F., Chandrawati, R., Cozzolino, D., & Ahmed, M. U. (2022). Nanozyme-based sensors for detection of food biomarkers: a review. *RSC advances*, 12(40), 26160-26175.
39. Jangi, A. R. H., Jangi, M. R. H., & Jangi, S. R. H. (2020). Detection mechanism and classification of design principles of peroxidase mimic based colorimetric sensors: A brief overview. *Chinese Journal of Chemical Engineering*, 28(6), 1492-1503.
40. Jangi, S. R. H., Akhond, M., & Absalan, G. (2020). A novel selective and sensitive multinanozyme colorimetric method for glutathione detection by using an indamine polymer. *Analytica Chimica Acta*, 1127, 1-8.
41. Jangi, S. R. H., Davoudli, H. K., Delshad, Y., Jangi, M. R. H., & Jangi, A. R. H. (2020). A novel and reusable multinanozyme system for sensitive and selective quantification of hydrogen peroxide and highly efficient degradation of organic dye. *Surfaces and Interfaces*, 21, 100771.
42. Xu, S., Zhang, S., Li, Y., & Liu, J. (2023). Facile Synthesis of Iron and Nitrogen Co-Doped Carbon Dot Nanozyme as Highly Efficient Peroxidase Mimics for Visualized Detection of Metabolites. *Molecules*, 28(16), 6064.
43. Jangi, S. R. H., & Akhond, M. (2020). Synthesis and characterization of a novel metal-organic framework called nanosized electroactive quasi-coral-340 (NEQC-

- 340) and its application for constructing a reusable nanozyme-based sensor for selective and sensitive glutathione quantification. *Microchemical Journal*, 158, 105328.
44. Ray, S., Biswas, R., Banerjee, R., & Biswas, P. (2020). A gold nanoparticle-intercalated mesoporous silica-based nanozyme for the selective colorimetric detection of dopamine. *Nanoscale Advances*, 2(2), 734-745.
45. Shen, Y., Wei, Y., Liu, Z., Nie, C., & Ye, Y. (2022). Engineering of 2D artificial nanozyme-based blocking effect-triggered colorimetric sensor for onsite visual assay of residual tetracycline in milk. *Microchimica Acta*, 189(6), 233.
46. Akhond, M., Hormozi Jangi, S. R., Barzegar, S., & Absalan, G. (2020). Introducing a nanozyme-based sensor for selective and sensitive detection of mercury (II) using its inhibiting effect on production of an indamine polymer through a stable n-electron irreversible system. *Chemical Papers*, 74, 1321-1330.
47. Chen, J., Wu, W., Huang, L., Ma, Q., & Dong, S. (2019). Self-indicative gold nanozyme for H₂O₂ and glucose sensing. *Chemistry–A European Journal*, 25(51), 11940-11944.
48. Hormozi Jangi, S. R., & Dehghani, Z. (2023). Spectrophotometric quantification of hydrogen peroxide utilizing silver nanozyme. *Chemical Research and Nanomaterials*, 2(1), 15-23.

49. Hormozi Jangi, S. R., Akhond, M., & Absalan, G. (2020). A field-applicable colorimetric assay for notorious explosive triacetone triperoxide through nanozyme-catalyzed irreversible oxidation of 3, 3'-diaminobenzidine. *Microchimica Acta*, 187, 1-10.
50. Singh, M., Weerathunge, P., Liyanage, P. D., Mayes, E., Ramanathan, R., & Bansal, V. (2017). Competitive inhibition of the enzyme-mimic activity of Gd-based nanorods toward highly specific colorimetric sensing of l-cysteine. *Langmuir*, 33(38), 10006-10015.
51. Wang, M., Zhang, J., Zhou, X., Sun, H., & Su, X. (2022). Fluorescence sensing strategy for xanthine assay based on gold nanoclusters and nanozyme. *Sensors and Actuators B: Chemical*, 358, 131488.
52. Heo, N. S., Song, H. P., Lee, S. M., Cho, H. J., Kim, H. J., Huh, Y. S., & Kim, M. I. (2020). Rosette-shaped graphitic carbon nitride acts as a peroxidase mimic in a wide pH range for fluorescence-based determination of glucose with glucose oxidase. *Microchimica Acta*, 187, 1-11.
53. Liang, C., Liu, B., Li, J., Lu, J., Zhang, E., Deng, Q., ... & Li, T. (2021). A nanoenzyme linked immunochromatographic sensor for rapid and quantitative detection of SARS-CoV-2 nucleocapsid protein in human blood. *Sensors and Actuators B: Chemical*, 349, 130718.

54. Fu, Z., Zeng, W., Cai, S., Li, H., Ding, J., Wang, C., ... & Yang, R. (2021). Porous Au@ Pt nanoparticles with superior peroxidase-like activity for colorimetric detection of spike protein of SARS-CoV-2. *Journal of colloid and interface science*, 604, 113-121.
55. Liu, D., Ju, C., Han, C., Shi, R., Chen, X., Duan, D., ... & Yan, X. (2021). Nanozyme chemiluminescence paper test for rapid and sensitive detection of SARS-CoV-2 antigen. *Biosensors and Bioelectronics*, 173, 112817.
56. Liu, B., Wu, Z., Liang, C., Lu, J., Li, J., Zhang, L., ... & Li, C. (2021). Development of a smartphone-based nanozyme-linked immunosorbent assay for quantitative detection of SARS-CoV-2 nucleocapsid phosphoprotein in blood. *Frontiers in Microbiology*, 12, 692831.
57. Ali, G. K., & Omer, K. M. (2022). Ultrasensitive aptamer-functionalized Cu-MOF fluorescent nanozyme as an optical biosensor for detection of C-reactive protein. *Analytical Biochemistry*, 658, 114928.
58. Zhao, X., Yang, Z., Niu, R., Tang, Y., Wang, H., Gao, R., ... & Meng, L. (2022). MIL-101 (CuFe) Nanozymes with Excellent Peroxidase-like Activity for Simple, Accurate, and Visual Naked-Eye Detection of SARS-CoV-2. *Analytical Chemistry*, 95(2), 1731-1738.

59. Wu, L., Wang, X., Wu, X., Xu, S., Liu, M., Cao, X., ... & Huang, H. (2022). MnO₂ nanozyme-mediated CRISPR-Cas12a system for the detection of SARS-CoV-2. *ACS Applied Materials & Interfaces*, 14(45), 50534-50542.
60. He, M., Xu, X., Wang, H., Wu, Q., Zhang, L., Zhou, D., ... & Liu, H. (2023). Nanozyme-Based Colorimetric SARS-CoV-2 Nucleic Acid Detection by Naked Eye. *Small*, 2208167.