

Identifiable specializations for ODE models

Alexey Ovchinnikov^a, Anand Pillay^b, Gleb Pogudin^c, Thomas Scanlon^d

^a*CUNY Queens College, Department of Mathematics, 65-30 Kissena Blvd, Queens, NY 11367, USA
CUNY Graduate Center, Mathematics and Computer Science, 365 Fifth Avenue, New York, NY 10016, USA*

^b*University of Notre Dame, Department of Mathematics, Notre Dame, IN 46556, USA*

^c*LIX, CNRS, École Polytechnique, Institut Polytechnique de Paris, 1 rue Honoré d'Estienne d'Orves, 91120, Palaiseau, France*

^d*University of California, Berkeley, Mathematics Department, Evans Hall, Berkeley, CA, 94720-3840*

Abstract

The parameter identifiability problem for a dynamical system is to determine whether the parameters of the system can be found from data for the outputs of the system. Verifying whether the parameters are identifiable is a necessary first step before a meaningful parameter estimation can take place. Non-identifiability occurs in practical models. To reparametrize a model to achieve identifiability is a challenge. The existing approaches have been shown to be useful for many important examples. However, these approaches are either limited to linear models and scaling parametrizations or are not guaranteed to find a reparametrization even if it exists. In the present paper, we prove that there always exists a locally identifiable model with the same input-output behaviour as the original one obtained from a given one by a partial specialization of the parameters. As an extra feature of our approach, the resulting (at least) locally identifiable reparameterization has the same shape: the monomials in the new state variables in the new model are formed in the same way as in the original model. Furthermore, we give a sufficient observability condition for the existence of a state space transformation from the original model to the new one. Our proof is constructive and can be translated to an algorithm, which we illustrate by several examples.

I. Introduction

A. Motivation

Scientists and engineers often model a process under investigation using a parametric ODE, which is a system of ordinary differential equations involving some unspecified parametric constants. The unknown parameters are usually determined (identified) from the input and measured output data. However, for some parametric ODEs, due to their intrinsic structure, it might not be possible to identify the parameters uniquely from the input and measured data (even noise-free). Therefore, while designing a model, it is crucial to make sure that the parametric ODE model is identifiable.

If the initially designed model has non-identifiable parameters, the next natural step would be to find another model with the same input-output behavior but with all parameters identifiable. This is a problem we study in the present paper.

B. Prior work

There exist efficient algorithms for finding reparametrizations of a specific form such as scaling transformations [12] or linear reparametrizations [15, 22]. More refined results have been obtained for scaling reparametrizations of linear compartmental models [1, 20]. Several approaches have been proposed

for producing locally identifiable reparametrizations [9, 16, 19] which succeed in finding nontrivial parametrizations for models from the literature but are not guaranteed to produce a reparametrization if it exists. The existence of an identifiable reparametrization was also not completely understood: Sussmann's theorem [30] guarantees the existence of an identifiable model with the same input-output behaviour at the cost of allowing the models to be defined on a manifold, in other words, if we allow to replace ODEs with differential-algebraic equations.

C. Our contribution

We prove that it is, in fact, always possible to replace the original ODE system with another one with the same input-output behaviour but all the parameters being locally identifiable by partially specializing parameters, in particular, without changing the “shape of the system” (Theorem 1). The latter means that

- the reparametrized system has the same number of equations and state variables as the old system and
- the monomials in the new system are obtained from the monomial of the old system by replacing the old state variable with the new state variables.

We also give an example showing that this statement is not true for global identifiability (see Section A). Under an additional observability condition, we also show that there exists a state space transformation between the original model and the new

Email addresses: aovchinnikov@qc.cuny.edu (Alexey Ovchinnikov), apillay@nd.edu (Anand Pillay), gleb.pogudin@polytechnique.edu (Gleb Pogudin), scanlon@math.berkeley.edu (Thomas Scanlon)

one (Theorem 1). Our proofs are constructive and can be directly translated into algorithms, which we showcase by several examples (Section IV). Our MAPLE code for these examples can be found here in [24].

II. Main result

A. Preliminaries and setup

In what follows, for a letter z , \bar{z} means that $\bar{z} = (z_1, \dots, z_n)$ is a tuple of length n (different tuples can have different lengths); we will write expressions like $\bar{z} \in A$ to mean that $z_i \in A$ for each component z_i of \bar{z} . Our main object will be an ODE system

$$\Sigma(\bar{\alpha}) := \begin{cases} \bar{x}'(t) = \bar{f}(\bar{x}(t), \bar{\alpha}, \bar{u}(t)) \\ \bar{y}(t) = \bar{g}(\bar{x}(t), \bar{\alpha}, \bar{u}(t)), \end{cases} \quad (1)$$

where $\bar{\alpha}$ is a vector of scalar parameters, $\bar{x}(t)$, $\bar{y}(t)$, and $\bar{u}(t)$ are the state, output, and input functions, respectively (in what follows we will omit the dependence on t for brevity). We will focus on rational ODE models, that is, the case of \bar{f} and \bar{g} being tuples of rational functions over \mathbb{Q} (or \mathbb{C}).

To formally define the main property of interest, input-output identifiability (IO-identifiability), we will introduce some notation from algebra.

1. A *differential ring* $(R, ')$ is a commutative ring with a derivation $' : R \rightarrow R$, that is, a map such that, for all $a, b \in R$, $(a + b)' = a' + b'$ and $(ab)' = a'b + ab'$.
2. The *ring of differential polynomials* in the variables x_1, \dots, x_n over a field K is the ring

$$K[x_j^{(i)} \mid i \geq 0, 1 \leq j \leq n]$$

with a derivation defined on the ring by $(x_j^{(i)})' := x_j^{(i+1)}$. This differential ring is denoted by $K\{x_1, \dots, x_n\}$.

3. For a differential polynomial $p \in K\{x_1, \dots, x_n\}$, its *order* in a variable x_i is the order of the highest derivative of x_i that appears in p . We denote this by $\text{ord}_{x_i} p$. The order of p in a tuple of variables \bar{x} is the maximal order of all x_i from \bar{x} that appear in p . We denote this by $\text{ord}_{\bar{x}} p$. When speaking about orders of equations, we will also mean orders of the corresponding differential polynomials.
4. An ideal I of a differential ring $(R, ')$ is called a *differential ideal* if, for all $a \in I$, $a' \in I$. For $F \subset R$, the smallest differential ideal containing the set F is denoted by $[F]$.
5. For an ideal I and element a in a ring R , we denote $I : a^\infty = \{r \in R \mid \exists \ell : a^\ell r \in I\}$. This set is also an ideal in R .
6. Given $\Sigma(\bar{\alpha})$ as in (1), we define the differential ideal of $\Sigma(\bar{\alpha})$ as

$$I_{\Sigma(\bar{\alpha})} := [Q\bar{x}' - Q\bar{f}, Q\bar{y} - Q\bar{g}] : Q^\infty \subset \mathbb{Q}(\bar{\alpha})\{\bar{x}, \bar{y}, \bar{u}\},$$

where Q is the common denominator of \bar{f} and \bar{g} . The relations between the inputs and outputs of the system can

be found by intersecting this ideal with the corresponding subring:

$$I_{\Sigma(\bar{\alpha})} \cap \mathbb{Q}(\bar{\alpha})\{\bar{y}, \bar{u}\}. \quad (2)$$

7. For a field K , let \bar{K} denote the algebraic closure of K .

Roughly speaking, input-output identifiability is a property for a parameter to be determined from inputs and outputs using IO-equations. Here is a precise formal definition:

- Definition 1** (IO-identifiability). 1. The smallest field k such that $\mathbb{Q} \subset k \subset \mathbb{Q}(\bar{\alpha})$ and $I_{\Sigma(\bar{\alpha})} \cap \mathbb{Q}(\bar{\alpha})\{\bar{y}, \bar{u}\}$ is generated (as an ideal or, equivalently, as a differential ideal) by $I_{\Sigma(\bar{\alpha})} \cap k\{\bar{y}, \bar{u}\}$ is called *the field of IO-identifiable functions*.
2. We call $h \in \mathbb{Q}(\bar{\alpha})$ *IO-identifiable* if $h \in k$. We also call $h \in \mathbb{Q}(\bar{\alpha})$ *locally IO-identifiable* if h is in the algebraic closure of the field k .
 3. The *IO-equations* are defined as the monic characteristic presentation of the differential ideal $I_{\Sigma(\bar{\alpha})} \cap \mathbb{Q}(\bar{\alpha})\{\bar{y}, \bar{u}\}$ (see [25, Definition 6 and Section 5.2] for more details). For a fixed differential ranking, such a monic characteristic presentation is unique [3, Theorem 3].

In many cases, IO-identifiability is equivalent to identifiability. See, e.g., a rigorously written definition of identifiability [10, Definition 2.5], [25, Section 4] for a sufficient condition for the equivalence, and [25, Examples 2.6 and 2.7] for simple examples of non-equivalence. Additionally, it turns out that IO-identifiability is equivalent to multi-experimental identifiability [23, Theorem 19]. Finally, several software packages check IO-identifiability [2, 17, 21, 28, 29] and find all IO-identifiable functions of the parameters [14].

Remark 1. Some authors prefer to work only with differential fields $(K, ')$ containing the field of complex numbers \mathbb{C} as a subfield of the field of constants, that is, the field of elements a of K satisfying $a' = 0$. With this convention, the field of IO-identifiable functions is taken to be the smallest field k for which $\mathbb{C} \subset k \subset \mathbb{C}(\bar{\alpha})$. For computational reasons, we prefer to work over \mathbb{Q} instead of \mathbb{C} . All of what we discuss in this paper may be generalized to the case of \mathbb{C} as the base with no essential changes to the arguments.

B. Statement of the main result

Theorem 1. *For every ODE system $\Sigma(\bar{\alpha})$ as in (1), there exists a tuple $\tilde{\alpha} \in \mathbb{Q}(\bar{\alpha})$ of the same length as $\bar{\alpha}$ such that*

- *the entries of $\tilde{\alpha}$ are locally IO-identifiable;*
- *the system $\Sigma(\tilde{\alpha})$ has the same input-output equations as the original $\Sigma(\bar{\alpha})$.*

Furthermore, let E_1, \dots, E_m be the IO-equations of $\Sigma(\tilde{\alpha})$. If $\sum_i \text{ord}_{\bar{y}} E_i$ is equal to the dimension of the model (n in the notation of (1)), then the state variables of $\Sigma(\tilde{\alpha})$ can be expressed as algebraic functions of \bar{x} and $\tilde{\alpha}$.

Thus, system $\Sigma(\tilde{\alpha})$ is a locally identifiable reparametrization of system $\Sigma(\bar{\alpha})$, and these systems have the same shape (see Section C).

C. On the existence of the state transformation

Let us make some remarks about the second part of the theorem.

Remark 2 (On the last condition and observability). The condition in Theorem 1 that the sum of the orders of the input-output equations w.r.t. \bar{y} be equal to the dimension of the system is, in fact, equivalent to the fact that all the states are locally observable (the initial conditions are locally identifiable) if the parameters are assumed to be *known* (see [7, Proposition 5]). This equivalence can also be deduced from [11, Corollary 4.11] and [10, Proposition 3.4]. In particular, this restriction is significantly milder than observability of all the states.

Example 1 (Non-existence of the state transformation). *Let us give an example showing that the condition for the orders of IO-equations summing up to n cannot be removed. Consider the system*

$$\begin{cases} x_1' = \alpha_1, \\ x_2' = \frac{x_2}{\alpha_2}, \\ y = x_1. \end{cases}$$

The IO-equation of this model is $y' - \alpha_1 = 0$, so only α_1 is (locally) IO-identifiable. Then any specialization described in Theorem 1 will be of the form

$$\begin{cases} w_1' = \alpha_1, \\ w_2' = f(\alpha_1)w_2, \\ y = x_1. \end{cases}$$

for some nonzero algebraic function $f \in \overline{\mathbb{Q}(\alpha_1)}$. Any solution of the new system will be of the form

$$(\alpha_1 t + c_1, c_2 e^{f(\alpha_1)t}),$$

while any solution of the original system was

$$(\alpha_1 t + c_3, c_4 e^{\alpha_2 t}).$$

The existence of an algebraic state space transformation as in the theorem would imply that $e^{f(\alpha_1)t}$ is algebraic over $\mathbb{Q}(\alpha_1, e^{\alpha_2 t})$, which is not the case.

Remark 3 (On possible preprocessing). While not all the models satisfy the condition of the second part of the theorem, one can use the approach from [6, Section 3] (see [26, Section 3.2] for the case with inputs) by constructing a realization of the input-output equation of the model of minimal dimension. The corresponding system (5) from our proof will be nonsingular in this case, providing a coordinate change. After that, Theorem 1 can be applied. Note that since the dimension of the model changes under this transformation, it is not possible to preserve the shape (see Section C) as in Theorem 1. We give an example of such a preprocessing in Section E.

III. Constructive proof of Theorem 1

We break down the proof into several **steps**, each of which can be viewed as a step in an algorithm to solve the problem:

1. We first fix $\bar{\beta} = (\beta_1, \dots, \beta_N) \in \mathbb{Q}(\bar{\alpha})$ a tuple generating the field of IO-identifiable functions of an ODE system $\Sigma(\bar{\alpha})$ and write it explicitly in terms of $\bar{\alpha}$:

$$\beta_1 = p_1(\bar{\alpha}), \dots, \beta_N = p_N(\bar{\alpha})$$

for some rational functions $p_1, \dots, p_N \in \mathbb{Q}(\bar{\alpha})$.

2. Let n be the order of the original ODE system, that is, the length of the tuple \bar{x} . Let $E_1(\bar{\alpha}, \bar{u})(\bar{y}), \dots, E_m(\bar{\alpha}, \bar{u})(\bar{y})$ be the input-output equations with respect to any ranking on \bar{y} . Then the orders of \bar{y} and \bar{u} in E_i do not exceed n for every $1 \leq i \leq m$.

Using Lie derivatives, we can write $\bar{y}, \dots, \bar{y}^{(m)}$ as rational functions R_0, \dots, R_m in $\bar{x}, \bar{\alpha}, \bar{u}, \dots, \bar{u}^{(m)}$ for any m . Therefore, for all $m \geq n$, $\bar{y}, \dots, \bar{y}^{(m)}$ are algebraically dependent over $\mathbb{Q}(\bar{\alpha})(\bar{u})$, where $F\langle a \rangle$ denotes the differential field generated by a over F , that is, $F(a, a', a'', \dots)$. Furthermore, by [10, Lemma 3.18], we have

$$\mathbb{Q}(\bar{\alpha})(\bar{y}, \bar{u}) = \mathbb{Q}(\bar{\alpha}, \bar{y}, \bar{y}', \dots, \bar{y}^{(m)})(\bar{u}). \quad (3)$$

Let M be the Jacobian matrix of $\bar{R} := R_0, \dots, R_n$ with respect to \bar{x} . Then, by the Jacobian criterion together with (3), its rank r will be equal to

$$\text{trdeg}_{\mathbb{Q}(\bar{\alpha})(\bar{u})} \mathbb{Q}(\bar{\alpha})(\bar{u})(\bar{R}) = \text{trdeg}_{\mathbb{Q}(\bar{\alpha})(\bar{u})} \mathbb{Q}(\bar{\alpha})(\bar{u})(\bar{y}).$$

Let D be the determinant of a nonsingular $r \times r$ -minor of M . We consider D as a rational function in $\bar{x}, \bar{u}, \bar{u}', \dots, \bar{u}^{(n)}$, and take any nonzero coefficient of its numerator, which we denote by $D_0(\bar{\alpha})$. We will use this coefficient in the next step.

3. We now form a system of algebraic equations and inequalities in $\bar{\alpha}$ over $\mathbb{Q}(\bar{\beta})$:

$$\begin{cases} \beta_1 = p_1(\bar{\alpha}), \\ \vdots \\ \beta_N = p_N(\bar{\alpha}), \\ D_0(\bar{\alpha}) \cdot C(\bar{\alpha}) \neq 0, \end{cases} \quad (4)$$

where C is the common denominator of all coefficients from $\mathbb{Q}(\bar{\alpha})$ in (1). System (4) has a solution $\bar{\alpha} = \bar{\alpha}$ and, thus, by Hilbert's Nullstellensatz, has a solution $\bar{\alpha}$ in the algebraic closure of the ground field, $\overline{\mathbb{Q}(\bar{\beta})}$. For this solution, we have $\bar{\beta} \in \mathbb{Q}(\bar{\alpha})$ and $\bar{\alpha} \in \overline{\mathbb{Q}(\bar{\beta})}$ by construction. Furthermore, since the rank of $M|_{\bar{\alpha}=\bar{\alpha}}$ is also r , the specialized ODE system has the same input-output equations.

Indeed, the specialization is possible because $C(\bar{\alpha}) \neq 0$. The specializations of the input-output equations $E_1(\bar{\alpha}, \bar{u})(\bar{y}), \dots, E_m(\bar{\alpha}, \bar{u})(\bar{y})$ belong to the specialized elimination ideal $J := I_{\Sigma(\bar{\alpha})} \cap k\{\bar{y}, \bar{u}\}$ and they still form

a characteristic presentation of some prime differential ideal, we denote it by J_0 . Therefore, if $J_0 = J$, then $E_1(\tilde{\alpha}, \tilde{u})(\tilde{y}), \dots, E_m(\tilde{\alpha}, \tilde{u})(\tilde{y})$ are indeed the input-output equations of the specialized system $\Sigma(\tilde{\alpha})$.

Assume the contrary, that J_0 is a proper subset of J . Since $J \cap k\{\tilde{u}\} = J_0 \cap k\{\tilde{u}\} = \{0\}$ by [10, Lemma 3.1], the sum of the orders of the characteristic presentation of J_0 with respect to \tilde{y} must be less than $\sum_{i=1}^m \text{ord}_{\tilde{y}} E_i = r$. This is because, if one prime ideal is contained in another one and they have the same dimension, then the ideals are equal. On the other hand, under the specialization to $\tilde{\alpha}$, the matrix rank is preserved (as shown above), the sum of the orders of the input-output equations of (1) after specializing $\tilde{\alpha} \rightarrow \tilde{\alpha}$ is also r . This provides us a contradiction with the assumption that $J_0 \neq J$.

4. Finally, let \tilde{w} denote the state variables of $\Sigma(\tilde{\alpha})$. Assume that the sum of the orders of IO-equations is equal to n . We will now see how \tilde{w} can be obtained from the original \tilde{x} . Consider the irreducible affine variety V defined by the IO-equations in the space with coordinates $\tilde{y}, \dots, \tilde{y}^{(m)}$ over the field $\mathbb{Q}(\tilde{\beta})\langle\tilde{u}\rangle$. The dimension of this variety is equal to sum of the orders of the IO-equations [11, Corollary 4.11], and thus it is equal to n . The equalities

$$\tilde{y}^{(i)} = R_i(\tilde{\alpha}, \tilde{w}, \tilde{u}), \quad i = 0, \dots, m,$$

define a dominant rational map ψ from the affine n -space with coordinates \tilde{w} to V over the field $\mathbb{Q}(\tilde{\beta})\langle\tilde{u}\rangle = \mathbb{Q}(\tilde{\alpha})\langle\tilde{u}\rangle$.

Consider the field $F := \mathbb{Q}(\tilde{\beta})\langle\tilde{u}\rangle(V) = \mathbb{Q}(\tilde{\alpha})\langle\tilde{u}\rangle(V)$ and the generic point $\tilde{Y} = (Y_0, \dots, Y_m)$ of the variety V with points whose components belong to the field F . Since the morphism ψ is dominant and \tilde{Y} is generic, the system of equations $\psi(\tilde{w}) = \tilde{Y}$ in the variables \tilde{w} has a solution in \overline{F} . On the other hand, the field

$$\overline{\mathbb{Q}(\tilde{\alpha})\langle\tilde{u}\rangle} (R_0(\tilde{\alpha}, \tilde{x}, \tilde{u}), \dots, R_m(\tilde{\alpha}, \tilde{x}, \tilde{u}))$$

is isomorphic to the extension of F by $\overline{\mathbb{Q}(\tilde{\alpha})}$ via

$$R_i(\tilde{\alpha}, \tilde{x}, \tilde{u}) \mapsto Y_i, \quad 0 \leq i \leq m.$$

Denote this isomorphism by φ . The desired correspondence between \tilde{w} and \tilde{x} can now be found by applying φ^{-1} to a solution of $\psi(\tilde{w}) = \tilde{Y}$. This way, we get algebraic functions $\tilde{w} = \tilde{w}(\tilde{\alpha}, \tilde{x}, \tilde{u}, \tilde{u}', \dots)$ such that

$$R_i(\tilde{\alpha}, \tilde{w}, \tilde{u}) = R_i(\tilde{\alpha}, \tilde{x}, \tilde{u}) \quad \text{for } i = 0, \dots, m. \quad (5)$$

We will now show that w'_i is indeed equal to $f_i(\tilde{w}, \tilde{\alpha}, \tilde{u})$. By differentiating (5), we establish that w'_1, \dots, w'_n satisfy the following linear system (cf. [6, Section 3] and [26, Section 3.2]):

$$\begin{aligned} \sum_{j=1}^n w'_j \frac{\partial R_i(\tilde{\alpha}, \tilde{w}, \tilde{u})}{\partial w_j} &= \sum_{j=1}^n f_j(\tilde{x}, \tilde{\alpha}, \tilde{u}) \frac{\partial R_i(\tilde{\alpha}, \tilde{x}, \tilde{u})}{\partial x_j} \\ &+ \sum_{j=1}^{\ell} u'_j \left(\frac{\partial R_i(\tilde{\alpha}, \tilde{x}, \tilde{u})}{\partial u_j} - \frac{\partial R_i(\tilde{\alpha}, \tilde{w}, \tilde{u})}{\partial u_j} \right) \end{aligned}$$

for $i = 0, \dots, m$. Since $\dim V = n$, the rank of the matrix of this linear system is also n , so it has a unique solution. On the other hand, $f_1(\tilde{w}, \tilde{\alpha}, \tilde{u}), \dots, f_n(\tilde{w}, \tilde{\alpha}, \tilde{u})$ is a solution of this system by construction of the R_i 's. Therefore,

$$w'_i = f_i(\tilde{w}, \tilde{\alpha}, \tilde{u}), \quad 1 \leq i \leq n,$$

for the found algebraic functions $\tilde{w} = \tilde{w}(\tilde{\alpha}, \tilde{x}, \tilde{u}, \tilde{u}', \dots)$.

Finally, we will show that $\tilde{w}(\tilde{\alpha}, \tilde{x}, \tilde{u}, \tilde{u}', \dots)$ in fact does not depend on \tilde{u} (or any of its derivatives). Let h be the largest order of \tilde{u} occurring in these expressions, say $u_1^{(h)}$ occurs in w_1 . Then w'_1 will depend nontrivially on $u_1^{(h+1)}$. On the other hand, $w'_1 = f_1(\tilde{w}, \tilde{\alpha}, \tilde{u})$, where $\tilde{\alpha}$ does not involve \tilde{u} , \tilde{u} is of order zero in \tilde{u} , and $\tilde{w}(\tilde{\alpha}, \tilde{x}, \tilde{u}, \tilde{u}', \dots)$ is of order at most h in \tilde{u} . Thus, we have arrived at a contradiction with our original assumption that $\tilde{w}(\tilde{\alpha}, \tilde{x}, \tilde{u}, \tilde{u}', \dots)$ depended in \tilde{u} .

To carry this out computationally, we solve the system of rational equations (5) for \tilde{w} over $\mathbb{Q}(\tilde{\alpha}, \tilde{x})\langle\tilde{u}\rangle$. This can be done, for instance, by computing a Gröbner bases of the numerator of (5) with an elimination monomial ordering $w_i > \tilde{x}$ for each i , $1 \leq i \leq n$, over $\mathbb{Q}(\tilde{\alpha})\langle\tilde{u}\rangle$ to find a polynomial equation $P(w_i, \tilde{x}, \tilde{\alpha}, \tilde{u}) = 0$ for each i (see Sections C, D, and E for concrete examples).

Remark 4. The equations given by p_1, \dots, p_N in (4) can be very complicated. Our MAPLE code <https://github.com/pogudingleb/AllIdentifiableFu> based on [23] can significantly simplify the system by applying the functions `FieldToIdeal` and then `FilterGenerators` to $p_1(\tilde{\alpha}), \dots, p_N(\tilde{\alpha})$.

Remark 5. For some models, IO-equations do not depend on a subset of parameters. These parameters are then definitely non-identifiable. From the measurement point of view, these parameters do not affect the differential equations satisfied by the inputs and outputs of the model but can (and typically will) affect the initial conditions of these equations. In particular, a change in these nonidentifiable parameters can typically be "compensated" (in the sense of preserving the input and output data) by a change in one of the initial conditions for the states. For example, scaling $b \rightarrow \lambda b$ in (14) can be compensated by additionally scaling $x_2(0) \rightarrow \frac{x_2(0)}{\lambda}$.

IV. Examples

In this section, we will illustrate the approach

- by a series of simple examples that explain what can and cannot be achieved in principle and how the algorithm works in practice as well as
- by systems from modeling, Lotka-Volterra with and without inputs, chemical reaction network system, a bilinear model with input, a rational Goodwin oscillator model, and a rational blood coagulation and inhibition model with multiple outputs.

A. *Not possible to achieve global identifiability with any method*

We will begin by showing that a globally IO-identifiable reparametrization does not always exist¹ (cf. [5, Theorem 3.5]). For this, consider the system

$$\begin{cases} x' = \frac{\alpha_2}{2\alpha_1}(x^2 + 1), \\ y = \frac{2x}{\alpha_2(1 + x^2)}. \end{cases} \quad (6)$$

The corresponding IO-equation is

$$\alpha_1^2(y')^2 + \alpha_2^2y^2 - 1 = 0, \quad (7)$$

and so the field of IO-identifiable functions is $\mathbb{Q}(\alpha_1^2, \alpha_2^2)$, and so neither α_1 nor α_2 are globally IO-identifiable. As in Theorem 1, we set $\beta_1 = \alpha_1^2$ and $\beta_2 = \alpha_2^2$. If system (6) had had a reparametrization over $\mathbb{Q}(\beta_1, \beta_2)$ (and so had been IO-identifiable), then the curve C (ellipse) defined by

$$\beta_1x^2 + \beta_2y^2 = 1$$

would have had a rational parametrization over $\mathbb{Q}(\beta_1, \beta_2)$ but it does not. If it had had a parametrization over $\mathbb{Q}(\beta_1, \beta_2)$, it would have had a point

$$(x_0, y_0) = (q_1(\beta_1, \beta_2), q_2(\beta_1, \beta_2)) \in \mathbb{Q}(\beta_1, \beta_2).$$

We write both q_1 and q_2 as Laurent series in β_1 over the field $\mathbb{Q}(\beta_2)$. Then $\beta_1q_1^2$ is a Laurent series of odd valuation. Since $\beta_2q_2^2$ and 1 are of even valuation their valuations must be equal and the dominating terms must cancel. Hence,

$$q_2 = c_0(\beta_2) + \mathcal{O}(\beta_1),$$

so the constant term of $1 - \beta_2q_2^2$ is $1 - \beta_2c_0(\beta_2)^2$. This cannot be equal to zero because $1/\beta_2$ is not a square, so we arrive at a contradiction.

B. *Not possible to achieve global identifiability with this method*

In this section, we give an example for which there is a reparametrization that gives global IO-identifiability, although our method only gives a reparameterization with local IO-identifiability. Consider the coupled by measurements exponential growth/decay system

$$\begin{cases} x'_1 = ax_1, \\ x'_2 = bx_2, \\ y = x_1 + x_2, \end{cases} \quad (8)$$

and so $\bar{x} = (x_1, x_2)$, $\bar{y} = y$, and $\bar{a} = (a, b)$. There is no \bar{u} . The IO-equation is

$$y'' - (a + b)y' + ab \cdot y = 0. \quad (9)$$

¹We are grateful to Sebastian Falkensteiner and Rafael Sendra for pointing out a mistake in a previous version of this example and offering a correction.

Therefore, $\bar{\beta} = (a + b, a \cdot b)$ and the identifiable functions are $K := \mathbb{Q}(a + b, a \cdot b)$, and so a and b are algebraic of degree 2 over K , therefore, are only locally identifiable. For $i = 0, 1, 2$, we will compute $y^{(i)}$ as a function $R_i(x_1, x_2, a, b)$:

$$\begin{aligned} y &= R_0(x_1, x_2, a, b) = x_1 + x_2, \\ y' &= R_1(x_1, x_2, a, b) = x'_1 + x'_2 = ax_1 + bx_2, \\ y'' &= R_2(x_1, x_2, a, b) = x''_1 + x''_2 = a^2x_1 + b^2x_2, \end{aligned} \quad (10)$$

The Jacobian with respect to \bar{x} is

$$M = \begin{pmatrix} 1 & 1 \\ a & b \\ a^2 & b^2 \end{pmatrix}$$

Let $r = \text{rank } M = 2$ and $D = \det M = \begin{vmatrix} 1 & 1 \\ a & b \end{vmatrix} = b - a$ be a non-singular minor of M . Considering D as a rational function in \bar{x} , we pick a (the only, in fact) non-zero coefficient $D_0(\bar{a})$ of its numerator as $b - a$. We now consider the following system of equations and inequations in $\bar{\alpha}$ over $\mathbb{Q}(\bar{\beta})$:

$$\begin{cases} a + b = \bar{\alpha}_1 + \bar{\alpha}_2 \\ a \cdot b = \bar{\alpha}_1 \cdot \bar{\alpha}_2 \\ \bar{\alpha}_1 - \bar{\alpha}_2 \neq 0, \end{cases}$$

which has two solutions: $(\bar{\alpha}_1 = a, \bar{\alpha}_2 = b)$ and $(\bar{\alpha}_1 = b, \bar{\alpha}_2 = a)$, neither of which make the locally identifiable model globally identifiable.

However, for example, the following reparametrization makes the model globally identifiable:

$$\begin{cases} w_1 = x_1 + x_2, \\ w_2 = ax_1 + bx_2, \end{cases}$$

resulting in this reparametrized ODE system:

$$\begin{cases} w'_1 = w_2, \\ w'_2 = (a + b) \cdot w_2 - a \cdot b \cdot w_1, \\ y = w_1, \end{cases}$$

whose IO-equation is also (9).

C. *Choosing different solutions of (4)*

Consider the simple linear harmonic oscillator system

$$\begin{cases} x'_1 = ax_2, \\ x'_2 = bx_1, \\ y = x_1, \end{cases} \quad (11)$$

so $\bar{x} = (x_1, x_2)$, $\bar{y} = (y)$, $\bar{a} = (a, b)$, and we have no \bar{u} . The IO-equation is

$$y'' - ab \cdot y = 0. \quad (12)$$

Therefore, $\bar{\beta} = (ab)$ and ab is globally identifiable but neither a nor b is identifiable. Let us begin by computing Lie derivatives of \bar{y} . We have

$$\begin{aligned} y &= R_0(x_1, x_2, a, b) = x_1, \\ y' &= R_1(x_1, x_2, a, b) = x'_1 = ax_2, \\ y'' &= R_2(x_1, x_2, a, b) = x''_1 = ax'_2 = abx_1. \end{aligned} \quad (13)$$

Following the the proof of Theorem 1, we have

$$\beta_1 = ab = p_1(\bar{\alpha}) = p_1(a, b).$$

We then find the Jacobian of (13) with respect to \bar{x} :

$$M = \begin{pmatrix} 1 & 0 \\ 0 & a \\ ab & 0 \end{pmatrix}.$$

Then $r := \text{rank } M = 2 = \text{trdeg } \mathbb{Q}(a, b)(\bar{y})/\mathbb{Q}(a, b)$. Let $D = \det \begin{pmatrix} 1 & 0 \\ 0 & a \end{pmatrix} = a$, a non-singular maximal minor of M . Considering D as a rational function of \bar{x} , we pick a non-zero coefficient $D_0(\bar{\alpha})$ of its numerator as a . We now consider the following system of equations and inequations over $\mathbb{Q}(\bar{\beta})$:

$$\begin{cases} ab = \bar{\alpha}_1 \cdot \bar{\alpha}_2, \\ \bar{\alpha}_1 \neq 0, \end{cases}$$

which has infinitely many solutions (over the algebraic closure of $\mathbb{Q}(ab)$), including this one: $\bar{\alpha}_1 = ab$, $\bar{\alpha}_2 = 1$. Specializing (11), we obtain the reparametrized system as:

$$\begin{cases} w'_1 = \beta_1 w_2, \\ w'_2 = w_1, \\ y = w_1, \end{cases}$$

whose input-output equation is still (12), and the corresponding change of variables, calculated by equating the old and new Lie derivatives, is

$$w_1 = x_1, \quad w_2 = \frac{x_2}{b},$$

which is a scaling reparametrization. We could choose a different solution for $\bar{\alpha}_1, \bar{\alpha}_2$, for example, $\bar{\alpha}_1 = 1, \bar{\alpha}_2 = ab$ which would yield

$$w'_1 = w_2, \quad w'_2 = \beta_1 w_1, \quad y = w_1.$$

D. Lotka-Volterra examples

D.1. Classical model

Consider the system

$$\begin{cases} x'_1 = ax_1 - bx_1x_2, \\ x'_2 = -cx_2 + dx_1x_2, \\ y = x_1 \end{cases} \quad (14)$$

with two state variables $\bar{x} = (x_1, x_2)$, four parameters $\bar{a} = (a, b, c, d)$, and one output $\bar{y} = y$. The input-output equation is

$$yy'' - y'^2 - dy^2y' + cyy' + ady^3 - acy^2 = 0. \quad (15)$$

So, we have that the field of IO-identifiable functions is $\mathbb{Q}(d, c, ad, ac) = \mathbb{Q}(a, c, d)$. The Lie derivatives of the y -variable are as follows:

$$\begin{aligned} y &= x_1, \\ y' &= -bx_1x_2 + ax_1, \\ y'' &= -bdx_1^2x_2 + b^2x_1x_2^2 + (bc - 2ab)x_1x_2 + a^2x_1 \end{aligned} \quad (16)$$

Following the proof of Theorem 1, we define

$$\beta_1 = p_1(\bar{\alpha}) = a, \quad \beta_2 = p_2(\bar{\alpha}) = c, \quad \beta_3 = p_3(\bar{\alpha}) = d.$$

The Jacobian of (16) w.r.t. \bar{x} is

$$M = \begin{pmatrix} 1 & 0 \\ a - bx_2 & -bx_1 \\ b^2x_2^2 - 2b(dx_1 + a - c/2)x_2 + a^2 & -bx_1(dx_1 - 2bx_2 + 2a - c) \end{pmatrix}.$$

Then

$$D = \det \begin{pmatrix} 1 & 0 \\ a - bx_2 & -bx_1 \end{pmatrix} = -bx_1$$

is a maximal non-zero minor of M . Considering D as a rational function of \bar{x} , we pick a non-zero coefficient $D_0(\bar{\alpha})$ of its numerator as $-b$. We now arrive at the following system in $\bar{\alpha}$ over $\mathbb{Q}(\bar{\beta})$:

$$\begin{cases} a = \bar{\alpha}_1, \\ c = \bar{\alpha}_3, \\ d = \bar{\alpha}_4, \\ -\bar{\alpha}_2 \neq 0 \end{cases}, \quad (17)$$

and we pick the following solution (out of infinitely many solutions): $\bar{\alpha}_1 = a, \bar{\alpha}_2 = 1, \bar{\alpha}_3 = c, \bar{\alpha}_4 = d$, which results in the following reparametrized system of equations

$$\begin{cases} w'_1 = aw_1 - w_1w_2, \\ w'_2 = -cw_2 + dw_1w_2, \\ y = w_1 \end{cases}$$

The corresponding change of variables, obtained by equating old and new Lie derivatives, is

$$w_1 = x_1, \quad w_2 = bx_2, \quad (18)$$

which is a scaling reparametrization in this case.

D.2. Version with input

Consider the following model from [13, eqs. (4)-(6)]:

$$\begin{cases} x'_1 = ax_1 - bx_1x_2 + ux_1, \\ x'_2 = -cx_2 + dx_1x_2 + ux_2, \\ y = x_1. \end{cases}$$

The IO-equation is

$$\begin{aligned} yy'' - y'^2 - dy^2y' - yuy' + cyy' + duy^3 + ady^3 - y^2u' \\ + y^2u^2 + au^2 - cuy^2 - acy^2 = 0. \end{aligned}$$

Therefore, the globally IO-identifiable parameters are $\mathbb{Q}(a, d, c)$. Following the steps from Section III, we consider the Lie derivatives of y :

$$\begin{aligned} y &= x_1, \\ y' &= x_1(-bx_2 + a + u), \\ y'' &= x_1(x_2^2b^2 - (dx_1 + 2a - c + 3u)bx_2 + (a + u)^2) \end{aligned} \quad (19)$$

The first 2×2 principal minor of the Jacobian of (19) w.r.t. \bar{x} is:

$$\begin{pmatrix} 1 & 0 \\ -bx_2 + a + u & -bx_1 \end{pmatrix}.$$

Then

$$D = \det \begin{pmatrix} 1 & 0 \\ -bx_2 + a + u & -bx_1 \end{pmatrix} = -bx_1$$

is a maximal non-zero minor of M . Considering D as a rational function of \bar{x} , we pick a non-zero coefficient $D_0(\bar{\alpha})$ of its numerator as $-b$. We again arrive at the system (17) and we pick the following solution: $\bar{\alpha}_1 = a, \bar{\alpha}_2 = 1, \bar{\alpha}_3 = c, \bar{\alpha}_4 = d$, which results in the following reparametrized system of equations

$$\begin{cases} w'_1 = aw_1 - w_1w_2 + uw_1, \\ w'_2 = -cw_2 + dw_1w_2 + uw_2. \end{cases}$$

The corresponding change of variables, obtained by equating old and new Lie derivatives, is again (18).

E. Chemical reaction network

Consider the following ODE model originating from a chemical reaction network, cf. [4, system (2.3)]

$$\begin{cases} X' = k_2 \cdot (A_{UX} + 2A_{XX} + A_{XU}) - k_1 \cdot X \cdot (A_{UX} + A_{XU} + 2A_{UU}), \\ A'_{UU} = k_2 \cdot (A_{UX} + A_{XU}) - 2k_1 \cdot X \cdot A_{UU}, \\ A'_{UX} = k_1 \cdot X \cdot (A_{UU} - A_{UX}) + k_2 \cdot (A_{XX} - A_{UX}), \\ A'_{XX} = k_1 \cdot X \cdot (A_{UX} + A_{XU}) - 2 \cdot k_2 \cdot A_{XX}, \\ A'_{XU} = k_1 \cdot X \cdot (A_{UU} - A_{XU}) + k_2 \cdot (A_{XX} - A_{XU}), \\ y = X. \end{cases}$$

We have $\bar{x} = (X, A_{UU}, A_{UX}, A_{XX}, A_{XU})$, $\bar{\alpha} = (k_1, k_2)$, $\bar{y} = (y)$, and there is no \bar{u} . A computation shows that

$$y'y''' - (y'')^2 + 2k_1(y')^3 = 0 \quad (20)$$

is the IO-equation, and so $\bar{\beta} = (k_1)$ generates the field of IO-identifiable parameters. Note that the order of the input-output equation is less than the dimension of the system, so we will first perform a preprocessing reduction as in Remark 3.

The Lie derivatives are as follows:

$$\begin{aligned} y &= X \\ y' &= k_2(A_{UX} + 2A_{XX} + A_{XU}) - k_1X(A_{UX} + 2A_{UU} + A_{XU}) \\ y'' &= -y' \cdot (k_1X + 2k_1A_{UU} + k_1A_{UX} + k_1A_{XU} + k_2) \\ y''' &= y' \cdot ((8Xk_1^2 + 4k_1(k_1A_{UX} + k_1A_{XU} + k_2))A_{UU} \\ &\quad + (4k_1^2A_{UX} + 4k_1^2A_{XU} + 2k_1k_2)X \\ &\quad - 4k_1k_2A_{XX} + k_2^2 + k_1^2(A_{UX}^2 + 2A_{UX}A_{XU} + A_{XU}^2 + 4A_{UU} + X^2)). \end{aligned}$$

(21)

We will now search for a three-dimensional system in only three variables w_1, w_2, w_3 having the same input-output equation. First we set up the desired Lie derivatives for this new system by replacing the original variables with arbitrary linear forms in w_1, w_2, w_3 , say:

$$X = w_1, \quad A_{UU} = w_2, \quad A_{UX} = w_3, \quad A_{XX} = A_{XU} = 0.$$

We apply this substitution to (21), equate the results before and after the substitution, and solve for w_1, w_2, w_3 (similarly to (5)). We get

$$w_1 = X, \quad w_2 = A_{UU} - A_{XX}, \quad w_3 = A_{UX} + 2A_{XX} + A_{XU}. \quad (22)$$

Then we can set up a linear system on w'_1, w'_2, w'_3 as in [6, Section 3] and find the reduced model:

$$\begin{cases} w'_1 = k_2w_3 - k_1w_1(w_3 + 2w_2), \\ w'_2 = -k_1w_1(w_3 + 2w_2) + k_2w_3, \\ w'_3 = k_1w_1(w_3 + 2w_2) - k_2w_3. \end{cases} \quad (23)$$

To this model, we can apply both parts of Theorem 1. We still have the same IO-equation (20), so $\beta_1 = k_1$. The first three Lie derivatives

$$\begin{aligned} y &= w_1, \\ y' &= k_2w_3 - k_1w_1(w_3 + 2w_2), \\ y'' &= -y'(k_1(w_1 + 2w_2 + w_3)k_1 + k_2) \end{aligned}$$

have nonsingular Jacobian, and one of the coefficients of its determinant is $k_1k_2^2$, so we set up a system

$$\begin{cases} k_1 = \bar{\alpha}_1, \\ \bar{\alpha}_1\bar{\alpha}_2 \neq 0. \end{cases}$$

We take a solution $\bar{\alpha}_1 = k_1$ and $\bar{\alpha}_2 = 1$ in $\overline{\mathbb{Q}(\bar{\beta})} = \overline{\mathbb{Q}(k_1)}$ and, substituting this solution into (23), obtain

$$\begin{cases} v'_1 = v_3 - k_1v_1(v_3 + 2v_2), \\ v'_2 = -k_1v_1(v_3 + 2v_2) + v_3, \\ v'_3 = k_1v_1(v_3 + 2v_2) - v_3. \end{cases}$$

The corresponding state transformation from the last step of the proof of Theorem 1 is

$$\begin{aligned} v_1 &= w_1, \quad v_3 = k_2(w_1 + w_3) - w_1, \\ v_2 &= ((-w_1 - w_3)k_2 + w_1 + 2w_2 + w_3)/2 + \frac{k_2 - 1}{2k_1}. \end{aligned}$$

The overall state transformation between the new and original system can be obtained by composing this with (22):

$$\begin{aligned} v_1 &= X, \\ v_2 &= X + 2A_{UU} + A_{UX} + A_{XU} \\ &\quad - (X + A_{UX} + A_{XU} + 2A_{XX})k_2 + \frac{k_2 - 1}{2k_1}, \\ v_3 &= \frac{X + A_{UX} + A_{XU} + 2A_{XX}}{k_2} - X. \end{aligned}$$

F. Bilinear model with input

Consider the model [18, Example 1]:

$$\begin{cases} x_1' = -p_1x_1 + p_2u, \\ x_2' = -p_3x_2 + p_4u, \\ x_3' = -(p_1 + p_3)x_3 + (p_4x_1 + p_2x_2)u, \\ y = x_3. \end{cases}$$

Complete details of the computation are available here: [24]. Computing the IO-equations, extracting the coefficients, and simplifying the IO-identifiable field generators, we obtain that the globally IO-identifiable functions are

$$k := \mathbb{Q}(p_1p_3, p_2p_4, p_1 + p_3).$$

Following the steps from Section III in our MAPLE code for this example (e.g., a non-zero maximal minor of the Jacobian matrix of Lie derivatives is $p_2p_4(p_1 - p_3)u^2$), we arrive at the following system of equations and inequations in the unknowns $\bar{p}_1, \bar{p}_2, \bar{p}_3, \bar{p}_4$:

$$\begin{cases} p_1p_3 = \bar{p}_1\bar{p}_3, \\ p_2p_4 = \bar{p}_2\bar{p}_4, \\ p_1 + p_3 = \bar{p}_1 + \bar{p}_3, \\ \bar{p}_2\bar{p}_4(\bar{p}_1 - \bar{p}_3) \neq 0. \end{cases}$$

with solutions sought in the field \bar{k} . We pick the tuple $(p_1, 1, p_3, p_2p_4)$ as a solution. In this tuple, $1, p_2p_4 \in k$, and p_1 and p_3 are algebraic over k since they satisfy a polynomial equation

$$Z^2 - (p_1 + p_3)Z + p_1p_3 = 0 \quad (24)$$

with coefficients in k . Thus, we arrive at the following ODE model, which is locally IO-identifiable:

$$\begin{cases} w_1' = -p_1w_1 + u, \\ w_2' = -p_3w_2 + p_2p_4u, \\ w_3' = -(p_1 + p_3)w_3 + (p_2p_4w_1 + w_2)u, \\ y = w_3, \end{cases} \quad (25)$$

in particular, all coefficients except for p_1 and p_3 are globally IO-identifiable (belong to k), and p_1 and p_3 are locally IO-identifiable because they are roots of the polynomial (24). Equating the old and new Lie derivatives of the output variable, we obtain that (25) corresponds to the following change of variables:

$$w_1 = \frac{x_1}{p_2}, \quad w_2 = p_2x_2, \quad w_3 = x_3.$$

G. Goodwin Oscillator

Consider the following rational ODE model [8]:

$$\begin{cases} x_1' = -bx_1 + \frac{1}{c + x_4}, \\ x_2' = \alpha x_1 - \beta x_2, \\ x_3' = \gamma x_2 - \delta x_3, \\ x_4' = \frac{\sigma x_4(\gamma x_2 - \delta x_3)}{x_3}, \\ y = x_1. \end{cases} \quad (26)$$

We have $\bar{x} = (x_1, x_2, x_3, x_4)$, $\bar{y} = (y)$, $\bar{\alpha} = (b, c, \alpha, \beta, \gamma, \delta, \sigma)$, and there is no \bar{u} . We follow the steps from Section III using MAPLE. The IO-equation is an order 4 ODE in y and is too big to display here. However, the field of IO-identifiable functions is

$$k := \mathbb{Q}(b, c, \sigma, \beta\delta, \beta + \delta).$$

In particular, b, c , and σ are globally IO-identifiable, β and δ are locally but not globally IO-identifiable, and α and γ are not locally IO-identifiable. The Lie derivatives are too big to display here too, but their Jacobian w.r.t. to \bar{x} has a 4×4 minor equal

$$\frac{\alpha x_1 x_4^2 \gamma^2 \sigma^2}{x_3^3 (c + x_4)^6}. \quad (27)$$

The numerator of fraction (27) has a non-zero coefficient $\alpha\gamma^2\sigma^2$. Therefore, we arrive at the following system of constraints:

$$\begin{cases} b = \bar{b}, \quad c = \bar{c}, \quad \sigma = \bar{\sigma}, \quad \beta\delta = \bar{\beta}\bar{\delta}, \quad \beta + \delta = \bar{\beta} + \bar{\delta}, \\ \bar{\alpha}\bar{\gamma}^2\bar{\sigma}^2 \neq 0. \end{cases}$$

Therefore, by the constructive proof of Theorem 1 (Section III), substituting any non-zero element of k into α and γ will turn (26) into a locally IO-identifiable model. For example, we can substitute $\alpha = 1$ and $\gamma = 1$:

$$\begin{cases} w_1' = -bw_1 + \frac{1}{c + w_4}, \\ w_2' = w_1 - \beta w_2, \\ w_3' = w_2 - \delta w_3, \\ w_4' = \frac{\sigma w_4(w_2 - \delta w_3)}{w_3}, \\ y = w_1, \end{cases}$$

and the corresponding change of variables can be found by equating the old and new Lie derivatives, and it is:

$$w_1 = x_1, \quad w_2 = \frac{x_2}{\alpha}, \quad w_3 = \frac{x_3}{\alpha\gamma}, \quad w_4 = x_4.$$

H. Blood coagulation and inhibition

Consider the following model from [27, eq. (1)]:

$$\begin{cases} [\text{IXa}]' = k_1 \cdot \beta - h_1 \cdot [\text{IXa}], \\ [\text{VIIIa}]' = k_2 \cdot [\text{IIa}] + \frac{k_3 \cdot [\text{APC}] \cdot [\text{VIIIa}]}{b_1 + [\text{VIIIa}]} - h_2 \cdot [\text{VIIIa}], \\ [\text{Xa}]' = \frac{k_5 \cdot [\text{IXa}] \cdot [\text{VIIIa}]}{b_2 + [\text{VIIIa}]} - h_3 \cdot [\text{Xa}], \\ [\text{Va}]' = k_6 \cdot [\text{IIa}] - \frac{k_7 \cdot [\text{APC}] \cdot [\text{Va}]}{b_3 + [\text{Va}]} - h_4 \cdot [\text{Va}], \\ [\text{APC}]' = k_8 \cdot [\text{IIa}] - h_5 \cdot [\text{APC}], \\ [\text{IIa}]' = \frac{k_9 \cdot [\text{Xa}] \cdot [\text{Va}]}{b_4 + [\text{Va}]} - h_6 \cdot [\text{IIa}], \\ y_1 = [\text{Xa}], \\ y_2 = [\text{IIa}], \\ y_3 = [\text{IXa}], \\ y_4 = [\text{APC}] \end{cases}$$

(28) reparametrization of (28):

The 4 IO-equations are too big to display here. The orders of the equations are 2, 2, 1, and 1, respectively. So, the sum of the orders is 6. The field of IO-identifiable functions of the parameters is

$$k := \mathbb{Q}\left(h_1, \dots, h_6, k_5, k_8, k_9, \frac{b_1}{k_3}, \frac{b_2}{k_3}, \frac{b_3}{k_7}, \frac{b_4}{k_7}, k_1\beta, \frac{k_2}{k_3}, \frac{k_6}{k_7}\right).$$

The Jacobian matrix M w.r.t. \bar{x} of the Lie derivatives of \bar{y} is too big to display as well. The first 6×6 principal leading minor of M is non-zero, and it has a non-zero coefficient w.r.t. \bar{x} of the numerator equal to

$$b_1^2 b_2^2 b_4^2 \beta k_1 k_5 k_7 k_9^2.$$

We therefore, following Section III, obtain the following system of constraints, after removing unnecessary exponents from the inequality product:

$$\begin{cases} h_i = \tilde{h}_i, 1 \leq i \leq 6, \\ k_5 = \tilde{k}_5, k_8 = \tilde{k}_8, k_9 = \tilde{k}_9, \\ \frac{b_1}{k_3} = \frac{\tilde{b}_1}{\tilde{k}_3}, \frac{b_2}{k_3} = \frac{\tilde{b}_2}{\tilde{k}_3}, \\ \frac{b_3}{k_7} = \frac{\tilde{b}_3}{\tilde{k}_7}, \frac{b_4}{k_7} = \frac{\tilde{b}_4}{\tilde{k}_7}, \\ k_1\beta = \tilde{k}_1\tilde{\beta}, \\ \frac{k_2}{k_3} = \frac{\tilde{k}_2}{\tilde{k}_3}, \frac{k_6}{k_7} = \frac{\tilde{k}_6}{\tilde{k}_7}, \\ \tilde{b}_1\tilde{b}_2\tilde{b}_4\tilde{\beta}k_1\tilde{k}_5\tilde{k}_7\tilde{k}_9 \neq 0. \end{cases} \quad (29)$$

Our calculation in MAPLE shows that the solution set

$$\begin{aligned} \tilde{b}_1 &= \frac{b_1\tilde{k}_3}{k_3}, \tilde{b}_2 = \frac{b_2\tilde{k}_3}{k_3}, \tilde{b}_3 = \frac{b_3\tilde{k}_7}{k_7}, \tilde{b}_4 = \frac{b_4\tilde{k}_7}{k_7}, \tilde{\beta} = \frac{k_1\beta}{\tilde{k}_1}, \\ \tilde{h}_i &= h_i, 1 \leq i \leq 6, \tilde{k}_1 = \tilde{k}_1, \tilde{k}_2 = \frac{\tilde{k}_3 k_2}{k_3}, \\ \tilde{k}_3 &= \tilde{k}_3, \tilde{k}_5 = k_5, \tilde{k}_6 = \frac{\tilde{k}_7 k_6}{k_7}, \tilde{k}_7 = \tilde{k}_7, \tilde{k}_8 = k_8, \tilde{k}_9 = k_9 \end{aligned}$$

of (29) can be parametrized by $(\tilde{k}_1, \tilde{k}_3, \tilde{k}_7)$, with none of $\tilde{k}_1, \tilde{k}_3,$ or \tilde{k}_7 being zero. As one of the reparametrization choices, we let $\tilde{k}_1 = \tilde{k}_3 = \tilde{k}_7 = 1$, obtaining the following locally identifiable

$$\begin{cases} w'_1 = K_1 - h_1 w_1, \\ w'_2 = \frac{(w_5 - B_1 h_2 + K_2 w_6) w_2 + K_2 B_1 w_6 - h_2 w_2^2}{B_1 + w_2}, \\ w'_3 = \frac{k_5 w_1 w_2 - h_3 w_2 w_3 - B_2 h_3 w_3}{B_2 + w_2}, \\ w'_4 = \frac{(K_6 w_6 - B_3 h_4 - w_5) w_4 + K_6 B_3 w_6 - h_4 w_4^2}{B_3 + w_4}, \\ w'_5 = k_8 w_6 - h_5 w_5, \\ w'_6 = \frac{k_9 w_3 w_4 - h_6 (B_4 + w_4) w_6}{B_4 + w_4}, \\ y_1 = w_1, \\ y_2 = w_3, \\ y_3 = w_5, \\ y_4 = w_6, \end{cases} \quad (30)$$

where

$$K_1 = k_1\beta, B_1 = \frac{b_1}{k_3}, K_2 = \frac{k_2}{k_3}, B_2 = \frac{b_2}{k_3}, K_6 = \frac{k_6}{k_7}, B_3 = \frac{b_3}{k_7}, B_4 = \frac{b_4}{k_7}$$

are all in the field k . By setting the Lie derivatives of the outputs from (28) equal to those of (30), we arrive at the following change of variables to achieve the reparametrization (30):

$$\begin{aligned} w_1 &= [\text{IXa}], & w_2 &= \frac{[\text{VIIIa}]}{k_3}, & w_3 &= [\text{Xa}], \\ w_4 &= \frac{[\text{Va}]}{k_7}, & w_5 &= [\text{APC}], & w_6 &= [\text{IIa}]. \end{aligned}$$

Acknowledgments

We are grateful to the CCiS at CUNY Queens College for the computational resources and to Julio Banga, Sebastian Falkensteiner, Gemma Massonis, Nikki Meshkat, Rafael Sendra and Alejandro Villaverde for useful discussions and the referees for careful reading and helpful comments.

References

- [1] J. A. Baaijens and J. Draisma. On the existence of identifiable reparametrizations for linear compartment models. *SIAM Journal on Applied Mathematics*, 76:1577–1605, 2015. URL <https://doi.org/10.1137/15M1038013>.
- [2] G. Bellu, M. P. Saccomani, S. Audoly, and L. D'Angiò. DAISY: A new software tool to test global identifiability of biological and physiological systems. *Computer Methods and Programs in Biomedicine*, 88(1):52–61, 2007. URL <http://dx.doi.org/10.1016/j.cmpb.2007.07.002>.
- [3] F. Boulier and F. Lemaire. Computing canonical representatives of regular differential ideals. In *Proceedings of the 2000 International Symposium on Symbolic and Algebraic Computation (ISSAC)*, pages 38–47. ACM, 2000. URL <https://doi.org/10.1145/345542.345571>.
- [4] A. Demin, E. Demitraki, and G. Pogudin. Exact linear reductions of dynamical models, 2023. URL <https://arxiv.org/pdf/2301.11653.pdf>.
- [5] S. Falkensteiner, D. Pavlov, and R. Sendra. On real and observable realizations of input-output equations, 2023. URL <https://arxiv.org/abs/2303.16799>.

- [6] K. Forsman. On rational state space realizations. *IFAC Proceedings Volumes*, 25(13):341–346, 1992. URL [https://doi.org/10.1016/S1474-6670\(17\)52305-8](https://doi.org/10.1016/S1474-6670(17)52305-8).
- [7] S. Glad. Differential algebraic modelling of nonlinear systems. In M. Kaashoek, J. van Schuppen, and A. Ran, editors, *Realization and Modelling in System Theory*, volume 3 of *Progress in Systems and Control Theory*, Boston, 1990. Birkhäuser. URL https://doi.org/10.1007/978-1-4612-3462-3_9.
- [8] B. C. Goodwin. Oscillatory behavior in enzymatic control processes. *Advances in enzyme regulation*, 3:425–437, 1965. URL [https://doi.org/10.1016/0065-2571\(65\)90067-1](https://doi.org/10.1016/0065-2571(65)90067-1).
- [9] R. N. Gunn, M. J. Chappell, and V. J. Cunningham. Reparameterisation of unidentifiable systems using the Taylor series approach. *IFAC Proceedings Volumes*, 30(2):247–252, 1997. URL [https://doi.org/10.1016/S1474-6670\(17\)44579-4](https://doi.org/10.1016/S1474-6670(17)44579-4).
- [10] H. Hong, A. Ovchinnikov, G. Pogudin, and C. Yap. Global identifiability of differential models. *Communications on Pure and Applied Mathematics*, 73(9):1831–1879, 2020. URL <https://doi.org/10.1002/cpa.21921>.
- [11] E. Hubert. Notes on triangular sets and triangulation-decomposition algorithms II: Differential Systems. In *Proceedings of the 2nd International Conference on Symbolic and Numerical Scientific Computation*, SNSC’01, pages 40–87, Berlin, Heidelberg, 2003. Springer-Verlag. ISBN 3-540-40554-2. URL https://doi.org/10.1007/3-540-45084-X_2.
- [12] E. Hubert and G. Labahn. Scaling invariants and symmetry reduction of dynamical systems. *Foundations of Computational Mathematics*, 13(4):479–516, 2013. URL <https://doi.org/10.1007/s10208-013-9165-9>.
- [13] A. Ibañez. Optimal control of the Lotka–Volterra system: turnpike property and numerical simulations. *Journal of Biological Dynamics*, 11(1):25–41, 2017. URL <https://doi.org/10.1080/17513758.2016.1226435>.
- [14] I. Ilmer, A. Ovchinnikov, and G. Pogudin. Web-based structural identifiability analyzer. In *Computational Methods in Systems Biology*, volume 12881 of *Lecture Notes in Computer Science*, pages 254–265, 2021. URL https://doi.org/10.1007/978-3-030-85633-5_17.
- [15] A. Jiménez-Pastor, J. P. Jacob, and G. Pogudin. Exact linear reduction for rational dynamical systems. In *Computational Methods in Systems Biology*, pages 198–216, 2022. URL https://doi.org/10.1007/978-3-031-15034-0_10.
- [16] D. Joubert, J. Stigter, and J. Molenaar. An efficient procedure to assist in the re-parametrization of structurally unidentifiable models. *Mathematical Biosciences*, 323:108328, 2020. URL <https://doi.org/10.1016/j.mbs.2020.108328>.
- [17] A. Kalami Yazdi, M. Nadjafikhah, and J. DiStefano. COMBOS2: an algorithm to the input–output equations of dynamic biosystems via Gaussian elimination. *Journal of Taibah University for Science*, 14(1):896–907, 2020. URL <https://doi.org/10.1080/16583655.2020.1776466>.
- [18] Y. Lecourtier, F. Lamnabhi-Lagarrigue, and E. Walter. A method to prove that nonlinear models can be unidentifiable. *26th IEEE Conference on Decision and Control*, pages 2144–2145, 1987. URL <https://doi.org/10.1109/CDC.1987.272467>.
- [19] G. Massonis, J. R. Banga, and A. F. Villaverde. Autorepar: A method to obtain identifiable and observable reparameterizations of dynamic models with mechanistic insights. *International Journal of Robust and Nonlinear Control*, 2021. URL <https://doi.org/10.1002/rnc.5887>.
- [20] N. Meshkat and S. Sullivant. Identifiable reparameterizations of linear compartment models. *Journal of Symbolic Computation*, 63:46–67, 2014. URL <https://doi.org/10.1016/j.jsc.2013.11.002>.
- [21] N. Meshkat, C. Kuo, and J. DiStefano. On finding and using identifiable parameter combinations in nonlinear dynamic systems biology models and COMBOS: A novel web implementation. *PLoS ONE*, 9(10):e110261, 2014. URL <https://doi.org/10.1371/journal.pone.0110261>.
- [22] A. Ovchinnikov, I. Pérez Verona, G. Pogudin, and M. Tribastone. CLUE: exact maximal reduction of kinetic models by constrained lumping of differential equations. *Bioinformatics*, 37(12):1732–1738, 2021. URL <https://doi.org/10.1093/bioinformatics/btab010>.
- [23] A. Ovchinnikov, A. Pillay, G. Pogudin, and T. Scanlon. Computing all identifiable functions of parameters for ODE models. *Systems & Control Letters*, 157:105030, 2021. URL <https://doi.org/10.1016/j.sysconle.2021.105030>.
- [24] A. Ovchinnikov, A. Pillay, G. Pogudin, and T. Scanlon. Maple code for the examples in this paper, 2023. URL <https://github.com/alexeyovchinnikov/Identifiable-special>.
- [25] A. Ovchinnikov, G. Pogudin, and P. Thompson. Parameter identifiability and input-output equations. *Applicable Algebra in Engineering, Communication and Computing*, 34:165–182, 2023. URL <https://doi.org/10.1007/s00200-021-00486-8>.
- [26] D. Pavlov and G. Pogudin. On realizing differential-algebraic equations by rational dynamical systems. In *Proceedings of the 2022 International Symposium on Symbolic and Algebraic Computation*, 2022. URL <https://doi.org/10.1145/3476446.3535492>.
- [27] Y. H. Qiao, C. Q. Xu, Y. J. Zeng, X. H. Xu, H. Zhao, and H. Xu. The kinetic model and simulation of blood coagulation—the kinetic influence of activated protein C. *Medical Engineering & Physics*, 26(4):341–347, 2004. URL <https://doi.org/10.1016/j.medengphy.2004.01.003>.
- [28] M. Saccomani and L. D’Angiò. Examples of testing global identifiability with the DAISY software. *IFAC Proceedings Volumes*, 42(10):48–53, 2009. URL <https://doi.org/10.3182/20090706-3-FR-2004.00007>.
- [29] M. P. Saccomani and G. Bellu. DAISY: An efficient tool to test global identifiability. Some case studies. In *2008 16th Mediterranean Conference on Control and Automation*, pages 1723–1728, 2008. URL <http://dx.doi.org/10.1109/MED.2008.4602152>.
- [30] H. J. Sussmann. Existence and uniqueness of minimal realizations of nonlinear systems. *Mathematical Systems Theory*, 10(1):263–284, 1976. URL <https://doi.org/10.1007/bf01683278>.