The known unknowns of the Hsp90 chaperone

Laura-Marie Silbermann^{1‡}, Benjamin Vermeer^{2‡}, Sonja Schmid^{2*}, Katarzyna (Kasia) Tych^{1*}

¹Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Nijenborgh 7,

9747 AG Groningen, the Netherlands

²Laboratory of Biophysics, Wageningen University & Research, Stippeneng 4, 6708 WE Wageningen,

The Netherlands

[‡]Equal contributions.

*Correspondence to: <u>k.m.tych@rug.nl</u>, <u>schmid@nanodynlab.org</u>.

Abstract

Molecular chaperones are vital proteins that maintain protein homeostasis by assisting in protein folding, activation, degradation, and stress protection. Among them, heat-shock protein 90 (Hsp90) stands out as an essential proteostasis hub in eukaryotes, chaperoning hundreds of "clients" (substrates). After decades of research, several "known unknowns" about the molecular function of Hsp90 remain unanswered, hampering rational drug design for the treatment of cancers, neurodegenerative and other diseases. We highlight three fundamental open questions, reviewing the current state of the field for each, and discuss new opportunities, including single-molecule technologies, to answer the known unknowns of the Hsp90 chaperone.

Proteins are the active workforce in the cell. Among them, chaperone proteins are responsible for maintaining proteostasis, i.e., sustaining a functional proteome adapted to ever-changing cellular and environmental conditions. In eukaryotic cells, the Heat shock protein 90 (Hsp90) plays a central role in proteostasis, as it provides a scaffold that binds substrate proteins (termed clients) as well as a variety of helper chaperones (known as cochaperones)¹⁻⁸. Hsp90 can therefore be regarded as a versatile workbench⁷ where cochaperones act as tools to customise the chaperone's function as needed for client recruitment⁹⁻¹¹, (re)folding^{12,13}, (de)activation^{14,15}, degradation, or protection from degradation^{7,16}. Due to the many clients relying on its function, Hsp90 constitutes a proteostatic hub in eukaryotes, while only a handful of prokaryotic Hsp90 clients are known^{10,11}. Among the hundreds of eukaryotic clients are transcription factors, signalling kinases, DNA replication proteins, cell division regulators, synapse proteins, and many more¹. Hsp90 is highly conserved throughout prokaryotes to higher eukaryotes¹⁷. and appears to have back-transferred from eukaryotes to their archaeal precursors through horizontal gene transfer¹⁸. In the absence of stress, Hsp90 constitutes 1 - 2% of the protein mass in eukaryotic cells¹⁹ and exists in super-stoichiometric ratios (~2:1) to its cochaperones while it is outcompeted tenfold in concentration by that of all of its clients²⁰. Under stress conditions, Hsp90 expression is strongly upregulated²¹. The active Hsp90 dimer²² proceeds through an intricate functional cycle involving ATP hydrolysis, multiple conformational rearrangements, and (transient) protein-protein interactions, whose functional roles have remained largely enigmatic despite over 40 years of Hsp90 research^{23,24}. This is surprising for an ATPase as prevalent as Hsp90, and clearly sets Hsp90 apart from many other important ATPases with very well understood molecular mechanisms. In this Review, we highlight the most important 'known unknowns' of Hsp90's molecular function, with the goal of inspiring new routes to address them and advance the field (Fig. 1).

Structurally, Hsp90 monomers consist of three domains (**Fig. 2A**): an N-terminal domain (NTD), a middle domain (MD), and a carboxy-terminal domain (CTD), as well as an unstructured charged linker region between the NTD and MD. The NTD contains an ATP-binding pocket and a flexible loop segment, the so-called lid, which can fold over the ATP pocket^{25,26}. The charged linker modulates Hsp90 function via NTD interactions^{27,28} and can influence client binding²⁹. Most known client binding sites are located in



Figure 1. Three central questions about Hsp90's functional mechanism. (a) The current combined knowledge describes a general functional cycle of eukaryotic Hsp90, where Hsp90 hydrolyses ATP at a slow rate, interacts with cochaperones and clients, and (by only partially understood mechanisms) matures clients to their functional form. (b) Several open questions remain, as illustrated: What is the role of Hsp90's ATPase activity? How does Hsp90, alone or in cooperation with cochaperones, process its vast client base? Which is the sequence of interactions by which the Hsp90 system operates to produce functional clients? And which are reversible processes as compared to unidirectionally driven ones? Which are fast or rate-limiting steps?

the MD^{2,30}, while specific MD residues also participate in nucleotide binding, alongside NTD residues³¹. The CTD provides the main dimerization interface³², and can participate in client interactions³³. In cytosolic eukaryotic Hsp90, the CTD contains an additional recognition motif (MEEVD) for tetratricopeptide repeat (TPR)-containing cochaperones, such as Sti1/Hop and Cpr6³⁴. The Hsp90 dimer undergoes large conformational rearrangements, involving conformational ensembles with several open (V-shaped), intermediate, and compacted dimer structures^{34–38}, as illustrated in **Fig. 1** (grey shading). Hsp90 is also assisted by cochaperones that either directly modify clients, or interact with Hsp90 to modulate client processing³⁹, resulting in a vast variety of transiently occurring chaperone complexes. Given this polymorphic behaviour, it has been impossible to define a single functional cycle for Hsp90, which – alongside other peculiarities discussed below – limits the current understanding of the Hsp90 functional mechanism.

Peculiarities of the Hsp90 protein system

For common substrate-binding ATPase proteins, such as proteases, helicases, motor proteins etc., the ATPase mechanism and function are well-defined. In stark contrast, Hsp90's function is less deterministic and (likely for that reason) less well-understood. A comparative overview of Hsp90 and a selection of well-known ATPases (**Table 1**) reveals several unique features of Hsp90.

First, Hsp90's ATP hydrolysis rate is very slow $(0.1 - 1.5 \text{ min}^{-1})^{22,40-42}$ – tens to hundreds of times slower than other ATPases in both presence or absence of substrate. This is a common feature of the so-called GHKL family (including the eponymous DNA *G*yrase, *H*sp90, Histidine *K*inase (HisK), and Mut*L*), which share slow ATPase rates (generally $0.02 - 10/\text{min}^{43-45}$, and DNA Gyrase with ~100/min^{46}). Second, the role of ATP hydrolysis is well understood in the molecular mechanism of many standard ATPases. Contrastingly, the function of ATP hydrolysis by Hsp90 has remained unclear, and several hypotheses have been posed by the Hsp90 field, which we discuss below.

Next, the *molecular mechanisms* by which Hsp90 chaperones its clients are not well understood, since often only the *outcomes* (e.g., a functional client) are measurable, but not the molecular process leading there. This contrasts with most ATPases, where the molecular mechanism is well-defined (e.g., myosin's power stroke⁴⁷, kinesin's hand-over-hand cargo transport⁴⁸, or protein threading through protease rings for degradation⁴⁹). Moreover, most ATPases show a specific reaction coordinate (e.g., directional

Functional feature	Eukaryotic Hsp90	Model ATPases
Mechanistic role of ATPase:	Unclear ^{66,79}	Understood [†]
ATPase rate [¥] :	Slow: 0.1 – 5 / minute ^{67,77,143,144}	Fast: 1 – 100 / second ^{145–149}
Substrate specificity:	Weak: diverse clients ¹	Specifics
Interaction affinities:	Low ^{51,52}	Medium to high ^{150–153}
ATPase functional cycle:	Non-deterministic65,66	Deterministic ^{49,145,154}

Table 1: Special features of Hsp90 compared to other well-known ATPase proteins: (i) the motor protein kinesin, (ii) the helicase DnaB. (iii) the protease ClpXP.

†: (i) hand-over-hand transport^{48,155}; (ii) DNA unwinding¹⁵⁶; (iii) unfolded protein translocation⁴⁹.

§: (i) microtubules¹⁵⁵; (ii) DNA¹⁵⁶; (iii) recognition motifs¹⁵⁷.

¥: In presence of substrate.

translocation of an unfolded protein⁴⁹, or sliding along DNA⁵⁰). For Hsp90, no well-defined reaction coordinate seems to exist. In fact, it interacts with a broad clientele, and with (client-specific) sets of cochaperones involved¹, as opposed to standard ATPases recognizing one specific substrate or motif. These numerous interactions occur with low affinity (up to the micromolar range^{30,51,52}) and across the entire protein, not just at one specific client-binding site. While some clients bind within the cleft between both monomers (referred to as the lumen^{2–4,7,53}) others bind outside of it^{1,51}, and cochaperones interact with all domains of Hsp90. Ultimately, Hsp90 achieves the seemingly impossible, namely, to chaperone structurally and functionally diverse clients in *client-specific* ways^{54,55}, which is assumed to be orchestrated via its many cochaperone interactions.

Lastly, for many proteins the timing of their functional cycle is well understood: which functional states are involved (including rare, potentially rate-limiting intermediates), at what rates they occur, whether they occur sequentially or reversibly, and which of them are deterministically driven by ATP hydrolysis (**Table 1**, ATPase mechanistic role & functional cycle). This quantitative information provides a mechanistic understanding of the molecular system which aids rational drug and therapy design. As illustrated in **Fig. 1**, for Hsp90, this detailed molecular level of understanding is yet to be achieved, which may have had an impact on the relatively low success of Hsp90-targeted drugs up to now^{56,57}. However, a vast body of existing biochemical and structural results about client-specific functional cycles exists and indicates possible routes forward.

The enigmatic role of Hsp90's ATPase function

The precise functional role of the slow ATPase of Hsp90 has remained unsolved despite intense research. We first discuss the structural basis of ATP binding and the hydrolysis mechanism. Hsp90 binds ATP in a unique Bergerat fold⁵⁸. The active site of ATP hydrolysis in yeast Hsp90 comprises five amino acid residues (**Fig. 2a**³¹): in the NTD R32, E33, and N37, and in the MD R380 and E381^{25,31,35}. N37 is involved in coordinating ATP via a magnesium ion⁵⁹. Hybrid quantum/classical (QM/MM) free-energy calculations²⁶, combined with large-scale atomistic molecular dynamics (MD) simulations, suggest that upon opening of the R32-E33 ion pair (**Fig. 2a**, rightmost panel) via long-range interaction and conformational switching, E33 deprotonates a water molecule in the active site, resulting in a hydroxide ion which is stabilized by R380, subsequently allowing nucleophilic attack of the bond linking the β- and γ-phosphate in the bound ATP, in which the magnesium ion coordinated by N37 pulls away electron density from the γ-phosphate to lower the hydrolysis energy barrier. After hydrolysis, R380



Figure 2. The role of Hsp90's ATPase function. (a) Structure of the yeast Hsp90 dimer (PDB ID: 2CG9)³⁵. The insert obtained from an MD simulation shows the active site with bound ATP. MD simulations indicate a role of ion pair opening in the ATP-hydrolysis reaction. Figure taken from ³¹ (CC BY 4.0) (b) Visual summary of Hsp90's ATPase rate modulation by interactors, ATPase rate differences among different species and, partially understood aspects of Hsp90's ATPase function, such as interactor dissociation. The notion that ATP hydrolysis is essential for yeast cell viability was challenged by recent evidence. Nucleotide binding and release may be sufficient to support yeast cell viability.

stabilizes the resultant cleaved inorganic phosphate³¹. The coordinated structural changes required for ATP hydrolysis are understood to be rate-limiting for Hsp90's slow ATPase cycle^{25,60–62}. In other words, the probability of being in the hydrolysis competent state⁶³ is low due to the general flexibility of Hsp90 and, in particular, its split ATP binding site³⁵ composed of residues of both the NTD and MD. Last but not least, the measured ca. 3-fold higher affinity for ADP than ATP^{59,64} implies a considerable product-inhibited fraction of ADP-bound Hsp90 even in the presence of excess ATP, which is expected to affect interactor binding and client maturation also *in vivo*³¹.

Interestingly, prokaryotic and eukaryotic Hsp90 show differences regarding the mechano-chemical coupling of ATP hydrolysis and large-scale structural rearrangements of the Hsp90 dimer (V-shaped opening/closing). Prokaryotic Hsp90 (subsequently referred to as HtpG) is described to function as a Brownian ratchet, where thermal fluctuations are biased by ATP-binding towards the closed state, providing a degree of directionality to its conformational cycle³⁸. This has not been observed for eukaryotic Hsp90 (Hsp90_{euk}) where thermal fluctuations dominate even in the presence of ATP^{65,66}, and its conformational equilibrium is more affected by cochaperone and client interactions, as well as molecular crowding^{52,63,67,68}. Nevertheless, Hsp90_{euk}'s nucleotide state modulates the cochaperone- and therefore client-interaction affinities. For example, ATP-binding induces NTD rotation in Hsp90_{euk}, affecting distal client binding sites through long-range allosteric communication, increasing the binding affinity of some (GR-LBD, MR-LBD) but not all studied clients (p53-DBD, Tau)^{51,67}.

A number of functional roles of Hsp90's ATPase activity have been discussed. For example, it was suggested that it facilitates the dissociation of interaction partners during GR maturation⁶⁹ (illustrated in **Fig. 2b**). Interestingly, also for MutL – another member of the GHKL class with a similar ATP binding site and hydrolysis rate – ATPase activity induces (DNA) substrate dissociation⁴⁵. In the Hsp90-Cdc37-kinase cycle, Cdc37 inhibits Hsp90's ATPase^{41,70} while clients are held in the complex⁵⁵, suggesting a role of ATP hydrolysis in regulated client dissociation – which is further supported by molecular dynamics simulations⁷¹. A similar case is the ATPase inhibition by p23 during GR processing⁷². While for some clients, ATP hydrolysis by Hsp90 was found to be necessary (e.g., folding of luciferase⁷³ and reversal

of Hsp70-induced inhibition of GR activation by Hsp90⁶⁹), in other cases, only ATP binding but not hydrolysis is required for client processing (e.g., cochaperone-independent p53 processing⁷⁴ and stabilization of the client kinase v-Src at elevated temperatures⁷⁵). In addition, Hsp90 and its clients *mutually* affect each other and some clients alter Hsp90's ATPase rate: interestingly, ATPase stimulation (α-synuclein and ribosomal protein L2^{76,77}) as well as reduction (GR-LBD client^{52,67}) have been observed. Both effects were attributed to conformational stabilization of either ATPase competent or non-competent states, respectively. In this way, some clients have been suggested to modulate their own residence time on Hsp90⁵². Other clients, such as the mineralocorticoid receptor (MR)-LBD, Tau, and the p53-DNA binding domain (DBD) seem not to affect Hsp90_{euk}'s ATPase activity⁶⁷.

Altogether, the large variation of reported effects of Hsp90's ATPase activity remains puzzling. This is even more so, in light of the unsolved question: is Hsp90's ATPase activity indeed essential for cell viability? While, up to now, most reports maintain the notion that ATPase activity is essential for Hsp90's function and viability of the eukaryotic cell^{40,78}, recent evidence challenges this view: a hydrolysis-dead yeast Hsp90 E33A mutant was unexpectedly found to still support yeast cell growth⁶⁶. Along with a recent study covering several Hsp90 orthologs with this mutation⁷⁹, these findings imply that the ability to adopt different conformations is enough for some functions of Hsp90, and that nucleotide association and dissociation may be sufficient for regulation of Hsp90's conformational equilibrium, while hydrolysis may play a different role. In summary, the large body of experimental evidence for (i) the mutual effects of cochaperones and clients on Hsp90's ATPase rate and for (ii) the role of ATP hydrolysis in client processing and cell viability, raise new questions and we discuss possible experiments to address them further below.

Client processing by Hsp90

Client processing by Hsp90_{euk} can be classified into refolding^{69,80}, sorting for degradation^{51,76,81}, ligand binding regulation^{69,82,83}, re-/de-activation^{14,15}, and regulation of assemblies^{84–86}. Hsp90's diverse clients differ both structurally and functionally^{11,87,88} and, related to that, Hsp90 does not have a well-defined client binding site, which is in contrast to other chaperones, such as Hsp70⁸⁹. Instead, Hsp90 clients form numerous low-affinity contacts through hydrophobic interactions and some client-specific charge-charge interactions⁷². Certain clients (e.g., GR and kinases) primarily bind to Hsp90's lumen formed by both MDs, while intrinsically disordered proteins (IDPs, e.g.: Tau, misfolded transthyretin, etc.) bind interfaces extending from the MD to the NTD^{90,91}. Despite the breadth of clients, Hsp90 is understood to exhibit client-specific chaperoning action, presumably through specific sets of cochaperones^{2–7,92}. Indeed, only certain cochaperones function as integral components of the Hsp90_{euk} system, while others are client specific and provide Hsp90 with the necessary plasticity^{93,94} (**Fig. 3a**), for example, through specific client recruitment¹¹ or by modulating the time for which a client is bound to Hsp90^{93,95–97}.



Figure 3: Convergent model of Hsp90's functional mechanism. (a) Hsp90 can be regarded as a workbench, bringing clients and cochaperones (coloured tools) into close proximity via interaction sites spread over the entire protein (blue spots). However, many details of recognition of the numerous clients by Hsp90 and its cochaperones remain elusive. (b) Various client processing modes are being discussed based on growing experimental evidence, such as sliding, ratcheting, stretching, shielding, and holding. These processing modes seem to be client-specific, and they may offer distinct functional outcomes (e.g., misfolding rescue, or targeting for or protection from degradation).

Exciting molecular-level details of client processing have recently been revealed for the GR, Hop/Sti, p23 system in 3D structures of these multi-component complexes, solved by the Agard lab^{98,99} with cryo-EM. They represent two snapshots along the complex client processing trajectory, termed the loading complex³ and the maturation complex². Together, they suggest a sliding or ratcheting movement of the client GR through the Hsp90 lumen. Likewise, three kinase clients (BRAF_{V600E}⁷, Cdk4⁴, RAF1⁸) and the aryl hydrocarbon receptor⁶ also thread through the Hsp90 lumen. Such molecular clamping by Hsp90 was suggested to keep the domains of multidomain clients separated, thus facilitating their independent folding² and preventing inter-domain misfolding (**Fig. 3b**). Destabilized kinases are thought to be held by Hsp90 to facilitate their reactivation⁵⁵, after local kinase motions poise the kinase for Hsp90 processing¹⁰⁰. During the processing of Tau, which binds to open Hsp90_{euk} in numerous conformations⁵¹, Hsp90 appears to act as a holdase, preventing the formation of toxic oligomeric species until Tau is targeted for degradation⁹¹. Overall, Hsp90's structural plasticity allows clients and cochaperones to individually orchestrate client-processing mechanisms, making it a flexible master chaperone.

The energetic driving forces for Hsp90's diverse molecular modes of action – folding, holding, stretching, etc. – result from different underlying kinetics of inter- and intra-molecular interactions. For example, the balance between client holding versus folding depends on the (forward and reverse) kinetic rate constants of the two processes: if the client-folding step is slower than client (re-)binding to the chaperone, the holdase function is kinetically favoured and folding is rate-limiting for that particular client¹⁰¹. The sequential refolding of individual structural elements of the client can decrease their local affinity for Hsp90, enabling gradual dissociation from Hsp90. Transient interactions are prerequisite for this dynamic model. Thermodynamically, transient interactions can result in avidity effects where multiple low-affinity contacts add up, resulting in high affinities with down to nanomolar dissociation constants (K_d)¹⁰². While such strong avidity is characterized by large interfaces (>~1500 Å²), the low affinities of Hsp90 to its clients (micromolar K_d) ^{51,52} likely result from much smaller interfaces.

11

For folding processes to occur, specific energy barriers need to be overcome. While small conformational changes have low energy barriers and occur spontaneously in thermal equilibrium, large rearrangements with higher energy barriers are not accessible by thermal energy alone (1 k_BT \approx 4.1x10⁻²¹ J \triangleq 0.6 kcal/mol). In other words, although folding is an exothermic process (cf. Anfinsen's dogma¹⁰³), energy barriers can kinetically trap un- or misfolded states and external energy is often required for efficient folding. This can be binding energy from molecular interactions (including cochaperone binding), ATP hydrolysis energy, or potential energy changes provided by covalent post-translational modifications (PTMs). Notably, for its client-specific function, Hsp90 exploits all of these modes of action in one unified system, and first efforts have been made to directly quantify the energies involved¹⁰⁴. Overall, Hsp90_{euk} has emerged as a highly multi-functional workbench that uses cochaperones as tools for the customized processing of its diverse clients. Many proposed models for client processing are yet to be experimentally proven. New routes to address this are discussed below and in **Box 1**.

The timing of Hsp90's functional cycle

The precise temporal progression through Hsp90's functional cycle is a matter of ongoing research (**Fig. 4**), complicated (amongst others) by Hsp90's client-specific behaviour, leading to many different functional cycles proposed in the literature^{4–6,69}. We focus here on two of the best-studied systems, the Hsp70-Hsp90 cascade and the kinase cycle, which both progress through three main phases: (i) client loading/recruitment, leading to (ii) client processing and maturation, followed by (iii) client release. Both examples show cochaperone-dependent client loading, but we note that several clients bind Hsp90 without cochaperone involvement, for example the IDPs Tau⁵¹ and alpha-synuclein¹⁰⁵. Also, due to the generally transient low-affinity interactions, numerous intermediate complexes and their interconversion kinetics are yet to be revealed and quantified. Therefore, as direct real-time observations are still lacking, the depicted cycles represent early models from which neither a strict order of events nor their timing can be inferred.



Figure 4: Hsp90's functional cycle timing illustrated by two literature examples. Schematic representation of the two discussed functional cycles of eukaryotic Hsp90, illustrated using six states each. Cryo-EM structures of two of the states exist, as indicated. Many transient interactions and intermediate complexes limit the current understanding of the order and timing of events, as well as their directionality or reversibility. (a) The Hsp70-Hsp90 cascade^{2,3,69,73,80} and (b) the Cdc37-kinase-Hsp90 cycle^{4,11}. Both cycles have in common that the client is (partially) unfolded prior to recruitment to Hsp90 in an intermediate loading complex. Next, by unknown mechanisms, clients are processed by Hsp90 through at least one intermediate maturation state. It is not completely understood, among other details, when ATP binding or hydrolysis by Hsp90 occurs, and how clients are released from these complexes.

The Hsp70-Hsp90 cascade (**Fig. 4a**) was found to process the transcription factors GR^{2,3,69,80} and p53¹⁰⁶, as well as Argonaute which regulates transcription through RNA silencing¹⁰⁷. Clients are prepared for Hsp90 interaction through (partial) unfolding by Hsp70 via hydrophobic interactions¹⁰⁸. Two recently resolved multi-partite intermediate states, the loading and the maturation complex^{2,3}, provide much awaited molecular-level structural details. The loading complex, Hsp70-Bridge-Client-Hsp90³, contains a 'bridging' protein (e.g., Hop⁸⁰, NudC¹⁰, Tomm34¹⁰⁹) that facilitates client transfer from Hsp70 to Hsp90. Interestingly, Hop was found not to be required *in vivo*¹¹⁰ which may reflect redundancy among bridging proteins. Another bridging protein, NudC, was found to directly recruit Hsp40-bound clients to Hsp90¹⁰. Following recruitment, the partially unfolded client is clamped within the closed Hsp90 dimer lumen, and whether this is a directional or reversible step still needs to be determined by time-resolved techniques. Next, the Hsp90-client-p23 maturation complex² is formed after dissociation of the bridging protein and Hsp70, where dissociation involves either nucleotide exchange or Hsp90 ATPase activity⁶⁹. p23 binds and stabilizes Hsp90's closed state¹¹¹ during GR processing, while it is not required for luciferase processing⁸⁰. Interestingly, a single-molecule FRET study showed directionality of an Hsp90/p23 system conferred by ATP binding and hydrolysis even in the absence of a client protein⁶⁹.

implying that cochaperones can steer the Hsp90 cycle. Additionally, p23 binding was found to relieve a Hop-inhibited Hsp70-Hsp40-Hop-GRLBD-Hsp90 complex and maximize GR hormone binding recovery beyond the levels achieved by the binding of either Hop or p23 alone¹¹², further indicating chaperoneendowed directionality in the Hsp70-Hsp90 cascade. The position of GR captured in the Hsp90 lumen shifts by several amino acids between the 3D-resolved loading and the maturation states, implying a sliding or ratcheting movement². Although not all molecular details are known yet (e.g., on client release), the Hsp70-Hsp90 cascade appears to enable client activation through assisted folding^{69,80}.

Another functional cycle has been described for kinase-specific processing^{4,11} (**Fig. 4b**). Here, Cdc37 plays the client-recruiting role, where the loading complex consists of Cdc37-Client-Hsp90. Cdc37 stabilizes a partially unfolded kinase state^{100,113}, in which the kinase N and C domain are separated⁴, and thereby primes the kinase client for subsequent interaction with residues in the Hsp90 dimer lumen⁵⁵. Cdc37 can bind Hsp90 in its open state, inhibiting Hsp90's ATPase through interaction with its ATP lid⁷⁰. In this loading complex, phosphorylation of serine 13 on Cdc37 (Cdc37- pSer13) by casein kinase 2¹¹⁴ is critical for kinase activation by Hsp90^{114,115}, and stabilizes the kinase-bound conformation⁴. In general, such post-translational modifications can affect the conformational dynamics, ATPase activity, client and cochaperone interactions, and other functions of Hsp90^{33,116}. Phosphatases and kinases regulate client maturation by changing the phosphorylation status of Hsp90^{117,118}: certain phosphorylations of Hsp90 represent allosteric regulation points¹¹⁹, others promote various interactions during the functional cycle¹²⁰. Cochaperone function is also regulated by phosphorylation, for example Hop phosphorylation at residue Y354 regulates its binding to Hsp70/90¹²¹.

The maturation complex is obtained by binding of Protein phosphatase 5 (Pp5), whereafter Pp5 dephosphorylates the kinase clients⁷ as well as Cdc37-pSer13 inducing client release^{122,123}. As Cdc37's pSer13 residue is buried in the closed, ATP-bound Hsp90 dimer⁷, the Pp5 dephosphorylation of Cdc37 likely happens upon Hsp90 transitioning from a compact ATP-bound state to a post-hydrolysis state, which is more flexible than the former¹²⁴. The Hsp90-Cdc37-kinase system has been described to process clients by "factory resetting" of phosphorylations⁷, as well as physical stabilization or shielding of clients⁵⁵.

The timing of functional cycles is known to be crucial for biomolecular systems and their precise regulation. Prominent examples include: the pausing mechanism of RNA polymerase^{125,126}, ribosome rotation, stalling, and frameshifting during translation¹²⁷, processive protein threading by proteases⁴⁹, etc. For Hsp90, this level of mechanistic understanding has not yet been reached. Up to now, only some of the state transitions of the functional cycle could be observed experimentally, leaving a range of questions unanswered: When and for what is ATP binding required? When does ATP hydrolysis take place? Is ATP binding or hydrolysis or rather cochaperone phosphorylation dominant in providing directionality – and hence processivity – to a given Hsp90 cycle? *In cellulo*, do cochaperones interact and dissociate from Hsp90 in a given sequence or randomly? What determines which processing? Clearly, the time-resolved direct observation of Hsp90's functional cycles (see below) would hold tremendous information about cycle timing, off-target waiting states, the client-specific differences, and the energetic driving forces of Hsp90's functional cycle.

Single-molecule observations complement the mechanistic understanding of Hsp90

Single-molecule methods have provided unique insight into Hsp90, and their abilities keep growing. By providing direct observations of individual proteins and protein complexes, they complement ensemble techniques (such as bulk enzymatic/activity assays, nuclear magnetic resonance, mass spectrometry, X-ray crystallography, etc.). Amongst others, single-molecule methods can uncover conformational heterogeneity normally hidden in ensemble averages, reveal reversible processes, elucidate the multiple reaction rates underlying the rate-limiting one, and quantitate the energies that drive molecular mechanisms. Amongst others, single-molecule Förster resonance energy transfer (smFRET)¹²⁸ and force spectroscopies¹²⁹ are popular techniques that provide time-resolved access to protein functional determinants, such as intra- and inter-molecular dynamics¹²⁸, molecular forces¹²⁹, enzymatic reactions¹³⁰, and movement⁴⁸ at single-molecule resolution.

Specifically, using smFRET, it was revealed that prokaryotic and eukaryotic Hsp90 are distinct regarding their conformational rearrangements, which are strongly ATP-hydrolysis dependent in the prokaryotic³⁸ but not eukaryotic^{37,131} homologue. Next, the conferral of directionality by ATP binding or hydrolysis in an Hsp90/p23 system⁶⁸ was uncovered by smFRET, and also how Hsp90's conformational kinetics are regulated by environmental conditions (e.g., the presence of point mutations, crowding agents, or cochaperones)⁶³. In addition, it could be shown that Hsp90 inhibitors have little effect on Hsp90's conformational dynamics¹³². Furthermore, smFRET revealed that Grp94 (a human Hsp90 isoform confined to the endoplasmic reticulum) exhibits two closed states, which was explained to arise from sequential ATP hydrolysis by the Hsp90 protomers¹³³. Also, smFRET elucidated that the yeast Hsp90 charged linker reduces the probability of the N-terminal dimerization conformation of the Hsp90 dimer as compared to Hsp90 mutants lacking charged residues in the linker²⁸. Moreover, smFRET showed that the p53 DNA-binding domain exhibits considerable dynamics in the presence of all required chaperoning components (i.e., Hsp70, Hsp40, ATP, HOP, Bag1, and Hsp90), suggesting that p53 conformation is constantly remodelled by Hsp70/Hsp90¹⁰⁶. In an integrative study, building on structural information, more than 100 FRET pairs on the Hsp90 dimer were used to disentangle local and global dynamics of the Hsp90 dimer, and it was found that the model client $\Delta 131\Delta$ only affects open state dynamics and does not affect open-closed interconversion dynamics¹³⁴. Another study connecting smFRET and molecular dynamics observations suggests that ATP hydrolysis leads to a conformational state in which the client binding site in the Hsp90 dimer lumen becomes constricted¹³⁵. A different fluorescence technique, single-molecule photo-induced electron transfer fluorescence correlation spectroscopy (PET-FCS) showed the degree of cooperative motion of Hsp90 domains as well as specific regions during various stages of Hsp90's catalytic cycle^{25,60}. Complementary to fluorescence, single-molecule force spectroscopy studies using optical tweezers have revealed the mechanism by which Hsp90 (itself) folds¹³⁶, and elucidated the functional role of its flexible charged linker region¹³⁷. Similar experiments have been used to perform a detailed comparison of Hsp90 orthologs, where large differences in the flexibility of the charged linker regions were identified¹³⁸. Also, Hsp90 isoforms were investigated by force spectroscopy, where despite a very high sequence similarity, differences in stability, refolding capacity and their conformational cycles were observed¹³⁹. Lastly, optical tweezer experiments revealed details of the ATP dependence of Hsp90's dimerization dynamics¹⁴⁰, of Hsp70mediated client unfolding¹¹², and the local compaction of the model client luciferase in an ATPdependent manner¹⁰⁴. Altogether, by building on a large body of existing biochemical work, singlemolecule techniques offer valuable complementary experiments that provided otherwise inaccessible insights through direct observations of single proteins at work.

Looking ahead, the future is bright for new approaches applied to Hsp90 – in part due to recent technological advances. For example, Hsp90's weak interactions with clients and cochaperones posed a major challenge of single-molecule experiments in the past. This regime can now be studied using nanophotonic approaches such as zero-mode waveguides (ZMW) – which overcome previous detection limitations for single-molecule fluorescence spectroscopy^{127,141}. Alternatively, tethered constructs, where the local concentration of a cochaperone, for example, is enhanced by connecting it to Hsp90 through an unstructured amino acid linker, are being used for both fluorescence spectroscopy and force-based measurements¹⁴². In fact, many of the urgent questions discussed herein could be clarified with the help of single-molecule experiments, and, in **Box 1**, we provide a non-exhaustive list with specific examples.

Box 1: New opportunities to elucidate Hsp90's enigmas with single-molecule resolution

(i) The order of interactions during Hsp90's functional cycle can be observed in real-time in singlemolecule fluorescence experiments, by recording the consecutive binding and dissociation of fluorescently labelled interactors to surface-tethered Hsp90. Zero-mode waveguides can be used to achieve single-molecule resolution at physiological concentrations.

(ii) Questions about Hsp90's ATPase and how it is coupled to Hsp90's conformational state can be elucidated by investigating the differences between prokaryotic and eukaryotic Hsp90 in this regard. smFRET and force spectroscopy can observe these differences in conformational dynamics and reveal the energies involved.

(iii) Client processing by Hsp90 can be observed at the single-molecule level in real-time, e.g., to probe the proposed sliding hypothesis for GR processing². Such nanometer distance changes induced by sliding or ratcheting can be observed by smFRET within one chaperone complex. Force spectroscopy can also be used to detect distance changes associated with client-remodelling.

(iv) Functional cycle timing, the order of individual events, and whether they occur directionally or reversibly can be observed by single-molecule fluorescence and dye-labelled interactors (e.g., Hsp90, client, Hsp70, p23, etc.) at physiological concentrations (using ZMW). The impact of clients and cochaperones on the conformational state of Hsp90 can also be probed by using force spectroscopy experiments combined with microfluidics.

Summary

In summary, the central chaperone Hsp90 is a peculiar ATPase differing drastically from other wellknown ATPases, and it has remained surprisingly enigmatic. We highlighted three fundamental unanswered questions on Hsp90. *First*, what is the purpose of Hsp90's ATPase activity? It has been revealed that Hsp90 can show widespread allostery in its hydrolysis mechanism. Additionally, Hsp90's ATP hydrolysis was proposed to induce dissociation of interaction partners, but *in cellulo*, it was found that nucleotide exchange (not hydrolysis) was sufficient for several Hsp90 chaperoning functions. *Second*, what are the molecular actions by which the Hsp90 system processes its clients? First molecular-level insights from cryo-EM structures provided static evidence for dynamic rearrangements, such as sliding, stretching, and holding – which represent valuable hypotheses to test with time-resolved observations, e.g.: using single-molecule techniques. *Third*, what are the kinetics of Hsp90's functional cycle, or rather its multiple client-specific cycles? I.e., what is the timing, order and interdependence of events that describe this multi-component system? With first molecular structures elucidating parts of the functional cycle plus ample biochemical results on the interaction partners involved, it is now possible to set up experiments to directly observe these functional cycles by following a single multipartite complex over time.

Altogether, mechanistic insight on the multifaceted chaperone function of Hsp90 is urgently needed to inform and accelerate biomedical and pharmaceutical advances, targeting several conditions ranging from cancer to neurodegenerative diseases and more. Compared to other intensely studied protein systems, the understanding of the Hsp90 chaperone lags behind, because its transient multipartite complexes have been challenging to study in molecular detail. Nevertheless, a large body of biochemical knowledge has accumulated over the years, 3D structural information on the transient Hsp90 complexes is growing, and recent technological advances enable now direct observation of Hsp90 at work. Equipped with these, the time is better than ever to address and uncover the remaining mechanistic known unknowns of the Hsp90 chaperone.

Acknowledgements

We thank Prof. Johannes Buchner for a critical reading of the manuscript and for his highly insightful comments and support. We regret not being able to cover all Hsp90 work related to our Review topic, and we acknowledge the work of all researchers working in the broad Hsp90 field. KT is supported by an MSCA-IF (NOTE, grant agreement number 101028366) and KT and SS are supported by the NWO grant (Solve90, grant agreement number OCENW.M.22.092).

References

- Picard, D. Curated list of Hsp90 Interactors. https://www.picard.ch/downloads/Hsp90interactors.pdf (2022).
- 2. Noddings, C. M., Wang, R. Y.-R., Johnson, J. L. & Agard, D. A. Structure of Hsp90–p23–GR reveals the Hsp90 client-remodelling mechanism. *Nature* **601**, 465–469 (2021).
- 3. Wang, R. Y.-R. *et al.* Structure of Hsp90–Hsp70–Hop–GR reveals the Hsp90 client-loading mechanism. *Nature* **601**, 460–464 (2021).
- 4. Verba, K. A. *et al.* Atomic structure of Hsp90-Cdc37-Cdk4 reveals that Hsp90 traps and stabilizes an unfolded kinase. *Science* **352**, 1542–1547 (2016).
- 5. Lee, K. *et al.* The structure of an Hsp90-immunophilin complex reveals cochaperone recognition of the client maturation state. *Mol Cell* **81**, 3496-3508.e5 (2021).
- 6. Wen, Z., Zhang, Y., Sun, F., Zhai, Y. & Zhu, Y. Cryo-EM structure of the cytosolic AhR complex. *Structure* **31**, 295–308 (2022).
- 7. Oberoi, J. *et al.* HSP90-CDC37-PP5 forms a structural platform for kinase dephosphorylation. *Nat Commun* **13**, 7343 (2022).
- 8. García-Alonso, S. *et al.* Structure of the RAF1-HSP90-CDC37 complex reveals the basis of RAF1 regulation. *Mol Cell* **82**, 3438-3452.e8 (2022).
- 9. Wegele, H., Wandinger, S. K., Schmid, A. B., Reinstein, J. & Buchner, J. Substrate transfer from the chaperone Hsp70 to Hsp90. *J Mol Biol* **356**, 802–811 (2006).
- 10. Biebl, M. M. *et al.* NudC guides client transfer between the Hsp40/70 and Hsp90 chaperone systems. *Mol Cell* **82**, 555-569.e7 (2022).
- 11. Taipale, M. *et al.* Quantitative analysis of Hsp90-client interactions reveals principles of substrate recognition. *Cell* **150**, 987–1001 (2012).
- 12. Pirkl, F. & Buchner, J. Functional analysis of the Hsp90-associated human peptidyl prolyl cis/trans isomerases FKBP51, FKBP52 and Cyp40. *J Mol Biol* **308**, 795–806 (2001).
- 13. Riggs, D. L. *et al.* Noncatalytic Role of the FKBP52 Peptidyl-Prolyl Isomerase Domain in the Regulation of Steroid Hormone Signaling. *Mol Cell Biol* **27**, 8658–8669 (2007).
- Broemer, M., Krappmann, D. & Scheidereit, C. Requirement of Hsp90 activity for IjB kinase (IKK) biosynthesis and for constitutive and inducible IKK and NF-jB activation. *Oncogene* 23, 5378–5386 (2004).

- Belova, L., Brickley, D. R., Ky, B., Sharma, S. K. & Conzen, S. D. Hsp90 regulates the phosphorylation and activity of serum- and glucocorticoid-regulated kinase-1. *J Biol Chem* 283, 18821–18831 (2008).
- 16. Dickey, C. A. *et al.* The high-affinity HSP90-CHIP complex recognizes and selectively degrades phosphorylated tau client proteins. *J Clin Investig* **117**, 648–658 (2007).
- 17. Chen, B., Zhong, D. & Monteiro, A. Comparative genomics and evolution of the HSP90 family of genes across all kingdoms of organisms. *BMC Genomics* **7**, 1–19 (2006).
- Rebeaud, M. E., Mallik, S., Goloubinoff, P. & Tawfik, D. S. On the evolution of chaperones and cochaperones and the expansion of proteomes across the Tree of Life. *Proc Natl Acad Sci U S A* 118, e2020885118 (2021).
- Lai, B. T., Chin, N. W., Stanek, A. E., Keh, W. & Lanks, K. W. Quantitation and intracellular localization of the 85K heat shock protein by using monoclonal and polyclonal antibodies. *Mol Cell Biol* 4, 2802–2810 (1984).
- 20. Finka, A. & Goloubinoff, P. Proteomic data from human cell cultures refine mechanisms of chaperone-mediated protein homeostasis. *Cell Stress Chaperones* **18**, 591–605 (2013).
- 21. Borkovich, K. A., Farrelly, F. W., Finkelstein, D. B., Taulien, J. & Lindquist', S. hsp82 is an essential protein that is required in higher concentrations for growth of cells at higher temperatures. *Mol Cell Biol* **9**, 3919–3930 (1989).
- 22. Wayne, N. & Bolon, D. N. Dimerization of Hsp90 is required for in vivo function: Design and analysis of monomers and dimers. *J Biol Chem* **282**, 35386–35395 (2007).
- 23. Ritossa, F. A new puffing pattern induced by temperature shock and DNP in drosophila. *Experientia* **18**, 571–573 (1962).
- 24. Finkelstein, D. B. & Strausberg, S. Identification and Expression of a Cloned Yeast Heat Shock Gene. *J Biol Chem* **258**, 1908–1913 (1983).
- 25. Schulze, A. *et al.* Cooperation of local motions in the Hsp90 molecular chaperone ATPase mechanism. *Nat Chem Biol* **12**, 628–635 (2016).
- 26. Henot, F. *et al.* Visualizing the transiently populated closed-state of human HSP90 ATP binding domain. *Nat Commun* **13**, 1–13 (2022).
- 27. Hainzl, O., Lapina, M. C., Buchner, J. & Richter, K. The Charged Linker Region Is an Important Regulator of Hsp90 Function. *J Biol Chem* **284**, 22559–22567 (2009).
- 28. Jahn, M. *et al.* The charged linker of the molecular chaperone Hsp90 modulates domain contacts and biological function. *Proc Natl Acad Sci U S A* **111**, 17881–17886 (2014).
- 29. López, A., Elimelech, A. R., Klimm, K. & Sattler, M. The Charged Linker Modulates the Conformations and Molecular Interactions of Hsp90. *ChemBioChem* **22**, 1084–1092 (2021).
- 30. Biebl, M. M. & Buchner, J. Structure, function, and regulation of the Hsp90 machinery. *Cold Spring Harb Perspect Biol* **11**, a034017 (2019).
- 31. Mader, S. L. *et al.* Conformational dynamics modulate the catalytic activity of the molecular chaperone Hsp90. *Nat Commun* **11**, 1–12 (2020).

- 32. Minami, Y., Kimura, Y., Kawasaki, H., Suzuki, K. & Yahara, I. The carboxy-terminal region of mammalian HSP90 is required for its dimerization and function in vivo. *Mol Cell Biol* **14**, 1459–1464 (1994).
- 33. Mayer, M. P. & le Breton, L. Hsp90: Breaking the symmetry. *Mol Cell* 58, 8–20 (2015).
- 34. Schopf, F. H., Biebl, M. M. & Buchner, J. The HSP90 chaperone machinery. *Nat Rev Mol Cell Biol* **18**, 345–360 (2017).
- 35. Ali, M. M. U. *et al.* Crystal structure of an Hsp90-nucleotide-p23/Sba1 closed chaperone complex. *Nature* **440**, 1013–1017 (2006).
- Shiau, A. K., Harris, S. F., Southworth, D. R. & Agard, D. A. Structural Analysis of E. coli hsp90 Reveals Dramatic Nucleotide-Dependent Conformational Rearrangements. *Cell* **127**, 329–340 (2006).
- 37. Mickler, M., Hessling, M., Ratzke, C., Buchner, J. & Hugel, T. The large conformational changes of Hsp90 are only weakly coupled to ATP hydrolysis. *Nat Struct Mol Biol* **16**, 281–286 (2009).
- 38. Ratzke, C., Nguyen, M. N. T., Mayer, M. P. & Hugel, T. From a ratchet mechanism to random fluctuations evolution of Hsp90's mechanochemical cycle. *J Mol Biol* **423**, 462–471 (2012).
- 39. Röhl, A., Rohrberg, J. & Buchner, J. The chaperone Hsp90: Changing partners for demanding clients. *Trends Biochem Sci* **38**, 253–262 (2013).
- 40. Panaretou, B. *et al.* ATP binding and hydrolysis are essential to the function of the Hsp90 molecular chaperone in vivo. *EMBO Journal* **17**, 4829–4836 (1998).
- 41. Siligardi, G. *et al.* Regulation of Hsp90 ATPase Activity by the Co-chaperone Cdc37p/p50 cdc37 *. *J Biol Chem* **277**, 20151–20159 (2002).
- 42. Richter, K., Muschler, P., Hainzl, O. & Buchner, J. Coordinated ATP Hydrolysis by the Hsp90 Dimer. *J Biol Chem* **276**, 33689–33696 (2001).
- 43. Surette, M. G. *et al.* Dimerization Is Required for the Activity of the Protein Histidine Kinase CheA That Mediates Signal Transduction in Bacterial Chemotaxis. *J Biol Chem* **271**, 939–945 (1996).
- 44. Childers, W. S. *et al.* Cell Fate Regulation Governed by a Repurposed Bacterial Histidine Kinase. *PLoS Biol* **12**, e1001979 (2014).
- 45. Yang, X. W. *et al.* MutS functions as a clamp loader by positioning MutL on the DNA during mismatch repair. *Nat Commun* **13**, 1–17 (2022).
- 46. Maxwell, A. & Gellert, M. The DNA dependence of the ATPase activity of DNA gyrase. *J Biol Chem* **259**, 14472–14480 (1984).
- 47. Piazzesi, G. *et al.* Mechanism of force generation by myosin heads in skeletal muscle. *Nature* **415**, 659–662 (2002).
- 48. Yildiz, A., Tomishige, M., Vale, R. D. & Selvin, P. R. Kinesin Walks Hand-Over-Hand. *Science* **303**, 676–678 (2004).
- 49. Cordova, J. C. *et al.* Stochastic but Highly Coordinated Protein Unfolding and Translocation by the ClpXP Proteolytic Machine. *Cell* **158**, 647–658 (2014).

- 50. Gradia, S. *et al.* hMSH2–hMSH6 Forms a Hydrolysis-Independent Sliding Clamp on Mismatched DNA. *Mol Cell* **3**, 255–261 (1999).
- 51. Karagöz, G. E. *et al.* Hsp90-tau complex reveals molecular basis for specificity in chaperone action. *Cell* **156**, 963–974 (2014).
- 52. Lorenz, O. R. *et al.* Modulation of the Hsp90 Chaperone Cycle by a Stringent Client Protein. *Mol Cell* **53**, 941–953 (2014).
- 53. Prodromou, C., Roe, S. M., Piper, P. W. & Pearl, L. H. A molecular clamp in the crystal structure of the N-terminal domain of the yeast Hsp90 chaperone. *Nat Struct Biol* **4**, 477–482 (1997).
- 54. Karagöz, G. E. *et al.* Hsp90-tau complex reveals molecular basis for specificity in chaperone action. *Cell* **156**, 963–974 (2014).
- 55. Verba, K. A. & Agard, D. A. How Hsp90 and Cdc37 Lubricate Kinase Molecular Switches. *Trends Biochem Sci* **42**, 799–811 (2017).
- 56. Xiao, Y. & Liu, Y. Recent Advances in the Discovery of Novel HSP90 Inhibitors: An Update from 2014. *Curr Drug Targets* **21**, 302–317 (2019).
- 57. Chiosis, G., Digwal, C. S., Trepel, J. B. & Neckers, L. Structural and functional complexity of HSP90 in cellular homeostasis and disease. *Nature Reviews Molecular Cell Biology*, (2023).
- 58. Dutta, R. & Inouye, M. GHKL, an emergent ATPase/kinase superfamily. *Trends Biochem Sci* **25**, 24–28 (2000).
- 59. Prodromou, C. *et al.* Identification and structural characterization of the ATP/ADP-binding site in the Hsp90 molecular chaperone. *Cell* **90**, 65–75 (1997).
- 60. Schubert, J., Schulze, A., Prodromou, C. & Neuweiler, H. Two-colour single-molecule photoinduced electron transfer fluorescence imaging microscopy of chaperone dynamics. *Nat Commun* **12**, (2021).
- 61. Richter, K. *et al.* Conserved conformational changes in the ATPase cycle of human Hsp90. *J Biol Chem* **283**, 17757–17765 (2008).
- 62. Hessling, M., Richter, K. & Buchner, J. Dissection of the ATP-induced conformational cycle of the molecular chaperone Hsp90. *Nat Struct Mol Biol* **16**, 287–293 (2009).
- 63. Schmid, S. & Hugel, T. Controlling protein function by fine-tuning conformational flexibility. *Elife* **9**, e57180 (2020).
- 64. Weikl, T. *et al.* C-terminal regions of Hsp90 are important for trapping the nucleotide during the ATPase cycle. *J Mol Biol* **303**, 583–592 (2000).
- 65. Ratzke, C., Berkemeier, F. & Hugel, T. Heat shock protein 90's mechanochemical cycle is dominated by thermal fluctuations. *Proc Natl Acad Sci U S A* **109**, 161–166 (2012).
- 66. Zierer, B. K. *et al.* Importance of cycle timing for the function of the molecular chaperone Hsp90. *Nat Struct Mol Biol* **23**, 1020–1028 (2016).
- 67. Lopez, A. *et al.* Client binding shifts the populations of dynamic Hsp90 conformations through an allosteric network. *Sci Adv* **7**, 1–19 (2021).

- 68. Ratzke, C., Hellenkamp, B. & Hugel, T. Four-colour FRET reveals directionality in the Hsp90 multicomponent machinery. *Nat Commun* **5**, (2014).
- 69. Kirschke, E., Goswami, D., Southworth, D., Griffin, P. R. & Agard, D. A. Glucocorticoid receptor function regulated by coordinated action of the Hsp90 and Hsp70 chaperone cycles. *Cell* **157**, 1685–1697 (2014).
- 70. Roe, S. M. *et al.* The Mechanism of Hsp90 Regulation by the Protein Kinase-Specific Cochaperone p50cdc37. *Cell* **116**, 87–98 (2004).
- 71. D'Annessa, I., Moroni, E. & Colombo, G. Visualizing the Dynamics of a Protein Folding Machinery: The Mechanism of Asymmetric ATP Processing in Hsp90 and its Implications for Client Remodelling. *J Mol Biol* **433**, (2021).
- 72. Radli, M. & Rüdiger, S. G. D. Dancing with the Diva: Hsp90–Client Interactions. *J Mol Biol* **430**, 3029–3040 (2018).
- 73. Morán Luengo, T., Mayer, M. P. & Rüdiger, S. G. D. The Hsp70–Hsp90 Chaperone Cascade in Protein Folding. *Trends Cell Biol* **29**, 164–177 (2019).
- 74. Walerych, D. *et al.* ATP Binding to Hsp90 Is Sufficient for Effective Chaperoning of p53 Protein. *J Biol Chem* **285**, 32020–32028 (2010).
- 75. Boczek, E. E. *et al.* Conformational processing of oncogenic v-Src kinase by the molecular chaperone Hsp90. *Proc Natl Acad Sci U S A* **112**, E3189–E3198 (2015).
- 76. Falsone, S. F., Kungl, A. J., Rek, A., Cappai, R. & Zangger, K. The molecular chaperone Hsp90 modulates intermediate steps of amyloid assembly of the Parkinson-related protein α-synuclein. J Biol Chem 284, 31190–31199 (2009).
- 77. Street, T. O., Lavery, L. A. & Agard, D. A. Substrate Binding Drives Large-Scale Conformational Changes in the Hsp90 Molecular Chaperone. *Mol Cell* **42**, 96–105 (2011).
- Obermann, W. M. J., Sondermann, H., Russo, A. A., Pavletich, N. P. & Hartl, F. U. In vivo function of Hsp90 is dependent on ATP binding and ATP hydrolysis. *Journal of Cell Biology* 143, 901–910 (1998).
- 79. Reidy, M., Garzillo, K. & Masison, D. C. Nucleotide exchange is sufficient for Hsp90 functions in vivo. *Nat Commun* **14**, (2023).
- 80. Morán Luengo, T., Kityk, R., Mayer, M. P. & Rüdiger, S. G. D. Hsp90 Breaks the Deadlock of the Hsp70 Chaperone System. *Mol Cell* **70**, 545–552 (2018).
- 81. Xu, Y. & Lindquist, S. Heat-shock protein hsp90 governs the activity of pp60v-src kinase. *Proc Natl Acad Sci U S A* **90**, 7074–7078 (1993).
- Pratt, W. B. & Dittmar, K. D. Studies with Purified Chaperones Advance the Understanding of the Mechanism of Glucocorticoid Receptor–hsp90 Heterocomplex Assembly. *Trends in Endocrinology & Metabolism* 9, 244–252 (1998).
- Ghosh, A. & Stuehr, D. J. Soluble guanylyl cyclase requires heat shock protein 90 for heme insertion during maturation of the NO-active enzyme. *Proc Natl Acad Sci U S A* **109**, 12998–13003 (2012).

- 84. Zhao, R. *et al.* Molecular chaperone Hsp90 stabilizes Pih1/Nop17 to maintain R2TP complex activity that regulates snoRNA accumulation. *Journal of Cell Biology* **180**, 563–578 (2008).
- 85. Manjarrez, J. R., Sun, L., Prince, T. & Matts, R. L. Hsp90-Dependent Assembly of the DBC2/RhoBTB2-Cullin3 E3-Ligase Complex. *PLoS One* **9**, e90054 (2014).
- 86. Sato, S., Fujita, N. & Tsuruo, T. Modulation of Akt kinase activity by binding to Hsp90. *Proc Natl Acad Sci U S A* **97**, 10832–10837 (2000).
- 87. Pearl, L. H. & Prodromou, C. Structure and Mechanism of the Hsp90 Molecular Chaperone Machinery. *Annu Rev Biochem* **75**, 271–294 (2006).
- 88. McClellan, A. J. *et al.* Diverse Cellular Functions of the Hsp90 Molecular Chaperone Uncovered Using Systems Approaches. *Cell* **131**, 121–135 (2007).
- 89. Rüdiger, S., Buchberger, A. & Bukau, B. Interaction of Hsp70 chaperones with substrates. *Nat Struct Biol* **4**, 342–349 (1997).
- 90. Oroz, J., Kim, J. H., Chang, B. J. & Zweckstetter, M. Mechanistic basis for the recognition of a misfolded protein by the molecular chaperone Hsp90. *Nat Struct Mol Biol* **24**, 407–413 (2017).
- 91. Rutledge, B. S., Choy, W. Y. & Duennwald, M. L. Folding or holding?—Hsp70 and Hsp90 chaperoning of misfolded proteins in neurodegenerative disease. *J Biol Chem* **298**, 101905 (2022).
- 92. Liu, Y. *et al.* Cryo-EM structures reveal a multistep mechanism of Hsp90 activation by cochaperone Aha1. *J Chem Inf Model* **53**, 1689–1699 (2020).
- 93. Sahasrabudhe, P., Rohrberg, J., Biebl, M. M., Rutz, D. A. & Buchner, J. The Plasticity of the Hsp90 Co-chaperone System. *Mol Cell* **67**, 947-961.e5 (2017).
- 94. Radli, M. & Rüdiger, S. G. D. Picky Hsp90—Every Game with Another Mate. *Mol Cell* **67**, 899– 900 (2017).
- 95. Prince, T. & Neckers, L. A Network of Its Own: The Unique Interactome of the Hsp90 Cochaperone, Sba1/p23. *Mol Cell* **43**, 159–160 (2011).
- 96. Koulov, A. v. *et al.* Biological and structural basis for Aha1 regulation of Hsp90 ATPase activity in maintaining proteostasis in the human disease cystic fibrosis. *Mol Biol Cell* **21**, 871–884 (2010).
- 97. Biebl, M. M., Riedl, M. & Buchner, J. Hsp90 Co-chaperones Form Plastic Genetic Networks Adapted to Client Maturation. *Cell Rep* **32**, 108063 (2020).
- 98. Wang, R. Y. R. *et al.* Structure of Hsp90–Hsp70–Hop–GR reveals the Hsp90 client-loading mechanism. *Nature* **601**, 460–464 (2022).
- 99. Noddings, C. M., Wang, R. Y. R., Johnson, J. L. & Agard, D. A. Structure of Hsp90–p23–GR reveals the Hsp90 client-remodelling mechanism. *Nature* **601**, 465–469 (2022).
- 100. Keramisanou, D., Kumar, M. V. V., Boose, N., Abzalimov, R. R. & Gelis, I. Assembly mechanism of early Hsp90-Cdc37-kinase complexes. *Sci Adv* **8**, 1–14 (2022).
- 101. Kawagoe, S., Ishimori, K. & Saio, T. Structural and Kinetic Views of Molecular Chaperones in Multidomain Protein Folding. *Int J Mol Sci* **23**, (2022).

- 102. Nooren, I. M. A. & Thornton, J. M. Structural characterisation and functional significance of transient protein-protein interactions. *J Mol Biol* **325**, 991–1018 (2003).
- 103. Anfinsen, C. B. Principles that Govern the Folding of Protein Chains. *Science* **181**, 223–230 (1973).
- 104. Mashaghi, A. *et al.* Direct observation of Hsp90-induced compaction in a protein chain. *Cell Rep* **41**, 111734 (2022).
- 105. Daturpalli, S., Waudby, C. A., Meehan, S. & Jackson, S. E. Hsp90 inhibits α-synuclein aggregation by interacting with soluble oligomers. *J Mol Biol* **425**, 4614–4628 (2013).
- 106. Dahiya, V. *et al.* Coordinated Conformational Processing of the Tumor Suppressor Protein p53 by the Hsp70 and Hsp90 Chaperone Machineries. *Mol Cell* **74**, 816-830.e7 (2019).
- Tsuboyama, K., Tadakuma, H. & Tomari, Y. Conformational Activation of Argonaute by Distinct yet Coordinated Actions of the Hsp70 and Hsp90 Chaperone Systems. *Mol Cell* 70, 722-729.e4 (2018).
- Rüdiger, S., Germeroth, L., Schneider-Mergener, J. & Bukau, B. Substrate specificity of the DnaK chaperone determined by screening cellulose-bound peptide libraries. *EMBO J* 16, 1501–1507 (1997).
- 109. Trcka, F. *et al.* The Assembly and Intermolecular Properties of the Hsp70-Tomm34-Hsp90 Molecular Chaperone Complex. *J Biol Chem* **289**, 9887–9901 (2014).
- 110. Bhattacharya, K. *et al.* The Hsp70-Hsp90 co-chaperone Hop/Stip1 shifts the proteostatic balance from folding towards degradation. *Nat Commun* **11**, 5975 (2020).
- 111. Richter, K., Walter, S. & Buchner, J. The Co-chaperone Sba1 Connects the ATPase Reaction of Hsp90 to the Progression of the Chaperone Cycle. *J Mol Biol* **342**, 1403–1413 (2004).
- 112. Dahiya, V. *et al.* The switch from client holding to folding in the Hsp70/Hsp90 chaperone machineries is regulated by a direct interplay between co-chaperones. *Mol Cell* **82**, 1543-1556.e6 (2022).
- 113. Keramisanou, D. *et al.* Molecular Mechanism of Protein Kinase Recognition and Sorting by the Hsp90 Kinome-Specific Cochaperone Cdc37. *Mol Cell* **62**, 260–271 (2016).
- 114. Miyata, Y. & Nishida, E. CK2 Controls Multiple Protein Kinases by Phosphorylating a Kinase-Targeting Molecular Chaperone, Cdc37. *Mol Cell Biol* **24**, 4065–4074 (2004).
- Shao, J., Prince, T., Hartson, S. D. & Matts, R. L. Phosphorylation of Serine 13 Is Required for the Proper Function of the Hsp90 Co-chaperone, Cdc37*. *J Biol Chem* 278, 38117–38120 (2003).
- Backe, S. J., Sager, R. A., Woodford, M. R., Makedon, A. M. & Mollapour, M. Post-translational modifications of Hsp90 and translating the chaperone code. *J Biol Chem* **295**, 11099–11117 (2020).
- 117. Duval, M., le Bœuf, F., Huot, J. & Gratton, J. P. Src-mediated phosphorylation of Hsp90 in response to vascular endothelial growth factor (VEGF) is required for VEGF receptor-2 signaling to endothelial NO synthase. *Mol Biol Cell* **18**, 4659–4668 (2007).

- 118. Zuehlke, A. D. *et al.* An Hsp90 co-chaperone protein in yeast is functionally replaced by sitespecific posttranslational modification in humans. *Nat Commun* **8**, 1–10 (2017).
- 119. Rehn, A. *et al.* Allosteric Regulation Points Control the Conformational Dynamics of the Molecular Chaperone Hsp90. *J Mol Biol* **428**, 4559–4571 (2016).
- 120. Xu, W. *et al.* Dynamic Tyrosine Phosphorylation Modulates Cycling of the HSP90-P50 CDC37-AHA1 Chaperone Machine. *Mol Cell* **47**, 434–443 (2012).
- 121. Castelli, M. *et al.* Phosphorylation of the Hsp90 Co-Chaperone Hop Changes its Conformational Dynamics and Biological Function. *J Mol Biol* **435**, (2023).
- 122. Vaughan, C. K. *et al.* Hsp90-Dependent Activation of Protein Kinases Is Regulated by Chaperone-Targeted Dephosphorylation of Cdc37. *Mol Cell* **31**, 886–895 (2008).
- 123. Oberoi, J. *et al.* Structural and functional basis of protein phosphatase 5 substrate specificity. *Proc Natl Acad Sci U S A* **113**, 9009–9014 (2016).
- 124. Graf, C., Stankiewicz, M., Kramer, G. & Mayer, M. P. Spatially and kinetically resolved changes in the conformational dynamics of the Hsp90 chaperone machine. *EMBO Journal* **28**, 602–613 (2009).
- 125. Lerner, E. *et al.* Backtracked and paused transcription initiation intermediate of Escherichia coli RNA polymerase. *Proc Natl Acad Sci U S A* **113**, E6562–E6571 (2016).
- 126. Janissen, R., Eslami-Mossallam, B., Artsimovitch, I., Depken, M. & Dekker, N. Accounting for RNA polymerase heterogeneity reveals state switching and two distinct long-lived backtrack states escaping through cleavage. *Biophys J* **118**, 543a (2020).
- Tsai, A., Puglisi, J. D. & Uemura, S. Probing the Translation Dynamics of Ribosomes Using Zero-Mode Waveguides. in *Progress in Molecular Biology and Translational Science* vol. 139 1–43 (Elsevier Inc., 2016).
- 128. Lerner, E. *et al.* Toward dynamic structural biology: Two decades of single-molecule förster resonance energy transfer. *Science* **359**, (2018).
- 129. Bustamante, C. J., Chemla, Y. R., Liu, S. & Wang, M. D. Optical tweezers in single-molecule biophysics. *Nature Reviews Methods Primers* **1**, (2021).
- 130. Lu, H. P., Xun, L. & Xie, X. S. Single-Molecule Enzymatic Dynamics. *Science* **282**, 1877–1882 (1998).
- 131. Ratzke, C., Berkemeier, F. & Hugel, T. Heat shock protein 90's mechanochemical cycle is dominated by thermal fluctuations. *Proc Natl Acad Sci U S A* **109**, 161–166 (2012).
- 132. Schmid, S., Götz, M. & Hugel, T. Effects of Inhibitors on Hsp90's Conformational Dynamics, Cochaperone and Client Interactions. *ChemPhysChem* **19**, 1716–1721 (2018).
- 133. Huang, B., Friedman, L. J., Sun, M., Gelles, J. & Street, T. O. Conformational Cycling within the Closed State of Grp94, an Hsp90-Family Chaperone. *J Mol Biol* **431**, 3312–3323 (2019).
- Hellenkamp, B., Wortmann, P., Kandzia, F., Zacharias, M. & Hugel, T. Multidomain structure and correlated dynamics determined by self-consistent FRET networks. *Nat Methods* 14, 176– 182 (2017).

- 135. Wolf, S. *et al.* Hierarchical dynamics in allostery following ATP hydrolysis monitored by single molecule FRET measurements and MD simulations. *Chemical and Engineering News* **68**, 42–61 (2021).
- Jahn, M., Buchner, J., Hugel, T. & Rief, M. Folding and assembly of the large molecular machine Hsp90 studied in single-molecule experiments. *Proc Natl Acad Sci U S A* **113**, 1232– 1237 (2016).
- 137. Jahn, M. *et al.* The charged linker of the molecular chaperone Hsp90 modulates domain contacts and biological function. *Proc Natl Acad Sci U S A* **111**, 17881–17886 (2014).
- 138. Jahn, M. *et al.* Folding and Domain Interactions of Three Orthologs of Hsp90 Studied by Single-Molecule Force Folding and Domain Interactions of Three Orthologs of Hsp90 Studied by Single-Molecule Force Spectroscopy. *Structure* **26**, 96–105 (2018).
- 139. Girstmair, H. *et al.* The Hsp90 isoforms from S. cerevisiae differ in structure, function and client range. *Nat Commun* **10**, (2019).
- 140. Tych, K. M. *et al.* Nucleotide-Dependent Dimer Association and Dissociation of the Chaperone Hsp90. *Journal of Physical Chemistry B* **122**, 11373–11380 (2018).
- 141. Eid, J. *et al.* Real-time DNA sequencing from single polymerase molecules. *Science* **323**, 133–138 (2009).
- 142. Mondol, T. *et al.* Aha1 regulates Hsp90's conformation and function in a stoichiometrydependent way. *Biophys J* **In Press**, (2023).
- 143. Lorenz, O. R. *et al.* Modulation of the Hsp90 chaperone cycle by a stringent client protein. *Mol Cell* **53**, 941–953 (2014).
- 144. Falsone, S. F., Kungl, A. J., Rek, A., Cappai, R. & Zangger, K. The molecular chaperone Hsp90 modulates intermediate steps of amyloid assembly of the Parkinson-related protein α-synuclein. *Journal of Biological Chemistry* **284**, 31190–31199 (2009).
- 145. Hunter, B. & Allingham, J. S. These motors were made for walking. *Protein Science* **29**, 1707–1723 (2020).
- 146. Hackney, D. D. Kinesin ATPase: Rate-limiting ADP release. *Proc Natl Acad Sci U S A* **85**, 6314–6318 (1988).
- Aubin-Tam, M. E., Olivares, A. O., Sauer, R. T., Baker, T. A. & Lang, M. J. Single-Molecule Protein Unfolding and Translocation by an ATP-Fueled Proteolytic Machine. *Cell* 145, 257–267 (2011).
- 148. Henderson, A., Erales, J., Hoyt, M. A. & Coffino, P. Dependence of proteasome processing rate on substrate unfolding. *J Biol Chem* **286**, 17495–17502 (2011).
- 149. Rajendran, S., Jezewska, M. J. & Bujalowski, W. Multiple-step kinetic mechanism of DNAindependent ATP binding and hydrolysis by Escherichia coli replicative helicase DnaB protein: quantitative analysis using the rapid quench-flow method. *J Mol Biol* **303**, 773–795 (2000).
- 150. Woehlke, G. *et al.* Microtubule interaction site of the kinesin motor. *Cell* **90**, 207–216 (1997).

- 151. Biswas, S. B. & Biswas-Fiss, E. E. Quantitative analysis of binding of single-stranded DNA by Escherichia coli DnaB helicase and the DnaB·DnaC complex. *Biochemistry* **45**, 11505–11513 (2006).
- 152. Singh, S. K., Grimaud, R., Hoskins, J. R., Wickner, S. & Maurizi, M. R. Unfolding and internalization of proteins by the ATP-dependent proteases ClpXP and ClpAP. *Proc Natl Acad Sci U S A* **97**, 8898–8903 (2000).
- 153. Baker, T. A. & Sauer, R. T. ClpXP, an ATP-powered unfolding and protein-degradation machine. *Biochim Biophys Acta* **1823**, 15–28 (2012).
- 154. Wiegand, T. *et al.* The conformational changes coupling ATP hydrolysis and translocation in a bacterial DnaB helicase. *Nat Commun* **10**, (2019).
- 155. Hancock, W. O. & Howard, J. Kinesin's processivity results from mechanical and chemical coordination between the ATP hydrolysis cycles of the two motor domains. *Proc Natl Acad Sci U S A* **96**, 13147–13152 (1999).
- 156. Biswas, E. E. & Biswas, S. B. Mechanism of DnaB helicase of Escherichia coli: Structural domains involved in ATP hydrolysis, DNA binding, and oligomerization. *Biochemistry* **38**, 10919–10928 (1999).
- 157. Chan, C. M., Hahn, E. & Zuber, P. Adaptor bypass mutations of Bacillus subtilis spx suggest a mechanism for YjbH-enhanced proteolysis of the regulator Spx by ClpXP. *Mol Microbiol* **93**, 426–438 (2014).