Strong solutions for the Navier-Stokes-Voigt equations with non-negative density

H.B. de Oliveira^{1,2}, Kh. Khompysh³, A.G. Shakir³

¹FCT - Universidade do Algarve, Faro, Portugal ²CMAFcIO - Universidade de Lisboa, Lisbon, Portugal ³Al-Farabi Kazakh National University, Almaty, Kazakhstan

ABSTRACT. The aim of this work is to study the Navier-Stokes-Voigt equations that govern flows with non-negative density of incompressible fluids with elastic properties. For the associated non-linear initial-and boundary-value problem, we prove the global-in-time existence of strong solutions (velocity, density and pressure). We also establish some other regularity properties of these solutions and find the conditions that guarantee the uniqueness of velocity and density. The main novelty of this work is the hypothesis that, in some subdomain of space, there may be a vacuum at the initial moment, that is, the possibility of the initial density vanishing in some part of the space domain.

Keywords: incompressible Navier-Stokes-Voigt equations; non-negative density; strong solutions; regularity; uniqueness.

MSC (2020): 35Q35; 76D03; 76A05; 76N10.

1. INTRODUCTION

In general terms, it can be said that the Navier-Stokes equations describe the evolution of the velocity field of an incompressible viscous fluid in the laminar regime. It is one of the most important system of equations in mathematical physics and therefore has been widely studied by several authors during the last 120 years, either from the mathematical viewpoint or from the applications - see e.g. [10, 11, 15, 17, 19]. However the developed theory is still incomplete, specially in 3d, where general existence and uniqueness results for smooth solutions are still partial. For this reason, many authors began to study slightly modified models of these equations for which it would be possible to prove the existence and uniqueness of smooth solutions. Among these models are the so-called Navier-Stokes-Voigt equations for which it is possible to answer the still open questions of the Navier-Stokes equations. In turn, the main feature of the Navier-Stokes-Voigt equations is that they can be used to model flows of viscous fluids with elastic properties, as for instance polymer solutions. These materials are part of a wider class of fluids, called viscoelastic fluids, and can exhibit all intermediate ranges of properties between an elastic solid and a viscous fluid. Typically, fluids that exhibit this behavior are macromolecular in nature and the most common examples are polymeric melts and solutions used to make plastic articles, food products, such as dough used to make bread and pasta, and biological fluids such as synovial fluids found in joints – see e.g. [10, 17]. In most applications, density is considered a constant parameter. However, this assumption is unrealistic, because in almost all fluids the density varies, either with time or with the position of

E-mail address: holivei@ualg.pt, konat_k@mail.ru, ajdossakir@gmail.com. *Date*: September 4, 2023.

the space where the fluid element is. These variations may or may not be significant. There are situations in which the density cannot be assumed as a constant parameter as, for example, in the study of multi-phase flows consisting of several immiscible and incompressible fluids.

In this work, we are interested in studying density-dependent flows (nonhomogeneous flows) of incompressible fluids with elastic properties. We assume the flow is governed by the following initial-and boundary-value problem,

$$\operatorname{div} u = 0 \qquad \text{in} \quad Q_T, \tag{1.1}$$

$$(\rho u)_t + \operatorname{div}(\rho u \otimes u) = \rho f - \nabla p + \mu \Delta u + \kappa \Delta u_t \quad \text{in} \quad Q_T, \tag{1.2}$$

$$\rho_t + \operatorname{div}(\rho u) = 0, \quad \rho \ge 0 \qquad \text{in} \quad Q_T, \tag{1.3}$$

$$\rho u = \rho_0 u_0, \quad \rho = \rho_0 \qquad \text{in} \quad \{0\} \times \Omega, \tag{1.4}$$

$$u = 0$$
 on Γ_T . (1.5)

Here, Q_T denotes the time-space cylinder $(0,T) \times \Omega$, where $\Omega \subset \mathbb{R}^3$ is a bounded domain and T is a given positive constant. The lateral boundary $(0,T) \times \partial \Omega$ of Q_T is denoted by Γ_T , and the other notation is defined as follows. The unknowns of the problem are the velocity $u = (u_1, u_2, u_3)$, density ρ and pressure p, while the external forces field f, initial velocity u_0 and initial density ρ_0 are given data. Equations (1.2) and (1.3) are derived from the classical principles of conservation of momentum and mass, while (1.1) expresses the incompressibility constraint of the fluid. The system of equations (1.1)-(1.3) shall be denoted in the sequel as the Navier-Stokes-Voigt equations, where μ corresponds to the dynamic viscosity and κ to the relaxation time, that is the characteristic time required for a viscoelastic fluid to relax from a deformed state to its equilibrium configuration. With this respect, we assume that they are both constant and such that

$$\mu > 0, \quad \kappa > 0. \tag{1.6}$$

We assume the initial momentum, say m_0 , is given by the product $\rho_0 u_0$, where u_0 is the given initial velocity. In this work, we are interested in the case of initial data u_0 and ρ_0 satisfying

$$\operatorname{div} u_0 = 0 \quad \text{in} \quad \Omega, \tag{1.7}$$

$$0 \le \rho_0 \le M < \infty \quad \text{in} \quad \Omega, \tag{1.8}$$

for some positive constant M. The main novelty of this work lies in the assumption (1.8), where it is understood that the initial density ρ_0 may eventually vanish in some domain $\omega \subset \Omega$, i.e. the possibility that, at the initial moment, there might be vacuum in some part of the space domain.

We strength hypothesis (1.7) with one of the following conditions,

$$u_0 \in V, \tag{1.9}$$

$$u_0 \in V \cap H^2(\Omega), \tag{1.10}$$

where $H^2(\Omega)$ is the usual $W^{2,2}(\Omega)$ Sobolev space and V is the function space defined below at (2.1). For the definitions and notations of the function spaces used throughout the paper, we address the reader to the monographs [15, 26]. In particular, given $m \in \mathbb{N}$ and $r \in [1, \infty]$, we denote by $L^r(\Omega)$ and $W^{m,r}(\Omega)$ the usual Lebesgue and Sobolev function spaces. As usual, when r = 2, we use the notation $H^m(\Omega) = W^{m,2}(\Omega)$. By $W_0^{m,r}(\Omega)$, we denote the closure of $C_0^{\infty}(\Omega)$ in the norm of $W^{m,r}(\Omega)$. The dual space of $W_0^{m,r}(\Omega)$ is denoted by $W^{-m,r'}(\Omega)$, where r' denotes the Hölder conjugate of r.

Condition (1.9) is enough to obtain weak solutions, but to obtain solutions with further regularity condition (1.10) is mandatory (see the authors works [2, 3, 4]). On the forcing term f, we shall

also assume two distinct situations,

$$f \in L^2(0, T; L^2(\Omega)),$$
 (1.11)

$$f \in L^{\infty}(0,T;L^2(\Omega)). \tag{1.12}$$

Of course that $L^{\infty}(0,T;L^2(\Omega)) \hookrightarrow L^2(0,T;L^2(\Omega))$, unless we consider the entire time interval $(0,\infty)$. But, as we shall see, condition (1.11) is sufficient for the results we aim to prove, being condition (1.12) only required to get some further L^{∞} -in time regularity.

The version of the problem (1.1)-(1.4), with constant density, was intensively studied by Oskolkov in a series of works (see [27] and references cited there in) who coined the name Kelvin-Voigt for the associated system of equations. However, as observed by Zvyagin and Turbin [34], neither Kelvin nor Voigt have suggested any stress-strain relation, or system of governing equations, for viscoelastic fluids. Currently, the Navier-Stokes-Voigt name for the associated system of equations seems to be the most accepted by the people working in this field, especially because this model is, in fact, an extension of the system of equations proposed by Voigt (for elastic materials that exhibit relaxation time) to model materials with viscoelastic properties. Mathematically speaking, the interesting feature of this system of equations, as noted first by Ladyzhenskaya [18], is that the relaxation term $\kappa \Delta u_t$ works as a regularization of the Navier-Stokes equations so that the corresponding problem has a unique global solution. Since then, the same problem, or some of its variants, have been studied by many authors, in many settings and under different conditions, with respect to the existence, uniqueness and asymptotic behaviour of the solutions. See for instance [3, 5, 34] and the references cited therein. On the other hand, the works by Titi and his collaborators [20, 30] make a clear relation between the homogeneous Navier-Stokes-Voigt equations and the turbulence modeling, in particular with Bardina turbulence models. The same relation was touched on by Lewandowski et al. [22, 21]. Existence of weak and strong solutions of nonlinear problems governed by the Navier-Stokes-Voigt equations (1.1)-(1.2) and for some of its generalisations, with p-Laplacian diffusion and damping terms, and for nonhomogeneous flows (non-constant density), were studied by the authors in [3, 5] in the case of a strictly positive initial density. For results on nonhomogeneous flows governed by the incompressible Navier-Stokes equations, which corresponds to take $\kappa = 0$ in the momentum equation (1.2), we address the reader to the works by Antontsev et al. [1] and by Ladyzhenskava and Solonnikov [19] in the case of a strictly positive initial density, and for Simon [31], Lions [24] and Desjardins [13] in the case of a initial density that vanishes in some part of the space domain. In all these works, the authors were primarily interested in the global-in-time existence of weak solutions in bounded domains of \mathbb{R}^d , d = 2, 3, or in the whole space \mathbb{R}^d , $d \ge 2$, and in the uniqueness of solution in the case of d = 2. Moreover, the initial data were considered so that $u_0 \in H^1(\Omega)$ and $\rho_0 \in L^{\infty}(\Omega)$. The case of a strictly positive initial density has been worked out also by many other authors during the last 20 years with respect to existence of weak and strong solutions, uniqueness, asymptotic stability and blow-up – see, for instance, [16, 33] and the references cited therein. We just want to point out the work by Paicu et al. in [28], where the authors have proved global-in-time existence and uniqueness in the whole space \mathbb{R}^d , d = 2, 3, for $u_0 \in H^s(\mathbb{R}^2)$, for s > 0, or $u_0 \in H^1(\mathbb{R}^3)$. When the density vanishes in some space subdomain, the momentum equation (1.2) degenerates into an elliptic equation, which makes it difficult to achieve the existence of strong solutions. To overcome this difficulty, Choe and Kim [8, 9] estimated $\|\nabla u_t\|_{L^2(\Omega)}$ by requiring that the initial data should satisfy a compatibility condition, expressed by the following Stokes problem,

$$\operatorname{div} u_0 = 0 \quad \text{in} \ \Omega, \tag{1.13}$$

$$-\mu\Delta u_0 = \sqrt{\rho_0}g - \nabla p_0 \quad \text{in} \quad \Omega, \tag{1.14}$$

$$u_0 = 0 \quad \text{on} \quad \partial\Omega, \tag{1.15}$$

for some $p_0 \in H^1(\Omega)$ and $g \in L^2(\Omega)$. This condition, combined with the assumption that $u_0 \in V \cap H^2(\Omega)$ and $\rho_0 \in L^{\infty}(\Omega)$, were of the utmost importance to prove the local-in-time existence of strong solutions in bounded domains of \mathbb{R}^3 . More recently, without requiring the compatibility condition (1.13)-(1.15), Lu *et al.* [25] have proved that, if the initial density decays not too slowly as $|x| \longrightarrow \infty$, then the 2d problem in the whole plane \mathbb{R}^2 admits a unique global-in-time strong solution. The decay condition on the initial density was written in the following form

$$\rho_0 \overline{x}^a \in L^1(\Omega) \cap H^1(\Omega) \cap W^{1,q}(\Omega), \qquad \overline{x} := \sqrt{e + |x|^2 \log^2\left(e + |x|^2\right)}, \tag{1.16}$$

for some a > 1 and q > 2 (see [25]). Around the same time, Li [23] has proved the existence of local-in-time strong solutions in bounded domains of \mathbb{R}^3 , assuming only that $u_0 \in V$ and $\rho_0 \in$ $L^{\infty}(\Omega) \cap W^{1,\gamma}(\Omega)$ for any $\gamma > 1$. Uniqueness was also proved in [23] but for all $\gamma \geq 2$. More recently, Danchin and Mucha [12] improved the results of [23], assuming only that $u_0 \in V$ and $\rho_0 \in L^{\infty}(\Omega)$ (without requiring any condition, be it regularity, strict positivity or compatibility condition that the initial density satisfies). The problem was considered in a spatial domain that can either be a bounded domain of \mathbb{R}^d , d = 2, 3, with a C^2 boundary $\partial \Omega$, or the torus \mathbb{T}^d , d = 2, 3. In these conditions, the authors [12] have proved global-in-time existence of strong solutions and their uniqueness in 2d, and also in 3d, in the last case only if $\|\nabla u_0\|_{L^2(\Omega)}$ is suitably small. About 2 years ago, He et al. [16] have proved global-in-time existence of strong solutions and its exponential stability in unbounded domains of \mathbb{R}^3 . These authors have considered, in addition, the difficult situation of a density-dependent viscosity. Last year, Zhang et al. [33] have extended the results of [12] requiring that the initial velocity can be in a larger function space: $u_0 \in H_0^s(\Omega)$ for s > 0. Nonhomogeneous flows, with initial vacuum, governed by the incompressible Navier-Stokes-Voigt equations were firstly studied by the authors in [2], where it was proved the existence of weak solutions in the whole space \mathbb{R}^d , d=2, 3, 4. There it were also proved some properties regarding the large-time behavior of the solutions in special unbounded domains. In the present, work we are interested in studying the existence and uniqueness of strong solutions for the Navier-Stokes-Voigt problem (1.1)-(1.5) with non-constant density. For the sake of mathematical generality, we shall assume throughout the rest of the work that the space dimension is $d \geq 2$, knowing in advance that there will be restrictions on the upper bound of d depending on the results we shall obtain. As we shall see in the sequel, the gain in regularity promoted by the presence of the relaxation term $\kappa \Delta u_t$ in the momentum equation shall allow us to prove the global-in-time existence, as well the uniqueness, of a strong solution without invoking any extra condition on the initial density.

We are interested in strong solutions to the problem (1.1)-(1.5) in the sense of the following definition.

Definition 1. Let $d \ge 2$ and assume the conditions (1.6), (1.8), (1.10) and (1.11) are fulfilled. If all the derivatives of ρ , u and p involved in (1.1)-(1.3) are regular distributions and the equations (1.1)-(1.3) hold almost everywhere in Q_T , and if still ρ and u satisfy the initial and boundary conditions (1.4)-(1.5), then the triple (ρ, u, p) is said to be a strong solution of the problem (1.1)-(1.5).

The main results of this work address the issue of existence of strong solutions for the problem (1.1)-(1.5). The first one is given by the following theorem and requires minimal assumptions on the regularity of the boundary domain.

Theorem 1. Let $2 \le d \le 4$ and assume that Ω is a bounded domain with $\partial\Omega$ Lipschitz-continuous. If the conditions (1.6), (1.8), (1.10) and (1.11) are fulfilled, then there exists, at least, a solution (ρ, u, p) for the problem (1.1)-(1.5) and such that:

(1) $0 \le \rho \le M$ in Q_T , $\rho \in C([0,T]; L^q(\Omega))$ for all $q \ge 1$ and $\rho_t \in L^2(0,T; W^{-1,2}(\Omega));$

- (2) $u \in L^{\infty}(0,T;V)$ and $\sqrt{\rho}u \in L^{\infty}(0,T;L^{2}(\Omega));$ (3) $u_{t} \in L^{2}(0,T;V)$ and $\sqrt{\rho}u_{t} \in L^{2}(0,T;L^{2}(\Omega));$ (4) $p \in C_{w}([0,T);L^{2}(\Omega));$
- If, instead of (1.11), is fulfilled (1.12), then: (5) $u_t \in L^{\infty}(0,T;V)$ and $\sqrt{\rho}u_t \in L^{\infty}(0,T;L^2(\Omega)).$

It is noteworthy that, in the Navier-Stokes setting (without the relaxation term $\kappa \Delta u_t$ in the momentum equation (1.2)) for nonhomogeneous flows, to prove that $u_t \in L^2(0,T;V)$ or $u_t \in L^{\infty}(0,T;V)$, extra assumptions are needed. In fact, to prove these results in [8], it was assumed that

$$f_t \in L^2(0, T; L^2(\Omega)),$$
 (1.17)

$$f \in L^2(0,T; H^1(\Omega)).$$
 (1.18)

Moreover, and in addition to (1.17)-(1.18), it was required the initial data u_0 and ρ_0 should satisfy the compatibility condition (1.13)-(1.15). With respect to assumption (1.17), it should be noted that, by the Sobolev imbedding $W^{1,2}(0,T) \hookrightarrow L^{\infty}(0,T)$ (see [7, Theoreme VIII.7]), the assumption $f_t \in L^2(0,T; L^2(\Omega))$ (whence $f \in W^{1,2}(0,T; L^2(\Omega))$) would imply (1.11).

This way, we can realize that for the nonhomogeneous Navier-Stokes-Voigt problem we can prove the same regularity results without assuming the extra conditions (1.17)-(1.18) on the forcing term f, nor requiring the compatibility problem (1.13)-(1.15), or condition (1.16). More importantly, our proof is technically much more accessible and much less time consuming. This is only possible due to the presence of the relaxation term $\kappa \Delta u_t$ in the momentum equation (1.2). Besides (6.6)-(6.7), see the proofs of (4.3) and (4.6) below.

To obtain more regularity in the solutions, not only a smoother boundary domain is needed, but also more restrictions on the space dimension. This is the aim of the next theorem.

Theorem 2. Let $2 \le d \le 3$ and assume that Ω is a bounded domain with $\partial\Omega$ supposed to be of class C^2 . If the conditions (1.6), (1.8), (1.10) and (1.11) are fulfilled, then there exists, at least, a solution (ρ, u, p) for the problem (1.1)-(1.5) and such that, in addition to (1)-(4) of Theorem 1, we have:

(1) $D^2 u \in L^{\infty}(0,T; L^2(\Omega))$ and $D^2 u_t \in L^2(0,T; L^2(\Omega));$ (2) $\nabla p \in L^2(0,T; L^2(\Omega));$

If, instead of (1.11), is fulfilled (1.12), then, in addition to (5) of Theorem 1, we have:

(3) $D^2 u_t \in L^{\infty}(0, T; L^2(\Omega));$ (4) $\nabla p \in L^{\infty}(0, T; L^2(\Omega)).$

The greatest gain in regularity, relatively to the nonhomogeneous Navier-Stokes equations (see again [8, 25]), is observed in the regularity results $D^2 u_t \in L^2(0, T; L^2(\Omega))$ and $D^2 u_t \in L^{\infty}(0, T; L^2(\Omega))$ (see (4.5) and (4.7) below), which cannot at all be achieved if we remove the relaxation term $\kappa \Delta u_t$ from the momentum equation (1.2).

The proofs of Theorems 1 and 2 will appear as a consequence of what is done later on in Sections 3-6.

Remark 1. In addition to the regularity results $D^2 u \in L^{\infty}(0,T;L^2(\Omega))$ and $D^2 u_t \in L^{\infty}(0,T;L^2(\Omega))$, we also have

$$u, \ \frac{\partial u}{\partial t} \in L^{\infty}(0,T; C^{0,\alpha}(\overline{\Omega})) \quad whenever \quad 0 < \alpha \le 2 - \frac{d}{2} \quad and \quad 2 \le d \le 3$$

This can be shown by combining the Sobolev inequalities (2.7) and (2.9) with the regularity results aforementioned. See also Remark 4 below.

As a complementary result to the existence of strong solutions (ρ, u, p) , we provide, in the following theorem, the conditions that allow us to prove the uniqueness of ρ and u.

Theorem 3. Let $(\hat{u}, \hat{p}, \hat{\rho})$ and $(\overline{u}, \overline{p}, \overline{\rho})$ be two solutions of the problem (1.1)-(1.5) with the same data and in the conditions of Theorem 2. If, in addition,

$$\rho_0 \in W^{1,\infty}(\Omega) \tag{1.19}$$

and

$$u_0 \in W^{2,r}(\Omega) \cap V, \tag{1.20}$$

$$f \in L^2(0, T; L^r(\Omega)),$$
 (1.21)

for

$$d < r \le 2^*. \tag{1.22}$$

then $\hat{\rho} = \overline{\rho}$ and $\hat{u} = \overline{u}$.

Here, 2^* denotes the Sobolev conjugate of 2, and note that (1.22) implies $2 \le d \le 3$.

The rest of the paper is organised as follows. In Section 2, we provide important auxiliary results that will be used throughout the following sections. Problem (1.1)-(1.5) is approximated by two cascade of problems, being Sections 3-5 devoted to proving the existence of the Galerkin approximations for the second cascade of approximate problems. Sections 4-5 also prove some results that make the solutions of the two cascade of approximate problems strong. In Section 6, we prove the existence of strong solutions for the original problem (1.1)-(1.5). Further regularity results are proved in Section 7, where we also establish the uniqueness of the velocity and density under additional assumptions on the problem data.

2. AUXILIARY RESULTS

In this section, we introduce important auxiliary results that we will be used in the course of our work. We recall the definition of the following function spaces,

$$\mathcal{V} := \{ u \in C_0^{\infty}(\Omega) : \operatorname{div} u = 0 \},\$$

$$H := \operatorname{closure} \text{ of } \mathcal{V} \text{ in the norm of } L^2(\Omega),\$$

$$V_q := \operatorname{closure} \text{ of } \mathcal{V} \text{ in the norm of } W^{1,q}(\Omega).$$
(2.1)

The particular case of q = 2 in (2.1) will be denoted only by V. In the sequel, we shall denote the inclusion $X \subset Y$ of two Banach spaces X and Y with a continuous imbedding $X \to Y$ by $X \hookrightarrow Y$. If the imbedding $X \to Y$ is compact, we denote the inclusion $X \subset Y$ by $X \hookrightarrow Y$.

We start by recalling the Sobolev, Moser and Morrey inequalities and the continuous and compact imbeddings that come with them.

Lemma 1. Let Ω be a bounded domain in \mathbb{R}^d with a Lipschitz-continuous boundary $\partial \Omega$. If $v \in W_0^{1,r}(\Omega)$, then

$$\|v\|_{L^{r^*}(\Omega)} \le C(r,d) \|\nabla v\|_{L^r(\Omega)}, \quad r^* = \frac{dr}{d-r}, \quad 1 \le r < d,$$
(2.2)

$$\|v\|_{L^{q}(\Omega)} \le C(r, q, d) \|\nabla v\|_{W^{1, r}(\Omega)}, \quad d \le q < \infty, \quad r = d,$$
(2.3)

$$[v]_{C^{0,\alpha}(\overline{\Omega})} \le C(r,d) \left\| \nabla v \right\|_{L^{r}(\Omega)}, \quad 0 < \alpha \le 1 - \frac{d}{r}, \quad r > d.$$

Moreover, $W^{1,r}(\Omega) \hookrightarrow L^q(\Omega)$ if $1 \le q \le r^*$ and $1 \le r < d$, $W^{1,r}(\Omega) \hookrightarrow L^q(\Omega)$ if $d \le q < \infty$ and r = d, and $W^{1,r}(\Omega) \hookrightarrow C^{0,\alpha}(\overline{\Omega})$ if $\alpha = 1 - \frac{r}{d}$ and r > d. In addition, $W^{1,r}(\Omega) \hookrightarrow L^q(\Omega)$ if $1 \le q < r^*$ and $1 \le r < d$, $W^{1,r}(\Omega) \hookrightarrow L^q(\Omega)$ if $1 \le q < \infty$ and r = d, and $W^{1,r}(\Omega) \hookrightarrow C^{0,\alpha}(\overline{\Omega})$ if $0 < \alpha \le 1 - \frac{d}{r}$ and r > d.

Proof. For the proof, we address the reader to Maz'ya [26, Chapter 2].

For the sake of simplifying the writing, in the sequel, we shall use the notation r^* with the broadest meaning that $r^* = \frac{rp}{d-r}$ if r < d, r^* is any real in the interval $[1, \infty)$ if r = d, or $r^* = \infty$ if r > d.

In the following lemma, we collect two important generalizations of the Aubin-Lions compactness lemma.

Lemma 2. If X, E and Y are Banach spaces such that $X \hookrightarrow E \hookrightarrow Y$, then

$$L^{r}(0,T;X) \cap \left\{ v : v_{t} \in L^{1}(0,T;Y) \right\} \hookrightarrow \hookrightarrow L^{r}(0,T;E) \quad if \quad 1 \le r \le \infty,$$

$$(2.4)$$

$$L^{\infty}(0,T;X) \cap \{v : v_t \in L^q(0,T;Y)\} \hookrightarrow \hookrightarrow C([0,T];E) \quad if \quad 1 < q \le \infty.$$

$$(2.5)$$

Proof. See Simon [32, Corollary 4].

The following variant of de Rham's lemma is of the utmost importance to recover the pressure, after the velocity field and the density are found. In what follows, by $\langle \cdot, \cdot \rangle$ we denote the duality paring between $W^{-1,q'}(\Omega)$ and $W_0^{1,q}(\Omega)$.

Lemma 3. Let $1 < q < \infty$ and $\varphi^* \in W^{-1,q'}(\Omega)$. If

$$\langle \varphi^*, \varphi \rangle = 0 \quad \forall \ \varphi \in V_q,$$

then there exists a unique $p \in L^q(\Omega)$, with $\int_{\Omega} p \, dx = 0$, such that

$$\langle \varphi^*, \varphi \rangle = \int_{\Omega} p \operatorname{div} \varphi \, dx \quad \forall \, \varphi \in W_0^{1,q}(\Omega).$$

Moreover, there exists a positive constant C such that

$$\|p\|_{L^q(\Omega)} \le C \|\varphi^*\|_{W^{-1,q'}(\Omega)}$$

Proof. The proof combines the results of Bogovskii [6] and Pileckas [29] (see also Theorems III.3.1 and III.5.3 of Galdi [15]). \Box

Next, some other auxiliary results that shall be used in the sequel are collected. We start by recalling some useful inequalities related with the Sobolev inequalities stated in Lemma 1. Here, the notation $|D^m u|$, where $m \in \mathbb{N}$, stands for

$$|D^m u| := \sum_{|\gamma|=m} \left| \frac{\partial^{|\gamma|} u}{\partial_{x_1}^{\gamma_1} \cdots \partial_{x_d}^{\gamma_d}} \right|, \quad |\gamma| = \gamma_1 + \cdots + \gamma_d.$$

In particular, $|Du|^2 = |\nabla u|^2$ and $|D^2 u|^2 = |\nabla u_{x_1}|^2 + \dots + |\nabla u_{x_d}|^2$. We say that the (bounded) boundary $\partial \Omega$ belongs to the class $C^{m,\alpha}$, with $0 \le \alpha \le 1$, if each point $x = (x_1, \dots, x_{d-1}, x_d) \in \partial \Omega$

has a neighborhood U such that the set $U \cap \Omega$ is represented by the inequality $x_d < f(x_1, \ldots, x_{d-1})$ in some Cartesian coordinate system and for some function f in the Hölder space $C^{m,\alpha}(\omega)$, where here ω is the projection of Ω onto \mathbb{R}^{d-1} .

Lemma 4. Let Ω be a bounded domain in \mathbb{R}^d and assume that $r \geq 1$. If the boundary $\partial \Omega$ is assumed to be of class $C^{0,1}$, then

$$\|\nabla u\|_{L^{r^*}(\Omega)} \le C(r,d) \left\| D^2 u \right\|_{L^r(\Omega)} \qquad \forall \ u \in W^{2,r}(\Omega) \cap W^{1,r}_0(\Omega),$$

$$(2.6)$$

$$\frac{1}{C(r,d)} \|\Delta u\|_{L^{r}(\Omega)} \leq \left\|D^{2}u\right\|_{L^{r}(\Omega)} \leq C(r,d) \|\Delta u\|_{L^{r}(\Omega)} \qquad \forall \ u \in W^{2,r}(\Omega) \cap W^{1,r}_{0}(\Omega).$$
(2.7)

If $\partial \Omega$ is assumed to be of class $C^{1,1}$, then

$$\|\nabla u\|_{L^{r^*}(\Omega)} \le C(d,r) \|\Delta u\|_{L^r(\Omega)} \qquad \forall \ u \in W^{2,r}(\Omega) \cap W^{1,r}_0(\Omega),$$

$$(2.8)$$

$$\|u\|_{C^{0,\alpha}(\overline{\Omega})} \le C(r,d) \|\Delta u\|_{L^{r}(\Omega)} \qquad \forall \ u \in W^{2,r}(\Omega) \cap W^{1,r}_{0}(\Omega), \qquad 0 < \alpha \le 1 - \frac{d}{r^{*}}, \quad 2r > d.$$
(2.9)

Proof. For the proof we address the reader to [3, Lemma 1] (see also Maz'ya [26, Chapters 1-2]).

As we will do throughout this work, the notation C = C(d, r), used in the previous lemma, emphasizes the fact that the positive constants C considered in (2.8)-(2.9), where they are supposed to be all distinct, depend on the parameters d and r.

In the next lemma, we recall some of the properties of the Stokes operator. The operator

where $\mathbb{P}: L^2(\Omega) \to H$ is the Leray projection, is called the Stokes operator. This operator establishes a correspondence between the solutions u of the stationary Stokes problems

$$\operatorname{div} u = 0 \qquad \text{in} \quad \Omega, \tag{2.11}$$

$$-\mu\Delta u = f - \nabla p \qquad \text{in} \quad \Omega, \tag{2.12}$$

$$u = 0$$
 on $\partial\Omega$, (2.13)

and the corresponding external forces f. Due to the symmetry of the Leray projection, it can be proved that

$$\mu \int_{\Omega} \nabla u : \nabla \varphi \, dx = \int_{\Omega} \mathbb{A}(u) \cdot \varphi \, dx \qquad \forall \ u \in W^{2,2}(\Omega) \cap V, \quad \forall \ \varphi \in V.$$
(2.14)

The following result follows from the existence and regularity theory for elliptic operators.

Lemma 5. Let Ω be a bounded domain in \mathbb{R}^d , with the boundary $\partial\Omega$ assumed to be of class C^2 . If $f \in L^r(\Omega)$, with $1 < r < \infty$, then there exist unique $u \in W^{2,r}(\Omega)$ and $p \in W^{1,r}(\Omega)$, with $\int_{\Omega} p(x) dx = 0$, such that (u, p) verify the Stokes system (2.11)-(2.12) a.e. in Ω and u satisfies (2.13) in the trace sense. Moreover, there exists a positive constant $C = C(\mu, r, \Omega)$ such that

$$\|u\|_{W^{2,r}(\Omega)} + \|p\|_{W^{1,r}(\Omega)} \le C \|f\|_{L^{r}(\Omega)}.$$
(2.15)

Proof. We address the proof to Galdi [15, Theorem IV.6.1].

From (2.15), we easily derive the following estimate,

$$\left\| D^2 u \right\|_{L^r(\Omega)} + \left\| \nabla p \right\|_{L^r(\Omega)} \le C \left\| f \right\|_{L^r(\Omega)}.$$
(2.16)

Using the correspondence between the Stokes operator (2.10) and the forces field f, and taking r = 2 in (2.16), we also obtain

$$\left\| D^2 u \right\|_{L^2(\Omega)} + \left\| \nabla p \right\|_{L^2(\Omega)} \le C \left\| \mathbb{A}(u) \right\|_{L^2(\Omega)}.$$
(2.17)

3. EXISTENCE OF APPROXIMATE SOLUTIONS

The proof of the existence of solutions for the problem (1.1)-(1.5) shall follow from the existence of suitable Galerkin approximations. Let us consider the following family of subsets of Ω ,

$$\Omega_n := \left\{ x \in \Omega : \operatorname{dist}(x, \partial \Omega) > \frac{1}{n}, \quad n \in \mathbb{N} \right\},$$

and the usual Friedrichs mollifying kernel $\eta_n(x) := \frac{1}{n^d} \eta\left(\frac{x}{n}\right)$. Recall that

$$\eta_n \in C^{\infty}(\mathbb{R}^d), \quad \operatorname{supp} \eta_n \subset B(0,n), \quad \int_{\mathbb{R}^d} \eta_n(x) \, dx = 1$$

We regularize the initial data u_0 and ρ_0 by considering its mollifying functions, say $u_{0,n}$ and $\rho_{0,n}$, defined by

$$u_{0,n}(x) := (\eta_n \star u_0)(x) = \int_{\Omega} \eta_n(x - y)u_0(y) \, dy, \quad x \in \Omega_n,$$
(3.1)

$$\rho_{0,n}(x) := (\eta_n \star \rho_0)(x) + \frac{1}{n} = \int_{\Omega} \eta_n(x-y)\rho_0(y)\,dy + \frac{1}{n}, \quad x \in \Omega_n.$$
(3.2)

It is well-known that for any $w \in L^1_{loc}(\Omega)$, its mollifying function $w_n = \eta_n \star w$ satisfies

$$w_n \in C^{\infty}(\Omega_n), \qquad w_n \xrightarrow[n \to \infty]{} w \text{ a.e. in } \Omega,$$
(3.3)

$$w_n \xrightarrow[n \to \infty]{} w \text{ in } L^p(\Omega), \text{ whenever } w \in L^p(\Omega), \quad 1 \le p < \infty.$$
 (3.4)

As supp $\eta_n \subset B(0,n)$, one has

 $\operatorname{supp} u_{0,n} \subset \Omega_n + B(0,n) \,.$

Moreover, in view of (1.8), (3.1)-(3.2) and (3.3)-(3.4), we have

$$0 < \frac{1}{n} \le \rho_{0,n} \le M^* := M + 1 < \infty \text{ in } \Omega,$$
 (3.5)

$$\|u_{0,n}\|_{L^2(\Omega)} \le \|u_0\|_{L^2(\Omega)} \,. \tag{3.6}$$

Moreover, provided u_0 is sufficiently regular, we also have

$$\|\nabla u_{0,n}\|_{L^{2}(\Omega)} \leq \|\nabla u_{0}\|_{L^{2}(\Omega)}, \qquad \left\|D^{2}u_{0,n}\right\|_{L^{2}(\Omega)} \leq \left\|D^{2}u_{0}\right\|_{L^{2}(\Omega)}.$$
(3.7)

Given a large enough, but arbitrary, $n \in \mathbb{N}$, we consider the following initial-and boundary-value problem,

$$\operatorname{div} u = 0 \qquad \text{in} \quad Q_T, \tag{3.8}$$

$$(\rho u)_t + \operatorname{div}(\rho u \otimes u) = \rho f - \nabla p + \mu \Delta u + \kappa \Delta u_t \quad \text{in} \quad Q_T,$$
(3.9)

$$\rho_t + \operatorname{div}(\rho u) = 0 \qquad \text{in} \quad Q_T, \tag{3.10}$$

$$\rho u = \rho_{0,n} u_{0,n}, \quad \rho = \rho_{0,n} \quad \text{in} \quad \{0\} \times \Omega,$$
(3.11)

$$u = 0 \qquad \text{on} \quad \Gamma. \tag{3.12}$$

We construct a solution to the problem (3.8)-(3.12) by using a semi-discrete Galerkin scheme. Since the Stokes operator is injective, self-adjoint and has a compact inverse (see e.g. [11, Propositions 4.2-4]), there exists an increasing sequence of positive eigenvalues λ_i and a sequence of corresponding eigenfunctions $\psi_i \in H^2(\Omega) \cap V$ such that

$$\mathbb{A}(\psi_i) = \lambda_i \psi_i. \tag{3.13}$$

Moreover, the family $\{\psi_i\}_{i\in\mathbb{N}}$ can be made orthogonal in H and orthonormal in V. Given $j\in\mathbb{N}$, let us consider the *j*-dimensional space X^j spanned by the first *j* eigenvalues given by (3.13): ψ_1 , ..., ψ_j . For each $j \in \mathbb{N}$, and proceeding as in the proof of [3, Theorem 1] (see also the proof of [2, Proposition 1), we can prove the existence of approximate solutions

$$u_n^j \in C^1([0,T); X^j), \quad u_n^j(x,t) = \sum_{i=1}^j c_i^j(t)\psi_i(x), \quad \psi_i \in X^j,$$
(3.14)

$$\rho_n^j \in C^1([0,T); C^1(\overline{\Omega})) \tag{3.15}$$

to the following system of j + 1 ordinary differential equations

$$\int_{\Omega} \rho_n^j(t) \left[\frac{\partial u_n^j(t)}{\partial t} + \left(u_n^j(t) \cdot \nabla \right) u_n^j(t) \right] \cdot \psi_i \, dx + \kappa \int_{\Omega} \frac{\partial \nabla u_n^j(t)}{\partial t} : \nabla \psi_i \, dx + \mu \int_{\Omega} \nabla u_n^j(t) : \nabla \psi_i \, dx = \int_{\Omega} \rho_n^j(t) f(t) \cdot \psi_i \, dx, \qquad i = 1, \dots, j,$$

$$\frac{\partial \rho_n^j}{\partial t} + u_n^j \cdot \nabla v_n^j = 0$$
(3.16)

$$\frac{\partial \rho_n^j}{\partial t} + u_n^j \cdot \nabla \rho_n^j = 0, \qquad (3.17)$$

System (3.16)-(3.17) is supplemented with the following initial conditions

$$\rho_n^j u_n^j = \rho_{0,n}^j u_{0,n}^j, \quad u_n^j = u_{0,n}^j, \quad \rho_n^j = \rho_{0,n}^j \quad \text{in } \{0\} \times \Omega, \tag{3.18}$$

where $u_{0,n}^j = P^j(u_{0,n})$, with P^j denoting the orthogonal projection $P^j: V \longrightarrow X^j$ so that

$$u_n^j(0,x) = \sum_{i=1}^j c_i^j(0)\psi_i(x), \quad c_i^j(0) = c_{i,0}^j := (u_{0,n},\psi_i), \quad i \in \{1,\dots,j\},$$
(3.19)

where (\cdot, \cdot) denotes the L^2 -scalar product. Since the operator P^j is uniformly continuous, we can assume that

$$u_{0,n}^j \xrightarrow[j \to \infty]{} u_{0,n} \quad \text{in } L^2(\Omega) \cap W^{1,2}(\Omega).$$
 (3.20)

About the approximate initial density $\rho_{0,n}^{j}$, we assume that

$$\rho_{0,n}^{j} \in C^{1}(\overline{\Omega}), \qquad \rho_{0,n}^{j} \xrightarrow{j \to \infty} \rho_{0,n} \text{ in } L^{p}(\Omega) \quad \forall \ p \in [1,\infty).$$
(3.21)

From (3.5)-(3.7) and (3.20)-(3.21), one readily has

$$\frac{1}{n} \le \rho_{0,n}^{j} \le M^{*} < \infty \quad \text{in} \quad \Omega,$$

$$\left\| u_{0,n}^{j} \right\|_{L^{2}(\Omega)} \le \| u_{0} \|_{L^{2}(\Omega)},$$
(3.22)
(3.23)

$$u_{0,n}^{j}\Big\|_{L^{2}(\Omega)} \le \|u_{0}\|_{L^{2}(\Omega)}, \tag{3.23}$$

$$\left\|\nabla u_{0,n}^{j}\right\|_{L^{2}(\Omega)} \leq \|\nabla u_{0}\|_{L^{2}(\Omega)}, \qquad \left\|D^{2} u_{0,n}^{j}\right\|_{L^{2}(\Omega)} \leq \left\|D^{2} u_{0}\right\|_{L^{2}(\Omega)}.$$
(3.24)

Observe that, due to the regularity of u_n^j and by linearity and continuity, one can derive from (3.16)

$$\int_{\Omega} \left[\rho_n^j(t) \frac{\partial u_n^j(t)}{\partial t} + \rho_n^j(t) \left(u_n^j(t) \cdot \nabla \right) u_n^j(t) - \mu \Delta u_n^j(t) - \kappa \frac{\partial \Delta u_n^j(t)}{\partial t} \right] \cdot \psi \, dx =$$

$$\int_{\Omega} \rho_n^j(t) f(t) \cdot \psi \, dx \qquad \forall \, \psi \in H^2(\Omega) \cap V$$
(3.25)

in the distribution sense on (0, T). Moreover, by using Lemma 3, it can also be proved the existence of a unique approximate pressure

$$p_n^j \in C_w([0,T); L^2(\Omega)), \quad \text{with } \int_{\Omega} p_n^j(t) \, dx = 0,$$
 (3.26)

so that

$$\int_{\Omega} \left[\rho_n^j(t) \frac{\partial u_n^j(t)}{\partial t} + \rho_n^j(t) \left(u_n^j(t) \cdot \nabla \right) u_n^j(t) - \mu \Delta u_n^j(t) - \kappa \frac{\partial \Delta u_n^j(t)}{\partial t} \right] \cdot \psi \, dx - \int_{\Omega} \rho_n^j(t) f(t) \cdot \psi \, dx = \int_{\Omega} \rho_n^j(t) \operatorname{div} \psi \, dx \qquad \forall \ \psi \in H^2(\Omega) \cap W_0^{1,2}(\Omega)$$

$$(3.27)$$

holds in the distribution sense on (0, T) – see [3, Theorem 2].

4. A priori estimates independent of j

In this section, we aim to obtain independent of j estimates for the approximate solutions u_n^j , ρ_n^j and p_n^j . We list the obtained estimates in several propositions according to the conditions that are imposed on the forcing term f. We highlight the dependence of these estimates on the viscous and relaxation parameters μ and κ to perceive the importance of the presence of the viscous term $\mu \Delta u$ and of the relaxation one $\kappa \Delta u_t$, in the momentum equation (1.2), for the results we achieve. It is important to note that these estimates are also independent of n.

Proposition 1. Let u_n^j , ρ_n^j and p_n^j be the approximate weak solutions of the problem (3.8)-(3.12) that have been found in (3.14), (3.15) and (3.26).

(1) If (1.8) holds true, then

$$0 < \frac{1}{n} \le \inf_{x \in \overline{\Omega}} \rho_{0,n}^j(x) \le \rho_n^j(x,t) \le \sup_{x \in \overline{\Omega}} \rho_{0,n}^j(x) \le M^* < \infty \qquad \forall \ (x,t) \in Q_T.$$
(4.1)

(2) If $2 \le d \le 4$ and (1.8) and (1.9) are verified, then there exists an independent of j (and n) positive constant K_1 such that

$$\sup_{t \in (0,T)} \left(\left\| \sqrt{\rho_n^j(t)} u_n^j(t) \right\|_{L^2(\Omega)}^2 + \kappa \left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)}^2 \right) + \mu \int_0^T \left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)}^2 dt \le K_1$$
(4.2)

(3) If $2 \le d \le 4$ and (1.6), (1.8), (1.9) and (1.11) hold, then there exists an independent of j (and n) positive constant K_2 such that

$$\int_{0}^{T} \left(\left\| \sqrt{\rho_{n}^{j}(t)} \frac{\partial u_{n}^{j}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} + \kappa \left\| \frac{\partial \nabla u_{n}^{j}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} \right) dt \leq K_{2}.$$

$$(4.3)$$

(4) If $2 \le d \le 3$ and (1.6), (1.8), (1.10) and (1.11) are verified, then there exists an independent of j (and n) positive constant K_3 such that

$$\kappa \sup_{t \in (0,T)} \left\| D^2 u_n^j(t) \right\|_{L^2(\Omega)}^2 + \mu \int_0^T \left\| D^2 u_n^j(t) \right\|_{L^2(\Omega)}^2 dt \le K_3.$$
(4.4)

(5) If $2 \le d \le 3$ and (1.6), (1.8), (1.10) and (1.11) hold, then there exists an independent of j (and n) positive constant K_4 such that

$$\kappa^2 \int_0^T \left\| \frac{\partial D^2 u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 dt \le K_4.$$
(4.5)

In the following proposition, we improve some of the estimates established in Proposition 1 by requiring that (1.12) is fulfilled, instead of (1.11).

Proposition 2. Let u_n^j , ρ_n^j and p_n^j be the approximate weak solutions of the problem (3.8)-(3.12) that have been found in (3.14), (3.15) and (3.26).

(1) If $2 \le d \le 4$ and (1.6), (1.8), (1.9) and (1.12) hold, then there exists an independent of j (and n) positive constant K'_2 such that

$$\sup_{t\in[0,T]} \left(\left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 + \kappa \left\| \frac{\partial \nabla u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 \right) \le K_2'.$$
(4.6)

(2) If $2 \le d \le 3$ and (1.6), (1.8), (1.10) and (1.12) hold, then there exists an independent of j (and n) positive constant K'_4 such that

$$\kappa^{2} \sup_{t \in [0,T]} \left\| \frac{\partial D^{2} u_{n}^{j}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} \leq K_{4}^{\prime}.$$

$$(4.7)$$

Before we proceed to the proof of Propositions 1 and 2, let us make some comments on the estimates obtained here, in particular the relation between these estimates and its counterparts of the Navier-Stokes setting, i.e. when considering $\kappa = 0$ in the momentum equation (1.2).

Remark 2. (1) Estimate (4.2) has already been established in [2, Lemma 4-(2)], but here we just have required that $f \in L^2(0,T; L^2(\Omega))$. Moreover, in view of (4.14) below (see also (4.32)), the relaxation parameter κ in (4.2) can be replaced by the viscous parameter μ .

(2) Estimate (4.4) is obtained regardless we consider the hypothesis of initial vacuum or not. In the context of considering an initial density ρ_0 that is always positive, this estimate was already proved in [3, Theorem 3], by using a different approach. In view of (4.32) below, we can also replace the relaxation parameter κ in (4.4) by the viscous parameter μ .

Proof. (Proposition 1) For the sake of simplifying the exposition, we shall split the proof into the several enumerated items.

(1) Arguing as we did in the proof of [3, Theorem 1] (see also [19, Lemma 1.2]), and using the maximum principle together with (3.22), we can show that (4.1) holds true. On the other hand, using the solenoidality of u_n^j , the fact that $u_n^j = 0$ on $\partial\Omega$, together with (3.17) and (3.18)₃, we can

prove that for all $t \ge 0$ and $1 \le q \le \infty$ one has

$$\begin{aligned} \left\| \rho_{n}^{j}(t) \right\|_{L^{q}(\Omega)}^{q} &= \int_{0}^{t} \frac{\partial}{\partial s} \left\| \rho_{n}^{j}(s) \right\|_{L^{q}(\Omega)}^{q} ds + \left\| \rho_{0,n}^{j} \right\|_{L^{q}(\Omega)}^{q} \\ &= -\int_{0}^{t} \int_{\Omega} \nabla \left(|\rho_{n}^{j}(s)|^{q} \right) \cdot u_{n}^{j}(s) \, dx ds + \left\| \rho_{0,n}^{j} \right\|_{L^{q}(\Omega)}^{q} = \left\| \rho_{0,n}^{j} \right\|_{L^{q}(\Omega)}^{q}. \end{aligned}$$

$$(4.8)$$

(2) Testing (3.25) with $\psi = u_n^j(t)$, integrating the resulting identity between 0 and $t \in (0,T)$, using the continuity equation (3.17) together with the solenoidality of u_n^j , and still using the initial conditions (3.18), we obtain

$$\frac{1}{2} \left\| \sqrt{\rho_n^j(t)} u_n^j(t) \right\|_{L^2(\Omega)}^2 + \frac{\kappa}{2} \left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)}^2 + \mu \int_0^t \left\| \nabla u_n^j(s) \right\|_{L^2(\Omega)}^2 ds = \frac{1}{2} \left\| \sqrt{\rho_{0,n}^j} u_{0,n}^j \right\|_{L^2(\Omega)}^2 + \frac{\kappa}{2} \left\| \nabla u_{0,n}^j \right\|_{L^2(\Omega)}^2 + \int_0^t \int_\Omega \rho_n^j f \cdot u_n^j dx ds$$

$$(4.9)$$

The last term of (4.9) is estimated by using the Hölder and Cauchy inequalities,

$$\int_{0}^{t} \int_{\Omega} \rho_{n}^{j}(s) f(s) \cdot u_{n}^{j}(s) dx ds \leq \frac{1}{4} \int_{0}^{t} \left\| \sqrt{\rho_{n}^{j}(s)} u_{n}^{j}(s) \right\|_{L^{2}(\Omega)}^{2} ds + \int_{0}^{t} \left\| \sqrt{\rho_{n}^{j}(s)} f(s) \right\|_{L^{2}(\Omega)}^{2} ds.$$
(4.10)

Plugging (4.10) into (4.9) and using (3.22)-(3.23) and $(3.24)_1$, together with (4.1), one has

$$\frac{1}{4} \left\| \sqrt{\rho_n^j(t)} u_n^j(t) \right\|_{L^2(\Omega)}^2 + \frac{\kappa}{2} \left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)}^2 + \mu \int_0^t \left\| \nabla u_n^j(s) \right\|_{L^2(\Omega)}^2 ds \le \frac{M^*}{2} \left\| u_0 \right\|_{2,\Omega}^2 + \frac{\kappa}{2} \left\| \nabla u_0 \right\|_{2,\Omega}^2 + \frac{1}{2} \int_0^t \left\| \sqrt{\rho_n^j(s)} u_n^j(s) \right\|_{L^2(\Omega)}^2 ds + M^* \int_0^t \left\| f(s) \right\|_{L^2(\Omega)}^2 ds$$

Using the Grönwall inequality and taking the supremum in (0,T) in the resulting inequality, we obtain (4.2) for some positive constant $K_1 = C\left(M^*, \kappa, \|u_0\|_{2,\Omega}, \|\nabla u_0\|_{2,\Omega}, \|f\|_{L^2(0,T;L^2(\Omega))}, T\right)$. (3) Testing (3.25) with $\psi = \frac{\partial u_n^j(t)}{\partial t}$, we obtain

$$\frac{\mu}{2}\frac{d}{dt}\left\|\nabla u_{n}^{j}(t)\right\|_{L^{2}(\Omega)}^{2}+\left\|\sqrt{\rho_{n}^{j}(t)}\frac{\partial u_{n}^{j}(t)}{\partial t}\right\|_{L^{2}(\Omega)}^{2}+\kappa\left\|\frac{\partial\nabla u_{n}^{j}(t)}{\partial t}\right\|_{L^{2}(\Omega)}^{2}=$$

$$J(t)+\int_{\Omega}\rho_{n}^{j}(t)f(t)\cdot\frac{\partial u_{n}^{j}(t)}{\partial t}\,dx,$$

$$(4.11)$$

where

$$J(t) := -\int_{\Omega} \rho_n^j(t) \left(u_n^j(t) \cdot \nabla \right) u_n^j(t) \cdot \frac{\partial u_n^j(t)}{\partial t} dx.$$

To estimate the last term of (4.11), we proceed as in (4.10) so that

$$\int_{\Omega} \rho_n^j(t) f(t) \cdot \frac{\partial u_n^j(t)}{\partial t} dx \le \frac{1}{2} \left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 + \frac{1}{2} \left\| \sqrt{\rho_n^j(t)} f(t) \right\|_{L^2(\Omega)}^2.$$
(4.12)

For the estimate of J(t), we use (4.1) together with the Hölder, Cauchy and Sobolev inequalities so that

$$\begin{aligned} |J(t)| &\leq M^* \left\| u_n^j(t) \right\|_{L^d(\Omega)} \left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)} \left\| \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^{2^*}(\Omega)}, \qquad 2 \leq d \leq 4 \\ &\leq C_1 \left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)}^2 \left\| \frac{\partial \nabla u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)} \\ &\leq \frac{\kappa}{2} \left\| \frac{\partial \nabla u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 + C_2 \left(\sup_{t \in (0,T)} \left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)}^2 \right)^2 \end{aligned}$$
(4.13)

for some positive constants $C_1 = C(d, M^*, \Omega)$ and $C_2 = C(M^*, \kappa, d, \Omega)$. Plugging (4.12) and (4.13) into (4.11), integrating the resulting inequality between 0 and $t \in (0, T)$ and using (3.18) and the estimate (4.2), we achieve to

$$\begin{split} & \mu \left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)}^2 + \int_0^t \left(\left\| \sqrt{\rho_n^j(s)} \frac{\partial u_n^j}{\partial s}(s) \right\|_{L^2(\Omega)}^2 + \kappa \left\| \frac{\partial \nabla u_n^j(s)}{\partial s} \right\|_{L^2(\Omega)}^2 \right) ds \leq \\ & \mu \left\| \nabla u_{0,n}^j \right\|_{L^2(\Omega)}^2 + \|f(t)\|_{L^2(\Omega)}^2 + C, \end{split}$$

for some positive constant $C = C(M^*, \kappa, d, K_1, \Omega, T)$. Taking the supremum in (0, T) and using $(3.24)_1$, there holds

$$\mu \sup_{t \in (0,T)} \left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)}^2 + \int_0^T \left(\left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 + \kappa \left\| \frac{\partial \nabla u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 \right) dt \le K_2$$
(4.14)

for some positive constant $K_2 = C\left(M^*, \mu, \kappa, d, \|f\|_{L^2(0,T;L^2(\Omega))}, \Omega, T, K_1\right)$. In view of (4.2), the relevant information to extract from the estimate (4.14) is given by (4.3).

(4) We start by testing (3.25) with $\psi = \mathbb{A}(u_n^j(t))$, where \mathbb{A} is the Stokes operator considered in (2.10),

$$-\kappa \int_{\Omega} \frac{\partial \Delta u_n^j(t)}{\partial t} \cdot \mathbb{A}(u_n^j(t)) \, dx - \mu \int_{\Omega} \Delta u_n^j(t) \cdot \mathbb{A}(u_n^j(t)) \, dx =$$

$$\int_{\Omega} \rho_n^j(t) f(t) \cdot \mathbb{A}(u_n^j(t)) \, dx - \int_{\Omega} \rho_n^j(t) \left[\frac{\partial u_n^j(t)}{\partial t} + \left(u_n^j(t) \cdot \nabla \right) u_n^j(t) \right] \cdot \mathbb{A}(u_n^j(t)) \, dx.$$

$$(4.15)$$

Writing u_n^j in the form (3.14) and using the linearity of the Stokes operator (2.10), together with (2.14) and (3.13), we can show that

$$-\kappa \int_{\Omega} \frac{\partial \Delta u_n^j(t)}{\partial t} \cdot \mathbb{A}(u_n^j(t)) \, dx = \frac{\kappa}{2} \frac{d}{dt} \int_{\Omega} |\mathbb{A}(u_n^j(t))|^2 \, dx,$$
$$-\mu \int_{\Omega} \Delta u_n^j(t) \cdot \mathbb{A}(u_n^j(t)) \, dx = \mu \int_{\Omega} |\mathbb{A}(u_n^j(t))|^2 \, dx.$$

Replacing in (4.15), we get

$$\frac{\kappa}{2}\frac{d}{dt}\left\|\mathbb{A}(u_n^j(t))\right\|_{L^2(\Omega)}^2 + \mu\left\|\mathbb{A}(u_n^j(t))\right\|_{L^2(\Omega)}^2 = \int_{\Omega}\rho_n^j(t)\left[f(t) - \frac{\partial u_n^j(t)}{\partial t} - \left(u_n^j(t) \cdot \nabla\right)u_n^j(t)\right] \cdot \mathbb{A}(u_n^j(t))\,dx.$$

Using the Hölder, Cauchy and Minkovski inequalities, together with (4.1), one has

$$\kappa \frac{d}{dt} \left\| \mathbb{A}(u_n^j(t)) \right\|_{L^2(\Omega)}^2 + \mu \left\| \mathbb{A}(u_n^j(t)) \right\|_{L^2(\Omega)}^2 \leq C \left(\left\| f(t) \right\|_{L^2(\Omega)}^2 + \left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 + \int_{\Omega} |u_n^j(t)|^2 |\nabla u_n^j(t)|^2 dx \right)$$

for some positive constant $C = C(\mu, M^*)$. Integrating between 0 and $t \in (0, T)$ and then taking the supremum in (0, T) in the resulting inequality,

$$\kappa \sup_{t \in (0,T)} \left\| \mathbb{A}(u_n^j(t)) \right\|_{L^2(\Omega)}^2 + \mu \int_0^T \left\| \mathbb{A}(u_n^j(t)) \right\|_{L^2(\Omega)}^2 dt \le \kappa \left\| \mathbb{A}(u_{0,n}^j) \right\|_{L^2(\Omega)}^2 + C\left(\int_0^T \|f(t)\|_{L^2(\Omega)}^2 dt + \int_0^T \left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 dt + \int_0^T \int_\Omega |u_n^j|^2 |\nabla u_n^j|^2 dx dt \right).$$
(4.16)

We now observe that, in view of (2.17) from one hand, and (2.7), (2.10) and (3.19) on the other, there holds

$$\left\| D^2(u_n^j(t)) \right\|_{L^2(\Omega)} \le C_1 \left\| \mathbb{A}(u_n^j(t)) \right\|_{L^2(\Omega)}^2, \tag{4.17}$$

$$\left\|\mathbb{A}(u_{0,n}^{j})\right\|_{L^{2}(\Omega)}^{2} \leq C_{2} \left\|\Delta u_{0,n}^{j}\right\|_{L^{2}(\Omega)}^{2} \leq C_{3} \left\|D^{2}u_{0,n}^{j}\right\|_{L^{2}(\Omega)}^{2}$$
(4.18)

for some positive constants $C_1 = C(\mu, \Omega)$, $C_2 = C(\mu)$ and $C_3 = C(\mu, d)$. Note that in the first inequality of (4.18) we have used the fact that the Leray projection \mathbb{P} commutes with the Laplacian for the eigenfunctions $\psi_i \in H^2(\Omega) \cap V$ (see (3.13)), which in turn can be proved by using the symmetry of \mathbb{P} , similarly to (3.13). Hence, by the application of (4.17)-(4.18) in (4.16), there holds

$$\kappa \sup_{t \in (0,T)} \left\| D^2 u_n^j(t) \right\|_{L^2(\Omega)}^2 + \mu \int_0^T \left\| D^2 u_n^j(t) \right\|_{L^2(\Omega)}^2 dt \le C \left(\left\| D^2 u_{0,n}^j \right\|_{L^2(\Omega)}^2 + \int_0^T \left\| f(t) \right\|_{L^2(\Omega)}^2 dt + \int_0^T \left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 dt + \int_0^T \int_\Omega |u_n^j|^2 |\nabla u_n^j|^2 dx dt \right)$$

$$(4.19)$$

for some positive constant $C = C(\mu, \kappa, M^*, d, \Omega)$. On the other hand, by combining the Hölder and Cauchy inequalities with the Sobolev inequalities (2.2)-(2.3) and (2.6), one has

$$\begin{split} \int_{0}^{T} \int_{\Omega} |u_{n}^{j}|^{2} |\nabla u_{n}^{j}|^{2} dx dt &\leq \int_{0}^{T} \|u_{n}^{j}(t)\|_{L^{2*}(\Omega)} \|u_{n}^{j}(t)\|_{L^{q}(\Omega)} \|\nabla u_{n}^{j}(t)\|_{L^{2}(\Omega)} \|\nabla u_{n}^{j}(t)\|_{L^{2*}(\Omega)} dt \\ & \left(\text{for } \frac{1}{2^{*}} + \frac{1}{q} + \frac{1}{2} + \frac{1}{2^{*}} = 1 \Leftrightarrow q = \frac{2d}{4-d}, \ 2 \leq d \leq 3 \right) \\ & \leq C_{1} \sup_{t \in (0,T)} \|\nabla u_{n}^{j}(t)\|_{L^{2}(\Omega)}^{2} \int_{0}^{t} \|\nabla u_{n}^{j}(t)\|_{L^{2}(\Omega)} \|D^{2}u_{n}^{j}(t)\|_{L^{2}(\Omega)} dt \quad (4.20) \\ & \leq C_{2} \int_{0}^{t} \|\nabla u_{n}^{j}(t)\|_{L^{2}(\Omega)} \|D^{2}u_{n}^{j}(t)\|_{L^{2}(\Omega)} dt \\ & \leq C_{3} \int_{0}^{t} \|\nabla u_{n}^{j}(t)\|_{L^{2}(\Omega)}^{2} dt + \frac{\mu}{2C} \int_{0}^{t} \|D^{2}u_{n}^{j}(t)\|_{L^{2}(\Omega)}^{2} dt, \end{split}$$

for some positive constants $C_1 = C(d, \Omega)$, $C_2 = C(\kappa, d, \Omega, K_1)$ and $C_3 = C(\mu, \kappa, M^*, d, \Omega, K_1)$, and where C is the positive constant from (4.19). Plugging (4.20) into (4.19) and using (3.24)₂, together with the estimates (4.2) and (4.3), we prove that (4.4) is verified for some positive constant $K_3 =$ $C(\mu, \kappa, M^*, d, \Omega, \left\| D^2 u_0 \right\|_{L^2(\Omega)}, \\ \|f\|_{L^2(0,T; L^2(\Omega))}, K_1, K_2).$

(5) In this case, we use the Hölder and Cauchy inequalities, together with (4.1), to estimate the r.h.s. terms of (4.11) as follows,

$$\int_{\Omega} \rho_n^j(t) f(t) \cdot \frac{\partial u_n^j(t)}{\partial t} dx \le \frac{1}{4} \left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 + \left\| \sqrt{\rho_n^j(t)} f(t) \right\|_{L^2(\Omega)}^2, \tag{4.21}$$

$$|J(t)| \leq \frac{1}{4} \left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 + M^* \int_{\Omega} |u_n^j(t)|^2 |\nabla u_n^j(t)|^2 dx.$$

$$(4.22)$$

Plugging (4.21) and (4.22) into (4.11), and integrating the resulting inequality between 0 and $t \in (0, T)$, we obtain

$$\mu \left\| \nabla u_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} + \int_{0}^{t} \left(\left\| \sqrt{\rho_{n}^{j}(s)} \frac{\partial u_{n}^{j}}{\partial s}(s) \right\|_{L^{2}(\Omega)}^{2} + \kappa \left\| \frac{\partial \nabla u_{n}^{j}(s)}{\partial s} \right\|_{L^{2}(\Omega)}^{2} \right) ds \leq$$

$$\mu \left\| \nabla u_{0,n}^{j} \right\|_{L^{2}(\Omega)}^{2} + 2 \int_{0}^{t} \|f(s)\|_{L^{2}(\Omega)}^{2} ds + 2M^{*} \int_{0}^{t} \int_{\Omega} |u_{n}^{j}|^{2} |\nabla u_{n}^{j}|^{2} dx ds.$$

$$(4.23)$$

On the other hand, observing that $\partial \Omega \in C^2$, we can use Lemma 5 to prove the existence of a unique weak solution $(w, p) : w \in H^2(\Omega)$ and $p \in H^1(\Omega)$, with $\int_{\Omega} p(x) dx = 0$, for the stationary Stokes problem

$$\operatorname{div} w = 0 \qquad \text{in} \quad \Omega, \tag{4.24}$$

$$-\mu\Delta w + \nabla p = \rho_n^j(t)f(t) - \left[\rho_n^j(t)\frac{\partial u_n^j(t)}{\partial t} - \rho_n^j(t)\left(u_n^j(t)\cdot\nabla\right)u_n^j(t)\right] \quad \text{in} \quad \Omega,$$
(4.25)

$$w = 0$$
 on $\partial\Omega$. (4.26)

Moreover, there exists a positive constant $C_2 = C(\mu, \Omega)$ such that

$$\|w\|_{H^{2}(\Omega)} + \|p\|_{H^{1}(\Omega)} \leq C_{2} \left\|\rho_{n}^{j}(t)f(t) - \left[\rho_{n}^{j}(t)\frac{\partial u_{n}^{j}(t)}{\partial t} - \rho_{n}^{j}(t)\left(u_{n}^{j}(t) \cdot \nabla\right)u_{n}^{j}(t)\right]\right\|_{L^{2}(\Omega)}.$$
 (4.27)

Note that for any $t \in (0,T)$, $w = u_n^j(t) + \sigma \frac{\partial u_n^j(t)}{\partial t}$ and $p = p_n^j(t)$, with $\sigma = \frac{\kappa}{\mu}$, satisfy the Stokes problem (4.24)-(4.26) and therefore (4.27) implies

$$\mu^{2} \left\| D^{2} u_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} + \mu \kappa \frac{d}{dt} \left\| D^{2} u_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} + \kappa^{2} \left\| \frac{\partial D^{2} u_{n}^{j}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} \leq C_{3} \left(M^{*} \left\| f(t) \right\|_{L^{2}(\Omega)}^{2} + \left\| \sqrt{\rho_{n}^{j}(t)} \frac{\partial u_{n}^{j}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} + M^{*} \int_{\Omega} |u_{n}^{j}(t)|^{2} |\nabla u_{n}^{j}(t)|^{2} dx \right),$$

$$(4.28)$$

where $C_3 = C(\mu, M^*, \Omega)$ is a positive constant. Integrating between 0 and $t \in (0, T)$, and using (3.18), one has

$$\mu^{2} \int_{0}^{t} \left\| D^{2} u_{n}^{j}(s) \right\|_{L^{2}(\Omega)}^{2} ds + \mu \kappa \left\| D^{2} u_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} + \kappa^{2} \int_{0}^{t} \left\| \frac{\partial D^{2} u_{n}^{j}(s)}{\partial s} \right\|_{L^{2}(\Omega)}^{2} ds \leq \mu \kappa \left\| D^{2} u_{0,n}^{j} \right\|_{L^{2}(\Omega)}^{2} + C_{3} \left(M^{*} \int_{0}^{t} \|f(s)\|_{L^{2}(\Omega)}^{2} ds + \int_{0}^{t} \left\| \sqrt{\rho_{n}^{j}(s)} \frac{\partial u_{n}^{j}}{\partial s}(s) \right\|_{L^{2}(\Omega)}^{2} ds + + M^{*} \int_{0}^{t} \int_{\Omega} |u_{n}^{j}|^{2} |\nabla u_{n}^{j}|^{2} dx ds \right).$$

$$(4.29)$$

Choosing $\delta > 0$ so small that $C_3 \delta < \frac{1}{2}$, we obtain from (4.23) and (4.29)

$$\mu \left\| \nabla u_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} + \int_{0}^{t} \left(\frac{1}{2} \left\| \sqrt{\rho_{n}^{j}(s)} \frac{\partial u_{n}^{j}}{\partial s}(s) \right\|_{L^{2}(\Omega)}^{2} + 2\kappa \left\| \frac{\partial \nabla u_{n}^{j}(s)}{\partial s} \right\|_{L^{2}(\Omega)}^{2} \right) ds + \delta \left(\mu^{2} \int_{0}^{t} \left\| D^{2} u_{n}^{j}(s) \right\|_{L^{2}(\Omega)}^{2} ds + \mu \kappa \left\| D^{2} u_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} + \kappa^{2} \int_{0}^{t} \left\| \frac{\partial D^{2} u_{n}^{j}(s)}{\partial s} \right\|_{L^{2}(\Omega)}^{2} ds \right) \leq \mu \left\| \nabla u_{0,n}^{j} \right\|_{L^{2}(\Omega)}^{2} + \delta \mu \kappa \left\| D^{2} u_{0,n}^{j} \right\|_{L^{2}(\Omega)}^{2} + C_{4} \int_{0}^{t} \| f(s) \|_{L^{2}(\Omega)}^{2} ds + C_{5} \int_{0}^{t} \int_{\Omega} |u_{n}^{j}(s)|^{2} |\nabla u_{n}^{j}(s)|^{2} dx ds$$

$$(4.30)$$

where $C_4 = C(M^*)$ and $C_5 = C(M^*)$ are distinct positive constants. To estimate the last term, we proceed as in (4.20), but using in the final part the Cauchy inequality with δ . Hence, we get

$$\int_{0}^{t} \int_{\Omega} |u_{n}^{j}(s)|^{2} |\nabla u_{n}^{j}(s)|^{2} dx ds \leq C_{6} \int_{0}^{t} \|\nabla u_{n}^{j}(s)\|_{L^{2}(\Omega)} \|D^{2} u_{n}^{j}(s)\|_{L^{2}(\Omega)} ds \\
\leq C_{6} \left(\frac{2C_{7}\mu^{2}}{\delta} \int_{0}^{t} \left\|\nabla u_{n}^{j}(s)\right\|_{L^{2}(\Omega)}^{2} ds + \frac{\delta\mu^{2}}{2C_{7}} \int_{0}^{t} \left\|D^{2} u_{n}^{j}(s)\right\|_{L^{2}(\Omega)}^{2} ds\right), \tag{4.31}$$

where $C_6 = \frac{C_2}{\mu}$ and $C_7 = C_6 C_5$, being C_2 given in (4.20) and C_5 in (4.30). Plugging (4.31) into (4.30), choosing δ in the above conditions, taking the supremum in (0, T) of the resulting inequality and using (3.24) and, again, (4.2), we obtain

$$\mu \sup_{t \in (0,T)} \left(\left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)}^2 + \kappa \left\| D^2 u_n^j(t) \right\|_{L^2(\Omega)}^2 \right)
+ \int_0^T \left(\left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 + \kappa \left\| \frac{\partial \nabla u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 \right) dt \qquad (4.32)
+ \mu^2 \int_0^T \left\| D^2 u_n^j(t) \right\|_{L^2(\Omega)}^2 dt + \kappa^2 \int_0^T \left\| \frac{\partial D^2 u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 dt \le K_4,$$

for some positive constant $K_4 = C\left(M^*, \mu, \kappa, d, \Omega, \|u_0\|_{2,\Omega}, \|\nabla u_0\|_{2,\Omega}, \|D^2 u_0\|_{L^2(\Omega)}, \|f\|_{L^2(0,T;L^2(\Omega))}, K_1\right).$

In view of (4.2), (4.3) and (4.4), the relevant information to extract from the estimate (4.32) is written in (4.5).

Remark 3. Estimate (4.5) and part of (4.4) can be obtained by using a slightly different approach. In fact, testing (3.25) with $\psi = \mathbb{A}\left(\frac{\partial u_n^j(t)}{\partial t}\right)$, where \mathbb{A} is the Stokes operator considered in (2.10), we obtain

$$\frac{\mu}{2}\frac{d}{dt}\left\|\mathbb{A}(u_n^j(t))\right\|_{L^2(\Omega)}^2 + \kappa \left\|\mathbb{A}\left(\frac{\partial u_n^j(t)}{\partial t}\right)\right\|_{L^2(\Omega)}^2 = \int_{\Omega} \rho_n^j(t) \left[f(t) - \frac{\partial u_n^j(t)}{\partial t} - \left(u_n^j(t) \cdot \nabla\right)u_n^j(t)\right] \cdot \mathbb{A}\left(\frac{\partial u_n^j(t)}{\partial t}\right) dx$$

Arguing as we did for (4.16), we have

$$\frac{\mu}{2} \sup_{t \in (0,T)} \left\| \mathbb{A}(u_n^j(t)) \right\|_{L^2(\Omega)}^2 + \frac{\kappa}{2} \int_0^T \left\| \mathbb{A}\left(\frac{\partial u_n^j(t)}{\partial t}\right) \right\|_{L^2(\Omega)}^2 dt \le C \left(\left\| A(u_{0,n}^j) \right\|_{L^2(\Omega)}^2 + \int_0^T \left\| f(t) \right\|_{L^2(\Omega)}^2 dt + \int_0^T \left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 dt + \int_0^T \int_\Omega |u_n^j|^2 |\nabla u_n^j|^2 dx dt \right),$$

for some positive constant $C = C(\kappa)$. Next, by using (4.17)-(4.18) and

$$\left\| D^2 \left(\frac{\partial u_n^j(t)}{\partial t} \right) \right\|_{L^2(\Omega)} \le C \left\| \mathbb{A} \left(\frac{\partial u_n^j(t)}{\partial t} \right) \right\|_{L^2(\Omega)}^2$$

where $C = C(\mu, \Omega)$ is a positive constant, we obtain

$$\mu \sup_{t \in (0,T)} \left\| D^2 \left(u_n^j(t) \right) \right\|_{L^2(\Omega)}^2 + \kappa \int_0^T \left\| \frac{\partial D^2 \left(u_n^j(t) \right)}{\partial t} \right\|_{L^2(\Omega)}^2 dt \le C \left(\left\| D^2 (u_{0,n}^j) \right\|_{L^2(\Omega)}^2 + \int_0^T \| f(t) \|_{L^2(\Omega)}^2 dt + \int_0^T \left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 dt + \int_0^T \int_\Omega |u_n^j|^2 |\nabla u_n^j|^2 dx dt \right).$$

$$(4.33)$$

To estimate the last term, we proceed in a slightly different way than in (4.20),

$$\int_{0}^{T} \int_{\Omega} |u_{n}^{j}|^{2} |\nabla u_{n}^{j}|^{2} dx dt \leq C_{1} \sup_{t \in (0,T)} \|\nabla u_{n}^{j}(t)\|_{L^{2}(\Omega)}^{2} \int_{0}^{T} \|\nabla u_{n}^{j}(t)\|_{L^{2}(\Omega)} \|D^{2}u_{n}^{j}(t)\|_{L^{2}(\Omega)} dt \\
\leq C_{2} \left(\int_{0}^{T} \|\nabla u_{n}^{j}(t)\|_{L^{2}(\Omega)}^{2} dt + \frac{\mu}{2CC_{2}T} \int_{0}^{T} \|D^{2}u_{n}^{j}(t)\|_{L^{2}(\Omega)}^{2} dt \right) \qquad (4.34)$$

$$\leq C_{2} \int_{0}^{T} \|\nabla u_{n}^{j}(t)\|_{L^{2}(\Omega)}^{2} dt + \frac{\mu}{2C} \sup_{t \in (0,T)} \|D^{2}u_{n}^{j}(t)\|_{L^{2}(\Omega)}^{2} dt$$

for some positive constants $C_1 = C(d, \Omega)$, $C_2 = C(\mu, d, \Omega, T, K_1)$, and where C is the constant from (4.33). Plugging (4.34) into (4.33), and using (3.24)₂, together with the estimates (4.2) and (4.3), we prove that

$$\mu \sup_{t \in (0,T)} \left\| D^2 u_n^j(t) \right\|_{L^2(\Omega)}^2 + \kappa \int_0^T \left\| \frac{\partial D^2 \left(u_n^j(t) \right)}{\partial t} \right\|_{L^2(\Omega)}^2 dt \le K$$

$$(4.35)$$

for some positive constant $K = C(\mu, M^*, d, \Omega, T, \left\| D^2 u_0 \right\|_{L^2(\Omega)}, \left\| f \right\|_{L^2(0,T;L^2(\Omega))}, K_1, K_2)$. Finally, that (4.5) follows from (4.35) is immediate.

Before proceeding with the proof of the estimates (4.6) and (4.7), it should be stressed that, despite the reasoning is more direct when the estimate (4.5) is obtained by the approach described in Remark 3, we lose some information, when we compare the estimate (4.35), which gives rise to it by this method, with the estimate (4.32), from which (4.5) follows by the approach used in the proof of Proposition 1. More importantly, the method used in the proof of Proposition 1 will be of the utmost importance to prove the estimate (4.7) below.

Proof. (Proposition 2) Here we also shall split the proof into the two enumerated items.

(1) We rewrite the identity (4.11) as follows,

$$\left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 + \kappa \left\| \frac{\partial \nabla u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 = -\mu \int_{\Omega} \nabla u_n^j(t) : \frac{\partial \nabla u_n^j(t)}{\partial t} \, dx + J(t) + \int_{\Omega} \rho_n^j(t) f(t) \cdot \frac{\partial u_n^j(t)}{\partial t} \, dx.$$
(4.36)

Proceeding as we did for (4.13), we get

$$|J(t)| \le \frac{\kappa}{4} \left\| \frac{\partial \nabla u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 + C \left(\sup_{t \in (0,T)} \left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)}^2 \right)^2, \tag{4.37}$$

for some positive constant $C = C(M^*, \kappa, d, \Omega)$. Using the Cauchy inequality, one has

$$\left|-\mu \int_{\Omega} \nabla u_n^j(t) : \frac{\partial \nabla u_n^j(t)}{\partial t} \, dx\right| \le \frac{\kappa}{4} \left\|\frac{\partial \nabla u_n^j(t)}{\partial t}\right\|_{L^2(\Omega)}^2 + \frac{\mu^2}{\kappa} \left\|\nabla u_n^j(t)\right\|_{L^2(\Omega)}^2. \tag{4.38}$$

Plugging (4.12) and (4.37)-(4.38) into (4.36), we get

$$\frac{1}{2} \left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 + \frac{\kappa}{2} \left\| \frac{\partial \nabla u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 \leq \frac{\mu^2}{\kappa} \left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)}^2 + C \left(\sup_{t \in (0,T)} \left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)}^2 \right)^2 + \frac{1}{2} \left\| \sqrt{\rho_n^j(t)} f(t) \right\|_{L^2(\Omega)}^2.$$

$$(4.39)$$

Taking the supremum in (0, T) of (4.39) and using the hypothesis (1.12), together with the estimates (4.1) and (4.2), we show that (4.6) holds true for some positive constant $K'_2 = C\left(M^*, \mu, \kappa, d, \Omega, \|f\|_{L^{\infty}(0,T;L^2(\Omega))}, K_1(\Omega)\right)$ (2) To prove this case, we first observe that (4.28) can be written as follows,

$$\mu^{2} \left\| D^{2} u_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} + \kappa^{2} \left\| \frac{\partial D^{2} u_{n}^{j}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} \leq -2\mu\kappa \int_{\Omega} D^{2} u_{n}^{j}(t) : \frac{\partial D^{2} u_{n}^{j}(t)}{\partial t} \, dx$$

$$+ C_{3} \left(M^{*} \left\| f(t) \right\|_{L^{2}(\Omega)}^{2} + \left\| \sqrt{\rho_{n}^{j}(t)} \frac{\partial u_{n}^{j}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} + M^{*} \int_{\Omega} |u_{n}^{j}(t)|^{2} |\nabla u_{n}^{j}(t)|^{2} dx \right).$$

$$(4.40)$$

Proceeding as we did for (4.20), we get

$$\kappa^{2} \left\| \frac{\partial D^{2} u_{n}^{j}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} \leq C_{4} \left(\left\| f(t) \right\|_{L^{2}(\Omega)}^{2} + \left\| \sqrt{\rho_{n}^{j}(t)} \frac{\partial u_{n}^{j}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} + \left\| \nabla u_{n}^{j}(s) \right\|_{L^{2}(\Omega)}^{2} + \left\| D^{2} u_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} \right)$$

$$(4.41)$$

for some positive constant $C_4 = C(\mu, \kappa, M^*, d, \Omega, K_1)$.

Taking the supremum of (4.41) in (0, T), and using the assumption (1.12), together with the estimates (4.1), (4.2), (4.4) and (4.6), we prove that (4.7) holds for some positive constant $K'_4 = C(\mu, \kappa, M^*, d, \Omega, K_1, K_3, K'_2)$.

Proposition 3. Assume $2 \le d \le 3$ and let u_n^j , ρ_n^j and p_n^j be the approximate weak solutions of the problem (3.8)-(3.12) that have been found in (3.14), (3.15) and (3.26).

(1) If (1.6), (1.8), (1.9) and (1.11) hold, then there exists an independent of j (and n) positive constant K_5 such that

$$\int_{0}^{T} \left\| \nabla p_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} dt \leq K_{5}$$
(4.42)

(2) If (1.6), (1.8), (1.9) and (1.12) hold, then there exists an independent of j (and n) positive constant K_6 such that

$$\sup_{t \in [0,T]} \left\| \nabla p_n^j(t) \right\|_{L^2(\Omega)}^2 \le K_6.$$
(4.43)

Proof. (1) Going back a little bit, we recall the Stokes problem (4.24)-(4.26) and the associated regularity result (4.27). Similarly to (4.28), (4.27) also implies

$$\mu^{2} \left\| D^{2} u_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} + \mu \kappa \frac{d}{dt} \left\| D^{2} u_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} + \kappa^{2} \left\| \frac{\partial D^{2} u_{n}^{j}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} + \left\| \nabla p_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} \leq C_{3} \left(\left\| f(t) \right\|_{L^{2}(\Omega)}^{2} + \left\| \sqrt{\rho_{n}^{j}(t)} \frac{\partial u_{n}^{j}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} + \int_{\Omega} |u_{n}^{j}(t)|^{2} |\nabla u_{n}^{j}(t)|^{2} dx \right)$$

$$(4.44)$$

for some positive constant $C_3 = C(\mu, M^*, \Omega)$. Departing from (4.44), and proceeding exactly in the same way as we did for (4.32), we can also show that

$$\mu \sup_{t \in (0,T)} \left(\left\| \nabla u_n^j(t) \right\|_{L^2(\Omega)}^2 + \kappa \left\| D^2 u_n^j(t) \right\|_{L^2(\Omega)}^2 \right)$$

$$+ \int_0^T \left(\left\| \sqrt{\rho_n^j(t)} \frac{\partial u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 + \kappa \left\| \frac{\partial \nabla u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 \right) dt$$

$$+ \mu^2 \int_0^T \left\| D^2 u_n^j(t) \right\|_{L^2(\Omega)}^2 dt + \kappa^2 \int_0^T \left\| \frac{\partial D^2 u_n^j(t)}{\partial t} \right\|_{L^2(\Omega)}^2 dt + \int_0^T \left\| \nabla p_n^j(t) \right\|_{L^2(\Omega)}^2 dt \le K_4,$$

$$(4.45)$$

for the same positive constant K_4 . Hence, (4.42) follows immediately from (4.45), with $K_5 = K_4$ and for K4 given in (4.5).

(2) In this case, and similarly to (4.40), we can write (4.44) as follows,

$$\mu^{2} \left\| D^{2} u_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} + \kappa^{2} \left\| \frac{\partial D^{2} u_{n}^{j}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} + \left\| \nabla p_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} \leq -2\mu\kappa \int_{\Omega} D^{2} u_{n}^{j}(t) : \frac{\partial D^{2} u_{n}^{j}(t)}{\partial t} dx + C_{3} \left(\left\| f(t) \right\|_{L^{2}(\Omega)}^{2} + \left\| \sqrt{\rho_{n}^{j}(t)} \frac{\partial u_{n}^{j}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} + \int_{\Omega} |u_{n}^{j}(t)|^{2} |\nabla u_{n}^{j}(t)|^{2} dx \right).$$

$$(4.46)$$

Using (4.46), and proceeding as we did for (4.41), one has

$$\left\|\nabla p_{n}^{j}(t)\right\|_{L^{2}(\Omega)}^{2} \leq C_{4}\left(\left\|f(t)\right\|_{L^{2}(\Omega)}^{2} + \left\|\sqrt{\rho_{n}^{j}(t)}\frac{\partial u_{n}^{j}(t)}{\partial t}\right\|_{L^{2}(\Omega)}^{2} + \left\|\nabla u_{n}^{j}(t)\right\|_{L^{2}(\Omega)}^{2} + \left\|D^{2}u_{n}^{j}(t)\right\|_{L^{2}(\Omega)}^{2}\right)$$

$$(4.47)$$

for some positive constant $C_4 = C(\mu, \kappa, M^*, d, \Omega, K_1)$.

Finally, justifying as we did to show that (4.7) is a consequence of (4.41), we can also prove that (4.47) implies (4.43), with $K_6 = K'_4$ and for K'_4 given in (4.7).

5. Passing to the limit $j \to \infty$

In this section, we perform the proof of Theorem 1, which in fact has been started in the previous sections.

Proof. (Theorem 1) Due to (4.2)-(4.7), we can use the Banach-Alaoglu theorem to extract subsequences (still labelled by the superscript j) such that

$$u_n^j \xrightarrow{j \to \infty} u_n \quad \text{in } L^2(0,T;V), \qquad \qquad u_n^j \xrightarrow{*}_{j \to \infty} u_n \quad \text{in } L^\infty(0,T;V), \tag{5.1}$$

$$u_n^j \xrightarrow{j \to \infty} u_n \quad \text{in } L^2(0,T; H^2(\Omega)), \qquad \qquad u_n^j \xrightarrow{*}_{j \to \infty} u_n \quad \text{in } L^\infty(0,T; H^2(\Omega)), \tag{5.2}$$

$$\frac{\partial u_n^j}{\partial t} \xrightarrow{j \to \infty} \frac{\partial u_n}{\partial t} \quad \text{in } L^2(0,T;V), \qquad \qquad \frac{\partial u_n^j}{\partial t} \xrightarrow{*} \frac{\partial u_n}{\partial t} \quad \text{in } L^\infty(0,T;V), \tag{5.3}$$

$$\frac{\partial u_n^j}{\partial t} \xrightarrow{j \to \infty} \frac{\partial u_n}{\partial t} \quad \text{in } L^2(0,T; H^2(\Omega)), \qquad \frac{\partial u_n^j}{\partial t} \xrightarrow{*} \frac{\partial u_n}{j \to \infty} \frac{\partial u_n}{\partial t} \quad \text{in } L^\infty(0,T; H^2(\Omega)), \tag{5.4}$$

$$p_n^j \xrightarrow{j \to \infty} p_n \quad \text{in } L^2(0,T; W^{1,2}(\Omega)), \qquad p_n^j \xrightarrow{*}_{j \to \infty} p_n \quad \text{in } L^\infty(0,T; W^{1,2}(\Omega)).$$
(5.5)

Then, we can use the Aubin-Lions compactness lemma (see (2.4) in Lemma 2), together with (5.1), (5.3) and the compact embedding $W_0^{1,2}(\Omega) \hookrightarrow L^2(\Omega)$, so that

$$u_n^j \xrightarrow[j \to \infty]{} u_n \quad \text{in } L^2(0, T; L^2(\Omega))$$

$$(5.6)$$

for some subsequence still labelled by u_n^j . On the other hand, in view of (4.1), the Banach-Alaoglu theorem also allows us to extract a subsequence (still labelled by ρ_n^j) such that

$$\rho_n^j \xrightarrow[j \to \infty]{\star} \rho_n \quad \text{in} \quad L^{\infty}(0, T; L^{\infty}(\Omega)).$$
(5.7)

Moreover, ρ_n satisfies

$$\frac{\partial \rho_n}{\partial t} + \operatorname{div}(\rho_n u_n) = 0 \quad \text{in} \quad Q_T, \tag{5.8}$$

$$\frac{1}{n} \le \rho_n \le M^* < \infty \qquad \text{in } Q_T.$$
(5.9)

In addition, using the fact that div $u_n = 0$ in Q_T , together with (5.8), and with the estimates (4.1) and (4.2), it can be proved that

$$\frac{\partial \rho_n^j}{\partial t} \text{ is uniformly bounded in } L^2(0,T;W^{-1,2^*}(\Omega)).$$
(5.10)

Moreover the following compact and continuous embeddings hold,

$$L^{\infty}(\Omega) \hookrightarrow W^{-1,\infty}(\Omega) \hookrightarrow W^{-1,2^*}(\Omega).$$
 (5.11)

Then, (4.1), (5.10) and (5.11) allow us to use the Aubin-Lions compactness lemma (see (2.5) in Lemma 2) so that, for some subsequence (still labelled by ρ_n^j)

$$\rho_n^j \xrightarrow[j \to \infty]{} \rho_n \quad \text{in } C([0,T]; W^{-1,\infty}(\Omega)).$$
(5.12)

On the other hand, it follows from $(3.11)_2$ and $(3.18)_3$, together with (4.8), that

$$\left\|\rho_{n}^{j}(t)\right\|_{L^{2}(\Omega)}^{2} = \left\|\rho_{0,n}\right\|_{L^{2}(\Omega)}^{2} \quad \text{and} \quad \left\|\rho_{n}(t)\right\|_{L^{2}(\Omega)}^{2} = \left\|\rho_{0,n}\right\|_{L^{2}(\Omega)}^{2} \quad \forall \ t \in [0,T].$$
(5.13)

Thus, applying (5.7) and (5.13), together with (5.12), we get for all $t \in [0, T]$

$$\left\|\rho_n^j(t) - \rho_n(t)\right\|_{L^2(\Omega)}^2 = \left\|\rho_n^j(t)\right\|_{L^2(\Omega)}^2 - \left\|\rho_n(t)\right\|_{L^2(\Omega)}^2 + 2\int_{\Omega} (\rho_n(t) - \rho_n^j(t))\rho(t) \, dx \xrightarrow{j \to \infty} 0.$$
(5.14)

As a consequence of (5.9) and (5.14), we have

$$\left\|\rho_n^j(t) - \rho_n(t)\right\|_{L^2(\Omega)}^2 \xrightarrow[j \to \infty]{} 0 \quad \forall \ q : 2 \le q < \infty.$$
(5.15)

Hence, (5.12) and (5.15) assure that

$$\rho_n^j \xrightarrow[j \to \infty]{} \rho_n \quad \text{in } C([0,T]; L^q(\Omega)) \quad \forall \ q: 2 \le q < \infty,$$

From this and (4.1), one has

$$\rho_n^j \xrightarrow[j \to \infty]{} \rho_n \quad \text{in } C([0,T]; L^q(\Omega)) \quad \forall \ q \ge 1.$$
(5.16)

By the application of (5.3) and (5.6), together with (5.9) and (5.16), we have

$$\rho_n^j \frac{\partial u_n^j}{\partial t} \xrightarrow{j \to \infty} \rho_n \frac{\partial u_n}{\partial t} \quad \text{in } L^2(0, T; L^2(\Omega)), \tag{5.17}$$

$$\rho_n^j u_n^j \xrightarrow{j \to \infty} \rho_n u_n \quad \text{in } L^r(0, T; L^r(\Omega)), \text{ with } 1 \le r < 2^*.$$
(5.18)

Gathering the information of (4.1), (4.2) and (4.3) with (5.6) and (5.16), we can prove that

$$\rho_n^j(u_n^j \cdot \nabla) u_n^j \xrightarrow[j \to \infty]{} \rho_n(u_n \cdot \nabla) u_n \quad \text{in} \quad L^1(0, T; L^1(\Omega)).$$
(5.19)

In addition, due to (5.2) and (5.4), we also have

$$\Delta u_n^j \xrightarrow{j \to \infty} \Delta u_n \quad \text{in } L^2(0, T; L^2(\Omega)), \tag{5.20}$$

$$\frac{\partial \Delta u_n^j}{\partial t} \xrightarrow[j \to \infty]{} \frac{\partial \Delta u_n}{\partial t} \quad \text{in } L^2(0, T; L^2(\Omega)).$$
(5.21)

Let now $\zeta \in C_0^{\infty}([0,T))$. Multiplying (3.25) by ζ and integrating the resulting equation between 0 and T, we obtain

$$\int_{Q_T} \left[\rho_n^j \frac{\partial u_n^j}{\partial t} + \rho_n^j \left(u_n^j \cdot \nabla \right) u_n^j - \mu \Delta u_n^j - \kappa \frac{\partial \Delta u_n^j}{\partial t} \right] \cdot \psi \zeta \, dx dt = \int_{Q_T} \rho_n^j f \cdot \psi \zeta \, dx dt \tag{5.22}$$

for all $\psi \in H^2(\Omega) \cap V$. Proceeding similarly for (3.27), we get

$$\int_{Q_T} \left[\rho_n^j \frac{\partial u_n^j}{\partial t} + \rho_n^j \left(u_n^j \cdot \nabla \right) u_n^j - \mu \Delta u_n^j - \kappa \frac{\partial \Delta u_n^j}{\partial t} \right] \cdot \psi \zeta \, dx dt - \int_{Q_T} \rho_n^j f \cdot \psi \zeta \, dx dt = \int_{Q_T} p_n^j \operatorname{div} \psi \zeta \, dx dt \tag{5.23}$$

for all $\psi \in H^2(\Omega) \cap W_0^{1,2}(\Omega)$. For the same function ζ , we multiply (3.17) by $\eta = \phi \zeta$, with $\phi \in C_0^{\infty}(\Omega)$, and integrate the resulting equation over Q so that

$$-\int_{Q_T} \rho_n^j \phi \,\zeta' \,dx dt - \int_{Q_T} \rho_n^j u_n^j \cdot \nabla \phi \,\zeta \,dx dt = \zeta(0) \int_{\Omega} \rho_{0,n}^j \phi \,dx \qquad \forall \ \phi \in C_0^\infty(\Omega).$$
(5.24)

Then, we use, for each corresponding term of (5.22), the convergence results (5.17), (5.19), (5.20)-(5.21), together with (5.16), to pass the equation (5.22) to the limit $j \to \infty$. To pass (5.23) to the limit $j \to \infty$, we use, in addition, (5.5).

The passage of the equation (5.24) to the limit $j \to \infty$ uses (5.16) and (5.18) together with (3.21). After all, we obtain

$$\int_{Q_T} \left[\rho_n \frac{\partial u_n}{\partial t} + \rho_n \left(u_n \cdot \nabla \right) u_n - \mu \Delta u_n - \kappa \frac{\partial \Delta u_n}{\partial t} \right] \cdot \psi \zeta \, dx = \int_{Q_T} \rho_n f \cdot \psi \zeta \, dx dt \tag{5.25}$$

for all $\psi \in H^2(\Omega) \cap V$ and all $\zeta \in C_0^{\infty}([0,T))$,

$$\int_{Q_T} \left[\rho_n \frac{\partial u_n}{\partial t} + \rho_n \left(u_n \cdot \nabla \right) u_n - \mu \Delta u_n - \kappa \frac{\partial \Delta u_n}{\partial t} \right] \cdot \psi \, dx dt - \int_{\Omega} \rho_n f \cdot \psi \, \zeta \, dx dt = \int_{Q_T} p_n \operatorname{div} \psi \, \zeta \, dx dt$$
(5.26)

for all $\psi \in H^2(\Omega) \cap W_0^{1,2}(\Omega)$ and all $\zeta \in C_0^{\infty}([0,T))$, and

$$-\int_{Q_T} \rho_n \eta \zeta' \, dx dt - \int_{Q_T} \rho_n u_n \cdot \nabla \eta \zeta \, dx dt = \zeta(0) \int_{Q_T} \rho_{0,n} \eta \, dx \tag{5.27}$$

for all $\phi \in C_0^{\infty}(\Omega)$ and all $\zeta \in C_0^{\infty}([0,T))$.

6. Passing to the limit $n \to \infty$

In the last section, we have proved that for each $n \in \mathbb{N}$ there exists, at least, a solution (ρ_n, u_n, p_n) such that

$$\int_{\Omega} \left[\rho_n \frac{\partial u_n}{\partial t} + \rho_n \left(u_n \cdot \nabla \right) u_n - \mu \Delta u_n - \kappa \frac{\partial \Delta u_n}{\partial t} \right] \cdot \psi \, dx = \int_{\Omega} \rho_n f \cdot \psi \, dx \tag{6.1}$$

for all $\psi \in H^2(\Omega) \cap V$, and

$$\int_{\Omega} \left[\rho_n \frac{\partial u_n}{\partial t} + \rho_n \left(u_n \cdot \nabla \right) u_n - \mu \Delta u_n - \kappa \frac{\partial \Delta u_n}{\partial t} \right] \cdot \psi \, dx - \int_{\Omega} \rho_n f \cdot \psi \, dx = \int_{\Omega} p_n \operatorname{div} \psi \, dx \qquad (6.2)$$

for all $\psi \in H^2(\Omega) \cap W_0^{1,2}(\Omega)$, and both in the distribution sense on (0,T). Moreover, from (5.8), we easily realize that ρ_n satisfies

$$\frac{\partial \rho_n}{\partial t} + u_n \cdot \nabla \rho_n = 0 \qquad \text{in} \quad Q.$$
(6.3)

Using the identities (6.1), (6.2) and (6.3), we can proceed as we did for (4.1), (4.2)-(4.7) and (4.42)-(4.43), using in this case (3.6) and (3.7), to show that

$$0 < \frac{1}{n} \le \inf_{x \in \overline{\Omega}} \rho_{0,n}(x) \le \rho_n(x,t) \le \sup_{x \in \overline{\Omega}} \rho_{0,n}(x) \le M^* < \infty \qquad \forall \ (x,t) \in Q_T,$$
(6.4)

$$\sup_{t \in (0,T)} \left(\left\| \sqrt{\rho_n(t)} u_n(t) \right\|_{L^2(\Omega)}^2 + \kappa \left\| \nabla u_n(t) \right\|_{L^2(\Omega)}^2 \right) + \mu \int_0^T \left\| \nabla u_n(t) \right\|_{L^2(\Omega)}^2 dt \le K_1, \tag{6.5}$$

$$\int_{0}^{T} \left(\left\| \sqrt{\rho_{n}(t)} \frac{\partial u_{n}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} + \kappa \left\| \frac{\partial \nabla u_{n}(t)}{\partial t} \right\|_{L^{2}(\Omega)}^{2} \right) dt \leq K_{2},$$
(6.6)

$$\sup_{t \in [0,T]} \left(\left\| \sqrt{\rho_n(t)} \frac{\partial u_n(t)}{\partial t} \right\|_{L^2(\Omega)}^2 + \kappa \left\| \frac{\partial \nabla u_n(t)}{\partial t} \right\|_{L^2(\Omega)}^2 \right) \le K_2'$$
(6.7)

$$\kappa \sup_{t \in (0,T)} \left\| D^2 u_n(t) \right\|_{L^2(\Omega)}^2 + \mu \int_0^T \left\| D^2 u_n(t) \right\|_{L^2(\Omega)}^2 dt \le K_3,$$
(6.8)

$$\kappa^2 \int_0^T \left\| \frac{\partial D^2 u_n(t)}{\partial t} \right\|_{L^2(\Omega)}^2 dt \le K_4 \tag{6.9}$$

$$\kappa^2 \sup_{t \in [0,T]} \left\| \frac{\partial D^2 u_n(t)}{\partial t} \right\|_{L^2(\Omega)}^2 \le K_4', \tag{6.10}$$

$$\int_{0}^{T} \left\| \nabla p_{n}^{j}(t) \right\|_{L^{2}(\Omega)}^{2} dt \le K_{7}, \tag{6.11}$$

$$\sup_{t \in [0,T]} \left\| \nabla p_n^j(t) \right\|_{L^2(\Omega)}^2 \le K_8.$$
(6.12)

and where the positive constants K_1 , K_2 , K'_2 , K_3 , K_4 and K'_4 do not depend on n. Then, due to (6.5), (6.6), (6.7), (6.8), (6.9), (6.10), (6.11) and (6.12), we can use the Banach-Alaoglu theorem to extract subsequences (still labelled by the subscript n) such that

$$u_n \xrightarrow[n \to \infty]{} u \quad \text{in } L^2(0, T; V), \qquad \qquad u_n \xrightarrow[n \to \infty]{} u \quad \text{in } L^\infty(0, T; V), \qquad (6.13)$$

$$u_n \xrightarrow[n \to \infty]{} u \quad \text{in } L^2(0, T; H^2(\Omega)), \qquad u_n \xrightarrow[n \to \infty]{} u \quad \text{in } L^\infty(0, T; H^2(\Omega)), \tag{6.14}$$

$$\frac{\partial u_n}{\partial t} \xrightarrow[n \to \infty]{} \frac{\partial u}{\partial t} \quad \text{in } L^2(0, T; V), \qquad \qquad \frac{\partial u_n}{\partial t} \xrightarrow[n \to \infty]{} \frac{\partial u}{\partial t} \quad \text{in } L^\infty(0, T; V), \tag{6.15}$$

$$\frac{\partial u_n}{\partial t} \xrightarrow[n \to \infty]{} \frac{\partial u}{\partial t} \quad \text{in } L^2(0, T; H^2(\Omega)), \qquad \frac{\partial u_n}{\partial t} \xrightarrow[n \to \infty]{} \frac{\partial u}{\partial t} \quad \text{in } L^\infty(0, T; H^2(\Omega)), \tag{6.16}$$

$$p_n \xrightarrow[n \to \infty]{} p \quad \text{in } L^2(0, T; W^{1,2}(\Omega)), \qquad p_n \xrightarrow[n \to \infty]{} p \quad \text{in } L^\infty(0, T; W^{1,2}(\Omega)).$$
 (6.17)

Observing (6.13) and (6.15), we can use the Aubin-Lions compactness lemma (see (2.4) in Lemma 2), so that for some subsequence (still labelled by the subscript n)

$$\nabla u_n \xrightarrow[n \to \infty]{} \nabla u \quad \text{in} \quad L^2(0, T; L^2(\Omega)).$$
 (6.18)

On the other hand, by using (5.8), together with (6.4) and (6.5), it can be proved that

$$\frac{\partial \rho_n}{\partial t}$$
 is uniformly bounded in $L^2(0,T;W^{-1,2}(\Omega)).$ (6.19)

Now, due to (6.4) and (6.19), we can use the Aubin-Lions compactness lemma (see (2.5) in Lemma 2) so that, for some subsequence (still labelled by ρ_n),

$$\rho_n \xrightarrow[n \to \infty]{} \rho \quad \text{in } C([0,T]; W^{-1,2}(\Omega)).$$
(6.20)

Hence,

$$\rho_n u_n \xrightarrow[n \to \infty]{} \rho u \quad \text{in } (C_0([0, T] \times \Omega))'$$
(6.21)

 ρ is a solution of (1.3) and

$$0 \le \rho \le M^* < \infty \quad \text{in } Q_T,$$

$$\|\rho(t)\|_{L^q(\Omega)} = \|\rho_0\|_{L^q(\Omega)} \quad \forall q: 1 \le q \le \infty.$$

$$(6.22)$$

However, contrary to the case when ρ_0 is bounded away from 0, in this work u_n is not bounded in $L^2(0,T; L^2(\Omega))$. This brings us much more difficulty in the passage to the limit $n \to \infty$, which is overcame by the following result.

Lemma 6. Let the conditions of Theorem 1 be fulfilled and assume that (6.20) and (6.21) hold. Then there exist subsequences (still labelled by the subscript n) such that

$$\rho_n \xrightarrow[n \to \infty]{} \rho \quad in \ C([0, T]; L^q(\Omega)) \quad \forall \ q \ge 1,$$
(6.23)

$$\sqrt{\rho_n} u_n \xrightarrow[n \to \infty]{} \sqrt{\rho} u \quad in \ L^2(0, T; L^2(\Omega)).$$
 (6.24)

Proof. The proof of Lemma 6 uses the DiPerna-Lions [14] theory for linear transport equations combined with renormalization arguments. For the proof see [24, Theorem 2.5] and Desjardins [13]. \Box

In order to apply this result, we observe that combining (5.8) and (3.8) with (5.26) and (5.27), we can show that

$$-\int_{Q_T} (\rho_n u_n - \kappa \Delta u_n) \cdot \frac{\partial \varphi}{\partial t} dx dt - \mu \int_{Q_T} \Delta u_n \cdot \varphi \, dx dt - \int_{Q_T} \rho_n u_n \otimes u_n : \nabla \varphi \, dx dt -\int_{Q_T} \rho_n f \cdot \varphi \, dx dt = \int_{\Omega} (\rho_{n,0} u_{n,0} \cdot \varphi(0) + \kappa \nabla u_{n,0} : \nabla \varphi(0)) \, dx + \int_{Q_T} p_n \operatorname{div} \varphi \, dx$$
(6.25)

for all $\varphi \in C_0^1([0,T); W_0^{1,2}(\Omega) \cap H^2(\Omega))$, and

$$-\int_{Q_T} \rho_n \frac{\partial \phi}{\partial t} \, dx dt - \int_{Q_T} \rho_n u_n \cdot \nabla \phi \, dx dt = \int_{\Omega} \rho_{0,n} \phi(0) \, dx \tag{6.26}$$

for all $\phi \in C_0^{\infty}([0,T) \times \Omega)$. We can proceed for (5.25) in the same way as we did for (6.25).

Now, we can use (6.13)-(6.14), (6.17), (6.18), (6.23) and (6.24), together with (3.1), (3.2) and (3.4) to pass the integral identity (6.25) to the limit $n \to \infty$ so that, in view of the regularity of ρ , u and p, (1.2) holds almost everywhere in Q_T . By using (6.23) and (6.24), together with (3.2) and (3.4), we can also pass the integral identity (6.26) to the limit $n \to \infty$ so that, in view of the regularity of ρ and u, (1.3) holds almost everywhere in Q_T .

Finally, combining the estimates (6.4) and (6.5)-(6.10) with the convergence results (6.13)-(6.17) and (6.23)-(6.24), we can show that the enumerated items (1)-(5) of Theorem 1 and (1)-(4) of Theorem 2 hold true.

Remark 4. In addition to the estimates (6.8) and (6.9), we can see that, by combining the Sobolev inequalities (2.7) and (2.9) with these estimate, we easily obtain

$$\kappa \sup_{t \in (0,T)} \|u_n(t)\|_{C^{0,\alpha}(\overline{\Omega})}^2 \le K_7, \quad 0 < \alpha \le 2 - \frac{d}{2}, \ 2 \le d \le 3,$$

$$\kappa^2 \int_0^T \left\|\frac{\partial u_n(t)}{\partial t}\right\|_{C^{0,\alpha}(\overline{\Omega})}^2 dt \le K_8, \quad 0 < \alpha \le 2 - \frac{d}{2}, \ 2 \le d \le 3,$$

for some positive constants $K_7 = C(d, K_3)$ $K_8 = C(d, K_4)$.

7. Proof of Theorem 3

In this section, we shall prove the uniqueness of the components u and ρ of a solution (u, ρ, p) of the problem (1.1)-(1.5). Before doing so, let us invoke some results that provides us with further regularity on the solutions (u, ρ, p) of the problem (1.1)-(1.5).

We firstly recall a result about the regularity of the density ρ for the nonhomogeneous Navier-Stokes equations. This result is still useful here, because the continuity equation is the same for both nonhomogeneous Navier-Stokes and Navier-Stokes-Voigt systems of equations.

Proposition 4. Let (u, ρ, p) be a solution of the problem (1.1)-(1.5) in the conditions of Theorem 2. If (1.19) and

$$u \in C([0,T]; W^{1,\infty}(\Omega)) \tag{7.1}$$

hold, then

$$\|\nabla\rho(t)\|_{L^{\infty}(\Omega)} \leq \sqrt{d} \|\nabla\rho_0\|_{L^{\infty}(\Omega)} \exp\left(\int_0^t \|\nabla u(s)\|_{L^{\infty}(\Omega)} \, ds\right),\tag{7.2}$$

$$\|\rho_t(t)\|_{L^{\infty}(\Omega)} \le \sqrt{d} \|\nabla\rho_0\|_{L^{\infty}(\Omega)} \|u(t)\|_{L^{\infty}(\Omega)} \exp\left(\int_0^t \|\nabla u(s)\|_{L^{\infty}(\Omega)} \, ds\right)$$
(7.3)

for all $t \in [0, T]$.

Proof. We address the proof of Proposition 4 to Ladyzhenskaya and Solonnikov [19, Lemma 1.3]. \Box

In the Navier-Stokes-Voigt setting, note that, in view of (7.4) below, assumption (7.1) can be shown to be satisfied if one assume that, in addition to (1.19), (1.20)-(1.21) and (1.22) are also verified. By assuming these hypotheses are satisfied altogether, one can show the boundedness of $\|\nabla \rho(t)\|_{L^{\infty}(\Omega)}$ and $\|\rho_t(t)\|_{L^{\infty}(\Omega)}$, which in a certain sense can replace (7.2) and (7.3) in the Navier-Stokes-Voigt case (see (7.5) below).

The next result shows us how a higher regularity of the solutions depends on the smoothness of the problem data ρ_0 , u_0 and f.

Proposition 5. Let (u, ρ, p) be a solution of the problem (1.1)-(1.5) in the conditions of Theorem 2. If, in addition to (1.19), is verified (1.20)-(1.21) and (1.22), then there exist positive constants K_9 and K_{10} such that

$$\sup_{t \in [0,T]} \left(\left\| D^2 u(t) \right\|_{L^r(\Omega)}^2 + \left\| \nabla u(t) \right\|_{L^{\infty}(\Omega)}^2 \right) + \left\| \nabla p \right\|_{L^2(0,T;L^r(\Omega))}^2 \le K_9, \tag{7.4}$$

$$\sup_{t \in [0,T]} \left(\|\nabla \rho\|_{L^{\infty}(\Omega)}^2 + \|\rho_t\|_{L^{\infty}(\Omega)}^2 \right) \le K_{10}.$$
(7.5)

Proof. Using the assumptions (1.19) and (1.20)-(1.21), we have proved in [3, Theorem 4] that

$$\sup_{t \in [0,T]} \left(\|\Delta u(t)\|_{L^{r}(\Omega)}^{2} + \|\nabla u(t)\|_{C^{0,\alpha}(\overline{\Omega})}^{2} \right) + \|\nabla p\|_{L^{2}(0,T;L^{r}(\Omega))}^{2} \le C_{1},$$
(7.6)

$$\sup_{t \in [0,T]} \left(\|\nabla \rho\|_{L^{\infty}(\Omega)}^{2} + \|\rho_{t}\|_{L^{\infty}(\Omega)}^{2} \right) \le C_{2}$$
(7.7)

for some positive constants $C_1 = C\left(\mu, \kappa, M^*, d, \Omega, T, \|\nabla u_0\|_{L^2(\Omega)}, \|\Delta u_0\|_{L^r(\Omega)}, \|f\|_{L^2(0,T;L^r(\Omega))}\right)$ and $C_2 = C\left(\mu, \kappa, M^*, d, \Omega, T, \|\nabla \rho_0\|_{L^\infty(\Omega)}^2, \|\nabla u_0\|_{L^2(\Omega)}, \|\Delta u_0\|_{L^2(\Omega)}, \|f\|_{L^2(0,T;L^r(\Omega))}\right)$. The restriction (1.22) results by the application of the Sobolev inequalities (2.2) and (2.6) as follows,

$$\begin{aligned} \|u_t(t)\|_{L^r(\Omega)} &\leq C(r, d, \Omega) \|\nabla u_t(t)\|_{L^2(\Omega)}, \quad r \leq 2^*, \\ \|\nabla u(t)\|_{L^\infty(\Omega)} &\leq C(r, d) \|D^2 u(t)\|_{L^r(\Omega)}, \quad r > d. \end{aligned}$$

It is immediate now that (7.4) and (7.5) is a consequence of the estimates (7.6)-(7.7) by the application of (2.7).

We are now in conditions to prove Theorem 3. In [3, Theorems 4-5] we already have proved a uniqueness result in the case of a strictly positive initial density and under the same assumptions (1.19), (1.20)-(1.21) and (1.22). The issue of wether or not the initial density may vanish in a subdomain of Ω does not matter for this result. Therefore, we can prove Theorem 3 telegraphically, addressing the details to [3].

Proof. (Theorem 3) Let $(\hat{u}, \hat{p}, \hat{\rho})$ and $(\overline{u}, \overline{p}, \overline{\rho})$ be two solutions of the problem (1.1)-(1.5) with the same data. By algebraic manipulations of the equations (1.1)-(1.5), satisfied by each of these two couple of solutions, one has

$$\hat{\rho}u_t + \hat{\rho}\left(\hat{u}\cdot\nabla\right)u - \mu\Delta u - \kappa\Delta u_t + \nabla p = \rho\left[f - \overline{u}_t - \left(\overline{u}\cdot\nabla\right)\overline{u}\right] - \hat{\rho}\left(u\cdot\nabla\right)\overline{u}$$
(7.8)

$$\rho_t + \nabla \hat{\rho} \cdot u + \nabla \rho \cdot \hat{u} = 0, \tag{7.9}$$

$$\operatorname{div} u = 0, \tag{7.10}$$

where $u = \hat{u} - \overline{u}$, $p = \hat{p} - \overline{p}$ and $\rho = \hat{\rho} - \overline{\rho}$. Multiplying (7.8) by u, next integrating the resulting identity over Ω and using (7.10), we have

$$\frac{1}{2}\frac{d}{dt}\left(\left\|\sqrt{\hat{\rho}(t)}u(t)\right\|_{L^{2}(\Omega)}^{2}+\kappa\left\|\nabla u(t)\right\|_{L^{2}(\Omega)}^{2}\right)+\mu\left\|\nabla u(t)\right\|_{L^{2}(\Omega)}^{2}=\int_{\Omega}\left(\rho(t)\left[f(t)-\overline{u}_{t}(t)-\left(\overline{u}(t)\cdot\nabla\right)\overline{u}(t)\right]-\hat{\rho}(t)\left(u(t)\cdot\nabla\right)\overline{u}(t)\right)\cdot u(t)dx$$
(7.11)

Multiplying now (7.9) by ρ , integrating the resulting equation over Ω , and using (7.10), we also have

$$\frac{1}{2}\frac{d}{dt}\|\rho(t)\|_{L^2(\Omega)}^2 = -\int_{\Omega}\rho(t)(u(t)\cdot\nabla)\overline{\rho}(t)dx.$$
(7.12)

Adding up (7.11) and (7.12), we have

$$\frac{1}{2}\frac{d}{dt}\left(\left\|\sqrt{\hat{\rho}(t)}u(t)\right\|_{L^{2}(\Omega)}^{2}+\kappa\left\|\nabla u(t)\right\|_{L^{2}(\Omega)}^{2}+\left\|\rho(t)\right\|_{L^{2}(\Omega)}^{2}\right)+\mu\left\|\nabla u(t)\right\|_{L^{2}(\Omega)}^{2}=\int_{\Omega}\left(\rho(t)\left[f(t)-\overline{u}_{t}(t)-\left(\overline{u}(t)\cdot\nabla\right)\overline{u}(t)\right]-\hat{\rho}(t)\left(u(t)\cdot\nabla\right)\overline{u}(t)\right)\cdot u(t)dx-\int_{\Omega}\rho(t)(u(t)\cdot\nabla)\overline{\rho}(t)dx$$
(7.13)

for all $t \in [0, T]$. To estimate the r.h.s. terms of (7.13), we proceed as we did in the proof of [3, Theorem 5], using in particular (1.19), (6.22) and (7.4)-(7.5). Proceeding so, we obtain

$$\frac{d}{dt} \left(\left\| \sqrt{\hat{\rho}(t)} u(t) \right\|_{L^{2}(\Omega)}^{2} + \kappa \left\| \nabla u(t) \right\|_{L^{2}(\Omega)}^{2} + \left\| \rho(t) \right\|_{L^{2}(\Omega)}^{2} \right) \leq A(t) \left(\left\| \sqrt{\hat{\rho}(t)} u(t) \right\|_{L^{2}(\Omega)}^{2} + \kappa \left\| \nabla u(t) \right\|_{L^{2}(\Omega)}^{2} + \left\| \rho(t) \right\|_{L^{2}(\Omega)}^{2} \right) \tag{7.14}$$

for all $t \in [0, T]$ and for some function A that, in view of Theorem 2, (1.19), (6.22), and (7.4)-(7.5), can be proven to belong to $L^1(0, T)$. Therefore we can apply the Grönwall inequality to (7.14), which will finally allow us to conclude that $\hat{\rho} = \overline{\rho}$ and $\hat{u} = \overline{u}$.

Acknowledgments

All authors were supported by the Grants no. AP19676624 of the Ministry of Education and Science of the Republic of Kazakhstan (Kazakhstan). The first author was also partially supported by the

Portuguese Foundation for Science and Technology (FCT) under the project no. UIDB/04561/2020 (Portugal).

References

- S.N. Antontsev, A.V. Kazhikhov and V.N. Monakhov. Boundary Value Problems in Mechanics of Nonhomogeneous Fluids. (Translation from the original Russian edition, Nauka, Novosibirsk, 1983), North-Holland, Amsterdam, 1990. 1
- [2] S.N. Antontsev and H.B. de Oliveira. Cauchy problem for the Navier-Stokes-Voigt model governing nonhomogeneous flows. *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM* **116** (2022), no. 4, Paper no. 158. 1, 1, 3, (1)
- [3] S.N. Antontsev, H.B. de Oliveira and Kh. Khompysh, The classical Kelvin-Voigt problem for nonhomogeneous and incompressible fluids: existence, uniqueness and regularity. *Nonlinearity* **34** (2021), no. 5, 3083–3111. 1, 1, 2, 3, 3, (2), (1), 7, 7, 7
- [4] S.N. Antontsev, H.B. de Oliveira and Kh. Khompysh. Existence and large time behavior for generalized Kelvin-Voigt equations governing nonhomogeneous and incompressible fluids. J. Phys. Conf. Ser. 1268 (2019), 012008.
 1
- [5] S.N. Antontsev, H.B. de Oliveira and Kh. Khompysh. Generalized Kelvin-Voigt equations for nonhomogeneous and incompressible fluids. *Commun. Math. Sci.* 17 (2019), no.7, 1915–1948.
- [6] M.E. Bogovskii. Solutions of some problems of vector analysis, associated with the operators div and grad. (Russian). Theory of cubature formulas and the application of functional analysis to problems of mathematical physics, 5–40, 149. Trudy Sem. S. L. Soboleva, no. 1, 1980. Akad. Nauk SSSR Sibirsk. Otdel., Inst. Mat., Novosibirsk, 1980. 2
- [7] H. Brezis. Analyse fonctionnelle. Masson, Paris, 1983. 1
- [8] H.J. Choe and H. Kim. Strong solutions of the Navier-Stokes equations for nonhomogeneous incompressible fluids. Comm. Partial Differential Equations 28 (2003), no. 5-6, 1183–1201. 1, 1, 1
- H.J. Choe and H. Kim. Strong solutions of the Navier-Stokes equations for isentropic compressible fluids. J. Differential Equations 190 (2003), no. 2, 504–523.
- [10] D. Cioranescu, V. Girault and K.R. Rajagopal. Mechanics and mathematics of fluids of the differential type. Springer, Swuitzerland, 2016. 1
- [11] P. Constantin and C. Foias, Navier-Stokes equations, The University of Chicago Press, Chicago, 1988. 1, 3
- [12] R. Danchin and P.B. Mucha. The incompressible Navier-Stokes equations in vacuum. Comm. Pure Appl. Math. 72 (2019), 1351–1385.
- [13] B. Desjardins. Global existence results for the incompressible density-dependent Navier-Stokes equations in the whole space. *Differential Integral Equations* 10 (1997), no. 3, 587–598. 1, 6
- [14] R. J. DiPerna and P.-L. Lions. Ordinary differential equations, transport theory and Sobolev spaces. Invent. Math. 98 (1989), no. 3, 511–547. 6
- [15] G.P. Galdi. An introduction to the Mathematical Theory of the Navier-Stokes Equations. Steady-State Problems. Springer, New York, 2011. 1, 1, 2, 2
- [16] C. He, J. Li and B. Lü. Global well-posedness and exponential stability of 3D Navier-Stokes equations with density-dependent viscosity and vacuum in unbounded domains. Arch. Ration. Mech. Anal. 239 (2021), no. 3, 1809–1835. 1, 1
- [17] D.D. Joseph. Fluid dynamics of viscoelastic liquids. Springer-Verlag, New York, 1990. 1
- [18] O.A. Ladyzenskaya. On certain nonlinear problems of the theory of continuous media. In Internat. Congress of Mathematicians at Moscow, Abstracts of Reports, 149, 1966.
- [19] O.A. Ladyzenskaya and V.A. Solonnikov. Unique solvability of an initial-and boundary-value problem for viscous incompressible nonhomogeneous fluids. Translated from the Russian in J. Soviet Math. 9 (1978), 697–749. 1, 1, (1), 7
- [20] B. Levant, F. Ramos, and E.S. Titi. On the statistical properties of the 3D incompressible Navier-Stokes-Voigt model. Commun. Math. Sci. 8 (2010), no. 1, pp. 277–293. 1
- [21] R. Lewandowski, L.C. Berselli. On the Bardina's model in the whole space. J. Math. Fluid Mech. 20 (2018), no. 3, pp. 1335–1351.
- [22] R. Lewandowski, W. Layton, On a well-posed turbulence model. Discrete Contin. Dyn. Syst. Ser. B 6 (2006), no. 1, pp. 111–128. 1
- [23] J. Li. Local existence and uniqueness of strong solutions to the Navier–Stokes equations with nonnegative density. J. Differ. Equ. 263 (2017), no. 10, 6512–6536. 1
- [24] P.-L. Lions. Mathematical Topics in Fluid Mechanics, Volume 1: Incompressible Models. Clarendon Press, Oxford, 1996. 1, 6

- [25] B. Lu, X. Shi and X. Zhong. Global existence and large time asymptotic behavior of strong solutions to the Cauchy problem of 2D density-dependent Navier-Stokes equations with vacuum. *Nonlinearity* **31** (2018), no. 6, 2617–2632. 1, 1, 1
- [26] V. Maz'ya. Sobolev Spaces with Applications to Elliptic Partial Differential Equations. Springer, Heidelberg, 2011. 1, 2, 2
- [27] A.P. Oskolkov. Nonlocal problems for equations of Kelvin-Voigt fluids and their ϵ -approximations. J. Math. Sci. 87 (1997), 3393–3408. 1
- [28] M. Paicu, P. Zhang and Z. Zhang. Global unique solvability of inhomogeneous Navier-Stokes equations with bounded density. Comm. Partial Differential Equations 38 (2013), no. 7, 1208–1234.
- [29] K. Pileckas. On spaces of solenoidal vectors (Russian). Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 96 (1980), 237–239. 2
- [30] F. Ramos, E.S. Titi, Invariant measures for the 3D Navier-Stokes-Voigt equations and their Navier-Stokes limit. Discrete Contin. Dyn. Syst. 28 (2010), no. 1, pp. 375–403. 1
- [31] J. Simon. Nonhomogeneous viscous incompressible fluids: existence of velocity, density, and pressure. SIAM J. Math. Anal. 21 (1990), no. 5, 1093–1117. 1
- [32] J. Simon. Compact sets in the space $L^{p}(0,T;B)$. Ann. Mat. Pura Appl. 146 (1987), no. 4, 65–96. 2
- [33] J. Zhang, W. Shi and H. Cao. Global unique solvability of inhomogeneous incompressible Navier-Stokes equations with nonnegative density. *Nonlinearity* 35 (2022), no. 9, 4795–4819. 1, 1
- [34] V. G Zvyagin and M. V. Turbin, Investigation of initial-boundary value problems for mathematical models of the motion of Kelvin-Voigt fluids, Translated from the Russian in J. Math. Sci. 168 (2010), no. 2, 157–308.