CONTINUITY ESTIMATES FOR DOUBLY DEGENERATE PARABOLIC EQUATIONS WITH LOWER ORDER TERMS VIA NONLINEAR POTENTIALS

QIFAN LI*

ABSTRACT. This article studies the continuity of bounded nonnegative weak solutions to inhomogeneous doubly nonlinear parabolic equations. A model equation is

$$\partial_t u - \operatorname{div}(u^{m-1}|Du|^{p-2}Du) = f$$
 in $\Omega \times (-T,0) \subset \mathbb{R}^{n+1}$.

Here, we consider the case m > 1 and 2 . We establish a continuity estimate for <math>u in terms of elliptic Riesz potentials of the right-hand side of the equation.

1. Introduction and main results

We are concerned in this paper with the doubly degenerate parabolic equation with lower order terms. More precisely, we consider quasilinear parabolic equations of the form

$$\partial_t u - \operatorname{div} A(x, t, u, Du) = f, \tag{1.1}$$

in a bounded domain $\Omega_T = \Omega \times (-T, 0)$, where $\Omega \subset \mathbb{R}^n$. Here, we assume that $f = f(x) \in L^1_{loc}(\Omega)$ and u is nonnegative. The vector field $A(x, t, u, \xi)$ is measurable in $\Omega_T \times \mathbb{R} \times \mathbb{R}^n$ and satisfies the growth and ellipticity conditions:

$$\begin{cases} \langle A(x,t,u,\xi),\xi\rangle \ge C_0 u^{m-1} |\xi|^p, \\ |A(x,t,u,\xi)| \le C_1 u^{m-1} |\xi|^{p-1} + g(x) u^{\frac{m-1}{p}}, \end{cases}$$
(1.2)

where C_0 , C_1 are given positive constants, $g \ge 0$ and $g = g(x) \in L^{p/(p-1)}_{loc}(\Omega)$. Throughout the paper, we keep m > 1 and 2 . In this case, the quasilinear equation (1.1) with the structure conditions (1.2) is classified as doubly degenerate. Doubly degenerate parabolic equations play an important role in the study of the non-stationary, polytropic flow of a fluid in a porous medium (see, for example [1]). The Hölder continuity of nonnegative weak solutions to (1.1)-(1.2) was proved in [9, 13]. In [4] the authors proved the local Hölder continuity of sign-changing solutions to doubly nonlinear parabolic equations in the range <math>p > 2 and m + p > 2. Furthermore, Harnack inequality for doubly degenerate parabolic equations was first established by Vespri [17]. This result was extended in [8] to the general quasilinear structure. Recently, Bögelein et al. [5] proved the Harnack inequality in the case m > 0, p > 1 and m(p - 1) > 1 which is the full range in slow diffusion case.

The aim of this paper is to prove continuity of weak solutions to (1.1) in terms of non-linear potentials of the functions f and g. The continuity problem related to the potentials can be traced back to the study of the Wiener criteria of p-Laplace equations (see [10]). In [10] the authors proved the local boundedness of weak solutions to p-Laplace equations

1

²⁰¹⁰ Mathematics Subject Classification. 35K10, 35K59, 35K65, 35K92.

Key words and phrases. Doubly degenerate parabolic equation, Quasilinear parabolic equation, Regularity theory, Potential theory.

with measure data. Furthermore, Liskevich and Skrypnik [11, 14] studied evolutionary *p*-Laplace equations with lower order terms from nonlinear Kato-type classes and established qualitative properties such as local boundedness, continuity and Harnack estimate for weak solutions. Subsequently, Liskevich, Skrypnik and Sobol [12] extended the local boundedness result via a time-dependent parabolic potential. Similar results for porous medium equations have been obtained in [2, 3, 18, 19]. However, limited work has been done for the doubly nonlinear equations. In [16] the author obtained a local boundedness result for the doubly degenerate parabolic equations in terms of the parabolic nonlinear potentials. Motivated by this work, we are interested in the continuity estimate for doubly degenerate parabolic equations with lower order terms from nonlinear Kato-type classes. Before formulating the main result, let us introduce the following quantities

$$F_1(R) = \sup_{x \in \Omega} \int_0^R \left(\frac{1}{r^{n-p}} \int_{B_r(x)} g(y)^{\frac{p}{p-1}} \, \mathrm{d}y \right)^{\frac{1}{p}} \frac{1}{r} \, \mathrm{d}r \tag{1.3}$$

and

$$F_2(R) = \sup_{x \in \Omega} \int_0^R \left(\frac{1}{r^{n-p}} \int_{B_r(x)} f(y) \, \mathrm{d}y \right)^{\frac{1}{p-1}} \frac{1}{r} \, \mathrm{d}r, \tag{1.4}$$

where R > 0 and $B_r(x) = \{y \in \Omega : |y - x| \le r\}$. These quantities are also termed as elliptic Riesz potentials (see [18]). Next, we give the definition of weak solutions to doubly nonlinear parabolic equations.

Definition 1.1. A nonnegative measurable function $u: \Omega_T \to \mathbb{R}$ is said to be a local weak solution to (1.1)-(1.2) if

$$u \in C_{\text{loc}}(-T, 0; L^2_{\text{loc}}(\Omega)), \quad u^{\alpha} \in L^p_{\text{loc}}(-T, 0; W^{1,p}_{\text{loc}}(\Omega)), \quad where \quad \alpha = \frac{m+p-2}{p-1}$$

and for every open set $U \subset \Omega$ and every subinterval $(t_1, t_2) \subset (-T, 0)$ the identity

$$\int_{U} u(\cdot, t)\varphi(\cdot, t)dx\Big|_{t=t_{1}}^{t_{2}} + \iint_{U\times(t_{1}, t_{2})} -u\partial_{t}\varphi + \langle A(x, t, u, Du), D\varphi \rangle dxdt$$

$$= \iint_{U\times(t_{1}, t_{2})} f\varphi dxdt$$
(1.5)

 $holds for \ all \ any function \ \varphi \in W^{1,2}_{\mathrm{loc}}(-T,0;L^2(U)) \cap L^p_{\mathrm{loc}}(-T,0;W^{1,p}_0(U)).$

We note that the gradient of u in the structure conditions (1.2) is interpreted as $Du = \alpha^{-1}u^{1-\alpha}\chi_{\{u>0\}}Du^{\alpha}$. We are now in a position to state our main theorem.

Theorem 1.2. Let u be a nonnegative, locally bounded, weak solution to the doubly degenerate parabolic equation (1.1) in the sense of Definition 1.1, where the vector field A fulfills the structure conditions (1.2). Furthermore, assume that

$$\lim_{R \downarrow 0} [F_1(R) + F_2(R)] = 0 \tag{1.6}$$

holds true. Then, u is locally continuous in Ω_T .

No attempt has been made here to study the regularity problem for the time-dependent parabolic potential. Our proof relies on the method of intrinsic scaling and follow the idea from [6, chapter III]. In the context of the potential estimate, it seems that neither De Giorgi nor Moser iteration technique work in the proof of De Giorgi type lemmas. Our proof is based on the Kilpeläinen-Malý technique [10, 11], but the arguments become more involved when we move from the *p*-Laplace to the doubly nonlinear setting. Contrary to [3], which established a continuity result for the porous medium equations with nonnegative

measure data, our result could handle the case when the right-hand side of the equation f is a sign-changing function.

The rest of the paper is organized as follows. In Sect. 2, we derive the Caccioppoli inequalities and set up an alternative argument. Subsequently Sect. 3 is devoted to the analysis of the first alternative. We obtain a decay estimate for the oscillation of the weak solution in terms of $F_1(R)$ and $F_2(R)$. In Sect. 4 we proceed with the study of the second alternative and obtain a similar estimate for the oscillation of the weak solutions via nonlinear potentials. Finally the proof of the main theorem is presented in Sect. 5.

2. Preliminary material and Caccioppoli inequalities

In this section, we provide some preliminary lemmas. To start with, we follow the notation used in [12] and introduce some auxiliary functions. Throughout the paper, we define

$$G(u) = \begin{cases} u, & \text{for } u > 1, \\ u^2, & \text{for } 0 \le u \le 1 \\ 0, & \text{for } u < 0. \end{cases}$$
 (2.1)

Let $s \in \mathbb{R}$ and $s_+ = \max\{s, 0\}$. We observe that $G(s) = \min\{s_+, s_+^2\}$. Fix $0 < \lambda < p - 1$, we introduce the function

$$\phi_{+}(s) = \int_{0}^{s_{+}} (1+\tau)^{-1-\lambda} d\tau = \frac{1}{\lambda} \left(1 - (1+s_{+})^{-\lambda} \right).$$

It can be easily seen that there exist γ_1 and γ_2 depending only upon λ such that $\gamma_1 < \gamma_2$,

$$\gamma_1 \min\{s_+, 1\} \le \phi_+(s) \le \gamma_2 \min\{s_+, 1\} \quad \text{and} \quad \gamma_1 \frac{s_+}{s_+ + 1} \le \phi_+(s) \le \gamma_2 \frac{s_+}{s_+ + 1}. \quad (2.2)$$

In this paper, we fix the parameter λ such that $\lambda < \min\{(p-1)^{-1}, n^{-1}\}$. The statement that a constant γ depends only upon the data means that it can be determined a priori only in terms of $\{n, m, p, C_0, C_1, \lambda\}$. Next, we set

$$\Phi(s) = \int_0^{s_+} \phi_+(\tau) \, d\tau$$

and it follows from (2.2) that

$$\gamma_1 G(s) = \gamma_1 \min\{s_+, s_+^2\} \le \Phi(s) \le \gamma_2 \min\{s_+, s_+^2\} = \gamma_2 G(s). \tag{2.3}$$

Furthermore, we introduce the function

$$\psi(s) = \int_0^{s_+} (1+\tau)^{-\frac{1}{p} - \frac{\lambda}{p}} d\tau = \frac{p}{p-1-\lambda} \left[(1+s_+)^{1-\frac{1+\lambda}{p}} - 1 \right].$$

Let d > 0, l > 0 and u be the weak solution as in the Definition 1.1. We define the quantities

$$\psi_{-} = \left(\frac{1}{d} \int_{u}^{l} \left(1 + \frac{l-s}{d}\right)^{-\frac{1}{p} - \frac{d}{p}} ds\right)_{+} = \psi\left(\frac{l-u}{d}\right)$$

$$\tag{2.4}$$

and

$$\psi_{+} = \left(\frac{1}{d} \int_{l}^{u} \left(1 + \frac{s - l}{d}\right)^{-\frac{1}{p} - \frac{\lambda}{p}} ds\right) = \psi\left(\frac{u - l}{d}\right). \tag{2.5}$$

At this point, we state the following two lemmas which give a characterization of the quantities ψ_- and ψ_+ .

Lemma 2.1. Let ψ_{-} be the quantity defined in (2.4). Then, we have

$$c_1(\varepsilon_1) \left(\frac{l-u}{d} \right)^{\frac{p-1-\lambda}{p}} \le \psi_- \le c_2(\varepsilon_1) \left(\frac{l-u}{d} \right)^{\frac{p-1-\lambda}{p}} \qquad if \qquad \frac{l-u}{d} \ge \varepsilon_1. \tag{2.6}$$

Lemma 2.2. Let ψ_+ be the quantity defined in (2.5). Then, we have

$$c_1(\varepsilon_1) \left(\frac{u-l}{d}\right)^{\frac{p-1-\lambda}{p}} \le \psi_+ \le c_2(\varepsilon_1) \left(\frac{u-l}{d}\right)^{\frac{p-1-\lambda}{p}} \qquad if \qquad \frac{u-l}{d} \ge \varepsilon_1 \tag{2.7}$$

The proofs of these two lemmas are not particularly difficult but will not be reproduced here. Let $\varrho > 0$, $\theta > 0$ and $(x_0, t_0) \in \Omega_T$ be a fixed point. We set $Q_{\varrho,\theta}(z_0) = B_\varrho(x_0) \times (t_0 - \theta, t_0) \subset \Omega_T$ and denote by φ a piecewise smooth function in $Q_{\varrho,\theta}(z_0)$ such that

$$0 \le \varphi \le 1$$
, $|D\varphi| < \infty$ and $\varphi = 0$ on $\partial B_{\rho}(x_0)$, (2.8)

where $\partial B_{\varrho}(x_0) = \{y : |y - x_0| = \varrho\}$. If x_0 is the origin, then we omit in our notation the point x_0 and write B_{ϱ} for $B_{\varrho}(x_0)$. Finally, let

$$\partial_P Q_{\rho,\theta}(z_0) = [\partial B_{\rho}(x_0) \times (t_0 - \theta, t_0)] \cup [B_{\rho}(x_0) \times \{t_0 - \theta\}]$$

denotes the parabolic boundary of $Q_{\varrho,\theta}(z_0)$. The crucial result in our development of regularity property of weak solutions will be the following Caccioppoli inequality.

Lemma 2.3. Let d > 0, l > 0 and let u be a nonnegative weak solution to the doubly degenerate parabolic equation (1.1) in the sense of Definition 1.1. There exists a constant γ depending only upon the data, such that for every piecewise smooth cutoff function φ satisfying (2.8), we have

$$\operatorname{ess\,sup}_{t_{0} < t < t_{1}} \int_{L^{-}(t)} G\left(\frac{l-u}{d}\right) \varphi^{k} \, \mathrm{d}x + d^{p-2} \iint_{L^{-}} u^{m-1} |D\psi_{-}|^{p} \varphi^{k} \, \mathrm{d}x \mathrm{d}t$$

$$\leq \gamma \int_{L^{-}(t_{0}-\theta)} G\left(\frac{l-u}{d}\right) \varphi^{k} \, \mathrm{d}x + \gamma \iint_{L^{-}} \frac{l-u}{d} |\partial_{t}\varphi| \, \varphi^{k-1} \, \mathrm{d}x \mathrm{d}t$$

$$+ \gamma d^{p-2} \iint_{L^{-}} u^{m-1} \left(\frac{l-u}{d}\right)^{(1+\lambda)(p-1)} \varphi^{k-p} |D\varphi|^{p} \, \mathrm{d}x \mathrm{d}t$$

$$+ \gamma \frac{\theta}{d^{2}} \int_{B_{\varrho}(x_{0})} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{\theta}{d} \int_{B_{\varrho}(x_{0})} |f| \, \mathrm{d}x,$$

$$(2.9)$$

where k > p, $L^- = Q_{\rho,\theta}(z_0) \cap \{u \le l\}$ and $L^-(t) = B_{\rho}(x_0) \cap \{u(\cdot,t) \le l\}$.

Proof. To simplify the notation, we set $B = B_{\varrho}(x_0)$ and $Q = Q_{\varrho,\theta}(z_0)$. In the weak formulation (1.5) we choose the testing function

$$\varphi_{-} = \left[\int_{u}^{l} \left(1 + \frac{l - s}{d} \right)^{-1 - \lambda} ds \right] \varphi^{k} = d\phi_{+} \left(\frac{l - u}{d} \right) \varphi^{k}.$$

The use of φ_{-} as a testing function is justified, modulus a time mollification technique (see [16, Lemma 2.7]). This gives

$$\iint_{Q} \varphi_{-} \partial_{t} u \, dx dt + \iint_{Q} \langle A(x, t, u, Du), D\varphi_{-} \rangle \, dx dt = \iint_{Q} f \varphi_{-} \, dx dt.$$

We now proceed formally for the term involving the time derivative. From (2.3), we see that for any $t \in (t_0 - \theta, t_0)$

$$\begin{split} &\int_{t_0-\theta}^t \int_B \varphi_- \partial_t u \, \mathrm{d}x \mathrm{d}t \\ &= -\int_{t_0-\theta}^t \int_B \frac{\partial}{\partial t} \left[\left(\int_u^l \, \mathrm{d}w \, \int_w^l \left(1 + \frac{l-s}{d} \right)^{-1-\lambda} \, \mathrm{d}s \right)_+ \right] \varphi^k \, \mathrm{d}x \mathrm{d}t \\ &\leq -\gamma d^2 \int_{L^-(t)} G\left(\frac{l-u}{d} \right) \varphi^k \, \mathrm{d}x \\ &+ \gamma d^2 \iint_{L^-} \frac{l-u}{d} \varphi^{k-1} \, |\partial_t \varphi| \, \, \mathrm{d}x \mathrm{d}t + \gamma d^2 \int_{L^-(t_0-\theta)} G\left(\frac{l-u}{d} \right) \varphi^k \, \mathrm{d}x. \end{split}$$

Next, we consider the second term on the left-hand side. To this end, we decompose

$$\iint_{Q} \langle A(x, t, u, Du), D\varphi_{-} \rangle \, dxdt$$

$$= k \iint_{Q} A(x, t, u, Du) \varphi^{k-1} D\varphi \left(\int_{u}^{l} \left(1 + \frac{l-s}{d} \right)^{-1-\lambda} \, ds \right)_{+} \, dxdt$$

$$- \iint_{L^{-}} A(x, t, u, Du) \cdot Du \left(1 + \frac{l-u}{d} \right)^{-1-\lambda} \varphi^{k} \, dxdt =: T_{1} + T_{2}.$$

We first consider the estimate for T_2 . From the first structure condition (1.2), we deduce that

$$T_{2} \leq -C_{0} \iint_{L^{-}} u^{m-1} |Du|^{p} \left(1 + \frac{l-u}{d}\right)^{-1-\lambda} \varphi^{k} \, dx dt$$
$$= -C_{0} d^{p} \iint_{L^{-}} u^{m-1} |D\psi_{-}|^{p} \varphi^{k} \, dx dt.$$

To estimate T_1 , we use the second structure condition (1.2) to deduce that

$$T_{1} \leq \iint_{Q} (|Du|^{p-1}u^{m-1} + u^{\frac{m-1}{p}}g)\varphi^{k-1}|D\varphi| \left(\int_{u}^{l} \left(1 + \frac{l-s}{d}\right)^{-1-\lambda} ds \right)_{+} dxdt$$

=: $T_{3} + T_{4}$,

with the obvious meanings of T_3 and T_4 . Next, we consider the estimate for T_3 . We use (2.2) and Young's inequality to conclude that for any fixed $\varepsilon > 0$ there holds

$$T_{3} \leq \gamma d \iint_{L^{-}} |Du|^{p-1} u^{m-1} \varphi^{k-1} |D\varphi| \left(\frac{l-u}{d}\right) \left(1 + \frac{l-u}{d}\right)^{-1} dxdt$$

$$\leq \varepsilon d^{p} \iint_{L^{-}} u^{m-1} |D\psi_{-}|^{p} \varphi^{k} dxdt$$

$$+ \gamma d^{p} \iint_{L^{-}} \varphi^{k-p} |D\varphi|^{p} u^{m-1} \left(1 + \frac{l-u}{d}\right)^{\lambda(p-1)-1} \left(\frac{l-u}{d}\right)^{p} dxdt$$

$$\leq \varepsilon d^{p} \iint_{L^{-}} u^{m-1} |D\psi_{-}|^{p} \varphi^{k} dxdt$$

$$+ \gamma(\lambda, \varepsilon) d^{p} \iint_{L^{-}} \varphi^{k-p} |D\varphi|^{p} u^{m-1} \left(\frac{l-u}{d}\right)^{(1+\lambda)(p-1)} dxdt,$$

since $\lambda < (p-1)^{-1}$. To estimate T_4 , we apply (2.2), $\lambda < (p-1)^{-1}$ and Young's inequality to conclude that

$$T_{4} \leq \gamma d \iint_{L^{-}} u^{\frac{m-1}{p}} g \varphi^{k-1} |D\varphi| \left(\frac{l-u}{d}\right) \left(1 + \frac{l-u}{d}\right)^{-1} dx dt$$

$$= \gamma \iint_{L^{-}} \left[du^{\frac{m-1}{p}} \varphi^{\frac{k}{p}-1} \left(\frac{l-u}{d}\right) \left(1 + \frac{l-u}{d}\right)^{-\frac{1}{p} + \frac{\lambda}{p'}} |D\varphi| \right]$$

$$\times \left[\left(1 + \frac{l-u}{d}\right)^{-\frac{1}{p'} - \frac{\lambda}{p'}} g \varphi^{\frac{k}{p'}} \right] dx dt$$

$$\leq \gamma d^{p} \iint_{L^{-}} \varphi^{k-p} |D\varphi|^{p} u^{m-1} \left(\frac{l-u}{d}\right)^{(1+\lambda)(p-1)} dx dt$$

$$+ \gamma \theta \iint_{B} g^{\frac{p}{p-1}} dx,$$

where the constant γ depends only upon the data. Taking into account that $\varphi_- \leq \lambda^{-1}d$, we obtain an estimate for the right-hand side of the equation by

$$\iint_{Q} f\varphi_{-} \, \mathrm{d}x \mathrm{d}t \le \gamma d\theta \int_{B} |f| \, \mathrm{d}x.$$

Combining the above estimates and dividing by d^2 , we obtain the desired estimate (2.9). We have thus proved the lemma.

Furthermore, by substituting

$$\varphi_{+} = \left[\int_{l}^{u} \left(1 + \frac{s - l}{d} \right)^{-1 - \lambda} ds \right]_{\perp} \varphi^{k} = d\phi_{+} \left(\frac{u - l}{d} \right) \varphi^{k}$$

into the weak formulation (1.5), we get the following lemma, the proof of which we omit.

Lemma 2.4. Let a > 0, d > 0, l > 0 and let u be a nonnegative weak solution to the doubly degenerate parabolic equation (1.1) in the sense of Definition 1.1. Let $\varrho > 0$ be such that $(t_0 - a^{1-m}d^{2-p}\varrho^p, t_0) \subset (-T,0)$ and $B_\varrho(x_0) \subset \Omega$. There exists a positive constant γ depending only upon the data, such that for every piecewise smooth cutoff function φ vanishing on $\partial B_\varrho(x_0) \times (t_0 - a^{1-m}d^{2-p}\varrho^p, t_0)$, there holds

$$\sup_{t_{0}-a^{1-m}d^{2-p}\varrho^{p} < t < t_{0}} \int_{L^{+}(t)} G\left(\frac{u-l}{d}\right) \varphi^{k} \, dx + d^{p-2} \iint_{L^{+}} u^{m-1} |D\psi_{+}|^{p} \varphi^{k} \, dx dt \\
\leq \gamma \int_{L^{+}(t_{0})} G\left(\frac{u-l}{d}\right) \varphi^{k} \, dx + \gamma \iint_{L^{+}} \frac{u-l}{d} |\partial_{t}\varphi| \, dx dt \\
+ \gamma d^{p-2} \iint_{L^{+}} u^{m-1} \left(\frac{u-l}{d}\right)^{(1+\lambda)(p-1)} \varphi^{k-p} |D\varphi|^{p} \, dx dt \\
+ \gamma \frac{\varrho^{p}}{d^{p}} a^{1-m} \int_{B_{\varrho}(x_{0})} g^{\frac{p}{p-1}} \, dx + \gamma \frac{\varrho^{p}}{d^{p-1}} a^{1-m} \int_{B_{\varrho}(x_{0})} |f| \, dx, \tag{2.10}$$

where k > p, $L^+ = [B_{\rho}(x_0) \times (t_0 - a^{1-m}d^{2-p}\varrho^p, t_0)] \cap \{u \ge l\}$ and $L^+(t) = B_{\rho}(x_0) \cap \{u(\cdot, t) \ge l\}$.

We now turn to consider the logarithmic estimates for nonnegative weak solutions. To this aim, we introduce the logarithmic function

$$\psi^{\pm}(u) = \ln^{+} \left(\frac{H_{k}^{\pm}}{H_{k}^{\pm} - (u - k)_{\pm} + c} \right), \qquad 0 < c < H_{k}^{\pm}, \tag{2.11}$$

where H_k^{\pm} is a constant chosen such that $H_k^{\pm} \ge \operatorname{ess\,sup}_{Q_{\varrho,\theta}}(u-k)_{\pm}$. We are now in a position to state the following lemma.

Lemma 2.5. Let u be a nonnegative weak solution to the doubly degenerate parabolic equation (1.1) in the sense of Definition 1.1. There exists a constant γ that can be determined a priori only in terms of the data such that

$$\operatorname{ess \,sup}_{t_{0}-\theta < t < t_{0}} \int_{B_{\varrho}(x_{0}) \times \{t\}} [\psi^{\pm}(u)]^{2} \zeta^{p} \, \mathrm{d}x \\
\leq \int_{B_{\varrho}(x_{0}) \times \{t_{0}-\theta\}} [\psi^{\pm}(u)]^{2} \zeta^{p} \, \mathrm{d}x + \gamma \iint_{Q_{\varrho,\theta}(z_{0})} u^{m-1} \psi^{\pm}(u) |D\zeta|^{p} \left[(\psi^{\pm})'(u) \right]^{2-p} \, \mathrm{d}x \mathrm{d}t \\
+ \gamma \ln \left(\frac{H}{c} \right) \left(\frac{1}{c} \iint_{Q_{\alpha,\theta}(z_{0})} |f| \, \mathrm{d}x \mathrm{d}t + \frac{1}{c^{2}} \iint_{Q_{\alpha,\theta}(z_{0})} g^{\frac{p}{p-1}} \, \mathrm{d}x \mathrm{d}t \right), \tag{2.12}$$

where $\zeta = \zeta(x)$ is independent of t and satisfies (2.8).

Proof. The proof of (2.12) follows in a similar manner as the arguments in [13, Proposition 2.1] and we sketch the proof. For simplicity of notation, we write ψ instead of $\psi^{\pm}(u)$ and set $B = B_{\varrho}(x_0)$. In the weak formulation (1.5) we use the testing function $\varphi = [\psi^2(u)]'\zeta^p$, where $\zeta = \zeta(x) \in C_0^{\infty}(B_{\varrho})$. The following formal computations can be made rigorous with a standard smoothing procedure with respect to time. Then, we find

$$\int_{t_0-\theta}^t \int_B \partial_t u [\psi^2(u)]' \zeta^p \, \mathrm{d}x \mathrm{d}t = \int_{B \times \{t\}} \psi^2(u) \zeta^p \, \mathrm{d}x - \int_{B \times \{t_0-\theta\}} \psi^2(u) \zeta^p \, \mathrm{d}x.$$

Next, we consider the term involving A(x, t, u, Du). To this end, we decompose

$$\begin{split} &\int_{t_0-\theta}^t \int_B \left\langle A(x,t,u,Du), D\varphi \right\rangle \mathrm{d}x \mathrm{d}t \\ &= \int_{t_0-\theta}^t \int_B A(x,t,u,Du) [\psi^2(u)]' p \zeta^{p-1} \cdot D\zeta \, \mathrm{d}x \mathrm{d}t \\ &+ \int_{t_0-\theta}^t \int_B A(x,t,u,Du) \cdot Du [\psi^2(u)]'' \zeta^p \, \mathrm{d}x \mathrm{d}t =: T_1 + T_2. \end{split}$$

We first estimate the second term T_2 . Noting that $[\psi^2(u)]'' = 2(1 + \psi)[\psi'(u)]^2$, we infer from (1.2) that

$$T_2 \ge 2C_0 \int_{t_0-\theta}^t \int_B u^{m-1} |Du|^p (1+\psi) [\psi'(u)]^2 \zeta^p \, \mathrm{d}x \mathrm{d}t.$$

To estimate T_1 , we use (1.2) to deduce that

$$T_1 \le 2C_1 \int_{t_0 - \theta}^t \int_B (u^{m-1} |Du|^{p-1} + gu^{\frac{m-1}{p}}) \psi \psi' \zeta^{p-1} |D\zeta| \, \mathrm{d}x \mathrm{d}t$$

=: $T_3 + T_4$,

with the obvious meanings of T_3 and T_4 . By Young's inequality, we conclude that

$$T_{3} \leq \varepsilon \int_{t_{0}-\theta}^{t} \int_{B} u^{m-1} |Du|^{p} \zeta^{p} [\psi'(u)]^{2} \psi \, \mathrm{d}x \mathrm{d}t$$
$$+ c(\varepsilon) \int_{t_{0}-\theta}^{t} \int_{B} u^{m-1} \psi |D\zeta|^{p} [\psi'(u)]^{2-p} \, \mathrm{d}x \mathrm{d}t,$$

since $p - \frac{2p}{p'} = 2 - p$. Next, we turn our attention to the estimate of T_4 . By Young's inequality, we see that

$$\begin{split} T_4 = & 2C_1 \int_{t_0 - \theta}^t \int_B \psi \left[\zeta^{p-1} \psi'(u)^{\frac{2}{p'}} g \right] \left[|D\zeta| \psi'(u)^{1 - \frac{2}{p'}} u^{\frac{m-1}{p}} \right] \mathrm{d}x \mathrm{d}t \\ \leq & \gamma \int_{t_0 - \theta}^t \int_B g^{p'} \zeta^p [\psi'(u)]^2 \psi \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \int_{t_0 - \theta}^t \int_B u^{m-1} \psi |D\zeta|^p [\psi'(u)]^{2-p} \, \mathrm{d}x \mathrm{d}t. \end{split}$$

Taking into account that $\psi \leq \ln(H/c)$ and $\psi' \leq 1/c$, we conclude that

$$\int_{t_0-\theta}^t \int_B g^{p'} \zeta^p [\psi'(u)]^2 \psi \, \mathrm{d}x \mathrm{d}t \le \ln\left(\frac{H}{c}\right) \frac{1}{c^2} \iint_{Q_{o,\theta}(z_0)} g^{\frac{p}{p-1}} \, \mathrm{d}x \mathrm{d}t$$

and

$$\int_{t_0-\theta}^t \int_B f\varphi \, \mathrm{d}x \mathrm{d}t \le \ln\left(\frac{H}{c}\right) \frac{1}{c} \iint_{O_0\theta(z_0)} |f| \, \mathrm{d}x \mathrm{d}t.$$

Collecting above estimates and taking the supremum over $t \in (t_0 - \theta, t_0)$, we obtain the desired logarithmic estimate (2.12).

Next, we state in Lemma 2.6 a Caccioppoli estimate, which will be used in the proof of Lemma 4.4.

Lemma 2.6. Let u be a nonnegative weak solution to the doubly degenerate parabolic equation (1.1) in the sense of Definition 1.1. There exists a positive constant γ depending only upon the data, such that for every piecewise smooth cutoff function φ and satisfying (2.8), there holds

$$\operatorname{ess \, sup}_{t_{0}-\theta < t < t_{0}} \int_{B_{\varrho}(x_{0}) \times \{t\}} (u - k)_{\pm}^{2} \varphi^{p} \, dx + \iint_{Q_{\varrho,\theta}(z_{0})} u^{m-1} |D(u - k)_{\pm} \varphi|^{p} \, dx dt \\
\leq \int_{B_{\varrho}(x_{0}) \times \{t_{0}\}} (u - k)_{\pm}^{2} \varphi^{p} \, dx + \gamma \iint_{Q_{\varrho,\theta}(z_{0})} u^{m-1} (u - k)_{\pm}^{p} |D\varphi|^{p} \, dx dt \\
+ \gamma \iint_{Q_{\varrho,\theta}(z_{0})} (u - k)_{\pm}^{2} |\partial_{t} \varphi| \, dx dt + \gamma \iint_{Q_{\varrho,\theta}(z_{0})} |f|(u - k)_{\pm} \, dx dt \\
+ \gamma \iint_{Q_{\varrho,\theta}(z_{0})} g^{\frac{p}{p-1}} \, dx dt. \tag{2.13}$$

This is a standard result that can be proved by choosing the testing function $\pm (u - k)_{\pm} \varphi^p$ into (1.5) and the proof will be omitted.

We now turn our attention to the proof of Theorem 1.2. The continuity of the weak solution u at a point \mathfrak{z}_0 will be a consequence of the following assertion. There exists a family of nested and shrinking cylinders $Q_{\varrho_n,\theta_n}(\mathfrak{z}_0)$, with vertex at \mathfrak{z}_0 , such that the essential oscillation of u in $Q_{\varrho_n,\theta_n}(\mathfrak{z}_0)$ converges to zero as the cylinders shrink to the point \mathfrak{z}_0 .

There is no loss of generality in assuming $\mathfrak{z}_0 = (0,0)$. Let $\varepsilon_0 > 0$ and $\hat{R} > 0$ be such that $Q_{\hat{R},\hat{R}^{p-\varepsilon_0}}(0) \subset \Omega_T$. For any $0 < R < \hat{R}$, we fix a cylinder $Q_R = B_R \times (-R^{p-\varepsilon_0},0)$ and set

$$\mu_+ = \operatorname{ess\,sup} u, \qquad \mu_- = \operatorname{ess\,inf} u \qquad \text{and} \qquad \operatorname{ess\,osc} u = \mu_+ - \mu_-.$$

Henceforth, let $\omega > 0$ be a parameter such that $\frac{1}{2}\omega \le \mu_+ - \mu_- \le \omega$. Here we assume that $\mu_+ > 0$, since otherwise, if $\mu_+ = 0$, then $\omega = 0$ and Theorem 1.2 holds trivially. Let A > 1

be a constant which will be determined later. We introduce the enlarged cylinder

$$\hat{Q} = B_R \times \left(-(\mu_+)^{1-m} \left(\frac{\omega}{A} \right)^{2-p} R^p, 0 \right). \tag{2.14}$$

In the case that \hat{Q} is not contained in $\widehat{Q}_R = B_R \times (-\frac{1}{2}R^{p-\varepsilon_0}, 0)$ for all $0 < R < \hat{R}$, we conclude that

$$-(\mu_{+})^{1-m} \left(\frac{\omega}{A}\right)^{2-p} \le -\frac{1}{2} R^{-\varepsilon_0}. \tag{2.15}$$

Since u is nonnegative, we have $\omega \le 2\mu_+$ and the inequality (2.15) implies that

$$\operatorname*{ess \ osc}_{O_{P}} u \leq \omega \leq 2^{\frac{m}{m+p-3}} A^{\frac{p-2}{m+p-3}} R^{\frac{\varepsilon_{0}}{m+p-3}} \tag{2.16}$$

holds for all $0 < R < \hat{R}$. In this case the weak solution is Hölder continuous and Theorem 1.2 holds immediately. Next, we concentrate on the case $\hat{Q} \subset \widehat{Q}_R$ for a fixed $R \in (0, \hat{R})$. For any fixed time level $-(A^{p-2}-1)(\mu_+)^{1-m}\omega^{2-p}R^p \le t \le 0$ we introduce the intrinsic cylinder of the type

$$Q_r^-(t) = B_r \times (t - (\mu_+)^{1-m} \omega^{2-p} r^p, t),$$

where $0 < r \le R$. Motivated by the work of DiBenedetto [6, chapter III], we consider two complementary cases. For a fixed constant $v_0 > 0$, we see that either

• The first alternative. There exists $-(A^{p-2}-1)(\mu_+)^{1-m}\omega^{2-p}R^p \le \bar{t} \le 0$ such that

$$\left| \left\{ (x,t) \in Q_{\frac{3}{4}R}^{-}(\bar{t}) : u < \mu_{-} + \frac{\omega}{4} \right\} \right| \le \nu_{0} |Q_{\frac{3}{4}R}^{-}(\bar{t})| \tag{2.17}$$

or this does not hold. Then, we have

• The second alternative. For any $-(A^{p-2}-1)(\mu_+)^{1-m}\omega^{2-p}R^p \le \bar{t} \le 0$, there holds

$$\left| \left\{ (x,t) \in Q_{\frac{3}{4}R}^{-}(\bar{t}) : u > \mu_{+} - \frac{\omega}{4} \right\} \right| \le (1 - \nu_{0}) |Q_{\frac{3}{4}R}^{-}(\bar{t})|. \tag{2.18}$$

The constant $v_0 > 0$ will be determined in the course of the proof of Lemma 3.5 in Sect.3, while the value of A will be fixed during the proof of Proposition 4.5 in Sect. 4.

3. The first alternative

The aim of this section is to establish a decay estimate of the essential oscillation for the first alternative. A key ingredient in the proof of the decay estimate is a De Giorgi type lemma. Before we prove this result, we first recall the following definition of the Lebesgue point.

Definition 3.1. [7] Let f be a locally integrable function in Ω_T . A point (x,t) for which

$$\lim_{r \to 0} \frac{1}{2r^p |B_r|} \int_{t-r^p}^{t+r^p} \int_{B_r(x)} |f(y,s) - f(x,t)| \, \mathrm{d}y \, \mathrm{d}s = 0$$

holds is called a Lebesgue point of f.

We now state and prove a couple of lemmas. The first lemma can be deduced from [15, section 1.3] and [7, Theorem 1.33].

Lemma 3.2. If $f \in L_{loc}(\Omega_T)$, then almost every point in Ω_T is a Lebesgue point of f.

In applications, the cylinders used here are not centered on the Lebesgue point, that is in fact the vertex. Moreover, we need the following lemma, which states that the Lebesgue point for f is also a Lebesgue point for f_+ .

Lemma 3.3. Let $f \in L_{loc}(\Omega_T)$ and $z_1 = (x_1, t_1) \in \Omega_T$ be a Lebesgue point for f. Then, we have

$$f^{+}(z_{1}) = \lim_{r \to 0} \frac{1}{|Q_{r,r^{p}}(z_{1})|} \iint_{Q_{r,r^{p}}(z_{1})} f^{+}(y,s) \, \mathrm{d}y \, \mathrm{d}s, \tag{3.1}$$

where $Q_{r,r^p}(z_1) = B_r(x_1) \times (t_1 - r^p, t_1)$.

Proof. We first observe that $|f^+(y,s) - f^+(x,t)| \le |f(y,s) - f(x,t)|$. Consequently, we conclude that

$$\lim_{r \to 0} \frac{1}{|Q_{r,r^{p}}(z_{1})|} \iint_{Q_{r,r^{p}}(z_{1})} |f^{+}(y,s) - f^{+}(x_{1},t_{1})| \, \mathrm{d}y \, \mathrm{d}s$$

$$\leq 2 \lim_{r \to 0} \frac{1}{2r^{p}|B_{r}|} \int_{t-r^{p}}^{t+r^{p}} \int_{B_{r}(x)} |f^{+}(y,s) - f^{+}(x_{1},t_{1})| \, \mathrm{d}y \, \mathrm{d}s = 0,$$

which proves (3.1). This completes the proof.

For technical reasons, we need the following result concerning the scaling property for function G defined in (2.1).

Lemma 3.4. Let $\varepsilon_1 > 0$ and $k \ge 1$ be the fixed constants. If $v \ge \varepsilon_1$, then

$$G(kv) \le c(k, \varepsilon_1)G(v).$$
 (3.2)

Proof. If $v \le 1$ and $kv \le 1$, then $G(kv) = k^2v^2 = k^2G(v)$. If $v \ge 1$ and $kv \ge 1$, then G(kv) = kv = kG(v). Finally, if $\varepsilon_1 \le v \le 1$ and kv > 1, then $G(kv) = \frac{k}{v}v^2 = \frac{k}{v}G(v) \le \frac{k}{\varepsilon_1}G(v)$.

With the help of these lemmas, we can now prove the following De Giorgi type lemma. We adopt the Kilpeläinen-Malý technique from [10, 11]. For the treatment of the doubly nonlinear problem, the Kilpeläinen-Malý technique should be suitably modified to handle the cutoff functions.

Lemma 3.5. Let u be a bounded nonnegative weak solution to (1.1)-(1.2) in Ω_T . There exist constants $v_0 \in (0, 1)$ and B > 1, depending only on the data, such that if

$$\left| \left\{ (x,t) \in Q_{\frac{3}{4}R}^{-}(\bar{t}) : u < \mu_{-} + \frac{\omega}{4} \right\} \right| \le \nu_{0} |Q_{\frac{3}{4}R}^{-}(\bar{t})|, \tag{3.3}$$

then either

$$u(x,t) > \mu_{-} + \frac{\omega}{25}$$
 for a.e. $(x,t) \in Q_{\frac{1}{2}R}^{-}(\bar{t})$ (3.4)

or

$$\omega \le B\left((\mu_+)^{\frac{1-m}{p}}F_1(2R) + (\mu_+)^{\frac{1-m}{p-1}}F_2(2R)\right). \tag{3.5}$$

Proof. Let B > 1 to be determined in the course of the proof. We first assume that (3.5) is violated, that is,

$$\omega > \frac{3}{4}B\left(\frac{1}{3B}\omega + (\mu_{+})^{\frac{1-m}{p}}F_{1}(2R)\right) + (\mu_{+})^{\frac{1-m}{p-1}}F_{2}(2R)\right). \tag{3.6}$$

If we can prove (3.4) then the lemma follows immediately. Fix $(x_1,t_1) \in Q_{\frac{1}{4}R}^-(\bar{t})$ and assume that (x_1,t_1) is a Lebesgue point of u. Let $r_j=4^{-j}C^{-1}R$ and $B_j=B_{r_j}(x_1)$ where C>4 is to be determined. For a sequence $\{l_j\}_{j=0}^\infty$ and a fixed l>0, we set

$$Q_j(l) = B_j \times (t_1 - (\mu_+)^{1-m} (l_j - l)^{2-p} r_i^p, t_1).$$

Moreover, we define $\varphi_j(l) = \phi_j(x)\theta_{j,l}(t)$, where $\phi_j \in C_0^{\infty}(B_j)$, $\phi_j = 1$ on B_{j+1} , $|D\phi_j| \le r_j^{-1}$ and $\theta_{j,l}(t)$ is a Lipschitz function satisfies

$$\theta_{j,l}(t) = 1$$
 in $t \ge t_1 - \frac{4}{9}(\mu_+)^{1-m}(l_j - l)^{2-p}r_j^p$, $\theta_{j,l}(t) = 0$ in $t \le t_1 - \frac{5}{9}(\mu_+)^{1-m}(l_j - l)^{2-p}r_j^p$

and

$$\theta_{j,l}(t) = \frac{t - t_1 - \frac{5}{9}(\mu_+)^{1-m}(l_j - l)^{2-p}r_j^p}{\frac{1}{9}(\mu_+)^{1-m}(l_j - l)^{2-p}r_j^p}$$

in $t_1 - \frac{5}{9}(\mu_+)^{1-m}(l_j - l)^{2-p}r_j^p \le t \le t_1 - \frac{4}{9}(\mu_+)^{1-m}(l_j - l)^{2-p}r_j^p$. From the definition of $\varphi_j(l)$, we see that $\varphi_j(l) = 0$ on $\partial_P Q_j(l)$. Next, for $j = -1, 0, 1, 2, \cdots$, we define the sequence $\{\alpha_j\}$ by

$$\alpha_{j} = \frac{4^{-j-100}}{3B}\omega + \frac{3}{4} \int_{0}^{r_{j}} \left(r^{p-n}(\mu_{+})^{1-m} \int_{B_{r}(x_{1})} g(y)^{\frac{p}{p-1}} dy\right)^{\frac{1}{p}} \frac{dr}{r} + \frac{3}{4} \int_{0}^{r_{j}} \left(r^{p-n}(\mu_{+})^{1-m} \int_{B_{r}(x_{1})} |f(y)| dy\right)^{\frac{1}{p-1}} \frac{dr}{r}.$$
(3.7)

From the definition of α_j , we see that $\alpha_j \to 0$ as $j \to \infty$ and there holds $B\alpha_{j-1} \le \omega$,

$$\alpha_{j-1} - \alpha_{j} \ge \frac{4^{-j-100}}{3B} \omega + \gamma \left(r_{j}^{p-n} (\mu_{+})^{1-m} \int_{B_{j}} g(y)^{\frac{p}{p-1}} \, \mathrm{d}y \right)^{\frac{1}{p}} + \gamma \left(r_{j}^{p-n} (\mu_{+})^{1-m} \int_{B_{j}} |f(y)| \, \mathrm{d}y \right)^{\frac{1}{p-1}}$$

$$(3.8)$$

and

$$\alpha_{j-1} - \alpha_{j} \leq \frac{4^{-j-100}}{3B} \omega + \gamma \left(r_{j-1}^{p-n} (\mu_{+})^{1-m} \int_{B_{j-1}} g(y)^{\frac{p}{p-1}} dy \right)^{\frac{1}{p}} + \gamma \left(r_{j-1}^{p-n} (\mu_{+})^{1-m} \int_{B_{j-1}} |f(y)| dy \right)^{\frac{1}{p-1}}$$

$$(3.9)$$

for all $j = 0, 1, 2, \dots$, where the constant γ depends only upon the data. Moreover, we define a quantity $A_i(l)$ by

$$A_{j}(l) = \frac{(\mu_{+})^{m-1}(l_{j} - l)^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}(l)} \left(\frac{l_{j} - u}{l_{j} - l}\right)^{(1+\lambda)(p-1)} \varphi_{j}(l)^{k-p} \, \mathrm{d}x \mathrm{d}t + \operatorname{ess} \sup_{t} \frac{1}{r_{j}^{n}} \int_{B_{j} \times \{t\}} G\left(\frac{l_{j} - u}{l_{j} - l}\right) \varphi_{j}(l)^{k} \, \mathrm{d}x,$$
(3.10)

where k > p, G is defined in (2.1) and $L_j(l) = Q_j(l) \cap \{u \le l_j\} \cap \Omega_T$. The proof of (3.4) will be divided into several steps.

Step 1: We claim that $A_j(l)$ is continuous in $l < l_j$. To prove this assertion, it suffices to show that the function

$$B_{j}(l) := \operatorname{ess \, sup}_{t} \theta_{j,l}(t)^{k} \frac{1}{r_{j}^{n}} \int_{B_{j}} G\left(\frac{l_{j} - u}{l_{j} - l}\right) \phi_{j}^{k} \, \mathrm{d}x$$

is continuous. For a fixed $l < l_j$, we take $|l' - l| < \delta < \frac{1}{2}(l_j - l)$. Observe that G is Lipschitz continuous, we have

$$\left|G\left(\frac{l_j-u}{l_j-l}\right)-G\left(\frac{l_j-u}{l_j-l'}\right)\right| \leq \gamma(l_j-u)_+\frac{|l-l'|}{(l_j-l)(l_j-l')} \leq \gamma l_j\frac{|l-l'|}{(l_j-l)^2},$$

since $l_j - l' = l_j - l + l - l' > \frac{1}{2}(l_j - l)$. It follows that

$$|B_{j}(l) - B_{j}(l')| \le \gamma \frac{|l - l'|}{(l_{j} - l)^{2}} l_{j} + \operatorname{ess \, sup} \frac{1}{r_{i}^{n}} \int_{B_{j}} G\left(\frac{l_{j} - u}{l_{j} - l}\right) \phi_{j}^{k} \, \mathrm{d}x \left(\operatorname{ess \, sup} |\theta_{j,l}(t)^{k} - \theta_{j,l'}(t)^{k}|\right)$$

and this implies that the function $B_i(l)$ is continuous for $l < l_i$.

Step 2: Determine the values of l_0 and l_1 . Initially, we set $l_0 = \mu_- + \frac{1}{4}\omega$ and $\bar{l} = \frac{1}{2}l_0 + \frac{1}{2}\mu_- + \frac{1}{16}B\alpha_0 + \frac{1}{32}\omega$. Recalling that $B\alpha_0 < \omega$, we deduce

$$l_0 - \bar{l} = \frac{1}{8}\omega - \frac{1}{16}B\alpha_0 - \frac{1}{32}\omega \ge \frac{1}{32}\omega \quad \text{and} \quad l_0 - \bar{l} \le \frac{3}{32}\omega.$$
 (3.11)

Then, we have $(\mu_+)^{1-m}(l_0-\bar{l})^{2-p}r_0^p \leq C^{-p}2^{5(p-2)}(\mu_+)^{1-m}\omega^{2-p}R^p < \frac{1}{100^p}(\mu_+)^{1-m}\omega^{2-p}R^p$, provided that we choose $C=2^{\frac{5(p-2)}{p}+100}$. This guarantees the inclusion of $Q_0(\bar{l})\subset Q_{\frac{3}{4}R}^-(\bar{t})$. In view of $l_0-u\leq \frac{1}{4}\omega$ in $L_0(\bar{l})$, we infer from (3.11) and (3.3) that the inequality

$$\frac{(\mu_{+})^{m-1}(l_{0} - \bar{l})^{p-2}}{r_{0}^{n+p}} \iint_{L_{0}(\bar{l})} \left(\frac{l_{0} - u}{l_{0} - \bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{0}(\bar{l})^{k-p} \, dxdt$$

$$\leq \frac{(\mu_{+})^{m-1}(l_{0} - \bar{l})^{p-2-(1+\lambda)(p-1)}}{r_{0}^{n+p}} \left(\frac{\omega}{4}\right)^{(1+\lambda)(p-1)} |L_{0}(\bar{l})|$$

$$\leq \gamma_{0}C^{n+p} \frac{(\mu_{+})^{m-1}\omega^{p-2}}{R^{n+p}} |L_{0}(\bar{l})| = \gamma_{0}C^{n+p} \frac{|Q_{0}(\bar{l}) \cap \{u \leq l_{0}\}|}{|Q_{\frac{3}{4}R}^{-}(\bar{t})|}$$

$$\leq \gamma_{0}C^{n+p} \frac{|Q_{\frac{3}{4}R}^{-}(\bar{t}) \cap \{u \leq \mu_{-} + \frac{\omega}{4}\}|}{|Q_{\frac{5}{2}R}^{-}(\bar{t})|} \leq \gamma_{0}C^{n+p}\nu_{0}$$
(3.12)

holds for a constant γ_0 depending only upon the data. Moreover, we apply Lemma 2.3 with (l, d, θ) replaced by $(l_0, l_0 - \bar{l}, (\mu_+)^{1-m}(l_0 - \bar{l})^{2-p}r_0^p)$ to obtain

$$\begin{split} & \operatorname{ess\,sup} \frac{1}{r_0^n} \int_{B_0 \times \{t\}} G\left(\frac{l_0 - u}{l_0 - \bar{l}}\right) \varphi_0(\bar{l})^k \, \mathrm{d}x \\ & \leq \gamma \frac{(\mu_+)^{m-1} (l_0 - \bar{l})^{p-2}}{r_0^{p+n}} \iint_{L_0(\bar{l})} \left(\frac{l_0 - u}{l_0 - \bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_0(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{1}{r_0^n} \iint_{L_0(\bar{l})} \frac{l_0 - u}{l_0 - \bar{l}} |\partial_t \varphi_0(\bar{l})| \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{r_0^{p-n}}{(l_0 - \bar{l})^p} (\mu_+)^{1-m} \int_{B_0} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{r_0^{p-n}}{(l_0 - \bar{l})^{p-1}} (\mu_+)^{1-m} \int_{B_0} |f| \, \mathrm{d}x, \end{split}$$

since $u \le \mu_+$ in $L_0(\bar{l})$, $|D\varphi_0(\bar{l})| \le r_0^{-1}$, $\varphi_0(\bar{l}) = 0$ on $\partial_P Q_0(\bar{l})$ and the first term on the right-hand side of (2.9) vanishes. In view of $l_0 - u \le \frac{1}{4}\omega$ in $L_0(\bar{l})$ and $|\partial_t \varphi_0(\bar{l})| \le 9(l_0 - u)$

 $\bar{l})^{p-2}(\mu_+)^{m-1}r_0^{-p}$, we infer from (3.11) and (3.12) that

$$\begin{split} &\frac{1}{r_0^n} \iint_{L_0(\bar{l})} \frac{l_0 - u}{l_0 - \bar{l}} |\partial_t \varphi_0(\bar{l})| \, \mathrm{d}x \mathrm{d}t \\ & \leq \gamma \frac{(\mu_+)^{m-1} (l_0 - \bar{l})^{p-3}}{r_0^{n+p}} \iint_{L_0(\bar{l})} (l_0 - u) \, \mathrm{d}x \mathrm{d}t \\ & \leq \gamma C^{n+p} \frac{(\mu_+)^{m-1} \omega^{p-2}}{R^{n+p}} |L_0(\bar{l})| \leq \gamma C^{n+p} \nu_0, \end{split}$$

where the constant γ depends only upon the data. Furthermore, we infer from (3.11) and (3.6) that there exists a constant γ depending only upon the data, such that

$$\frac{r_0^{p-n}}{(l_0 - \bar{l})^p} (\mu_+)^{1-m} \int_{B_0} g^{\frac{p}{p-1}} dx + \frac{r_0^{p-n}}{(l_0 - \bar{l})^{p-1}} (\mu_+)^{1-m} \int_{B_0} |f| dx
\leq \gamma \frac{r_0^{p-n}}{\omega^p} (\mu_+)^{1-m} \int_{B_0} g^{\frac{p}{p-1}} dx + \gamma \frac{r_0^{p-n}}{\omega^{p-1}} (\mu_+)^{1-m} \int_{B_0} |f| dx
\leq \gamma C^{n-p} (B^{-p} + B^{1-p}).$$

Collecting above estimates, we conclude with

$$\operatorname{ess\,sup} \frac{1}{r_0^n} \int_{B_0 \times \{t\}} G\left(\frac{l_0 - u}{l_0 - \bar{l}}\right) \varphi_0(\bar{l})^k \, \mathrm{d}x \le \gamma_1 C^{n+p} \nu_0 + \gamma_1 C^{n-p} (B^{-p} + B^{1-p}),$$

where $\gamma_1 = \gamma_1(\text{data})$. Consequently, we arrive at

$$A_0(\bar{l}) \le (\gamma_0 + \gamma_1)C^{n+p}\nu_0 + \gamma_1C^{n-p}(B^{-p} + B^{1-p}).$$

At this stage, we fix a number $\chi \in (0,1)$ which will be determined in the course of the proof. Then, we choose $v_0 = v_0(\text{data}, \chi) < 1$ and $B = B(\text{data}, \chi) > 1$ be such that

$$(\gamma_0 + \gamma_1)C^{n+p}\nu_0 = \frac{\chi}{4}$$
 and $\gamma_1C^{n-p}(B^{-p} + B^{1-p}) < \frac{\chi}{4}$. (3.13)

This implies that $A_0(\bar{l}) \leq \frac{1}{2}\chi$. Our task now is to determine the value of l_1 . We first consider the case

$$\iint_{B_{r_0}(x_1)\times\{t_1-r_0^p< t< t_1\}} (l_0-u)_+^{(1+\lambda)(p-1)}\,\mathrm{d}x\,\mathrm{d}t=0.$$

According to Lemma 3.3, we conclude that $u(x_1, t_1) \ge l_0 = \mu_- + \frac{1}{4}\omega$, which proves the desired estimate (3.4). In the case

$$\iint_{B_{r_0}(x_1)\times\{t_1-r_0^p< t< t_1\}} (l_0-u)_+^{(1+\lambda)(p-1)} \,\mathrm{d}x\,\mathrm{d}t > 0,$$

we see that $A_0(l) \to +\infty$ as $l \to l_0$. Noting that $A_0(l)$ is continuous and increasing, then there exists a number $\tilde{l} \in (\bar{l}, l_0)$ such that $A_0(\tilde{l}) = \chi$. From (3.11) and (3.6), we infer that for B > 8 there holds

$$l_0 - \bar{l} \ge \frac{1}{32}\omega \ge \frac{1}{4B}\omega > \frac{1}{4}(\alpha_{-1} - \alpha_0),$$
 (3.14)

since $B\alpha_{-1} < \omega$. At this point, we set

$$l_{1} = \begin{cases} \tilde{l}, & \text{if } \tilde{l} < l_{0} - \frac{1}{4}(\alpha_{-1} - \alpha_{0}), \\ l_{0} - \frac{1}{4}(\alpha_{-1} - \alpha_{0}), & \text{if } \tilde{l} \ge l_{0} - \frac{1}{4}(\alpha_{-1} - \alpha_{0}). \end{cases}$$
(3.15)

Moreover, we define $Q_0 = Q_0(l_1)$ and $d_0 = l_0 - l_1$. In view of (3.14), we see that the definition of l_1 is justified. Since $B\alpha_0 < \omega$, we have $l_1 \ge \overline{l} > \mu_- + \frac{1}{8}B\alpha_0 + \frac{1}{16}\omega$.

Step 3: Determine the sequence $\{l_j\}_{j=0}^{+\infty}$. Assume that we have chosen two sequences l_1, \dots, l_j and d_0, \dots, d_{j-1} such that for $i = 1, \dots, j$, there holds

$$\frac{1}{2}\mu_{-} + \frac{1}{32}\omega + \frac{1}{2}l_{i-1} + \frac{1}{16}B\alpha_{i-1} < l_{i} \le l_{i-1} - \frac{1}{4}(\alpha_{i-2} - \alpha_{i-1}),\tag{3.16}$$

$$A_{i-1}(l_i) \le \chi,\tag{3.17}$$

$$l_i > \mu_- + \frac{1}{8}B\alpha_{i-1} + \frac{1}{16}\omega.$$
 (3.18)

Then, we set $Q_i = Q_i(l_{i+1})$, for $i = 1, 2, \dots, j-1$ and claim that

$$A_j(\bar{l}) \le \frac{1}{2}\chi,$$
 where $\bar{l} = \frac{1}{2}l_j + \frac{1}{16}B\alpha_j + \frac{1}{32}\omega + \frac{1}{2}\mu_-.$ (3.19)

To prove (3.19), we first assert that the inclusion $Q_i \subset \hat{Q}$ holds for $i = 0, 1, \dots, j-1$, where \hat{Q} is defined in (2.14). In view of (3.8), we see that $\alpha_{i-1} - \alpha_i \ge \frac{1}{3B} 4^{-i-100} \omega$. From (3.16), we can verify the inequality

$$\begin{split} (\mu_+)^{1-m}(l_i-l_{i+1})^{2-p}r_i^p &\leq (\mu_+)^{1-m}4^{p-2}(\alpha_{i-1}-\alpha_i)^{2-p}r_i^p \\ &\leq 4^{-2i}4^{101(p-2)}(\mu_+)^{1-m}(3B)^{p-2}\omega^{2-p}R^p \leq (\mu_+)^{1-m}\left(\frac{\omega}{A}\right)^{2-p}R^p, \end{split}$$

provided that we choose

$$A > 4^{102}B. (3.20)$$

This implies the inclusion $Q_i \subset \hat{Q}$ for $i = 0, 1, \dots, j-1$ and hence, $\mu_- \le u \le \mu_+$ on Q_i for $i = 0, 1, \dots, j-1$. We now turn our attention to the proof of (3.19). To start with, we introduce a cutoff function $\tilde{u} = \max \left\{ u, \frac{1}{128} \omega \right\}$ and decompose $L_j(\bar{l}) = L'_i(\bar{l}) \cup L''_i(\bar{l})$, where

$$L'_{j}(\bar{l}) = L_{j}(\bar{l}) \cap \left\{ \frac{l_{j} - \tilde{u}}{l_{i} - \bar{l}} \le \varepsilon_{1} \right\} \quad \text{and} \quad L''_{j}(\bar{l}) = L_{j}(\bar{l}) \setminus L'_{j}(\bar{l}). \tag{3.21}$$

Moreover, we observe from (3.16) that $\{u \le l_j\} = \{\tilde{u} \le l_j\}$ and $l_j - \frac{1}{128}\omega > \frac{1}{2}l_j$. This implies that

$$\iint_{L_{j}(\overline{l})} \left(\frac{l_{j} - u}{l_{j} - \overline{l}} \right)^{(1+\lambda)(p-1)} \varphi_{j}(\overline{l})^{k-p} \, dxdt \\
\leq \iint_{L_{j}(\overline{l}) \cap \{u > \frac{1}{128}\omega\}} \left(\frac{l_{j} - \widetilde{u}}{l_{j} - \overline{l}} \right)^{(1+\lambda)(p-1)} \varphi_{j}(\overline{l})^{k-p} \\
+ \iint_{L_{j}(\overline{l}) \cap \{u \leq \frac{1}{128}\omega\}} \left(\frac{l_{j}}{l_{j} - \overline{l}} \right)^{(1+\lambda)(p-1)} \varphi_{j}(\overline{l})^{k-p} \, dxdt \\
\leq \iint_{L_{j}(\overline{l}) \cap \{u > \frac{1}{128}\omega\}} \left(\frac{l_{j} - \widetilde{u}}{l_{j} - \overline{l}} \right)^{(1+\lambda)(p-1)} \varphi_{j}(\overline{l})^{k-p} \\
+ 2^{(1+\lambda)(p-1)} \iint_{L_{j}(\overline{l}) \cap \{u \leq \frac{1}{128}\omega\}} \left(\frac{l_{j} - \frac{1}{128}\omega}{l_{j} - \overline{l}} \right)^{(1+\lambda)(p-1)} \varphi_{j}(\overline{l})^{k-p} \, dxdt \\
\leq 2^{(1+\lambda)(p-1)} \iint_{L_{j}(\overline{l})} \left(\frac{l_{j} - \widetilde{u}}{l_{j} - \overline{l}} \right)^{(1+\lambda)(p-1)} \varphi_{j}(\overline{l})^{k-p} \, dxdt$$

and we can rewrite $A_i(\bar{l})$ by

$$\begin{split} A_j(\bar{l}) \leq & \gamma \frac{(\mu_+)^{m-1}(l_j - \bar{l})^{p-2}}{r_j^{n+p}} \iint_{L_j(\bar{l})} \left(\frac{l_j - \tilde{u}}{l_j - \bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_j(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\ & + \mathrm{ess} \sup_t \frac{1}{r_j^n} \int_{B_j \times \{t\}} G\left(\frac{l_j - u}{l_j - \bar{l}}\right) \varphi_j(\bar{l})^k \, \mathrm{d}x, \end{split}$$

where the constant γ depends only upon the data. Furthermore, we infer from (3.16) and (3.18) that

$$\begin{split} l_{j} - \bar{l} &= \frac{1}{2} l_{j} - \frac{1}{16} B \alpha_{j} - \frac{1}{32} \omega - \frac{1}{2} \mu_{-} \\ &\geq \frac{1}{4} (l_{j-1} - l_{j}) + \frac{1}{4} l_{j} + \frac{1}{64} \omega + \frac{1}{32} B \alpha_{j-1} - \frac{1}{16} B \alpha_{j} - \frac{1}{32} \omega - \frac{1}{4} \mu_{-} \\ &\geq \frac{1}{4} (l_{j-1} - l_{j}) + \frac{1}{32} \left(B \alpha_{j-1} + \frac{1}{2} \omega \right) + \frac{1}{32} B \alpha_{j-1} - \frac{1}{16} B \alpha_{j} - \frac{1}{64} \omega \\ &= \frac{1}{4} (l_{j-1} - l_{j}) + \frac{1}{16} B (\alpha_{j-1} - \alpha_{j}). \end{split}$$
(3.23)

For simplicity of notation, we write $\varphi_i = \varphi_i(l_{i+1})$ and observe from (3.23) that

$$(\mu_+)^{1-m}(l_j-\bar{l})^{2-p}r_j^p \leq (\mu_+)^{1-m}\frac{r_{j-1}^p}{4^p}\left(\frac{l_{j-1}-l_j}{4}\right)^{2-p} = \frac{1}{16}(\mu_+)^{1-m}(l_{j-1}-l_j)^{2-p}r_{j-1}^p,$$

which yields $Q_j(\bar{l}) \subset Q_{j-1} \subset \hat{Q}$ and $\varphi_{j-1}(x,t) = 1$ for $(x,t) \in Q_j(\bar{l})$. Since $u \leq l_j$ on $L_j(\bar{l})$, we use (3.17) to deduce

$$\frac{(\mu_{+})^{m-1}(l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}}|L_{j}(\bar{l})| \leq \frac{1}{r_{j}^{n}} \operatorname{ess sup} \int_{L_{j}(t)} \varphi_{j-1}^{k}(\cdot,t) \, \mathrm{d}x$$

$$\leq \frac{4^{n}}{r_{j-1}^{n}} \operatorname{ess sup} \int_{B_{j-1}} G\left(\frac{l_{j-1}-u}{l_{j-1}-l_{j}}\right) \varphi_{j-1}^{k} \, \mathrm{d}x \leq 4^{n}\chi, \tag{3.24}$$

where $L_i(t) = \{x \in B_i : u(\cdot, t) \le l_i\}$. In view of (3.24), we conclude that

$$\frac{(\mu_{+})^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L'_{j}(\bar{l})} \left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\
\leq \frac{(\mu_{+})^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{i}^{n+p}} \varepsilon_{1}^{(1+\lambda)(p-1)} |L_{j}(\bar{l})| \leq 4^{n} \varepsilon_{1}^{(1+\lambda)(p-1)} \chi.$$
(3.25)

Moreover, for any fixed $\varepsilon_2 < 1$, we apply the Young's inequality to obtain

$$\begin{split} & \frac{(\mu_{+})^{m-1}(l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}''(\bar{l})} \left(\frac{l_{j}-\tilde{u}}{l_{j}-\bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\ & \leq \varepsilon_{2} \frac{(\mu_{+})^{m-1}(l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} |L_{j}(\bar{l})| \\ & + \gamma(\varepsilon_{2}) \frac{(\mu_{+})^{m-1}(l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}''(\bar{l})} \left(\frac{l_{j}-\tilde{u}}{l_{j}-\bar{l}}\right)^{p\frac{n+h}{nh}} \varphi_{j}(\bar{l})^{(k-p)q} \, \mathrm{d}x \mathrm{d}t \\ & =: T_{1} + T_{2}, \end{split}$$

with the obvious meanings of T_1 and T_2 . Here, we set

$$h = \frac{p}{p - 1 - \lambda} > 1$$
 and $q = p \frac{n + h}{nh(1 + \lambda)(p - 1)} = \frac{p - 1 - \lambda + \frac{p}{n}}{p - 1 + \lambda p - \lambda} > 1$, (3.26)

since $\lambda < \min \{ p - 1, \frac{1}{n} \}$. According to (3.24), we see that $T_1 \le 4^n \varepsilon_2 \chi$. To estimate T_2 , we set

$$\tilde{\psi}_{j}(x,t) = \frac{1}{l_{j} - \bar{l}} \left[\int_{\bar{u}}^{l_{j}} \left(1 + \frac{l_{j} - s}{l_{j} - \bar{l}} \right)^{-\frac{1}{p} - \frac{\lambda}{p}} ds \right]_{+}$$
(3.27)

and apply Lemma 2.1 with (l, d) replaced by $(l_j, l_j - \bar{l})$ to conclude with

$$T_2 \le \gamma \frac{(\mu_+)^{m-1} (l_j - \bar{l})^{p-2}}{r_j^{n+p}} \iint_{L_j''(\bar{l})} \tilde{\psi}_j^{p \frac{n+h}{n}} \varphi_j(\bar{l})^{(k-p)q} \, \mathrm{d}x \mathrm{d}t. \tag{3.28}$$

Let $v = \tilde{\psi}_j \varphi_j^{k_1}$, where $k_1 = \frac{(k-p)nq}{p(n+h)}$. Recalling that p < n, we use Hölder's inequality with $r = \frac{n}{n-p}$ and $r' = \frac{n}{p}$, Sobolev's inequality and Lemma 2.1 with (l,d) replaced by $(l_j, l_j - \bar{l})$ to deduce

$$\iint_{L''_{j}(\bar{l})} \widetilde{\psi}_{j}^{p\frac{n+h}{n}} \varphi_{j}(\bar{l})^{(k-p)q} \, dxdt = \iint_{L''_{j}(\bar{l})} v^{p+\frac{ph}{n}} \, dxdt
\leq \int_{t_{1}-(\mu_{+})^{1-m}(l_{j}-\bar{l})^{2-p}r_{j}^{p}}^{t_{1}} \left(\int_{B_{j}} v^{\frac{np}{n-p}} \, dx \right)^{\frac{n-p}{n}} \left(\int_{L''_{j}(t)} v^{h} \, dx \right)^{\frac{p}{n}} \, dt
\leq \gamma \operatorname{ess} \sup_{t} \left(\int_{L''_{i}(t)} \frac{l_{j}-\tilde{u}}{l_{j}-\bar{l}} \varphi_{j}(\bar{l})^{k_{1}h} \, dx \right)^{\frac{p}{n}} \iint_{Q_{j}(\bar{l})} |Dv|^{p} \, dxdt,$$
(3.29)

where

$$L_j''(t) = \left\{x \in B_j : u(\cdot,t) \le l_j\right\} \cap \left\{x \in B_j : \frac{l_j - \tilde{u}(\cdot,t)}{l_i - \overline{l}} > \varepsilon_1\right\}.$$

According to Lemma 3.4, we see that

$$\int_{L_{j}''(t)} \frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}} \varphi_{j}(\bar{l})^{k_{1}h} dx \leq \int_{L_{j}''(t)} \varepsilon_{1} G\left(\frac{1}{\varepsilon_{1}} \frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}}\right) \varphi_{j}(\bar{l})^{k_{1}h} dx
\leq c(\varepsilon_{1}) \int_{L_{j}''(t)} G\left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}}\right) \varphi_{j}(\bar{l})^{k_{1}h} dx \leq c(\varepsilon_{1}) \int_{B_{j}} G\left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right) \varphi_{j}(\bar{l})^{k_{1}h} dx.$$
(3.30)

At this point, we apply Lemma 2.3 with (l, d, θ) replaced by $(l_j, l_j - \bar{l}, (\mu_+)^{1-m}(l_j - \bar{l})^{2-p}r_j^p)$ to conclude that

$$\begin{split} & \operatorname{ess\,sup} \frac{1}{r_{j}^{n}} \int_{B_{j}} G\left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right) \varphi_{j}(\bar{l})^{k_{1}h} \, \mathrm{d}x \\ & \leq \gamma \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{p+n}} \iint_{L_{j}(\bar{l})} u^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k_{1}h - p} \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{1}{r_{j}^{n}} \iint_{L_{j}(\bar{l})} \frac{l_{j} - u}{l_{j} - \bar{l}} |\partial_{t}\varphi_{j}(\bar{l})| \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{r_{j}^{p-n}}{(l_{j} - \bar{l})^{p}} (\mu_{+})^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{r_{j}^{p-n}}{(l_{j} - \bar{l})^{p-1}} (\mu_{+})^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x \\ & = : T_{3} + T_{4} + T_{5} + T_{6}, \end{split}$$

with the obvious meanings of T_3 - T_7 . To estimate T_3 , we apply (3.23) and (3.17) to conclude that

$$T_{3} \leq \gamma \frac{(l_{j} - \bar{l})^{p-2-(1+\lambda)(p-1)}(\mu_{+})^{m-1}}{r_{j}^{p+n}} \iint_{L_{j}(\bar{l})} (l_{j} - u)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k_{1}h-p} \, dxdt$$

$$\leq \gamma \frac{(l_{j-1} - l_{j})^{p-2}(\mu_{+})^{m-1}}{r_{j-1}^{p+n}} \iint_{L_{j-1}(l_{j})} \left(\frac{l_{j-1} - u}{l_{j-1} - l_{j}}\right)^{(1+\lambda)(p-1)} \varphi_{j-1}^{k-p} \, dxdt$$

$$\leq \gamma A_{j-1}(l_{j}) \leq \gamma_{1} \chi, \tag{3.31}$$

where the constant γ_1 depends only upon the data. Noting that $(1 + \lambda)(p - 1) > 1$ and $|\partial_t \varphi_j(\bar{l})| \le 9(l_j - \bar{l})^{p-2}(\mu_+)^{m-1}r_j^{-p}$, we infer from (3.24) and (3.17) that the inequality

$$\begin{split} T_{4} \leq & \gamma \frac{(l_{j} - \bar{l})^{p-2} (\mu_{+})^{m-1}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \frac{l_{j} - u}{l_{j} - \bar{l}} \, \mathrm{d}x \mathrm{d}t \\ \leq & \gamma \frac{(l_{j} - \bar{l})^{p-2} (\mu_{+})^{m-1}}{r_{j}^{n+p}} |L_{j}(\bar{l})| + \gamma \frac{(l_{j} - \bar{l})^{p-2} (\mu_{+})^{m-1}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \, \mathrm{d}x \mathrm{d}t \\ \leq & \gamma \chi + A_{j-1}(l_{j}) \leq \gamma_{2} \chi, \end{split}$$

holds for the constant γ_2 depending only upon the data. Finally, we deduce from (3.8) and (3.23) that $T_5 + T_6 \le \gamma (B^{-p} + B^{-(p-1)})$. Combining the above estimates, we infer that the estimate

$$\operatorname{ess\,sup} \frac{1}{r_{j}^{n}} \int_{B_{j}} G\left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right) \varphi_{j}(\bar{l})^{k_{1}h} \, \mathrm{d}x \le \gamma \chi + \gamma (B^{-p} + B^{-(p-1)})$$
 (3.32)

holds for a constant γ depending only upon the data. We now turn our attention to the estimate of T_2 . With the help of (3.28)-(3.32), we can rewrite the upper bound for T_2 by

$$T_{2} \leq \gamma \frac{(\mu_{+})^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{j}^{n}} (\chi + B^{-p} + B^{-(p-1)})^{\frac{p}{n}} \times \left[\iint_{Q_{j}(\bar{l})} \varphi_{j}(\bar{l})^{k_{1}p} |D\tilde{\psi}_{j}|^{p} dxdt + \iint_{Q_{j}(\bar{l})} \varphi_{j}(\bar{l})^{(k_{1}-1)p} \tilde{\psi}_{j}^{p} |D\varphi_{j}|^{p} dxdt \right]$$

$$= : \gamma(\chi + B^{-p} + B^{-(p-1)})^{\frac{p}{n}} (T_{7} + T_{8}),$$

with the obvious meanings of T_7 and T_8 . We first consider the estimate for T_7 . In the case $\mu_- \le \frac{1}{2}\mu_+$. We have $\mu_+ \le \mu_- + \omega \le \frac{1}{2}\mu_+ + \omega$, which implies that $\mu_+ \le 2\omega$. Therefore, we deduce

$$\tilde{u} \ge \frac{1}{128}\omega \ge \frac{1}{256}\mu_+.$$

In the case $\mu_- > \frac{1}{2}\mu_+$. We get $\tilde{u} \ge u \ge \mu_- > \frac{1}{2}\mu_+$ on $Q_i(\bar{l})$. In both cases, we see that

$$\tilde{u}^{m-1} \ge \left(\frac{1}{256}\mu_+\right)^{m-1} \quad \text{on} \quad Q_j(\bar{l}).$$
 (3.33)

In view of (3.33), we conclude that

$$\begin{split} T_7 &= \frac{(\mu_+)^{m-1}(l_j - \bar{l})^{p-2}}{r_j^n} \iint_{Q_j(\bar{l})} \varphi_j(\bar{l})^{k_1 p} |D\tilde{\psi}_j|^p \,\mathrm{d}x\mathrm{d}t \\ &\leq \gamma \frac{(l_j - \bar{l})^{p-2}}{r_j^n} \iint_{Q_j(\bar{l})} \tilde{u}^{m-1} |D\tilde{\psi}_j|^p \varphi_j(\bar{l})^{k_1 p} \,\mathrm{d}x\mathrm{d}t. \end{split}$$

Taking into account that $\{\tilde{u} < l_j\} = \{u < l_j\}$ and $D\tilde{u} = 0$ on $\{u \le \frac{1}{128}\omega\}$, we conclude from (3.27) that

$$\begin{split} \tilde{u}^{\frac{m-1}{p}}|D\tilde{\psi}_{j}| &= \tilde{u}^{\frac{m-1}{p}}|D\tilde{u}|\frac{1}{l_{j}-\bar{l}}\left(1+\frac{l_{j}-\tilde{u}}{l_{j}-\bar{l}}\right)^{-\frac{1}{p}-\frac{\lambda}{p}}\chi_{\left\{\bar{u}< l_{j}\right\}} \\ &= u^{\frac{m-1}{p}}|Du|\frac{1}{l_{j}-\bar{l}}\left(1+\frac{l_{j}-u}{l_{j}-\bar{l}}\right)^{-\frac{1}{p}-\frac{\lambda}{p}}\chi_{\left\{\frac{1}{128}\omega< u< l_{j}\right\}} \\ &\leq u^{\frac{m-1}{p}}|Du|\frac{1}{l_{j}-\bar{l}}\left(1+\frac{l_{j}-u}{l_{j}-\bar{l}}\right)^{-\frac{1}{p}-\frac{\lambda}{p}}\chi_{\left\{u< l_{j}\right\}} = u^{\frac{m-1}{p}}|D\psi_{j}|, \end{split}$$

where

$$\psi_j(x,t) = \frac{1}{l_j - \bar{l}} \left[\int_u^{l_j} \left(1 + \frac{l_j - s}{l_j - \bar{l}} \right)^{-\frac{1}{p} - \frac{\lambda}{p}} ds \right].$$

At this point, we apply Lemma 2.3 with (l, d, θ) replaced by $(l_j, l_j - \bar{l}, (\mu_+)^{1-m}(l_j - \bar{l})^{2-p}r_j^p)$. Taking into account the estimates for T_3 - T_6 , we conclude that

$$\begin{split} T_{7} \leq & \gamma \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{n}} \iint_{Q_{j}(\bar{l})} u^{m-1} |D\psi_{j}|^{p} \varphi_{j}(\bar{l})^{k_{1}p} \, \mathrm{d}x \mathrm{d}t \\ \leq & \gamma \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{p}} \iint_{L_{j}(\bar{l})} u^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{(k_{1}-1)p} \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \iint_{L_{j}(\bar{l})} \frac{l_{j} - u}{l_{j} - \bar{l}} |\partial_{t}\varphi_{j}(\bar{l})| \, \mathrm{d}x \mathrm{d}t + \gamma \frac{r_{j}^{p}}{(l_{j} - \bar{l})^{p}} (\mu_{+})^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x \\ & + \gamma \frac{r_{j}^{p}}{(l_{j} - \bar{l})^{p-1}} (\mu_{+})^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x \\ \leq & \gamma (\chi + B^{-p} + B^{-(p-1)}). \end{split}$$

We now turn our attention to the estimate of T_8 . To this end, we use Hölder inequality to deduce an upper bound for $\tilde{\psi}_i$

$$\tilde{\psi}_{j}(x,t) \leq \frac{1}{l_{j} - \bar{l}} \left[\int_{\tilde{u}}^{l_{j}} \left(1 + \frac{l_{j} - s}{l_{j} - \bar{l}} \right)^{-1 - \lambda} ds \right]^{\frac{1}{p}} (l_{j} - \tilde{u})_{+}^{\frac{1}{p'}} \leq \frac{(l_{j} - \tilde{u})_{+}^{\frac{1}{p'}}}{(l_{i} - \bar{l})_{p'}^{\frac{1}{p'}}}.$$
(3.34)

Since $|D\varphi_j| \le r_j^{-1}$, we use Young's inequality, (3.24) and (3.31) to conclude that

$$\begin{split} T_{8} & \leq \frac{(\mu_{+})^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{j}^{n}} \iint_{Q_{j}(\bar{l})} \varphi_{j}(\bar{l})^{(k_{1}-1)p} \tilde{\psi}_{j}^{p} |D\varphi_{j}|^{p} \, \mathrm{d}x \mathrm{d}t \\ & \leq \frac{(\mu_{+})^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}}\right)^{p-1} \varphi_{j}(\bar{l})^{(k_{1}-1)p} \, \mathrm{d}x \mathrm{d}t \\ & \leq \frac{(\mu_{+})^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} |L_{j}(\bar{l})| \\ & + \frac{(\mu_{+})^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{(k_{1}-1)p} \, \mathrm{d}x \mathrm{d}t \\ & \leq \gamma \chi \end{split}$$

and hence we arrive at $T_2 \le \gamma (\chi + B^{-p} + B^{-(p-1)})^{1+\frac{p}{n}}$ for a constant γ depending only upon the data. Finally, we arrive at

$$\frac{(\mu_{+})^{m-1}(l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}^{m}(\bar{l})} \left(\frac{l_{j}-\tilde{u}}{l_{j}-\bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \,\mathrm{d}x\mathrm{d}t \\
\leq 4^{n} \varepsilon_{2} \chi + \gamma(\varepsilon_{2})(\chi + B^{-p} + B^{-(p-1)})^{1+\frac{p}{n}}.$$
(3.35)

This also implies that

$$\frac{(\mu_{+})^{m-1}(l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \left(\frac{l_{j}-\tilde{u}}{l_{j}-\bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\
\leq 4^{n} \varepsilon_{1}^{(1+\lambda)(p-1)} \chi + 4^{n} \varepsilon_{2} \chi + \gamma(\varepsilon_{2}) (\chi + B^{-p} + B^{-(p-1)})^{1+\frac{p}{n}}.$$
(3.36)

Our next goal is to refine the estimate (3.32). To this aim, we apply Lemma 2.3 with (l, d, θ) replaced by $(l_j, l_j - \bar{l}, (\mu_+)^{1-m}(l_j - \bar{l})^{2-p}r_i^p)$ to obtain

$$\operatorname{ess\,sup} \frac{1}{r_{j}^{n}} \int_{B_{j}} G\left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right) \varphi_{j}(\bar{l})^{k} \, \mathrm{d}x \\
\leq \gamma \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{p+n}} \iint_{L_{j}(\bar{l})} u^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\
+ \gamma \frac{1}{r_{j}^{n}} \iint_{L_{j}(\bar{l})} \frac{l_{j} - u}{l_{j} - \bar{l}} \varphi_{j}(\bar{l})^{k-1} |\partial_{t}\varphi_{j}(\bar{l})| \, \mathrm{d}x \mathrm{d}t \\
+ \gamma \frac{r_{j}^{p-n}}{(l_{j} - \bar{l})^{p}} (\mu_{+})^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{r_{j}^{p-n}}{(l_{j} - \bar{l})^{p-1}} (\mu_{+})^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x \\
= : S_{1} + S_{2} + S_{3} + S_{4} + S$$

with the obvious meanings of S_1 - S_4 . To estimate S_1 , we apply (3.36) and (3.22) to deduce

$$\begin{split} S_1 & \leq 2^{(1+\lambda)(p-1)} \gamma \frac{(l_j - \bar{l})^{p-2} (\mu_+)^{m-1}}{r_j^{p+n}} \iint_{L_j(\bar{l})} \left(\frac{l_j - \tilde{u}}{l_j - \bar{l}} \right)^{(1+\lambda)(p-1)} \varphi_j(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\ & \leq 2^{(1+\lambda)(p-1)} \gamma \left[4^n \varepsilon_1^{(1+\lambda)(p-1)} \chi + 4^n \varepsilon_2 \chi + \gamma(\varepsilon_2) (\chi + B^{-p} + B^{-(p-1)})^{1+\frac{p}{n}} \right]. \end{split}$$

Next, we consider the estimate for S_2 . Similar to (3.22), we infer from $l_j - \frac{1}{128}\omega > \frac{1}{2}l_j$ that

$$\begin{split} S_2 = & \gamma \frac{1}{r_j^n} \iint_{L_j(\bar{l}) \cap \left\{u > \frac{1}{128}\omega\right\}} \frac{l_j - \tilde{u}}{l_j - \bar{l}} \varphi_j(\bar{l})^{k-1} |\partial_t \varphi_j(\bar{l})| \, \mathrm{d}x \mathrm{d}t + \\ & + \gamma \frac{1}{r_j^n} \iint_{L_j(\bar{l}) \cap \left\{u \le \frac{1}{128}\omega\right\}} \frac{l_j}{l_j - \bar{l}} \varphi_j(\bar{l})^{k-1} |\partial_t \varphi_j(\bar{l})| \, \mathrm{d}x \mathrm{d}t \\ \leq & 2\gamma \frac{1}{r_j^n} \iint_{L_j(\bar{l})} \frac{l_j - \tilde{u}}{l_j - \bar{l}} \varphi_j(\bar{l})^{k-1} |\partial_t \varphi_j(\bar{l})| \, \mathrm{d}x \mathrm{d}t. \end{split}$$

Furthermore, we decompose $L_j(\bar{l}) = L'_j(\bar{l}) \cup L''_j(\bar{l})$, where $L'_j(\bar{l})$ and $L''_j(\bar{l})$ satisfy (3.21). Since $|\partial_t \varphi_j(\bar{l})| \leq 9(l_j - \bar{l})^{p-2} (\mu_+)^{m-1} r_j^{-p}$, we use (3.24) and (3.35) to conclude that

$$\begin{split} S_2 \leq & \gamma \frac{(l_j - \bar{l})^{p-2} (\mu_+)^{m-1}}{r_j^{n+p}} \iint_{L_j(\bar{l})} \frac{l_j - \tilde{u}}{l_j - \bar{l}} \varphi_j(\bar{l})^{k-1} \, \mathrm{d}x \mathrm{d}t \\ \leq & \gamma \varepsilon_1 \frac{(\mu_+)^{m-1} (l_j - \bar{l})^{p-2}}{r_j^{n+p}} |L_j'(\bar{l})| \\ & + \gamma \frac{(l_j - \bar{l})^{p-2} (\mu_+)^{m-1}}{r_j^{p+n}} \iint_{L_j'(\bar{l})} \left(\frac{l_j - \tilde{u}}{l_j - \bar{l}} \right)^{(1+\lambda)(p-1)} \varphi_j(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\ \leq & 4^n \varepsilon_1 \chi + \gamma \left[4^n \varepsilon_2 \chi + \gamma (\varepsilon_2) (\chi + B^{-p} + B^{-(p-1)})^{1+\frac{p}{n}} \right]. \end{split}$$

Similar to the estimates of T_5 and T_6 , we see that $S_3 + S_4 \le \gamma (B^{-p} + B^{-(p-1)})$. Plugging the estimates for S_1 - S_4 into (3.37) and taking into account (3.36), we arrive at

$$A_{j}(\bar{l}) \leq \gamma (B^{-p} + B^{-(p-1)}) + 2^{(1+\lambda)(p-1)} \gamma 4^{n} (\varepsilon_{1} + \varepsilon_{2}) \chi + \gamma (\varepsilon_{1}, \varepsilon_{2}) \left(\chi^{1+\frac{p}{n}} + (B^{-p} + B^{-(p-1)})^{1+\frac{p}{n}} \right).$$
(3.38)

At this point, we first choose ε_1 and ε_2 be such that

$$\varepsilon_1 = \varepsilon_2 = \frac{1}{2^{3+2n+(1+\lambda)(p-1)\gamma}}. (3.39)$$

Next, we determine the value of χ by

$$\chi = \frac{1}{100^{\frac{n}{p}} \gamma(\varepsilon_1, \varepsilon_2)^{\frac{n}{p}}}.$$
(3.40)

Finally, with the choices of ε_1 , ε_2 and χ , we set B so large that

$$B^{-p} + B^{1-p} < \min \left\{ \frac{1}{100\gamma} \chi, \left(\frac{1}{100\gamma(\varepsilon_1, \varepsilon_2)} \chi \right)^{\frac{1}{1+\frac{p}{n}}} \right\}.$$

With the choices of ε_1 , ε_2 , χ and B, we get $A_j(\bar{l}) \leq \frac{1}{2}\chi$, which completes the proof of (3.19). We remark that the choice of χ in (3.40) determines the value of v_0 via (3.13), i.e.,

$$\nu_0 = \frac{\chi}{4(\gamma_0 + \gamma_1)C^{n+p}},\tag{3.41}$$

where γ_0 and γ_1 are the constants in (3.13). Our task now is to determine the value of l_{j+1} . To this end, we first consider the case

$$\iint_{B_{r_j}(x_1) \times \left\{t_1 - r_j^p < t < t_1\right\}} (l_j - u)_+^{(1+\lambda)(p-1)} \, \mathrm{d}x \, \mathrm{d}t = 0.$$

Then, we apply Lemma 3.3 and (3.18) to conclude that

$$u(x_1, t_1) \ge l_j \ge \mu_- + \frac{1}{8}B\alpha_{i-1} + \frac{1}{16}\omega > \mu_- + \frac{1}{16}\omega,$$

which proves (3.4). Next, we consider the case

$$\iint_{B_{r_i}(x_1) \times \left\{t_1 - r_i^p < t < t_1\right\}} (l_j - u)_+^{(1+\lambda)(p-1)} \, \mathrm{d}x \, \mathrm{d}t > 0.$$

We see that $A_j(l) \to +\infty$ as $l \to l_j$. Noting that $A_j(l)$ is continuous and increasing, then there exists a number $\tilde{l} \in (\bar{l}, l_j)$ such that $A_j(\tilde{l}) = \chi$. At this point, we choose l_{j+1} via

$$l_{j+1} = \begin{cases} \tilde{l}, & \text{if } \tilde{l} < l_j - \frac{1}{4}(\alpha_{j-1} - \alpha_j), \\ l_j - \frac{1}{4}(\alpha_{j-1} - \alpha_j), & \text{if } \tilde{l} \ge l_j - \frac{1}{4}(\alpha_{j-1} - \alpha_j). \end{cases}$$
(3.42)

According to (3.23), we see that $\bar{l} < l_j - \frac{1}{4}(\alpha_{j-1} - \alpha_j)$ and the definition of l_{j+1} is justified. Step 4: *Proof of the inequality* (3.4). To start with, we set $Q_j = Q_j(l_{j+1})$, $L_j = L_j(l_{j+1})$ and $d_j = l_j - l_{j+1}$. In view of (3.42), we observe that (3.16) and (3.17) hold with i = j + 1. Moreover, from (3.18) i, we find that

$$\begin{split} l_{j+1} > \bar{l} &= \frac{1}{2} l_j + \frac{1}{16} B \alpha_j + \frac{1}{32} \omega + \frac{1}{2} \mu_- \\ &> \frac{1}{2} \left(\mu_- + \frac{1}{8} B \alpha_{j-1} + \frac{1}{16} \omega \right) + \frac{1}{16} B \alpha_j + \frac{1}{32} \omega + \frac{1}{2} \mu_- \\ &= \mu_- + \frac{1}{16} B (\alpha_j + \alpha_{j-1}) + \frac{1}{16} \omega \ge \mu_- + \frac{1}{8} B \alpha_j + \frac{1}{16} \omega, \end{split}$$

since $\alpha_{j-1} \geq \alpha_j$. This implies that $(3.18)_{j+1}$ holds. Repeating the arguments above, we continue to define l_{j+2} and therefore we can construct a sequence of numbers $\{l_i\}_{i=0}^{\infty}$ satisfying (3.16)-(3.18). Since the sequence $\{l_i\}_{i=0}^{\infty}$ is decreasing, we infer from (3.18) that the limitation of l_i exists. This also implies that $d_i \to 0$ as $i \to \infty$. Set

$$\hat{l} = \lim_{i \to \infty} l_i$$

and we claim that $\hat{l} = u(x_1, t_1)$. Noting that $\mu_+ > 0$ and (3.17) holds for any $i = 1, 2, \dots$, we conclude that

$$\frac{1}{r_i^{n+p}} \iint_{B_{r_i}(x_1) \times \{t_1 - r_i^p < t < t_1\}} (\hat{l} - u)_+^{(1+\lambda)(p-1)} \, \mathrm{d}x \, \mathrm{d}t \le \frac{4^{n+p}}{r_{i-1}^{n+p}} \iint_{L_{i-1}} (l_{i-1} - u)^{(1+\lambda)(p-1)} \, \varphi_{i-1}^{k-p} \, \mathrm{d}x \, \mathrm{d}t \\
\le 4^{n+p} (\mu_+)^{1-m} A_{i-1} (l_i) d_{i-1}^{(1+\lambda)(p-1)-(p-2)} \le 4^{n+p} (\mu_+)^{1-m} \chi d_{i-1}^{(1+\lambda)(p-1)-(p-2)} \to 0$$

as $i \to \infty$. According to Lemma 3.3, we have $\hat{l} = u(x_1, t_1)$. Next, we show that for any $j \ge 1$ there holds

$$d_{j} \leq \frac{1}{4} d_{j-1} + \gamma \frac{4^{-j-100}}{B} \omega + \gamma \left(r_{j-1}^{p-n} (\mu_{+})^{1-m} \int_{B_{j-1}} g(y)^{\frac{p}{p-1}} dy \right)^{\frac{1}{p}} + \gamma \left(r_{j-1}^{p-n} (\mu_{+})^{1-m} \int_{B_{j-1}} |f(y)| dy \right)^{\frac{1}{p-1}}.$$

$$(3.43)$$

To start with, for any fixed $j \ge 1$, we first assume that

$$d_j > \frac{1}{4}d_{j-1}$$
 and $d_j > \frac{1}{4}(\alpha_{j-1} - \alpha_j),$ (3.44)

since otherwise (3.43) holds immediately. From $d_j > \frac{1}{4}(\alpha_{j-1} - \alpha_j)$, we infer from (3.42) that $A_i(l_{j+1}) = A_i(\tilde{l}) = \chi$. In view of $d_i > \frac{1}{4}d_{j-1}$, we find that

$$(\mu_{+})^{1-m}d_{j}^{2-p}r_{j}^{p} \leq (\mu_{+})^{1-m}\frac{r_{j-1}^{p}}{4^{p}}\left(\frac{d_{j-1}}{4}\right)^{2-p} = \frac{1}{16}(\mu_{+})^{1-m}d_{j-1}^{2-p}r_{j-1}^{p},$$

which yields $Q_j \subset Q_{j-1} \subset \hat{Q}$ and $\varphi_{j-1}(x,t) = 1$ for $(x,t) \in Q_j$. Taking into account that $d_j > \frac{1}{4}d_{j-1}$, we can repeat the arguments from Step 3. Then, we deduce the estimate similar

to (3.38),

$$\begin{split} \chi &= A_j(l_{j+1}) \leq \gamma \frac{r_j^{p-n}}{d_j^p} (\mu_+)^{1-m} \int_{B_j} g^{\frac{p}{p-1}} \, \mathrm{d} x + \gamma \frac{r_j^{p-n}}{d_j^{p-1}} (\mu_+)^{1-m} \int_{B_j} |f| \, \mathrm{d} x \\ &+ 2^{(1+\lambda)(p-1)} \gamma 4^n (\varepsilon_1 + \varepsilon_2) \chi + \gamma (\varepsilon_1, \varepsilon_2) \chi^{1+\frac{p}{n}} \\ &+ \gamma \left[\frac{r_j^{p-n}}{d_j^p} (\mu_+)^{1-m} \int_{B_j} g^{\frac{p}{p-1}} \, \mathrm{d} x + \gamma \frac{r_j^{p-n}}{d_j^{p-1}} (\mu_+)^{1-m} \int_{B_j} |f| \, \mathrm{d} x \right]^{1+\frac{p}{n}}. \end{split}$$

Recalling that the choices of ε_1 , ε_2 and χ in (3.39)-(3.40) yields that

$$2^{(1+\lambda)(p-1)}\gamma 4^n(\varepsilon_1+\varepsilon_2)\chi+\gamma(\varepsilon_1,\varepsilon_2)\chi^{1+\frac{p}{n}}\leq \frac{1}{2}\chi.$$

Consequently, we conclude that either

$$d_{j} \leq \gamma \left(r_{j}^{p-n} (\mu_{+})^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} dx \right)^{\frac{1}{p}} \quad \text{or} \quad d_{j} \leq \gamma \left(r_{j}^{p-n} (\mu_{+})^{1-m} \int_{B_{j}} |f| dx \right)^{\frac{1}{p-1}},$$

which proves the inequality (3.43). Let J > 1 be a fixed integer. We sum up the inequality (3.43) for $j = 1, \dots, J-1$ and there holds

$$\begin{split} l_1 - l_J &\leq \frac{1}{3} d_0 + \gamma \frac{1}{B} \sum_{j=1}^{J-1} 4^{-j-100} \omega + \gamma \sum_{j=1}^{J-1} \left(r_{j-1}^{p-n} (\mu_+)^{1-m} \int_{B_{j-1}} |f(y)| \, \mathrm{d}y \right)^{\frac{1}{p-1}} \\ &+ \gamma \sum_{j=1}^{J-1} \left(r_{j-1}^{p-n} (\mu_+)^{1-m} \int_{B_{j-1}} g(y)^{\frac{p}{p-1}} \, \mathrm{d}y \right)^{\frac{1}{p}}, \end{split}$$

where the constant γ depends only upon the data. Since $d_0 = l_0 - l_1$ and $l_0 = \mu_- + \frac{1}{4}\omega$, we use (3.6) to obtain

$$\mu_{-} + \frac{1}{4}\omega \leq \frac{4}{3}d_{0} + l_{J} + \gamma \frac{1}{B}\omega + \gamma(\mu_{+})^{\frac{1-m}{p-1}} \int_{0}^{4C^{-1}R} \left(\frac{1}{r^{n-p}} \int_{B_{r}(x_{1})} f(y) \, \mathrm{d}y\right)^{\frac{1}{p-1}} \frac{1}{r} \, \mathrm{d}r$$

$$+ \gamma(\mu_{+})^{\frac{1-m}{p}} \int_{0}^{4C^{-1}R} \left(\frac{1}{r^{n-p}} \int_{B_{r}(x_{1})} g(y)^{\frac{p}{p-1}} \, \mathrm{d}y\right)^{\frac{1}{p}} \frac{1}{r} \, \mathrm{d}r$$

$$\leq \frac{4}{3}d_{0} + l_{J} + \gamma \frac{1}{B}\omega$$

$$(3.45)$$

for a constant γ depending only upon the data. Recalling that $d_0 \leq l_0 - \bar{l}$ where $\bar{l} = \frac{1}{2}l_0 + \frac{1}{2}\mu_- + \frac{1}{16}B\alpha_0 + \frac{1}{32}\omega$. According to (3.11), we have $d_0 \leq \frac{1}{8}\omega$. Letting $J \to \infty$ in (3.45), we infer that

$$u(x_1, t_1) > \mu_- + \frac{1}{24}\omega,$$

provided that we choose $B > 24\gamma$. Since (x_1, t_1) is a fixed Lebesgue point of u, we conclude that the inequality (3.4) holds for almost everywhere point in $Q_{\frac{1}{4}R}^-(\bar{t})$, provided that ν_0 satisfies (3.41). The proof of the lemma is now complete.

We now define a time level $\hat{t} = \bar{t} - (\mu_+)^{1-m} \omega^{2-p} \left(\frac{1}{4}R\right)^p$ and it follows from Lemma 3.5 that

$$u(x,\hat{t}) > \mu_{-} + \frac{\omega}{2^{5}}$$
 for $x \in B_{\frac{R}{4}}$. (3.46)

Furthermore, there exists a constant $A_1 > 0$ such that $4^{-\frac{p}{p-2}} < A_1 < A$ and

$$\hat{t} = -(\mu_+)^{1-m} A_1^{p-2} \omega^{2-p} R^p.$$

Next, we establish the following result regarding the time propagation of positivity.

Lemma 3.6. Let u be a bounded nonnegative weak solution to (1.1)-(1.2) in Ω_T . Given $v_* \in (0, 1)$, there exists a constant $s_* = s_*(data, v_*) > 5$, such that either

$$\omega \le 2^{\frac{2}{p}S_*}(\mu_+)^{\frac{1-m}{p}}F_1(2R) + 2^{\frac{1}{p-1}S_*}(\mu_+)^{\frac{1-m}{p-1}}F_2(2R)$$
(3.47)

or

$$\left| \left\{ x \in B_{\frac{R}{8}} : u(x,t) < \mu_{-} + \frac{\omega}{2^{s_{*}}} \right\} \right| \le \nu_{*} |B_{\frac{R}{8}}| \tag{3.48}$$

holds for any $t \in (\hat{t}, 0)$.

Proof. We first recall that

$$\psi^{-} = \ln^{+} \left(\frac{H_{k}^{-}}{H_{k}^{-} - (u - k)_{-} + c} \right) = \begin{cases} \ln \left(\frac{H_{k}^{-}}{H_{k}^{-} + u - k + c} \right), & k - H_{k}^{-} \le u < k - c, \\ 0, & u \ge k - c, \end{cases}$$
(3.49)

where

$$H_k^- \ge \operatorname{ess\,sup}(u - k)_- \qquad \text{ and } \qquad 0 < c < H_k^-.$$

Next, we let $k = \mu_- + \frac{1}{2^3}\omega$, $c = \frac{1}{2^{3+l}}\omega$ where $l \ge 1$ will be determined later. Moreover, we choose $H_k^- = \frac{1}{32}\omega$, which is admissible since $(u - k)_- \le \frac{1}{32}\omega$. In view of (3.49), we see that

$$\psi^{-}(u) \le l \ln 2$$
 and $|(\psi^{-})'(u)|^{2-p} \le \omega^{p-2}$. (3.50)

We now apply the logarithmic estimate (2.12) over the cylinder $B_{\frac{R}{2}} \times [\hat{t}, 0]$. This yields that

$$\begin{split} & \operatorname{ess\,sup} \int_{B_{\frac{R}{4}} \times \{i\}} [\psi^{-}(u)]^2 \zeta^p \, \mathrm{d}x \\ & \leq \int_{B_{\frac{R}{4}} \times \{\hat{i}\}} [\psi^{-}(u)]^2 \zeta^p \, \mathrm{d}x + \gamma \iint_{B_{\frac{R}{4}} \times [\hat{i}, 0]} u^{m-1} \psi^{-}(u) |D\zeta|^p \left[(\psi^{-})'(u) \right]^{2-p} \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \ln \left(\frac{H}{c} \right) \left(\frac{1}{c} \iint_{B_{\frac{R}{4}} \times [\hat{i}, 0]} |f| \, \mathrm{d}x \mathrm{d}t + \frac{1}{c^2} \iint_{B_{\frac{R}{4}} \times [\hat{i}, 0]} g^{\frac{p}{p-1}} \, \mathrm{d}x \mathrm{d}t \right) \\ & = : I_1 + I_2 + I_3. \end{split}$$

We take the cutoff function $\zeta = \zeta(x)$, which satisfies $0 \le \zeta \le 1$ in $B_{\frac{R}{4}}$, $\zeta \equiv 1$ in $B_{\frac{R}{8}}$ and $|D\zeta| \le 4R^{-1}$. Observe from (3.46) that $\psi^-(x,\hat{t}) = 0$ for all $x \in B_{\frac{R}{4}}$, we get $I_1 = 0$. To estimate I_2 , we note that $-\hat{t} \le A^{p-2}(\mu_+)^{1-m}\omega^{2-p}R^p$. In view of (3.50), we conclude that

$$I_2 \le 4l(\ln 2)(-\hat{t})|B_{\frac{R}{4}}|(\mu_+)^{m-1}R^{-p}\omega^{p-2} \le \gamma lA^{p-2}|B_{\frac{R}{8}}|.$$

Finally, we consider the estimate for I_3 . Using the inequality $-\hat{t} \leq A^{p-2}(\mu_+)^{1-m}\omega^{2-p}R^p$ again, we infer that

$$\begin{split} I_{3} \leq & l(\ln 2) \frac{2^{5+l} A^{p-2}}{\omega^{p-1}} (\mu_{+})^{1-m} R^{p} \int_{B_{R}} |f| \, \mathrm{d}x + l(\ln 2) \frac{2^{10+2l} A^{p-2}}{\omega^{p}} (\mu_{+})^{1-m} R^{p} \int_{B_{R}} g^{\frac{p}{p-1}} \, \mathrm{d}x \\ \leq & \gamma (lA^{p-2}) \frac{2^{l}}{\omega^{p-1}} \left((\mu_{+})^{1-m} R^{p-n} \int_{B_{R}} |f| \, \mathrm{d}x \right) |B_{\frac{R}{8}}| \\ & + \gamma (lA^{p-2}) \frac{2^{2l}}{\omega^{p}} \left((\mu_{+})^{1-m} R^{p-n} \int_{B_{R}} g^{\frac{p}{p-1}} \, \mathrm{d}x \right) |B_{\frac{R}{8}}|. \end{split}$$

At this point, we assume that

$$\omega > 2^{\frac{2}{p}l}(\mu_{+})^{\frac{1-m}{p}}F_{1}(2R) + 2^{\frac{1}{p-1}l}(\mu_{+})^{\frac{1-m}{p-1}}F_{2}(2R)$$
(3.51)

and this yields $I_3 \leq \gamma l A^{p-2} |B_{\frac{R}{5}}|$. Therefore, we arrive at

$$\int_{B_{\frac{R}{4}} \times \{t\}} [\psi^{-}(u)]^{2} \zeta^{p} \, \mathrm{d}x \le \gamma l A^{p-2} |B_{\frac{R}{8}}|$$
(3.52)

for any $t \in (\hat{t}, 0)$. On the other hand, the left-hand side of (3.52) is estimated below by integrating over the smaller set

$$\left\{x\in B_{\frac{R}{8}}: u<\mu_-+\frac{\omega}{2^{l+5}}\right\}\subset B_{\frac{R}{4}}.$$

On such a set, $\zeta \equiv 1$ and

$$\psi^{-} = \ln^{+} \left(\frac{\frac{1}{2^{5}} \omega}{\frac{1}{2^{5}} \omega - (u - k)_{-} + \frac{1}{2^{l+5}} \omega} \right) \ge \ln 2^{l-1} > (l-1) \ln 2.$$

This yields

$$\int_{B_{\frac{R}{8}} \times \{t\}} (\psi^{-})^{2} \zeta^{p} dx \ge (l-1)^{2} (\ln 2)^{2} \left| \left\{ x \in B_{\frac{R}{8}} : u < \mu_{-} + \frac{\omega}{2^{l+5}} \right\} \right|, \tag{3.53}$$

for all $t \in (\hat{t}, 0)$. Combining (3.52) and (3.53), we deduce

$$\left| \left\{ x \in B_{\frac{R}{8}} : u < \mu_{-} + \frac{\omega}{2^{l+5}} \right\} \right| \leq \gamma \frac{l}{(l-1)^{2}} A^{p-2} |B_{\frac{R}{8}}|.$$

For $v_* \in (0, 1)$, we set l be a fixed integer such that $l > 1 + \gamma v_*^{-1} A^{p-2}$ and choose $s_* = l + 5$. This proves the inequality (3.48) under the assumption (3.51). On the other hand, if (3.51) is violated, then we obtain (3.47) for such a choice of s_* . The proof of the lemma is now complete.

According to the proof of Lemma 3.6, we conclude that for any fixed $v_* \in (0, 1)$, we can choose

$$s_* = s_*(\nu_*) = 2\nu_*^{-1} A^{p-2}$$
(3.54)

and Lemma 3.6 follows for such a choice of s_* . Moreover, for R > 0, we set

$$\tilde{Q}_1 = B_{\frac{R}{8}} \times (\hat{t}, 0) \qquad \tilde{Q}_2 = B_{\frac{R}{16}} \times \left(\frac{\hat{t}}{2}, 0\right).$$

We now establish a variant De Giorgi type lemma which concerns the initial data. The proof is similar to that of Lemma 3.5, but the arguments need some nontrivial modifications.

Lemma 3.7. Let $\xi = 2^{-s_*}$ and $0 < \xi < 2^{-5}$. Let u be a bounded nonnegative weak solution to (1.1)-(1.2) in Ω_T . Then, there exist $v_1 \in (0,1)$ and $\tilde{B} > 1$ which can be quantitatively determined a priori only in terms of the data, such that if

$$\left|\left\{(x,t)\in \tilde{Q}_1: u<\mu_-+\xi\omega\right\}\right|\leq \nu_1|\tilde{Q}_1|,$$

then either

$$u(x,t) > \mu_{-} + \frac{1}{2}\xi\omega$$
 for a.e. $(x,t) \in \tilde{Q}_{2}$ (3.55)

or

$$\xi\omega \le \tilde{B}\left((\mu_{-} + \xi\omega)^{\frac{1-m}{p}}F_{1}(2R) + (\mu_{-} + \xi\omega)^{\frac{1-m}{p-1}}F_{2}(2R)\right).$$
 (3.56)

Proof. For simplicity of notation, we write $\tilde{\mu} = \mu_- + \xi \omega$ and observe that $\tilde{\mu} \le \mu_+$. We first assume that (3.56) is violated, that is,

$$\xi \omega > \tilde{B}\left(\tilde{\mu}^{\frac{1-m}{p}}F_1(2R) + \tilde{\mu}^{\frac{1-m}{p-1}}F_2(2R)\right),$$
 (3.57)

since otherwise there is nothing to prove. Our problem reduces to prove (3.55). Fix $(x_1,t_1)\in \tilde{Q}_{\frac{R}{16}}$ and assume that (x_1,t_1) is a Lebesgue point of u in the sense of Definition 3.1. Set $r_j=4^{-j-4}R$ and $B_j=B_{r_j}(x_1)$. Next, we set $Q_j=B_j\times(\hat{t},t_1)$ and $\phi_j\in C_0^\infty(B_j)$ where $\phi_j=1$ on B_{j+1} and $|D\phi_j|\leq r_j^{-1}$. For a sequence $\{l_j\}_{j=0}^\infty$ and a fixed l>0, we set

$$Q_{j}(l) = B_{j} \times (t_{1} - \tilde{\mu}^{1-m}(l_{j} - l)^{2-p}r_{j}^{p}, t_{1}).$$

Furthermore, we define $\varphi_j(l) = \phi_j(x)\theta_{j,l}(t)$, where $\phi_j \in C_0^{\infty}(B_j)$, $\phi_j = 1$ on B_{j+1} , $|D\phi_j| \le r_j^{-1}$ and $\theta_{j,l}(t)$ is a Lipschitz function satisfies

$$\theta_{j,l}(t) = 1$$
 in $t \ge t_1 - \frac{4}{9}\tilde{\mu}^{1-m}(l_j - l)^{2-p}r_j^p$, $\theta_{j,l}(t) = 0$ in $t \le t_1 - \frac{5}{9}\tilde{\mu}^{1-m}(l_j - l)^{2-p}r_j^p$

and

$$\theta_{j,l}(t) = \frac{t - t_1 - \frac{5}{9}\tilde{\mu}^{1-m}(l_j - l)^{2-p}r_j^p}{\frac{1}{9}\tilde{\mu}^{1-m}(l_j - l)^{2-p}r_i^p}$$

in $t_1 - \frac{5}{9}\tilde{\mu}^{1-m}(l_j - l)^{2-p}r_j^p \le t \le t_1 - \frac{4}{9}\tilde{\mu}^{1-m}(l_j - l)^{2-p}r_j^p$. Next, for $j = -1, 0, 1, 2, \cdots$, we introduce the sequence $\{\alpha_j\}$ by

$$\alpha_{j} = \int_{0}^{r_{j}} \left(r^{p-n} \tilde{\mu}^{1-m} \int_{B_{r}(x_{1})} g(y)^{\frac{p}{p-1}} dy \right)^{\frac{1}{p}} \frac{dr}{r} + \int_{0}^{r_{j}} \left(r^{p-n} \tilde{\mu}^{1-m} \int_{B_{r}(x_{1})} |f(y)| dy \right)^{\frac{1}{p-1}} \frac{dr}{r}.$$
(3.58)

In view of (3.58), we see that $\alpha_j \to 0$ as $j \to \infty$ and there exists a constant γ depending only upon the data such that

$$\alpha_{j-1} - \alpha_j \ge \gamma \left(r_j^{p-n} \tilde{\mu}^{1-m} \int_{B_i} g(y)^{\frac{p}{p-1}} \, \mathrm{d}y \right)^{\frac{1}{p}} + \gamma \left(r_j^{p-n} \tilde{\mu}^{1-m} \int_{B_i} |f(y)| \, \mathrm{d}y \right)^{\frac{1}{p-1}}$$
(3.59)

and

$$\alpha_{j-1} - \alpha_j \le \gamma \left(r_{j-1}^{p-n} \tilde{\mu}^{1-m} \int_{B_{j-1}} g(y)^{\frac{p}{p-1}} \, \mathrm{d}y \right)^{\frac{1}{p}} + \gamma \left(r_{j-1}^{p-n} \tilde{\mu}^{1-m} \int_{B_{j-1}} |f(y)| \, \mathrm{d}y \right)^{\frac{1}{p-1}}. \tag{3.60}$$

We introduce a cutoff function $\tilde{u} = \max \left\{ u, \frac{1}{128} \xi \omega \right\}$. Moreover, for a sequence $\{l_j\}$ and a fixed l > 0 we define a quantity $A_j(l)$ as follows.

• In the case $Q_i(l) \subseteq \tilde{Q}_1$, we define $A_i(l)$ by

$$A_{j}(l) = \frac{(l_{j} - l)^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}(l)} \tilde{u}^{m-1} \left(\frac{l_{j} - u}{l_{j} - l}\right)^{(1+\lambda)(p-1)} \varphi_{j}(l)^{k-p} \, dxdt + \operatorname{ess \, sup}_{t} \frac{1}{r_{j}^{n}} \int_{B_{j} \times \{t\}} G\left(\frac{l_{j} - u}{l_{j} - l}\right) \varphi_{j}(l)^{k} \, dx.$$
(3.61)

• In the case $Q_j(l) \nsubseteq \tilde{Q}_1$, i.e., $t_1 - \tilde{\mu}^{1-m}(l_j - l)^{2-p}r_i^p < \hat{t}$, we define $A_j(l)$ by

$$A_{j}(l) = \frac{(l_{j} - l)^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}} \tilde{u}^{m-1} \left(\frac{l_{j} - u}{l_{j} - l}\right)^{(1+\lambda)(p-1)} \phi_{j}^{k-p} \, dxdt + \operatorname{ess \, sup}_{\hat{l} < t < l_{1}} \frac{1}{r_{j}^{n}} \int_{B_{j} \times \{t\}} G\left(\frac{l_{j} - u}{l_{j} - l}\right) \phi_{j}^{k} \, dx.$$
(3.62)

Here, the function G is defined in (2.1), $L_j(l) = Q_j(l) \cap \{u \le l_j\}$ and $L_j = Q_j \cap \{u \le l_j\}$. From the proof of Lemma 3.5, we conclude that $A_j(l)$ is continuous for $l < l_j$. The proof of the lemma will be divided into several steps.

Step 1: Determine the values of l_0 and l_1 . To start with, we set $l_0 = \mu_- + \xi \omega$ and $\bar{l} = \frac{1}{2}l_0 + \frac{1}{2}\mu_- + \frac{1}{4}\tilde{B}\alpha_0 + \frac{1}{8}\xi\omega$. Observe that $\tilde{B}\alpha_0 < \xi\omega$ and this implies

$$l_0 - \bar{l} = \frac{1}{2}\mu_- + \frac{1}{2}\xi\omega - \frac{1}{2}\mu_- - \frac{1}{4}\tilde{B}\alpha_0 - \frac{1}{8}\xi\omega \ge \frac{1}{8}\xi\omega \quad \text{and} \quad l_0 - \bar{l} \le \frac{3}{8}\xi\omega. \quad (3.63)$$

Let $A_p > 1$ be a constant such that for any $X > A_p$ there holds $X \le 2^{(p-2)X}$. At this point, we choose

$$A > \max\left\{ \left(\frac{100p}{p-2} \right)^{\frac{1}{p-2}}, A_p^{\frac{1}{p-2}} \right\}. \tag{3.64}$$

We infer from (3.54) and (3.64) that

$$\xi^{p-2} = 2^{-s_*(p-2)} = 2^{-2(p-2)\nu_*^{-1}A^{p-2}} \le 2^{-2(p-2)A^{p-2}} \le 2^{-100p}A^{2-p}, \tag{3.65}$$

since $\nu_* < 1$. We now claim that $Q_0(\bar{l}) \nsubseteq \tilde{Q}_1$. Taking into account that $\tilde{\mu} \leq \mu_+$, we use (3.63) and (3.65) to obtain

$$\tilde{\mu}^{1-m}(l_0 - \bar{l})^{2-p}r_0^p \ge 4^{-4p}(\mu_+)^{1-m}(\xi\omega)^{2-p}R^p \ge t_1 + (\mu_+)^{1-m}A^{p-2}\omega^{2-p}R^p,$$

which proves $Q_0(\bar{l}) \nsubseteq \tilde{Q}_1$. Our task now is to deduce an upper bound for $A_0(\bar{l})$. Noting that $\tilde{u} \leq \mu_+$ and $l_0 - u \leq \xi \omega$ in L_0 , we infer from (3.63) and (3.65) that there exists a constant γ_1 depending only upon the data such that

$$\begin{split} &\frac{(l_0 - \bar{l})^{p-2}}{r_0^{n+p}} \iint_{L_0} \tilde{u}^{m-1} \left(\frac{l_0 - u}{l_0 - \bar{l}} \right)^{(1+\lambda)(p-1)} \phi_0^{k-p} \, \mathrm{d}x \mathrm{d}t \\ &\leq \frac{(\mu_+)^{m-1} (l_0 - \bar{l})^{p-2 - (1+\lambda)(p-1)}}{r_0^{n+p}} \left(\xi \omega \right)^{(1+\lambda)(p-1)} |L_0| \\ &\leq \gamma_1 \frac{(\mu_+)^{m-1} (\xi \omega)^{p-2}}{R^{n+p}} |L_0| \leq \gamma_1 \frac{(\mu_+)^{m-1} A^{2-p} \xi \omega^{p-2}}{R^{n+p}} R^n (-\hat{t}) \frac{|Q_0 \cap \{u \leq l_0\}|}{|\tilde{Q}_1|} \\ &\leq \gamma_1 \frac{\left| \left\{ (x, t) \in \tilde{Q}_1 : u < \mu_- + \xi \omega \right\} \right|}{|\tilde{Q}_1|} \leq \gamma_1 \gamma_1, \end{split}$$

since $-\hat{t} \leq A^{p-2}(\mu_+)^{1-m}\omega^{2-p}R^p$. In view of (3.46), we see that $u(x,\hat{t}) < \mu_- + \xi\omega = l_0$ for all $x \in B_{\frac{R}{4}}$. We now apply Lemma 2.3 with (l,d,θ) replaced by $(l_0,l_0-\bar{l},t_1-\hat{t})$, (3.57) and (3.63) to deduce

$$\begin{split} & \operatorname{ess\,sup} \frac{1}{r_0^n} \int_{B_0 \times \{t\}} G\left(\frac{l_0 - u}{l_0 - \bar{l}}\right) \phi_0^k \, \mathrm{d}x \\ \leq & \gamma \frac{(l_0 - \bar{l})^{p-2}}{r_0^{p+n}} \iint_{L_0} u^{m-1} \left(\frac{l_0 - u}{l_0 - \bar{l}}\right)^{(1+\lambda)(p-1)} \phi_0^{k-p} \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{r_0^{p-n}}{(\xi \omega)^p} (\mu_+)^{1-m} \int_{B_0} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{r_0^{p-n}}{(\xi \omega)^{p-1}} (\mu_+)^{1-m} \int_{B_0} |f| \, \mathrm{d}x \\ \leq & \gamma_1 \nu_1 + \gamma \frac{(\mu_+)^{1-m}}{\tilde{B}^p \tilde{u}^{1-m}} + \gamma \frac{(\mu_+)^{1-m}}{\tilde{B}^{p-1} \tilde{u}^{1-m}}, \end{split}$$

since $\phi_0 = \phi_0(x)$ is time independent and the first two terms on the right-hand side of (2.9) vanishes. Since $\tilde{\mu} \leq \mu_+$, we conclude that there exists a constant γ_1 depending only upon the data such that

$$\operatorname{ess\,sup}_{\hat{t} < t < t_1} \frac{1}{r_0^n} \int_{B_0 \times \{t\}} G\left(\frac{l_0 - u}{l_0 - \bar{l}}\right) \phi_0^k \, \mathrm{d}x \le \gamma_1 \nu_1 + \gamma_1 (\tilde{B}^{-p} + \tilde{B}^{1-p}).$$

Consequently, we conclude that $A_0(\bar{l}) \leq 2\gamma_1\nu_1 + \gamma_1(\tilde{B}^{-p} + \tilde{B}^{1-p})$. At this stage, we fix a number $\chi \in (0,1)$ which will be determined later. Now, we choose $\nu_1 < 1$ and $\tilde{B} > 1$ be such that

$$2\gamma_1 \nu_1 = \frac{\chi}{4}$$
 and $\gamma_1(\tilde{B}^{-p} + \tilde{B}^{1-p}) < \frac{\chi}{4}$. (3.66)

This yields that $A_0(\bar{l}) \leq \frac{1}{2}\chi$. The next thing to do is to determine the value of l_1 . We first consider the case

$$\iint_{B_{r_0}(x_1)\times\{t_1-r_0^p< t< t_1\}} \tilde{u}^{m-1} (l_0-u)_+^{(1+\lambda)(p-1)} \,\mathrm{d}x\,\mathrm{d}t = 0.$$

Then, we apply Lemma 3.3 to conclude that $u(x_1, t_1) \ge l_0 = \mu_- + \xi \omega$, which proves (3.55). Next, we consider the case

$$\iint_{B_{r_0}(x_1)\times\{t_1-r_0^p< t< t_1\}} \tilde{u}^{m-1} (l_0-u)_+^{(1+\lambda)(p-1)} \,\mathrm{d}x\,\mathrm{d}t > 0.$$

We find that $A_0(l) \to +\infty$ as $l \to l_0$. Noting that $A_0(l)$ defined in (3.62) is continuous and increasing, then there exists a number $\tilde{l} \in (\bar{l}, l_0)$ such that $A_0(\tilde{l}) = \chi$. In view of (3.63) and (3.57), we infer that for B > 8 there holds

$$l_0 - \bar{l} \ge \frac{1}{8}\xi\omega > \frac{1}{4}(\alpha_{-1} - \alpha_0).$$
 (3.67)

At this point, we define

$$l_{1} = \begin{cases} \tilde{l}, & \text{if } \tilde{l} < l_{0} - \frac{1}{4}(\alpha_{-1} - \alpha_{0}), \\ l_{0} - \frac{1}{4}(\alpha_{-1} - \alpha_{0}), & \text{if } \tilde{l} \ge l_{0} - \frac{1}{4}(\alpha_{-1} - \alpha_{0}), \end{cases}$$
(3.68)

and $d_0 = l_0 - l_1$. Since $\tilde{B}\alpha_0 < \xi\omega$, we have $l_1 \ge \bar{l} > \mu_- + \frac{1}{2}\tilde{B}\alpha_0 + \frac{1}{4}\xi\omega$.

Step 2: Determine the sequence $\{l_j\}_{j=0}^{\infty}$. Suppose that we have chosen two sequences l_1, \dots, l_j and d_0, \dots, d_{j-1} such that for $i = 1, \dots, j$, there holds

$$\frac{1}{2}\mu_{-} + \frac{1}{8}\xi\omega + \frac{1}{2}l_{i-1} + \frac{1}{4}\tilde{B}\alpha_{i-1} < l_{i} \le l_{i-1} - \frac{1}{4}(\alpha_{i-2} - \alpha_{i-1}),\tag{3.69}$$

$$A_{i-1}(l_i) \le \chi, \tag{3.70}$$

$$l_i > \mu_- + \frac{1}{2}\tilde{B}\alpha_{i-1} + \frac{1}{4}\xi\omega.$$
 (3.71)

Then, we claim that

$$A_{j}(\bar{l}) \le \frac{1}{2}\chi, \quad \text{where} \quad \bar{l} = \frac{1}{2}l_{j} + \frac{1}{4}\tilde{B}\alpha_{j} + \frac{1}{8}\xi\omega + \frac{1}{2}\mu_{-}.$$
 (3.72)

To prove (3.72), we first observe from (3.69) and (3.71) that $l_j - \bar{l}$ can be bounded below by

$$\begin{split} l_{j} - \bar{l} &= \frac{1}{2} l_{j} - \frac{1}{4} \tilde{B} \alpha_{j} - \frac{1}{8} \xi \omega - \frac{1}{2} \mu_{-} \\ &\geq \frac{1}{4} (l_{j-1} - l_{j}) + \frac{1}{4} l_{j} + \frac{1}{16} \xi \omega + \frac{1}{8} \tilde{B} \alpha_{j-1} - \frac{1}{4} \tilde{B} \alpha_{j} - \frac{1}{8} \xi \omega - \frac{1}{4} \mu_{-} \\ &\geq \frac{1}{4} (l_{j-1} - l_{j}) + \frac{1}{4} \left(\mu_{-} + \frac{1}{2} \tilde{B} \alpha_{j-1} + \frac{1}{4} \xi \omega \right) \\ &+ \frac{1}{16} \xi \omega + \frac{1}{8} \tilde{B} \alpha_{j-1} - \frac{1}{4} \tilde{B} \alpha_{j} - \frac{1}{8} \xi \omega - \frac{1}{4} \mu_{-} \\ &= \frac{1}{4} (l_{j-1} - l_{j}) + \frac{1}{4} \tilde{B} (\alpha_{j-1} - \alpha_{j}). \end{split}$$

$$(3.73)$$

To proceed further, we distinguish two cases. In the case $Q_j(\bar{l}) \nsubseteq \tilde{Q}_1$. We infer from (3.73) that

$$\tilde{\mu}^{1-m}(l_{j-1}-l_j)^{2-p}r_{j-1}^p \ge 16\tilde{\mu}^{1-m}(l_j-\bar{l})^{2-p}r_j^p > t_1-\hat{t}, \tag{3.74}$$

since $t_1 - \tilde{\mu}^{1-m}(l_j - \bar{l})^{2-p}r_j^p < \hat{t}$. This yields that $Q_{j-1}(l_j) \nsubseteq \tilde{Q}_1$. We now decompose $L_j = L'_j \cup L''_j$, where

$$L'_{j} = L_{j} \cap \left\{ \frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}} \le \varepsilon_{1} \right\} \quad \text{and} \quad L''_{j} = L_{j} \setminus L'_{j}. \tag{3.75}$$

Furthermore, we deduce from (3.71) that $\{u \leq l_j\} = \{\tilde{u} \leq l_j\}$ and $l_j - \frac{1}{128}\xi\omega > \frac{1}{2}l_j$, where $\tilde{u} = \max\{u, \frac{1}{128}\xi\omega\}$. It follows that

$$\iint_{L_{j}} \tilde{u}^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}} \right)^{(1+\lambda)(p-1)} \phi_{j}^{k-p} \, dxdt \\
\leq \iint_{L_{j} \cap \left\{ u > \frac{1}{128} \xi \omega \right\}} \tilde{u}^{m-1} \left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}} \right)^{(1+\lambda)(p-1)} \phi_{j}^{k-p} \\
+ 2^{(1+\lambda)(p-1)} \iint_{L_{j} \cap \left\{ u \leq \frac{1}{128} \xi \omega \right\}} \tilde{u}^{m-1} \left(\frac{l_{j} - \frac{1}{128} \xi \omega}{l_{j} - \bar{l}} \right)^{(1+\lambda)(p-1)} \phi_{j}^{k-p} \, dxdt \\
\leq 2^{(1+\lambda)(p-1)} \iint_{L_{i}} \tilde{u}^{m-1} \left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}} \right)^{(1+\lambda)(p-1)} \phi_{j}^{k-p} \, dxdt$$
(3.76)

and hence

$$\begin{split} A_j(\bar{l}) \leq & \gamma \frac{(l_j - \bar{l})^{p-2}}{r_j^{n+p}} \iint_{L_j} \tilde{u}^{m-1} \left(\frac{l_j - \tilde{u}}{l_j - \bar{l}} \right)^{(1+\lambda)(p-1)} \phi_j^{k-p} \, \mathrm{d}x \mathrm{d}t \\ & + \underset{\hat{l} < l < l_1}{\mathrm{ess}} \sup_{\hat{r}} \frac{1}{r_j^n} \int_{B_j \times \{t\}} G\left(\frac{l_j - u}{l_j - \bar{l}} \right) \phi_j^k \, \mathrm{d}x, \end{split}$$

where the constant γ depends only upon p and λ . Our task now is to establish an estimate for $A_i(\bar{l})$. Noting that $u \leq l_i$ on L_i , we have

$$\tilde{u} \le \mu_- + \xi \omega = \tilde{\mu}$$
 and $\frac{l_{j-1} - u}{l_{j-1} - l_j} \ge 1$ on L_j . (3.77)

Taking into account that $\phi_{j-1}(x) = 1$ for $(x,t) \in Q_j$, we use (3.74), (3.70) and (3.77) to deduce

$$\frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}} \tilde{u}^{m-1} \, dx dt \leq \frac{(l_{j} - \bar{l})^{p-2} \tilde{\mu}^{m-1}}{r_{j}^{n+p}} |L_{j}|
\leq \gamma \frac{(l_{j} - \bar{l})^{p-2} \tilde{\mu}^{m-1} (t_{1} - \hat{t})}{r_{j}^{n+p}} \underset{\hat{l} < t < t_{1}}{\operatorname{ess}} \sup_{\hat{l} < t < t_{1}} \int_{B_{j-1}} \phi_{j-1}^{k} \, dx
\leq \gamma \frac{1}{r_{j-1}^{n}} \underset{\hat{l} < t < t_{1}}{\operatorname{ess}} \sup_{\hat{l} < t < t_{1}} \int_{B_{j-1}} G\left(\frac{l_{j-1} - u}{l_{j-1} - l_{j}}\right) \phi_{j-1}^{k} \, dx \leq \gamma A_{j-1}(l_{j}) \leq \gamma \chi, \tag{3.78}$$

since $Q_{j-1}(l_j) \nsubseteq \tilde{Q}_1$ and $A_{j-1}(l_j)$ is defined via (3.62) with $l = l_j$. According to (3.78), we have

$$\frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L'_{j}} \tilde{u}^{m-1} \left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \phi_{j}^{k-p} \, \mathrm{d}x \mathrm{d}t \\
\leq \frac{\tilde{\mu}^{m-1} (l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \varepsilon_{1}^{(1+\lambda)(p-1)} |L_{j}| \leq \gamma \varepsilon_{1}^{(1+\lambda)(p-1)} \chi.$$
(3.79)

On the other hand, for any fixed $\varepsilon_2 < 1$, we apply Young's inequality to conclude that

$$\begin{split} &\frac{(l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}''} \tilde{u}^{m-1} \left(\frac{l_{j}-\tilde{u}}{l_{j}-\bar{l}}\right)^{(1+\lambda)(p-1)} \phi_{j}^{k-p} \, \mathrm{d}x \mathrm{d}t \\ &\leq \varepsilon_{2} \frac{\tilde{\mu}^{m-1} (l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} |L_{j}| \\ &+ \gamma(\varepsilon_{2}) \frac{\tilde{\mu}^{m-1} (l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}''} \left(\frac{l_{j}-\tilde{u}}{l_{j}-\bar{l}}\right)^{p\frac{n+h}{nh}} \phi_{j}^{(k-p)q} \, \mathrm{d}x \mathrm{d}t \\ &=: T_{1}+T_{2}, \end{split}$$

with the obvious meanings of T_1 and T_2 . Here, parameters h and q are chosen via (3.26). In view of (3.78), we deduce that $T_1 \le \gamma \varepsilon_2 \chi$. To estimate T_2 , we set

$$\tilde{\psi}_{j}(x,t) = \frac{1}{l_{j} - \bar{l}} \left[\int_{\tilde{u}}^{l_{j}} \left(1 + \frac{l_{j} - s}{l_{j} - \bar{l}} \right)^{-\frac{1}{p} - \frac{1}{p}} ds \right]_{+}$$
(3.80)

and use Lemma 2.1 with (l, d) replaced by $(l_j, l_j - \bar{l})$ to deduce that

$$T_2 \le \gamma \frac{\tilde{\mu}^{m-1} (l_j - \bar{l})^{p-2}}{r_j^{n+p}} \iint_{L_j'} \tilde{\psi}_j^{p\frac{n+h}{n}} \phi_j^{(k-p)q} \, \mathrm{d}x \mathrm{d}t.$$
 (3.81)

To proceed further, we set $v = \tilde{\psi}_j \phi_j^{k_1}$, where $k_1 = \frac{(k-p)nq}{p(n+h)}$. Using the similar arguments as in the proof of Step 3 of Lemma 3.5, we can deduce the following estimate

$$\iint_{L''_{j}} \tilde{\psi}_{j}^{\frac{n+h}{n}} \phi_{j}^{(k-p)q} \, dxdt = \iint_{L''_{j}} v^{p+\frac{ph}{n}} \, dxdt
\leq \int_{\hat{I}}^{t_{1}} \left(\int_{B_{j}} v^{\frac{np}{n-p}} \, dx \right)^{\frac{n-p}{n}} \left(\int_{L''_{j}(t)} v^{h} \, dx \right)^{\frac{p}{n}} \, dt
\leq \gamma \operatorname{ess sup}_{\hat{I} < t < t_{1}} \left(\int_{L''_{j}(t)} \frac{l_{j} - \tilde{u}}{l_{j} - \tilde{l}} \phi_{j}^{k_{1}h} \, dx \right)^{\frac{p}{n}} \iint_{Q_{j}} |Dv|^{p} \, dxdt,$$
(3.82)

where

$$L_{j}^{\prime\prime}(t) = \left\{ x \in B_{j} : u(\cdot, t) \le l_{j} \right\} \cap \left\{ x \in B_{j} : \frac{l_{j} - \tilde{u}(\cdot, t)}{l_{i} - \bar{l}} > \varepsilon_{1} \right\}. \tag{3.83}$$

In view of (3.2), we conclude with

$$\int_{L_{j}''(t)} \frac{l_{j} - \tilde{u}}{l_{j} - \tilde{l}} \phi_{j}^{k_{1}h} dx$$

$$\leq c(\varepsilon_{1}) \int_{L_{i}''(t)} G\left(\frac{l_{j} - \tilde{u}}{l_{j} - \tilde{l}}\right) \phi_{j}^{k_{1}h} dx \leq c(\varepsilon_{1}) \int_{B_{i}} G\left(\frac{l_{j} - u}{l_{j} - \tilde{l}}\right) \phi_{j}^{k_{1}h} dx. \tag{3.84}$$

At this stage, we use Lemma 2.3 with (l, d, θ) replaced by $(l_j, l_j - \bar{l}, t_1 - \hat{t})$ to deduce that

$$\begin{split} & \operatorname*{ess\,sup}_{\hat{l} < t < t_{1}} \frac{1}{r_{j}^{n}} \int_{B_{j}} G\left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right) \phi_{j}^{k_{1}h} \, \mathrm{d}x \\ \leq & \gamma \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{p+n}} \iint_{L_{j}} u^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \phi_{j}^{k_{1}h-p} \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{t_{1} - \hat{t}}{(l_{j} - \bar{l})^{2} r_{j}^{n}} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{t_{1} - \hat{t}}{(l_{j} - \bar{l}) r_{j}^{n}} \int_{B_{j}} |f| \, \mathrm{d}x \\ & = : T_{3} + T_{4} + T_{5}, \end{split}$$

with the obvious meanings of T_3 - T_5 . We first consider the estimate for T_3 . We apply (3.78), (3.73) and (3.70) to find that

$$T_{3} \leq \gamma \frac{(l_{j} - \bar{l})^{p-2-(1+\lambda)(p-1)}}{r_{j}^{p+n}} \iint_{L_{j}} \tilde{u}^{m-1} (l_{j} - u)^{(1+\lambda)(p-1)} \phi_{j}^{k_{1}h-p} \, dxdt$$

$$\leq \gamma \frac{(l_{j-1} - l_{j})^{p-2}}{r_{j-1}^{p+n}} \iint_{L_{j-1}} \tilde{u}^{m-1} \left(\frac{l_{j-1} - u}{l_{j-1} - l_{j}}\right)^{(1+\lambda)(p-1)} \phi_{j-1}^{k-p} \, dxdt$$

$$\leq \gamma A_{j-1}(l_{j}) \leq \gamma_{1} \chi, \tag{3.85}$$

where the constant γ_1 depends only upon the data. Finally, we infer from (3.59), (3.73) and (3.74) that

$$T_4 + T_5 \leq \gamma \frac{\tilde{\mu}^{1-m}}{(l_j - \bar{l})^p} r_j^{p-n} \int_{B_j} g^{\frac{p}{p-1}} dx + \gamma \frac{\tilde{\mu}^{1-m}}{(l_j - \bar{l})^{p-1}} r_j^{p-n} \int_{B_j} |f| dx \leq \gamma (\tilde{B}^{-p} + \tilde{B}^{-(p-1)}).$$

Consequently, we infer that

$$\operatorname{ess\,sup}_{\hat{i} < t < l_1} \frac{1}{r_i^n} \int_{B_j} G\left(\frac{l_j - u}{l_j - \bar{l}}\right) \phi_j^{k_1 h} \, \mathrm{d}x \le \gamma \chi + \gamma (\tilde{B}^{-p} + \tilde{B}^{-(p-1)}). \tag{3.86}$$

We now turn our attention to the estimate of T_2 . Combining the estimates (3.81)-(3.86), we can rewrite the upper bound for T_2 by

$$T_{2} \leq \gamma \frac{\tilde{\mu}^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{j}^{n}} (\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)})^{\frac{p}{n}} \times \left[\iint_{Q_{j}} \phi_{j}^{k_{1}p} |D\tilde{\psi}_{j}|^{p} dxdt + \iint_{Q_{j}} \phi_{j}^{(k_{1}-1)p} \tilde{\psi}_{j}^{p} |D\phi_{j}|^{p} dxdt \right]$$

$$= : \gamma (\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)})^{\frac{p}{n}} (T_{6} + T_{7}),$$

with the obvious meanings of T_6 and T_7 . We first consider the estimate for T_6 . Noting that

$$\tilde{\mu}^{m-1} \le \gamma_m (\mu_-^{m-1} + (\xi \omega)^{m-1}) \le \gamma \tilde{u}^{m-1}$$
 on $\hat{Q} = B_R \times (-(\mu_+)^{1-m} A^{p-2} \omega^{2-p} R^p, 0)$, (3.87)

we conclude that

$$T_{6} = \frac{\tilde{\mu}^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{i}^{n}} \iint_{Q_{j}} \phi_{j}^{k_{1}p} |D\tilde{\psi}_{j}|^{p} dxdt \leq \gamma \frac{(l_{j} - \bar{l})^{p-2}}{r_{i}^{n}} \iint_{Q_{j}} \tilde{u}^{m-1} |D\tilde{\psi}_{j}|^{p} \phi_{j}^{k_{1}p} dxdt.$$

Taking into account that $\{\tilde{u} < l_j\} = \{u < l_j\}$, we infer from (3.80) that

$$\tilde{u}^{\frac{m-1}{p}}|D\tilde{\psi}_{j}| = u^{\frac{m-1}{p}}|Du|\frac{1}{l_{j}-\bar{l}}\left(1 + \frac{l_{j}-u}{l_{j}-\bar{l}}\right)^{-\frac{1}{p}-\frac{\lambda}{p}}\chi_{\left\{\frac{1}{128}\xi\omega < u < l_{j}\right\}} \\
\leq u^{\frac{m-1}{p}}|Du|\frac{1}{l_{j}-\bar{l}}\left(1 + \frac{l_{j}-u}{l_{j}-\bar{l}}\right)^{-\frac{1}{p}-\frac{\lambda}{p}}\chi_{\left\{u < l_{j}\right\}} = u^{\frac{m-1}{p}}|D\psi_{j}|, \tag{3.88}$$

where

$$\psi_j(x,t) = \frac{1}{l_j - \bar{l}} \left[\int_u^{l_j} \left(1 + \frac{l_j - s}{l_j - \bar{l}} \right)^{-\frac{1}{p} - \frac{\lambda}{p}} ds \right]_+.$$

At this stage, we use Lemma 2.3 with (l, d, θ) replaced by $(l_j, l_j - \bar{l}, t_1 - \hat{t})$ and taking into account the estimates for T_3 - T_5 . We conclude that

$$\begin{split} T_6 \leq & \gamma \frac{(l_j - \bar{l})^{p-2}}{r_j^n} \iint_{Q_j} u^{m-1} |D\psi_j|^p \phi_j^{k_1 p} \, \mathrm{d}x \mathrm{d}t \\ \leq & \gamma \frac{(l_j - \bar{l})^{p-2}}{r_j^p} \iint_{L_j} u^{m-1} \left(\frac{l_j - u}{l_j - \bar{l}} \right)^{(1+\lambda)(p-1)} \phi_j^{(k_1 - 1)p} \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{t_1 - \hat{t}}{(l_j - \bar{l})^2 r_j^n} \iint_{B_j} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{t_1 - \hat{t}}{(l_j - \bar{l}) r_j^n} \iint_{B_j} |f| \, \mathrm{d}x \\ \leq & \gamma \frac{(l_{j-1} - l_j)^{p-2}}{r_{j-1}^{p+n}} \iint_{L_{j-1}} \tilde{u}^{m-1} \left(\frac{l_{j-1} - u}{l_{j-1} - l_j} \right)^{(1+\lambda)(p-1)} \phi_{j-1}^{k-p} \, \mathrm{d}x \mathrm{d}t \\ & + \gamma (\tilde{B}^{-p} + \tilde{B}^{-(p-1)}) \\ \leq & \gamma (\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)}). \end{split}$$

Next, we consider the estimate for T_7 . Similar to the estimate (3.34), we have

$$\tilde{\psi}_{j}(x,t) \le \frac{(l_{j} - \tilde{u})_{p'}^{\frac{1}{p'}}}{(l_{j} - \bar{l})^{\frac{1}{p'}}}.$$
(3.89)

Consequently, we apply (3.87), (3.89) and (3.85) to conclude that

$$T_{7} \leq \frac{\tilde{\mu}^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}} \left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}}\right)^{p-1} \phi_{j}^{(k_{1}-1)p} \, \mathrm{d}x \mathrm{d}t$$

$$\leq \frac{\tilde{\mu}^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} |L_{j}|$$

$$+ \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}} \tilde{u}^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \phi_{j}^{(k_{1}-1)p} \, \mathrm{d}x \mathrm{d}t$$

$$\leq \gamma \chi.$$

Then, we arrive at $T_2 \le \gamma (\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)})^{1+\frac{p}{n}}$ and hence

$$\frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}^{"}} \tilde{u}^{m-1} \left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \phi_{j}^{k-p} \, \mathrm{d}x \mathrm{d}t
\leq \gamma \varepsilon_{2} \chi + \gamma (\varepsilon_{2}) (\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)})^{1+\frac{p}{n}}.$$
(3.90)

This yields that

$$\frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}} \tilde{u}^{m-1} \left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \phi_{j}^{k-p} \, \mathrm{d}x \mathrm{d}t
\leq \gamma \varepsilon_{1}^{(1+\lambda)(p-1)} \chi + \gamma \varepsilon_{2} \chi + \gamma (\varepsilon_{2}) (\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)})^{1+\frac{p}{n}}.$$
(3.91)

Furthermore, we apply Lemma 2.3 to obtain

$$\operatorname{ess\,sup} \frac{1}{r_{j}^{n}} \int_{B_{j}} G\left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right) \phi_{j}^{k} \, \mathrm{d}x \\
\leq \gamma \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{p+n}} \iint_{L_{j}} u^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \phi_{j}^{k-p} \, \mathrm{d}x \mathrm{d}t \\
+ \gamma \frac{t_{1} - \hat{t}}{(l_{j} - \bar{l})^{2} r_{j}^{n}} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{t_{1} - \hat{t}}{(l_{j} - \bar{l}) r_{j}^{n}} \int_{B_{j}} |f| \, \mathrm{d}x \\
= : S_{1} + S_{2} + S_{3}, \tag{3.92}$$

with the obvious meanings of S_1 - S_3 . To estimate S_1 , we use (3.91) and (3.76) to infer that

$$\begin{split} S_1 &\leq \gamma \frac{(l_j - \bar{l})^{p-2}}{r_j^{p+n}} \iint_{L_j} \tilde{u}^{m-1} \left(\frac{l_j - \tilde{u}}{l_j - \bar{l}} \right)^{(1+\lambda)(p-1)} \phi_j^{k-p} \, \mathrm{d}x \mathrm{d}t \\ &\leq \gamma \left[\varepsilon_1^{(1+\lambda)(p-1)} \chi + \varepsilon_2 \chi + \gamma(\varepsilon_2) (\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)})^{1+\frac{p}{n}} \right]. \end{split}$$

Similar to the estimates of T_4 and T_5 , we obtain $S_2 + S_3 \le \gamma(\tilde{B}^{-p} + \tilde{B}^{-(p-1)})$. Inserting the estimates for S_1 - S_3 into (3.92) and taking (3.91) into consideration, we conclude that there exist constants $\gamma_1 = \gamma_1(\text{data})$ and $\gamma_2 = \gamma_2(\text{data}, \varepsilon_1, \varepsilon_2)$ such that

$$A_{i}(\bar{l}) \leq \gamma_{1}(\tilde{B}^{-p} + \tilde{B}^{-(p-1)}) + \gamma_{1}(\varepsilon_{1} + \varepsilon_{2})\chi + \gamma_{2}(\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)})^{1 + \frac{p}{n}}.$$
 (3.93)

We now turn our attention to consider the case $Q_j(\bar{l}) \subseteq \tilde{Q}_1$. For simplicity, we may take $\varphi_{j-1} = \varphi_{j-1}(l_j)$. We decompose $L_j(\bar{l}) = L'_i(\bar{l}) \cup L''_i(\bar{l})$, where

$$L'_{j}(\bar{l}) = L_{j}(\bar{l}) \cap \left\{ \frac{l_{j} - \tilde{u}}{l_{i} - \bar{l}} \le \varepsilon_{1} \right\} \quad \text{and} \quad L''_{j}(\bar{l}) = L_{j}(\bar{l}) \setminus L'_{j}(\bar{l}). \tag{3.94}$$

Similar to (3.76), we have

$$\iint_{L_{j}(\bar{l})} \tilde{u}^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}} \right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\
\leq 2^{(1+\lambda)(p-1)} \iint_{L_{j}(\bar{l})} \tilde{u}^{m-1} \left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}} \right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \tag{3.95}$$

and hence

$$\begin{split} A_{j}(\bar{l}) \leq & \gamma \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \tilde{u}^{m-1} \left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}} \right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\ &+ \mathrm{ess} \sup_{t} \frac{1}{r_{j}^{n}} \int_{B_{j} \times \{t\}} G\left(\frac{l_{j} - u}{l_{j} - \bar{l}} \right) \varphi_{j}(\bar{l})^{k} \, \mathrm{d}x, \end{split}$$

where the constant γ depends only upon p and λ . The next thing to do in the proof is to obtain an upper bound for $A_i(\bar{l})$. To this end, we first observe that $u \leq l_i$ on $L_i(\bar{l})$ and hence

$$\frac{l_{j-1} - u}{l_{j-1} - l_j} \ge 1 \qquad \text{on} \quad L_j(\bar{l}).$$

In the case $Q_{j-1}(l_j) \nsubseteq \tilde{Q}_1$, we have $Q_j(\bar{l}) \subset Q_{j-1}$ and $\phi_{j-1}(x) = 1$ for $(x, t) \in Q_j(\bar{l})$. Then, we use (3.70) to deduce

$$\frac{(l_{j} - \bar{l})^{p-2} \tilde{\mu}^{m-1}}{r_{j}^{n+p}} |L_{j}(\bar{l})| \leq \gamma \frac{1}{r_{j-1}^{n}} \underset{\hat{l} < t < l_{1}}{\text{ess sup}} \int_{B_{j-1}} G\left(\frac{l_{j-1} - u}{l_{j-1} - l_{j}}\right) \phi_{j-1}^{k} \, \mathrm{d}x$$
$$\leq \gamma A_{j-1}(l_{j}) \leq \gamma \chi.$$

Moreover, we apply (3.70) and (3.73) to conclude that

$$\begin{split} &\frac{(l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \tilde{u}^{m-1} \left(\frac{l_{j}-u}{l_{j}-\bar{l}}\right)^{(1+\lambda)(p-1)} \, \mathrm{d}x \mathrm{d}t \\ &\leq \gamma \frac{(l_{j-1}-l_{j})^{p-2-(1+\lambda)(p-1)}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \tilde{u}^{m-1} (l_{j-1}-u)^{(1+\lambda)(p-1)} \, \mathrm{d}x \mathrm{d}t \\ &\leq \gamma \frac{(l_{j-1}-l_{j})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j-1}} \tilde{u}^{m-1} \left(\frac{l_{j-1}-u}{l_{j-1}-l_{j}}\right)^{(1+\lambda)(p-1)} \phi_{j-1}^{k-p} \, \mathrm{d}x \mathrm{d}t \\ &\leq \gamma A_{j-1}(l_{j}) \leq \gamma \chi. \end{split}$$

Next, we consider the case $Q_{j-1}(l_j) \subseteq \tilde{Q}_1$. In view of (3.73), we see that $\tilde{\mu}^{1-m}(l_j-\bar{l})^{2-p}r_j^p \le \frac{1}{16}\tilde{\mu}^{1-m}(l_{j-1}-l_j)^{2-p}r_{j-1}^p$. This yields that $Q_j(\bar{l}) \subset Q_{j-1}(l_j)$ and $\varphi_{j-1}(x,t) = 1$ for $(x,t) \in Q_j(\bar{l})$. It follows that

$$\frac{(l_{j} - \bar{l})^{p-2} \tilde{\mu}^{m-1}}{r_{j}^{n+p}} |L_{j}(\bar{l})| \le \gamma \frac{1}{r_{j-1}^{n}} \operatorname{ess \, sup} \int_{B_{j-1}} G\left(\frac{l_{j-1} - u}{l_{j-1} - l_{j}}\right) \varphi_{j-1}^{k} \, \mathrm{d}x$$

$$\le \gamma A_{j-1}(l_{j}) \le \gamma \chi$$

and

$$\begin{split} \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} & \iint_{L_{j}(\bar{l})} \tilde{u}^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \, \mathrm{d}x \mathrm{d}t \\ & \leq \gamma \frac{(l_{j-1} - l_{j})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j-1}(l_{j})} \tilde{u}^{m-1} \left(\frac{l_{j-1} - u}{l_{j-1} - l_{j}}\right)^{(1+\lambda)(p-1)} \varphi_{j-1}^{k-p} \, \mathrm{d}x \mathrm{d}t \\ & \leq \gamma A_{j-1}(l_{j}) \leq \gamma \chi. \end{split}$$

Consequently, we conclude that the estimates

$$\frac{(l_j - \bar{l})^{p-2} \tilde{\mu}^{m-1}}{r_j^{n+p}} |L_j(\bar{l})| \le \gamma \chi$$
 (3.96)

and

$$\frac{(l_j - \bar{l})^{p-2}}{r_j^{n+p}} \iint_{L_j(\bar{l})} \tilde{u}^{m-1} \left(\frac{l_j - u}{l_j - \bar{l}}\right)^{(1+\lambda)(p-1)} dxdt \le \gamma \chi \tag{3.97}$$

hold for either $Q_{j-1}(l_j) \nsubseteq \tilde{Q}_1$ or $Q_{j-1}(l_j) \subseteq \tilde{Q}_1$. According to (3.96) and (3.77), we deduce that

$$\frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L'_{j}(\bar{l})} \tilde{u}^{m-1} \left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t$$

$$\leq \frac{\tilde{\mu}^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \varepsilon_{1}^{(1+\lambda)(p-1)} |L_{j}(\bar{l})| \leq \gamma \varepsilon_{1}^{(1+\lambda)(p-1)} \chi. \tag{3.98}$$

Moreover, for any fixed $\varepsilon_2 < 1$, we apply Young's inequality and (3.77) to conclude that

$$\begin{split} &\frac{(l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}^{\prime\prime}(\bar{l})} \tilde{u}^{m-1} \left(\frac{l_{j}-\tilde{u}}{l_{j}-\bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\ &\leq \varepsilon_{2} \frac{\tilde{\mu}^{m-1}(l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} |L_{j}(\bar{l})| \\ &+ \gamma(\varepsilon_{2}) \frac{\tilde{\mu}^{m-1}(l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}^{\prime\prime}(\bar{l})} \left(\frac{l_{j}-\tilde{u}}{l_{j}-\bar{l}}\right)^{p\frac{n+h}{nh}} \varphi_{j}(\bar{l})^{(k-p)q} \, \mathrm{d}x \mathrm{d}t \\ &=: \tilde{T}_{1} + \tilde{T}_{2}, \end{split}$$

with the obvious meanings of \tilde{T}_1 and \tilde{T}_2 . In view of (3.96), we obtain $\tilde{T}_1 \leq \gamma \varepsilon_2 \chi$. Next, we use Lemma 2.1 with (l,d) replaced by $(l_j,l_j-\bar{l})$ to deduce that

$$\tilde{T}_{2} \leq \gamma \frac{\tilde{\mu}^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}^{"}(\bar{l})} \tilde{\psi}_{j}^{p\frac{n+h}{n}} \varphi_{j}(\bar{l})^{(k-p)q} \, \mathrm{d}x \mathrm{d}t, \tag{3.99}$$

where $\tilde{\psi}_j$ is the function defined in (3.80). Let $v = \tilde{\psi}_j \varphi_j^{k_1}$, where $k_1 = \frac{(k-p)nq}{p(n+h)}$. We adopt the same procedure as in the proof of Step 3 of Lemma 3.5. This yields that

$$\iint_{L''_{j}(\bar{l})} \tilde{\psi}_{j}^{p\frac{n+h}{n}} \varphi_{j}(\bar{l})^{(k-p)q} \, dxdt \\
\leq \int_{t_{1}-(\mu_{+})^{1-m}(l_{j}-\bar{l})^{2-p}r_{j}^{p}}^{t_{1}} \left(\int_{B_{j}} v^{\frac{np}{n-p}} \, dx\right)^{\frac{n-p}{n}} \left(\int_{L''_{j}(t)} v^{h} \, dx\right)^{\frac{p}{n}} \, dt \\
\leq \gamma \operatorname{ess} \sup_{t} \left(\int_{L''_{j}(t)} \frac{l_{j}-\tilde{u}}{l_{j}-\bar{l}} \varphi_{j}(\bar{l})^{k_{1}h} \, dx\right)^{\frac{p}{n}} \iint_{Q_{j}(\bar{l})} |Dv|^{p} \, dxdt, \tag{3.100}$$

where $L_i''(t)$ is the set defined in (3.83). According to Lemma 3.4, we deduce that

$$\int_{L_i''(t)} \frac{l_j - \tilde{u}}{l_j - \bar{l}} \varphi_j(\bar{l})^{k_1 h} \, \mathrm{d}x \le c(\varepsilon_1) \int_{B_j} G\left(\frac{l_j - u}{l_j - \bar{l}}\right) \varphi_j(\bar{l})^{k_1 h} \, \mathrm{d}x. \tag{3.101}$$

At this point, we apply Lemma 2.3 with (l, d, θ) replaced by $(l_j, l_j - \bar{l}, \tilde{\mu}^{1-m}(l_j - \bar{l})^{2-p}r_j^p)$ to conclude that

$$\begin{split} & \operatorname{ess\,sup} \frac{1}{r_{j}^{n}} \int_{B_{j}} G\left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right) \varphi_{j}(\bar{l})^{k_{1}h} \, \mathrm{d}x \\ \leq & \gamma \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{p+n}} \iint_{L_{j}(\bar{l})} u^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k_{1}h - p} \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{1}{r_{j}^{n}} \iint_{L_{j}(\bar{l})} \frac{l_{j} - u}{l_{j} - \bar{l}} |\partial_{t}\varphi_{j}(\bar{l})| \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{r_{j}^{p-n}}{(l_{j} - \bar{l})^{p}} \tilde{\mu}^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{r_{j}^{p-n}}{(l_{j} - \bar{l})^{p-1}} \tilde{\mu}^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x \\ & = : \tilde{T}_{3} + \tilde{T}_{4} + \tilde{T}_{5} + \tilde{T}_{6}, \end{split}$$

with the obvious meanings of \tilde{T}_3 - \tilde{T}_6 . In view of (3.97), we find that $\tilde{T}_3 \leq \gamma \chi$, since $u \leq \tilde{u}$ on $L_j(\bar{l})$. Noting that $(1 + \lambda)(p - 1) > 1$ and $|\partial_t \varphi_j(\bar{l})| \leq 9(l_j - \bar{l})^{p-2} \tilde{\mu}^{m-1} r_j^{-p}$, we infer from (3.87), (3.96) and (3.97) that

$$\begin{split} \tilde{T}_{4} \leq & \gamma \frac{(l_{j} - \bar{l})^{p-2} \tilde{\mu}^{m-1}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \frac{l_{j} - u}{l_{j} - \bar{l}} \, \mathrm{d}x \mathrm{d}t \\ \leq & \gamma \frac{(l_{j} - \bar{l})^{p-2} \tilde{\mu}^{m-1}}{r_{j}^{n+p}} |L_{j}(\bar{l})| + \gamma \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \tilde{u}^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \, \mathrm{d}x \mathrm{d}t \\ \leq & \gamma \chi + A_{i-1}(l_{i}) \leq \gamma_{2} \chi, \end{split}$$

where the constant γ_2 depends only upon the data. Finally, we infer from (3.59) and (3.73) that $\tilde{T}_5 + \tilde{T}_6 \leq \gamma(\tilde{B}^{-p} + \tilde{B}^{-(p-1)})$. Combining the above estimates, we arrive at

$$\operatorname{ess\,sup} \frac{1}{r_i^n} \int_{B_j} G\left(\frac{l_j - u}{l_i - \bar{l}}\right) \varphi_j(\bar{l})^{k_1 h} \, \mathrm{d}x \le \gamma \chi + \gamma (\tilde{B}^{-p} + \tilde{B}^{-(p-1)}). \tag{3.102}$$

We now turn our attention to the estimate of \tilde{T}_2 . Combining (3.28)-(3.32), we can rewrite the upper bound for \tilde{T}_2 by

$$\begin{split} \tilde{T}_{2} \leq & \gamma \frac{\tilde{\mu}^{m-1} (l_{j} - \bar{l})^{p-2}}{r_{j}^{n}} (\chi + B^{-p} + B^{-(p-1)})^{\frac{p}{n}} \\ & \times \left[\iint_{Q_{j}(\bar{l})} \varphi_{j}(\bar{l})^{k_{1}p} |D\tilde{\psi}_{j}|^{p} \, \mathrm{d}x \mathrm{d}t + \iint_{Q_{j}(\bar{l})} \varphi_{j}(\bar{l})^{(k_{1}-1)p} \tilde{\psi}_{j}^{p} |D\varphi_{j}|^{p} \, \mathrm{d}x \mathrm{d}t \right] \\ & = : \gamma (\chi + B^{-p} + B^{-(p-1)})^{\frac{p}{n}} (\tilde{T}_{7} + \tilde{T}_{8}), \end{split}$$

with the obvious meanings of \tilde{T}_7 and \tilde{T}_8 . We first consider the estimate for \tilde{T}_7 . In view of (3.87) and (3.88), we see that

$$\begin{split} \tilde{T}_7 &\leq \gamma \frac{(l_j - \bar{l})^{p-2}}{r_j^n} \iint_{Q_j(\bar{l})} \tilde{u}^{m-1} |D\tilde{\psi}_j|^p \varphi_j(\bar{l})^{k_1 p} \, \mathrm{d}x \mathrm{d}t \\ &\leq \gamma \frac{(l_j - \bar{l})^{p-2}}{r_i^n} \iint_{Q_j(\bar{l})} u^{m-1} |D\psi_j|^p \varphi_j(\bar{l})^{k_1 p} \, \mathrm{d}x \mathrm{d}t. \end{split}$$

To proceed further, we use Lemma 2.3 and take into account the estimates for \tilde{T}_3 - \tilde{T}_6 . We proceed to estimate \tilde{T}_7 by

$$\begin{split} \tilde{T}_{7} \leq & \gamma \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{p}} \iint_{L_{j}(\bar{l})} u^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}} \right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{(k_{1}-1)p} \, \mathrm{d}x \mathrm{d}t \\ &+ \gamma \iint_{L_{j}(\bar{l})} \frac{l_{j} - u}{l_{j} - \bar{l}} |\partial_{t} \varphi_{j}(\bar{l})| \, \mathrm{d}x \mathrm{d}t + \gamma \frac{r_{j}^{p}}{(l_{j} - \bar{l})^{p}} \tilde{\mu}^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x \\ &+ \gamma \frac{r_{j}^{p}}{(l_{j} - \bar{l})^{p-1}} \tilde{\mu}^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x \\ \leq & \gamma (\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)}), \end{split}$$

where the constant γ depends only upon the data. To estimate \tilde{T}_8 , we use (3.89), (3.87), (3.96), (3.97) and Young's inequality to obtain

$$\begin{split} \tilde{T}_{8} &\leq \frac{\tilde{\mu}^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}}\right)^{p-1} \varphi_{j}(\bar{l})^{(k_{1}-1)p} \, \mathrm{d}x \mathrm{d}t \\ &\leq \frac{\tilde{\mu}^{m-1}(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} |L_{j}(\bar{l})| \\ &+ \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \tilde{u}^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{(k_{1}-1)p} \, \mathrm{d}x \mathrm{d}t \\ &\leq \gamma \chi \end{split}$$

and hence we arrive at $\tilde{T}_2 \le \gamma (\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)})^{1+\frac{p}{n}}$. This also yields that

$$\frac{\tilde{\mu}^{m-1}(l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}''(\bar{l})} \left(\frac{l_{j}-\tilde{u}}{l_{j}-\bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \,\mathrm{d}x\mathrm{d}t$$

$$\leq \gamma \varepsilon_{2} \chi + \gamma(\varepsilon_{2})(\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)})^{1+\frac{p}{n}}.$$
(3.103)

Moreover, we conclude with

$$\frac{\tilde{\mu}^{m-1}(l_{j}-\bar{l})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \left(\frac{l_{j}-\tilde{u}}{l_{j}-\bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\
\leq \gamma \varepsilon_{1}^{(1+\lambda)(p-1)} \chi + \gamma \varepsilon_{2} \chi + \gamma (\varepsilon_{2}) \eta^{1-m} (\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)})^{1+\frac{p}{n}}.$$
(3.104)

In order to derive an upper bound for $A_j(\bar{l})$, we need to estimate the second term on the right-hand side of (3.61). To this end, we apply Lemma 2.3 with (l, d, θ) replaced by $(l_j, l_j - \bar{l}, \tilde{\mu}^{1-m}(l_j - \bar{l})^{2-p}r_i^p)$ to obtain

$$\operatorname{ess\,sup} \frac{1}{r_{j}^{n}} \int_{B_{j}} G\left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right) \varphi_{j}(\bar{l})^{k} \, \mathrm{d}x$$

$$\leq \gamma \frac{(l_{j} - \bar{l})^{p-2}}{r_{j}^{p+n}} \iint_{L_{j}(\bar{l})} u^{m-1} \left(\frac{l_{j} - u}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t$$

$$+ \gamma \frac{1}{r_{j}^{n}} \iint_{L_{j}(\bar{l})} \frac{l_{j} - u}{l_{j} - \bar{l}} |\partial_{t} \varphi_{j}(\bar{l})| \varphi_{j}(\bar{l})^{k-1} \, \mathrm{d}x \mathrm{d}t$$

$$+ \gamma \frac{r_{j}^{p-n}}{(l_{j} - \bar{l})^{p}} \tilde{\mu}^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{r_{j}^{p-n}}{(l_{j} - \bar{l})^{p-1}} \tilde{\mu}^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x$$

$$= : \tilde{S}_{1} + \tilde{S}_{2} + \tilde{S}_{3} + \tilde{S}_{4},$$
(3.105)

with the obvious meanings of \tilde{S}_1 - \tilde{S}_4 . To estimate \tilde{S}_1 , we apply (3.104) and (3.95) to deduce

$$\begin{split} \tilde{S}_{1} & \leq 2^{(1+\lambda)(p-1)} \gamma \frac{(l_{j} - \bar{l})^{p-2} \tilde{\mu}^{m-1}}{r_{j}^{p+n}} \iint_{L_{j}(\bar{l})} \left(\frac{l_{j} - u}{l_{j} - \bar{l}} \right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\ & \leq 2^{(1+\lambda)(p-1)} \gamma \left[\varepsilon_{1}^{(1+\lambda)(p-1)} \chi + \varepsilon_{2} \chi + \gamma(\varepsilon_{2}) \eta^{1-m} (\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)})^{1+\frac{p}{n}} \right]. \end{split}$$

Next, we consider the estimate for \tilde{S}_2 . Similar to (3.76), we note from $l_j - \frac{1}{128}\xi\omega > \frac{1}{2}l_j$ that

$$\begin{split} \tilde{S}_2 = & \gamma \frac{1}{r_j^n} \iint_{L_j(\bar{l}) \cap \left\{u > \frac{1}{128} \xi \omega\right\}} \frac{l_j - \tilde{u}}{l_j - \bar{l}} |\partial_t \varphi_j(\bar{l})| \varphi_j(\bar{l})^{k-1} \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{1}{r_j^n} \iint_{L_j(\bar{l}) \cap \left\{u \leq \frac{1}{128} \xi \omega\right\}} \frac{l_j}{l_j - \bar{l}} |\partial_t \varphi_j(\bar{l})| \varphi_j(\bar{l})^{k-1} \, \mathrm{d}x \mathrm{d}t \\ \leq & 2\gamma \frac{1}{r_i^n} \iint_{L_j(\bar{l})} \frac{l_j - \tilde{u}}{l_j - \bar{l}} |\partial_t \varphi_j(\bar{l})| \varphi_j(\bar{l})^{k-1} \, \mathrm{d}x \mathrm{d}t. \end{split}$$

Furthermore, we decompose $L_j(\bar{l}) = L'_j(\bar{l}) \cup L''_j(\bar{l})$, where $L'_j(\bar{l})$ and $L''_j(\bar{l})$ satisfy (3.94). In view of $|\partial_t \varphi_j(\bar{l})| \le 9(l_j - \bar{l})^{p-2} \tilde{\mu}^{m-1} r_j^{-p}$, we use (3.96) and (3.103) to conclude that

$$\begin{split} \tilde{S}_{2} \leq & \gamma \frac{(l_{j} - \bar{l})^{p-2} \tilde{\mu}^{m-1}}{r_{j}^{n+p}} \iint_{L_{j}(\bar{l})} \frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}} \varphi_{j}(\bar{l})^{k-1} \, \mathrm{d}x \mathrm{d}t \\ \leq & \gamma \varepsilon_{1} \frac{\tilde{\mu}^{m-1} (l_{j} - \bar{l})^{p-2}}{r_{j}^{n+p}} |L'_{j}(\bar{l})| \\ & + \gamma \frac{(l_{j} - \bar{l})^{p-2} \tilde{\mu}^{m-1}}{r_{j}^{p+n}} \iint_{L'_{j}(\bar{l})} \left(\frac{l_{j} - \tilde{u}}{l_{j} - \bar{l}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\ \leq & \varepsilon_{1} \chi + \gamma \left[\varepsilon_{2} \chi + \gamma(\varepsilon_{2})(\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)})^{1+\frac{p}{n}}\right]. \end{split}$$

Similar to the estimates of \tilde{T}_5 and \tilde{T}_6 , we see that $\tilde{S}_3 + \tilde{S}_4 \leq \gamma(\tilde{B}^{-p} + \tilde{B}^{-(p-1)})$. Next, we substitute the estimates for S_1 - S_4 into (3.37). Taking into account (3.36), we infer that there exist constants $\gamma_1' = \gamma_1'(\text{data})$ and $\gamma_2' = \gamma_2'(\text{data}, \varepsilon_1, \varepsilon_2)$ such that

$$A_{j}(\bar{l}) \leq \gamma'_{1}(\tilde{B}^{-p} + \tilde{B}^{-(p-1)}) + \gamma'_{1}(\varepsilon_{1} + \varepsilon_{2})\chi + \gamma'_{2}(\chi + \tilde{B}^{-p} + \tilde{B}^{-(p-1)})^{1 + \frac{p}{n}}.$$

At this point, we first choose ε_1 and ε_2 be such that

$$\varepsilon_1 = \varepsilon_2 = \frac{1}{2} \min \left\{ \frac{1}{8\gamma_1}, \frac{1}{8\gamma_1'} \right\}$$
 (3.106)

and then we determine the value of χ by

$$\chi = \frac{1}{2} \min \left\{ \frac{1}{100^{\frac{n}{p}} \gamma_2^{\frac{n}{p}}}, \frac{1}{100^{\frac{n}{p}} (\gamma_2')^{\frac{n}{p}}} \right\}, \tag{3.107}$$

where γ_1 and γ_2 are the constants in the estimate (3.93). Finally, with the choices of ε_1 , ε_2 and χ , we set B so large that

$$\tilde{B}^{-p} + \tilde{B}^{1-p} < \min \left\{ \frac{1}{100\gamma_1} \chi, \left(\frac{1}{100\gamma_2} \chi \right)^{\frac{1}{1+\frac{p}{n}}}, \frac{1}{100\gamma_1'} \chi, \left(\frac{1}{100\gamma_2'} \chi \right)^{\frac{1}{1+\frac{p}{n}}} \right\}.$$

With the choices of ε_1 , ε_2 , χ and \tilde{B} , we get $A_j(\bar{l}) \leq \frac{1}{2}\chi$ holds for either $Q_j(\bar{l}) \nsubseteq \tilde{Q}_1$ or $Q_j(\bar{l}) \subseteq \tilde{Q}_1$, which completes the proof of (3.72). Our next goal is to determine the value of l_{j+1} . We first consider the case

$$\iint_{B_{r_i}(x_1)\times\left\{t_1-r_i^p< t< t_1\right\}} \tilde{u}^{m-1}(l_j-u)_+^{(1+\lambda)(p-1)}\,\mathrm{d}x\,\mathrm{d}t=0.$$

According to Lemma 3.3 and (3.71), we conclude that

$$u(x_1, t_1) \ge l_j \ge \mu_- + \frac{1}{2}\tilde{B}\alpha_{i-1} + \frac{1}{4}\xi\omega > \mu_- + \frac{1}{4}\xi\omega,$$

which proves (3.55). We now consider the case

$$\iint_{B_{r_j}(x_1)\times\left\{t_1-r_j^p< t< t_1\right\}} \tilde{u}^{m-1}(l_j-u)_+^{(1+\lambda)(p-1)}\,\mathrm{d}x\,\mathrm{d}t>0.$$

It follows that $A_j(l) \to +\infty$ as $l \to l_j$. Since $A_j(l)$ is continuous and increasing, we infer that there exists a number $\tilde{l} \in (\bar{l}, l_j)$ such that $A_j(\tilde{l}) = \chi$. At this point, we set

$$l_{j+1} = \begin{cases} \tilde{l}, & \text{if } \tilde{l} < l_j - \frac{1}{4}(\alpha_{j-1} - \alpha_j), \\ l_j - \frac{1}{4}(\alpha_{j-1} - \alpha_j), & \text{if } \tilde{l} \ge l_j - \frac{1}{4}(\alpha_{j-1} - \alpha_j). \end{cases}$$
(3.108)

Recalling from (3.73) that $l_j - \bar{l} > \frac{1}{4}\tilde{B}(\alpha_{j-1} - \alpha_j) > \frac{1}{4}(\alpha_{j-1} - \alpha_j)$, we find that the definition of l_{j+1} is justified.

Step 4: *Proof of the estimate* (3.55). For simplicity of notation, we write $Q_j = Q_j(l_{j+1})$, $L_j = L_j(l_{j+1})$ and $d_j = l_j - l_{j+1}$. According to (3.108), we find that (3.69) and (3.70) hold with i = j + 1. In view of (3.71), we conclude from $\alpha_{j-1} \ge \alpha_j$ that

$$\begin{split} l_{j+1} > \bar{l} &= \frac{1}{2} l_j + \frac{1}{4} \tilde{B} \alpha_j + \frac{1}{8} \xi \omega + \frac{1}{2} \mu_- \\ &> \frac{1}{2} \left(\mu_- + \frac{1}{2} \tilde{B} \alpha_{j-1} + \frac{1}{4} \xi \omega \right) + \frac{1}{4} \tilde{B} \alpha_j + \frac{1}{8} \xi \omega + \frac{1}{2} \mu_- \\ &= \mu_- + \frac{1}{4} \tilde{B} (\alpha_j + \alpha_{j-1}) + \frac{1}{4} \xi \omega \ge \mu_- + \frac{1}{2} \tilde{B} \alpha_j + \frac{1}{4} \xi \omega. \end{split}$$

This yields that $(3.71)_{j+1}$ holds. Repeating the arguments as in Step 3, we construct l_{j+2} and hence we can determine a sequence of numbers $\{l_i\}_{i=0}^{\infty}$ satisfying (3.69)-(3.71). Noting that the sequence $\{l_i\}_{i=0}^{\infty}$ is decreasing, we infer from (3.71) that the limitation of l_i exists. This also yields that $d_i \to 0$ as $i \to \infty$. Define

$$\hat{l} = \lim_{i \to \infty} l_i$$

and we assert that $\hat{l} = u(x_1, t_1)$. Recalling that (3.70) holds for any $i = 1, 2, \dots$, we deduce that

$$\begin{split} \frac{1}{r_{i}^{n+p}} & \iint_{B_{r_{i}}(x_{1}) \times \left\{t_{1} - r_{i}^{p} < t < t_{1}\right\}} (\hat{l} - u)_{+}^{(1+\lambda)(p-1)} \, \mathrm{d}x \, \mathrm{d}t \leq \frac{4^{n+p}}{r_{i-1}^{n+p}} \iint_{L_{i-1}} (l_{i-1} - u)^{(1+\lambda)(p-1)} \, \varphi_{i-1}^{k-p} \, \mathrm{d}x \mathrm{d}t \\ & \leq 4^{n+p} (\xi \omega)^{1-m} A_{i-1}(l_{i}) d_{i-1}^{(1+\lambda)(p-1)-(p-2)} \leq 4^{n+p} (\xi \omega)^{1-m} \chi d_{i-1}^{(1+\lambda)(p-1)-(p-2)} \to 0 \end{split}$$

as $i \to \infty$. By Lemma 3.3, we have $\hat{l} = u(x_1, t_1)$. Next, we claim that for any $j \ge 1$ there holds

$$d_{j} \leq \frac{1}{4} d_{j-1} + \gamma \left(r_{j-1}^{p-n} \tilde{\mu}^{1-m} \int_{B_{j-1}} g(y)^{\frac{p}{p-1}} dy \right)^{\frac{1}{p}} + \gamma \left(r_{j-1}^{p-n} \tilde{\mu}^{1-m} \int_{B_{j-1}} |f(y)| dy \right)^{\frac{1}{p-1}}.$$
(3.109)

To start with, for any fixed $j \ge 1$, we first assume that

$$d_j > \frac{1}{4}d_{j-1}$$
 and $d_j > \frac{1}{4}(\alpha_{j-1} - \alpha_j),$ (3.110)

since otherwise (3.43) holds immediately. In view of $d_j > \frac{1}{4}(\alpha_{j-1} - \alpha_j)$, we infer from (3.42) that $A_j(l_{j+1}) = A_j(\tilde{l}) = \chi$. According to (3.110), we repeat the arguments from Step 3 and this gives

$$\begin{split} \chi &= A_{j}(l_{j+1}) \leq \gamma \frac{r_{j}^{p-n}}{d_{j}^{p}} \tilde{\mu}^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{r_{j}^{p-n}}{d_{j}^{p-1}} \tilde{\mu}^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x \\ &+ \gamma(\varepsilon_{1} + \varepsilon_{2}) \chi + \gamma(\varepsilon_{2}) \chi^{1+\frac{p}{n}} \\ &+ \gamma(\varepsilon_{2}) \left[\frac{r_{j}^{p-n}}{d_{j}^{p}} \tilde{\mu}^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{r_{j}^{p-n}}{d_{j}^{p-1}} \tilde{\mu}^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x \right]^{1+\frac{p}{n}}. \end{split}$$

According to the choices of ε_1 , ε_2 and χ in (3.106)-(3.107), we conclude with

$$(\varepsilon_1 + \varepsilon_2)\chi + \gamma(\varepsilon_2)\chi^{1+\frac{p}{n}} \leq \frac{1}{4}\chi.$$

Consequently, we infer that either

$$d_j \le \gamma \left(r_j^{p-n} \tilde{\mu}^{1-m} \int_{B_j} g^{\frac{p}{p-1}} \, \mathrm{d}x \right)^{\frac{1}{p}} \qquad \text{or} \qquad d_j \le \gamma \left(r_j^{p-n} \tilde{\mu}^{1-m} \int_{B_j} |f| \, \mathrm{d}x \right)^{\frac{1}{p-1}},$$

which proves the inequality (3.109). Let J > 1 be a fixed integer. We sum up the inequality (3.109) for $j = 1, \dots, J - 1$ and obtain

$$\begin{split} l_1 - l_J &\leq \frac{1}{3} d_0 + \gamma \sum_{j=1}^{J-1} \left(r_{j-1}^{p-n} \tilde{\mu}^{1-m} \int_{B_{j-1}} |f(y)| \, \mathrm{d}y \right)^{\frac{1}{p-1}} \\ &+ \gamma \sum_{j=1}^{J-1} \left(r_{j-1}^{p-n} \tilde{\mu}^{1-m} \int_{B_{j-1}} g(y)^{\frac{p}{p-1}} \, \mathrm{d}y \right)^{\frac{1}{p}} \, . \end{split}$$

Recalling that $d_0 = l_0 - l_1$ and $l_0 = \mu_- + \xi \omega$, we apply (3.57) to obtain

$$\mu_{-} + \xi \omega \leq \frac{4}{3} d_{0} + l_{J} + \gamma \tilde{\mu}^{\frac{1-m}{p-1}} \int_{0}^{2R} \left(\frac{1}{r^{n-p}} \int_{B_{r}(x_{1})} f(y) \, \mathrm{d}y \right)^{\frac{1}{p-1}} \frac{1}{r} \, \mathrm{d}r$$

$$+ \gamma \tilde{\mu}^{\frac{1-m}{p}} \int_{0}^{2R} \left(\frac{1}{r^{n-p}} \int_{B_{r}(x_{1})} g(y)^{\frac{p}{p-1}} \, \mathrm{d}y \right)^{\frac{1}{p}} \frac{1}{r} \, \mathrm{d}r$$

$$\leq \frac{4}{3} d_{0} + l_{J} + \gamma \frac{1}{\tilde{B}} \xi \omega. \tag{3.111}$$

Taking into account that $l_1 \ge \bar{l}$ where $\bar{l} = \frac{1}{2}l_0 + \frac{1}{2}\mu_- + \frac{1}{4}\tilde{B}\alpha_0 + \frac{1}{8}\xi\omega$. We infer from (3.63) that $d_0 \le l_0 - \bar{l} \le \frac{3}{8}\xi\omega$. Passing to the limit $J \to \infty$, we conclude from (3.111) that

$$u(x_1, t_1) > \mu_- + \frac{1}{2}\xi\omega,$$

provided that we choose $\tilde{B} > 4\gamma$. This shows that the inequality (3.55) holds for almost everywhere point in \tilde{Q}_2 , since (x_1, t_1) is the Lebesgue point of u. The constant v_1 is determined via (3.66), where the quantity χ in (3.66) is fixed in terms of (3.107). We have thus proved the lemma.

With the help of the proceeding lemmas we can now establish a decay estimate for the oscillation of the weak solution in a smaller cylinder, and the following proposition is our main result in this section.

Proposition 3.8. Let $\tilde{Q}_0 = Q\left(\frac{1}{16}R, (\mu_+)^{1-m}\omega^{2-p}\left(\frac{1}{16}R\right)^p\right)$ and let u be a bounded nonnegative weak solution to (1.1)-(1.2) in Ω_T . There exist $0 < \xi_1 < 2^{-5}$ and $B_1 > 1$ depending only upon the data and A such that

$$\operatorname*{ess\,osc}_{\tilde{O}_{0}}u\leq (1-2^{-1}\xi_{1})\omega+B_{1}\xi_{1}^{-1}(F_{1}(2R)^{\frac{p}{p+m-1}}+F_{2}(2R)^{\frac{p-1}{p+m-2}}).$$

Proof. To start with, we first assume that (3.5), (3.47) and (3.56) are violated. We take $v_* = v_1$ according to (3.66) in Lemma 3.6. Moreover, we choose $s_* = 2v_*^{-1}A^{p-2}$ and $\xi_1 = 2^{-s_*}$. It follows from Lemma 3.6 that

$$\left| \left\{ (x,t) \in \tilde{Q}_1 : u < \mu_- + \xi_1 \omega \right\} \right| \le \nu_1 |\tilde{Q}_1|,$$

where $\hat{Q}_1 = B_{\frac{R}{8}} \times (\hat{t}, 0)$. This implies that the condition for Lemma 3.7 is satisfied and it follows that

$$\underset{\tilde{O}_0}{\text{ess osc }} u \le (1 - 2^{-1} \xi_1) \omega.$$

On the other hand, if either (3.5), (3.47) or (3.56) holds, then we conclude that either $\omega \leq (\mu_+)^{\frac{1-m}{p}} F_1(2R), \ \omega \leq (\mu_+)^{\frac{1-m}{p-1}} F_2(2R), \ \omega \leq \xi_1^{-\frac{2}{p}} (\mu_+)^{\frac{1-m}{p}} F_1(2R), \ \omega \leq \xi_1^{-\frac{1}{p-1}} (\mu_+)^{\frac{1-m}{p-1}} F_2(2R),$

$$\xi_1\omega \leq \tilde{B}(\mu_- + \xi_1\omega)^{\frac{1-m}{p}}F_1(2R)$$
 or $\xi_1\omega \leq \tilde{B}(\mu_- + \xi_1\omega)^{\frac{1-m}{p-1}}F_2(2R)$.

Noting that $\mu_+ \geq \frac{1}{2}\omega$, $(\mu_- + \xi_1\omega)^{\frac{1-m}{p}} \leq (\xi_1\omega)^{\frac{1-m}{p}}$ and $(\mu_- + \xi_1\omega)^{\frac{1-m}{p-1}} \leq (\xi_1\omega)^{\frac{1-m}{p-1}}$, we deduce that the inequality

$$\operatorname{ess \ osc}_{\tilde{O}_0} u \le \omega \le B_1 \xi_1^{-1} (F_1(2R)^{\frac{p}{p+m-1}} + F_2(2R)^{\frac{p-1}{p+m-2}}).$$

holds for a constant $B_1 = B_1(\text{data}, A) > 1$. It is now obvious that the proposition holds. \Box

4. The second alternative

In this section, we will establish the decay estimate of the essential oscillation for the second alternative. We begin with the following lemma which is a standard result that can be found in [13, Lemma 7.1].

Lemma 4.1. Let $-(A^{p-2}-1)(\mu_+)^{1-m}\omega^{2-p}R^p \le \bar{t} \le 0$ and let u be a bounded nonnegative weak solution to (1.1)-(1.2) in Ω_T . Then there exists a time level

$$t^* \in \left[\bar{t} - (\mu_+)^{1-m} \omega^{2-p} \left(\frac{3}{4} R \right)^p, \bar{t} - \frac{1}{2} \nu_0(\mu_+)^{1-m} \omega^{2-p} \left(\frac{3}{4} R \right)^p \right]$$

such that

$$\left| \left\{ x \in B_{\frac{3}{4}R} : u(x, t^*) > \mu_+ - \frac{\omega}{4} \right\} \right| \le \left(\frac{1 - \nu_0}{1 - \frac{1}{2}\nu_0} \right) |B_{\frac{3}{4}R}|, \tag{4.1}$$

where v_0 is the constant claimed by Lemma 3.5.

Next. we provide the following lemma regarding the time propagation of positivity.

Lemma 4.2. Let u be a bounded nonnegative weak solution to (1.1)-(1.2) in Ω_T . There exists a positive constant s_1 that can be determined a priori only in terms of the data such that either

$$\omega \le 2^{\frac{s_1+n}{p-1}}(\mu_+)^{\frac{1-m}{p-1}}F_2(2R) + 2^{\frac{2s_1+n}{p}}(\mu_+)^{\frac{1-m}{p}}F_1(2R) \tag{4.2}$$

or

$$\left| \left\{ x \in B_{\frac{3}{4}R} : u(x,t) > \mu_{+} - \frac{\omega}{2^{s_{1}}} \right\} \right| \le \left(1 - \left(\frac{\nu_{0}}{2} \right)^{2} \right) |B_{\frac{3}{4}R}| \tag{4.3}$$

for all $t \in \left[\bar{t} - \frac{1}{2}v_0(\mu_+)^{1-m}\omega^{2-p}(\frac{3}{4}R)^p, \bar{t}\right].$

Proof. For simplicity, we write $\varrho = \frac{3}{4}R$. Let $k = \mu_+ - \frac{1}{4}\omega$ and $c = 2^{-2-l}\omega$ where $l \ge 2$ is to be determined later. Moreover, we set $H_k^+ = \frac{1}{4}\omega$ and it follows that

$$H_k^+ \ge \operatorname{ess\,sup}_{B_\varrho \times [t^*, \bar{t}]} |(u - k)_+|.$$

We now consider the logarithmic function defined by

$$\psi^{+} = \ln^{+} \left(\frac{\frac{1}{4}\omega}{\frac{1}{4}\omega - (u - k)_{+} + c} \right).$$

Next, we take a smooth cutoff function $0 < \zeta(x) \le 1$, defined in B_{ϱ} , and satisfying $\zeta \equiv 1$ in $B_{(1-\sigma)\varrho}$ and $|D\zeta| \le (\sigma\varrho)^{-1}$, where $\sigma \in (0,1)$ is to be determined. With these choices, we

use the logarithmic estimate (2.12) to obtain

$$\int_{B_{(1-\sigma)\varrho}\times\{t\}} [\psi^{+}(u)]^{2} dx
\leq \int_{B_{\varrho}\times\{t^{+}\}} [\psi^{+}(u)]^{2} dx + \frac{\gamma}{\sigma^{p}\varrho^{p}} \iint_{B_{\varrho}\times[t^{+},\bar{t}]} u^{m-1}\psi^{+}(u)[(\psi^{+})'(u)]^{2-p} dxdt
+ \gamma(\ln 2)l \frac{2^{2+l}}{\omega^{p-1}} (\mu_{+})^{1-m}\varrho^{p} \int_{B_{\varrho}} |f| dx + \gamma(\ln 2)l \frac{2^{4+2l}}{\omega^{p}} (\mu_{+})^{1-m}\varrho^{p} \int_{B_{\varrho}} g^{\frac{p}{p-1}} dx
=: I_{1} + I_{2} + I_{3} + I_{4}$$
(4.4)

for all $t \in [t^*, \bar{t}]$. Taking into account that $\psi^+(u) \le l \ln 2$, we conclude from (4.1) that

$$I_1 \le (l \ln 2)^2 \left| \left\{ x \in B_\varrho : u(x, t^*) > \mu_+ - \frac{\omega}{4} \right\} \right| \le l^2 (\ln^2 2) \left(\frac{1 - \nu_0}{1 - \frac{1}{2} \nu_0} \right) |B_\varrho|.$$

In view of $[(\psi^+)'(u)]^{2-p} \le (4^{-1}\omega)^{p-2}$, we have

$$I_2 \leq \gamma \frac{1}{\sigma^p o^p} (\mu_+)^{m-1} l(\ln 2) \left(\frac{\omega}{4}\right)^{p-2} (\bar{t} - t^*) |B_{\varrho}| \leq \gamma \frac{l}{\sigma^p} |B_{\varrho}|,$$

where the constant γ depends only upon the data. At this stage, we assume that

$$\omega > 2^{\frac{1}{p-1}} (\mu_+)^{\frac{1-m}{p-1}} \left(\varrho^{p-n} \int_{B_\rho} |f| \, \mathrm{d}x \right)^{\frac{1}{p-1}} + 2^{\frac{2l}{p}} (\mu_+)^{\frac{1-m}{p}} \left(\varrho^{p-n} \int_{B_\rho} g^{\frac{p}{p-1}} \, \mathrm{d}x \right)^{\frac{1}{p}}$$
(4.5)

and hence

$$I_3 + I_4 \le \gamma l |B_{\varrho}| \le \gamma \frac{l}{\sigma^p} |B_{\varrho}|.$$

Combining the estimates for I_1 - I_4 , we arrive at

$$\int_{B_{(1-\sigma)\varrho}\times\{t\}} [\psi^+(u)]^2 \, \mathrm{d}x \le l^2 (\ln^2 2) \left(\frac{1-\nu_0}{1-\frac{1}{2}\nu_0} \right) |B_{\varrho}| + \gamma \frac{l}{\sigma^p} |B_{\varrho}|.$$

On the other hand, the left-hand side of (4.4) can be estimated below by integrating over the smaller set

$$S = \left\{x \in B_{(1-\sigma)\varrho} : u(x,t) > \mu_+ - \frac{\omega}{2^{l+2}}\right\}$$

and this implies that

$$\int_{B_{(1-\sigma)o}\times\{t\}} [\psi^+(u)]^2 \, \mathrm{d}x \ge (l-1)^2 (\ln^2 2) |S|.$$

Consequently, we infer that the estimate

$$\left|\left\{x \in B_{\varrho} : u(x,t) > \mu_{+} - \frac{\omega}{2^{l+2}}\right\}\right| \leq |S| + |B_{\varrho} \setminus B_{(1-\sigma)\varrho}|$$

$$\leq \left[\left(\frac{l}{l-1}\right)^{2} \left(\frac{1-\nu_{0}}{1-\frac{1}{2}\nu_{0}}\right) + \gamma \frac{l}{\sigma^{p}(l-1)^{2}} + \gamma \sigma\right] |B_{\varrho}|$$

holds for all $t \in [t^*, \bar{t}]$. At this point, we choose $\gamma \sigma \leq \frac{3}{8} v_0^2$ and then l so large that

$$\left(\frac{l}{l-1}\right)^2 < \left(1 - \frac{1}{2}\nu_0\right)(1 + \nu_0) \quad \text{and} \quad \gamma \frac{1}{\sigma^p l} \le \frac{3}{8}\nu_0^2.$$

This proves the inequality (4.3) with $s_1 = l + 2$. Moreover, if (4.5) is violated, then we get (4.2) for such a choice of s_1 . We have thus proved the lemma.

Our task now is to establish a De Giorgi type lemma for the second alternative. To this end, we introduce the concentric parabolic cylinders

$$Q_A^{(1)} = B_{\frac{3}{4}R} \times \left(-\frac{1}{2} A^{p-2} (\mu_+)^{1-m} \omega^{2-p} R^p, 0 \right)$$

and

$$Q_A^{(2)} = B_{\frac{1}{2}R} \times \left(-\frac{1}{4} A^{p-2} (\mu_+)^{1-m} \omega^{2-p} R^p, 0 \right),$$

where A satisfies (3.20) and (3.64). The proof of the De Giorgi type lemma is based on the Kilpeläinen-Malý technique. Contrary to Lemma 3.5, we deal with the estimates for u near the supremum μ_+ and our proof makes no use of the cutoff function.

Lemma 4.3. Let $0 < \xi < 4^{-1}p$ and let u be a bounded nonnegative weak solution to (1.1)-(1.2) in Ω_T . There exist constants $\nu_2 \in (0,1)$ and $\hat{B} > 1$, depending only upon the data, such that if

$$\left|\left\{(x,t)\in Q_A^{(1)}: u\geq \mu_+ -\xi\omega\right\}\right|\leq \nu_2|Q_A^{(1)}|,$$

then either

$$u(x,t) < \mu_{+} - \frac{1}{6}\xi\omega$$
 for a.e. $(x,t) \in Q_{A}^{(2)}$ (4.6)

or

$$\xi\omega \le \hat{B}\left((\mu_{+})^{\frac{1-m}{p}}F_{1}(2R) + (\mu_{+})^{\frac{1-m}{p-1}}F_{2}(2R)\right). \tag{4.7}$$

Here $A = \hat{B}\xi^{-1}$ and satisfies (3.20) and (3.64).

Proof. To start with, we first assume that

$$\hat{B} > \max\left\{4^{102}B, \left(\frac{100p}{p-2}\right)^{\frac{1}{p-2}}, A_p^{\frac{1}{p-2}}\right\},$$
 (4.8)

where $A_p > 1$ is the constant such that for any $X > A_p$ there holds $X \le 2^{(p-2)X}$. For such a choice of \hat{B} , the constant $A = \hat{B}\xi^{-1}$ satisfies (3.20) and (3.64). Next, we assume that (3.5) is violated, that is,

$$\xi \omega > \frac{3}{4} \hat{B} \left(\frac{1}{3\hat{B}} \xi \omega + (\mu_+)^{\frac{1-m}{p}} F_1(2R) \right) + (\mu_+)^{\frac{1-m}{p-1}} F_2(2R) \right). \tag{4.9}$$

We are reduced to proving (3.4). Let $(x_1, t_1) \in Q_A^{(2)}$ be a fixed point and assume that (x_1, t_1) is a Lebesgue point of the function u. We set $r_j = 4^{-j}C^{-1}R$ and $B_j = B_{r_j}(x_1)$ where C > 40 is to be determined. For a sequence $\{l_j\}_{j=0}^{\infty}$ and a fixed l > 0, we define

$$Q_j(l) = B_j \times (t_1 - (\mu_+)^{1-m} (l - l_j)^{2-p} r_j^p, t_1).$$

Next, we set $\varphi_j(l) = \phi_j(x)\theta_{j,l}(t)$, where $\phi_j \in C_0^{\infty}(B_j)$, $\phi_j = 1$ on B_{j+1} , $|D\phi_j| \le r_j^{-1}$ and $\theta_{j,l}(t)$ is a Lipschitz function satisfies

$$\theta_{j,l}(t) = 1$$
 in $t \ge t_1 - \frac{4}{9}(\mu_+)^{1-m}(l-l_j)^{2-p}r_j^p$

$$\theta_{j,l}(t) = 0$$
 in $t \le t_1 - \frac{5}{9}(\mu_+)^{1-m}(l-l_j)^{2-p}r_j^p$

and

$$\theta_{j,l}(t) = \frac{t - t_1 - \frac{5}{9}(\mu_+)^{1-m}(l - l_j)^{2-p}r_j^p}{\frac{1}{9}(\mu_+)^{1-m}(l - l_j)^{2-p}r_i^p}$$

in $t_1 - \frac{5}{9}(\mu_+)^{1-m}(l-l_j)^{2-p}r_j^p \le t \le t_1 - \frac{4}{9}(\mu_+)^{1-m}(l-l_j)^{2-p}r_j^p$. It is easy to show that $\varphi_j(l) = 0$ on $\partial_P Q_j(l)$. Moreover, for $j = -1, 0, 1, 2, \cdots$, we construct the sequence $\{\alpha_j\}$ similar to (3.7), that is,

$$\alpha_{j} = \frac{4^{-j-100}}{3\hat{B}} \xi \omega + \frac{3}{4} \int_{0}^{r_{j}} \left(r^{p-n} (\mu_{+})^{1-m} \int_{B_{r}(x_{1})} g(y)^{\frac{p}{p-1}} \, dy \right)^{\frac{1}{p}} \frac{dr}{r} + \frac{3}{4} \int_{0}^{r_{j}} \left(r^{p-n} (\mu_{+})^{1-m} \int_{B_{r}(x_{1})} |f(y)| \, dy \right)^{\frac{1}{p-1}} \frac{dr}{r}.$$

$$(4.10)$$

According to the definition of α_j , we see that $\alpha_j \to 0$ as $j \to \infty$ and the sequence of α_j satisfy

$$\alpha_{j-1} - \alpha_{j} \ge \frac{4^{-j-100}}{\hat{B}} \xi \omega + \gamma \left(r_{j}^{p-n} (\mu_{+})^{1-m} \int_{B_{j}} g(y)^{\frac{p}{p-1}} dy \right)^{\frac{1}{p}} + \gamma \left(r_{j}^{p-n} (\mu_{+})^{1-m} \int_{B_{j}} |f(y)| dy \right)^{\frac{1}{p-1}},$$

$$(4.11)$$

$$\alpha_{j-1} - \alpha_{j} \leq \frac{4^{-j-100}}{\hat{B}} \xi \omega + \gamma \left(r_{j-1}^{p-n} (\mu_{+})^{1-m} \int_{B_{j-1}} g(y)^{\frac{p}{p-1}} \, \mathrm{d}y \right)^{\frac{1}{p}} + \gamma \left(r_{j-1}^{p-n} (\mu_{+})^{1-m} \int_{B_{j-1}} |f(y)| \, \mathrm{d}y \right)^{\frac{1}{p-1}}$$

$$(4.12)$$

and $\hat{B}\alpha_{j-1} \le \xi \omega$ for all $j = 0, 1, 2, \cdots$. Furthermore, we introduce a quantity $K_j(l)$ by

$$\begin{split} K_{j}(l) = & \frac{(\mu_{+})^{m-1}(l-l_{j})^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}(l)} \left(\frac{u-l_{j}}{l-l_{j}}\right)^{(1+\lambda)(p-1)} \varphi_{j}(l)^{k-p} \, \mathrm{d}x \mathrm{d}t \\ & + \operatorname{ess\,sup} \frac{1}{r_{j}^{n}} \int_{B_{j} \times \{t\}} G\left(\frac{u-l_{j}}{l-l_{j}}\right) \varphi_{j}(l)^{k} \, \mathrm{d}x, \end{split}$$

where G is the function defined in (2.1) and $L_j(l) = Q_j(l) \cap \{u \ge l_j\} \cap \Omega_T$. We observe that $K_j(l)$ is continuous in $l > l_j$ and $K_j(l) \downarrow 0$ as $l \to \infty$. Let $l_0 = \mu_+ - \xi \omega$ and $\kappa > 0$ is to be determined later. We construct the sequence l_j in the following way. If $K_j(l_j + \frac{1}{4}(\alpha_{j-1} - \alpha_j)) < \kappa$, then we set $l_{j+1} = l_j + \frac{1}{4}(\alpha_{j-1} - \alpha_j)$. If $K_j(l_j + \frac{1}{4}(\alpha_{j-1} - \alpha_j)) \ge \kappa$, then we choose $l_{j+1} > l_j + \frac{1}{4}(\alpha_{j-1} - \alpha_j)$ such that $K_j(l_{j+1}) = \kappa$. From the definition of l_j , we find that

$$K_j(l_{j+1}) \le \kappa \tag{4.13}$$

for any $j \ge 0$. We proceed similarly as in Lemma 3.5 and divide the proof of (4.6) into three steps.

Step 1: We establish an iteration scheme. To start with, we set $\bar{l} = \frac{1}{2}(l_0 + \mu_+) - \frac{1}{4}B\alpha_0$ and assert that $l_1 \leq \bar{l}$. Observe that $\hat{B}\alpha_0 < \xi\omega$ and hence

$$\bar{l} - l_0 = \frac{1}{2}\xi\omega - \frac{1}{4}\hat{B}\alpha_0 \ge \frac{1}{4}\xi\omega \ge \frac{1}{4}(\alpha_{-1} - \alpha_0)$$
 and $\bar{l} - l_0 \le \frac{1}{2}\xi\omega$. (4.14)

It follows that

$$(\mu_{+})^{1-m}(\bar{l}-l_{0})^{2-p}r_{0}^{p} \leq (\mu_{+})^{1-m}4^{p-2}\hat{B}^{2-p}A^{p-2}\omega^{2-p}C^{-p}R^{p} < \frac{1}{16}(\mu_{+})^{1-m}A^{p-2}\omega^{2-p}R^{p},$$

since C > 40. This also yields that $Q_0(\bar{l}) \subset Q_A^{(1)}$. Since $u - l_0 \le \xi \omega$ in $L_0(\bar{l})$, we conclude that

$$\begin{split} &\frac{(\mu_{+})^{m-1}(\bar{l}-l_{0})^{p-2}}{r_{0}^{n+p}} \iint_{L_{0}(\bar{l})} \left(\frac{u-l_{0}}{\bar{l}-l_{0}}\right)^{(1+\lambda)(p-1)} \varphi_{0}(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\ &\leq \frac{(\mu_{+})^{m-1}(\bar{l}-l_{0})^{p-2-(1+\lambda)(p-1)}}{r_{0}^{n+p}} \left(\xi\omega\right)^{(1+\lambda)(p-1)} |L_{0}(\bar{l})| \\ &\leq \hat{\gamma}_{0} C^{n+p} \frac{(\mu_{+})^{m-1}(\xi\omega)^{p-2}}{R^{n+p}} |L_{0}(\bar{l})| \\ &\leq \hat{\gamma}_{0} C^{n+p} \hat{B}^{p-2} \frac{\left|Q_{A}^{(1)} \cap \{u \geq \mu_{+} - \xi\omega\}\right|}{|Q_{A}^{(1)}|} \leq \hat{\gamma}_{0} C^{n+p} \hat{B}^{p-2} \nu_{2}, \end{split}$$

where the constant $\hat{\gamma}_0$ depends only upon the data. Moreover, we apply Lemma 2.4 with (a, d, l) replaced by $(\mu_+, \bar{l} - l_0, l_0)$ to obtain

$$\begin{split} & \operatorname{ess\,sup} \, \frac{1}{r_0^n} \int_{B_0 \times \{t\}} G\left(\frac{u - l_0}{\bar{l} - l_0}\right) \varphi_0(\bar{l})^k \, \mathrm{d}x \\ \leq & \gamma \frac{(\bar{l} - l_0)^{p-2}}{r_0^{p+n}} \int_{L_0(\bar{l})} u^{m-1} \left(\frac{u - l_0}{\bar{l} - l_0}\right)^{(1+\lambda)(p-1)} \varphi_0(\bar{l})^{k-p} \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{1}{r_0^n} \int_{L_0(\bar{l})} \frac{u - l_0}{\bar{l} - l_0} |\partial_t \varphi_0(\bar{l})| \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{r_0^{p-n}}{(\bar{l} - l_0)^p} (\mu_+)^{1-m} \int_{B_0} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{r_0^{p-n}}{(\bar{l} - l_0)^{p-1}} (\mu_+)^{1-m} \int_{B_0} |f| \, \mathrm{d}x, \end{split}$$

since $\varphi_0(\bar{l}) \in C_0^\infty(Q_0(\bar{l}))$ and the first term on the right-hand side of (2.10) vanishes. Taking into account that $u \le \mu_+$ in $L_0(\bar{l})$ and $|\partial_t \varphi_0(\bar{l})| \le 9(\bar{l} - l_0)^{p-2}(\mu_+)^{m-1}r_0^{-p}$, an argument similar to the one used in the proof of Lemma 3.5 shows that there exists a constant $\hat{\gamma}_1$ depending only upon the data such that

$$\operatorname{ess\,sup} \frac{1}{r_0^n} \int_{B_0 \times \{t\}} G\left(\frac{u - l_0}{\bar{l} - l_0}\right) \varphi_0(\bar{l})^k \, \mathrm{d}x \le \hat{\gamma}_1 C^{n+p} \hat{B}^{p-2} \nu_2 + \hat{\gamma}_1 C^{n-p} (\hat{B}^{-p} + \hat{B}^{1-p}).$$

Consequently, we infer that

$$K_0(\bar{l}) \leq (\hat{\gamma}_0 + \hat{\gamma}_1)C^{n+p}\hat{B}^{p-2}\nu_2 + \hat{\gamma}_1C^{n-p}(\hat{B}^{-p} + \hat{B}^{1-p}).$$

At this point, we choose $v_2 < 1$ and $\hat{B} > 1$ be such that

$$(\hat{\gamma}_0 + \hat{\gamma}_1)C^{n+p}\hat{B}^{p-2}\nu_2 = \frac{\kappa}{4} \quad \text{and} \quad \hat{\gamma}_1C^{n-p}(\hat{B}^{-p} + \hat{B}^{1-p}) < \frac{\kappa}{4}. \tag{4.15}$$

Consequently, we deduce that $K_0(\bar{l}) \le \frac{1}{2}\kappa$. It follows from (4.13) that $l_1 \le \bar{l}$. Next, we set $d_j = l_{j+1} - l_j$ and $Q_j = B_j \times (t_1 - (\mu_+)^{1-m} d_j^{2-p} r_j^p, t_1)$. We claim that for any $j \ge 1$ there holds

$$d_{j} \leq \frac{1}{4}d_{j-1} + \gamma \frac{4^{-j}}{\hat{B}} \xi \omega + \gamma \left(r_{j-1}^{p-n} (\mu_{+})^{1-m} \int_{B_{j-1}} g(y)^{\frac{p}{p-1}} dy \right)^{\frac{1}{p}} + \gamma \left(r_{j-1}^{p-n} (\mu_{+})^{1-m} \int_{B_{j-1}} |f(y)| dy \right)^{\frac{1}{p-1}}.$$

$$(4.16)$$

Step 2: *Proof of the inequality* (4.16). To start with, we first assume that for any fixed $j \ge 1$ there holds

$$d_j > \frac{1}{4}d_{j-1}$$
 and $d_j > \frac{1}{4}(\alpha_{j-1} - \alpha_j),$ (4.17)

since otherwise (4.16) holds immediately. Since $d_j > \frac{1}{4}(\alpha_{j-1} - \alpha_j)$, we infer from the construction of $K_j(l_{j+1})$ that $K_j(l_{j+1}) = \kappa$. To simplify the notation, we set $\varphi_i = \varphi_i(l_{i+1})$. In view of $d_j > \frac{1}{4}d_{j-1}$, we deduce that

$$(\mu_{+})^{1-m}d_{j}^{2-p}r_{j}^{p} \leq (\mu_{+})^{1-m}\frac{r_{j-1}^{p}}{4^{p}}\left(\frac{d_{j-1}}{4}\right)^{2-p} = \frac{1}{16}(\mu_{+})^{1-m}d_{j-1}^{2-p}r_{j-1}^{p}$$

which yields $Q_j \subset Q_{j-1}$ and $\varphi_{j-1}(x,t) = 1$ for $(x,t) \in Q_j$. We also remark that for any $i \ge 0$, the inclusion $Q_i \subset \hat{Q}$ holds, where \hat{Q} is defined in (2.14). Taking into account that $A = \hat{B}\mathcal{E}^{-1}$, we infer from (4.11) that

$$\begin{split} (\mu_+)^{1-m} d_i^{2-p} r_i^p & \leq (\mu_+)^{1-m} 4^{p-2} (\alpha_{i-1} - \alpha_i)^{2-p} r_i^p \\ & \leq (\mu_+)^{1-m} 4^{-2i} 4^{101(p-2)} \left(\frac{\xi \omega}{\hat{B}}\right)^{2-p} C^{-p} R^p \leq (\mu_+)^{1-m} \left(\frac{\omega}{A}\right)^{2-p} R^p, \end{split}$$

provided that we choose $C=4^{101}$. This proves the inclusion $Q_i\subset \hat{Q}$ for any $i\geq 0$. We now turn our attention to the proof of (4.16). To this end, we set $L_j=Q_j\cap \{u\geq l_j\}\cap \Omega_T$ and decompose $L_j=L_i'\cup L_i''$, where

$$L'_{j} = L_{j} \cap \left\{ \frac{u - l_{j}}{d_{j}} \le \varepsilon_{1} \right\}$$
 and $L''_{j} = L_{j} \setminus L'_{j}$. (4.18)

In view of $u \le l_j$ on L_j , we use (4.13) to deduce that

$$\frac{(\mu_{+})^{m-1}d_{j}^{p-2}}{r_{j}^{n+p}}|L_{j}| \le \frac{4^{n}}{r_{j-1}^{n}}\operatorname{ess\,sup}\int_{B_{j-1}}G\left(\frac{u-l_{j-1}}{d_{j-1}}\right)\varphi_{j-1}^{k}\,\mathrm{d}x \le 4^{n}\kappa. \tag{4.19}$$

It follows from (4.19) that

$$\frac{(\mu_{+})^{m-1}d_{j}^{p-2}}{r_{j}^{n+p}} \iint_{L'_{j}} \left(\frac{u-l_{j}}{d_{j}}\right)^{(1+\lambda)(p-1)} \varphi_{j}^{k-p} \, \mathrm{d}x \mathrm{d}t
\leq \frac{(\mu_{+})^{m-1}d_{j}^{p-2}}{r_{j}^{n+p}} \varepsilon_{1}^{(1+\lambda)(p-1)} |L_{j}| \leq 4^{n} \varepsilon_{1}^{(1+\lambda)(p-1)} \kappa.$$
(4.20)

Let parameters h and q be as in (3.26). For a fixed $\varepsilon_2 < 1$, we apply Young's inequality to conclude that

$$\begin{split} &\frac{(\mu_{+})^{m-1}d_{j}^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}'} \left(\frac{u-l_{j}}{d_{j}}\right)^{(1+\lambda)(p-1)} \varphi_{j}^{k-p} \, \mathrm{d}x \mathrm{d}t \\ &\leq \varepsilon_{2} \frac{(\mu_{+})^{m-1}d_{j}^{p-2}}{r_{j}^{n+p}} |L_{j}| + \gamma(\varepsilon_{2}) \frac{(\mu_{+})^{m-1}d_{j}^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}'} \left(\frac{u-l_{j}}{d_{j}}\right)^{p\frac{n+h}{nh}} \varphi_{j}^{(k-p)q} \, \mathrm{d}x \mathrm{d}t \\ &=: T_{1} + T_{2}, \end{split}$$

with the obvious meanings of T_1 and T_2 . According to (4.19), we see that $T_1 \le 4^n \varepsilon_2 \kappa$. To estimate T_2 , we use Lemma 2.2 with (l,d) replaced by (l_j,d_j) to deduce that

$$T_2 \le \gamma \frac{(\mu_+)^{m-1} (l_j - \bar{l})^{p-2}}{r_j^{n+p}} \iint_{L_j''} \psi_j^{p \frac{n+h}{n}} \varphi_j^{(k-p)q} \, \mathrm{d}x \mathrm{d}t, \tag{4.21}$$

where

$$\psi_j(x,t) = \frac{1}{d_j} \left[\int_{l_j}^u \left(1 + \frac{s - l_j}{d_j} \right)^{-\frac{1}{p} - \frac{\lambda}{p}} \, \mathrm{d}s \right]_\perp.$$

Let $v = \psi_j \varphi_j^{k_1}$, where $k_1 = \frac{(k-p)nq}{p(n+h)}$. Recalling that p < n, we use Hölder's inequality, Sobolev's inequality and Lemma 2.2 to deduce

$$\iint_{L''_{j}} \psi_{j}^{\frac{n+h}{n}} \varphi_{j}^{(k-p)q} \, \mathrm{d}x \mathrm{d}t$$

$$\leq \int_{t_{1}-(\mu_{+})^{1-m} d_{j}^{2-p} r_{j}^{p}}^{t_{1}} \left(\int_{B_{j}} v^{\frac{np}{n-p}} \, \mathrm{d}x \right)^{\frac{n-p}{n}} \left(\int_{L''_{j}(t)} v^{h} \, \mathrm{d}x \right)^{\frac{p}{n}} \, \mathrm{d}t$$

$$\leq \gamma \operatorname{ess} \sup_{t} \left(\int_{L''_{j}(t)} \frac{u - l_{j}}{d_{j}} \varphi_{j}^{k_{1}h} \, \mathrm{d}x \right)^{\frac{p}{n}} \iint_{Q_{j}} |Dv|^{p} \, \mathrm{d}x \mathrm{d}t, \tag{4.22}$$

where

$$L_j''(t) = \{x \in B_j : u(\cdot, t) \ge l_j\} \cap \left\{x \in B_j : \frac{u(\cdot, t) - l_j}{d_j} > \varepsilon_1\right\}.$$

According to Lemma 3.4, we find that

$$\int_{L_i''(t)} \frac{u - l_j}{d_j} \varphi_j^{k_1 h} \, \mathrm{d}x \le c(\varepsilon_1) \int_{L_j(t)} G\left(\frac{u - l_j}{d_j}\right) \varphi_j^{k_1 h} \, \mathrm{d}x. \tag{4.23}$$

At this point, we introduce the quantities M_1 and M_2 by

$$M_1 = \frac{r_j^{p-n}}{d_j^p} (\mu_+)^{1-m} \int_{B_j} g^{\frac{p}{p-1}} dx$$
 and $M_2 = \frac{r_j^{p-n}}{d_j^{p-1}} (\mu_+)^{1-m} \int_{B_j} |f| dx$.

We now apply Lemma 2.4 with (a, d, l) replaced by (μ_+, d_i, l_i) to conclude that

$$\begin{aligned} & \operatorname{ess\,sup} \frac{1}{r_{j}^{n}} \int_{B_{j}p} G\left(\frac{u - l_{j}}{d_{j}}\right) \varphi_{j}^{k_{1}h} \, \mathrm{d}x \\ & \leq \gamma \frac{d_{j}^{p-2}}{r_{j}^{p+n}} \iint_{L_{j}} u^{m-1} \left(\frac{u - l_{j}}{d_{j}}\right)^{(1+\lambda)(p-1)} \varphi_{j}^{k_{1}h-p} \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{1}{r_{j}^{n}} \iint_{L_{j}} \frac{u - l_{j}}{d_{j}} |\partial_{t}\varphi_{j}| \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{r_{j}^{p-n}}{d_{j}^{p}} (\mu_{+})^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{r_{j}^{p-n}}{d_{j}^{p-1}} (\mu_{+})^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x \\ & = : T_{3} + T_{4} + \gamma M_{1} + \gamma M_{2}, \end{aligned}$$

since $\varphi_j = 0$ on $\partial_P Q_j$. We first consider the estimate for T_3 . According to (4.13) and (4.17), we conclude that

$$T_{3} \leq \gamma \frac{d_{j}^{p-2-(1+\lambda)(p-1)}(\mu_{+})^{m-1}}{r_{j}^{p+n}} \iint_{L_{j}} (u - l_{j})^{(1+\lambda)(p-1)} \varphi_{j}^{k_{1}h-p} \, \mathrm{d}x \mathrm{d}t$$

$$\leq \gamma \frac{d_{j-1}^{p-2}(\mu_{+})^{m-1}}{r_{j-1}^{p+n}} \iint_{L_{j-1}} \left(\frac{u - l_{j-1}}{d_{j-1}}\right)^{(1+\lambda)(p-1)} \varphi_{j-1}^{k-p} \, \mathrm{d}x \mathrm{d}t$$

$$\leq \gamma K_{i-1}(l_{i}) \leq \gamma \kappa.$$

Taking into account that $(1 + \lambda)(p-1) > 1$, $l_{j-1} \le l_j$, $d_j > 4^{-1}d_{j-1}$ and $|\partial_t \varphi_j| \le 9d_j^{p-2}(\mu_+)^{m-1}r_j^{-p}$, we infer from (4.19) that

$$\begin{split} T_4 \leq & \gamma \frac{d_j^{p-2}(\mu_+)^{m-1}}{r_j^{n+p}} \iint_{L_j} \frac{u - l_j}{d_j} \, \mathrm{d}x \mathrm{d}t \\ \leq & \gamma \frac{d_j^{p-2}(\mu_+)^{m-1}}{r_j^{n+p}} |L_j| + \gamma \frac{d_j^{p-2}(\mu_+)^{m-1}}{r_j^{n+p}} \iint_{L_j} \left(\frac{u - l_j}{d_j}\right)^{(1+\lambda)(p-1)} \, \mathrm{d}x \mathrm{d}t \\ \leq & \gamma \kappa + K_{i-1}(l_i) \leq \gamma \kappa. \end{split}$$

Combining the above estimates, we infer that there exists a constant γ depending only upon the data such that

$$\operatorname{ess\,sup} \frac{1}{r_{j}^{n}} \int_{B_{j}} G\left(\frac{u - l_{j}}{d_{j}}\right) \varphi_{j}^{k_{1}h} \, \mathrm{d}x \le \gamma \kappa + \gamma (M_{1} + M_{2}). \tag{4.24}$$

We now turn our attention to the estimate of T_2 . Combining (4.21)-(4.24), we can rewrite the upper bound for T_2 by

$$T_{2} \leq \gamma \frac{(\mu_{+})^{m-1} d_{j}^{p-2}}{r_{j}^{n}} (\kappa + M_{1} + M_{2})^{\frac{p}{n}} \times \left[\iint_{Q_{j}} \varphi_{j}^{k_{1}p} |D\psi_{j}|^{p} dxdt + \iint_{Q_{j}} \varphi_{j}^{(k_{1}-1)p} \psi_{j}^{p} |D\varphi_{j}|^{p} dxdt \right]$$

$$= : \gamma (\kappa + M_{1} + M_{2})^{\frac{p}{n}} (T_{7} + T_{8}),$$

with the obvious meanings of T_7 and T_8 . We first consider the estimate for T_7 . Noting that $\xi < 4^{-1}$ and $\omega \le 2\mu_+$, we have $u \ge l_j = \mu_+ - \xi\omega \ge \mu_+ - 2\xi\mu_+ > \frac{1}{2}\mu_+$ on L_j and hencep

$$T_7 = \frac{(\mu_+)^{m-1} d_j^{p-2}}{r_j^n} \iint_{L_j} \varphi_j^{k_1 p} |D\psi_j|^p \, \mathrm{d}x \mathrm{d}t \le \gamma \frac{d_j^{p-2}}{r_j^n} \iint_{Q_j} u^{m-1} |D\psi_j|^p \varphi_j^{k_1 p} \, \mathrm{d}x \mathrm{d}t.$$

To proceed further, we use Lemma 2.3 with (a, d, l) replaced by (μ_+, d_j, l_j) and taking into account the estimates for T_3 - T_6 . Since $\varphi_j = 0$ on $\partial_P Q_j$, we conclude that

$$\begin{split} T_{7} \leq & \gamma \frac{d_{j}^{p-2}}{r_{j}^{n}} \iint_{Q_{j}} u^{m-1} |D\psi_{j}|^{p} \varphi_{j}^{k_{1}p} \, \mathrm{d}x \mathrm{d}t \\ \leq & \gamma \frac{d_{j}^{p-2}}{r_{j}^{p}} \iint_{L_{j}} u^{m-1} \left(\frac{u-l_{j}}{d_{j}}\right)^{(1+\lambda)(p-1)} \varphi_{j}^{(k_{1}-1)p} \, \mathrm{d}x \mathrm{d}t + \gamma \iint_{L_{j}} \frac{u-l_{j}}{d_{j}} |\partial_{t}\varphi_{j}| \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{r_{j}^{p}}{d_{j}^{p}} (\mu_{+})^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{r_{j}^{p}}{d_{j}^{p-1}} (\mu_{+})^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x \\ \leq & \gamma \frac{d_{j-1}^{p-2} (\mu_{+})^{m-1}}{r_{j-1}^{p+n}} \iint_{L_{j-1}} \left(\frac{u-l_{j-1}}{d_{j-1}}\right)^{(1+\lambda)(p-1)} \varphi_{j-1}^{k-p} \, \mathrm{d}x \mathrm{d}t \\ & + \gamma \frac{d_{j}^{p-2} (\mu_{+})^{m-1}}{r_{j}^{n+p}} |L_{j}| + M_{1} + M_{2} \\ \leq & \gamma (\kappa + M_{1} + M_{2}). \end{split}$$

To estimate T_8 , we first note that

$$\psi_j(x,t) = \frac{1}{d_j} \left[\int_{l_j}^u \left(1 + \frac{s - l_j}{d_j} \right)^{-1 - \lambda} ds \right]^{\frac{1}{p}} (u - l_j)_+^{\frac{1}{p'}} \le \frac{(u - l_j)_+^{\frac{1}{p'}}}{d_j^{\frac{1}{p'}}}.$$

Consequently, we infer that

$$\begin{split} T_{8} &\leq \frac{(\mu_{+})^{m-1}d_{j}^{p-2}}{r_{j}^{n}} \iint_{Q_{j}} \varphi_{j}^{(k_{1}-1)p} \psi_{j}^{p} |D\varphi_{j}|^{p} \, \mathrm{d}x \mathrm{d}t \\ &\leq \frac{(\mu_{+})^{m-1}d_{j}^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}} \left(\frac{u-l_{j}}{d_{j}}\right)^{p-1} \varphi_{j}^{(k_{1}-1)p} \, \mathrm{d}x \mathrm{d}t \\ &\leq \frac{(\mu_{+})^{m-1}d_{j}^{p-2}}{r_{j}^{n+p}} |L_{j}| + \frac{(\mu_{+})^{m-1}d_{j}^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}} \left(\frac{u-l_{j}}{d_{j}}\right)^{(1+\lambda)(p-1)} \varphi_{j}^{(k_{1}-1)p} \, \mathrm{d}x \mathrm{d}t \\ &\leq \gamma_{K} \end{split}$$

and hence we arrive at $T_2 \le \gamma(\kappa + M_1 + M_2)^{1+\frac{p}{n}}$. Combining the above estimates, we conclude with

$$\frac{(\mu_{+})^{m-1}d_{j}^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}^{\prime\prime}} \left(\frac{u-l_{j}}{d_{j}}\right)^{(1+\lambda)(p-1)} \varphi_{j}^{k-p} \, \mathrm{d}x \mathrm{d}t$$

$$\leq 4^{n} \varepsilon_{2} \kappa + \gamma(\varepsilon_{2})(\kappa + M_{1} + M_{2})^{1+\frac{p}{n}}.$$
(4.25)

This also yields that

$$\frac{(\mu_{+})^{m-1}d_{j}^{p-2}}{r_{j}^{n+p}} \iint_{L_{j}} \left(\frac{u-l_{j}}{d_{j}}\right)^{(1+\lambda)(p-1)} \varphi_{j}^{k-p} \, \mathrm{d}x \mathrm{d}t
\leq 4^{n} \varepsilon_{1}^{(1+\lambda)(p-1)} \kappa + 4^{n} \varepsilon_{2} \kappa + \gamma(\varepsilon_{2})(\kappa + M_{1} + M_{2})^{1+\frac{p}{n}}.$$
(4.26)

Furthermore, we aim to improve the estimate (4.24). To this end, we apply Lemma 2.4 to obtain

$$\operatorname{ess\,sup} \frac{1}{r_{j}^{n}} \int_{L_{j}(t)} G\left(\frac{u - l_{j}}{d_{j}}\right) \varphi_{j}^{k} \, \mathrm{d}x$$

$$\leq \gamma \frac{d_{j}^{p-2}}{r_{j}^{p+n}} \iint_{L_{j}} u^{m-1} \left(\frac{u - l_{j}}{d_{j}}\right)^{(1+\lambda)(p-1)} \varphi_{j}^{k-p} \, \mathrm{d}x \, \mathrm{d}t$$

$$+ \gamma \frac{1}{r_{j}^{n}} \iint_{L_{j}} \frac{u - l_{j}}{d_{j}} \varphi_{j}^{k-1} |\partial_{t} \varphi_{j}| \, \mathrm{d}x \, \mathrm{d}t$$

$$+ \gamma \frac{r_{j}^{p-n}}{d_{j}^{p}} (\mu_{+})^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x + \gamma \frac{r_{j}^{p-n}}{d_{j}^{p-1}} (\mu_{+})^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x$$

$$=: S_{1} + S_{2} + \gamma M_{1} + \gamma M_{2},$$

$$(4.27)$$

with the obvious meanings of S_1 and S_2 . To estimate S_1 , we apply (4.26) to deduce that

$$\begin{split} S_{1} &\leq 2^{(1+\lambda)(p-1)} \gamma \frac{d_{j}^{p-2} (\mu_{+})^{m-1}}{r_{j}^{p+n}} \iint_{L_{j}} \left(\frac{u - l_{j}}{d_{j}} \right)^{(1+\lambda)(p-1)} \varphi_{j}^{k-p} \, \mathrm{d}x \mathrm{d}t \\ &\leq 2^{(1+\lambda)(p-1)} \gamma \left[4^{n} \varepsilon_{1}^{(1+\lambda)(p-1)} \kappa + 4^{n} \varepsilon_{2} \kappa + \gamma(\varepsilon_{2}) (\kappa + M_{1} + M_{2})^{1 + \frac{\rho}{n}} \right]. \end{split}$$

Finally, we consider the estimate for S_2 . To this end, we decompose $L_j = L'_j \cup L''_j$, where L'_j and L''_j satisfy (4.18). In view of $|\partial_t \varphi_j| \le 9d_j^{p-2}(\mu_+)^{m-1}r_j^{-p}$, we use (4.19) and (4.25) to conclude that

$$\begin{split} S_2 \leq & \gamma \frac{d_j^{p-2}(\mu_+)^{m-1}}{r_j^{n+p}} \iint_{L_j} \frac{u-l_j}{d_j} \varphi_j^{k-1} \, \mathrm{d}x \mathrm{d}t \\ \leq & \gamma \varepsilon_1 \frac{(\mu_+)^{m-1} d_j^{p-2}}{r_j^{n+p}} |L_j'| + \gamma \frac{d_j^{p-2}(\mu_+)^{m-1}}{r_j^{p+n}} \iint_{L_j''} \left(\frac{u-l_j}{d_j} \right)^{(1+\lambda)(p-1)} \varphi_j^{k-p} \, \mathrm{d}x \mathrm{d}t \\ \leq & 4^n \varepsilon_1 \kappa + \gamma \left[4^n \varepsilon_2 \kappa + \gamma (\varepsilon_2) (\kappa + M_1 + M_2)^{1+\frac{p}{n}} \right]. \end{split}$$

Inserting the estimates for S_1 and S_2 into (4.27) and taking into account (4.26), we arrive at

$$K_i(l_{i+1}) \leq \gamma (M_1 + M_2) + 2^{(1+\lambda)(p-1)} \gamma 4^n (\varepsilon_1 + \varepsilon_2) \kappa + \gamma (\varepsilon_2) (\kappa + M_1 + M_2)^{1+\frac{p}{n}}.$$

Recalling that $\kappa = K_j(l_{j+1})$, we conclude that there exist constants $\hat{\gamma}_2 = \hat{\gamma}_2(\text{data})$ and $\hat{\gamma}_3 = \hat{\gamma}_3(\text{data}, \varepsilon_1, \varepsilon_2)$ such that

$$\begin{split} \kappa &= K_{j}(l_{j+1}) \leq \hat{\gamma}_{3} \frac{r_{j}^{p-n}}{d_{j}^{p}} (\mu_{+})^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x + \hat{\gamma}_{3} \frac{r_{j}^{p-n}}{d_{j}^{p-1}} (\mu_{+})^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x \\ &+ \hat{\gamma}_{2} (\varepsilon_{1} + \varepsilon_{2}) \kappa + \hat{\gamma}_{3} \kappa^{1+\frac{p}{n}} \\ &+ \hat{\gamma}_{3} \left[\frac{r_{j}^{p-n}}{d_{j}^{p}} (\mu_{+})^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x + \frac{r_{j}^{p-n}}{d_{j}^{p-1}} (\mu_{+})^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x \right]^{1+\frac{p}{n}}. \end{split}$$

At this point, we first determine the values of ε_1 and ε_2 by $\varepsilon_1 = \varepsilon_2 = (8\hat{\gamma}_2)^{-1}$. The choices of ε_1 and ε_2 determine the value of $\hat{\gamma}_3$. Moreover, we choose κ be such that

$$\kappa = 4^{-\frac{n}{p}} \hat{\gamma}_3^{-\frac{n}{p}}. \tag{4.28}$$

The choices of ε_1 , ε_2 and χ ensuring $\hat{\gamma}_2(\varepsilon_1 + \varepsilon_2)\kappa + \hat{\gamma}_3\kappa^{1+\frac{p}{n}} \leq \frac{1}{2}\kappa$ and we conclude that either

$$d_{j} \leq \gamma \left(r_{j}^{p-n} (\mu_{+})^{1-m} \int_{B_{j}} g^{\frac{p}{p-1}} \, \mathrm{d}x \right)^{\frac{1}{p}} \qquad \text{or} \qquad d_{j} \leq \gamma \left(r_{j}^{p-n} (\mu_{+})^{1-m} \int_{B_{j}} |f| \, \mathrm{d}x \right)^{\frac{1}{p-1}}$$

holds, which proves the inequality (4.17).

Step 3: *Proof of the inequality* (4.6). Let J > 1 be a fixed integer. We sum up the inequality (4.17) for $j = 1, \dots, J-1$ and obtain

$$\begin{split} l_{J} - l_{1} &\leq \frac{1}{3} d_{0} + \gamma \frac{\xi \omega}{\hat{B}} \sum_{j=1}^{J-1} 4^{-j} + \gamma \sum_{j=1}^{J-1} \left(r_{j-1}^{p-n} (\mu_{+})^{1-m} \int_{B_{j-1}} |f(y)| \, \mathrm{d}y \right)^{\frac{1}{p-1}} \\ &+ \gamma \sum_{j=1}^{J-1} \left(r_{j-1}^{p-n} (\mu_{+})^{1-m} \int_{B_{j-1}} g(y)^{\frac{p}{p-1}} \, \mathrm{d}y \right)^{\frac{1}{p}} \, . \end{split}$$

In view of $d_0 = l_1 - l_0$ and $l_0 = \mu_+ - \xi \omega$, we apply (4.9) to conclude that there exists a constant $\hat{\gamma}_4 = \hat{\gamma}_4(\text{data}) > 1$ such that

$$l_{J} \leq \frac{4}{3}d_{0} + \mu_{+} - \xi\omega + \gamma \frac{\xi\omega}{\hat{B}} + \gamma(\mu_{+})^{\frac{1-m}{p-1}} \int_{0}^{4C^{-1}R} \left(\frac{1}{r^{n-p}} \int_{B_{r}(x_{1})} f(y) \, dy\right)^{\frac{1}{p-1}} \frac{1}{r} \, dr + \gamma(\mu_{+})^{\frac{1-m}{p}} \int_{0}^{4C^{-1}R} \left(\frac{1}{r^{n-p}} \int_{B_{r}(x_{1})} g(y)^{\frac{p}{p-1}} \, dy\right)^{\frac{1}{p}} \frac{1}{r} \, dr \leq \frac{4}{3}d_{0} + \mu_{+} - \xi\omega + \hat{\gamma}_{4} \frac{\xi\omega}{\hat{B}}.$$

$$(4.29)$$

According to (4.14), we find that $d_0 \le \overline{l} - l_0 \le \frac{1}{2} \xi \omega$. Passing to the limit $J \to \infty$, we infer from (4.29) that

$$u(x_1, t_1) < \mu_+ - \frac{1}{6} \xi \omega,$$

provided that we choose $\hat{B} > 6\hat{\gamma}_4$. This shows that the estimate (4.6) holds for almost everywhere point in $Q_A^{(2)}$, since (x_1, t_1) is the Lebesgue point of u. Taking into account that $C = 4^{101}$, (4.8) and (4.15), we determine the constant \hat{B} by

$$\hat{B} = \max \left\{ \left(\frac{\kappa}{4^{101(n-p)+2} \hat{\gamma}_1} \right)^{\frac{1}{1-p}}, 6\hat{\gamma}_4, 4^{102} B, \left(\frac{100p}{p-2} \right)^{\frac{1}{p-2}}, A_p^{\frac{1}{p-2}} \right\},$$
(4.30)

where the constant κ is determined via (4.28). For such a choice of \hat{B} , we determine the value of ν_2 via (4.15) by

$$\nu_2 = \frac{\kappa}{4^{101(n-p)+1}(\hat{\gamma}_0 + \hat{\gamma}_1)\hat{B}^{p-2}}.$$
(4.31)

We find that the constant v_2 depends only upon the data and independent of ξ . The proof of the Lemma is now complete.

The crucial step to obtain the main result proved in this section is the following lemma concerning the estimate of the measure of level sets.

Lemma 4.4. Let u be a bounded nonnegative weak solution to (1.1)-(1.2) in Ω_T . For every $\bar{v} \in (0, 1)$, there exists a positive integer $q_* = q_*(data, \bar{v})$ such that either

$$\omega \le 2^{s_1 + q_*} (\mu_+)^{\frac{1 - m}{p}} F_1(2R) + 2^{s_1 + q_*} (\mu_+)^{\frac{1 - m}{p - 1}} F_2(2R) \tag{4.32}$$

or

$$\left| \left\{ (x,t) \in Q_A^{(1)} : u \ge \mu_+ - \frac{\omega}{2^{s_1 + q_*}} \right\} \right| \le \bar{\nu} |Q_A^{(1)}|, \tag{4.33}$$

where $s_1 > 2$ is the constant claimed by Lemma 4.2 and $A = 2^{s_1+q_*}\hat{B}$.

Proof. For simplicity, we abbreviate $Q_A = Q_A^{(1)}$. To start with, we first assume that (4.2) is violated, that is,

$$\omega > 2^{\frac{s_1+n}{p-1}}(\mu_+)^{\frac{1-m}{p-1}}F_2(2R) + 2^{\frac{2s_1+n}{p}}(\mu_+)^{\frac{1-m}{p}}F_1(2R). \tag{4.34}$$

According to Lemma 4.2, we infer that the slice-wise estimate

$$\left| \left\{ x \in B_{\frac{3}{4}R} : u(x,t) > \mu_{+} - \frac{\omega}{2^{s_{1}}} \right\} \right| \le \left(1 - \left(\frac{\nu_{0}}{2} \right)^{2} \right) |B_{\frac{3}{4}R}| \tag{4.35}$$

holds for all $t \in \left(-\frac{1}{2}A^{p-2}(\mu_+)^{1-m}\omega^{2-p}R^p, 0\right]$. Let $q_* > 1$ to be determined in the course of the proof. For j > 2 and $A = 2^{s_1+q_*}\hat{B}$, we define

$$Q'_{A} = B_{R} \times \left(-A^{p-2} (\mu_{+})^{1-m} \omega^{2-p} R^{p}, 0 \right],$$

$$A_{j}(t) = \left\{ x \in B_{\frac{3}{4}R} : u(x, t) > \mu_{+} - \frac{\omega}{2^{j}} \right\}$$

and

$$Q_j^+ = \left\{ (x, t) \in Q_A : u(x, t) > \mu_+ - \frac{\omega}{2j} \right\}.$$

Now for $q_* \in \mathbb{N}$, we set $k_j = \mu_+ - 2^{-j}\omega$ for $j = s_1, s_1 + 1, \dots, s_1 + q_*$. Take a cutoff function $0 \le \varphi \le 1$, such that $\varphi = 1$ in Q_A , $\varphi = 0$ on $\partial_P Q'_A$,

$$|D\varphi| \le \frac{4}{R}$$
 and $0 \le \frac{\partial \varphi}{\partial t} \le 2A^{2-p}(\mu_+)^{m-1}\omega^{p-2}R^{-p}$.

We consider the Caccioppoli estimate (2.13) for the truncated functions $(u - k_j)_+$ over the cylinder Q'_A and obtain

$$\iint_{\mathcal{Q}_{A}'} u^{m-1} |D(u-k_{j})_{+}\varphi|^{p} \, \mathrm{d}x \mathrm{d}t
\leq \gamma \iint_{\mathcal{Q}_{A}'} u^{m-1} (u-k_{j})_{+}^{p} |D\varphi|^{p} \, \mathrm{d}x \mathrm{d}t + \gamma \iint_{\mathcal{Q}_{A}'} (u-k_{j})_{+}^{2} |\partial_{t}\varphi| \, \mathrm{d}x \mathrm{d}t
+ \gamma \iint_{\mathcal{Q}_{A}'} |f|(u-k_{j})_{+} \, \mathrm{d}x \mathrm{d}t + \gamma \iint_{\mathcal{Q}_{A}'} g^{\frac{p}{p-1}} \, \mathrm{d}x \mathrm{d}t.$$

Taking into accound that j > 2 and $\omega < 2\mu_+$, we have $u \ge k_j > \mu_+ - \frac{1}{4}\omega > \frac{1}{2}\mu_+$ on the set $\{u \ge k_j\}$. Then, we deduce

$$\iint_{\mathcal{Q}_A'} u^{m-1} |D(u-k_j)_+ \varphi|^p \, \mathrm{d}x \mathrm{d}t \ge \left(\frac{\mu_+}{2}\right)^{m-1} \iint_{\mathcal{Q}_j^+} |Du|^p \, \mathrm{d}x \mathrm{d}t.$$

In view of $(u - k_i)_+ \le 2^{-j}\omega$ and $A = 2^{s_1 + q_*}\hat{B}$, we conclude that

$$\iint_{Q_A'} u^{m-1} (u-k_j)_+^p |D\varphi|^p \,\mathrm{d}x\mathrm{d}t \leq \gamma \frac{(\mu_+)^{m-1}}{R^p} \left(\frac{\omega}{2^j}\right)^p |Q_A|$$

and

$$\begin{split} \iint_{Q_A'} (u-k_j)_+^2 |\partial_t \varphi| \, \mathrm{d}x \mathrm{d}t &\leq \gamma \frac{(\mu_+)^{m-1} \omega^{p-2}}{A^{p-2} R^p} \left(\frac{\omega}{2^j}\right)^2 |Q_A| \\ &= \gamma \frac{(\mu_+)^{m-1}}{R^p} \left(\frac{\omega}{2^{s_1+q_*} \hat{R}}\right)^{p-2} \left(\frac{\omega}{2^j}\right)^2 |Q_A| \leq \gamma \frac{(\mu_+)^{m-1}}{R^p} \left(\frac{\omega}{2^j}\right)^p |Q_A|, \end{split}$$

since $j \le s_1 + q_*$ and $\hat{B} > 1$. Furthermore, we deal with the estimates for the lower order terms. In view of $u - k_j \le 2^{-j}\omega$, we find that

$$\begin{split} \iint_{\mathcal{Q}_{A}^{\prime}} |f|(u-k_{j})_{+} \, \mathrm{d}x \mathrm{d}t &\leq \gamma \left(\frac{\omega}{2^{j}R^{p}}\right) \left(R^{p-n} \int_{B_{R}} |f| \, \mathrm{d}x\right) |Q_{A}| \\ &= \gamma \left(\frac{(\mu_{+})^{m-1}}{R^{p}} \left(\frac{\omega}{2^{j}}\right)^{p} |Q_{A}|\right) \left[\left(\frac{\omega}{2^{j}}\right)^{1-p} (\mu_{+})^{1-m}R^{p-n} \int_{B_{R}} |f| \, \mathrm{d}x\right] \\ &\leq \gamma \frac{(\mu_{+})^{m-1}}{R^{p}} \left(\frac{\omega}{2^{j}}\right)^{p} |Q_{A}|, \end{split}$$

provided that we assume

$$\omega \ge 2^{j}(\mu_{+})^{\frac{1-m}{p-1}} \left(R^{p-n} \int_{B_{p}} |f| \, \mathrm{d}x \right)^{\frac{1}{p-1}}. \tag{4.36}$$

Finally, we deduce that

$$\begin{split} \iint_{Q_A'} g^{\frac{p}{p-1}} \, \mathrm{d}x \mathrm{d}t &\leq \gamma \left(\frac{(\mu_+)^{m-1}}{R^p} \left(\frac{\omega}{2^j} \right)^p |Q_A| \right) \left[\left(\frac{\omega}{2^j} \right)^{-p} (\mu_+)^{1-m} R^{p-n} \int_{B_R} g^{\frac{p}{p-1}} \, \mathrm{d}x \right] \\ &\leq \gamma \frac{(\mu_+)^{m-1}}{R^p} \left(\frac{\omega}{2^j} \right)^p |Q_A|, \end{split}$$

provided that we assume

$$\omega \ge 2^{j} (\mu_{+})^{\frac{1-m}{p}} \left(R^{p-n} \int_{B_{R}} g^{\frac{p}{p-1}} \, \mathrm{d}x \right)^{\frac{1}{p}}. \tag{4.37}$$

Combining the above estimates, we conclude that there exists a constant γ depending only upon the data such that

$$\iint_{Q_{i}^{+}} |Du|^{p} \, \mathrm{d}x \mathrm{d}t \le \gamma \frac{1}{R^{p}} \left(\frac{\omega}{2^{j}}\right)^{p} |Q_{A}|. \tag{4.38}$$

To proceed further, we apply a De Giorgi type Lemma (see [6, Lemma 2.2]) to the function $u(\cdot,t)$, for all $-\frac{1}{2}A^{p-2}(\mu_+)^{1-m}\omega^{2-p}R^p \le t \le 0$, and with $l=\mu_+-2^{-(j+1)}\omega$ and $k=\mu_+-2^{-j}\omega$. According to (4.35), we deduce that

$$\left|\left\{x \in B_{\frac{3}{4}R} : u(x,t) \leq \mu_+ - \frac{\omega}{2^j}\right\}\right| = |B_{\frac{3}{4}R}| - |A_j(t)| \geq \left(\frac{\nu_0}{2}\right)^2 |B_{\frac{3}{4}R}|$$

holds for all $t \in \left(-\frac{1}{2}A^{p-2}(\mu_+)^{1-m}\omega^{2-p}R^p, 0\right]$. This yields that the estimate

$$\frac{\omega}{2^{j+1}}|A_{j+1}(t)| \leq \frac{\gamma}{\nu_0^2} \frac{R^{n+1}}{|B_R|} \int_{A_s(t) \setminus A_{s+1}(t)} |Du| \, \mathrm{d}x \leq \gamma \frac{R}{\nu_0^2} \int_{A_s(t) \setminus A_{s+1}(t)} |Du| \, \mathrm{d}x$$

holds for any $-\frac{1}{2}A^{p-2}(\mu_+)^{1-m}\omega^{2-p}R^p \le t \le 0$. Integrating the preceding inequality over the interval $\left(-\frac{1}{2}A^{p-2}(\mu_+)^{1-m}\omega^{2-p}R^p,0\right]$, we conclude from (4.38) that

$$\begin{split} \frac{\omega}{2^{j+1}} |Q_{j+1}^+| &\leq \frac{\gamma R}{v_0^2} \iint_{Q_j^+ \setminus Q_{j+1}^+} |Du| \, \mathrm{d}x \mathrm{d}t \\ &\leq \frac{\gamma R}{v_0^2} \left(\iint_{Q_j^+ \setminus Q_{j+1}^+} |Du|^p \, \mathrm{d}x \mathrm{d}t \right)^{\frac{1}{p}} |Q_j^+ \setminus Q_{j+1}^+|^{\frac{p-1}{p}} \\ &\leq \frac{\gamma}{v_0^2} \left(\frac{\omega}{2^j} \right) |Q_A|^{\frac{1}{p}} |Q_j^+ \setminus Q_{j+1}^+|^{\frac{p-1}{p}}. \end{split}$$

It follows that

$$|Q_{j+1}^+|^{\frac{p}{p-1}} \leq \gamma \nu_0^{-\frac{2p}{p-1}} |Q_A|^{\frac{1}{p-1}} |Q_j^+ \setminus Q_{j+1}^+|.$$

Furthermore, we add up these inequalities for $j = s_1, s_1 + 1, \dots, s_1 + q_* - 1$ and deduce that

$$(q_*-1)|Q_{s_1+q_*}^+|^{\frac{p}{p-1}} \le \gamma v_0^{-\frac{2p}{p-1}}|Q_A|^{\frac{p}{p-1}},$$

where the constant γ depends only upon the data. From the definition of Q_j^+ , the above inequality reads

$$\left|\left\{(x,t)\in Q_A: u(x,t)>\mu_+-\frac{\omega}{2^{s_1+q_*}}\right\}\right|\leq \frac{\gamma}{\nu_0^2}\frac{1}{(q_*-1)^{\frac{p-1}{p}}}|Q_A|.$$

At this point, we take $q_* = q_*(\text{data}, \bar{\nu}) \ge 2$ according to

$$q_* = q_*(\bar{\nu}) = n + 2 + \left(\frac{\gamma}{\nu_0^2 \bar{\nu}}\right)^{\frac{p}{p-1}}.$$
 (4.39)

For such a choice of q_* , the inequality (4.33) holds true. We also determine the value of j in (4.36) and (4.37) by $j = s_1 + q_*$. On the other hand, if the inequality (4.34), (4.36) or (4.37) is violated, then we deduce the inequality (4.32). We have thus proved the lemma.

We are now in a position to prove the decay estimate for the oscillation of the weak solution for the second alternative. The next proposition is our main result in this section.

Proposition 4.5. Let $\tilde{Q}_0 = Q\left(\frac{1}{16}R, (\mu_+)^{1-m}\omega^{2-p}\left(\frac{1}{16}R\right)^p\right)$ and let u be a bounded nonnegative weak solution to (1.1)-(1.2) in Ω_T . There exist $0 < \xi_2 < 2^{-5}$ and $B_2 > 1$ depending only upon the data such that

ess osc
$$u \le (1 - 6^{-1}\xi_2)\omega + B_2\xi_2^{-1}(F_1(2R)^{\frac{p}{p+m-1}} + F_2(2R)^{\frac{p-1}{p+m-2}}).$$

Proof. Let v_2 be the constant claimed in Lemma 4.3. We take $\bar{v} = v_2$ in Lemma 4.4 and this fixes q_* via (4.39). Moreover, we fix $\xi_2 = 2^{-(s_1+q_*)}$ and $A = \xi_2^{-1}\hat{B}$. We first assume that (4.2), (4.7) and (4.32) are violated, then we infer from Lemma 4.3 that

$$\operatorname{ess osc}_{\tilde{O}_0} u \le (1 - 6^{-1} \xi_2) \omega.$$

On the other hand, if either (4.2), (4.7) or (4.32) holds, then we deduce that either

$$\omega \leq 2^{\frac{2s_1+n}{p}}(\mu_+)^{\frac{1-m}{p}}F_1(2R), \quad \omega \leq 2^{\frac{s_1+n}{p-1}}(\mu_+)^{\frac{1-m}{p-1}}F_2(2R), \quad \xi_2\omega \leq \hat{B}(\mu_+)^{\frac{1-m}{p}}F_1(2R),$$

$$\xi_2\omega \leq \hat{B}(\mu_+)^{\frac{1-m}{p-1}}F_2(2R), \quad \omega \leq 2^{s_1+q_*}(\mu_+)^{\frac{1-m}{p}}F_1(2R) \quad \text{or} \quad \omega \leq 2^{s_1+q_*}(\mu_+)^{\frac{1-m}{p-1}}F_2(2R).$$

In view of $\mu_+ \ge \frac{1}{2}\omega$, we conclude that there exists a constant $B_2 > 1$ depending only upon the data such that

$$\operatorname{ess \ osc}_{\bar{O}_0} u \le \omega \le B_2 \xi_2^{-1} (F_1(2R)^{\frac{p}{p+m-1}} + F_2(2R)^{\frac{p-1}{p+m-2}}).$$

The proof of the proposition is now complete.

5. Proof of the main result

This section is devoted to give the final part of the proof of Theorem 1.2. Our proof follows the idea from [3] and [13] but the argument is considerably more delicate in our setting. First, we set up an iteration scheme. Let $\{\xi_1, B_1\}$ and $\{\xi_2, B_2\}$ are the constants claimed by Proposition 3.8 and Proposition 4.5, respectively. Let $\omega_0 = \omega$, $\mu_0 = \mu_+$, $R_0 = R$ and $Q_0 = Q_R = B_R \times (-R^{p-\varepsilon_0}, 0)$. Moreover, we set

$$\omega_1 = \eta \omega_0 + \gamma [F_1(2R_0)^{\frac{p}{p+m-1}} + F_2(2R_0)^{\frac{p-1}{p+m-2}}] \quad \text{and} \quad \mu_1 = \operatorname{ess\,sup}_{O_1} u, \tag{5.1}$$

where $\eta = \max \left\{ 1 - 2^{-1} \xi_1, \ 1 - 6^{-1} \xi_2 \right\} < 1$,

$$\gamma = \max \left\{ B_1 \xi_1^{-1}, \ B_2 \xi_2^{-1} \right\}, \quad \text{and} \quad Q_1 = B_{\frac{1}{16}R_0} \times \left(-\mu_0^{1-m} \omega_0^{2-p} \left(\frac{1}{16} R_0 \right)^p, 0 \right).$$

According to Proposition 3.8 and Proposition 4.5, we conclude that

$$\operatorname{ess \, osc}_{Q_1} u = \operatorname{ess \, sup}_{Q_1} u - \operatorname{ess \, inf}_{Q_1} u \le \omega_1. \tag{5.2}$$

Our task now is to construct a smaller cylinder Q_2 under the assumption that $\operatorname{ess} \operatorname{osc}_{Q_1} u \geq \frac{1}{2}\omega_1$. We first claim that $\mu_0 \leq \frac{4}{\eta}\mu_1$. In the case $\mu_- \leq \frac{1}{2}\mu_+$, we have $\mu_+ \leq \omega + \mu_- \leq \omega + \frac{1}{2}\mu_+$ and hence $\mu_+ \leq 2\omega = 2\omega_0$. In view of (5.1), we find that $\mu_0 = \mu_+ \leq \frac{2}{\eta}\omega_1 \leq \frac{4}{\eta}\mu_1$. In the case $\mu_- > \frac{1}{2}\mu_+$, we have $\mu_1 \geq \mu_- > \frac{1}{2}\mu_+$. At this point, we set

$$\delta = \frac{1}{16} 2^{-\frac{3m}{p}} \eta^{\frac{m+p-3}{p}} A^{\frac{2-p}{p}}, \qquad R_1 = \delta R_0 \qquad \text{and} \qquad Q_2 = B_{\frac{1}{16}R_1} \times \left(-\mu_1^{1-m} \omega_1^{2-p} \left(\frac{1}{16}R_1\right)^p, 0\right).$$

In view of $\mu_0 \le \frac{4}{n}\mu_1$ and $\omega_1 \ge \eta\omega_0$, we conclude that

$$\widehat{Q}_{2} := B_{R_{1}} \times \left(-\mu_{1}^{1-m} \left(\frac{\omega_{1}}{A}\right)^{2-p} R_{1}^{p}, 0\right) \subseteq B_{\frac{1}{16}R_{0}} \times \left(-\frac{1}{2} \mu_{0}^{1-m} \omega_{0}^{2-p} \left(\frac{1}{16} R_{0}\right)^{p}, 0\right). \tag{5.3}$$

Next, we claim that

ess osc
$$u \le \omega_2$$
, where $\omega_2 = \eta \omega_1 + \gamma [F_1(2R_1)^{\frac{p}{p+m-1}} + F_2(2R_1)^{\frac{p-1}{p+m-2}}].$ (5.4)

The claim can be proved in the same way as shown before. In fact, we observe that the nonuniformly parabolic cylinders

$$Q_{j}(\bar{l}) = B_{j} \times (t_{1} - (\mu_{+})^{1-m}(l_{j} - \bar{l})^{2-p}r_{j}^{p}, t_{1}), \qquad \bar{l} = \frac{1}{2}l_{j} + \frac{1}{16}B\alpha_{j} + \frac{1}{32}\omega + \frac{1}{2}\mu_{-}$$
and
$$Q_{j} = Q_{j}(l_{j+1}) = B_{j} \times (t_{1} - (\mu_{+})^{1-m}(l_{j} - l_{j+1})^{2-p}r_{j}^{p}, t_{1})$$

constructed in the proof of Lemma 3.5 are contained in the cylinder \hat{Q} , where $\hat{Q} = B_R \times (-(\mu_+)^{1-m}A^{p-2}\omega^{2-p}R^p, 0)$. Similarly, we find that the cylinders

$$Q_0(\bar{l}) = B_0 \times (t_1 - (\mu_+)^{1-m}(\bar{l} - l_0)^{2-p}r_0^p, t_1), \qquad \bar{l} = \mu_+ - \frac{1}{2}\xi\omega - \frac{1}{4}\hat{B}\alpha_0$$
and
$$Q_j = Q_j(l_{j+1}) = B_j \times (t_1 - (\mu_+)^{1-m}(l_{j+1} - l_j)^{2-p}r_j^p, t_1)$$

considered in the proof of Lemma 4.3 are also contained in the cylinder \hat{Q} . Consequently, the statements of Proposition 3.8 and Proposition 4.5 are still valid if we modify the settings of μ_+ and μ_- by

$$\mu_+ = \operatorname{ess\,sup} u$$
 and $\mu_- = \operatorname{ess\,inf} u$.

Thus we arrive at the conclusion that (5.4) holds. At this point, we define $R_k = \delta R_{k-1} = \delta^k R_0$ and

$$\omega_k = \eta \omega_{k-1} + \gamma [F_1(2R_{k-1})^{\frac{p}{p+m-1}} + F_2(2R_{k-1})^{\frac{p-1}{p+m-2}}].$$

We first observe that $\omega_{k+1} \ge \frac{1}{2}\omega_k$ for any $k \ge 0$. Another step in the proof is to show that $\omega_k \le \omega_0$ and $\omega_k \to 0$ as $k \to \infty$. For simplicity of notation, we write $F_k = F_1(2R_k)^{\frac{p}{p+m-1}} + F_2(2R_k)^{\frac{p-1}{p+m-2}}$. We see that $F_k \downarrow 0$ as $k \to \infty$. For any fixed integer $k \ge 0$, we infer that for n > k there holds

$$\omega_n = \eta^{n-k} \omega_k + \gamma \eta^n \sum_{j=k+1}^n \eta^{-j} F_{j-1} \le \eta^{n-k} \omega_k + \gamma \frac{1}{1-\eta} F_k.$$
 (5.5)

Then, we have $\omega_n \le \omega_0$, provided that we choose R > 0 so small that $F_0 \le (1 - \eta)\omega_0$. Passing to the limit $n \to \infty$, we conclude from (5.5) that

$$\lim_{n \to \infty} \omega_n \le \gamma \frac{1}{1 - \eta} F_k \tag{5.6}$$

holds for any fixed k > 1. Passing to the limit $k \to \infty$ in (5.6), we conclude that $\omega_k \to 0$ as $k \to \infty$. Next, we consider the case ess $\operatorname{osc}_{Q_1} u < \frac{1}{2}\omega_1$. Let $k_1 \ge 1$ be a smallest integer such that

$$\frac{1}{2}\omega_{k_1} < \operatorname*{ess\,osc}_{O_1} u < \omega_{k_1}. \tag{5.7}$$

At this point, we first define Q_{k_1+1} by

$$Q_{k_1+1} = B_{\frac{1}{16}R_{k_1}} \times \left(-\mu_1^{1-m}\omega_{k_1}^{2-p}\left(\frac{1}{16}R_{k_1}\right)^p, 0\right).$$

Taking into account that $\omega_{k_1} \ge \eta^{k_1} \omega_0$, $R_{k_1} = \delta^{k_1} R_0$ and

$$\begin{cases} \mu_0 \le 2\omega_0 \le 2\eta^{-k_1}\omega_{k_1} < 4\eta^{-k_1}\mu_1 & \text{if} \qquad \mu_- \le \frac{1}{2}\mu_+, \\ \mu_0 < 2\mu_- \le 2\mu_1 & \text{if} \qquad \mu_- > \frac{1}{2}\mu_+. \end{cases}$$

It follows that

$$\mu_{1}^{1-m} \left(\frac{\omega_{k_{1}}}{A}\right)^{2-p} R_{k_{1}}^{p} \leq 4^{m-1} \eta^{k_{1}(1-m)} \eta^{k_{1}(2-p)} A^{p-2} \mu_{0}^{1-m} \omega_{0}^{2-p} \left(\delta^{k_{1}} R_{0}\right)^{p}$$

$$\leq \frac{1}{2} \mu_{0}^{1-m} \omega_{0}^{2-p} \left(\frac{1}{16} R_{0}\right)^{p},$$

$$(5.8)$$

since $\delta = \frac{1}{16} 2^{-\frac{3m}{p}} \eta^{\frac{m+p-3}{p}} A^{\frac{2-p}{p}}$. Consequently, we infer that

$$\widehat{Q}_{k_1+1} := B_{R_{k_1}} \times \left(-\mu_1^{1-m} \left(\frac{\omega_{k_1}}{A} \right)^{2-p} R_{k_1}^p, 0 \right) \subseteq B_{\frac{1}{16}R_0} \times \left(-\frac{1}{2} \mu_0^{1-m} \omega_0^{2-p} \left(\frac{1}{16} R_0 \right)^p, 0 \right).$$

Taking into account (5.7) and (5.8), and using the same argument as in the proof of (5.4), we deduce that $\operatorname{ess} \operatorname{osc}_{Q_{k_1+1}} u \leq \omega_{k_1+1}$. Furthermore, we introduce the cylinders Q_2, \dots, Q_{k_1}

by

$$Q_{2} = B_{\frac{1}{16}R_{1}} \times \left(-(\mu^{(2)})^{1-m} (\omega^{(2)})^{2-p} \left(\frac{1}{16}R_{1} \right)^{p}, 0 \right),$$

$$Q_{3} = B_{\frac{1}{16}R_{2}} \times \left(-(\mu^{(3)})^{1-m} (\omega^{(3)})^{2-p} \left(\frac{1}{16}R_{2} \right)^{p}, 0 \right),$$

$$Q_{k_1} = B_{\frac{1}{16}R_{k_1-1}} \times \left(-(\mu^{(k_1)})^{1-m} (\omega^{(k_1)})^{2-p} \left(\frac{1}{16}R_{k_1-1} \right)^p, 0 \right),$$

where $\mu^{(2)} = \eta \mu_0$, $\mu^{(3)} = \eta^2 \mu_0$, \cdots , $\mu^{(k_1)} = \eta^{k_1-1} \mu_0$ and $\omega^{(2)} = \eta \omega_0$, $\omega^{(3)} = \eta^2 \omega_0$, \cdots , $\omega^{(k_1)} = \eta^{k_1-1} \omega_0$. It is easy to check that $Q_{k_1+1} \subset Q_{k_1} \subset \cdots \subset Q_1$. Taking into account that $k_1 \geq 1$ is the smallest integer such that (5.7) holds and $\omega_{j+1} \geq \frac{1}{2}\omega_j$ for any $j \geq 0$, we conclude that ess $\operatorname{osc}_{Q_1} u < \frac{1}{2}\omega_1 \leq \omega_2$ and hence

$$\operatorname{ess \, osc}_{O_1} u \leq \min\{\omega_1, \omega_2, \cdots, \omega_{k_1}\}.$$

Consequently, we infer that

$$\operatorname{ess \ osc} u \leq \operatorname{ess \ osc} u \leq \omega_{2}, \quad \operatorname{ess \ osc} u \leq \operatorname{ess \ osc} u \leq \omega_{3},$$

$$\cdots, \quad \operatorname{ess \ osc} u \leq \operatorname{ess \ osc} u \leq \omega_{k_{1}}.$$

$$Q_{1}$$

We may now repeat the same arguments as in the previous proof and conclude that there exist two sequences $\omega^{(n)}$, $\mu^{(n)}$ and a sequence of cylinders

$$Q_n = B_{\frac{1}{16}R_{n-1}} \times \left(-(\mu^{(n)})^{1-m} (\omega^{(n)})^{2-p} \left(\frac{1}{16}R_{n-1} \right)^p, 0 \right)$$

such that $\omega^{(n)} \leq \omega_0$, $\mu^{(n)} \leq \mu_0$, $Q_0 \supset Q_1 \supset \cdots \supset Q_n \supset \cdots$ and there holds

$$\operatorname{ess} \operatorname{osc} u \leq \omega_n$$

for all $n=0,1,\cdots$. We are now in a position to establish a continuity estimate for the weak solutions. To this end, we fix $\bar{\varrho} \in (0,R_0)$, $\bar{r} \in (0,\bar{\varrho})$ and $0 < b < \min\{\delta,\eta\}$. Then, there exist two integers $k \in \mathbb{N}$ and $l \in \mathbb{N}$ such that

$$k-1 < \frac{\ln\frac{\varrho}{R_0}}{\ln\delta} \le k$$
 and $l-1 < \frac{\ln\frac{\bar{l}}{\varrho}}{\ln b} \le l.$ (5.9)

Moreover, we set n = k + l and observe that

$$\eta^{n-k} = \exp(\alpha l \ln b) = b^{\alpha l} \le \left(\frac{\bar{r}}{\bar{\varrho}}\right)^{\alpha}, \quad \text{where} \quad \alpha = \frac{\ln \eta}{\ln b}.$$

It follows from (5.5) that the estimate

$$\omega_n \le \left(\frac{\bar{r}}{\bar{\varrho}}\right)^{\alpha} \omega_0 + \gamma \frac{1}{1 - \eta} \left(F_1(2\bar{\varrho})^{\frac{p}{p + m - 1}} + F_2(2\bar{\varrho})^{\frac{p - 1}{p + m - 2}} \right). \tag{5.10}$$

holds for any $0 < \bar{r} < \bar{\varrho} < R_0$ and n = k + l, where k and l satisfy (5.9). Furthermore, we take $0 < \bar{r} < R_0$ and specify $\bar{\varrho} = R_0^{1-\beta} \bar{r}^{\beta}$. This also determines k_* and l_* via (5.9) by

$$k_* - 1 < \frac{\beta}{\ln \delta} \ln \frac{\bar{r}}{R_0} \le k_* \quad \text{and} \quad l_* - 1 < \frac{(1 - \beta)}{\ln b} \ln \frac{\bar{r}}{R_0} \le l_*.$$

For $n_* = k_* + l_*$, we find that the estimate (5.10) reads

$$\omega_{n_*} \leq \left(\frac{\bar{r}}{R_0}\right)^{\alpha(1-\beta)} \omega_0 + \gamma \left(F_1(2R_0^{1-\beta}\bar{r}^\beta)^{\frac{p}{p+m-1}} + F_2(2R_0^{1-\beta}\bar{r}^\beta)^{\frac{p-1}{p+m-2}}\right)$$

and n_* satisfies

$$n_* - 2 < \left(\frac{\beta}{\ln \delta} + \frac{1 - \beta}{\ln b}\right) \ln \frac{\bar{r}}{R_0} \le n_*. \tag{5.11}$$

At this stage, we set $a_0 = \mu_0^{1-m} \omega_0^{2-p}$ and $\bar{Q}_n = B_{R_n} \times (-a_0 R_n^p, 0)$. Recalling that $\omega^{(n)} \le \omega_0$ and $\mu^{(n)} \le \mu_0$, we see that $\bar{Q}_n \subset Q_n$. Moreover, we define

$$\sigma = \beta + (1 - \beta) \frac{\ln \delta}{\ln b}$$
 and $r = \delta^2 R_0 \left(\frac{\bar{r}}{R_0}\right)^{\sigma}$.

This yields that

$$\bar{r} = R_0 \left(\delta^{-2} R_0^{-1} r \right)^{\frac{1}{\sigma}} \quad \text{and} \quad R_0^{1-\beta} \bar{r}^{\beta} = \delta^{-2 \frac{\beta}{\beta + \frac{1-\beta}{\ln b} \ln \delta}} r^{\frac{\beta}{\sigma}} R_0^{\frac{(1-\beta) \ln \delta}{\sigma \ln b}} \leq c(\delta) r^{\frac{\beta}{\sigma}}.$$

In view of (5.11), we conclude with

$$r = R_0 \exp\left(\sigma \ln \frac{\bar{r}}{R_0} + 2\ln \delta\right) < R_0 e^{n_* \ln \delta} = R_{n_*}$$

and hence $Q(r, a_0 r^p) \subset \bar{Q}_{n_*} \subset Q_{n_*}$. Consequently, we arrive at the continuity estimate as follows

$$\operatorname{ess \ osc}_{Q(r,a_0r^p)} u \leq \operatorname{ess \ osc}_{\bar{Q}_{n_*}} u \leq \operatorname{ess \ osc}_{Q_{n_*}} u \leq \omega_{n_*}$$

$$\leq \gamma \left(\frac{r}{R_0}\right)^{\frac{\alpha(1-\beta)}{\sigma}} \omega_0 + \gamma \left(F_1(cr^{\frac{\beta}{\sigma}})^{\frac{p}{p+m-1}} + F_2(cr^{\frac{\beta}{\sigma}})^{\frac{p-1}{p+m-2}}\right).$$

Thus we are led to the conclusion that the weak solution to (1.1)-(1.2) is locally continuous in Ω_T .

ACKNOWLEDGEMENTS

The author wishes to thank the Institute for Advanced Study in Mathematics of Harbin Institute of Technology for the kind hospitality during his visit in July 2023.

References

- [1] G. I. Barenblatt, V. M. Entov and V. M. Rizhnik: Motion of fluids and gases in natural strata, Nedra. Moscow, (1984).
- [2] V. Bögelein, F. Duzaar and U. Gianazza: Porous medium type equations with measure data and potential estimates, SIAM J. Math. Anal., 45(6), 3283-3330, (2013).
- [3] V. Bögelein, F. Duzaar and U. Gianazza: Continuity estimates for porous medium type equations with measure data, J. Funct. Anal., 267, 3351-3396, (2014).
- [4] V. Bögelein, F. Duzaar, N. Liao and L. Schätzler: On the Hölder regularity of signed solutions to a doubly nonlinear equation. Part II, Rev. Mat. Iberoam., 39(3), 1005-1037, (2022).
- [5] V. Bögelein, A. Heran, L. Schätzler and T. Singer: Harnack's inequality for doubly nonlinear equations of slow diffusion type, Calc. Var., 60, 215 (2021).
- [6] E. DiBenedetto: Degenerate Parabolic Equations. Universitext, Springer, New York (1993).
- [7] L. C. Evans and R. F. Gariepy: Measure theory and fine properties of functions, Textbooks in Mathematics, Revised edition. CRC Press (2015).

- [8] S. Fornaro and M. Sosio: Intrinsic Harnack estimates for some doubly nonlinear degenerate parabolic equations, Adv. Differ. Equ., 13(1-2), 139-168, (2008).
- [9] A. V. Ivanov: Hölder estimates for quasilinear doubly degenerate parabolic equations, Zap. Nauchn. Sem. LOMI., 171, 70-105, (1989).
- [10] T. Kilpeläinen and J. Malý: The Wiener test and potential estimates for quasilinear elliptic equations, Acta Math., 172, 137-161, (1993).
- [11] V. Liskevich and I. I. Skrypnik: Harnack inequality and continuity of solutions to quasi-linear degenerate parabolic equations with coefficients from Kato-type classes, J. Diff. Equa., 247(10), 2740-2777, (2009).
- [12] V. Liskevich, I. I. Skrypnik and Z. Sobol: Estimates of solutions for the parabolic *p*-Laplacian equation with measure via parabolic nonlinear potentials, Commun. Pure Appl. Anal., 12(4), 1731-1744, (2013).
- [13] M. M. Porzio and V. Vespri: Hölder estimates for local solutions of some doubly nonlinear degenerate parabolic equations, J. Differ. Equ., 103 146-178, (1993).
- [14] I. I. Skrypnik: Continuity of solutions to singular parabolic equations with coefficients from Kato-type classes. Annali di Matematica, 195, 1153-1176, (2016).
- [15] E. Stein: Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals, Princeton Math. Series, vol. 43. Princeton University Press, Princeton (1993).
- [16] S. Sturm: Pointwise estimates via parabolic potentials for a class of doubly nonlinear parabolic equations with measure data, Manuscr. Math., 157, 295-322, (2018).
- [17] V. Vespri: Harnack type inequalities for solutions of certain doubly nonlinear parabolic equations, J. Math. Anal. Appl., 181(1), 104-131, (1994).
- [18] Y. Zozulia: Pointwise estimates of solutions to weighted porous medium and fast diffusion equations via weighted Riesz potentials, Ukr. Mat. Bull., 17(1), 116-144, (2020).
- [19] Y. Zozulia: On the continuity of solutions of the equations of a porous medium and the fast diffusion with weighted and singular lower-order terms, J. Math. Sci., 256, 803-830, (2021).

* DEPARTMENT OF MATHEMATICS
SCHOOL OF SCIENCES
WUHAN UNIVERSITY OF TECHNOLOGY
430070, 122 Luoshi Road, Wuhan, Hubei
P. R. China

Email address: qifan_li@yahoo.com, qifan_li@whut.edu.cn