

# Linearized Analysis of Adiabatic Oscillations of Rotating Gaseous Stars

Tetu Makino \*

February 27, 2024

## Abstract

We study adiabatic oscillations of rotating self-gravitating gaseous stars in mathematically rigorous manner. The internal motion of the star is supposed to be governed by the Euler-Poisson equations with rotation of constant angular velocity under the equation of state of the ideal gas. The motion is supposed to be adiabatic, but not to be barotropic in general. This causes a free boundary problem to gas-vacuum interface. Existence of solutions to the linearized equation in the Lagrangian coordinates of the perturbations around a fixed stationary solution, the eigenvalue problem with concept of quadratic pencil of operators, and the stability problem with a new concept of stability introduced in this article are discussed.

*Key Words and Phrases.* Gaseous star, adiabatic oscillation, Euler-Poisson equations, Hille-Yoshida theory, quadratic pencil, stability

*2020 Mathematical Subject Classification Numbers.* 3610, 35Q65, 35P05, 35R35, 35B35, 47N20, 76N15, 76U05, 85A30.

## 1 Introduction

We consider the motion of a rotating gaseous star governed by the Euler-Poisson equation:

$$\frac{D\rho}{Dt} + \rho \operatorname{div}_{\mathbf{x}} \mathbf{v} = 0, \quad (1.1a)$$

$$\rho \left[ \frac{D\mathbf{v}}{Dt} + 2\boldsymbol{\Omega} \times \mathbf{v} + \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{x}) \right] + \nabla_{\mathbf{x}} P + \rho \nabla_{\mathbf{x}} \Phi = 0, \quad (1.1b)$$

$$\rho \frac{DS}{Dt} = 0, \quad (1.1c)$$

$$\Delta \Phi = 4\pi G \rho \quad (1.1d)$$

---

\*Professor Emeritus at Yamaguchi University, Japan; E-mail: makino@yamaguchi-u.ac.jp

on  $(t, \mathbf{x}) = (t, x^1, x^2, x^3) \in [0, +\infty[ \times \mathbb{R}^3$ . We are denoting

$$\nabla_{\mathbf{x}} = \begin{bmatrix} \frac{\partial}{\partial x^1} \\ \frac{\partial}{\partial x^2} \\ \frac{\partial}{\partial x^3} \end{bmatrix}, \quad \frac{D}{Dt} = \frac{\partial}{\partial t} + \sum_{k=1}^3 v^k \frac{\partial}{\partial x^k}, \quad \Delta = \sum_{k=1}^3 \left( \frac{\partial}{\partial x^k} \right)^2. \quad (1.2)$$

The unknown variables  $\rho, P, S, \mathbf{v} = (v^1, v^2, v^3)^\top$  are density, the pressure, the specific entropy, and the velocity field, while

$$\mathbf{\Omega} = \Omega \frac{\partial}{\partial x^3}, \quad (1.3)$$

$\Omega$  being a constant, the angular velocity of the rotation around the  $x^3$ -axis. Hereafter we denote

$$B = 2\Omega \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad (1.4)$$

so that

$$B\mathbf{v} = 2\mathbf{\Omega} \times \mathbf{v} = 2\Omega \begin{bmatrix} -v^2 \\ v^1 \\ 0 \end{bmatrix}. \quad (1.5)$$

On the other hand  $\Phi$  is the gravitational potential and  $G$  is a positive constant, while we suppose that the support of  $\rho(t, \cdot)$  is compact for  $\forall t$  and replace the Poisson equation (1.1d) by the Newton potential

$$\Phi(t, \mathbf{x}) = -4\pi G \mathcal{K}[\rho(t, \cdot)](\mathbf{x}), \quad (1.6)$$

where

$$\mathcal{K}[f](\mathbf{x}) := \frac{1}{4\pi} \int \frac{f(\mathbf{x}')}{\|\mathbf{x} - \mathbf{x}'\|} d\mathbf{x}'. \quad (1.7)$$

The pressure  $P$  is supposed to be a prescribed function  $\mathcal{P}(\rho, S)$  of  $\rho, S$ , where  $\mathcal{P} \in C^1([0, +\infty[ \times \mathbb{R})$  and  $\mathcal{P} > 0$ ,  $\frac{\partial \mathcal{P}}{\partial \rho} > 0$  for  $\rho > 0$  and  $\mathcal{P}(0, S) = 0$ . So we read

$$P = \mathcal{P}(\rho, S) \quad (1.8)$$

in (1.1b).

We consider the initial data  $\rho^0, S^0, \mathbf{v}^0$  for which the motion enjoys

$$\rho(0, \mathbf{x}) = \rho^0(\mathbf{x}), \quad S(0, \mathbf{x}) = S^0(\mathbf{x}), \quad \mathbf{v}(0, \mathbf{x}) = \mathbf{v}^0(\mathbf{x}). \quad (1.9)$$

We denote

$$P^0(\mathbf{x}) = \mathcal{P}(\rho^0(\mathbf{x}), S^0(\mathbf{x})), \quad \Phi^0(\mathbf{x}) = -4\pi G \mathcal{K}[\rho^0](\mathbf{x}). \quad (1.10)$$

We suppose that there is fixed an axially symmetric stationary solution  $(\rho, S, \mathbf{v}) = (\rho_b(\mathbf{x}), S_b(\mathbf{x}), \mathbf{v}_b(\mathbf{x}))$ , which satisfy (1.1a), (1.1b), (1.1c), (1.6), such that  $\mathfrak{R}_b = \{\mathbf{x} \in \mathbb{R}^3 | \rho_b(\mathbf{x}) > 0\}$  is a bounded domain. The existence of stationary solutions is discussed in Section 2.

We consider small perturbations at this fixed stationary solution. The linearized equation which governs the perturbations, which will be called “the equation of LASO ( linear adiabatic stellar oscillations )”, turns out to be

$$\frac{\partial^2 \mathbf{u}}{\partial t^2} + B \frac{\partial \mathbf{u}}{\partial t} + \mathcal{L} \mathbf{u} = 0 \quad \text{on} \quad [0, +\infty[ \times \mathfrak{R}_b, \quad (1.11)$$

where

$$\mathcal{L} \mathbf{u} = \frac{1}{\rho_b} \nabla \mathfrak{d} P - \frac{\mathfrak{d} \rho}{\rho_b^2} \nabla P_b + \nabla \mathfrak{d} \Phi, \quad (1.12)$$

with

$$\mathfrak{d} \rho = -\text{div}(\rho_b \mathbf{u}), \quad (1.13a)$$

$$\mathfrak{d} S = -(\mathbf{u} | \nabla S_b), \quad (1.13b)$$

$$\mathfrak{d} P = \left( \frac{\partial P}{\partial \rho} \right)_b \mathfrak{d} \rho + \left( \frac{\partial P}{\partial S} \right)_b \mathfrak{d} S, \quad (1.13c)$$

$$\mathfrak{d} \Phi = -4\pi G \mathcal{K}[\mathfrak{d} \rho]. \quad (1.13d)$$

We are denoting

$$\left( \frac{\partial P}{\partial \rho} \right)_b = \partial_\rho \mathcal{P}(\rho_b, S_b), \quad \left( \frac{\partial P}{\partial S} \right)_b = \partial_S \mathcal{P}(\rho_b, S_b).$$

The unknown variable  $\mathbf{u}$  means

$$\mathbf{u}(t, \mathbf{x}) = \boldsymbol{\varphi}(t, \mathbf{x}) - \boldsymbol{\varphi}_b(t, \mathbf{x}) + \mathbf{u}^0 \quad (1.14)$$

where  $\mathbf{u}^0$  is a given vector field on  $\mathfrak{R}_b$  and  $\boldsymbol{\varphi}(t, \mathbf{x})$  is the solution of

$$\frac{\partial}{\partial t} \boldsymbol{\varphi}(t, \mathbf{x}) = \mathbf{v}(t, \boldsymbol{\varphi}(t, \mathbf{x})), \quad \boldsymbol{\varphi}(0, \mathbf{x}) = \mathbf{x}, \quad (1.15)$$

while  $\boldsymbol{\varphi}_b(t, \mathbf{x})$  is the solution of

$$\frac{\partial}{\partial t} \boldsymbol{\varphi}_b(t, \mathbf{x}) = \mathbf{v}_b(\boldsymbol{\varphi}_b(t, \mathbf{x})), \quad \boldsymbol{\varphi}_b(0, \mathbf{x}) = \mathbf{x}. \quad (1.16)$$

Here we suppose

$$\{\mathbf{x} \mid \rho^0(\mathbf{x}) > 0\} = \mathfrak{R}_b \quad (1.17)$$

and  $\mathbf{u}^0$  enjoys

$$\rho^0(\mathbf{x}) - \rho_b(\mathbf{x}) = -\text{div}(\rho_b(\mathbf{x}) \mathbf{u}^0(\mathbf{x})), \quad (1.18a)$$

$$S^0(\mathbf{x}) - S_b(\mathbf{x}) = -(\mathbf{u}^0(\mathbf{x}) | \nabla S_b(\mathbf{x})). \quad (1.18b)$$

The equation (1.11) is considered with the initial condition

$$\mathbf{u}|_{t=0} = \mathbf{u}^0, \quad \frac{\partial \mathbf{u}}{\partial t} \Big|_{t=0} = \mathbf{v}^0 \quad \text{on } \mathfrak{R}_b. \quad (1.19)$$

In Section 3 the derivation of this equation of LASO (1.11) is discribed.

In Section 4 we give the basic existence result to the initial value problem (1.11), (1.19), be realizing the integrodifferential operator  $\mathcal{L}$  as a selfadjoint operator  $\mathbf{L}$  in the Hilbert space  $\mathfrak{H} = L^2(\rho_b d\mathbf{x}, \mathfrak{R}_b)$ . This is done by applying the Hille-Yosida theory. The applied theorem is formulated and proved in **Appendix A** for the sake of selfcontainedness.

In Section 5 we discuss about the eigenvalue problem to the equation (1.11). The ‘eigenvalue’ of (1.11) is the square of the usual eigenvalue of  $-\mathbf{L}$  only when  $\Omega = 0, B = O$ . But, when  $\Omega \neq 0$ , we need the concept ‘quadratic pencil’ (after Bogner [3]) and its ‘spectrum’. The result of the structure of the ‘spectrum’ of the quadratic pencil by J. Dyson and B. F. Schultz [6] is justified.

In Section 6 we discuss about the stability of solutions of (1.11)(1.19). We propose a new concept of stability based on seminorms on the space of the values  $\mathbf{u}(t, \cdot)$  of the solution at instant  $t$ , taking into account that the magnitude of  $\mathbf{u}(t, \cdot)$  itself is not essential but the magnitude of the perturbation  $\mathfrak{d}\rho = -\text{div}(\rho_b \mathbf{u}), \mathfrak{d}\mathbf{v} = \partial \mathbf{u} / \partial t - \mathbf{v}_b^L$  is essential in the discussion of stability. Examples of the seminorms are presented with an open question.

In Section 7 the definition of Lagrange displacement and the equation of linear adiabatic stellar oscillations, traditionally used by astrophysicists, “equation of LASO ( $\blacklozenge$ )”, which are slightly deifferent from those, (1.14) (1.11), presented in this article, are described and their properties are discussed.

## 2 Existence of stationary solutions

In this section we establish the existence of stationary solutions which enjoy good properties used in the following consideration. In this section we suppose

**Assumption 1** *There are a function  $\mathcal{F} = \mathcal{F}(v, S) \in C^\infty([0, +\infty[ \times \mathbb{R})$  and a number  $\gamma$  such that  $1 < \gamma < 2$ ,  $\mathcal{F}(v, S) > 0$  for  $v \geq 0, |S| < \infty$ ,  $\mathcal{F} + \frac{\gamma-1}{\gamma} v \frac{\partial \mathcal{F}}{\partial v} > 0$  for  $v > 0, |S| < \infty$ ,  $\left| \frac{\partial}{\partial v} \log \mathcal{F} \right| \leq C$ , for which*

$$\mathcal{P}(\rho, S) = \rho^\gamma \mathcal{F}(\rho^{\gamma-1}, S) \quad \text{for } \rho \geq 0, |S| < \infty. \quad (2.1)$$

We keep in mind the case of  $\mathcal{F}(v, S) = \exp(S/C_V)$ ,  $C_V$  being a positive

constant, namely, the case when the equation of state (1.8) turns out to be

$$P = \rho^\gamma \exp\left(\frac{S}{C_V}\right). \quad (2.2)$$

Let us put the following

**Definition 1** *A triple of  $t$ -independent axially symmetric functions  $(\rho_b, S_b, \mathbf{v}_b) \in C_0^1(\mathbb{R}^3; [0, +\infty[) \times C^1(\mathbb{R}^3; \mathbb{R}) \times C^1(\mathbb{R}^3; \mathbb{R}^3)$  which satisfies (1.1a)(1.1b)(1.1c) with  $\Phi = \Phi_b$ ,  $P = P_b$  determined by (1.6) (1.8) with  $\rho = \rho_b$ ,  $S = S_b$  is called an admissible stationary solution, if*

0) *there is a function  $\omega_b(\varpi, z)$  of  $(\varpi, z) \in [0, +\infty[ \times \mathbb{R}$  such that*

$$\mathbf{v}_b(\mathbf{x}) = \omega_b(\varpi, z) \frac{\partial}{\partial x^3} \times \mathbf{x} = \omega_b(\varpi, z) \begin{bmatrix} -x^2 \\ x^1 \\ 0 \end{bmatrix}, \quad (2.3)$$

where

$$\varpi = \sqrt{(x^1)^2 + (x^2)^2}, \quad z = x^3 \quad \text{for } \mathbf{x} = (x^1, x^2, x^3), \quad (2.4)$$

and the function  $\mathbf{x} \mapsto \omega_b(\varpi, z)$  belongs to  $C^1(\mathbb{R}^3)$ ;

1)  $\rho_b(\mathbf{x}), S_b(\mathbf{x})$  are functions of  $(r, \zeta)$ , and  $\mathfrak{R}_b = \{\mathbf{x} | \rho_b(\mathbf{x}) > 0\}$  is of the form  $\{\mathbf{x} | r < R_b(\zeta)\}$ , where

$$r = \|\mathbf{x}\| = \sqrt{(x^1)^2 + (x^2)^2 + (x^3)^2}, \quad \zeta = \frac{x^3}{r} \quad \text{for } \mathbf{x} = (x^1, x^2, x^3), \quad (2.5)$$

and  $R_b(\zeta)$  is a continuous function of  $\zeta \in [-1, 1]$  such that  $R_b(\zeta) > 0 \forall \zeta \in [-1, 1]$ ;

2) *The boundary  $\partial\mathfrak{R}_b$  is of  $C^{3,\alpha}$ -class and the functions  $\rho_b^{\gamma-1}, S_b, \Phi_b = -4\pi G K[\rho_b] \in C^\infty(\mathfrak{R}_b) \cap C^{3,\alpha}(\mathfrak{R}_b \cup \partial\mathfrak{R}_b)$ ,  $\alpha$  being a positive number such that  $0 < \alpha < \left(\frac{1}{\gamma-1} - 1\right) \wedge 1 \left( := \min \left\{ \frac{1}{\gamma-1} - 1, 1 \right\} \right)$  ;*

3)  $\frac{\partial \rho_b}{\partial r}, \frac{\partial P_b}{\partial r} < 0$  in  $\mathfrak{R}_b$  and

$$\frac{\partial \rho_b}{\partial r} \leq -\frac{r}{C}, \quad \frac{\partial P}{\partial r} \leq -\frac{r}{C} \quad \text{for } 0 < r \ll 1; \quad (2.6)$$

4) *The boundary  $\partial\mathfrak{R}_b$ , on which  $\rho_b = 0$ , is a physical vacuum boundary, that is,*

$$0 < \frac{\partial}{\partial \mathbf{n}} \sqrt{\left(\frac{\partial P}{\partial \rho}\right)_b} < +\infty \quad \text{on } \partial\mathfrak{R}_b, \quad (2.7)$$

where  $\mathbf{n}$  stands for the unit normal vector at the boundary point directed inward to  $\mathfrak{R}_b$ . Here we note that  $\partial P / \partial \rho$  is the square of the sound speed.

Note that the condition 2) requires  $\|\nabla \rho_b\|, \|\nabla P_b\| \neq 0$  for  $r \neq 0$ .

**Definition 2** The admissible stationary solution  $(\rho_b, S_b, \mathbf{v}_b)$  is said to be ‘barotropic’, if  $\omega_b$  is independent of  $z$ , that is,

$$\mathbf{v}_b(\mathbf{x}) = \omega_b(\varpi) \begin{bmatrix} -x^2 \\ x^1 \\ 0 \end{bmatrix},$$

and there is a function  $\Pi \in C^{1,\alpha}(\mathbb{R})$  such that  $d\Pi(v)/v > 0$  and it holds

$$P_b(\mathbf{x}) = \Pi(\rho_b(\mathbf{x})^{\gamma-1}) \quad \text{for } \forall \mathbf{x} \in \mathfrak{R}_b. \quad (2.8)$$

Note that, then,

$$(\nabla P_b | \nabla \rho_b) = (\gamma - 1)v \frac{d\Pi}{dv} \Big|_{v=\rho_b^{\gamma-1}} \|\nabla \rho_b\|^2 > 0 \quad \text{for } r \neq 0. \quad (2.9)$$

**Remark 1** It is well-known as the Poincaré-Wavre theorem ([17, CHAPITRE II, 20, p.33]) that  $\frac{\partial \Omega_b}{\partial z} = 0$  if and only if  $P_b = f(\rho_b)$ , where  $\Omega_b(\varpi, z) = \Omega + \omega_b(\varpi, z)$ .

For admissible stationary solutions, let us put

**Definition 3** We denote

$$\Gamma := \frac{\rho}{P} \frac{\partial P}{\partial \rho} = \rho \frac{\partial_\rho \mathcal{P}(\rho, S)}{\mathcal{P}(\rho, S)}, \quad (2.10a)$$

$$\begin{aligned} \mathfrak{a} &:= -\left(\rho \frac{\partial P}{\partial \rho}\right)^{-1} \left(\frac{\partial P}{\partial S}\right) \nabla S = -\left(\frac{1}{\Gamma P} \frac{\partial P}{\partial S}\right) \nabla S \\ &= -\frac{1}{\Gamma P} \nabla P + \frac{1}{\rho} \nabla \rho, \end{aligned} \quad (2.10b)$$

$$\begin{aligned} \mathcal{A} &:= \frac{1}{\Gamma P} \frac{(\nabla P | \nabla \rho)}{\|\nabla \rho\|} - \frac{\|\nabla \rho\|}{\rho} \\ &= -(\mathfrak{a} | \mathbf{n}), \quad \text{with } \mathbf{n} := \frac{\nabla \rho}{\|\nabla \rho\|} \quad \text{wherever } \|\nabla \rho\| \neq 0. \end{aligned} \quad (2.10c)$$

and

$$\Gamma_b := \left(\frac{\rho}{P} \frac{\partial P}{\partial \rho}\right)_b = \rho_b \frac{\partial_\rho \mathcal{P}(\rho_b, S_b)}{\mathcal{P}(\rho_b, S_b)}, \quad (2.11a)$$

$$\begin{aligned} \mathfrak{a}_b &:= -\left(\rho \frac{\partial P}{\partial \rho}\right)_b^{-1} \left(\frac{\partial P}{\partial S}\right)_b \nabla S_b = -\left(\frac{1}{\Gamma_b P} \frac{\partial P}{\partial S}\right)_b \nabla S_b \\ &= -\frac{1}{\Gamma_b P_b} \nabla P_b + \frac{1}{\rho_b} \nabla \rho_b, \end{aligned} \quad (2.11b)$$

$$\begin{aligned} \mathcal{A}_b &:= \frac{1}{\Gamma_b P_b} \frac{(\nabla P_b | \nabla \rho_b)}{\|\nabla \rho_b\|} - \frac{\|\nabla \rho_b\|}{\rho_b} \\ &= -(\mathfrak{a}_b | \mathbf{n}), \quad \text{with } \mathbf{n} := \frac{\nabla \rho_b}{\|\nabla \rho_b\|} \quad \text{on } \mathfrak{R}_b \setminus \{O\}. \end{aligned} \quad (2.11c)$$

Since

$$\Gamma = \gamma + (\gamma - 1)v \frac{\partial}{\partial v} \log \mathcal{F}, \quad (2.12)$$

we have  $\Gamma_b \in C^\infty(\mathfrak{R}_b) \cap C^{3,\alpha}(\mathfrak{R}_b \cup \partial\mathfrak{R}_b)$ . Since

$$\frac{1}{\Gamma P} \frac{\partial P}{\partial S} = \frac{\partial_S \log \mathcal{F}}{\gamma + (\gamma - 1)v \frac{\partial}{\partial v} \log \mathcal{F}}, \quad (2.13)$$

we see  $\mathbf{a}_b \in C^\infty(\mathfrak{R}_b) \cap C^{2,\alpha}(\mathfrak{R}_b \cup \partial\mathfrak{R}_b)$ . And we see, at least,  $\mathcal{A}_b \in \bigcap_{0 < \delta \ll 1} C^{1,\alpha}(\mathfrak{R}_b \cup \partial\mathfrak{R}_b \setminus \{r < \delta\}) \cap L^\infty(\mathfrak{R}_b)$ . When  $(\rho_b, S_b)$  is equatorially symmetric moreover, we can claim  $\mathcal{A}_b \in C(\mathfrak{R} \cup \partial\mathfrak{R}_b)$  by evaluating  $\mathcal{A}_b(O) = 0$ .

If the admissible stationary soltion is barotropic, then we have

$$\mathbf{a}_b = -\mathcal{A}_b \mathbf{n} \quad \text{in } \mathfrak{R}_b, \quad (2.14)$$

and  $\mathcal{A}_b \in C^\alpha(\mathfrak{R} \cup \partial\mathfrak{R})$ , since

$$\mathcal{A}_b = \frac{\gamma - 1}{\Gamma P} v \frac{d\Pi}{dv} \Big|_{v=\rho_b^{\gamma-1}} \|\nabla \rho_b\| - \frac{\|\nabla \rho_b\|}{\rho_b}.$$

**Remark 2** When the stationary solution is barotropic, this does not mean that the perturbed motion to be considered should be barotropic. The perturbed state  $\rho = \rho_b + \mathfrak{d}\rho$ ,  $P = P_b + \mathfrak{d}P$  is approximately

$$\rho = \rho_b - \operatorname{div}(\rho_b \mathbf{u}), \quad P = P_b - \left( \frac{\Gamma P}{\rho} \right)_b \operatorname{div}(\rho_b \mathbf{u}) + (\Gamma P)_b ((\mathbf{u} | \mathbf{a}_b),$$

and  $\nabla \rho, \nabla P$  are not necessarily parallel.

**Remark 3** After [6] we can call  $\mathbf{a}_b$  the “vector Schwarzschild discriminens”, and  $\mathcal{A}_b$  the “generalized Schwarzschild discriminant”. But, in spite of the definition [6, (3.7)], we would like to define the square of the “generalized Brunt-Väisälä frequency (local bouyancy frequency)”  $\mathcal{N}^2$  by

$$\mathcal{N}^2 := -\mathcal{A}_b \left( \frac{1}{\rho_b} \nabla P_b \Big| \mathbf{n} \right) = -\mathcal{A}_b \frac{(\nabla P_b | \nabla \rho_b)}{\rho_b \|\nabla \rho_b\|}. \quad (2.15)$$

Then, at  $\|\mathbf{x}\| \neq 0$ ,  $\mathcal{N}_b(\mathbf{x}) \in \mathbb{R}$  if and only if  $\mathcal{A}_b(\mathbf{x}) \leq 0$ , since  $(\nabla P_b | \nabla \rho_b) > 0$ . It seems curious that the Lagrangian displacemnt of perturbation  $\boldsymbol{\xi}$  is involved in the definition [6, (3.7)]. Anyway, if  $\Omega = \omega_b = 0$  and the background is spherically symmetric, then it turns out to be  $\mathbf{n} = -\mathbf{e}_r = -\frac{\mathbf{x}}{\|\mathbf{x}\|}$  and the definitions (2.14), (2.15) of  $\mathcal{A}$ ,  $\mathcal{N}^2$  coincide with those [16, (1.11), (1.12)]. In **Appendix B** we derive the definition (2.15) of the Brunt-Väisälä frequency  $\mathcal{N}$  here by generalizing that found in meteorological texts, e.g. [5, Chapter II, Section 21], [8, Chapter 2, Section 2.7.3].

We are going to construct a barotropic admissible stationary solution.  
Suppose

$$\mathbf{v}_b(\mathbf{x}) = \omega_b(\varpi) \frac{\partial}{\partial x^3} \times \mathbf{x} = \omega_b(\varpi) \begin{bmatrix} -x^2 \\ x^1 \\ 0 \end{bmatrix}, \quad (2.16)$$

where  $\omega_b \in C^1([0, +\infty[)$ .

The equation (1.1a) is reduced to

$$(\mathbf{v}_b | \nabla \rho_b) + \rho_b \operatorname{div} \mathbf{v}_b = 0. \quad (2.17)$$

But

$$(\mathbf{v}_b | \nabla \rho_b) = \omega(\varpi) \left( -x^2 \frac{\partial \rho_b}{\partial x^1} + x^1 \frac{\partial \rho_b}{\partial x^2} \right) = \omega(\varpi) \frac{\partial \rho_b}{\partial \phi} = 0,$$

if  $\rho_b$  is axially symmetric. Moreover we have  $\operatorname{div} \mathbf{v}_b = 0$  for (2.16). Thus (2.17) holds for axially symmetric  $\rho_b$ .

The equation (1.1c) is reduced to

$$(\mathbf{v}_b | \nabla S_b) = 0. \quad (2.18)$$

This holds provided (2.16) and axially symmetricity of  $\rho_b$ .

Thus the equations are reduced to

$$(\mathbf{v}_b | \nabla) \mathbf{v}_b + 2\boldsymbol{\Omega} \times \mathbf{v}_b + \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{x}) + \frac{1}{\rho_b} \nabla P_b + \nabla \Phi_b = 0 \quad (2.19)$$

with

$$\Phi_b = -4\pi \mathbf{G} \mathcal{K}[\rho_b], \quad P_b = \mathcal{P}(\rho_b, S_b).$$

We claim

**Theorem 1** *Let a smooth function  $\omega_b$  on  $[0, +\infty[$ , a smooth function  $\Sigma$  on  $\mathbb{R}$  such that*

$$\gamma + (\gamma - 1) \left[ v \frac{\partial}{\partial v} + v \frac{d\Sigma}{dv} \frac{\partial}{\partial S} \right] \mathcal{F} > 0 \quad \text{for } v > 0 \quad (2.20)$$

*and a positive number  $\rho_O$  be given. If  $\frac{6}{5} < \gamma < 2$ , and  $\frac{1}{\rho_O} \|\boldsymbol{\Omega} + \omega_b\|_{L^\infty}^2$  is sufficiently small, then there exists a barotropic admissible stationary solution  $(\rho_b, S_b, \mathbf{v}_b)$  such that  $S_b = \Sigma(\rho_b^{\gamma-1})$ ,  $\rho_b(O) = \rho_O$ , and  $\mathbf{v}_b = \omega_b(\varpi) \frac{\partial}{\partial x^3} \times \mathbf{x}$ .*

Proof . Consider the functions  $f^P, f^{\mathcal{F}}$  defined by

$$f^P(\rho) := \rho^\gamma \mathcal{F}(\rho^{\gamma-1}, \Sigma(\rho^{\gamma-1})), \quad (2.21)$$

$$f^{\mathcal{F}}(\rho) := \int_0^\rho \frac{Df^P(\rho')}{\rho'} d\rho' \quad (2.22)$$



for  $\rho > 0$ . Thanks to the assumption (2.20) we have  $Df^P(\rho) > 0$  for  $\rho > 0$ , and there exists a smooth function  $\Lambda$  on  $\mathbb{R}$  such that  $\Lambda(0) = 0$  and

$$f^P(\rho) = A\rho^\gamma(1 + \Lambda(\rho^{\gamma-1})) \quad (2.23)$$

for  $\rho > 0$ . Here  $A := \mathcal{F}(0, \Sigma(0))$  is a positive constant. Then we have

$$\mathcal{R} = f^{\mathcal{R}}(\rho) = \frac{\gamma A}{\gamma - 1} \rho^{\gamma-1} (1 + \Lambda_1(\rho^{\gamma-1})) \quad (2.24)$$

for  $\rho > 0$ , where  $\Lambda_1$  is a smooth function on  $\mathbb{R}$  such that  $\Lambda_1(0) = 0$ , and the inverse function  $f^\rho$  of  $f^{\mathcal{R}}$

$$f^\rho(\mathcal{R}) = \left( \frac{\gamma - 1}{\gamma A} \right)^{\frac{1}{\gamma-1}} (\mathcal{R} \vee 0)^{\frac{1}{\gamma-1}} (1 + \Lambda_2(\mathcal{R})) \quad (2.25)$$

is given so that  $\rho = f^\rho(\mathcal{R}) \Leftrightarrow \mathcal{R} = f^{\mathcal{R}}(\rho)$  for  $\mathcal{R} > 0 (\rho > 0)$ . Here  $\mathcal{R} \vee 0$  stands for  $\max(\mathcal{R}, 0)$  and  $\Lambda_2$  is a smooth functions on  $\mathbb{R}$  such that  $\Lambda_2(0) = 0$ .

As shown in [10] with the inertial coordinate system, the problem is reduced to :

$$\mathcal{R}_b(\mathbf{x}) + \Phi_b(\mathbf{x}) - \mathfrak{B}(\varpi) = \mathcal{R}_O + \Phi_b(O) \quad \text{on} \quad \mathfrak{D}_0 \quad (2.26)$$

with

$$\begin{aligned} \Phi_b(\mathbf{x}) &= -4\pi\mathcal{G}\mathcal{K}[\rho_b], \\ \rho_b(\mathbf{x}) &= f^\rho(\mathcal{R}_b(\mathbf{x})) = \left( \frac{\gamma - 1}{\gamma A} \right)^{\frac{1}{\gamma-1}} (\mathcal{R}_b(\mathbf{x}) \vee 0)^{\frac{1}{\gamma-1}} (1 + \Lambda_2(\mathcal{R}_b(\mathbf{x}))). \end{aligned} \quad (2.27)$$

and

$$\mathfrak{B}(\varpi) := \int_0^\varpi (\Omega + \omega_b(\varpi'))^2 \varpi' d\varpi'. \quad (2.28)$$

Here  $\mathfrak{D}_0$  is a domain such that  $\mathcal{R}_b \in C^{2,\alpha}(\mathfrak{D}_0 \cup \partial\mathfrak{D}_0)$  and  $\{\mathcal{R}_b \geq 0\} \subset \mathfrak{D}_0$ .

Put

$$\mathcal{R}_O := f^{\mathcal{R}}(\rho_O) = \frac{\gamma A}{\gamma - 1} \rho_O^{\gamma-1} (1 + \Lambda_1(\rho_O^{\gamma-1})) \quad (2.29)$$

by the given  $\rho_O$ , and put

$$\mathfrak{a} = \frac{1}{\sqrt{4\pi\mathcal{G}}} \left( \frac{A\gamma}{\gamma - 1} \right)^{\frac{1}{2(\gamma-1)}} \mathcal{R}_O^{-\frac{2-\gamma}{2(\gamma-1)}}, \quad (2.30)$$

$$\mathfrak{b}(\varpi) = \mathcal{R}_O^{-1} \mathfrak{B}(\mathfrak{a}\varpi). \quad (2.31)$$

We put

$$\|\mathfrak{b}\|_1 = \sup_{\varpi \leq r_\infty} |\mathfrak{b}| + \sup_{\varpi \leq r_\infty} \left| \frac{d\mathfrak{b}}{d\varpi} \right|,$$

where  $\mathfrak{D}_0 \subset \{\|\mathbf{x}\| \leq r_\infty\}$ , and note

$$\|\mathfrak{b}\|_1 \leq \frac{1}{4\pi\mathbf{G}} \left( \frac{\mathbf{A}\gamma}{\gamma-1} \right)^{\frac{1}{\gamma-1}} \mathcal{R}_O^{-\frac{1}{\gamma-1}} \|\Omega + \omega_b\|_{L^\infty(0, ar_\infty)} \left( \frac{r_\infty^2}{2} + r_\infty \right).$$

We suppose

**Assumption 2**  $\frac{6}{5} < \gamma < 2$ , and  $\mathfrak{b}$  is sufficiently small, say,  $\|\mathfrak{b}\|_1 \leq \beta^0$ ,  $\beta^0$  being a positive number depending on  $\gamma, \Lambda$ .

Then we put

$$\rho_b(\mathbf{x}) = f^\rho(\mathcal{R}_O\Theta) = \left( \frac{\gamma-1}{\mathbf{A}\gamma} \right)^{\frac{1}{\gamma-1}} (\mathcal{R}_O\Theta \vee 0)^{\frac{1}{\gamma-1}} (1 + \Lambda_2(\mathcal{R}_O\Theta))$$

with

$$\Theta = \Theta\left(\frac{r}{a}, \zeta; \frac{1}{\gamma-1}, \mathfrak{b}\right),$$

and

$$S_b(\mathbf{x}) = \Sigma(\rho_b(\mathbf{x})^{\gamma-1}). \quad (2.32)$$

Here  $\Theta(w, \zeta; \frac{1}{\gamma-1}, \mathfrak{b})$  is the "distorted Lane-Emden function", which is the solution of the integral equation

$$\Theta(r, \zeta) = 1 + \mathfrak{b}(r\sqrt{1-\zeta^2}) + \mathcal{K}g(r, \zeta) - \mathcal{K}g(0, 0)$$

with

$$\begin{aligned} \mathcal{K}g(r, \zeta) &= \frac{1}{4\pi} \int_{-1}^1 \int_0^{+\infty} K(r, \zeta, r', \zeta') g(r', \zeta') r'^2 dr' d\zeta', \\ K(r, \zeta, r', \zeta') &= \int_0^{2\pi} \frac{d\phi}{\sqrt{r^2 + r'^2 - 2rr'(\sqrt{1-\zeta^2}\sqrt{1-\zeta'^2}\cos\phi + \zeta\zeta')}}, \\ g(r, \zeta) &= (\Theta(r, \zeta) \vee 0)^{\frac{1}{\gamma-1}} (1 + \Lambda_2(\mathcal{R}_O\Theta(r, \zeta))), \end{aligned}$$

and enjoys the following properties:

1) The function  $\boldsymbol{\xi} \mapsto \Theta(\|\boldsymbol{\xi}\|, \frac{\xi^3}{\|\boldsymbol{\xi}\|}; \frac{1}{\gamma-1}, \mathfrak{b})$  belongs to  $C^{2,\alpha}(\{\|\boldsymbol{\xi}\| \leq \Xi_0\})$  with  $0 < \alpha < (\frac{1}{\gamma-1} - 1) \wedge 1$ ,  $\Xi_0 = 4\xi_1(\frac{1}{\gamma-1})$ ,  $\xi_1(\frac{1}{\gamma-1})$  being the zero of the Lane-Emden function  $\theta(\xi; \frac{1}{\gamma-1})$  of the index  $\frac{1}{\gamma-1}$ , that is, the solution of

$$-\frac{1}{\xi^2} \frac{d}{d\xi} \xi^2 \frac{d\theta}{d\xi} = (\theta \vee 0)^{\frac{1}{\gamma-1}}, \quad \theta|_{\xi=0} = 1.$$

2)  $\Theta(0, \zeta; \frac{1}{\gamma-1}, \mathfrak{b}) = 1$  and there is a curve  $\zeta \in [-1, 1] \mapsto \Xi_1(\zeta; \frac{1}{\gamma-1}, \mathfrak{b})$  such that

$$\xi_1\left(\frac{1}{\gamma-1}\right) \leq \Xi_1\left(\zeta; \frac{1}{\gamma-1}, \mathfrak{b}\right) < \frac{1}{2}\Xi_0 = 2\xi_1\left(\frac{1}{\gamma-1}\right)$$

and, for  $0 \leq \xi < \Xi_0$ ,

$$0 < \Theta(\xi, \zeta; \frac{1}{\gamma-1}, \mathbf{b}) \Leftrightarrow 0 \leq \xi < \Xi_1(\zeta; \frac{1}{\gamma-1}, \mathbf{b}). \quad (2.33)$$

$$3) \frac{\partial \Theta}{\partial \xi} \leq -\frac{\xi}{C} \text{ with a positive number } C = C(\frac{1}{\gamma-1}, \mathbf{b}).$$

For proof of the existence of the distorted Lane-Emden function  $\Theta$ , see [10].

For the sake of simplicity we consider  $\Theta(\xi, \zeta) < 0$  for  $\xi \geq \Xi_0$  by modifying the values of  $\Theta$  on  $\xi > \Xi_0$ , so that (2.33) is valid for  $0 \leq \xi < \infty, |\zeta| \leq 1$ . Let us denote

$$\mathfrak{D}_0 = \{\mathbf{x} \in \mathbb{R}^3 \mid \|\mathbf{x}\| < R_{b0}\}, \quad R_{b0} = \mathbf{a}\Xi_0 = 4\mathbf{a}\xi_1(\frac{1}{\gamma-1}). \quad (2.34)$$

Then

$$\gamma_b(\mathbf{x}) = \gamma_O \Theta\left(\frac{\|\mathbf{x}\|}{\mathbf{a}}, \frac{x^3}{\|\mathbf{x}\|}, \frac{1}{\gamma-1}, \mathbf{b}\right), \quad (2.35)$$

which belongs to  $C^{2,\alpha}(\mathfrak{D}_0 \cup \partial\mathfrak{D}_0)$ , satisfies

$$\gamma_b(\mathbf{x}) + \Phi_b(\mathbf{x}) - \mathfrak{B}(\varpi) = \gamma_O + \Phi_b(O) \quad \text{on } \mathfrak{D}_0. \quad (2.36)$$

We denote

$$\mathfrak{R}_b = \{\rho_b > 0\} = \{\Theta > 0\} = \{r < R_{1b}(\zeta)\}, \quad (2.37a)$$

$$\partial\mathfrak{R}_b = \{\Theta = 0\} = \{r = R_{1b}(\zeta)\}, \quad (2.37b)$$

where we put  $R_{1b}(\zeta) = \mathbf{a}\Xi_1(\zeta)$ .

Since

$$P_b = \Pi(\rho_b^{\gamma-1}), \quad (2.38)$$

where

$$\Pi(v) = (v \vee 0)^{\frac{\gamma}{\gamma-1}} \mathcal{F}(v, \Sigma(v)), \quad (2.39)$$

the stationary solution is barotropic. Here  $\Pi \in C^{1,\alpha}$  for  $\frac{\gamma}{\gamma-1} > 1 + \alpha$ .

As for  $\mathbf{a}_b = -\left(\frac{1}{\Gamma P} \frac{\partial P}{\partial S}\right)_b \nabla S_b$ , since  $S_b = \Sigma(\rho_b^{\gamma-1})$  with  $\Sigma \in C^\infty$ , we see

$$\mathbf{a}_b = -\left(\frac{1}{\Gamma P} \frac{\partial P}{\partial S}\right)_b \nabla S_b = -\left(\frac{1}{\Gamma P} \frac{\partial P}{\partial S}\right)_b \frac{d\Sigma}{dv} \nabla v \Big|_{v=\rho_b^{\gamma-1}}, \quad (2.40)$$

and

$$\mathcal{A}_b = \left(\frac{1}{\Gamma P} \frac{\partial P}{\partial S}\right)_b v \frac{d\Sigma}{dv} \|\nabla v\| \Big|_{v=\rho_b^{\gamma-1}}. \quad (2.41)$$

Since  $\rho_b^{\gamma-1} \in C^{2,\alpha}(\mathfrak{R}_b \cup \partial\mathfrak{R}_b)$ , we have  $\mathcal{A}_b \in C^{1,\alpha}(\mathfrak{R}_b \cup \partial\mathfrak{R}_b)$ . Recall (2.13).

Summing up, we get an admissible barotropic stationary solution  $(\rho_b, S_b, \mathbf{v}_b)$ .

□

**Remark 4** Let us note that, by (2.41), we see

$$\left(\frac{d\Upsilon}{d\rho}\right)_b = Df^{\Upsilon}(\rho_b) = \left(\frac{\Gamma P}{\rho^2}\right)_b \left(1 + \mathcal{A}_b \frac{\rho_b}{\|\rho_b\|}\right).$$

Therefore, if  $\frac{d\Sigma}{dv} = 0 \quad \forall v$ , that is,  $\Sigma$  is constant, then  $\mathcal{A}_b = 0$  and it holds

$$\left(\frac{d\Upsilon}{d\rho}\right)_b = \left(\frac{\Gamma P}{\rho^2}\right)_b. \quad (2.42)$$

We shall say that the background is ‘isentropic’ in this case. Then, if  $\mathcal{F}(v, S) = \exp(S/C_V)$ , we have  $\Gamma = \gamma$  and

$$P = A\rho^\gamma, \quad \Upsilon = \frac{\gamma A}{\gamma - 1} \rho^{\gamma-1}; \quad S = C_V \log A.$$

### 3 Derivatin of the equation of LASO (1.11)

Let us fix a compactly supported axially symmetric stationary solution  $(\rho_b, S_b, \mathbf{v}_b)$  with  $\mathbf{v}_b(\mathbf{x}) = \omega_b(\varpi, z) \frac{\partial}{\partial x^3} \times \mathbf{x}$ . In order to investigate solutions  $(\rho, S, \mathbf{v})$  near the fixed stationary solution, for which the domain  $\mathfrak{R}(t) = \{\mathbf{x} \mid \rho(t, \mathbf{x}) > 0\}$  should move with the free boundary  $\Sigma(t) = \partial\mathfrak{R}(t)$ , we introduce the Lagrangian coordinate to describe the equations.

Let us look at a solution  $(\rho, S, \mathbf{v})$  which belongs to  $C^1([0, T] \times \mathbb{R}^3)$  with  $\mathfrak{R}(t) = \{\mathbf{x} \mid \rho(t, \mathbf{x}) > 0\}$ , provided that it exists. Suppose that  $(\mathfrak{R}(t), \partial\mathfrak{R}(t))$  is diffeomorphic onto  $(\mathfrak{R}_b, \partial\mathfrak{R}_b)$ . Let us suppose

$$\mathfrak{R}(0) = \{\mathbf{x} \mid \rho^0(\mathbf{x}) > 0\} = \mathfrak{R}_b. \quad (3.1)$$

However we do not suppose that  $\rho^0 = \rho_b$ .

We can consider the flow mapping  $\varphi \in C^1([0, T] \times \mathbb{R}^3)$  defined by

$$\frac{\partial}{\partial t} \varphi(t, \bar{\mathbf{x}}) = \mathbf{v}(t, \varphi(t, \bar{\mathbf{x}})), \quad \varphi(0, \bar{\mathbf{x}}) = \bar{\mathbf{x}}, \quad (3.2)$$

and  $\varphi_b(t, \bar{\mathbf{x}}) \in C^1([0, +\infty[; \mathbb{R}^3)$  defined by

$$\frac{\partial}{\partial t} \varphi_b(t, \bar{\mathbf{x}}) = \mathbf{v}_b(\varphi_b(t, \bar{\mathbf{x}})), \quad \varphi_b(0, \bar{\mathbf{x}}) = \bar{\mathbf{x}}, \quad (3.3)$$

which can be written explicitly as

$$\varphi_b(t, \bar{\mathbf{x}}) = \begin{bmatrix} \cos(\omega_b t) & -\sin(\omega_b t) & 0 \\ \sin(\omega_b t) & \cos(\omega_b t) & 0 \\ 0 & 0 & 1 \end{bmatrix} \bar{\mathbf{x}}, \quad (3.4)$$

where  $\omega_b = \omega_b(\bar{\varpi}, \bar{z})$ ,  $\bar{\varpi} = \sqrt{(\bar{x}^1)^2 + (\bar{x}^2)^2}$ ,  $\bar{z} = \bar{x}^3$ .

Through the mapping  $\mathbf{x} = \boldsymbol{\varphi}(t, \bar{\mathbf{x}})$  the equations on  $(t, \mathbf{x})$  can be written as equations on  $(t, \bar{\mathbf{x}})$ . This is the Lagrangian description.

Hereafter we denote

$$(D_{\bar{\mathbf{x}}} \mathbf{x})(t, \bar{\mathbf{x}}) = D\boldsymbol{\varphi}(t, \bar{\mathbf{x}}) = \left( \frac{\partial}{\partial \bar{\mathbf{x}}^\alpha} \left( \boldsymbol{\varphi}(t, \bar{\mathbf{x}}) \right)^j \right)_{j, \alpha}. \quad (3.5)$$

Then it hold that  $D_{\bar{\mathbf{x}}} \mathbf{x}(0, \bar{\mathbf{x}}) = I$  and

$$D_{\bar{\mathbf{x}}} \mathbf{x}(t, \bar{\mathbf{x}}) = D\boldsymbol{\varphi}(t, \bar{\mathbf{x}}) = \exp \left[ \int_0^t (D_{\mathbf{x}} \mathbf{v})(t', \boldsymbol{\varphi}(t', \bar{\mathbf{x}})) dt' \right] \quad (3.6)$$

Here  $D_{\mathbf{x}} \mathbf{v}$  stands for the matrix field  $\left( \frac{\partial v^k}{\partial x^j} \right)_{k, j}$ .

Let us denote

$$\begin{aligned} \rho^L(t, \bar{\mathbf{x}}) &= \rho(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})), & P^L(t, \bar{\mathbf{x}}) &= P(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})), \\ S^L(t, \bar{\mathbf{x}}) &= S(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})), & \mathbf{v}^L(t, \bar{\mathbf{x}}) &= \mathbf{v}(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) \end{aligned} \quad (3.7)$$

for  $(t, \bar{\mathbf{x}}) \in [0, +\infty[ \times \mathfrak{R}_b$ .

The equations (1.1a), (1.1b), (1.1c) read

$$\frac{\partial \rho^L}{\partial t} + (\operatorname{div}_{\mathbf{x}} \mathbf{v})(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) = 0, \quad (3.8a)$$

$$\begin{aligned} \frac{\partial}{\partial t} (\mathbf{v}^L - \mathbf{v}_b^L) + B(\mathbf{v}^L - \mathbf{v}_b^L) + \left( \frac{1}{\rho} (\nabla_{\mathbf{x}} P) - \frac{1}{\rho_b} (\nabla_{\mathbf{x}} P_b) + \right. \\ \left. + (\nabla_{\mathbf{x}} (\Phi - \Phi_b)) \right)^L = 0, \end{aligned} \quad (3.8b)$$

$$\frac{\partial S^L}{\partial t} = 0. \quad (3.8c)$$

Here, for function  $Q(t, \mathbf{x})$ , we denote

$$Q^L(t, \bar{\mathbf{x}}) = Q(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) \quad (3.9)$$

in general.

Integrating (3.8c), we have

$$S^L(t, \bar{\mathbf{x}}) = S^0(\bar{\mathbf{x}}). \quad (3.10)$$

Integrating (3.8a), we have

$$\rho^L(t, \bar{\mathbf{x}}) = \rho^0(\bar{\mathbf{x}}) \exp \left[ - \int_0^t (\operatorname{div}_{\mathbf{x}} \mathbf{v})(s, \boldsymbol{\varphi}(s, \bar{\mathbf{x}})) ds \right] \quad (3.11)$$

for  $(t, \bar{\mathbf{x}}) \in [0, +\infty[ \times \mathfrak{R}_b$ .

Let us denote

$$\mathfrak{d}\rho(t, \bar{\mathbf{x}}) = \rho^L(t, \bar{\mathbf{x}}) - \rho_b^L(t, \bar{\mathbf{x}}) = \rho(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) - \rho_b(\boldsymbol{\varphi}(t, \bar{\mathbf{x}})), \quad (3.12a)$$

$$\mathfrak{d}P(t, \bar{\mathbf{x}}) = P^L(t, \bar{\mathbf{x}}) - P_b^L(t, \bar{\mathbf{x}}) = P(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) - P_b(\boldsymbol{\varphi}(t, \bar{\mathbf{x}})), \quad (3.12b)$$

$$\begin{aligned} \mathfrak{d}S(t, \bar{\mathbf{x}}) &= S^L(t, \bar{\mathbf{x}}) - S_b^L(t, \bar{\mathbf{x}}) = S(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) - S_b(\boldsymbol{\varphi}(t, \bar{\mathbf{x}})) = \\ &= S^0(\bar{\mathbf{x}}) - S_b(\boldsymbol{\varphi}(t, \bar{\mathbf{x}})), \end{aligned} \quad (3.12c)$$

$$\mathfrak{d}\Phi(t, \bar{\mathbf{x}}) = \Phi^L(t, \bar{\mathbf{x}}) - \Phi_b^L(t, \bar{\mathbf{x}}) = \Phi(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) - \Phi_b(\boldsymbol{\varphi}(t, \bar{\mathbf{x}})), \quad (3.12d)$$

$$\mathfrak{d}\mathbf{v}(t, \bar{\mathbf{x}}) = \mathbf{v}^L(t, \bar{\mathbf{x}}) - \mathbf{v}_b^L(t, \bar{\mathbf{x}}), \quad (3.12e)$$

and

$$\delta\rho(t, \mathbf{x}) = \rho(t, \mathbf{x}) - \rho_b(t, \mathbf{x}), \quad (3.13a)$$

$$\delta P(t, \mathbf{x}) = P(t, \mathbf{x}) - P_b(t, \mathbf{x}), \quad (3.13b)$$

$$\delta S(t, \mathbf{x}) = S(t, \mathbf{x}) - S_b(t, \mathbf{x}), \quad (3.13c)$$

$$\delta\Phi(t, \mathbf{x}) = \Phi(t, \mathbf{x}) - \Phi_b(t, \mathbf{x}), \quad (3.13d)$$

$$\delta\mathbf{v}(t, \mathbf{x}) = \mathbf{v}(t, \mathbf{x}) - \mathbf{v}_b(t, \mathbf{x}). \quad (3.13e)$$

Then the equations (3.8b) and (2.19) read

$$\begin{aligned} &\frac{\partial}{\partial t} \mathfrak{d}\mathbf{v} + 2\boldsymbol{\Omega} \times \mathfrak{d}\mathbf{v} + \\ &+ \left( \frac{1}{\rho_b} \nabla_{\mathbf{x}} P - \frac{1}{\rho_b} \nabla_{\mathbf{x}} P_b + \nabla_{\mathbf{x}} (\Phi - \Phi_b) \right)^L = - \left( (D_{\mathbf{x}} \mathbf{v}_b) \delta\mathbf{v} \right)^L. \end{aligned} \quad (3.14)$$

• Now let us derive the linearized approximation of (3.11), (3.14). Considering small  $\varepsilon$ , a quantity  $Q$  will be denoted by  $\mathcal{O}(\varepsilon)$  if  $Q$  and its derivatives are of order  $\mathcal{O}(\varepsilon)$  uniformly on  $t \in [0, T]$ ,  $\mathbf{x} \in \mathfrak{R}(t)$ ,  $\text{dist}(\mathbf{x}, \partial\mathfrak{R}(t)) \geq \delta$  for  $\forall$  fixed  $T, \delta$ ,  $0 < \delta \ll 1$ .

We assume that  $\rho - \rho_b, S - S_b, \mathbf{v}_b, \mathbf{v} - \mathbf{v}_b$  are of  $\mathcal{O}(\varepsilon)$ .

First of all we note that (3.6) shows

$$D_{\bar{\mathbf{x}}} \mathbf{x}(t, \bar{\mathbf{x}}) = I + \int_0^t (D_{\mathbf{x}} \mathbf{v})(s, \boldsymbol{\varphi}(s, \bar{\mathbf{x}})) ds + \mathcal{O}(\varepsilon^2) \quad (3.15)$$

$$= I + \mathcal{O}(\varepsilon), \quad (3.16)$$

since we are assuming  $\mathbf{v} = \mathbf{v}_b + (\mathbf{v} - \mathbf{v}_b) = \mathcal{O}(\varepsilon)$ . Of course

$$\begin{aligned} \boldsymbol{\varphi}(t, \bar{\mathbf{x}}) &= \bar{\mathbf{x}} + \int_0^t \mathbf{v}(s, \boldsymbol{\varphi}(s, \bar{\mathbf{x}})) ds \\ &= \bar{\mathbf{x}} + \mathcal{O}(\varepsilon). \end{aligned} \quad (3.17)$$

Therefore, if  $\mathbf{Q} = \mathbf{Q}(t, \mathbf{x}) = \mathcal{O}(\varepsilon)$ , then

$$\operatorname{div}_{\mathbf{x}} \mathbf{Q}(t, \mathbf{x}) = \operatorname{div}_{\bar{\mathbf{x}}} \mathbf{Q}^L(t, \bar{\mathbf{x}}) + \mathcal{O}(\varepsilon^2), \quad (3.18)$$

and, if  $Q = Q(t, \mathbf{x}) = \mathcal{O}(\varepsilon)$ , then

$$\nabla_{\mathbf{x}} Q(t, \mathbf{x}) = \nabla_{\bar{\mathbf{x}}} Q^L(t, \bar{\mathbf{x}}) + \mathcal{O}(\varepsilon^2), \quad (3.19)$$

while  $\mathbf{x} = \boldsymbol{\varphi}(t, \bar{\mathbf{x}})$ . In other words

$$\left( \operatorname{div}_{\mathbf{x}} \mathbf{Q} \right)^L = \operatorname{div}_{\bar{\mathbf{x}}} \mathbf{Q}^L + \mathcal{O}(\varepsilon^2), \quad (3.20)$$

$$\left( \nabla_{\mathbf{x}} Q \right)^L = \nabla_{\bar{\mathbf{x}}} Q^L + \mathcal{O}(\varepsilon^2). \quad (3.21)$$

**Remark 5** *If we assume only  $\mathbf{v} - \mathbf{v}_b = \mathcal{O}(\varepsilon)$ , but do not assume  $\mathbf{v}_b = \mathcal{O}(\varepsilon)$ , then (3.16) can be not the case, but we have*

$$\begin{aligned} D_{\bar{\mathbf{x}}} \boldsymbol{\varphi}(t, \bar{\mathbf{x}}) &= D_{\bar{\mathbf{x}}} \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}}) + \mathcal{O}(\varepsilon), \\ D_{\bar{\mathbf{x}}} \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}}) &= \begin{bmatrix} \cos(\omega_b t) & -\sin(\omega_b t) & 0 \\ \sin(\omega_b t) & \cos(\omega_b t) & 0 \\ 0 & 0 & 1 \end{bmatrix} + \frac{1}{\varpi} \frac{\partial \omega_b}{\partial \varpi} t \begin{bmatrix} -x_b^2 \bar{x}^1 & -x_b^1 \bar{x}^2 & 0 \\ x_b^1 \bar{x}^1 & x_b^1 \bar{x}^2 & 0 \\ 0 & 0 & 0 \end{bmatrix} \\ &\quad + \frac{\partial \omega_b}{\partial z} t \begin{bmatrix} 0 & 0 & -x_b^2 \\ 0 & 0 & x_b^1 \\ 0 & 0 & 0 \end{bmatrix}, \end{aligned}$$

where  $\mathbf{x}_b = \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})$ ,  $\omega_b = \omega_b(\bar{\varpi}, \bar{z}) = \omega_b(\varpi, z)$ ,  $\varpi = \sqrt{(x_b^1)^2 + (x_b^2)^2}$ ,  $\bar{\varpi} = \sqrt{(\bar{x}^1)^2 + (\bar{x}^2)^2}$ ,  $z = x_b^3$ ,  $\bar{z} = \bar{x}^3$ . Thus, when  $\omega_b \neq 0$ , we have  $D_{\bar{\mathbf{x}}} \mathbf{x} \neq I + \mathcal{O}(\varepsilon)$ , and the formulae (3.20), (3.21) do not work.

Let us look at (3.11), which can be written as

$$\rho(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) = \rho^0(\bar{\mathbf{x}}) \exp \left[ - \int_0^t (\operatorname{div}_{\mathbf{x}} \delta \mathbf{v})(s, \boldsymbol{\varphi}(s, \bar{\mathbf{x}})) ds \right]. \quad (3.22)$$

Let us verify it. Look at

$$\begin{aligned} \exp \left[ - \int_0^t (\operatorname{div}_{\mathbf{x}} \mathbf{v})(s, \boldsymbol{\varphi}(s, \bar{\mathbf{x}})) ds \right] &= \\ &= \exp \left[ - \int_0^t (\operatorname{div}_{\mathbf{x}} \mathbf{v}_b)(s, \boldsymbol{\varphi}(s, \bar{\mathbf{x}})) ds \right] \cdot \exp \left[ - \int_0^t (\operatorname{div}_{\mathbf{x}} \delta \mathbf{v})(s, \boldsymbol{\varphi}(s, \bar{\mathbf{x}})) ds \right]. \end{aligned}$$

Since

$$\mathbf{v}_b(\mathbf{x}) = \omega_b(\varpi, z) \begin{bmatrix} -x^2 \\ x^1 \\ 0 \end{bmatrix},$$

we see  $\operatorname{div}_{\bar{\mathbf{x}}} \mathbf{v}_b = 0$ . Therefore

$$\exp \left[ - \int_0^t (\operatorname{div}_{\bar{\mathbf{x}}} \mathbf{v}_b)(s, \boldsymbol{\varphi}(s, \bar{\mathbf{x}})) ds \right] = I$$

and (3.22) follows.

Recalling

$$\frac{\partial}{\partial t} \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}}) = \mathbf{v}_b(t, \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})), \quad \boldsymbol{\varphi}_b(0, \bar{\mathbf{x}}) = \bar{\mathbf{x}},$$

we have

$$\boldsymbol{\varphi}(t, \bar{\mathbf{x}}) - \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}}) = \int_0^t (\delta \mathbf{v})(s, \boldsymbol{\varphi}(s, \bar{\mathbf{x}})) ds + \mathcal{O}(\varepsilon^2), \quad (3.23)$$

since

$$\begin{aligned} \mathbf{v}(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) - \mathbf{v}_b(t, \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})) &= \\ &= (\delta \mathbf{v})(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) + \left( \mathbf{v}_b(\boldsymbol{\varphi}(t, \bar{\mathbf{x}})) - \mathbf{v}_b(\boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})) \right) \\ &= (\delta \mathbf{v})(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) + \mathcal{O}(\varepsilon^2). \end{aligned}$$

Then we have

$$\begin{aligned} \operatorname{div}_{\bar{\mathbf{x}}}(\boldsymbol{\varphi}(t, \bar{\mathbf{x}}) - \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})) &= \int_0^t \operatorname{div}_{\bar{\mathbf{x}}}(\delta \mathbf{v}(s, \boldsymbol{\varphi}(s, \bar{\mathbf{x}}))) ds + \mathcal{O}(\varepsilon^2) \\ &= \int_0^t (\operatorname{div}_{\bar{\mathbf{x}}}(\delta \mathbf{v}))(s, \boldsymbol{\varphi}(s, \bar{\mathbf{x}})) ds + \mathcal{O}(\varepsilon^2). \end{aligned}$$

Thus (3.22) reads

$$\rho(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) = \rho^0(\bar{\mathbf{x}}) \exp \left[ - \operatorname{div}_{\bar{\mathbf{x}}}(\boldsymbol{\varphi}(t, \bar{\mathbf{x}}) - \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})) \right] + \mathcal{O}(\varepsilon^2),$$

or we can claim

$$\rho(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) = \rho^0(\bar{\mathbf{x}}) - \rho^0(\bar{\mathbf{x}}) \operatorname{div}_{\bar{\mathbf{x}}}(\boldsymbol{\varphi}(t, \bar{\mathbf{x}}) - \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})) + \mathcal{O}(\varepsilon^2). \quad (3.24)$$

Further we have, in modulo  $\mathcal{O}(\varepsilon^2)$ ,

$$\begin{aligned} \mathfrak{d}\rho(t, \bar{\mathbf{x}}) &= \rho(t, \boldsymbol{\varphi}(t, \bar{\mathbf{x}})) - \rho_b(\boldsymbol{\varphi}(t, \bar{\mathbf{x}})) \\ &\equiv \rho^0(\bar{\mathbf{x}}) - \rho_b(\boldsymbol{\varphi}(t, \bar{\mathbf{x}})) - \rho^0(\bar{\mathbf{x}}) \operatorname{div}_{\bar{\mathbf{x}}}(\boldsymbol{\varphi}(t, \bar{\mathbf{x}}) - \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})) \\ &\equiv \rho^0(\bar{\mathbf{x}}) - \rho_b(\boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})) - \left( \nabla \rho_b(\boldsymbol{\varphi}_b(t, \bar{\mathbf{x}}) - \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})) \Big| \boldsymbol{\varphi}(t, \bar{\mathbf{x}}) \right) \\ &\quad - \operatorname{div}_{\bar{\mathbf{x}}} \rho^0(\bar{\mathbf{x}})(\boldsymbol{\varphi}(t, \bar{\mathbf{x}}) - \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})) + \left( \nabla \rho^0(\bar{\mathbf{x}}) \Big| \boldsymbol{\varphi}(t, \bar{\mathbf{x}}) - \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}}) \right) \\ &= \rho^0(\bar{\mathbf{x}}) - \rho_b(\boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})) - \operatorname{div}_{\bar{\mathbf{x}}} \left( \rho^0(\bar{\mathbf{x}})(\boldsymbol{\varphi}(t, \bar{\mathbf{x}}) - \boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})) \right). \end{aligned}$$

Since it holds that

$$\rho_b(\boldsymbol{\varphi}_b(t, \bar{\mathbf{x}})) = \rho_b(\bar{\mathbf{x}}), \quad (3.25)$$



we can claim

$$\mathfrak{d}\rho(t, \bar{x}) = \rho^0(\bar{x}) - \rho_b(\bar{x}) - \operatorname{div}_{\bar{x}} \left( \rho_b(\bar{x})(\varphi(t, \bar{x}) - \varphi_b(t, \bar{x})) \right) + \mathcal{O}(\varepsilon^2), \quad (3.26)$$

since we are assuming that  $\rho_b - \rho^0 = \mathcal{O}(\varepsilon)$ .

Now let us introduce  $\mathbf{u}$  by

$$\mathbf{u}(t, \bar{x}) = \varphi(t, \bar{x}) - \varphi_b(t, \bar{x}) + \mathbf{u}^0(\bar{x}), \quad (3.27)$$

where  $\mathbf{u}^0 = \mathcal{O}(\varepsilon)$ . Then we can claim

$$\mathfrak{d}\rho(t, \bar{x}) = -\operatorname{div}_{\bar{x}}(\rho_b(\bar{x})\mathbf{u}(t, \bar{x})) + \mathcal{O}(\varepsilon^2), \quad (3.28)$$

provided that

$$\rho^0(\bar{x}) - \rho_b(\bar{x}) = -\operatorname{div}_{\bar{x}}(\rho_b(\bar{x})\mathbf{u}^0(\bar{x})). \quad (3.29)$$

We note that

$$\frac{\partial \mathbf{u}}{\partial t}(t, \bar{x}) = \mathfrak{d}\mathbf{v}(t, \bar{x}) + \mathcal{O}(\varepsilon^2), \quad (3.30)$$

where we recall  $\mathfrak{d}\mathbf{v}(t, \bar{x}) = (\delta\mathbf{v})(t, \varphi(t, \bar{x}))$  and  $\mathbf{v}(t, \varphi(t, \bar{x})) - \mathbf{v}_b(\varphi_b(t, \bar{x})) = \mathfrak{d}\mathbf{v}(t, \bar{x}) + \mathcal{O}(\varepsilon^2)$ .

Let us look at  $\mathfrak{d}S$ . We can claim

$$\mathfrak{d}S(t, \bar{x}) = -\left( \mathbf{u}(t, \bar{x}) \Big| \nabla_{\bar{x}} S_b(\bar{x}) \right) + \mathcal{O}(\varepsilon^2), \quad (3.31)$$

provided that

$$S^0(\bar{x}) - S_b(\bar{x}) = -(\mathbf{u}^0(\bar{x}) \Big| \nabla_{\bar{x}} S_b(\bar{x})). \quad (3.32)$$

In fact, we see

$$\begin{aligned} \mathfrak{d}S(t, \bar{x}) &= S^0(\bar{x}) - S_b(\varphi(t, \bar{x})) \\ &= S^0(\bar{x}) - \left[ S_b(\varphi_b(t, \bar{x})) + \left( \varphi(t, \bar{x}) - \varphi_b(t, \bar{x}) \Big| \nabla_{\bar{x}} S_b(\varphi_b(t, \bar{x})) \right) + \mathcal{O}(\varepsilon^2) \right] \\ &= S^0(\bar{x}) - S_b(\bar{x}) - \left( \mathbf{u} - \mathbf{u}^0 \Big| \nabla_{\bar{x}} S_b(\bar{x}) \right) + \mathcal{O}(\varepsilon^2) \\ &= -(\mathbf{u} \Big| \nabla_{\bar{x}} S_b(\bar{x})) + \mathcal{O}(\varepsilon^2), \end{aligned}$$

provided (3.32). Here we have used the identity

$$S_b(\varphi_b(t, \bar{x})) = S_b(\bar{x}), \quad (3.33)$$

which holds thanks to the axial symmetricity of  $S_b$ .

Moreover we can replace (3.32) by

$$\rho^0(\bar{x})S^0(\bar{x}) - \rho_b(\bar{x})S_b(\bar{x}) = -\operatorname{div}_{\bar{x}} \left( \rho_b(\bar{x})S_b(\bar{x})\mathbf{u}^0(\bar{x}) \right) \quad (3.34)$$

in modulo  $\mathcal{O}(\varepsilon^2)$ , since

$$\begin{aligned}\rho^0 S^0 - \rho_b S_b + \operatorname{div}(\rho_b S_b \mathbf{u}^0) &= \rho^0 (S^0 - S_b + (\mathbf{u}^0 | \nabla S_b)) + \\ &+ S_b \operatorname{div}((\rho^0 - \rho_b) \mathbf{u}^0) + (\rho^0 - \rho_b)(S^0 - S_b),\end{aligned}$$

provided (3.29). Note that (3.29) and (3.34) imply

$$\int_{\mathfrak{R}_b} \rho^0(\mathbf{x}) d\mathbf{x} = \int_{\mathfrak{R}_b} \rho_b(\mathbf{x}) d\mathbf{x}, \quad (3.35)$$

$$\int_{\mathfrak{R}_b} \rho^0(\mathbf{x}) S^0(\mathbf{x}) d\mathbf{x} = \int_{\mathfrak{R}_b} \rho_b(\mathbf{x}) S_b(\mathbf{x}) d\mathbf{x}, \quad (3.36)$$

which mean that the total mass and the total entropy are the same for the perturbed configuration and the unperturbed background.

Let us look at  $\mathfrak{d}P$ . We have

$$\begin{aligned}\mathfrak{d}P(t, \bar{\mathbf{x}}) &= \left( \frac{\partial P}{\partial \rho} \right) \Big|_{(\rho, S) = (\rho_b^L, S_b^L)(t, \bar{\mathbf{x}})} \mathfrak{d}\rho(t, \bar{\mathbf{x}}) + \\ &+ \left( \frac{\partial P}{\partial S} \right) \Big|_{(\rho, S) = (\rho_b^L, S_b^L)(t, \bar{\mathbf{x}})} \mathfrak{d}S(t, \bar{\mathbf{x}}) + \mathcal{O}(\varepsilon^2).\end{aligned}$$

Since

$$\begin{aligned}\left( \frac{\partial P}{\partial \rho} \right) \Big|_{(\rho, S) = (\rho_b^L, S_b^L)(t, \bar{\mathbf{x}})} &= \left( \frac{\partial P}{\partial \rho} \right) \Big|_{(\rho, S) = (\rho_b, S)(\bar{\mathbf{x}})} + \mathcal{O}(\varepsilon), \\ \left( \frac{\partial P}{\partial S} \right) \Big|_{(\rho, S) = (\rho_b, S_b)(\bar{\mathbf{x}})} &= \left( \frac{\partial P}{\partial S} \right) \Big|_{(\rho, S) = (\rho_b, S_b)(\bar{\mathbf{x}})} + \mathcal{O}(\varepsilon),\end{aligned}$$

we see

$$\begin{aligned}\mathfrak{d}P(t, \bar{\mathbf{x}}) &= \left( \frac{\partial P}{\partial \rho} \right) \Big|_{(\rho, S) = (\rho_b, S_b)(\bar{\mathbf{x}})} \mathfrak{d}\rho(t, \bar{\mathbf{x}}) + \\ &+ \left( \frac{\partial P}{\partial S} \right) \Big|_{(\rho, S) = (\rho_b, S_b)(\bar{\mathbf{x}})} \mathfrak{d}S(t, \bar{\mathbf{x}}) + \mathcal{O}(\varepsilon^2).\end{aligned}$$

Let us look at

$$\begin{aligned}\Phi(t, \mathbf{x}) - \Phi_b(\mathbf{x}) &= -G \int \frac{\rho(t, \mathbf{x}') - \rho_b(\mathbf{x}')}{\|\mathbf{x} - \mathbf{x}'\|} d\mathbf{x}' \\ &= -G \int_{\mathfrak{R}_b} \frac{\mathfrak{d}\rho(t, \bar{\mathbf{x}}')}{\|\bar{\mathbf{x}} - \bar{\mathbf{x}}'\|} d\bar{\mathbf{x}}' + \mathcal{O}(\varepsilon^2)\end{aligned} \quad (3.37)$$

In fact, applying the Gronwall argument to the identity

$$\varphi(t, \bar{\mathbf{x}}) - \varphi(t, \bar{\mathbf{x}}') = \bar{\mathbf{x}} - \bar{\mathbf{x}}' + \int_0^t \left[ \mathbf{v}(s, \varphi(s, \bar{\mathbf{x}})) - \mathbf{v}(s, \varphi(s, \bar{\mathbf{x}}')) \right] ds,$$

we can derive the estimate

$$\frac{1}{1 + C_1|\varepsilon|T} \|\varphi(t, \bar{x}) - \varphi(t, \bar{x}')\| \leq \|\bar{x} - \bar{x}'\| \leq \frac{1}{1 - C_1|\varepsilon|T} \|\varphi(t, \bar{x}) - \varphi(t, \bar{x}')\|$$

for  $t \in [0, T]$ ,  $\bar{x}, \bar{x}' \in \mathfrak{R}_b$ , provided that  $C_1|\varepsilon|T < 1$ , where

$$C_1 = \frac{1}{|\varepsilon|} \sup \left\{ \|D_{\mathbf{x}} \mathbf{v}(t, \mathbf{x})\| \mid 0 \leq t \leq T, \mathbf{x} \in \mathfrak{D}_0 \right\}$$

so that

$$\left| \int_0^t [\mathbf{v}(s, \varphi(s, \bar{x})) - \mathbf{v}(s, \varphi(s, \bar{x}'))] ds \right| \leq C_1|\varepsilon| \int_0^t \|\varphi(s, \bar{x}) - \varphi(s, \bar{x}')\| ds,$$

and we have

$$d\mathbf{x}' = \left| \det D_{\bar{x}} \mathbf{x}(t, \bar{x}') \right| d\bar{x}' = (1 + \mathcal{O}(\varepsilon)) d\bar{x}'.$$

Hence

$$\frac{\rho(t, \mathbf{x}') - \rho_b(\mathbf{x}')}{\|\mathbf{x} - \mathbf{x}'\|} d\mathbf{x}' = \frac{\mathfrak{d}\rho(t, \bar{x}')}{\|\bar{x} - \bar{x}'\|} d\bar{x}' + \mathcal{O}(\varepsilon^2)$$

For  $\mathbf{x} = \varphi(1, \bar{x})$ ,  $\mathbf{x}' = \varphi(t, \bar{x}')$ , therefore (3.37) holds.

Combining (3.28) and (3.37), we see

$$\Phi(t, \mathbf{x}) - \Phi_b(\mathbf{x}) = \mathbb{G} \int_{\mathfrak{R}_b} \frac{\operatorname{div}_{\bar{x}}(\rho_b \mathbf{u}(t, \bar{x}'))}{\|\bar{x} - \bar{x}'\|} d\bar{x}' + \mathcal{O}(\varepsilon^2). \quad (3.38)$$

We note

$$\begin{aligned} \frac{1}{\rho} \nabla_{\mathbf{x}} P - \frac{1}{\rho_b} \nabla_{\mathbf{x}} P_b &= \frac{1}{\rho_b} \nabla_{\mathbf{x}} (P - P_b) - \frac{\rho - \rho_b}{\rho_b^2} \nabla_{\mathbf{x}} P_b \\ &= \frac{1}{\rho_b^L} \nabla_{\bar{x}} (\mathfrak{d}P) - \frac{\mathfrak{d}\rho}{(\rho_b^L)^2} \nabla_{\bar{x}} P_b^L + \mathcal{O}(\varepsilon^2) \end{aligned} \quad (3.39)$$

with  $\mathbf{x} = \varphi(t, \bar{x})$ .

Summing up, we get the linearized approximation of (3.14):

$$\frac{\partial \mathfrak{d}\mathbf{v}}{\partial t} + B \mathfrak{d}\mathbf{v} + \mathcal{L} \mathbf{u} = 0, \quad (3.40)$$

with

$$\mathcal{L} \mathbf{u} = \frac{1}{\rho_b} \nabla_{\bar{x}} (\mathfrak{d}P) - \frac{\nabla_{\bar{x}} P_b}{\rho_b^2} \mathfrak{d}\rho + \nabla_{\bar{x}} (4\pi \mathbb{G} \mathcal{K} [-\mathfrak{d}\rho]), \quad (3.41)$$

$$\mathfrak{d}\rho = -\operatorname{div}_{\bar{x}}(\rho_b \mathbf{u}), \quad (3.42)$$

$$\mathfrak{d}S = -(\mathbf{u} | \nabla_{\bar{x}} S_b(\bar{x})), \quad (3.43)$$

$$\mathfrak{d}P(t, \bar{x}) = \left( \frac{\partial P}{\partial \rho} \right) \Big|_{(\rho, S) = (\rho_b, S_b)(\bar{x})} \mathfrak{d}\rho + \quad (3.44)$$

$$+ \left( \frac{\partial P}{\partial S} \right) \Big|_{(\rho, S) = (\rho_b, S_b)(\bar{x})} \mathfrak{d}S, \quad (3.45)$$

where we read  $\rho_b = \rho_b(\bar{\mathbf{x}})$ ,  $P_b = P_b(\bar{\mathbf{x}})$ ,  $S_b = S_b(\bar{\mathbf{x}})$ .

Since  $\mathfrak{d}\mathbf{v} = \partial\mathbf{u}/\partial t + \mathcal{O}(\varepsilon^2)$  ( (3.30) ), (3.40) reads

$$\frac{\partial^2 \mathbf{u}}{\partial t^2} + B \frac{\partial \mathbf{u}}{\partial t} + \mathcal{L}\mathbf{u} = 0. \quad (3.46)$$

This equation (3.46) is the linearized equation for perturbations described by the Lagrangian coordinate to be analyzed.

Let us recall that we are assuming (3.1), (3.29) and (3.32).

**Remark 6** *Strictly speaking, we should introduce  $[\mathfrak{d}\rho]$ ,  $[\mathfrak{d}S]$ ,  $[\mathfrak{d}P]$ ,  $[\mathfrak{d}\Phi]$  by defining*

$$[\mathfrak{d}\rho](t, \bar{\mathbf{x}}) = -\operatorname{div}_{\bar{\mathbf{x}}}(\rho_b(\bar{\mathbf{x}})\mathbf{u}(t, \bar{\mathbf{x}})), \quad (3.47a)$$

$$[\mathfrak{d}S](t, \bar{\mathbf{x}}) = -(\mathbf{u}(t, \bar{\mathbf{x}})|\nabla_{\bar{\mathbf{x}}}S_b(\bar{\mathbf{x}})), \quad (3.47b)$$

$$[\mathfrak{d}P](t, \bar{\mathbf{x}}) = \left(\frac{\partial P}{\partial \rho}\right)_b(\bar{\mathbf{x}})[\mathfrak{d}\rho](t, \bar{\mathbf{x}}) + \left(\frac{\partial P}{\partial S}\right)_b(\bar{\mathbf{x}})[\mathfrak{d}S](t, \bar{\mathbf{x}}), \quad (3.47c)$$

$$[\mathfrak{d}\Phi](t, \bar{\mathbf{x}}) = 4\pi\mathcal{G}\mathcal{K}[-[\mathfrak{d}\rho]](t, \bar{\mathbf{x}}). \quad (3.47d)$$

Then (3.41) should read

$$\mathbf{L}\mathbf{u} = \frac{1}{\rho_b}\nabla_{\bar{\mathbf{x}}}[\mathfrak{d}P] - \frac{\nabla_{\bar{\mathbf{x}}}P_b}{\rho_b^2}[\mathfrak{d}\rho] + \nabla_{\bar{\mathbf{x}}}[\mathfrak{d}\Phi], \quad (3.48)$$

while

$$\mathfrak{d}\rho = [\mathfrak{d}\rho] + \mathcal{O}(\varepsilon^2),$$

$$\mathfrak{d}S = [\mathfrak{d}S] + \mathcal{O}(\varepsilon^2),$$

$$\mathfrak{d}P = [\mathfrak{d}P] + \mathcal{O}(\varepsilon^2),$$

$$\mathfrak{d}\Phi = [\mathfrak{d}\Phi] + \mathcal{O}(\varepsilon^2).$$

**Remark 7** *The derivation of the linearized approximation of the equations in Lagrangian co-ordinate system can be found [14, Sect. 56], [1, pp. 139-140.], [2, p.11, (A)], [15], [13, p.500, (1)] and so on. But there was considered only the case of  $\rho^0 = \rho_b$ ,  $S^0 = S_b$ . Moreover, the definition of Lagrange displacement and the equation of LASO, the linearized equation of adiabatic perturbations, derived by them and traditionally used by astrophysicists are slightly different from those formulated here, when  $\mathbf{v}_b = \mathcal{O}(\varepsilon)$  is not supposed. For the details see Section 7.*

## 4 Basic existence theorem for the linearized equation of perturbations

We are going to discuss on the existence of solutions to the equation of LASO (1.11).

We formulate the initial boundary value problem to be considered as:

$$\frac{\partial^2 \mathbf{u}}{\partial t^2} + B \frac{\partial \mathbf{u}}{\partial t} + \mathcal{L} \mathbf{u} = \mathbf{f}(t, \mathbf{x}) \quad \text{on} \quad [0, +\infty[ \times \mathfrak{R}_b, \quad (4.1a)$$

$$\mathbf{u}|_{t=0} = \mathbf{u}^0, \quad \left. \frac{\partial \mathbf{u}}{\partial t} \right|_{t=0} = \mathbf{v}^0 \quad \text{on} \quad \mathfrak{R}_b, \quad (4.1b)$$

where  $\mathbf{u}^0, \mathbf{v}^0, \mathbf{f}$  are given data.

Let us rewrite

$$\mathcal{L}_0 \mathbf{u} := \frac{1}{\rho_b} \nabla \mathfrak{d}P - \frac{\nabla P_b}{\rho_b^2} \mathfrak{d}\rho,$$

using  $\Gamma_b, \mathbf{a}_b$  defined in Definition 3. Since

$$\mathfrak{d}P = \left( \frac{\Gamma P}{\rho} \right)_b \mathfrak{d}\rho + (\Gamma P)_b (\mathbf{u} | \mathbf{a}_b),$$

we have

$$\begin{aligned} \mathcal{L}_0 \mathbf{u} &= \nabla \left[ - \left( \frac{\Gamma P}{\rho^2} \right)_b \operatorname{div}(\rho_b \mathbf{u}) + \left( \frac{\Gamma P}{\rho} \right)_b (\mathbf{u} | \mathbf{a}_b) \right] + \\ &+ \left( \frac{\Gamma P}{\rho^2} \right)_b \left[ - \mathbf{a}_b \operatorname{div}(\rho_b \mathbf{u}) + (\mathbf{u} | \mathbf{a}_b) \nabla \rho_b \right]. \end{aligned}$$

Consequently we can write

$$\mathcal{L} \mathbf{u} = \mathcal{L}_0 \mathbf{u} + 4\pi \mathbb{G} \mathcal{L}_1 \quad (4.2)$$

with

$$\begin{aligned} \mathcal{L}_0 \mathbf{u} &= \nabla \left[ - \left( \frac{\Gamma P}{\rho^2} \right)_b \operatorname{div}(\rho_b \mathbf{u}) + \left( \frac{\Gamma P}{\rho} \right)_b (\mathbf{u} | \mathbf{a}_b) \right] + \\ &+ \left( \frac{\Gamma P}{\rho^2} \right)_b \left[ - \mathbf{a}_b \operatorname{div}(\rho_b \mathbf{u}) + (\mathbf{u} | \mathbf{a}_b) \nabla \rho_b \right] \end{aligned} \quad (4.3)$$

$$\mathcal{L}_1 \mathbf{u} = \nabla \mathcal{K}(\operatorname{div}(\rho_b \mathbf{u})). \quad (4.4)$$

Now we suppose the Assumption 1 and the background  $(\rho_b, S_b, \mathbf{v}_b)$  is a barotropic admissible stationary solution, and we denote  $\mathfrak{R}_b, \rho_b(\mathbf{x}), P_b(\mathbf{x}), S_b(\mathbf{x}), \mathbf{a}_b, \mathcal{A}_b$  by  $\mathfrak{R}, \rho(\mathbf{x}), P(\mathbf{x}), S(\mathbf{x}), \mathbf{a}, \mathcal{A}$  for the sake of simplicity of symbols.

Let us consider the integro-differential operator  $\mathcal{L}$  in the Hilbert space  $\mathfrak{H} = L^2(\mathfrak{R}, \rho d\mathbf{x}; \mathbb{C}^3)$  endowed with the inner product

$$(\mathbf{u}_1 | \mathbf{u}_2)_{\mathfrak{H}} = \int_{\mathfrak{R}} (\mathbf{u}_1(\mathbf{x}) | \mathbf{u}_2(\mathbf{x})) \rho(\mathbf{x}) d\mathbf{x}. \quad (4.5)$$

Of course

$$(\mathbf{u}_1(\mathbf{x})|\mathbf{u}_2(\mathbf{x})) := \sum_k u_1^k(\mathbf{x})u_2^k(\mathbf{x})^* \quad \text{for} \quad \mathbf{u}_\mu(\mathbf{x}) = \begin{bmatrix} u_\mu^1(\mathbf{x}) \\ u_\mu^2(\mathbf{x}) \\ u_\mu^3(\mathbf{x}) \end{bmatrix}, \mu = 1, 2.$$

Here and hereafter  $Z^*$  denotes the complex conjugate  $X - iY$  of  $Z = X + iY$ , while  $i$  stands for the imaginary unit,  $\sqrt{-1}$ .

First we observe  $\mathcal{L}$  restricted on  $C_0^\infty(\mathfrak{R}; \mathbb{C}^3)$ .

We have the following formula by integration by parts:

$$\begin{aligned} (\mathcal{L}_0 \mathbf{u}_1 | \mathbf{u}_2) &= \int \frac{\Gamma P}{\rho^2} \operatorname{div}(\rho \mathbf{u}_1) \operatorname{div}(\rho \mathbf{u}_2)^* + \\ &\quad + \int \frac{\Gamma P}{\rho} \left( -(\mathbf{u}_1 | \mathbf{a}) \operatorname{div}(\rho \mathbf{u}_2)^* - \operatorname{div}(\rho \mathbf{u}_1) (\mathbf{u}_2 | \mathbf{a})^* \right) + \\ &\quad + \int \frac{\Gamma P}{\rho} (\mathbf{u}_1 | \mathbf{a}) (\nabla \rho | \mathbf{u}_2) \\ &= \int \frac{\Gamma P}{\rho^2} \operatorname{div}(\rho \mathbf{u}_1) \operatorname{div}(\rho \mathbf{u}_2)^* + \\ &\quad + \int \frac{\Gamma P \mathcal{A}}{\rho} \left( (\mathbf{u}_1 | \mathbf{n}) \operatorname{div}(\rho \mathbf{u}_2)^* + (\operatorname{div}(\rho \mathbf{u}_1) (\mathbf{u}_2 | \mathbf{n})^* \right) \\ &\quad - \int \frac{\Gamma P \mathcal{A}}{\rho} \|\nabla \rho\|^2 (\mathbf{u}_1 | \mathbf{n}) (\mathbf{u}_2 | \mathbf{n})^*, \end{aligned}$$

where we recall

$$\mathbf{a} = -\mathcal{A} \mathbf{n}, \quad \mathbf{n} = \frac{\nabla \rho}{\|\nabla \rho\|}, \quad (4.6)$$

and

$$(\mathcal{L}_1 \mathbf{u}_1 | \mathbf{u}_2) = - \int \mathcal{K} [\operatorname{div}(\rho \mathbf{u}_1)] \operatorname{div}(\rho \mathbf{u}_2)^*. \quad (4.7)$$

Hence

$$\begin{aligned} (\mathcal{L} \mathbf{u}_1 | \mathbf{u}_2) &= \int \frac{\Gamma P}{\rho^2} \operatorname{div}(\rho \mathbf{u}_1) \operatorname{div}(\rho \mathbf{u}_2)^* + 2\Re \left[ \int \frac{\Gamma P \mathcal{A}}{\rho} (\mathbf{u}_1 | \mathbf{n}) \operatorname{div}(\rho \mathbf{u}_2)^* \right] \\ &\quad - \int \frac{\Gamma P \mathcal{A}}{\rho} \|\nabla \rho\|^2 (\mathbf{u}_1 | \mathbf{n}) (\mathbf{u}_2 | \mathbf{n})^* + \\ &\quad - 4\pi \mathbf{G} \int \mathcal{K} [\operatorname{div}(\rho \mathbf{u}_1)] \operatorname{div}(\rho \mathbf{u}_2)^*. \end{aligned} \quad (4.8)$$

Then it is clear that

$$(\mathcal{L} \mathbf{u}_1 | \mathbf{u}_2) = (\mathbf{u}_1 | \mathcal{L} \mathbf{u}_2) \quad \text{for} \quad \forall \mathbf{u}_1, \mathbf{u}_2 \in C_0^\infty(\mathfrak{R}), \quad (4.9)$$

that is,  $\mathcal{L}$  restricted on  $C_0^\infty(\mathfrak{R})$  is a symmetric operator.

**Remark 8** *When the background is not barotropic, then  $\mathcal{L}_0$  is not symmetric. In fact, we consider*

$$\begin{aligned} (\mathcal{L}_0 \mathbf{u}_1 | \mathbf{u}_2)_{\mathfrak{H}} - (\mathbf{u}_1 | \mathcal{L}_0 \mathbf{u}_2)_{\mathfrak{H}} &= \int \frac{\Gamma P}{\rho} \left( (\mathbf{u}_1 | \mathbf{a})(\nabla \rho | \mathbf{u}_2) - (\mathbf{u}_1 | \nabla \rho)(\mathbf{a} | \mathbf{u}_2) \right) d\mathbf{x} \\ &= \int \frac{1}{\rho} \left( -(\mathbf{u}_1 | \nabla P)(\nabla \rho | \mathbf{u}_2) + (\mathbf{u}_1 | \nabla \rho)(\nabla P | \mathbf{u}_2) \right) d\mathbf{x} \\ &=: \clubsuit \end{aligned}$$

for  $\mathbf{u}_1, \mathbf{u}_2 \in C_0^\infty(\mathfrak{R})$ . Put

$$c(\mathbf{x}) := \frac{|(\nabla \rho(\mathbf{x}) | \nabla P(\mathbf{x}))|}{\|\nabla \rho(\mathbf{x})\| \|\nabla P(\mathbf{x})\|}.$$

If  $\nabla P$  is not parallel to  $\nabla \rho$  at a point  $\mathbf{x}_1 \in \mathfrak{R}$ , then there is a neighborhood  $\mathcal{V}$  of  $\mathbf{x}_1$ , say a small ball with center  $\mathbf{x}_1$  and radius  $\mathfrak{d}$ , such that

$$\mathcal{V} \Subset \mathfrak{R}, \quad c(\mathbf{x})^2 \leq 1 - \kappa \quad \forall \mathbf{x} \in \mathcal{V}$$

with  $0 < \kappa < 1$ . Consider  $\mathbf{u}_1, \mathbf{u}_2 \in C_0^\infty(\mathfrak{R})$  such that

$$\mathbf{u}_1 = \nabla \rho, \quad \mathbf{u}_2 = \nabla P \quad \text{on } \mathcal{V}. \quad (4.10)$$

Then

$$\begin{aligned} \clubsuit &= \int_{\mathcal{V}} \frac{1}{\rho} \|\nabla \rho\|^2 \|\nabla P\|^2 (1 - c(\mathbf{x})^2) d\mathbf{x} + I \\ &\geq \varepsilon + I, \end{aligned}$$

where

$$\begin{aligned} \varepsilon &= \kappa \inf_{\mathcal{V}} \frac{1}{\rho} \|\nabla \rho\|^2 \|\nabla P\|^2 \cdot \frac{4\pi}{3} \mathfrak{d}^3 > 0, \\ I &= \int_{\mathfrak{R} \setminus \mathcal{V}} \frac{1}{\rho} \left( -(\mathbf{u}_1 | \nabla P)(\nabla \rho | \mathbf{u}_2) + (\mathbf{u}_1 | \nabla \rho)(\nabla P | \mathbf{u}_2) \right) d\mathbf{x}. \end{aligned}$$

Clearly we can find  $\mathbf{u}_1, \mathbf{u}_2 \in C_0^\infty(\mathfrak{R})$  which enjoy (4.10) and  $|I| \leq \frac{\varepsilon}{2}$ . Then  $\clubsuit \geq \frac{\varepsilon}{2} > 0$ , that is,  $(\mathcal{L}_0 \mathbf{u}_1 | \mathbf{u}_2)_{\mathfrak{H}} \neq (\mathbf{u}_1 | \mathcal{L}_0 \mathbf{u}_2)_{\mathfrak{H}}$ .

Moreover we have

$$\begin{aligned} (\mathcal{L}_0 \mathbf{u} | \mathbf{u}) &= \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} + 2\Re \left[ \int \frac{\Gamma P \mathcal{A}}{\rho} (u | \mathbf{n}) \operatorname{div}(\rho \mathbf{u})^* \right] + \\ &\quad - \int \frac{\Gamma P \mathcal{A}}{\rho} |(\mathbf{u} | \mathbf{n})|^2 \|\nabla \rho\|. \end{aligned} \quad (4.11)$$

Since  $\mathcal{A} \in C^\alpha(\mathfrak{R} \cup \partial\mathfrak{R})$ , we have

$$\kappa_1 := \sup_{\mathfrak{R}} \left\{ |\mathcal{A}| \sqrt{\frac{\Gamma P}{\rho}} \right\} < \infty. \quad (4.12)$$

Therefore

$$\begin{aligned} \left| \int \frac{\Gamma P \mathcal{A}}{\rho} (\mathbf{u} | \mathbf{n}) \operatorname{div}(\rho \mathbf{u})^* \right| &\leq \frac{\kappa_1}{2} \left[ \frac{1}{\epsilon} \int |(\mathbf{u} | \mathbf{n})|^2 \rho d\mathbf{x} + \epsilon \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} \right] \\ &\leq \frac{\kappa_1}{2} \left[ \frac{1}{\epsilon} \|\mathbf{u}\|_{\mathfrak{H}}^2 + \epsilon \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} \right]. \end{aligned}$$

On the other hand, we have

$$\kappa_2 := \sup_{\mathfrak{R}} \left\{ \frac{\Gamma P |\mathcal{A}|}{\rho^2} \|\nabla \rho\| \right\} < \infty. \quad (4.13)$$

Then

$$\left| \int \frac{\Gamma P \mathcal{A}}{\rho} |(\mathbf{u} | \mathbf{n})|^2 \|\nabla \rho\| \right| \leq \kappa_2 \|\mathbf{u}\|_{\mathfrak{H}}^2.$$

Taking  $\epsilon = \frac{1}{2\kappa_1}$ , we have

$$\begin{aligned} \frac{1}{2} \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} - (2\kappa_1^2 + \kappa_2) \|\mathbf{u}\|_{\mathfrak{H}}^2 &\leq (\mathcal{L}_0 \mathbf{u} | \mathbf{u}) \leq \\ &\leq \frac{3}{2} \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} + (2\kappa_1^2 + \kappa_2) \|\mathbf{u}\|_{\mathfrak{H}}^2 \end{aligned} \quad (4.14)$$

and, taking  $\epsilon = \frac{2}{\kappa_1}$ , we have

$$-\left(\frac{\kappa_1^2}{4} + \kappa_2\right) \|\mathbf{u}\|_{\mathfrak{H}}^2 \leq (\mathcal{L}_0 \mathbf{u} | \mathbf{u}) \leq 2 \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} + \left(\frac{\kappa_1^2}{4} + \kappa_2\right) \|\mathbf{u}\|_{\mathfrak{H}}^2. \quad (4.15)$$

As for  $\mathcal{L}_1$ , on the other hand, we have

**Proposition 1** *It holds that*

$$-\rho_O \|\mathbf{u}\|_{\mathfrak{H}}^2 \leq (\mathcal{L}_1 \mathbf{u} | \mathbf{u})_{\mathfrak{H}} \leq 0. \quad (4.16)$$

Proof. Look at

$$\Psi := -\mathcal{K}[g], \quad \mathbf{C} := \rho \mathbf{u} - \nabla \Psi. \quad (4.17)$$

Since  $\triangle \Psi = g$ , we have

$$\operatorname{div} \mathbf{C} = 0.$$

Keeping in mind that  $\Psi = O(\frac{1}{r})$ ,  $\nabla \Psi, \mathbf{C} = O(\frac{1}{r^2})$  as  $r \rightarrow +\infty$ , we derive from this that

$$\int_{\mathbb{R}^3} (\nabla \Psi | \mathbf{C}) d\mathbf{x} = 0 \quad (4.18)$$



Then we see

$$\begin{aligned}
\int_{\mathfrak{R}} \mathcal{K}[g]g^* d\mathbf{x} &= \int_{\mathfrak{R}} (\nabla \Psi | \rho \mathbf{u}) \\
&= \int_{\mathbb{R}^3} (\nabla \Psi | \mathbf{C} + \nabla \Psi) d\mathbf{x} \\
&= \int_{\mathbb{R}^3} \|\nabla \Psi\|^2 d\mathbf{x} \quad (\text{by (4.18)}) \\
&= \int_{\mathfrak{R}} \|\rho \mathbf{u}\|^2 d\mathbf{x} - \int_{\mathbb{R}^3} \|\mathbf{C}\|^2 d\mathbf{x} \quad (\text{again by (4.18)}) \\
&\leq \int_{\mathfrak{R}} \|\rho \mathbf{u}\|^2 d\mathbf{x} \\
&\leq \rho_O \int_{\mathfrak{R}} \|u\|^2 \rho d\mathbf{x} = \rho_O \|\mathbf{u}\|_{\mathfrak{H}}^2.
\end{aligned}$$

and

$$\int_{\mathfrak{R}} \mathcal{K}[g]g^* d\mathbf{x} = \int_{\mathbb{R}^3} \|\nabla \Psi\|^2 d\mathbf{x} \geq 0.$$

□

This proof is due to Juhi Jang. See [11, Proposition 2].

Summing up, we have

$$\begin{aligned}
\frac{1}{2} \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} - (2\kappa_1^2 + \kappa_2 + 4\pi \mathsf{G} \rho_O) \|\mathbf{u}\|_{\mathfrak{H}}^2 &\leq (\mathcal{L} \mathbf{u} | \mathbf{u}) \leq \\
&\leq \frac{3}{2} \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} + (2\kappa_1^2 + \kappa_2) \|\mathbf{u}\|_{\mathfrak{H}}^2.
\end{aligned} \tag{4.19}$$

and, on the other hand, thanks to (4.15), we have

$$-\left(\frac{\kappa_1^2}{4} + \kappa_2 + 4\pi \mathsf{G} \rho_O\right) \|\mathbf{u}\|_{\mathfrak{H}}^2 \leq (\mathcal{L} \mathbf{u} | \mathbf{u}) \leq 2 \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} + \left(\frac{\kappa_1^2}{4} + \kappa_2\right) \|\mathbf{u}\|_{\mathfrak{H}}^2. \tag{4.20}$$

Therefore the operator  $\mathcal{L}$  restricted on  $C_0^\infty(\mathfrak{R})$  is symmetric, and is bounded from below as an operator in  $\mathfrak{H}$ . Applying the Stone-Friedrichs theorem, (e.g., [12, Chapter VI, Section 2.3]), we can claim:

**Theorem 2**  $\mathcal{L} \upharpoonright C_0^\infty(\mathfrak{R})$  admits the Friedrichs extension  $\mathbf{L}$ , which is a self-adjoint operator in  $\mathfrak{H}$ , whose domain is

$$\mathbf{D}(\mathbf{L}) = \{\mathbf{u} \in \mathfrak{G}_0 \mid \mathcal{L} \mathbf{u} \in \mathfrak{H}\}. \tag{4.21}$$

Here  $\mathfrak{G}$  is the Hilbert space of all  $\mathbf{u} \in \mathfrak{H}$  such that  $\int_{\mathfrak{R}} \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} < \infty$  endowed with the inner product

$$(\mathbf{u}_1 | \mathbf{u}_2)_{\mathfrak{G}} = (\mathbf{u}_1 | \mathbf{u}_2)_{\mathfrak{H}} + \int_{\mathfrak{R}} \frac{\Gamma P}{\rho^2} (\operatorname{div}(\rho \mathbf{u}_1) (\operatorname{div}(\rho \mathbf{u}_2))^* d\mathbf{x}, \tag{4.22}$$

and  $\mathfrak{G}_0$  is the closure of  $C_0^\infty(\mathfrak{R})$  in  $\mathfrak{G}$ .

In fact, we define the quadratic form  $Q_0$  by

$$\begin{aligned} Q_0[\mathbf{u}] = Q_0(\mathbf{u}, \mathbf{u}) &= \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} + 2\Re \left[ \int \frac{\Gamma P \mathcal{A}}{\rho} (\mathbf{u} | \mathbf{n}) \operatorname{div}(\rho \mathbf{u})^* d\mathbf{x} \right] + \\ &\quad - \int \frac{\Gamma P \mathcal{A}}{\rho} |(\mathbf{u} | \mathbf{n})|^2 \|\nabla \rho\| d\mathbf{x} - 4\pi \mathbf{G} \int \mathcal{K}[\operatorname{div}(\rho \mathbf{u})] \operatorname{div}(\rho \mathbf{u})^* d\mathbf{x}. \end{aligned} \quad (4.23)$$

Then it holds that

$$(\mathcal{L} \mathbf{u}_1 | \mathbf{u}_2)_{\mathfrak{H}} = Q_0(\mathbf{u}_1, \mathbf{u}_2) \quad (4.24)$$

for  $\forall \mathbf{u}_1, \mathbf{u}_2 \in C_0^\infty(\mathfrak{R})$ . We define

$$Q_a(\mathbf{u}_1, \mathbf{u}_2) = Q_0(\mathbf{u}_1, \mathbf{u}_2) + a(\mathbf{u}_1 | \mathbf{u}_2)_{\mathfrak{H}} \quad (4.25)$$

with

$$a := 2\kappa_1^2 + \kappa_2 + 4\pi \mathbf{G} \rho_O. \quad (4.26)$$

Since

$$\frac{1}{2} \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} \leq Q_a[\mathbf{u}] \leq \frac{3}{2} \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} + (2\kappa_1^2 + \kappa_2 + a) \|\mathbf{u}\|_{\mathfrak{H}}^2, \quad (4.27)$$

we have

$$0 \leq Q_a[\mathbf{u}] \quad \text{for } \mathbf{u} \in \mathfrak{G}. \quad (4.28)$$

Since

$$\frac{1}{2} \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} \leq Q_a[\mathbf{u}] \leq \frac{3}{2} \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} + (2\kappa_1^2 + \kappa_2 + a) \|\mathbf{u}\|_{\mathfrak{H}}^2 \quad (4.29)$$

for  $\mathbf{u} \in \mathfrak{G}$ , we have

$$\frac{1}{C} \|\mathbf{u}\|_{\mathfrak{G}} \leq [\|\mathbf{u}\|] \leq C \|\mathbf{u}\|_{\mathfrak{G}}, \quad (4.30)$$

where

$$[\|\mathbf{u}\|] := \left[ \|\mathbf{u}\|_{\mathfrak{H}}^2 + Q_a[\mathbf{u}] \right]^{\frac{1}{2}}. \quad (4.31)$$

Thus the norm  $\|\cdot\|_{\mathfrak{G}}$  can be replaced by  $[\|\cdot\|]$  equivalently. We have

$$((\mathbf{L} + a) \mathbf{u}_1 | \mathbf{u}_2)_{\mathfrak{H}} = Q_a(\mathbf{u}_1, \mathbf{u}_2) \quad (4.32)$$

for  $\forall \mathbf{u}_1 \in \mathbf{D}(\mathbf{L}), \forall \mathbf{u}_2 \in \mathfrak{G}_0$ .

In this situation we can apply the Hille-Yosida theory, as described in **Appendix A**. The conclusion is:

**Theorem 3** Suppose  $\mathbf{u}^0 \in \mathbf{D}(\mathbf{L})$ ,  $\mathbf{v}^0 \in \mathfrak{G}_0$  and  $\mathbf{f} \in C([0, +\infty[; \mathfrak{H})$ . Then the initial-boundary value problem (4.1a) (4.1b) admits a unique solution  $\mathbf{u} = \mathbf{u}(t, \mathbf{x})$  in

$$C^2([0, +\infty[, \mathfrak{H}) \cap C^1([0, +\infty[, \mathfrak{G}_0) \cap C([0, +\infty[, \mathbf{D}(\mathbf{L}))$$

and the energy

$$\begin{aligned} E(t, \mathbf{u}) &:= \|\mathbf{u}\|^2 + \left\| \frac{\partial \mathbf{u}}{\partial t} \right\|_{\mathfrak{H}}^2 \\ &= \|\mathbf{u}\|_{\mathfrak{H}}^2 + Q_a[\mathbf{u}] + \left\| \frac{\partial \mathbf{u}}{\partial t} \right\|_{\mathfrak{H}}^2 = (1+a)\|\mathbf{u}\|_{\mathfrak{H}}^2 + Q_0[\mathbf{u}] + \left\| \frac{\partial \mathbf{u}}{\partial t} \right\|_{\mathfrak{H}}^2 \end{aligned} \quad (4.33)$$

enjoys the estimate

$$\sqrt{E(t, \mathbf{u})} \leq e^{\kappa t} \sqrt{E(0, \mathbf{u})} + \int_0^t e^{\kappa(t-s)} \|\mathbf{f}(s)\|_{\mathfrak{H}} ds \quad (4.34)$$

for

$$\kappa = 1 + a + \max_{i,j} \|B_j^i\|_{L^\infty(\mathfrak{R})}.$$

Let us claim the following

**Theorem 4** 0 is an eigenvalue of  $\mathbf{L}$  and  $\dim \text{Ker}[\mathbf{L}] = \infty$ .

Proof. 1) Suppose  $\mathcal{A} \neq 0$ , that is,  $\nabla S_b(\mathbf{x}) \neq 0$  for  $\exists \mathbf{x} \in \mathfrak{R}$ . Then

$$\mathbf{u}^0(\mathbf{x}) = \frac{1}{\rho_b(\mathbf{x})} \nabla S_b(\mathbf{x}) \times \nabla f(\mathbf{x}), \quad (4.35)$$

where  $f \in C_0^\infty(\mathfrak{R}; \mathbb{R})$  is arbitrary, enjoys

$$\begin{aligned} [\mathfrak{d}\rho; \mathbf{u}^0] &= -\text{div}(\rho_b \mathbf{u}^0) = 0, \\ [\mathfrak{d}S; \mathbf{u}^0] &= -(\mathbf{u}^0 | \nabla S_b) = 0, \\ [\mathfrak{d}P; \mathbf{u}^0] &= -\left(\frac{\Gamma P}{\rho}\right)_b [\mathfrak{d}\rho; \mathbf{u}^0] + \left(\frac{\partial P}{\partial S}\right)_b [\mathfrak{d}S; \mathbf{u}^0] = 0, \\ [\mathfrak{d}\Phi; \mathbf{u}^0] &= -4\pi \text{GK}[[\mathfrak{d}\rho; \mathbf{u}^0]] = 0, \end{aligned}$$

therefore  $\mathbf{L}\mathbf{u}^0 = \mathbf{0}$ . (Recall the notations of Remark 6.) If  $\nabla f(\mathbf{x}_1) \neq 0$  and  $\nabla S_b(\mathbf{x}_1) \neq \mathbf{0}$  at  $\exists \mathbf{x}_1 \in \mathfrak{R}$ , then  $\mathbf{u}^0 \neq 0$  and  $\mathbf{u}^0$  is an eigenvector of the eigenvalue 0.

2) Suppose  $\mathcal{A} = 0$ , that is,  $\nabla S_b(\mathbf{x}) = 0$  for  $\forall \mathbf{x} \in \mathfrak{R}$ , or, the background is isentropic. Then, for any vector field  $\mathbf{f} \in C_0^\infty(\mathfrak{R}; \mathbb{R}^3)$ , the vector field  $\mathbf{u}^0$  defined by

$$\mathbf{u}^0(\mathbf{x}) = \frac{1}{\rho_b(\mathbf{x})} \text{rot} \mathbf{f}(\mathbf{x}) \quad (4.36)$$

enjoys  $\mathfrak{d}\rho = 0, \mathfrak{d}S = 0, \mathfrak{d}P = 0, \mathfrak{d}\Phi = 0$ , so that  $\mathbf{L}\mathbf{u}^0 = \mathbf{0}$ . If  $\text{rot} \mathbf{f} \neq \mathbf{0}$  somewhere, then  $\mathbf{u}^0 \neq \mathbf{0}$  and it is an eigenvector.  $\square$

Roughly or formally speaking, the boundary condition for  $\mathbf{u} \in \mathcal{D}(\mathbf{L})$  is:

$$\operatorname{div}(\rho \mathbf{u}) = 0 \quad \text{on} \quad \partial \mathfrak{R},$$

or

$$(\mathbf{n}|\mathbf{u}) = 0 \quad \text{on} \quad \partial \mathfrak{R},$$

since, formally,

$$\operatorname{div}(\rho \mathbf{u}) = \rho \operatorname{div} \mathbf{u} + (\nabla \rho | \mathbf{u})$$

with  $\rho = 0$  and  $\mathbf{n} = \nabla \rho / \|\nabla \rho\|$  on the boundary.

## 5 Eigenvalue problem

We are considering the equation

$$\frac{\partial^2 \mathbf{u}}{\partial t^2} + \mathbf{B} \frac{\partial \mathbf{u}}{\partial t} + \mathbf{L} \mathbf{u} = 0. \quad (5.1)$$

Keeping in mind  $\mathbf{B} \mathbf{v} = 2\Omega \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \mathbf{v}$ , we suppose that  $\mathbf{B}$  is an anti-symmetric bounded linear operator in  $\mathcal{B}(\mathfrak{H})$ . Therefore  $(\mathbf{B} \mathbf{v} | \mathbf{v}) = -(\mathbf{v} | \mathbf{B} \mathbf{v})$  and  $\Re[(\mathbf{B} \mathbf{v} | \mathbf{v})] = 0 \quad \forall \mathbf{v}$ . Put

$$\beta := \|\mathbf{B}\|_{\mathcal{B}(\mathfrak{H})} = \sup \frac{\|\mathbf{B} \mathbf{v}\|_{\mathfrak{H}}}{\|\mathbf{v}\|_{\mathfrak{H}}}. \quad (5.2)$$

$\mathbf{L}$  is a self-adjoint operator in  $\mathfrak{H}$  and

$$(\mathbf{L} \mathbf{u} | \mathbf{u})_{\mathfrak{H}} \geq -\mu_* \|\mathbf{u}\|_{\mathfrak{H}}^2 \quad \forall \mathbf{u} \in \mathcal{D}(\mathbf{L}), \quad (5.3)$$

$\mu_*$  being a nonnegative number, say,

$$\mu_* := \left( - \inf_{\mathbf{u} \in \mathcal{D}(\mathbf{L})} \frac{(\mathbf{L} \mathbf{u} | \mathbf{u})_{\mathfrak{H}}}{\|\mathbf{u}\|_{\mathfrak{H}}^2} \right) \vee 0. \quad (5.4)$$

We know

$$\mu_* \leq \frac{\kappa_1^2}{4} + \kappa_2 + 4\pi \mathbb{G} \rho_O$$

by (4.20).

Now we look for nontrivial solution of (5.1) of the form  $\mathbf{u}(t, \mathbf{x}) = e^{\lambda t} \mathbf{u}^0(\mathbf{x})$ , where  $\lambda \in \mathbb{C}$ ,  $\mathbf{u}^0 \in \mathcal{D}(\mathbf{L})$ . Namely

**Definition 4** *If there is a  $\mathbf{u}^0 \in \mathcal{D}(\mathbf{L})$  such that  $\mathbf{u}^0 \neq \mathbf{0}$ , and*

$$\left[ \lambda^2 + \lambda \mathbf{B} + \mathbf{L} \right] \mathbf{u}^0 = \mathbf{0} \quad (5.5)$$

*holds, then  $\lambda$  is called an eigenvalue of (5.1) and  $\mathbf{u}^0$  is called an eigenvector associated with the eigenvalue  $\lambda$ .*

If  $\lambda$  is an eigenvalue of (5.1) and  $\mathbf{u}^0$  is an associated eigenvector, then

1)  $\mathbf{u} : (t, \mathbf{x}) \mapsto e^{\lambda t} \mathbf{u}^0(\mathbf{x})$  is a solution of (5.1) with the initial condition

$$\mathbf{u}|_{t=0} = \mathbf{u}^0(\mathbf{x}), \quad \frac{\partial \mathbf{u}}{\partial t} \Big|_{t=0} = \lambda \mathbf{u}^0(\mathbf{x}), \quad (5.6)$$

2)  $\mathbf{u}^{\Re} : (t, \mathbf{x}) \mapsto \Re[e^{\lambda t} \mathbf{u}^0(\mathbf{x})]$  is a solution of (5.1) with the initial condition

$$\mathbf{u}^{\Re}|_{t=0} = \Re[\mathbf{u}^0(\mathbf{x})], \quad \frac{\partial \mathbf{u}^{\Re}}{\partial t} \Big|_{t=0} = \Re[\lambda] \Re[\mathbf{u}^0(\mathbf{x})] - \Im[\lambda] \Im[\mathbf{u}^0(\mathbf{x})], \quad (5.7)$$

3)  $\lambda$  satisfies the quadratic equation

$$a\lambda^2 + ib\lambda + c = 0 \quad (5.8)$$

where the coefficients  $a, b, c$  are given by

$$a = \|\mathbf{u}^0\|_{\mathfrak{H}}^2, \quad b = -i(B\mathbf{u}^0|\mathbf{u}^0)_{\mathfrak{H}}, \quad c = (L\mathbf{u}^0|\mathbf{u}^0)_{\mathfrak{H}}, \quad (5.9)$$

and  $a, b, c$  are real numbers.

If  $L$  is the operator defined by Theorem 2, it follows from Theorem 4 that 0 is an eigenvalue of (5.1) and the multiplicity is infinite. The solution  $\mathbf{u}(t, \mathbf{x}) = \mathbf{u}^0(\mathbf{x})$  with  $L\mathbf{u}^0 = \mathbf{0}$  may be called a ‘trivial perturbation’ after [7].

By the operator  $A$  in  $\mathfrak{E} = \mathfrak{G}_0 \times \mathfrak{H}$  defined by

$$AU = \begin{bmatrix} O & -I \\ L & B \end{bmatrix} U, \quad D(A) = D(L) \times \mathfrak{G}_0, \quad (5.10)$$

the equation (5.1) reads

$$\frac{dU}{dt} + AU = \mathbf{0}$$

with

$$U = \begin{bmatrix} \mathbf{u} \\ \frac{\partial \mathbf{u}}{\partial t} \end{bmatrix}.$$

**Proposition 2**  $\lambda \in \mathbb{C}$  is an eigenvalue of (5.1) if and only if  $\lambda$  is an eigenvalue of the operator  $-A$ .

Following [6], we use

**Definition 5** The family of operators  $\mathfrak{L} = (\mathfrak{L}(\lambda))_{\lambda \in \mathbb{C}}$ , where

$$\mathfrak{L}(\lambda) = \lambda^2 + \lambda \mathbf{B} + \mathbf{L}, \quad (5.11)$$

is called the ‘quadratic pencil’ of the equation (5.1).

The resolvent set  $\mathbf{P}(\mathfrak{L})$  is the set of  $\lambda \in \mathbb{C}$  such that  $\mathfrak{L}(\lambda)$  has the bounded inverse in  $\mathcal{B}(\mathfrak{H})$ .  $\mathbb{C} \setminus \mathbf{P}(\mathfrak{L})$  is the spectrum of  $\mathfrak{L}$ , and is denoted by  $\Sigma(\mathfrak{L})$ . The set of all eigenvalues of (5.1) is denoted by  $\Sigma_p(\mathfrak{L})$ .

If  $\lambda$  is an eigenvalue of (5.1), then it belongs to the spectrum of  $\mathfrak{L}$ , that is,  $\Sigma_p(\mathfrak{L}) \subset \Sigma(\mathfrak{L})$ . However we cannot say that any  $\lambda \in \Sigma(\mathfrak{L})$  is an eigenvalue of (5.1), that is,  $\Sigma_p(\mathfrak{L}) = \Sigma(\mathfrak{L})$ , a priori. We are going to study the structure of  $\Sigma(\mathfrak{L})$  as much as possible at the moment.

**Proposition 3**  $\mathbf{P}(\mathfrak{L})$  is an open subset of  $\mathbb{C}$  and  $\Sigma(\mathfrak{L})$  is closed .

Proof. Let us consider  $\lambda \in \mathbf{P}(\mathfrak{L})$ . Then

$$\mathfrak{L}(\lambda + \Delta\lambda) = \mathfrak{L}(\lambda) \left[ I + \mathfrak{L}(\lambda)^{-1}(\Delta\lambda(2\lambda + \mathbf{B} + \Delta\lambda)) \right]$$

admits the bounded inverse and  $\lambda + \Delta\lambda \in \mathbf{P}(\mathfrak{L})$ , if

$$\| \mathfrak{L}(\lambda)^{-1} \Delta\lambda(2\lambda + \mathbf{B} + \Delta\lambda) \|_{\mathcal{B}(\mathfrak{H})} < 1.$$

For this inequality, it is sufficient that

$$\| \mathfrak{L}(\lambda)^{-1} \|_{\mathcal{B}(\mathfrak{H})} \cdot |\Delta\lambda| \cdot (2|\lambda| + \beta + |\Delta\lambda|) < 1,$$

or

$$|\Delta\lambda| < -\left(|\lambda| + \frac{\beta}{2}\right) + \sqrt{\left(|\lambda| + \frac{\beta}{2}\right)^2 + \frac{1}{\| \mathfrak{L}(\lambda)^{-1} \|_{\mathcal{B}(\mathfrak{H})}}}.$$

This means  $\mathbf{P}(\mathfrak{L})$  is open.  $\square$

**Proposition 4**  $\mathbf{P}(\mathfrak{L})$  and  $\Sigma(\mathfrak{L})$  are symmetric about the imaginary axis  $i\mathbb{R}$ .

In fact we see  $\mathfrak{L}(\lambda)^* = \mathfrak{L}(-\lambda^*)$ .

**Proposition 5** It hold

$$\mathbf{P}(\mathfrak{L}) = \mathbf{P}(-\mathbf{A}), \quad \Sigma(\mathfrak{L}) = \Sigma(-\mathbf{A}), \quad \Sigma_p(\mathfrak{L}) \subset \Sigma_p(-\mathbf{A}).$$

Here  $\mathbf{P}(-\mathbf{A}), \Sigma(-\mathbf{A}), \Sigma_p(-\mathbf{A})$  stand for the usual resolvent set, spectrum, point spectrum (the set of all eigenvalues) of the operator  $-\mathbf{A}$  in  $\mathfrak{G}_0 \times \mathfrak{H}$ .

Proof. Let  $\lambda \in \mathbf{P}(\mathfrak{L})$ . Consider the equation

$$\mathbf{A}U + \lambda U = F = \begin{bmatrix} \mathbf{g} \\ \mathbf{f} \end{bmatrix} \in \mathfrak{G}_0 \times \mathfrak{H},$$

or,

$$\begin{cases} -\mathbf{v} + \lambda \mathbf{u} = \mathbf{g} \\ \mathbf{L}\mathbf{u} + \mathbf{B}\mathbf{u} + \lambda \mathbf{v} = \mathbf{f} \end{cases}$$

This system of equations can be solved as

$$\begin{cases} \mathbf{u} = \mathfrak{L}(\lambda)^{-1}(\mathbf{f} + \mathbf{B}\mathbf{g} - \lambda \mathbf{f}) \\ \mathbf{v} = \lambda \mathfrak{L}(\lambda)^{-1}(\mathbf{f} + \mathbf{B}\mathbf{g} - \lambda \mathbf{g}) - \mathbf{f} \end{cases}$$

since  $\lambda \in \mathbf{P}(\mathfrak{L})$ , while  $F \mapsto U$  is continuous. Therefore  $\lambda \in \mathbf{P}(-\mathbf{A})$ .

Inversely let  $\lambda \in \mathbf{P}(-\mathbf{A})$ . Consider the equation

$$(\lambda^2 + \lambda \mathbf{B} + \mathbf{L})\mathbf{u} = \mathbf{f} \in \mathfrak{H},$$

which is equivalent to the system of equations

$$\begin{cases} \lambda \mathbf{v} + \mathbf{B}\mathbf{v} + \mathbf{L}\mathbf{u} = \mathbf{f}, \\ \mathbf{v} = \lambda \mathbf{u} \end{cases}$$

But this is nothing but

$$\mathbf{A}U + \lambda U = \begin{bmatrix} \mathbf{0} \\ \mathbf{f} \end{bmatrix}$$

for  $U = (\mathbf{u}, \mathbf{v})^\top$ . Since  $\lambda \in \mathbf{P}(-\mathbf{A})$  is supposed, this admits the solution

$$U = \begin{bmatrix} \mathbf{u} \\ \mathbf{v} \end{bmatrix} = (\mathbf{A} + \lambda)^{-1} \begin{bmatrix} \mathbf{0} \\ \mathbf{f} \end{bmatrix},$$

and  $\mathbf{f} \mapsto \mathbf{u}$  is continuous, that is,  $\lambda \in \mathbf{P}(\mathfrak{L})$ .  $\square$

Let us consider the case of  $\mathbf{B} = O$ , or  $\Omega = 0$ . Then

$$\Sigma(\mathfrak{L}) = \left\{ \lambda \in \mathbb{C} \mid -\lambda^2 \in \Sigma(\mathbf{L}) \right\},$$

where  $\Sigma(\mathbf{L})$  is the spectrum of the operator  $\mathbf{L}$  in  $\mathfrak{H}$ . We know  $\Sigma(\mathbf{L}) \subset [-\mu_*, +\infty[$ . Therefore we see

$$\Sigma(\mathfrak{L}) \subset i\mathbb{R} \cup [-\sqrt{\mu_*}, \sqrt{\mu_*}].$$

But, when  $\mathbf{B} \neq 0, \Omega \neq 0$ , the situation is not so simple. Note that, when  $\mathbf{B} \neq 0, \Omega \neq 0$ ,  $\mathbf{A}$  is not self-adjoint, since

$$(\mathbf{A}U|U)_{\mathfrak{E}} - (U|\mathbf{A}U)_{\mathfrak{E}} = (\mathbf{B}\mathbf{v}|\mathbf{v})_{\mathfrak{H}} - (\mathbf{v}|\mathbf{B}\mathbf{v})_{\mathfrak{H}} = 2(\mathbf{B}\mathbf{v}|\mathbf{v})_{\mathfrak{H}} \quad \text{for } U = \begin{bmatrix} \mathbf{u} \\ \mathbf{v} \end{bmatrix}.$$

At least we can claim

**Proposition 6** *It holds*

$$]-\infty, -A[ \cup ]A, +\infty[ \subset \mathbf{P}(\mathfrak{L}) = \mathbf{P}(-\mathbf{A}) \quad (5.12)$$

$$\text{with } A = \frac{\beta}{2} + \sqrt{\frac{\beta^2}{4} + \mu_*}.$$

For proof see Proposition **A2** of **Appendix A**.

**Proposition 7** *For  $\lambda \in \mathbf{P}(\mathfrak{L})$ , we have*

$$\|\mathfrak{L}(\lambda)^{-1}\|_{\mathcal{B}(\mathfrak{H})} \geq \frac{1}{d(2|\lambda| + \beta + d)} \quad (5.13)$$

where  $d := \text{dist.}(\lambda, \Sigma(\mathfrak{L}))$ .

Proof. Let  $\lambda \in \mathbf{P}(\mathfrak{L})$ . Then, for  $\Delta\lambda \in \mathbb{C}$ , the operator

$$\mathfrak{L}(\lambda + \Delta\lambda) = \mathfrak{L}(\lambda) \left[ I + \mathfrak{L}(\lambda)^{-1} \Delta\lambda (2\lambda + \mathbf{B} + \Delta\lambda) \right]$$

admits the bounded inverse in  $\mathcal{B}(\mathfrak{H})$  and  $\lambda + \Delta\lambda \in \mathbf{P}(\mathfrak{L})(\mathfrak{L})$ , if

$$\|\mathfrak{L}(\lambda)^{-1} \Delta\lambda (2\lambda + \mathbf{B} + \Delta\lambda)\|_{\mathcal{B}(\mathfrak{H})} < 1.$$

For this inequality, it is sufficient that

$$\|\mathfrak{L}(\lambda)^{-1}\|_{\mathcal{B}(\mathfrak{H})} \cdot |\Delta\lambda| \cdot (2|\lambda| + \beta + |\Delta\lambda|) < 1.$$

In other words, if  $\lambda + \Delta\lambda \in \Sigma(\mathfrak{L})$ , then it should hold

$$\|\mathfrak{L}(\lambda)^{-1}\|_{\mathcal{B}(\mathfrak{H})} \cdot |\Delta\lambda| \cdot (2|\lambda| + \beta + |\Delta\lambda|) \geq 1.$$

If  $d < +\infty$ , then there is a sequence  $\lambda + (\Delta\lambda)_n \in \Sigma(\mathfrak{L})$  such that  $|(\Delta\lambda)_n| \rightarrow d$ , and the assertion follows.  $\square$

**Theorem 5** ([6, Theorem 1])  $\Sigma(\mathfrak{L})$  is a subset of

$$S := i\mathbb{R} \cup \left\{ \lambda \in \mathbb{C} \mid |\lambda| \leq \sqrt{\mu_*}, \quad |\Im \lambda| \leq \frac{\beta}{2} \right\}. \quad (5.14)$$



Proof. First we consider  $\lambda_\infty \in \partial\Sigma(\mathfrak{L})$ .

Let us take a sequence  $(\lambda_n)_n$  such that  $\lambda_n \in \mathbf{P}(\mathfrak{L})$  and  $\lambda_n \rightarrow \lambda_\infty$  as  $n \rightarrow \infty$ . By Proposition 7 we have  $\|(\mathfrak{L}(\lambda_n))^{-1}\|_{\mathcal{B}(\mathfrak{H})} \rightarrow +\infty$ , therefore there are  $\mathbf{f}_n \in \mathfrak{H}$  such that  $\|\mathbf{f}_n\|_{\mathfrak{H}} = 1$  and  $\|\mathfrak{L}(\lambda_n))^{-1}\mathbf{f}_n\|_{\mathfrak{H}} \rightarrow +\infty$  as  $n \rightarrow \infty$ . Put  $\mathbf{u}_n = \mathfrak{L}(\lambda_n)^{-1}\mathbf{f}_n (\in \mathbf{D}(\mathbf{L}))$  and  $\mathbf{u}_n = \mathbf{u}_n / \|\mathbf{u}_n\|_{\mathfrak{H}}$ . Then  $\|\mathbf{u}_n\|_{\mathfrak{H}} = 1$  and

$$\left| (\mathfrak{L}(\lambda_n)\mathbf{u}_n | \mathbf{u}_n)_{\mathfrak{H}} \right| = \left| \frac{1}{\|\mathbf{u}_n\|_{\mathfrak{H}}^2} (\mathbf{f}_n | \mathbf{u}_n)_{\mathfrak{H}} \right| \leq \frac{1}{\|\mathbf{u}_n\|_{\mathfrak{H}}} \rightarrow 0.$$

But we see

$$(\mathfrak{L}(\lambda_n)\mathbf{u}_n | \mathbf{u}_n)_{\mathfrak{H}} = \lambda_n^2 + i\lambda_n b_n + c_n,$$

where

$$b_n := -i(\mathbf{B}\mathbf{u} | \mathbf{u})_{\mathfrak{H}}, \quad c_n = (\mathbf{L}\mathbf{u}_n | \mathbf{u}_n)_{\mathfrak{H}}.$$

Here  $b_n, c_n \in \mathbb{R}$  and  $|b_n| \leq \beta, c_n \geq -\mu_*$ . Hence, by taking a subsequence if necessary, we can suppose that  $b_n$  tends to a limit  $b_\infty$  such that  $b_\infty \in \mathbb{R}, |b_\infty| \leq \beta$ . Put  $c_\infty := -\lambda_\infty^2 - i\lambda_\infty b_\infty$ . Then we see  $c_n \rightarrow c_\infty$ . Hence  $c_\infty \in \mathbb{R}$  and  $c_\infty \geq -\mu_*$ , and  $\lambda_\infty$  turns out to enjoy the quadratic equation

$$\lambda_\infty^2 + ib_\infty \lambda_\infty + c_\infty = 0.$$

Consequently,

$$\lambda_\infty = i \left[ -\frac{b_\infty}{2} \pm \sqrt{\frac{b_\infty^2}{4} + c_\infty} \right].$$

If  $\frac{b_\infty^2}{4} + c_\infty \geq 0$ , then  $\lambda_\infty \in i\mathbb{R}$ . If  $\frac{b_\infty^2}{4} + c_\infty \leq 0$ , then  $|\lambda_\infty|^2 = -c_\infty \leq \mu_*$  and  $|\Im[\lambda_\infty]| = \left| -\frac{b_\infty}{2} \right| \leq \frac{\beta}{2}$ . Hence  $\lambda_\infty \in S$ .

Let us consider  $\lambda_0 \in \Sigma(\mathfrak{L})$ . We claim  $\lambda_0 \in S$ . Suppose  $\lambda_0 \notin S$ . By the symmetricity, we suppose  $\Re[\lambda_0] > 0$ . Then there would exist a curve  $\Gamma : t \in [0, 1] \mapsto \lambda(t) \in \mathbb{C} \setminus S$  such that  $\lambda(0) = \lambda_0$  and  $\lambda(1) \in \left] \frac{\beta}{2} + \sqrt{\frac{\beta^2}{4} + \mu_*}, +\infty \right[$ . Recall  $\lambda(1) \in \mathbf{P}(\mathfrak{L})$  by Proposition 6. The time

$$\bar{t} = \sup \left\{ t \in [0, 1] \mid \lambda(t) \in \Sigma(\mathfrak{L}) \right\}$$

would enjoy  $\bar{t} \in [0, 1[, \lambda(\bar{t}) \in \partial\Sigma(\mathfrak{L}), \lambda(\bar{t}) \notin S$ , a contradiction to  $\partial\Sigma(\mathfrak{L}) \subset S$ . Therefore we can claim  $\lambda_0 \in S$ .  $\square$

We want to clarify the structure of  $\Sigma(\mathfrak{L})$  more concretely. In [6, p.405] J. Dyson and B. F. Schutz say that the following assumption about  $\Sigma(\mathfrak{L})$  is very reasonable and provable:

**Assumption 3** Let  $S = S_{(i)} \cup S_{(ii)} \cup S_{(iii)}$ , where

$$\begin{aligned} S_{(i)} &= \{ \lambda \in i\mathbb{R} \mid |\Im[\lambda]| > \frac{\beta}{2} \} \\ S_{(ii)} &= \{ \lambda \in S \mid \lambda \notin i\mathbb{R} \}, \\ S_{(iii)} &= \{ \lambda \in i\mathbb{R} \mid |\Im[\lambda]| \leq \frac{\beta}{2} \}. \end{aligned}$$

Then

- (i)  $\Sigma(\mathfrak{L}) \cap S_{(i)} \subset \Sigma_p(\mathfrak{L})$ ,
- (ii)  $\Sigma(\mathfrak{L}) \cap S_{(ii)} \subset \Sigma_p(\mathfrak{L})$ , and  $\partial(\Sigma_p(\mathfrak{L}) \cap S_{(ii)}) \cap (i\mathbb{R}) = \emptyset$ ,
- (iii)  $(\Sigma(\mathfrak{L}) \cap S_{(iii)}) \setminus \Sigma_p(\mathfrak{L}) \neq \emptyset$ .

However it seems that this assumption has not yet been justified mathematically. Note that the Assumption 3 implies that, when  $\Omega = 0$ , it holds that  $\Sigma(\mathfrak{L}) = \Sigma_p(\mathfrak{L})$  and that, when  $\Omega \neq 0$ , it holds that  $\Sigma(\mathfrak{L}) \setminus \Sigma_p(\mathfrak{L}) \neq \emptyset$ .

## 6 Stability problem

Let us discuss on the stability problem.

D. Lynden-Bell and J. P. Ostriker, [15, p.301, line 18], say:

Equation (36) shows that the system is stable if  $c$  is positive for each eigen  $\boldsymbol{\xi}$  [ read  $\mathbf{u}^0$  ]. This assured if  $\mathbf{C}$  [ read  $\mathbf{L}$  ] is positive definite. Thus:

A sufficient condition for stability is that  $\mathbf{C}$  [read  $\mathbf{L}$  ] is positive definite. This is *the* condition for secular stability.

We should careful to understand the meaning of the saying ‘the system is stable’ and ‘ $\mathbf{L}$  is positive definite’. As for ‘stability’ C. Hunter [9] says:

A general system is said to be ordinarily or dynamically unstable if the amplitude of some mode grows exponentially in time, but ordinarily stable if every mode is oscillatory in time. An ordinarily stable system can be said to be secularly unstable if small additional dissipative forces can cause some perturbation to grow. Otherwise, the system is said to be secularly stable.

So ‘(ordinary) stability’ means the condition:

**(ST.1)** For any eigenvalue  $\lambda \in \Sigma_p(\mathfrak{L})$  of (5.1) and its associated solution  $\mathbf{u}(t, \mathbf{x}) = e^{\lambda t} \mathbf{u}^0(\mathbf{x})$ ,  $\mathbf{u}^0 \in \mathcal{D}(\mathbf{L})$ ,  $\mathbf{u}^0 \neq \mathbf{0}$ , it holds

$$\|U(t, \cdot)\|_{\mathfrak{H} \times \mathfrak{H}} \leq C \quad \text{for } \forall t \in [0, +\infty[, \quad (6.1)$$

Here and hereafter

$$U = \begin{bmatrix} \mathbf{u} \\ \dot{\mathbf{u}} \end{bmatrix}, \quad \dot{\mathbf{u}} = \frac{\partial \mathbf{u}}{\partial t}. \quad (6.2)$$

Since

$$\|\mathbf{u}(t, \cdot)\|_{\mathfrak{H}} = e^{\Re[\lambda]t} \|\mathbf{u}^0\|_{\mathfrak{H}}, \quad \|\dot{\mathbf{u}}(t, \cdot)\|_{\mathfrak{H}} = |\lambda| e^{\Re[\lambda]t} \|\mathbf{u}^0\|_{\mathfrak{H}},$$

the following condition is necessary and sufficient for the stability in the sense of **(ST.1)**:

**(\*)** For any eigenvalue  $\lambda \in \Sigma_p(\mathfrak{L})$  it holds that  $\Re[\lambda] = 0$ , that is,  $\Sigma_p(\mathfrak{L}) \subset i\mathbb{R}$ .

On the other hand, any  $\lambda \in \Sigma_p(\mathfrak{L})$  satisfies the quadratic equation

$$\lambda^2 + ib\lambda + c = 0 \tag{6.3}$$

where the coefficients  $b, c$  are given by

$$b = -i(B\mathbf{u}^0|\mathbf{u}^0)_{\mathfrak{H}}, \quad c = (L\mathbf{u}^0|\mathbf{u}^0)_{\mathfrak{H}}, \tag{6.4}$$

$\mathbf{u}^0$  being an associated eigen vector such that  $\|\mathbf{u}^0\|_{\mathfrak{H}} = 1$ , and  $b, c$  being real numbers, therefore

$$\lambda = i \left[ \frac{b}{2} \pm \sqrt{\frac{b^2}{4} + c} \right]. \tag{6.5}$$

Hence

$$\Re[\lambda] = 0 \quad \Leftrightarrow \quad \frac{b^2}{4} + c \geq 0 \quad \Leftarrow c \geq 0. \tag{6.6}$$

Therefore, considering the condition

**(PD.1)**: It holds

$$(L\mathbf{u}|\mathbf{u})_{\mathfrak{H}} \geq 0 \quad \forall \mathbf{u} \in D(L), \tag{6.7}$$

we can claim

**Proposition 8** If  $L$  is ‘positive definite’ in the sense of **(PD.1)**, then the background  $(\rho, S, \mathbf{v}) = (\rho_b, S_b, \mathbf{v}_b)$  is ‘stable’ in the sense of **(ST.1)**.

Thus the saying of [15] can be interpreted as this Proposition.

Note that **(PD.1)** implies

$$\inf_{\mathbf{u} \in D(L)} \frac{(L\mathbf{u}|\mathbf{u})_{\mathfrak{H}}}{\|\mathbf{u}\|_{\mathfrak{H}}^2} \geq 0,$$

so that

$$\mu_* = \left( \inf_{\mathbf{u} \in D(L)} \frac{(L\mathbf{u}|\mathbf{u})_{\mathfrak{H}}}{\|\mathbf{u}\|_{\mathfrak{H}}^2} \right) \vee 0 = 0,$$

and  $\Sigma(\mathfrak{L}) \subset i\mathbb{R}$  by Theorem 5.

The concept of ‘stability’ in the sense of **(ST.1)** is concerned with only ‘every modes’, say, waves associated with eigenvalues. We may consider all solutions  $\mathbf{u}$  of (5.1) and the wider stability concept:

**(ST.2)** : For all solution  $\mathbf{u} \in C^2([0, +\infty[, \mathfrak{H}) \cap C^1([0, +\infty[; \mathfrak{G}_0) \cap C([0, +\infty[; \mathcal{D}(\mathbf{L}))$  of (5.1) it holds

$$\|U(t, \cdot)\|_{\mathfrak{H} \times \mathfrak{H}} \leq C \quad \text{for } \forall t \in [0, +\infty[. \quad (6.8)$$

There is a gap between **(ST.1)** and **(ST.2)**. Of course **(ST.2)**  $\Rightarrow$  **(ST.1)**, but the inverse is not obvious, since we do not have an answer to the question:

**Question 1** Is the set of all eigenvectors of (5.1) dense in  $\mathfrak{H}$ ?

or

**Question 2** Is  $\Sigma_p(\mathfrak{L}) = \Sigma(\mathfrak{L})$ ?

Let us note that, if  $\Omega = \omega_b = 0$  and the background is spherically symmetric, then it holds that  $\Sigma(\mathbf{L}^G) = \Sigma_p(\mathbf{L}^G)$ , where  $\mathbf{L}^G$  is the Friedrichs extension of  $\mathcal{L} \upharpoonright C_0^\infty$  in the Hilbert space  $\mathfrak{G} = \left\{ \mathbf{u} \in \mathfrak{H} \mid \operatorname{div}(\rho \mathbf{u}) \in L^2\left(\frac{\Gamma P}{\rho^2} d\mathbf{x}\right) \right\}$ . See [11]. But, even in this non-rotating case, we do not know whether  $\Sigma(\mathbf{L}) = \Sigma_p(\mathbf{L})$  or not,  $\mathbf{L}$  being the Friedrichs extension in  $\mathfrak{H}$ . Moreover, when  $\Omega \neq 0$ ,  $B$  is not a bounded linear operator in  $\mathfrak{G}$  so that the application of Hille-Yosida theory for the basic existence proof in Section 4 does not work if we take  $\mathfrak{G}$  as the basic space instead of  $\mathfrak{H}$ .

Also let us note that, if  $\Omega = 0$ ,  $\mathbf{B} = \mathbf{0}$ , when (5.1) reduces

$$\frac{\partial^2 \mathbf{u}}{\partial t^2} + \mathbf{L} \mathbf{u} = 0, \quad (6.9)$$

**(PD.1)** implies that

$$\mathbf{u} = \int_0^{+\infty} \cos(\sqrt{\sigma} t) dE(\sigma) \mathbf{a} + \int_0^{+\infty} \sin(\sqrt{\sigma} t) dE(\sigma) \mathbf{b} \quad (6.10)$$

solves (6.9) with  $\mathbf{u}^0 = \mathbf{a}$ ,  $\mathbf{v}^0 = \sqrt{\mathbf{L}} \mathbf{b}$ , provided that  $\mathbf{a}, \mathbf{b} \in \mathcal{D}(\mathbf{L})$ . Here  $(E(\sigma))_{\sigma \in \mathbb{R}}$  is the spectral decomposition of the self-adjoint operator  $\mathbf{L}$ , which enjoys  $E(-0) = 0$  thanks to **(PD.1)**. In this case we have

$$\|U\|_{\mathfrak{H} \times \mathfrak{H}} \leq C \left( \|\mathbf{a}\|_{\mathcal{D}(\sqrt{\mathbf{L}})} + \|\mathbf{b}\|_{\mathcal{D}(\sqrt{\mathbf{L}})} \right).$$

Here

$$\sqrt{\mathbf{L}} = \int_0^{+\infty} \sqrt{\sigma} dE(\sigma)$$

so that  $(\sqrt{\mathbf{L}})^2 = \mathbf{L}$  and

$$\|\mathbf{u}\|_{\mathbf{D}(\sqrt{\mathbf{L}})} = \left[ \|\mathbf{u}\|_{\mathfrak{H}}^2 + \|\sqrt{\mathbf{L}}\mathbf{u}\|_{\mathfrak{H}}^2 \right]^{\frac{1}{2}}.$$

• Now let us introduce more general concepts of stability. Namely, let  $\mathfrak{N}$  be a semi-norm on  $\mathbf{D}(\mathbf{L}) \times \mathfrak{G}_0$ , and consider the condition:

**(ST.3):** For any solution  $\mathbf{u}$  of (5.1) it holds

$$\mathfrak{N}(U(t, \cdot)) \leq C \quad \text{for } \forall t \in [0, +\infty[. \quad (6.11)$$

Here  $C$  is a bound depending on  $U^0 = U(0, \cdot) = (\mathbf{u}^0, \mathbf{v}^0)^\top$ .

Recall

$$\mathfrak{N}_0(U(t, \cdot)) \leq e^{\Lambda t} \mathfrak{N}_0(U^0)$$

for

$$\mathfrak{N}_0(U) := \|U\|_{\mathfrak{E}} = \left[ \|\mathbf{u}\|_{\mathfrak{G}}^2 + \|\dot{\mathbf{u}}\|_{\mathfrak{H}}^2 \right]^{\frac{1}{2}} \quad \text{for } U = \begin{bmatrix} \mathbf{u} \\ \dot{\mathbf{u}} \end{bmatrix}$$

Here  $\Lambda = 1 + \mu_* + \beta \geq 1$  so this does not give the stability **(ST.3)**. We are keeping in mind the following situation.

Suppose a seminorm  $\mathfrak{n}$  on  $\mathbf{D}(\mathbf{L})$  satisfy

**(PD.2):** There is a positive number  $\delta$  such that

$$(\mathbf{L}\mathbf{u}|\mathbf{u}) \geq \delta \mathfrak{n}(\mathbf{u})^2 \quad \forall \mathbf{u} \in \mathbf{D}(\mathbf{L}). \quad (6.12)$$

Put

$$\mathfrak{N}(U) := \left[ \delta \mathfrak{n}(\mathbf{u})^2 + \|\dot{\mathbf{u}}\|_{\mathfrak{H}}^2 \right]^{\frac{1}{2}}. \quad (6.13)$$

Then **(ST.3)** holds with

$$C = \left[ \|\mathbf{v}^0\|_{\mathfrak{H}}^2 + (\mathbf{L}\mathbf{u}^0|\mathbf{u}^0)_{\mathfrak{H}} \right]^{\frac{1}{2}}.$$

In fact, multiplying (5.1) by  $\dot{\mathbf{u}}$  by the  $\mathfrak{H}$ -inner product, taking the real part, using  $\Re[(B\dot{\mathbf{u}}|\dot{\mathbf{u}})_{\mathfrak{H}}] = 0$ , we have

$$\frac{d}{dt} \left[ \|\dot{\mathbf{u}}\|_{\mathfrak{H}}^2 + (\mathbf{L}\mathbf{u}|\mathbf{u})_{\mathfrak{H}} \right] = 0.$$

Of course **(PD.2)** implies **(PD.1)** but is much stronger concept of ‘positive definiteness’.

Now we are going to find a seminorm  $\mathfrak{n}$  which enjoys **(PD.2)**, under the following situation:

**Situation 1** The equation of state is (2.2):  $P = \rho^\gamma \exp(S/C_V)$ . 1)  $\Omega = \omega_b = 0$  and the background  $(\rho, S, \mathbf{0})$  is spherically symmetric, 2) and is isentropic, that is,  $S = S_0$ ,  $S_0$  being a constant, and 3)  $\frac{4}{3} < \gamma < 2$ .

Then  $\Upsilon = \Upsilon_O \theta\left(\frac{r}{a}; \frac{1}{\gamma-1}\right)$  and  $\mathfrak{R} = \{\mathbf{x} | r = \|\mathbf{x}\| < R\}$  with  $R := a\xi_1\left(\frac{1}{\gamma-1}\right)$ , where  $\theta(\cdot, \frac{1}{\gamma-1})$  is the Lane-Emden function and  $\xi_1(\frac{1}{\gamma-1})$  is its zero. It holds

$$P = A\rho^\gamma, \quad \Upsilon = \frac{A\gamma}{\gamma-1}\rho^{\gamma-1}, \quad \frac{\gamma P}{\rho^2} = \frac{d\Upsilon}{d\rho} = A\rho^{-(2-\gamma)},$$

and

$$(\mathbf{L}\mathbf{u}|\mathbf{u})_{\mathfrak{H}} = \int_{\mathfrak{R}} \left( \frac{d\Upsilon}{d\rho} |g|^2 - 4\pi G\mathcal{K}[g]g^* \right) d\mathbf{x}, \quad (6.14)$$

with  $g = \operatorname{div}(\rho\mathbf{u})$ . ( Recall Remark 4. )

Denote

$$\mathfrak{g}_L := \left\{ g \in L^2\left(\frac{d\Upsilon}{d\rho} d\mathbf{x}\right) \middle| \exists \mathbf{u} \in \mathbf{D}(\mathbf{L}) : g = \operatorname{div}(\rho\mathbf{u}) \right\}, \quad (6.15)$$

and put

$$\Psi = -\mathcal{K}[g] \quad (6.16)$$

for  $g \in \mathfrak{g}_L$ . Then, for  $g \in \mathfrak{g}_L$ ,  $g \in C(\mathfrak{R})$ ,  $\Psi \in C(\mathbb{R}^3)$ , and the expansion with respect to spherical harmonics

$$g(\mathbf{x}) = \sum_{0 \leq \ell, |m| \leq \ell} g_{\ell m}(r) Y_{\ell m}(\vartheta, \phi), \quad (6.17a)$$

$$\Psi(\mathbf{x}) = \sum_{0 \leq \ell, |m| \leq \ell} \Psi_{\ell m}(r) Y_{\ell m}(\vartheta, \phi) \quad (6.17b)$$

can be used. Here

$$\mathbf{x} = (x^1, x^2, x^3) = (r \sin \vartheta \cos \phi, r \sin \vartheta \sin \phi, r \cos \vartheta)$$

and

$$\begin{aligned} Y_{\ell m}(\vartheta, \phi) &= \sqrt{\frac{2\ell+1}{4\pi} \frac{(\ell-m)!}{(\ell+m)!}} P_{\ell}^m(\cos \vartheta) e^{im\phi}, \\ Y_{\ell, -m} &= (-1)^m Y_{\ell m}^* \quad (0 \leq m \leq \ell), \end{aligned} \quad (6.18)$$

$$g_{\ell m}(r) = \int_0^{2\pi} \int_0^{\pi} g(\mathbf{x}) Y_{\ell m}(\vartheta, \phi)^* \sin \vartheta d\vartheta d\phi, \quad (6.19)$$

$$\Psi_{\ell m}(r) = \int_0^{2\pi} \int_0^{\pi} \Psi(\mathbf{x}) Y_{\ell m}(\vartheta, \phi)^* \sin \vartheta d\vartheta d\phi. \quad (6.20)$$

See [11, Lemma 5]. Note that

$$\Delta^{(\ell)}\Psi_{\ell m} = g_{\ell m}, \quad \Psi_{\ell m} = -\mathcal{H}_\ell[g_{\ell m}], \quad (6.21)$$

where

$$\Delta^{(\ell)}w = \frac{1}{r^2} \frac{d}{dr} r^2 \frac{dw}{dr} - \frac{\ell(\ell+1)}{r^2} w, \quad (6.22)$$

$$\mathcal{H}_\ell[\check{g}](r) = \frac{1}{2\ell+1} \left[ \int_0^r \check{g}(s) \left(\frac{r}{s}\right)^{-\ell-1} s ds + \int_r^{+\infty} \check{g}(s) \left(\frac{r}{s}\right)^\ell s ds \right], \quad (6.23)$$

and

$$\|\nabla\Psi\|^2 = \sum_{\ell m} \|\nabla^{(\ell)}\Psi_{\ell m}\|^2, \quad (6.24)$$

where

$$\|\nabla^{(\ell)}w\|^2 = \left| \frac{dw}{dr} \right|^2 + \frac{\ell(\ell+1)}{r^2} |w|^2. \quad (6.25)$$

We are observing

$$(\mathbf{L}\mathbf{u}|\mathbf{u})_{\mathfrak{H}} = 4\pi \sum_{\ell, m} Q_{\ell m}, \quad (6.26)$$

where

$$Q_{\ell m} = \int_0^{+\infty} \left( \frac{d\Upsilon}{d\rho} |\Delta^{(\ell)}\Psi_{\ell m}|^2 - 4\pi \mathbb{G} \|\nabla^{(\ell)}\Psi_{\ell m}\|^2 \right) r^2 dr. \quad (6.27)$$

First we consider  $\ell = 0$ . Then we have

$$Q_{00} = \int_0^R (\mathcal{L}^{\text{ss}}y)y^* \rho r^4 dr, \quad (6.28)$$

where

$$y = -\frac{1}{r\rho} \frac{d}{dr} \Psi_{00}, \quad g_{00} = \frac{1}{r^2} \frac{d}{dr} (r^3 \rho y), \quad (6.29)$$

$$\mathcal{L}^{\text{ss}}y = -\frac{1}{\rho r^4} \frac{d}{dr} \left( \gamma r^4 \rho \frac{dy}{dr} \right) - (3\gamma - 4) \frac{1}{r} \frac{d\Upsilon}{dr} y, \quad (6.30)$$

$$\int_0^R (\mathcal{L}^{\text{ss}}y)y^* \rho r^4 dr = \int_0^R \left( \gamma \left| \frac{dy}{dr} \right|^2 - (3\gamma - 4) \frac{1}{r} \frac{d\Upsilon}{dr} |y|^2 \right) \rho r^4 dr. \quad (6.31)$$

Since we are supposing  $\frac{4}{3} < \gamma$ , we have

$$\delta_* := \inf_{0 < r < R} - (3\gamma - 4) \frac{1}{r} \frac{d\Upsilon}{dr} > 0, \quad (6.32)$$

and

$$Q_{00} \geq \delta_* \int_0^R |y|^2 \rho r^4 dr = \delta_* \int_0^R \frac{1}{\rho} \left| \frac{d\Psi_{00}}{dr} \right|^2 r^2 dr. \quad (6.33)$$

But

$$\frac{d\Psi_{00}}{dr} = \frac{1}{r^2} \int_0^r g_{00}(s) s^2 ds = 0 \quad \text{for } r > R,$$

since  $g_{00} = 0$  on  $]R, +\infty[$  and

$$\int_0^R g_{00}(r) r^2 dr = \frac{1}{\sqrt{4\pi}} \int_{\mathfrak{R}} g(\mathbf{x}) d\mathbf{x} = 0$$

for  $g \in \mathfrak{g}_L$ . On the other hand,

$$\frac{1}{\rho} \left| \frac{d\Psi_{00}}{dr} \right|^2 \in L^\infty(0, R),$$

since

$$\frac{d\Psi_{00}}{dr} = -\frac{1}{r^2} \int_r^R g_{00}(s) s^2 ds \quad \text{for } 0 < r < R$$

enjoys

$$\begin{aligned} \left| \frac{d\Psi_{00}}{dr} \right| &\leq \frac{1}{r^2} \sqrt{\int_r^R \frac{d\rho}{dY} s^2 ds} \sqrt{\int_0^R \frac{dY}{d\rho} |g_{00}(s)|^2 s^2 ds} \\ &\leq C(R-r)^{\frac{1}{2(\gamma-1)}} \quad \text{for } \frac{R}{2} \leq r < R, \end{aligned}$$

for  $\frac{d\rho}{dY} = O((R-r)^{\frac{2-\gamma}{\gamma-1}})$ .

Therefore, under the convention that  $\frac{1}{\rho} \left| \frac{d\Psi_{00}}{dr} \right|^2$  means 0 for  $r > R$ , we can write

$$Q_{00} \geq \delta_* \int_0^{+\infty} \frac{1}{\rho} \left| \frac{d\Psi_{00}}{dr} \right|^2 r^2 dr. \quad (6.34)$$

In other words, if we adopt the decomposition

$$u^r(\mathbf{x}) = \sum_{\ell m} u_{\ell m}^r(r) Y_{\ell m}(\vartheta, \phi), \quad (6.35)$$

$$u_{\ell m}^r(r) = \int_0^{2\pi} \int_0^\pi u^r(\mathbf{x}) Y_{\ell m}(\vartheta, \phi)^* \sin \vartheta d\vartheta d\phi, \quad (6.36)$$

while

$$\mathbf{u}(\mathbf{x}) = u^r(\mathbf{x}) \mathbf{e}_r + u^\vartheta(\mathbf{x}) \mathbf{e}_\vartheta + u^\phi(\mathbf{x}) \mathbf{e}_\phi, \quad (6.37)$$

where

$$\mathbf{e}_r = \frac{\partial}{\partial r}, \quad \mathbf{e}_\vartheta = \frac{1}{r} \frac{\partial}{\partial \vartheta}, \quad \mathbf{e}_\phi = \frac{1}{r \sin \vartheta} \frac{\partial}{\partial \phi}. \quad (6.38)$$



we see

$$\frac{1}{r^2} \frac{d}{dr} (r^2 \rho u_{00}^r) = \frac{1}{r^2} \frac{d}{dr} (r^2 g_{00}), \quad (6.39)$$

for

$$g = \operatorname{div}(\rho \mathbf{u}) = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \rho u^r) + \frac{1}{r \sin \vartheta} \frac{\partial}{\partial \vartheta} (\sin \vartheta u^\vartheta) + \frac{1}{r \sin \vartheta} \frac{\partial u^\phi}{\partial \phi}.$$

Hence we have

$$\rho(r) u_{00}^r(r) = \int_0^r g_{00}(s) s^2 ds = -\frac{d}{dr} \Psi_{00}(r) \quad \text{for } 0 \leq r < R,$$

therefore we can write

$$\int_0^{+\infty} \frac{1}{\rho} \left| \frac{d\Psi_{00}}{dr} \right|^2 r^2 dr = \int_0^{+\infty} |u_{00}^r|^2 \rho(r) r^2 dr.$$

Thus

$$Q_{00} \geq \delta_* \int_0^{+\infty} |u_{00}^r|^2 \rho(r) r^2 dr. \quad (6.40)$$

Moreover there is a positive number  $\mu_0$  such that

$$\int_0^R \frac{1}{\rho} \left| \frac{d\Psi_{00}}{dr} \right|^2 r^2 dr \geq \mu_0 \int_0^R |\Psi_{00}(r)|^2 r^2 dr. \quad (6.41)$$

Proof. Looking at

$$\mu := \inf_{\psi \in \mathcal{F}} \frac{\int_0^R \frac{1}{\rho} \left| \frac{d\Psi_{00}}{dr} \right|^2 r^2 dr}{\int_0^R |\Psi_{00}(r)|^2 r^2 dr}, \quad (6.42)$$

where

$$\mathcal{F} = \left\{ \psi \in C^1[0, R] \mid \psi(R) = 0, |\psi(r)| \leq \|\Psi_{00}\|_{L^\infty} \right\},$$

we prove  $\mu > 0$ .

Suppose  $\mu = 0$  for reductio ad absurdum. Then there is a sequence  $(\psi_n)_{n=1,2,\dots}$ , in  $\mathcal{F}$  such that  $\int_0^R |\psi_n|^2 r^2 dr = 1$  and

$$M_n = \int_0^R \frac{1}{\rho} \left| \frac{d\psi_n}{dr} \right|^2 r^2 dr \searrow 0 \quad \text{as } n \rightarrow \infty.$$

For  $0 < r \leq r' \leq R$  it hold

$$\begin{aligned}
|\psi_n(r)| &= \left| - \int_r^R D\psi_n(s) ds \right| \\
&\leq \sqrt{\int_r^R \frac{\rho(s)}{s^2} ds} \sqrt{\int_r^R \frac{1}{\rho(s)} |D\psi_n(s)|^2 s^2 ds} \\
&\leq \sqrt{\rho_O} \sqrt{\frac{R-r}{Rr}} M_n \leq \frac{\sqrt{\rho_O}}{r} M_1,
\end{aligned}$$

and

$$|\psi_n(r') - \psi_n(r)| \leq \sqrt{\rho_O} \sqrt{\frac{r'-r}{r'r}} M_n \leq \sqrt{\rho_O} \sqrt{\frac{r'-r}{r'r}} M_1.$$

Therefore by the Ascoli-Arzelà theorem we can suppose  $\psi_n$  tends to a limit  $\psi_\infty$  uniformly on any compact subinterval of  $]0, R]$ , by taking a subsequence. Since  $M_n \rightarrow 0$ , we have  $\psi_\infty = 0$ . For any  $0 < r_* \ll 1$  we have

$$\int_0^{r_*} |\psi_n(r)|^2 r^2 dr \leq \|\Psi_{00}\|_{L^\infty}^2 \frac{r_*^3}{3}.$$

Therefore

$$\begin{aligned}
1 &= \int_0^R |\psi_n(r)|^2 r^2 dr \\
&\leq \|\Psi_{00}\|_{L^\infty}^2 \frac{r_*^3}{3} + \int_{r_*}^R |\psi_n(r)|^2 r^2 dr \\
&\rightarrow \|\Psi_{00}\|_{L^\infty}^2 \frac{r_*^3}{3} + \int_{r_*}^R |\psi_\infty(r)|^2 r^2 dr \\
&= \|\Psi_{00}\|_{L^\infty}^2 \frac{r_*^3}{3}.
\end{aligned}$$

Taking  $r_*$  so small that  $\|\Psi_{00}\|_{L^\infty}^2 \frac{r_*^3}{3} < 1$ , we see a contradiction.  $\square$

Next we consider  $\ell \geq 1$ . We claim

$$Q_{\ell m} \geq 4\pi G \int_0^{+\infty} \left( \|\nabla^{(\ell)} \Psi_{\ell m}\|^2 - 4\pi G \frac{d\rho}{d\mathcal{R}} |\Psi_{\ell m}|^2 \right) r^2 dr. \quad (6.43)$$

For  $0 < \epsilon \ll 1$ , we put  $\left(\frac{d\mathcal{R}}{d\rho}\right)_\epsilon = \gamma A(\rho + \epsilon)^{\gamma-2}$ . Then, as  $\epsilon \searrow +0$ ,  $\left(\frac{d\mathcal{R}}{d\rho}\right)_\epsilon \nearrow \frac{d\mathcal{R}}{d\rho}$  on  $0 < r < R$ , and

$$\begin{aligned}
&\int_0^{+\infty} \left( \left(\frac{d\mathcal{R}}{d\rho}\right)_\epsilon |g_{\ell m}|^2 - 4\pi G \|\nabla^{(\ell)} \Psi_{\ell m}\|^2 \right) r^2 dr \\
&\nearrow Q_{\ell m} = \int_0^{+\infty} \left( \left(\frac{d\mathcal{R}}{d\rho}\right) |g_{\ell m}|^2 - 4\pi G \|\nabla^{(\ell)} \Psi_{\ell m}\|^2 \right) r^2 dr.
\end{aligned}$$

But we see

$$\begin{aligned}
& \int_0^{+\infty} \left( \left( \frac{d\Upsilon}{d\rho} \right)_\epsilon |g_{\ell m}|^2 - 4\pi\mathbf{G} \|\nabla^{(\ell)} \Psi_{\ell m}\|^2 \right) r^2 dr \\
&= \int_0^{+\infty} \left| \sqrt{\left( \frac{d\Upsilon}{d\rho} \right)_\epsilon} \Delta^{(\ell)} \Psi_{\ell m} + 4\pi\mathbf{G} \sqrt{\left( \left( \frac{d\Upsilon}{d\rho} \right)_\epsilon \right)^{-1}} \Psi_{\ell m} \right|^2 r^2 dr \\
&\quad - 4\pi\mathbf{G} \int_0^{+\infty} 2\Re[(\Delta^{(\ell)} \Psi_{\ell m}) \Psi_{\ell m}^*] + \|\nabla^{(\ell)} \Psi_{\ell m}\|^2 + 4\pi\mathbf{G} \left( \left( \frac{d\Upsilon}{d\rho} \right)_\epsilon \right)^{-1} |\Psi_{\ell m}|^2 r^2 dr \\
&\geq 4\pi\mathbf{G} \int_0^{+\infty} \left( \|\nabla^{(\ell)} \Psi_{\ell m}\|^2 - 4\pi\mathbf{G} \left( \left( \frac{d\Upsilon}{d\rho} \right)_\epsilon \right)^{-1} |\Psi_{\ell m}|^2 \right) r^2 dr,
\end{aligned}$$

for

$$\int_0^{+\infty} (\Delta^{(\ell)} \Psi_{\ell m}) \Psi_{\ell m}^* r^2 dr = - \int_0^{+\infty} \|\nabla^{(\ell)} \Psi_{\ell m}\|^2 r^2 dr,$$

since  $r^2 \frac{d\Psi_{\ell m}}{dr} \Psi_{\ell m}^* = O\left(\frac{1}{r}\right)$  as  $r \rightarrow +\infty$ . Since  $\left( \left( \frac{d\Upsilon}{d\rho} \right)_\epsilon \right)^{-1} = \frac{(\rho + \epsilon)^{2-\gamma}}{\gamma\mathbf{A}} \searrow \frac{d\rho}{d\Upsilon} = \frac{\rho^{2-\gamma}}{\gamma\mathbf{A}}$  as  $\epsilon \searrow 0$ , we have (6.43).

Let us show that

$$\mathcal{Q}_1[w] := \int_0^{+\infty} \left( \|\nabla^{(1)} w\|^2 - 4\pi\mathbf{G} \frac{d\rho}{d\Upsilon} |w|^2 \right) r^2 dr \geq 0 \quad (6.44)$$

for  $w = \Psi_{\ell m}$ .

Fixing  $N \geq R$ , put

$$\mathcal{Q}_1^N[w] := \int_0^N \left( \|\nabla^{(1)} w\|^2 - 4\pi\mathbf{G} \frac{d\rho}{d\Upsilon} |w|^2 \right) r^2 dr. \quad (6.45)$$

This is the quadratic form associated with the operator

$$\mathcal{P}w = \left[ -\Delta^{(1)} - 4\pi\mathbf{G} \frac{d\rho}{d\Upsilon} \right] w. \quad (6.46)$$

We consider  $\mathcal{P}$  in  $\mathfrak{X}_0^N = L^2([0, N]; r^2 dr)$ . We have the Friedrichs extension  $\mathbf{P}^N$  of  $\mathcal{P} \upharpoonright C_0^\infty(]0, N[)$ , keeping in mind that

$$4\pi\mathbf{G} \frac{d\rho}{d\Upsilon} \leq \frac{4\pi\mathbf{G}}{\gamma\mathbf{A}} \rho_O^{2-\gamma}.$$

We see that  $\mathbf{P}^N$  is of the Sturm-Liouville type. the boundary condition at  $r = N$  is the Dirichlet boundary condition  $w|_{r=N} = 0$ .

Let  $\mu_1$  be the least eigenvalue of  $\mathbf{P}^N$ . We are going to show  $\mu_1 \geq 0$ . Let  $\phi_1$  be an eigenfunction. We can suppose  $\phi_1(r) > 0$  for  $0 < r < N$ . Consider  $\Upsilon' := \frac{d\Upsilon}{dr}$ , which satisfies  $\mathcal{P}\Upsilon' = 0$  on  $[0, +\infty[$ . Here we consider

$$\Upsilon(r) = -K \left( \frac{1}{R} - \frac{1}{r} \right), \quad \Upsilon'(r) = -\frac{K}{r^2} \quad \text{for } R < r$$

with  $K = -\left(r^2 \frac{d\mathcal{Y}}{dr}\right)_{r=R-0} (> 0)$ . we look at

$$\begin{aligned} 0 &= (\mathcal{P}\mathcal{Y}'|\phi_1)_{\mathfrak{X}_0^N} = (\mathcal{Y}'|\mathcal{P}\phi_1)_{\mathfrak{X}_0^N} + r^2 \mathcal{Y}' \frac{d\phi_1}{dr} \Big|_{r=N-0} \\ &= \mu_1(\mathcal{Y}'|\phi_1) + r^2 \mathcal{Y}' \frac{d\phi_1}{dr} \Big|_{r=N-0} \end{aligned}$$

Since  $\mathcal{Y}' < 0$  on  $]0, N]$ , we have  $(\mathcal{Y}'|\phi_1)_{\mathfrak{X}_0^N} < 0$ . But  $r^2 \mathcal{Y}' \frac{d\phi_1}{dr} \Big|_{r=N-0} \geq 0$ , since  $\frac{d\phi_1}{dr} \Big|_{r=N-0} \leq 0$ . Consequently  $\mu_1 \geq 0$  and

$$\mathcal{Q}_1^N[w] \geq 0 \quad \text{for } \forall w \in \mathbf{D}(\mathbf{P}^N).$$

Since we cannot say  $w = \Psi_{\ell m} \in \mathbf{D}(\mathbf{P}^N)$ , for maybe  $\Psi_{\ell m}(N) \neq 0$ , we decompose it as

$$w = \Psi_{\ell m} = W^N + \Pi^N, \quad (6.47)$$

where

$$\Pi^N = -C_{\ell m} N^{-\ell-1} \left(\frac{r}{N}\right)^\ell, \quad C_{\ell m} = \frac{1}{2\ell+1} \int_0^R g_{\ell m}(s) s^{\ell+2} ds. \quad (6.48)$$

Then

$$\Delta^{(\ell)} W^N = \Delta^{(\ell)} w, \quad W^N(N) = 0. \quad (6.49)$$

Note

$$\mathcal{P}W^N = -g_{\ell m} - \left(\frac{\ell(\ell+1)-2}{r^2} + 4\pi\mathbf{G} \frac{d\rho}{d\mathcal{Y}}\right)(w - \Pi^N) \in \mathfrak{X}_0^N,$$

since  $g_{\ell m} \in C([0, R], w = \Psi_{\ell m} = O(r^2)$  for  $\ell \geq 2$ ,  $\Pi^N = O(r^\ell)$ . Therefore  $W^N \in \mathbf{D}(\mathbf{P}^N)$  and

$$\mathcal{Q}_1^N[W^N] \geq 0.$$

Since

$$\begin{aligned} \|\nabla^{(1)} \Pi^N\|^2 &\leq 3C_{\ell m} N^{-2\ell-4}, \\ 4\pi\mathbf{G} \frac{d\rho}{d\mathcal{Y}} |\Pi^N|^2 &\leq 4\pi\mathbf{G} \frac{\rho_O^{2-\gamma}}{\gamma\mathbf{A}} C_{\ell m}^2 N^{-2\ell-2} \end{aligned}$$

on  $[0, N]$ , we see

$$\begin{aligned} \mathcal{Q}_1^N[w] &\geq \mathcal{Q}_1^S[W^N] + O(N^{-2\ell+1}) \\ &\geq O(N^{-2\ell+1}), \end{aligned}$$

Taking the limit as  $N \rightarrow +\infty$ , we get (6.44) for  $w = \Psi_{\ell m}$ .

Now

$$Q_{\ell m} \geq 4\pi\mathbf{G} \mathcal{Q}_1[\Psi_{\ell m}] \geq 0, \quad (6.50)$$

and, for  $\ell \geq 2$ , we have

$$\begin{aligned}
Q_{\ell m} &\geq 4\pi G \int_0^{+\infty} \left( \|\nabla^{(\ell)} \Psi_{\ell m}\|^2 - 4\pi G \frac{d\rho}{dY} |\Psi_{\ell m}|^2 \right) r^2 dr \\
&= 4\pi G \left( \mathcal{Q}_1[\Psi_{\ell m}] + \int_0^{+\infty} (\ell(\ell+1) - 2) |\Psi_{\ell m}|^2 dr \right) \\
&\geq 4\pi G \int_0^{+\infty} (\ell(\ell+1) - 2) |\Psi_{\ell m}|^2 dr.
\end{aligned} \tag{6.51}$$

Note that

$$(\ell(\ell+1) - 2) \int_0^{+\infty} |\Psi_{\ell m}(r)|^2 dr \geq \frac{4}{R^2} \int_0^R |\Psi_{\ell m}(r)|^2 r^2 dr \tag{6.52}$$

for  $\ell \geq 2$ .

Consequently

$$\begin{aligned}
(\mathbf{L}\mathbf{u}|\mathbf{u})_{\mathfrak{H}} &= 4\pi \sum_{0 \leq \ell, |m| \leq \ell} Q_{\ell m} \\
&\geq \mathfrak{d} \left( \int_0^{+\infty} \frac{1}{\rho} \left| \frac{d\Psi_{00}}{dr} \right|^2 r^2 dr + \sum_{2 \leq \ell, |m| \leq \ell} (\ell(\ell+1) - 2) \int_0^{+\infty} |\Psi_{\ell m}|^2 dr \right),
\end{aligned} \tag{6.53}$$

with  $\delta := 4\pi(\delta_* \wedge G)$ .

Therefore

**Theorem 6** *Under Situation 1 , (PD.2) holds for  $\mathbf{n} = \mathbf{n}_1, \mathbf{n}_{10}, \mathbf{n}_1 \geq \mathbf{n}_{10}$ , defined by*

$$\begin{aligned}
\mathbf{n}_1(\mathbf{u}) &= \left[ \int_0^{+\infty} \frac{1}{\rho} \left| \frac{d\Psi_{00}}{dr} \right|^2 r^2 dr + \sum_{2 \leq \ell, |m| \leq \ell} (\ell(\ell+1) - 2) \int_0^{+\infty} |\Psi_{\ell m}|^2 dr \right]^{\frac{1}{2}} \\
&= \left[ \int_0^{+\infty} |u_{00}^r|^2 \rho(r) r^2 dr + \sum_{2 \leq \ell, |m| \leq \ell} (\ell(\ell+1) - 2) \int_0^{+\infty} |\Psi_{\ell m}|^2 dr \right]^{\frac{1}{2}},
\end{aligned} \tag{6.54a}$$

$$\mathbf{n}_{10}(\mathbf{u}) = \sqrt{\mu_1} \left[ \int_0^R |\Psi_{00}(r)|^2 r^2 dr + \sum_{2 \leq \ell, |m| \leq \ell} \int_0^R |\Psi_{\ell m}(r)|^2 r^2 dr \right]^{\frac{1}{2}} \tag{6.54b}$$

$$\text{with } \mu_1 = \mu_0 \wedge \frac{4}{R^2}.$$

Here

$$\Psi_{\ell m}(r) = - \int_0^{2\pi} \int_0^\pi \mathcal{K}[g](\mathbf{x}) Y_{\ell m}(\vartheta, \phi)^* \sin \vartheta d\vartheta d\phi, \tag{6.55}$$

$$u_{00}^r(r) = \sqrt{4\pi} \int_0^{2\pi} \int_0^\pi \left( \mathbf{u}(\mathbf{x}) \middle| \frac{\mathbf{x}}{\|\mathbf{x}\|} \right) \sin \vartheta d\vartheta d\phi, \tag{6.56}$$

in which

$$\mathbf{x} = (x^1, x^2, x^3) = (r \sin \vartheta \cos \phi, r \sin \vartheta \sin \phi, r \cos \vartheta)$$

and  $g = \operatorname{div}(\rho \mathbf{u})$ .

Note that the seminorm  $\mathbf{n} = \mathbf{n}_1, \mathbf{n}_{10}$  and the associated  $\mathfrak{N}$  are not norm, that is,  $\mathfrak{N}(U) = 0$  does not imply  $U = \mathbf{0}$ , say, does not imply  $\mathbf{u} = 0$ . But  $\mathfrak{N}(U)$  can control the amplitude of  $\Psi = \mathcal{K}[\operatorname{div}(\rho \mathbf{u})]$ , except for the  $\ell = 1$  components, and  $\dot{\mathbf{u}}$ . Here  $-4\pi \mathbf{G}\Psi = \mathfrak{D}\Phi = \Phi_{\rho_b + \mathfrak{D}\rho} - \Phi_b, \mathfrak{D}\rho = -\operatorname{div}(\rho \mathbf{u})$  and  $\mathbf{v} = \dot{\mathbf{u}}$  are the quantities essential for the perturbation, and the control of the magnitude of  $\mathbf{u} = \boldsymbol{\varphi}(t, \mathbf{x}) - \mathbf{x} + \mathbf{u}^0$  itself is not essential in the discussion of the stability of the background. (Recall  $\boldsymbol{\varphi}_b(t, \bar{\mathbf{x}}) = \bar{\mathbf{x}}$  for  $\omega_b = 0$ .) In fact  $\mathbf{n}_1(\mathbf{u}) = 0$  implies that  $\Psi_{\ell m} = 0, g_{\ell m} = 0$  on  $[0, +\infty[$  for  $\ell \neq 1$ , and  $\mathbf{n}_{10}(\mathbf{u}) = 0$  implies that  $\Psi_{\ell m} = 0, g_{\ell m} = 0$  on  $[0, +R]$  for  $\ell \neq 1$ , and

$$g(\mathbf{x}) = \sqrt{\frac{3}{4\pi}} g_{1,0}(r) \cos \vartheta - \sqrt{\frac{3}{8\pi}} g_{1,1}(r) \sin \vartheta e^{i\phi} + \sqrt{\frac{3}{8\pi}} g_{1,-1}(r) \sin \vartheta e^{-i\phi}.$$

**Remark 9** *It seems impossible to control the magnitude of  $\Psi_{1m}, m = 0, \pm 1$  by the following reason. The differential operator  $\mathcal{P} \upharpoonright C_0^\infty([0, +\infty[)$  admits the Friedrichs extension  $\mathbf{P}$  in  $\mathfrak{X} = L^2([0, +\infty[, r^2 dr)$ , and  $\mathcal{Q}_1$  is the quadratic form associated with  $\mathbf{P}$ . But  $\mathbf{P}$  is not of the Sturm-Liouville type. In fact, let  $\phi \in \mathcal{D}(\mathbf{P})$  and  $\mathcal{P}\phi = \lambda\phi$  with  $\lambda > 0$ . Since  $-\Delta^{(1)}\phi = 0$  on  $]R, +\infty[$ , there are constants  $C_\pm$  such that*

$$\phi = C_+ \phi_+ + C_- \phi_- \quad \text{on } ]R, +\infty[,$$

where

$$\phi_\pm(r) = \sqrt{r} J_{\pm \frac{3}{2}}(\sqrt{\lambda} r), \quad J_{\pm \frac{3}{2}} \quad \text{being the Bessel function.}$$

Since

$$J_{\pm \frac{3}{2}}(r) \sim \sqrt{\frac{2}{\pi r}} \cos\left(r \pm \frac{3\pi}{4} - \frac{\pi}{4}\right) \quad \text{as } r \rightarrow +\infty,$$

we see  $\phi_\pm \notin L^2([R, +\infty[, r^2 dr)$ , and  $\phi \in L^2([R, +\infty[, r^2 dr)$  requires  $C_\pm = 0$ ,  $\phi = 0$  on  $]R, +\infty[$ . By the uniqueness of solutions of ODE, it follows that  $\phi = 0$  on  $[0, +\infty[$ . In other words, any positive real number cannot be an eigenvalue of  $\mathbf{P}$ , so  $\mathbf{P}$  is not of the Sturm-Liouville type.

However, if we restrict ourselves to axially and equatorially symmetric perturbations, functions of  $(r, |\zeta|)$ , we have  $\Psi_{\ell m} = 0$  for odd  $\ell$  a priori, since  $\int_0^{2\pi} e^{im\phi} d\phi = 0$  for  $m \neq 0$  and  $P_\ell^0(-\zeta) = -P_\ell^0(\zeta)$  for odd  $\ell$ , we need not control the magnitudes of  $\Psi_{1m}, m = 0, \pm 1$ . In this situation  $\mathbf{n}(\mathbf{u}) = 0$  implies  $\Psi = 0, g = 0$ , or,  $\mathfrak{D}\Phi = 0, \mathfrak{D}\rho = 0$ , or, more precisely, we have

$$\mathbf{n}_{10}(\mathbf{u}) \geq \sqrt{\frac{\mu_1}{4\pi}} \|\Psi\|_{L^2(\mathfrak{X})} = \sqrt{\frac{\mu_1}{4\pi}} \|\mathcal{K}[\mathfrak{D}\rho]\|_{L^2(\mathfrak{X})}, \quad (6.57)$$

since

$$\|\Psi\|_{L^2(\mathfrak{X})}^2 = 4\pi \sum_{\substack{\ell: \text{even} \\ |m| \leq \ell}} \|\Psi_{\ell m}\|_{L^2([0, R], r^2 dr)}^2$$

in this case.

• Now we are going to try to derive **(PD.2)** with another seminorm

$$\mathbf{n}(\mathbf{u}) := \|\operatorname{div}(\rho\mathbf{u})\|_{L^2\left(\frac{\Gamma P}{\rho^2}d\mathbf{x}\right)} = \left[\int_{\mathfrak{R}} \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho\mathbf{u})|^2 d\mathbf{x}\right]^{\frac{1}{2}} \quad (6.58)$$

for

$$\begin{aligned} (\mathcal{L}\mathbf{u}|\mathbf{u})_{\mathfrak{H}} &= \int_{\mathfrak{R}} \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho\mathbf{u})|^2 d\mathbf{x} + 2\Re\left[\int_{\mathfrak{R}} \frac{\Gamma P \mathcal{A}}{\rho} (\mathbf{u}|\mathbf{n}) \operatorname{div}(\rho\mathbf{u})^* d\mathbf{x}\right] \\ &\quad - \int_{\mathfrak{R}} \frac{\Gamma P \mathcal{A}}{\rho} \|\nabla \rho\|^2 |(\mathbf{u}|\mathbf{n})|^2 d\mathbf{x} + \\ &\quad - 4\pi G \int_{\mathfrak{R}} \mathcal{K}[\operatorname{div}(\rho\mathbf{u})] \operatorname{div}(\rho\mathbf{u})^* d\mathbf{x}. \end{aligned} \quad (6.59)$$

First we claim:

**Proposition 9** *Let  $\varepsilon$  be a positive number. If*

$$-\varepsilon \frac{\|\nabla \rho\|^2}{\rho} \leq \mathcal{A} \leq 0 \quad \text{on } \mathfrak{R}, \quad (6.60)$$

then

$$(\mathcal{L}\mathbf{u}|\mathbf{u}) \geq (1 - \varepsilon) \int_{\mathfrak{R}} \frac{\Gamma P}{\rho^2} |g|^2 d\mathbf{x} - 4\pi G \int_{\mathfrak{R}} \mathcal{K}[g] g^* d\mathbf{x}. \quad (6.61)$$

Here  $g = \operatorname{div}(\rho\mathbf{u})$ .

Proof. We see

$$\begin{aligned} \left| 2\Re\left[\int_{\mathfrak{R}} \frac{\Gamma P \mathcal{A}}{\rho} (\mathbf{u}|\mathbf{n}) g^* d\mathbf{x}\right] \right| &\leq \varepsilon \int_{\mathfrak{R}} \int_{\mathfrak{R}} \frac{\Gamma P}{\rho^2} |g|^2 d\mathbf{x} + \frac{1}{\varepsilon} \int_{\mathfrak{R}} \Gamma P \mathcal{A}^2 |(\mathbf{u}|\mathbf{n})|^2 d\mathbf{x} \\ &\leq \varepsilon \int_{\mathfrak{R}} \frac{\Gamma P |\mathcal{A}|}{\rho^2} |g|^2 d\mathbf{x} + \int_{\mathfrak{R}} \frac{\Gamma P \mathcal{A}}{\rho} \|\nabla \rho\|^2 |(\mathbf{u}|\mathbf{n})|^2 d\mathbf{x} \end{aligned}$$

by (6.60), where  $g = \operatorname{div}(\rho\mathbf{u})$ , since  $-|\mathcal{A}| - \mathcal{A} = 0$  for  $\mathcal{A} \leq 0$ , we have (6.61).  $\square$

Therefore, if we adopt the Cowling approximation, which neglect the perturbed self-gravitation term  $-4\pi G \int_{\mathfrak{R}} \mathcal{K}[g] g^* d\mathbf{x}$ , we have **(PD.2)** with  $0 < \varepsilon < 1$ . Namely we consider the following

**Situation 2** *The equation of state is (2.2):  $P = \rho^\gamma \exp(S/C_V)$ . Instead of  $\mathcal{L}$ , we consider  $\mathcal{L}_0$ :*

$$\begin{aligned} \mathcal{L}_0 \mathbf{u} &= \nabla \left( -\frac{\gamma P}{\rho^2} \operatorname{div}(\rho\mathbf{u}) + \frac{\gamma P}{\rho} (\mathbf{u}|\mathbf{a}) \right) + \\ &\quad + \frac{\gamma P}{\rho} \left( -\mathbf{a} \operatorname{div}(\rho\mathbf{u}) + (\mathbf{u}|\mathbf{a}) \nabla \rho \right). \end{aligned} \quad (6.62)$$

We can claim

**Theorem 7** *Under the Situation 2 suppose (6.60) with  $0 < \varepsilon < 1$ . Put  $\delta := 1 - \varepsilon$ . Then **(PD.2)** holds for  $\mathbf{L}$  replaced by  $\mathbf{L}_0$  with  $\delta$  and  $\mathbf{n} = \mathbf{n}_2$  defined by*

$$\mathbf{n}_2(\mathbf{u}) = \left[ \int_{\mathfrak{R}} \frac{\gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} \right]^{\frac{1}{2}}. \quad (6.63)$$

So let us look at the perturbed self-gravitation term in order to justify the Cowling approximation in this context.

Put

$$K^\spadesuit := \sup_{g \in \mathfrak{g}_L} \frac{4\pi \mathbf{G} \int_{\mathfrak{R}} \mathcal{K}[g] g^* d\mathbf{x}}{\int_{\mathfrak{R}} \frac{\gamma P}{\rho^2} |g|^2 d\mathbf{x}}, \quad (6.64)$$

where

$$\mathfrak{g}_L := \left\{ g \in L^2 \left( \frac{\gamma P}{\rho^2} d\mathbf{x}, \mathfrak{R} \right) \mid g = \operatorname{div}(\rho \mathbf{u}) \quad \text{for} \quad \exists \mathbf{u} \in \mathbf{D}(\mathbf{L}) \right\}. \quad (6.65)$$

We can claim  $K^\spadesuit < \infty$  by the following

**Proposition 10** *There exists a constant  $C$  such that*

$$0 \leq 4\pi \mathbf{G} \int_{\mathfrak{R}} \mathcal{K}[g] g^* d\mathbf{x} \leq C \int_{\mathfrak{R}} \frac{\gamma P}{\rho^2} |g|^2 d\mathbf{x}. \quad (6.66)$$

Proof. We have

$$\begin{aligned} |\mathcal{K}[g](\mathbf{x})| &\leq \frac{1}{4\pi} \sqrt{\int_{\mathfrak{R}} \frac{d\mathbf{x}'}{\|\mathbf{x} - \mathbf{x}'\|}} \sqrt{\int_{\mathfrak{R}} |g|^2 d\mathbf{x}} \\ &\leq \sqrt{\frac{R}{2\pi}} \|g\|_{L^2(\mathfrak{R})}, \end{aligned}$$

where  $R := \sup\{\|\mathbf{x}\| \mid \mathbf{x} \in \mathfrak{R}\}$ . Therefore

$$\begin{aligned} \int_{\mathfrak{R}} \mathcal{K}[g] g^* d\mathbf{x} &\leq \sqrt{\frac{R}{2\pi}} \|g\|_{L^2} \int_{\mathfrak{R}} |g| d\mathbf{x} \\ &\leq \sqrt{\frac{2}{3}} R^2 \|g\|_{L^2}^2 \end{aligned}$$

On the other hand,

$$\nu_* := \inf_{\mathfrak{R}} \frac{\gamma P}{\rho^2} > 0, \quad (6.67)$$

since

$$\frac{1}{C} d^{-\frac{2-\gamma}{\gamma-1}} \leq \frac{\gamma P}{\rho^2} \leq C d^{-\frac{2-\gamma}{\gamma-1}} \quad \text{on} \quad \mathfrak{R},$$



where  $\mathbf{d} := \text{dist}(\mathbf{x}, \partial\mathfrak{R})$  and  $1 < \gamma < 2$ . Then

$$\int_{\mathfrak{R}} |g|^2 d\mathbf{x} \leq \frac{1}{\nu_*} \int_{\mathfrak{R}} \frac{\gamma P}{\rho^2} |g|^2 d\mathbf{x},$$

therefore (6.66) holds with  $C = \sqrt{\frac{2}{3}} \frac{R^2}{\nu_*}$ .  $\square$

Consequently we can claim

Suppose  $K^\spadesuit < 1$  and let  $0 < \varepsilon < 1 - K^\spadesuit$  so that  $\delta := 1 - \varepsilon - K^\spadesuit > 0$ . If (6.60) holds with  $\varepsilon$ , then **(PD.2)** holds with  $\delta$  and

$$\mathbf{n}(\mathbf{u}) := \|\text{div}(\rho \mathbf{u})\|_{L^2\left(\frac{\gamma P}{\rho^2} d\mathbf{x}\right)} = \left[ \int_{\mathfrak{R}} \frac{\gamma P}{\rho^2} |\text{div}(\rho \mathbf{u})|^2 d\mathbf{x} \right]^{\frac{1}{2}}.$$

Thus we have the following question, which is still open:

**Question 3** When the background realizes the condition  $K^\spadesuit < 1$  ?

Let us note the nondimensionalization of the estimate of  $K^\spadesuit$ . We have

$$\frac{\gamma P}{\rho^2} = 4\pi \mathbf{Ga}^2 (\gamma - 1) \Theta^{-\frac{2-\gamma}{\gamma-1}} (1 + \omega_1)(1 + \omega_2),$$

where

$$\begin{aligned} \Theta &= \Theta\left(\frac{\|\mathbf{x}\|}{\mathbf{a}}, \frac{x^3}{\|\mathbf{x}\|}; \frac{1}{\gamma-1}, \mathbf{b}\right), \\ \omega_1 &= O(\mathcal{R}_O), \quad \omega_2 = O\left(\frac{\gamma-1}{\gamma \mathbf{A}} \mathcal{R}_O\right), \end{aligned}$$

and

$$\int_{\mathfrak{R}} \frac{\gamma P}{\rho^2} |g|^2 d\mathbf{x} = 4\pi \mathbf{Ga}^5 \int_{\mathfrak{R}_1} (\gamma - 1) \Theta^{-\frac{2-\gamma}{\gamma-1}} (1 + \omega_1)(1 + \omega_2) |\hat{g}(\underline{\mathbf{x}})|^2 d\underline{\mathbf{x}},$$

where

$$\begin{aligned} \mathfrak{R}_1 &= \left\{ \underline{\mathbf{x}} \mid \|\underline{\mathbf{x}}\| < U_1\left(\frac{x^3}{\|\underline{\mathbf{x}}\|}\right) \right\}, \\ \Theta &= \Theta\left(\|\underline{\mathbf{x}}\|, \frac{x^3}{\|\underline{\mathbf{x}}\|}; \frac{1}{\gamma-1}, \mathbf{b}\right), \quad \hat{g}(\underline{\mathbf{x}}) = g(\mathbf{a}\underline{\mathbf{x}}). \end{aligned}$$

On the other hand, we see

$$4\pi \mathbf{G} \int_{\mathfrak{R}} \mathcal{K}[g] g^* d\mathbf{x} = 4\pi \mathbf{Ga}^5 \int_{\mathfrak{R}_1} \mathcal{K}[\hat{g}](\underline{\mathbf{x}}) \hat{g}^*(\underline{\mathbf{x}}) d\underline{\mathbf{x}}.$$

Therefore we have the reduction

$$K^\spadesuit = \sup_{\hat{g}} \frac{\int_{\mathfrak{R}_1} \mathcal{K}[\hat{g}] \hat{g}^* d\mathbf{x}}{\int_{\mathfrak{R}_1} (\gamma - 1) \Theta^{-\frac{2-\gamma}{\gamma-1}} (1 + \omega_1)(1 + \omega_2) |\hat{g}|^2 d\mathbf{x}}. \quad (6.68)$$

If the background is isentropic, we have  $\omega_1 = \omega_2 = 0$ . If  $\Omega = 0$  and the background is isentropic and spherically symmetric, then

$$K^\spadesuit = \sup_{\hat{g}} \frac{\int_{\mathfrak{R}_1} \mathcal{K}[\hat{g}] \hat{g}^* d\mathbf{x}}{\int_{\mathfrak{R}_1} (\gamma - 1) \theta^{-\frac{2-\gamma}{\gamma-1}} |\hat{g}|^2 d\mathbf{x}}.$$

where  $\theta = \theta(\|\mathbf{x}\|, \frac{1}{\gamma-1})$  is the Lane-Emden function.

## 7 The equation LASO used by astrophysicists

### 7.1

The equation of linearized adiabatic stellar oscillations traditionally used by astrophysicists is little bit different from the equation LASO (1.11) with (1.12), (1.13a), (1.13b), (1.13c), (1.13d) derived in Section 3. In this section we derive the equation used by astrophysicists after V. R. Lebovitz 1961 [13], D. Lynden-Bell and J. P. Ostriker 1967 [15] and J. L. Friedman and B. F. Schutz 1978 [7].

We consider a pair of solutions  $(\rho, S, \mathbf{v})$  and  $(\rho_b, S_b, \mathbf{v}_b)$  of the Euler-Poisson equations described in a rotating frame (1.1a), (1.1b), (1.1c), (1.6) with the equation of state (1.8). Here  $(\rho_b, S_b, \mathbf{v}_b)$  is a compactly supported axially symmetric stationary solution with

$$\mathbf{v}_b(t, \mathbf{x}) = \omega_b(\varpi, z) \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \times \mathbf{x} = \omega_b(\varpi, z) \begin{bmatrix} -x^2 \\ x^1 \\ 0 \end{bmatrix}.$$

We define the Lagrangian displacement  $\boldsymbol{\xi}$  in the Lebovitz sense ([13]) by

$$\boldsymbol{\xi}(t, \mathbf{x}) = \boldsymbol{\varphi}(t|0, \boldsymbol{\varphi}(0|t, \mathbf{x}|\mathbf{v}_b)|\mathbf{v}) - \mathbf{x}. \quad (7.1)$$

Here, for  $\mathbf{V} = \mathbf{v}, \mathbf{v}_b$ ,  $\boldsymbol{\varphi}(t|s, \mathbf{y}|\mathbf{V})$  stands for the solution  $\mathbf{x} = \boldsymbol{\varphi}(t|s, \mathbf{y}|\mathbf{V})$  of the problem

$$\frac{d\mathbf{x}}{dt} = \mathbf{V}(t, \mathbf{x}), \quad \mathbf{x}|_{t=s} = \mathbf{y}. \quad (7.2)$$

Under the assumption

$$\rho^0 := \rho(0, \cdot) = \rho_b, \quad (7.3)$$

$$S^0 := S(0, \cdot) = S_b, \quad (7.4)$$

the linearized equation for  $\xi$  turns out to be

$$\frac{\partial^2 \xi}{\partial t^2} + \mathcal{B}_\diamond \frac{\partial \xi}{\partial t} + \mathcal{L}_\diamond \xi = 0 \quad \text{in } [0, +\infty[ \times \mathfrak{R}_b, \quad (7.5)$$

where

$$\mathcal{B}_\diamond \dot{\xi} = 2\Omega \times \dot{\xi} + \mathcal{B}_{\diamond 1} \dot{\xi} \quad (7.6)$$

with

$$\mathcal{B}_{\diamond 1} \dot{\xi} = 2(D\dot{\xi})v_b = 2\omega_b \frac{\partial \dot{\xi}}{\partial \phi}, \quad (7.7)$$

and

$$\mathcal{L}_\diamond = \mathcal{L}_{\diamond 2} + \mathcal{L}_{\diamond 1} + \mathcal{L}_{\diamond 0} + \mathcal{L}, \quad (7.8)$$

with

$$\mathcal{L}_{\diamond 2} \xi = D((D\xi)v_b)v_b = \omega_b^2 \frac{\partial^2 \xi}{\partial \phi^2}, \quad (7.9a)$$

$$\mathcal{L}_{\diamond 1} \xi = 2\Omega \times ((D\xi)v_b) = 2\Omega\omega_b \frac{\partial}{\partial \phi} \begin{bmatrix} -\xi^2 \\ \xi^1 \\ 0 \end{bmatrix}, \quad (7.9b)$$

$$\mathcal{L}_{\diamond 0} \xi = -\left(D((Dv_b)v_b) + D(2\Omega \times v_b)\right)\xi = W_b \xi, \quad (7.9c)$$

with

$$W_b := -D((Dv_b)v_b) - D(2\Omega \times v_b) \quad (7.9d)$$

$$= (2\Omega + \omega_b) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} + 2(\Omega + \omega_b) \begin{bmatrix} \frac{1}{\varpi} \frac{d\omega_b}{d\varpi} (x^1)^2 & \frac{1}{\varpi} \frac{d\omega_b}{d\varpi} x^1 x^2 & \frac{\partial \omega_b}{\partial z} x^1 \\ \frac{1}{\varpi} \frac{d\omega_b}{d\varpi} x^1 x^2 & \frac{1}{\varpi} \frac{d\omega_b}{d\varpi} (x^2)^2 & \frac{\partial \omega_b}{\partial z} x^2 \\ 0 & 0 & 0 \end{bmatrix},$$

$$\mathcal{L}\xi = \frac{1}{\rho_b} \nabla [\delta P] - \frac{P_b}{\rho_b^2} [\delta \rho] - 4\pi G \nabla \mathcal{K} [[\delta \rho]], \quad (7.9e)$$

with

$$[\delta \rho] = -\text{div}(\rho_b \xi), \quad [\delta S] = -(\xi | \nabla S_b),$$

$$[\delta P] = \left(\frac{\Gamma P}{\rho}\right)_b [\delta \rho] + \left(\frac{\partial P}{\partial S}\right)_b [\delta S] = \left(\frac{\Gamma P}{\rho}\right)_b [\delta \rho] + (\Gamma P)_b (\xi | \mathfrak{a}_b). \quad (7.9f)$$

Here and hereafter  $\nabla, D$  stand for  $\nabla_{\mathbf{x}}, D_{\mathbf{x}}$ , namely,

$$\nabla f = \begin{bmatrix} \frac{\partial f}{\partial x^1} \\ \frac{\partial f}{\partial x^2} \\ \frac{\partial f}{\partial x^3} \end{bmatrix},$$

$$DQ = \begin{bmatrix} \frac{\partial Q^1}{\partial x^1} & \frac{\partial Q^1}{\partial x^2} & \frac{\partial Q^1}{\partial x^3} \\ \frac{\partial Q^2}{\partial x^1} & \frac{\partial Q^2}{\partial x^2} & \frac{\partial Q^2}{\partial x^3} \\ \frac{\partial Q^3}{\partial x^1} & \frac{\partial Q^3}{\partial x^2} & \frac{\partial Q^3}{\partial x^3} \end{bmatrix},$$

and  $\frac{\partial \mathbf{u}}{\partial \phi}$  means  $-x^2 \frac{\partial \mathbf{u}}{\partial x^1} + x^1 \frac{\partial \mathbf{u}}{\partial x^2}$ .

The equation (7.5) with (7.6) (7.7) (7.8) with (7.9a) - (7.9f) is the equation of LASO used by astrophysicists after [13] [7] [15]. Hereafter we call this equation the ‘equation of LASO(♦)’.

Note that, if  $\mathbf{u}^0 = \mathbf{0}$ , we have the relation

$$\boldsymbol{\xi}(t, \mathbf{x}) = \mathbf{u}(t, \varphi(0|t, \mathbf{x}||\mathbf{v}_b)), \quad (7.10)$$

since the Lagrange displacement  $\mathbf{u}$  in our equation (1.11) was defined by (3.27), which reads

$$\mathbf{u}(t, \bar{\mathbf{x}}) = \varphi(t|0, \bar{\mathbf{x}}||\mathbf{v}) - \varphi(t|0, \bar{\mathbf{x}}||\mathbf{v}_b) + \mathbf{u}^0(\bar{\mathbf{x}}).$$

Therefore  $\boldsymbol{\xi} = \mathbf{u}$  when  $\mathbf{v}_b = \mathbf{0}$  so that  $\varphi(t|s, \mathbf{x}||\mathbf{v}_b) = \mathbf{x}$ , but  $\boldsymbol{\xi} \neq \mathbf{u}$  in general.

## 7.2 Derivation of the equation of LASO(♦) (7.5)

Let us explain how to derive the equation (7.5) following [13], [7] and [15].

We shall use the identities

$$D_{\mathbf{y}}\varphi(t|s, \mathbf{y}||\mathbf{V}) = \left( D_{\mathbf{y}}\varphi(s|t, \varphi(t|s, \mathbf{y}||\mathbf{V})||\mathbf{V}) \right)^{-1}, \quad (7.11)$$

$$\partial_s \varphi(t|s, \mathbf{y}||\mathbf{V}) = -D_{\mathbf{y}}\varphi(t|s, \mathbf{y}||\mathbf{V})\mathbf{V}(s, \mathbf{y}). \quad (7.12)$$

In fact, (7.11) can be deduced by differentiating the identity

$$\mathbf{y} = \varphi(s|t, \varphi(t|s, \mathbf{y}||\mathbf{V})||\mathbf{V}) \quad (7.13)$$

by  $\mathbf{y}$ , and (7.12) can be deduced by differentiating (7.13) by  $s$  and using (7.11).

Note that, for  $\mathbf{x} = \mathbf{x}(t, \bar{\mathbf{x}}) = \boldsymbol{\varphi}(t|0, \bar{\mathbf{x}}|\mathbf{V})$ , it holds that

$$D_{\bar{\mathbf{x}}}\mathbf{x}(t, \bar{\mathbf{x}}) = \exp \left[ \int_0^t D_{\mathbf{x}}\mathbf{V}(t', \boldsymbol{\varphi}(t'|0, \bar{\mathbf{x}}|\mathbf{V}))dt' \right], \quad (7.14)$$

$$\det D_{\bar{\mathbf{x}}}\mathbf{x}(t, \bar{\mathbf{x}}) = \exp \left[ \int_0^t \operatorname{div}_{\mathbf{x}}\mathbf{V}(t', \boldsymbol{\varphi}(t'|0, \bar{\mathbf{x}}|\mathbf{V}))dt' \right], \quad (7.15)$$

For a pair  $(Q, Q_b)$  of scalar fields, we define the scalar field  $\Delta Q$ , the Lagrangian perturbation, and  $\delta Q$ , the Eulerian perturbation, by

$$\Delta Q(t, \mathbf{x}) = Q(t, \mathbf{x} + \boldsymbol{\xi}(t, \mathbf{x})) - Q_b(t, \mathbf{x}), \quad (7.16)$$

$$\delta Q(t, \mathbf{x}) = Q(t, \mathbf{x}) - Q_b(t, \mathbf{x}). \quad (7.17)$$

For a pair  $(\mathbf{Q}, \mathbf{Q}_b)$  of vector fields, we define

$$\Delta \mathbf{Q}(t, \mathbf{x}) = \mathbf{Q}(t, \mathbf{x} + \boldsymbol{\xi}(t, \mathbf{x})) - \mathbf{Q}_b(t, \mathbf{x}), \quad (7.18)$$

$$\delta \mathbf{Q}(t, \mathbf{x}) = \mathbf{Q}(t, \mathbf{x}) - \mathbf{Q}_b(t, \mathbf{x}). \quad (7.19)$$

We suppose that  $\delta \rho, \delta S, \delta \mathbf{v} = \mathcal{O}(\varepsilon)$ .

Then we have  $\boldsymbol{\xi} = \mathcal{O}(\varepsilon)$ .

In fact, writing

$$\boldsymbol{\xi} = \underline{\mathbf{x}} - \mathbf{x}, \quad \underline{\mathbf{x}} = \boldsymbol{\varphi}(t|0, \bar{\mathbf{x}}|\mathbf{v}), \quad \mathbf{x} = \boldsymbol{\varphi}(t|0, \bar{\mathbf{x}}|\mathbf{v}_b),$$

we see that  $\boldsymbol{\psi}(t) = \boldsymbol{\xi}(t, \mathbf{x})$  satisfies

$$\frac{d\boldsymbol{\psi}}{dt} = \mathbf{v}(t, \mathbf{x} + \boldsymbol{\psi}) - \mathbf{v}(t, \mathbf{x}) + \mathbf{v}(t, \mathbf{x}) - \mathbf{v}_b(t, \mathbf{x}).$$

Hence

$$\left\| \frac{d\boldsymbol{\psi}}{dt} \right\| \leq C_0 + C_1 \|\boldsymbol{\psi}\|, \quad \boldsymbol{\psi}(0) = \mathbf{0},$$

where

$$C_0 = \|\delta \mathbf{v}\|_{L^\infty}, \quad C_1 = \|D_{\mathbf{x}}\mathbf{v}\|_{L^\infty}.$$

The Gronwall argument implies

$$\|\boldsymbol{\psi}(t)\| \leq C_0(e^{C_1 t} - 1).$$

Thus  $\boldsymbol{\xi} = \mathcal{O}(\varepsilon)$ . As for the derivatives  $\psi_j(t) = \frac{\partial}{\partial x^j} \boldsymbol{\xi}(t, \mathbf{x})$ , we see  $\psi_j = \mathcal{O}(\varepsilon)$  by the same argument, so we have  $\frac{\partial}{\partial x^j} \boldsymbol{\xi} = \mathcal{O}(\varepsilon)$ .

Consequently, when  $\delta Q = \mathcal{O}(\varepsilon)$  for scalar fields paired, we can claim

$$\Delta Q = \delta Q + (\boldsymbol{\xi} | \nabla_{\mathbf{x}} Q_b) + \mathcal{O}(\varepsilon^2). \quad (7.20)$$

and when  $\delta \mathbf{Q} = \mathcal{O}(\varepsilon)$  for vector fields, we have

$$\Delta \mathbf{Q} = \delta \mathbf{Q} + (D \mathbf{Q}_b) \boldsymbol{\xi} + \mathcal{O}(\varepsilon^2). \quad (7.21)$$

- Note that it holds that

$$\operatorname{div} \mathbf{v}_b = 0, \quad (\mathbf{v}_b | \nabla \rho_b) = 0, \quad (\mathbf{v}_b | \nabla S_b) = 0. \quad (7.22)$$

- Integration of the equation (1.1c) gives

$$S(t, \boldsymbol{\varphi}(t|0, \bar{\mathbf{x}} | \mathbf{v})) = S^0(\bar{\mathbf{x}}) := S(0, \bar{\mathbf{x}}). \quad (7.23)$$

Then

$$\Delta S(t, \mathbf{x}) = S^0(\bar{\mathbf{x}}) - S_b(\mathbf{x}).$$

Here  $\bar{\mathbf{x}} = \boldsymbol{\varphi}(0|t, \mathbf{x} | \mathbf{v}_b)$ .

We suppose

$$S^0 = S_b. \quad (7.24)$$

Since  $S_b$  is supposed to be axially symmetric, and the values of coordinates  $\varpi, z$  of  $\boldsymbol{\varphi}(t|s, \mathbf{y} | \mathbf{v}_b)$  are invariant in  $t$ , we see  $S_b(\bar{\mathbf{x}}) = S_b(\mathbf{x})$ . Therefore

$$\Delta S = 0, \quad (7.25)$$

provided (7.24). Thanks to (7.20), this implies

$$\delta S = -(\boldsymbol{\xi} | \nabla S_b) + \mathcal{O}(\varepsilon^2). \quad (7.26)$$

- Integration of the equation of continuity (1.1a) gives

$$\rho(t, \boldsymbol{\varphi}(t|0, \bar{\mathbf{x}} | \mathbf{v})) = \rho^0(\bar{\mathbf{x}}) \exp \left[ - \int_0^t (\operatorname{div}_{\mathbf{x}} \mathbf{v})(t', \boldsymbol{\varphi}(t'|0, \bar{\mathbf{x}} | \mathbf{v})) dt' \right], \quad (7.27)$$

where  $\rho^0 = \rho(0, \cdot)$ .

We denote

$$\mathfrak{R}(t) = \{\mathbf{x} \mid \rho(t, \mathbf{x}) > 0\}, \quad \mathfrak{R}_b = \{\mathbf{x} \mid \rho_b(\mathbf{x}) > 0\}. \quad (7.28)$$

Of course

$$\mathfrak{R}(t) = \{\mathbf{x} = \boldsymbol{\varphi}(t|0, \bar{\mathbf{x}} | \mathbf{v}) \mid \bar{\mathbf{x}} \in \mathfrak{R}(0)\},$$

and the equation (7.27) is valid for  $\forall(t, \bar{\mathbf{x}}) \in [0, T] \times \mathbb{R}^3$ .

We suppose

$$\rho^0 = \rho_b. \quad (7.29)$$

Then  $\mathfrak{R}(0) = \mathfrak{R}_b$  and

$$\mathfrak{R}(t) = \{\mathbf{x} = \boldsymbol{\varphi}(t|0, \bar{\mathbf{x}}|\mathbf{v}) \mid \bar{\mathbf{x}} \in \mathfrak{R}_b\},$$

that is,

$$\underline{\mathbf{x}} = \mathbf{x} + \boldsymbol{\xi}(t, \mathbf{x}) \in \mathfrak{R}(t) \iff \mathbf{x} \in \mathfrak{R}_b.$$

Here we note that  $\rho_b(\boldsymbol{\varphi}(0|t, \mathbf{x}|\mathbf{v}_b)) = \rho_b(\mathbf{x})$  by the axial symmetry.

We claim

$$\Delta\rho + \rho_b \operatorname{div} \boldsymbol{\xi} = \mathcal{O}(\varepsilon^2). \quad (7.30)$$

Thanks to (7.20), this implies

$$\delta\rho = -\operatorname{div}(\rho_b \boldsymbol{\xi}) + \mathcal{O}(\varepsilon^2) \quad (7.31)$$

Proof of (7.30). Proof is divided into several steps. 1) Let  $(Q, Q_b)$  be a pair of scalar fields such that  $\delta Q = \mathcal{O}(\varepsilon)$ . Let  $D$  be an arbitrary domain in  $\mathbb{R}^3$ . Put

$$\begin{aligned} \underline{D}(t) &= \{\underline{\mathbf{x}} = \mathbf{x} + \boldsymbol{\xi}(t, \mathbf{x}) \mid \mathbf{x} \in D\}, \\ I &= \int_{\underline{D}(t)} Q(t, \underline{\mathbf{x}}) d\underline{\mathbf{x}}, \quad I_b = \int_D Q_b(t, \mathbf{x}) d\mathbf{x}, \\ \delta I &= I - I_b. \end{aligned} \quad (7.32)$$

Then we have

$$\delta I = \int_D q(t, \mathbf{x}) d\mathbf{x}, \quad (7.33)$$

where

$$q(t, \mathbf{x}) = \Delta Q(t, \mathbf{x}) + Q(t, \underline{\mathbf{x}})(\det(I + D_{\mathbf{x}} \boldsymbol{\xi}(t, \mathbf{x})) - I). \quad (7.34)$$

Note that

$$q = \Delta Q + Q_b \operatorname{div} \boldsymbol{\xi} + \mathcal{O}(\varepsilon^2).$$

2) Let  $\mathbf{V} = \mathbf{v}$  or  $\mathbf{v}_b$ . Let  $Q$  be a scalar field, and  $E$  a domain in  $\mathbb{R}^3$ . Put

$$\begin{aligned} E(t) &= \{\mathbf{z} = \boldsymbol{\varphi}(t|0, \mathbf{y}) \mid \mathbf{y} \in E\}, \\ I(t) &= \int_{E(t)} Q(t, \mathbf{z}) d\mathbf{z}. \end{aligned} \quad (7.35)$$

Here we denote  $\boldsymbol{\varphi}(t|s, \mathbf{y}) = \boldsymbol{\varphi}(t|s, \mathbf{y}|\mathbf{V})$ . Then

$$I(t) = \int_E Q(t, \mathbf{z}) \det D_{\mathbf{y}} \boldsymbol{\varphi}(t|0, \mathbf{y}) d\mathbf{y}. \quad (7.36)$$

Note that

$$\det D_{\mathbf{y}}\varphi(t|0, \mathbf{y}) = \exp \left[ \int_0^t (\operatorname{div}_{\mathbf{x}} \mathbf{V})(t', \varphi(t'|0, \mathbf{y})) dt' \right] > 0$$

and

$$\frac{\partial}{\partial t} \det D_{\mathbf{y}}\varphi(t|0, \mathbf{y}) = (\operatorname{div}_{\mathbf{x}} \mathbf{V})(t, \mathbf{z}) \cdot \det D_{\mathbf{y}}\varphi(t|0, \mathbf{y}).$$

Differentiating (7.36) by  $t$ , we get

$$\frac{dI}{dt} = \int_E \left( \frac{\partial Q}{\partial t} + (\mathbf{V} | \nabla_{\mathbf{x}} Q) + Q \operatorname{div}_{\mathbf{x}} \mathbf{V} \right) (t, \mathbf{z}) \det D_{\mathbf{y}}\varphi d\mathbf{y} \quad (7.37)$$

Therefore, if

$$\frac{\partial Q}{\partial t} + (\mathbf{V} | \nabla_{\mathbf{x}} Q) + Q \operatorname{div}_{\mathbf{x}} \mathbf{V} = 0 \quad (7.38)$$

holds everywhere, then  $dI/dt = 0$  and

$$I(t) = \int_E Q(0, \mathbf{y}) d\mathbf{y}. \quad (7.39)$$

Note that (7.38) holds for  $Q = \rho$  or  $\rho_b$  respectively with  $\mathbf{V} = \mathbf{v}$  or  $\mathbf{v}_b$ , that is, (1.1a).

3) Let  $D$  be an arbitrary domain in  $\mathbb{R}^3$ , and  $(Q, Q_b)$  be a pair of scalar fields such that (7.38) holds everywhere for  $Q$  with  $\mathbf{V} = \mathbf{v}$  and for  $Q_b$  with  $\mathbf{v}_b$ .

Then, keeping in mind

$$\begin{aligned} \underline{\mathbf{x}} &= \mathbf{x} + \boldsymbol{\xi}(t, \mathbf{x}) = \varphi(t|0, \bar{\mathbf{x}}|\mathbf{v}), \\ \mathbf{x} &= \varphi(t|0, \bar{\mathbf{x}}|\mathbf{v}_b) \quad \Leftrightarrow \quad \bar{\mathbf{x}} = \varphi(0|t, \mathbf{x}|\mathbf{v}_b), \end{aligned}$$

we see

$$\mathbf{x} \in D \quad \Leftrightarrow \quad \bar{\mathbf{x}} \in \bar{D} \quad \Leftrightarrow \quad \underline{\mathbf{x}} \in D(t),$$

where

$$\begin{aligned} \bar{D} &= \{ \bar{\mathbf{x}} = \varphi(0|t, \mathbf{x}|\mathbf{v}_b) \mid \mathbf{x} \in D \}, \\ \underline{D}(t) &= \{ \underline{\mathbf{x}} = \varphi(t|0, \bar{\mathbf{x}}|\mathbf{v}) \mid \bar{\mathbf{x}} \in \bar{D} \}. \end{aligned}$$

We consider

$$\begin{aligned} I &= \int_{\underline{D}} Q(t, \underline{\mathbf{x}}) d\underline{\mathbf{x}}, \quad I_b = \int_D Q(t, \mathbf{x}) d\mathbf{x}, \\ \delta I &= I - I_b. \end{aligned}$$

Thanks to 2) we have

$$I = \int_{\bar{D}} Q(0, \bar{\mathbf{x}}) d\bar{\mathbf{x}}, \quad I_b = \int_{\bar{D}} Q_b(0, \bar{\mathbf{x}}) d\bar{\mathbf{x}}.$$



Therefore

$$\delta I = \int_{\bar{D}} (Q(0, \bar{\mathbf{x}}) - Q_b(0, \bar{\mathbf{x}})) d\bar{\mathbf{x}}.$$

On the other hand, thanks to 1), we have

$$\delta I = \int_D q(t, \mathbf{x}) d\mathbf{x}.$$

Hence

$$\int_{\bar{D}} (Q(0, \bar{\mathbf{x}}) - Q_b(0, \bar{\mathbf{x}})) d\bar{\mathbf{x}} = \int_D q(t, \mathbf{x}) d\mathbf{x}.$$

The change of variable  $\bar{D} \rightarrow D : \bar{\mathbf{x}} \mapsto \mathbf{x} = \varphi(t|0, \bar{\mathbf{x}}|\mathbf{v}_b)$ , for which we have  $\det D_{\bar{\mathbf{x}}}\mathbf{x} = 1$  since  $\operatorname{div} \mathbf{v}_b = 0$ , we get

$$\int_{\bar{D}} (Q(0, \bar{\mathbf{x}}) - Q_b(0, \bar{\mathbf{x}}) - q(t, \mathbf{x})) d\bar{\mathbf{x}} = 0.$$

Since  $D$ , therefore  $\bar{D}$  is arbitrary, we can claim

$$Q(0, \bar{\mathbf{x}}) - Q_b(0, \bar{\mathbf{x}}) = q(t, \mathbf{x}) \quad \forall \mathbf{x} \in \mathbb{R}^3.$$

4) Applying the above discussion to  $Q = \rho, Q_b = \rho_b$ , we can claim

$$\rho^0(\bar{\mathbf{x}}) - \rho_b(\bar{\mathbf{x}}) = \Delta \rho(t, \mathbf{x}) + \rho(t, \underline{\mathbf{x}}) \left( \operatorname{der}(I + D_{\mathbf{x}} \boldsymbol{\xi}(t, \mathbf{x})) - 1 \right). \quad (7.40)$$

Supposing  $\rho^0 = \rho_b$ , we can claim

$$\Delta \rho + \rho_b \operatorname{div} \boldsymbol{\xi} = \mathcal{O}(\varepsilon^2).$$

Proof is done.  $\square$

• As for the equation of motion (1.1b), we divide it by  $\rho$  and  $\rho_b$ :

$$\frac{\partial \mathbf{v}}{\partial t} + \sum_k v^k \frac{\partial \mathbf{v}}{\partial x^k} + 2\boldsymbol{\Omega} \times \mathbf{v} + \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{x}) + \frac{1}{\rho} \nabla P + \nabla \Phi = 0, \quad (7.41)$$

$$\frac{\partial \mathbf{v}_b}{\partial t} + \sum_k v_b^k \frac{\partial \mathbf{v}_b}{\partial x^k} + 2\boldsymbol{\Omega} \times \mathbf{v}_b + \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{x}) + \frac{1}{\rho_b} \nabla P_b + \nabla \Phi_b = 0, \quad (7.42)$$

Here (7.41) is valid on  $\mathfrak{R}(t)$  and (7.42) is valid on  $\mathfrak{R}_b$ . So we can apply the operator  $\Delta$  on  $\mathfrak{R}_b$ , for which  $\underline{\mathbf{x}} = \mathbf{x} + \boldsymbol{\xi} \in \mathfrak{R}(t)$  while  $\mathbf{x} \in \mathfrak{R}_b$ , to consider

$$\Delta \left[ \frac{\partial \mathbf{v}}{\partial t} + \sum_k v^k \frac{\partial \mathbf{v}}{\partial x^k} + 2\boldsymbol{\Omega} \times \mathbf{v} + \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{x}) + \frac{1}{\rho} \nabla P + \nabla \Phi \right] = 0, \quad (7.43)$$

In order to analyze  $\Delta \left[ \frac{\partial \mathbf{v}}{\partial t} + \sum_k v^k \frac{\partial \mathbf{v}}{\partial x^k} \right]$ , we introduce the operator  $\frac{D}{Dt} \Big|_{\mathbf{V}}$ ,  $\mathbf{V} = \mathbf{v}, \mathbf{v}_b$ . Namely, for a scalar field  $Q$ , we define

$$\frac{D}{Dt} \Big|_{\mathbf{V}} Q(t, \mathbf{x}) = \partial_t Q(t, \mathbf{x}) + (\mathbf{V}(t, \mathbf{x}) | \nabla_{\mathbf{x}} Q(t, \mathbf{x})). \quad (7.44)$$

Here  $\partial_t = \partial/\partial t$ . For a vector field  $\mathbf{Q}$ , we define

$$\left. \frac{D}{Dt} \right|_{\mathbf{V}} \mathbf{Q}(t, \mathbf{x}) = \partial_t \mathbf{Q}(t, \mathbf{x}) + (D_{\mathbf{x}} \mathbf{Q}(t, \mathbf{x})) \mathbf{V}(t, \mathbf{x}). \quad (7.45)$$

Then

$$\left( \left. \frac{D}{Dt} \right|_{\mathbf{V}} \mathbf{Q} \right)^j = \left. \frac{D}{Dt} \right|_{\mathbf{V}} Q^j,$$

since  $((D_{\mathbf{x}} \mathbf{Q}) \mathbf{V})^j = (\mathbf{V} | \nabla_{\mathbf{x}} Q^j)$ .

We claim the formula for scalar field  $Q$  :

$$\Delta \left. \frac{D}{Dt} \right|_{\mathbf{v}} Q = \left. \frac{D}{Dt} \right|_{\mathbf{v}_b} \Delta Q, \quad (7.46)$$

and the same formula for vector field  $\mathbf{Q}$ :

$$\Delta \left. \frac{D}{Dt} \right|_{\mathbf{v}} \mathbf{Q} = \left. \frac{D}{Dt} \right|_{\mathbf{v}_b} \Delta \mathbf{Q}. \quad (7.47)$$

Here we mean

$$\left( \left. \frac{D}{Dt} \right|_{\mathbf{v}} Q \right)_b = \left. \frac{D}{Dt} \right|_{\mathbf{v}_b} Q_b,$$

so we mean

$$\Delta \left. \frac{D}{Dt} \right|_{\mathbf{v}} Q(t, \mathbf{x}) = \left( \left. \frac{D}{Dt} \right|_{\mathbf{v}} Q \right)(t, \mathbf{x} + \boldsymbol{\xi}) - \left. \frac{D}{Dt} \right|_{\mathbf{v}_b} Q_b(t, \mathbf{x}).$$

Proof. It is sufficient to verify (7.46). By definition we see

$$\begin{aligned} \clubsuit &:= \Delta \left. \frac{D}{Dt} \right|_{\mathbf{v}} Q(t, \mathbf{x}) - \left. \frac{D}{Dt} \right|_{\mathbf{v}_b} \Delta Q_b(t, \mathbf{x}) \\ &= (\mathbf{v}(t, \mathbf{x} + \boldsymbol{\xi}) | \nabla_{\mathbf{x}} Q(t, \mathbf{x} + \boldsymbol{\xi}) - \left( \nabla_{\mathbf{x}} Q(t, \mathbf{x} + \boldsymbol{\xi}) \middle| \frac{\partial \boldsymbol{\xi}}{\partial t} \right) \\ &\quad - (\mathbf{v}_b(\mathbf{x}) | (D_{\mathbf{x}} \boldsymbol{\xi})^\top \nabla_{\mathbf{x}} Q(t, \mathbf{x} + \boldsymbol{\xi})) - (\mathbf{v}_b(\mathbf{x}) | \nabla_{\mathbf{x}} Q(t, \mathbf{x} + \boldsymbol{\xi})) \end{aligned}$$

Using (7.12), we have

$$\frac{\partial \boldsymbol{\xi}}{\partial t} = \mathbf{v}(t, \mathbf{x} + \boldsymbol{\xi}) - D_{\mathbf{y}} \boldsymbol{\varphi}(t|0, \bar{\mathbf{x}}) D_{\mathbf{y}} \boldsymbol{\varphi}_b(0|t, \mathbf{x}) \mathbf{v}_b(\mathbf{x}). \quad (7.48)$$

Here we denote

$$\begin{aligned} \boldsymbol{\varphi}(t|s, \mathbf{y}) &= \boldsymbol{\varphi}(t|s, \mathbf{y} || \mathbf{v}), \quad \boldsymbol{\varphi}_b(t|s, \mathbf{y}) = \boldsymbol{\varphi}(t|s, \mathbf{y} || \mathbf{v}_b), \\ \bar{\mathbf{x}} &= \boldsymbol{\varphi}_b(0|t, \mathbf{x}). \end{aligned}$$

On the other hand, we have

$$D_{\mathbf{x}} \boldsymbol{\xi}(t, \mathbf{x}) = D_{\mathbf{y}} \boldsymbol{\varphi}(t|0, \bar{\mathbf{x}}) D_{\mathbf{y}} \boldsymbol{\varphi}_b(0|t, \mathbf{x}) - I. \quad (7.49)$$

Hence

$$\begin{aligned}
\clubsuit &= (\nabla_{\mathbf{x}} Q(t, \mathbf{x} + \boldsymbol{\xi}) | D_{\mathbf{y}} \varphi(t|0, \bar{\mathbf{x}}) D_{\mathbf{y}} \varphi_b(0|t, \mathbf{x}) \mathbf{v}_b(t, \mathbf{x})) \\
&\quad - \left( \mathbf{v}_b(\mathbf{x}) \middle| \left( D_{\mathbf{y}} \varphi(t|0, \bar{\mathbf{x}}) D_{\mathbf{y}} \varphi_b(0|t, \mathbf{x}) \right)^\top \nabla_{\mathbf{x}} Q(t, \mathbf{x} + \boldsymbol{\xi}) \right) \\
&= 0.
\end{aligned}$$

□

We have

$$\Delta \mathbf{v} = \frac{D}{Dt} \Big|_{\mathbf{v}_b} \boldsymbol{\xi}. \quad (7.50)$$

Proof. Consider

$$\clubsuit := \frac{D}{Dt} \Big|_{\mathbf{v}_b} \boldsymbol{\xi}(t, \mathbf{x}) = \partial_t \boldsymbol{\xi}(t, \mathbf{x}) + D_{\mathbf{x}} \boldsymbol{\xi}(t, \mathbf{x}) \mathbf{v}_b(\mathbf{x}).$$

Recalling (7.48), (7.49), we see

$$\begin{aligned}
\clubsuit &= \mathbf{v}(t, \mathbf{x} + \boldsymbol{\xi}) - D_{\mathbf{y}} \varphi(t|0, \bar{\mathbf{x}}) D_{\mathbf{y}} \varphi_b(0|t, \mathbf{x}) \mathbf{v}_b(\mathbf{x}) \\
&\quad + D_{\mathbf{x}} \varphi(t|0, \bar{\mathbf{x}}) D_{\mathbf{y}} \varphi_b(0|t, \mathbf{x}) \mathbf{v}_b(\mathbf{x}) - \mathbf{v}_b(\mathbf{x}) \\
&= \mathbf{v}(t, \mathbf{x} + \boldsymbol{\xi}) - \mathbf{v}_b(\mathbf{x}) = \Delta \mathbf{v}(t, \mathbf{x}).
\end{aligned}$$

□

Consequently we have

$$\begin{aligned}
\Delta \frac{D}{Dt} \Big|_{\mathbf{v}} \mathbf{v} &= \frac{D}{Dt} \Big|_{\mathbf{v}_b} \Delta \mathbf{v} = \left( \frac{D}{Dt} \Big|_{\mathbf{v}_b} \right)^2 \boldsymbol{\xi} \\
&= \frac{\partial^2 \boldsymbol{\xi}}{\partial t^2} + 2 \left( D_{\mathbf{x}} \frac{\partial \boldsymbol{\xi}}{\partial t} \right) \mathbf{v}_b + D_{\mathbf{x}} ((D_{\mathbf{x}} \boldsymbol{\xi}) \mathbf{v}_b) \mathbf{v}_b
\end{aligned} \quad (7.51)$$

and

$$\begin{aligned}
\Delta[2\boldsymbol{\Omega} \times \mathbf{v}] &= 2\boldsymbol{\Omega} \times \frac{D}{Dt} \Big|_{\mathbf{v}_b} \boldsymbol{\xi} \\
&= 2\boldsymbol{\Omega} \times \frac{\partial \boldsymbol{\xi}}{\partial t} + 2\boldsymbol{\Omega} \times ((D_{\mathbf{x}} \boldsymbol{\xi}) \mathbf{v}_b).
\end{aligned} \quad (7.52)$$

Since  $\Delta \mathbf{x} = \boldsymbol{\xi}$  clearly, we have

$$\Delta[\boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{x})] = \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \boldsymbol{\xi}). \quad (7.53)$$

Summing up, we can write

$$\begin{aligned} \Delta \left[ \frac{\partial \mathbf{v}}{\partial t} + \sum_k v^k \frac{\partial \mathbf{v}}{\partial x^k} + 2\boldsymbol{\Omega} \times \mathbf{v} + \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{x}) \right] = \\ = \frac{\partial^2 \boldsymbol{\xi}}{\partial t^2} + 2\boldsymbol{\Omega} \times \frac{\partial \boldsymbol{\xi}}{\partial t} + \mathcal{B}_{\blacklozenge 1} \frac{\partial \boldsymbol{\xi}}{\partial t} + \mathcal{L}_{\blacklozenge 2} \boldsymbol{\xi} + \mathcal{L}_{\blacklozenge 1} \boldsymbol{\xi} + L_{\blacklozenge 0} \boldsymbol{\xi}, \end{aligned} \quad (7.54)$$

with

$$\mathcal{B}_{\blacklozenge 1} \dot{\boldsymbol{\xi}} = 2(D_{\mathbf{x}} \dot{\boldsymbol{\xi}}) \mathbf{v}_b, \quad (7.55a)$$

$$\mathcal{L}_{\blacklozenge 2} \boldsymbol{\xi} = D_{\mathbf{x}}((D_{\mathbf{x}} \boldsymbol{\xi}) \mathbf{v}_b) \mathbf{v}_b, \quad (7.55b)$$

$$\mathcal{L}_{\blacklozenge 1} \boldsymbol{\xi} = 2\boldsymbol{\Omega} \times ((D_{\mathbf{x}} \boldsymbol{\xi}) \mathbf{v}_b), \quad (7.55c)$$

$$L_{\blacklozenge 0} \boldsymbol{\xi} = \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \boldsymbol{\xi}). \quad (7.55d)$$

- Let us analyze  $\Delta \left( \frac{1}{\rho} \nabla P \right)$ . It is easy to see

$$\Delta \left( \frac{1}{\rho} \nabla P \right) = -\frac{\Delta \rho}{\rho_b^2} \nabla P_b + \frac{1}{\rho_b} \nabla \delta P + \frac{1}{\rho_b} (D_{\mathbf{x}} \nabla P_b) \boldsymbol{\xi} + \mathcal{O}(\varepsilon^2)$$

We can write this as

$$\Delta \left( \frac{1}{\rho} \nabla P \right) = -\frac{\delta \rho}{\rho_b^2} \nabla P_b - \frac{(\boldsymbol{\xi} | \nabla \rho_b)}{\rho_b^2} \nabla P_b + \frac{1}{\rho_b} \nabla \delta P + \frac{1}{\rho_b} (D_{\mathbf{x}} \nabla P_b) \boldsymbol{\xi} + \mathcal{O}(\varepsilon^2),$$

since

$$\Delta \rho = \delta \rho + (\boldsymbol{\xi} | \nabla \rho_b) + \mathcal{O}(\varepsilon^2).$$

Moreover we can write

$$\Delta \left( \frac{1}{\rho} \nabla P \right) = -\frac{\delta \rho}{\rho_b^2} \nabla P_b + \frac{1}{\rho_b} \nabla \delta P + D_{\mathbf{x}} \left( \frac{1}{\rho_b} \nabla P_b \right) + \mathcal{O}(\varepsilon^2), \quad (7.56)$$

since

$$D_{\mathbf{x}} \left( \frac{1}{\rho_b} \nabla P_b \right) \boldsymbol{\xi} = \frac{1}{\rho_b} (D_{\mathbf{x}} \nabla P_b) \boldsymbol{\xi} - \frac{(\boldsymbol{\xi} | \nabla \rho_b)}{\rho_b^2} \nabla P_b.$$

Now let us consider  $\delta P$ . Since  $\Delta S = 0$ , we have

$$\Delta P = \left( \frac{\partial P}{\partial \rho} \right)_b \Delta \rho + \mathcal{O}(\varepsilon^2) \quad (7.57)$$

By (7.20), this means

$$\begin{aligned} \delta P &= \left( \frac{\partial P}{\partial \rho} \right)_b \delta \rho + \left( \boldsymbol{\xi} \left| \left( \frac{\partial P}{\partial \rho} \right)_b \nabla \rho_b - \nabla P_b \right. \right) + \mathcal{O}(\varepsilon^2) \\ &= \left( \frac{\partial P}{\partial \rho} \right)_b \delta \rho - \left( \boldsymbol{\xi} \left| \left( \frac{\partial P}{\partial S} \right)_b \nabla S_b \right. \right) + \mathcal{O}(\varepsilon^2) \\ &= \left( \frac{\partial P}{\partial \rho} \right)_b \delta \rho + \left( \frac{\partial P}{\partial S} \right)_b \delta S + \mathcal{O}(\varepsilon^2). \end{aligned}$$

Here we have used (7.26).

Summing up, we have

$$\Delta\left(\frac{1}{\rho}\nabla P\right) = \mathcal{L}_0\xi + \left(D_x\left(\frac{\nabla P_b}{\rho_b}\right)\right)\xi + \mathcal{O}(\varepsilon^2), \quad (7.58)$$

with

$$\mathcal{L}_0\xi = -\frac{\nabla P_b}{\rho_b^2}[\delta\rho] + \frac{1}{\rho_b}\nabla[\delta P], \quad (7.59a)$$

$$[\delta\rho] = -\operatorname{div}(\rho_b\xi), \quad (7.59b)$$

$$[\delta S] = -(\xi|\nabla S_b), \quad (7.59c)$$

$$[\delta P] = \left(\frac{\Gamma P}{\rho}\right)_b[\delta\rho] + \left(\frac{\partial P}{\partial S}\right)_b[\delta S]. \quad (7.59d)$$

Here we recall  $\frac{\partial P}{\partial\rho} = \frac{\Gamma P}{\rho}$  by the definition of  $\Gamma$ , Definition 3.

Moreover we can avoid to mention  $\left(\frac{\partial P}{\partial S}\right)_b$  in the approximation of  $\delta P$ . In fact, since

$$\begin{aligned} \delta P &= \Delta P - (\xi|\nabla P_b) + \mathcal{O}(\varepsilon^2), \\ \Delta\rho &= -\rho_b\operatorname{div}\xi + \mathcal{O}(\varepsilon^2), \end{aligned}$$

(7.57) implies

$$\delta P = -(\Gamma P)_b\operatorname{div}\xi - (\xi|\nabla P_b) + \mathcal{O}(\varepsilon^2). \quad (7.60)$$

We note that [7] gives another expression of  $\Delta\left(\frac{1}{\rho}\nabla P\right)$ , namely,

$$\Delta\left(\frac{1}{\rho}\nabla P\right) = \frac{1}{\rho_b}\left[-\nabla(\Gamma_b P_b\operatorname{div}\xi) - ((D_x\xi)\nabla P_b)^\top + (\operatorname{div}\xi)\nabla P_b\right] + \mathcal{O}(\varepsilon^2). \quad (7.61)$$

This expression (7.61) can be derived by applying the formula

$$\begin{aligned} \Delta\nabla Q &= \nabla\Delta Q + (D\nabla Q)\xi - \nabla(\xi|\nabla Q) + \mathcal{O}(\varepsilon^2) \\ &= \nabla\Delta Q - ((D\xi)\nabla Q)^\top + \mathcal{O}(\varepsilon^2) \end{aligned} \quad (7.62)$$

to  $Q = P$  and noting

$$\Delta P = \left(\frac{\Gamma P}{\rho}\right)_b\Delta\rho + \mathcal{O}(\varepsilon^2) = \Gamma_b P_b\operatorname{div}\xi + \mathcal{O}(\varepsilon^2).$$

and

$$\frac{\Delta\rho}{\rho_b} = -\operatorname{div}\xi + \mathcal{O}(\varepsilon^2).$$

- Let us look at

$$\Delta \nabla \Phi(\mathbf{x}) = \nabla \Phi(\mathbf{x} + \boldsymbol{\xi}) - \nabla \Phi_b(\mathbf{x}),$$

where

$$\begin{aligned} \Phi &= -4\pi G \mathcal{K}[\rho], & \Phi_b &= -4\pi G \mathcal{K}[\rho_b], \\ \mathcal{K}[g](\mathbf{x}) &= \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{g(\mathbf{x}')}{\|\mathbf{x} - \mathbf{x}'\|} d\mathbf{x}'. \end{aligned}$$

We introduce the integral operator  $\mathcal{K}^\nabla$  by

$$\mathcal{K}^\nabla[g](\mathbf{x}) = -\frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{\mathbf{x} - \mathbf{x}'}{\|\mathbf{x} - \mathbf{x}'\|^3} g(\mathbf{x}') d\mathbf{x}'. \quad (7.63)$$

When  $g \in C_0^\alpha(\mathbb{R}^3)$ ,  $0 < \alpha < 1$ , then  $\mathcal{K}^\nabla[g] \in C^{1+\alpha}(\mathbb{R}^3)$  and  $\nabla \mathcal{K}[g] = \mathcal{K}^\nabla[g]$ . So

$$\begin{aligned} \Delta \nabla \Phi(\mathbf{x}) &= -4\pi G \left( \mathcal{K}^\nabla[\delta\rho](\underline{\mathbf{x}}) + \nabla \Phi_b(\underline{\mathbf{x}}) - \nabla \Phi_b(\mathbf{x}) \right) \\ &= -4\pi G \mathcal{K}^\nabla[\delta\rho] + (D_{\mathbf{x}} \nabla \Phi_b(\mathbf{x})) \boldsymbol{\xi} + \mathcal{O}(\varepsilon^2). \end{aligned}$$

Here  $\underline{\mathbf{x}} = \mathbf{x} + \boldsymbol{\xi}$ , and we note  $\mathcal{K}^\nabla[\delta\rho] \in C^{1+\alpha}$  and  $\mathcal{O}(\varepsilon)$ .

Consequently we can claim

$$\Delta \nabla \Phi = 4\pi G \mathcal{L}_1 \boldsymbol{\xi} + (D_{\mathbf{x}} \nabla \Phi_b) \boldsymbol{\xi} + \mathcal{O}(\varepsilon^2) \quad (7.64)$$

with

$$\mathcal{L}_1 \boldsymbol{\xi} = \nabla \mathcal{K}[\text{div}(\rho_b \boldsymbol{\xi})]. \quad (7.65)$$

- Summing up, we have derived the equation used by astrophysicists:

$$\frac{\partial^2 \boldsymbol{\xi}}{\partial t^2} + \mathcal{B}_\diamond \frac{\partial w}{\partial t} + \mathcal{L}_\diamond \boldsymbol{\xi} = 0, \quad (7.66)$$

where

$$\mathcal{B}_\diamond \dot{\boldsymbol{\xi}} = 2\boldsymbol{\Omega} \times \dot{\boldsymbol{\xi}} + \mathcal{B}_{\diamond 1} \dot{\boldsymbol{\xi}}, \quad (7.67)$$

with

$$\mathcal{B}_{\diamond 1} \dot{\boldsymbol{\xi}} = 2(D\dot{\boldsymbol{\xi}}) \mathbf{v}_b, \quad (7.68)$$

and

$$\mathcal{L}_\diamond = \mathcal{L}_{\diamond 2} + \mathcal{L}_{\diamond 1} + L_{\diamond 0} + \mathcal{L}_0 + L_{0\diamond} + 4\pi G \mathcal{L}_1 + L_{1\diamond} \quad (7.69)$$

with

$$\mathcal{L}_{\blacklozenge 2}\boldsymbol{\xi} = D((D\boldsymbol{\xi})\mathbf{v}_b)\mathbf{v}_b, \quad (7.70a)$$

$$\mathcal{L}_{\blacklozenge 1}\boldsymbol{\xi} = 2\boldsymbol{\Omega} \times ((D\boldsymbol{\xi})\mathbf{v}_b), \quad (7.70b)$$

$$L_{\blacklozenge 0}\boldsymbol{\xi} = \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \boldsymbol{\xi}), \quad (7.70c)$$

$$\mathcal{L}_0\boldsymbol{\xi} = -\frac{[\delta\rho]}{\rho_b^2} + \frac{1}{\rho_b}\nabla[\delta P], \quad (7.70d)$$

with

$$\begin{aligned} [\delta\rho] &= -\operatorname{div}(\rho_b\boldsymbol{\xi}), \quad [\delta S] = -(\boldsymbol{\xi}|\nabla S_b), \\ [\delta P] &= \left(\frac{\Gamma P}{\rho}\right)_b[\delta\rho] + \left(\frac{\partial P}{\partial S}\right)_b[\delta S] = \left(\frac{\Gamma P}{\rho}\right)_b[\delta\rho] + \Gamma_b P_b(\boldsymbol{\xi}|\mathbf{a}_b) \end{aligned} \quad (7.70e)$$

$$L_{0\blacklozenge}\boldsymbol{\xi} = \left(D\left(\frac{\nabla P_b}{\rho_b}\right)\right)\boldsymbol{\xi}, \quad (7.70f)$$

$$\mathcal{L}_1\boldsymbol{\xi} = -\nabla\mathcal{K}[[\delta\rho]] = \nabla\mathcal{K}[\operatorname{div}(\rho_b\boldsymbol{\xi})], \quad (7.70g)$$

$$L_{1\blacklozenge}\boldsymbol{\xi} = (D\nabla\Phi_b)\boldsymbol{\xi}. \quad (7.70h)$$

We denote

$$\mathcal{L} = \mathcal{L}_0 + 4\pi G\mathcal{L}_1.$$

However we would like to rewrite  $\mathcal{L}_{\blacklozenge}$  further. Since  $(\rho_b, S_b, \mathbf{v}_b)$  satisfies the equation of motion

$$(D\mathbf{v}_b)\mathbf{v}_b + 2\boldsymbol{\Omega} \times \mathbf{v}_b + \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{x}) + \frac{\nabla P_b}{\rho_b} + \nabla\Phi_b = 0$$

on  $\mathfrak{R}_b$ , we see

$$L_{0\blacklozenge}\boldsymbol{\xi} + L_{1\blacklozenge}\boldsymbol{\xi} = -D\left((D\mathbf{v}_b)\mathbf{v}_b + 2\boldsymbol{\Omega} \times \mathbf{v}_b + \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{x})\right)\boldsymbol{\xi}.$$

But we have

$$D\left(\boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{x})\right)\boldsymbol{\xi} = \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \boldsymbol{\xi}).$$

Thus

$$L_{0\blacklozenge} + L_{1\blacklozenge} + L_{\blacklozenge 0} = \mathcal{L}_{\blacklozenge 00} + \mathcal{L}_{\blacklozenge 01}, \quad (7.71)$$

where

$$\mathcal{L}_{\blacklozenge 00}\boldsymbol{\xi} = -\left(D((D\mathbf{v}_b)\mathbf{v}_b)\right)\boldsymbol{\xi}, \quad \mathcal{L}_{\blacklozenge 01}\boldsymbol{\xi} = -\left(D(2\boldsymbol{\Omega} \times \mathbf{v}_b)\right)\boldsymbol{\xi}. \quad (7.72)$$

We define

$$\mathcal{L}_{\blacklozenge 0} := \mathcal{L}_{\blacklozenge 00} + \mathcal{L}_{\blacklozenge 01}. \quad (7.73)$$

Consequently we have the equation (7.5) with (7.6) (7.7) (7.8) with (7.9a) - (7.9f), as the equation of linear adiabatic stellar oscillations used by astrophysicists after [13] [7] [15].

We note that the derivation above does not require that  $\mathbf{v}_b$  is small, say,  $\mathcal{O}(\varepsilon)$  during the linear approximation procedure, while we supposed  $\mathbf{v}_b = \mathcal{O}(\varepsilon)$  to derive the equation (1.11) in Section 3. But we recall that (7.29):  $\rho^0 = \rho_b$  and (7.24):  $S^0 = S_b$  are supposed, while only (3.1):  $\{\rho^0 > 0\} = \{\rho_b > 0\}$  was supposed in Section 3. Moreover we note that  $\boldsymbol{\xi}(0, \mathbf{x}) = \mathbf{0}$  by the definition (7.1), while the Lagrange displacement  $\mathbf{u}$  in our equation (1.11) which was defined by (1.14) or (3.27):

$$\mathbf{u}(t, \bar{\mathbf{x}}) = \boldsymbol{\varphi}(t|0, \bar{\mathbf{x}}|\mathbf{v}) - \boldsymbol{\varphi}(t|0, \bar{\mathbf{x}}|\mathbf{v}_b) + \mathbf{u}^0(\bar{\mathbf{x}})$$

enjoys  $\mathbf{u}(0, \bar{\mathbf{x}}) = \mathbf{u}^0(\bar{\mathbf{x}})$ , which  $\mathcal{O}(\varepsilon)$  was arbitrary as long as

$$\rho^0 - \rho_b = -\operatorname{div}(\rho_b \mathbf{u}^0), \quad S^0 - S_b = -(\nabla S_b | \mathbf{u}^0)$$

and can be  $\neq \mathbf{0}$ .

### 7.3 Trivial displacements

By the definition (7.1) it holds that  $\boldsymbol{\xi}(0, \mathbf{x}) = \mathbf{0}$ . But it is possible to consider solutions  $\tilde{\boldsymbol{\xi}} = \boldsymbol{\xi} + \boldsymbol{\eta}$  of the equation LASO ( $\blacklozenge$ ) (7.5). After [7] let us call  $\boldsymbol{\eta}(t, \mathbf{x})$  a trivial displacement if it satisfies

$$0 = -\operatorname{div}(\rho_b \boldsymbol{\eta}), \quad (7.74)$$

$$0 = -(\nabla S_b | \boldsymbol{\eta}), \quad (7.75)$$

$$\mathbf{0} = \partial_t \boldsymbol{\eta} + (D\boldsymbol{\eta})\mathbf{v}_b - (D\mathbf{v}_b)\boldsymbol{\eta}. \quad (7.76)$$

Here  $\partial_t = \partial/\partial t$ . Then

$$[\delta\rho || \boldsymbol{\eta}] := -\operatorname{div}(\rho_b \boldsymbol{\eta}) = 0, \quad (7.77)$$

$$[\delta S || \boldsymbol{\eta}] := -(\nabla S_b | \boldsymbol{\eta}) = 0, \quad (7.78)$$

$$[\delta P || \boldsymbol{\eta}] := \left(\frac{\Gamma P}{\rho}\right)_b [\delta\rho || \boldsymbol{\eta}] - \left(\frac{\partial P}{\partial S}\right)_b (\nabla S_b | \boldsymbol{\eta}) = 0, \quad (7.79)$$

$$[\delta \mathbf{v} || \boldsymbol{\eta}] := \partial_t \boldsymbol{\eta} + (D\boldsymbol{\eta})\mathbf{v}_b - (D\mathbf{v}_b)\boldsymbol{\eta} = \mathbf{0}. \quad (7.80)$$

Here we note that

$$\begin{aligned} \delta \mathbf{v} &= \Delta \mathbf{v} - (D\mathbf{v}_b)\boldsymbol{\xi} + \mathcal{O}(\varepsilon^2) \quad (\text{see } (7.21)), \\ &= \frac{D}{Dt} \Big|_{\mathbf{v}_b} \boldsymbol{\xi} - (D\mathbf{v}_b)\boldsymbol{\xi} + \mathcal{O}(\varepsilon^2) \quad (\text{see } (7.50)) \\ &= \partial_t \boldsymbol{\xi} + (D\boldsymbol{\xi})\mathbf{v}_b - (D\mathbf{v}_b)\boldsymbol{\xi} + \mathcal{O}(\varepsilon^2), \quad (\text{see } (7.45)), \end{aligned}$$



so

$$[\delta \mathbf{v} || \boldsymbol{\xi}] = \partial_t \boldsymbol{\xi} + (D\boldsymbol{\xi})\mathbf{v}_b - (D\mathbf{v}_b)\boldsymbol{\xi}$$

approximates  $\delta \mathbf{v}$  .

Let  $\boldsymbol{\eta}(t, \mathbf{x})$  be a trivial displacement. Then (7.74) and (7.76) imply  $\mathcal{L}_0 \boldsymbol{\eta} = \mathbf{0}$  and  $\mathcal{L}_1 \boldsymbol{\eta} = \mathbf{0}$ . Moreover we have

$$\partial_t^2 \boldsymbol{\eta} + \mathcal{B}_{\blacklozenge 1} \partial_t \boldsymbol{\eta} + (\mathcal{L}_{\blacklozenge 2} + \mathcal{L}_{\blacklozenge 00}) \boldsymbol{\eta} = \mathbf{0}, \quad (7.81)$$

and

$$2\boldsymbol{\Omega} \times \partial_t \boldsymbol{\eta} + (\mathcal{L}_{\blacklozenge 1} + \mathcal{L}_{\blacklozenge 01}) \boldsymbol{\eta} = \mathbf{0}. \quad (7.82)$$

Therefore we can claim that  $\boldsymbol{\eta}$  satisfies the equation (7.5), so  $\tilde{\boldsymbol{\xi}} = \boldsymbol{\xi} + \boldsymbol{\eta}$  should be a solution of (7.5).

Proof of (7.81). Eliminating  $\eta_t \boldsymbol{\eta}, \partial_t^2 \boldsymbol{\eta}$  by (7.76), we see that the left-hand side of (7.81) reduces to

$$\begin{aligned} & (D((D\mathbf{v}_b)\boldsymbol{\eta}) - (D\mathbf{v}_b)(D\boldsymbol{\eta}))\mathbf{v}_b - ((D\mathbf{v}_b)(D\mathbf{v}_b) - D((D\mathbf{v}_b)\mathbf{v}_b))\boldsymbol{\eta} \\ &= \langle D^2 \mathbf{v}_b | \boldsymbol{\eta}, \mathbf{v}_b \rangle - \langle D^2 \mathbf{v}_b | \boldsymbol{\eta}, \mathbf{v}_b \rangle = \mathbf{0}. \end{aligned}$$

Here we denote

$$\langle D^2 \mathbf{f} | \mathbf{X}, \mathbf{Y} \rangle^i = \sum_{k,\ell} \frac{\partial^2 f^i}{\partial x^k \partial x^\ell} X^k Y^\ell. \quad (7.83)$$

□

Proof of (7.82). Eliminating  $\partial_t \boldsymbol{\eta}$  by (7.76), we see that the left-hand side of (7.82) reduces to

$$2\boldsymbol{\Omega} \times (D\mathbf{v}_b)\boldsymbol{\eta} - (D(2\boldsymbol{\Omega} \times \mathbf{v}_b))\boldsymbol{\eta} = \mathbf{0}.$$

□

Of course  $\boldsymbol{\eta} = \mathbf{0}$  is a trivial displacement, which could be called the trivial trivial displacement, and we are interested in non-trivial trivial displacements in this sense.

Let us consider

$$\boldsymbol{\eta} = \frac{1}{\rho_b} \nabla S_b \times \nabla f, \quad (7.84)$$

where  $f$  is a scalar fields wich satisfies

$$\left. \frac{D}{Dt} \right|_{\mathbf{v}_b} f = \partial_t f + (\mathbf{v}_b | f) = 0. \quad (7.85)$$

Then  $\eta$  is a trivial displacement.

Proof. Clearly (7.74), (7.75) are satisfied, for any  $f$ . Usng the properties of the background

$$\sum_k \mathbf{v}_b \partial_k \rho_b = 0, \quad \sum_k \mathbf{v}_b \partial_k S_b = 0, \quad (7.86)$$

where  $\partial_k = \partial/\partial x^k$ , we can verify

$$((D\boldsymbol{\eta})\mathbf{v}_b - (D\mathbf{v}_b)\boldsymbol{\eta})^j = \frac{1}{\rho_b} \left( - \sum_{\alpha, \beta} E^{j\alpha\beta} (\partial_\alpha S_b) (\partial_\beta f) + \nabla S_b \times \nabla(\mathbf{v}_b|f) \right), \quad (7.87)$$

where

$$E^{j\alpha\beta} = \sum_k \left( \epsilon^{jk\beta} \partial_k v_b^\alpha + \epsilon^{j\alpha k} \partial_k v_b^\beta + \epsilon^{k\alpha\beta} \partial_k v_b^j \right), \quad \epsilon^{\lambda\mu\nu} = \text{sign} \begin{pmatrix} 1 & 2 & 3 \\ \lambda & \mu & \nu \end{pmatrix}. \quad (7.88)$$

But we can verify

$$E^{j\alpha\beta} = \epsilon^{j\alpha\beta} \sum_k \partial_k v_b^k, \quad (7.89)$$

therefore  $E^{j\alpha\beta} = 0$  for  $\forall j, \alpha, \beta$ , since  $\text{div} \mathbf{v}_b = 0$ . Thus

$$(D\boldsymbol{\eta})\mathbf{v}_b - (D\mathbf{v}_b)\boldsymbol{\eta} = \frac{1}{\rho_b} \nabla S_b \times \nabla(\mathbf{v}_b|f)$$

so that (7.76) is satisfied, since

$$\partial_t \boldsymbol{\eta} = \frac{1}{\rho_b} \nabla S_b \times \nabla \partial_t f.$$

□

General solution of (7.85) is

$$f(t, \mathbf{x}) = f^0(\varphi(0|t, \mathbf{x}||\mathbf{v}_b)), \quad (7.90)$$

where  $\bar{\mathbf{x}} \mapsto f^0(\bar{\mathbf{x}})$  is an arbitrary function on  $\mathfrak{R}_b$ . Then

$$\boldsymbol{\eta}(0, \mathbf{x}) = \frac{1}{\rho_b(\mathbf{x})} \nabla S_b \times \nabla_{\bar{\mathbf{x}}} f^0(\mathbf{x}). \quad (7.91)$$

When  $\nabla S_b \neq \mathbf{0}$ , that is,  $S_b$  is not a constant, then this construction gives many non-trivial trivial solutions. However when  $\nabla S_b = \mathbf{0}$ ,  $\boldsymbol{\eta}$  given by (7.84) is trivial,  $= \mathbf{0}$ . If  $\mathbf{v}_b = \mathcal{O}(\varepsilon)$  the displacement given by

$$\boldsymbol{\eta}(t, \mathbf{x}) = \frac{1}{\rho_b(\bar{\mathbf{x}})} \text{rot}_{\bar{\mathbf{x}}} \mathbf{f}(\bar{\mathbf{x}}), \quad \bar{\mathbf{x}} = \varphi(0|t, \mathbf{x}||\mathbf{v}_b) \quad (7.92)$$

satisfies (7.74), (7.76) approximately with errors  $\mathcal{O}(\varepsilon)$ , provided that the function  $\mathbf{f}$  is small  $\mathcal{O}(\varepsilon)$ . We compare these observations with the discussion of Theorem 4 in Section 4.

#### 7.4 Functional analysis of the equation of LASO(♦) (7.5)

Let us consider the equation (7.5) in the functional space  $\mathfrak{H} = L^2(\mathfrak{R}_b, \rho_b d\mathbf{x}; \mathbb{C}^3)$ . We suppose Assumption 1 and the background  $(\rho_b, S_b, \mathbf{v}_b)$  is an admissible

barotropic stationary solution.

The operator  $\mathcal{B} : \dot{\xi} \mapsto 2\Omega \times \dot{\xi}$  can be considered as a bounded linear operator in  $\mathcal{B}(\mathfrak{H})$ , and  $\mathcal{L} = \mathcal{L}_0 + 4\pi\mathbb{G}\mathcal{L}_1$  can be extended to a self-adjoint operator bounded from below in  $\mathfrak{H}$ , by following the discussion of Section 4. We look at the resting terms  $\mathcal{B}_{\blacklozenge 1}, \mathcal{L}_{\blacklozenge 2}, \mathcal{L}_{\blacklozenge 1}, \mathcal{L}_{\blacklozenge 0}$ .

First, we see that  $\mathcal{B}_{\blacklozenge 1}$  is antisymmetric considered on  $C_0^\infty(\mathfrak{R}_b)$ . In fact, for  $\mathbf{v}_1, \mathbf{v}_2 \in C_0^\infty$ , we have

$$\begin{aligned} (\mathcal{B}_{\blacklozenge 1}\mathbf{v}_1|\mathbf{v}_2)_{\mathfrak{H}} &= 2 \int \omega_b \left( \frac{\partial \mathbf{v}_1}{\partial \phi} \middle| \mathbf{v}_2 \right) \rho_b d\mathbf{x} = \\ &= -2 \int \omega_b \left( \mathbf{v}_1 \middle| \frac{\partial \mathbf{v}_2}{\partial \phi} \right) \rho_b d\mathbf{x} = -(\mathbf{v}_1|\mathcal{B}_{\blacklozenge 1}\mathbf{v}_2)_{\mathfrak{H}}. \end{aligned}$$

However this operator is not bounded in  $\mathfrak{H}$ -norm, if  $\mathbf{v}_b \neq \mathbf{0}$ , say,  $\omega_b \neq 0$ .

$\mathcal{L}_{\blacklozenge 2}, \mathcal{L}_{\blacklozenge 1}$  are symmetric on  $C_0^\infty(\mathfrak{R}_b)$ . In fact we have, for  $\mathbf{u}_1, \mathbf{u}_2 \in C_0^\infty$ ,

$$\begin{aligned} (\mathcal{L}_{\blacklozenge 2}\mathbf{u}_1|\mathbf{u}_2)_{\mathfrak{H}} &= - \int \omega_b^2 \left( \frac{\partial \mathbf{u}_1}{\partial \phi} \middle| \frac{\partial \mathbf{u}_2}{\partial \phi} \right) \rho_b d\mathbf{x} \\ &= (\mathbf{u}_1|\mathcal{L}_{\blacklozenge 2}\mathbf{u}_2)_{\mathfrak{H}}, \\ (\mathcal{L}_{\blacklozenge 1}\mathbf{u}_1|\mathbf{u}_2)_{\mathfrak{H}} &= 2\Omega \int \omega_b \left( \frac{\partial u_1^1}{\partial \phi} (u_2^2)^* + u_1^2 \left( \frac{\partial u_2^1}{\partial \phi} \right)^* \right) \rho_b d\mathbf{x} \\ &= (\mathbf{u}_1|\mathcal{L}_{\blacklozenge 1}\mathbf{u}_2)_{\mathfrak{H}}. \end{aligned}$$

However  $\mathcal{L}_{\blacklozenge 2} + \mathcal{L}_{\blacklozenge 1}$  cannot be bounded from below, if  $\mathbf{v}_b \neq 0$ . In fact, we see

$$(\mathcal{L}_{\blacklozenge 2}\mathbf{u}|\mathbf{u})_{\mathfrak{H}} = - \int \omega_b^2 \left\| \frac{\partial \mathbf{u}}{\partial \phi} \right\|^2 \rho_b d\mathbf{x}, \quad (7.93)$$

$$(\mathcal{L}_{\blacklozenge 1}\mathbf{u}|\mathbf{u})_{\mathfrak{H}} = 4\Omega \int \omega_b \Re \left[ \frac{\partial u^1}{\partial \phi} (u^2)^* \right] \rho_b d\mathbf{x}. \quad (7.94)$$

Therefore

$$((\mathcal{L}_{\blacklozenge 2} + \mathcal{L}_{\blacklozenge 1})\mathbf{u}|\mathbf{u})_{\mathfrak{H}} = -m^2 \int \omega_b^2 \|\mathbf{u}\|^2 \rho_b d\mathbf{x},$$

when

$$\mathbf{u}(\mathbf{x}) = \mathbf{U}(r, \vartheta) e^{im\phi}, \quad \mathbf{U} = \begin{bmatrix} U \\ U \\ 0 \end{bmatrix}, \quad \text{while } U \in C_0^\infty(\mathfrak{R}_b; \mathbb{R}).$$

Suppose  $\omega_b \neq 0$ , and we suppose  $\omega_b^2 \geq \kappa^2 > 0$ ,  $\kappa$  being a constant, on a domain  $\mathcal{V}(\neq \emptyset), \text{supp}[U] \subset \mathcal{V}$ . Then we have

$$((\mathcal{L}_{\blacklozenge 2} + \mathcal{L}_{\blacklozenge 1})\mathbf{u}|\mathbf{u})_{\mathfrak{H}} \leq -m^2 \kappa^2 \|\mathbf{u}\|_{\mathfrak{H}}^2.$$

Therefore

$$\inf_{\mathbf{u} \in C_0^\infty} \frac{((\mathcal{L}_{\blacklozenge 2} + \mathcal{L}_{\blacklozenge 1})\mathbf{u}|\mathbf{u})_{\mathfrak{H}}}{\|\mathbf{u}\|_{\mathfrak{H}}^2} \leq -m^2 \kappa^2$$

for  $\forall m$  so that

$$\inf_{\mathbf{u} \in C_0^\infty} \frac{((\mathcal{L}_{\blacklozenge 2} + \mathcal{L}_{\blacklozenge 1})\mathbf{u}|\mathbf{u})_{\mathfrak{H}}}{\|\mathbf{u}\|_{\mathfrak{H}}^2} = -\infty.$$

Thus  $\mathcal{L}_{\blacklozenge 2} + \mathcal{L}_{\blacklozenge 1}$  is not bounded from below, if  $\omega_b \neq 0$ .

Finally, as for the operator  $\mathcal{L}_{\blacklozenge 0} = \mathcal{L}_{\blacklozenge 00} + \mathcal{L}_{\blacklozenge 01}$ , it is a simple multiplication by the real-valued matrix function  $W_b \in C(\mathbb{R}^3; \mathbb{R}^{3 \times 3})$ , therefore  $\mathcal{L}_{\blacklozenge 0}$  can be considered as a symmetric bounded operator on  $\mathfrak{H}$ .

Summing up, when  $\mathbf{v}_b \neq 0$ , or at least, when  $\mathbf{v}_b \neq \mathcal{O}(\varepsilon)$ , by which we cannot neglect terms  $\mathcal{B}_{\blacklozenge 1} \partial_t \xi, \mathcal{L}_{\blacklozenge 2} \xi, \mathcal{L}_{\blacklozenge 1} \xi, \mathcal{L}_{\blacklozenge 0} \xi$  as  $\mathcal{O}(\varepsilon^2)$ , the application of the Hille-Yosida theory as discussed in Section 4 does not work for the unboundedness of  $\mathcal{B}_{\blacklozenge 1}$ , which implies the unboundedness of the total  $\mathcal{B}_{\blacklozenge}$ , and the unboundedness from below of  $\mathcal{L}_{\blacklozenge 2} + \mathcal{L}_{\blacklozenge 1}$ , which implies the unboundedness from below of the total  $\mathcal{L}_{\blacklozenge}$ . However, if we can suppose that  $\mathbf{v}_b = \mathcal{O}(\varepsilon)$ , then the terms  $\mathcal{B}_{\blacklozenge 1} \partial_t \xi, \mathcal{L}_{\blacklozenge 2} \xi, \mathcal{L}_{\blacklozenge 1} \xi, \mathcal{L}_{\blacklozenge 0} \xi$  are of  $\mathcal{O}(\varepsilon^2)$  and negligible. In this situation,  $\mathcal{B}_{\blacklozenge}, \mathcal{L}_{\blacklozenge}$  reduce to  $\mathcal{B}$  and  $\mathcal{L} = \mathcal{L}_0 + 4\pi G \mathcal{L}_1$ , that is, the astrophysicists equation of LASO ( $\blacklozenge$ ) (7.5) reduces to our equation of LASO (1.11).

• However we want to say something for the initial value problem to the equation of LASO ( $\blacklozenge$ ):

$$\frac{\partial^2 \xi}{\partial t^2} + \mathcal{B}_{\blacklozenge} \frac{\partial \xi}{\partial t} + \mathcal{L}_{\blacklozenge} \xi = \mathbf{f}(t, \mathbf{x}), \quad (7.95)$$

$$\xi|_{t=0} = \xi^0, \quad \frac{\partial \xi}{\partial t} \Big|_{t=0} = \mathbf{v}^0, \quad (7.96)$$

where  $\mathbf{f}, \xi^0, \mathbf{v}^0$  are given data. Suppose that the background is a barotropic admissible stationary solution.

In order to fix the idea to consider the integro-differential operator  $\mathcal{L}_{\blacklozenge}$  in the Hilbert space  $\mathfrak{H} = L^2(\mathfrak{R}_b, \rho_b d\mathbf{x}; \mathbb{C}^3)$ , we define  $\mathbf{L}_{\blacklozenge}$  as

$$\begin{aligned} \mathbf{D}(\mathbf{L}_{\blacklozenge}) &= \{\mathbf{u} \in (\mathfrak{G}_{\blacklozenge})_0 \mid \text{Both } \mathcal{L}_{\blacklozenge 2} \mathbf{u} \text{ and } \mathcal{L} \mathbf{u} \\ &\text{belong to } \mathfrak{H} \text{ in distribution sense}\}, \end{aligned} \quad (7.97)$$

$$\mathbf{L}_{\blacklozenge} \mathbf{u} = \mathcal{L}_{\blacklozenge} \mathbf{u} \quad \text{for } \mathbf{u} \in \mathbf{D}(\mathbf{L}_{\blacklozenge}), \quad (7.98)$$

where  $\mathfrak{G}_{\blacklozenge}$  stands for the functional space

$$\mathfrak{G}_{\blacklozenge} = \left\{ \mathbf{u} \in \mathfrak{H} \mid \operatorname{div}(\rho_b \mathbf{u}) \in L^2\left(\left(\frac{\Gamma P}{\rho^2}\right)_b d\mathbf{x}\right), \quad \omega_b \frac{\partial \mathbf{u}}{\partial \phi} \in \mathfrak{H} \right\} \quad (7.99)$$

endowed with tht norm

$$\|\mathbf{u}\|_{\mathfrak{G}_{\blacklozenge}} = \left[ \|\mathbf{u}\|_{\mathfrak{H}}^2 + \left\| \operatorname{div}(\rho_b \mathbf{u}); L^2\left(\left(\frac{\Gamma P}{\rho^2}\right)_b d\mathbf{x}\right) \right\|^2 + \left\| \omega_b \frac{\partial \mathbf{u}}{\partial \phi} \right\|_{\mathfrak{H}}^2 \right]^{\frac{1}{2}}, \quad (7.100)$$

and  $(\mathfrak{G}_\diamond)_0$  stands for the closure of  $C_0^\infty(\mathfrak{R}_b)$  in  $\mathfrak{G}_\diamond$ .

Note that, if  $\mathbf{u} \in \mathfrak{G}_\diamond$ , then we have, a priori,  $\mathcal{L}_{\diamond 1}\mathbf{u} \in \mathfrak{H}$  and  $\mathcal{L}_{\diamond 0}\mathbf{u} \in \mathfrak{H}$ . Thus, if  $\mathbf{u} \in \mathbf{D}(\mathbf{L}_\diamond)$ , then  $\mathcal{L}_\diamond\mathbf{u} \in \mathfrak{H}$  in distribution sense. But, if  $\mathbf{v}_b \neq 0$ , we may not say that  $\mathcal{L}_\diamond\mathbf{u} \in \mathfrak{H}$  guarantees  $\mathcal{L}_{\diamond 2}\mathbf{u}, \mathcal{L}\mathbf{u} \in \mathfrak{H}$  separately.

This formulation comes from the following observation.

Put, for  $\mathbf{u} \in \mathfrak{G}_\diamond$ ,

$$Q_{\diamond 2}[\mathbf{u}] := - \int \omega_b^2 \left\| \frac{\partial \mathbf{u}}{\partial \phi} \right\|^2 \rho_b d\mathbf{x} = - \left\| \omega_b \frac{\partial \mathbf{u}}{\partial \phi} \right\|_{\mathfrak{H}}^2, \quad (7.101)$$

$$Q_{\diamond 1}[\mathbf{u}] := 4\Omega \int \omega_b \Re \left[ \frac{\partial u^1}{\partial \phi} (u^2)^* \right] \rho_b d\mathbf{x}, \quad (7.102)$$

$$Q_{\diamond 0}[\mathbf{u}] := - \int (W_b \mathbf{u} | \mathbf{u}) \rho_b d\mathbf{x}, \quad (7.103)$$

$$\begin{aligned} Q_0[\mathbf{u}] &:= \int \frac{\Gamma P}{\rho^2} |\operatorname{div}(\rho \mathbf{u})|^2 d\mathbf{x} + 2\Re \left[ \int \frac{\Gamma P \mathcal{A}}{\rho} (\mathbf{u} | \mathbf{n}) \operatorname{div}(\rho \mathbf{u})^* d\mathbf{x} \right] + \\ &\quad - \int \frac{\Gamma P \mathcal{A}}{\rho} |(\mathbf{u} | \mathbf{n})|^2 \|\nabla \rho\| d\mathbf{x} 4\pi G \int \mathcal{K}[\operatorname{div}(\rho \mathbf{u})] \operatorname{div}(\rho \mathbf{u})^* d\mathbf{x}. \end{aligned} \quad (7.104)$$

Then we have the estimates

$$|Q_{\diamond 1}[\mathbf{u}]| \leq 2|\Omega| \left( \|\mathbf{u}\|_{\mathfrak{H}}^2 + \left\| \omega_b \frac{\partial \mathbf{u}}{\partial \phi} \right\|_{\mathfrak{H}}^2 \right), \quad (7.105)$$

$$|Q_{\diamond 0}[\mathbf{u}]| \leq \|W_b\|_{L^\infty} \|\mathbf{u}\|_{\mathfrak{H}}^2. \quad (7.106)$$

Put, for  $\mathbf{u} \in \mathfrak{G}_\diamond$ ,

$$Q_0^\diamond[\mathbf{u}] := \sum_{\nu=0,1,2} Q_{\diamond \nu}[\mathbf{u}] + Q_0[\mathbf{u}], \quad (7.107)$$

Then, if  $\mathbf{u} \in C_0^\infty$ , we have

$$Q_{\diamond 2}[\mathbf{u}] = (\mathcal{L}_{\diamond 2} \mathbf{u} | \mathbf{u})_{\mathfrak{H}}, \quad (7.108)$$

$$Q_{\diamond 1}[\mathbf{u}] = (\mathcal{L}_{\diamond 1} \mathbf{u} | \mathbf{u})_{\mathfrak{H}}, \quad (7.109)$$

$$Q_{\diamond 0}[\mathbf{u}] = (\mathcal{L}_{\diamond 0} \mathbf{u} | \mathbf{u})_{\mathfrak{H}}, \quad (7.110)$$

$$Q_0[\mathbf{u}] = (\mathcal{L} \mathbf{u} | \mathbf{u})_{\mathfrak{H}}, \quad (7.111)$$

and

$$Q_0^\diamond[\mathbf{u}] = (\mathcal{L}_\diamond \mathbf{u} | \mathbf{u})_{\mathfrak{H}}. \quad (7.112)$$

Let us introduce an equivalent norm  $[\|\cdot\|]_\diamond$  on  $\mathfrak{G}_\diamond$  as follows.

Putting

$$K_1 := 2 + 4|\Omega|, \quad (7.113)$$

$$K_0 := 2 + 4|\Omega| + 2\|W_b\|_{L^\infty}, \quad (7.114)$$

we have

$$\begin{aligned} & \left(\frac{K_0}{2} + 1\right) \|\mathbf{u}\|_{\mathfrak{H}}^2 + \frac{K_1}{2} \left\| \omega_b \frac{\partial \mathbf{u}}{\partial \phi} \right\|_{\mathfrak{H}}^2 \leq \\ & \sum_{\nu=0,1,2} Q_{\blacklozenge \nu}[\mathbf{u}] + K_0 \|\mathbf{u}\|_{\mathfrak{H}}^2 + K_1 \left\| \omega_b \frac{\partial \mathbf{u}}{\partial \phi} \right\|_{\mathfrak{H}}^2 \leq \\ & \leq \left(\frac{3K_0}{2} - 1\right) \|\mathbf{u}\|_{\mathfrak{H}}^2 + \frac{3K_1}{2} \left\| \omega_b \frac{\partial \mathbf{u}}{\partial \phi} \right\|_{\mathfrak{H}}^2. \end{aligned} \quad (7.115)$$

As for  $\mathcal{L}$  and  $Q_0$  we know (4.29):

$$\begin{aligned} & \frac{1}{2} \int \left(\frac{\Gamma P}{\rho^2}\right)_b |\operatorname{div}(\rho_b \mathbf{u})|^2 d\mathbf{x} \leq Q_0[\mathbf{u}] + \|\mathbf{u}\|_{\mathfrak{H}}^2 \leq \\ & \leq \frac{3}{2} \int \left(\frac{\Gamma P}{\rho^2}\right)_b |\operatorname{div}(\rho_b \mathbf{u})|^2 d\mathbf{x} + 2a \|\mathbf{u}\|_{\mathfrak{H}}^2, \end{aligned} \quad (7.116)$$

with the positive number  $a$  defined by (4.26). Therefore, putting

$$Q_{\blacklozenge}[\mathbf{u}] := Q_0^{\blacklozenge}[\mathbf{u}] + (a + K_0) \|\mathbf{u}\|_{\mathfrak{H}}^2 + K_1 \left\| \omega_b \frac{\partial \mathbf{u}}{\partial \phi} \right\|_{\mathfrak{H}}^2, \quad (7.117)$$

when

$$\begin{aligned} Q_{\blacklozenge}[\mathbf{u}] &= (K_1 - 1) \left\| \omega_b \frac{\partial \mathbf{u}}{\partial \phi} \right\|_{\mathfrak{H}}^2 + (a + K_0) \|\mathbf{u}\|_{\mathfrak{H}}^2 + Q_{\blacklozenge 1}[\mathbf{u}] + Q_{\blacklozenge 0}[\mathbf{u}] + Q_0[\mathbf{u}] \\ &\geq \frac{K_1}{2} \left\| \omega_b \frac{\partial \mathbf{u}}{\partial \phi} \right\|_{\mathfrak{H}}^2 + \frac{K_0 - 2}{2} \|\mathbf{u}\|_{\mathfrak{H}}^2 + \frac{1}{2} \int \left(\frac{\Gamma P}{\rho^2}\right)_b |\operatorname{div}(\rho_b \mathbf{u})|^2 d\mathbf{x} \geq 0, \end{aligned} \quad (7.118)$$

we have

$$\frac{1}{C} \|\mathbf{u}\|_{\mathfrak{G}_{\blacklozenge}}^2 \leq Q_{\blacklozenge}[\mathbf{u}] + \|\mathbf{u}\|_{\mathfrak{H}}^2 \leq C \|\mathbf{u}\|_{\mathfrak{G}_{\blacklozenge}}^2, \quad (7.119)$$

that is,  $[\|\cdot\|]_{\blacklozenge} := \sqrt{Q_{\blacklozenge} + \|\cdot\|_{\mathfrak{H}}^2}$  is equivalent to  $\|\cdot\|_{\mathfrak{G}_{\blacklozenge}}$ .

Although we do not know whether this operator  $\mathbf{L}_{\blacklozenge}$  is a closed operator in  $\mathfrak{H}$  or not, we can claim, at least, the following

**Proposition 11** *If a sequence  $(\mathbf{u}_n)_{n=0,1,2,\dots}$  in  $\mathbf{D}(\mathbf{L}_{\blacklozenge})$  converges to a limit  $\mathbf{u}$  in  $\mathfrak{H}$  and  $\mathcal{L}_{\blacklozenge 2} \mathbf{u}_n$  and  $\mathcal{L} \mathbf{u}_n$  converge in  $\mathfrak{H}$ , then the limit  $\mathbf{u}$  belongs to  $\mathbf{D}(\mathbf{L}_{\blacklozenge})$  and  $\mathbf{L}_{\blacklozenge} \mathbf{u}_n$  converge to  $\mathbf{L}_{\blacklozenge} \mathbf{u}$ .*

Proof. Let  $\mathcal{L}_{\blacklozenge 2} \mathbf{u}_n \rightarrow \mathbf{f}_{\blacklozenge 2}$ ,  $\mathcal{L} \mathbf{u} \rightarrow \mathbf{f}$  in  $\mathfrak{H}$ . Of course  $\mathcal{L}_{\blacklozenge 0} \mathbf{u}_n \rightarrow \mathcal{L}_{\blacklozenge 0} \mathbf{u}$  in  $\mathfrak{H}$ . Put  $\mathbf{f}_{\blacklozenge 0} := \mathcal{L}_{\blacklozenge 0} \mathbf{u}$ . Since  $\mathcal{L}_{\blacklozenge 2} \mathbf{u}_n$  are supposed to converge, we have

$$\begin{aligned} & - \left\| \omega_b \frac{\partial}{\partial \phi} (\mathbf{u}_{n+p} - \mathbf{u}_n) \right\|_{\mathfrak{H}}^2 = Q_{\blacklozenge 2} [\mathbf{u}_{n+p} - \mathbf{u}_n] \\ & = (\mathcal{L}_{\blacklozenge 2} \mathbf{u}_{n+p} - \mathcal{L}_{\blacklozenge 2} \mathbf{u}_n | \mathbf{u}_{n+p} - \mathbf{u}_n)_{\mathfrak{H}} \rightarrow 0 \end{aligned}$$

as  $n + p \geq n \rightarrow \infty$ . For any  $\psi \in C_0^\infty$ , we have

$$\begin{aligned} & |(\mathcal{L}_{\blacklozenge 1} \mathbf{u}_{n+p} - \mathcal{L}_{\blacklozenge 1} \mathbf{u}_n | \psi)_{\mathfrak{H}}| = |Q_{\blacklozenge 1} (\mathbf{u}_{n+p} - \mathbf{u}_n, \psi)| \\ & \leq 2|\Omega| \left\| \omega_b \frac{\partial}{\partial \phi} (\mathbf{u}_{n+p} - \mathbf{u}_n) \right\|_{\mathfrak{H}} \|\psi\|_{\mathfrak{H}}. \end{aligned}$$

Therefore

$$\|\mathcal{L}_{\blacklozenge 1} \mathbf{u}_{n+p} - \mathcal{L}_{\blacklozenge 1} \mathbf{u}_n\|_{\mathfrak{H}} \leq 2|\Omega| \left\| \omega_b \frac{\partial}{\partial \phi} (\mathbf{u}_{n+p} - \mathbf{u}_n) \right\|_{\mathfrak{H}} \rightarrow 0,$$

that is,  $\mathcal{L}_{\blacklozenge 1} \mathbf{u}_n$  converge in  $\mathfrak{H}$ . Put  $\mathbf{f}_{\blacklozenge 1} := \mathcal{L}_{\blacklozenge 1} \mathbf{u}$ . Since  $\mathcal{L} \mathbf{u}_n$  converge in  $\mathfrak{H}$ ,

$$Q_0 [\mathbf{u}_{n+p} - \mathbf{u}_n] = (\mathcal{L} \mathbf{u}_{n+p} - \mathcal{L} \mathbf{u}_n | \mathbf{u}_{n+p} - \mathbf{u}_n)_{\mathfrak{H}} \rightarrow 0.$$

Summing up, we have that  $Q_{\blacklozenge} [\mathbf{u}_{n+p} - \mathbf{u}_n] \rightarrow 0$  and  $[\|\mathbf{u}_{n+p} - \mathbf{u}_n\|_{\blacklozenge}]_{\blacklozenge} \rightarrow 0$ . This implies that  $\mathbf{u}_n$  converge in  $\mathfrak{G}_{\blacklozenge}$  to the limit  $\mathbf{u} \in \mathfrak{G}_{\blacklozenge}$ . Since  $\mathbf{u}_n \in (\mathfrak{G}_{\blacklozenge})_0$ ,  $\mathbf{u} \in (\mathfrak{G}_{\blacklozenge})_0$ . We see  $\mathbf{u} \in \mathcal{D}(\mathbf{L}_{\blacklozenge})$  and  $\mathcal{L}_{\blacklozenge} \mathbf{u}_n \rightarrow \mathcal{L} \mathbf{u} = \mathbf{f}_{\blacklozenge 2} + \mathbf{f}_{\blacklozenge 1} + \mathbf{f}_{\blacklozenge 0} + \mathbf{f}$ .  $\square$

If the condition could be weakend to that the total  $\mathbf{L}_{\blacklozenge} \mathbf{u}_n$  converge, then it would be told that  $\mathbf{L}_{\blacklozenge}$  is a closed operator, which is the case when  $\omega_b = 0$ .

Since we do not know whether  $\mathbf{L}_{\blacklozenge}$  is a self-adjoint operator in  $\mathfrak{H}$  or not, we try to consider its action on vector fields  $\mathbf{u}$  of the form  $\mathbf{u}(\mathbf{x}) = \check{\mathbf{u}}(r, \vartheta) e^{im\phi}$ , where  $m$  is a fixed integer.

Precisely speaking, we put

$$\check{\mathfrak{R}}_b := \{(r, \vartheta) | (r \sin \vartheta, 0, r \cos \vartheta) \in \mathfrak{R}_b\} \subset [0, +\infty[ \times \mathbb{R} / 2\pi \mathbb{Z}. \quad (7.120)$$

Since  $\mathfrak{R}_b$  is axially symmetric, it holds that

$$\check{\mathfrak{R}}_b = \{(r, \vartheta) | (r \sin \vartheta \cos \phi, r \sin \vartheta \sin \phi, r \cos \vartheta) \in \mathfrak{R}_b \quad \forall \phi\}.$$

While  $\rho_b(\mathbf{x}) = \rho_b^b(r, \vartheta)$ ,  $\check{\mathfrak{R}}_b = \{(r, \vartheta) | \rho_b^b(r, \vartheta) > 0\}$ .

Let  $m$  be an integer.

We denote by  $\mathfrak{H}_{|m} = \mathfrak{H}_{|m}(\mathfrak{R}_b; \mathbb{C})$  the set of all functions  $u \in \mathfrak{H} = L^2(\mathfrak{R}_b, \rho_b d\mathbf{x}; \mathbb{C})$  such that there exists a function  $\check{u}$  defined on  $\check{\mathfrak{R}}_b$  for which it holds that  $u(\mathbf{x}) = \check{u}(r, \vartheta) e^{im\phi}$ .  $\mathfrak{H}_{|m}(\mathfrak{R}_b; \mathbb{C}^3)$  is defined in the same manner. We denote  $\mathfrak{D}_{|m} := C_0^\infty(\mathfrak{R}_b) \cap \mathfrak{H}_{|m}$ . Clearly,  $u(\mathbf{x}) = \check{u}(r, \vartheta) e^{im\phi}$  belongs to  $\mathfrak{D}_{|m}$  if and only if there exists  $\check{u}^\sharp \in C_0^\infty(\check{\mathfrak{R}}_b^\sharp)$  such that  $\check{u}(r, \vartheta) = \check{u}^\sharp(r \sin \vartheta, 0, r \cos \vartheta)$ , where  $\check{\mathfrak{R}}_b^\sharp := \{(r \sin \vartheta, 0, r \cos \vartheta) | (r, \vartheta) \in \check{\mathfrak{R}}_b\}$ .

The following calculus can be easily verified:

- 1) If  $q \in \mathfrak{H}$  is axially symmetric, that is,  $\in \mathfrak{H}_{|0}$ , like  $\rho_b, S_b, P_b$ , and if  $\mathbf{u} \in \mathfrak{H}_{|m}$ ,  $q\mathbf{u} \in \mathfrak{H}$ , then  $q\mathbf{u} \in \mathfrak{H}_{|m}$  and  $(q\mathbf{u})^\sim = q^\flat \check{\mathbf{u}}$ .
- 2) If  $\mathbf{Q} \in \mathfrak{H}_{|m}$  and  $\operatorname{div} \mathbf{Q} \in \mathfrak{H}$ , then  $\operatorname{div} \mathbf{Q} \in \mathfrak{H}_{|m}$ . In fact

$$(\operatorname{div} \mathbf{Q})^\sim(r, \vartheta) = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \check{Q}^r) + \frac{1}{r \sin \vartheta} \frac{\partial}{\partial \vartheta} (\sin \vartheta \check{Q}^\vartheta) + \frac{im}{r \sin \vartheta} \check{Q}^\phi$$

for  $\check{\mathbf{Q}} = \check{Q}^r \mathbf{e}_r + \check{Q}^\vartheta \mathbf{e}_\vartheta + \check{Q}^\phi \mathbf{e}_\phi$ .

- 3) If  $\mathbf{Q} \in \mathfrak{H}_{|m}$  and  $\nabla \mathbf{Q} \in \mathfrak{H}$ , then  $\nabla \mathbf{Q} \in \mathfrak{H}_{|m}$ . In fact

$$(\nabla \mathbf{Q})^\sim = \frac{\partial \check{Q}}{\partial r} \mathbf{e}_r + \frac{1}{r} \frac{\partial \check{Q}}{\partial \vartheta} \mathbf{e}_\vartheta + \frac{im}{r \sin \vartheta} \check{Q} \mathbf{e}_\phi.$$

- 4) If  $\mathbf{Q} \in \mathfrak{H}_{|m}$ , then  $(\mathbf{Q} | \nabla S_b) \in \mathfrak{H}_{|m}$ . In fact

$$(\mathbf{Q} | \nabla S_b)^\sim = \check{Q}^r \frac{\partial S_b}{\partial r} + \check{Q}^\vartheta \frac{\partial S_b}{\partial \vartheta}.$$

- 5) If  $q \in C^\alpha(\mathfrak{R}_b \cap \partial \mathfrak{R}_b) \cap \mathfrak{H}_{|m}$ , then  $\mathcal{K}[q] \in \mathfrak{H}_{|m}$ . In fact

$$\mathcal{K}[q]^\sim(r, \vartheta) = \int \int_{\mathfrak{R}_b} \check{q}(r', \vartheta') K_{|m}(r, \vartheta, r', \vartheta') (r')^2 \sin \vartheta' dr' d\vartheta',$$

where

$$K_{|m}(r, \vartheta, r', \vartheta') := \frac{1}{2\pi} \int_0^\pi \frac{\cos(m\phi) d\phi}{\sqrt{r^2 + (r')^2 - 2rr'(\sin \vartheta \sin \vartheta' \cos \phi + \cos \vartheta \cos \vartheta')}}.$$

and so on.

Thanks to these calculus, we can claim that the operators

$$B_\diamond_{|m} := \mathcal{B}_\diamond \upharpoonright \mathfrak{H}_{|m}, \quad (7.121)$$

$$L_\diamond_{|m} := \mathcal{L}_\diamond \upharpoonright \mathfrak{H}_{|m} \quad (7.122)$$

can be considered as operators from dense subspace of  $\mathfrak{H}_{|m}$  into  $\mathfrak{H}_{|m}$ .

Let us consider  $\mathbf{u} \in \mathfrak{G}_{\diamond| m} := \mathfrak{G}_\diamond \cap \mathfrak{H}_{|m}$ . Then, since  $\frac{\partial \mathbf{u}}{\partial \phi} = im\mathbf{u}$ , we see

$$0 \leq Q_\diamond[\mathbf{u}] \leq Q_0^\diamond[\mathbf{u}] + a(|m|) \|\mathbf{u}\|_{\mathfrak{H}}^2, \quad (7.123)$$

where

$$a(|m|) := a + K_0 + K_1 \|\omega_b\|_{L^\infty}^2 |m|^2. \quad (7.124)$$

Thus  $Q_\diamond \upharpoonright \mathfrak{G}_{\diamond| m}$  is equivalent to the norm of

$$\mathfrak{G}_{\diamond| m} = \left\{ \mathbf{u} \in \mathfrak{H}_{|m} \mid \operatorname{div}(\rho_b \mathbf{u}) \in L^2 \left( \left( \frac{\Gamma P}{\rho^2} \right)_b dx \right) \right\}.$$



Of course, for  $\mathbf{u} \in \mathfrak{D}_{|m|}$ , we have

$$Q_0^\diamond[u] = (\mathcal{L}_\diamond \mathbf{u} | \mathbf{u})_{\mathfrak{H}} \geq -a(|m|) \|\mathbf{u}\|_{\mathfrak{H}}^2, \quad (7.125)$$

that is,  $\mathcal{L}_\diamond \upharpoonright \mathfrak{D}_{|m|}$  is bounded from below, and symmetric. Therefore  $\mathbf{L}_{\diamond|m|} := \mathbf{L} \upharpoonright \mathfrak{H}_{|m|}$  is self-adjoint, while

$$\mathcal{D}(\mathbf{L}_{\diamond|m|}) = \{\mathbf{u} \in (\mathfrak{G}_{\diamond|m|})_0 \mid \mathcal{L}_\diamond \mathbf{u} \in \mathfrak{H}\}.$$

Here  $(\mathfrak{G}_{\diamond|m|})_0$  stands for the closure of  $\mathfrak{D}_{|m|} = C_0^\infty(\mathfrak{R}_b) \cap \mathfrak{H}_{|m|}$  in  $\mathfrak{G}_{\diamond|m|}$ . Note that it holds  $(\mathfrak{G}_{\diamond|m|})_0 = (\mathfrak{G}_\diamond)_0 \cap \mathfrak{H}_{|m|}$ , and, for  $\mathbf{u} \in \mathfrak{G}_{\diamond|m|}$ , we have

$$\mathcal{L}_{\diamond 2} \mathbf{u} = -m^2 \omega_b^2 \mathbf{u}, \quad \mathcal{L}_{\diamond 1} \mathbf{u} = -2mi\Omega\omega_b \begin{bmatrix} -u^2 \\ u^1 \\ 0 \end{bmatrix},$$

which belong to  $\mathfrak{H}$  automatically.

On the other hand,  $\mathbf{B}_{\diamond|m|} = \mathcal{B}_\diamond \upharpoonright \mathfrak{H}_{|m|}$  is an operator in  $\mathfrak{H}_{|m|}$ . Actually, for  $\mathbf{v} \in \mathfrak{H}_{|m|}$ , it turns out to be that

$$\mathbf{B}_{\diamond|m|} \mathbf{v} = 2\Omega \times \mathbf{v} + 2im\omega_b \mathbf{v}.$$

therefore  $\mathbf{B}_{\diamond|m|}$  is a bounded linear operator in  $\mathfrak{H}_{|m|}$  with

$$\|\mathbf{B}_{\diamond|m|}\|_{\mathcal{B}(\mathfrak{H}_{|m|})} \leq \beta(|m|), \quad (7.126)$$

where

$$\beta(|m|) := 2|\Omega| + 2\|\omega_b\|_{L^\infty} |m|. \quad (7.127)$$

Consequently we can apply the Hille-Yosida theory to the initial value problem (7.95)(7.96) to claim:

**Proposition 12** *If  $\xi^0 \in \mathcal{D}(\mathbf{L}_{\diamond|m|})$ ,  $\mathbf{v}^0 \in (\mathfrak{G}_{\diamond|m|})_0$ ,  $\mathbf{f} \in C([0, +\infty[; \mathfrak{H}_{|m|})$ , then the problem (7.95)(7.96) admits a unique solution  $\xi = \xi(t, \mathbf{x})$  in*

$$C^2([0, +\infty[; \mathfrak{H}_{|m|}) \cap C^1([0, +\infty[; (\mathfrak{G}_{\diamond|m|})_0) \cap C([0, +\infty[; \mathcal{D}(\mathbf{L}_{\diamond|m|})).$$

Moreover the energy

$$E(t, \mathbf{u}) = \|\mathbf{u}\|_{\mathfrak{H}}^2 + Q_\diamond[\mathbf{u}] + \left\| \frac{\partial \mathbf{u}}{\partial t} \right\|_{\mathfrak{H}}^2$$

of the solution enjoys the inequality

$$\sqrt{E(t, \xi)} \leq e^{\kappa(|m|)t} \sqrt{E(0, \xi)} + \int_0^t e^{\kappa(|m|)(t-s)} \|\mathbf{f}(s)\|_{\mathfrak{H}} ds, \quad (7.128)$$

where

$$\kappa(|m|) := 1 + a(|m|) + \beta(|m|). \quad (7.129)$$

Note that

$$E(0, \xi) = (1 + a(|m|)) \|\xi^0\|_{\mathfrak{H}}^2 + (\mathbf{L}_\diamond \xi^0 | \xi^0)_{\mathfrak{H}} + \|v^0\|_{\mathfrak{H}}^2.$$

Now, given a positive integer  $m^\vee = 1, 2, \dots$ , we consider the Hilbert space  $\mathfrak{H}_{||m^\vee}$  of all linear combinations

$$\mathbf{u} = \sum_{|m| \leq m^\vee} \mathbf{u}_m = \sum_{|m| \leq m^\vee} \check{\mathbf{u}}_m(r, \vartheta) e^{im\phi},$$

where  $\mathbf{u}_m \in \mathfrak{H}_{|m|}$ . Put

$$\mathfrak{G}_{\diamond||m^\vee} := \left\{ \mathbf{u} \in \mathfrak{H}_{||m^\vee} \mid \operatorname{div}(\rho_b \mathbf{u}) \in L^2\left(\left(\frac{\Gamma P}{\rho^2}\right)_b dx\right) \right\}, \quad (7.130)$$

$$(\mathfrak{G}_{\diamond||m^\vee})_0 := \text{the closure of } C_0^\infty(\mathfrak{R}_b) \cap \mathfrak{H}_{||m^\vee} \text{ in } \mathfrak{G}_{\diamond||m^\vee}. \quad (7.131)$$

Then  $\mathbf{L}_{\diamond||m^\vee} := \mathbf{L}_\diamond \upharpoonright \mathfrak{H}_{||m^\vee}$  turns out to be a self-adjoint operator in  $\mathfrak{H}_{||m^\vee}$  with

$$\mathbf{D}(\mathbf{L}_{\diamond||m^\vee}) = \{\mathbf{u} \in (\mathfrak{G}_{\diamond||m^\vee})_0 \mid \mathcal{L}_\diamond \mathbf{u} \in \mathfrak{H}\}$$

and  $\mathbf{B}_{\diamond||m^\vee} := \mathbf{B}_\diamond \upharpoonright \mathfrak{H}_{||m^\vee}$  turns out to be a bounded linear operator in  $\mathfrak{H}_{||m^\vee}$ . We can claim the following

**Theorem 8** *Let  $m^\vee$  be a positive integer. If  $\xi^0 \in \mathbf{D}(\mathbf{L}_{\diamond||m^\vee})$ ,  $v^0 \in (\mathfrak{G}_{\diamond||m^\vee})_0$ ,  $\mathbf{f} \in C([0, +\infty[; \mathfrak{H}_{||m^\vee})$ , then the problem (7.95)(7.96) admits a unique solution  $\xi = \xi(t, \mathbf{x})$  in*

$$C^2([0, +\infty[; \mathfrak{H}_{||m^\vee}) \cap C^1([0, +\infty[; (\mathfrak{G}_{\diamond||m^\vee})_0) \cap C([0, +\infty[; \mathbf{D}(\mathbf{L}_{\diamond||m^\vee})).$$

Moreover the energy

$$E(t, \mathbf{u}) = \|\mathbf{u}\|_{\mathfrak{H}}^2 + Q_\diamond[\mathbf{u}] + \left\| \frac{\partial \mathbf{u}}{\partial t} \right\|_{\mathfrak{H}}^2$$

of the solution enjoys the inequality

$$\sqrt{E(t, \xi)} \leq e^{\kappa(m^\vee)t} \sqrt{E(0, \xi)} + \int_0^t e^{\kappa(m^\vee)(t-s)} \|\mathbf{f}(s)\|_{\mathfrak{H}} ds. \quad (7.132)$$

Proof. First we note that, if  $m \neq m'$ ,  $\mathbf{u} \in C^1([0, +\infty[; \mathfrak{G}_{\diamond|m})$ ,  $\mathbf{u}' \in C^1([0, +\infty[; \mathfrak{G}_{\diamond|m'})$ , then  $(\mathbf{u} | \mathbf{u}')_{\mathfrak{H}} = 0$ ,  $Q_\diamond(\mathbf{u}, \mathbf{u}') = 0$ ,  $(\partial_t \mathbf{u} | \partial_t \mathbf{u}')_{\mathfrak{H}} = 0$ , therefore  $E(t, \mathbf{u} + \mathbf{u}') = E(t, \mathbf{u}) + E(t, \mathbf{u}')$ . Therefore the assertion is a direct consequence of Proposition 12, since  $\kappa(|m|) \leq \kappa(m^\vee)$  for  $|m| \leq m^\vee$ .  $\square$

When  $\|\omega_b\|_{L^\infty} \neq 0$ , then we see  $\kappa(m^\vee)t \geq 2\|\omega_b\|_{L^\infty}^2(m^\vee)^2 t \rightarrow \infty$  as  $m^\vee \rightarrow \infty$  for  $\forall t > 0$ . Therefore we can not deduce from this theorem the existence of solutions to the problem (7.95)(7.96) for non-trivial data  $(\xi^0, v^0) \in \mathbf{D}(\mathbf{L}_\diamond) \times$

$(\mathfrak{G}_\diamond)_0$  by considering the sequence of the solutions  $\boldsymbol{\xi} = \boldsymbol{\xi}_{m^\vee}(t, \mathbf{x})$  to the data  $(\boldsymbol{\xi}_{m^\vee}^0, \mathbf{v}_{m^\vee}^0) \in \mathcal{D}(\mathbf{L}_\diamond \|_{m^\vee}) \times (\mathfrak{G}_\diamond \|_{m^\vee})_0$  which approximate the data  $(\boldsymbol{\xi}^0, \mathbf{v}^0)$  and passing to the limit of  $m^\vee \rightarrow \infty$ .

## Appendix A

We consider the initial value problem:

$$\begin{aligned} \frac{d^2 \mathbf{u}}{dt^2} + \mathbf{B} \frac{d\mathbf{u}}{dt} + \mathbf{L}\mathbf{u} &= 0, \\ \mathbf{u}(0) &= \mathring{\mathbf{u}}, \quad \frac{d\mathbf{u}}{dt}(0) = \mathring{\mathbf{v}}. \end{aligned}$$

Here the unknown is a function  $\mathbf{u} : t \mapsto \mathbf{u}(t) : [0, +\infty[ \rightarrow \mathbf{X}$ ,  $\mathbf{X}$  being a Hilbert space endowed with an inner product  $(\cdot | \cdot)_{\mathbf{X}}$ .

We assume:

$\mathbf{Y}$  is a Hilbert space endowed with an inner product  $(\cdot | \cdot)_{\mathbf{Y}}$ , which is continuously and densely imbedded in  $\mathbf{X}$ .  $\mathbf{Y}_0$  is a closed subspace of  $\mathbf{Y}$ .

$\mathbf{L}$  is a self-adjoint operator in  $\mathbf{X}$  whose domain  $\mathcal{D}(\mathbf{L})$  is included in  $\mathbf{Y}_0$ .

There is a quadratic form  $Q$  in  $\mathbf{Y}$  such that

$$\begin{aligned} (\mathbf{u}_1 | \mathbf{u}_2)_{\mathbf{Y}} &= (\mathbf{u}_1 | \mathbf{u}_2)_{\mathbf{X}} + Q(\mathbf{u}_1, \mathbf{u}_2) \quad \text{for } \forall \mathbf{u}_1, \mathbf{u}_2 \in \mathbf{Y}, \\ Q[\mathbf{u}] &= Q(\mathbf{u}, \mathbf{u}) \geq 0 \quad \text{for } \forall \mathbf{u} \in \mathbf{Y}, \end{aligned}$$

and

$$(\mathbf{L}\mathbf{u}_1 | \mathbf{u}_2)_{\mathbf{X}} + a(\mathbf{u}_1 | \mathbf{u}_2)_{\mathbf{X}} = Q(\mathbf{u}_1, \mathbf{u}_2) \quad \text{for } \forall \mathbf{u}_1 \in \mathcal{D}(\mathbf{L}), \forall \mathbf{u}_2 \in \mathbf{Y}_0.$$

Here  $a$  is a non-negative number.

$\mathbf{B}$  is a bounded linear operator in  $\mathbf{X}$  such that  $\|\mathbf{B}\|_{\mathcal{B}(\mathbf{X})} \leq \beta$ .

**Proposition A1** *For any  $c \in \mathbb{R}$  and  $\lambda > a + |c|\beta$  the operator  $\mathbf{L} + c\mathbf{B} + \lambda$  has the bounded linear inverse operator  $(\mathbf{L} + c\mathbf{B} + \lambda)^{-1}$  defined on the whole space  $\mathbf{X}$  such that*

$$\|(\mathbf{L} + c\mathbf{B} + \lambda)^{-1}\|_{\mathcal{B}(\mathbf{X})} \leq \frac{1}{\lambda - a - |c|\beta}.$$

Proof. First we see  $\mathbf{L} + c\mathbf{B} + \lambda$  is invertible. In fact, if

$$(\mathbf{L} + c\mathbf{B} + \lambda)\mathbf{u} = \mathbf{f}, \quad \mathbf{u} \in \mathcal{D}(\mathbf{L}), \quad \mathbf{f} \in \mathbf{X},$$

then we see

$$(\lambda - a - |c|\beta)\|\mathbf{u}\|_{\mathbf{X}}^2 \leq Q[\mathbf{u}] + (\lambda - a)\|\mathbf{u}\|_{\mathbf{X}} + (c\mathbf{B}\mathbf{u} | \mathbf{u})_{\mathbf{X}} + \lambda\|\mathbf{u}\|_{\mathbf{X}}^2 = (\mathbf{u} | \mathbf{f})_{\mathbf{X}} \leq \|\mathbf{u}\|_{\mathbf{X}} \|\mathbf{f}\|_{\mathbf{X}},$$

therefore we have

$$\|\mathbf{u}\|_{\mathbf{X}} \leq \frac{1}{\lambda - a - |c|\beta} \|\mathbf{f}\|_{\mathbf{X}}.$$

We claim that the range  $\mathbf{R}(\mathbf{L} + c\mathbf{B} + \lambda)$  is dense in  $\mathbf{X}$ . In fact, suppose

$$((\mathbf{L} + c\mathbf{B} + \lambda)\mathbf{u}|\mathbf{f}) = 0 \quad \forall \mathbf{u} \in \mathbf{D}(\mathbf{L}).$$

Then

$$\begin{aligned} (\mathbf{L}\mathbf{u}|\mathbf{f}) &= -((c\mathbf{B} + \lambda)\mathbf{u}|\mathbf{f}) \\ &= (\mathbf{u}|(-c\mathbf{B}^* - \lambda)\mathbf{f}) \end{aligned}$$

for  $\forall \mathbf{u} \in \mathbf{D}(\mathbf{L})$ . Hence  $\mathbf{f} \in \mathbf{D}(\mathbf{L}^*)$  and

$$\mathbf{L}^*\mathbf{f} = -c\mathbf{B}^* - \lambda\mathbf{f}.$$

Since  $\mathbf{L} = \mathbf{L}^*$ , this means that  $\mathbf{f} \in \mathbf{D}(\mathbf{L})$  and

$$(\mathbf{L} + c\mathbf{B}^* + \lambda)\mathbf{f} = 0.$$

Since  $\mathbf{L} + c\mathbf{B}^* + \lambda$  is invertible, for  $\|\mathbf{B}^*\| = \|\mathbf{B}\| \leq \beta$ , it follows that  $\mathbf{f} = 0$ . Summing up, we have the assertion.  $\square$

We are going to apply the Hille-Yosida theory to the initial-boundary value problem (1) :

$$\begin{aligned} \frac{d^2\mathbf{u}}{dt^2} + \mathbf{B}\frac{d\mathbf{u}}{dt} + \mathbf{L}\mathbf{u} &= \mathbf{0}, \\ \mathbf{u}(t) &\in \mathbf{D}(\mathbf{L}) \quad \text{for } \forall t \geq 0, \\ \mathbf{u} &= \mathring{\mathbf{u}}, \quad \frac{d\mathbf{u}}{dt} = \mathring{\mathbf{v}} \quad \text{at } t = 0, \end{aligned}$$

We put

$$\begin{aligned} U &= \begin{bmatrix} \mathbf{u} \\ \dot{\mathbf{u}} \end{bmatrix}, \quad \dot{\mathbf{u}} = \frac{d\mathbf{u}}{dt}, \\ \mathbf{A}U &= \begin{bmatrix} O & -I \\ \mathbf{L} & \mathbf{B} \end{bmatrix} U = \begin{bmatrix} -\dot{\mathbf{u}} \\ \mathbf{B}\dot{\mathbf{u}} + \mathbf{L}\mathbf{u} \end{bmatrix}, \\ Z &= \mathbf{Y}_0 \times \mathbf{X} \end{aligned}$$

with

$$\begin{aligned} (U_1|U_2)_Z &= (\mathbf{u}_1|\mathbf{u}_2)_{\mathbf{Y}} + (\dot{\mathbf{u}}_1|\dot{\mathbf{u}}_2)_{\mathbf{X}} = \\ &= Q(\mathbf{u}_1, \mathbf{u}_2) + (\mathbf{u}_1|\mathbf{u}_2)_{\mathbf{X}} + (\dot{\mathbf{u}}_1|\dot{\mathbf{u}}_2)_{\mathbf{X}}, \\ \mathbf{D}(\mathbf{A}) &= \mathbf{D}(\mathbf{L}) \times \mathbf{Y}_0. \end{aligned}$$

Then the initial-boundary value problem (1) can be written as the problem (0):

$$\frac{dU}{dt} + \mathbf{A}U = \mathbf{0}, \quad U|_{t=0} = U_0,$$

where

$$U_0 = \begin{bmatrix} \mathring{\mathbf{u}} \\ \mathring{\mathbf{v}} \end{bmatrix}.$$

Applying [4, Theorem 7.4], we can claim

**Proposition A2** *If  $U_0 \in \mathbf{D}(\mathbf{A})$ , say, if  $\mathring{\mathbf{u}} \in \mathbf{D}(\mathbf{L})$  and  $\mathring{\mathbf{v}} \in \mathbf{Y}_0$ , then there exists a unique solution  $U \in C^1([0, +\infty[, \mathbf{Z}) \cap C([0, +\infty[, \mathbf{D}(\mathbf{A}))$  to the problem (0). Moreover  $E(t) = \|U(t)\|_{\mathbf{Z}}^2$  enjoys*

$$\sqrt{E(t)} \leq e^{\Lambda t} \sqrt{E(0)},$$

where  $\Lambda = 1 + a + \beta$ .

Here we consider that  $\mathbf{D}(\mathbf{A})$  is equipped with the operator norm  $(\|U\|_{\mathbf{Z}}^2 + \|\mathbf{A}U\|_{\mathbf{Z}}^2)^{1/2}$ .

Proof of Proposition A2. Firstly  $\mathbf{A} + 1 + a + \beta$  is monotone, that is, for  $\forall U \in \mathbf{D}(\mathbf{A})$  we have

$$\begin{aligned} \Re[(\mathbf{A}U|U)_{\mathbf{Z}}] + (1 + a + \beta)\|U\|_{\mathbf{Z}}^2 &= \Re \left[ -Q(\dot{\mathbf{u}}, \mathbf{u}) - (\dot{\mathbf{u}}|\mathbf{u})_{\mathbf{X}} + (\mathbf{L}\mathbf{u}|\dot{\mathbf{u}})_{\mathbf{X}} + (\mathbf{B}\dot{\mathbf{u}}|\dot{\mathbf{u}})_{\mathbf{X}} \right] + \\ &\quad + (1 + a + \beta)(Q[\mathbf{u}] + \|\mathbf{u}\|_{\mathbf{X}}^2 + \|\dot{\mathbf{u}}\|_{\mathbf{X}}^2) \\ &\geq -(1 + a)\Re[(\dot{\mathbf{u}}|\mathbf{u})_{\mathbf{X}}] - \beta\|\dot{\mathbf{u}}\|_{\mathbf{X}}^2 + \\ &\quad + (1 + a + \beta)\|\mathbf{u}\|_{\mathbf{X}}^2 + (1 + a + \beta)\|\dot{\mathbf{u}}\|_{\mathbf{X}}^2 \\ &\geq (1 + a) \left[ \|\dot{\mathbf{u}}\|_{\mathbf{X}}^2 - \Re[(\dot{\mathbf{u}}|\mathbf{u})_{\mathbf{X}}] + \|\mathbf{u}\|_{\mathbf{X}}^2 \right] \\ &\geq 0, \end{aligned}$$

since  $((\mathbf{L} + a)\mathbf{u}|\dot{\mathbf{u}}) = Q(\mathbf{u}, \dot{\mathbf{u}}) = Q(\dot{\mathbf{u}}, \mathbf{u})^*$  and  $|\Re[(\mathbf{B}\dot{\mathbf{u}}|\dot{\mathbf{u}})_{\mathbf{X}}]| \leq \beta\|\dot{\mathbf{u}}\|_{\mathbf{X}}^2$ .

If  $\Lambda > \frac{\beta}{2} + \sqrt{\frac{\beta^2}{4}} + a$ , then the operator  $\mathbf{A} + \Lambda$  has the bounded inverse defined on  $\mathbf{Z}$ . Actually the equation

$$\mathbf{A}U + \Lambda U = F = \begin{bmatrix} \mathbf{f} \\ \mathbf{g} \end{bmatrix} \in \mathbf{Z}$$

means

$$\begin{cases} -\dot{\mathbf{u}} + \Lambda \mathbf{u} = \mathbf{f} \in \mathbf{Y}_0 \\ \mathbf{B}\dot{\mathbf{u}} + \mathbf{L}\mathbf{u} + \Lambda \dot{\mathbf{u}} = \mathbf{g} \in \mathbf{X}, \end{cases}$$

which can be solved as

$$\begin{cases} \mathbf{u} = (\mathbf{L} + \Lambda \mathbf{B} + \Lambda^2)^{-1}(\mathbf{B}\mathbf{f} + \Lambda \mathbf{f} + \mathbf{g}) \in \mathbf{D}(\mathbf{L}), \\ \dot{\mathbf{u}} = (\mathbf{L} + \Lambda \mathbf{B} + \Lambda^2)^{-1}(\mathbf{B}\mathbf{f} + \Lambda \mathbf{f} + \mathbf{g}) - \mathbf{f} \in \mathbf{Y}_0, \end{cases}$$

thanks to Proposition A1, since  $\Lambda^2 > a + |\Lambda|\beta$  holds for  $\Lambda > \frac{\beta}{2} + \sqrt{\frac{\beta^2}{4} + a}$ .  $\square$

Therefore, considering the problem (1):

$$\begin{aligned} \frac{d^2 \mathbf{u}}{dt^2} + \mathbf{B} \frac{d\mathbf{u}}{dt} + \mathbf{L}\mathbf{u} &= \mathbf{0}, \\ \mathbf{u}(t) &\in \mathbf{D}(\mathbf{L}) \quad \text{for } \forall t \geq 0, \\ \mathbf{u} &= \mathring{\mathbf{u}}, \quad \frac{d\mathbf{u}}{dt} = \mathring{\mathbf{v}} \quad \text{at } t = 0, \end{aligned}$$

we can claim

**Theorem A1** *Suppose  $\mathring{\mathbf{u}} \in \mathbf{D}(\mathbf{L})$  and  $\mathring{\mathbf{v}} \in \mathbf{Y}_0$ . Then the initial-boundary value problem (1) admits a unique solution*

$$\mathbf{u} \in C^2([0, +\infty[, \mathbf{X}) \cap C^1([0, +\infty[, \mathbf{Y}_0) \cap C([0, +\infty[, \mathbf{D}(\mathbf{L}))$$

and the energy

$$\begin{aligned} E(t) &= \|\mathbf{u}\|_{\mathbf{Y}}^2 + \|\dot{\mathbf{u}}\|_{\mathbf{X}}^2 \\ &= \|\mathbf{u}\|_{\mathbf{X}}^2 + Q[\mathbf{u}] + \left\| \frac{d\mathbf{u}}{dt} \right\|_{\mathbf{X}}^2 \end{aligned}$$

enjoys the estimate

$$\sqrt{E(t)} \leq e^{\Lambda t} \cdot \sqrt{E(0)},$$

where  $\Lambda = 1 + a + \beta$ .

Here  $\mathbf{D}(\mathbf{L})$  is equipped with the norm  $(\|\mathbf{u}\|_{\mathbf{Y}}^2 + \|\mathbf{L}\mathbf{u}\|_{\mathbf{X}}^2)^{1/2}$ .

Correspondingly we may consider the inhomogeneous initial-boundary value problem (2):

$$\frac{dU}{dt} + \mathbf{A}U = F(t), \quad U|_{t=0} = U_0.$$

We can claim

**Proposition A3** *If  $U_0 \in \mathbf{D}(\mathbf{A})$  and  $F \in C([0, +\infty[; \mathbf{Z})$ , then there exists a unique solution*

$$U \in C^1([0, +\infty[; \mathbf{Z}) \cap C([0, +\infty[; \mathbf{D}(\mathbf{A}))$$

to the problem **(2)** , and it enjoys the estimate

$$\|U(t)\|_{\mathbf{Z}} \leq e^{\Lambda t} \left( \|U_0\|_{\mathbf{Z}} + \int_0^t e^{-\Lambda s} \|F(s)\|_{\mathbf{Z}} ds \right),$$

where  $\Lambda = 1 + a + \beta$ .

Therefore, considering the problem **(3)**:

$$\begin{aligned} \frac{d^2 \mathbf{u}}{dt^2} + \mathbf{B} \frac{d\mathbf{u}}{dt} + \mathbf{L}\mathbf{u} &= \mathbf{f}(t, \mathbf{x}), \\ \mathbf{u} &= \mathring{\mathbf{u}}, \quad \frac{d\mathbf{u}}{dt} = \mathring{\mathbf{v}} \quad \text{at } t = 0, \\ \mathbf{u}(t) &\in \mathbf{D}(\mathbf{L}) \quad \text{for } \forall t \geq 0., \end{aligned}$$

we can claim:

**Theorem A2** Suppose  $\mathring{\mathbf{u}} \in \mathbf{D}(\mathbf{L})$ ,  $\mathring{\mathbf{v}} \in \mathbf{Y}_0$  and  $\mathbf{f} \in C([0, +\infty[; \mathbf{X})$ . Then the initial-boundary value problem **(3)** admits a unique solution

$$\mathbf{u} \in C^2([0, +\infty[; \mathbf{X}) \cap C^1([0, +\infty[; \mathbf{Y}_0) \cap C([0, +\infty[; \mathbf{D}(\mathbf{L}))$$

and the energy

$$\begin{aligned} E(t) = E(t, \mathbf{u}) &:= \|\mathbf{u}\|_{\mathbf{Y}}^2 + \|\dot{\mathbf{u}}\|_{\mathbf{X}}^2 \\ &= \|\mathbf{u}\|_{\mathbf{X}}^2 + Q[\mathbf{u}] + \left\| \frac{d\mathbf{u}}{dt} \right\|_{\mathbf{X}}^2 \end{aligned}$$

enjoys the estimate

$$\sqrt{E(t)} \leq e^{\Lambda t} \left( \sqrt{E(0)} + \int_0^t e^{-\Lambda s} \|\mathbf{f}(s)\|_{\mathbf{X}} ds \right)$$

for  $\Lambda = 1 + a + \beta$ .

## Appendix B

Let us imagine the motion  $\mathbf{x} = \mathbf{x}_p(t)$  of an infenitesimally small percel in the background  $(\rho_b, S_b)$ . The density  $\rho_p(\mathbf{x}(t))$ , pressure  $P_p(\mathbf{x}(t))$ , entropy density  $S_p(\mathbf{x}(t))$  of the percel obeys the same EOS:  $P = \mathcal{P}(\rho, S)$  as the background. Suppose that  $P_p(\mathbf{x}(t)) = P_b(\mathbf{x}(t))$ .

Here we note the following: Let  $\rho = \mathcal{R}(\rho, S)$  be the inverse function of  $P = \mathcal{P}(\rho, S)$ , so

$$\rho = \mathcal{R}(\mathcal{P}(\rho, S), S);$$

then it holds

$$\begin{aligned}\partial_S \mathcal{R}(P, S) &= -(\partial_\rho \mathcal{R}(P, S))(\partial_S \mathcal{P}(\mathcal{R}(P, S), S)) \\ &= -\frac{\partial_S \mathcal{P}(\rho, S)}{\partial_\rho \mathcal{P}(\rho, S)} \Big|_{\rho=\mathcal{R}(P, S)};\end{aligned}$$

therefore, by Definition 3, (2.10b), we have

$$\partial_S \mathcal{R}(P, S) \nabla S = \rho \mathbf{a}.$$

Now the equation of motion of the parcel is

$$\frac{d^2 \mathbf{x}_p}{dt^2} = -\frac{\nabla P_b}{\rho_p} + \mathbf{F}(t, \mathbf{x}_p(t)),$$

while the background satisfies

$$0 = -\frac{\nabla P_b}{\rho_b} + \mathbf{F}(t, \mathbf{x})$$

for some vector field of force. ( In our situation, we are concerned with

$$\mathbf{F} = -\nabla \Phi_b - (D\mathbf{v}_b)\mathbf{v}_b - 2\boldsymbol{\Omega} \times \mathbf{v}_b - \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{x}).$$

But we need not the specification of  $\mathbf{F}$ , wich contains both real and virtual forces.) Therefore the equation of motion is

$$\frac{d^2 \mathbf{x}_p}{dt^2} = \left( \frac{1}{\rho_b} - \frac{1}{\rho_p} \right) \nabla P_b(\mathbf{x}_p(t)).$$

On the other hand we have approximately

$$\begin{aligned}\frac{1}{\rho_b} - \frac{1}{\rho_p} &\cong \frac{1}{\rho_b^2} \partial_S \mathcal{R}(P_b, S_b) (\nabla S_b | \mathbf{x}_p(t) - \mathbf{x}_p(0)) \\ &= -\frac{1}{\rho_b} (\mathbf{a}_b | \mathbf{x}_p(t) - \mathbf{x}_p(0)).\end{aligned}$$

Since  $\nabla P_b$  is parallel to  $\mathbf{n} \approx \mathbf{n}_0 = \mathbf{n}(\mathbf{x}_p(0))$ , we suppose that

$$\mathbf{x}_p(t) = X(t) \mathbf{n}_0.$$

Then

$$\frac{1}{\rho_b} - \frac{1}{\rho_p} \cong \frac{1}{\rho_b} \mathcal{A}_b \Big|_{\mathbf{x}=\mathbf{x}_p(0)} (X(t) - X(0),$$

and the equation of motion is

$$\begin{aligned}\frac{d^2 X}{dt^2} &\cong \frac{1}{\rho_b} \mathcal{A}_b (\nabla P_b | \mathbf{n}_0) (X(t) - X(0)) \\ &\cong -\mathcal{N}_b^2 \Big|_{\mathbf{x}=\mathbf{x}_p(0)} (X(t) - X(0)).\end{aligned}$$



The solution of this equation is

$$X(t) = X(0) + C_+ e^{iNt} + C_- e^{-iNt} \quad \text{where} \quad N := \mathcal{N}_b(\mathbf{x}_p(0)).$$

Clearly  $|X(t) - X(0)| = O(1)$  for non-trivial  $X(t)$  if and only if  $N \in \mathbb{R}$ . Therefore the definition (2.15) is justifiable.

## Acknowledgment

To do this study has been caused by helpful discussions with Professors Akitaka Matsumura (Osaka University) and Masao Takata (the University of Tokyo), to whom sincere thanks are expressed. This work was supported by Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for Scientific Research (KAKENHI) grant JP21K03311, and by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University.

## The data availability statement

No new data were created or analyzed in this study.

## References

- [1] G. K. Batchelor, An Introduction to Fluid Dynamics, Cambridge UP, 1967.
- [2] V. Bjerknes, Über die hydrodynamischen Gleichungen in Lagrangescher und Eulerscher Form und ihre Linearisierung für das Studium kleiner Störungen, Geof. Publ., 5(1929), 3-43.
- [3] J. Bognar, Indefinite inner-product spaces, Springer, Berlin-Hidelberg-New York, 1974.
- [4] H. Brezis, Functional Analysis, Sobolev Spaces and Partial Differential Equations, Springer, New York-Dordrecht-Heidelberg-London, 2011.
- [5] D. Brunt, Physical and Dynamical Meteorology, Cambridge Univ. Press, 1934.

- [6] J. Dyson and B. F. Schutz, Perturbation and stability of rotating stars, I. Completeness of normal modes, *Proc. Roy. Soc. London, A* 368(1979), 389-410.
- [7] J. L. Friedman and B. F. Schutz, Lagrangian perturbation theory of non-relativistic fluids, *Astrophys. J.*, 221(1978), 937-957.
- [8] J. R. Holton, *An Introduction to Dynamical Meteorology*, 5th ed., Academic Press, 2013.
- [9] C. Hunter, On secular stability, secular instability, and points of bifurcation of rotating gaseous masses, *Astrophys. J.*, 213(1977), 497-517.
- [10] Juhi Jang and T. Makino, On rotating axially symmetric solutions of the Euler-Poisson equations, *J. Differential equations*, 266(2019), 3942-3972.
- [11] Juhi Jang and T. Makino, Linearized analysis of barotropic perturbations around spherically symmetric gaseous stars governed by the Euler-Poisson equations, *J. Math. Phys.*, 61(2020), 051508.
- [12] T. Kato, *Perturbation Theory for Linear Operators*, Springer, 1980.
- [13] N. R. Lebovitz, The virial tensor and its application to self-gravitating fluids, *Astrophys. J.*, 134(1961), 500-536.
- [14] P. Ledoux and Th. Walraven, Variable stars, in *Handbuch der Physik*, Band LI : Sternaufbau, Springer, Berlin, Göttingen, Heidelberg, 1958.
- [15] D. Lynden-Bell and J. P. Ostriker, On the stability of differentially rotating bodies, *Mon. Not. Astr. Soc.*, 136(1967), 293-310.
- [16] T. Makino, On linear adiabatic perturbations of spherically symmetric gaseous stars governed by the Euler-Poisson equations, *Kyoto J. Math.*, 63(2023), 353-420.
- [17] R. Wavre, *Figures Planétaires et Géodésie*, Gauthier-Villard, Paris, 1932.