

On blow-up conditions for nonlinear higher order evolution inequalities

A. A. Kon'kov^{1,2*} and A. E. Shishkov^{2†}

^{1*}Department of Differential Equations, Faculty of Mechanics and Mathematics, Lomonosov Moscow State University, Vorobyovy Gory, Moscow, 119991, Russia.

²Center of Nonlinear Problems of Mathematical Physics, RUDN University, Miklukho-Maklaya str. 6, Moscow, 117198, Russia.

*Corresponding author(s). E-mail(s): konkov@mech.math.msu.su;

Contributing authors: aeshkv@yahoo.com;

†These authors contributed equally to this work.

Abstract

For the problem

$$\begin{cases} \partial_t^k u - \sum_{|\alpha|=m} \partial^\alpha a_\alpha(x, t, u) \geq f(|u|) & \text{in } \mathbb{R}_+^{n+1} = \mathbb{R}^n \times (0, \infty), \\ u(x, 0) = u_0(x), \partial_t u(x, 0) = u_1(x), \dots, \partial_t^{k-1} u(x, 0) = u_{k-1}(x) \geq 0, \end{cases}$$

where $u_i \in L_{1,loc}(\mathbb{R}^n)$ and a_α are Caratheodory functions such that

$$|a_\alpha(x, t, \zeta)| \leq A|\zeta|^p, \quad A, p = \mathit{const} > 0,$$

for almost all $(x, t) \in \mathbb{R}_+^{n+1}$ and for all $\zeta \in \mathbb{R}$, we obtain exact conditions on the function f guaranteeing that any global weak solution is identically zero.

Keywords: Higher order differential evolution inequalities, Nonlinearity, Blow-up

MSC Classification: 35B44 , 35B08 , 35J30 , 35J70

1 Introduction

We study solutions of the inequality

$$\partial_t^k u - \mathcal{L}u \geq f(|u|) \quad \text{in } \mathbb{R}_+^{n+1} = \mathbb{R}^n \times (0, \infty) \quad (1)$$

satisfying the initial conditions

$$u(x, 0) = u_0(x), \partial_t u(x, 0) = u_1(x), \dots, \partial_t^{k-1} u(x, 0) = u_{k-1}(x) \geq 0, \quad (2)$$

where $k \geq 1$ and $n \geq 1$ are integers and

$$\mathcal{L}u = \sum_{|\alpha|=m} \partial^\alpha a_\alpha(x, t, u)$$

is a differential operator of order $m \geq 1$ whose coefficients are Caratheodory functions such that

$$|a_\alpha(x, t, \zeta)| \leq A|\zeta|^p, \quad |\alpha| = m,$$

with some constants $A > 0$ and $p > 0$ for almost all $(x, t) \in \mathbb{R}_+^{n+1}$ and for all $\zeta \in \mathbb{R}$. As is customary, by α we mean a multi-index $\alpha = (\alpha_1, \dots, \alpha_n)$. In so doing, $|\alpha| = \alpha_1 + \dots + \alpha_n$ and $\partial^\alpha = \partial^{|\alpha|}/(\partial_{x_1}^{\alpha_1} \dots \partial_{x_n}^{\alpha_n})$.

Let us denote by B_r the open ball in \mathbb{R}^n of radius $r > 0$ and center at zero. Also let $Q_r^\varkappa = B_r \times (0, r^\varkappa)$, $\varkappa > 0$. We say that a function u belongs to $L_{\lambda,loc}(\mathbb{R}^n)$, $0 < \lambda \leq \infty$, if $u \in L_\lambda(B_r)$ for all real numbers $r > 0$. In its turn, $u \in L_{\lambda,loc}(\mathbb{R}_+^{n+1})$ if $u \in L_\lambda(Q_r^1)$ for all real numbers $r > 0$.

We assume that $u_i \in L_{1,loc}(\mathbb{R}^n)$, $i = 0, \dots, k-1$. Also let $f(\zeta)$ and $f(\zeta^{1/p})$ are non-decreasing convex functions on the interval $[0, \infty)$ such that $f(\zeta) > 0$ for all $\zeta \in (0, \infty)$.

A function $u \in L_{1,loc}(\mathbb{R}_+^{n+1}) \cap L_{p,loc}(\mathbb{R}_+^{n+1})$ is called a global weak solution of problem (1), (2) if $f(|u|) \in L_{1,loc}(\mathbb{R}_+^{n+1})$ and

$$\begin{aligned} & \int_{\mathbb{R}_+^{n+1}} (-1)^k u \partial_t^k \varphi dx dt - \int_{\mathbb{R}_+^{n+1}} \sum_{|\alpha|=m} (-1)^m a_\alpha(x, t, u) \partial^\alpha \varphi dx dt \\ & \geq \int_{\mathbb{R}^n} \sum_{i=0}^{k-1} (-1)^i u_{k-1-i}(x) \partial_t^i \varphi(x, 0) dx + \int_{\mathbb{R}_+^{n+1}} f(|u|) \varphi dx dt \end{aligned} \quad (3)$$

for any non-negative function $\varphi \in C_0^\infty(\mathbb{R}^{n+1})$. Analogously, a function $u \in L_{1,loc}(\mathbb{R}_+^{n+1}) \cap L_{p,loc}(\mathbb{R}_+^{n+1})$ is a global weak solution of the equation

$$\partial_t^k u - \mathcal{L}u = f(|u|) \quad \text{in } \mathbb{R}_+^{n+1} \quad (4)$$

satisfying conditions (2) if $f(|u|) \in L_{1,loc}(\mathbb{R}_+^{n+1})$ and

$$\begin{aligned} & \int_{\mathbb{R}_+^{n+1}} (-1)^k u \partial_t^k \varphi dx dt - \int_{\mathbb{R}_+^{n+1}} \sum_{|\alpha|=m} (-1)^m a_\alpha(x, t, u) \partial^\alpha \varphi dx dt \\ &= \int_{\mathbb{R}^n} \sum_{i=0}^{k-1} (-1)^i u_{k-1-i}(x) \partial_t^i \varphi(x, 0) dx + \int_{\mathbb{R}_+^{n+1}} f(|u|) \varphi dx dt \end{aligned}$$

for any $\varphi \in C_0^\infty(\mathbb{R}^{n+1})$. It is obvious that every global weak solution of (4), (2) is also a global weak solution of (1), (2).

As an example of (4), we can take the nonlinear diffusion equation

$$u_t = \Delta u^p + f(u) \quad \text{in } \mathbb{R}_+^{n+1}$$

which can be represented in the form

$$u_t = \operatorname{div}(D \nabla u) + f(u) \quad \text{in } \mathbb{R}_+^{n+1}, \quad (5)$$

where $D = pu^{p-1}$ is a diffusion coefficient depending on the density $u > 0$ in a power-law manner and $f(u)$ is a source-density function. In the case of $p \geq 1$, it is customary to say that (5) is the slow diffusion equation. In its turn, if $0 < p < 1$, then one says that (5) is the fast diffusion equation.

The absence of non-trivial solutions of differential equations and inequalities or, in other words, the blow-up phenomenon was studied by many authors [1–23]. In most cases, these studies were limited to power-law nonlinearities in the right-hand side of inequality (1) or dealt with second-order differential operators. In our paper, we deal with a much wider class of differential inequalities. Our aim is to obtain exact conditions guaranteeing that any global weak solution of (1), (2) is trivial, i.e. is equal to zero almost everywhere in \mathbb{R}_+^{n+1} .

In paper [15], in the partial case where $k = 1$ and $\mathcal{L} = \Delta$, it was obtained a blow-up condition of a Dini-type structure that strengthens the well-known result of Fujita–Hayakawa. Moreover, it was shown in [15] that the validity of this condition is necessary for the non-existence of non-trivial global solutions. Our general condition (12) coincides with that in [15] in the mentioned partial case. Let us note that, in contrast to [15], we impose no ellipticity conditions on the differential operator \mathcal{L} and no growth conditions on the initial data of the Cauchy problem and on a solution of this problem itself. In so doing, we have reason to believe that (12) is exact in much more general case. For example, in the case of the Cauchy problem for the wave equation, i.e. in the case where $k = 2$ and $\mathcal{L} = \Delta$, condition (12) is more general than the well-known Kato blow-up condition [8]. In accordance with [20, Theorem 2] the Kato blow-up condition is exact in the scale of power-law nonlinearities for weak solutions of the Cauchy problem for the wave equation.

For $f(t) = t^\lambda$, i.e. in the case of power-law nonlinearities, Theorem 2 proved in our paper implies the result obtained in [3, Theorem 1] (see Example 1). In this case, conditions (10) and (11) take the simple form $\lambda > 1$ and $\lambda > p$, respectively.

We also note that (11) coincides with the analogous condition in [14, Theorem 2.1], where the blow-up phenomenon was studied for the stationary inequality

$$-\mathcal{L}u \geq f(|u|) \quad \text{in } \mathbb{R}^n. \quad (6)$$

This seems natural because the existence of a non-trivial solution of the last inequality implies the existence of a non-trivial solution of problem (1), (2) that does not depend on t . However, condition (12) is stronger than the similar blow-up condition for solutions of inequality (6). This effect can be detected even in the case of power-law nonlinearities. For instance, if $f(t) = t^\lambda$ and, at the same time, $p \geq 1$ and $n > m$, then in accordance with [16, Theorems 4.1 and 29.1] the blow-up of solutions of inequality (6) is guaranteed by the condition

$$p < \lambda \leq \frac{n}{n-m}p, \quad (7)$$

whereas the blow-up of solutions of the problem

$$u_t - \mathcal{L}u \geq |u|^\lambda \quad \text{in } \mathbb{R}_+^{n+1}, \quad u(x, 0) = u_0(x) \geq 0, \quad (8)$$

occurs if

$$p < \lambda \leq p + \frac{m}{n}. \quad (9)$$

In so doing, both conditions (7) and (9) are exact. Moreover, if $p = 1$ and $m = 2$, then (9) coincides with the well-known Fujita–Hayakawa condition. As mentioned above, the first inequality in (9) follows from (11), while the second inequality is a consequence of (12).

Now, let us say a few words about the structure of this paper. Our main results are Theorems 1–4 formulated in Section 2. This section also contains some examples and remarks demonstrating the exactness of these results.

Sections 3 and 4 are devoted to the proof of Theorems 1–4. The main idea of the proof is to estimate the function E defined by formula (60). The final aim is to show that the integrand in the right-hand sides of (60) is equal to zero. This is obviously equivalent to the triviality of a solution of (1), (2).

The key statements of Section 3 are Lemmas 4–7. They imply Lemmas 8 and 9 that contain estimates of the function E . Lemmas 1–3 are of a technical nature. We need them to prove Lemmas 4–7.

Theorems 1–4 are proved in Section 4. According to Lemmas 4 and 5, in the case of $n \leq m(1 - 1/k)$, conditions (10) and (11) guarantee that $E(r)$ tends to zero as $r \rightarrow \infty$. This is obviously enough to prove Theorem 1. To prove Theorems 2 and 3, we assume by contradiction that (1), (2) has a non-trivial solution. This, in turn, allows one to assert that $E(r) > 0$ for all sufficiently large $r > 0$. Applying Lemmas 4, 5, 8, and 9, we obtain estimates of $E(r)$ divided by a suitable power of r in a neighborhood of zero. These estimates lead us to a contradiction with conditions of Theorems 2 and 3. Finally, we note that Theorem 4 is a consequence of Theorems 1–3.

2 Main results

Theorem 1. Let $n \leq m(1 - 1/k)$ and, moreover,

$$\int_1^\infty \frac{\zeta^{1/k-1} d\zeta}{f^{1/k}(\zeta)} < \infty \quad (10)$$

and

$$\int_1^\infty \frac{\zeta^{p/m-1} d\zeta}{f^{1/m}(\zeta)} < \infty. \quad (11)$$

Then any global weak solution of (1), (2) is trivial.

Theorem 2. Let $n > m(1 - 1/k)$ and $p \in (0, \infty) \cap (1 - m/n, \infty)$. If (10) and (11) are valid and

$$\int_0^1 \frac{f(r) dr}{r^{1+\gamma}} = \infty, \quad (12)$$

where

$$\gamma = \frac{np + m/k}{n - (1 - 1/k)m}, \quad (13)$$

then any global weak solution of (1), (2) is trivial.

Theorem 3. Let $n > m(1 - 1/k)$ and $p \in (0, 1 - m/n]$. If (10) and (11) are valid and

$$\lim_{r \rightarrow +0} \frac{f(r)}{r^\mu} = \infty \quad (14)$$

for some real number $\mu < p/(1 - m/n)$, then any global weak solution of (1), (2) is trivial.

Theorem 4. Let (10) and (11) be valid and $f(0) > 0$. Then (1), (2) has no global weak solutions.

Proof of Theorems 1–4 is given in Section 4.

Remark 1. Inequality (10) coincides with the Kiguradze–Kvinikadze blow-up condition for solutions of the Cauchy problem

$$w^{(k)}(r) = f(w), \quad w(0) > 0, \quad w^{(i)}(0) = 0, \quad i = 1, \dots, k-1, \quad (15)$$

for the ordinary differential equation [10, Theorem 1.2]. In the case where (10) is not fulfilled, it can be shown that (15) has a solution defined on the whole interval $[0, \infty)$. Therefore, taking $u(x, t) = w(t)$, we obtain a non-trivial global weak solution of (1), (2) which does not depend on spatial variables.

Analogously, (11) coincides with the Kiguradze–Kvinikadze blow-up condition for the Cauchy problem

$$w^{(m)}(r) = f(w^{1/p}), \quad w(0) > 0, \quad w^{(i)}(0) = 0, \quad i = 1, \dots, m-1. \quad (16)$$

If (11) is not valid, then (16) has a solution defined on the whole interval $[0, \infty)$. Since $w(r) > w(0)$ for all $r \in (0, \infty)$, setting $v = w - w(0)$, we obtain a solution of the inequality

$$v^{(m)}(r) \geq f(v^{1/p})$$

for all $r \in [0, \infty)$ satisfying the conditions $v^{(i)}(0) = 0$, $i = 0, \dots, m-1$. Thus, the function $u(x, t) = v^{1/p}(|x_1|)$ is a non-trivial global weak solution of (1), (2) with

$$\mathcal{L}u = -\frac{\partial^m |u|^p (\text{sign } x_1)^m}{\partial x_1^m}.$$

In the case where $k = p = 1$ and $m = 2$, we obviously have $\gamma = 1 + 2/n$. In this case, (12) coincides with the blow-up condition for solutions of the Cauchy problem for the heat equation obtained in [15]. Note that, in paper [15], it is assumed that the initial values of the Cauchy problem belongs to $L_\infty(\mathbb{R}^n)$ and the solutions belong to $L_\infty(\mathbb{R}^n \times (0, T))$ for all real number $T > 0$. We do not need such assumptions.

For the problem

$$u_{tt} - \Delta u \geq f(|u|) \quad \text{in } \mathbb{R}_+^{n+1}, \quad u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x) \geq 0. \quad (17)$$

Theorems 1 and 2 imply the following statements.

Corollary 1. *Let $n = 1$ and*

$$\int_1^\infty (f(\zeta)\zeta)^{-1/2} d\zeta < \infty. \quad (18)$$

Then any global weak solution of (17) is trivial.

Corollary 2. *Let $n \geq 2$, condition (18) be valid, and*

$$\int_0^1 \frac{f(r)dr}{r^{2+2/(n-1)}} = \infty. \quad (19)$$

Then any global weak solution of (17) is trivial.

To prove Corollaries 1 and 2, we note that, for $p = 1$ and $k = m = 2$, condition (18) is equivalent to (10) and (11). At the same time, (19) is equivalent to (12). Thus, Corollary 1 follows from Theorem 1, while Corollary 2 is a consequence of Theorem 2.

Remark 2. *As mentioned above, condition (18) is essential. If (18) is not valid, then problem (17) has a positive global weak solution which depends only on t . For power-law nonlinearities $f(t) = t^\lambda$, Corollary 2 leads to the Kato blow-up condition*

$$1 < \lambda \leq 1 + \frac{2}{n-1} \quad (20)$$

obtained in paper [8]. Indeed, the first inequality in (20) is equivalent to (18), whereas the second inequality is equivalent to (19). It is well-known that both inequalities in (20) are exact. To verify the exactness of the first inequality, it is sufficient to check that, in the case of $\lambda \leq 1$, the function $u(x, t) = e^t$ is a positive global weak solution of the problem

$$u_{tt} - \Delta u \geq u^\lambda \quad \text{in } \mathbb{R}_+^{n+1}, \quad u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x) \geq 0.$$

In accordance with [20, Theorem 2] the second inequality in (20) is also exact.

Example 1. Let us consider problem (8). It is easy to see that condition (10) is equivalent to $\lambda > 1$, while (11) is equivalent to $\lambda > p$. In so doing, (12) is fulfilled if

$$\lambda \leq p + \frac{m}{n}.$$

Thus, in accordance with Theorem 2 if

$$\max\{1, p\} < \lambda \leq p + \frac{m}{n}, \quad (21)$$

then any global weak solution of (8) is trivial.

Note that condition (21) was earlier obtained in [3, Theorem 1].

Example 2. We examine the critical exponent $\lambda = p + m/n > 1$ in (21). Namely, consider the problem

$$u_t - \mathcal{L}u \geq u^{p+m/n} \log^\mu \left(e + \frac{1}{u} \right) \quad \text{in } \mathbb{R}_+^{n+1}, \quad u(x, 0) = u_0(x) \geq 0, \quad (22)$$

where μ is a real number. For $u = 0$, we assume by continuity that the right-hand side of inequality (22) is equal to zero. By Theorem 2, if

$$\mu \geq -1,$$

then any global weak solution of (22) is trivial.

3 Preliminary and auxiliary statements

In this section, it is assumed that u is a global weak solution of (1), (2) and, moreover, conditions (10) and (11) are valid. Let us recall that we have agreed to denote by Q_r^\varkappa the cylinder $Q_r^\varkappa = B_r \times (0, r^\varkappa)$, where $\varkappa > 0$ is a real number and B_r is the open ball of radius $r > 0$ centered at zero. Below, by C and σ we mean various positive constants that can depend only on A, k, m, n, p , and \varkappa . Put

$$F_\varkappa(s) = \left(\int_s^\infty \frac{\zeta^{1/k-1} d\zeta}{f^{1/k}(\zeta)} \right)^{1/\varkappa} + \int_s^\infty \frac{\zeta^{p/m-1} d\zeta}{f^{1/m}(\zeta)}.$$

Lemma 1. Let $R > 0$ and $\varkappa > 0$ be real numbers. Then

$$\begin{aligned} \frac{1}{(r_2^\varkappa - r_1^\varkappa)^k} \int_{r_1^\varkappa}^{r_2^\varkappa} \int_{B_{r_2}} |u| dx dt + \frac{1}{(r_2 - r_1)^m} \int_0^{r_2^\varkappa} \int_{B_{r_2} \setminus B_{r_1}} |u|^p dx dt \\ \geq C \int_{Q_{r_1}^\varkappa} f(|u|) dx dt \end{aligned} \quad (23)$$

for all real numbers $R \leq r_1 < r_2 \leq 2R$.

Proof. Take a non-decreasing function $\varphi_0 \in C^\infty(\mathbb{R})$ such that

$$\varphi_0|_{(-\infty,0]} = 0 \quad \text{and} \quad \varphi_0|_{[1,\infty)} = 1.$$

We put

$$\varphi(x, t) = \varphi_0\left(\frac{r_2 - |x|}{r_2 - r_1}\right) \varphi_0\left(\frac{r_2^\varkappa - t}{r_2^\varkappa - r_1^\varkappa}\right). \quad (24)$$

It is easy to see that

$$\left| \int_{\mathbb{R}_+^{n+1}} u \partial_t^k \varphi dx dt \right| \leq \frac{C}{(r_2^\varkappa - r_1^\varkappa)^k} \int_{r_1^\varkappa}^{r_2^\varkappa} \int_{B_{r_2}} |u| dx dt,$$

$$\left| \int_{\mathbb{R}_+^{n+1}} \sum_{|\alpha|=m} (-1)^m a_\alpha(x, t, u) \partial^\alpha \varphi dx dt \right| \leq \frac{C}{(r_2 - r_1)^m} \int_0^{r_2^\varkappa} \int_{B_{r_2} \setminus B_{r_1}} |u|^p dx dt,$$

and

$$\int_{\mathbb{R}_+^{n+1}} f(|u|) \varphi dx dt \geq \int_0^{r_1^\varkappa} \int_{B_{r_1}} f(|u|) dx dt = \int_{Q_{r_1}^\varkappa} f(|u|) dx dt.$$

Thus, taking (24) as a test function in (3), we readily obtain (23). \square

Lemma 2. *Let $p \geq 1$, then for all real numbers $R > 0$, $\varkappa > 0$, and $R \leq r_1 < r_2 \leq 2R$ at least one of the following two inequalities is valid:*

$$I(r_2) - I(r_1) \geq C(r_2 - r_1)^{kp} R^{kp(\varkappa-1)} f^p(I^{1/p}(r_1)), \quad (25)$$

$$I(r_2) - I(r_1) \geq C(r_2 - r_1)^m f(I^{1/p}(r_1)), \quad (26)$$

where

$$I(r) = \frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_r^\varkappa} |u|^p dx dt, \quad R < r < 2R. \quad (27)$$

Proof. At first, assume that

$$\frac{1}{(r_2^\varkappa - r_1^\varkappa)^k} \int_{r_1^\varkappa}^{r_2^\varkappa} \int_{B_{r_2}} |u| dx dt \geq \frac{1}{(r_2 - r_1)^m} \int_0^{r_2^\varkappa} \int_{B_{r_2} \setminus B_{r_1}} |u|^p dx dt. \quad (28)$$

Then Lemma 1 implies the estimate

$$\frac{1}{(r_2^\varkappa - r_1^\varkappa)^k} \int_{r_1^\varkappa}^{r_2^\varkappa} \int_{B_{r_2}} |u| dx dt \geq C \int_{Q_{r_1}^\varkappa} f(|u|) dx dt.$$

Since

$$\int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u| dx dt \geq \int_{r_1^\varkappa}^{r_2^\varkappa} \int_{B_{r_2}} |u| dx dt,$$

this yields

$$\int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u| dxdt \geq C(r_2^\varkappa - r_1^\varkappa)^k \int_{Q_{r_1}^\varkappa} f(|u|) dxdt. \quad (29)$$

Evaluating the left-hand side of the last expression by Hölder's inequality

$$(\text{mes } Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa)^{(p-1)/p} \left(\int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u|^p dxdt \right)^{1/p} \geq \int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u| dxdt,$$

we obtain

$$\left(\int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u|^p dxdt \right)^{1/p} \geq \frac{C(r_2^\varkappa - r_1^\varkappa)^k}{(\text{mes } Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa)^{(p-1)/p}} \int_{Q_{r_1}^\varkappa} f(|u|) dxdt,$$

whence it follows that

$$\begin{aligned} \frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u|^p dxdt &\geq C(r_2^\varkappa - r_1^\varkappa)^{kp} \left(\frac{\text{mes } Q_{2R}^\varkappa}{\text{mes } Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} \right)^{p-1} \\ &\quad \times \left(\frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_1}^\varkappa} f(|u|) dxdt \right)^p. \end{aligned} \quad (30)$$

Since $f(\zeta^{1/p})$ is a non-decreasing convex function, we have

$$\begin{aligned} \frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_1}^\varkappa} f(|u|) dxdt &\geq \frac{1}{2^{n+\varkappa} \text{mes } Q_{r_1}^\varkappa} \int_{Q_{r_1}^\varkappa} f(|u|) dxdt \\ &\geq \frac{1}{2^{n+\varkappa}} f \left(\left(\frac{1}{\text{mes } Q_{r_1}^\varkappa} \int_{Q_{r_1}^\varkappa} |u|^p dxdt \right)^{1/p} \right) \\ &\geq \frac{1}{2^{n+\varkappa}} f \left(\left(\frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_1}^\varkappa} |u|^p dxdt \right)^{1/p} \right). \end{aligned} \quad (31)$$

In view of (30), this implies the estimate

$$\begin{aligned} \frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u|^p dxdt &\geq C(r_2^\varkappa - r_1^\varkappa)^{kp} \left(\frac{\text{mes } Q_{2R}^\varkappa}{\text{mes } Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} \right)^{p-1} \\ &\quad \times f^p \left(\left(\frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_1}^\varkappa} |u|^p dxdt \right)^{1/p} \right), \end{aligned}$$

combining which with the evident inequalities

$$\left(\frac{\text{mes } Q_{2R}^\varkappa}{\text{mes } Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} \right)^{p-1} \geq 1$$

and

$$r_2^\varkappa - r_1^\varkappa \geq C(r_2 - r_1)R^{\varkappa-1}, \quad (32)$$

we arrive at (25).

Now, let (28) not hold. In this case, Lemma 1 implies that

$$\frac{1}{(r_2 - r_1)^m} \int_0^{r_2^\varkappa} \int_{B_{r_2} \setminus B_{r_1}} |u|^p dx dt \geq C \int_{Q_{r_1}^\varkappa} f(|u|) dx dt, \quad (33)$$

whence in accordance with (31) and the inequality

$$\int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u|^p dx dt \geq \int_0^{r_2^\varkappa} \int_{B_{r_2} \setminus B_{r_1}} |u|^p dx dt \quad (34)$$

we obtain

$$\frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u|^p dx dt \geq C(r_2 - r_1)^m f \left(\left(\frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_1}^\varkappa} |u|^p dx dt \right)^{1/p} \right).$$

To complete the proof, it remains to note that the last expression is equivalent to (26). \square

Lemma 3. *Let $0 < p < 1$, then for all real numbers $R > 0$, $\varkappa > 0$, and $R \leq r_1 < r_2 \leq 2R$ at least one of the following two inequalities is valid:*

$$J(r_2) - J(r_1) \geq C(r_2 - r_1)^k R^{k(\varkappa-1)} f(J(r_1)), \quad (35)$$

$$J(r_2) - J(r_1) \geq C(r_2 - r_1)^{m/p} f^{1/p}(J(r_1)), \quad (36)$$

where

$$J(r) = \frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_r^\varkappa} |u| dx dt, \quad R < r < 2R. \quad (37)$$

Proof. At first, assume that (28) is valid. As in the proof of Lemma 2, we can obviously assert that (29) holds. At the same time, since f is a non-decreasing convex function, we have

$$\begin{aligned} \frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_1}^\varkappa} f(|u|) dx dt &\geq \frac{1}{2^{n+\varkappa} \text{mes } Q_{r_1}^\varkappa} \int_{Q_{r_1}^\varkappa} f(|u|) dx dt \\ &\geq \frac{1}{2^{n+\varkappa}} f \left(\frac{1}{\text{mes } Q_{r_1}^\varkappa} \int_{Q_{r_1}^\varkappa} |u| dx dt \right) \end{aligned}$$

$$\geq \frac{1}{2^{n+\varkappa}} f \left(\frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_1}^\varkappa} |u| dxdt \right). \quad (38)$$

Thus, (29) implies the estimate

$$\frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u| dxdt \geq C(r_2^\varkappa - r_1^\varkappa)^k f \left(\frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_1}^\varkappa} |u| dxdt \right),$$

whence in accordance with (32) inequality (35) follows at once.

In its turn, if (28) is not valid, then Lemma 1 implies (33), whence in accordance with (34) and (38) we obtain

$$\frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u|^p dxdt \geq C(r_2 - r_1)^m f \left(\frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_1}^\varkappa} |u| dxdt \right). \quad (39)$$

By Hölder's inequality,

$$\int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u|^p dxdt \leq (\text{mes } Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa)^{1-p} \left(\int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u| dxdt \right)^p;$$

therefore, (39) yields

$$\begin{aligned} \frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} |u| dxdt &\geq C(r_2 - r_1)^{m/p} \left(\frac{\text{mes } Q_{2R}^\varkappa}{\text{mes } Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} \right)^{1/p-1} \\ &\quad \times f^{1/p} \left(\frac{1}{\text{mes } Q_{2R}^\varkappa} \int_{Q_{r_1}^\varkappa} |u| dxdt \right). \end{aligned}$$

Since

$$\left(\frac{\text{mes } Q_{2R}^\varkappa}{\text{mes } Q_{r_2}^\varkappa \setminus Q_{r_1}^\varkappa} \right)^{1/p-1} \geq 1,$$

this readily implies (36). \square

Lemma 4. *Suppose that $p \geq 1$, then*

$$F_\varkappa \left(\left(\frac{\sigma}{R^{n+\varkappa}} \int_{Q_R^\varkappa} |u|^p dxdt \right)^{1/p} \right) \geq CR, \quad (40)$$

for any real number $R > 0$ and $\varkappa > 0$ such that $\text{mes}\{(x, t) \in Q_R^\varkappa : u(x, t) \neq 0\} > 0$.

Proof. Take the minimal positive integer l such that $2^l I(R) \geq I(2R)$, where the function I is defined by (27). We construct a finite sequence of real numbers r_i , $i = 0, \dots, l$,

by putting $r_0 = R$,

$$r_i = \sup\{r \in (r_{i-1}, 2R) : I(r) \leq 2I(r_{i-1})\}, \quad 0 < i \leq l-1,$$

and $r_l = 2R$. It can easily be seen that

$$I(r_i) = 2I(r_{i-1}) \quad \text{for all } 0 < i \leq l-1 \quad \text{and} \quad I(r_l) \leq 2I(r_{l-1}). \quad (41)$$

According to Lemma 2, for any integer $0 < i \leq l$ at least one of the following two inequalities is valid:

$$I(r_i) - I(r_{i-1}) \geq C(r_i - r_{i-1})^{kp} R^{kp(\alpha-1)} f^p(I^{1/p}(r_{i-1})), \quad (42)$$

$$I(r_i) - I(r_{i-1}) \geq C(r_i - r_{i-1})^m f(I^{1/p}(r_{i-1})). \quad (43)$$

We denote by Ξ_1 the set of all integers $0 < i \leq l$ for which (42) holds. Also let Ξ_2 be the set of all other integers $0 < i \leq l$. If $i \in \Xi_1$, then (41) and (42) imply that

$$\frac{I^{1/(kp)}(r_{i-1})}{f^{1/k}(I^{1/p}(r_{i-1}))} \geq C(r_i - r_{i-1}) R^{\alpha-1},$$

whence in accordance with the evident inequality

$$\int_{I(r_{i-1})}^{2I(r_{i-1})} \frac{\zeta^{1/(kp)-1} d\zeta}{f^{1/k}(2^{-1/p}\zeta^{1/p})} \geq \frac{2^{1/(kp)-1} I^{1/(kp)}(r_{i-1})}{f^{1/k}(I^{1/p}(r_{i-1}))}$$

we obtain

$$\int_{I(r_{i-1})}^{2I(r_{i-1})} \frac{\zeta^{1/(kp)-1} d\zeta}{f^{1/k}(2^{-1/p}\zeta^{1/p})} \geq C(r_i - r_{i-1}) R^{\alpha-1}, \quad (44)$$

In its turn, if $i \in \Xi_2$, then (41) and (43) allow us to assert that

$$\frac{I^{1/m}(r_{i-1})}{f^{1/m}(I^{1/p}(r_{i-1}))} \geq C(r_i - r_{i-1}).$$

Since

$$\int_{I(r_{i-1})}^{2I(r_{i-1})} \frac{\zeta^{1/m-1} d\zeta}{f^{1/m}(2^{-1/p}\zeta^{1/p})} \geq \frac{2^{1/m-1} I^{1/m}(r_{i-1})}{f^{1/m}(I^{1/p}(r_{i-1}))},$$

this readily implies the estimate

$$\int_{I(r_{i-1})}^{2I(r_{i-1})} \frac{\zeta^{1/m-1} d\zeta}{f^{1/m}(2^{-1/p}\zeta^{1/p})} \geq C(r_i - r_{i-1}). \quad (45)$$

At first, we assume that

$$\sum_{i \in \Xi_1} (r_i - r_{i-1}) \geq \frac{R}{2}. \quad (46)$$

Summing (44) over all $i \in \Xi_1$, we obtain

$$\int_{I(R)}^{\infty} \frac{\zeta^{1/(kp)-1} d\zeta}{f^{1/k}(2^{-1/p}\zeta^{1/p})} \geq CR^{\varkappa}.$$

By the change of variable $\xi = 2^{-1/p}\zeta^{1/p}$, the last expression is reduced to the form

$$\int_{(I(R)/2)^{1/p}}^{\infty} \frac{\xi^{1/k-1} d\xi}{f^{1/k}(\xi)} \geq CR^{\varkappa}.$$

This, in its turn, implies (40).

Now let (46) not hold. Then

$$\sum_{i \in \Xi_2} (r_i - r_{i-1}) \geq \frac{R}{2}; \quad (47)$$

therefore, summing (45) over all $i \in \Xi_2$, we have

$$\int_{I(R)}^{\infty} \frac{\zeta^{1/m-1} d\zeta}{f^{1/m}(2^{-1/p}\zeta^{1/p})} \geq CR.$$

By the change of variable $\xi = 2^{-1/p}\zeta^{1/p}$, this can be transformed to the inequality

$$\int_{(I(R)/2)^{1/p}}^{\infty} \frac{\xi^{p/m-1} d\xi}{f^{1/m}(\xi)} \geq CR,$$

whence (40) follows again. The proof is completed. \square

Lemma 5. *Suppose that $0 < p < 1$, then*

$$F_{\varkappa} \left(\frac{\sigma}{R^{n+\varkappa}} \int_{Q_R^{\varkappa}} |u| dx dt \right) \geq CR \quad (48)$$

for any real number $R > 0$ and $\varkappa > 0$ such that $\text{mes}\{(x, t) \in Q_R^{\varkappa} : u(x, t) \neq 0\} > 0$.

Proof. We take the minimal positive integer l such that $2^l J(R) \geq J(2R)$, where the function J is defined by (37). Consider the sequence of real numbers r_i , $i = 0, \dots, l$, constructed in the proof of Lemma 4 with I replaced by J . We obviously have

$$J(r_i) = 2J(r_{i-1}) \quad \text{for all } 0 < i \leq l-1 \quad \text{and} \quad J(r_l) \leq 2J(r_{l-1}). \quad (49)$$

In view of Lemma 3, for any integer $0 < i \leq l$ at least one of the following two inequalities is valid:

$$J(r_i) - J(r_{i-1}) \geq C(r_i - r_{i-1})^k R^{k(\varkappa-1)} f(J(r_{i-1})), \quad (50)$$

$$J(r_i) - J(r_{i-1}) \geq C(r_i - r_{i-1})^{m/p} f^{1/p}(J(r_{i-1})). \quad (51)$$

We denote by Ξ_1 the set of integers $0 < i \leq l$ such that (50) holds and let Ξ_2 be the set of all other positive integers $0 < i \leq l$. For any $i \in \Xi_1$, it follows from (49) and (50) that

$$\frac{J^{1/k}(r_{i-1})}{f^{1/k}(J(r_{i-1}))} \geq C(r_i - r_{i-1})R^{\alpha-1}.$$

Combining this with the evident inequality

$$\int_{J(r_{i-1})}^{2J(r_{i-1})} \frac{\zeta^{1/k-1} d\zeta}{f^{1/k}(\zeta/2)} \geq \frac{2^{1/k-1} J^{1/k}(r_{i-1})}{f^{1/k}(J(r_{i-1}))},$$

we obtain

$$\int_{J(r_{i-1})}^{2J(r_{i-1})} \frac{\zeta^{1/k-1} d\zeta}{f^{1/k}(\zeta/2)} \geq C(r_i - r_{i-1})R^{\alpha-1}. \quad (52)$$

In its turn, for any $i \in \Xi_2$ in accordance with (49) and (51) we have

$$\frac{J^{p/m}(r_{i-1})}{f^{1/m}(J(r_{i-1}))} \geq C(r_i - r_{i-1}).$$

In view of the estimate

$$\int_{J(r_{i-1})}^{2J(r_{i-1})} \frac{\zeta^{p/m-1} d\zeta}{f^{1/m}(\zeta/2)} \geq \frac{2^{p/m-1} J^{p/m}(r_{i-1})}{f^{1/m}(J(r_{i-1}))},$$

this implies that

$$\int_{J(r_{i-1})}^{2J(r_{i-1})} \frac{\zeta^{p/m-1} d\zeta}{f^{1/m}(\zeta/2)} \geq C(r_i - r_{i-1}). \quad (53)$$

At first, let (46) be fulfilled. Summing (52) over all $i \in \Xi_1$, we obtain

$$\int_{J(R)}^{\infty} \frac{\zeta^{1/k-1} d\zeta}{f^{1/k}(\zeta/2)} \geq CR^{\alpha}.$$

After the change of variable $\xi = \zeta/2$, the last expression takes the form

$$\int_{J(R)/2}^{\infty} \frac{\xi^{1/k-1} d\xi}{f^{1/k}(\xi)} \geq CR^{\alpha},$$

whence (48) immediately follows.

Now, assume that (46) is not valid. In this case, (47) holds and, summing (53) over all $i \in \Xi_2$, we arrive at the inequality

$$\int_{J(R)}^{\infty} \frac{\zeta^{p/m-1} d\zeta}{f^{1/m}(\zeta/2)} \geq CR$$

which, by the change of variable $\xi = \zeta/2$, can be transformed to

$$\int_{J(R)/2}^{\infty} \frac{\xi^{p/m-1} d\xi}{f^{1/m}(\xi)} \geq CR.$$

This again leads us to (48). \square

Lemma 6. *Suppose that $p \geq 1$, then*

$$\begin{aligned} \frac{1}{R^{k\kappa-m}} \left(\frac{1}{\text{mes } Q_R^\kappa \setminus Q_{R/2}^\kappa} \int_{Q_R^\kappa \setminus Q_{R/2}^\kappa} |u|^p dxdt \right)^{1/p} &+ \frac{1}{\text{mes } Q_R^\kappa \setminus Q_{R/2}^\kappa} \int_{Q_R^\kappa \setminus Q_{R/2}^\kappa} |u|^p dxdt \\ &\geq \frac{C}{R^{n+\kappa-m}} \int_{Q_{R/2}^\kappa} f(|u|) dxdt \end{aligned} \quad (54)$$

for all real numbers $R > 0$ and $\kappa > 0$.

Proof. Lemma 1 with $r_1 = R/2$ and $r_2 = R$ yields

$$\frac{1}{R^{k\kappa}} \int_{Q_R^\kappa \setminus Q_{R/2}^\kappa} |u| dxdt + \frac{1}{R^m} \int_{Q_R^\kappa \setminus Q_{R/2}^\kappa} |u|^p dxdt \geq C \int_{Q_{R/2}^\kappa} f(|u|) dxdt \quad (55)$$

for all real numbers $R > 0$ and $\kappa > 0$, whence in accordance with Hölder's inequality

$$(\text{mes } Q_R^\kappa \setminus Q_{R/2}^\kappa)^{(p-1)/p} \left(\int_{Q_R^\kappa \setminus Q_{R/2}^\kappa} |u|^p dxdt \right)^{1/p} \geq \int_{Q_R^\kappa \setminus Q_{R/2}^\kappa} |u| dxdt$$

and the fact that

$$\sigma R^{n+\kappa} \leq \text{mes } Q_R^\kappa \setminus Q_{R/2}^\kappa \leq CR^{n+\kappa} \quad (56)$$

estimate (54) follows at once. \square

Lemma 7. *Assume that $0 < p < 1$, then*

$$\begin{aligned} \frac{1}{\text{mes } Q_R^\kappa \setminus Q_{R/2}^\kappa} \int_{Q_R^\kappa \setminus Q_{R/2}^\kappa} |u| dxdt + \frac{1}{R^{m-k\kappa}} \left(\frac{1}{\text{mes } Q_R^\kappa \setminus Q_{R/2}^\kappa} \int_{Q_R^\kappa \setminus Q_{R/2}^\kappa} |u| dxdt \right)^p \\ \geq \frac{C}{R^{n-(k-1)\kappa}} \int_{Q_{R/2}^\kappa} f(|u|) dxdt \end{aligned} \quad (57)$$

for all real numbers $R > 0$ and $\kappa > 0$.

Proof. In view of Lemma 1, relation (55) holds for all real numbers $R > 0$ and $\kappa > 0$. Evaluating the second summand on the left in (55) by Hölder's inequality

$$(\text{mes } Q_R^\kappa \setminus Q_{R/2}^\kappa)^{1-p} \left(\int_{Q_R^\kappa \setminus Q_{R/2}^\kappa} |u| dxdt \right)^p \geq \int_{Q_R^\kappa \setminus Q_{R/2}^\kappa} |u|^p dxdt$$

and using estimate (56), we readily obtain (57). \square

Lemma 8. *Let $p \geq 1$, then for all real numbers $R > 0$ and $\varkappa > 0$ at least one of the following two systems of inequalities is valid:*

$$\left\{ \begin{array}{l} \left(\frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} |u|^p dxdt \right)^{1/p} \geq \frac{\sigma}{R^{n-(k-1)\varkappa}} \int_{Q_{R/2}^\varkappa} f(|u|) dxdt, \\ E(R) - E(R/2) \geq CR^{n+\varkappa} f \left(\frac{\sigma E(R/2)}{R^{n-(k-1)\varkappa}} \right), \end{array} \right. \quad (58)$$

$$\left\{ \begin{array}{l} \frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} |u|^p dxdt \geq \frac{\sigma}{R^{n+\varkappa-m}} \int_{Q_{R/2}^\varkappa} f(|u|) dxdt, \\ E(R) - E(R/2) \geq CR^{n+\varkappa} f \left(\left(\frac{\sigma E(R/2)}{R^{n+\varkappa-m}} \right)^{1/p} \right), \end{array} \right. \quad (59)$$

where

$$E(r) = \int_{Q_r^\varkappa} f(|u|) dxdt, \quad r > 0. \quad (60)$$

Proof. Let

$$\begin{aligned} & \frac{1}{R^{k\varkappa-m}} \left(\frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} |u|^p dxdt \right)^{1/p} \\ & \geq \frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} |u|^p dxdt. \end{aligned} \quad (61)$$

Applying Lemma 6, we have the first inequality in (58), whence it follows that

$$f \left(\left(\frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} |u|^p dxdt \right)^{1/p} \right) \geq f \left(\frac{\sigma}{R^{n-(k-1)\varkappa}} \int_{Q_{R/2}^\varkappa} f(|u|) dxdt \right)$$

since f is a non-decreasing function. Combining this with the estimate

$$\begin{aligned} & \frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} f(|u|) dxdt \\ & \geq f \left(\left(\frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} |u|^p dxdt \right)^{1/p} \right) \end{aligned} \quad (62)$$

which is valid as $f(\zeta^{1/p})$ is a convex function, we obtain

$$\frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} f(|u|) dxdt \geq f \left(\frac{\sigma}{R^{n-(k-1)\varkappa}} \int_{Q_{R/2}^\varkappa} f(|u|) dxdt \right).$$

In view of (56), the last expression is equivalent to the second inequality in (58).

Now, assume that (61) is not satisfied. In this case, Lemma 6 implies the first inequality in (59) which, in turn, allows us to assert that

$$\begin{aligned} & f \left(\left(\frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} |u|^p dxdt \right)^{1/p} \right) \\ & \geq f \left(\left(\frac{\sigma}{R^{n+\varkappa-m}} \int_{Q_{R/2}^\varkappa} f(|u|) dxdt \right)^{1/p} \right). \end{aligned}$$

Combining this with (62), we have

$$\frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} f(|u|) dxdt \geq f \left(\left(\frac{\sigma}{R^{n+\varkappa-m}} \int_{Q_{R/2}^\varkappa} f(|u|) dxdt \right)^{1/p} \right),$$

whence in accordance with (56) the second inequality in (59) follows at once. \square

Lemma 9. *Let $0 < p < 1$, then for all real numbers $R > 0$ and $\varkappa > 0$ at least one of the following two systems of inequalities is valid:*

$$\begin{cases} \frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} |u| dxdt \geq \frac{\sigma}{R^{n-(k-1)\varkappa}} \int_{Q_{R/2}^\varkappa} f(|u|) dxdt, \\ E(R) - E(R/2) \geq CR^{n+\varkappa} f \left(\frac{\sigma E(R/2)}{R^{n-(k-1)\varkappa}} \right), \end{cases}$$

$$\begin{cases} \left(\frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} |u| dxdt \right)^p \geq \frac{\sigma}{R^{n+\varkappa-m}} \int_{Q_{R/2}^\varkappa} f(|u|) dxdt, \\ E(R) - E(R/2) \geq CR^{n+\varkappa} f \left(\left(\frac{\sigma E(R/2)}{R^{n+\varkappa-m}} \right)^{1/p} \right), \end{cases}$$

where the function E is defined by (60).

Proof. The proof is completely analogous to the proof of Lemma 8. The only difference is that Lemma 6 should be replaced by Lemma 7. \square

4 Proof of Theorems 1–4

In this section, as in the previous one, by C and σ we denote various positive constants that can depend only on A , k , m , n , and p , unless otherwise stated.

Proof of Theorem 1. Let us take $\varkappa = m/k$. At first, we consider the case where $p \geq 1$. For any non-trivial global weak solution of (1), (2) in accordance with the evident estimate

$$\sigma R^{n+\varkappa} \leq \text{mes } Q_R^\varkappa \leq CR^{n+\varkappa}$$

Lemma 4 implies that

$$\lim_{R \rightarrow \infty} \frac{1}{\text{mes } Q_R^\varkappa} \int_{Q_R^\varkappa} |u|^p dx dt = 0.$$

In doing, Lemma 6 yields

$$\begin{aligned} \left(\frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} |u|^p dx dt \right)^{1/p} &+ \frac{1}{\text{mes } Q_R^\varkappa \setminus Q_{R/2}^\varkappa} \int_{Q_R^\varkappa \setminus Q_{R/2}^\varkappa} |u|^p dx dt \\ &\geq \frac{C}{R^{n-m(1-1/k)}} \int_{Q_{R/2}^\varkappa} f(|u|) dx dt \end{aligned}$$

for all real numbers $R > 0$. Passing in the last expression to the limit as $R \rightarrow \infty$, we have

$$\int_{\mathbb{R}_+^{n+1}} f(|u|) dx dt = 0.$$

Since $f(\zeta) > 0$ for all $\zeta \in (0, \infty)$, this implies that u is equal to zero almost everywhere in \mathbb{R}_+^{n+1} . Thus, we arrive at a contradiction with our assumption that u is a non-trivial solution.

For $0 < p < 1$, all the arguments are completely analogous. The only difference is that Lemmas 4 and 6 should be replaced by Lemmas 5 and 7, respectively. \square

Proof of Theorem 2. Assume the converse. Let u be a non-trivial global weak solution of (1), (2). We put

$$\varkappa = \frac{n(p-1) + m}{1 + (k-1)p}. \quad (63)$$

Since $p > 1 - m/n$ by the hypotheses of the theorem, one can assert that $\varkappa > 0$. It can also be verified that \varkappa satisfies the system of equations

$$\begin{cases} n + \varkappa - \gamma(n - (k-1)\varkappa) = 0, \\ n + \varkappa - \frac{\gamma}{p}(n + \varkappa - m) = 0, \end{cases} \quad (64)$$

where γ is defined by (13). In particular, from (64), it follows that

$$\frac{n + \varkappa - m}{p} = n - (k-1)\varkappa. \quad (65)$$

We denote

$$h(\zeta) = \frac{f(\zeta)}{\zeta^\gamma}, \quad \zeta > 0. \quad (66)$$

Take a real number $r_0 > 0$ such that $E(r_0) > 0$, where the function E is defined by (60). Also let $r_i = 2^i r_0$, $i = 1, 2, \dots$

From now on, we assume that the constants C and σ can also depend on $E(r_0)$.

At first, we consider the case of $p \geq 1$. Let us establish the validity of the estimate

$$\frac{E(r_{i+1}) - E(r_i)}{E^{1/p}(r_i)} \geq Ch \left(\frac{\sigma E^{1/p}(r_i)}{r_i^{n-(k-1)\varkappa}} \right) \quad (67)$$

for all $i = 0, 1, 2, \dots$. Indeed, by Lemma 8, for any integer $i \geq 0$ at least one of the following two systems of inequalities is satisfied:

$$\left\{ \begin{array}{l} \left(\frac{1}{\text{mes } Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} \int_{Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} |u|^p dxdt \right)^{1/p} \geq \frac{\sigma}{r_i^{n-(k-1)\varkappa}} \int_{Q_{r_i}^\varkappa} f(|u|) dxdt, \\ E(r_{i+1}) - E(r_i) \geq Cr_i^{n+\varkappa} f \left(\frac{\sigma E(r_i)}{r_i^{n-(k-1)\varkappa}} \right), \end{array} \right. \quad (68)$$

$$\left\{ \begin{array}{l} \frac{1}{\text{mes } Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} \int_{Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} |u|^p dxdt \geq \frac{\sigma}{r_i^{n+\varkappa-m}} \int_{Q_{r_i}^\varkappa} f(|u|) dxdt, \\ E(r_{i+1}) - E(r_i) \geq Cr_i^{n+\varkappa} f \left(\left(\frac{\sigma E(r_i)}{r_i^{n+\varkappa-m}} \right)^{1/p} \right). \end{array} \right. \quad (69)$$

Let (68) be valid for some integer $i \geq 0$. Since f is a non-decreasing function, the second inequality in (68) implies that

$$E(r_{i+1}) - E(r_i) \geq Cr_i^{n+\varkappa} f \left(\frac{\sigma E^{1-1/p}(r_0) E^{1/p}(r_i)}{r_i^{n-(k-1)\varkappa}} \right),$$

whence in accordance with the first equation in (64), we readily obtain (67).

In its turn, if (69) is valid for some integer $i \geq 0$, then the second inequality in (69) yields

$$E(r_{i+1}) - E(r_i) \geq Cr_i^{n+\varkappa} f \left(\frac{\sigma E^{1/p}(r_i)}{r_i^{(n+\varkappa-m)/p}} \right),$$

whence in accordance with (65) and the second equation in (64) we have (67).

It can be seen that

$$\left(\frac{1}{\text{mes } Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} \int_{Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} |u|^p dxdt \right)^{1/p} \geq \frac{\sigma E^{1/p}(r_i)}{r_i^{n-(k-1)\varkappa}} \quad (70)$$

for all $i = 0, 1, 2, \dots$. Indeed, in the case where (68) holds, estimate (70) follows from the first inequality in (68) and the fact that

$$\int_{Q_{r_i}^\varkappa} f(|u|) dxdt \geq E^{1-1/p}(r_0) E^{1/p}(r_i).$$

In its turn, if (69) holds, then (70) follows from (65) and the first inequality in (69).

By Lemma 4 and estimate (56), the left-hand side of (70) tends to zero as $i \rightarrow \infty$. Thus, passing in (70) to the limit as $i \rightarrow \infty$, we obtain

$$\lim_{i \rightarrow \infty} \frac{E^{1/p}(r_i)}{r_i^{n-(k-1)\varkappa}} = 0. \quad (71)$$

Let us denote $\lambda = n - (k-1)\varkappa$ and

$$\zeta_i = E^{1/p}(r_i), \quad i = 0, 1, 2, \dots$$

Taking into account (63) and the condition $n > m(1 - 1/k)$, we have

$$\lambda = \frac{k(n - m(1 - 1/k))}{1 + (k-1)p} > 0.$$

There are sequences of integers $0 \leq s_i < l_i \leq s_{i+1}$, $i = 1, 2, \dots$, such that

$$\frac{\zeta_j}{r_j^\lambda} > \frac{\zeta_{j+1}}{r_{j+1}^\lambda}$$

if

$$j \in \bigcup_{i=1}^{\infty} [s_i, l_i) \quad (72)$$

and

$$\frac{\zeta_j}{r_j^\lambda} \leq \frac{\zeta_{j+1}}{r_{j+1}^\lambda}$$

if (72) is not fulfilled. Since $E(r_{j+1}) \geq E(r_j)$ for all $j = 1, 2, \dots$, we have

$$\frac{2^\lambda \zeta_{j+1}}{r_{j+1}^\lambda} \geq \frac{\zeta_j}{r_j^\lambda} > \frac{\zeta_{j+1}}{r_{j+1}^\lambda} \quad (73)$$

for all integers j satisfying (72). It can also be seen that

$$2^\lambda \zeta_j > \zeta_{j+1} \geq \zeta_j \quad (74)$$

for all integers j satisfying (72).

Estimate (67) can obviously be written as

$$\frac{\zeta_{j+1}^p - \zeta_j^p}{\zeta_j^\gamma} \geq Ch \left(\frac{\sigma \zeta_j}{r_j^\lambda} \right), \quad j = 1, 2, \dots, \quad (75)$$

whence in accordance with (74) it follows that

$$\frac{\zeta_{j+1} - \zeta_j}{\zeta_j^{\gamma-p+1}} \geq Ch \left(\frac{\sigma \zeta_j}{r_j^\lambda} \right)$$

for all integers j satisfying (72). Multiplying this by the inequality

$$1 \geq \frac{\zeta_j/r_j^\lambda - \zeta_{j+1}/r_{j+1}^\lambda}{\zeta_j/r_j^\lambda},$$

we obtain

$$\frac{\zeta_{j+1} - \zeta_j}{\zeta_j^{\gamma-p+1}} \geq C \frac{h(\sigma\zeta_j/r_j^\lambda)}{\zeta_j/r_j^\lambda} \left(\frac{\zeta_j}{r_j^\lambda} - \frac{\zeta_{j+1}}{r_{j+1}^\lambda} \right) \quad (76)$$

for all integers j satisfying (72).

In view of (74), we have

$$\int_{\zeta_j}^{\zeta_{j+1}} \frac{d\zeta}{\zeta^{\gamma-p+1}} \geq C \frac{\zeta_{j+1} - \zeta_j}{\zeta_j^{\gamma-p+1}}$$

for all integers j satisfying (72). In turn, (73) allows us to assert that

$$\frac{h(\sigma\zeta_j/r_j^\lambda)}{\zeta_j/r_j^\lambda} \left(\frac{\zeta_j}{r_j^\lambda} - \frac{\zeta_{j+1}}{r_{j+1}^\lambda} \right) \geq C \int_{\zeta_{j+1}/r_{j+1}^\lambda}^{\zeta_j/r_j^\lambda} \frac{\tilde{h}(\sigma\zeta)}{\zeta} d\zeta$$

for all integers j satisfying (72), where

$$\tilde{h}(\zeta) = \inf_{(\zeta, 2^\lambda\zeta)} h.$$

Thus, (76) implies the estimate

$$\int_{\zeta_j}^{\zeta_{j+1}} \frac{d\zeta}{\zeta^{\gamma-p+1}} \geq C \int_{\zeta_{j+1}/r_{j+1}^\lambda}^{\zeta_j/r_j^\lambda} \frac{\tilde{h}(\sigma\zeta)}{\zeta} d\zeta$$

for all integers j satisfying (72), whence it follows that

$$\sum_{i=1}^{\infty} \int_{\zeta_{s_i}}^{\zeta_{l_i}} \frac{d\zeta}{\zeta^{\gamma-p+1}} \geq C \sum_{i=1}^{\infty} \int_{\zeta_{l_i}/r_{l_i}^\lambda}^{\zeta_{s_i}/r_{s_i}^\lambda} \frac{\tilde{h}(\sigma\zeta)}{\zeta} d\zeta.$$

Combining this with (71) and the inequalities $\zeta_{s_{i+1}} \geq \zeta_{l_i}$ and $\zeta_{l_i}/r_{l_i}^\lambda \leq \zeta_{s_{i+1}}/r_{s_{i+1}}^\lambda$, $i = 1, 2, \dots$, we obtain

$$\int_{\zeta_{s_1}}^{\infty} \frac{d\zeta}{\zeta^{\gamma-p+1}} \geq C \int_0^{\zeta_{s_1}/r_{s_1}^\lambda} \frac{\tilde{h}(\sigma\zeta)}{\zeta} d\zeta. \quad (77)$$

In view of the choice of the real number r_0 , one can assert that $E(r_{s_1}) \geq E(r_0) > 0$. At the same time, $\gamma > p$ in accordance with (13). Hence,

$$\int_{\zeta_{s_1}}^{\infty} \frac{d\zeta}{\zeta^{\gamma-p+1}} < \infty.$$

In so doing, from the monotonicity of the function f , it follows that

$$\tilde{h}(\zeta) \geq \frac{1}{2^{\lambda\gamma}} \frac{f(\zeta)}{\zeta^\gamma}, \quad \zeta > 0. \quad (78)$$

Thus, (77) implies the relation

$$\int_0^{\zeta_{s_1}/r_{s_1}^\lambda} \frac{f(\sigma\zeta)}{\zeta^{1+\gamma}} d\zeta < \infty$$

which contradicts (12).

Now, we consider the case of $1 - m/n < p < 1$. Let us show that

$$\frac{E(r_{i+1}) - E(r_i)}{E^\gamma(r_i)} \geq Ch \left(\frac{\sigma E(r_i)}{r_i^{n-(k-1)\varkappa}} \right) \quad (79)$$

for all $i = 0, 1, 2, \dots$. Indeed, by Lemma 9, for any integer $i \geq 0$ at least one of the following two systems of inequalities is satisfied:

$$\left\{ \begin{array}{l} \frac{1}{\text{mes } Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} \int_{Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} |u| dx dt \geq \frac{\sigma}{r_i^{n-(k-1)\varkappa}} \int_{Q_{r_i}^\varkappa} f(|u|) dx dt, \\ E(r_{i+1}) - E(r_i) \geq Cr_i^{n+\varkappa} f \left(\frac{\sigma E(r_i)}{r_i^{n-(k-1)\varkappa}} \right), \end{array} \right. \quad (80)$$

$$\left\{ \begin{array}{l} \left(\frac{1}{\text{mes } Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} \int_{Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} |u| dx dt \right)^p \geq \frac{\sigma}{r_i^{n+\varkappa-m}} \int_{Q_{r_i}^\varkappa} f(|u|) dx dt, \\ E(r_{i+1}) - E(r_i) \geq Cr_i^{n+\varkappa} f \left(\left(\frac{\sigma E(r_i)}{r_i^{n+\varkappa-m}} \right)^{1/p} \right). \end{array} \right. \quad (81)$$

In the case where (80) is valid for some integer $i \geq 0$, the second inequality in (80) and the first equation in (64) immediately imply (79). In its turn, if (81) is valid for some integer $i \geq 0$, then (65) and the second inequality in (81) yield

$$E(r_{i+1}) - E(r_i) \geq Cr_i^{n+\varkappa} f \left(\frac{\sigma E^{1/p}(r_i)}{r_i^{n-(k-1)\varkappa}} \right),$$

whence in accordance with the monotonicity of the function f and the fact that

$$E^{1/p}(r_i) \geq E^{1/p-1}(r_0)E(r_i) \quad (82)$$

we obtain

$$E(r_{i+1}) - E(r_i) \geq Cr_i^{n+\varkappa} f\left(\frac{\sigma E(r_i)}{r_i^{n-(k-1)\varkappa}}\right).$$

According to the first equation in (64) and the definition of function h , this again leads us to (79).

It can also be shown that

$$\frac{1}{\text{mes } Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} \int_{Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} |u| dx dt \geq \frac{\sigma E(r_i)}{r_i^{n-(k-1)\varkappa}} \quad (83)$$

for all $i = 0, 1, 2, \dots$. Indeed, if (80) is satisfied for some integer $i \geq 0$, then (83) follows directly from the first inequality in (80). In the case where (81) holds, the first inequality in (81) implies that

$$\frac{1}{\text{mes } Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} \int_{Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} |u| dx dt \geq \frac{\sigma E^{1/p}(r_i)}{r_i^{(n+\varkappa-m)/p}},$$

whence in accordance with (65) and (82) we again arrive at (83).

By Lemma 5 and estimate (56), the left-hand side of (83) tends to zero as $i \rightarrow \infty$. Thus, passing in (83) to the limit as $i \rightarrow \infty$, one can conclude that

$$\lim_{i \rightarrow \infty} \frac{E(r_i)}{r_i^{n-(k-1)\varkappa}} = 0. \quad (84)$$

We take

$$\zeta_i = E(r_i), \quad i = 0, 1, 2, \dots,$$

this time. In so doing, let $\lambda = n - (k-1)\varkappa$ as before. Then inequality (79) can obviously be written in the form

$$\frac{\zeta_{i+1} - \zeta_i}{\zeta_i^\gamma} \geq Ch \left(\frac{\sigma \zeta_i}{r_i^\lambda} \right), \quad i = 0, 1, 2, \dots \quad (85)$$

Thus, repeating the proof of estimate (77) with (75) replaced by (85), we obtain

$$\int_{\zeta_{s_1}}^\infty \frac{d\zeta}{\zeta^\gamma} \geq C \int_0^{\zeta_{s_1}/r_{s_1}^\lambda} \frac{\tilde{h}(\sigma\zeta)}{\zeta} d\zeta.$$

The left-hand side of the last expression is less than infinity as

$$\gamma = \frac{n + \varkappa}{n - \varkappa(k-1)} > 1$$

according to the first equation in (64), whereas

$$\int_0^{\zeta_{s_1}/r_{s_1}^\lambda} \frac{\tilde{h}(\sigma\zeta)}{\zeta} d\zeta \geq \frac{1}{2^{\lambda\gamma}} \int_0^{\zeta_{s_1}/r_{s_1}^\lambda} \frac{f(\sigma\zeta)}{\zeta^{1+\gamma}} d\zeta = \infty \quad (86)$$

in view of (78) and condition (12). This contradiction proves the theorem. \square

Proof of Theorem 3. Assume the converse. Let u be a non-trivial global weak solution of (1), (2) and let q be some real number such that $p \leq q < p/(1 - m/n)$. We put

$$\varkappa = \frac{(p/q - 1)n + m}{1 + (k - 1)p/q} \quad (87)$$

and

$$\gamma = \frac{np + mq/k}{n - m(1 - 1/k)}. \quad (88)$$

It is easy to see that $\varkappa > 0$ and $\gamma > 0$. In so doing, \varkappa and γ satisfy the system of equations

$$\begin{cases} n + \varkappa - \frac{\gamma}{q}(n - (k - 1)\varkappa) = 0, \\ n + \varkappa - \frac{\gamma}{p}(n + \varkappa - m) = 0. \end{cases} \quad (89)$$

In particular, this system implies that

$$\frac{n + \varkappa - m}{p} = \frac{n - (k - 1)\varkappa}{q}. \quad (90)$$

Take a real number $r_0 > 0$ such that $E(r_0) > 0$, where the function E is defined by (60). Also let $r_i = 2^i r_0$, $i = 1, 2, \dots$. As in the proof of Theorem 2, we assume that the constants C and σ below can also depend on $E(r_0)$.

Let us show that relation (84) remains valid. Indeed, by Lemma 9, for any integer $i \geq 0$ at least one of the two systems of inequalities (80) or (81) is satisfied. In the case where (80) holds, the first inequality in (80) implies (83). In its turn, if (81) holds, then the first inequality in (81) yields

$$\left(\frac{1}{\text{mes } Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} \int_{Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} |u| dx dt \right)^q \geq \frac{\sigma E^{q/p}(r_i)}{r_i^{q(n+\varkappa-m)/p}},$$

whence in accordance with (90) and the fact that $E^{q/p}(r_i) \geq E^{q/p-1}(r_0)E(r_i)$, we have

$$\left(\frac{1}{\text{mes } Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} \int_{Q_{r_{i+1}}^\varkappa \setminus Q_{r_i}^\varkappa} |u| dx dt \right)^q \geq \frac{\sigma E(r_i)}{r_i^{n-(k-1)\varkappa}}. \quad (91)$$

Hence, to establish the validity of (84), it sufficient to note that the left-hand sides of (83) and (91) tend to zero as $i \rightarrow \infty$ by Lemma 5 and relation (56).

Now, we show the validity of the estimate

$$\frac{E(r_{i+1}) - E(r_i)}{E^{\gamma/q}(r_i)} \geq Ch \left(\sigma \left(\frac{E(r_i)}{r_i^{n-(k-1)\varkappa}} \right)^{1/q} \right) \quad (92)$$

for all $i = 0, 1, 2, \dots$, where h is defined by (66). Indeed, in view of (84), there exists an integer $i_0 \geq 0$ such that

$$\frac{E(r_i)}{r_i^{n-(k-1)\varkappa}} \leq 1 \quad (93)$$

for all $i \geq i_0$. Without loss of generality, it can be assumed that $i_0 = 0$; otherwise we replace r_0 by r_{i_0} .

If (80) holds, then in accordance with (93) and the second inequality in (80) we obtain

$$E(r_{i+1}) - E(r_i) \geq Cr_i^{n+\varkappa} f \left(\sigma \left(\frac{E(r_i)}{r_i^{n-(k-1)\varkappa}} \right)^{1/q} \right). \quad (94)$$

By the first equation in (89), this immediately implies (92). In its turn, if (81) is satisfied, then (90) and the second inequality in (81) yield

$$E(r_{i+1}) - E(r_i) \geq Cr_i^{n+\varkappa} f \left(\frac{\sigma E^{1/p}(r_i)}{r_i^{(n-(k-1)\varkappa)/q}} \right).$$

Since $E^{1/p}(r_i) \geq E^{1/p-1/q}(r_0)E^{1/q}(r_i)$, this again leads us to (94), whence (92) follows at once.

Let us denote $\lambda = (n - (k-1)\varkappa)/q$ and

$$\zeta_i = E^{1/q}(r_i), \quad i = 0, 1, 2, \dots$$

Taking into account (87) and the condition $n > m(1 - 1/k)$, we have

$$\lambda = \frac{k(n - m(1 - 1/k))}{q + (k-1)p} > 0.$$

In the above notation, (92) can obviously be written as

$$\frac{\zeta_{i+1}^q - \zeta_i^q}{\zeta_i^\gamma} \geq Ch \left(\frac{\sigma \zeta_i}{r_i^\lambda} \right), \quad i = 0, 1, 2, \dots \quad (95)$$

Thus, repeating the proof of estimate (77) with (75) replaced by (95), we obtain

$$\int_{\zeta_{s_1}}^{\infty} \frac{d\zeta}{\zeta^{\gamma-q+1}} \geq C \int_0^{\zeta_{s_1}/r_{s_1}^\lambda} \frac{\tilde{h}(\sigma\zeta)}{\zeta} d\zeta.$$

From (88), it follows that $\gamma \rightarrow p/(1 - m/n)$ as $q \rightarrow p/(1 - m/n)$. Therefore, the real number $q \in [p, p/(1 - m/n))$ can be chosen such that $\gamma > \mu$. In view of (14) and (78), this implies (86). On the other hand, from the first equation in (89), it follows that $\gamma > q$; therefore,

$$\int_{\zeta_{s_1}}^{\infty} \frac{d\zeta}{\zeta^{\gamma-q+1}} < \infty.$$

This contradiction completes the proof. \square

Proof of Theorem 4. It is obvious that (1), (2) can have only trivial solutions. This follows from Theorem 1 if $n \leq m(1 - 1/k)$ or from Theorems 2 and 3 if $n > m(1 - 1/k)$. At the same time, problem (1), (2) can not have trivial solutions. Really, if $f(0) > 0$, then for any non-negative function $\varphi \in C_0^\infty(\mathbb{R}_+^{n+1})$ with a non-empty support the right-hand side of (3) is positive, whereas the left-hand side is equal to zero. \square

Remark 3. *Theorems 1–4 remain valid if, instead of the non-negativity of the initial value u_{k-1} , one requires that*

$$U(r) = \int_{B_r} u_{k-1} dx$$

is a non-decreasing non-negative function in a neighborhood of infinity. In this case, we have

$$U'(r) = \int_{S_r} u_{k-1} dS_r \geq 0$$

for almost all r in a neighborhood of infinity, where S_r is the sphere of radius $r > 0$ centered at zero and dS_r is the volume element on S_r . Therefore, Lemma 1 and all subsequent statements remain valid if the real number R in their conditions is large enough. Indeed, taking the test function φ such as in the proof of Lemma 1, we obtain that the first summand on the right in (3) is non-negative. This is sufficient for the validity of inequality (23).

Acknowledgments The work of the first author was supported by the Russian Ministry of Education and Science as part of the program of the Moscow Center for Fundamental and Applied Mathematics under the agreement 075-15-2022-284 (critical exponents), and by the Russian Science Foundation, project 20-11-20272-II (estimates of global solutions). The work of the second author was supported by the Russian Science Foundation, project 23-11-00056 (asymptotic properties of solutions).

Data availability Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

Declarations

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

- [1] Cai, G., Pan, H., Xing, R.: A note on parabolic Liouville theorems and blow-up rates for a higher-order semilinear parabolic system. *Inter. J. Diff. Eq.* Article ID 896427 (2011). <https://doi.org/10.1155/2011/896427>
- [2] Caristi, G.: Nonexistence theorems for systems of quasilinear parabolic inequalities. *Adv. Nonlinear Stud.* **1**(1), 143–158 (2001).
- [3] Egorov, Yu.V., Galaktionov, V.A., Kondratiev, V.A., Pohozaev, S.I.: On the necessary conditions of global existence of solutions to a quasilinear inequality in the half-space. *C. R. Acad. Sci. Paris, Sér. 1: Math.* **330**, 93–98 (2000).
- [4] Egorov, Yu.V., Galaktionov, V.A., Kondratiev, V.A., Pohozaev, S.I.: Conditions for the existence of solutions to a quasilinear inequality in half-space. *Math. Notes* **67**(1), 119–121 (2000).
- [5] Fujita, H.: On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **13**, 109–124 (1966).
- [6] Galaktionov, V.A., Shishkov, A.E.: Higher-order quasilinear parabolic equations with singular initial data. *Comm. Cont. Math.* **8**(3), 1–24 (2006).
- [7] Hayakawa, K.: On nonexistence of global solutions of some semilinear parabolic differential equations. *Proc. Japan Acad.* **49**, 503–505 (1973).
- [8] Kato, T.: Blow-up of solutions of some nonlinear hyperbolic equations. *Commun. Pure Appl. Math.* **33**, 501–505 (1980).
- [9] Keller, J.B.: On solution of $\Delta u = f(u)$. *Comm. Pure. Appl. Math.* **10**, 503–510 (1957).
- [10] Kiguradze, I.T., Kvinikadze, G.G.: *On strongly increasing solutions of nonlinear ordinary differential equations*, *Ann. Mat. Pure Appl.* **130**, 67–87 (1982).
- [11] Kondratiev, V.A., Véron, L.: Asymptotic behavior of solutions of some nonlinear parabolic or elliptic equations. *Asympt. Anal.* **14**, 117–156 (1997).
- [12] Kon'kov, A.A.: On the asymptotic behaviour of solutions of nonlinear parabolic equations. *Proc. Royal Soc. Edinburgh* **136**, 365–384 (2006).
- [13] Kon'kov, A.A.: On solutions of non-autonomous ordinary differential equations. *Izv. Math.* **65**, 285–327 (2001).
- [14] Kon'kov, A.A., Shishkov, A.E.: Generalization of the Keller-Osserman theorem for higher order differential inequalities. *Nonlinearity* **32**, 3012–3022 (2019).

- [15] Laister R, Sierżęga M.: A blow-up dichotomy for semilinear fractional heat equations. *Math. Ann.* **381**, 75–90 (2021).
- [16] Mitidieri, E., Pohozaev, S.I.: A priori estimates and blow-up of solutions to nonlinear partial differential equations and inequalities. *Proc. V.A. Steklov Inst. Math.* **234**, 3–383 (2001).
- [17] Osserman, R.: On the inequality $\Delta u \geq f(u)$. *Pacific J. Math.* **7**, 1641–1647 (1957).
- [18] Palmieri, A., Takamura, H.: A blow-up result for a Nakao-type weakly coupled system with nonlinearities of derivative-type. *Math. Ann.* **387**, 111–132 (2023).
- [19] Pohozaev, S.I., Tesei, A.: Blow-up of nonnegative solutions to quasilinear parabolic inequalities. *Atti Accad. Naz. Lincei. Cl. Sci. Fis. Mat. Natur. Rend. Lincei. Ser. 9.* **11**, 99–109 (2000).
- [20] Pohozaev, S.I., Véron, L.: Blow-up results for nonlinear hyperbolic inequalities. *Ann. Scuola Norm. Super. Pisa. Cl. Ser. 4.* **29**(2), 393–420 (2001).
- [21] Pucci, P., Serrin, J.: Global nonexistence for abstract evolution equations with positive initial energy. *J. Diff. Equat.* **150**(1), 203–214 (1998).
- [22] Samarskii, A.A., Galaktionov, V.A., Kurdyumov, S.P. Mikhailov, A.P.: *Blow-Up in Quasilinear Parabolic Equations*. De Gruyter, Berlin, New York (1995).
- [23] Shishkov, A., Véron, L.: The balance between diffusion and absorption in semilinear parabolic equations. *Rend. Lincei Mat. Appl.* **18**, 59–96 (2007).