## Model Selection with Baryonic Acoustic Oscillations in the Lyman- $\alpha$ Forest

F. Melia

Department of Physics, The Applied Math Program, and Department of Astronomy,

Department of Physics, The Applied Math Program, and L The University of Arizona, AZ 85721, USA PACS 98.80.Cq – Inflation PACS 98.80.Cq – Inflation PACS 98.80.Jk – Cosmology PACS 98.80.Jk – Relativistic Astrophysics Abstract –The recent release of the final, complete cillation measurements provides the most significant a geometry at an effective redshift  $z_{eff} = 2.334$ , which is we use these data to select among four distinct cosmol the Milne universe and Einstein-de Sitter. Given the this BAO measurement alone provides a strong mod studies that combined Lyman-α data with measurem proaches are useful, the latter tends to dilute the di observations. We therefore examine how the models of BAO scale measured in the Lyman-α forest and bac Einstein-de Sitter are strongly ruled out by these dat ing the standard model. The Lyman-α measurements geometry predicted by  $R_h = ct$ . As such, evidence c condition from general relativity ought to be an essent the radiation began dissociating from baryonic matter. Believed to represent a sonic horizon,  $r_d$ , this comoving distance is apparently associated with sound waves cre-Abstract – The recent release of the final, complete survey of Lyman- $\alpha$  baryonic acoustic oscillation measurements provides the most significant and accurate data base for studying cosmic geometry at an effective redshift  $z_{\text{eff}} = 2.334$ , which is inaccessible to other sources. In this Letter, we use these data to select among four distinct cosmologies: Planck ACDM, the  $R_{\rm h} = ct$  universe, the Milne universe and Einstein-de Sitter. Given the breadth and depth of the Lyman- $\alpha$  study, this BAO measurement alone provides a strong model comparison, complementary to previous studies that combined Lyman- $\alpha$  data with measurements at lower redshifts. Though both approaches are useful, the latter tends to dilute the disparity between model predictions and the observations. We therefore examine how the models compare to each other strictly based on the BAO scale measured in the Lyman- $\alpha$  forest and background quasars. We find that Milne and Einstein-de Sitter are strongly ruled out by these data. There is also strong evidence disfavoring the standard model. The Lyman- $\alpha$  measurements are completely consistent with the cosmic geometry predicted by  $R_{\rm h} = ct$ . As such, evidence continues to grow that the zero active mass condition from general relativity ought to be an essential ingredient in  $\Lambda$ CDM.

The radiation began dissociating from baryonic matter. Believed to represent a sonic horizon,  $r_d$ , this comoving distance is apparently associated with sound waves created at or near the Planck regime, which then propagated across the baryon-photon fluid until the CMB was pro-duced [1-3].

In a Friedmann-Lemaître-Robertson-Walker (FLRW) cosmology, in which all proper (or physical) distances grow in proportion to a universal scale factor, a(t), while comoving distances remain constant, the scale  $r_d$  is expected to have remained unchanged, reappearing subsequently as a characteristic length in the matter correlation function. In this context, the sonic scale is more commonly inferred from baryonic acoustic oscillations (BAO) seen in the large-scale structure, the first measurement of which was carried out using the auto-correlation of galaxy positions [4] at  $z \sim 0.35$ , and the galaxy power spectrum [5] at  $z \sim 0.1$ . More generally, the BAO scale at  $z \leq 2$  has been studied using a variety of discrete tracers, such as galaxy clusters [6], and quasars [7], in addition to the aforementioned galaxies [8–15].

Beyond  $z \sim 2$ , however, a measurement of the BAO scale must to be handled differently because the number density of observable dicrete tracers is too low for high precision clustering studies. The method of choice for measuring  $r_d$  in the high-redshift Universe instead rests on the observation of opacity fluctuations in the Lyman- $\alpha$ forest irradiated by background quasars. The first such studies, focusing on the Lyman- $\alpha$  auto-correlation function, were carried out by refs. [16], [17], [18], [19], [20] and [21]. Complementary results based on the Lyman- $\alpha$  and quasar cross-correlation function have also been reported by refs. [22], [23] and [24].

The BAO feature is a powerful diagnostic in cosmology because it yields both angular-diameter distances and the expansion rate normalized to the sound horizon  $r_d$ . And while the luminosity distance to type Ia supernovae has already provided strong evidence for the existence of dark energy [25–27], these transient events are difficult to observe at redshifts  $z \gtrsim 1.8$ .

The BAO measurements in the Lyman- $\alpha$  forest, at an effective  $z \sim 2.334$ , may therefore be used in several distinct tests of the geometry of the Universe at intermediate redshifts not accessible to local surveys focusing on supernovae ( $z \leq 1.8$ ) and instruments designed to study the CMB at  $z \gg 1$ .

A common approach is to combine all of the BAO data, those from the galaxy surveys at low redshifts and those from the Lyman- $\alpha$  forest farther away, under the assumption that  $r_d$  is independent of z. This can be done either in the context of  $\Lambda CDM$ , where one includes the predicted value of  $r_d$  in this model to extract the redshift-dependent angular-diameter distance and expansion rate, or in model selection by examining which cosmology is preferred by the BAO measurements. For the latter, one considers  $r_d$  to be 'unanchored,' since its value may not be the same from one model to the next. In this case, one either optimizes  $r_d$ along with the other model parameters individually for each cosmology being tested, or avoids it altogether by considering ratios of the angular-diameter and Hubble distances, both of which are proportional to the sonic horizon (see Eqs. 3 and 4 below).

In previous work [28–30], we have followed the latter approach using older, less complete BAO catalogs than those available now to compare the standard model with one of its principal competitors known as the  $R_{\rm h} = ct$  universe [31,32]. Interestingly, the aggregated BAO data have tended to favour the latter model rather than  $\Lambda$ CDM. But the BAO measurements using the lower redshift galaxy surveys are less discerning than their higher redshift counterparts, so while the outcome of these studies has been suggestive, it has not necessarily been compelling.

In view of the significantly more complete eBOSS catalog available now (see section labeled 'Data' below), however, the BAO data at z = 2.334 by themselves should constitute an important constraint on the geometry of the Universe sampled solely by the quasars in this catalog (distributed at  $z \gtrsim 1.77$ ) and their Lyman- $\alpha$  forests. Since it is well known that a comparison of the model predictions and the BAO measurements tends to become more discordant with increasing redshift, it is desirable to use the Lyman- $\alpha$  data by themselves for model selection purposes, in parallel to the already completed studies based on the BAO observations at various redshifts.

In this Letter, we therefore complement our previous model selection analysis by restricting the comparison to just the Lyman- $\alpha$  measurements. As noted, we do this for two principal reasons. First, the model contrast at  $z \gtrsim 2$  is significantly greater than that at  $z \sim 0$ . Second, we now have the final BAO constraints from the Lyman- $\alpha$  auto-correlation and Lyman- $\alpha$ -quasar cross-correlation functions produced from the sixteenth (DR16) and final release [33] of the fourth generation Sloan Digital Sky Survey (SDSS-IV), containing all of the clustering and Lyman- $\alpha$  data from the completed 'extended' Baryonic Oscillation Spectroscopic Survey (eBOSS) [34]. This catalog of quasars and Lyman- $\alpha$  profiles is so large (see 'Data' below) that the BAO scale measured from it at an effective redshift ~ 2.334 constitutes a crucial probe of the cosmology on its own merit. As we shall see, these data alone provide an important comparison of cosmological models and their predictions without having to pre-assume  $r_d$  and  $H_0$ . The latter is especially desirable in view of the growing disparity between the measurements of  $H_0$  at low and high redshifts, creating a ~  $4\sigma$  uncertainty in its value [35].

We shall find that the depth and extent of the final Lyman- $\alpha$  BAO analysis [36], along with the option of altogether avoiding the use of  $r_d$  and  $H_0$ , provides us with a very clean and compelling test of the various FLRW cosmologies.

**Data.** – The analysis in this Letter utilizes the baryonic acoustic oscillations (BAO) measured in the Lyman- $\alpha$  absorption and background quasars, with an effective (enemble) redshift z = 2.334. The data are taken from the complete extended Baryonic Oscillation Spectroscopic Survey (eBOSS) [36], which includes the Lyman- $\alpha$  absorption profiles of 210,005 background quasars distributed at  $z_q > 2.10$ . The BAO scale has been measured in both the auto-correlation of the Lyman- $\alpha$  absorbers and their crosscorrelation with 341,468 quasars at  $z_q > 1.77$ . This data release represents several advances over previously published Lyman- $\alpha$  BAO measurements, including improved statistics from a larger quasar catalog and deeper observations, and a more accurate modeling of the systematics.

**Model Comparisons.** – The Lyman- $\alpha$  BAO survey measures the BAO scale in the Lyman- $\alpha$  forest of absorption of light from distant quasars, both via the forestforest correlation function [19], and in the forest-quasar cross-correlation [22]. The scale is measured along the line-of-sight,

$$\Delta z = \frac{r_d}{(1+z) d_H(z)} , \qquad (1)$$

and in the transverse direction,

$$\Delta \theta = \frac{r_d}{d_M(z)} \tag{2}$$

where, as previously noted,  $r_d$  is the length corresponding to the peak of the matter two-point function in comoving coordinates. In addition, the quantity

$$d_H(z) \equiv \frac{c}{H(z)} \tag{3}$$

is the Hubble scale at z, while  $d_M(z)$  is given in terms of the angular-diameter distance via the relation

$$d_M(z) \equiv (1+z) d_A(z)$$
. (4)

We mention here that surveys often also define the distance

$$d_V(z) \equiv z^{1/3} d_M^{2/3}(z) \, d_H^{1/3}(z) \,, \tag{5}$$

Data/Model	$d_M/d_H$ (P-value)
Data	$4.17\pm0.18$
<ol> <li>R<sub>h</sub> = ct</li> <li>Planck-ΛCDM</li> <li>Milne</li> <li>Einstein-de Sitter</li> </ol>	$\begin{array}{rrrr} 4.02 & (0.39) \\ 4.56 & (0.03) \\ 5.058 & (< 0.00001) \\ 5.507 & (< 0.00001) \end{array}$

Table 1: Model comparison using the Lyman- $\alpha$  BAO data

an angle-weighted average of  $d_M(z)$  and  $d_H(z)$ , but this quantity was not derived independently of  $d_M$  and  $d_H$  for these data [36]. Thus, though it tends to be the best determined scale in measurements of BAO from galaxy surveys, we will not find it useful for our analysis in this particular instance, and we shall instead focus our attention on  $d_M$ and  $d_H$  themselves.

Given that these BAO observables are proportional to  $r_d H_0$  (see below for the model-dependent functional form of H[z]), the ratios of the various BAO scales are completely independent of the sonic horizon and Hubble constant. Indeed, as long as we restrict our comparisons to these ratios, three of the models we compare here have no free parameters at all, while the standard model,  $\Lambda$ CDM (see Eq. 11 below), has just one: the scaled matter density  $\Omega_{\rm m}$ . But to avoid unduly 'punishing' this cosmology by treating  $\Omega_{\rm m}$  as an unknown variable when assessing its likelihood, we shall simply adopt its *Planck* optimized value,  $\Omega_{\rm m} = 0.315 \pm 0.007$ , with a spatial curvature constant k = 0 [3].

We shall compare fits to the Lyman- $\alpha$  data using four different cosmological models, each with its own prediction of the angular-diameter and Hubble distances. The quantity  $\Omega_i$  is the energy density of species *i*, scaled to the critical density

$$\rho_{\rm c} \equiv \frac{3c^2 H_0^2}{8\pi G} \ , \tag{6}$$

in terms of the Hubble constant,  $H_0 = 67.4 \pm 0.5$  km s<sup>-1</sup> Mpc<sup>-1</sup>. In principle, we could completely avoid any reliance on the *Planck* measurements by selecting the value of  $\Omega_{\rm m}$  that optimizes  $\Lambda$ CDM's fit to the Lyman- $\alpha$  BAO data, but the improvement is too small to justify the introduction of an additional free parameter. The four models we compare are:

1. The  $R_{\rm h} = ct$  universe, a Friedmann-Lemaître-Robertson-Walker cosmology with zero active mass,  $\rho + 3p = 0$ , in terms of the total energy density  $(\rho)$ and pressure (p) in the cosmic fluid [31, 32]. In this case,

$$d_M^{(1)}(z) = \frac{c}{H_0} \ln(1+z) , \qquad (7)$$

and

$$d_H^{(1)}(z) = \frac{c}{H_0(1+z)} \,. \tag{8}$$

2. Flat *Planck*- $\Lambda$ CDM, with  $\Omega_{\Lambda} = 1 - \Omega_{\rm m}$  and a darkenergy equation of state parameter  $w_{\rm de} = -1$ . For this model,

$$d_M^{(2)}(z) = \frac{c}{H_0} \int_0^z \frac{du}{E(u)} , \qquad (9)$$

and

$$d_H^{(2)}(z) = \frac{c}{H_0 E(z)} , \qquad (10)$$

where

$$E(z) \equiv \left[\Omega_{\rm m}(1+z)^3 + \Omega_{\Lambda}\right]^{1/2} . \tag{11}$$

At these redshifts, the contribution to E(z) from radiation is negligible, so we have ignored it for the calculation of  $d_H(z)$  and  $d_M(z)$ .

3. The Milne universe. This well studied solution is also a Friedmann-Lemaître-Robertson-Walker cosmology, but its energy density, pressure and cosmological constant are all zero. Its expansion instead derives from a non-zero spatial curvature, with k = -1. Like the  $R_{\rm h} = ct$  universe, the Milne scale factor, a(t), is also linear in time [37,38], but we include it here primarily because the observable signatures in these two models are very different. In the Milne universe,

$$d_M^{(3)}(z) = \frac{c}{H_0} \sinh\left[\ln(1+z)\right] , \qquad (12)$$

and

$$d_H^{(3)}(z) = \frac{c}{H_0(1+z)} \,. \tag{13}$$

4. Einstein-de Sitter (i.e., Eqs. 9 and 10 with  $\Omega_{\rm m} = 1$ and  $\Omega_{\Lambda} = 0$ ):

$$d_M^{(4)}(z) = \frac{2c}{H_0} \left( 1 - \frac{1}{\sqrt{1+z}} \right) , \qquad (14)$$

and

$$d_H^{(4)}(z) = \frac{c}{H_0(1+z)^{3/2}} \,. \tag{15}$$

This cosmology is already heavily disfavoured by many other observations, but we include it here because of its relevance as a former 'standard' model.

The combined BAO measurements from the auto- and cross-correlation in the final, complete eBOSS release [36] yield the following constraints:

$$d_H(z = 2.334)/r_d = 8.99 \pm 0.19$$
  
$$d_M(z = 2.334)/r_d = 37.5 \pm 1.1.$$
(16)

Though a 'fiducial' cosmology is employed in the calculation of these distances, the ratios shown in Equation (16) are model independent, as studied in detail by ref. [39] in the context of galaxy correlations, and confirmed for the Lyman- $\alpha$  observations by ref. [36].

The model predictions are compared with these data in Table 1, prioritized in terms of the p-values estimated from the various fits. Given a model's prediction for  $R^{\rm th} \equiv d_M/d_H$  and the standard deviation (in this case,  $\sigma_R = 0.18$ ) of the measurement, this p-value represents the probability of observing a difference  $|R^{obs} - \bar{R}^{th}|$  greater than  $|4.17 - R^{\text{th}}|$  under the assumption that the null hypothesis is true and that the distribution of  $R^{\rm obs}$  is normal. The error quoted for the measured value of  $d_M/d_H$ includes the correlation between  $d_M/r_d$  and  $d_H/r_d$ , characterized by the correlation coefficient  $C(d_H, d_M) = -0.45$ [36]. Small p-values provide evidence against the assumed model, with the evidence getting stronger as the p-value approaches zero. Typical guidelines suggest the following hierarchy: (p > 0.10) weak or no evidence; (0.05 moderate evidence; <math>(0.01strong evidence;  $(p \le 0.01)$  very strong evidence.

On the basis of these comparisons, it is clear that the Lyman- $\alpha$  BAO measurements very strongly rule out the Milne and Einstein-de Sitter cosmologies, affirming an already established conclusion drawn from many other comparative tests [32]. There is also strong evidence disfavoring the standard model, an outcome confirmed by several previous studies, including those reported in refs. [36] and [40]. But the principal result of this work is that the Lyman- $\alpha$  BAO measurements are completely consistent with the geometry of the cosmos predicted by the  $R_{\rm h} = ct$  universe.

Pursuing this head-to-head comparison further, we note that if  $\Omega_{\rm m}$  were to differ from its *Planck* value by  $3\sigma$ , i.e., if  $\Omega_{\rm m} = 0.294$ , then the p-value for  $\Lambda$ CDM would improve somewhat to 0.065, so the evidence against the standard model in that case would be 'moderate' instead of 'strong.' Nevertheless, to make the standard model equally likely with  $R_{\rm h} = ct$ , i.e., to improve its p-value to 0.39,  $\Omega_{\rm m}$  would have to be smaller than ~ 0.236, a value different by many standard deviations from that inferred by *Planck*.

It is also straightforward to perform a Bayesian analysis [41, 42] of the head-to-head comparison between  $\Lambda$ CDM and  $R_{\rm h} = ct$  based on the Lyman- $\alpha$  data at z = 2.334, specifically, the measurement of  $d_M/d_H = 4.17 \pm 0.18$  shown in Table 1. Under the assumption that this ratio follows a normal distribution, and adopting a point model for both the null and alternative hypotheses (remember that we are fixing the value of  $\Omega_{\rm m}$  at the *Planck* measurement to avoid unduly 'punishing' the standard model for having an additional free parameter), one infers a Bayes factor of 7.39. The marginal likelihoods at the measured value of  $d_M/d_H$ , from which the Bayes factor is estimated, are shown in Figure 1.

A Bayes factor between 3 and 10 (as we have here) indicates 'moderate' or 'substantial' evidence in favor of the alternative hypothesis [43–45]. As such, the Bayesian analysis complements and confirms our earlier conclusion, derived from the p-values, that the Lyman- $\alpha$  BAO data



Fig. 1: Estimation of the Bayes factor from the marginal prediction of the  $R_{\rm h} = ct$  and  $\Lambda {\rm CDM}$  models versus the measurement of the ratio  $d_M/d_H$  from the Lyman- $\alpha$  forest at an effective redshift z = 2.334.

at z = 2.334 favor the  $R_{\rm h} = ct$  cosmology over the current standard model.

**Conclusion.** – The strong rejection of the Milne and Einstein-de Sitter cosmologies by the final Lyman- $\alpha$  BAO data release is hardly surprising in view of their similarly strong rejection by other observations. Our main conclusion from this study instead refocuses our attention on the fact that the data tend to favour the  $R_{\rm h} = ct$  cosmology over the current standard model.

Indeed, the BAO measurements offer a new perspective on this comparison, following an earlier examination by ref. [40], who attempted to identify the reason behind the BAO anomaly in the context of  $\Lambda$ CDM. Their analysis showed that the BAO data at z > 0.43 are in tension with the standard model, whether or not the *Planck* optimized parameters (e.g., for  $\Omega_{\rm m}$ ) are assumed. They concluded that this tension arises not from the  $\Lambda$ CDM parameters, but instead from the dark energy evolution at 0.57 < z <2.334. If one further sets  $r_d$  equal to the acoustic scale measured in the CMB, a cosmological constant for dark energy is firmly rejected.

The  $R_{\rm h} = ct$  cosmology is essentially  $\Lambda$ CDM, though with the crucial additional constraint of zero active mass from general relativity [32]. This constraint features an equation-of-state,  $\rho + 3p = 0$ , in terms of the total energy density  $\rho$  and pressure p in the cosmic fluid. Dark energy is therefore dynamic in this model, evolving along with the other constituents, presumably an extension to the standard model of particle physics.

The current standard model suffers from several major conflicts and inconsistencies that continue to defy serious attempts at mitigation [46]. Together with the continued success of  $R_{\rm h} = ct$  to account for the data better than  $\Lambda$ CDM [47], as affirmed by the work reported in this *Letter*, the prospects for further development of this alternative FLRW cosmology look very promising. An especially exciting future observation to anticipate over the coming decade is the real-time measurement of redshift drift [48,49], which should provide an unambiguous yes/no answer to the question of whether or not the cosmic fluid is in fact driven by a zero active mass equation-of-state.

## REFERENCES

- [1] PEEBLES P. J. E. and YU J. T., ApJ, **162** (1970) 815.
- [2] SUNYAEV R. A. and ZELDOVICH Y. B., Astrophys Sp Science, 7 (1970) 3.
- [3] PLANCK COLLABORATION, AGHANIM N., AKRAMI Y., ASHDOWN M., AUMONT J., BACCIGALUPI C., BALLAR-DINI M., BANDAY A. J., BARREIRO R. B., BARTOLO N., BASAK S. and ET AL., A&A, 641 (2020) A6.
- [4] EISENSTEIN D. J., ZEHAVI I., HOGG D. W., SCOCCI-MARRO R., BLANTON M. R., NICHOL R. C., SCRANTON R., SEO H.-J., TEGMARK M., ZHENG Z., ANDERSON S. F., ANNIS J., BAHCALL N., BRINKMANN J., BURLES S., CASTANDER F. J., CONNOLLY A., CSABAI I., DOI M., FUKUGITA M., FRIEMAN J. A., GLAZEBROOK K., GUNN J. E., HENDRY J. S. and ET AL., *ApJ*, **633** (2005) 560.
- [5] COLE S., PERCIVAL W. J., PEACOCK J. A., NORBERG P., BAUGH C. M., FRENK C. S., BALDRY I., BLAND-HAWTHORN J., BRIDGES T., CANNON R., COLLESS M., COLLINS C., COUCH W., CROSS N. J. G., DALTON G., EKE V. R., DE PROPRIS R., DRIVER S. P., EFSTATHIOU G., ELLIS R. S., GLAZEBROOK K., JACKSON C. and ET AL., MNRAS, 362 (2005) 505.
- [6] HONG T., HAN J. L. and WEN Z. L., ApJ, 826 (2016) 154.
- [7] ATA M., BAUMGARTEN F., BAUTISTA J., BEUTLER F., BIZYAEV D., BLANTON M. R., BLAZEK J. A., BOLTON A. S., BRINKMANN J., BROWNSTEIN J. R., BURTIN E. and ET AL., MNRAS, 473 (2018) 4773.
- [8] PERCIVAL W. J., COLE S., EISENSTEIN D. J., NICHOL R. C., PEACOCK J. A., POPE A. C. and SZALAY A. S., *MNRAS*, **381** (2007) 1053.
- [9] BLAKE C., KAZIN E. A., BEUTLER F., DAVIS T. M., PARKINSON D., BROUGH S., COLLESS M., CONTRERAS C., COUCH W., CROOM S., CROTON D., DRINKWA-TER M. J., FORSTER K., GILBANK D., GLADDERS M., GLAZEBROOK K. and ET AL., MNRAS, 418 (2011) 1707.
- [10] BEUTLER F., BLAKE C., COLLESS M., JONES D. H., STAVELEY-SMITH L., CAMPBELL L., PARKER Q., SAUN-DERS W. and WATSON F., MNRAS, 416 (2011) 3017.
- [11] PADMANABHAN N., XU X., EISENSTEIN D. J., SCALZO R., CUESTA A. J., MEHTA K. T. and KAZIN E., MNRAS , 427 (2012) 2132.
- [12] ANDERSON L., AUBOURG E., BAILEY S., BIZYAEV D., BLANTON M., BOLTON A. S., BRINKMANN J., BROWN-STEIN J. R., BURDEN A., CUESTA A. J. and ET AL., MNRAS, 427 (2012) 3435.
- [13] Ross A. J., SAMUSHIA L., HOWLETT C., PERCIVAL W. J., BURDEN A. and MANERA M., MNRAS, 449 (2015) 835.
- [14] ALAM S., ATA M., BAILEY S., BEUTLER F., BIZYAEV D., BLAZEK J. A., BOLTON A. S., BROWNSTEIN J. R., BUR-

DEN A., CHUANG C.-H., COMPARAT J., CUESTA A. J., DAWSON K. S., EISENSTEIN D. J., ESCOFFIER S., GIL-MARÍN H., GRIEB J. N., HAND N., HO S., KINEMUCHI K., KIRKBY D. and ET AL., *MNRAS*, **470** (2017) 2617.

- [15] BAUTISTA J. E., VARGAS-MAGAÑA M., DAWSON K. S., PERCIVAL W. J., BRINKMANN J., BROWNSTEIN J., CA-MACHO B., COMPARAT J., GIL-MARÍN H., MUELLER E.-M., NEWMAN J. A., PRAKASH A., ROSS A. J., SCHNEI-DER D. P., SEO H.-J., TINKER J., TOJEIRO R., ZHAI Z. and ZHAO G.-B., *ApJ*, 863 (2018) 110.
- [16] BUSCA N. G., DELUBAC T., RICH J., BAILEY S., FONT-RIBERA A., KIRKBY D., LE GOFF J. M., PIERI M. M., SLOSAR A., AUBOURG É., BAUTISTA J. E., BIZYAEV D. and ET AL., A&A, 552 (2013) A96.
- [17] SLOSAR A., IRŠIČ V., KIRKBY D., BAILEY S., BUSCA N. G., DELUBAC T., RICH J., AUBOURG É., BAUTISTA J. E., BHARDWAJ V., BLOMQVIST M. and ET AL., *JCAP*, **2013** (2013) 026.
- [18] KIRKBY D., MARGALA D., SLOSAR A., BAILEY S., BUSCA N. G., DELUBAC T., RICH J., BAUTISTA J. E., BLOMQVIST M., BROWNSTEIN J. R., CARITHERS B. and ET AL., JCAP, **2013** (2013) 024.
- [19] DELUBAC T., BAUTISTA J. E., BUSCA N. G., RICH J., KIRKBY D., BAILEY S., FONT-RIBERA A., SLOSAR A., LEE K.-G., PIERI M. M., HAMILTON J.-C., AUBOURG É., BLOMQVIST M., BOVY J., BRINKMANN J., CARITHERS W., DAWSON K. S., EISENSTEIN D. J. and ET AL., A&A, 574 (2015) A59.
- [20] BAUTISTA J. E., BUSCA N. G., GUY J., RICH J., BLOMQVIST M., DU MAS DES BOURBOUX H., PIERI M. M., FONT-RIBERA A., BAILEY S., DELUBAC T., KIRKBY D., LE GOFF J.-M., MARGALA D., SLOSAR A., VAZQUEZ J. A., BROWNSTEIN J. R., DAWSON K. S. and ET AL., A&A, 603 (2017) A12.
- [21] DE SAINTE AGATHE V., BALLAND C., DU MAS DES BOURBOUX H., BUSCA N. G., BLOMQVIST M., GUY J., RICH J., FONT-RIBERA A., PIERI M. M., BAUTISTA J. E., DAWSON K. and ET AL., A&A, 629 (2019) A85.
- [22] FONT-RIBERA A., KIRKBY D., BUSCA N., MIRALDA-ESCUDÉ J., ROSS N. P., SLOSAR A., RICH J., AUBOURG É., BAILEY S., BHARDWAJ V., BAUTISTA J., BEUT-LER F., BIZYAEV D., BLOMQVIST M., BREWINGTON H., BRINKMANN J., BROWNSTEIN J. R., CARITHERS B., DAWSON K. S., DELUBAC T., EBELKE G., EISENSTEIN D. J., GE J., KINEMUCHI K. and ET AL., JCAP, 2014 (2014) 027.
- [23] DU MAS DES BOURBOUX H., LE GOFF J.-M., BLOMQVIST M., BUSCA N. G., GUY J., RICH J., YÈCHE C., BAUTISTA J. E., BURTIN É., DAWSON K. S., EISENSTEIN D. J. and ET AL., A&A, 608 (2017) A130.
- [24] BLOMQVIST M., DU MAS DES BOURBOUX H., BUSCA N. G., DE SAINTE AGATHE V., RICH J., BALLAND C., BAUTISTA J. E., DAWSON K., FONT-RIBERA A., GUY J., LE GOFF J.-M. and ET AL., A&A, 629 (2019) A86.
- [25] RIESS A. G., FILIPPENKO A. V., CHALLIS P., CLOC-CHIATTI A., DIERCKS A., GARNAVICH P. M., GILLILAND R. L., HOGAN C. J., JHA S., KIRSHNER R. P., LEI-BUNDGUT B., PHILLIPS M. M., REISS D., SCHMIDT B. P., SCHOMMER R. A., SMITH R. C., SPYROMILIO J., STUBBS C., SUNTZEFF N. B. and TONRY J., Astronomical Journal, **116** (1998) 1009.
- [26] Schmidt B. P., Suntzeff N. B., Phillips M. M.,

SCHOMMER R. A., CLOCCHIATTI A., KIRSHNER R. P., GARNAVICH P., CHALLIS P., LEIBUNDGUT B., SPY-ROMILIO J., RIESS A. G., FILIPPENKO A. V., HAMUY M., SMITH R. C., HOGAN C., STUBBS C., DIERCKS A., REISS D., GILLILAND R., TONRY J., MAZA J., DRESSLER A., WALSH J. and CIARDULLO R., *ApJ*, **507** (1998) 46.

- [27] PERLMUTTER S., ALDERING G., GOLDHABER G., KNOP R. A., NUGENT P., CASTRO P. G., DEUSTUA S., FABBRO S., GOOBAR A., GROOM D. E., HOOK I. M., KIM A. G., KIM M. Y., LEE J. C., NUNES N. J., PAIN R., PENNY-PACKER C. R., QUIMBY R., LIDMAN C., ELLIS R. S., IRWIN M., MCMAHON R. G., RUIZ-LAPUENTE P., WAL-TON N., SCHAEFER B., BOYLE B. J., FILIPPENKO A. V., MATHESON T., FRUCHTER A. S., PANAGIA N., NEW-BERG H. J. M., COUCH W. J. and PROJECT T. S. C., ApJ, 517 (1999) 565.
- [28] MELIA F. and LÓPEZ-CORREDOIRA M., International Journal of Modern Physics D, 26 (2017) 1750055.
- [29] MELIA F., QIN J. and ZHANG T.-J., MNRAS, 499 (2020) L36.
- [30] MELIA F. and LÓPEZ-CORREDOIRA M., International Journal of Modern Physics D, 31 (2022) 2250065.
- [31] MELIA F. and SHEVCHUK A. S. H., MNRAS, 419 (2012) 2579.
- [32] MELIA F., The Cosmic Spacetime (Taylor and Francis, Oxford) 2020.
- [33] AHUMADA R., ALLENDE PRIETO C., ALMEIDA A., AN-DERS F., ANDERSON S. F., ANDREWS B. H., ANGUIANO B., ARCODIA R., ARMENGAUD E., AUBERT M. and ET AL., arXiv e-prints, (2019) arXiv:1912.02905.
- [34] DAWSON K. S., KNEIB J.-P., PERCIVAL W. J., ALAM S., ALBARETI F. D., ANDERSON S. F., ARMENGAUD E., AUBOURG É., BAILEY S., BAUTISTA J. E., BERLIND A. A., BERSHADY M. A., BEUTLER F., BIZYAEV D., BLANTON M. R., BLOMQVIST M., BOLTON A. S., BOVY J., BRANDT W. N., BRINKMANN J., BROWNSTEIN J. R., BURTIN E. and ET AL., AJ, **151** (2016) 44.
- [35] RIESS A. G., CASERTANO S., YUAN W., BOWERS J. B., MACRI L., ZINN J. C. and SCOLNIC D., *ApJ Letters*, 908 (2021) L6.
- [36] DU MAS DES BOURBOUX H., RICH J., FONT-RIBERA A., DE SAINTE AGATHE V., FARR J., ETOURNEAU T., LE GOFF J.-M., CUCEU A., BALLAND C., BAUTISTA J. E., BLOMQVIST M., BRINKMANN J., BROWNSTEIN J. R., CHABANIER S., CHAUSSIDON E., DAWSON K., GONZÁLEZ-MORALES A. X., GUY J., LYKE B. W., DE LA MACORRA A. and ET AL., *ApJ*, **901** (2020) 153.
- [37] VISHWAKARMA R. G., Physica Scripta, 87 (2013) 055901.
- [38] CHASHCHINA O. and SILAGADZE Z., Universe, 1 (2015) 307.
- [39] CARTER P., BEUTLER F., PERCIVAL W. J., DEROSE J., WECHSLER R. H. and ZHAO C., *MNRAS*, **494** (2020) 2076.
- [40] EVSLIN J., JCAP, 2017 (2017) 024.
- [41] DIENES Z., Journal of Mathematical Psychology, 75 (2016) 78.
- [42] KRUSCHKE J. K. and LIDDELL T. M., Psychonomic Bulletin & Review, 2017 (2017) 1.
- [43] JEFFREYS H., Theory of probability (Oxford University Press, Oxford, UK) 1961.
- [44] KASS R. E. and RAFTERY A. E., Journal of the American Statistical Association, 90 (1995) 773.

- [45] WETZELS R., MATZKE D., LEE M. D., ROUDER J., IVER-SON G. and WAGENMAKERS E. J., Perspectives on Psychological Science, 6 (2011) 291.
- [46] MELIA F., Pub Astron Soc Pacific, 134 (2022) 121001.
- [47] MELIA F., MNRAS, **481** (2018) 4855.
- $[48]\,$  Melia F., MNRAS ,  ${\bf 463}$  (2016) L61.
- [49] MELIA F., European Journal of Physics, 43 (2022) 035601.